

Physical Drivers and Parameter Sensitivities of Pearl
River-derived Sediment Dispersal on the Northern South China
Sea Shelf: A modeling study
~~Modeling dispersal of the Pearl River-derived sediment over the~~
~~Shelf~~

Guang Zhang^{1,2,3}, Suan Hu^{1,2,3}, Xiaolong Yu^{1,2,3}, Heng Zhang^{1,2,3}, Wenping Gong^{1,2,3*}

¹ School of Marine Sciences, Sun Yat-sen University, Zhuhai, 519082, China

² Guangdong Provincial Key Laboratory of Marine Resources and Coastal Engineering, Zhuhai, 519082, China

³ Pearl River Estuary Marine Ecosystem Research Station, Ministry of Education, Zhuhai 519082, China

Corresponding author: Wenping Gong (gongwp@mail.sysu.edu.cn)

Abstract

This study employs the Coupled-Ocean-Atmosphere-Wave-Sediment-Transport (COAWST) modeling system to quantitatively assess the seasonal suspension, transport, and annual fate of Pearl River-derived sediment (riverine slow-settling single fine grains and high-settling flocs) on the northern continental shelf of the South China Sea (SCS). Following careful model validation, a series of sensitivity experiments were conducted to investigate the effects of tides, waves, background circulation, sediment settling velocity, critical shear stress, and sediment spin-up durations. The results reveal strong seasonal variations in sediment dynamics driven by the East Asian monsoon. During the wet summer, weaker hydrodynamic conditions promote the initial deposition of riverine sediment via the surface buoyant plume. In contrast, stronger winds and waves during winter enhance sediment resuspension and southwestward transport, particularly toward the Beibu Gulf. Spatially, approximately two-thirds of the annual Pearl River-derived sediment load is retained near the estuary. About 9% reaches the continental shelf east of the PRE, while similar proportions accumulate in the Beibu Gulf and south of Hainan Island. Sensitivity experiments highlight the distinct and significant roles of different physical forcings in sediment dispersal. ~~tidal~~ Tidal dynamics strongly enhance ~~influence~~ sediment mobilization and transport behavior ~~within the estuary by~~ increasing, ~~where the absence of tidal forcing reduces~~ bottom shear stress, which promotes ~~leading to increased local deposition and reduced~~ offshore sediment export transport and limits local deposition. Wave forcing plays a dominant role in sediment

resuspension near the river mouth and along the coast, especially during winter, facilitating sediment redistribution across the shelf. ~~Excluding waves leads to greater sediment retention near the estuary and diminished transport toward distant regions.~~ The remotely forced (large-scale, non-local forcing) ambient shelf ~~Ambient~~ circulation, ~~particularly in summer,~~ drives ~~is essential for~~ eastward sediment transport, enabling sediment to spread widely across the shelf, ~~when it is omitted, northeastward dispersal is greatly diminished.~~ Model outcomes are also sensitive to sediment parameterization. The natural seasonal increase in critical shear stress for erosion during winter counteracts part of the wave-enhanced resuspension capacity, thereby reducing resuspension and erosion on the continental shelf east of ~~Using non-seasonal critical shear stress for erosion increases wintertime sediment mobility east of~~ the Leizhou Peninsula. Higher settling velocities decrease suspended sediment concentrations and promote near-source retention, limiting long-distance transport. Spin-up duration experiments indicate that Pearl River-derived sediment, which enters and accumulates in various regions of the model domain during the first year, continues to migrate southwestward in the second year under the influence of the mean annual flow field. In contrast, the spin-up duration of seabed sediment has little impact on the retentions of Pearl River-derived sediment on the shelf. Overall, this study reveals the transport pathway and fate of the Pearl River-derived sediment and provides a model-based assessment of its seasonal behavior and the sensitivity of suspended sediment dispersal to physical drivers and sediment parameters or conditions, ~~dispersal mechanisms~~ on the northern SCS shelf. It identifies key physical

drivers regulating sediment transport and deposition patterns, offering new insight into sediment fate in a monsoon-dominated shelf system.

Keywords

Riverine sediment transport; Sediment retention; Numerical modeling; Pearl River Estuary

1. Introduction

The transport process of suspended sediment from river source to ocean sink is an important link in the global material cycle ([Geyer et al., 2004](#); [McKee et al., 2004](#); [Kuehl et al., 2016](#); [Liu et al., 2016](#); [Cao et al., 2019](#)). Much of the riverine sediment is trapped on the shallow shoals in estuaries, while the rest is transported by buoyant plume out of the estuary ([Meade, 1969](#); [Burchard et al., 2018](#); [Zhang et al., 2019](#)). The riverine sediment carried by the buoyant plume has a significant impact on the water quality, ecology, and geomorphology of the estuaries and continental shelves ([Wright and Coleman, 1973](#); [Turner and Millward, 2002](#)).

The transport and deposition of riverine sediments from river source to estuarine, coastal, and shelf environments are controlled by diverse physical processes, including tidal forces, wave action, and shelf circulation dynamics ([Dalyander et al., 2013](#); [Gao and Collins, 2014](#); [Xu et al., 2016](#); [Warner et al., 2017](#); [Zang et al., 2019](#); [Wang et al., 2020](#)). Tides play a critical role in sediment transport dynamics in estuarine and shelf regions, as spring tides typically produce higher bed shear stress, enhanced sediment resuspension, and greater offshore sediment transport flux compared to neap tides ([Bever and MacWilliams, 2013](#); [Zhang et al., 2019](#); [Wang et](#)

79 [al., 2020](#)). In nearshore regions, wave-induced bed shear stress often exceeds
80 current-induced stress by an order of magnitude ([Xue et al., 2012](#); [Dalyander et al.,](#)
81 [2013](#)). Furthermore, wave-driven sediment resuspension frequently surpasses, and is
82 often several times greater than, the peak levels achieved by current-induced
83 resuspension ([Sanford, 1994](#); [Harris et al., 2008](#); [Brand et al., 2010](#); [Xu et al., 2016](#)).
84 In shelf regions, circulation patterns significantly modulate sediment transport, with
85 the magnitude of along-shelf transport substantially exceeding the cross-shelf
86 component in most areas ([Nittrouer and Wright, 1994](#); [Geyer et al., 2004](#); [Gao and](#)
87 [Collins, 2014](#); [Wang et al., 2020](#)).

88 Furthermore, sediment properties, including settling velocity ([Xia et al., 2004](#);
89 [Chen et al., 2010](#); [Cheng et al., 2013](#)), critical shear stress for erosion ([Dong et al.,](#)
90 [2020](#)), and bed grain size distribution ([Xue et al., 2012](#); [Bever and MacWilliams,](#)
91 [2013](#)), significantly influence sediment transport dynamics and
92 deposition/resuspension processes. Settling velocity can influence the location of
93 sediment depocenters, with higher settling velocities leading to more proximal
94 entrapment and vice versa ([Ralston and Geyer, 2017](#)). Similarly, critical shear stress
95 for erosion can affect the resuspension of deposited sediment, with higher critical
96 shear stress resulting in less resuspension and more deposition especially during neap
97 tides and weak wind wave periods ([Dong et al., 2020](#); [Choi et al., 2023](#)).

98 A comprehensive understanding of sediment transport and deposition from river
99 source to ocean sink requires the integrated consideration of both physical forcing
100 factors and inherent sediment characteristics. Here, we present the transport and

deposition of the Pearl River-derived sediments on the continental shelf as a case study. The Pearl River, ranking as China's second-largest river in terms of freshwater discharge (Hu et al., 2011), forms the Pearl River Estuary (PRE) in its lower reaches (Figures 1 and S1). Its freshwater and sediment discharge are primarily delivered through eight major outlets (Figure S1b; Wu et al., 2016; Zhang et al., 2019; Zhang et al., 2025), forming distinct buoyant plumes that extend across the northern South China Sea (SCS) shelf (Zhang et al., 2025). The present average annual (2001-2022) freshwater and riverine sediment loads are $2.74 \times 10^{11} \text{ m}^3$ and 2.84×10^7 tons, respectively, ~~as reported by the Ministry of Water Resources of the People's Republic of China~~ (Ministry of Water Resources of the PRC, ~~2022~~<http://www.mwr.gov.cn/sj/#tjgb>). The distribution of these inputs shows significant seasonal variability: approximately 80% of the freshwater and 95% of the sediment load are transported during the wet summer season (April to September), while the remaining portion is discharged during the dry winter season (Xia et al., 2004).

The northern SCS, ~~under the influence of~~~~shaped by~~ the East Asian Monsoon, displays marked seasonal contrasts, featuring winter monsoon winds averaging 7-10 m s^{-1} and summer winds typically below 6 m s^{-1} (Su, 2004; Ou et al., 2009). This seasonal shift drives coastal currents: northeastward in summer and southwestward in winter (Gan et al., 2009; Gan et al., 2013). Beyond the coastal zone, the consistent SCS Warm Current (~~SCSWC~~) flows northeastward along the shelf break and inner continental slope toward the Taiwan Strait, originating near Hainan Island and

persisting year-round, even during the winter northeast monsoon, across a remarkable distance of 600-700 km to the southern tip of the Taiwan Strait (Su, 2004; Yang et al., 2008).

The PRE is situated in the central part of the northern South China Sea boundary, positioned between the Taiwan Banks and Hainan Island. The PRE has a micro-tidal and mixed semi-diurnal regime, with daily tidal inequality ~~in the range and in the time between the high and low tides~~ (Mao et al., 2004). The ~~neap and spring tides alternately influence the water elevation downstream of the estuary, with~~ tidal ranges vary~~ing~~ from approximately 0.7 m during neap tides to over 2 m during spring tides (Chen et al., 2016; Gong et al., 2018b). ~~The PRE and the nearby shelf exhibit strong seasonal variation in w~~Water column ~~stratification is strong~~stability and are highly stratified during the wet summer~~season, but weakens in the dry winter when~~ while the PRE becomes partially mixed or vertically well-mixed~~during the dry winter season~~ (Dong et al., 2004). Offshore ~~of the PRE region,~~ wave conditions are seasonally variable, being ~~display distinct seasonal patterns: the waves are~~ mild during~~in~~ summer, and~~become~~ stronger during~~in~~ winter, ~~dominated~~marked by larger southeasterly waves (Gong et al., 2018a; Gong et al., 2018b; Zhang et al., 2021).

Previous studies have focused on sediment transport within the PRE (Zhang et al., 2019; Zhang et al., 2021; Ma et al., 2024). Most Pearl River-derived sediments are deposited within the estuary, and neglecting tidal effects can lead to higher deposition rates and lower offshore sediment flux when compared to those with tides (Hu et al., 2011). Pearl River-derived sediment behavior ~~The depositional dynamics of sediments~~

~~from different PRE outlets are~~ is regulated by outlet location, topography, and tidal
~~phase conditions~~, with neap tides favoring sediment accumulation on shoals and spring
tides driving erosion and enhancing erosion and export~~offshore sediment transport~~
([Zhang et al., 2019](#)). Waves further intensify both lateral trapping within the PRE and
offshore sediment transport ([Liu and Cai, 2019](#); [Zhang et al., 2021](#)).

However, numerical studies on the transport of the Pearl River-derived sediments
across the continental shelf remain scarce, even amidst the widespread adoption of
computer modeling approaches. Previous research ~~on the distribution of these~~
~~sediments~~ has primarily relied on analyses of seismic profiles, gravity cores, and
laboratory-based radiometric dating of sediment samples ([Ge et al., 2014](#); [Liu et al.,](#)
[2014](#); [Cao et al., 2019](#); [Lin et al., 2020](#); [Chen et al., 2023](#)). Outside the PRE, ~~gravity~~
~~core and seismic surveys~~such data ~~were used to examine the Holocene sedimentary~~
~~processes~~, revealing two Holocene ~~distinct~~ mud depo-centers: an eastward proximal
depo-center extending southeastward from the PRE's mouth and a southwestward
distal mud belt ([Ge et al., 2014](#); [Liu et al., 2014](#); [Chen et al., 2023](#)). However, seismic
and drilling data cannot confirm transport of ~~that the~~ Pearl River sediments can be
~~transported~~ to the Beibu Gulf ([Ge et al., 2014](#)), and limited sampling makes it. ~~Due to~~
~~the lack of sufficient gravity core samples and seismic data, it is~~ difficult to quantify
their contribution~~quantitatively attribute the sediment in the Beibu Gulf to the Pearl~~
~~River-derived sediment~~ ([Cao et al., 2019](#)). Using radionuclide endmember
models ~~Afterward~~, [Lin et al. \(2020\)](#) estimated that ~15% of nearshore surface
sediments in the Beibu Gulf originate from the PRE, but their study did not address

transport pathways, fluxes, or annual deposition. ~~used the ^{226}Ra — ^{238}U and ^{232}Th — ^{238}U~~
~~endmembers model based on measurements of radionuclides in the surface sediment~~
~~samples. They found that approximately 15% of the surface sediment in the nearshore~~
~~area of the Beibu Gulf originates from the PRE region. However, their studies only~~
~~address the proportion of the PRE sediment in the surface sediment of the Beibu Gulf,~~
~~without directly indicating the seasonal transport pathways, flux, and annual~~
~~deposition of the sediment from the Pearl River.~~

A gap persists in understanding how physical drivers ~~processes~~ (such as tides,
waves, and remotely forced (large-scale, non-local forcing) ambient shelf circulations)
and sediment characteristics (such as critical shear stress for erosion, settling velocity)
and sediment initial conditions influence the seasonal suspension, transport, and
annual deposition of the Pearl River-derived sediment on the shelf. ~~Specifically, this~~
~~study focuses on sediment classes 4 and 5 in Table 1, which represent the Pearl~~
~~River derived components: slow settling single fine grains (Class 4) and fast settling~~
~~flocs (Class 5), in contrast to the background seabed sediments represented by classes~~
~~1 to 3.~~ To address this, we utilize numerical modeling calibrated and validated with
field observations and seabed grain size data, ~~complemented by extensive collection~~
~~of field observations and seabed grain size distribution data for model calibration and~~
~~validation—a highly effective approach for exploring mechanisms and testing~~
~~hypotheses derived from limited observational datasets.~~ This is an effective approach
for investigating processes and testing hypotheses where observations are limited.
This study ~~aims to~~ systematically examines ~~investigate~~ the dispersal dynamics of ~~the~~

Pearl River-derived sediment ~~over~~on the northern South China Sea shelf,
focusing~~with particular emphasis~~ on the following objectives:

(1) Quantify the seasonal dispersal and annual deposition of ~~the~~ Pearl
River-derived sediment (~~classes 4 and 5 in Table 1~~) on ~~over~~ the continental shelf.

(2) Examine the relative roles of physical forcings (tides, waves, and ambient
circulations), sediment characteristics (critical shear stress for erosion, settling
velocity) and (Pearl River-derived versus Seabed) sediment spin-up durations on its
~~the dispersal of the Pearl River-derived sediment.~~




Figure 1. Bathymetry (shading) and isobath contours of the study area, with the
ROMS/SWAN model grid domain outlined by black-to-white dashed lines. Circled
numbers ① - ⑧ denote the eight regions: "Proximal", "Southern", "Eastern",
"Southeastern", "Western", "Southwestern", "Gulf", and "Distal" regions, as defined

by transects and detailed in Section 2.1. The abbreviations TWI and TWS mean Taiwan Island and Taiwan Strait, respectively. The gray contours represent 30-180 m isobaths at 30 m intervals, a consistent feature maintained in all subsequent figures that include these isobath contours.

2. Methods

2.1 Model coupling

This study employed the Coupled Ocean Atmosphere Wave Sediment Transport (COAWST, version 3.4) modeling system ([Warner et al., 2005](#); [Warner et al., 2008](#); [Warner et al., 2010](#)), which includes a Model Coupling Toolkit (MCT) to facilitate data exchange among different modules ([Jacob et al., 2005](#); [Larson et al., 2005](#)). The COAWST system consists of several modeling components, mainly comprises a hydrodynamic module (Regional Ocean Modeling System; ROMS) ([Shchepetkin and McWilliams, 2005](#); [Haidvogel et al., 2008](#)), an atmospheric module (Advanced Research Weather Research and Forecasting; WRF) ([Skamarock et al., 2005](#)), a wave module (Simulating Waves Nearshore; SWAN) ([Booij et al., 1999](#)), and a sediment transport module (Community Sediment Transport Modeling System; CSTM) ([Warner et al., 2008](#)).

In this study, we established a coupling between ROMS, SWAN, and CSTM. The model grid covered the northern continental shelf of the South China Sea, including the PRE (Figure 1). The regional model was configured with 170×482 horizontal grid cells, with horizontal resolution varying from approximately 0.1 km near the PRE

to about 10 km at outer open boundaries ([Hu et al., 2024](#); [Zhang et al., 2025](#)). The model grid bathymetry data was obtained from nautical charts compiled by the China Maritime Safety Administration and the General Bathymetric Chart of the Oceans (GEBCO) ([Weatherall et al., 2015](#)). The vertical grid used a terrain-following S-coordinate system ([Song and Haidvogel, 1994](#)) with 20 layers and a stretching transformation for higher resolution near the surface and bottom. For model validations, please refer to the Supplementary Material (Supplement Figures S1-S10).

To improve the understanding of the spatial-temporal variabilities in the riverine sediment dispersal, and the estimation of the fate of the Pearl River sediment during the wet summer season, dry winter season, and throughout the year, we partitioned the model domain into eight distinct regions delineated by various transects as illustrated in Figure 1. The division criteria are mainly based on the distance from the estuary and the natural separation by the Leizhou Peninsula and Hainan Island (Figure 1). These regions include:

- ① Proximal region: Proximity to the estuary,
- ② Southern region: Located deeper in the southern part of the estuary,
- ③ Eastern region: Eastern side of the estuary, closer to the shoreline,
- ④ Southeastern region: Further offshore on the eastern side of the estuary,
- ⑤ Western region: Western side of the estuary, closer to the shoreline,
- ⑥ Southwestern region: Offshore on the western side of the estuary,
- ⑦ Gulf region: Mainly the Beibu Gulf,
- ⑧ Distal region: South of the Hainan Island.

By dividing the model domain into these delineated regions, we calculated the riverine sediment flux for each transect, thereby determining the total riverine sediment volume retained in each region.

2.2 ROMS model setup

For the ROMS model, we utilized the Generic Length Scale turbulence closure scheme ([Warner et al., 2005](#)) for vertical turbulence parameterization. The method of [Smagorinsky \(1963\)](#) was employed to calculate the horizontal eddy viscosity and diffusivity. The Flather and Chapman boundary conditions were applied to barotropic current and water elevation at open boundaries, respectively ([Flather, 1976](#); [Chapman, 1985](#)). Meanwhile, the open-boundary conditions for temperature, salinity, and sediment concentration were imposed by radiation methods ([Orlanski, 1976](#); [Raymond and Kuo, 1984](#)). Surface forcing (including wind, net shortwave radiation, air temperature, atmospheric pressure, specific/relative humidity, and rain, etc.) data were sourced from the Climate Forecast System Reanalysis of the National Centers for Environmental Prediction (NCEP) ([Saha et al., 2014](#)), with a temporal resolution of 1 h and a spatial resolution of $0.3^{\circ} \times 0.3^{\circ}$. Water level and current [velocity](#) open-boundary conditions comprised two components: tidal and subtidal. The tidal component was obtained from the Oregon State University Tidal Prediction Software database ([Egbert and Erofeeva, 2002](#)), while the subtidal component was interpolated from the HYbrid Coordinate Ocean Model (HYCOM) outputs ([Chassignet et al., 2007](#)).

2.3 Wave model setup

The SWAN model was executed and coupled to the same grid as the ROMS model ([Warner et al., 2010](#)). It was driven by surface atmospheric forces, real-time water level, and current fields from the ROMS and boundary reanalysis data. Wave boundary conditions were specified using nonstationary wave parameters from outputs of the NOAA WAVEWATCH III global ocean wave model solutions ([Tolman et al., 2016](#)). Information was exchanged at 15-minute intervals to introduce wave-current interaction (WCI) between the ROMS and SWAN models ([McWilliams et al., 2004](#); [Kumar et al., 2012](#)). This exchange included significant wave height (H_{sig}), surface peak wave period, mean wave direction and length, wave energy dissipation, and the percentage of breaking waves from SWAN to ROMS, as well as water level and current from ROMS to SWAN.

Additionally, the wave-current bottom boundary module based on [Madsen \(1994\)](#), was activated to simulate the wave-current bottom boundary layer. The vortex force module of wave forces was also activated to compute the wave-induced momentum flux, utilizing the method proposed by [McWilliams et al. \(2004\)](#) and implemented in COAWST by [Kumar et al. \(2012\)](#). The bottom friction was computed based on a logarithmic velocity profile ([Warner et al., 2008](#)).

2.4 Specifications of riverine input and sediment model

The freshwater discharge for the Pearl River was specified at the northern boundary using daily measured data from the Pearl River Water Resources Commission, while downstream precipitation within the Pearl River Basin was neglected. The full simulation model was initialized on the first day of January 2016


using temperature, salinity, and current fields interpolated from the HYCOM model, and it concluded on March 31, 2018. This study primarily analyzes the last 12 months, specifically from April 1, 2017, to March 31, 2018. This year was selected because the freshwater discharge and sediment load of the Pearl River closely approximated the average values of the past two decades, with a runoff of $3.35 \times 10^{11} \text{ m}^3$ and a sediment load of 3.45×10^7 tons, closely resembling the averages from 2001 to 2022.

Since the daily riverine sediment loads were unavailable, we modified the previous research results on sediment rating curves ([Zhang et al., 2012](#)) to suit for our study, as expressed by

$$C_s = 0.00002263Q^{1.792} \quad (1)$$

where C_s is the Pearl River-derived suspended sediment concentration (mg L^{-1}), Q is the Pearl River freshwater discharge rate ($\text{m}^3 \text{ s}^{-1}$). Based on this relationship, the total amount of Pearl River sediment input over our 12-months study period (Figure 3b) was 34.52 million tons, aligning closely with the annual load reported in 2017 by the Pearl River Water Resources Commission. The riverine sediment input, derived from the river discharge, was allocated across the eight outlets along the north boundary (Figure S1b) based on the distribution approach of [Hu et al. \(2011\)](#). The subsequent step involved establishing the proportion of seabed sediment particle size components. Sediments are typically categorized into three grain-size classes: clay (0-4 μm), silt (4-63 μm), and sand (63-2000 μm), as outlined by [Shepard \(1954\)](#). Data on sediment particle size composition for the northern continental shelf of the South China Sea and the PRE area were acquired through multiple voyage observations

(Zhang et al., 2013; Zhang et al., 2019). Furthermore, publicly available data from published literature were compiled (Gao et al., 2007; Kirby et al., 2008; Gao et al., 2010; Huang et al., 2013; Liu et al., 2014; Wang et al., 2014; Wang et al., 2015; Wang et al., 2016; Ge et al., 2017; Lu et al., 2017; Zhong et al., 2017; Yang et al., 2018; Ge et al., 2019). Finally, component distribution data for different particle size classes of seabed sediment were obtained from a total of 1,981 measured stations (Figure 2a-c).






Figure 2. Row 1 presents the spatial distribution patterns of seabed sediment fractions derived from 1,981 sampling sites, while Row 2 demonstrates the initial spatial distribution prototype of seabed sediment fractions developed based on the observational data presented in Row 1. Row 3 shows the spatial distribution patterns of seabed sediment fractions following the completion of spin-up phase in the Control run case on April 1st, 2017, with Columns 1, 2, and 3 representing the fractions of clay, silt, and sand, respectively.

As illustrated in Figure 2a-c, the measured stations exhibit a widespread distribution, offering comprehensive coverage of the entire northern continental shelf of the South China Sea, including the PRE. Particularly dense distribution is observed in the PRE and the coastal areas of western Guangdong. These regions represent the primary scope of transport and deposition associated with the Pearl River-derived

sediment. Hence, the stations utilized in this study well represent the distribution of bed sediment particle size components in these study areas. It is evident that among the stations in the offshore area of the northern continental shelf of the South China Sea, silt dominates, followed by clay, while sand with the largest particle size is the least abundant. This suggests a significant presence of terrestrial sediment or Pearl River sediment in the offshore area of the northern continental shelf of the South China Sea. It should be noted that the lack of in situ grain size distribution data in specific regions of the model domain, especially in the Beibu Gulf area, may lead to uncertainties in sediment transport predictions. We will address these limitations and quantify their potential errors in the discussion part of this study.

To derive the component proportions of the initial prototype field on the model grid, this study employed the Kriging method ([Krige, 1951](#)), widely recognized for spatially interpolating various types of observational data. The sediment distribution pattern obtained through interpolation (Figure 2d-f) closely resembles the original 1,981 measured sediment particle size distribution patterns (Figure 2a-c), suggesting the suitability of this interpolation method for the study area.

The initial prototype field underwent a 15-month spin-up period (from January 1, 2016, to March 31, 2017), during which the bottom sediment composition evolved [in response to](#) ~~under realistic~~ hydrodynamic forcings [through](#) ~~from~~ the [coupled](#) ROMS, SWAN, and CSTM models. This method has been utilized in numerous previous studies, including those by [Bever et al. \(2009\)](#), [van der Wegen et al. \(2010\)](#), and [Zhang et al. \(2021\)](#). This process allows the initially [estimated](#) ~~idealized~~ sediment

distribution to evolve under ~~realistic dynamic forcings, including~~ tides, waves, and currents, thereby minimizing unreasonable spatial patterns introduced by the Kriging, ~~sparse or problematic data interpolation method~~. Such unreasonable spatial patterns may arise due to limitations in the number, representativeness, and timing of field sediment samples relative to the model start date. As a result, the sediment field after the spin-up period (Figures 2g–i) is thought to exhibit spatial patterns that are ~~more physically plausible and~~ better aligned with the hydrodynamic conditions of the study region. During both the 15-month spin-up period and the subsequent 12-month formal model experiments (see Section 2.6 and Table 2), the CSTM utilized five sediment classes (Table 1), representing a range of sediment sizes and characteristics. These included three types of seabed sediments (clay, silt, and sand, corresponding to sediment Classes 1 to 3 in Table 1) and two types of Pearl River-derived sediments (Class 4 and Class 5 in Table 1). The riverine sediments consisted of slow-settling single fine grains (Class 4) and high-settling flocs (Class 5), which were delivered into the model domain during both the 15-month spin-up period and the subsequent 12-month formal model experiments. The riverine flocs correspond to the flocculated fractions of clay and silt, whereas the single fine grains represent the non-flocculated components within the Pearl River-derived sediments, following the setting of [Bever and MacWilliams \(2013\)](#). To clarify, at the start of the 12-month formal model experiments, the retained Pearl River-derived sediments (Classes 4–5 in Table 1) that entered the model during the 15-month spin-up period were added ~~as to~~ Class 1 and Class 2, respectively, to avoid contaminating the data analysis of the formal

experiments. This approach allows for a better distinction between Pearl River sediment and seabed sediment, enabling separate analysis of the suspension, transport, and deposition of Pearl River-derived sediment (Harris et al., 2008; Zhang et al., 2019). Specifically, the fractions of the two types of Pearl River-derived sediments were set at 40% and 60%, respectively, following Zhang et al. (2019) and Zhang et al. (2021). The parameters for all five sediment classes are summarized in Table 1. Sediment density, porosity, and erosion rate for all sediment classes were set to 2650 kg m⁻³, 0.672 (Zhang et al., 2019; Zhang et al., 2021), and 1×10^{-4} kg m⁻² s⁻¹ (Ralston et al., 2012), respectively. Settling velocities (w_s) were guided by the ranges reported by Xia et al. (2004) and Warner et al. (2017), and further refined through extensive sensitivity testing and model calibration to optimally reproduce observed suspended sediment concentration (SSC). ~~The settling velocity (w_s),~~ Critical shear stresses for erosion (τ_{ce}), and other parameters were set following previous studies or were based on model calibration (Ralston et al., 2012; Warner et al., 2017; Zhang et al., 2019; Dong et al., 2020; Zhang et al., 2021; Cao et al., 2025).

Our model configuration incorporates seasonal variations in τ_{ce} , supported by multiple lines of evidence from field observations, laboratory experiments, and numerical analyses (Dong et al., 2020; Cao et al., 2025). Previous studies have established a distinct seasonal pattern in the PRE, with winter τ_{ce} values significantly exceeding those in summer. Dong et al. (2020)'s laboratory experiments using the UMCES-Gust Erosion Microcosm System (U-GEMS) on 2017-winter sediment samples yielded a τ_{ce} of 0.26 Pa, which effectively reproduced observed ~~suspended~~

~~sediment concentration (SSC)~~ in winter simulations. However, this value proved excessive for summer conditions, when a τ_{ce} of 0.15 Pa provided better agreement with field observations in summer simulations, indicating a winter-to-summer τ_{ce} ratio of 1.73. Recent 2020-summer in situ measurements by [Cao et al. \(2025\)](#) using a benthic quadrapod-mounted 3D Profiling Sonar revealed a two-layer erosion threshold system: a surface "fluffy layer" with $\tau_{ce} = 0.06$ Pa overlying a consolidated seabed with $\tau_{ce} = 0.13$ Pa. The latter value aligns with [Dong et al. \(2020\)](#)'s summer calibration, suggesting that [Dong et al. \(2020\)](#)'s laboratory measurements, potentially affected by sediment consolidation during sample transport, might have missed the lower τ_{ce} of the surface fluffy layer. Based on these consistent findings, we implemented a seasonal τ_{ce} adjustment factor of 1.73 (winter/summer) in our model configuration (Table 1).

Table 1. CSTM model Sediment Properties

Source	Seabed			Pearl River	
Class	1	2	3	4	5
Sediment Type	Clay	Silt	Sand	Single grains	Flocs
w_s (mm s ⁻¹)	0.02 ^c	1.2 ^e	57 ^d	0.005 ^c	0.6 ^e
Summer τ_{ce} (Pa)	0.14 ^e	0.03	0.27 ^d	0.15 ^{abef}	0.05 ^{abe}
Winter τ_{ce} (Pa)	0.24 ^f	0.05 ^f	0.47 ^{df}	0.26 ^{abf}	0.09 ^{abf}
Fraction	Spatially variable, see Figure 2g-i			40% ^{ab}	60% ^{ab}

^a[Zhang et al. \(2019\)](#), ^b[Zhang et al. \(2021\)](#), ^cCalibrated, ^d[Warner et al. \(2017\)](#), ^e[Cao et al. \(2025\)](#), and

^f([Dong et al., 2020](#)).


2.5 Wet and dry season regimes

The study area exhibits pronounced seasonal variability, which can be distinctly categorized into two primary seasons ([Dong et al., 2004](#); [Su, 2004](#); [Liu et al., 2014](#); [Zhang et al., 2021](#)). This seasonal classification is supported by multiple environmental parameters, including river freshwater discharge ([Figure 3a](#)), riverine sediment load ([Figure 3a](#)), wind patterns ([Figure 3b](#)), air temperature ([Figure 3c](#)), and modeled wave conditions ([Figure 3d-f](#)) at a representative site (21.5°N, 114°E; corresponding to station W1 in [Figure S1a](#), located immediately south of the PRE). [All values presented in Figure 3 correspond to the period from April 1, 2017, to March 31, 2018, consistent with the 1-year model simulation. The daily measured freshwater discharge for the Pearl River was obtained from the Pearl River Water Resources Commission, while the riverine sediment load was estimated using the method described in Section 2.4.](#) The meteorological data for wind and air temperature were obtained from the NCEP reanalysis dataset, while wave parameters were derived from numerical model simulations. These comprehensive indicators collectively characterize the distinct seasonal patterns observed in the study area ([Figure 3](#)). The entire year (from April 1, 2017, to March 31, 2018) is typically divided into two main seasons: wet summer (from April 1, 2017, to September 30, 2017) and dry winter (from October 1, 2017, to March 31, 2018).

During the wet summer season, freshwater discharge tends to be notably high,

often exceeding $10,000 \text{ m}^3 \text{ s}^{-1}$ and reaching a maximum of $53,000 \text{ m}^3 \text{ s}^{-1}$, with an average value of $15,266 \text{ m}^3 \text{ s}^{-1}$. This discharge constitutes a significant portion of the entire year, accounting for 72.06% of the annual total. During this period, the river carries a substantial sediment load of 32.85 megatons, constituting 95.17% of the total annual sediment transport. Prevailing winds predominantly blow from the south. For example, Figure 3b depicts the average monthly wind vector direction during the summer months as northward, with weak southeasterly winds in April, May, and September, and moderate southeasterly winds in July. June and August experience moderate southwesterly winds. The 2 m height air temperatures typically range between 20°C and 30°C . The daily average Hsig remains relatively low, with the monthly average Hsig less than 1 m. The wave propagation direction is generally consistent with the wind direction, being easterly in April and May, and southerly from June to September.

In stark contrast, the dry winter season demonstrates markedly lower runoff, typically falling below $10,000 \text{ m}^3 \text{ s}^{-1}$, with an average value of $5,953 \text{ m}^3 \text{ s}^{-1}$. The sediment load during this period is significantly reduced to merely 1.67 megatons, marking a substantial decrease compared to the wet summer season. Prevailing winds during the dry winter are predominantly northeasterly, with relatively high wind speeds. Except for moderate wind intensity in March, the monthly average wind speed in other months exceeds 5 m s^{-1} . The 2 m height air temperatures typically range between 10°C and 25°C during this season. The wave propagation direction aligns with the prevailing northeasterly winds of the season, predominantly northeasterly.



461

462 **Figure 3.** Time series of (a) the daily Pearl River freshwater discharge and sediment
 463 load, (b) daily (black vectors) and monthly (red vectors) averaged 10-meter height
 464 winds, (c) hourly 2-meter height air temperature, daily (black) and monthly (red)
 465 averaged (d) significant wave height (H_{sig} , lines) and (f) wave propagation direction
 466 (vectors) weighted by H_{sig}^2 . Two distinct seasons are delineated by the dashed
 467 magenta line.

468

469 2.6 Model experiments

470 To assess the relative importance of tides, waves, ambient shelf currents and
 471 residual water levels, seasonal variation in critical shear stress for erosion, the settling

velocity, and the spin-up duration of Pearl River-derived sediment (Classes 4-5 in Table 1) in the transport and dispersal of Pearl River-derived sediments, we conducted seven simulation experiments (Table 2). In all experiments, we implemented the Charnock approach within COAWST's bulk air-sea flux parameterization scheme to calculate surface wind stress using the NCEP 10-m wind product ([Charnock, 1955](#); [Fairall et al., 1996](#)), ensuring consistency in wind stress forcing across all simulations.

Exp 1 (the **Control** run) incorporated all the aforementioned forcing agents (including winds) and accounted for the seasonal variation in critical shear stress for erosion, with the winter critical shear stress for erosion set to be 1.73 times of that in summer. Exp 2 (**NTS** hereafter) was identical to Exp 1 but excluded tides, while Exp 3 (**NWS** hereafter) excluded waves. In Exp 4 (**NAS** hereafter), waves, tides, and the seasonal variation in critical shear stress for erosion were included, but the [remotely forced \(large-scale, non-local forcing\) ambient](#) shelf current and residual [\(non-tidal\)](#) water levels were omitted (i.e., no subtidal circulation forcing at open boundaries) to examine the influence of the South China Sea circulation. Exp 5 (**NVS** hereafter) replicated the setup of Experiment 1, but with one modification: it used a constant critical shear stress for erosion (τ_{ce}) across both seasons, specifically adopting the summer τ_{ce} value from Table 1 throughout the simulation (i.e., no seasonal adjustment between winter and summer). Exp 6 (**DSV** hereafter) was identical to Exp 1, except that it set a double sediment settling velocity of the Exp 1. Finally, to assess the model's sensitivity to the spin-up duration of Pearl River-derived sediment, particularly regarding the retention of riverine sediments in both the water column and

the seabed, we adopted the sediment distributions (Classes 1 to 5) from the [final state of the](#) Control run on March 31, 2018, as the alternative initial conditions for the Cycle experiment (designated as Exp 7, **Cycle** hereafter). This setup carries over the full year's evolution of riverine sediment transport and deposition from the Control run (Exp 1), including changes in all sediment classes, into the start of Exp 7. As a result, Exp 7 mainly evaluates how the presence of previously deposited riverine sediments influences subsequent sediment transport estimates.

Table 2. Experiment Settings

Experiments	Tides	Waves	Ambients	τ_{ce}	w_s	Re-run
Exp 1 (Control)	✓	✓	✓	Variable	Original	✗
Exp 2 (NTS)	✗	✓	✓	Variable	Original	✗
Exp 3 (NWS)	✓	✗	✓	Variable	Original	✗
Exp 4 (NAS)	✓	✓	✗	Variable	Original	✗
Exp 5 (NVS)	✓	✓	✓	Constant	Original	✗
Exp 6 (DSV)	✓	✓	✓	Variable	Double	✗
Exp 7 (Cycle)	✓	✓	✓	Variable	Original	✓

The term 'Ambients' denotes [remotely forced \(large-scale, non-local forcing\)](#) ambient shelf currents and residual [\(non-tidal\)](#) water levels. Variable indicates simulations employing seasonally varying τ_{ce} values (from Table 1), while 'Constant' refers to runs using exclusively the summer τ_{ce} value throughout the entire experiment. 'Original'

designates cases utilizing the settling velocities specified in Table 1, whereas 'Double' indicates simulations with these values doubled."

3 Results

3.1 Seasonal hydrodynamics and transport patterns of the Pearl

River-derived sediment

We quantified the spatial distributions of seasonal mean wind stress, H_{sig} , wave bottom orbital velocity (WBOV), and bottom shear stress for both the wet summer and dry winter periods (as defined in Section 2.5). These distributions serve as representative hydrodynamic conditions for typical summer and winter scenarios, respectively (Figure 4).

During summer, the prevailing winds predominantly originate from the south, with the average wind stress generally below 0.03 Pa, except in the eastern coastal waters of Hainan Island, where localized values reach up to 0.05 Pa (Figure 4a). In contrast, during the dry winter season, the prevailing winds shift to a northeasterly direction, resulting in generally higher average wind stress compared to summer (Figure 4b), with values typically exceeding 0.1 Pa in areas deeper than 40 m and surpassing 0.2 Pa in the offshore eastern Guangdong Coast near the Taiwan Bank (see Figure 1).

Corresponding to the seasonal wind stress (Figures 4a-b), the seasonally averaged wave characteristics in the PRE and the adjacent northern continental shelf of the South China Sea exhibit significant seasonal variations (Figures 4c-d).

During the wet summer season, the Hsig in the studied area is relatively low, with waves predominantly coming from the southeast (Figure 4c). The seasonal average Hsig across the entire shelf remains below 1 m, with areas deeper than 60 m showing Hsig values above 0.8 m, while in shallower nearshore regions (water depth < 20 m), Hsig is less than 0.6 m (Figure 4c). Corresponding to the lower Hsig in the wet summer, the seasonally-averaged WBOV is also relatively small, generally less than 1 cm s⁻¹ in areas deeper than 40 m, except in some nearshore shallow water regions where it reaches up to 10 cm s⁻¹ (Figure 4e). The seasonally-averaged bottom shear stress during the wet summer is relatively high in the PRE, nearshore regions, and the Taiwan Bank, where tidal dissipation is strong (Figure 4g).

In the dry winter season, the Hsig increases significantly compared to the wet summer, with waves primarily coming from the northeast, although refraction occurs in some nearshore regions, changing the wave direction to southeasterly (Figure 4d). The area with water depths exceeding 60 m has a Hsig greater than 1.5 m, while in the 20-meter depth region, the Hsig reaches approximately 1 m (Figure 4d). Compared to the wet summer, the WBOV increases significantly in the PRE mouth and many nearshore regions, reaching up to 10-20 cm s⁻¹ (Figure 4d). The average bottom shear stress on the continental shelf outside the estuary also increases significantly during the dry winter compared to the wet summer (Figure 4f).




Figure 4. The patterns of variables averaged for the wet summer season (April 1st to September 30th, 2017; Column 1) and the dry winter season (October 1st to March 31st, 2018; Column 2) in the Control case. Row 1 (a-b) depicts wind stress (color) and direction (vectors), Row 2 (c-d) shows Hsig (color) and propagation direction (vectors) weighted by Hsig², Row 3 (e-f) presents wave bottom orbital velocity (WBOV), and Row 4 (g-h) displays bottom shear stress magnitude.

The patterns of residual sediment dispersal, flux, and deposition over the simulation period provide clear information on the mechanisms for sediment redistribution on both annual and seasonal timescales. The following section presents

a detailed analysis of the seasonally averaged fields of salinity, flow, riverine SSC, depth-integrated riverine sediment flux, and riverine sediment deposition patterns during the wet summer season (Figure 5) and dry winter season (Figure 6) on the continental shelf.

During the wet summer season, when freshwater discharge is high and water column stratification is strong, riverine SSC (“riverine” means only Pearl River-derived sediment, classes 4-5 in Table 1, as follows) is primarily influenced by advection from the buoyant river plume (salinity less than 33.5 in Figure 3a, as follows) into the shelf sea, primarily in the surface layer (Figures 5a-b), high SSC regions closely align with the buoyant plume, as sediment is efficiently transported by the low-salinity, high-momentum freshwater outflow (Figures 5a-d). The buoyant plume extends both northeastward and southwestward along the coastline (Figure 5a). Due to the influence of southerly winds (Figure 4a) and ambient shelf currents, the extent of the buoyant plume extending northeastward is significantly higher than that extending southwestward. In terms of riverine sediment suspension, its estuarine turbidity maxima (ETM) zone ($\sim 100 \text{ mg L}^{-1}$) is situated in the shallow water area within the estuary (water depth $< 10 \text{ m}$) (Figure 5c-d). Beyond the estuary, suspended riverine sediment disperses across the shelf through the buoyant plume. Further away from the estuary, its distribution aligns with that of the buoyant plume, with concentrations diminishing as dispersal distance increases. The depth-integrated advective horizontal flux (without including vertical processes such as settling, resuspension, or diffusion, which are handled separately within the model) of riverine

582 sediment offers a clear indication of the primary net transport pathway of the riverine
583 sediment (Figure 5e). The riverine sediment exhibits both southwestward and
584 northeastward fluxes (Figure 5e). Southwestward coastal transport can extend as far
585 as the Leizhou Peninsula and Hainan Island. On the eastern side, the northeastward
586 transport extends toward the Taiwan Bank. However, the primary transport pathway
587 there is diverted southward (Figure 5e) due to the obstruction caused by summer
588 upwelling near the Guangdong east coast ([Chen et al., 2017a](#); [Chen et al., 2017b](#)), as
589 evidenced by the cross-shore current in the bottom layer (Figure 5b). The
590 southwestward transport pathway follows the region where the water depth is
591 shallower than 30 m, with a riverine sediment flux of $10\text{--}20\text{ g m}^{-1}\text{ s}^{-1}$. In contrast, the
592 northeastward transport pathway occurs in the 30–60 m depth range, but the riverine
593 sediment flux is below $10\text{ g m}^{-1}\text{ s}^{-1}$. Throughout the wet summer season, substantial
594 amounts of riverine sediment are deposited near the estuary (Figure 5d), particularly
595 leading to notably high deposition of riverine sediment near the river mouth (> 100
596 mm). Outside the estuary, the thickness of riverine sediment is comparatively lower,
597 but it can reach approximately ~ 0.5 mm during the wet summer season in certain
598 areas off the western Guangdong coast.




Figure 5. Patterns averaged over the entire wet summer season in the Control case: (a) surface and (b) bottom salinity (color, psu) and flow (arrows, m s^{-1}); (c) surface and (d) bottom riverine (classes 4 and 5 in Table 1, as follows) SSC (mg L^{-1}); (e) depth-integrated horizontal riverine sediment transport rate (color, $\text{g m}^{-1} \text{s}^{-1}$) and direction (arrows); and (f) riverine sediment deposition thickness (mm) on the seabed during the wet summer season. Flow vectors in regions with water depths exceeding 100 m are masked for clarity.

In contrast, during winter, when river discharge is low and vertical mixing is more intense, the correlation between the buoyant plume and riverine SSC is much weaker, and the riverine SSC is largely governed by resuspension processes driven by strong northeasterly winds and waves, rather than by freshwater transport. The expansion of the Pearl River buoyant plume is constrained to the southwestward

direction by strong northeasterly winds (Figure 6a), resulting in a narrow cross-shore width of the buoyant plume and the formation of a strong horizontal salinity gradient (i.e., a salinity front, particularly within the 30–33.5 psu range shown in Figure 6a) outside the estuary (Figure 6a). Flow velocity increases near this salinity front, facilitating the westward extension of the buoyant plume through the Qiongzhou Strait into the "Gulf" region. The riverine SSC is significantly lower than in the wet summer: in the ETM zone inside the PRE, riverine SSC falls from roughly 100 mg L^{-1} in summer to about 10 mg L^{-1} , while on the offshore shelf, it decreases from approximately 5 mg L^{-1} to around 2 mg L^{-1} (Figures 6c-d vs. 5c-d). During the dry winter, following the coastal current, the riverine suspended sediment primarily moves southwestward along the coast, deflecting southward along the topography near the Leizhou Peninsula (Figure 6c). It then bifurcates near the east entrance of the Qiongzhou Strait, with one branch continuing into the "Gulf" region, and the other one proceeding southward along the east coast of Hainan Island. Stronger winds and waves in the dry winter lead to the resuspension of a considerable amount of riverine sediments, originally deposited in "Proximal", "Western", and "Eastern" regions during summer. The resuspended sediments are then transported to coastal bays as well as to the sides and rears of the islands (Figure 6d). Additionally, a portion of the riverine sediment transported to the "Gulf" region gets deposited on the seabed during the dry winter season.




Figure 6. Same as Figure 5, but for the dry winter season in the Control case. Notably, (f) illustrates the changes in riverine sediment (classes 4 and 5 in Table 1) deposition on the seabed at the end of the dry winter season compared to the end of the wet summer season.

3.2 Riverine sediment budgets and annual deposition over the shelf

We present the sediment fluxes and retention amounts in different regions. Figure 7a-c illustrates the proportion of riverine sediment retention budget within each region, expressed as a percentage of the total annual river sediment load input (Figures 3a), for the wet summer season (Figure 7a), the dry winter season (Figure 7b), and the entire year (Figure 7c), based on under the Control run-case, respectively. Meanwhile, Figure 7d illustrates the annual deposition over the shelf.

The retention of Pearl River sediment on the continental shelf exhibits significant

seasonal variations (Figures 7a-c). During the wet summer (characterized by high discharge and relatively calm ~~low~~ wind/waves), the PRE and continental shelf receive 95.17% of the annual sediment input ~~load from the Pearl River~~ (Figures 3a and 7a). Of this, about ~~Approximately~~ two-thirds ~~of this sediment~~ is retained in the "Proximal" region (Figure 7a). Influenced ~~Additionally, influenced~~ by the prevailing southerly winds and northeastward shelf currents ~~circulation~~, 13.01% of the annual sediment load is retained in the "Eastern" and "Southeastern" regions (Figure 7a). Meanwhile, the shelf west of the PRE (⑤-⑧ regions) retains 15.87% of the annual load, with the "Western" region alone ~~being the primary receiver~~, accounting for 8.48% (Figure 7a). In contrast, only ~~Only~~ 0.92% and 2.3% ~~of the annual load~~ enter the more remote "Gulf" and "Distal" regions, respectively, ~~during summer~~ (Figure 7a). The "Southern" region retains a mere 1.22% of the sediment (Figure 7a).

In the dry winter (characterized by low discharge and energetic winds/waves), the PRE and the continental shelf receive only 4.83% of the annual sediment load (Figures 3a and 7b). The sediment distribution during this season ~~is~~ primarily reflects a ~~result of the dynamic~~ reworking of previously retained ~~the~~ sediments ~~of from~~ summer ~~deposition~~ (Figure 7b). Retention in ~~While~~ the "Proximal" region ~~continues to receive~~ ~~sediment, with a 1.38%~~ increases slightly (+1.38%) in retention, while retention ~~decreases in~~ the ②-⑥ regions ~~experience a decrease in sediment retention~~. Much of this remobilized sediment ~~This sediment~~ is ~~predominantly~~ transported farther offshore and retained in the ~~more distant~~ "Gulf" and "Distal" regions (Figure 7b).

The annual sediment budget reveals that 66.45% of the Pearl River sediment is

retained in the "Proximal" region (Figure 7c). Additionally, 9.2% is retained in the "Eastern" and "Southeastern" regions (Figure 7c), primarily during summer (Figures 7a vs. 7c), while 24.12% is retained on the shelf west of the PRE (⑤–⑧ regions), with most of that occurring ~~the majority~~ in the "Gulf" and "Distal" regions, ~~mainly~~ during winter (Figures 7b vs. 7c).

The annual deposition thickness of the Pearl River derived sediments (Figure 7d), ~~as illustrated in Figure 7d~~, reveals significant deposition within the "Proximal" region, with many areas exceeding 10 mm despite winter time resuspension and redistribution ~~transport~~. ~~Additionally, deposition on~~ In the "Eastern" region, deposition ~~reached~~ reaches a magnitude of 0.1 mm, while the ~~western~~ inner shelf west of the PRE ("Western" and "Gulf" regions) ~~exhibited~~ exhibits significantly greater accumulation. For instance, the deposition west of the Chuanshan Islands reached a magnitude of 0.5 mm. In the "Gulf" region, deposition ~~was~~ is primarily concentrated in the northeastern part, extending southwestward along the 30-60 m isobaths. Sediments transported southwestward along the east coast of Hainan Island and into ~~retained in~~ the "Distal" regions ~~did not predominantly settle on the seabed due to the greater water depth but~~ remained largely suspended in the water column due to the greater water depth, with limited deposition on the seabed.

(see Figure 3a). The black percentage values represent the combined total of riverine sediment Class 4 and Class 5, while the red and blue values denote sediment Class 4 and Class 5, respectively. Arrows indicate the direction of net riverine sediment flux at each transect during the specified period.

3.3 Model sensitivity experiments: relative roles of physical drivers, processes, sediment properties, and spin-up durations

Six sensitivity simulations, namely Exp 2-7 (NTS, NWS, NAS, NVS, DSV, and Cycle), were conducted (Table 2). As the latter three experiments do not impact hydrodynamics, we focus on presenting the seasonal mean differences in bottom shear stress between the Control run and the first three cases (NTS-Control, NWS-Control, NAS-Control) for both summer and winter (Figure 8).

In the NTS case, bottom shear stress is reduced relative to the Control run by a similar amount in both summer and winter due to the minimal seasonal variation in tidal intensity. This reduction primarily occurs in the PRE, around the Taiwan Bank, and near the Leizhou Peninsula (Figures 8a-b). In contrast, in the NWS case, the reduction in bottom shear stress is greater in winter than in summer, reflecting the intense seasonal variability of wind and wave activities (Figures 3b, 3d-e, and 4a-f). Unlike the NTS case, the NWS-induced decrease occurs mainly in the nearshore areas outside the PRE, although similar declines are also found around the Taiwan Bank and along the eastern side of the Leizhou Peninsula (Figures 8c-d). For the NAS case, the impact on bottom stress is minimal compared to the NTS and NWS cases. The effect

is almost negligible on the inner shelf at depths less than 100 m, with widespread impacts generally below 0.02 Pa. Some pronounced deviations are noted in localized deeper areas near the southern boundary of the domain (Figures 8e-f). These deviations, likely arising from boundary condition effects, are situated far from the Pearl River-derived sediment distribution areas (Figures 5-6). Consequently, they do not influence the dynamics of the Pearl River-derived sediment transport over the continental shelf (Figures 8e-f).




Figure 8. The seasonal mean differences in bottom shear stress between the Control run and the following cases: (a-b) NTS (NTS minus Control), (c-d) NWS (NWS minus Control), and (e-f) NAS (NAS minus Control). The first column represents the wet summer season, while the second column corresponds to the dry winter season.

731

732 Then, we analyzed seasonal riverine sediment transport and deposition patterns
733 (“riverine” means only the Pearl River-derived sediment, classes 4-5 in Table 1, as
734 follows) by comparing the control run with six sensitivity experiments (NTS-Control,
735 NWS-Control, NAS-Control, NVS-Control, DSV-Control, and Cycle-Control)
736 (Figures 9-11). The study focuses on the Pearl River-derived sediment dynamics,
737 indicated by surface circulation and riverine SSC distribution patterns (Figures 5 and
738 6). Specifically, Figures 9 and 10 present seasonal surface currents and SSC
739 differences between control and sensitivity runs, complemented by deposition pattern
740 differences in Figure 11.

741 Compared to the Control case, the NTS case demonstrates that ~~while~~-tides
742 significantly enhance bottom stress (Figures 8a-b), while ~~but~~ have minimal impact on
743 the mean circulation (Figures 5a, 6a, and 9a-b), and their exclusion reduces bottom
744 shear stress by over 0.2 Pa in the PRE and near the Leizhou Peninsula. Consequently,
745 increased deposition of the Pearl River-derived sediments occurs inside the PRE, its
746 adjacent areas, and on both sides of the Leizhou Peninsula (Figure 11a). During
747 summer, riverine SSC notably decreases in the ①–⑥ regions (Figure 9a). This
748 reduction pattern persists in winter, particularly in the PRE and on both sides of the
749 Leizhou Peninsula (Figure 9b).

750 Like the NTS, NWS has a relatively minor impact on circulation (Figures 5a, 6a,
751 and 9c-d). However, NWS leads to more Pearl River-derived sediment being
752 deposited in the nearshores of “Western” and “Eastern” regions (Figure 11b).

Consequently, the riverine SSC in summer is much lower in the downstream of the PRE and in ②–⑥ regions (Figure 9c). This similar reduction pattern persists in the winter, but is slightly in more western regions (Figure 9d).

For the NAS case, the impact on bottom stress is minimal compared to the NTS and NWS cases. However, NAS has a relatively large impact on the mean circulation (Figures 5a, 6a, and 9e-f). It mainly influences the summer circulation. Specifically, ignoring these factors would cause the relatively strong northeastward flow along the Guangdong coast to become very weak (Figure 9e). When it comes to winter, the influence of NAS on circulation is relatively small. That is, in the absence of the background residual water level and residual circulation, due to the strong northeasterly winds in winter, the overall circulation is still southwestward (Figure 9f). The decreased northeastward flow in summer leads to the Riverine SSC being scarcely transported to the vicinity of the “Eastern” and “Southeastern” regions. Consequently, the Riverine SSC there is decreased (Figure 9e) and sediment deposition is significantly reduced (Figure 11c). Most of the suspended Riverine sediment is transported southwestward, resulting in an increase in the Riverine SSC along the “Western” region. In winter, since most of the suspended Riverine sediment has been transported southwestward in summer, the Riverine SSC decreases compared to the Control run (Figure 9f). Ultimately, NAS mainly causes a significant reduction in sediment deposition in the “Eastern” region, while sediment deposition increases in the "Gulf" and the "Distal" regions (Figure 11c).




Figure 9. Same as Figure 8, but for seasonal mean differences in surface riverine SSC between the Control run and the following cases: (a-b) NTS-Control, (c-d) NWS-Control, and (e-f) NAS-Control. The first column represents the wet summer season, while the second column corresponds to the dry winter season. Vectors show the seasonal mean surface current fields in each experiment. Note that the riverine SSC values shown in the figure correspond to classes 4 and 5 as defined in Table 1.

For the NVS case, the summer conditions of NVS are precisely the same as those of the Control run (Figure 10a). Since the critical shear stress for erosion in winter is lower than that in the Control run, this leads to an increase in re-suspension in the “Proximal”, “Western” and the “Gulf” regions, increasing Riverine SSC (Figure 10b).

Eventually, this causes a reduction in the deposition thickness of the Pearl River-derived sediments in these regions (Figure 11d).

In the DSV case, significant reductions occur in the primarily high SSC areas in the Control run in both summer and winter (Figures 5c, 6c, 10c-d). The enhanced settling velocity results in an increased deposition of Pearl River-derived sediments along the Guangdong coastline ("Western" and "Eastern" regions) and the eastern "Gulf" region, accompanied by a reduced deposition thickness in the western "Gulf" region (Figure 11e).

In the Cycle case, the new riverine sediment input and its transport processes during the Cycle experiment are nearly identical to those in the Control run. Therefore, compared to the Control run, the Cycle experiment specifically focuses on examining the impact of the presence of pre-existing Pearl River-derived sediments on estimating the riverine SSC and the annual seabed riverine sediment budget in the second year. Consequently, during the summer period, the Cycle case experiences elevated riverine SSC in the primary depocenters identified in the Control run (Figures 7d and 10e), while this effect is diminished by winter (Figure 10f). Figure 11f thus captures the transport trends of pre-existing riverine sediments in the second year, demonstrating that riverine sediments deposited during the first year can be resuspended and transported further southwestward during the second year. This migration is driven by the annually averaged net alongshore coastal current, which remains predominantly directed toward the southwest. The current becomes stronger during the winter monsoon under the influence of prevailing northeasterly winds, whereas the opposing

summer southerly winds are comparatively weaker, indicating a persistent southwestward sediment transport trend over multi-year timescales.




Figure 10. Same as Figure 9, but for the latter three experiments (NVS, DSV, Cycle). (a-b) NVS-Control, (c-d) DSV-Control, and (e-f) Cycle-Control.




Figure 11. The differences in annual deposition patterns of Pearl River-derived sediment (classes 4-5, Table 1) on the seabed between the Control run and the following cases: (a) NTS-Control, (b) NWS-Control, (c) NAS-Control, (d) NVS-Control, (e) DSV-Control, and (f) Cycle-Control.

3.4 Modeled regional retention budgets in sensitivity experiments

Finally, we analyze the impact of various factors on the annual riverine sediment retention budget across different regions. Specifically, Figure 12 illustrates the annual riverine sediment retention budget in various regions under six sensitivity simulations, namely Exp 2-7 (including NTS, NWS, NAS, NVS, DSV, and Cycle). It should be

noted that the retention percentages budget and their variations discussed hereinafter are all relative to the annual riverine sediment load (Figure 3a).

As shown in Figure 12, tides and sediment settling velocity have the most significant impact on the retention in the "Proximal" region. In the NTS case and the DSV case, the retention in the "Proximal" region is 70.92% and 71.57%, respectively (Figures 12a and 12e), which is higher than 66.45% in the Control run (Figure 7c). This indicates that ignoring tides will cause the "Proximal" region to capture more riverine sediments, and a larger settling velocity will result in more riverine sediments being retained within the "Proximal" region. In these two cases, compared with the Control run, the retention in the "Gulf" and "Distal" regions decreases. Meanwhile, the DSV case causes the greatest increase in retention in the "Western" region, with an increase of +1.91%.

Furthermore, the NWS also leads to a 2.2% increase in retention in the "Proximal" region (Figure 12b), which is lower than that in the NTS case. This shows that tides dominate resuspension versus deposition in the "Proximal" region more than waves do. However, for the "Western" region, compared with the NTS case, the NWS causes a greater increase in retention, indicating that waves dominate the resuspension of Pearl River-derived sediments in these nearshore areas more than tides do.

For the "Eastern" and "Southeastern" regions, NAS brings about the most dramatic changes, the retention of Pearl River-derived sediments in these regions drops from 9.1% to 0.84% compared to the Control run (Figure 12c). Meanwhile, ignoring these background circulations results in a substantial increase in the retention

in the "Distal" region, with an increase of 6.49%.

The NVS case leads to a decrease in the retention of the Pearl River-derived sediments in ①–⑥ regions compared to the Control run. The reduction ranges from -0.05% to -0.85% (Figure 12d), which in turn causes the retention in the "Gulf" and "Distal" regions to increase by 0.7% and 1.47%, respectively. Overall, compared with scenarios that ignore physical drivers ~~processes~~ and alter sediment settling velocity (NTS, NWS, NAS, and DSV), the NVS scenario has a relatively smaller impact on the retention of the Pearl River-derived sediments.

Finally, in the Cycle case, to isolate the pre-existing Pearl River-derived sediments, the initial retentions (the end conditions of the Control run on March 31, 2018) were subtracted before calculating the retention in the Cycle case (Figure 12f). The retention of Pearl River-derived sediments in ①–⑥ regions shows little variation, with values ranging from -0.98% to +0.24% compared to the Control run (Figure 12f). The most significant changes are the decreases and increases in retention in the "Gulf" and "Distal" regions, which are -2.17% and +3.54%, respectively. This demonstrates the long-term trend of southwestward transport of Pearl River-derived sediments on the shelf (relative to the Control run).




Figure 12. Same as Figure 7c, but for the other six cases: (a) NTS, (b) NWS, (c) NAS, (d) NVS, (e) DSV, and (f) Cycle, respectively. All percentages shown in the figure are expressed relative to the annual riverine sediment load (see Figure 3a). Magenta values denote the differences in retention percentage of riverine sediments (Classes 4 and 5; Table 1) between the Control run and each sensitivity case. Black values represent the combined retention of Classes 4 + 5, while blue values indicate Class 5 alone. To obtain the retention percentage for Class 4, simply subtract the Class 5 percentage (blue) from the combined Classes 4 + 5 percentage (black).

4. Discussions

4.1 Fidelity of our model results

We simulated the suspension, transport, and deposition of the Pearl River-derived sediment over the shelf from April 2017 to March 2018. Comparisons with multiple types of observations demonstrated that the model simulation reasonably well captured the regional patterns and temporal variability of water levels (Figures S2-S4), surface waves (Figures S5-S6), estuarine and shelf currents (Figures S8a-b and S9-S10), salinity and temperature (Figures S7a-b and S8c-d), and SSC (Figures S7c and S8e-f). Even though we have made substantial efforts to collect observational data and conduct calibration and validation, the large scope of the study area and the fact that the study covers an entire year mean that there are some inherent challenges to achieving a complete analysis. The real-world situation may be extremely complex, and these validations may still not be sufficient to address all issues (such as the accurate parameterization of sediment characteristics and their seasonal variations, as well as the proportion of slow-settling fine grains and flocculated flocs in riverine sediments). Therefore, in this section, we intend to discuss the fidelity of our results.

Studies have demonstrated a seasonal dependence of sediment critical shear stress ([Xu et al., 2014](#); [Briggs et al., 2015](#)). On the Louisiana shelf, seabed erodibility is controlled by grain size, sediment age, proximity to river sources, bioturbation, and flood deposits, and is higher during the wet season than the dry season ([Xu et al., 2014](#)). It is also linked to seasonal hypoxia: sites experiencing hypoxic (hypoxia greater than 75% of the time, hypoxia defined as $< 2 \text{ mg O}_2 \text{ L}^{-1}$) conditions exhibit

greater erodibility, whereas the normoxic (hypoxia less than 25% of the time) site shows the lowest erodibility ([Briggs et al., 2015](#)). These findings imply that sediment models should incorporate seasonal variations in critical shear stress, parameterized using field observations or seasonal sampling data. Similarly, in the PRE, field observations, laboratory experiments, and numerical sensitivity analyses have shown that the critical shear stress for erosion of sediments in the PRE is higher in winter/dry season than in summer/wet season ([Dong et al., 2020](#); [Cao et al., 2025](#)). The summer period in the PRE also coincides with seasonal hypoxia ([Cui et al., 2018](#); [Cui et al., 2022](#)), which likely contributes to this seasonal variation in erodibility. Therefore, we considered the seasonal variation of critical shear stress for erosion in a suite of model experiments, that is, increasing the critical shear stress for erosion in winter. This setting was also applied during the 15-month spin-up before conducting the model experiments ([the spin-up employed the same forcing conditions and model settings as the Control run to maintain consistency](#)~~the driving force of the spin-up is realistic, i.e., the settings are consistent with those of the Control run~~). After ~~realistic reworking~~ during the spin-up, the bed sediment grain size distribution (used as initial conditions in the Control run and all Sensitivity cases except the Cycle case) is quite close to the initial prototype (Figures 2d-f vs. 2g-i).

When the seasonal variation of critical shear stress for erosion is omitted (NVS case), the model results suggest a reduced retention of Pearl River-derived sediments in ①–⑥ regions during winter, alongside an increased retention in the "Gulf" and "Distal" regions. However, these differences are relatively small in magnitude

compared to the annual load (Figure 12d). Thus, accounting for seasonal variations in critical shear stress for erosion has a limited influence on the annual-scale retention patterns. The dispersal distance of fluvial sediments on continental shelves is strongly influenced by settling velocity ([Harris et al., 2008](#)). For example, Apennine-derived sediments, characterized by lower settling velocities, travel farther before deposition than Po River sediments, which are predominantly flocculated and settle more rapidly ([Fox et al., 2004](#); [Harris et al., 2008](#)). Likewise, our results suggest that selecting an appropriate settling velocity parameter exerts a greater control on sediment dispersal patterns than accounting for seasonal variations in critical shear stress for erosion (Figure 12d vs. 12e). The results of the DSV case show that a sediment settling velocity twice that of the Control run leads to the highest retention in the "Proximal" and "Western" regions across all experiments (Figure 12e), while reducing the retention in the "Distal" region (Figure 12e). Although the settling velocity we adopted is based on previous studies ([Xia et al., 2004](#)) and model calibrations (Figures S7c and S8e-f), with due consideration given to the presence of slow-settling single fine grains and high-settling flocs in riverine sediments, certain discrepancies might still exist in this setting. These discrepancies are contingent upon the actual magnitude of the low settling velocity of fine grains. In almost all cases, all flocs [in the model](#) are retained in the "Proximal" region (Figure 12a-e), and only in the Cycle case, flocs accounting for 0.01% of the annual load are retained in the "Western" region west of the "Proximal" region (Figure 12f, blue values), indicating that high-settling flocs hardly leave the "Proximal" region. This finding shows close alignment with, yet

exhibits minor distinctions from, the observed patterns in the Mekong Shelf ([Xue et al., 2012](#)). [Xue et al. \(2012\)](#) found that while the preponderance of flocs is deposited on the Mekong delta front precisely at the river mouth, a quantity equivalent to 1.6% of the annual riverine sediment load of flocs is deposited on the downdrift delta front further downstream from the river mouth. This is mainly because the estuarine bay of the PRE is wider and there are numerous islands near the river mouth. The overwhelming majority of flocs are either deposited within the estuarine bay or captured by the surrounding islands. In conclusion, our results are affected by the settling velocity of fine grains. More field observations and studies on model parameterization regarding the settling velocity of fine grains are urgently needed.

As previously noted, we classified riverine sediments into two categories based on established research: 40% slow-settling fine grains and 60% fast-settling flocs. This 40%/60% distribution is consistent with the setting from earlier studies ([Zhang et al., 2019](#); [Zhang et al., 2021](#)), as summarized in Table 1. While such assumptions are necessary for modeling purposes, the actual composition of riverine sediments in natural environments remains uncertain. To evaluate the sensitivity of our results to this uncertainty, we conducted a conceptual analysis. If all riverine sediments were hypothetically composed entirely of fast-settling flocs, they would be completely retained near the source, with no transport to the "Gulf" region. However, this scenario is inconsistent with the radionuclide measurements obtained from "Gulf" region surface sediment samples ([Lin et al., 2020](#)). On the other hand, if all sediments were considered slow-settling fine grains, only 16.13% would be retained in

“Proximal” region under normal conditions (or 28.9% in the DSV case), a result that diverges significantly from established research.

[Chen et al. \(2023\)](#) analyzed high-resolution seismic data and demonstrated that approximately 35% of the Pearl River-derived sediment has been transported to offshore shelf areas over the past 6,500 years, suggesting that 65% was deposited proximally. Our findings are in close agreement, indicating that 66.45% of the Pearl River sediments are retained in the proximal region, while 33.55% are transported elsewhere. This consistency with [Chen et al. \(2023\)](#) supports the validity of our approach. Taken together, these analyses confirm that the 40%/60% fraction assumption is a reasonable approximation for modeling purposes.

Furthermore, our model results demonstrate reasonable reliability in other aspects. [Liu et al. \(2009\)](#) and [Ge et al. \(2014\)](#) using chirp sonar profiles from the inner shelf of the South China Sea combined with [Zong et al. \(2009\)](#)’s onshore borehole data, found that the thickness of Pearl River-derived sediments within the PRE since the Holocene is over 20 m, while the mud thickness in the shallow waters west of the Chuanshan Islands (see Figure 1) is approximately 5-10 m. Our calculated annual sediment thicknesses for these two regions are approximately 2 mm and 0.3 mm (Figure 7d), respectively. Given our model’s annual riverine sediment load of 34.52 megatons, which has been significantly reduced due to recent human activities ([Dai et al., 2008](#)), compared to the widely accepted Holocene average of around 90 megatons ([Liu et al., 2009](#)), we estimate the total sediment thickness over the past 7500 years to be roughly 39 meters and 6 meters for these depositional zones, consistent with previous studies

([Liu et al., 2009](#); [Zong et al., 2009](#)).

Furthermore, our results reveal that 8.86% of the riverine sediment derived from the Pearl River is transported to the "Gulf" region (Figure 7c), primarily during the winter season (Figure 7b). This finding is not only consistent with the earlier speculation proposed by [Ge et al. \(2014\)](#) but also supplements the conclusions drawn by [Lin et al. \(2020\)](#). From a hydrodynamic perspective, [Shi et al. \(2002\)](#) found that the net flux of currents in the Qiongzhou Strait is westward throughout the year. Our results for both wet summer (Figures 5a-b) and dry winter currents (Figures 6a-b) in the Qiongzhou Strait are consistent with [Shi et al. \(2002\)](#). This westward flow contributes to the westward transport of Pearl River sediment to the "Gulf" region.

4.2 Implications of our model results

The fate of sediment dispersed from the river into the coastal ocean involves at least four processes: supply via buoyant plumes; initial deposition; resuspension and transport by marine processes; and long-term net accumulation ([Wright and Nittrouer, 1995](#)). In general, a significant proportion of river sediment tends to deposit in the estuary and its vicinity ([Walsh and Nittrouer, 2009](#); [Hanebuth et al., 2015](#)).

[Walsh and Nittrouer \(2009\)](#) present a hierarchical decision tree designed to predict the marine dispersal system at a river mouth based on fundamental oceanographic and morphological characteristics. Within this framework, riverine sediment deposition is characterized using key factors, including riverine sediment discharge (greater or less than 2 megatons), shelf width (greater or less than 12 km), and wave and tidal range conditions (greater or less than 2 m) ([Walsh and Nittrouer,](#)

2009).

We aim to analyze our PRE simulation results using the framework established by [Walsh and Nittrouer \(2009\)](#). Although the Pearl River's [annual](#) riverine sediment discharge (Figure 2a) exceeds the [Walsh and Nittrouer \(2009\)](#)'s 2 megatons per year threshold, most of the sediment still remains deposited near the estuary (Figure 7c), indicating an estuarine accumulation-dominated (EAD) system. [This behavior deviates from the predictions of](#), ~~unlike~~ the hierarchical decision tree proposed by [Walsh and Nittrouer \(2009\)](#). Outside the estuary, the continental shelf, spanning 200-220 km in width ([Liu et al., 2014](#)), significantly exceeds [Walsh and Nittrouer \(2009\)](#)'s 12 km threshold. As a result, most escaped riverine sediments tend to accumulate on the shelf rather than being captured by submarine canyons (Figures 7c-d). This wide, shallow shelf promotes sediment deposition and limits the direct transport of fine sediments into deeper waters ([Walsh and Nittrouer, 2009](#)). Subsequently, given that the annual mean tidal range ([Chen et al., 2016](#); [Gong et al., 2018b](#)) and Hsig (Figure 3d) near the PRE are both below the 2 m threshold established by [Walsh and Nittrouer \(2009\)](#), the majority of escaped riverine sediments are predominantly deposited in the proximal depo-center. Our findings demonstrate that most of our outcomes are consistent with the hierarchical decision tree proposed by [Walsh and Nittrouer \(2009\)](#), except for the application of the 2 megatons per year threshold for riverine sediment discharge. This phenomenon can primarily be attributed to the unique geomorphological characteristics of the PRE, including its broad mouth (Figures 1 and S1), extensive accommodation space encompassing

approximately 2385 km² of water area ([Wu et al., 2018](#)), the presence and sheltering effect of numerous adjacent islands ([Li et al., 2024b](#)), and the division of fluvial sediment discharge through eight distinct outlets ([Hu et al., 2011](#)).

The monsoonal nature of the northern SCS (Figures 4a-b) induces pronounced seasonal variations in Pearl River-derived sediment transport and deposition (Figures 5 and 6). During the summer wet season, the Pearl River delivers approximately 95.17% of its annual sediment load to the PRE and the adjacent shelf (Figure 7a) under relatively calm wind and wave conditions (first column of Figure 4), leading to predominant proximal deposition (Figure 5f). In contrast, the winter dry season is characterized by strong northeasterly monsoon winds that generate high-energy waves (second column of Figure 4), significantly increasing bottom shear stress (Figures 4g-h). This process resuspends previously deposited sediments and facilitates their redistribution, particularly toward regions such as the "Gulf" region (Figure 6f).

The PRE exhibits distinctive geomorphological features, yet dispersal of its fine-grained sediment transport on the continental shelf conforms to general patterns observed offshore of other monsoon-influenced estuarine systems. Similar multiple-stage sediment delivery and dispersal mechanisms have been documented offshore of various major estuaries and their adjacent shelves, including the Yellow River Shelf ([Bian et al., 2013](#); [Zeng et al., 2015](#)), Changjiang River Shelf ([Zeng et al., 2015](#)), and Mekong River Shelf ([Xue et al., 2012](#); [Eidam et al., 2017](#)), demonstrating comparable sedimentary processes under monsoon climatic influences. In these systems, sediment transport is not confined to a single process but rather occurs in

stages, influenced by seasonal variations in hydrodynamic conditions. Like the PRE Shelf, the Mekong Shelf experiences distinct phases of sediment deposition, with fine sediments being delivered during periods of high river discharge and then redistributed by waves and tidal forces, particularly during monsoonal shifts ([Xue et al., 2012](#); [Eidam et al., 2017](#)). These complex patterns highlight the interplay between riverine inputs, coastal morphology, and oceanographic processes in shaping sediment dynamics.

These sediment delivery patterns have implications beyond sediment fate, particularly for carbon cycling. Sediment deposition in coastal and shelf areas plays a significant role in trapping organic carbon, influencing long-term carbon burial rates ([LaRowe et al., 2020](#)). Sediment dynamics directly influence the fate of organic carbon (OC) in marine environments, where sediments function as both a sink and a source of OC, playing a pivotal role in global carbon cycling ([Repasch et al., 2021](#)). The multiple-step transport mechanisms can lead to varying carbon storage locations, affecting the sequestration potential of these systems. Additionally, resuspension and redistribution of sediments, especially during high-energy events, may expose previously buried organic material, leading to carbon remineralization and influencing coastal nutrient cycles and ecosystem health ([Ståhlberg et al., 2006](#); [Moriarty et al., 2018](#)). Therefore, understanding these patterns is crucial for assessing the broader impacts on carbon cycling and coastal biogeochemical processes.

4.3 Limitations and future work

This study focuses on analyzing simulation results from a typical year,

encompassing both wet and dry seasons from 2017 to 2018, to understand the seasonal variations and annual patterns of suspension, transport, and deposition of sediment in the PRE and adjacent shelf. However, it's essential to recognize that the long-term sediment transport and deposition dynamics in the Pearl River are influenced by numerous complex factors. These include changes in sea level and coastal line ([Church and White, 2006](#); [Harff et al., 2010](#); [Hong et al., 2020](#); [Lin et al., 2022](#); [Ma et al., 2023](#)), alterations in wind field and precipitation ([Ning and Qian, 2009](#); [Young et al., 2011](#)), natural sedimentation within the Pearl River Delta ([Wu et al., 2010](#)), modifications in sediment load and underwater volume of the estuary caused by anthropogenic impact ([Wu et al., 2014](#); [Wu et al., 2018](#); [Lin et al., 2022](#)), interannual variations of the shelf circulations ([Liu et al., 2020](#); [Deng et al., 2022](#)) and Kuroshio intrusions ([Caruso et al., 2006](#); [Nan et al., 2015](#); [Sun et al., 2020](#)). Therefore, while this study sheds light on seasonal and annual timescale patterns, it cannot fully represent the short or long-term transport and deposition trends of the Pearl River sediment. Yet for many shelf systems, a lot of sediment transport happens during short-lived events such as hurricanes ([Xu et al., 2016](#); [Warner et al., 2017](#); [Georgiou et al., 2024](#)). Consideration of the episodicity of transport would be helpful for future studies ([Xu et al., 2016](#); [Warner et al., 2017](#); [Georgiou et al., 2024](#)).

Additionally, it's important to note that this article primarily focuses on the fate of the Pearl River sediment on the inner shelf. However, within the expansion range of the Pearl River buoyant plume, a number of smaller rivers, including the Jiulong River, Han River, Moyang River, Jian River, Nanliu River, Changhua River and

Nandu River, also contribute freshwater and sediment to the northern South China Sea (Milliman and Farnsworth, 2011; Zhang et al., 2012; Liu et al., 2016). Although these rivers contribute significantly less freshwater and sediment compared to the Pearl River, they still impact seawater salinity, suspended sediment concentration, and seabed geomorphology (Liu et al., 2016; Wang et al., 2023; Zong et al., 2024). Since the 1950s, South China delivers approximately 102 Mt/year of fluvial sediment to the SCS, with the Pearl River alone accounting for 84.3 Mt/year—about 83% of the total sediment load (Milliman and Farnsworth, 2011; Zhang et al., 2012; Liu et al., 2016).

~~The specific contributions of each river are detailed in Table 3.~~ While the data highlight the Pearl River's plays a dominant role in sediment delivery to the northern South China Sea, a comprehensive understanding of the region's sedimentary processes and impacts ~~in the northern South China Sea~~ also requires a systematic investigation of the contributions from ~~into the roles of the~~ smaller ~~contributing~~ rivers.

~~**Table 3.** Annual mean runoff and annual suspended sediment load of major rivers in South China that flow directly into the northern South China Sea since the 1950s (Milliman and Farnsworth, 2011; Zhang et al., 2012; Liu et al., 2016).~~

River name	Runoff ($\text{m}^3 \text{s}^{-1}$)	Suspended sediment load (Mt/year)
Pearl River	9075	84.3
Jiulong River	476	3.1
Han River	825	10
Moyang River	269	0.8

Jian River	174	1.5
Nanliu River	162	1.1
Changhua River	120	0.08
Nandu River	179	0.4

Then, while the model used in this study performs well in~~has shown good~~ validation~~results~~, further progress relies on additional observational data to better constrain key parameters such as settling velocity and critical shear stress for erosion. ~~conducting more sensitivity experiments on sediment parameters, such as settling velocity and critical shear stress for erosion, would be beneficial.~~ Direct measurements under varying conditions would help refine these inputs and improve model accuracy. For instance, ~~Settling~~ settling velocity can~~influences~~ the location of sediment depocenters, with higher settling velocities leading to more proximal entrapment and vice versa ([Harris et al., 2008](#)). Similarly, critical erosion stress ~~can~~ affects the sediment resuspension~~of deposited sediment, particularly with higher critical erosion stress resulting in less resuspension and more deposition especially~~ during neap tides and weak wind wave events ([Dong et al., 2020](#); [Choi et al., 2023](#)). Targeted~~Conducting such~~ sensitivity analyses, supported by such data, would enhance our understanding of sediment dynamics in estuaries and shelves. Besides, the model does not account for cohesive processes, such as consolidation and flocculation, which can significantly impact sediment behavior ([Sherwood et al., 2018](#)). Our model does not incorporate wave and current-supported gravity flows, which are important

factors influencing sediment transport in submarine canyon areas ([Harris et al., 2005](#); [Ma et al., 2010](#); [Zhang et al., 2020](#)). Since our study area primarily focuses on the continental shelf and the simulated results indicate that sediment transport occurs mainly in the shallow inner shelf, where canyons are relatively rare, this omission has a relatively minor impact on our results.

Lastly, we employ the COAWST model, ~~which uses~~ with an S-coordinate vertical system ~~in the vertical direction with that enhances increased~~ resolution near the surface and bottom layers ([Song and Haidvogel, 1994](#)). This vertical layering allows cell heights to vary, enabling finer resolution in dynamically important regions and improving performance in areas with sloping bathymetry compared to traditional sigma-coordinate systems ([Bryan, 1969](#); [Song and Haidvogel, 1994](#)). ~~In addition, our model includes h~~Horizontal grid refinement in the PRE further enhances, ~~enhancing the model's its~~ ability to resolve estuarine features, successfully capturing. ~~As a result, the model effectively captures~~ estuarine turbidity maxima (ETM) and ~~horizontal~~ salinity fronts (Figures S11 and S12, see Supplement). ~~During summer, multiple ETMs appear near the estuary bottom (Figure S11b), and while these features persist in winter, their concentrations vary (Figure S12b),~~ consistent with the findings of [Wang et al. \(2018\)](#), [Zhan et al. \(2019\)](#), [Zhang et al. \(2021\)](#), [Ma et al. \(2022\)](#), and [Ma et al. \(2024\)](#). ~~Horizontal salinity fronts are also well represented, showing an upstream shift from the high discharge summer season to the low discharge winter season (Figures S11e-f and S12e-f), in agreement with previous studies by Zhang et al. (2021) and Ma et al. (2024).~~ Nonetheless, compared with the S-coordinate system,

models that employ vertically adaptive layering (e.g., SCHISM, the Semi-implicit Cross-scale Hydrosience Integrated System Model, [Zhang et al. \(2016\)](#)) or Cartesian vertical coordinates (e.g., MITgcm, the MIT General Circulation Model, [Marshall et al. \(1997a\)](#); [Marshall et al. \(1997b\)](#)) generally perform better in regions with steep topographic gradients ([Bijvelds, 2001](#)). ~~Future studies~~ ~~Therefore, future research~~ could benefit from ~~such approaches combined with finer~~ ~~adopting models with higher~~ horizontal resolution and Cartesian vertical coordinates to improve ~~the simulation of~~ Pearl River-derived sediment dynamics ~~simulations~~ ~~across the estuary and the adjacent shelf~~.

5. Conclusions

This study utilizes the COAWST model to quantitatively analyze the seasonal suspension, transport, and annual fate of Pearl River-derived sediment (classes 4-5 in Table 1) on the continental shelf over a typical year, capturing key marine variables such as water level, wave height, flow velocity, salinity, temperature, and SSC.

The monsoonal nature of the northern SCS (Figures 4a-b) induces pronounced seasonal variations in Pearl River-derived sediment transport and deposition (Figures 5 and 6). During the wet summer, calm conditions foster initial Pearl River-derived sediment deposition via the buoyant plume (Figures 5 and 7a). Conversely, winter's stronger winds and waves resuspend and transport Pearl River-derived sediments into "Gulf" region (Figures 6 and 7b). Our quantitative assessment reveals distinct spatial patterns in the annual fate of riverine sediments: approximately two-thirds of the Pearl

River-derived sediment is retained within the estuarine vicinity (“Proximal” region), while about 9% reaches the continental shelf east of the PRE (“Eastern” and “Southeastern” regions), while similar proportions are transported to and retained in “Gulf” and “Distal” regions, respectively (Figure 7c). Furthermore, we evaluated the contributions of different physical drivers ~~processes~~ by comparing the Control run with the reduced-physics sensitivity experiments. Our analysis reveals distinct roles of tidal forces, wave action, and background circulation in governing the transport and deposition of Pearl River-derived sediments (Figures 8, 9, 11a-c, and 12a-c).

Tidal dynamics play a primary role in ~~primarily governing~~ sediment behavior within and offshore of the PRE. In the presence of ~~Relative to the Control run,~~ ~~neglecting~~ tides, ~~leads to lower~~ bottom shear stress in the PRE is enhanced (Figures 8a-b), promoting higher Pearl River-derived sediment resuspension and reducing excessive sediment deposition ~~deposition~~ in the PRE (Figure 11a), while facilitating ~~and reduced~~ sediment retention patterns in “Gulf” and “Distal” regions (Figure 12a). Wave activity primarily controls Pearl River-derived sediment resuspension in three critical dimensions: (1) the river mouth, (2) the nearshore of “Eastern” and “Western” regions outside the estuary, and (3) periods characterized by high wave energy during winter (Figures 9c-d). ~~Excluding waves results in increased modeled Pearl River-derived~~ These wave-driven resuspension processes regulate sediment deposition and accumulation patterns in these areas and seasons ~~times~~ (Figure 11b), contributing to sediment export from the nearshore to offshore regions ~~such as~~ ~~while reducing accumulation in~~ the “Gulf” ~~region~~ (Figure 11b). Background

circulation exerts its strongest influence~~is most influential~~ in summer (Figure 5a vs. Figure 9e), with a strong northeastward current transporting Pearl River-derived sediments toward regions "Eastern" and "Southeastern"~~③-④~~. When this current is weak or absent, sediment delivery to these regions drops significantly, with only 0.84% of the total reaching them, while deposition increases in the "Gulf" and "Distal" regions~~Without this current and under weak southerly winds, only 0.84% of Pearl River-derived sediments are transported to regions ③-④, while deposition in "Gulf" and "Distal" regions is greater compared to the Control run~~ (Figure 12c). The sediment model solutions are also highly sensitive to the parameterization of sediment characteristics and spin-up durations (riverine or seabed sediments) (Figures 10, 11d-f, and 12d-f). The natural seasonal increase in critical shear stress for erosion during winter counteracts part of the wave-enhanced resuspension capacity, thereby reducing resuspension and erosion on the continental shelf east of the Leizhou Peninsula (Figure 10b).~~Neglecting the seasonal increase in critical shear stress for erosion leads to enhanced resuspension and erosion in regions ①-⑥ during winter (Figure 10b), which in turn shifts more sediment toward "Gulf" and "Distal" regions compared to the Control run (Figure 12d).~~ Increasing the settling velocity reduces the overall riverine SSC (Figures 10c-d) and results in a spatial redistribution pattern characterized by greater retention mainly in "Proximal" and "Western"~~in ①-③ and ⑤~~ regions and reduced riverine sediment presence in "Gulf",~~and~~ "Distal" and "Southeastern" regions relative to the Control simulation (Figure 12e). Additionally, the modeled riverine SSC is influenced by pre-existing Pearl River-derived sediments,

as shown in the Cycle experiment (Figure 10e-f). The experiment highlights the effect of riverine sediment spin-up, showing that first-year retained Pearl River-derived sediments are predominantly redistributed from the “Gulf” region toward the more distant “Distal” region during the second year. ~~In contrast, the model exhibits only minor sensitivity to the duration of seabed sediment spin-up, as demonstrated in the Cycle2 experiment (see Supplement), in which riverine sediments present during the spin-up period were added to seabed sediment classes 1–2 at the start of the Cycle2.~~

Acknowledgments

This research was funded by the National Natural Science Foundation of China (grant numbers 42306015 and 42276169), the China Postdoctoral Science Foundation (grant number 2023M743988). Wenping Gong is supported by the Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai) (SML2023SP238). The authors would like to thank the crew of the R/V Changhe Ocean for their valuable contribution during the collection of the field data. We express our gratitude to the Co-editors-in-chief, Dr. Mario Hoppema, and three anonymous reviewers for their valuable suggestions in enhancing and improving this article.

Data availability

The figure data and model configuration files used in this paper can be downloaded from: <https://doi.org/10.5281/zenodo.15013448>. The HYbrid Coordinate Ocean Model (HYCOM) outputs are from: <http://hycom.org/hycom>. The NCEP Climate Forecast System Version 2 (CFSv2) reanalysis data can be obtained at the following website: <https://rda.ucar.edu/datasets/ds094.1/dataaccess/>. The NOAA

WAVEWATCH III global ocean wave model output fields can be downloaded from:
<ftp://polar.ncep.noaa.gov/pub/history/waves>. Hourly water-level data observed at
Quarry Bay station are provided by the Hong Kong Observatory website:
<https://www.hko.gov.hk/sc/tide/marine/realtide.htm?s=QUB&t=TABLE>. Hourly
water-level data from Zhapo and Qinglan stations, provided by the Flanders Marine
Institute (VLIZ), are part of the UNESCO/IOC Global Sea Level Observing System
(GLOSS) and accessible at <http://www.ioc-sealevelmonitoring.org>. The mooring data
for the M1 and M2 stations are sourced from [Liu et al. \(2023\)](#) and [Li et al. \(2024a\)](#).

~~The figure data and model configuration files used in this paper can be downloaded
from: <https://doi.org/10.5281/zenodo.15013448>.~~

Declaration of Competing interests

The authors declare that they have no known competing financial interests or
personal relationships that could have appeared to influence the work reported in this
paper.

CRedit authorship contribution statement

GZ: Conceptualization, Numerical modeling, Validation, Data visualization,
Writing-original draft, and Funding acquisition. **SH:** Writing-review & editing,
Validation. **XY:** Writing-review & editing. **HZ:** Writing-review & editing. **WG:**
Writing-review & editing, and Funding acquisition

Supplement:

The Supplement includes validation and analysis of the model's water levels,
Hsig, flow velocities, salinity, temperature, and SSC. It provides additional text and

figures that support the model validation and supplementary analyses, which could not be fully presented in the main article due to space limitations.

References

- Bever, A. J., Harris, C. K., Sherwood, C. R., and Signell, R. P.: Deposition and flux of sediment from the Po River, Italy: An idealized and wintertime numerical modeling study, *Marine Geology*, 260, 69-80, 10.1016/j.margeo.2009.01.007, 2009.
- Bever, A. J., and MacWilliams, M. L.: Simulating sediment transport processes in San Pablo Bay using coupled hydrodynamic, wave, and sediment transport models, *Marine Geology*, 345, 235-253, 10.1016/j.margeo.2013.06.012, 2013.
- Bian, C., Jiang, W., and Greatbatch, R. J.: An exploratory model study of sediment transport sources and deposits in the Bohai Sea, Yellow Sea, and East China Sea, *Journal of Geophysical Research: Oceans*, 118, 5908-5923, <https://doi.org/10.1002/2013JC009116>, 2013.
- Bijvelds, M. D. J. P.: Numerical modelling of estuarine flow over steep topography, Doctoral, Delft University of Technology, 2001.
- Booij, N., Ris, R. C., and Holthuijsen, L. H.: A third-generation wave model for coastal regions: 1. Model description and validation, *Journal of Geophysical Research: Oceans*, 104, 7649-7666, <https://doi.org/10.1029/98JC02622>, 1999.
- Brand, A., Lacy, J. R., Hsu, K., Hoover, D., Gladding, S., and Stacey, M. T.: Wind-enhanced resuspension in the shallow waters of South San Francisco Bay: Mechanisms and potential implications for cohesive sediment transport, *Journal of Geophysical Research*, 115, 10.1029/2010jc006172, 2010.
- Briggs, K. B., Cartwright, G., Friedrichs, C. T., and Shivarudruppa, S.: Biogenic effects on cohesive sediment erodibility resulting from recurring seasonal hypoxia on the Louisiana shelf, *Continental Shelf Research*, 93, 17-26, <https://doi.org/10.1016/j.csr.2014.11.005>, 2015.
- Bryan, K.: A numerical method for the study of the circulation of the world ocean, *Journal of Computational Physics*, 4, 347-376, [https://doi.org/10.1016/0021-9991\(69\)90004-7](https://doi.org/10.1016/0021-9991(69)90004-7), 1969.
- Burchard, H., Schuttelaars, H. M., and Ralston, D. K.: Sediment Trapping in Estuaries, *Annual Review of Marine Science*, 10, 371-395, 10.1146/annurev-marine-010816-060535, 2018.

1294 Cao, L., Liu, J., Shi, X., He, W., and Chen, Z.: Source-to-sink processes of fluvial sediments in the
 1295 northern South China Sea: Constraints from river sediments in the coastal region of South China,
 1296 Journal of Asian Earth Sciences, 185, 104020, 10.1016/j.jseaes.2019.104020, 2019.

1297 Cao, Z., Ren, J., Deng, Z., Ye, L., and Wu, J.: Small-scale spatial variability in erosion threshold and
 1298 bedform for cohesive sediment measured by 3D Sonar, Journal of Hydrology, 650, 132513,
 1299 10.1016/j.jhydrol.2024.132513, 2025.

1300 Caruso, M. J., Gawarkiewicz, G. G., and Beardsley, R. C.: Interannual variability of the Kuroshio
 1301 intrusion in the South China Sea, Journal of Oceanography, 62, 559-575,
 1302 10.1007/s10872-006-0076-0, 2006.

1303 Chapman, D. C.: Numerical Treatment of Cross-Shelf Open Boundaries in a Barotropic Coastal Ocean
 1304 Model, Journal of Physical Oceanography, 15, 1060-1075, 1985.

1305 Charnock, H.: Wind stress on a water surface, Quarterly Journal of the Royal Meteorological Society,
 1306 81, 639-640, <https://doi.org/10.1002/qj.49708135027>, 1955.

1307 Chassignet, E. P., Hurlburt, H. E., Smedstad, O. M., Halliwell, G. R., Hogan, P. J., Wallcraft, A. J.,
 1308 Baraille, R., and Bleck, R.: The HYCOM (HYbrid Coordinate Ocean Model) data assimilative
 1309 system, Journal of Marine Systems, 65, 60-83, <https://doi.org/10.1016/j.jmarsys.2005.09.016>,
 1310 2007.

1311 Chen, S.-N., Geyer, W. R., Sherwood, C. R., and Ralston, D. K.: Sediment transport and deposition on
 1312 a river-dominated tidal flat: An idealized model study, Journal of Geophysical Research, 115,
 1313 10.1029/2010jc006248, 2010.

1314 Chen, Y., Deng, B., Saito, Y., Wang, Z., Yang, X., and Wu, J.: Pearl River sediment dispersal over its
 1315 associated delta–estuary–shelf system during the Holocene, Sedimentology, 70, 2331-2354,
 1316 10.1111/sed.13123, 2023.

1317 Chen, Z., Pan, J., and Jiang, Y.: Role of pulsed winds on detachment of low salinity water from the
 1318 Pearl River Plume Upwelling and mixing processes, Journal of Geophysical Research: Oceans,
 1319 121, 2769-2788, 10.1002/2015JC011337, 2016.

1320 Chen, Z., Gong, W., Cai, H., Chen, Y., and Zhang, H.: Dispersal of the Pearl River plume over
 1321 continental shelf in summer, Estuarine, Coastal and Shelf Science, 194, 252-262,
 1322 10.1016/j.ecss.2017.06.025, 2017a.

- 1323 Chen, Z., Pan, J., Jiang, Y., and Lin, H.: Far-reaching transport of Pearl River plume water by
1324 upwelling jet in the northeastern South China Sea, *Journal of Marine Systems*, 173, 60-69,
1325 10.1016/j.jmarsys.2017.04.008, 2017b.
- 1326 Cheng, P., Li, M., and Li, Y.: Generation of an estuarine sediment plume by a tropical storm, *Journal of*
1327 *Geophysical Research: Oceans*, 118, 856-868, 10.1002/jgrc.20070, 2013.
- 1328 Choi, S. M., Seo, J. Y., and Ha, H. K.: Contribution of local erosion enhanced by winds to sediment
1329 transport in intertidal flat, *Marine Geology*, 465, 107171, 10.1016/j.margeo.2023.107171, 2023.
- 1330 Church, J. A., and White, N. J.: A 20th century acceleration in global sea-level rise, *Geophysical*
1331 *Research Letters*, 33, n/a-n/a, 10.1029/2005gl024826, 2006.
- 1332 Cui, Y., Wu, J., Ren, J., and Xu, J.: Physical dynamics structures and oxygen budget of summer
1333 hypoxia in the Pearl River Estuary, *Limnology and Oceanography*, 64, 131-148,
1334 10.1002/lno.11025, 2018.
- 1335 Cui, Y., Wu, J., Tan, E., and Kao, S. J.: Role of Particle Resuspension in Maintaining Hypoxic Level in
1336 the Pearl River Estuary, *Journal of Geophysical Research: Oceans*, 127, 10.1029/2021jc018166,
1337 2022.
- 1338 Dai, S. B., Yang, S. L., and Cai, A. M.: Impacts of dams on the sediment flux of the Pearl River,
1339 southern China, *Catena*, 76, 36-43, 10.1016/j.catena.2008.08.004, 2008.
- 1340 Dalyander, P. S., Butman, B., Sherwood, C. R., Signell, R. P., and Wilkin, J. L.: Characterizing wave-
1341 and current- induced bottom shear stress: U.S. middle Atlantic continental shelf, *Continental Shelf*
1342 *Research*, 52, 73-86, <https://doi.org/10.1016/j.csr.2012.10.012>, 2013.
- 1343 Deng, Y., Liu, Z., Zu, T., Hu, J., Gan, J., Lin, Y., Li, Z., Quan, Q., and Cai, Z.: Climatic Controls on the
1344 Interannual Variability of Shelf Circulation in the Northern South China Sea, *Journal of*
1345 *Geophysical Research: Oceans*, 127, e2022JC018419, <https://doi.org/10.1029/2022JC018419>,
1346 2022.
- 1347 Dong, H., Jia, L., He, Z., Yu, M., and Shi, Y.: Application of parameters and paradigms of the erosion
1348 and deposition for cohesive sediment transport modelling in the Lingdingyang Estuary, China,
1349 *Applied Ocean Research*, 94, 101999, 10.1016/j.apor.2019.101999, 2020.
- 1350 Dong, L., Su, J., Wong, L., Cao, Z., and Chen, J.: Seasonal variation and dynamics of the Pearl River
1351 plume, *Continental Shelf Research*, 24, 1761-1777, 10.1016/j.csr.2004.06.006, 2004.

1352 Egbert, G. D., and Erofeeva, S. Y.: Efficient inverse Modeling of barotropic ocean tides, Journal of
 1353 Atmospheric and Oceanic Technology, 19, 183-204, Doi
 1354 10.1175/1520-0426(2002)019<0183:Eimobo>2.0.Co;2, 2002.
 1355 Eidam, E. F., Nittrouer, C. A., Ogston, A. S., DeMaster, D. J., Liu, J. P., Nguyen, T. T., and Nguyen, T.
 1356 N.: Dynamic controls on shallow clinoform geometry: Mekong Delta, Vietnam, Continental Shelf
 1357 Research, 147, 165-181, 10.1016/j.csr.2017.06.001, 2017.
 1358 Fairall, C. W., Bradley, E. F., Rogers, D. P., Edson, J. B., and Young, G. S.: Bulk parameterization of
 1359 air-sea fluxes for Tropical Ocean-Global Atmosphere Coupled-Ocean Atmosphere Response
 1360 Experiment, Journal of Geophysical Research: Oceans, 101, 3747-3764,
 1361 <https://doi.org/10.1029/95JC03205>, 1996.
 1362 Flather, R. A.: A tidal model of the north-west European continental shelf, Memoires Societe Royale
 1363 des Sciences de Liege, 10, 141-164, 1976.
 1364 Fox, J. M., Hill, P. S., Milligan, T. G., Ogston, A. S., and Boldrin, A.: Flocculation in the waters of the
 1365 Po River prodelta, Continental Shelf Research, 24, 1699-1715,
 1366 <https://doi.org/10.1016/j.csr.2004.05.009>, 2004.
 1367 Gan, J., Cheung, A., Guo, X., and Li, L.: Intensified upwelling over a widened shelf in the northeastern
 1368 South China Sea, Journal of Geophysical Research, 114, 10.1029/2007jc004660, 2009.
 1369 Gan, J., San Ho, H., and Liang, L.: Dynamics of Intensified Downwelling Circulation over a Widened
 1370 Shelf in the Northeastern South China Sea, Journal of Physical Oceanography, 43, 80-94,
 1371 10.1175/jpo-d-12-02.1, 2013.
 1372 Gao, S., and Collins, M. B.: Holocene sedimentary systems on continental shelves, Marine Geology,
 1373 352, 268-294, 10.1016/j.margeo.2014.03.021, 2014.
 1374 Gao, X., Chen, S., Xie, X., Long, A., and Ma, F.: Non-aromatic hydrocarbons in surface sediments near
 1375 the Pearl River estuary in the South China Sea, Environmental Pollution, 148, 40-47,
 1376 10.1016/j.envpol.2006.11.001, 2007.
 1377 Gao, X., Arthur Chen, C.-T., Wang, G., Xue, Q., Tang, C., and Chen, S.: Environmental status of Daya
 1378 Bay surface sediments inferred from a sequential extraction technique, Estuarine, Coastal and
 1379 Shelf Science, 86, 369-378, 10.1016/j.ecss.2009.10.012, 2010.
 1380 Ge, Q., Liu, J. P., Xue, Z., and Chu, F.: Dispersal of the Zhujiang River (Pearl River) derived sediment

1381 in the Holocene, *Acta Oceanologica Sinica*, 33, 1-9, 10.1007/s13131-014-0407-8, 2014.

1382 Ge, Q., Xue, Z., Yao, Z., Zang, Z., and Chu, F.: Anti-phase relationship between the East Asian winter
 1383 monsoon and summer monsoon during the Holocene?, *Journal of Ocean University of China*, 16,
 1384 175-183, 10.1007/s11802-017-3098-x, 2017.

1385 Ge, Q., Xu, D., Ye, L., Yang, K., and Yao, Z.: Linking Monsoon Activity with River-Derived Sediments
 1386 Deposition in the Northern South China Sea, *Journal of Ocean University of China*, 18, 1098-1104,
 1387 10.1007/s11802-019-4155-4, 2019.

1388 Georgiou, I. Y., FitzGerald, D. M., Sakib, M. M., Messina, F., Kulp, M. A., and Miner, M. D.: Storm
 1389 Dynamics Control Sedimentation and Shelf-Bay-Marsh Sediment Exchange Along the Louisiana
 1390 Coast, *Geophysical Research Letters*, 51, e2024GL111344,
 1391 <https://doi.org/10.1029/2024GL111344>, 2024.

1392 Geyer, W. R., Hill, P. S., and Kineke, G. C.: The transport, transformation and dispersal of sediment by
 1393 buoyant coastal flows, *Continental Shelf Research*, 24, 927-949, 10.1016/j.csr.2004.02.006, 2004.

1394 Gong, W., Chen, Y., Zhang, H., and Chen, Z.: Effects of Wave–Current Interaction on Salt Intrusion
 1395 During a Typhoon Event in a Highly Stratified Estuary, *Estuaries and Coasts*, 41, 1904-1923,
 1396 10.1007/s12237-018-0393-8, 2018a.

1397 Gong, W., Lin, Z., Chen, Y., Chen, Z., Shen, J., and Zhang, H.: Effect of waves on the dispersal of the
 1398 Pearl River plume in winter, *Journal of Marine Systems*, 186, 47-67,
 1399 10.1016/j.jmarsys.2018.05.003, 2018b.

1400 Haidvogel, D. B., Arango, H., Budgell, W. P., Cornuelle, B. D., Curchitser, E., Di, L. E., Fennel, K.,
 1401 Geyer, W. R., Hermann, A. J., Lanerolle, L., Levin, J., McWilliams, J. C., Miller, A. J., Moore, A.
 1402 M., Powell, T. M., Shchepetkin, A. F., Sherwood, C. R., Signell, R. P., Warner, J. C., and Wilkin, J.:
 1403 Ocean forecasting in terrain-following coordinates: Formulation and skill assessment of the
 1404 Regional Ocean Modeling System, *Journal of Computational Physics*, 227, 3595-3624,
 1405 10.1016/j.jcp.2007.06.016, 2008.

1406 Hanebuth, T. J. J., Lantzsch, H., and Nizou, J.: Mud depocenters on continental shelves—appearance,
 1407 initiation times, and growth dynamics, *Geo-Marine Letters*, 35, 487-503,
 1408 10.1007/s00367-015-0422-6, 2015.

1409 Harff, J., Leipe, T., and Zhou, D.: Pearl River Estuary related sediments as response to Holocene

1410 climate change and anthropogenic impact (PECAI), *Journal of Marine Systems*, 82, S1-S2,
1411 10.1016/j.jmarsys.2010.02.008, 2010.

1412 Harris, C. K., Traykovski, P. A., and Geyer, W. R.: Flood dispersal and deposition by near-bed
1413 gravitational sediment flows and oceanographic transport: A numerical modeling study of the Eel
1414 River shelf, northern California, *Journal of Geophysical Research: Oceans*, 110,
1415 10.1029/2004jc002727, 2005.

1416 Harris, C. K., Sherwood, C. R., Signell, R. P., Bever, A. J., and Warner, J. C.: Sediment dispersal in the
1417 northwestern Adriatic Sea, *Journal of Geophysical Research*, 113, 10.1029/2006jc003868, 2008.

1418 Hong, B., Liu, Z., Shen, J., Wu, H., Gong, W., Xu, H., and Wang, D.: Potential physical impacts of
1419 sea-level rise on the Pearl River Estuary, China, *Journal of Marine Systems*, 201, 103245,
1420 10.1016/j.jmarsys.2019.103245, 2020.

1421 Hu, J., Li, S., and Geng, B.: Modeling the mass flux budgets of water and suspended sediments for the
1422 river network and estuary in the Pearl River Delta, China, *Journal of Marine Systems*, 88, 252-266,
1423 10.1016/j.jmarsys.2011.05.002, 2011.

1424 Hu, S., Li, Y., Hu, P., Zhang, H., Zhang, G., and Gong, W.: The Impacts of Far-Field Typhoon-
1425 Generated Coastal Trapped Waves on the Hydrodynamics in the Northern South China Sea: A
1426 Case Study of Typhoon In - Fa, *Journal of Geophysical Research: Oceans*, 129,
1427 10.1029/2024jc021359, 2024.

1428 Huang, D., Du, J., Deng, B., and Zhang, J.: Distribution patterns of particle-reactive radionuclides in
1429 sediments off eastern Hainan Island, China: Implications for source and transport pathways,
1430 *Continental Shelf Research*, 57, 10-17, 10.1016/j.csr.2012.04.019, 2013.

1431 Jacob, R., Larson, J., and Ong, E.: $M \times N$ Communication and Parallel Interpolation in Community
1432 Climate System Model Version 3 Using the Model Coupling Toolkit, *IJHPCA*, 19, 293-307,
1433 10.1177/1094342005056116, 2005.

1434 Kirby, M. F., Devoy, B., Law, R. J., Ward, A., and Aldridge, J.: The use of a bioassay based approach to
1435 the hazard/risk assessment of cargo derived toxicity during shipping accidents: a case study--the
1436 MSC Napoli, *Marine Pollution Bulletin*, 56, 781-786, 10.1016/j.marpolbul.2008.01.006, 2008.

1437 Krige, D. G.: A Statistical Approach to Some Basic Mine Valuation Problems on the Witwatersrand,

1438 Chemical, Metallurgical and Mining Society of South Africa, 1951.

1439 Kuehl, S. A., Alexander, C. R., Blair, N. E., Harris, C. K., Marsaglia, K. M., Ogston, A. S., Orpin, A. R.,
 1440 Roering, J. J., Bever, A. J., Bilderback, E. L., Carter, L., Cerovski-Darriau, C., Childress, L. B.,
 1441 Reide Corbett, D., Hale, R. P., Leithold, E. L., Litchfield, N., Moriarty, J. M., Page, M. J., Pierce,
 1442 L. E. R., Upton, P., and Walsh, J. P.: A source-to-sink perspective of the Waipaoa River margin,
 1443 Earth-Science Reviews, 153, 301-334, 10.1016/j.earscirev.2015.10.001, 2016.

1444 Kumar, N., Voulgaris, G., Warner, J. C., and Olabarrieta, M.: Implementation of the vortex force
 1445 formalism in the coupled ocean-atmosphere-wave-sediment transport (COAWST) modeling
 1446 system for inner shelf and surf zone applications, Ocean Modelling, 47, 65-95,
 1447 10.1016/j.ocemod.2012.01.003, 2012.

1448 LaRowe, D. E., Arndt, S., Bradley, J. A., Estes, E. R., Hoarfrost, A., Lang, S. Q., Lloyd, K. G.,
 1449 Mahmoudi, N., Orsi, W. D., Shah Walter, S. R., Steen, A. D., and Zhao, R.: The fate of organic
 1450 carbon in marine sediments - New insights from recent data and analysis, Earth-Science Reviews,
 1451 204, 103146, 10.1016/j.earscirev.2020.103146, 2020.

1452 Larson, J., Jacob, R., and Ong, E.: The Model Coupling Toolkit: A New Fortran90 Toolkit for Building
 1453 Multiphysics Parallel Coupled Models, IJHPCA, 19, 277-292, 2005.

1454 Li, J., Li, M., and Xie, L.: Observations of near-inertial oscillations trapped at inclined front on
 1455 continental shelf of the northwestern South China Sea, EGU sphere, 2024, 1-25,
 1456 10.5194/egusphere-2024-3909, 2024a.

1457 Li, X., Chrysagi, E., Klingbeil, K., and Burchard, H.: Impact of Islands on Tidally Dominated River
 1458 Plumes: A High-Resolution Modeling Study, Journal of Geophysical Research: Oceans, 129,
 1459 e2023JC020272, <https://doi.org/10.1029/2023JC020272>, 2024b.

1460 Lin, S., Niu, J., Liu, G., Wei, X., and Cai, S.: Variations of suspended sediment transport caused by
 1461 changes in shoreline and bathymetry in the Zhujiang (Pearl) River Estuary in the wet season, Acta
 1462 Oceanologica Sinica, 41, 54-73, 10.1007/s13131-022-2017-1, 2022.

1463 Lin, W., Feng, Y., Yu, K., Lan, W., Wang, Y., Mo, Z., Ning, Q., Feng, L., He, X., and Huang, Y.:
 1464 Long-lived radionuclides in marine sediments from the Beibu Gulf, South China Sea: Spatial
 1465 distribution, controlling factors, and proxy for transport pathway, Marine Geology, 424, 106157,
 1466 10.1016/j.margeo.2020.106157, 2020.

1467 Liu, G., and Cai, S.: Modeling of suspended sediment by coupled wave-current model in the Zhujiang
1468 (Pearl) River Estuary, *Acta Oceanologica Sinica*, 38, 22-35, 10.1007/s13131-019-1455-3, 2019.

1469 Liu, H., Ye, L., Zhou, W., and Wu, J.: Salt-wedge intrusion-retreat cycle induced sediment flocculation
1470 dynamics in bottom boundary layer (BBL) of a micro-tidal estuary, *Marine Geology*, 466, 107175,
1471 10.1016/j.margeo.2023.107175, 2023.

1472 Liu, J. P., Xue, Z., Ross, K., Yang, Z., and Gao, S.: Fate of Sediments Delivered to the Sea by Asian
1473 Large Rivers: Long-Distance Transport and Formation of Remote Alongshore Clinothems,
1474 *Sediment. Rec.*, 7, 10.2110/sedred.2009.4.4, 2009.

1475 Liu, N., Geng, B., Xue, H., Xiu, P., Wang, Q., and Wang, D.: Interannual Variability of Shelf and Slope
1476 Circulations in the Northern South China Sea, *Journal of Ocean University of China*, 19,
1477 1005-1016, 10.1007/s11802-020-4446-9, 2020.

1478 Liu, Y., Gao, S., Wang, Y. P., Yang, Y., Long, J., Zhang, Y., and Wu, X.: Distal mud deposits associated
1479 with the Pearl River over the northwestern continental shelf of the South China Sea, *Marine
1480 Geology*, 347, 43-57, 10.1016/j.margeo.2013.10.012, 2014.

1481 Liu, Z., Zhao, Y., Colin, C., Stattegger, K., Wiesner, M. G., Huh, C.-A., Zhang, Y., Li, X.,
1482 Sompongchaiyakul, P., You, C.-F., Huang, C.-Y., Liu, J. T., Siringan, F. P., Le, K. P., Sathiamurthy,
1483 E., Hantoro, W. S., Liu, J., Tuo, S., Zhao, S., Zhou, S., He, Z., Wang, Y., Bunsomboonsakul, S.,
1484 and Li, Y.: Source-to-sink transport processes of fluvial sediments in the South China Sea,
1485 *Earth-Science Reviews*, 153, 238-273, 10.1016/j.earscirev.2015.08.005, 2016.

1486 Lu, X., Wang, Z., Guo, X., Gu, Y., Liang, W., and Liu, L.: Impacts of metal contamination and
1487 eutrophication on dinoflagellate cyst assemblages along the Guangdong coast of southern China,
1488 *Marine Pollution Bulletin*, 120, 239-249, 10.1016/j.marpolbul.2017.05.032, 2017.

1489 Ma, C., Zhao, J., Ai, B., Sun, S., and Yang, Z.: Machine Learning Based Long-Term Water Quality in
1490 the Turbid Pearl River Estuary, China, *Journal of Geophysical Research: Oceans*, 127,
1491 10.1029/2021jc018017, 2022.

1492 Ma, M., Zhang, W., Chen, W., Deng, J., and Schrum, C.: Impacts of morphological change and
1493 sea-level rise on stratification in the Pearl River Estuary, *Frontiers in Marine Science*, Volume 10 -
1494 2023, 10.3389/fmars.2023.1072080, 2023.

1495 Ma, M., Porz, L., Schrum, C., and Zhang, W.: Physical mechanisms, dynamics and interconnections of

multiple estuarine turbidity maximum in the Pearl River estuary, *Frontiers in Marine Science*, Volume 11 - 2024, 10.3389/fmars.2024.1385382, 2024.

Ma, Y., Friedrichs, C. T., Harris, C. K., and Wright, L. D.: Deposition by seasonal wave- and current-supported sediment gravity flows interacting with spatially varying bathymetry: Waiapu shelf, New Zealand, *Marine Geology*, 275, 199-211, <https://doi.org/10.1016/j.margeo.2010.06.001>, 2010.

Madsen, O. S.: Spectral Wave-Current Bottom Boundary Layer Flows, *Coastal Engineering* 1, 384-398, 1994.

Mao, Q., Shi, P., Yin, K., Gan, J., and Qi, Y.: Tides and tidal currents in the Pearl River Estuary, *Continental Shelf Research*, 24, 1797-1808, 10.1016/j.csr.2004.06.008, 2004.

Marshall, J., Adcroft, A., Hill, C., Perelman, L., and Heisey, C.: A finite-volume, incompressible Navier Stokes model for studies of the ocean on parallel computers, *Journal of Geophysical Research: Oceans*, 102, 5753-5766, <https://doi.org/10.1029/96JC02775>, 1997a.

Marshall, J., Hill, C., Perelman, L., and Adcroft, A.: Hydrostatic, quasi-hydrostatic, and nonhydrostatic ocean modeling, *Journal of Geophysical Research: Oceans*, 102, 5733-5752, <https://doi.org/10.1029/96JC02776>, 1997b.

McKee, B. A., Aller, R. C., Allison, M. A., Bianchi, T. S., and Kineke, G. C.: Transport and transformation of dissolved and particulate materials on continental margins influenced by major rivers: benthic boundary layer and seabed processes, *Continental Shelf Research*, 24, 899-926, <https://doi.org/10.1016/j.csr.2004.02.009>, 2004.

McWilliams, J. C., Restrepo, J. M., and Lane, E. M.: An asymptotic theory for the interaction of waves and currents in coastal waters, *Journal of Fluid Mechanics*, 511, 135-178, 10.1017/s0022112004009358, 2004.

Meade, R. H.: Landward Transport of Bottom Sediments in Estuaries of the Atlantic Coastal Plain, *Journal of Sedimentary Petrology*, 39, 222-234, 1969.

Milliman, J., and Farnsworth, K. L.: River discharge to the coastal ocean: A global synthesis. UK: Cambridge University Press; ISBN 978-0-521-87987-3, 2011.

Ministry of Water Resources of the PRC. 2022. *Bulletin of River Sediment in China*. <http://www.mwr.gov.cn/sj/#tjgb>.

1525 Moriarty, J. M., Harris, C. K., Friedrichs, M. A. M., Fennel, K., and Xu, K.: Impact of Seabed
 1526 Resuspension on Oxygen and Nitrogen Dynamics in the Northern Gulf of Mexico: A Numerical
 1527 Modeling Study, *Journal of Geophysical Research: Oceans*, 123, 7237-7263,
 1528 <https://doi.org/10.1029/2018JC013950>, 2018.
 1529 Nan, F., Xue, H., and Yu, F.: Kuroshio intrusion into the South China Sea: A review, *Progress in*
 1530 *Oceanography*, 137, 314-333, <https://doi.org/10.1016/j.pocean.2014.05.012>, 2015.
 1531 Ning, L., and Qian, Y.: Interdecadal change in extreme precipitation over South China and its
 1532 mechanism, *Advances in Atmospheric Sciences*, 26, 109-118, 10.1007/s00376-009-0109-x, 2009.
 1533 Nittrouer, C. A., and Wright, L. D.: Transport of particles across continental shelves, *Reviews of*
 1534 *Geophysics*, 32, 85-113, <https://doi.org/10.1029/93RG02603>, 1994.
 1535 Orlanski, I.: A simple boundary condition for unbounded hyperbolic flows, *Journal of Computational*
 1536 *Physics*, 21, 251-269, 10.1016/0021-9991(76)90023-1, 1976.
 1537 Ou, S., Zhang, H., and Wang, D.: Dynamics of the buoyant plume off the Pearl River Estuary in
 1538 summer, *Environmental Fluid Mechanics*, 9, 471-492, 10.1007/s10652-009-9146-3, 2009.
 1539 Ralston, D. K., Geyer, W. R., and Warner, J. C.: Bathymetric controls on sediment transport in the
 1540 Hudson River estuary: Lateral asymmetry and frontal trapping, *Journal of Geophysical Research:*
 1541 *Oceans*, 117, 10.1029/2012jc008124, 2012.
 1542 Ralston, D. K., and Geyer, W. R.: Sediment Transport Time Scales and Trapping Efficiency in a Tidal
 1543 River, *Journal of Geophysical Research: Earth Surface*, 122, 2042-2063, 10.1002/2017jfr004337,
 1544 2017.
 1545 Raymond, W. H., and Kuo, H. L.: A radiation boundary condition for multi-dimensional flows,
 1546 *Quarterly Journal of the Royal Meteorological Society*, 110, 535-551, 1984.
 1547 Repasch, M., Scheingross, J. S., Hovius, N., Lupker, M., Wittmann, H., Haghipour, N., Gröcke, D. R.,
 1548 Orfeo, O., Eglinton, T. I., and Sachse, D.: Fluvial organic carbon cycling regulated by sediment
 1549 transit time and mineral protection, *Nature Geoscience*, 14, 842-848,
 1550 10.1038/s41561-021-00845-7, 2021.
 1551 Saha, S., Moorthi, S., Wu, X., Wang, J., Nadiga, S., Tripp, P., Behringer, D., Hou, Y.-T., Chuang, H.-y.,
 1552 Iredell, M., Ek, M., Meng, J., Yang, R., Mendez, M. P., van den Dool, H., Zhang, Q., Wang, W.,
 1553 Chen, M., and Becker, E.: The NCEP Climate Forecast System Version 2, *Journal of Climate*, 27,

2185-2208, <https://doi.org/10.1175/JCLI-D-12-00823.1>, 2014.

Sanford, L. P.: Wave-forced resuspension of upper Chesapeake Bay muds, *Estuaries*, 17, 148-165, 1994.

Shchepetkin, A. F., and McWilliams, J. C.: The regional oceanic modeling system (ROMS): a split-explicit, free-surface, topography-following-coordinate oceanic model, *Ocean Modelling*, 9, 347-404, <https://doi.org/10.1016/j.ocemod.2004.08.002>, 2005.

Shepard, F. P.: Nomenclature Based on Sand-silt-clay Ratios, *Journal of Sedimentary Research*, 24, 151-158, 1954.

Sherwood, C. R., Aretxabaleta, A. L., Harris, C. K., Rinehimer, J. P., Verney, R., and Ferré, B.: Cohesive and mixed sediment in the Regional Ocean Modeling System (ROMS v3.6) implemented in the Coupled Ocean–Atmosphere–Wave–Sediment Transport Modeling System (COAWST r1234), *Geoscientific Model Development*, 11, 1849-1871, [10.5194/gmd-11-1849-2018](https://doi.org/10.5194/gmd-11-1849-2018), 2018.

Shi, M., Chen, C., Xu, Q., Lin, H., Liu, G., Wang, H., Wang, F., and Yan, J.: The Role of Qiongzhou Strait in the Seasonal Variation of the South China Sea Circulation, *Journal of Physical Oceanography*, 32, 103-121, 2002.

Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Wang, W., and Powers, J. G.: A Description of the Advanced Research WRF Version 2, 2005,

Smagorinsky, J.: GENERAL CIRCULATION EXPERIMENTS WITH THE PRIMITIVE EQUATIONS, *Monthly Weather Review*, 91, 99-164, [10.1175/1520-0493\(1963\)091<0099:gcewtp>2.3.co;2](https://doi.org/10.1175/1520-0493(1963)091<0099:gcewtp>2.3.co;2), 1963.

Song, Y., and Haidvogel, D.: A Semi-implicit Ocean Circulation Model Using a Generalized Topography-Following Coordinate System, *Journal of Computational Physics*, 115, 228-244, <https://doi.org/10.1006/jcph.1994.1189>, 1994.

Ståhlberg, C., Bastviken, D., Svensson, B. H., and Rahm, L.: Mineralisation of organic matter in coastal sediments at different frequency and duration of resuspension, *Estuarine, Coastal and Shelf Science*, 70, 317-325, <https://doi.org/10.1016/j.ecss.2006.06.022>, 2006.

Su, J.: Overview of the South China Sea circulation and its influence on the coastal physical oceanography outside the Pearl River Estuary, *Continental Shelf Research*, 24, 1745-1760,

1583 10.1016/j.csr.2004.06.005, 2004.

1584 Sun, Z., Zhang, Z., Qiu, B., Zhang, X., Zhou, C., Huang, X., Zhao, W., and Tian, J.: Three-Dimensional
1585 Structure and Interannual Variability of the Kuroshio Loop Current in the Northeastern South
1586 China Sea, *Journal of Physical Oceanography*, 50, 2437-2455,
1587 <https://doi.org/10.1175/JPO-D-20-0058.1>, 2020.

1588 Tolman, H., Accensi, M., Alves, J.-H., Ardhuin, F., Barbariol, F., Benetazzo, A., Bennis, A.-C., Bidlot,
1589 J., Booij, N., Boutin, G., Campbell, T., Chalikov, D., Chawla, A., Cheng, S., Collins Iii, C., Filipot,
1590 J.-F., Foreman, M., Janssen, P., Leckler, F., and Westhuysen, A.: User manual and system
1591 documentation of WAVEWATCH III (R) version 5.16, 2016.

1592 Turner, A., and Millward, G. E.: Suspended Particles: Their Role in Estuarine Biogeochemical Cycles,
1593 *Estuarine, Coastal and Shelf Science*, 55, 857-883, 10.1006/ecss.2002.1033, 2002.

1594 van der Wegen, M., Dastgheib, A., Jaffe, B. E., and Roelvink, D.: Bed composition generation for
1595 morphodynamic modeling: case study of San Pablo Bay in California, USA, *Ocean Dynamics*, 61,
1596 173-186, 10.1007/s10236-010-0314-2, 2010.

1597 Walsh, J. P., and Nittrouer, C. A.: Understanding fine-grained river-sediment dispersal on continental
1598 margins, *Marine Geology*, 263, 34-45, 10.1016/j.margeo.2009.03.016, 2009.

1599 Wang, C., Li, W., Chen, S., Li, D., Wang, D., and Liu, J.: The spatial and temporal variation of total
1600 suspended solid concentration in Pearl River Estuary during 1987-2015 based on remote sensing,
1601 *Science of The Total Environment*, 618, 1125-1138, 10.1016/j.scitotenv.2017.09.196, 2018.

1602 Wang, C., Liu, Z., Harris, C. K., Wu, X., Wang, H., Bian, C., Bi, N., Duan, H., and Xu, J.: The Impact
1603 of Winter Storms on Sediment Transport Through a Narrow Strait, Bohai, China, *Journal of*
1604 *Geophysical Research: Oceans*, 125, e2020JC016069, <https://doi.org/10.1029/2020JC016069>,
1605 2020.

1606 Wang, S., Zhang, N., Chen, H., Li, L., and Yan, W.: The surface sediment types and their rare earth
1607 element characteristics from the continental shelf of the northern south China sea, *Continental*
1608 *Shelf Research*, 88, 185-202, 10.1016/j.csr.2014.08.005, 2014.

1609 Wang, S., Wu, S., Yan, W., Huang, W., Miao, L., Lu, J., Chen, Z., and Liu, F.: Rare metal elements in
1610 surface sediment from five bays on the northeastern coast of the South China Sea, *Environmental*
1611 *Earth Sciences*, 74, 4961-4971, 10.1007/s12665-015-4504-6, 2015.

1612 Wang, S., Li, J., Wu, S., Yan, W., Huang, W., Miao, L., and Chen, Z.: The distribution characteristics of
 1613 rare metal elements in surface sediments from four coastal bays on the northwestern South China
 1614 Sea, *Estuarine, Coastal and Shelf Science*, 169, 106-118, 10.1016/j.ecss.2015.12.001, 2016.

1615 Wang, Y., Wang, Y., Wan, X., Huang, C., Wang, R., Liu, X., Yi, J., and Zhang, Y.: Influence of the
 1616 Hanjiang River's Inlet Sediment Decrease on Modern Sedimentation in the Underwater Delta,
 1617 *Applied Sciences*, 13, 8039, 10.3390/app13148039, 2023.

1618 Warner, J. C., Sherwood, C. R., Arango, H. G., and Signell, R. P.: Performance of four turbulence
 1619 closure models implemented using a generic length scale method, *Ocean Modelling*, 8, 81-113,
 1620 10.1016/j.ocemod.2003.12.003, 2005.

1621 Warner, J. C., Sherwood, C. R., Signell, R. P., Harris, C. K., and Arango, H. G.: Development of a
 1622 three-dimensional, regional, coupled wave, current, and sediment-transport model, *Computers &*
 1623 *Geosciences*, 34, 1284-1306, 10.1016/j.cageo.2008.02.012, 2008.

1624 Warner, J. C., Armstrong, B., He, R., and Zambon, J. B.: Development of a Coupled
 1625 Ocean–Atmosphere–Wave–Sediment Transport (COAWST) Modeling System, *Ocean Modelling*,
 1626 35, 230-244, 10.1016/j.ocemod.2010.07.010, 2010.

1627 Warner, J. C., Schwab, W. C., List, J. H., Safak, I., Liste, M., and Baldwin, W.: Inner-shelf ocean
 1628 dynamics and seafloor morphologic changes during Hurricane Sandy, *Continental Shelf Research*,
 1629 138, 1-18, 10.1016/j.csr.2017.02.003, 2017.

1630 Weatherall, P., Marks, K. M., Jakobsson, M., Schmitt, T., Tani, S., Arndt, J. E., Rovere, M., Chayes, D.,
 1631 Ferrini, V., and Wigley, R.: A new digital bathymetric model of the world's oceans, *Earth and*
 1632 *Space Science*, 2, 331-345, <https://doi.org/10.1002/2015EA000107>, 2015.

1633 Wright, L. D., and Coleman, J. M.: Variations in Morphology of Major River Deltas as Functions of
 1634 Ocean Wave and River Discharge Regimes, *AAPG Bulletin*, 57, 370-398, 1973.

1635 Wright, L. D., and Nittrouer, C. A.: Dispersal of river sediments in coastal seas: Six contrasting cases,
 1636 *Estuaries*, 18, 494-508, 10.2307/1352367, 1995.

1637 Wu, C., Xing, W., Jie, R., Yun, B., Zhigang, H., Yiaping, L., Heyin, S., and Wenyan, Z.:
 1638 Morphodynamics of the rock-bound outlets of the Pearl River estuary, South China — A
 1639 preliminary study, *Journal of Marine Systems*, 82, S17-S27, 10.1016/j.jmarsys.2010.02.002, 2010.

1640 Wu, Z., Milliman, J. D., Zhao, D., Zhou, J., and Yao, C.: Recent geomorphic change in LingDing Bay,

1641 China, in response to economic and urban growth on the Pearl River Delta, Southern China,
 1642 Global and Planetary Change, 123, 1-12, 10.1016/j.gloplacha.2014.10.009, 2014.

1643 Wu, Z., Milliman, J. D., Zhao, D., Cao, Z., Zhou, J., and Zhou, C.: Geomorphologic changes in the
 1644 lower Pearl River Delta, 1850–2015, largely due to human activity, Geomorphology, 314, 42-54,
 1645 10.1016/j.geomorph.2018.05.001, 2018.

1646 Wu, Z. Y., Saito, Y., Zhao, D. N., Zhou, J. Q., Cao, Z. Y., Li, S. J., Shang, J. H., and Liang, Y. Y.:
 1647 Impact of human activities on subaqueous topographic change in Lingding Bay of the Pearl River
 1648 estuary, China, during 1955-2013, Scientific Reports, 6, 37742, 10.1038/srep37742, 2016.

1649 Xia, X. M., Li, Y., Yang, H., Wu, C. Y., Sing, T. H., and Pong, H. K.: Observations on the size and
 1650 settling velocity distributions of suspended sediment in the Pearl River Estuary, China,
 1651 Continental Shelf Research, 24, 1809-1826, 10.1016/j.csr.2004.06.009, 2004.

1652 Xu, K., Corbett, D. R., Walsh, J. P., Young, D., Briggs, K. B., Cartwright, G. M., Friedrichs, C. T.,
 1653 Harris, C. K., Mickey, R. C., and Mitra, S.: Seabed erodibility variations on the Louisiana
 1654 continental shelf before and after the 2011 Mississippi River flood, Estuarine, Coastal and Shelf
 1655 Science, 149, 283-293, <https://doi.org/10.1016/j.ecss.2014.09.002>, 2014.

1656 Xu, K., Mickey, R. C., Chen, Q., Harris, C. K., Hetland, R. D., Hu, K., and Wang, J.: Shelf sediment
 1657 transport during hurricanes Katrina and Rita, Computers & Geosciences, 90, 24-39,
 1658 <https://doi.org/10.1016/j.cageo.2015.10.009>, 2016.

1659 Xue, Z., He, R., Liu, J. P., and Warner, J. C.: Modeling transport and deposition of the Mekong River
 1660 sediment, Continental Shelf Research, 37, 66-78, 10.1016/j.csr.2012.02.010, 2012.

1661 Yang, B., Liu, S.-M., and Zhang, G.-L.: Geochemical characteristics of phosphorus in surface
 1662 sediments from the continental shelf region of the northern South China Sea, Marine Chemistry,
 1663 198, 44-55, 10.1016/j.marchem.2017.11.001, 2018.

1664 Yang, J., Wu, D., and Lin, X.: On the dynamics of the South China Sea Warm Current, Journal of
 1665 Geophysical Research: Oceans, 113, <https://doi.org/10.1029/2007JC004427>, 2008.

1666 Young, I. R., Zieger, S., and Babanin, A. V.: Global trends in wind speed and wave height, Science, 332,
 1667 451-455, 10.1126/science.1197219, 2011.

1668 Zang, Z., Xue, Z. G., Xu, K., Bentley, S. J., Chen, Q., D'Sa, E. J., and Ge, Q.: A Two Decadal
 1669 (1993–2012) Numerical Assessment of Sediment Dynamics in the Northern Gulf of Mexico,

1670 Water, 11, 938, 2019.

1671 Zeng, X., He, R., Xue, Z., Wang, H., Wang, Y., Yao, Z., Guan, W., and Warrillow, J.: River-derived
 1672 sediment suspension and transport in the Bohai, Yellow, and East China Seas: A preliminary
 1673 modeling study, *Continental Shelf Research*, 111, 112-125,
 1674 <https://doi.org/10.1016/j.csr.2015.08.015>, 2015.

1675 Zhan, W., Wu, J., Wei, X., Tang, S., and Zhan, H.: Spatio-temporal variation of the suspended sediment
 1676 concentration in the Pearl River Estuary observed by MODIS during 2003–2015, *Continental
 1677 Shelf Research*, 172, 22-32, 10.1016/j.csr.2018.11.007, 2019.

1678 Zhang, G., Cheng, W., Chen, L., Zhang, H., and Gong, W.: Transport of riverine sediment from
 1679 different outlets in the Pearl River Estuary during the wet season, *Marine Geology*, 415, 105957,
 1680 10.1016/j.margeo.2019.06.002, 2019.

1681 Zhang, G., Chen, Y., Cheng, W., Zhang, H., and Gong, W.: Wave Effects on Sediment Transport and
 1682 Entrapment in a Channel-Shoal Estuary: The Pearl River Estuary in the Dry Winter Season,
 1683 *Journal of Geophysical Research: Oceans*, 126, 10.1029/2020jc016905, 2021.

1684 Zhang, G., Hu, P., Hu, S., Zhang, H., and Gong, W.: Tidal effects on the dispersal and water age of the
 1685 plumes from eight outlets of the Pearl river during the wet summer, *Ocean & Coastal Management*,
 1686 266, 107704, <https://doi.org/10.1016/j.ocecoaman.2025.107704>, 2025.

1687 Zhang, J., Jiang, Q., Jeng, D., Zhang, C., Chen, X., and Wang, L.: Experimental Study on Mechanism
 1688 of Wave-Induced Liquefaction of Sand-Clay Seabed, *Journal of Marine Science and Engineering*,
 1689 8, 66, 10.3390/jmse8020066, 2020.

1690 Zhang, W., Wei, X., Zheng, J., Zhu, Y., and Zhang, Y.: Estimating suspended sediment loads in the
 1691 Pearl River Delta region using sediment rating curves, *Continental Shelf Research*, 38, 35-46,
 1692 10.1016/j.csr.2012.02.017, 2012.

1693 Zhang, W., Zheng, J., Ji, X., Hoitink, A. J. F., van der Vegt, M., and Zhu, Y.: Surficial sediment
 1694 distribution and the associated net sediment transport pattern retain-->in the Pearl River Estuary,
 1695 South China, *Continental Shelf Research*, 61-62, 41-51, 10.1016/j.csr.2013.04.011, 2013.

1696 Zhang, Y. J., Ye, F., Stanev, E. V., and Grashorn, S.: Seamless cross-scale modeling with SCHISM,
 1697 *Ocean Modelling*, 102, 64-81, <https://doi.org/10.1016/j.ocemod.2016.05.002>, 2016.

1698 Zhong, Y., Chen, Z., Li, L., Liu, J., Li, G., Zheng, X., Wang, S., and Mo, A.: Bottom water

1699 hydrodynamic provinces and transport patterns of the northern South China Sea: Evidence from
 1700 grain size of the terrigenous sediments, *Continental Shelf Research*, 140, 11-26,
 1701 10.1016/j.csr.2017.01.023, 2017.

1702 Zong, X., Cheng, X., Zhang, S., Lian, Q., Deng, F., and Chen, Z.: Tidal effects on dynamics and
 1703 freshwater transport of a medium-scale river plume with multiple outlets, *Ocean Modelling*, 188,
 1704 102338, <https://doi.org/10.1016/j.ocemod.2024.102338>, 2024.

1705 Zong, Y., Huang, K., Switzer, A., yu, F., and Yim, W.: An evolutionary model for the Holocene
 1706 formation of the Pearl River delta, China, *The Holocene*, 19, 129-142,
 1707 10.1177/0959683608098957, 2009.

1708