

Reviews and Syntheses: Trait-based approach to constrain controls on planktic foraminiferal ecology: key trade-offs and current knowledge gaps

Kirsty M. Edgar^{1,2}, Maria Grigoratou^{2*}, Fanny M. Monteiro^{3,4}, Ruby Barrett², Rui Ying^{2^} and Daniela N. Schmidt^{1,2}

¹School of Geography, Earth and Environmental Sciences, University of Birmingham, B15 2TT, UK

²School of Earth Sciences, University of Bristol, Wills Memorial Building, Queen's Road, Bristol, BS8 1RJ, UK

³BRIDGE, School of Geographical Sciences, University of Bristol, Bristol, BS8 1SS, UK

⁴[iC3, Department for Geosciences, The Arctic University of Norway, 9010 Tromsø, Norway](#)

^{*}Now at European Polar Board, A Working Lab Universum, Universums gränd 8, 907 36 Umeå, SWEDEN

[^]Now at School of Environmental Sciences, University of East Anglia, Norwich, NR4 7TJ, [UK](#)

Correspondence to: Kirsty M Edgar (k.m.edgar@bham.ac.uk)

Abstract. Planktic foraminifera are a major contributor to global marine inorganic carbon production. They leave abundant calcium carbonate shells on the seafloor~~sea floor~~, ~~which~~that serve as prime proxies for the physical and chemical attributes of past oceans. Despite ~~at~~their well-preserved fossil record and wide spread use in palaeoceanography, our understanding of their ecology ~~remains~~is limited due to their low ~~-~~standing stocks in the modern ~~surface~~-ocean and the challenges in culturing multiple generations under laboratory conditions, even after decades of data collection. This limitation affects our ability to ~~interpret~~use their fossil remains to describe past ecosystems and predict their responses to ~~ongoing~~modern environmental changes. Trait-based ecology ~~offers a powerful framework to~~can be particularly useful at characterising how and why foraminifera ~~might~~ interact with their environment. Here, we review the current state of knowledge ~~on~~f key planktic foraminifera ~~key~~-traits, including morphological, physiological, behavioural and life history traits. Most spinose taxa are carnivorous, host to dinoflagellate photosymbionts, and are abundant and diverse in oligotrophic environments. In contrast, non-spinose taxa are typically herbivorous and most common in high-productivity regions. We highlight the potential of trait modelling to generate ~~hypotheses~~-testable hypotheses in the field. ~~Integration of~~We propose that together, trait-based modelling ~~with~~, metabarcoding, ~~environmental~~ DNA, and enhanced standardised data collection ~~made openly available, released in open data, can~~will help fill critical gaps in our understanding of planktic foraminiferal ~~trait-based~~ ecology, and allow us to use foraminifera as a key model organism for addressing fundamental~~testing~~ ecological questions.

1 Introduction

Planktic foraminifera (heterotrophic marine protists, Fig. 1), along with phytoplankton coccolithophores and planktic snail pteropods, are the dominant calcifying plankton in the modern ocean (Deuser et al., 1981). ~~Among~~Of these, planktic foraminifera constitute 23-56 % of the global CaCO₃ flux from the top 100 m of surface waters and between 32-80 % of the

32 total deep-marine calcite budget, with their carbonate shells covering much of the seafloor (Neukermans et al., 2023). Thus,
33 these organisms have pronounced impacts on biogeochemical cycling, in particular the inorganic carbon cycle, by transferring
34 carbon from surface to deep waters, and over longer timescales, modulating ocean alkalinity, carbonate chemistry and
35 ultimately climate (Ridgwell and Zeebe, 2005). Moreover, in part due to their calcareous shells that can be preserved in marine
36 sediments for millions of years, planktic foraminifera have provide one of the best plankton-fossil records of any plankton
37 group (Aze et al., 2011; Fenton et al., 2021). This, makesing them a critical group for studying the influence of climate change
38 in the past and improving our understanding offer the future climate (Strack et al., 2024; Woodhouse et al., 2023; Fenton et
39 al., 2023). Their relative abundances, biometry, and a wealth of geochemical proxies based on shell chemistry are extensively
40 used to reconstruct past climate, including temperature and ocean chemistry (Kucera, 2007).




Figure 1. Selection of living and recent planktic foraminifera. (a) Assorted shells of planktic foraminifera from sea floor sediment (Image credit: Wilfried Rönfeld). (b) *Orbulina universa* with symbionts attached along thin radial spines (Fig. 2 of Topa et al., 2017). (c) *Orbulina universa* eating a small copepod (Image credit: Oscar Branson). (d) *Globorotalia truncatulinoides* in culture with pseudopodial network (a web-like structure formed of pseudopodia [see below] filaments, that interacts with the environment and provides physiological functions) extending from shell (<https://www.usgs.gov/media/images/live-foraminifera-globorotalia-truncatulinoides>). (e) *Neogloboquadrina dutertrei* suspended with pseudopodia (cytoplasmic projections that assist locomotion, feeding and other physiological functions) extending from shell (Image credit: Kate Darling). (f) *Hastigerinella digitata* with triradiate spines and bubble capsule (Fig. 1A of Hull et al., 2011).

Despite their importance in understanding past oceans, planktic foraminifera ecology has received relatively little attention from modern ecologists due to their small contribution to the total plankton biomass (Michaels et al., 1995; Buitenhuis et al., 2013) and the challenges associated with culturing them (Del Campo et al., 2024; Meilland et al., 2024). Developing a mechanistic understanding of the controls on planktic foraminifera diversity and distribution is essential to generating accurate predictions of how changing environmental conditions will impact ~~their planktic foraminifera~~ communities and, ultimately, biogeochemical cycles (Dutkiewicz et al., 2020).

An improved mechanistic understanding can be supported by ~~will~~ can contribute to ~~T~~ trait-based approaches (Dutkiewicz et al., 2020), ~~allow to which~~ describe how species interact with each other and their environment based on measurable organismal characteristics or traits (e.g. size, resource acquisition mode and defence). Rather than focusing on individual species, these approaches group organisms with similar traits into functional groups, simplifying the characterisation of highly diverse ecosystems like plankton communities, where millions of specimens from thousands of species interact (De Vargas et al., 2015). This approach effectively links individual-level traits to larger ecosystem processes, helping to explain patterns of biodiversity, species distributions and how communities respond to environmental changes. However, a key challenge is identifying the trade-offs between traits (i.e., costs and benefits) (e.g., Litchman et al., 2013; Barton et al., 2013; Violle et al., 2007; Westoby, 2024). Trade-offs arise when the optimisation of one trait occurs at the expense of another (Kjørboe et al., 2018). Overall, trait-based approaches offer a mechanistic yet computationally efficient means of explaining large-scale patterns of diversity and abundance across microbial, planktonic and nekton communities (e.g., Barton et al., 2013; Dutkiewicz et al., 2009; Follows et al., 2007; Grigoratou et al., 2019; Naidoo-Bagwell et al., 2024; Ying, 2024; Ward, 2013; Monteiro et al., 2016; Litchman et al., 2021).

Trait-based approaches are particularly a promising ~~tool~~ for making predictions beyond the sampling domain and ~~to testing~~ different (and often complex) hypotheses over longer timescales - both past and future (e.g., Barton et al., 2016; Grigoratou et al., 2022; Ying, 2024). This makes this approach ~~is~~ particularly valuable ~~promising~~ in macroevolutionary studies spanning over millions of years, where new traits evolve in response to changing environments and to facilitate adaptation to the novel climates and environments. By permitting focusing on, as it permits ‘species’ or functional groups ~~(i.e., species with similar~~

traits) with novel trait combinations, trait-based models can simulate ~~of traits to~~ emergent taxa as an environment evolves that may not be present in the modern ocean but could have existed in the past or evolve in the future. For instance, the model of Knoll and Follows (2016) ~~model~~ shows that how the rise of the mixotrophy trait in the Mesozoic increases ~~the~~ energy transfer efficiency upward to higher trophic levels in the food web. Similarly Using a similar trait framework processes, Gibbs et al. (2020) used trait-based models to explore ecological selectivity ~~in the aftermath of the Cretaceous-Paleogene mass extinction~~ in marine plankton following the Cretaceous-Paleogene mass extinction.

While trait-based models have significant potential, they have inherent limitations. Like any models, they are a highly simplified version of natural systems. They may simplify complex ecological interactions and environmental influences, and the expression and importance of traits may vary by environment impacting generalisation. Additionally, trade-offs between traits are not always well understood or quantified, as for example, the absence or presence of symbionts in foraminifera. The eco-evolutionary model used by Gibbs et al. (2020), for instance, resulted in modelled trait evolution at rates orders of magnitude faster than observed in the fossil dataset, highlighting the need for caution when interpreting these model results. Despite these caveats, trait-based approaches present an exciting opportunity to leverage the exceptional fossil record of foraminifera to test our understanding outside of the modern range of environmental conditions, assess the universal applicability of traits and trade-offs through time and ultimately improve our understanding of evolutionary processes.

Trait-based modelling has already provided novel insights into ~~the potential to vastly improve our understanding of the driver(s) of~~ planktic foraminifera ecology and fitness. ~~For example, the first~~ non-species specific or “trait-based general” planktic foraminifera ~~trait-based~~ modelling study by Grigoratou et al. (2019) highlighted ~~provided novel insights regarding~~ the cost and benefits of calcification, and the influence of resource competition among planktic foraminifera and other zooplankton. ~~This approach was expanded by~~ Ying et al. (2023) further expanded this approach to characterise the main ecogroups of planktonic foraminifera (Figs 1 and 2). In this study, we aim to ~~[1]~~ present a comprehensive review of our current knowledge of planktic foraminifera traits and trade-offs, ~~and [2]~~ identify critical key knowledge gaps ~~in our understanding, and propose future research directions to advance the application of trait-based ecology in foraminifera research to facilitate future studies.~~




Figure 2: Schematic representation of a trait-based ecosystem model incorporating planktic foraminifers. Key functional traits include size, calcification, spinosity and symbiosis, and are assumed to influence foraminiferal biotic-interactions with other components in the food web bythrough modifying maximum growth rate, mortality rate, feeding strategy, and other predator-prey interactions.

2. Planktic foraminifera traits and trade-offs

Planktic foraminifera have been predominantly studied by geologists for their paleoecology and evolution, and as proxy carriers for paleoenvironmental studies, while most biological studies until relatively recently, were from have been from seminal papers by Allan Bé, and co-workers in the 1970s and '80s. Classification of planktic foraminifera is primarily based on adult morphology, with ~45 morphologically distinct species or 'morphospecies' in the modern ocean (Brummer and Kučera, 2022; Schiebel and Hemleben, 2017), many but not all of which consist of cryptic species (i.e., organisms that look identical but represent distinct evolutionary lineages; Morard et al., 2024; Morard et al., 2016).

Planktic foraminifera spend their lives in the open surface-ocean, predominantly the upper ~200 m of the water column (Table 1), with very few individuals or species found below ~1 km water depth (Rebotim et al., 2019; Vincent, 1981), which results in their absence in shallow marine seas and on coastal shelves (Schiebel and Hemleben, 2017). Some taxa have unique ecologies, e.g. *Neogloboquadrina pachyderma* may spend parts of the year in sea ice (Dieckmann et al., 1991).

122 **Table 1 Modern planktic foraminiferal species list and associated key traits.** Species [are](#) listed [here](#) alphabetically [here](#) and correspond
123 to those listed in the ForCenS database of Siccha and Kucera (2017) and underpin the trait distribution maps shown in Fig. 3. Summary [of](#)
124 [ecologically](#) information from Schiebel and Hemleben (2017), Takagi et al. (2019) and Aze et al. (2011). Obligate symbiosis=essential for
125 optimal host fitness, and observed in the majority of individuals of a species. Facultative [symbiosis](#)=(not essential [for](#)to the host success and
126 [thus](#), only associated with ~~350~~ some individuals of the species) (Table 1; Takagi et al., 2019). Determining whether symbiosis is obligate or
127 facultative is challenging as photosymbiosis is a spectrum as proposed by Stoecker et al (2009) from non-symbiosis to robust symbiosis.
128 [*=bacterial endobionts *Synechococcus* reported by Bird et al. \(2017\) but unclear if symbionts.](#)
129

SPECIES	SPINOSITY	SYMBIONT TYPE	SYMBIONT RELATIONSHIP	DEPTH HABITAT
<i>Beella digitata</i>	spinose	none	none	sub-thermocline
<i>Berggrenia pumilio</i>	non-spinose	none	none	sub-thermocline
<i>Candeiina nitida</i>	non-spinose	chrysophyte	facultative	thermocline
<i>Dentigloborotalia anfracta</i>	non-spinose	none	none	surface
<i>Globigerina bulloides</i>	spinose	none	none	surface
<i>Globigerina falconensis</i>	spinose	chrysophyte	facultative	surface
<i>Globigerinella adamsi</i>	spinose	none	none	sub-thermocline
<i>Globigerinella calida</i>	spinose	chrysophyte	facultative	thermocline
<i>Globigerinella siphonifera</i>	spinose	chrysophyte	obligate	thermocline
<i>Globigerinita glutinata</i>	non-spinose	chrysophyte	facultative	surface
<i>Globigerinita minuta</i>	non-spinose	none	none	surface
<i>Globigerinita uvula</i>	non-spinose	dinoflagellate	obligate	surface
<i>Globigerinoides conglobatus</i>	spinose	dinoflagellate	obligate	surface
<i>Globigerinoides ruber</i>	spinose	dinoflagellate	obligate	surface
<i>Globigerinoides tenellus</i>	spinose	dinoflagellate	obligate	surface
<i>Globigerinoides white</i>	spinose	dinoflagellate	obligate	surface
<i>Globoconella inflata</i>	non-spinose	chrysophyte	facultative	thermocline
<i>Globoquadrina conglomerata</i>	non-spinose	none	none	surface
<i>Globorotalia cavernula</i>	non-spinose	none	none	sub-thermocline
<i>Globorotalia crassaformis</i>	non-spinose	none	none	thermocline
<i>Globorotalia hirsuta</i>	non-spinose	none	none	sub-thermocline
<i>Globorotalia menardii</i>	non-spinose	chrysophyte/prymnesiophyte	facultative	thermocline
<i>Globorotalia scitula</i>	non-spinose	none	none	sub-thermocline
<i>Globorotalia theyeri</i>	non-spinose	none	none	surface
<i>Globorotalia truncatulinoides</i>	non-spinose	none	none	sub-thermocline
<i>Globorotalia tumida</i>	non-spinose	none	none	thermocline
<i>Globorotalia unguolata</i>	non-spinose	none	none	thermocline
<i>Globorotaloides hexagonus</i>	spinose	none	none	sub-thermocline
<i>Globoturborotalita rubescens</i>	spinose	dinoflagellate	obligate	surface
<i>Hastigerina pelagica</i>	spinose	none	none	sub-thermocline
<i>Hastigerinella digitata</i>	spinose	none	none	sub-thermocline
<i>Neogloboquadrina dutertrei</i>	non-spinose	chrysophyte, pelagophyte	facultative	thermocline
<i>Neogloboquadrina incompta</i>	non-spinose	none	none	thermocline
<i>Neogloboquadrina pachyderma</i>	non-spinose	none	none	thermocline
<i>Orbulina universa</i>	spinose	dinoflagellate	obligate	thermocline
<i>Pulleniatina obliquiloculata</i>	non-spinose	chrysophyte	facultative	surface
<i>Sphaeroidinella dehiscens</i>	non-spinose	dinoflagellate	obligate	thermocline
<i>Tenuitella iota</i>	non-spinose	none	none	surface
<i>Trilobatus sacculifer</i>	spinose	dinoflagellate	obligate	surface
<i>Turborotalita humilis</i>	spinose	chrysophyte, haptophyte, dinoflagellate	obligate	surface
<i>Turborotalita quinqueloba</i>	spinose	none	none	surface

130
131

SPECIES	SPINOSITY	SYMBIONT TYPE	SYMBIONT RELATIONSHIP	DEPTH HABITAT
<i>Beella digitata</i>	Spinose	None	None	Sub-thermocline
<i>Berggrenia pumilio</i>	Non-spinose	None	None	Sub-thermocline
<i>Candeina nitida</i>	Non-spinose	Chrysophyte	Facultative	Thermocline
<i>Dentigloborotalia anfracta</i>	Non-spinose	None	None	Surface
<i>Globigerina bulloides</i>	Spinose	None*	None	Surface
<i>Globigerina falconensis</i>	Spinose	Chrysophyte	Facultative	Surface
<i>Globigerinella adamsi</i>	Spinose	None	None	Sub-thermocline
<i>Globigerinella calida</i>	Spinose	Chrysophyte	Facultative	Thermocline
<i>Globigerinella siphonifera</i>	Spinose	Chrysophyte	Obligate	Thermocline
<i>Globigerinita glutinata</i>	Non-spinose	Chrysophyte	Facultative	Surface
<i>Globigerinita minuta</i>	Non-spinose	None	None	Surface
<i>Globigerinita uvula</i>	Non-spinose	Dinoflagellate	Obligate	Surface
<i>Globigerinoides conglobatus</i>	Spinose	Dinoflagellate	Obligate	Surface
<i>Globigerinoides ruber</i>	Spinose	Dinoflagellate	Obligate	Surface
<i>Globigerinoides tenellus</i>	Spinose	Dinoflagellate	Obligate	Surface
<i>Globigerinoides white</i>	Spinose	Dinoflagellate	Obligate	Surface
<i>Globoconella inflata</i>	Non-spinose	Chrysophyte	Facultative	Thermocline
<i>Globobulimina conglomerata</i>	Non-spinose	None	None	Surface
<i>Globorotalia cavernula</i>	Non-spinose	None	None	Sub-thermocline
<i>Globorotalia crassaformis</i>	Non-spinose	None	None	Thermocline
<i>Globorotalia hirsuta</i>	Non-spinose	None	None	Sub-thermocline
<i>Globorotalia menardii</i>	Non-spinose	Chrysophyte/Prymnesiophyte	Facultative	Thermocline
<i>Globorotalia scitula</i>	Non-spinose	None	None	Sub-thermocline
<i>Globorotalia theyeri</i>	Non-spinose	None	None	Surface
<i>Globorotalia truncatulinoides</i>	Non-spinose	None	None	Sub-thermocline
<i>Globorotalia tumida</i>	Non-spinose	None	None	Thermocline
<i>Globorotalia unguolata</i>	Non-spinose	None	None	Thermocline
<i>Globorotaloides hexagonus</i>	Spinose	None	None	Sub-thermocline
<i>Globoturborotalita rubescens</i>	Spinose	Dinoflagellate	Obligate	Surface
<i>Hastigerina pelagica</i>	Spinose	None	None	Sub-thermocline
<i>Hastigerinella digitata</i>	Spinose	None	None	Sub-thermocline
<i>Neogloboquadrina dutertrei</i>	Non-spinose	Chrysophyte, Pelagophyte	Facultative	Thermocline
<i>Neogloboquadrina incompta</i>	Non-spinose	None	None	Thermocline
<i>Neogloboquadrina pachyderma</i>	Non-spinose	None	None	Thermocline
<i>Orbulina universa</i>	Spinose	Dinoflagellate	Obligate	Thermocline
<i>Pulleniatina obliquiloculata</i>	Non-spinose	Chrysophyte	Facultative	Surface
<i>Sphaeroidinella dehiscens</i>	Non-spinose	Dinoflagellate	Obligate	Thermocline
<i>Tenuitella iota</i>	Non-spinose	None	None	Surface
<i>Trilobatus sacculifer</i>	Spinose	Dinoflagellate	Obligate	Surface
<i>Turborotalita humilis</i>	Spinose	Chrysophyte, Haptophyte, Dinoflagellate	Obligate	Surface
<i>Turborotalita quinqueloba</i>	Spinose	None	None	Surface

Here we characterise planktic foraminiferal traits according to their type and function, following the approach of Litchman et al. (2013) (Table 2). We discuss each identified key traits, along with their individual impacts on foraminiferal fitness, as well as their main associated trade-offs. We recognise two levels of traits: (1) those that are common to all planktic foraminifera, referred to as ‘foraminiferal’ traits and (2) those that are specific to individual species or groups of species as ‘Group/Species-specific’ traits.

Table 2. Summary of planktic foraminifera traits and associated trade-offs described in Sections 2.1-2.3. Traits with ‘*’ are group/species specific traits, all other traits are common to all planktic foraminifera. Rhizopodia refers to the network of cytoplasmic strands around the shell that help to capture, transport, and break-up prey items and carry waste out of the cell.

<i>Trait Type</i>	<i>Trait</i>	<i>Cost(s)</i>	<i>Benefit(s)</i>
<i>Physiological /Morphological</i>	Calcification (shell)	- Energy investment (~10-60% of energy budget) - Negative buoyancy	- Protection from predation and/or the environment
	Calcification (spines)*	- Energetic investment (<60% of total energy budget) - Reduction in growth	- Tool for controlling active prey - Rhizopodial support - Increased surface area to volume - Increased protection from predation - Positive buoyancy
<i>Behavioural</i>	Photosymbiotic host*	- Light limitation - Influenced by symbiont availability & health	- Nutrient exchange for calcification - Favorable ambient pH for calcification - Food for terminal, sick or starved hosts - Food for reproduction - Energy conservation by not actively swimming - Reduced predator encounters
	Immotility	- Reliant on surrounding available resources - Inability to actively escape predation - Sensitive to environmental conditions	
	Vertical migration (diel or seasonal)	- Sensitive to predation and environmental conditions	- Energy gain
	Passive feeding	- Reliant on availability of surrounding resources - Low feeding efficiency	- Advantage of prey migration - Energy conservation by not actively searching for food - Reduced predator encounters
<i>Life History</i>	Reproduction	- Only reproduce once during their life cycle - Number of gametes produced depends on shell size	- Energy gain as not actively searching for mate - Reduced predator exposure

+

<i>Trait Type</i>	<i>Trait</i>	<i>Cost(s)</i>	<i>Benefit(s)</i>
<i>Physiological /Morphological</i>	Calcification (shell)	- Energy investment (~10-60% of energy budget) - Negative buoyancy	- Protection from predation and/or the environment
	Calcification (spines)	- Energy investment (<60% of total energy budget) - Reduction in growth	- Tool for controlling active prey - Rhizopodial support - Increased prey capture area - Increased protection from predation - Positive buoyancy
<i>Behavioural</i>	Photosymbiotic host	- Light limitation - Influenced by symbiont availability & health	- Nutrient exchange for calcification - Favorable ambient pH for calcification - Food for terminal, sick or starved hosts - Food for reproduction - Energy conservation by not actively swimming - Reduced predator encounters
	Immotility	- Reliant on surrounding available resources - Inability to actively escape predation - Sensitive to environmental conditions	
	Vertical migration (diel or seasonal)	- Sensitive to predation and environmental conditions	- Energy gain - Advantage of prey migration
	Passive feeding	- Reliant on availability of surrounding resources - Low feeding efficiency	- Energy conservation by not actively searching for food - Reduced predator encounters
<i>Life History</i>	Reproduction	- Dominantly sexual reproduction, once in life cycle - Number of gametes produced depends on shell size	- Energy gain as not actively searching for mate - Reduced predator exposure by not actively searching for mate

2.1 Physiological/Morphological traits

2.1.1 Body Size (foraminiferal trait)

Size is often considered as a master trait because it influences most relationships between organisms in an ecosystem, and how they respond to and influence their environment (Brown et al., 2004; Peters, 1983). Body size directly impacts physiological and ecological aspects like metabolic rates, e.g., growth, diet, predator-prey relationships, abundance, biomass and reproduction in organisms (e.g., Kiørboe et al., 2018; Mckinney, 1990). For instance, the primary mode of resource acquisition for organisms shifts from osmotrophy in the smallest single-celled organisms, to phototrophy, mixotrophy, and ultimately heterotrophy as body size increases (Andersen et al., 2016; Haldane, 1926). Body size also influences predator-prey relationships by impacting the range of prey sizes an organism can consume, with prey being typically smaller than the predator (Gaskell et al., 2019; Wirtz, 2012).

Planktic foraminifera grow their organic cell by adding multiple chambers in their calcite shell (Brummer et al., 1987; Caromel et al., 2016). Depending on the species, the shell size of adult individuals can vary from ~100 to 1500 µm (Schmidt et al., 2004). At the individual level, organisms typically reach their largest shell size under specific optimal environmental growth conditions (Chernihovsky et al., 2023; Schmidt et al., 2004), although some show no relationships between size and environments (Rillo et al., 2020). Temperature and suitable prey availability have been identified as the primary environmental factors influencing shell growth, followed by pH, salinity, and light conditions—particularly for species that maintain symbiotic relationships with algae. As a group, planktic foraminifera reach large sizes in the tropics, while smaller individuals dominate at the higher latitudes and in equatorial or upwelling regions (Schmidt et al., 2004). This size distribution contrasts with most zooplankton species, which exhibit their largest sizes in polar regions (Brandão et al., 2021; Horne et al., 2017). The size distribution of foraminifera may reflect a combination of factors including higher temperatures and thus, metabolic rates promoting growth, higher carbonate saturation and light intensity promoting calcification, more energy subsidy from algal photosymbionts, increased volume for prey encounter, or enhanced niche diversity through stratification (Schmidt et al., 2004; Burke et al., 2025). Shell size not only responds to environmental conditions but also directly reflects individual fitness and reproduction (Hemleben, 1989). Larger foraminifera have a greater amount of cytoplasm available to generate gametes and hence potentially higher reproductive success (Bé and Anderson, 1976; Weinkauff et al., 2022). In the open ocean, a (semi)lunar periodicity in shell fluxes (i.e. shells exported to the sea floor after death) of several species is interpreted as the presence of a circadian clock (Bijma et al., 1990; Jonkers and Kučera, 2015; Lončarić et al., 2005). This periodicity is modulated by seasonal changes in shell fluxes (Jonkers et al., 2015) with intervals characterised by smaller body sizes and extended life cycles during unfavourable conditions and food scarcity contrasting with times of high foraminiferal fluxes and larger shell sizes (Chernihovsky et al., 2023). For organisms with a circadian clock and therefore a fixed life duration, large sizes may reflect


increased growth rates (rather than prolonged growth), versus rapid reproduction and smaller body size as seen in many phytoplankton (Schmidt et al., 2006).

2.1.2 Calcification (foraminiferal trait)

The primary function of producing a calcite shell in planktic foraminifera is still debated. However, one potential advantage is that the shell (and, to a lesser extent, the spines in spinose species) provide the cytoplasm with support and protection, increasing the individual's potential to survive to reproductive maturity. This is achieved by (1) reducing their palatability to predators, (2) increasing their body size to reduce predation pressure, and (3) forming a potential barrier from environmental conditions such as harmful UV rays or pathogens (Armstrong, 2004).

The ~~mineralised shell/skeleton~~ can comprise a significant proportion of an organism's total mass, making the construction and maintenance ~~of a mineralised skeleton~~ energetically expensive (Sanders et al., 2018). However, it is difficult to quantify the energetic costs of biomineralisation and foraminifera (as ~~with~~ many organisms) can modulate their energy use between different activities depending on the environment. ~~An unclear / Lack of a mechanistic~~ understanding of how biomineralization occurs for many groups further exacerbates the challenge (Gaylord et al., 2015). No laboratory assessments of the energy budget allotted to calcification in planktic foraminifera currently exists. However, theoretical trait-based models indicate a likely investment of ~10-60 % of foraminifera's total energy budget ~~is apportioned~~ to calcification (Grigoratou et al., 2021; Grigoratou et al., 2022; Grigoratou et al., 2019).

A calcite shell ~~reduces the~~ ~~increases the negative~~ buoyancy of foraminifera, i.e., it increases their tendency to sink through the water column (Armstrong, 2004; Caromel et al., 2014), ~~although~~ this is counterbalanced by low-density fibrillar bodies within the cytoplasm (Hemleben, 1989). Spinose foraminifera taxa have long calcite spines that may extend up to several ~~cm~~ centimeters from the shell surface (2-3x the shell size; Fig. 1b, c and f), which increases their drag and thus, ~~reduces their~~ sinking rate (Takahashi and Be, 1984). Other benefits of spines include ~~an~~ increased capture area for prey acquisition, ~~including~~ reduced palatability to predators, a rigid skeleton to anchor pseudopodia to enable acquisition of higher quality (active) prey, and a means of organising photosymbionts more effectively around the shell. Logically, ~~the~~ development of these compensatory features should increase the relative cost of calcification. However, trait-based modelling suggests similar calcification costs for both spinose and non-spinose foraminifera species (Grigoratou et al., 2021; Grigoratou et al., 2022; Grigoratou et al., 2019). This suggests that spine formation either does not significantly increase energy demand or is offset by nutritional benefits. Spinose species are predominantly carnivorous ~~as adults~~, which provides more energy, and many also host algal photosymbionts providing an additional carbon subsidy, while non-spinose species are ~~omnivorous, they prefer~~ ~~mainly~~ herbivorous diet (Hemleben et al., 1985) (see Section 2.2).



213

214

215

216

217

218

219

220

221

222

223

224

225

Figure 3. Modern distribution and relative abundance of planktic foraminifera by ecological group in surface sediment samples. Underlying data from ForCenS database (Siccha and Kucera, 2017) and traits coded as presented in Table 1 (this study).

The range of estimated energetic calcification cost for foraminifera is broadly consistent with those of other calcifying groups (although estimates vary dramatically), including: coccolithophores at ~30 % (Monteiro et al., 2016), marine benthic gastropods at <60% (Palmer, 1992; Sanders et al., 2018), and <10% across gastropods and bivalves globally (Watson et al., 2017). In the calcareous algae coccolithophores, the high energetic cost of calcification varies between species and environments, and protection from predation may be the primary benefit (Monteiro et al., 2016). However, the role of predation on foraminifera is unknown. Whilst adult foraminiferal shells have been found in pteropods, salps, shrimps and many other metazooplankton, we do not know of any specialised predators (Berger, 1971). Nevertheless, the role of predation on

populations should not be underestimated, as many juveniles do not reach maturity (Schiebel et al., 1995) and are selectively found in faecal pellets of salp (Bé, 1977). Juvenile forms may have a higher palatability due to their higher ratio of cytoplasm to calcite (i.e., lack of numerous spines, thick calcite walls and gametogenic calcite) than adult forms (Meilland et al., 2016). In culture, damaged specimens or those undergoing gametogenesis were highly susceptible to digestion by other smaller protists (bacteria, sporozoans, and other parasitic organisms) that entered the foraminifera via the aperture (Hemleben, 1989).

Despite numerous data on the weight of foraminiferal shells (see Barrett et al., 2025 for a recent review), there are still large gaps in our understanding of the metabolic and energetic needs for calcification. However, it is clear that some specific traits within ecogroups, for example symbiosis (see Section 2.2.3), can positively impact the calcification response due to modification of the calcifying microenvironment and provision of an energy subsidy at a group level (e.g., Köhler-Rink and Kühl, 2005). Biomineralization pathways in planktic foraminifera are poorly known but are inferred to be extracellular with calcification occurring predominantly via endocytosis based on analogy to benthic foraminifer, e.g., *Amphistegina lobifera* (Bentov et al., 2009; Schiebel and Hemleben, 2007). Briefly, seawater is engulfed in vacuoles by the foraminifers plasma membrane, modified in composition to become “alkalinized” via assorted biological and chemical processes, and transported to the site of calcification. Enhanced respiration near the site of calcification locally increases the CO₂ in the foraminifers cytoplasm and ultimately the pool of total inorganic carbon available for calcification (Bentov et al., 2009). Alternative foraminiferal biomineralization pathways are suggested (Erez, 2003; de Nooijer et al., 2013; Bentov and Erz, 2006; Nehrke et al., 2013). This includes the benthic foraminiferal group, the miliolids, which precipitate an imperforate porcelaneous shell intracellularly, but this is very different in structure to planktic foraminifera shells (Hemleben et al., 1986). Many studies though identify a species-specific calcification responses and sensitivities associated with a range of environmental drivers (Barker and Elderfield, 2002; Béjard et al., 2024; Weinkauff, 2016; Pallacks et al., 2023). Understanding drivers is complicated by differential environmental preferences of cryptic species (De Vargas et al., 2001), regional plasticity, and a lack of understanding of the controls on the thickness of gametogenic calcite, or the biomineralization pathway (De Nooijer et al., 2023; Erez, 2003; Lekieffre et al., 2018; Nehrke et al., 2013). Furthermore, the variety of methods used to collect individuals for foraminifera weight (i.e., the use of plankton tow, sediment trap and core-top data, as well as differences in the weighing technique itself) (Beer et al., 2010) and inadequate metadata reporting lead to further difficulty in understanding calcification. Our understanding of what drives calcification could be enhanced by resolving the aforementioned biological factors (i.e. resolving cryptic species, understanding regional plasticity, what controls the thickness of gametogenic calcite, and determining the biomineralization pathway) and improving data collection and data management practices, e.g. by following FAIR principles (Wilkinson et al., 2016). In turn, this will result in better modelling of the potential changes to in the pelagic carbonate production under future climate change (Barrett et al., 2024, 2025).

The cost of calcification increases under ocean acidification scenarios in a wide range of (though not all) marine calcifiers (Hoppit and Schmidt, 2022), e.g., corals, molluscs, coccolithophores (Leung et al., 2022),. Impacted groups which typically

show evidence of reduced growth, reduced calcification, muscle wastage or weakened shells (e.g., Alma et al., 2020; Hill and Hoogenboom, 2022; Swezey et al., 2020), supporting that the metabolic costs of calcification are at the expense of other life processes. While the allotment of the foraminifera energy budget to calcification is not well constrained, its cost is evidenced in the smaller terminal body sizes (e.g., maintaining calcification at the expense of growth) in field and laboratory studies (Schmidt et al., 2003; Russell et al., 2004; Lombard et al., 2010). ~~I-In the short-term,~~ this results in a reduced chance of individuals reaching sufficient size to obtain reproductive maturity (Bé et al., 1981), potentially a smaller number of gametes based on the relationship between gamete number and body size reducing reproductive success. ~~This and~~ ultimately impacts ~~individuals' fitness~~ ~~individuals fitness~~ by increasing their vulnerability to further environmental change or predation etc. that may prove fatal.

2.2 Behavioural traits

2.2.1 Feeding (group-specific traits)

The biggest challenge facing marine plankton is the acquisition of food from a very dilute suspension without being eaten, a challenge exacerbated in low prey density environments such as oligotrophic gyres. In these environments, organisms must maximize opportunities for finding food and capturing their prey. Marine plankton adopt a range of different feeding strategies to deal with the challenge, each with specific trade-offs that extend beyond an impact on feeding behaviour alone.

All planktic foraminifera are passive ambush feeders (Fenchel, 1986; Kiørboe, 2011). They do not actively detect or seek out their prey. Acquisition of food relies solely on the motility of their prey and the likelihood of direct interception, a function of prey density and foraminifera's ability to 'capture' prey (Kiørboe, 2011). The benefits of passive food capture are its low energy investment, as organisms do not need to move to acquire their prey, and reduced predation risk due to lower predator encounter rates. However, the cost of the passive ambush feeding strategy is a low feeding efficiency and potentially low mate encounter rate (Kiørboe, 2011). This trade-off is very well constrained in copepod populations, with passive ambush feeders having lower metabolic and mortality rates than their more active counterparts (Eiane and Ohman, 2004).

The importance of passive feeding as the main nutritional mode is supported by the correlation between growth rate and feeding frequency (Bé et al., 1981). Foraminifera are capable of digesting most organic materials, presumably a survival strategy for environments where one or more food sources may not be available and mitigating environmental change. They have an extensive network of thin, sticky strands of cytoplasm (rhizopodia) around their shell that are used to help capture, transport, and break-up prey items, as well as carry waste products out of the cell (Hemleben, 1989). During the early life stages (prolocular-juvenile, ~15-100 µm shell diameter (Spindler et al., 1984; Caron et al., 1987), foraminifera are omnivorous, although primarily consuming an herbivorous diet.

As some foraminiferal species transition from the juvenile to neanic stage (>100 µm shell diameter), they may become host to photosymbiotic algae and/or develop spines. Spines provide a rigid framework for rhizopodia to attach and increase foraminifera's capture area (Gaskell et al., 2019), allowing them to capture and hold active prey facilitating a ~~and~~ switch from a predominantly herbivorous to carnivorous diet (Fig. 1c; Anderson et al., 1979; Schiebel and Hemleben, 2017). For example, *O. universa* shifts from herbivorous during its early and juvenile phase, to a carnivorous diet in the spherical adult stage (Schiebel and Hemleben, 2017). A wide variety of animal prey can be consumed by spinose planktic foraminifera including copepods, hyperiid amphipods, tunicates, ostracods, pteropods, gastropod larvae, ciliates, radiolarians, acanthurids and polychaete larvae (Anderson et al., 1979). Observational evidence suggests that copepods account for >40% of spinose taxa diets, with an additional ~20-30% from ciliates and the remainder made up of a variety of prey items (Schiebel and Hemleben, 2017).

In a laboratory environment, spinose taxa exhibited higher acceptance rates of calanoid over cyclopoid copepods (Spindler et al., 1984), indicating some degree of prey selectivity. ~~Notably It is noteworthy that, compared to most zooplankton, s~~ Spinose planktic foraminifera are capable of capturing and digesting prey items 2-3 times bigger than their body size (Schiebel and Hemleben, 2017). Carnivory is advantageous in a food-poor environment for organisms relying on chance encounters because individual prey items have a higher calorific value relative to a similar sized phytoplankton (Boyd, 1971).

Non-spinose planktic foraminifera are omnivorous but prefer ~~a~~ primarily herbivorous diet throughout their life cycle (Hemleben et al., 1985; Schiebel and Hemleben, 2017), preying on diatoms, dinoflagellates, and eukaryotic algae. Copepods (dead and alive) have been given in culture to several non-spinose species (*Globorotalia truncatulinoides*, *G. hirsuta*, *G. inflata*, *Globigerinita glutinata* and *Pulleniatina obliquiloculata*), but it was found these taxa are unable to capture and hold live zooplankton, and are only able to consume zooplankton if they are immobilized or dead (Spindler et al., 1984; Anderson et al., 1979). In the ocean, non-spinose species such as *G. menardii* can capture and control small ciliates using their rhizopodia, as evidenced by muscle and other animal tissues in food vacuoles (Anderson et al., 1979). Metabarcoding studies have shown that the Arctic non-spinose species *Neogloboquadrina pachyderma* include animal tissue in their cytoplasm (Greco et al., 2021); however, it remains unclear whether this tissue comes from live or dead prey. Hence, the predominantly herbivorous diet may largely be because of their difficulty in capturing living zooplankton. Under laboratory conditions, non-spinose species exhibit ~~show~~ cannibalism but whether is unknown if they cannibalise in ~~thea~~ natural habitat is unknown and ~~and is not~~ considered unlikely due to very low foraminiferal abundance ~~standing stocks~~ (Schiebel and Hemleben, 2017; Westgård et al., 2023).

The temporal and spatial distribution of prey items is a major ~~drive~~ cause of the regional distribution of planktic foraminifera species influencing their growth and fecundity in addition to after sea surface temperature (Schiebel and Hemleben, 2017;

[Schmidt et al., 2004; Adebayo et al., 2017; Lombard et al., 2011](#)). Spinose taxa are most abundant in (sub)tropical oligotrophic gyres (Fig. 3), where copepods are most abundant (Grice and Hart, 1962), whereas non-spinose species are most abundant in upwelling and coastal waters and at higher latitudes (>30°), which are rich in phytoplankton and small zooplankton (Grice and Hart, 1962).

~~Model~~ ~~Fe~~ ~~simulations of~~ trophic dynamics ~~requires with mathematical models (especially dynamic mechanistic models),~~ ~~observational~~ data on prey preferences, ~~e.g., such as~~ prey acceptance rates, protein acquisition from zooplankton versus phytoplankton, and average digestion and capture times ~~are needed~~. Observational data can be collected by analysing feeding vacuoles and metabarcoding analyses of in-situ samples. For planktic foraminifera, this information is scarce, and limited to a few taxa (e.g., the spinose species *Trilobatus sacculifer*, *Globigerinoides ruber* and *Orbulina universa*), which are most easily cultured in the laboratory. It is unclear, though, how representative laboratory environments are of natural feeding behaviours given the stress responses of some species, e.g., shortening or spine loss of *G. ruber* (Bijma et al., 1990). Furthermore, many critical parameters are known only from a single experiment. There is a clear need to capitalise on the recent success in culturing of a wider range of taxa such as the eutrophic *G. bulloides* (Sykes et al., 2024) or the polar *N. pachyderma* (Meilland et al., 2022) to explore these questions across a wide range of foraminifera eco-groups. Such multigenerational experiments would also allow closure of the important gap of quantitative data on the herbivory preferences or digestion rates of juveniles or the role of bacteria and organic matter ~~in foraminifers in foraminiferal~~ diet.

There are many further gaps in knowledge ~~of around about~~ the costs and benefits of being a passive ambush feeder, competition with other foraminifera or their close relatives, the siliceous radiolarians, all of which impact our ability to consider interactions ~~between with~~ zooplankton in trait models. To advance trait models, data is needed on prey preferences, prey-predator optimum length ratio, and encounter rates (successful and unsuccessful) to cover the energetic needs for calcification and other metabolic costs.

2.2.2 Starvation tolerance/[dormancy](#) (group-specific traits)

Feeding experiments indicate that when food is available at the optimum frequency, foraminifera reproduce quickly but the reverse is also true. ~~There is no evidence at present for a specialist adaptation in planktic foraminifera that enables certain groups to tolerate food deprivation for longer, e.g., lipid acquisition or diapause as evident in benthic foraminifera (Alve and Goldstein, 2002; Alve and Goldstein, 2010) or dinoflagellates (Tang et al., 2021). These resting stages are also associated with bloom formation when conditions become favourable again.~~ In culture, planktic foraminifera can survive and grow for 16-46 days with little or no food after initial capture, i.e., in some cases beyond their ‘normal’ lifespan in the ocean (Anderson et al., 1979; Bé et al., 1982). However, [associated](#) low growth leads ultimately to lower standing stocks, and/or smaller body sizes, threatening their ability to undergo gametogenesis or, as volume is linked to number of gametes, fewer gametes (Bé et al.,

1981). The taxa that have survived longest in culture (>230 days) are asymbiotic herbivorous taxa [such as *Neogloboquadrina pachyderma*](#) with growth optimised for low temperatures (Lombard et al., 2010) with presumably lower metabolic rates (Spindler, 1996).

[N. pachyderma](#) is the dominant planktic foraminifera at high latitudes and thrives in upwelling and (sub)polar oceans, able to survive low temperatures (-2 to +15°C) and a wider range of pH and salinities (Westgard et al., 2023). This is the only planktic foraminifera species for which dormancy is known. ~~One extreme example of a species with periods of low metabolism is the overwintering in sea ice of the non-spinose non-symbiotic *Neogloboquadrina pachyderma* (s~~Specifically Antarctic genotype IV [of *N. pachyderma*](#) of Darling and Wade (2008), ~~which dominates polar planktic foraminifera assemblages)~~ is observed to overwinter in brine channels and pockets within Antarctic sea ice (Spindler and Dieckmann, 1986; Dieckmann et al., 1991; Berberich, 1996; Spindler, 1996). This species can survive ~~the lack of algal biomass on which to feed during the austral winter by surviving and growing in sea ice (but does not reproduce) in brine channels and pockets within Antarctic sea ice (Spindler and Dieckmann, 1986; Dieckmann et al., 1991; Berberich, 1996; Spindler, 1996) in brine channels and pockets within the ice with.~~ Brine pockets and channels can have salinities up to 177 psu and temperatures as low as -15°C. ~~However, at >50 psu, though, individuals though individuals grew were found to be able to grow slowly and reached smaller overall body sizes and were be~~unable to undergo gametogenesis. ~~At >73 psu, pseudopodal activity and movement ceased, but they were able to survive for up to several weeks at up to ~82 psu without feeding (Spindler, 1996). This finding is supported by more recent culturing work suggesting tolerance to a wide range of salinities but that specimens are less active and less likely to add new chambers at very high or low salinities or reproduce (Bertlich et al., 2021, Westgard et al., 2023). Indeed, in culture, specimens that were inactive (e.g., no growth, limited/no rhizopodal activity) or dormant (appeared to have empty shells/decaying cytoplasm and did not feed) that appeared dead were able to recover as conditions became more favourable (Westgard et al., 2023).~~ The same ~~brine pockets and channels habitats~~ also contain dense populations of phytoplankton, predominantly diatoms, a rich [potential](#) food source [for *N. pachyderma*](#) (Spindler and Dieckmann, 1986; Dieckmann et al., 1991). Overwintering via dormancy is thus advantageous because [it allows *N. pachyderma* to suppress metabolic activity to reduce energy consumption \(e.g., buoyancy compensation, growth\) and survive unfavourable environmental circumstance but also, provides](#) ~~of a~~ rich food source [for when conditions are favourable, and](#) protection from predators; ~~and/or a means of conserving energy by reducing the need for buoyancy compensation (Dieckmann et al., 1991).~~ [Dormancy ultimately means that taxa inhabiting sea ice can significantly extend their 'normal' lifespan, potentially surviving for up to one year from the formation of sea ice in the autumn until the following spring/summer \(Spindler, 1996\).](#)

2.2.3 Photosymbioses (group-specific traits)

Symbiosis between different biological organisms is a common ecological strategy in the ocean from shallow benthic marine ecosystems (e.g., coral reefs) through to the nekton and plankton, a major source of evolutionary innovation and hence

biodiversity (Decelle et al., 2015; Margulis, 1993). Planktic symbiotic relationships in the open ocean are relatively poorly constrained in contrast to benthic shallow-water counterparts, however, symbiosis in planktic foraminifera appears to represent an adaptation to nutrient-poor, sunlit waters (Fig. 3d; Bé, 1971).

Many of the largest benthic and planktic foraminifera in the modern ocean tend to host algal photosymbionts (Kucera, 2007), highlighting the important role of symbiosis in providing additional energy to support the energetic costs of building a larger skeleton. In foraminifera, species that occupy the photic zone are commonly host to algal photoendosymbionts, either chrysophycophytes or dinoflagellates (Fig. 1b and c). Symbionts ~~which~~ are acquired by juvenile foraminifera from the water column [following sexual reproduction](#) (Hemleben, 1989) [but potentially via the parent by direct vertical transmission during the more poorly known asexual reproductive cycle](#) (Takagi et al., 2020). Whilst many spinose taxa are host to dinoflagellate photosymbionts, there are exceptions, [e.g., *G. bulloides* is associated with bacterial endobionts, e.g., *Synechococcus* in the eastern Pacific Ocean](#) (Bird et al., 2017) or none, [e.g., *Hastigerina pelagica* \(Takagi et al., 2019\) and *G. bulloides*](#). [Whereas](#) ~~s~~Some non-spinose taxa such as *G. glutinata* or *Globigerinella siphonifera* may also host photosymbionts, mostly chrysophytes (Takagi et al., 2019). However, whether these species have a truly symbiotic relationship with foraminifera, or just utilise waste products, is currently unknown (Hemleben, 1989). Available data suggest that the type of symbiont dictates photophysiology rather than host size or spines (Hoadley et al., 2019; Takagi et al., 2019). This conclusion is supported by the [likely](#) evolution of dinoflagellate symbiosis in the Cretaceous before spines or spine-like structures evolved in the Palaeocene (Hoadley et al., 2019; Pearson et al., 2001; Bornemann and Norris, 2007), suggesting that whilst spines may help to optimise photosymbiont activities, they are not essential (and therefore, [this is](#) not the primary function of spines).

Symbiosis can be described as obligate (essential for optimal host fitness and observed in the majority of individuals of a species, making them functionally mixotrophic) or facultative (not essential to the host success and thus, only associated with some individuals of the species) (Table 1; Takagi et al., 2019). Determining whether symbiosis is obligate or facultative is challenging as photosymbiosis is a spectrum, as proposed by Stoecker et al (2009), from non-symbiosis to robust symbiosis. We have direct observations of symbiotic presence or absence in more than 30 species (see Table 1), typically recognised via microscopic observations (Anderson and Be, 1976) and/or molecular work (Gast and Caron, 1996; Yonathan and Colombari, 2006). However, it is difficult to clearly differentiate between digested and active symbiosis with these methods. Active chlorophyll fluorescence, a non-destructive and invasive approach that allows assessment of fluorescence through ontogeny, is a powerful technique helping to close this knowledge gap and advance modelling traits related to symbioses by enabling determination of ~~C~~chlorophyll—a content of specimens, health of symbionts and their light-level adaptation (Takagi et al., 2019; Takagi et al., 2016).

Within planktic foraminifera, taxa with obligate (dinoflagellate) photosymbionts tend to dominate oligotrophic regions with expanded mixed layers and high light penetration (Fig. 3). Symbionts are rarer at cold, low light high latitude areas, in deep

waters >200 m and in eutrophic regions of the ocean, presumably where they are either unable to survive or unnecessary (Hemleben, 1989). No symbiont bearing taxa are found at very high latitudes (>~50 degrees; Fig. 3).

Photosymbionts provide an important energy supplement to their host in the form of photosynthetically fixed carbon, aiding growth, longevity, calcification and reproductive potential (e.g., Lekieffre et al., 2018; Bé et al., 1982), but photosymbionts alone do not provide sufficient carbon subsidy to entirely support foraminiferal life processes. ~~Photosymbionts provide an additional energy source (Lekieffre et al., 2018), aiding foraminifera growth and calcification (Bé et al., 1982).~~ At the same time, symbionts preferentially use metabolites from the foraminifera for photosynthesis (Takagi et al., 2018). Algal photosymbionts also aid calcification by ~~increasing~~ modifying the pH of the foraminifera's immediate microenvironment above ambient seawater by utilising CO₂ during photosynthesis and therefore potentially enhancing calcification (Rink et al., 1998; Kohler-Rink and Kuhl, 2005).

Photosymbionts are acquired as juveniles, or, as new evidence suggests, provided to gametes during gametogenesis (Takagi et al., 2020) and are typically arranged in the external rhizopodial net during the day and brought into the shell at night (e.g., Anderson and Be, 1976; Lekieffre et al., 2018). Symbiont biomass rises and falls during the host's life (Takagi et al., 2016), with the symbionts eventually being consumed immediately prior to gametogenesis (Bé, 1983). Digestion of photosymbionts may help to meet the energy demands of gametogenesis or provide energy during periods of prolonged darkness (e.g., Spero and Parker, 1985; Bé, 1983; Takagi et al., 2016). However, eEven under starvation, some symbionts are retained in culture until gametogenesis (Takagi et al., 2018).

~~It is unclear what triggers the need to change from asymbiotic to symbiotic in these facultatively symbiont bearing taxa, making it impossible to model them~~ as the costs and benefits cannot be determined sufficiently. Takagi et al. (2019) showed that photosynthetic activity in symbiont-facultative species tends to be weaker than in symbiont-obligate species and thus, nutritional benefit to the host may be smaller. However, the benefit of ~~this~~ facultative symbiosis trait is still unclear. It may be opportunistic, and may overall support a more flexible range of nutritional sources for the host. A lower reliance on symbiont activity in organisms might also allow these ~~taxa~~ symbiont-facultative foraminifera to explore ~~more~~ low-light eutrophic regions, including the deep-water layers and the turbulent upwelling regions. A recent study developing the ForamECOGEniE model to incorporate the different symbiosis types (Ying et al., 2023) did a good job of replicating patterns of asymbiotic and obligate symbiotic global distributions but underestimated the abundance of the non-spinose symbiont-facultative group particularly in the eastern equatorial Pacific. The challenge of modelling this groups is the lack of information on what drives the symbiont-facultative group to acquire to lose their symbionts. Without a clear ecological/physiological understanding, trade-offs cannot be incorporated in the model. Intriguingly, if the abundance of the obligate and facultative symbiotic taxa are combined there is a good fit between the model and observations (Ying et al., 2023). This may suggest that the groups overall exploits the same benefits from symbiosis. However, determining the mechanistic underpinning is critical

to understand the data, particularly facultative symbiosis trait, its trade-offs and the environmental/biological conditions under which symbiosis is active or not, and the benefits of the relationship to the host. Thus, culture experiments and further observation of this phenomenon are still required under, including variable environmental conditions and resulting physiological changes.

There are several costs associated with hosting photosymbionts [are](#), [1] they restrict foraminifera's depth habitat to the euphotic zone (generally <200 m water depth; Caron et al., 1987; Schiebel and Hemleben, 2017), and [2] the host is dependent ~~to~~ on changes in the availability and health of the symbionts, which responds to environmental change. Studies suggesting that foraminifera symbionts are not favoured in low pH waters (Henehan et al., 2017) or in eutrophic areas due to light limitations (Ortiz et al., 1995). Reduced photosynthetic activity results in smaller final body sizes and shorter survival times (Ortiz et al., 1995; Bé et al., 1981; Caron et al., 1982; Faber et al., 1989). During extreme temperatures in the geological record, planktic foraminiferal populations of 'bleached' individuals may [have](#) persisted for thousands of years with smaller sizes and lower population abundances (Edgar et al., 2013; Wade et al., 2008), which may have impacted fitness and increased their susceptibility to extinction (e.g., Wade et al., 2008).

The knowledge gap relating to symbioses, beyond the fundamental knowledge of which symbiont each species hosts, includes what triggers changes in symbiont [activity and mode of symbiosis hosting](#), the degree of energy subsidy symbionts provide to their host and further understanding of the use of symbionts as prey items. [For *G. bulloides*, the nature of their association with bacterial endobionts also needs further exploration to understand how widespread the association is between genetic types, and within the ocean.](#) Without this understanding, and the ecological impact on the symbiont, the future fitness of planktic foraminifera cannot be explored fully, nor [can](#) the importance of [the](#) loss of symbiosis in the fossil record after extinctions (Birch et al., 2016).

2.3 Reproduction

2.3.1 Sexual reproduction (foraminiferal/group-specific traits)

Planktic foraminifera were traditionally [observed to only considered to](#) reproduce sexually (e.g., Bé and Anderson, 1976, Ketten and Edmond, 1979, and references in Schiebel and Hemleben, 2017) differing from their benthic counterparts that can [alternate between sexual and asexual reproductive modes \(Goldstein, 1999\).](#) ~~only, however~~ but recent culture breakthroughs [have shown that some non-spinose foraminiferal species \(*N. pachyderma* and *G. glutinata* and *G. uvula*\) can generate via both asexual and sexual reproduction \(Davis et al., 2020; Kimoto and Tsuchiya, 2006; Meilland et al., 2023; Meilland et al., 2024\).](#) [With most shallow dwelling planktic foraminifera reproducing on a semi-lunar or lunar synodic cycle \(~2-4 weeks\), whereas intermediate to deeper dwelling species may live for up to one year \(Spindler, 1979; Hemleben, 1989\). Foraminifera migrate](#)

to reproduce at depth close to the deep chlorophyll maximum, where there are optimum feeding and grazing protection opportunities for offspring (Hemleben, 1989). Notably most observational work was on spinose species. During gametogenesis, foraminifera undergo a suite of morphological, physiological and ecological changes. They retract their rhizopodia, shorten and shed their spines by dissolution at the tips and resorption at the base, consume any photosymbionts and precipitate an additional outer layer of calcite (gametogenic calcite) over the shell (Schiebel and Hemleben, 2017). Gametogenesis ends with the conversion of all cytoplasm into gametes via vacuolisation, which are released directly into the water column (broadcast spawning). Gametogenesis takes ~1-3 days from the formation of the final chamber (Bé, 1983), resulting in an empty adult shell which sinks to the seafloor (Seibold, 1993).

~~Most shallow dwelling planktic foraminifera reproduce on a semi-lunar or lunar synodic cycle (~2-4 weeks), whereas intermediate to deeper dwelling species may live for up to one year (Spindler, 1979; Hemleben, 1989). Foraminifera migrate to reproduce at depth close to the deep chlorophyll maximum, where there are optimum feeding and grazing protection opportunities for offspring (Hemleben, 1989). As they only grow for short periods and typically undergo reproduction without producing viable offspring, multi generational experiments of planktic foraminifera have only recently been achieved (Meilland et al., 2024; Sykes et al., 2024).~~

Sexual reproduction allows populations to maintain higher genetic diversity and select for advantageous mutations or conversely eradicate unfavourable mutations (Otto and Lenormand, 2002). Therefore, sexual reproduction provides a definite (if though difficult to quantify) advantage for survival in the dynamic surface waters with constantly changing selective pressures (Lynch, 1991). In addition, the broadcast spawning strategy presumably confers energy savings compared to organisms actively searching for mates and reduced predator encounters. However, the typically low concentration of foraminifera in the water column (~1 specimen per m³ in the open ocean) makes this a risky strategy. Further, any mismatches in the timing (or place) of reproduction between individuals would reduce the chance of fertilization.

As such, planktic foraminifera have developed a number of strategies to maximise the successful fertilization of gametes from different parents in the water column, including (1) the synchronization of the timing and depth of reproduction between multiple individuals of the same species, (2) the release of a large number of gametes (200,000-400,000 individuals; dependent on both shell and gamete size) from each adult, thereby increasing encounter and survival rates, and (3) the development of motile gametes (see Weinkauf et al., 2022a and references therein).

~~Being able to change reproductive strategies is consistent with the ability to respond to short-lived optimal conditions in the ocean (Davis et al., 2020). Sexual reproduction allows populations to maintain higher genetic diversity and select for advantageous mutations or conversely eradicate unfavourable mutations (Otto and Lenormand, 2002). Therefore, sexual reproduction provides a definite (if though difficult to quantify) advantage for survival in the dynamic surface waters with~~

constantly changing selective pressures (Lynch, 1991). In addition, the broadcast spawning strategy presumably confers energy savings compared to organisms actively searching for mates and reduced predator encounters. However, the typically low concentration of foraminifera in the water column (~1 specimen per m³ in the open ocean blue waters) makes this a risky strategy. Delayed reproduction potentially reduces the chance of fertilization by putting because individuals are out of sync with the rest of the population and/or because the organism is smaller and thus, produces fewer gametes.

As such, planktic foraminifera have developed a number of strategies to maximise the successful fertilization of gametes from different parents in the water column, including (1) the synchronization of the timing and depth of reproduction between multiple individuals of the same species, (2) the release of a large number of gametes (200,000–400,000 individuals; dependent on both shell and gamete size) from each adult, thereby increasing encounter and survival rates, and (3) the development of motile gametes (see Weinkauf et al., 2022a and references therein).

Modelling the development in of foraminifera, in a way that resembling resembles their accretionary growth has not yet been achieved to date. The main challenge is the lack of information on the different trades and traits and trade-offs in their development, such as changes in metabolic rates, food uptake, etc. The new culturing breakthrough, which enables us to explore the full life cycle of planktic foraminifera, opens the door to explore these questions and close our knowledge gaps.

2.3.2 Asexual reproduction (species/group-specific traits)

Recent culture breakthroughs have shown that some non-spinose foraminiferal species (*N. pachyderma*, *G. glutinata* and *G. uvula*) can reproduce both asexually and sexually- (Davis et al., 2020; Kimoto and Tsychiya, 2006; Meilland et al., 2023; Meilland et al., 2024, Takagi et al., 2020). In *N. pachyderma*, precursors to asexual reproduction include feeding in large quantities, the development of bright red cytoplasm, and maintenance of the rhizopodial network, the latter in contrast to pre-gametogenic changes (Greco et al., 2023, Meilland et al., 2024). There also remains a positive relationship between the size of the foraminifera shell and number of offspring under asexual reproduction, albeit with fewer offspring per individual (~80–300) than via sexual reproduction (Meilland et al., 2024). However, there is generally considered to be higher survival of offspring compared to sexual reproduction hence this reproductive mode supports rapid growth of planktic foraminifera populations when conditions are optimal. Asexual reproduction also provides a mechanism to explain the survival of and rapid population growth of *N. pachyderma* at high latitudes (Davies et al., 2020, Meilland et al., 2024). *N. pachyderma* returns to the water column after overwintering in sea ice by migration towards the base of the ice (Berberich, 1996) and/or, as the ice melts in the polar spring/summer (Spindler and Dieckmann, 1986).

The ability of, at least some species, of planktic foraminifera to switch between sexual and asexual reproduction helps to reconcile rapid population growth in response to temporally and spatially optimal conditions, and conditions and represents a

distinct advantage for survival of polar species in particular (Davies et al., 2020). It is highly likely that future targeted experimental work will expand the evidence base for the number of planktic foraminifera species that can reproduce asexually.

~~The multigenerational experiments were achieved with *N. pachyderma*, which, as discussed above, can overwinter in sea ice. After overwintering in sea ice, *N. pachyderma* either return to the water column by migrating towards the base of the ice (Berberich, 1996) and/or, as the ice melts, are released to seed a new population (Spindler and Dieckmann, 1986). This ultimately means that taxa inhabiting sea ice can significantly extend their ‘normal’ lifespan, potentially surviving for up to one year from the formation of sea ice in the autumn until the following spring/summer (Spindler, 1996). The and ability of to *N. pachyderma* to switch/change between asexual and sexual reproductive modes on might be the way of achieving this unique life cycle with asexual reproduction facilitating reproduction and facilitating need for rapid blooms under optimal environmental conditions in the short Antarctic summer (Meilland et al., 2023). However, specimens grown under high salinity conditions are small, resulting in reduced gamete production.~~

Modelling the development of foraminifera, in a way that resembles their accretionary growth has not been achieved to date. The main challenge is the lack of information on the different trades and traits and trade-offs in their development, such as changes in metabolic rates, food uptake, etc. The new culturing breakthrough, which enables us to explore the full life cycle of planktic foraminifera, opens the door to explore these questions and close our knowledge gaps.

3. Suggestions for further research ~~Conclusions~~

~~Planktic foraminifera are an ideal target group for testing trait-based approaches, as a~~ a relatively small number of functional traits ~~can be necessary to define their~~ ecological niches of planktic foraminifera in the modern ocean ~~and they are thus, an ideal target group for testing trait-based approaches.~~ ~~However~~ Unfortunately, many of the traits identified here remain poorly qualified and quantitatively constrained, requiring ~~further~~ new observational and experimental ~~and/or~~ laboratory-based/experimental investigations work.

A key trait requiring further study is calcification. Its trade-offs (Table 2), particularly its hypothesised benefit of protection against grazing, are fundamental to current trait-based models for calcifying plankton but still lacks direct evidence (Barrett et al. 2025; Monteiro et al. 2016). We also require an understanding of the energetic costs of calcification and the mechanisms controlling biomineralization. This requires moving away from analogies largely based on physiological calcification studies of benthic foraminifera (e.g., De Nooijer et al 2009, De Nooijer et al 2014) to similar experiments on planktic foraminifera. At present, the poorly constrained relationship between calcification and seawater carbonate chemistry, limits our ability to

predict the impact of ocean acidification on foraminiferal growth and carbonate production. These knowledge gaps also hamper accurate modelling of ocean alkalinity distribution and carbonate production in response to a changing climate.

In other organisms, food availability can support metabolism to reduce the impacts of anthropogenic climate change on other physiological processes (Thomsen et al., 2013). Further understanding of foraminiferal predator-prey dynamics is therefore urgently needed. For example, cultures and *in-situ* observations can improve our understanding of feeding, by providing quantitative data on (1) the role of bacteria, (2) the role of organic matter in foraminiferal diet - specifically protein acquisition from zooplankton versus phytoplankton, (3) prey-predator optimum length ratio, (4) prey encounter rates (successful and unsuccessful), and (5) average digestion and capture time of prey. These data could (in part) be collected by analysing feeding vacuoles and metabarcoding analyses of samples.

Photosymbiosis, a major trait among symbiotic foraminifera, also requires further investigation. Culture and observational studies should focus on the impacts of environmental change on symbiosis. These include understanding what triggers change in symbiont hosting (e.g., bleaching, and hosting for symbiont-facultative species), which symbiont each species hosts (facilitated by eDNA analyses), and the energy the symbiont provides its host.

Experiments that exploit the new breakthrough in multigenerational planktic foraminifera culturing will improve our knowledge of their life cycle, through understanding the traits and trade-offs associated with their development, e.g., their metabolic rates, food uptake, and reproduction, and importantly which species are capable of asexual reproduction and under what conditions this mode dominates. All of which will contribute to better data for improved modelling of planktic foraminifers.

Environmental DNA (eDNA) metabarcoding, which analyses the genetic material present in the environment, such as sediment, or water, is a powerful new tool for identifying and monitoring biodiversity, biogeography and reconstructing ecosystems and ecologies (Ruppert et al., 2019). This technique can also provide insights into community composition over timescales spanning several hundreds of thousands of years or longer, improving our understanding of the relationships between biodiversity, environment and climate (e.g., Ambrecht et al., 2019). However, the bulk of eDNA foraminiferal studies to date have focussed on benthic foraminifera. For instance, this technique has significantly increased the diversity of organic walled and “naked” foraminifera that are rarely observed, have few morphological characters for traditional species delimitation, and don’t preserve well in the fossil record (Pawlowski et al., 2014). But eDNA holds great promise for investigating marine plankton, as it has the potential to overcome many of the data limitations that we currently face in this group with typically low standing stocks (de Vargas et al., 2015). It is a potentially more effective means of detecting species presence in an environment than observations alone (Malviya et al., 2016; Ser-Giacomi et al., 2018; Barrenechea Angeles et

al., 2020) but can also contribute much more broadly. For instance, providing insights into plankton population size (Andres et al., 2023), response to environmental change (Cao et al., 2022) or predator-prey dynamics (Ruppert et al., 2019). A combination of molecular and microscopic approaches can also yield new insights, e.g., in benthic foraminifera it allowed rapid determination of multiple different feeding strategies driving diversity and abundance in several foraminiferal taxa (Schweizer et al., 2022), a question which is typically restricted to analysis of feeding vacuoles and laboratory experiments. However, further method development is still required as some groups are not as well represented by eDNA technique as others, e.g., specimens may visually be present in sediments not found in the eDNA analysis, likely because of limitations of the primer to detect certain groups (Barrenechea Angeles et al., 2020, Hoshino and Inagaki, 2024).

~~In general, Further~~ there is a need to improve data reporting practices ~~infor the aforementioned studies~~ by following guidance on data publishing (e.g. FAIR principles) and developing standardized, community agreed protocols ~~akin to other fields (e.g., Riebesell, et al., 2010)~~ for measuring traits to enhance our understanding of planktic foraminifera and how they are best modelled.

~~Ultimately, dynamic-trait-based models,~~ are a useful tool to create theoretical frameworks to quantify ecosystem functions of foraminiferal traits and explore questions at scale and outside of those possible in laboratory or natural environments. ~~For instance, ecological parameters measured in laboratory environments could be used in trait-based models to quantify their impacts on the global biomass and calcite production, both are poorly constrained in the modern carbon cycle (Ying et al. 2023).~~ ~~In addition, paleoceanography models could help to elucidate the relationship between foraminiferal evolution (i.e., the emergence of new traits) and the background climate change, e.g., the evolution of deeper-water taxa as the oceans cooled over the past 15 Myrs (Boscolo & Galazzo 2021).~~ For modern species with high mortality rates in laboratory culture or difficulties maintaining multigenerational experiments (and high time and financial costs), trait-based models provide realistic setups to estimate potential physiological ecology in the absence of physical organisms (Grigoratou et al., 2019).

~~However, whilst a powerful tool, numerical models are ultimately just that, models. They are by necessity based on assumptions and simplifications of the natural world. Model reliability and performance is highly dependent on the specific research question asked and the observations available against which models can be validated.~~ For planktic foraminifera, the most abundant data for model calibration is sediment core top data and most foraminiferal models can reproduce the observed global biogeography of main species/ecogroups in sediment cores. However, less empirical data can be used for foraminiferal physiological model parameterisation. The growth rate and temperature dependency data from Lombard et al. (2009) are adopted by most plankton functional type models (e.g., PLAFOM, FORAMCLIM, PLANKTOM). However, less constraint exists for food preference and grazing efficiency, hampering us to simulate the observed low foraminiferal biomass. Trait-based models can dynamically calculate grazing rate based on body size (allometric law) and temperature/food availability. However, a recent study shows that planktic foraminifera have lower allometric scaling of energetic needs compared to other

plankton when considered in the context of organic density and catchment volume (Burke et al. 2025). Therefore, key parameters needed to improve foraminiferal model performance include more quantitative data of fundamental parameters such as: growth rate, respiration rate, half-saturation constant, and grazing preferences.

Despite existing data gaps and model challenges, distinct groupings of planktic foraminifera traits identified in the modern ocean allow us to explore key drivers of past biogeography, changes in metabolic traits and trade-offs, and responses to extreme environments with moderate confidence (e.g. Deutsch et al. 2020). Closer collaboration between modellers and data scientists is essential to optimise the application of trait-based approaches to planktic foraminifera. Targeted data collection to fill specific critical knowledge gaps via models, laboratory culturing, field observations, molecular techniques and more inclusive reporting, will improve confidence in predictions from trait-based models and help to realise the full potential of this approach. Ultimately, understanding the associations of different traits and suites of traits with particular environments provides the basis to understand how environmental factor's structure planktic foraminifera communities in the past and in the future. This knowledge will be crucial for assessing the vulnerability of these communities to ongoing and future environmental changes.

~~For foraminifera, these models can address laboratory limitations, such as high mortality rates in multispecies experiments (and high time and financial costs) and provide realistic setups even in the absence of physical organisms. Beyond utilizing existing observational data, trait-based models are a powerful tool to generate new data that enhance our mechanistic understanding of the observations we gather, allowing us to move beyond correlation to uncover the underlying causative relationships.~~

Author contribution

DNS, FM and KME conceived the idea for the manuscript. KME and MG prepared the original draft of the manuscript. KME, MG and RY created the figures and tables. All authors contributed to reviewing and editing of the manuscript. DNS supervised the work.

Competing Interests

The authors declare that they have no conflict of interest

Acknowledgements

Financial support was provided to KME in the form of a Leverhulme Early Career Fellowship (ECF-2013-608) and NERC grant NE/X000567/1 to KME, MG the European Research Council "PALEOGENiE" Project (ERC-2013- CoG617313), RB by NE/S007504/1, RY by a Chinese scholarship Council grant (202006380070), F.M.M. by NE/X001261/1, NE/V01823X/1 and DNS NERC grant NE/P019439/1.

References

- [Adebayo, M.B., Bolton, C.T., Marchant, R., Bassinot, F., Conrod, S and de Garidel-Thoron, T.: Environmental controls of size distribution of modern planktonic foraminifera in the tropical Indian Ocean, *Geochemistry, Geophysics, Geosystems*, 24, e2022GC010586, 2023.](#)
- Alma, L., Kram, K. E., Holtgrieve, G. W., Barbarino, A., Fiamengo, C. J., and Padilla-Gamiño, J. L.: Ocean acidification and warming effects on the physiology, skeletal properties, and microbiome of the purple-hinge rock scallop, *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 240, 110579, <https://doi.org/10.1016/j.cbpa.2019.110579>, 2020.
- Alve, E. and Goldstein, S. T.: Resting stage in benthic foraminiferal propagules: a key feature for dispersal? Evidence from two shallow-water species, *J. Micropalaeontol.*, 21, 95-96, 10.1144/jm.21.1.95, 2002.
- Alve, E. and Goldstein, S. T.: Dispersal, survival and delayed growth of benthic foraminiferal propagules, *Journal of Sea Research*, 63, 36-51, <https://doi.org/10.1016/j.seares.2009.09.003>, 2010.
- Andersen, K. H., Blanchard, J. L., Fulton, E. A., Gislason, H., Jacobsen, N. S., and van Kooten, T.: Assumptions behind size-based ecosystem models are realistic, *ICES Journal of Marine Science*, 73, 1651-1655, 10.1093/icesjms/fsv211, 2016.
- Anderson, O. R. and Be, A. W. H.: The ultrastructure of a planktonic foraminifer, *Globigerinoides sacculifer* (Brady), and its symbiotic dinoflagellates, *Journal of Foraminiferal Research*, 6, 1-21, 10.2113/gsjfr.6.1.1, 1976.
- Anderson, O. R., Spindler, M., Bé, A. W. H., and Hemleben, C.: Trophic activity of planktonic foraminifera, *Journal of the Marine Biological Association of the United Kingdom*, 59, 791-799, 10.1017/S002531540004577X, 1979.
- [Armbrecht, L.H., Coolen, M.J.L., Lejzerowicz, F., George, S.C., Negandhi, K., Suzuki, Y., Young J., Foster N.R., Armand, L.K., Cooper, A., Ostrowski, M., Focardi, A., Stat, M., Moreau, J.W., Weyrich L.S.: Ancient DNA from marine sediments: precautions and considerations for seafloor coring, sample handling and data generation. *Earth-Sci Rev* 196:102887, 2019.](#)
- Armstrong, H. A. a. B., M.D.: Foraminifera, in: *Microfossils*, Blackwell Publishing, 142-187, 2004.

Aze, T., Ezard, T. H., Purvis, A., Coxall, H. K., Stewart, D. R., Wade, B. S., and Pearson, P. N.: A phylogeny of Cenozoic macroperforate planktonic foraminifera from fossil data, *Biol Rev Camb Philos Soc*, 86, 900-927, 10.1111/j.1469-185X.2011.00178.x, 2011.

Barker, S. and Elderfield, H.: Foraminiferal calcification response to glacial-interglacial changes in atmospheric CO₂, *Science*, 297, 833-836, 10.1126/science.1072815, 2002.

Barrett, R., de Vries, J., and Schmidt, D. N.: What controls planktic foraminiferal calcification?, *EGU sphere Biogeosciences*, 22, 791-807, <http://doi.org/10.5194/eguspherebg-22-791-2025-2024-2405>, 2025.

Barton, A. D., Irwin, A. J., Finkel, Z. V., and Stock, C. A.: Anthropogenic climate change drives shift and shuffle in North Atlantic phytoplankton communities, *Proceedings of the National Academy of Sciences*, 113, 2964-2969, doi:10.1073/pnas.1519080113, 2016.

Barton, A. D., Pershing, A. J., Litchman, E., Record, N. R., Edwards, K. F., Finkel, Z. V., Kiørboe, T., and Ward, B. A.: The biogeography of marine plankton traits, *Ecology Letters*, 16, 522-534, <https://doi.org/10.1111/ele.12063>, 2013.

Beer, C. J., Schiebel, R., and Wilson, P. A.: Technical Note: On methodologies for determining the size-normalised weight of planktic foraminifera, *Biogeosciences*, 7, 2193-2198, 10.5194/bg-7-2193-2010, 2010.

Berberich, D.: Die planktische Foraminifere *Neogloboquadrina pachyderma* (Ehrenberg) im Weddellmeer, Antarktis, *Berichte zur Polarforschung*, 195, 10.2312/BzP_0195_1996, 1996.

Berger, W. H.: Planktonic foraminifera; sediment production in an oceanic front, *Journal of Foraminiferal Research*, 1, 95-118, 10.2113/gsjfr.1.3.95, 1971.

Bijma, J., Erez, J., and Hemleben, C.: Lunar and semi-lunar reproductive cycles in some spinose planktonic foraminifers, *Journal of Foraminiferal Research*, 20, 117-127, 10.2113/gsjfr.20.2.117, 1990.

Birch, H. S., Coxall, H. K., Pearson, P. N., Kroon, D., and Schmidt, D. N.: Partial collapse of the marine carbon pump after the Cretaceous-Paleogene boundary, *Geology*, 44, 287-290, 10.1130/g37581.1, 2016.

- Bird, C., Darling, K.F., Russell, A.D., Davis, C.V., Fehrenbachers, J., Wyman, M. and Ngwenya, B.T. Cyanobacterial endobionts within a major marine planktonic calcifier (*Globigerina bulloides*, Foraminifera) revealed by 16S rRNS metabarcoding, *Biogeosciences*, 14, 901-920, 10.5194/bg-14-901-2017, 2017.
- Bornemann, A. and Norris, R. D.: Size-related stable isotope changes in Late Cretaceous planktic foraminifera: Implications for paleoecology and photosymbiosis, *Marine Micropaleontology*, 65, 32-42, <https://doi.org/10.1016/j.marmicro.2007.05.005>, 2007.
- Boyd, C. E. a. G., C.P.: Nutritive quality of food in ecological systems, *Arch. Hydrobiol.*, 69, 256-270, 1971.
- Brandão, M. C., Benedetti, F., Martini, S., Soviadan, Y. D., Irisson, J.-O., Romagnan, J.-B., Elineau, A., Desnos, C., Jalabert, L., Freire, A. S., Picheral, M., Guidi, L., Gorsky, G., Bowler, C., Karp-Boss, L., Henry, N., de Vargas, C., Sullivan, M. B., Acinas, S. G., ..., and Tara Oceans Consortium, C.: Macroscale patterns of oceanic zooplankton composition and size structure, *Scientific Reports*, 11, 15714, 10.1038/s41598-021-94615-5, 2021.
- Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., and West, G. B.: Toward a metabolic theory of ecology, *Ecology*, 85, 1771-1789, <https://doi.org/10.1890/03-9000>, 2004.
- Brummer, G.-J. A., Hemleben, C., and Spindler, M.: Ontogeny of extant spinose planktonic foraminifera (Globigerinidae): A concept exemplified by *Globigerinoides sacculifer* (Brady) and *G. ruber* (d'Orbigny), *Marine Micropaleontology*, 12, 357-381, [https://doi.org/10.1016/0377-8398\(87\)90028-4](https://doi.org/10.1016/0377-8398(87)90028-4), 1987.
- Brummer, G. J. A. and Kučera, M.: Taxonomic review of living planktonic foraminifera, *J. Micropalaeontol.*, 41, 29-74, 10.5194/jm-41-29-2022, 2022.
- Buitenhuis, E. T., Hashioka, T., and Quéré, C. L.: Combined constraints on global ocean primary production using observations and models, *Global Biogeochemical Cycles*, 27, 847-858, <https://doi.org/10.1002/gbc.20074>, 2013.
- Bé, A. W. and Anderson, O. R.: Gametogenesis in planktonic foraminifera, *Science*, 192, 890-892, 10.1126/science.946914, 1976.
- Bé, A. W. H.: An ecological, zoogeographic and taxonomic review of Recent planktonic foraminifera, in: *Oceanic Micropaleontology*, edited by: Ramsay, A. T. S., Academic Press, London, 1-100, 1977.

794 Bé, A. W. H., Caron, D. A., and Anderson, O. R.: Effects of feeding frequency on life processes of the planktonic
795 foraminifer *Globigerinoides sacculifer* in laboratory culture, *Journal of the Marine Biological Association of the United*
796 *Kingdom*, 61, 257-277, 10.1017/S002531540004604X, 1981.

797

798 Bé, A. W. H., Spero, H. J., and Anderson, O. R.: Effects of symbiont elimination and reinfection on the life processes of the
799 planktonic foraminifer *Globigerinoides sacculifer*, *Marine Biology*, 70, 73-86, 10.1007/BF00397298, 1982.

800

801 Bé, A. W. H., Anderson, O.R., Faber, W.W., Jr. and Caron, D.A.: Sequence of morphological and cytoplasmic changes
802 during gametogenesis in the planktonic foraminifer *Globigerinoides sacculifer* (Brady), *Micropaleontology*, 29, 310-325,
803 1983.

804

805 Bé, A. W. H. a. T., D.S.: Distribution and ecology of living planktonic foraminifera in surface waters of the Atlantic and
806 Indian oceans, in: *Micropaleontology of marine bottom sediments*, edited by: Riedel, B. M. F. a. W. R., Cambridge
807 University Press, Cambridge, 105-149, 1971.

808

809 Béjard, T. M., Rigual-Hernández, A. S., Tarruella, J. P., Flores, J. A., Sanchez-Vidal, A., Llamas-Cano, I., and Sierro, F. J.:
810 Planktonic foraminifera assemblage composition and flux dynamics inferred from an annual sediment trap record in the
811 central Mediterranean Sea, *Biogeosciences*, 21, 4051-4076, 10.5194/bg-21-4051-2024, 2024.

812

813 [Cao, Y., Lei, Y., Fang, J.K.H. and Li, T., Molecular diversity of foraminiferal eDNA in sediments and their correlations with](#)
814 [environmental factors from the Yellow Sea. *Ecological Indicators*, 142: 109294, 2022.](#)

815

816 Caromel, A. G. M., Schmidt, D. N., Fletcher, I., and Rayfield, E. J.: Morphological Change During The Ontogeny Of The
817 Planktic Foraminifera, *J. Micropalaeontol.*, 35, 2-19, 10.1144/jmpaleo2014-017, 2016.

818

819 Caromel, A. G. M., Schmidt, D. N., Phillips, J. C., and Rayfield, E. J.: Hydrodynamic constraints on the evolution and
820 ecology of planktic foraminifera, *Marine Micropaleontology*, 106, 69-78, <https://doi.org/10.1016/j.marmicro.2014.01.002>,
821 2014.

822

823 Caron, D. A., Bé, A. W. H., and Anderson, O. R.: Effects of variations in light intensity on life processes of the planktonic
824 foraminifer *Globigerinoides sacculifer* in laboratory culture, *Journal of the Marine Biological Association of the United*
825 *Kingdom*, 62, 435-451, 10.1017/S0025315400057374, 1982.

826

827 Caron, D. A., Faber, W. W., and Bé, A. W. H.: Effects of temperature and salinity on the growth and survival of the
 828 planktonic foraminifer *Globigerinoides sacculifer*, Journal of the Marine Biological Association of the United Kingdom, 67,
 829 323-341, 10.1017/S0025315400026643, 1987.
 830
 831 Chernihovsky, N., Torfstein, A., and Almogi-Labin, A.: Daily timescale dynamics of planktonic foraminifera shell-size
 832 distributions, Frontiers in Marine Science, 10, 10.3389/fmars.2023.1126398, 2023.
 833
 834 Darling, K. F. and Wade, C. M.: The genetic diversity of planktic foraminifera and the global distribution of ribosomal RNA
 835 genotypes, Marine Micropaleontology, 67, 216-238, <https://doi.org/10.1016/j.marmicro.2008.01.009>, 2008.
 836
 837 Davis, C. V., Livsey, C. M., Palmer, H. M., Hull, P. M., Thomas, E., Hill, T. M., and Benitez-Nelson, C. R.: Extensive
 838 morphological variability in asexually produced planktic foraminifera, Science Advances, 6, eabb8930,
 839 doi:10.1126/sciadv.abb8930, 2020.
 840
 841 de Nooijer, L. J., Pacho Sampedro, L., Jorissen, F. J., Pawlowski, J., Rosenthal, Y., Dissard, D., and Reichart, G. J.: 500
 842 million years of foraminiferal calcification, Earth-Science Reviews, 243, 104484,
 843 <https://doi.org/10.1016/j.earscirev.2023.104484>, 2023.
 844
 845 de Vargas, C., Renaud, S., Hilbrecht, H., and Pawlowski, J.: Pleistocene adaptive radiation in *Globorotalia truncatulinoides*:
 846 genetic, morphologic, and environmental evidence, Paleobiology, 27, 104-125, 10.1666/0094-
 847 8373(2001)027<0104:PARIGT>2.0.CO;2, 2001.
 848
 849 de Vargas, C., Audic, S., Henry, N., Decelle, J., Mahé, F., Logares, R., Lara, E., Berney, C., Le Bescot, N., Probert, I.,
 850 Carmichael, M., Poulain, J., Romac, S., Colin, S., Aury, J. M., Bittner, L., Chaffron, S., Dunthorn, M., Engelen, S., ..., and
 851 Karsenti, E.: Ocean plankton. Eukaryotic plankton diversity in the sunlit ocean, Science, 348, 1261605,
 852 10.1126/science.1261605, 2015.
 853
 854 Decelle, J., Romac, S., Stern, R. F., Bendif, E. M., Zingone, A., Audic, S., Guiry, M. D., Guillou, L., Tessier, D., Le Gall, F.,
 855 Gourvil, P., Dos Santos, A. L., Probert, I., Vaultot, D., de Vargas, C., and Christen, R.: PhytoREF: a reference database of the
 856 plastidial 16S rRNA gene of photosynthetic eukaryotes with curated taxonomy, Molecular Ecology Resources, 15, 1435-
 857 1445, <https://doi.org/10.1111/1755-0998.12401>, 2015.
 858

del Campo, J., Carlos-Oliveira, M., Čepička, I., Hehenberger, E., Horák, A., Karnkowska, A., Kolisko, M., Lara, E., Lukeš, J., Pánek, T., Piwosz, K., Richter, D. J., Škaloud, P., Sutak, R., Tachezy, J., and Hampl, V.: The protist cultural renaissance, Trends in Microbiology, 32, 128-131, <https://doi.org/10.1016/j.tim.2023.11.010>, 2024.

Deuser, W. G., Ross, E. H., Hemleben, C., and Spindler, M.: Seasonal changes in species composition, numbers, mass, size, and isotopic composition of planktonic foraminifera settling into the deep sargasso sea, Palaeogeography, Palaeoclimatology, Palaeoecology, 33, 103-127, [https://doi.org/10.1016/0031-0182\(81\)90034-1](https://doi.org/10.1016/0031-0182(81)90034-1), 1981.

[Deutsch, C., Penn, J.L. and Seibel, B.: Metabolic trait diversity shapes marine biogeography. Nature, 585, 557-562, 2020.](#)

Dieckmann, G. S., Spindler, M., Lange, M. A., Ackley, S. F., and Eicken, H.: Antarctic sea ice; a habitat for the foraminifer Neoglobobulimina pachyderma, Journal of Foraminiferal Research, 21, 182-189, 10.2113/gsjfr.21.2.182, 1991.

Dutkiewicz, S., Follows, M. J., and Bragg, J. G.: Modeling the coupling of ocean ecology and biogeochemistry, Global Biogeochemical Cycles, 23, <https://doi.org/10.1029/2008GB003405>, 2009.

Dutkiewicz, S., Cermen, P., Jahn, O., Follows, M. J., Hickman, A. E., Taniguchi, D. A. A., and Ward, B. A.: Dimensions of marine phytoplankton diversity, Biogeosciences, 17, 609-634, 10.5194/bg-17-609-2020, 2020.

Edgar, K. M., Bohaty, S. M., Gibbs, S. J., Sexton, P. F., Norris, R. D., and Wilson, P. A.: Symbiont ‘bleaching’ in planktic foraminifera during the Middle Eocene Climatic Optimum, Geology, 41, 15-18, 10.1130/g33388.1, 2013.

Eiane, K. and Ohman, M., D. : Stage-specific mortality of Calanus finmarchicus, Pseudocalanus elongatus and Oithona similis on Fladen Ground, North Sea, during a spring bloom, Marine Ecology Progress Series, 268, 183-193, 2004.

Erez, J.: The Source of Ions for Biomineralization in Foraminifera and Their Implications for Paleooceanographic Proxies, Reviews in Mineralogy and Geochemistry, 54, 115-149, 10.2113/0540115, 2003.

Faber, W. W., Anderson, O. R., and Caron, D. A.: Algal-foraminiferal symbiosis in the planktonic foraminifer Globobulimina aequilateralis; II, Effects of two symbiont species on foraminiferal growth and longevity, Journal of Foraminiferal Research, 19, 185-193, 10.2113/gsjfr.19.3.185, 1989.

Fenchel, T.: The Ecology of Heterotrophic Microflagellates, in: Advances in Microbial Ecology, edited by: Marshall, K. C., Springer US, Boston, MA, 57-97, 10.1007/978-1-4757-0611-6_2, 1986.

893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926

Fenton, I. S., Aze, T., Farnsworth, A., Valdes, P., and Saupe, E. E.: Origination of the modern-style diversity gradient 15 million years ago, *Nature*, 614, 708-712, 10.1038/s41586-023-05712-6, 2023.

Fenton, I. S., Woodhouse, A., Aze, T., Lazarus, D., Renaudie, J., Dunhill, A. M., Young, J. R., and Saupe, E. E.: Triton, a new species-level database of Cenozoic planktonic foraminiferal occurrences, *Sci Data*, 8, 160, 10.1038/s41597-021-00942-7, 2021.

Follows, M. J., Dutkiewicz, S., Grant, S., and Chisholm, S. W.: Emergent biogeography of microbial communities in a model ocean, *Science*, 315, 1843-1846, 10.1126/science.1138544, 2007.

Gaskell, D. E., Ohman, M. D., and Hull, P. M.: Zooglider-Based Measurements of Planktonic Foraminifera in the California Current System, *Journal of Foraminiferal Research*, 49, 390-404, 10.2113/gsjfr.49.4.390, 2019.

Gast, R. J. and Caron, D. A.: Molecular phylogeny of symbiotic dinoflagellates from planktonic foraminifera and radiolaria, *Molecular Biology and Evolution*, 13, 1192-1197, 10.1093/oxfordjournals.molbev.a025684, 1996.

Gaylord, B., Kroeker, K. J., Sunday, J. M., Anderson, K. M., Barry, J. P., Brown, N. E., Connell, S. D., Dupont, S., Fabricius, K. E., Hall-Spencer, J. M., Klinger, T., Milazzo, M., Munday, P. L., Russell, B. D., Sanford, E., Schreiber, S. J., Thiagarajan, V., Vaughan, M. L. H., Widdicombe, S., and Harley, C. D. G.: Ocean acidification through the lens of ecological theory, *Ecology*, 96, 3-15, <https://doi.org/10.1890/14-0802.1>, 2015.

Gibbs, S. J., Bown, P. R., Ward, B. A., Alvarez, S. A., Kim, H., Archontikis, O. A., Sauterey, B., Poulton, A. J., Wilson, J., and Ridgwell, A.: Algal plankton turn to hunting to survive and recover from end-Cretaceous impact darkness, *Sci Adv*, 6, 10.1126/sciadv.abc9123, 2020.

[Goldstein, S.T.: Foraminifera: A biological review in Modern Foraminifera, Sen Gupta, B.K. Eds., Kluwer Academic Publishers, Dordrecht, 37-55, 1999.](#)

Greco, M., Morard, R., and Kucera, M.: Single-cell metabarcoding reveals biotic interactions of the Arctic calcifier ^with the eukaryotic pelagic community, *Journal of Plankton Research*, 43, 113-125, 10.1093/plankt/fbab015, 2021.

Grice, G. D. and Hart, A. D.: The Abundance, Seasonal Occurrence and Distribution of the Epizooplankton between New York and Bermuda, *Ecological Monographs*, 32, 287-309, 10.2307/1942377, 1962.

927

928 Grigoratou, M., Monteiro, F. M., Ridgwell, A., and Schmidt, D. N.: Investigating the benefits and costs of spines and diet on
 929 planktonic foraminifera distribution with a trait-based ecosystem model, *Marine Micropaleontology*, 166, 102004,
 930 <https://doi.org/10.1016/j.marmicro.2021.102004>, 2021.

931

932 Grigoratou, M., Monteiro, F. M., Wilson, J. D., Ridgwell, A., and Schmidt, D. N.: Exploring the impact of climate change on
 933 the global distribution of non-spinose planktonic foraminifera using a trait-based ecosystem model, *Glob Chang Biol*, 28,
 934 1063-1076, 10.1111/gcb.15964, 2022.

935

936 Grigoratou, M., Monteiro, F. M., Schmidt, D. N., Wilson, J. D., Ward, B. A., and Ridgwell, A.: A trait-based modelling
 937 approach to planktonic foraminifera ecology, *Biogeosciences*, 16, 1469-1492, 10.5194/bg-16-1469-2019, 2019.

938

939 Haldane, J. B. S.: *On Being the Right Size*, 1926.

940

941 [Hemleben, C., Spindler, M., Breiting, I., and Deuser, W.: Field and laboratory studies on the ontogeny and ecology of
 942 some globorotaliid species from the Sargasso Sea off Bermuda, *Journal of Foraminiferal Research*, 15, 254-272,
 943 \[10.2113/gsjfr.15.4.254\]\(https://doi.org/10.2113/gsjfr.15.4.254\), 1985.](#)

944

945 Hemleben, C., Spindler, M., Anderson, R.O.: *Modern Planktonic Foraminifera*, Springer-Verlag, New York, Berlin,
 946 Heidelberg, 363 pp., 1989.

947

948 Henchan, M. J., Evans, D., Shankle, M., Burke, J. E., Foster, G. L., Anagnostou, E., Chalk, T. B., Stewart, J. A., Alt, C. H.
 949 S., Durrant, J., and Hull, P. M.: Size-dependent response of foraminiferal calcification to seawater carbonate chemistry,
 950 *Biogeosciences*, 14, 3287-3308, 10.5194/bg-14-3287-2017, 2017.

951

952 Hill, T. S. and Hoogenboom, M. O.: The indirect effects of ocean acidification on corals and coral communities, *Coral*
 953 *Reefs*, 41, 1557-1583, 10.1007/s00338-022-02286-z, 2022.

954

955 Hoadley, K. D., Lewis, A. M., Wham, D. C., Pettay, D. T., Grasso, C., Smith, R., Kemp, D. W., LaJeunesse, T. C., and
 956 Warner, M. E.: Host–symbiont combinations dictate the photo-physiological response of reef-building corals to thermal
 957 stress, *Scientific Reports*, 9, 9985, 10.1038/s41598-019-46412-4, 2019.

958

959 Hoppit, G. and Schmidt, D. N.: A Regional View of the Response to Climate Change: A Meta-Analysis of European Benthic
 960 Organisms’ Responses, *Frontiers in Marine Science*, 9, 10.3389/fmars.2022.896157, 2022.

961

962 Horne, C. R., Hirst, A. G., and Atkinson, D.: Seasonal body size reductions with warming covary with major body size

963 gradients in arthropod species, *Proceedings of the Royal Society B: Biological Sciences*, 284, 20170238,

964 doi:10.1098/rspb.2017.0238, 2017.

965

966 [Hoshino, T. and Inagaki, F., Distribution of eukaryotic environmental DNA in global subseafloor sediments, *Progress in*](#)

967 [Earth and Planetary Science](#), 11, <https://doi.org/10.1186/s40645-024000621-2>, 2024.

968

969 Hull, P. M., Osborn, K. J., Norris, R. D., and Robison, B. H.: Seasonality and depth distribution of a mesopelagic

970 foraminifer, *Hastigerinella digitata*, in Monterey Bay, California, *Limnology and Oceanography*, 56, 562-576,

971 <https://doi.org/10.4319/lo.2011.56.2.0562>, 2011.

972

973 Jonkers, L. and Kučera, M.: Global analysis of seasonality in the shell flux of extant planktonic Foraminifera,

974 *Biogeosciences*, 12, 2207-2226, 10.5194/bg-12-2207-2015, 2015.

975

976 Jonkers, L., Reynolds, C. E., Richey, J., and Hall, I. R.: Lunar periodicity in the shell flux of planktonic foraminifera in the

977 Gulf of Mexico, *Biogeosciences*, 12, 3061-3070, 10.5194/bg-12-3061-2015, 2015.

978

979 Kimoto, K.: The "unusual" reproduction of planktonic foraminifera: an Asexual reproductive phase of *Neogloboquadrina*

980 *pachyderma* (Ehrenberg), *Anuario do Instituto de Geociencias, Universidade Federal do Rio de Janeiro*, 29, 461, 2006.

981

982 Kjørboe, T.: How zooplankton feed: mechanisms, traits and trade-offs, *Biological Reviews*, 86, 311-339,

983 <https://doi.org/10.1111/j.1469-185X.2010.00148.x>, 2011.

984

985 Kjørboe, T., Visser, A., and Andersen, K. H.: A trait-based approach to ocean ecology, *ICES Journal of Marine Science*, 75,

986 1849-1863, 10.1093/icesjms/fsy090, 2018.

987

988 Knoll, A. H. and Follows, M. J.: A bottom-up perspective on ecosystem change in Mesozoic oceans, *Proceedings of the*

989 *Royal Society B: Biological Sciences*, 283, 20161755, doi:10.1098/rspb.2016.1755, 2016.

990

991 Kucera, M.: Chapter Six Planktonic Foraminifera as Tracers of Past Oceanic Environments, *Developments in Marine*

992 *Geology*, Volume 1, 213-262, 10.1016/S1572-5480(07)01011-1, 2007.

993

994 Köhler-Rink, S. and Kühl, M.: The chemical microenvironment of the symbiotic planktonic foraminifer *Orbulina universa*,
 995 Marine Biology Research, 1, 68-78, 10.1080/17451000510019015, 2005.

996

997 LeKieffre, C., Spero, H. J., Russell, A. D., Fehrenbacher, J. S., Geslin, E., and Meibom, A.: Assimilation, translocation, and
 998 utilization of carbon between photosynthetic symbiotic dinoflagellates and their planktic foraminifera host, Marine Biology,
 999 165, 104, 10.1007/s00227-018-3362-7, 2018.

000

001 Leung, J. Y. S., Zhang, S., and Connell, S. D.: Is Ocean Acidification Really a Threat to Marine Calcifiers? A Systematic
 002 Review and Meta-Analysis of 980+ Studies Spanning Two Decades, Small, 18, 2107407,
 003 <https://doi.org/10.1002/sml.202107407>, 2022.

004

005 Litchman, E., Edwards, K. F., and Boyd, P. W.: Toward trait-based food webs: Universal traits and trait matching in
 006 planktonic predator–prey and host–parasite relationships, Limnology and Oceanography, 66, 3857-3872,
 007 <https://doi.org/10.1002/lno.11924>, 2021.

008

009 Litchman, E., Ohman, M. D., and Kiørboe, T.: Trait-based approaches to zooplankton communities, Journal of Plankton
 010 Research, 35, 473-484, 10.1093/plankt/fbt019, 2013.

011

012 Lombard, F., da Rocha, R. E., Bijma, J., and Gattuso, J. P.: Effect of carbonate ion concentration and irradiance on
 013 calcification in planktonic foraminifera, Biogeosciences, 7, 247-255, 10.5194/bg-7-247-2010, 2010.

014

015 Lončarić, N., Brummer, G.-J. A., and Kroon, D.: Lunar cycles and seasonal variations in deposition fluxes of planktic
 016 foraminiferal shell carbonate to the deep South Atlantic (central Walvis Ridge), Deep Sea Research Part I: Oceanographic
 017 Research Papers, 52, 1178-1188, <https://doi.org/10.1016/j.dsr.2005.02.003>, 2005.

018

019 Lynch, M.: The Genetic interpretation of inbreeding depression and outbreeding depression, Evolution, 45, 622-629,
 020 10.1111/j.1558-5646.1991.tb04333.x, 1991.

021

022 [Malviya, S., Scalco, E., Audic, S., Vincent, F., Veluchamy, A., Poulain, J., Wincker, P., Iudicone, D., de Vargas, C., Bittner,](#)
 023 [L., Zingone, A. and Bowler, C., Insights into global diatom distribution and diversity in the world’s ocean. Proceedings of](#)
 024 [the National Academy of Sciences, 113\(11\): E1516-E1525, 2016.](#)

025

026 Margulis, L.: Origins of species: acquired genomes and individuality, *Biosystems*, 31, 121-125,
027 [https://doi.org/10.1016/0303-2647\(93\)90039-F](https://doi.org/10.1016/0303-2647(93)90039-F), 1993.

028

029 McKinney, M. L.: Trends in body-size evolution, in: *Evolutionary Trends*, edited by: McNamara, K. J., Belhaven Press,
030 London, 75-118, 1990.

031

032 Meilland, J., Howa, H., Lo Monaco, C., and Schiebel, R.: Individual planktic foraminifer protein-biomass affected by trophic
033 conditions in the Southwest Indian Ocean, 30°S–60°S, *Marine Micropaleontology*, 124, 63-74,
034 <https://doi.org/10.1016/j.marmicro.2016.02.004>, 2016.

035

036 Meilland, J., Siccha, M., Morard, R., and Kucera, M.: Continuous reproduction of planktonic foraminifera in laboratory
037 culture, *Journal of Eukaryotic Microbiology*, 71, e13022, <https://doi.org/10.1111/jeu.13022>, 2024.

038

039 Meilland, J., Ezat, M. M., Westgård, A., Manno, C., Morard, R., Siccha, M., and Kucera, M.: Rare but persistent asexual
040 reproduction explains the success of planktonic foraminifera in polar oceans, *Journal of Plankton Research*, 45, 15-32,
041 10.1093/plankt/fbac069, 2023~~2~~.

042

043 Michaels, A. F., Caron, D. A., Swanberg, N. R., Howse, F., and Michaels, C. M.: Planktonic sarcodines (Acantharia,
044 Radiolaria, Foraminifera) in surface waters near Bermuda: abundance, biomass and vertical flux, *Journal of Plankton*
045 *Research*, 17, 131-163, 1995.

046

047 Monteiro, F. M., Bach, L. T., Brownlee, C., Bown, P., Rickaby, R. E. M., Poulton, A. J., Tyrrell, T., Beaufort, L.,
048 Dutkiewicz, S., Gibbs, S., Gutowska, M. A., Lee, R., Riebesell, U., Young, J., and Ridgwell, A.: Why marine phytoplankton
049 calcify, *Science Advances*, 2, e1501822, doi:10.1126/sciadv.1501822, 2016.

050

051 Morard, R., Escarguel, G., Weiner, A. K., André, A., Douady, C. J., Wade, C. M., Darling, K. F., Ujjié, Y., Seears, H. A.,
052 Quillévéré, F., de Garidel-Thoron, T., de Vargas, C., and Kucera, M.: Nomenclature for the Nameless: A Proposal for an
053 Integrative Molecular Taxonomy of Cryptic Diversity Exemplified by Planktonic Foraminifera, *Syst Biol*, 65, 925-940,
054 10.1093/sysbio/syw031, 2016.

055

056 Morard, R., Darling, K. F., Weiner, A. K. M., Hassenrück, C., Vanni, C., Cordier, T., Henry, N., Greco, M., Vollmar, N. M.,
057 Milivojevic, T., Rahman, S. N., Siccha, M., Meilland, J., Jonkers, L., Quillévéré, F., Escarguel, G., Douady, C. J., de
058 Garidel-Thoron, T., de Vargas, C., and Kucera, M.: The global genetic diversity of planktonic foraminifera reveals the
059 structure of cryptic speciation in plankton, *Biological Reviews*, 99, 1218-1241, <https://doi.org/10.1111/brv.13065>, 2024.

060

061 Naidoo-Bagwell, A. A., Monteiro, F. M., Hendry, K. R., Burgan, S., Wilson, J. D., Ward, B. A., Ridgwell, A., and Conley,
 062 D. J.: A diatom extension to the cGEnIE Earth system model – EcoGEnIE 1.1, *Geosci. Model Dev.*, 17, 1729-1748,
 063 10.5194/gmd-17-1729-2024, 2024.

064

065 Nehrke, G., Keul, N., Langer, G., de Nooijer, L. J., Bijma, J., and Meibom, A.: A new model for biomineralization and trace-
 066 element signatures of Foraminifera tests, *Biogeosciences*, 10, 6759-6767, 10.5194/bg-10-6759-2013, 2013.

067

068 Neukermans, G., Bach, L. T., Butterley, A., Sun, Q., Claustre, H., and Fournier, G. R.: Quantitative and mechanistic
 069 understanding of the open ocean carbonate pump - perspectives for remote sensing and autonomous in situ observation,
 070 *Earth-Science Reviews*, 239, 104359, <https://doi.org/10.1016/j.earscirev.2023.104359>, 2023.

071

072 Ortiz, J. D., Mix, A. C., and Collier, R. W.: Environmental control of living symbiotic and asymbiotic foraminifera of the
 073 California Current, *Paleoceanography*, 10, 987-1009, <https://doi.org/10.1029/95PA02088>, 1995.

074

075 Otto, S. P. and Lenormand, T.: Resolving the paradox of sex and recombination, *Nature Reviews Genetics*, 3, 252-261,
 076 10.1038/nrg761, 2002.

077

078 Pallacks, S., Ziveri, P., Schiebel, R., Vonhof, H., Rae, J. W. B., Littley, E., Garcia-Orellana, J., Langer, G., Grelaud, M., and
 079 Martrat, B.: Anthropogenic acidification of surface waters drives decreased biogenic calcification in the Mediterranean Sea,
 080 *Communications Earth & Environment*, 4, 301, 10.1038/s43247-023-00947-7, 2023.

081

082 Palmer, A. R.: Calcification in marine molluscs: how costly is it?, *Proc Natl Acad Sci U-S-A*, 89, 1379-1382,
 083 10.1073/pnas.89.4.1379, 1992.

084

085 [Pawlowski, J., Lejzerowicz, F. and Esling, P.: Next-Generation Environmental Diversity Surveys of Foraminifera: Preparing](#)
 086 [the Future, *The Biological Bulletin*, 227\(2\): 93-106, 2014.](#)

087

088 Pearson, P. N., Ditchfield, P. W., Singano, J., Harcourt-Brown, K. G., Nicholas, C. J., Olsson, R. K., Shackleton, N. J., and
 089 Hall, M. A.: Warm tropical sea surface temperatures in the Late Cretaceous and Eocene epochs, *Nature*, 413, 481-487,
 090 10.1038/35097000, 2001.

091

092 Peters, R. H.: *The Ecological Implications of Body Size*, Cambridge Studies in Ecology, Cambridge University Press,
 093 Cambridge, DOI: 10.1017/CBO9780511608551, 1983.

094

095 Rebotim, A., Voelker, A. H. L., Jonkers, L., Wanick, J. J., Schulz, M., and Kucera, M.: Calcification depth of deep-dwelling
096 planktonic foraminifera from the eastern North Atlantic constrained by stable oxygen isotope ratios of shells from stratified
097 plankton tows, *J. Micropalaeontol.*, 38, 113-131, 10.5194/jm-38-113-2019, 2019.

098

099 Ridgwell, A. and Zeebe, R. E.: The role of the global carbonate cycle in the regulation and evolution of the Earth system,
100 *Earth and Planetary Science Letters*, 234, 299-315, <https://doi.org/10.1016/j.epsl.2005.03.006>, 2005.

101

102 [Riebesell, U., Fabry, V.J., Hansson, L. and Gattuso, J.-P. \(Editors\): Guide to best practices for ocean acidification research](#)
103 [and data reporting. Publications Office of the European Union, Luxembourg, 260, 2010.](#)

104

105 Rillo, M. C., Miller, C. G., Kucera, M., and Ezard, T. H. G.: Intraspecific size variation in planktonic foraminifera cannot be
106 consistently predicted by the environment, *Ecology and Evolution*, 10, 11579-11590, <https://doi.org/10.1002/ece3.6792>,
107 2020.

108

109 Rink, S., Kühl, M., Bijma, J., and Spero, H. J.: Microsensor studies of photosynthesis and respiration in the symbiotic
110 foraminifer *Orbulina universa*, *Marine Biology*, 131, 583-595, 10.1007/s002270050350, 1998.

111

112 [Ruppert, K.M., Kline, R.J. and Rahman, M.S., Past, present, and future perspectives of environmental DNA \(eDNA\)](#)
113 [metabarcoding: A systematic review in methods, monitoring, and applications of global eDNA. *Global Ecology and*](#)
114 [Conservation, 17: e00547, 2019.](#)

115

116 Russell, A. D., Hönisch, B., Spero, H. J., and Lea, D. W.: Effects of seawater carbonate ion concentration and temperature
117 on shell U, Mg, and Sr in cultured planktonic foraminifera, *Geochimica et Cosmochimica Acta*, 68, 4347-4361,
118 <https://doi.org/10.1016/j.gca.2004.03.013>, 2004.

119

120 Sanders, T., Schmittmann, L., Nascimento-Schulze, J. C., and Melzner, F.: High Calcification Costs Limit Mussel Growth at
121 Low Salinity, *Frontiers in Marine Science*, 5, 10.3389/fmars.2018.00352, 2018.

122

123 Schiebel, R. and Hemleben, C.: *Planktic Foraminifers in the Modern Ocean*, Springer, Heidelberg 2017.

124

125 Schiebel, R., Hiller, B., and Hemleben, C.: Impacts of storms on Recent planktic foraminiferal test production and CaCO₃
126 flux in the North Atlantic at 47 °N, 20 °W (JGOFS), *Marine Micropaleontology*, 26, 115-129, [https://doi.org/10.1016/0377-](https://doi.org/10.1016/0377-8398(95)00035-6)
127 [8398\(95\)00035-6](https://doi.org/10.1016/0377-8398(95)00035-6), 1995.

- Schmidt, D. N., Renaud, S., and Bollmann, J.: Response of planktic foraminiferal size to late Quaternary climate change, *Paleoceanography*, 18, <https://doi.org/10.1029/2002PA000831>, 2003.
- Schmidt, D. N., Lazarus, D., Young, J. R., and Kucera, M.: Biogeography and evolution of body size in marine plankton, *Earth-Science Reviews*, 78, 239-266, <https://doi.org/10.1016/j.earscirev.2006.05.004>, 2006.
- Schmidt, D. N., Renaud, S., Bollmann, J., Schiebel, R., and Thierstein, H. R.: Size distribution of Holocene planktic foraminifer assemblages: biogeography, ecology and adaptation, *Marine Micropaleontology*, 50, 319-338, [https://doi.org/10.1016/S0377-8398\(03\)00098-7](https://doi.org/10.1016/S0377-8398(03)00098-7), 2004.
- [Schweizer, M., Jauffrais, T., Choquel, C., Méléder, V., Quinchar, S. and Geslin, E., Trophic strategies of intertidal foraminifera explored with single-cell microbiome metabarcoding and morphological methods: What is on the menu? Ecology and Evolution, 12\(11\): e9437, 2022.](#)
- Seibold, E. a. B., W.H.: *The Sea Floor*, Springer Verlag, Berlin, Heidelberg, New York, 356 pp.1993.
- Siccha, M. and Kucera, M.: ForCenS, a curated database of planktonic foraminifera census counts in marine surface sediment samples, *Scientific Data*, 4, 170109, 10.1038/sdata.2017.109, 2017.
- [Ser-Giacomi, E., Zinger, L., Malviya, S., De Vargas, C., Karsenti, E., Bowler, C. and De Monte, S., Ubiquitous abundance distribution of non-dominant plankton across the global ocean. Nature Ecology & Evolution, 2\(8\): 1243-1249, 2018.](#)
- Spero, H. J. and Parker, S. L.: Photosynthesis in the symbiotic planktonic foraminifer *Orbulina universa*, and its potential contribution to oceanic primary productivity, *Journal of Foraminiferal Research*, 15, 273-281, 10.2113/gsjfr.15.4.273, 1985.
- Spindler, M.: On the salinity tolerance of the planktonic foraminifer *Neogloboquadrina pachyderma* from Antarctic sea ice, *Polar Biology*, 9, 85-91, 1996.
- Spindler, M. and Dieckmann, G. S.: Distribution and abundance of the planktic foraminifer *Neogloboquadrina pachyderma* in sea ice of the Weddell Sea (Antarctica), *Polar Biology*, 5, 185-191, 10.1007/BF00441699, 1986.
- Spindler, M., Hemleben, C., Salomons, J. B., and Smit, L. P.: Feeding behavior of some planktonic foraminifers in laboratory cultures, *Journal of Foraminiferal Research*, 14, 237-249, 10.2113/gsjfr.14.4.237, 1984.

- Spindler, M., Hernleben, C., Bayer, U., Bé, A. W. H., Anderson, O.R.: Lunar Periodicity of Reproduction in the Planktonic Foraminifer *Hastigerina pelagica*, Marine Ecology Progress Series, 1, 61-64, 1979.
- Stoecker, D. K., Johnson, M. D., deVargas, C., and Not, F.: Acquired phototrophy in aquatic protists, Aquatic Microbial Ecology, 57, 279-310, 2009.
- Strack, T., Jonkers, L., C. Rillo, M., Baumann, K.-H., Hillebrand, H., and Kucera, M.: Coherent response of zoo- and phytoplankton assemblages to global warming since the Last Glacial Maximum, Global Ecology and Biogeography, 33, e13841, <https://doi.org/10.1111/geb.13841>, 2024.
- Swezey, D. S., Boles, S. E., Aquilino, K. M., Stott, H. K., Bush, D., Whitehead, A., Rogers-Bennett, L., Hill, T. M., and Sanford, E.: Evolved differences in energy metabolism and growth dictate the impacts of ocean acidification on abalone aquaculture, Proc Natl Acad Sci U S A, 117, 26513-26519, 10.1073/pnas.2006910117, 2020.
- Sykes, F. E., Meilland, J., Westgård, A., Chalk, T. B., Chierici, M., Foster, G. L., and Ezat, M. M.: Large-scale culturing of the subpolar foraminifera *Globigerina bulloides* reveals tolerance to a large range of environmental parameters associated to different life-strategies and an extended lifespan, Journal of Plankton Research, 46, 403-420, 10.1093/plankt/fbae029, 2024.
- Takagi, H., Kurasawa, A., and Kimoto, K.: Observation of asexual reproduction with symbiont transmission in planktonic foraminifera, Journal of Plankton Research, 42, 403-410, 10.1093/plankt/fbaa033, 2020.
- Takagi, H., Kimoto, K., Fujiki, T., and Moriya, K.: Effect of nutritional condition on photosymbiotic consortium of cultured *Globigerinoides sacculifer* (Rhizaria, Foraminifera), Symbiosis, 76, 25-39, 10.1007/s13199-017-0530-3, 2018.
- Takagi, H., Kimoto, K., Fujiki, T., Kurasawa, A., Moriya, K., and Hirano, H.: Ontogenetic dynamics of photosymbiosis in cultured planktic foraminifers revealed by fast repetition rate fluorometry, Marine Micropaleontology, 122, 44-52, <https://doi.org/10.1016/j.marmicro.2015.10.003>, 2016.
- Takagi, H., Kimoto, K., Fujiki, T., Saito, H., Schmidt, C., Kucera, M., and Moriya, K.: Characterizing photosymbiosis in modern planktonic foraminifera, Biogeosciences, 16, 3377-3396, 10.5194/bg-16-3377-2019, 2019.
- Takahashi, K. and Be, A. W. H.: Planktonic foraminifera: factors controlling sinking speeds, Deep Sea Research Part A. Oceanographic Research Papers, 31, 1477-1500, [https://doi.org/10.1016/0198-0149\(84\)90083-9](https://doi.org/10.1016/0198-0149(84)90083-9), 1984.

196 Tang, Y. Z., Gu, H., Wang, Z., Liu, D., Wang, Y., Lu, D., Hu, Z., Deng, Y., Shang, L., and Qi, Y.: Exploration of resting
 197 cysts (stages) and their relevance for possibly HABs-causing species in China, *Harmful Algae*, 107, 102050,
 198 <https://doi.org/10.1016/j.hal.2021.102050>, 2021.

199

200 [Thomsen, J., Casties, I., Pansch, C., Körtzinger, A. and Melzner, F.: Food availability outweighs ocean acidification effects](#)
 201 [in juvenile *Mytilus edulis*: laboratory and field experiments. *Global Change Biology*, 19\(4\): 1017-1027, 2013.](#)

202

203 Topa, P., Faber, Ł., Tyszk, J., and Komosinski, M.: Modelling ecology and evolution of Foraminifera in the agent-oriented
 204 distributed platform, *Journal of Computational Science*, 18, 69-84, <https://doi.org/10.1016/j.jocs.2016.07.009>, 2017.

205

206 Vincent, E. a. B., W.H.: Planktonic foraminifera and their use in paleoceanography, in: *The Sea*, edited by: Emiliani, C.,
 207 John Wiley & Sons, New York, 1035-1119, 1981.

208

209 Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., and Garnier, E.: Let the concept of trait be
 210 functional!, *Oikos*, 116, 882-892, <https://doi.org/10.1111/j.0030-1299.2007.15559.x>, 2007.

211

212 Wade, B. S., Al-Sabouni, N., Hemleben, C., and Kroon, D.: Symbiont bleaching in fossil planktonic foraminifera,
 213 *Evolutionary Ecology*, 22, 253-265, 10.1007/s10682-007-9176-6, 2008.

214

215 Ward, B. B.: Oceans. How nitrogen is lost, *Science*, 341, 352-353, 10.1126/science.1240314, 2013.

216

217 Watson, S.-A., Morley, S. A., and Peck, L. S.: Latitudinal trends in shell production cost from the tropics to the poles,
 218 *Science Advances*, 3, e1701362, doi:10.1126/sciadv.1701362, 2017.

219

220 Weinkauff, M. F. G., Siccha, M., and Weiner, A. K. M.: Reproduction dynamics of planktonic microbial eukaryotes in the
 221 open ocean, *Journal of The Royal Society Interface*, 19, 20210860, doi:10.1098/rsif.2021.0860, 2022.

222

223 Weinkauff, M. F. G., Kunze, J.G., Waniek, J.J. and Kučera, M.: Seasonal Variation in Shell Calcification of Planktonic
 224 Foraminifera in the NE Atlantic Reveals Species-Specific Response to Temperature, Productivity, and Optimum Growth
 225 Conditions, *PLOS ONE*, 11, e0148363, 2016.

226

227 [Westgård, A., Ezat, M.E., Chalk, T.B., Chierici, M., Foster, G.L. and Meilland, J. Large-scale culturing of](#)
228 [Neogloboquadrina pachyderma, its growth in, and tolerance of, variable environmental conditions, Journal of Plankton](#)
229 [Research, 45, 732-745, 2023.](#)
230
231 Westoby, M.: Trait-based ecology, trait-free ecology, and in between, New Phytol, 10.1111/nph.20197, 2024.
232
233 Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da
234 Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S.,
235 Evelo, C. T., Finkers, R., ..., and Mons, B.: The FAIR Guiding Principles for scientific data management and stewardship,
236 Scientific Data, 3, 160018, 10.1038/sdata.2016.18, 2016.
237
238 Wirtz, K. W.: Who is eating whom? Morphology and feeding type determine the size relation between planktonic predators
239 and their ideal prey, Marine Ecology Progress Series, 445, 1-12, 2012.
240
241 Woodhouse, A., Swain, A., Fagan, W. F., Fraass, A. J., and Lowery, C. M.: Late Cenozoic cooling restructured global
242 marine plankton communities, Nature, 614, 713-718, 10.1038/s41586-023-05694-5, 2023.
243
244 Ying, R., Monteiro, F. M., Wilson, J. D., and Schmidt, D. N.: ForamEcoGENIE 2.0: incorporating symbiosis and spine traits
245 into a trait-based global planktic foraminiferal model, Geosci. Model Dev., 16, 813-832, 10.5194/gmd-16-813-2023, 2023.
246
247 Ying, R., Monteiro, F.M., Wilson, J.D., Ödalen, M. and Schmidt, D.N.: Past foraminiferal acclimation capacity is limited
248 during future climate warming, Nature, 2024.
249
250 Yonathan, S. and Colomban de, V.: Pelagic photosymbiosis: rDNA assessment of diversity and evolution of dinoflagellate
251 symbionts and planktonic foraminiferal hosts, Marine Ecology Progress Series, 325, 59-71, 2006.
252