

Contribution of sulfate to aerobic methane oxidation in upland soils: a mini-review

Rui Su¹, Kexin Li^{2,3}, Nannan Wang², Fenghui Yuan², Ying Zhao¹, Yunjiang Zuo², Ying Sun², Liyuan He⁴, Xiaofeng Xu^{4*}, Lihua Zhang^{1,2,5*}

¹College of Life and Environmental Sciences, Minzu University of China, Beijing, 100081, China

²Key Laboratory of Wetland Ecology and Environment, Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, Changchun, Jilin, 130102, China

³University of Chinese Academy of Sciences, Beijing, 100049, China

⁴Biology Department, San Diego State University, San Diego, CA, 92182, USA

⁵State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China

Corresponding authors: L.Z. (zhanglihua@muc.edu.cn); X.X. (xxu@sdsu.edu)

Abstract. Methane (CH₄) is a potent greenhouse gas, and its global warming potential is 25 times higher than carbon dioxide (CO₂). Various environmental factors influence aerobic CH₄ oxidation in soil. Sulfate (SO₄²⁻) ion is the main component of atmospheric deposition and has been increasing in recent years. It promotes CH₄ production and anaerobic CH₄ oxidation, however, the impact of SO₄²⁻ on aerobic CH₄ oxidation in soils has not yet been comprehensively summarized. We synthesize current research on the effects of SO₄²⁻ on aerobic CH₄ oxidation, examining both its macroscopic manifestations and microscale pathways. Through a literature review, we found that SO₄²⁻ enhances aerobic CH₄ oxidation by up to 0–42%, moreover, it has been found that various physicochemical properties and processes in the soil are influenced by the addition of SO₄²⁻, which in turn affects aerobic CH₄ oxidation. This review enhances our understanding of the role of SO₄²⁻ in promoting aerobic CH₄ oxidation and lays the foundation for future studies aimed at validating these findings by quantifying CH₄ flux and oxidation rates, as well as elucidating the underlying microbial processes through experimental research, while also providing directions for further investigation of SO₄²⁻'s impact on aerobic CH₄ oxidation.

1 Introduction

删除了: Reviews and syntheses:

删除了: ,

删除了: and

删除了: v

删除了: ,

删除了: i

删除了: remains inconclusive

删除了: Due to the limited research on the effects of SO₄²⁻ on CH₄ oxidation,

删除了: w

删除了: direct impact and its influence on the dynamics of soil substances, and the potential indirect effects of SO₄²⁻ on CH₄ oxidation.

删除了:

删除了: identified that SO₄²⁻ facilitates CH₄ oxidation within a range of 3

删除了: -

删除了: via

删除了: .

设置了格式: 下标

设置了格式: 上标

删除了: This review deepens the comprehension of atmospheric CH₄ flux and the global CH₄ cycle, particularly in the context of potential global environmental changes.

设置了格式: 下标

CH₄ is an important greenhouse gas, and its atmospheric concentration has increased since pre-industrial times (Place, 2024; Praeg et al., 2016). Its global warming potential is 28 times higher than carbon dioxide (CO₂), owing to its superior heat absorption efficiency (IPCC, 2013). Methanotrophs (aerobic methanotrophs) consume CH₄ under certain conditions (Le Mer and Roger, 2001), reducing CH₄ atmospheric concentration (Singh et al., 2010). Consequently, methanotrophs are crucial microbes that play an indispensable role in regulating and mitigating the greenhouse effect on Earth. Soil aerobic CH₄ oxidation is the sole known biological sink for atmospheric CH₄ (Ho et al., 2019; Murguia-Flores et al., 2018), contributing to 5%–7% of the global annual atmospheric CH₄ uptake (Saunois et al., 2020). Upland soils are the primary biological CH₄ sink (Bodelier, 2011; Guo et al., 2023), owing to methanotroph-mediated CH₄ consumption (Song et al., 2024). This represents the second-largest atmospheric CH₄ consumption sink, surpassed only by hydroxyl radical depletion (Deng et al., 2019). Aerobic CH₄ oxidation in soils are, influenced by many factors, such as soil water content, soil texture, soil type, temperature, soil pH, soil inorganic nitrogen content, metal availability, etc., many of these factors have been extensively reviewed (Shukla et al., 2013; Mishra et al., 2018). However, the effect of SO₄²⁻, a significant ion component of acid deposition, on aerobic CH₄ oxidation has not yet been reviewed. Acid rain, characterized by the deposition of SO₄²⁻ and other acidic compounds, has been a significant environmental issue (Chen et al., 2020; Qi et al., 2022). SO₄²⁻ is the major ion in acid rain (Wright and Henriksen, 1978) and has profound impacts on substances and biochemical processes in soils. As a crucial component of terrestrial ecosystems, soils serve as the ultimate receptor of acid deposition. SO₄²⁻ deposition induces soil acidification (Huang et al., 2019), alters soil plant diversity (Li et al., 2022), affects microbial properties (Wang et al., 2018), and limits grass yield potential (Klessa et al., 1989), as well as a reduction in the activities of soil enzymes such as cellulase, invertase, and polyphenol oxidase (Tie et al., 2020). SO₄²⁻ can inhibit CH₄ production (methanogenesis) and promote anaerobic CH₄ oxidation, playing a crucial role in the CH₄ cycle. SO₄²⁻ suppresses methanogenesis primarily due to its thermodynamic and

删除了: 25

删除了: Yang et al., 2023

删除了: Murguia-Flores et al., 2021

删除了: reaching approximately 30 teragrams per year (Tg yr⁻¹)

删除了: radicals

删除了: is

删除了: (Shukla et al., 2013)

删除了: The ion

删除了: may also significantly impact CH₄ oxidation.

设置了格式: 下标

移动了(插入) [1]

删除了: pollution

删除了: has

删除了: Sulfur is transferred into the earth's surface through dry and wet deposition, with SO₄²⁻ being the dominant form.

删除了: , and influences greenhouse gas emissions (Fan et al., 2017; Gauci et al., 2004; Schimel, 2004)

删除了: The cycling of

删除了: and CH₄ exhibits a significant inter-relationship.

删除了: CH₄ production (

删除了:)

kinetic preference as an electron acceptor (Granberg et al., 2001; Schimel, 2004), leading to decreased CH₄ emissions (Gauci et al., 2004). SO₄²⁻ has been shown to facilitate anaerobic CH₄ oxidation by anaerobic methanotrophic archaea in diverse ecosystems, such as oceans (Boetius et al., 2000), wetlands (La et al., 2022), and paddy fields (Fan et al., 2021), acting as a crucial electron acceptor. Despite these well-documented effects on anaerobic CH₄ cycling, the influence of SO₄²⁻ on aerobic CH₄ oxidation, particularly in upland soils, remains underexplored. Given the increasing global deposition of SO₄²⁻ due to industrial activities, understanding its impact on aerobic CH₄ oxidation is essential for predicting future CH₄ dynamics and developing effective climate mitigation strategies.

In this review, we have analyzed the literature on the effects of SO₄²⁻ on aerobic CH₄ oxidation. Our analysis not only reveals evidence suggesting that SO₄²⁻ promotes aerobic CH₄ oxidation but also identifies supporting evidence from related studies. In this review, we reviewed references about the influence of SO₄²⁻ on soil properties, substances, or biochemical processes, aiming to elucidate any microscale pathways on aerobic CH₄ oxidation through variations in soil substances or processes. Our analysis reveals that SO₄²⁻ may affect aerobic CH₄ oxidation. Based on the available literature, we infer that SO₄²⁻ favors aerobic CH₄ oxidation. This review summarizes the microscale pathways by which SO₄²⁻ influences aerobic CH₄ oxidation and highlights the importance of future research in this area. By providing a comprehensive synthesis of existing knowledge, this work serves as a valuable reference for future experimental studies. Furthermore, the findings of this review will contribute to a deeper understanding of global CH₄ cycling, particularly in the context of increasing SO₄²⁻ deposition. Moving forward, we aim to experimentally validate the impact of aerobic CH₄ oxidation following SO₄²⁻ addition and elucidate the underlying microbial mechanisms involved.

2 The microbial aerobic CH₄ oxidation processes

删除了：.

设置了格式：下标

删除了：However, there is still a lack of a comprehensive review that summarizes the impact of SO₄²⁻ on soil CH₄ oxidation, a process that significantly contributes to reducing atmospheric CH₄ concentrations.

删除了：Previous studies indicated that SO₄²⁻ addition may promote CH₄ oxidation by modulating the community structure or activity of methanotrophs in soils (Bradford et al., 2001b; Sitaula et al., 1995),

删除了：while some studies indicated no significant impact on CH₄ oxidation (Bradford et al., 2001a). Upon reviewing the literature, we found that the enhancement of SO₄²⁻ on CH₄ oxidation is prominent in numerous studies. Furthermore, SO₄²⁻ facilitates CH₄ oxidation within a range of 3-42% (Table 1). Thus, we hypothesize that SO₄²⁻ may stimulate CH₄ oxidation. Nonetheless, the scarcity of data precludes a definitive conclusion regarding the direct effect of SO₄²⁻ on CH₄ oxidation.

删除了：Further,

删除了：indirect effect

删除了：Indeed, we find that

删除了：both directly and indirectly

设置了格式：下标

设置了格式：上标

设置了格式：下标

删除了：This review provides a comprehensive summary of the direct and potential indirect impacts of SO₄²⁻ on CH₄ oxidation. The review underscores the viability of investigating the effect of SO₄²⁻ on CH₄ oxidation,

删除了：providing a valuable reference for future experimental research.

设置了格式：下标

设置了格式：下标

删除了：We will experimentally identify the enhancement of CH₄ oxidation following SO₄²⁻ addition, along with elucidating the underlying microbial processes. This will contribute to a deeper understanding of the global CH₄ cycling in the context of increasing acid deposition in the future.

设置了格式：字体：四号，加粗

2.1 Aerobic CH₄ oxidation processes

Aerobic CH₄ oxidation is mediated by methanotrophs, a group of specialized microorganisms (Chistoserdova et al., 2005). In soils, aerobic CH₄ oxidation can be classified into two distinct forms based on the concentration of CH₄ (Walsh et al., 2009). The first form, known as high-affinity oxidation, occurs at CH₄ concentrations close to atmospheric levels (<2ppm) and is carried out by high-affinity methanotrophs (Chowdhury and Dick, 2013). This process is commonly observed in upland soils, particularly in environments with high NH₄⁺ concentrations (Ho et al., 2019; Le Mer and Robért, 2001). The second form, referred to as low-affinity oxidation, occurs at CH₄ concentrations exceeding 40 ppm and is mediated by low-affinity methanotrophs (Chowdhury and Dick, 2013). This form is typically found in wetland environments, where CH₄ concentrations are significantly higher than atmospheric levels (Bechtold et al., 2025). Aerobic CH₄ oxidation processes can be further classified into assimilatory and dissimilatory pathways. In the dissimilatory pathways, CH₄ is sequentially oxidized to CO₂ by multiple enzymes (Fig. 1②). (Mancinelli, 1995). In the assimilation pathways, methanotrophs convert formaldehyde, an intermediate product of aerobic CH₄ oxidation, into biomass and other organic compounds mainly through the ribulose monophosphate pathway (RuMP pathway) (Fig. 1③), serine pathway (Fig. 1④), and xylulose monophosphate pathway (XyMP pathway) (Fig. 1①)(Yang et al., 2023).

2.2 Methanotrophs

Methanotrophs constitute a distinct subset of methylotrophs, primarily dependent on the one-carbon compound CH₄ as their sole source of carbon and energy (Hanson and Hanson, 1996). In the traditional classification system, Proteobacterial methanotrophs were categorized into type I (*Methylococcaceae* and *Crenotrichaceae*), type II (*Methylocystaceae* and *Beijerinckiaceae*), and type X (*Methylococcaceae*) (Li et al., 2020), based on their cell membrane arrangement, chemotaxonomic properties, physiological characteristics, and phylogenetic location. However, due to the discovery of non-canonical methanotrophs, the traditional classification system has become

删除了:

设置了格式: 字体: 非加粗

设置了格式: 字体: 非加粗

设置了格式: 字体: 非加粗

设置了格式: 字体: 非加粗

设置了格式: 字体: 非加粗, 下标

设置了格式: 字体: 非加粗

设置了格式: 字体: 非加粗

设置了格式: 字体: 非加粗

设置了格式: 字体: 非加粗

设置了格式: 字体: 非加粗

设置了格式: 字体: 非加粗

设置了格式: 字体: 非加粗, 下标

设置了格式: 字体: 非加粗, 上标

设置了格式: 字体: 非加粗

设置了格式: 字体: 非加粗

设置了格式: 字体: 非加粗

设置了格式: 字体: 非加粗

设置了格式: 下标

删除了:

设置了格式: 下标

删除了: Upland soils serve as sinks for atmospheric CH₄, contributing about 6% to the total atmospheric CH₄ consumption (Liu and Greaver, 2009). CH₄ oxidation, mediated by methanotrophs (Chistoserdova et al., 2005), occurs in uplands, including grasslands (Kou et al., 2017; Zhang et al., 2020), forests (Jang et al., 2011; Mohanty et al., 2007), and agricultural soils (Ho et al., 2019), with oxygen serving as final electron acceptor. Soil CH₄ oxidation contributes the largest biological sink for atmospheric CH₄ (Pratscher et al., 2018). Moreover, the highest methanotroph

删除了: CH₄ is ultimately oxidized to CO₂ through the

设置了格式: 下标

删除了: CH₄ undergoes sequential oxidation: first, it is

删除了: M

删除了: uniquely

删除了: Methanotrophs belong to various phyla, including

删除了: are

删除了: classified

设置了格式: 字体: 倾斜

删除了: ,

删除了: .

outdated. Consequently, methanotrophs are now classified into seven categories based on phylogenetic analysis: Type I-A (*Methylomonadaceae*), I-B (*Methylococcaceae*), I-C (*Methylothermaceae*), I-D (*Crenotrichaceae*), II-A (*Methylocystaceae*), II-B (*Beijerinckiaceae*), III (*Methylacidiphilaceae*), and NC10 (Fenibo et al., 2023). *Methylomonadaceae*, *Methylococcaceae*, *Methylothermaceae*, and *Crenotrichaceae* belong to the class *Gammaproteobacteria*, while *Methylocystaceae* and *Beijerinckiaceae* are classified under *Alphaproteobacteria*. *Methylacidiphilaceae* belongs to the phylum *Verrucomicrobia*. The composition of different types of methanotrophs is shown in Figure 1 (Fenibo et al., 2023). Notably, only four genera—*Methylocella*, *Methyacidimicrobium*, *Methylacidiphilum*, and *Methanomirabilis*—are capable of carbon fixation via the Calvin-Benson-Bassham (CBB) cycle (Fenibo et al., 2023; Op den Camp et al., 2009). Among Actinobacterial methanotrophs, *Candidatus* *Mycobacterium methanotrophicum* is classified with the *Mycobacterium* genus (van Spanning et al., 2022). Methanotrophs utilize two forms of methane monooxygenase (MMOs): soluble cytoplasmic monooxygenase (sMMO) and particulate membrane-bound monooxygenase (pMMO). The expression of these enzymes is regulated by copper (Cu) concentration (Hakemian and Rosenzweig, 2007).

3 Soil CH₄ oxidation in response to SO₄²⁻ addition

Sulfates, including SO₄²⁻ and sulfuric acid (H₂SO₄), enhance aerobic CH₄ oxidation within a range of 0–42% (Table 1), thus, we hypothesize that SO₄²⁻ may stimulate aerobic CH₄ oxidation. For example, in a temperate mixed deciduous woodland, the cumulative uptake of aerobic CH₄ oxidation was 25% higher in the experimental group with H₂SO₄ addition compared to the control group during the final quarter of the study period (Bradford et al., 2001b). Similar results were reported by Sitaula et al. (1995). In another study, King and Schell (1998) found that adding SO₄²⁻ (Na₂SO₄) increased aerobic CH₄ oxidation by 3% at a CH₄ concentration of 250 ppm compared to the control group, although this result was not statistically significant. The lack of significance may be attributed to the insufficient concentration gradient of Na₂SO₄ in

删除了:

设置了格式

删除了: Type I and type X methanotrophs belong to the γ -Proteobacteria class, while type II methanotrophs belong to the α -Proteobacteria class (Hanson and Hanson, 1996). Type I methanotrophs assimilate carbon using the RuMP pathway, type II methanotrophs using the serine pathway, and type X methanotrophs can use both pathways (Hanson and Hanson, 1996; Mancinelli, 1995). Verrucomicrobial methanotrophs, a newly discovered methanotrophs group (Op den Camp et al., 2009), are acidophilic and thermophilic extremophiles (Schmitz et al., 2021). They lack intracytoplasmic membranes, possess primarily saturated phospholipids, an

设置了格式

删除了: d use the serine and Calvin-Benson-Bassham for carbon fixation... (Fenibo et al., 2023; Koo and Rosenzweig, 2021; ...p den Camp et al., 2009). Among Actinobacterial methanotrophs The actinobacterial methanotrophs... *Candidatus* *Mycobacterium methanotrophicum*,... is classified with the *Mycobacterium* genus (van Spanning et al., 2022) has only recently been discovered and is preferred in low-pH and high-CH₄ environments (van Spanning et al., 2022). This group of methanotrophs expresses a

删除了:

Type I and type X methanotrophs belong to the

删除了: There are two forms of MMOs,...soluble cytoplasmic type...monooxygenase (sMMO) and particulate membrane-bound type...monooxygenase (pMMO). The

上移了 [1]: Acid rain pollution has been a significant (Chen et al., 2020; Qi et al., 2022). SO₄²⁻ is the major ion in acid rain (Wright and Henriksen, 1978) and has profound

删除了:

2.3 The influence of sulfate

Acid rain pollution has been a significant environmental

设置了格式

带格式的: 缩进: 首行缩进: 0 字符

删除了:

删除了: This increase in CH₄ oxidation could be attributed to alterations in the activity or community structure of methanotrophs caused by the addition of H₂SO₄ (Bradford et

设置了格式

the experimental setup, which limited the ability to fully assess the effects of SO_4^{2-} on aerobic CH_4 oxidation. Therefore, we propose that the observed enhancement of aerobic CH_4 oxidation following H_2SO_4 addition is primarily due to the increase in SO_4^{2-} concentration.

The promotional effect of SO_4^{2-} on aerobic CH_4 oxidation is further supported by comparisons with other anions under similar cationic conditions. Benstead and King (2001) observed that HNO_3 exerted a stronger inhibitory effect on aerobic CH_4 oxidation under equivalent soil acidic conditions than H_2SO_4 . This finding is consistent with the results of Bradford et al. (2001a), who experimentally confirmed the inhibitory effect of nitrate (NO_3^-) on aerobic CH_4 oxidation (Dunfield and Knowles, 1995; Wang and Ineson, 2003). When H_2SO_4 and HNO_3 were added to the soil to achieve H^+ concentrations of 10 and 1 $\mu\text{mol H}^+$ per gram of fresh weight (gfw), respectively, both acids inhibited aerobic CH_4 oxidation to a similar extent. However, H_2SO_4 exhibited a lesser inhibitory effect than HNO_3 . We hypothesize that SO_4^{2-} may promote aerobic CH_4 oxidation, as evidenced by the findings of Benstead and King (2001) and Bradford et al. (2001a). Consequently, when H_2SO_4 and HNO_3 are added to the soil, resulting in equivalent acidic conditions, the inhibitory effect of H_2SO_4 is less pronounced than that of HNO_3 .

However, not all studies support the hypothesis that SO_4^{2-} promotes aerobic CH_4 oxidation. For instance, Bradford et al. (2001a) observed no significant difference in aerobic CH_4 oxidation between low (564 μM) and high (1408 μM) concentrations of H_2SO_4 compared to the control group. This discrepancy may be due to differences in H_2SO_4 concentration across studies. Similarly, Hu et al. (2018) reported no significant effect of SO_4^{2-} on aerobic CH_4 oxidation. Based on the available evidence, SO_4^{2-} promotes aerobic CH_4 oxidation within a range of 0–42%. Although the mechanisms by which SO_4^{2-} influences aerobic CH_4 oxidation are not yet fully understood, we have identified potential microscopic pathways through which SO_4^{2-} may affect this aerobic process by reviewing relevant literature.

设置了格式: 下标

设置了格式: 下标

设置了格式: 下标

删除了: Consequently, more SO_4^{2-} is desorbed and transported in acidic soils, which could further augment CH_4 oxidation (Prietzel et al., 2004).

删除了:

删除了: can be further evidenced by comparing its effects with those of other anions under the same cationic conditions

删除了: When NH_4Cl and $(\text{NH}_4)_2\text{SO}_4$ were added to the soil at concentrations of 1 μmol and 10 $\mu\text{mol NH}_4^+$ per gram of fresh weight, respectively, both compounds caused the same level of inhibition on net CH_4 oxidation. However, the inhibition observed at the lower concentration (1 μmol) was significantly lower than that at the higher concentration (10 μmol) (Adamsen and King, 1993; Bradford et al., 2001a; King and Schnell, 1998). NH_4^+ has been found to inhibit CH_4 oxidation (Bronson and Mosier, 1994; Dunfield and Knowles, 1995). However, the inhibitory effect of NH_4Cl is greater than that of $(\text{NH}_4)_2\text{SO}_4$, as SO_4^{2-} may enhance the adsorption of NH_4^+ onto cation exchange sites in the soil (Bradford et al., 2001b; Gullledge and Schimel, 1998; King and Schnell, ...)

删除了: , under equivalent soil acidic conditions, HNO_3 exerted a greater inhibitory effect on CH_4 oxidation

设置了格式: 下标

删除了: Similar results have been reported by Bradford et al.

删除了: ,

删除了: albeit with

删除了: ing

删除了: Nonetheless, future experiments are required to verify the direct effect of SO_4^{2-} on CH_4 oxidation.

删除了: Another possibility is that the

批注 [芮苏 1]: Hu et al.
(<https://doi.org/10.1080/02757540.2018.1464153>)

删除了: ,

删除了: relative to the previous study, led to SO_4^{2-} - stimulating CH_4 oxidation. Alternatively, Bradford et al. ...

删除了: Due to the scarcity of studies investigating the direct effect of SO_4^{2-} on CH_4 oxidation, no definitive ...

4 Microscale pathways by which SO_4^{2-} addition influences aerobic CH_4 oxidation

At present, the impact of SO_4^{2-} on aerobic CH_4 oxidation is not fully understood, and its role in this process remains unclear, particularly regarding its potential enhancing effects. To address this knowledge gap, we conducted a literature review focusing on studies involving the addition of SO_4^{2-} . Through this review, we have identified several potential microscopic pathways through which SO_4^{2-} may promote aerobic CH_4 oxidation. One possible pathway involves changes in methanotroph activity and community structure in response to SO_4^{2-} (Fig. 2 path d) (Bradford et al., 2001b; Sitaula et al., 1995). Alternatively, SO_4^{2-} may affect aerobic CH_4 oxidation by altering soil physical properties (Fan et al., 2017), microbial substrates availability (Bjorneras et al., 2019; Palmer et al., 2013; Xu et al., 2017), and soil nutrition content (Islam, 2012) (Fig. 2).

First, the addition of SO_4^{2-} alters soil physical properties (Fig. 2 path a), i.e., particularly by reducing soil pH (Fig. 2 ①). Soil acidification increases due to enhanced base cation leaching associated with SO_4^{2-} addition (Hu et al., 2013), leading to a decrease in the pH of forest soils (Fasth et al., 1991; Tie et al., 2020). The addition of H_2SO_4 has been shown to promote aerobic CH_4 oxidation by altering the activity or community structure of methanotrophs (Bradford et al., 2001b; Sitaula et al., 1995). However, in experiments involving H_2SO_4 addition, it remains unclear whether the observed enhancement in aerobic CH_4 oxidation is primarily due to the decreased pH (Fig. 2 path e) or the increase in SO_4^{2-} concentration (Fig. 2 path d). Generally, CH_4 consumption is greater at higher pH conditions in forest soils (Brumme and Borken, 1999; Silver et al., 1999). Therefore, the reduction in soil pH caused by SO_4^{2-} addition may lead to a decrease in aerobic CH_4 oxidation. However, in acidic soils, a decrease in pH has been shown to

删除了： Indirect impact pathway of SO_4^{2-} on CH_4 oxidation

删除了： The influence of SO_4^{2-} on CH_4 oxidation has been relatively understudied in research. Thus, we conducted a literature review focused on studies involving the addition of

删除了： .

删除了： Our objective was to demonstrate that SO_4^{2-} can indirectly influence CH_4 oxidation through its effects on soil physicochemical properties and changes in other soil constituents.

删除了： Indeed, our review uncovered several significant findings that support this hypothesis. CH_4 oxidation is influenced by various environmental factors (Kou et al., 2017; Shukla et al., 2013), including SO_4^{2-} . However,

删除了： may affect CH_4 oxidation directly (Fig. 2 path c), stemming from variations in methanotrophs activity and

删除了： e

删除了： it

删除了： indirectly

删除了： influencing

删除了： , and soil bacterial metabolism (Hu et al., 2013)

删除了： ed

带格式的：制表位：不在 8.95 字符

删除了： specifically reducing

删除了： path

删除了： (Fasth et al., 1991; Tie et al., 2020) and elevation

删除了： An increase in

删除了： s

删除了： , resulting from the augmented base cation

删除了： led

设置了格式：下标

设置了格式：下标

删除了： general

删除了： due to

删除了： result

删除了： in

删除了： observed

increase aerobic CH₄ oxidation (Sitaula et al., 1995). Consequently, when evaluating the impact of SO₄²⁻ addition on aerobic CH₄ oxidation, it is essential to consider the initial soil pH (Fig. 2 path e), as methanotrophs exhibit different pH preferences in acidic and alkaline environments (Shukla et al., 2013).

Second, SO₄²⁻ addition can alter the soil microbial substrate (Fig. 2 path b), particularly by decreasing soil di-O-alkyl C content (Fig. 2 ②) (Xu et al., 2017). In a subtropical forest, SO₄²⁻ addition has been shown to increase the activity of gram-negative bacteria in soil by reducing the litter di-O-alkyl carbon (di-O-alkyl C) (Fig. 2 ② and path g) (Xu et al., 2017). Di-O-alkyl C is a component of soil organic carbon (SOC). SOC degradation is accelerated when the percentage of di-O-alkyl C is high (Huang et al., 2021). Conversely, when the content of di-O-alkyl C is low, SOC degradation slows down, leading to a greater availability of substrates for microorganisms, including methanotrophs. Methanotrophs, which are gram-negative bacteria (Schimel and Gullledge, 1998), may exhibit increased activity in response to SO₄²⁻ addition. This enhancement of methanotrophs activity (Fig. 2 path h), can ultimately promote aerobic CH₄ oxidation (Fig. 2 path g).

Third, SO₄²⁻ can alter soil nutrition content (Fig. 2 path e), specifically increasing soil Cu availability (Fig. 2 ③) (Islam, 2012), phosphorus (P) content (Fig. 2 ④) by enhancing acid phosphatase activity (Lv et al., 2014; Veraart et al., 2015), (aluminum ion) Al³⁺ toxicity (Fig. 2 ⑤) (Hu et al., 2013; Sogn and Abrahamsen, 1998), and NH₄⁺ absorption (Bradford et al., 2001b; Gullledge and Schimel, 1998; King and Schnell, 1998) (Fig. 2 ⑥). Cu is a crucial component in aerobic CH₄ oxidation processes, as it is utilized by methanotrophs in their molecular machinery, synthesized from metabolized CH₄ through the secretion of methanobactin into the environment. This process facilitates the oxidation of CH₄ to methanol (Dassama et al., 2016). It was anticipated that methanobactin secreted by methanotrophs during aerobic CH₄ oxidation would facilitate Cu uptake (Knapp et al., 2007); however, the specific

删除了: considering ...he impact of SO₄²⁻ addition on aerobic CH₄ oxidation, it is essential crucial ...o consider account for ...he initial soil pHacid-base status of the study soils...(Fig. 2 path i...), as methanotrophs exhibit different pH preferences in acidic and alkaline environments (Shukla et al., 2013). SO₄²⁻ addition has been found to increase soil Eh and soil OC in subtropical forest soil (Fan et al., 2017). Furthermore, the amount of O₂ in the soil was closely correlated with soil Eh, and a decrease in O₂ content can lead to a reduction in soil Eh (Zausig et al., 1993). Therefore, an increase in O₂ content results in an elevation of soil Eh (Fig. 2 path g). Methanotrophs are obligate aerobes (Teh et al., 2005), requiring MMO to bind O₂ for the initial step of CH₄ oxidation (Mancinelli, 1995; Shukla et al., 2013). Additionally, CH₄ oxidation exhibited a significant positive correlation with O₂ levels (Mancinelli, 1995). Consequently, the increase in soil Eh and OC, resulting from the elevated O₂ content due to SO₄²⁻ addition, may enhance CH₄ oxidation (Fig. 2 path h).

设置了格式: 字体: 非加粗

删除了: ould

删除了: specifically ...y decreasing soil DOC concentration (Fig. 2 path ③) (Bjorneras et al., 2019; Palmer et al., 2013) and ...i-O-alkyl C content (Fig. 2 path ...④... (Xu et al., 2017). Sullivan et al. (2013) reported that DOC stimulates CH₄ oxidation in semiarid soils; however, SO₄²⁻ addition reduces DOC concentrations (Bjorneras et al., 2019; Palmer et al., 2013). Consequently, it can be inferred that...

删除了: C

删除了: path ...④...and path j...) (Xu et al., 2017). Di-O-alkyl C is a component of soil organic carbon (SOC). SOC degradation is acceleratedoccurs...when the percentage ofmore readily with a higher percentage of...di-O-alkyl C...

删除了: d...), specifically increasing soil Cu availability (Fig. 2 path...③⑤... (Islam, 2012), phosphorus (P) content (Fig. 2 ④)path ⑥... by enhancing acid phosphatase activity (Lv et al., 2014; Veraart et al., 2015), and...(aluminum ion)...

设置了格式: 下标

删除了: when it is utilized ...y methanotrophs in their molecular machinery, synthesized from metabolized CH₄ through the secretion of methanobactin into the environment. This process facilitates the oxidationization

mechanism by which methanobactin affects Cu uptake remains unclear (Fig. 2 path j). For methanotrophs capable of expressing both sMMO and pMMO, the expression of these enzymes is regulated by the availability of Cu, a phenomenon known as the classic "copper switch" (Stanley et al., 1983). Under Cu-deficient conditions, these methanotrophs express sMMO. However, as the ratio of Cu to biomass increases, the expression of sMMO significantly decreases, while the expression of pMMO increases (Semrau et al., 2018). Notably, nearly all methanotrophs possess pMMO (Koo and Rosenzweig, 2021); therefore, increased Cu availability can enhance the expression of pMMO. Research indicates that Cu can serve as a promoter of aerobic CH₄ oxidation (Ho et al., 2013). Therefore, SO₄²⁻ addition may promote aerobic CH₄ oxidation by increasing the availability of soil Cu, thereby enhancing the expression of pMMO (Fig. 2 path j and k).

A positive correlation has been found between P and aerobic CH₄ oxidation in soils (Veraart et al., 2015; Zhang et al., 2020). P can potentially enhance the activity of soil methanotrophs (Fig. 2 path n) (Zhang et al., 2011), with an increase in soil P content achieved through the hydrolysis of organic compounds, including nucleic acids, phospholipids, and phosphate esters, by acid and alkaline phosphatases (Veraart et al., 2015). The addition of SO₄²⁻ accelerated acid phosphatase activity, thereby increasing soil P content (Lv et al., 2014). Therefore, we hypothesize that SO₄²⁻ may indirectly enhance aerobic CH₄ oxidation through the augmentation of soil P content, subsequently promoting the activity of methanotrophs in the soil (Fig. 2 path n and o). It is well-established that Al³⁺ inhibits aerobic CH₄ oxidation (Tamai et al., 2007; Tamai et al., 2003). Additionally, soil acidification resulting from SO₄²⁻ addition has been shown to intensify the toxicity of Al³⁺ in forest soils (Fig. 2 ⑤) (Hu et al., 2013; Sogn and Abrahamsen, 1998). The increase in Al³⁺ can inhibit the activity of methanotrophs (Nanba and King, 2000; Shukla et al., 2013) (Fig. 2 path l), thereby inhibiting aerobic CH₄ oxidation (Fig. 2 path m). Therefore, SO₄²⁻ addition may directly affect methanotrophs by enhancing the toxicity of Al³⁺ in the soil, thereby inhibiting aerobic CH₄ oxidation (Fig. 2 path o). When NH₄Cl and (NH₄)₂SO₄ were added to the soil at

删除了: of how methanobactin affects Cu uptake remains unclear

删除了: n

删除了: An increase in Cu availability results in the overexpression of protein-mediated steps in the conversion of CH₄ to CO₂, including the synthesis of cell walls, lipids, and membranes (DiSpirito et al., 2016).

设置了格式: 下标

删除了: Furthermore, Cu may act as a stimulant for CH₄ oxidation (Ho et al., 2013).

设置了格式: 下标

设置了格式: 上标

设置了格式: 下标

删除了: Therefore, the addition of SO₄²⁻ may indirectly enhance CH₄ oxidation by augmenting the availability of soil Cu

删除了: m

删除了: p

删除了: infer

删除了: p

删除了: q

删除了:

删除了: .

设置了格式: 上标

删除了: We hypothesize that the addition of SO₄²⁻ may indirectly inhibit CH₄ oxidation by enhancing the toxicity of soil Al³⁺ (Fig. 2 path o).

设置了格式: 下标

the same molar concentration of NH_4^+ , the inhibitory effect of $(\text{NH}_4)_2\text{SO}_4$ on aerobic CH_4 oxidation was weaker than that of NH_4Cl (Adamsen and King, 1993; Bradford et al., 2001a; King and Schnell, 1998). NH_4^+ has been found to inhibit aerobic CH_4 oxidation (Bronson and Mosier, 1994; Dunfield and Knowles, 1995), and the key mechanism is the competition between CH_4 and NH_4^+ for the same MMO enzyme (Gulledge et al., 2004). Due to the similar molecular structures of CH_4 and NH_4^+ , MMO can oxidize both CH_4 (to CH_3OH) and NH_4^+ (to NO_2^-). The inhibitory effect of NH_4Cl is greater than that of $(\text{NH}_4)_2\text{SO}_4$, as SO_4^{2-} may enhance the adsorption of NH_4^+ onto cation exchange sites in the soil (Bradford et al., 2001b; Gulledge and Schimel, 1998; King and Schnell, 1998) (Fig. 2 ⑥). This reduced availability of NH_4^+ limits its ability to compete with methanotrophs for MMO enzymes, thereby increasing the availability of MMO (Fig. 2 path p), promoting aerobic CH_4 oxidation (Fig. 2 path k), and further intensifying the inhibitory effect of NH_4Cl compared to $(\text{NH}_4)_2\text{SO}_4$. In conclusion, SO_4^{2-} served as a facilitator of aerobic CH_4 oxidation, mitigating the inhibitory effects of NH_4^+ on this process.

5 Conclusions

SO_4^{2-} plays a pivotal role in global acid deposition, with annual deposition rates ranging from $141.64 \pm 120.04 \text{ TgS a}^{-1} \text{ year}^{-1}$ (Gao et al., 2022). By synthesizing the available literature and exploring both its macroscopic effects and microscopic mechanisms, we investigated how SO_4^{2-} affects aerobic CH_4 oxidation. We observed that SO_4^{2-} enhances aerobic CH_4 oxidation by up to 0–42% on a macro scale. At the microscopic mechanism level, SO_4^{2-} can influence methanotrophs or MMO by modulating pH, di-O-alkyl C content, Cu availability, P content, Al^{3+} toxicity, and NH_4^+ absorption, thereby promoting or inhibiting aerobic CH_4 oxidation. Based on these findings, we hypothesize that SO_4^{2-} would promote aerobic CH_4 oxidation. If this hypothesis is validated in the future, it could provide significant benefits for CH_4 mitigation, particularly in the context of increasing global sulfur deposition. Therefore, future research in this field

设置了格式: 下标

设置了格式: 上标

批注 [芮苏2]: This content was initially placed in Section 3, aiming to illustrate that, under the same NH_4^+ concentration conditions, Na_2SO_4 exhibits a weaker inhibitory effect on aerobic CH_4 oxidation compared to NH_4Cl . However, due to concerns raised by Referee#2, we reconsidered and moved this section to Section 4. The revised argument now emphasizes that SO_4^{2-} promotes the adsorption of NH_4^+ in the soil, thereby alleviating the inhibitory effect on aerobic CH_4 oxidation.

设置了格式: 下标

设置了格式: 下标

设置了格式: 上标

设置了格式: 下标

设置了格式: 下标

设置了格式: 下标

设置了格式: 上标

设置了格式: 下标

设置了格式: 上标

删除了:

删除了: Finally, the addition of SO_4^{2-} alters soil microbial metabolism (Fig. 2 path f), specifically including an up-regulation of the tricarboxylic acid (TCA) cycle (Fig. 2 path ⑧) (Wang et al., 2021). Methanotrophs utilize the TCA cycle during the metabolism of the serine cycle to assimilate CH_4 ...

删除了: Implication

删除了: The microbiological processes of methanogenesis...

删除了:

删除了: (

删除了:)

删除了: Various studies have demonstrated that SO_4^{2-} ...

设置了格式: 上标

删除了: By synthesizing the available literature,

删除了: w

删除了: , suggesting that increased SO_4^{2-} deposition ma...

删除了: Mechanistically

删除了:

设置了格式: 下标

删除了: SO_4^{2-} can either directly enhance CH_4 oxidation or...

1225 should focus on investigating the response of aerobic CH₄ oxidation and its influencing factors under increasing SO₄²⁻ conditions, as well as clarifying the underlying microbial mechanisms.

删除了：which is beneficial for CH₄ mitigation in the context of heightened acid deposition. Moreover, the relationship between SO₄²⁻ and CH₄ oxidation may be mediated by intermediate substances, specifically, SO₄²⁻ affects CH₄ oxidation by modulating methanotrophs or including alterations in other soil components, requiring further experimental validation.

Data availability

All raw data can be provided by the corresponding authors upon request.

1230 Author contribution

Rui Su finished writing; Kexin Li, Nannan Wang, Fenghui Yuan, Yunjiang Zuo, Ying Sun, and Liyuan He gave constructive comments and revised the structure and content of the article; Xiaofeng Xu and Lihua Zhang reviewed and edited the manuscript.

Competing interests

1235 The authors declare that they have no conflict of interest.

Acknowledgments

1240 This study was partially supported by the National Natural Science Foundation of China (32471777; 32271681) and the Joint Funds of China's National Natural Science Foundation (U2006215), and by Key Laboratory of Ecology and Environment in Minority Areas (Minzu University of China), National Ethnic Affairs Commission (KLEEMA202206). X.X. acknowledged the financial assistance provided by the National Science Foundation (2145130) and SPRUCE and NGEE Arctic projects, supported by the Office of Biological and Environmental Research in the Department of Energy Office of Science.

1245





Figure 1: Diagram of the complete aerobic methane oxidation process, including methane assimilation and dissimilation, and classification of methanotrophs.

①Xylulose monophosphate pathway (XyMP pathway) of formaldehyde assimilation;
 ②CH₄ is oxidized to CO₂ under the sequential action of multiple enzymes; ③Ribulose monophosphate pathway (RuMP pathway) of formaldehyde assimilation; ④ Serine pathway of formaldehyde assimilation.



删除了：

删除了：MMO-containing status of identified methanotrophs.

设置了格式：字体：加粗

带格式的：到齐到网格，制表位：不在 8.95 字符

删除了：① CH₄ is oxidized to methanol (CH₃OH) by MMO; ② CH₃OH is oxidized to formaldehyde (HCOH) by methanol dehydrogenase (MDH); ③ HCOH is oxidized to formate (HCOOH) by formaldehyde dehydrogenase (FADH); ④ HCOOH is oxidized to CO₂ by formate dehydrogenase (FDH); ⑤ HCOH is assimilated into cellular biomass via the RuMP cycle; ⑥ HCOH is assimilated into cellular biomass via the Serine cycle; ⑦ Cu controls two MMOs expression (Hakemian and Rosenzweig, 2007); ⑧ High Cu concentration regulates pMMO expression in soil (Hakemian and Rosenzweig, 2007); ⑨ Low Cu concentration regulates sMMO expression in soil (Hakemian and Rosenzweig, 2007). pMMO and sMMO have different structures and mechanisms, and sMMO has broader substrate specificity than pMMO. Most methanotrophs possess pMMO, so the *pmoA* gene is a key gene for detecting methanotrophs in environment samples.





Figure 2. Conceptual diagram of the potential microscopic mechanisms by which sulfate influences aerobic methane oxidation in upland soil.

① SO_4^{2-} decreases soil pH (Fasth et al., 1991; Tie et al., 2020); ② SO_4^{2-} decreases soil di-O-alkyl C amount (Xu et al., 2017); ③ SO_4^{2-} increases soil Cu availability (Islam, 2012); ④ SO_4^{2-} increases soil P content by increasing soil acid phosphatase activity (Lv et al., 2014; Veraart et al., 2015); ⑤ SO_4^{2-} increases soil Al^{3+} toxicity (Hu et al., 2013; Sogn and Abrahamsen, 1998); ⑥ SO_4^{2-} increases NH_4^+ absorption (Bradford et al., 2001b; Gullledge and Schimel, 1998; King and Schnell, 1998); a. Changes in soil physical properties due to increased soil SO_4^{2-} content; b. Changes in soil microbial substrate due to increased soil SO_4^{2-} content; c. SO_4^{2-} may promote CH_4 oxidation; d. SO_4^{2-} affects the activity or community size of methanotrophs in soils (Bradford et al., 2001b; Sitaula et al., 1995); e. Decreased pH may inhibit or stimulate soil CH_4 oxidation (Sitaula et al., 1995); f. Decreased pH may inhibit or stimulate soil CH_4 oxidation (Sitaula et al., 1995); g. Decreased di-O-alkyl C amount increases soil gram-negative bacteria activity (Xu et al., 2017); h. The increased activity of gram-negative bacteria may stem from the enhanced activity of methanotrophs; i. Elevated Cu availability stimulates soil aerobic CH_4 oxidation (Ho et al., 2013); j. mb (methanobactin) is expected to accelerate Cu uptake (Knapp et al., 2007); k. Enhanced MMO activity facilitates aerobic CH_4 oxidation; l. Elevated Al^{3+} toxicity inhibits soil methanotrophs activity (Nanba and King, 2000; Shukla et al., 2013); m. Decreased methanotrophs activity inhibits soil CH_4 oxidation; n. Elevated P content increases soil methanotrophs



删除了： Conceptual diagram illustrating SO_4^{2-} affected CH_4 oxidation through direct or indirect ways in soil.

删除了： SO_4^{2-} increases soil Eh (redox potential) and soil OC (oxidation capacity) (Fan et al., 2017); ③ SO_4^{2-} decreases soil DOC concentration (Bjorneras et al., 2019; Palmer et al., 2013); ④

删除了： ⑤

删除了： ⑥

删除了： ⑦

删除了： ⑧

删除了： induced soil TCA cycle up-regulate (Wang et al., 2021)

删除了： Changes in soil nutrition content due to increased soil SO_4^{2-} content; e.

删除了： f. Change in soil bacterial metabolism due to increased soil SO_4^{2-} content; g. Elevated O_2 content increases soil Eh (Zausig et al., 1993); h. Elevated O_2 level stimulates soil CH_4 oxidation (Mancinelli, 1995);

删除了： i

删除了： j

删除了： k

删除了： Increased activity of gram-negative bacteria may increase activity of methanotrophs

删除了： l. Decreased DOC concentration may inhibit soil CH_4 oxidation (Sullivan et al., 2013);

删除了： m

删除了： n

删除了： o

删除了： CH_4 oxidation

删除了： (Tamai et al., 2007; Tamai et al., 2003)

删除了： p

activity (Zhang et al., 2011); p. Elevated methanotrophs activity stimulates soil CH₄ oxidation (Bradford et al., 2001b; Sitaula et al., 1995); p. The increased adsorption of NH₄⁺ enhances the availability of MMO to soil methanotrophs.

删除了: q

删除了: r

删除了: Increased CH₄ oxidation may increase CH₄ assimilation (Roslev et al., 1997); s. Methanotrophs undergo the TCA cycle during CH₄ metabolism and may cause up-regulation of the TCA cycle in soil.

设置了格式: 下标

设置了格式: 上标

Study site	Sulfate concentration	CH ₄ concentration	Effect	Reference
Perridge Forest	H ₂ SO ₄ (50 Kg S ha ⁻¹)	Ambient air	25 % increased	Bradford et al., 2001b
Perridge Forest	H ₂ SO ₄ (5mM)	Ambient air	no effect	Bradford et al., 2001a
	(NH ₄) ₂ SO ₄ (5mM)	Ambient air	no effect	
Maine forest	Na ₂ SO ₄ 0.5µg S g ⁻¹ soil	250ppm	3% increased	King and Schell, 1998
Norway Scots Pine forest	H ₂ SO ₄ pH3	Ambient air	42% increased	Sitaula et al., 1995
Birch taiga	Na ₂ SO ₄ 2.8 µmol S g ⁻¹ soil	4ppm	no effect	Gulledge and Schimel, 1998
	K ₂ SO ₄ 2.8 µmol S g ⁻¹ soil	4ppm	no effect	

Table 1. Promotion effect of sulfates on methane oxidation in diverse upland soils.

Reference

Adamsen, A.P.S., King, G.M., Methane consumption in temperate and subarctic forest soils rates, vertical zonation, and responses to water and nitrogen. *Applied and Environmental Microbiology* 59, 485-490. <http://doi.org/10.1128/aem.59.2.485-490.1993>, 1993.

Bechtold, E.K., Ellenbogen, J.B., Villa, J.A., Ferreira, D.K.d.M., Oliverio, A.M., Kostka, J.E., Rich, V.I., Varner, R.K., Bansal, S., Ward, E.J., Bohrer, G., Borton, M.A., Wrighton, K.C., Wilkins, M.J., Metabolic interactions underpinning high methane fluxes across terrestrial freshwater wetlands. *Nature Communications* 16. <http://doi.org/10.1038/s41467-025-56133-0>, 2025.

Bjorneras, C., Skerlep, M., Floudas, D., Persson, P., Kritzberg, E.S., High sulfate concentration enhances iron mobilization from organic soil to water. *Biogeochemistry* 144, 245-259. <http://doi.org/10.1007/s10533-019-00581-6>, 2019.

Bodelier, P.L.E., Interactions between nitrogenous fertilizers and methane cycling in wetland and upland soils. *Current Opinion in Environmental Sustainability* 3, 379-388. <http://doi.org/10.1016/j.cosust.2011.06.002>, 2011.

- Boetius, A., Ravenschlag, K., Schubert, C., Rickert, D., Widdel, F., Gieseke, A., Amann, R., Jorgensen, B., Witte, U., Pfannkuche, O., A marine microbial consortium apparently mediating anaerobic oxidation of methane. *Nature* 407, 623-626. <http://doi.org/10.1038/35036572>, 2000.
- Bradford, M.A., Ineson, P., Wookey, P.A., Lappin-Scott, H.M., The effects of acid nitrogen and acid sulphur deposition on CH₄ oxidation in a forest soil: a laboratory study. *Soil Biology and Biochemistry* 33, 1695-1702. [http://doi.org/10.1016/S0038-0717\(01\)00091-8](http://doi.org/10.1016/S0038-0717(01)00091-8), 2001a.
- Bradford, M.A., Wookey, P.A., Ineson, P., Lappin-Scott, H.M., Controlling factors and effects of chronic nitrogen and sulphur deposition on methane oxidation in a temperate forest soil. *Soil Biology and Biochemistry* 33, 93-102. [http://doi.org/10.1016/S0038-0717\(00\)00118-8](http://doi.org/10.1016/S0038-0717(00)00118-8), 2001b.
- Bronson, K.F., Mosier, A.R., Suppression of methane oxidation in aerobic soil by nitrogen fertilizers, nitrification inhibitors, and urease inhibitors. *Biology and Fertility of Soils* 17, 263-268. <http://doi.org/10.1007/bf00383979>, 1994.
- Brumme, R., Borken, W., Site variation in methane oxidation as affected by atmospheric deposition and type of temperate forest ecosystem. *Global Biogeochemical Cycles* 13, 493-501. <http://doi.org/10.1029/1998gb900017>, 1999.
- Chistoserdova, L., Vorholt, J.A., Lidstrom, M.E., A genomic view of methane oxidation by aerobic bacteria and anaerobic archaea. *Genome Biology* 6, 1-6. <http://doi.org/10.1186/gb-2005-6-2-208>, 2005.
- Chowdhury, T.R., Dick, R.P., Ecology of aerobic methanotrophs in controlling methane fluxes from wetlands. *Applied Soil Ecology* 65, 8-22. <http://doi.org/10.1016/j.apsoil.2012.12.014>, 2013.
- Dassama, L.M., Kenney, G.E., Ro, S.Y., Zielazinski, E.L., Rosenzweig, A.C., Methanobactin transport machinery. *Proceedings of the National Academy of Sciences of the United States of America* 113, 13027-13032. <http://doi.org/10.1073/pnas.1603578113>, 2016.
- Deng, Y., Che, R., Wang, F., Conrad, R., Dumont, M., Yun, J., Wu, Y., Hu, A., Fang, J., Xu, Z., Cui, X., Wang, Y., Upland Soil Cluster Gamma dominates methanotrophic communities in upland grassland soils. *Science of the Total Environment* 670, 826-836. <http://doi.org/10.1016/j.scitotenv.2019.03.299>, 2019.
- Dunfield, P., Knowles, R., Kinetics of inhibition of methane oxidation by nitrate, nitrite, and ammonium in a humisol. *Applied and Environmental Microbiology* 61, 3129-3135. <http://doi.org/10.1128/aem.61.8.3129-3135.1995>, 1995.
- Fan, J.L., Xu, Y.H., Chen, Z.M., Xiao, J., Liu, D.Y., Luo, J.F., Bolan, N., Ding, W.X., Sulfur deposition suppressed nitrogen-induced soil N₂O emission from a subtropical forestland in southeastern China. *Agricultural and Forest Meteorology* 233, 163-170. <http://doi.org/10.1016/j.agrformet.2016.11.017>, 2017.
- Fan, L.C., Schneider, D., Dippold, M.A., Poehlein, A., Wu, W.C., Gui, H., Ge, T.D., Wu, J.S., Thiel, V., Kuzyakov, Y., Dorodnikov, M., Active metabolic pathways of anaerobic methane oxidation in paddy soils. *Soil Biology and Biochemistry* 156, 108215. <http://doi.org/10.1016/j.soilbio.2021.108215>, 2021.
- Fasth, W.J., David, M.B., Vance, G.F., Sulfate retention and cation leaching of forest

删除了：Bowman, J., The Methanotrophs—The families Methylococcaceae and Methylocystaceae, in: Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K., Stackebrandt, E. (Eds.), The Prokaryotes. Springer, New York, NY, pp. 266-289. http://doi.org/10.1007/0-387-30745-1_15, 2006.

删除了：Cai, Y.F., Yun, J.L., Jia, Z.J., Phylogeny and metabolic potential of the methanotrophic lineage MO3 in Beijerinckiaaceae from the paddy soil through metagenome-assembled genome reconstruction. *Microorganisms* 10, 955. <http://doi.org/10.3390/microorganisms10050955>, 2022.

设置了格式：字体颜色：自动设置，非加宽量 / 紧缩量
带格式的：缩进：左侧：0 厘米，悬挂缩进：2 字符
设置了格式：字体颜色：自动设置，非加宽量 / 紧缩量
设置了格式：字体颜色：自动设置，非加宽量 / 紧缩量

删除了：DiSpirito, A.A., Semrau, J.D., Murrell, J.C., Gallagher, W.H., Dennison, C., Vuilleumier, S., Methanobactin and the link between copper and bacterial methane oxidation. *Microbiology and Molecular Biology Reviews* 80, 387-409. <http://doi.org/10.1128/mmbr.00058-15>, 2016.

删除了：Eriksson, T., Oquist, M.G., Nilsson, M.B., Production and oxidation of methane in a boreal mire after a decade of increased temperature and nitrogen and sulfur deposition. *Global Change Biology* 16, 2130-2144. <http://doi.org/10.1111/j.1365-2486.2009.02097.x>, 2010.

soils in response to acid additions. Canadian Journal of Forest Research 21, 32-41.
<http://doi.org/10.1139/x91-005>, 1991.

Fenibo, E.O., Selvarajan, R., Wang, H.Q., Wang, Y., Abia, A.L.K., Untapped talents: insight into the ecological significance of methanotrophs and its prospects. Science of the Total Environment 903, 166145.
<http://doi.org/10.1016/j.scitotenv.2023.166145>, 2023.

Gao, J., Zhou, W., Liu, Y., Sha, L., Song, Q., Lin, Y., Yu, G., Zhang, J., Zheng, X., Fang, Y., Grace, J., Zhao, J., Xu, J., Gui, H., Sinclair, F., Zhang, Y., Litter-derived nitrogen reduces methane uptake in tropical rainforest soils. Science of the Total Environment 849, 157891. <http://doi.org/10.1016/j.scitotenv.2022.157891>, 2022.

Gauci, V., Matthews, E., Dise, N., Walter, B., Koch, D., Granberg, G., Vile, M., Sulfur pollution suppression of the wetland methane source in the 20th and 21st centuries. Proceedings of the National Academy of Sciences of the United States of America 101, 12583-12587. <http://doi.org/10.1073/pnas.0404412101>, 2004.

Granberg, G., Sund, I., Svensson, B.H., Nilsson, M., Effect of temperature, and nitrogen and sulfur deposition, on methane emission from a boreal mire. Ecology 82, 1982-1998. 2001.

Gulledge, J., Hrywna, Y., Cavanaugh, C., Steudler, P.A., Effects of long-term nitrogen fertilization on the uptake kinetics of atmospheric methane in temperate forest soils. FEMS Microbiology Ecology 49, 389-400. <http://doi.org/10.1016/j.femsec.2004.04.013>, 2004.

Gulledge, J., Schimel, J., Low-concentration kinetics of atmospheric CH₄ oxidation in soil and mechanism of NH₄⁺ inhibition. Applied and Environmental Microbiology 64, 4291-4298. 1998.

Guo, J.H., Feng, H.L., Peng, C.H., Chen, H., Xu, X., Ma, X.H., Li, L., Kneeshaw, D., Ruan, H.H., Yang, H.Q., Wang, W.F., Global climate change increases terrestrial soil CH₄ emissions. Global Biogeochemical Cycles 37, e2021GB007255. <http://doi.org/10.1029/2021gb007255>, 2023.

Hakemian, A.S., Rosenzweig, A.C., The biochemistry of methane oxidation. Annual Review of Biochemistry 76, 223-241. <http://doi.org/10.1146/annurev.biochem.76.061505.175355>, 2007.

Hanson, R.S., Hanson, T.E., Methanotrophic bacteria. Microbiological Reviews 60, 439-471. <http://doi.org/10.1128/mmbr.60.2.439-471.1996>, 1996.

Ho, A., Lee, H.J., Reumer, M., Meima-Franke, M., Raaijmakers, C., Zweers, H., de Boer, W., Van der Putten, W.H., Bodelier, P.L.E., Unexpected role of canonical aerobic methanotrophs in upland agricultural soils. Soil Biology and Biochemistry 131, 1-8. <http://doi.org/10.1016/j.soilbio.2018.12.020>, 2019.

Ho, A., Lüke, C., Reim, A., Frenzel, P., Selective stimulation in a natural community of methane oxidizing bacteria: effects of copper on *pmoA* transcription and activity. Soil Biology and Biochemistry 65, 211-216. <http://doi.org/10.1016/j.soilbio.2013.05.027>, 2013.

Hu, M.J., Wilson, B.J., Sun, Z.G., Huang, J.F., Tong, C., Effects of nitrogen and sulphate addition on methane oxidation in the marsh soil of a typical subtropical estuary (Min River) in China. Chemistry and Ecology 34, 610-623.

- <http://doi.org/10.1080/02757540.2018.1464153>, 2018.
- Hu, Y.L., Jung, K., Zeng, D.H., Chang, S.X., Nitrogen- and sulfur-deposition-altered soil microbial community functions and enzyme activities in a boreal mixedwood forest in western Canada. *Canadian Journal of Forest Research* 43, 777-784. <http://doi.org/10.1139/cjfr-2013-0049>, 2013.
- Huang, J., Zhou, K.J., Zhang, W., Liu, J.X., Ding, X., Cai, X.A., Mo, J.M., Sulfur deposition still contributes to forest soil acidification in the Pearl River Delta, South China, despite the control of sulfur dioxide emission since 2001. *Environmental Science and Pollution Research* 26, 12928-12939. <http://doi.org/10.1007/s11356-019-04831-w>, 2019.
- Huang, R.L., Crowther, T.W., Sui, Y.Y., Sun, B., Liang, Y.T., High stability and metabolic capacity of bacterial community promote the rapid reduction of easily decomposing carbon in soil. *Communications Biology* 4, 1376. <http://doi.org/10.1038/s42003-021-02907-3>, 2021.
- IPCC, 2013. *Climate change 2013: the physical science basis*. In: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Islam, M., The effect of different rates and forms of sulfur on seed yield and micronutrient uptake by chickpea. *Plant, Soil and Environment* 58, 399-404. <http://doi.org/10.17221/145/2012-pse>, 2012.
- King, G.M., Schnell, S., Effects of ammonium and non-ammonium salt additions on methane oxidation by *Methylosinus trichosporium* OB3b and Maine forest soils. *Applied and Environmental Microbiology* 64, 253-257. 1998.
- Klessa, D.A., Frame, J., Golightly, R.D., Harkess, R.D., THE EFFECT OF FERTILIZER SULFUR ON GRASS PRODUCTION FOR SILAGE. *Grass and Forage Science* 44, 277-281. <http://doi.org/10.1111/j.1365-2494.1989.tb02165.x>, 1989.
- Knapp, C.W., Fowle, D.A., Kulczycki, E., Roberts, J.A., Graham, D.W., Methane monooxygenase gene expression mediated by methanobactin in the presence of mineral copper sources. *Proceedings of the National Academy of Sciences of the United States of America* 104, 12040-12045. <http://doi.org/10.1073/pnas.0702879104>, 2007.
- Koo, C.W., Rosenzweig, A.C., Biochemistry of aerobic biological methane oxidation. *Chemical Society Review* 50, 3424-3436. <http://doi.org/10.1039/d0cs01291b>, 2021.
- La, W., Han, X., Liu, C.Q., Ding, H., Liu, M., Sun, F., Li, S., Lang, Y., Sulfate concentrations affect sulfate reduction pathways and methane consumption in coastal wetlands. *Water Research* 217, 118441. <http://doi.org/10.1016/j.watres.2022.118441>, 2022.
- Le Mer, J., Roger, P., Production, oxidation, emission and consumption of methane by soil: a review. *European Journal of Soil Biology* 37, 25-50. [http://doi.org/10.1016/S1164-5563\(01\)01067-6](http://doi.org/10.1016/S1164-5563(01)01067-6), 2001.

删除了: Jang, I., Lee, S., Zoh, K.-D., Kang, H., Methane concentrations and methanotrophic community structure influence the response of soil methane oxidation to nitrogen content in a temperate forest. *Soil Biology and Biochemistry* 43, 620-627. <http://doi.org/10.1016/j.soilbio.2010.11.032>, 2011.

Jeremiason, J.D., Engstrom, D.R., Swain, E.B., Nater, E.A., Johnson, B.M., Almendinger, J.E., Monson, B.A., Kolka, R.K., Sulfate addition increases methylmercury production in an experimental wetland. *Environmental Science & Technology* 40, 3800-3806. <http://doi.org/10.1021/es0524144>, 2006.

删除了: Kou, Y.P., Li, J.B., Wang, Y.S., Li, C.N., Tu, B., Yao, M.J., Li, X.Z., Scale-dependent key drivers controlling methane oxidation potential in Chinese grassland soils. *Soil Biology and Biochemistry* 111, 104-114. <http://doi.org/10.1016/j.soilbio.2017.04.005>, 2017.

Li, C.H., Wang, B., Fang, Z., Yu, H.L., Huang, J.Y., Plant species diversity is driven by soil base cations under acid deposition in desert coal-mining region in northwestern China. *Ecological Indicators* 145, 109682. <http://doi.org/10.1016/j.ecolind.2022.109682>, 2022.

Li, Y., Liu, Y.W., Pan, H., Hernández, M., Guan, X.M., Wang, W., Zhang, Q.C., Luo, Y., Di, H.J., Xu, J.M., Impact of grazing on shaping abundance and composition of active methanotrophs and methane oxidation activity in a grassland soil. *Biology and Fertility of Soils* 56, 799-810. <http://doi.org/10.1007/s00374-020-01461-0>, 2020.

Lv, Y., Wang, C.Y., Jia, Y.Y., Wang, W.W., Ma, X., Du, J.J., Pu, G.Z., Tian, X.J., Effects of sulfuric, nitric, and mixed acid rain on litter decomposition, soil microbial biomass, and enzyme activities in subtropical forests of China. *Applied Soil Ecology* 79, 1-9. <http://doi.org/10.1016/j.apsoil.2013.12.002>, 2014.

Mancinelli, R., The regulation of methane oxidation in soil. *Annual Review of Microbiology* 49, 581-605. 1995.

Mishra, V.K., Shukla, R., Shukla, P.N., Inhibition of Soil Methane Oxidation by Fertilizer Application: an Intriguing but Persistent Paradigm. *Environmental Pollution and Protection* 3, 57-69. <http://doi.org/10.22606/epp.2018.32001>, 2018.

Murguia-Flores, F., Arndt, S., Ganesan, A.L., Murray-Tortarolo, G., Hornibrook, E.R.C., Soil Methanotrophy Model (MeMo v1.0): a process-based model to quantify global uptake of atmospheric methane by soil. *Geoscientific Model Development* 11, 2009-2032. <http://doi.org/10.5194/gmd-11-2009-2018>, 2018.

Nanba, K., King, G., Response of atmospheric methane consumption by maine forest soils to exogenous aluminum salts. *Applied and Environmental Microbiology* 66, 3674-3679. <http://doi.org/10.1128/AEM.66.9.3674-3679.2000>, 2000.

Op den Camp, H.J., Islam, T., Stott, M.B., Harhangi, H.R., Hynes, A., Schouten, S., Jetten, M.S., Birkeland, N.K., Pol, A., Dunfield, P.F., Environmental, genomic and taxonomic perspectives on methanotrophic *Verrucomicrobia*. *Environmental Microbiology Reports* 1, 293-306. <http://doi.org/10.1111/j.1758-2229.2009.00022.x>, 2009.

Palmer, S.M., Clark, J.M., Chapman, P.J., van der Heijden, G.M.F., Bottrell, S.H., Effects of acid sulphate on DOC release in mineral soils: the influence of SO₄²⁻ retention and Al release. *European Journal of Soil Science* 64, 537-544. <http://doi.org/10.1111/ejss.12048>, 2013.

Place, S.E., Examining the role of ruminants in sustainable food systems. *Grass and Forage Science* 79, 135-143. <http://doi.org/10.1111/gfs.12673>, 2024.

Praeg, N., Wagner, A.O., Illmer, P., Plant species, temperature, and bedrock affect net methane flux out of grassland and forest soils. *Plant and Soil* 410, 193-206. <http://doi.org/10.1007/s11104-016-2993-z>, 2016.

Saunois, M., Stavert, A.R., Poulter, B., Bousquet, P., Canadell, J.G., Jackson, R.B., Raymond, P.A., Dlugokencky, E.J., Houweling, S., Patra, P.K., Ciais, P., Arora, V.K., Bastviken, D., Bergamaschi, P., Blake, D.R., Brailsford, G., Bruhwiler, L., Carlson, K.M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P.M., Covey, K., Curry, C.L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M.I.,

删除了：Liu, L., Greaver, T.L., A review of nitrogen enrichment effects on three biogenic GHGs: the CO₂ sink may be largely offset by stimulated N₂O and CH₄ emission. *Ecology Letters* 12, 1103-1117. <http://doi.org/10.1111/j.1461-0248.2009.01351.x>, 2009.

删除了：Mohanty, S.R., Bodelier, P.L., Conrad, R., Effect of temperature on composition of the methanotrophic community in rice field and forest soil. *FEMS Microbiology Ecology* 62, 24-31. <http://doi.org/10.1111/j.1574-6941.2007.00370.x>, 2007.

Munthe, J., Bodaly, R.A., Branfireun, B.A., Driscoll, C.T., Gilmour, C.C., Harris, R., Horvat, M., Lucotte, M., Malm, O., Recovery of mercury-contaminated fisheries. *Ambio* 36, 33-44. [http://doi.org/10.1579/0044-7447\(2007\)36\[33:Romf\]2.0.Co;2](http://doi.org/10.1579/0044-7447(2007)36[33:Romf]2.0.Co;2), 2007.

删除了：Murguia-Flores, F., Ganesan, A.L., Arndt, S., Hornibrook, E.R.C., Global uptake of atmospheric methane by soil from 1900 to 2100. *Global Biogeochemical Cycles* 35, e2020GB006774. <http://doi.org/10.1029/2020gb006774>, 2021.

删除了：Pratscher, J., Vollmers, J., Wiegand, S., Dumont, M.G., Kaster, A.-K., Unravelling the identity, metabolic potential and global biogeography of the atmospheric methane-oxidizing upland soil cluster *α*. *Environmental Microbiology* 20, 1016-1029. <http://doi.org/10.1111/1462-2920.14036>, 2018.

Prietz, J., Mayer, B., Legge, A.H., Cumulative impact of 40 years of industrial sulfur emissions on a forest soil in west-central Alberta (Canada). *Environmental Pollution* 132, 129-144. <http://doi.org/10.1016/j.envpol.2004.03.016>, 2004.

Roslev, P., Iversen, N., Henriksen, K., Oxidation and assimilation of atmospheric methane by soil methane oxidizers. *Applied and Environmental Microbiology* 63, 874-880. <http://doi.org/10.1128/aem.63.3.874-880.1997>, 1997.

1620 [Hoglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K.M., Joos, F., Kleinen, T., Krummel, P.B., Langenfelds, R.L., Laruelle, G.G., Liu, L., Machida, T., Maksyutov, S., McDonald, K.C., McNorton, J., Miller, P.A., Melton, J.R., Morino, I., Muller, J., Murguia-Flores, F., Naik, V., Niwa, Y., Noce, S., Doherty, S.O., Parker, R.J., Peng, C., Peng, S., Peters, G.P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W.J., Rosentreter, J.A., Segers, A., Simpson, I.J., Shi, H., Smith, S.J., Steele, L.P., Thornton, B.F., Tian, H., Tohjima, Y., Tubiello, F.N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T.S., van Weele, M., van der Werf, G.R., Weiss, R.F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., Zhuang, Q., The Global Methane Budget 2000-2017. *Earth System Science Data* 12, 1561-1623. <http://doi.org/10.5194/essd-12-1561-2020>, 2020.](#)

1625 Schimel, J., Playing scales in the methane cycle: from microbial ecology to the globe. *Proceedings of the National Academy of Sciences of the United States of America* 101, 12400-12401. <http://doi.org/10.1073/pnas.0405075101>, 2004.

1630 Schimel, J., Guldge, J., Microbial community structure and global trace gases. *Global Change Biology* 4, 745-758. <http://doi.org/10.1046/j.1365-2486.1998.00195.x>, 1998.

1635 [Semrau, J.D., DiSpirito, A.A., Gu, W., Yoon, S., Metals and Methanotrophy. *Applied and Environmental Microbiology* 84. <http://doi.org/10.1128/aem.02289-17>, 2018.](#)

Shukla, P.N., Pandey, K.D., Mishra, V.K., Environmental determinants of soil methane oxidation and methanotrophs. *Critical Reviews in Environmental Science and Technology* 43, 1945-2011. <http://doi.org/10.1080/10643389.2012.672053>, 2013.

1640 Silver, W.L., Lugo, A.E., Keller, M., Soil oxygen availability and biogeochemistry along rainfall and topographic gradients in upland wet tropical forest soils. *Biogeochemistry* 44, 301-328. <http://doi.org/10.1007/bf00996995>, 1999.

Singh, B.K., Bardgett, R.D., Smith, P., Reay, D.S., Microorganisms and climate change: terrestrial feedbacks and mitigation options. *Nature Reviews Microbiology* 8, 779-790. <http://doi.org/10.1038/nrmicro2439>, 2010.

1645 Sitaula, B.K., Bakken, L.R., Abrahamsen, G., CH₄ uptake by temperate forest soil: effect of N input and soil acidification. *Soil Biology and Biochemistry* 27, 870-880. [http://doi.org/10.1016/0038-0717\(95\)00017-9](http://doi.org/10.1016/0038-0717(95)00017-9), 1995.

Sogn, T., Abrahamsen, G., Effects of N and S deposition on leaching from an acid forest soil and growth of Scots pine (*Pinus sylvestris* L.) after 5 years of treatment. *Forest Ecology and Management* 103, 177-190. [http://doi.org/10.1016/S0378-1127\(97\)00188-6](http://doi.org/10.1016/S0378-1127(97)00188-6), 1998.

1650 [Song, H., Peng, C., Zhu, Q., Chen, Z., Blanchet, J.-P., Liu, Q., Li, T., Li, P., Liu, Z., Quantification and uncertainty of global upland soil methane sinks: Processes, controls, model limitations, and improvements. *Earth-Science Reviews* 252. <http://doi.org/10.1016/j.earscirev.2024.104758>, 2024.](#)

1655 [Stanley, S.H., Prior, S.D., Leak, D.J., Dalton, H., COPPER STRESS UNDERLIES THE FUNDAMENTAL CHANGE IN INTRACELLULAR LOCATION OF METHANE MONO-OXYGENASE IN METHANE-OXIDIZING ORGANISMS - STUDIES IN BATCH AND CONTINUOUS CULTURES. *Biotechnology*](#)

1660

删除了: Schmitz, R.A., Peeters, S.H., Versantvoort, W., Picone, N., Pol, A., Jetten, M.S.M., Op den Camp, H.J.M., Verrucomicrobial methanotrophs: ecophysiology of metabolically versatile acidophiles. *FEMS Microbiology Reviews* 45, fuab007. <http://doi.org/10.1093/femsre/fuab007>, 2021.

Schnell, S., King, G.M., Mechanistic analysis of ammonium inhibition of atmospheric methane consumption in forest soils. *Applied and Environmental Microbiology* 60, 3514-3521. <http://doi.org/10.1128/aem.60.10.3514-3521.1994>, 1994.

- Letters 5, 487-492. <http://doi.org/10.1007/bf00132233>, 1983.
- 1675 Tamai, N., Takenaka, C., Ishizuka, S., Water-soluble Al inhibits methane oxidation at atmospheric concentration levels in Japanese forest soil. *Soil Biology and Biochemistry* 39, 1730-1736. <http://doi.org/10.1016/j.soilbio.2007.01.029>, 2007.
- Tamai, N., Takenaka, C., Ishizuka, S., Tezuka, T., Methane flux and regulatory variables in soils of three equal-aged Japanese cypress (*Chamaecyparis obtusa*) forests in central Japan. *Soil Biology and Biochemistry* 35, 633-641. [http://doi.org/10.1016/s0038-0717\(03\)00010-5](http://doi.org/10.1016/s0038-0717(03)00010-5), 2003.
- 1680 Tie, L.H., Fu, R., Penuelas, J., Sardans, J., Zhang, S.B., Zhou, S.X., Hu, J.X., Huang, C.D., The additions of nitrogen and sulfur synergistically decrease the release of carbon and nitrogen from litter in a subtropical forest. *Forests* 11, 1280. <http://doi.org/10.3390/f11121280>, 2020.
- 1685 van Spanning, R.J.M., Guan, Q., Melkonian, C., Gallant, J., Polarecky, L., Flot, J.F., Brandt, B.W., Braster, M., Iturbe Espinoza, P., Aerts, J.W., Meima-Franke, M.M., Piersma, S.R., Bunduc, C.M., Ummels, R., Pain, A., Fleming, E.J., van der Wel, N.N., Gherman, V.D., Sarbu, S.M., Bodelier, P.L.E., Bitter, W., Methanotrophy by a *Mycobacterium* species that dominates a cave microbial ecosystem. *Nature Microbiology* 7, 2089-2100. <http://doi.org/10.1038/s41564-022-01252-3>, 2022.
- 1690 Veraart, A.J., Steenbergh, A.K., Ho, A., Kim, S.Y., Bodelier, P.L.E., Beyond nitrogen: the importance of phosphorus for CH₄ oxidation in soils and sediments. *Geoderma* 259-260, 337-346. <http://doi.org/10.1016/j.geoderma.2015.03.025>, 2015.
- 1695 Walsh, C., O'Regan, B., Moles, R., Incorporating methane into ecological footprint analysis: A case study of Ireland. *Ecological Economics* 68, 1952-1962. <http://doi.org/10.1016/j.ecolecon.2008.07.008>, 2009.
- Wang, Q., Kwak, J.-H., Choi, W.-J., Chang, S.X., Decomposition of trembling aspen leaf litter under long-term nitrogen and sulfur deposition: effects of litter chemistry and forest floor microbial properties. *Forest Ecology and Management* 412, 53-61. <http://doi.org/10.1016/j.foreco.2018.01.042>, 2018.
- 1700 Wang, Z.P., Ineson, P., Methane oxidation in a temperate coniferous forest soil: effects of inorganic N. *Soil Biology and Biochemistry* 35, 427-433. [http://doi.org/10.1016/s0038-0717\(02\)00294-8](http://doi.org/10.1016/s0038-0717(02)00294-8), 2003.
- Xu, Y.H., Fan, J.L., Ding, W.X., Gunina, A., Chen, Z.M., Bol, R., Luo, J.F., Bolan, N., Characterization of organic carbon in decomposing litter exposed to nitrogen and sulfur additions: links to microbial community composition and activity. *Geoderma* 286, 116-124. <http://doi.org/10.1016/j.geoderma.2016.10.032>, 2017.
- 1705 Yang, C., Chen, Y.T., Zhang, Q., Qie, X.H., Chen, J.X., Che, Y.J., Lv, D.T., Xu, X.Y., Gao, Y.X., Wang, Z.Y., Sun, J., Mechanism of microbial regulation on methane metabolism in saline-alkali soils based on metagenomics analysis. *Journal of Environmental Management* 345, 118771. <http://doi.org/10.1016/j.jenvman.2023.118771>, 2023.
- 1710 Zhang, L.H., Yuan, F.H., Bai, J.H., Duan, H.T., Gu, X.Y., Hou, L.Y., Huang, Y., Yang, M.G., He, J.S., Zhang, Z.H., Yu, L.J., Song, C.C., Lipson, D.A., Zona, D., Oechel, W., Janssens, I.A., Xu, X.F., Phosphorus alleviation of nitrogen-suppressed methane sink in global grasslands. *Ecology Letters* 23, 821-830.
- 1715

删除了: Sullivan, B.W., Selmants, P.C., Hart, S.C., Does dissolved organic carbon regulate biological methane oxidation in semiarid soils? *Glob Chang Biol* 19, 2149-2157. <http://doi.org/10.1111/gcb.12201>, 2013.

删除了: Tate, K.R., Soil methane oxidation and land-use change - from process to mitigation. *Soil Biology and Biochemistry* 80, 260-272. <http://doi.org/10.1016/j.soilbio.2014.10.010>, 2015.

Teh, Y.A., Silver, W.L., Conrad, M.E., Oxygen effects on methane production and oxidation in humid tropical forest soils. *Global Change Biology* 11, 1283-1297. <http://doi.org/10.1111/j.1365-2486.2005.00983.x>, 2005.

删除了: Trotsenko, Y.A., Murrell, J.C., Metabolic aspects of aerobic obligate methanotrophy. *Advances in Applied Microbiology* 63, 183-229. [http://doi.org/10.1016/S0065-2164\(07\)00005-6](http://doi.org/10.1016/S0065-2164(07)00005-6), 2008.

删除了: Wang, M., Wang, L., Shi, H., Liu, Y., Chen, S., Soil bacteria, genes, and metabolites stimulated during sulfur cycling and cadmium mobilization under sodium sulfate stress. *Environmental Research* 201, 111599. <http://doi.org/10.1016/j.envres.2021.111599>, 2021.

删除了: Yun, J., Zhuang, G., Ma, A., Guo, H., Wang, Y., Zhang, H., Community structure, abundance, and activity of methanotrophs in the Zoige wetland of the Tibetan Plateau. *Microbial Ecology* 63, 835-843. <http://doi.org/10.1007/s00248-011-9981-x>, 2012.

Zausig, J., Stepniewski, W., Horn, R., Oxygen concentration redox potential gradients in unsaturated model soil aggregates. *Soil Science Society of America Journal* 57, 908-916. <http://doi.org/10.2136/sssaj1993.03615995005700040005x>, 1993.

<http://doi.org/10.1111/ele.13480>, 2020.

1750 Zhang, T., Zhu, W., Mo, J., Liu, L., Dong, S., Increased phosphorus availability
mitigates the inhibition of nitrogen deposition on CH₄ uptake in an old-growth
tropical forest, southern China. *Biogeosciences* 8, 2805-2813.
| <http://doi.org/10.5194/bg-8-2805-2011>, 2011.