

Sea Ice Freeboard Extrapolation from ICESat-2 to Sentinel-1

Karl Kortum¹, Suman Singha^{2, 3}, and Gunnar Spreen⁴

¹Remote Sensing Technology Institute, German Aerospace Center (DLR), Bremen, Germany

²National Center for Climate Research, Danish Meteorological Institute (DMI), Copenhagen, Denmark

³Department of Geography, University of Calgary, Calgary, Canada

⁴Institute of Environmental Physics, University of Bremen, Bremen, Germany

Correspondence: karl.kortum@dlr.de

Abstract. The Ice, Cloud and Land Elevation Satellite (ICESat-2) laser altimeter can capture sea ice freeboard along-track at both high vertical and high spatial resolution. The measurement occurs along three strong and three weak parallel beams. Thus, the across-track direction is only very sparsely covered, and capturing the two-dimensional spatial distribution of freeboard at high resolution by this instrument alone is not possible. This work shows how in the early Arctic winter months of October and November, Sentinel-1 synthetic aperture radar (SAR) acquisitions help bridge this gap. Freeboard measurements are shown to be meaningfully extrapolated to a full two-dimensional mapping. To achieve this, it is sufficient to use the SAR HV backscatter to sort the pixels by intensity and then map freeboards measured from altimetry in the area via the cumulative distribution functions. With the presented algorithm, the snow and ice freeboard derived from altimetry can be extrapolated to Sentinel-1 SAR scenes, unlocking an additional dimension of Arctic freeboard monitoring at high spatial resolution, with ice freeboard errors between 6 cm and 10.5 cm for spatial resolutions between 100 m and 400 m.

1 Introduction

Due to prevalent feedback loops, amplification in the Arctic makes it Earth's most affected region by climate change (Serreze and Barry (2011); Wendisch et al. (2023) present thorough overviews of the observed amplification). Along with its critical role in Earth's response to global warming, it is also one of the hardest places to monitor consistently. The remoteness and hostility of the environment to the human organism make in situ measurements difficult to obtain. As a result, the global community relies on remote sensing to observe change in the polar regions continuously. Space-borne photography in the optical spectrum is only feasible during polar day for approximately half a year. Passive microwave and active remote sensing techniques thus move to the forefront of operational monitoring of the polar regions. Passive microwave radiometer instruments and corresponding retrieval algorithms deliver robust data products (e.g. Spreen et al. (2008); Markus and Cavalieri (2000)) at the five- to twenty-kilometre scale. Observation of processes at finer spatial scales can only be carried out by active sensors. One such instrument capable of higher-resolution observations is synthetic aperture radar (SAR), delivering year-round backscatter measurements that are sensitive to changes in the ice cover. Due to the diverse backscattering properties that sea ice admits during its diverse development from frazil to perennial ice (see, e.g. Onstott (1992); Kortum et al. (2024)) the corresponding

25 data are more difficult to interpret than optical satellite imagery. This complex relationship between radar backscatter and the physical state of sea ice is a central complication of retrieval algorithms. Continuously operational SAR missions, such as ESA's Sentinel-1, provide SAR data in two polarisation channels. A co-pol channel with horizontal send and receive polarisation (denoted HH) and a cross-pol channel, with horizontal send and vertical receive polarisation (denoted HV). The combination of these two channels grants additional information about the sea ice, yet is still by far not sufficient to solve the inverse problem.

30 An alternative approach to high-resolution monitoring of sea ice is the use of altimeters, which detect the distance to the ground in nadir. In the case of the laser altimeter on ICESat-2, footprint sizes of the measurement are on the order of tens of metres, as detailed in Neumann et al. (2019). Altimeter measurements have low uncertainties of only a few centimetres in their height retrievals and thus allow precise measurements of the distance between the satellite and the scatterer on the ground. If cracks and leads open in the ice cover and open water or thin ice is detected, this distance can be used as a reference for the sea surface

35 height. Thus, measurements of the surrounding sea ice surface are converted to a freeboard measurement, as described, e.g., in Kwok et al. (2022). This is the total height of the ice and snow above the sea surface. Not only is the freeboard indicative of the ice development, series of such measurements can be combined into a topographic understanding of the surface, describing roughness at various scales (Mchedlishvili et al. (2023)). A large blind spot of the altimetry measurement is given by its spatial sparsity in the transversal/across-track direction of the flight path, as measurement takes place only along thin lines over the

40 Arctic. Tracks from multiple flights can be combined to give a large-scale overview on a monthly basis. However, resulting gridded products (Petty et al. (2020)) are constrained to a regional scale (25 km grid cell size) and have to be aggregated for about one month to achieve pan-Arctic coverage.

SAR and altimetry data both yield valuable insights into the Arctic system. At the same time, they are complementary in a variety of aspects: SAR has large 2-dimensional coverage, whilst altimetry coverage is sparse. However, converting radar

45 backscatter data into key measurements of the sea ice is very challenging, whilst laser altimetry measures the sea ice height very precisely, is easy to interpret, and gives concrete information about the sea ice topography. Because of that, some research has already been conducted concerning the combination of both instruments. Karvonen et al. (2022) combined Sentinel-1 SAR and CryoSat-2 radar altimeter measurements of ice thickness, seeking to leverage the advantages of each technique. The technique they developed uses the SAR data to interpolate between the altimetry data at kilometre scale, by segmenting the SAR image

50 and assigning CryoSat-2 measured ice thicknesses to segments. Recently Macdonald et al. (2024) published a study over landfast ice in the Canadian Arctic Archipelago, in which correlations of altimeter measurements (roughness, freeboard) and C-Band SAR HV backscatter appeared stronger than those with HH backscatter. Their research also suggests that roughness retrieval from SAR HV data is feasible. Concerning roughness and SAR HH/VV backscatter, strong correlations ($R_p = 0.82$) were found by Cafarella et al. (2019) under shallow incidence angles for first-year ice and similar correlations ($R_p = 0.74$)

55 were observed in Segal et al. (2020) also over the Canadian Arctic Archipelago. Meaningful correlations of surface roughness at smaller scales could not be observed in Kortum et al. (2024) for spaceborne X-Band SAR and airborne LiDAR data, with ($R_{\text{Pearson}} < 0.3$) over mixed, multiyear ice, second-year ice, and first-year ice over a small area of sea ice in the central Arctic. In this work, we present correlations of freeboard and roughness with C-Band SAR at a near pan-Arctic scope and demonstrate an algorithm to extrapolate ICESat-2 altimetry-derived freeboard to Sentinel-1 SAR scenes at up to 100 m resolution.




Figure 1. Location of all Sentinel-1 scenes from October or November (2018-2022) with near coincident ICESat-2 coverage. These acquisitions are the main source of data for this study.

60 In this study, we are not proposing that SAR backscatter is a direct indicator of sea ice thickness (which might be questionable). We are only using the backscatter intensity in the vicinity of actual ICESat-2 ice freeboard measurements to extrapolate them in space. Locally, one can expect that the ice thickness to SAR backscatter relationship is stable enough to retrieve sea ice freeboard for the whole SAR scene. We extrapolate ICESat-2 freeboard heights to coincident Sentinel-1 SAR scenes, which were acquired within 24 hours of the ICESat-2 overflight. This enables observations near the spatial scope and frequency of
 65 the Sentinel-1 constellation, which is considerably larger than the altimeter coverage alone, but the errors are higher than for the altimetry data, because of the limited correlation of sea ice backscatter and freeboard. A freeboard product with a spatial resolution of up to 100 m and time intervals and coverage of the Sentinel-1 mission, as proposed here, is a useful resource for polar research and stakeholders.

2 Data

70 An overview of all data products used in this study is given below.

The first source data product we use are Sentinel-1 SAR acquisitions, captured in EW (Extended Wide) mode. These scenes have a footprint of approximately 400 km by 400 km with an individual pixel size of 40 metres. We use the Ground Range Detected (GRD) product, which projects the measurement to geo coordinates using an earth ellipsoid model. The terrain

correction in the Sentinel-1 toolbox in SNAP (SNAP (2022)) is used to correct these measurements with a geoid model, which is close to the ocean height and reduces the geolocation error. The incidence angle ranges of the scenes is between 20 and 50 degrees. Thermal noise reduction, scalloping mitigation, and calibration to σ_0 is performed using the SNAP (2022) library and corrections developed by the Nansen Center and described in Park et al. (2018, 2019); Korosov et al. (2022). These mitigation measures help minimise the effect of sensor artefacts on the study. To allow more ICESat-2 footprints to fit into one pixel and thus to derive more meaningful statistics, the scenes are then resampled to 100 m x 100 m pixel spacing. This also mitigates speckle effects. The footprints of all scenes used in this study are plotted in Fig. 1 for an overview.

On the altimetry side, we are using ICESat-2's ATL-10 sea ice freeboard measurement. ICESat-2 is an optical laser altimeter that operates at a wavelength of 532 nm and is highly pulsed at 10.000 pulses a second. The resulting altimetry measurement is accurate to approximately 2 cm. Because the freeboard segments are dependent on the scattering conditions of the surface (a certain number of photons are collected per segment), the spacing of the ATL-10 product is variable and on the scale of tens of metres. At these intervals, segments of freeboard height and expected variance are retrieved. To have as many data points as possible, we use the three weak beams as well as the three strong ones, giving us a maximum of six beams from which data can be used. The benefit of including the less accurate weak beams is investigated later in the manuscript. Due to atmospheric conditions and the requirement of nearby open leads, a freeboard measurement is not always available when the instrument is measuring.

The bulk data in the study consist of 59 Sentinel-1 EW scenes, along with all ICESat-2 ATL-10 freeboard data, within 24 hours of the SAR acquisition, over the same footprint. The specific SAR scenes are selected because there exists an ICESat-2 overflight that is near-coincident with the SAR measurement (the time difference is less than 10 minutes) and the ATL-10 freeboard tracks overlap with at least 300 pixels of the SAR scene, each of $100 \times 100 \text{ m}^2$ size. The near-coincident flights are important for observing the correlations between the measurements and later for validation of the extrapolation results. October and November are selected for two reasons: Firstly, there exist comparatively many near-coincident acquisitions in this time period. This is likely due to atmospheric conditions, i.e., less clouds, as ICESat-2's laser at 532 nm does not penetrate these. Secondly, first-year ice is still quite young at this point and can therefore more easily be distinguished from older ice in both SAR and altimetry missions. As a result, the correlations between freeboard and backscatter are expected to be highest during this time of the Arctic sea ice cycle.

Setting the maximum time difference for a 'near-coincident' measurement at 10 minutes and with pixel sizes of 100 metres, significant decorrelation of both measurements can start to occur if the ice drifts faster than 50 metres in 10 minutes (= 300 m/h). Such high drift speeds are reached occasionally, but this constraint to 10 minutes is sufficient to make sure the vast majority of data points are still valuable. The data are matched using the geocoding of both products and no ice drift correction is applied. For Sentinel-1, the geolocation uncertainties reported by Schubert et al. (2017) are around 5 m over land, which we can use as a baseline error. Additionally, the geoid model used for the ground range projection will have an error relative to the real sea surface height, which should be of a similar scale as the local sea surface height anomaly. Skourup et al. (2017) investigated the model and observational differences and found differences in the central Arctic up to 0.5 m. Thus, we can assume the Sentinel-1 geocoding error is generally below 10 m. The geolocation errors of ICESat-2 are reported to be

around 2.5 m to 4.4 m by Luthcke et al. (2021). As the pixel sizes of 100 m are significantly larger than the uncertainties of the geocoding, we can obtain meaningful overlap between the SAR and freeboard products at this scale.

All ICESat-2 ATL-10 segments in one Sentinel-1 pixel are considered equally: To obtain a local freeboard, the mean of all freeboard segment heights from ATL-10 pertaining to a pixel is taken. For roughness, we investigate two different considerations that describe different roughness correlation lengths (scales). The ATL-10 product gives an expected variance for each freeboard segment, determined by local photon statistics, and thus approximately at metre scale. For the first roughness observation, all freeboard segments' respective variances are summed, and from the square root of their mean a final standard deviation for each pixel is obtained giving a roughness at approximately the metre scale. Alternatively, a larger scale roughness can be obtained by calculating the standard deviation of all ATL-10 freeboard segment heights within one 100 m x 100 m SAR pixel. The correlation length/scale of this roughness measure is equivalent to the spacing of the segments, i.e. on the order of 10s of metres. Both of these roughness measures use the variance of freeboard heights as a proxy for roughness of the ice surface.

2.1 Correlations

We will first investigate the statistical connections of the altimetry and SAR data. In this case, we are mainly interested in the correlations of these variables, as that will be of importance for the extrapolation measures described later.

Heatmaps of both freeboard and roughness from all 597,565 data points are plotted in Fig. 2, along with the respective Spearman correlations. We use Spearman correlations here, as we do not expect the values to be correlated linearly, but we are interested in how accurately we could construct a monotonic mapping from one to the other - which is exactly what the Spearman correlation coefficient captures. In table 1 the Spearman correlation coefficients are listed. The split into multiyear (MYI) and first-year (FYI) ice is performed for 51 of these scenes (with 392,364 data points), which admitted a clearly bimodal freeboard distribution, allowing to differentiate between the two ice types via thresholding the freeboard. The other eight scenes did not show such a split distribution and thus were not considered.

There are three main studies from the Canadian Arctic Archipelago we can compare these results with, all of which focus on fast ice. Cafarella et al. (2019) investigated the statistical relationship of high-resolution C- and L-band SAR data (resolution \approx 10 and 3 m, respectively) with airborne LiDAR-derived sea ice roughness (resolution = 1.2 m) over first-year ice. From two scenes acquired in the late winter season (March, April), they found a high correlation (Pearson's R) of 0.86 for high incidence angles (46 deg) and a low correlation of 0.30 for low incidence angles, for the HH backscatter and roughness. The correlation of the HV backscatter and roughness was found to be more similar across the two scenes, around 0.81 for high and 0.68 for low incidence angles. Segal et al. (2020) observed the correlations of LiDAR-derived roughness, a roughness proxy from the MISR optical satellite and Sentinel-1 C-Band SAR over first-year and multiyear ice in late winter (April). The roughness was derived from 1 m resolution LiDAR data and the grid cells were 1.2 km by 0.4 km large. They found a high correlation (Pearson's) for roughness and HH backscatter at 0.74 across their whole dataset, with 0.76 on only first-year and 0.12 on only multiyear ice. Recently, Macdonald et al. (2024) published a study comparing SAR and altimetry measurements for three ICESat-2 overflights in the Canadian Arctic Archipelago in March. It is also worth noting that they computed roughnesses




Figure 2. 2D Histograms of ICESat-2 freeboard and roughness versus Sentinel-1 HH and HV backscatter measurements, and the respective Spearman correlation coefficients. Brighter colours correspond to higher data density, whilst darker blueish colors correspond to lower density. Some banding effects are visible in the HV channel.

from the University of Maryland supersampled ICESat-2 product, described in Duncan and Farrell (2022); Farrell et al. (2020). As the source for SAR data they used the Radarsat Constellation Mission (RCM) in a low noise mode unique to the instrument and found (Spearman) correlations for first-year ice roughness and SAR backscatter at 0.42 for the HV and 0.31 for the HH channel. The correlations with mutli-year sea ice height and backscatter were 0.49 in the HV and 0.41 in the HH channel. They also demonstrated an accurate roughness retrieval at 800 m scale. The differences of these previous studies and ours are the spatial scales, seasons, and location. While these previous studies were looking at a more regional scale, we use satellite overflights from more diverse Arctic regions. However, our roughness measures are not as fine-scale or accurate as the airborne LiDAR data or the University of Maryland ICESat-2 product. Additionally, we are focusing on the early rather than the late winter season.

The freeboard correlations with the HV channel across our entire dataset are remarkably strong at 0.62. The correlation for MYI and the HV channel is the same as in the Macdonald et al. (2024) study at 0.49. However, the correlations with the roughness are weaker, especially in the HV channel, than in all previous studies. Causes for this could be the ice development, the difference in ice seasons, or the roughness measures used. Comparably low correlations were also found in Kortum et al. (2024) for sea ice roughness at length scales of 0.5 m with the HH and VV channels of X-band SAR. The correlations for freeboard across the entire dataset ($R = 0.62$) might be slightly stronger in this study in contrast with the Macdonald et al.

Table 1. Spearman correlations coefficients of ICESat-2 and Sentinel-1 measurements. The correlations for HH and HV are calculated from all 59 available flights. Of these 51 admitted a bimodal freeboard distribution, allowing the separation of first-year ice (FYI) and multiyear ice (MYI).

	HH	HH(FYI)	HH(MYI)	HV	HV(FYI)	HV(MYI)
freeboard	0.49	0.18	0.34	0.62	0.32	0.49
roughness (1 m)	0.27	0.24	0.20	0.36	0.26	0.31
roughness (10 m)	0.33	0.19	0.22	0.48	0.36	0.36

(2024) study, because of the rescaling to 100 m x 100 m, which should lead to an increase of correlations as quasi-random speckle effects average out. Additionally, the study area and time might be a cause for this, with both very thin first-year ice and the oldest, thickest perennial ice being captured within this study’s dataset. This should also lead to an increase in correlation.

3 Methods

3.1 Algorithm Structure

The structure of the proposed freeboard extrapolation method using SAR backscatter is as follows.

1. For the SAR scene to be used as basis for the extrapolation, all ATL-10 measurements within the last 24 hours are retrieved.
2. A mapping is constructed from the HV SAR data to non-coincident ATL-10 freeboard in the area via the cumulative distribution functions of the HV SAR measurement and the altimeter freeboard product.
3. The mapping is applied to the HV channel of the entire scene from step 1.

This extrapolation using the cumulative distribution functions entirely relies on the correlations of sea ice aging processes and its freeboard, illustrated in Fig. 3. As young ice freezes up, a brine expulsion on top of the ice leads to wet and saline surface and possibly wetted snow, as investigated by e.g. Drinkwater and Crocker (1988). This lossy material is quite absorbent and backscatter is typically quite low, especially for double bounces required for HV returns. Whatever backscatter is measured probably originates from surface roughness features, which also increase freeboard. As the ice gets older and desalinates (Cox and Weeks (1974)), the bulk ice becomes less opaque to the radar waves, thus increasing the volume scattering from bubbles and empty brine channels. In turn, the HV signal becomes stronger. Finally, large topographical features, such as ridges, can accommodate double-bounce backscatter returns and can again increase the HV backscatter return. It is important to note that there is no direct physical connection between the backscatter and ice freeboard, i.e. there is no physical reason why a ridge 1.5 m high should have a stronger HV backscatter response than one only 1 m high and this is therefore the strongest limitation of this approach. However, we propose that, in the vicinity of a measured freeboard distribution from ICESat-2, the backscatter




Figure 3. Illustration of the connection between freeboard and ice development responsible for the increase in HV backscatter (mainly desalination and surface roughness increase).

is a reasonable predictor of relative freeboard heights and can therefore be used to extrapolate the freeboard measurements. This is possible because in the most cases the freeboard distribution does not change drastically on a 100 km scale or within 24 hours. Of course, using coincident flights rather than those within 24 hours would yield better extrapolation results. However, these cases are extremely rare and the coverage of such a product would be extremely sparse.

185 3.2 Cumulative distribution function (CDF) mapping

To create the mapping from Sentinel-1 backscatter to ICESat-2 freeboard using the cumulative distribution functions (CDFs) all ATL-10 data from the last 24 hours within the footprint of the SAR scene are collected. They are resampled to match the 100 m pixel spacing from SAR. For our scenes, the resampling factor was approximately 10, which is used in the following. Their cumulative distribution function CDF_{fb} is formed from all measurements taken. For the CDF of the HV channel CDF_{HV} ,
 190 all pixels within 1000 m of an ICESat-2 track are considered. Because the ice has drifted in between the measurements, it is not the same exact ice that forms both CDFs. However, restriction to the approximate area ensures that the distribution of the underlying ice is similar. The constructed CDF map is illustrated in figure 4 and can be expressed as

$$\begin{aligned} \Phi : \{\sigma_{HV}\} &\mapsto \{fb\} \\ \Phi(\sigma_{HV}) &= (CDF_{fb}^{-1} \circ CDF_{HV})(\sigma_{HV}) \end{aligned} \quad (1)$$

With this mapping constructed, pixels can be mapped from HV backscatter to freeboard for the entirety of the Sentinel-1
 195 acquisition. It is worth noting that the Spearman correlation coefficient is invariant under such a monotonic transformation. Thus, all the improvement between the Spearman correlations of the predicted freeboard and the measured freeboard in contrast to the HV backscatter and the measured freeboard comes from the different CDF mappings for each scene.




Figure 4. Visualisation of the mapping constructed from the cumulative distribution functions of freeboard and HV backscatter. The black path illustrates a mapping from an HV backscatter value to a freeboard value.

3.3 Validation

To validate the results of the method, the procedure described above is performed for all 59 SAR scenes that additionally have a coincident ICESat-2 overflight. To form the cumulative distribution function CDF_{fb} for the freeboard, all ATL-10 data within 24 hours of the SAR acquisition are used, except for the validation flight within ten minutes. Then the extrapolated freeboard is compared with the near-coincident validation overflight over the same scene. This ensures the constructed mapping and extrapolated results are entirely independent of the validation data. Therefore, the validation results are representative of the algorithm performance in ice conditions in October and November. Validating with coincident ICESat-2, instead of helicopter-borne measurements such as collected during MOSAiC (Nicolaus et al. (2022)) or Operation Ice Bridge (MacGregor et al. (2021)), ensures that errors arising from the difference in measurement techniques do not need to be accounted for.

4 Results

Fig. 5 shows the central results of the predicted algorithm. At 100 m resolution a Pearson correlation of 0.68 between the measured and extrapolated freeboard shows that the relationship of HV backscatter and freeboard can be used to make meaningful extrapolation possible. At just above 10 cm, however, the errors are still significantly greater than the uncertainties of the underlying ATL-10 product. Judging also by the heatmap in Fig. 5, at 100 m resolution this technique enables the separation of ice into approximate classes such as first-year or multiyear ice and to detect ridges. As the resolution is reduced, the retrieval method becomes increasingly accurate, as is illustrated by the narrowing of the heatmap. At 400 m resolution, with




Figure 5. Results at different spatial resolutions (100, 200, 400 m) of the extrapolated freeboard data on all 59 scenes (597,565) data points, with Pearson correlation coefficients R_p and mean absolute errors (MAE) shown in the figures. Brighter areas indicate a higher density.

Pearson correlation $R_p = 0.82$ and errors of 6 cm, the retrieval method shows promising results that can unlock comprehensive
 215 freeboard surveys of the Arctic in two dimensions.

An example scene is shown in Fig. 6, where qualitatively the extrapolated freeboard aligns well with the overlaid ATL-
 10 measurements. The bottom track is shown in more detail below in Fig. 7, where it becomes clear that in most cases the
 characteristics are captured well ($R_{\text{Pearson}} = 0.67$), but the exact height (especially for ridged areas) cannot be accurately
 220 approximated (RMSE = 0.08). Occasionally, some younger ice areas are shown to be significantly thinner than assumed from
 the extrapolation. These areas have also posed problems in sea ice classification algorithms in the past, for example, described
 in Guo et al. (2022).

5 Error Analysis

The approach detailed in this manuscript is heavily based on the statistical relationship between SAR HV backscatter and
 freeboard as measured by ICESat-2. Whilst we suspect the limitations of such a purely statistical relationship to be the greatest
 225 source of error for the extrapolation, we can measure the effect of various other contributions to the error directly. In this
 section, we investigate the influence of thermal noise, incidence angle effects, and strong and weak ICESat-2 beam selection
 on the accuracy of the final product. To do so, we split the dataset in a variety of ways. The combined results are presented in
 Tab. 2.

To measure the influence of thermal noise on the freeboard product, we split the Sentinel-1 scenes into two disjoint subsets,
 230 according to the height of the noise floor. As a divisive criterion, we use -30 dB as the limit of the noise floor in the low-noise
 dataset. All data where the noise floor is higher than -30 dB are placed in the high-noise dataset. We then execute our algorithm
 exactly as before and compare the two datasets.

The incidence angle effect of sea ice for the HV channel is not well investigated in contrast to the effect on HH backscatter.
 In Aldenhoff et al. (2020) the slopes are found to be roughly half as steep in the HV as in the HH channel. Kortum et al.




Figure 6. Example scene from the 29th of November 2021 with both the extrapolated freeboard at 100 m resolution and overlaid ICESat-2 ATL-10 data in subfigure A. The ATL-10 data were thickened artificially (using nearest neighbour extrapolation) to allow easier visualisation and are shown within the white contour. The three visible tracks are made up of one strong and one weak beam each. Subfigure B shows the SAR image in false colour. The composition (HV, HH, HV/HH) is chosen for the respective (R, G, B) channels.




Figure 7. Track of southernmost (bottom) beam from figure 6 with measured (ICESat-2, blue) and extrapolated (Sentinel-1, orange) freeboard values at 100 m spacing.

Dataset	R_p (100 m)	MAE [m] (100 m)	R_p (400 m)	MAE [m] (400 m)
High Noise	0.68	0.104	0.83	0.063
Low Noise	0.69	0.102	0.83	0.054
Weak Beams	0.67	0.102	0.81	0.061
Strong Beams	0.67	0.107	0.81	0.063
Slope Corrected	0.67	0.100	0.82	0.055
Control	0.68	0.103	0.82	0.06

Table 2. Investigating the influences of noise, weak/strong beams, and SAR incidence angle on the freeboard extrapolation method through splitting the dataset along various criteria. Results are Pearson’s R (R_p) and mean absolute error (MAE), between extrapolated and measured (coincident) freeboard. We show results for the 100 m product and the 400 m resampling. The control dataset uses the results described in the methods section, with both strong and weak beams and all incidence angles included, but no incidence angle correction carried out. Best in category results are highlighted in bold font.

235 (2023) also find weaker HV slopes in their investigation. Despite the effect being smaller, it still influences the brightnesses and therefore the extrapolation of freeboard. To measure the effect of an incidence angle mitigation strategy on the freeboard extrapolation, we use a Gaussian clustering approach by Cristea et al. (2020). Thus, we obtain HV backscatter versus incidence angle slopes for every pixel in the scene and then use these to correct the entire image to 30 degree incidence angle. We can then compare the accuracy of the freeboard extrapolation with and without the incidence angle correction.

240 Finally, we investigate the inclusion of weak beams of the freeboard measurement by constructing two additional datasets with only weak and only strong beams and comparing with the original one which included both.

From the results in Tab. 2 we can infer the following:

1. Restricting to low SAR backscatter noise areas slightly improves the correlation of extrapolated and measured freeboard. In addition, the errors are lower in the low noise areas (noise floor lower than 30 dB) by $\approx 14\%$. However, such a
245 correction would significantly reduce the coverage of obtained extrapolated data.
2. Restricting to weak or strong beams makes only a small difference. Including both gives the best results.



3. Slope correction improves or matches the results of the control dataset for the investigated measures, and the amount of extrapolated data is not negatively affected.

6 Comparison with Upward Looking Sonar Data


250 To gain additional insight into the extrapolation performance and to demonstrate the usefulness of such an approach, we compare the extrapolated freeboard product to upward looking sonar data at mooring B (78 N, 150 W), acquired by Krishfield et al. (2023). Upward looking sonar measures the ice draft from below. In theory, having both measurements, the freeboard from above and draft from below, available allows us to characterise the ice (and snow) thickness. The advantage of upward looking sonar is that it is constantly acquiring and therefore we can evaluate all scenes that capture the location of the sensor for
255 comparison. We conducted this comparison for the entire November of 2022, using 29 SAR scenes. The results are presented in Fig. 8.

The greatest challenge in bringing these two measurements together is the difference in scales. We are working with a 100-400 m freeboard product, but the footprint of the sonar is only approximately 2 m. To enable sensible comparison, we use ten minutes of sonar data around the acquisition time of the satellite and the 400 m SAR product. As a result, we have
260 a 2 m thick line sampled by the sonar (with the length depending on the drift speed) being compared with 400 m x 400 m area of extrapolated freeboard. This means that the distribution of the ice sampled in the freeboard maps should overlap the sonar coverage, but the area sampled by the satellite is much larger. This is a circumstance we cannot mitigate further. The scatterplot in the top left of Fig. 8 shows that there exists a clear statistical correlation between freeboard and ice draft - as one would expect. It also shows that the relationship breaks down below around 0.5 m of ice draft. For such relatively young ice,
265 the freeboard values are probably not accurate. We investigate this further in two outlier cases A and B. From the freeboard map, it becomes obvious that the dynamic range of the HV measurement is not able to capture the subtleties of the backscatter response, as we are too close to the noise floor. This is apparent from the strong edges of the low ice areas in the freeboard maps. Another outlier, C, shows a high ice draft but medium freeboard. The freeboard map reveals that we are in a rather young ice area, but with signs of ridging, as can be seen from the linear features with higher freeboard. In fact, such a ridge area is
270 right in the measured area. Therefore, we suspect that the sonar sampled a large part of that ridge's keel, while the contribution is only small in SAR. I.e. the difference in sampling scales/footprint sizes is the reason for this strong disagreement. In the final outlier D, we have the opposite scenario, where the freeboard is large, yet the ice draft is not. The freeboard map shows a highly diverse ice area. Again, it is likely that the two distributions sampled by the two measurements are quite different, due to their differences in scales and limited overlap.


275 This brief excursion showed how freeboard extrapolation enables the comparison and combination of altimeter-derived freeboards with additional measurements. It also revealed that thin ice areas below 0.5 m ice thickness with low HV backscatter cannot be accurately extrapolated with the proposed CDF-based mapping. As expected, sampling scales are a considerable challenge with combining upward looking sonar data with satellite sea ice measurements.




A.
freeboard = 0.01 m
sonar = 0.5 m



B.
freeboard = 0.03 m
sonar = 0.44 m



C.
freeboard = 0.12 m
sonar = 1.4 m



D.
freeboard = 0.26 m
sonar = 0.85 m





Figure 8. Comparison of extrapolated freeboard and upward looking sonar data. Four statistical outliers A-D are investigated in more detail, with freeboard extrapolation in the area shown on the right. The centre 16 pixels used for comparison are indicated by the white brackets.

7 Discussion

280 The correlations between the SAR backscatter and altimetry freeboard and roughness data found in this study join those observed in previous studies by Cafarella et al. (2019); Segal et al. (2020); Macdonald et al. (2024) to form a more complete picture of the variability and correlations of SAR and topographic ice properties. As mentioned earlier, the study area and time are probably the main reason for the differences in observed correlations. The data studied here suggest that relating roughness and backscatter is more difficult in the early winter season than in the late winter season investigated by previous studies.
285 However, correlations with freeboard are still significant, which reinforce the notion that they can be successfully related to one another.

This study is the first time these correlations between satellite laser altimeter freeboard and SAR backscatter were observed for drifting sea ice across a large area in the Arctic. The correlations of 0.68 (100 m scale) to 0.82 (400 m scale) of freeboard and the SAR HV channel are remarkably high, considering that there is no direct physical connection between backscatter and
290 freeboard.

The results reveal that the proposed algorithm enables meaningful extrapolation of ice freeboard as measured by ICESat-2, capturing key features and revealing the spatial variability of freeboard in two dimensions at 100 m to 400 m resolution and for the coverage of full 400 km Sentinel-1 scenes. The accuracy of the retrieval is difficult to judge in relation to other methods as no comparable products exist. The algorithm performs accurately enough to separate ice types and ridges at 100 m resolution
295 with errors around 10 cm. At 400 m resolution the method is even more accurate given an error of approximately 6 cm.

We also demonstrated a comparison with other sea ice measurements in the case of upward looking sonar (ULS), which is enabled by the extrapolation effort. The comparison of satellite ice freeboard and ULS ice draft reveals a reasonable correlation of 0.58 between the dataset and that the correspondence breaks down below about 0.5 m ice thickness. However, the difference in measurement scales limits the information that can be derived from such a combination. With additional effort, cases with
300 two satellite acquisitions and largely homogenous drift could be found. In this case, the displacement between the two scenes can be derived, and the drift between points can be assumed to be a straight line. Then the overlap between two measurements would greatly increase and in part the difference of scales could be mitigated.

As previously mentioned, the main source of the remaining retrieval uncertainties is the absence of a physical connection between topography and SAR backscatter, something that cannot be circumvented. Additional sources of error also exist. For
305 example, the footprints of ICESat-2 do not cover the entire pixel they are being mapped to, meaning the ground truth we use for freeboard in every pixel is already contaminated by this undersampling. In addition to the existing uncertainty of the ATL-10 products, SAR noise and speckle effects also contribute to the error. Furthermore, the overlap of the validation overflights is limited by the accuracy of the georeferencing of the sensors. In the case of Sentinel-1, the GRD product uses an ellipsoid model, which can vary up to 10s of metres from the real ocean surface height.

310 Investigations into the incidence angle effect have shown that a brightness correction using slopes derived from a clustering method is a successful measure to mitigate the influence of incidence angle on backscatter, and thus the extrapolation algorithm.

It was also shown in Tab. 2, that restricting to weak beams yielded slightly better results than restricting to strong beams, which is counter-intuitive. The weak beam segments are derived from the same number of photons. As these take longer to accumulate for the weaker beams, the segments become longer. Keeping in mind that the strong extrapolations were evaluated
315 against measurements from strong beams and vice versa for weak beams, we offer two possible explanations for this. Firstly, the beams are not always available (or unavailable) at the same time, so it is possible that the correlation between freeboards and backscatter is stronger in the weak beam dataset simply by chance. The other possibility is that the matching of the pixels via geolocation is not quite pixel perfect, and the longer weak segments align better, as they smooth the validation measurements a little.

320 Overall, this purely statistical mapping is rather simple, given the complexity of the physical relationship between sea ice properties, such as freeboard, and the radar backscatter of a SAR sensor. However, we believe there is great merit in having such a simple and explainable method to advance scientific work in this field. For future work, it is very valuable to have such a baseline algorithm available to compare with or use as a basis for more complex methods.

So far, the extrapolation has been limited to only a certain season in the year, i.e. October/November, where older and
325 younger ice have significantly different freeboards, which increase the correlation with SAR backscatter. Expanding this approach to other seasons and the marginal ice zone will be more challenging. Part of the reason is, that the amount of overlapping data at 10 minutes of time difference, needed to validate the results, is sparser in other months and non-existing inside the marginal ice zone.

We have validated the approach with independent, near-coincident ICESat-2 flights. Comparison with CryoSat-2 radar al-
330 timeter measurements would be the next logical step. Because of the different dominant scattering surface of that radar instrument, however, the freeboard measured by CryoSat-2 is different from that measured by ICESat-2, as shown in Fredensborg Hansen et al. (2024) using the Cryo2Ice data. Therefore, it is less useful as validation data. It would be very interesting to investigate the possibility of extrapolating CryoSat-2 and future CRISTAL measurements using the same method and comparing the results. Additionally, the new surface water and ocean topography (SWOT) altimeter allows for 2D freeboard retrieval
335 that would be a great candidate for validation, or extrapolation. Work by Kacimi et al. (2025) has shown good correlations with ATL-10 freeboard used here. However, SWOTs coverage is restricted to 78° North/South, and therefore it's use for sea ice applications is unfortunately limited, but a case study comparison might be possible.

Whilst we worked with extrapolating ICESat-2s ATL-10 product from NASA, other current or future altimetry products might also be able to be extrapolated with SAR. For example, the University of Maryland product by Farrell et al. (2020);
340 Duncan and Farrell (2022) mentioned earlier would be worth using instead of the ATL-10 data for the roughness approximation as was done in Macdonald et al. (2024). As the main focus in this study was on freeboard, this was not considered.

The uses of a medium to high resolution freeboard product are manifold. The data can be used as a good proxy for sea ice thickness in terms of variability in two dimensions, something that has so far eluded consistent observation. Maritime stakeholders might also profit from these data, as well as weather and climate models, the former of which could be initialised
345 with observations in near-real time. High-resolution digital twin earth models, such as those currently in development by Hoffmann et al. (2023) at ECMWF, might especially benefit from these observations, due to their km-scale grid spacing.

8 Conclusions

Our work presented in this manuscript shows how ICESat-2 derived freeboard measurements can be meaningfully extrapolated with Sentinel-1 SAR measurements at resolutions up to 100 m for the entire 400 km SAR scene with up to a 24 hour time difference between SAR and altimetry acquisitions and an freeboard extrapolation error lower than 10 cm. This algorithm opens up an opportunity to monitor Arctic wide sea ice freeboard in two dimensions, capturing its spatial variability at previously unattainable coverage and making an important step towards monitoring ice thickness. It has yet to be shown that this approach can also work throughout all seasons and regions of the Arctic.

Acknowledgements. This study was funded by Deutsche Forschungsgemeinschaft (DFG) under project name 'MOSAICmicrowaveRS' (SI 2564/1-1 and SP 1128/8-1).

We would like to thank all people involved at ESA's Copernicus program for the acquisition and provision of Sentinel-1 SAR data. Additionally we express our thanks to NASA and everyone involved there with the acquisition and distribution of ICESat-2 data products.

Author contributions. Karl Kortum - Conceptualisation, Formal Analysis, Investigation, Methodology, Writing - original draft.
Suman Singha - Funding acquisition, Project administration, Supervision, Writing - review and editing.
Gunnar Spreen - Funding acquisition, Supervision, Writing - review and editing.

Competing interests. The authors have no competing interests to declare.

References

- Aldenhoff, W., Eriksson, L. E. B., Ye, Y., and Heuzé, C.: First-Year and Multiyear Sea Ice Incidence Angle Normalization of Dual-Polarized Sentinel-1 SAR Images in the Beaufort Sea, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 13, 1540–1550, <https://doi.org/10.1109/JSTARS.2020.2977506>, 2020.
- Cafarella, S. M., Scharien, R., Geldsetzer, T., Howell, S., Haas, C., Segal, R., and Nasonova, S.: Estimation of Level and Deformed First-Year Sea Ice Surface Roughness in the Canadian Arctic Archipelago from C- and L-Band Synthetic Aperture Radar, *Canadian Journal of Remote Sensing*, 45, 457–475, <https://doi.org/10.1080/07038992.2019.1647102>, 2019.
- Cox, G. F. N. and Weeks, W. F.: Salinity Variations in Sea Ice, *Journal of Glaciology*, 13, 109–120, <https://doi.org/10.3189/S0022143000023418>, 1974.
- Cristea, A., van Houtte, J., and Doulgeris, A. P.: Integrating Incidence Angle Dependencies Into the Clustering-Based Segmentation of SAR Images, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 13, 2925–2939, <https://doi.org/10.1109/JSTARS.2020.2993067>, 2020.
- Drinkwater, M. R. and Crocker, G.: Modelling Changes in Scattering Properties of the Dielectric and Young Snow-Covered Sea Ice at GHz Frequencies, *Journal of Glaciology*, 34, 274–282, <https://doi.org/10.3189/S0022143000007012>, 1988.
- Duncan, K. and Farrell, S. L.: Determining Variability in Arctic Sea Ice Pressure Ridge Topography With ICESat-2, *Geophysical Research Letters*, 49, e2022GL100272, <https://doi.org/https://doi.org/10.1029/2022GL100272>, e2022GL100272 2022GL100272, 2022.
- Farrell, S. L., Duncan, K., Buckley, E. M., Richter-Menge, J., and Li, R.: Mapping Sea Ice Surface Topography in High Fidelity With ICESat-2, *Geophysical Research Letters*, 47, e2020GL090708, <https://doi.org/https://doi.org/10.1029/2020GL090708>, e2020GL090708 2020GL090708, 2020.
- Fredensborg Hansen, R. M., Skourup, H., Rinne, E., Høyland, K. V., Landy, J. C., Merkouriadi, I., and Forsberg, R.: Arctic Freeboard and Snow Depth From Near-Coincident CryoSat-2 and ICESat-2 (CRYO2ICE) Observations: A First Examination of Winter Sea Ice During 2020–2022, *Earth and Space Science*, 11, e2023EA003313, <https://doi.org/https://doi.org/10.1029/2023EA003313>, e2023EA003313 2023EA003313, 2024.
- Guo, W., Itkin, P., Singha, S., Paul Doulgeris, A., Johansson, M., and Spreen, G.: Sea ice classification of TerraSAR-X ScanSAR images for the MOSAiC expedition incorporating per-class incidence angle dependency of image texture, *The Cryosphere Discussions*, 2022, 1–29, <https://doi.org/10.5194/tc-2022-86>, 2022.
- Hoffmann, J., Bauer, P., Sandu, I., Wedi, N., Geenen, T., and Thiemert, D.: Destination Earth – A digital twin in support of climate services, *Climate Services*, 30, 100394, <https://doi.org/https://doi.org/10.1016/j.cliser.2023.100394>, 2023.
- Kacimi, S., Jaruwatanadilok, S., and Kwok, R.: SWOT Observations Over Sea Ice: A First Look, *Geophysical Research Letters*, 52, e2025GL116079, <https://doi.org/https://doi.org/10.1029/2025GL116079>, e2025GL116079 2025GL116079, 2025.
- Karvonen, J., Rinne, E., Sallila, H., Uotila, P., and Mäkynen, M.: Kara and Barents sea ice thickness estimation based on CryoSat-2 radar altimeter and Sentinel-1 dual-polarized synthetic aperture radar, *The Cryosphere*, 16, 1821–1844, <https://doi.org/10.5194/tc-16-1821-2022>, 2022.
- Korosov, A., Demchev, D., Miranda, N., Franceschi, N., and Park, J.-W.: Thermal Denoising of Cross-Polarized Sentinel-1 Data in Interferometric and Extra Wide Swath Modes, *IEEE Transactions on Geoscience and Remote Sensing*, 60, 1–11, <https://doi.org/10.1109/TGRS.2021.3131036>, 2022.

- Kortum, K., Singha, S., and Spreen, G.: A Physics-Constrained GAN for Incidence Angle Dependence Estimation of Arctic Sea Ice C-Band Backscatter, <https://doi.org/10.36227/techrxiv.24681153.v1>, 2023.
- Kortum, K., Singha, S., Spreen, G., Hutter, N., Jutila, A., and Haas, C.: SAR deep learning sea ice retrieval trained with airborne laser scanner measurements from the MOSAiC expedition, *The Cryosphere*, 18, 2207–2222, <https://doi.org/10.5194/tc-18-2207-2024>, 2024.
- Krishfield, R., Timmermans, M.-L., Bras, I. L., and Proshutinsky, A.: Beaufort Gyre Observing System (BGOS) - Mooring B 2021-2022, <https://doi.org/doi:10.18739/A2XD0R018.>, 2023.
- 405 Kwok, R., Petty, A. A., Cunningham, G., Markus, T., Hancock, D., Ivanoff, A., Wimert, J., Bagnardi, M., Kurtz, N., and the ICESat-2 Science Team. 2021.: ATLAS/ICESat-2 L3A Sea Ice Freeboard, Version 5, NSIDC: National Snow and Ice Data Center, Boulder, Colorado, <https://doi.org/https://doi.org/10.5067/ATLAS/ATL10.005.>, 2022.
- Luthcke, S. B., Thomas, T. C., Pennington, T. A., Rebold, T. W., Nicholas, J. B., Rowlands, D. D., Gardner, A. S., and Bae, S.: ICESat-2 Pointing Calibration and Geolocation Performance, *Earth and Space Science*, 8, e2020EA001494, <https://doi.org/https://doi.org/10.1029/2020EA001494>, e2020EA001494 2020EA001494, 2021.
- 410 Macdonald, G. J., Scharien, R. K., Duncan, K., Farrell, S. L., Rezia, P., and Tavri, A.: Arctic Sea Ice Topography Information From RADARSAT Constellation Mission (RCM) Synthetic Aperture Radar (SAR) Backscatter, *Geophysical Research Letters*, 51, e2023GL107261, <https://doi.org/https://doi.org/10.1029/2023GL107261>, e2023GL107261 2023GL107261, 2024.
- MacGregor, J. A., Boisvert, L. N., Medley, B., Petty, A. A., Harbeck, J. P., Bell, R. E., Blair, J. B., Blanchard-Wigglesworth, E., Buckley, E. M., Christoffersen, M. S., Cochran, J. R., Csathó, B. M., De Marco, E. L., Dominguez, R. T., Fahnestock, M. A., Farrell, S. L., Gogineni, S. P., Greenbaum, J. S., Hansen, C. M., Hofton, M. A., Holt, J. W., Jezek, K. C., Koenig, L. S., Kurtz, N. T., Kwok, R., Larsen, C. F., Leuschen, C. J., Locke, C. D., Manizade, S. S., Martin, S., Neumann, T. A., Nowicki, S. M., Paden, J. D., Richter-Menge, J. A., Rignot, E. J., Rodríguez-Morales, F., Siegfried, M. R., Smith, B. E., Sonntag, J. G., Studinger, M., Tinto, K. J., Truffer, M., Wagner, T. P., Woods, J. E., Young, D. A., and Yungel, J. K.: The Scientific Legacy of NASA's Operation IceBridge, *Reviews of Geophysics*, 59, e2020RG000712, <https://doi.org/https://doi.org/10.1029/2020RG000712>, e2020RG000712 2020RG000712, 2021.
- 420 Markus, T. and Cavalieri, D.: An enhancement of the NASA Team sea ice algorithm, *IEEE Transactions on Geoscience and Remote Sensing*, 38, 1387–1398, <https://doi.org/10.1109/36.843033>, 2000.
- Mchedlishvili, A., Lüpkes, C., Petty, A., Tsamados, M., and Spreen, G.: New estimates of pan-Arctic sea ice–atmosphere neutral drag coefficients from ICESat-2 elevation data, *The Cryosphere*, 17, 4103–4131, <https://doi.org/10.5194/tc-17-4103-2023>, 2023.
- 425 Neumann, T. A., Martino, A. J., Markus, T., Bae, S., Bock, M. R., Brenner, A. C., Brunt, K. M., Cavanaugh, J., Fernandes, S. T., Hancock, D. W., Harbeck, K., Lee, J., Kurtz, N. T., Luers, P. J., Luthcke, S. B., Magruder, L., Pennington, T. A., Ramos-Izquierdo, L., Rebold, T., Skoog, J., and Thomas, T. C.: The Ice, Cloud, and Land Elevation Satellite – 2 mission: A global geolocated photon product derived from the Advanced Topographic Laser Altimeter System, *Remote Sensing of Environment*, 233, 111325, <https://doi.org/https://doi.org/10.1016/j.rse.2019.111325>, 2019.
- 430 Nicolaus, M., Perovich, D. K., Spreen, G., Granskog, M. A., von Albedyll, L., Angelopoulos, M., Anhaus, P., Arndt, S., Belter, H. J., Bessonov, V., Birnbaum, G., Brauchle, J., Calmer, R., Cardellach, E., Cheng, B., Clemens-Sewall, D., Dadic, R., Damm, E., de Boer, G., Demir, O., Dethloff, K., Divine, D. V., Fong, A. A., Fons, S., Frey, M. M., Fuchs, N., Gabarró, C., Gerland, S., Goessling, H. F., Gradinger, R., Haapala, J., Haas, C., Hamilton, J., Hannula, H.-R., Hendricks, S., Herber, A., Heuzé, C., Hoppmann, M., Høyland, K. V., Huntemann, M., Hutchings, J. K., Hwang, B., Itkin, P., Jacobi, H.-W., Jaggi, M., Jutila, A., Kaleschke, L., Katlein, C., Kolabutin, N., Krampe, D., Kristensen, S. S., Krumpen, T., Kurtz, N., Lampert, A., Lange, B. A., Lei, R., Light, B., Linhardt, F., Liston, G. E., Loose, B., Macfarlane, A. R., Mahmud, M., Matero, I. O., Maus, S., Morgenstern, A., Naderpour, R., Nandan, V., Niubom, A., Oggier, M., Oppelt, N., Pätzold,
- 435

- F., Perron, C., Petrovsky, T., Pirazzini, R., Polashenski, C., Rabe, B., Raphael, I. A., Regnery, J., Rex, M., Ricker, R., Riemann-Campe, K., Rinke, A., Rohde, J., Salganik, E., Scharien, R. K., Schiller, M., Schneebeili, M., Semmling, M., Shimanchuk, E., Shupe, M. D., Smith, M. M., Smolyanitsky, V., Sokolov, V., Stanton, T., Stroeve, J., Thielke, L., Timofeeva, A., Tonboe, R. T., Tavri, A., Tsamados, M., Wagner, D. N., Watkins, D., Webster, M., and Wendisch, M.: Overview of the MOSAiC expedition: Snow and sea ice, *Elementa: Science of the Anthropocene*, 10, <https://doi.org/10.1525/elementa.2021.000046>, 2022.
- 440 Onstott, R. G.: SAR and Scatterometer Signatures of Sea Ice, chap. 5, pp. 73–104, American Geophysical Union (AGU), <https://doi.org/https://doi.org/10.1029/GM068p0073>, 1992.
- Park, J., Korosov, A. A., Babiker, M., Sandven, S., and Won, J.: Efficient Thermal Noise Removal for Sentinel-1 TOPSAR Cross-Polarization Channel, *IEEE Transactions on Geoscience and Remote Sensing*, 56, 1555–1565, <https://doi.org/10.1109/TGRS.2017.2765248>, 2018.
- 445 Park, J., Won, J., Korosov, A. A., Babiker, M., and Miranda, N.: Textural Noise Correction for Sentinel-1 TOPSAR Cross-Polarization Channel Images, *IEEE Transactions on Geoscience and Remote Sensing*, 57, 4040–4049, <https://doi.org/10.1109/TGRS.2018.2889381>, 2019.
- Petty, A. A., R. Kwok, M. B., Ivanoff, A., Kurtz, N., Lee, J., Wimert, J., and Hancock, D.: ATLAS/ICESat-2 L3B Daily and Monthly Gridded Sea Ice Freeboard, Boulder, Colorado USA, <https://doi.org/https://doi.org/10.5067/ATLAS/ATL20.001>, 2020.
- 450 Schubert, A., Miranda, N., Geudtner, D., and Small, D.: Sentinel-1A/B Combined Product Geolocation Accuracy, *Remote Sensing*, 9, <https://doi.org/10.3390/rs9060607>, 2017.
- Segal, R. A., Scharien, R. K., Cafarella, S., and Tedstone, A.: Characterizing winter landfast sea-ice surface roughness in the Canadian Arctic Archipelago using Sentinel-1 synthetic aperture radar and the Multi-angle Imaging SpectroRadiometer, *Annals of Glaciology*, 61, 284–298, <https://doi.org/10.1017/aog.2020.48>, 2020.
- 455 Serreze, M. C. and Barry, R. G.: Processes and impacts of Arctic amplification: A research synthesis, *Global and Planetary Change*, 77, 85–96, <https://doi.org/https://doi.org/10.1016/j.gloplacha.2011.03.004>, 2011.
- Skourup, H., Farrell, S. L., Hendricks, S., Ricker, R., Armitage, T. W. K., Ridout, A., Andersen, O. B., Haas, C., and Baker, S.: An Assessment of State-of-the-Art Mean Sea Surface and Geoid Models of the Arctic Ocean: Implications for Sea Ice Freeboard Retrieval, *Journal of Geophysical Research: Oceans*, 122, 8593–8613, <https://doi.org/https://doi.org/10.1002/2017JC013176>, 2017.
- 460 SNAP: SNAP - ESA Sentinel Application Platform, <http://step.esa.int/>, software available from tensorflow.org, 2022.
- Spreen, G., Kaleschke, L., and Heygster, G.: Sea ice remote sensing using AMSR-E 89-GHz channels, *Journal of Geophysical Research: Oceans*, 113, <https://doi.org/https://doi.org/10.1029/2005JC003384>, 2008.
- Wendisch, M., Brückner, M., Crewell, S., Ehrlich, A., Notholt, J., Lüpkes, C., Macke, A., Burrows, J. P., Rinke, A., Quaas, J., Maturilli, M., Schemann, V., Shupe, M. D., Akansu, E. F., Barrientos-Velasco, C., Bärfuss, K., Blechschmidt, A.-M., Block, K., Bougoudis, I., Bozem, H., Böckmann, C., Bracher, A., Bresson, H., Bretschneider, L., Buschmann, M., Chechin, D. G., Chylik, J., Dahlke, S., Deneke, H., Dethloff, K., Donth, T., Dorn, W., Dupuy, R., Ebell, K., Egerer, U., Engelmann, R., Eppers, O., Gerdes, R., Gierens, R., Gorodetskaya, I. V., Gottschalk, M., Griesche, H., Gryanik, V. M., Handorf, D., Harm-Altstädter, B., Hartmann, J., Hartmann, M., Heinold, B., Herber, A., Herrmann, H., Heygster, G., Höschel, I., Hofmann, Z., Hölemann, J., Hünerbein, A., Jafariserajehlou, S., Jäkel, E., Jacobi, C., Janout, M., Jansen, F., Jourdan, O., Jurányi, Z., Kalesse-Los, H., Kanzow, T., Käthner, R., Kliesch, L. L., Klingebiel, M., Knudsen, E. M., Kovács, T., Körtke, W., Krampe, D., Kretzschmar, J., Kreyling, D., Kulla, B., Kunkel, D., Lampert, A., Lauer, M., Lelli, L., von Lerber, A., Linke, O., Löhnert, U., Lonardi, M., Losa, S. N., Losch, M., Maahn, M., Mech, M., Mei, L., Mertes, S., Metzner, E., Mewes, D., Michaelis, J., Mioche, G., Moser, M., Nakoudi, K., Neggers, R., Neuber, R., Nomokonova, T., Oelker, J., Papakonstantinou-Presvelou, I., Pätzold, F., Pefanis, V., Pohl, C., van Pinxteren, M., Radovan, A., Rhein, M., Rex, M., Richter, A., Risse, N., Ritter, C., Rostosky, P., Rozanov,
- 470

475 V. V., Donoso, E. R., Garfias, P. S., Salzmann, M., Schacht, J., Schäfer, M., Schneider, J., Schnierstein, N., Seifert, P., Seo, S., Siebert, H., Soppa, M. A., Spreen, G., Stachlewska, I. S., Stapf, J., Stratmann, F., Tegen, I., Viceto, C., Voigt, C., Vountas, M., Walbröl, A., Walter, M., Wehner, B., Wex, H., Willmes, S., Zanatta, M., and Zeppenfeld, S.: Atmospheric and Surface Processes, and Feedback Mechanisms Determining Arctic Amplification: A Review of First Results and Prospects of the (AC)3 Project, *Bull. Am. Meteorol. Soc.*, 104, E208–E242, <https://doi.org/10.1175/bams-d-21-0218.1>, 2023.