

Simulating the drought response of European tree species with the dynamic vegetation model LPJ-GUESS (v4.1, 97c552c5)

Benjamin F. Meyer^{1, *}, João P. Darela-Filho¹, Konstantin Gregor¹, Allan Buras¹, Qiao-Lin Gu¹, Andreas Krause¹, Daijun Liu², Phillip Papastefanou³, Sijeh Asuk⁴, Thorsten E. E. Grams⁵, Christian S. Zang⁶, and Anja Rammig¹

¹Technical University of Munich, Professorship of Land Surface-Atmosphere Interactions, TUM School of Life Sciences, Freising, Germany

²Department of Botany and Biodiversity Research, University of Vienna, Rennweg 14, 1030 Vienna, Austria

³Department Biogeochemical Signals, Max-Planck-Institute for Biogeochemistry, Hans-Knoll-Str., 10, Jena, 07745, Thuringia, Germany

⁴Department of Geography and Environment, School of Social Sciences and Humanities, Loughborough University, Loughborough, UK. LE11 3TU

⁵Technical University of Munich, Professorship of Land Surface-Atmosphere Interactions, Ecophysiology of Plants, TUM School of Life Sciences, Freising, Germany

⁶University of Applied Sciences Weihenstephan-Triesdorf, Professorship of Forests and Climate Change, Freising, Germany

Correspondence: Benjamin F. Meyer (ben.meyer@tum.de)

Abstract. Due to climate change severe drought events have become increasingly commonplace across Europe in recent decades with future projections indicating that this trend will likely continue, posing questions about the continued viability of European forests. Observations from the most recent pan-European droughts suggest that these types of "hotter droughts" may acutely alter the carbon balance of European forest ecosystems. Yet, substantial uncertainty remains regarding the possible future impacts of severe drought on the European forest carbon sink. Dynamic vegetation models can help to shed light on such uncertainties, however, the inclusion of dedicated plant hydraulic architecture modules in these has only recently become more widespread. Such developments intended to improve model performance also tend to add substantial complexity, yet, the sensitivity of the models to newly introduced processes is often left undetermined. Here, we describe and evaluate the recently developed mechanistic plant hydraulic architecture version of LPJ-GUESS and provide a parameterization for 12 common European forest tree species. We quantify the uncertainty introduced by the new processes using a variance-based global sensitivity analysis. Additionally, we evaluate the model against water and carbon fluxes from a network of eddy covariance flux sites across Europe. Our results indicate that the new model is able to capture drought-induced patterns of evapotranspiration along an isohydric gradient and manages to reproduce flux observations during drought better than standard LPJ-GUESS. Further, the sensitivity analysis suggests that hydraulic process related to hydraulic failure and stomatal regulation play the largest roles in shaping model response to drought.

1 Introduction

For the past decades, the face of European forests has been increasingly marred by heatwaves and droughts – effects of anthropogenic climate change (Ciais et al., 2005; European Environment Agency, 2019; Bigler and Vitasse, 2021; Fink et al., 2004). Severe pan-European droughts in 2003, 2018, and 2022 in combination with record high temperatures (so-called "hotter droughts") caused record reductions in forest growth and productivity as a result of defoliation, higher susceptibility to biotic agents, and mortality (Buras et al., 2023; Ciais et al., 2005; Schuldt et al., 2020; van der Woude et al., 2023). Concerningly, the most recent carbon losses induced by the 2022 hotter-drought have turned central European forests from a carbon sink to a carbon source (van der Woude et al., 2023). With more frequent and intense droughts looming on the horizon, the future of the European forest carbon sink remains uncertain (Brodribb et al., 2020; Cook et al., 2020; Pan et al., 2024, 2011). While dynamic vegetation models (DVMs) are popular tools commonly used to shed light on such uncertainties and estimate possible future impacts on the vegetation carbon sink, many of the established models display strongly diverging simulations in regard to the effects of drought and heat (Tschumi et al., 2023). In an attempt to ensure that future vegetation changes and the associated feedbacks on the water and carbon cycles can be simulated confidently, the latest generation of dynamic vegetation models features increasingly detailed representations of plant hydraulic architecture (Xu et al., 2016; Yao et al., 2022; Kennedy et al., 2019; Xu et al., 2023; Eller et al., 2018, 2020; Christoffersen et al., 2016).

In the simplest terms, ~~such representation~~ these representations of hydraulic architecture ~~considers~~ tend to consider two distinct drivers of drought induced stress: insufficient water availability in the soil and increased atmospheric demand for water (Papastefanou et al., 2020). The balance between supply and demand determines whether a tree will experience drought stress or not. The link between these two ends of the system is the hydraulic architecture of the tree which utilizes the xylem to transport water from the the roots through the stem and ultimately to the leaves where it is transpired through the stomata into the atmosphere (Lambers and Oliveira, 2019). Disruptions of this pipeline due to cavitation or stomatal closure trigger symptoms commonly associated with drought stress. As the ability of trees to transport water declines, other processes such as photosynthetic assimilation and growth cease (Lambers and Oliveira, 2019; Choat et al., 2012). Ultimately, critical dehydration – either directly or through predisposing affected trees to pathogens or insect attack – leads to tree death (Anderegg et al., 2012; McDowell et al., 2008; Hajek et al., 2022; Bigler et al., 2006).

Earlier DVMs generally included simple mechanisms to simulate drought stress, frequently opting for empirical approaches to reduce photosynthetic assimilation during periods of low water availability (Powell et al., 2013; Smith et al., 2001; Zhou et al., 2013). This strategy does not account for the mechanistic links between species-specific hydraulic traits, such as xylem vulnerability to cavitation, stomatal response to atmospheric drying, and xylem conductivity, which have been shown to play a key role in modulating the impact of drought conditions on forests in terms of both productivity and mortality (Hajek et al., 2022; Anderegg et al., 2016, 2015). To account for this behavior, current DVMs are increasingly including mechanistic, process-based representations of plant hydraulic

architecture with functional diversity in regards to stomatal control, water-potential regulation, water-flow through the soil-plant-atmosphere continuum, and hydraulic failure under drought conditions (Xu et al., 2016, 2023; Eller et al., 2018, 2020; Yao et al., 2022; Kennedy et al., 2019; Christoffersen et al., 2016; De Kauwe et al., 2020; Papastefanou et al., 2020, 2024).

55 While these improvements prove valuable in predicting the response of forests to present and future drought, they add further complexity to already complex models by introducing new parameters and processes potentially contributing to increased uncertainty between projections from various models (Oberpriller et al., 2022; Zaehle et al., 2005). Identifying the causes of uncertainty can help guide future model development, highlight the need for more observations of key traits, and determine which model processes may be over- or underrepresented compared to
60 reality (Zaehle et al., 2005; Dietze et al., 2018). In this context, global sensitivity analysis is commonly used to detect the sensitivity of model outputs to model parameters (Saltelli, 2008). Due to the complexity of DVMs and the associated computational demand in performing a comprehensive global sensitivity analysis, such analyses are rare and not consistently applied each time new processes are implemented and new parameters are introduced (Oberpriller et al., 2022). Nevertheless, these analyses remain paramount for enhancing our understanding of the
65 internal model processes and are invaluable in allowing solid interpretation of model results (Oberpriller et al., 2022; Zaehle et al., 2005; Pappas et al., 2013).

Here, we describe and examine the recently developed mechanistic hydraulic architecture in LPJ-GUESS, termed LPJ-GUESS-HYD, intended to more accurately capture tree drought responses based on the theoretical framework of isohydricity (Papastefanou et al., 2024). The concept of isohydricity has been used to classify the response
70 patterns of trees to drought (Tardieu et al., 2015; Jones and Sutherland, 1991) based in part on the sensitivity of leaf water potential to changes in canopy conductance (Klein, 2014). LPJ-GUESS-HYD builds upon a previous version of LPJ-GUESS with mechanistic plant hydraulic architecture which, although ~~not implementing it did not~~ implement the impact of xylem cavitation and stomatal regulation related to isohydricity, nevertheless was able to reproduce patterns of potential natural vegetation (Hickler et al., 2006). LPJ-GUESS-HYD expands upon this earlier
75 version by including a dynamic representation of species-specific water-potential regulation related to the concept of isohydricity (Papastefanou et al., 2020) and explicitly coupling the model representation of evapotranspiration to canopy conductance governed by plant hydraulic processes (Papastefanou et al., 2024) in contrast to the standard version of LPJ-GUESS which only does so during periods of limited water availability.

To thoroughly evaluate the implemented processes related to drought induced stress and the sensitivity of the model
80 to the model parameters governing these processes we conduct a variance-based global sensitivity analysis (Saltelli, 2008; Saltelli et al., 2010). To forego the limitations associated with the complexity of DVMs and the computational demand of running a sensitivity analysis, we focus on the newly introduced parameters governing the plant drought response. Accordingly, we compiled parameter ranges for 12 major European forest tree species from observations and analyzed their sensitivities by simulating a network of 34 eddy covariance flux sites throughout Europe (Warm

85 Winter 2020 Team and ICOS Ecosystem Thematic Centre, 2022). Furthermore, we establish viable parameterizations for our set of 12 species to compare simulated and observed evapotranspiration and gross primary productivity across the European forest sites.

We aim to answer the following questions:

- 90 1. Which of the seven newly introduced parameters related to hydraulic architecture introduce the most uncertainty to LPJ-GUESS-HYD?
2. Does the inclusion of hydraulic architecture reflect species-specific drought responses along an isohydricity gradient in the model, that is, under increasing drought will anisohydric species continue to transpire more than isohydric species?
- 95 3. Does LPJ-GUESS-HYD represent an improvement over LPJ-GUESS in depicting the drought response as represented by changes in GPP and evapotranspiration in European forest ecosystems when compared to observational data from eddy-covariance flux towers?

2 Methods

2.1 Description of standard version of LPJ-GUESS

LPJ-GUESS is a dynamic vegetation model simulating terrestrial ecosystem dynamics on a regional to global scale
100 driven by atmospheric CO₂, gridded meteorological inputs, nitrogen deposition, and soil physical properties (Smith et al., 2001, 2014). The model has been successfully applied and evaluated on global (e.g. Seiler et al., 2022) and regional scale (e.g. Hickler et al., 2012) for a wide range of applications in both managed (e.g. Lindeskog et al., 2021) and natural forest ecosystems (e.g. Ahlström et al., 2012). The following sections will provide an overview of LPJ-GUESS with particular focus on ~~modeled processes which are~~ the model processes critical to the representation
105 of drought effects on individual trees.

2.1.1 Representation of vegetation in LPJ-GUESS

Within each simulated gridcell or site, replicate patches serve as random samples of the entire landscape to account for disturbance- and ~~stand-development-related~~ stand-development-related differences between vegetation stands. Vegetation dynamics in each patch emerge from the competition of different age cohorts of plant functional types
110 (PFTs) or species for space, light, water, and nutrients. Individuals within a cohort are identical in age and size. Typically, PFTs represent classes of tree species with similar attributes in regards to characteristics such as phenology, shade-tolerance, bioclimatic limits, etc., that are described by a common set of parameters. Here, we use the parameterization developed by Hickler et al. (2012) and expanded upon by Lindeskog et al. (2021) to simulate a subset of the most pertinent European tree species. Except for the newly introduced hydraulic parameters (Table 2),
115 all species parameters are identical to those in Lindeskog et al. (2021).

LPJ-GUESS simulates photosynthesis and stomatal conductance based on the BIOME3 model (Sykes and Prentice, 1996) along with respiration, and phenology on a daily basis. At the end of each simulation year, accumulated net primary productivity (NPP) is allocated to leaves, roots, and sapwood following allometric constraints (Sitch et al., 2003). Population dynamics (establishment and mortality) and patch-destroying disturbances are simulated stochastically on a yearly time-step. Soil carbon and nitrogen cycles are simulated based on the CENTURY model (Parton et al., 2010; Kirschbaum and Paul, 2002; Parton et al., 1993; Comins and McMurtrie, 1993).

2.1.2 Soil hydrology

Soil hydrology is represented ~~by~~^{as} a “leaky bucket” model with percolation between layers based on Gerten et al. (2004) albeit with 15 soil layers (each 10 cm thick) instead of the original two (Zhou et al., 2024). The first five soil layers are considered “surface” layers and the remaining ten are referred to as “deep” layers. For each soil layer, l (1 to 15), the available water holding capacity (awc_l ; mm) is determined by the volumetric water content at wilting point (wp_l ; mm mm⁻¹), the (volumetric) field capacity (fc_l ; mm mm⁻¹) and the soil layer thickness (Dz_l ; mm) as:

$$awc_l = (fc_l - wp_l) * Dz_l \quad (1)$$

Field capacity and wilting point are determined by physical soil texture properties (e.g. clay-, sand- and silt-fraction, soil carbon content, bulk density, etc.) provided as input to the model and are the same for all layers. The dimensionless ratio of awc_l to the actual available liquid water in the soil (aw_l ; mm) is defined as the water content ($wcont \in [0, 1]$):

$$wcont_l = \frac{aw_l}{awc_l} \quad (2)$$

which indicates the amount of water available to plants in any given soil layer. Water input to soil comes from rainfall and snowmelt which are initially distributed among the five surface layers and subsequently percolate to the deeper layers. Water leaves the soil via evapotranspiration – where evaporation occurs from the fraction of soil not covered by vegetation and transpiration is dependent on vegetation characteristics – and runoff.

2.1.3 Water availability dynamics

In the standard version of LPJ-GUESS only a few processes are limited by water availability, but the plant hydraulic architecture is not explicitly modeled. Nevertheless, certain processes are affected by limited water availability reflecting plant responses to drought. Initially, low water availability – drought – constrains the establishment of new plant individuals. Each species is assigned a drought tolerance level from 0 (extremely drought tolerant) to 1 (extremely drought intolerant) based on the water content as a fraction of the available water holding capacity

required for that species to establish. This tolerance level is compared to the growing season average water content integrated over the upper five soil layers:

$$145 \quad establish = \begin{cases} \text{false,} & \text{drought_tolerance} > wcont \\ \text{true,} & \text{drought_tolerance} \leq wcont \end{cases} \quad (3)$$

Additionally, drought can limit photosynthetic assimilation by downregulating canopy conductance (g_c ; mm s^{-1}) and restricting the ratio (χ_{CO_2}) of inter-cellular CO_2 (c_i ; ppm) to ambient CO_2 (c_a ; ppm) (Haxeltine and Prentice, 1996). Photosynthesis is modeled based on the Collatz simplification of the Farquhar model (Collatz et al. 1991) described in detail in Haxeltine and Prentice (1996) and Sitch et al., (2003). When water supply is ample, the optimal canopy
150 conductance for photosynthesis is calculated as:

$$g_{cp} = g_{min} + \frac{1.6 * A_{dt}}{c_a * (1 - \lambda_{max})} \quad (4)$$

where g_{min} is the species-specific minimum canopy conductance (a parameter), A_{dt} is the daytime net assimilation, and λ_{max} is a species-specific parameter. Conversely, when water supply is limited, photosynthesis is calculated using *actual* rather than *maximum potential* canopy conductance, g_{cp} , and χ_{CO_2} where the actual canopy conductance, g_c ,
155 is calculated as:

$$g_c = -g_{min} * \ln\left[\frac{1 - E_{su}}{E_q * \alpha_m}\right] \quad (5)$$

where E_q is the equilibrium transpiration (mm s^{-1}), E_{su} is the water supply (mm s^{-1}), and α_m is an empirical parameter (Haxeltine and Prentice, 1996). This calculation is triggered under water-stressed conditions, i. e. when the supply of water from the soil (E_{su} ; mm s^{-1}) determined by the species-specific maximum transpiration rate (E_{max} ,
160 a species-specific parameter; mm s^{-1}) and the the soil moisture availability in the rooting zone (W_r , mm s^{-1}), that is, the fraction of soil water content accessible to an individual based on the parameterized species-specific root distribution across all soil layers (Haxeltine and Prentice, 1996):

$$E_{su} = E_{max} * W_r \quad (6)$$

is not sufficient to satisfy demand indicated by E_{de} :

$$165 \quad E_{de} = \frac{E_q * \alpha_m * g_c}{g_c + g_m} \quad (7)$$

Consequently, g_c is reduced to ensure that plant transpiration (E , mm s⁻¹) matches the supply (E_{su}) such that:

$$E = \min\{E_{su}, E_{de}\} \quad (8)$$

2.2 Description of hydraulic architecture as implemented in LPJ-GUESS-HYD

LPJ-GUESS-HYD provides a more ~~in-depth~~ in-depth implementation of plant physiological processes related to
 170 water availability (Papastefanou et al., 2024). Strategies for water-potential regulation along the isohydric spectrum
 determine how species react to changes in soil water availability (Papastefanou et al., 2020). The resulting water
 potential gradient governs the flow of water through the plant and, based on Darcy's law (Whitehead, 1998),
 determines the supply of water available for transpiration (Hickler et al., 2006). Atmospheric demand for water
 is driven by vapor pressure deficit (~~VDP~~ VPD) and, together with the supply of water, ultimately governs canopy
 175 conductance for photosynthetic assimilation. Lastly, to model the impact of drought on tree mortality, LPJ-GUESS-
 HYD includes an empirical representation of hydraulic failure mortality based on xylem cavitation. These new
 processes seamlessly integrate into the existing structure of LPJ-GUESS and primarily replace empirical relationships
 between soil hydrology and photosynthetic assimilation (Fig. 1).

2.2.1 Water-potential regulation

180 LPJ-GUESS-HYD incorporates the dynamic model for water-potential regulation introduced by Papastefanou et al.
 (2020). This model operates on the principle that water transport from the roots through the stem to the leaves and
 into the atmosphere is dictated by a dynamically changing forcing pressure ($\Delta\psi(t)$, MPa):

$$\Delta\psi(t) = \psi_l(t) - \psi_s(t) - \rho g h \quad (9)$$

where, $\psi_s(t)$ (MPa) and $\psi_l(t)$ (MPa) are the respective soil and leaf water potential at time t . The gravitational pull
 185 is defined by $\rho * g * h$, with ρ (kg m⁻³) referring to the density of liquid water, g (m s⁻²) the gravitational acceleration,
 and h (m) the canopy height. In situations with ample soil water supply $\Delta\psi$ is denoted as $\Delta\psi_{max}$, a parameter
 describing the average forcing potential under well-watered conditions.

Soil water potential is initially calculated as a function of soil water content according to (Saxton et al., 1986) for
 each soil layer ly :




Figure 1. Flow chart displaying the model structure of LPJ-GUESS-HYD including links to standard LPJ-GUESS processes. Objects in blue are introduced by LPJ-GUESS-HYD while objects outlined in black are part of the standard LPJ-GUESS structure. Lines between boxes identify links between individual process, drivers, and parameters. Arrows indicate directionality. Dotted lines highlight links between processes in LPJ-GUESS which are replaced by an alternative structure in LPJ-GUESS-HYD. The light blue diamonds indicate the hydraulic parameters introduced by LPJ-GUESS-HYD and defined in Table 1.

$$190 \quad \psi_{s_{ly}} = A * \text{wcont}_{tot,ly}^B \quad (10)$$

where A and B are functions of soil physical properties such as the clay and sand content (~~see Saxton et al. (1986);~~ Eqs. A1 and A2; and see Saxton et al. (1986);) and wcont_{tot} (mm) is the sum of plant available soil water and the soil water content at wilting point.

Subsequently, $\psi_{s_{ly}}$ is weighted by the fraction of roots in that layer ($r_{f_{ly}}$) to give the integrated soil water potential

$$195 \quad \psi_s:$$

$$\psi_s = \sum_{l=0}^{N_l-1} \psi_{s_l} * r_{f_l} \quad (11)$$

where N_l is the number of soil layers.

The model assumes that the change in ψ_l over time depends on the difference between ψ_l and ψ_s such that:

$$\frac{d\psi_L}{dt} = \alpha((1 - \lambda) * \psi_s(t) - \psi_l(t)) - \Delta\psi_{max} \quad (12)$$

200 where, $\lambda \in [0, 1]$ (Table 1) is a component of the isohydricity of water potential regulation, with higher λ contributing to more isohydric behavior and α (d^{-1}) is a rate parameter controlling how quickly ψ_L adjusts to changes in ψ_s . As LPJ-GUESS-HYD runs on a daily timestep, α is set to 1 (Papastefanou et al., 2024). To account for summergreen phenology we expand upon Equation 12 to include the daily phenological status:

$$\hat{\psi}_l = \min\{\psi_l * \text{phen}, \psi_s\} \quad (13)$$

205 where phen is the leaf phenological status as a fraction of full leaf cover from 0 (no leaves) to 1 (full leaf cover). Subsequently, $\hat{\psi}_l$ equals ψ_s during winter dormancy. For evergreens, phen is always 1 and thus $\hat{\psi}_l = \psi_l$.

Next, we assume that the ~~xylem/stem-stem-xylem~~ water potential (ψ_x ; MPa) is a function of $\hat{\psi}_l$ following Fisher et al. (2006):

$$\psi_x = b * (\hat{\psi}_l - \psi_s) + \psi_s + \rho gh \quad (14)$$

210 where b represents the ratio of resistance belowground (R_{bg} ; $\text{m}^2 \text{ MPa s kg}^{-1}$) to total plant resistance (R_p ; $\text{m}^2 \text{ MPa s kg}^{-1}$):

$$b = \frac{R_{bg}}{R_p} \quad (15)$$

2.2.2 Water supply in LPJ-GUESS-HYD

LPJ-GUESS-HYD simulates the effect of hydraulic architecture on water transport through the plant by using
215 alternative formulations of E_{su} and E_{de} (from Equation 8).

The calculation of E_{su} is adopted from Hickler et al. (2006):

$$E_{su} = \frac{-\Delta\psi}{R_r + R_s + R_l} \quad (16)$$

where R_r , R_s , and R_l are the hydraulic resistances of roots, stem, and leaves in $\text{m}^2 \text{ MPa s kg}^{-1}$, respectively and are defined as:

$$220 \quad R_r = \frac{1}{k_{r,max} * (1 - \text{plc}_r) * \frac{a_r}{h_{soil}} * \eta_s} \quad (17)$$

$$R_s = \frac{1}{k_{s,max} * (1 - \text{plc}_s) * \frac{a_s}{h} * \eta_a} \quad (18)$$

and

$$R_l = \frac{1}{k_{l,max} * (1 - \text{plc}_l) * a_l * M_{H_2O}} \quad (19)$$

where $k_{r,max}$ ($\text{kg m}^{-1} \text{ s}^{-1} \text{ MPa}^{-1}$), $k_{s,max}$ ($\text{kg m}^{-1} \text{ s}^{-1} \text{ MPa}^{-1}$), and $k_{l,max}$ ($\text{mmol m}^{-2} \text{ s}^{-1} \text{ MPa}^{-1}$) are species-specific
225 parameters describing the maximum potential conductance of each compartment ([Table 1](#)), plc_r , plc_s , and plc_l are the fraction of cavitated vessels of each compartment, a_s , a_r and a_l are the cross-sectional area of sapwood, roots and leaves in $\text{m}^2 \text{ m}^{-2}$, respectively, η_a and η_n are the viscosity of water in the stem and soil, respectively, h (m) is the tree height, h_{soil} is the depth of the simulated soil column, fpc is the individual's foliar projective cover and M_{H_2O} is the molar mass of water (mol kg^{-1}).

230 The sum of resistances, denoted as R_p , represents the total plant hydraulic resistance:

$$R_p = R_r + R_s + R_l \quad (20)$$

2.2.3 Water demand in LPJ-GUESS-HYD

The updated representation of E_{de} is based on the instructive form of the Penman-Monteith equation [presented](#) [described](#) by Köstner et al. (1992) as:

$$235 \quad E_{de} = \Omega * E_q + (1 - \Omega) * E_{imp} \quad (21)$$

E_{imp} is the transpiration rate imposed by the effects of VPD, defined as:

$$E_{imp} = \frac{g_c * VPD}{\rho * G_v * T_{air}} \quad (22)$$

where G_v ($\text{m}^3 \text{kPa kg}^{-1} \text{K}^{-1}$) is the gas constant for water vapor and T_{air} (K) is the ambient air temperature. The term Ω is the degree of coupling between the canopy and the atmosphere (i.e. VPD) representing the leaf/canopy
240 boundary layer, defined as:

$$\Omega = \frac{1 + \varepsilon}{1 + \varepsilon + \frac{g_a}{g_c}} \quad (23)$$

where ε is the change of latent heat relative to the change in sensible heat in air at 10 °C and g_a (m s^{-1}) is the aerodynamic conductance. Consistent with the new formulations of E_{de} and E_{su} , the calculation of g_c is also updated. The assumption of the supply-demand principle underlying the original calculation of g_c remains but the new
245 definition reflects the dependence of plant water transport on VPD and hydraulic architecture. This is obtained by equating E_{su} (Eq. 16) and E_{imp} (Eq. 22) and solving for g_c resulting in:

$$g_c = \frac{\lambda_{lvh} * \frac{\gamma}{c_{p_{air}}} * \frac{\rho}{\rho_{air}} * \frac{\Delta\psi}{R_p}}{VPD} \quad (24)$$

where λ_{lvh} (kJ kg^{-1}) is the latent heat of vaporization of water, γ (kPa K^{-1}) is the psychrometric constant, $c_{p_{air}}$ ($\text{kJ kg}^{-1} \text{K}^{-1}$) is the specific heat of air, and ρ_{air} (kg m^{-3}) is the density of air.

250 Subsequently, when the canopy conductance constrained by plant hydraulic processes (Eq. 24) is less than the
non-stressed canopy conductance (Eq. 4) trees experience water limitation:

$$\text{water_limitation} = \begin{cases} \text{true,} & g_c < g_p \\ \text{false,} & g_c \geq g_p \end{cases} \quad (25)$$

In this case, g_c (Eq. 24) rather than g_p (Eq. 4) is used in the photosynthesis calculation.

Through this representation of water supply and demand the integrity of the plant's water transport system can
 255 directly affect the canopy conductance and, subsequently, carbon assimilation through photosynthesis.

2.2.4 Cavitation and mortality

The transport of water from the soil through the plant and into the atmosphere described by (Eqs. 9 - 24) is
 susceptible to partial or total collapse when soil water availability (Eq. 2) does not suffice to satisfy the transpiration
 demand (Eq. 21). During periods of water limitation when evapotranspiration outweighs water availability, soil water
 260 potential declines (Eq. 10). Modulated by the species-specific hydraulic strategy (λ , $\Delta\psi_{max}$), leaf and xylem water
 potential react, as well (Equation 12). As ψ_s , ψ_x , and ψ_l decrease, conductance through the tree (Eqs. 17 - 19)
 is attenuated through higher resistance, stemming from the onset of cavitation. Cavitation is represented as the
 percentage loss of conductance (plc) in dependence on ψ_x , modeled as a sigmoidal curve (cf. Tyree et al., 1994; Tyree
 and Sperry, 1989; Sperry et al., 1998; Pammenter and Van Der Willigen, 1998):

$$265 \quad plc = \frac{1}{\frac{\psi_x}{\psi_{50}}^{m_{cav}} + 1} \quad (26)$$

where ψ_{50} (MPa) and m_{cav} (MPa) are species-specific parameters indicating the xylem water potential at which 50 %
 of conductance is lost and the slope of the vulnerability curve, respectively (Table 1). The slope parameter, m_{cav} , is
 calculated as:

$$m_{cav} = \frac{2}{\log_{10}\left(\frac{\psi_{50}}{\psi_{88}}\right)} \quad (27)$$

270 where ψ_{88} (MPa) is the water potential at which 88 % of conductance is lost. To curb drought-induced cavitation
 during winter when processes related to hydraulic failure are assumed to play only a minor role, cavitation is
 only allowed to occur when g_c is greater than g_{min} , the component of canopy conductance not associated with
 photosynthesis. With rising plc the ability of plants to transport water is increasingly inhibited and eventually reaches

a point of no return at which the inability to move water becomes lethal (Hammond et al., 2019; Wagner et al., 2023). The probability of fatal hydraulic failure (p_{mort}) is modeled as a Weibull function following the results from Hammond et al. (2019):

$$p_{mort} = 1 - e^{-\left(\frac{plc}{k_w}\right)^{\lambda_w}}$$

where k_w is a shape parameter and λ_w is a scale parameter. As plc approaches 100 %, i.e. total hydraulic failure, the probability of mortality tends toward 1.

2.3 Global Sensitivity Analysis

The new processes integral to LPJ-GUESS-HYD introduce seven new input parameters. To ascertain how these additions contribute to uncertainty in the model output, we perform a global sensitivity analysis on the new parameters. LPJ-GUESS simulates a large number of outputs suited for sensitivity analysis. Similarly to Oberpriller et al. (2022), we examine carbon- and water- related outputs (evapotranspiration, canopy conductance, NPP, and biomass) due to the importance of forests in the carbon cycle both in governing fluxes and contributing to the carbon sink, and in the water cycle (Bonan, 2008; Pan et al., 2011; Pugh et al., 2019). We place a strong focus on water-related outputs due to the role of water use in modulating forest productivity, particularly under drought conditions (Lambers and Oliveira, 2019; Sulman et al., 2016). Sensitivities were calculated by sampling parameter sets from the multivariate parameter space using Latin Hypercube Sampling (LHS) (Helton and Davis, 2003; Mckay et al.). LHS is a sampling technique which stratifies a parameter into equal, non-repeating intervals across its entire range. By randomly sampling with each interval, LHS reduces bias and efficiently ensures full coverage of the parameter space. Compared to other sampling techniques (e.g. Quasi-Random Numbers) LHS requires fewer samples to depict the “true” mean of the parameter range. Consequently, fewer simulations must be run, substantially reducing the computational effort required when working with complex models such as LPJ-GUESS-HYD (Saltelli, 2008). For each of the seven parameters we estimated the potential parameter range based on previous studies using all values for species classified as trees in the corresponding data sources (Table 1).

Subsequently, we created 6000 parameter sets via LHS covering the entire multivariate parameter space. The parameter sets were recycled for each of the 12 species and 34 sites.

We chose Sobol’ indices to analyze the influence of parameter variations on the model output. This variance-based method can capture non-linear processes and is particularly suitable for non-additive models, i.e., models with interaction effects between the individual parameters such as the one (i.e. LPJ-GUESS-HYD) investigated here (Saltelli, 2008) Saltelli (2008). To calculate the sensitivity indices, LPJ-GUESS outputs needed to be condensed to a singular value per simulation (i.e. per parameter set). Flux variables (gross primary productivity (GPP), evapotranspiration, canopy conductance) were averaged over all years in the simulation period while the last year of

Parameter	Unit	Min	Max	Data reference	Definition
ψ_{50}	MPa	-14.20	-0.11	Choat et al. (2012)	Xylem pressure inducing 50% loss of conductance
m_{cav}	MPa	-69.25	-0.84	Choat et al. (2012)	Slope of vulnerability curve between ψ_{50} and ψ_{88}
$k_{r,max}$	$\text{kg m}^{-1} \text{s}^{-1} \text{MPa}^{-1}$	0.07	32.76	Choat et al. (2012)	Maximum specific root conductivity
$k_{s,max}$	$\text{kg m}^{-1} \text{s}^{-1} \text{MPa}^{-1}$	0.10	49.00	Choat et al. (2012)	Maximum specific stem conductivity
$k_{l,max}$	$\text{mmol m}^{-2} \text{s}^{-1} \text{MPa}^{-1}$	0.94	43.10	Multiple sources ¹	Maximum specific leaf conductivity
λ	-	-0.30	1.00	Papastefanou et al. (2020)	Isohyricity scalar
$\Delta\psi_{max}$	MPa	0.26	4.46	Papastefanou et al. (2020)	Forcing pressure under well watered conditions

Table 1. Definitions of the 7 new hydraulic parameters introduced in LPJ-GUESS-HYD and the parameter ranges used in the sensitivity analysis. These ranges extend beyond the observed values for the 12 species used in this study in order to explore the ~~models-model's~~ reaction to as wide of a parameter range as plausible. The data reference column indicates the source of the compiled ranges for each parameter. ¹Flexas et al. (2013); Méndez-Alonzo et al. (2019); Johnson et al. (2009); Scoffoni et al. (2011); Johnson et al. (2016); Nolf et al. (2015); Blackman et al. (2010)

the simulation was used for biomass. We calculated three sensitivity indices for each combination of output variable, species, and site. First and second order estimates were calculated using the estimator method introduced by (Saltelli et al., 2010). Total order indices were computed following the method by ~~(Jansen, 1999)~~[Jansen \(1999\)](#). First order indices measure the contribution of a single parameter to the variance in the model output excluding any interactions with other parameters. Similarly, second order indices measure the contribution of the interaction between two parameters to the variation in model output. Lastly, total order indices measure the contribution of a single parameter, including all its interactions with other parameters, to variation in the model output (Saltelli, 2008). In practical terms, these interactions refer to instances where separate parameters jointly affect a given model process or a given model output. For example, leaf water potential regulation in LPJ-GUESS-HYD (Eq. 12) is driven in part by both λ and $\Delta\psi_{max}$. In this case, the first order index for each parameter quantifies that parameter's individual contribution to Equation 12. The second-order index then quantifies the joint effect of the two parameters on Equation 11. This concept also extends beyond single, self-contained processes. That is, since, for example, both the water potential gradient between leaf and soil (governed by λ and $\Delta\psi_{max}$, Eq. 9, Eq. 11, Eq. 12) and the total plant resistance (governed by $k_{r,max}$, $k_{s,max}$, and $k_{l,max}$, Eq. 17)-19 affect canopy conductance the joint effect of any combination of these five parameters on canopy conductance can be quantified using either the second-order or total-order indices. The sensitivity indices range between 0 (least influential) and 1 (most influential) and depict the proportion of variance in the model output attributed to variations in a a given parameter or interaction of parameters. [By sampling the parameters independently of one another, i.e. allowing each parameter to vary independent of any other parameter in the same parameter set, we avoid collinearity biasing the sensitivity indices.](#) To establish significance we calculated sensitivity indices for a dummy parameter (i.e. a parameter that has relationship to the model). First and second order indices for our analyzed parameters were considered significant only if their value was higher than the indices

Species	Ψ_{50}	m_{cav}	$\Delta\psi_{ww}$	λ	$k_{r,max}$	$k_{s,max}$	$k_{l,max}$	Sites
<i>Abies alba</i>	-3.65	-10.7	0.4	0.4	0.86	0.38	33.1	4
<i>Betula pendula</i>	-2.23	-10.96	1.15	0.4	1.12	1.86	19.54	1
<i>Carpinus betulus</i>	-3.75	-13.75	0.89	0.07	1.8	2.7	19.54	2
<i>Fagus sylvatica</i>	-2.6	-9	1.47	-0.08	1.22	1.83	34.2	8
<i>Fraxinus excelsior</i>	-2.8	-7.95	0.78	0.45	0.47	0.7	8.88	1
<i>Picea abies</i>	-3.7	-12	1.15	0.4	0.29	0.43	33.1	17
<i>Pinus halapensis</i>	-3.57	-10.95	0.47	0.44	0.35	0.52	12.5	1
<i>Pinus sylvestris</i>	-3.14	-6.96	0.63	0.8	0.3	0.45	12.5	9
<i>Populus tremula</i>	-1.65	-6.67	0.86	0.53	0.61	0.92	25.39	1
<i>Quercus ilex</i>	-3.27	-4.77	1.14	0.16	1.3	1.95	7.95	5
<i>Quercus pubescens</i>	-2.475	-3.88	1.71	0.18	1.05	1.65	7.3	1
<i>Quercus robur</i>	-2.8	-9.45	1.6	0.075	2.05	2.34	9.9	9

Table 2. Best estimate species values for the 7 hydraulic parameters introduced in LPJ-GUESS-HYD used in the comparison of LPJ-GUESS-HYD with the eddy covariance flux variables. For each species, the used value is the mean of all values present for that species extracted from the relevant database (see Table 1). Where no observation for a given species was available, the genus mean was used instead.

for the dummy parameter. We used the `sensobol` R package to sample the 6000 parameter sets and compute the sensitivity indices (Puy et al., 2022).

2.4 Simulation Protocol and model evaluation

To test the functionality of LPJ-GUESS-HYD across a wide range of species we selected 12 common forest tree species from boreal, temperate, and ~~mediterranean~~-Mediterranean ecosystems and extracted the relevant parameters from available plant trait databases (Table 2). Using data from plant trait databases for parameterizing models can carry potential pitfalls due to various methods used in the original analyses contributing the data (Cochard et al., 2013). To account for this we ran an additional simulation using the same parameters as displayed in Table 2 but with ψ_{50} values from ~~?~~-Martin-StPaul et al. (2017) where such artifacts have been removed.

We chose sites to simulate from the ICOS Warm Winter 2020 ecosystem eddy covariance flux due to the availability of observational data for evaluation of the model at those sites (Warm Winter 2020 Team and ICOS Ecosystem Thematic Centre, 2022). We selected sites at which at least one of the 12 target species was present. This yielded 34 individual sites, each of which included a varying number of species, yielding a total of 55 unique species-site combinations. To avoid confounding effects brought on by competition between species each species at each site was simulated separately. For the sensitivity analysis we repeated the simulation of each species-site combination for all 6000 parameter sets. For evaluation of the model against the eddy-covariance flux data we used a set of best estimate

parameters compiled from published literature for each species ([Table 2](#)). The forcing data and general simulation procedure was the same for both sets of simulations.

The simulation period was from 1989 to 2020. To ensure a near-equilibrium state of the simulated ecosystem at the start of the simulation period we spun up the model for 1000 years by recycling the first 30 years of the climate inputs following standard procedure for LPJ-GUESS.

We forced both LPJ-GUESS and LPJ-GUESS-HYD with ERA-Interim daily mean surface temperature, precipitation sum, shortwave radiation, average windspeed, pressure, and specific humidity which were downscaled to the specific site coordinates and provided with the eddy-covariance flux data (Warm Winter 2020 Team and ICOS Ecosystem Thematic Centre, 2022; Pastorello et al., 2020). Atmospheric CO₂ concentration were taken from NOAA (Lan et al., 2023) and nitrogen deposition data were taken from Lamarque et al. (2011). Physical soil properties (e.g. clay-, sand-, and, silt-fraction, soil carbon content, bulk density, etc.) were taken from the Harmonized World Soil Database v2.0 and aggregated by mode to match the 0.5° by 0.5° spatial resolution of the climate inputs (Har, 2023).

From the ICOS Warm Winter 2020 dataset we extracted the daily GPP averaged from half-hourly data and partitioned via the night time partitioning method and daily evapotranspiration derived from observed latent heat flux (Allen et al., 1998) to evaluate simulated GPP and evapotranspiration against (Pastorello et al., 2020).

3 Results

3.1 Sensitivity analysis

Of the seven parameters introduced in LPJ-GUESS-HYD, only two (ψ_{50} and $\Delta\psi_{max}$) consistently contributed to variance across various model outputs (Figure 2). Carbon mass in vegetation was most sensitive to variations in ψ_{50} . Across all sites and species, the median contribution of ψ_{50} to variation in carbon mass in vegetation, including all interactions with other parameters, was 93.2% (Fig. 2a). Excluding any interactions with other parameters, 75% of the variance in carbon mass in vegetation was attributable solely to ψ_{50} (Fig. 3a). Considering all possible interactions, $\Delta\psi_{max}$ and $k_{l,max}$ were the second- (37.7%) and third-most (9%) influential parameters for carbon mass in vegetation, respectively. However, no substantial first-order influence of either $\Delta\psi_{max}$ or $k_{l,max}$ was found (Figure 3a). Generally, the analysis revealed similar patterns of total order sensitivity for GPP and evapotranspiration. In all cases, ψ_{50} contributed the most to variability in the output. Larger differences only manifested themselves in the sensitivity of canopy conductance. While canopy conductance only showed significant first-order sensitivity to ψ_{50} , it displayed a number of significant second-order sensitivities (Figure 3c). Additionally, all sensitivity indices (total, first, and second) displayed a larger spread across species and sites for canopy conductance than for any of the other variables (Figure 2d; Figure 3d). Importantly, while the sensitivity indices for psi_{50} by far outweighed

those of the other parameters for GPP, evapotranspiration and vegetation carbon, the relative sensitivity of canopy conductance to psi_{50} compared to the other parameters was more balanced.

Although the total-order indices indicated that m_{cav} contributed only marginally to output variance, the first-order indices revealed that m_{cav} on its own did, in fact, lead to significant albeit low variance in all model outputs (Figure 3). For all considered output variables, second-order interactions consistently included ψ_{50} and $\Delta\psi_{max}$ (Figure 3), while only two other parameters, $k_{l,max}$ and $k_{r,max}$, occasionally featured in the second-order indices (Figure 3).

3.2 Evapotranspiration response to VPD

In LPJ-GUESS-HYD, evapotranspiration patterns of individual species were largely governed by the species-specific response to VPD (Figure 4, B). With increasing VPD classes, i.e. higher atmospheric demand for water, the spread of evapotranspiration patterns between species increased. While more isohydric species (e.g. *Pinus sylvestris*, *Abies alba*, *Populus tremuloides*) only marginally increased their evapotranspiration rate under higher VPD, more anisohydric species (e.g. *Fagus sylvatica*, *Quercus spec.*) tended to increase their evapotranspiration rates under higher VPD. In contrast, in LPJ-GUESS (Figure 4, C), although some species-specific differences in evapotranspiration rate were simulated, the general VPD response pattern was the same across all species; evapotranspiration increased with increasing VPD up to ~ 1.5 kPa and subsequently leveled off even as VPD continued to increase (Figure 4, right panel). Additionally, no clear pattern related to isohydricity was seen in LPJ-GUESS. Under high VPD the highest evapotranspiration rate was seen in an ostensibly more isohydric species, *Pinus sylvestris*, while the second highest rate was exhibited by *Quercus pubescens*, a relatively anishoydric species. Monospecific eddy-covariance flux sites were only available for a limited number of species (Figure 4, A). Here, more anisohydric species tended to continue transpiring even as VPD increased while more isohydric species reached maximum transpiration rates at relatively low levels of VPD and displayed decreasing evapotranspiration as VPD continued to increase. Under high VPD (~ 3 kPa) evapotranspiration simulated by LPJ-GUESS-HYD ranged from 0.9 mm d^{-1} to 7 mm d^{-1} . The range in LPJ-GUESS was considerably smaller ranging from 1.4 mm d^{-1} to 3.2 mm d^{-1} . For the eddy-covariance flux data observations at a VPD of ~ 3 kPa were only available for *Fagus sylvatica* which transpired 5.8 mm d^{-1} at that VPD level.

3.3 Comparison of model results with observational data from eddy-covariance towers

The comparison of evapotranspiration simulated by LPJ-GUESS(-HYD) with evapotranspiration from the eddy covariance flux product in three pan-european drought years revealed contrasting results (Figure 5). Across all sites, species, and all drought years, the observed ~~monthly~~-daily growing season evapotranspiration ranged from ~~$\sim 37 \text{ mm month}^{-1}$ to $\sim 106 \text{ mm month}^{-1}$~~ $\sim 1.20 \text{ mm d}^{-1}$ to $\sim 3.54 \text{ mm d}^{-1}$. LPJ-GUESS-HYD simulated a similar range (~~$\sim 33 \text{ mm month}^{-1}$ to $\sim 136 \text{ mm month}^{-1}$~~ $\sim 1.14 \text{ mm d}^{-1}$ to $\sim 4.45 \text{ mm d}^{-1}$) while LPJ-GUESS simulated a narrower range (~~$\sim 45 \text{ mm month}^{-1}$ to $\sim 71 \text{ mm month}^{-1}$~~ $\sim 1.50 \text{ mm d}^{-1}$ to $\sim 2.33 \text{ mm d}^{-1}$). Compared to the eddy covariance product both LPJ-GUESS and LPJ-GUESS-HYD displayed




Figure 2. Total order sensitivity indices for the seven parameters introduced in LPJ-GUESS-HYD. Total-order indices indicate the sensitivity of model output to variation of a given parameter including any and all interactions with other parameters. Each point represents the sensitivity index for a single species-site combination. The boxplots indicate the median and interquartile range of the sensitivity indices across species-site combinations. Each panel shows the sensitivity indices for a single model output (A) mean annual canopy conductance , (B) mean annual evapotranspiration, (C) mean annual gross primary productivity, (D) carbon mass in vegetation.




Figure 3. First and second order sensitivity indices for the seven parameters introduced in LPJ-GUESS-HYD. First-order indices indicate the sensitivity of model output solely due to variations of a single parameter. Second-order indices only consider variation in the output attributable to interactions between two parameters. First- and second-order indices are only shown for parameters with a median sensitivity greater than the median sensitivity of a dummy parameter (see methods for details). Each point represents the sensitivity index for a single species-site combination. The boxplots indicate the median and interquartile range of the sensitivity indices across species-site combinations. Each panel shows the sensitivity indices for a single model output (A) mean annual canopy conductance, (B) mean annual evapotranspiration, (C) mean annual gross primary productivity, (D) carbon mass in vegetation.




Figure 4. Species-specific daily evapotranspiration rates under differing levels of vapor pressure deficit from A) eddy-covariance flux towers, B) LPJ-GUESS-HYD, and C) standard LPJ-GUESS. The colors are ranked according to the λ of each species (see Fig. 2) from high λ (light) to low λ (dark). Daily VPD was binned into 6 equally-sized classes representing increasing levels of drought. Species-specific responses to drought remain constant in LPJ-GUESS while clear differences between more anisohydric and more isohydric species are seen in LPJ-GUESS-HYD.

a similar level of mismatch with RMSEs of ~~22.2 and 26.9 mm month~~0.70 and 0.84 mm d⁻¹, respectively. However,
405 while LPJ-GUESS consistently underestimated the observed evapotranspiration (Mean signed deviation (MSD):
~~-15.6~~-0.44), LPJ-GUESS-HYD showed a less negative bias (MSD: ~~-9.3~~-0.06). For GPP both LPJ-GUESS and LPJ-
GUESS-HYD show similar patterns broadly matching the observations. The RMSE for GPP was ~~0.06~~similarly low
for both LPJ-GUESS and LPJ-GUESS-HYD at 0.0017 and 0.0021, respectively. For both model versions the MSD
indicated no substantial over- or underestimation of the observations (LPJ-GUESS: ~~-0.006~~-0.0003; LPJ-GUESS-HYD:
410 ~~-0.02~~-0.0007)




Figure 5. Comparison of measured eddy-covariance flux values of ET and GPP with modeled values from LPJ-GUESS-HYD (left column) and LPJ-GUESS (right column). LPJ-GUESS-HYD (A) matches observed ET patterns better than standard LPJ-GUESS (B) during three pan-European ~~droughts~~ drought years while simulated GPP remains similar between both versions of the model (eC, dD). The dotted black line indicates perfect agreement between model and observations. Values above the dotted line represent instances where the model overestimates ET (GPP) compared to the observations and vice versa.

4 Discussion

We conducted an evaluation of the newly developed plant hydraulic architecture version of LPJ-GUESS, LPJ-GUESS-HYD, through a variance-based global sensitivity analysis and model evaluation for carbon and water fluxes at 34 eddy-covariance flux sites across Europe.

4.1 Relevance of hydraulic parameters

The results of our sensitivity analysis showed that of the seven newly introduced parameters (Table 1), two (ψ_{50} , $\Delta\psi_{max}$) consistently contributed substantially to the variance in model outputs either directly (Figure 3) or indirectly (Figure 2). Similarly, second-order interactions for all outputs included primarily those aforementioned parameters. Substantial differences in parameter importance were only seen for mean annual canopy conductance. Although, the two previously mentioned parameters still contributed the most to variance in simulated canopy conductance, nearly all other parameters played a substantial role as well (Figure 2, panel A). Additionally, across all sites and species, the sensitivity indices varied to a greater extent in the case of canopy conductance than in the other outputs (Figure 3, Figure 2). ~~Perhaps unsurprisingly, this~~ This pattern suggests that ~~model processes which are more the relative influence of the new parameters is most evenly spread in model processes~~ closely related (e.g., canopy conductance; see Figure 1) to the newly implemented plant hydraulic architecture ~~are more sensitive to the same factors affecting those processes~~. That is, while processes like carbon allocation to biomass which are further downstream from the new implementations are primarily affected by a single parameter (ψ_{50}), processes like canopy conductance which are directly affected by the new implementations are more sensitive to a greater number of the newly implemented parameters because the influence of these parameters is less diluted by other contributing model processes (e.g. plant demography) as is the case with the carbon allocation.

Strikingly, LPJ-GUESS-HYD output was by far most sensitive to variations in ψ_{50} , with roughly 75% of the variance in ET, GPP, and vegetation carbon being attributable to changes ψ_{50} alone (Figure 3). While not directly comparable, this aligns both with a previous meta-analysis which suggested that ψ_{50} was the single most effective predictor of tree drought mortality (Anderegg et al., 2016) and previous modeling efforts which indicated ψ_{50} to substantially influence modeled xylem embolism Cochard et al. (2021). The meta-analysis additionally indicated that $k_{s,max}$ played little role in determining tree mortality due to drought (Anderegg et al., 2016), a result partially supported by our sensitivity analysis that showed $k_{s,max}$ has negligible influence on drought mortality. Although our analysis focused on water and carbon fluxes rather than outright mortality, these findings complement each other as they suggest that the traits that are responsible for impairing water transport and assimilation under drought stress are the same traits that ultimately determine whether a tree will experience drought damage or eventually die under prolonged drought.

The model's strong sensitivity to the maximum possible soil-to-leaf water-potential difference, $\Delta\psi_{max}$, is less intuitive. Along with the conductivity of roots, stem, and leaves, the soil-to-leaf water-potential difference, also referred to

as the forcing pressure, plays a role in regulating the supply of water through the tree (Joshi et al., 2022; Da Sois et al., 2024). Why, then, does the model sensitivity to $\Delta\psi_{max}$ overshadow the sensitivity to the parameters which govern conductivity, namely, $k_{r,max}$, $k_{s,max}$, and $k_{l,max}$? This divergent response can be explained by the relationship of $\Delta\psi_{max}$ and ψ_{50} in LPJ-GUESS-HYD. Primarily, $\Delta\psi_{max}$ determines how tightly (or loosely) simulated leaf water potential is coupled to simulated soil water potential (Equation 12), affecting the degree of isohydricity. At a given soil water potential, species with a higher $\Delta\psi_{max}$ (i.e. looser coupling) will have a lower leaf water potential than species with a lower $\Delta\psi_{max}$ (i.e. stronger coupling). Due to the relationship between leaf water potential and xylem water potential in LPJ-GUESS-HYD (Equation 14), this means that the value of $\Delta\psi_{max}$, which influences leaf water potential (Equation 12), indirectly determines the xylem water potential and therefore affects the process of xylem cavitation. This is backed up by the significant second-order interactions between ψ_{50} and $\Delta\psi_{max}$ (Figure 3). As McDowell et al. (2008) point out, the soil-to-leaf water potential difference, $\Delta\psi$, tends to increase with increasing transpiration until a critical xylem tension is reached leading to cavitation and, consequently, the hydraulic conductance approaches zero. It follows that as actual hydraulic conductance approaches zero, the maximum possible hydraulic conductance specified by $k_{r,max}$, $k_{s,max}$, and $k_{l,max}$ loses relevance. Indeed, this finding matches existing evidence from model sensitivity analyses indicating that parameters related to xylem safety and stomatal regulation explained a substantial fraction of the model variability, while whole plant conductance (i.e. $k_{r,max}$, $k_{s,max}$, and $k_{l,max}$ in our study) played a lesser role ~~Ruffault et al. (2022)~~([Ruffault et al., 2022](#)).

To reiterate, the results of the sensitivity analysis indicate that two of the hydraulic parameters introduced in LPJ-GUESS-HYD, namely $\Delta\psi_{max}$ and ψ_{50} substantially shape long-term model behavior. These results imply that accurate estimations or, in the best case, measurements of these two parameters are paramount to reliably modeling plant hydraulics with LPJ-GUESS-HYD. Indeed, although using ~~alternative the, arguably better,~~ parameterizations for ψ_{50} [from Martin-StPaul et al. \(2017\)](#) (Figure A2) did not alter the general pattern that modeled evapotranspiration from LPJ-GUESS-HYD reflects the anisohydric-isohydric continuum, the evapotranspiration response to VPD of individual species was affected by this alternative parameterization (e.g. *Quercus ilex*; A2, panel B). What our sensitivity analysis cannot provide answers to, however, is how model sensitivity may change under stressed vs. non-stressed conditions. That is, does the pattern of influential parameters remain the same during drought as during non-drought conditions? To answer this, subsequent modeling endeavors specifically contrasting various climatic conditions are required.

4.2 Role of hydraulic architecture for carbon and water fluxes

The results of our sensitivity analysis show that simulated water and carbon fluxes from LPJ-GUESS-HYD are primarily influenced by hydraulic function – via ψ_{50} – and secondarily by stomatal regulation – via $\Delta\psi_{max}$. These results are largely in line with findings from experiments and observations that repeatedly and consistently identify

475 hydraulic failure as the preeminent factor governing tree drought mortality (Anderegg et al., 2016, 2015; Choat et al., 2012; Hammond et al., 2019; Adams et al., 2017).

However, the importance of $\Delta\psi_{max}$ in our model analysis also aligns with the ample evidence that stomatal regulation is critical in mediating drought responses of forests (Körner, 2019; Hajek et al., 2022; Mcdowell et al., 2008). The sensitivity of LPJ-GUESS-HYD to these widely supported mechanisms of tree drought response suggests that
480 LPJ-GUESS-HYD should be able to correctly simulate drought and its associated impacts across a range of different species and hydraulic strategies.

To demonstrate the ability of LPJ-GUESS-HYD to model drought responses across hydraulic strategies, we analyzed the effect of increasing VPD on simulated evapotranspiration in both LPJ-GUESS-HYD and standard LPJ-GUESS (Figure 4). This analysis effectively showed that while LPJ-GUESS displayed nearly identical VPD response trajectories
485 across all species, LPJ-GUESS-HYD exhibits distinct trajectories. This can be explained for one by the absence of VPD as a direct driver of evapotranspiration in standard LPJ-GUESS. Yet, it also shows the importance of the inclusion of dynamic stomatal regulation strategies as exhibited by the larger range in simulated evapotranspiration rates in LPJ-GUESS-HYD. More anisohydric species (i.e. lower λ_{iso} , higher $\Delta\psi_{max}$, see Table 2) tended to keep transpiring even under high VPD while more isohydric species displayed plateauing evapotranspiration as VPD increased. Our
490 simulations revealed no distinct clustering of evapotranspiration responses to VPD, but rather a gradation of responses dependent on the relevant parameters. This simulated behavior is congruent with the established notion of the anisohydric-isohydric continuum (Klein, 2014; Martínez-Vilalta et al., 2014; Martínez-Vilalta and Garcia-Forner, 2017). Similarly, the species-specific responses of evapotranspiration to VPD simulated by LPJ-GUESS-HYD reflect results from experiments identifying VPD as the most potent driver of both canopy conductance and evapotranspiration
495 (Schönbeck et al., 2022; Flo et al., 2022). In particular, the order of the evapotranspiration-VPD response simulated by LPJ-GUESS-HYD (Figure 4) for *Fagus sylvatica*, *Quercus pubescens*, and *Quercus ilex* are comparable to the results from Schönbeck et al. (2022).

Lastly, to evaluate the efficacy of LPJ-GUESS-HYD at simulating the real world response of water and carbon fluxes to drought we compared simulated evapotranspiration and GPP with eddy-covariance fluxes from 34 sites across Europe
500 during three pan-European drought years – 2003, 2015, and 2018 (Figure 5). Compared to LPJ-GUESS, LPJ-GUESS-HYD represents an improvement in terms of simulated evapotranspiration under drought. Since eddy-covariance flux data integrates the response of all species at a given site, our ability to conduct species-specific comparisons of modeled and observed evapotranspiration was limited. Nevertheless, the limited available data suggest that LPJ-GUESS-HYD is better at capturing the observed evapotranspiration patterns of more anisohydric species than of relatively isohydric
505 species (Figure A1). This may also partially explain the underestimation of evapotranspiration by LPJ-GUESS-HYD seen at some sites (Figure 5, A), yet the limited availability of species-specific comparisons does not allow for a conclusive explanation. In any case, this indication together with the fact that $\Delta\psi_{max}$, which largely governs modeled

stomatal regulation, was one of the most influential parameters, suggests that well-constrained estimates of $\Delta\psi_{max}$ are crucial for model performance.

510 Contrastingly, no meaningful difference was seen between LPJ-GUESS and LPJ-GUESS-HYD for simulated GPP under drought. Considering that the sensitivity analysis revealed that modeled GPP is sensitive to variations in ψ_{50} , the lack of differences between LPJ-GUESS-HYD and standard LPJ-GUESS may seem surprising. Yet, these results must be interpreted carefully. The control of ψ_{50} on GPP in the sensitivity analysis stems from the fact that with high values of ψ_{50} (i.e. low resistance to embolism) few viable parameter combinations remain, that is,
515 ψ_{50} represents a limiting factor which can override the effect of the other parameters. In the evaluation using the best estimate parameter sets (Table 2) the values of ψ_{50} remain within a viable range. Additionally, despite lacking a mechanistic representation of photosynthetic response to drought the empirical relationships of photosynthesis to low water availability implemented in LPJ-GUESS – and, in fact, in a host of other DVMs – are rooted in reality and have been shown to be sufficient in reproducing past droughts and their effect on carbon uptake (Ciais et al.,
520 2005; van der Woude et al., 2023; Gampe et al., 2021). Yet, the improved representation of evapotranspiration (based explicitly on canopy conductance) in LPJ-GUESS-HYD paves the way for implementing further hydraulic processes, such as capacitance, and improving existing ones, such as cavitation. Such advancements, coupled with sink-driven mechanisms (e.g. turgor-limited growth), are paramount to modeling carbon and water cycles in future climates where existing empirical relationships become less dependable (Körner, 2015; Torres-Ruiz et al., 2024).

525 4.3 Limitations of the modeling approach and ways forward

Despite the improvements in modeling plant-water relations offered by LPJ-GUESS-HYD, further improvements will be necessary in subsequent iterations of the model. Considering the hydraulic processes implemented in LPJ-GUESS-HYD (Figure 1), it is obvious that in the current state they are directed towards the water rather than the carbon cycle. As such, the path forward for LPJ-GUESS-HYD must focus on physiological processes connecting
530 plant water-usage with plant carbon-usage, both in terms of carbon assimilation and carbon losses. One major source of carbon loss due to drought is tree mortality (Allen et al., 2010). In the current version of LPJ-GUESS-HYD, drought mortality is implemented based on xylem cavitation but not on the downstream ramifications of hydraulic failure (e.g. higher susceptibility to insects and other biotic agents) although these are generally considered to be significant secondary-drivers in drought induced mortality (Senf et al., 2020; Desprez-Loustau et al., 2006; Rouault
535 et al., 2006; Bigler et al., 2006; Anderegg et al., 2015). Linking existing models dealing with biotic and non-biotic disturbance agents (Lagergren et al., 2012; Jönsson et al., 2012) to LPJ-GUESS-HYD could provide a pathway to better capture observed mortality associated with droughts. In this context, emphasis must be placed on mechanisms governing how drought stress increases vulnerability to these secondary processes. However, carbon losses due to drought are not confined only to tree mortality. Across the globe, an increase in drought-induced tree canopy dieback
540 has been observed (Allen et al., 2010, 2015; Lloret et al., 2004; Frei et al., 2022; Carnicer et al., 2011; Hartmann et al.,

2022). Evidence suggests that such dieback is caused primarily by hydraulic failure (Arend et al., 2022; Kannenberg et al., 2021; Walthert et al., 2021; Nolan et al., 2021), although a disruption of the soil-root interface (Körner, 2019; Carminati and Javaux, 2020) and preceding growth trends (Neycken et al., 2022) have been identified as potential drivers as well. Regardless of the underlying cause, crown dieback reduces the leaf area, altering canopy water demand and growth even once the drought has subsided (Arend et al., 2022; Guada et al., 2016). While early leaf senescence in response to drought has been widely observed in beech and other temperate broad-leaved species (Schuldt et al., 2020 and references therein), evidence suggests that coniferous species, such as spruce, may die from hydraulic failure before such protective measures can occur (Arend et al., 2021). Additionally, the relationship between drought intensity, hydraulic failure and early leaf senescence is difficult to quantify and studies establishing concrete thresholds for leaf senescence are scarce and focused on single species (~~Walthert et al., 2021, e.g.~~) (e.g. [Walthert et al., 2021](#)). Nevertheless, early leaf senescence plays an important role in governing tree drought response (Nadal-Sala et al., 2024). Yet, currently LPJ-GUESS(-HYD) neither includes any mechanistic nor empirical representation of this process. While the exact mechanisms may be too detailed for a model such as LPJ-GUESS-HYD, some relationship between hydraulic failure and reduced leaf area should be a part of future developments to ensure that the actual leaf area matches that which is able to be supported by the diminished sapwood area due to xylem cavitation.

Additionally, a better representation of drought-associated carbon losses (e.g. mortality, dieback, lost productivity, etc.) is only part of the puzzle. Most DVMs, including LPJ-GUESS-HYD, primarily model carbon allocation and tree growth as source-limited (Cabon and Anderegg, 2022; Eckes-Shephard et al., 2021). In LPJ-GUESS-HYD reduced carbon uptake under drought follows this pattern. As stomates close and gas exchange is reduced photosynthetic assimilation slows as well. Yet, emerging evidence emphasizes the importance of including sink limitations in models as they a crucial factor in modulating tree growth, particularly during drought as cambial cell formation is limited by turgor (Körner, 2015; Peters et al., 2021; Cabon et al., 2020). While mechanistic turgor-driven growth models exist (Steppe et al., 2006; Génard et al., 2001; Peters et al., 2021) they are too complex, both temporally and physiologically, for direct implementation into LPJ-GUESS-HYD (Potkay et al., 2022). To bridge this gap, including plant water storage and hydraulic capacitance could be a starting point for a simple approximation of the more complex process underlying turgor-driven growth limitations. Observations from dendrometers suggest that little to no growth occurs during periods of stem shrinkage, i.e., when plant water storage recedes (Zweifel et al., 2016). In contrast to dedicated turgor-driven growth models, the dynamics of plant water storage more easily lend themselves to implementation in DVMs and could nonetheless present a viable proxy for more complex sink-limitations under drought.

Lastly, if and when further developments to LPJ-GUESS-HYD are made subsequent sensitivity analyses should be conducted. In this study, the sensitivity analysis focused on long-term model outputs such as annual water and carbon fluxes which are also commonly used for benchmarking DVMs (~~Seiler et al., 2022; Collier et al., 2018, e.g.~~) (e.g. [Seiler et al., 2022; Collier et al., 2018](#)). However, as future developments discussed above (e.g. turgor-driven growth, drought-induced leaf-shedding) will likely focus on specific aspects of tree drought response, sensitivity analyses

575 on finer temporal scales may be both more practical and useful than the larger yet coarser sensitivity analysis used here. To this end, future analyses could also consider not only relying on direct model output variables but creating specific indices or metrics related to individual model processes (~~Ruffault et al., 2022, e.g.~~)(e.g. Ruffault et al., 2022)

5 Conclusion

580 In this study, we evaluated LPJ-GUESS-HYD for use with European tree species along an isohydricity gradient. The model was shown to simulate species-specific responses of evapotranspiration to increasing VPD in accordance with both results from experiments and current understanding of the anisohydric-isohydric continuum. A comparison of simulated ET and GPP with observations from eddy-covariance flux sites in three pan-European drought years (2003, 2015, 2018) revealed that LPJ-GUESS-HYD improved evapotranspiration compared to the standard version
585 of LPJ-GUESS although both versions of the model displayed a similar fit of simulated to observed GPP. These results not only emphasize the importance of including mechanistic representations of plant hydraulic architecture in dynamic vegetation models but also highlight the fact that simulating both water- and carbon fluxes based on canopy conductance provides improvements in model performance compared to only using canopy conductance for the calculation of carbon fluxes. In this context, future developments of LPJ-GUESS-HYD should continue to
590 focus on the connection between plant water-use and plant carbon-use, potentially under the aspect of sink-limited growth. Plant hydraulics are a crucial extension of current DVMs for modeling the effect of drought on altering ecosystem scale water-usage and continued refinements may be essential in providing robust estimates of future drought responses under a changing climate.

Code and data availability. LPJ-GUESS is publicly available at: <https://doi.org/10.5281/zenodo.8065736>. The version of
595 LPJ-GUESS used in this study is publicly available at: <https://doi.org/10.5281/zenodo.14000805>. The model version presented here is identified by the commit hash 97c552c5. The analysis code used to produce the results and figures of this study is available at: <https://doi.org/10.5281/zenodo.14001089>

Author contributions. BM, AR, and CZ conceived the study. BM wrote the manuscript, conducted the model simulation runs, and conducted the data analysis. JD gathered and prepared input data for the model runs, contributed model code, and
600 contributed to the draft writing. PP and KG contributed to model development. AB, QG, TG, and AK contributed to the interpretation of the results. DL and SA gathered an prepared parameter values for use in the sensitivity analysis. All authors edited the manuscript.

Competing interests. The authors declare no competing interests

References

- 605 Harmonized World Soil Database Version 2.0, FAO; International Institute for Applied Systems Analysis (IIASA);, <https://doi.org/10.4060/cc3823en>, 2023.
- Adams, H. D., Zeppel, M. J. B., Anderegg, W. R. L., Hartmann, H., Landhäusser, S. M., Tissue, D. T., Huxman, T. E., Hudson, P. J., Franz, T. E., Allen, C. D., Anderegg, L. D. L., Barron-Gafford, G. A., Beerling, D. J., Breshears, D. D., Brodrigg, T. J., Bugmann, H., Cobb, R. C., Collins, A. D., Dickman, L. T., Duan, H., Ewers, B. E., Galiano, L., Galvez, D. A., Garcia-Forner, N., Gaylord, M. L., Germino, M. J., Gessler, A., Hacke, U. G., Hakamada, R., Hector, A., Jenkins, M. W., Kane, J. M., Kolb, T. E., Law, D. J., Lewis, J. D., Limousin, J.-M., Love, D. M., Macalady, A. K., Martínez-Vilalta, J., Mencuccini, M., Mitchell, P. J., Muss, J. D., O'Brien, M. J., O'Grady, A. P., Pangle, R. E., Pinkard, E. A., Piper, F. I., Plaut, J. A., Pockman, W. T., Quirk, J., Reinhardt, K., Ripullone, F., Ryan, M. G., Sala, A., Sevanto, S., Sperry, J. S., Vargas, R., Vennetier, M., Way, D. A., Xu, C., Yezpe, E. A., and McDowell, N. G.: A Multi-Species Synthesis of Physiological Mechanisms in Drought-Induced Tree Mortality, *Nature Ecology & Evolution*, 1, 1285–1291, <https://doi.org/10.1038/s41559-017-0248-x>, 2017.
- 615 Ahlström, A., Schurgers, G., Arneth, A., and Smith, B.: Robustness and Uncertainty in Terrestrial Ecosystem Carbon Response to CMIP5 Climate Change Projections, *Environmental Research Letters*, 7, 044 008, <https://doi.org/10.1088/1748-9326/7/4/044008>, 2012.
- 620 Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Kitzberger, T., Rigling, A., Breshears, D. D., Hogg, E. H. T., Gonzalez, P., Fensham, R., Zhang, Z., Castro, J., Demidova, N., Lim, J.-H., Allard, G., Running, S. W., Semerci, A., and Cobb, N.: A Global Overview of Drought and Heat-Induced Tree Mortality Reveals Emerging Climate Change Risks for Forests, *Forest Ecology and Management*, 259, 660–684, <https://doi.org/10.1016/j.foreco.2009.09.001>, 2010.
- 625 Allen, C. D., Breshears, D. D., and McDowell, N. G.: On Underestimation of Global Vulnerability to Tree Mortality and Forest Die-off from Hotter Drought in the Anthropocene, *Ecosphere*, 6, art129, <https://doi.org/10.1890/ES15-00203.1>, 2015.
- Allen, R. G., Pereira, L. S., Raes, D., and Smith, M.: Crop Evapotranspiration - Guidelines for Computing Crop Water Requirements, FAO Irrigation and Drainage Paper 56, Food and Agriculture Organization of the United Nations, 1998.
- Anderegg, W. R. L., Berry, J. A., Smith, D. D., Sperry, J. S., Anderegg, L. D. L., and Field, C. B.: The Roles of Hydraulic and Carbon Stress in a Widespread Climate-Induced Forest Die-Off, *Proceedings of the National Academy of Sciences*, 109, 233–237, <https://doi.org/10.1073/pnas.1107891109>, 2012.
- 630 Anderegg, W. R. L., Hicke, J. A., Fisher, R. A., Allen, C. D., Aukema, J., Bentz, B., Hood, S., Lichstein, J. W., Macalady, A. K., McDowell, N., Pan, Y., Raffa, K., Sala, A., Shaw, J. D., Stephenson, N. L., Tague, C., and Zeppel, M.: Tree Mortality from Drought, Insects, and Their Interactions in a Changing Climate, *New Phytologist*, 208, 674–683, <https://doi.org/10.1111/nph.13477>, 2015.
- 635 Anderegg, W. R. L., Klein, T., Bartlett, M., Sack, L., Pellegrini, A. F. A., Choat, B., and Jansen, S.: Meta-Analysis Reveals That Hydraulic Traits Explain Cross-Species Patterns of Drought-Induced Tree Mortality across the Globe, *Proceedings of the National Academy of Sciences*, 113, 5024–5029, <https://doi.org/10.1073/pnas.1525678113>, 2016.
- Arend, M., Link, R. M., Patthey, R., Hoch, G., Schuldt, B., and Kahmen, A.: Rapid Hydraulic Collapse as Cause of Drought-Induced Mortality in Conifers, *Proceedings of the National Academy of Sciences*, 118, e2025251118, <https://doi.org/10.1073/pnas.2025251118>, 2021.
- 640

- Arend, M., Link, R. M., Zahnd, C., Hoch, G., Schuldt, B., and Kahmen, A.: Lack of Hydraulic Recovery as a Cause of Post-drought Foliage Reduction and Canopy Decline in European Beech, *New Phytologist*, 234, 1195–1205, <https://doi.org/10.1111/nph.18065>, 2022.
- 645 Bigler, C. and Vitasse, Y.: Premature Leaf Discoloration of European Deciduous Trees Is Caused by Drought and Heat in Late Spring and Cold Spells in Early Fall, *Agricultural and Forest Meteorology*, 307, 108492, <https://doi.org/10.1016/j.agrformet.2021.108492>, 2021.
- Bigler, C., Bräker, O. U., Bugmann, H., Dobbertin, M., and Rigling, A.: Drought as an Inciting Mortality Factor in Scots Pine Stands of the Valais, Switzerland, *Ecosystems*, 9, 330–343, <https://doi.org/10.1007/s10021-005-0126-2>, 2006.
- 650 Blackman, C. J., Brodribb, T. J., and Jordan, G. J.: Leaf Hydraulic Vulnerability Is Related to Conduit Dimensions and Drought Resistance across a Diverse Range of Woody Angiosperms, *New Phytologist*, 188, 1113–1123, <https://doi.org/10.1111/j.1469-8137.2010.03439.x>, 2010.
- Bonan, G. B.: Forests and Climate Change: Forcings, Feedbacks, and the Climate Benefits of Forests, *Science*, 320, 1444–1449, <https://doi.org/10.1126/science.1155121>, 2008.
- 655 Brodribb, T. J., Powers, J., Cochard, H., and Choat, B.: Hanging by a Thread? Forests and Drought, *Science*, 368, 261–266, <https://doi.org/10.1126/science.aat7631>, 2020.
- Buras, A., Meyer, B., and Rammig, A.: Record Reduction in European Forest Canopy Greenness during the 2022 Drought, Tech. Rep. EGU23-8927, Copernicus Meetings, <https://doi.org/10.5194/egusphere-egu23-8927>, 2023.
- Cabon, A. and Anderegg, W. R. L.: Turgor-Driven Tree Growth: Scaling-up Sink Limitations from the Cell to the Forest, *Tree Physiology*, 42, 225–228, <https://doi.org/10.1093/treephys/tpab146>, 2022.
- 660 Cabon, A., Peters, R. L., Fonti, P., Martínez-Vilalta, J., and De Cáceres, M.: Temperature and Water Potential Co-Limit Stem Cambial Activity along a Steep Elevational Gradient, *New Phytologist*, 226, 1325–1340, <https://doi.org/10.1111/nph.16456>, 2020.
- Carminati, A. and Javaux, M.: Soil Rather Than Xylem Vulnerability Controls Stomatal Response to Drought, *Trends in Plant Science*, 25, 868–880, <https://doi.org/10.1016/j.tplants.2020.04.003>, 2020.
- 665 Carnicer, J., Coll, M., Ninyerola, M., Pons, X., Sánchez, G., and Peñuelas, J.: Widespread Crown Condition Decline, Food Web Disruption, and Amplified Tree Mortality with Increased Climate Change-Type Drought, *Proceedings of the National Academy of Sciences*, 108, 1474–1478, <https://doi.org/10.1073/pnas.1010070108>, 2011.
- Choat, B., Jansen, S., Brodribb, T. J., Cochard, H., Delzon, S., Bhaskar, R., Bucci, S. J., Feild, T. S., Gleason, S. M., Hacke, U. G., Jacobsen, A. L., Lens, F., Maherali, H., Martínez-Vilalta, J., Mayr, S., Mencuccini, M., Mitchell, P. J., Nardini, A., Pittermann, J., Pratt, R. B., Sperry, J. S., Westoby, M., Wright, I. J., and Zanne, A. E.: Global Convergence in the Vulnerability of Forests to Drought, *Nature*, 491, 752–755, <https://doi.org/10.1038/nature11688>, 2012.
- Christoffersen, B. O., Gloor, M., Fauset, S., Fyllas, N. M., Galbraith, D. R., Baker, T. R., Kruijt, B., Rowland, L., Fisher, R. A., Binks, O. J., Sevanto, S., Xu, C., Jansen, S., Choat, B., Mencuccini, M., McDowell, N. G., and Meir, P.: Linking
- 675 Hydraulic Traits to Tropical Forest Function in a Size-Structured and Trait-Driven Model (TFS v.1-Hydro), *Geoscientific Model Development*, 9, 4227–4255, <https://doi.org/10.5194/gmd-9-4227-2016>, 2016.
- Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogée, J., Allard, V., Aubinet, M., Buchmann, N., Bernhofer, C., Carrara, A., Chevallier, F., Noblet, N. D., Friend, A. D., Friedlingstein, P., Grünwald, T., Heinesch, B., Keronen, P., Knohl, A., Krinner, G., Loustau, D., Manca, G., Matteucci, G., Miglietta, F., Ourcival, J. M., Papale, D., Pilegaard, K., Rambal, S., Seufert, G.,

- 680 Soussana, J. F., Sanz, M. J., Schulze, E. D., Vesala, T., and Valentini, R.: Europe-Wide Reduction in Primary Productivity
Caused by the Heat and Drought in 2003, *Nature*, 437, 529–533, <https://doi.org/10.1038/nature03972>, 2005.
- Cochard, H., Badel, E., Herbette, S., Delzon, S., Choat, B., and Jansen, S.: Methods for Measuring Plant Vulnerability to
Cavitation: A Critical Review, *Journal of Experimental Botany*, 64, 4779–4791, <https://doi.org/10.1093/jxb/ert193>, 2013.
- Cochard, H., Pimont, F., Ruffault, J., and Martin-StPaul, N.: SurEau: A Mechanistic Model of Plant Water Relations under
685 Extreme Drought, *Annals of Forest Science*, 78, 55, <https://doi.org/10.1007/s13595-021-01067-y>, 2021.
- Collier, N., Hoffman, F. M., Lawrence, D. M., Keppel-Aleks, G., Koven, C. D., Riley, W. J., Mu, M., and Randerson, J. T.:
The International Land Model Benchmarking (ILAMB) System: Design, Theory, and Implementation, *Journal of Advances
in Modeling Earth Systems*, 10, 2731–2754, <https://doi.org/10.1029/2018MS001354>, 2018.
- Comins, H. N. and McMurtrie, R. E.: Long-Term Response of Nutrient-Limited Forests to CO² Enrichment; Equilibrium
690 Behavior of Plant-Soil Models, *Ecological Applications*, 3, 666–681, <https://doi.org/10.2307/1942099>, 1993.
- Cook, B. I., Mankin, J. S., Marvel, K., Williams, A. P., Smerdon, J. E., and Anchukaitis, K. J.: Twenty-First Century Drought
Projections in the CMIP6 Forcing Scenarios, *Earth's Future*, 8, e2019EF001461, <https://doi.org/10.1029/2019EF001461>,
2020.
- Da Sois, L., Mencuccini, M., Castells, E., Sanchez-Martinez, P., and Martínez-Vilalta, J.: How Are Physiological Responses to
695 Drought Modulated by Water Relations and Leaf Economics' Traits in Woody Plants?, *Agricultural Water Management*,
291, 108613, <https://doi.org/10.1016/j.agwat.2023.108613>, 2024.
- De Kauwe, M. G., Medlyn, B. E., Ukkola, A. M., Mu, M., Sabot, M. E. B., Pitman, A. J., Meir, P., Cernusak, L. A., Rifai, S. W.,
Choat, B., Tissue, D. T., Blackman, C. J., Li, X., Roderick, M., and Briggs, P. R.: Identifying Areas at Risk of Drought-Induced
Tree Mortality across South-Eastern Australia, *Global Change Biology*, 26, 5716–5733, <https://doi.org/10.1111/gcb.15215>,
700 2020.
- Desprez-Loustau, M.-L., Marçais, B., Nageleisen, L.-M., Piou, D., and Vannini, A.: Interactive Effects of Drought and Pathogens
in Forest Trees, *Annals of Forest Science*, 63, 597–612, <https://doi.org/10.1051/forest:2006040>, 2006.
- Dietze, M. C., Fox, A., Beck-Johnson, L. M., Betancourt, J. L., Hooten, M. B., Jarnevich, C. S., Keitt, T. H., Kenney, M. A.,
Laney, C. M., Larsen, L. G., Loescher, H. W., Lunch, C. K., Pijanowski, B. C., Randerson, J. T., Read, E. K., Tredennick,
705 A. T., Vargas, R., Weathers, K. C., and White, E. P.: Iterative Near-Term Ecological Forecasting: Needs, Opportunities, and
Challenges, *Proceedings of the National Academy of Sciences*, 115, 1424–1432, <https://doi.org/10.1073/pnas.1710231115>,
2018.
- Eckes-Shephard, A. H., Tiavlovsky, E., Chen, Y., Fonti, P., and Friend, A. D.: Direct Response of Tree Growth
to Soil Water and Its Implications for Terrestrial Carbon Cycle Modelling, *Global Change Biology*, 27, 121–135,
710 <https://doi.org/10.1111/gcb.15397>, 2021.
- Eller, C. B., Rowland, L., Oliveira, R. S., Bittencourt, P. R. L., Barros, F. V., da Costa, A. C. L., Meir, P., Friend, A. D.,
Mencuccini, M., Sitch, S., and Cox, P.: Modelling Tropical Forest Responses to Drought and El Niño with a Stomatal
Optimization Model Based on Xylem Hydraulics, *Philosophical Transactions of the Royal Society B: Biological Sciences*,
373, 20170315, <https://doi.org/10.1098/rstb.2017.0315>, 2018.
- 715 Eller, C. B., Rowland, L., Mencuccini, M., Rosas, T., Williams, K., Harper, A., Medlyn, B. E., Wagner, Y., Klein, T., Teodoro,
G. S., Oliveira, R. S., Matos, I. S., Rosado, B. H. P., Fuchs, K., Wohlfahrt, G., Montagnani, L., Meir, P., Sitch, S., and Cox,

- P. M.: Stomatal Optimization Based on Xylem Hydraulics (SOX) Improves Land Surface Model Simulation of Vegetation Responses to Climate, *New Phytologist*, 226, 1622–1637, <https://doi.org/10.1111/nph.16419>, 2020.
- European Environment Agency: Global and European Temperature, Tech. rep., 2019.
- 720 Fink, A. H., Brücher, T., Krüger, A., Leckebusch, G. C., Pinto, J. G., and Ulbrich, U.: The 2003 European Summer Heatwaves and Drought – Synoptic Diagnosis and Impacts, *Weather*, 59, 209–216, <https://doi.org/10.1256/wea.73.04>, 2004.
- Fisher, R. A., Williams, M., Do Vale, R. L., Da Costa, A. L., and Meir, P.: Evidence from Amazonian Forests Is Consistent with Isohydic Control of Leaf Water Potential, *Plant, Cell & Environment*, 29, 151–165, <https://doi.org/10.1111/j.1365-3040.2005.01407.x>, 2006.
- 725 Flexas, J., Scoffoni, C., Gago, J., and Sack, L.: Leaf Mesophyll Conductance and Leaf Hydraulic Conductance: An Introduction to Their Measurement and Coordination, *Journal of Experimental Botany*, 64, 3965–3981, <https://doi.org/10.1093/jxb/ert319>, 2013.
- Flo, V., Martínez-Vilalta, J., Granda, V., Mencuccini, M., and Poyatos, R.: Vapour Pressure Deficit Is the Main Driver of Tree Canopy Conductance across Biomes, *Agricultural and Forest Meteorology*, 322, 109 029, 730 <https://doi.org/10.1016/j.agrformet.2022.109029>, 2022.
- Frei, E. R., Gossner, M. M., Vitasse, Y., Queloz, V., Dubach, V., Gessler, A., Ginzler, C., Hagedorn, F., Meusburger, K., Moor, M., Samblás Vives, E., Rigling, A., Uitentuis, I., von Arx, G., and Wohlgemuth, T.: European Beech Dieback after Premature Leaf Senescence during the 2018 Drought in Northern Switzerland, *Plant Biology*, 24, 1132–1145, <https://doi.org/10.1111/plb.13467>, 2022.
- 735 Gampe, D., Zscheischler, J., Reichstein, M., O’Sullivan, M., Smith, W. K., Sitch, S., and Buermann, W.: Increasing Impact of Warm Droughts on Northern Ecosystem Productivity over Recent Decades, *Nature Climate Change*, 11, 772–779, <https://doi.org/10.1038/s41558-021-01112-8>, 2021.
- Génard, M., Fishman, S., Vercambre, G., Huguet, J.-G., Bussi, C., Besset, J., and Habib, R.: A Biophysical Analysis of Stem and Root Diameter Variations in Woody Plants, *Plant Physiology*, 126, 188–202, 2001.
- 740 Gerten, D., Schaphoff, S., Haberlandt, U., Lucht, W., and Sitch, S.: Terrestrial Vegetation and Water Balance—Hydrological Evaluation of a Dynamic Global Vegetation Model, *Journal of Hydrology*, 286, 249–270, <https://doi.org/10.1016/j.jhydrol.2003.09.029>, 2004.
- Guada, G., Camarero, J. J., Sánchez-Salguero, R., and Cerrillo, R. M. N.: Limited Growth Recovery after Drought-Induced Forest Dieback in Very Defoliated Trees of Two Pine Species, *Frontiers in Plant Science*, 7, <https://doi.org/10.3389/fpls.2016.00418>, 745 2016.
- Hajek, P., Link, R. M., Nock, C. A., Bauhus, J., Gebauer, T., Gessler, A., Kovach, K., Messier, C., Paquette, A., Saurer, M., Scherer-Lorenzen, M., Rose, L., and Schuldt, B.: Mutually Inclusive Mechanisms of Drought-induced Tree Mortality, *Global Change Biology*, 28, 3365–3378, <https://doi.org/10.1111/gcb.16146>, 2022.
- Hammond, W. M., Yu, K., Wilson, L. A., Will, R. E., Anderegg, W. R. L., and Adams, H. D.: Dead or Dying? Quantifying 750 the Point of No Return from Hydraulic Failure in Drought-induced Tree Mortality, *New Phytologist*, 223, 1834–1843, <https://doi.org/10.1111/nph.15922>, 2019.
- Hartmann, H., Bastos, A., Das, A. J., Esquivel-Muelbert, A., Hammond, W. M., Martínez-Vilalta, J., McDowell, N. G., Powers, J. S., Pugh, T. A. M., Ruthrof, K. X., and Allen, C. D.: Climate Change Risks to Global Forest Health:

- Emergence of Unexpected Events of Elevated Tree Mortality Worldwide, *Annual Review of Plant Biology*, 73, 673–702, <https://doi.org/10.1146/annurev-arplant-102820-012804>, 2022.
- 755 Haxeltine, A. and Prentice, I. C.: BIOME3: An Equilibrium Terrestrial Biosphere Model Based on Ecophysiological Constraints, Resource Availability, and Competition among Plant Functional Types, *Global Biogeochemical Cycles*, 10, 693–709, <https://doi.org/10.1029/96GB02344>, 1996.
- Helton, J. C. and Davis, F. J.: Latin Hypercube Sampling and the Propagation of Uncertainty in Analyses of Complex Systems, *Reliability Engineering & System Safety*, 81, 23–69, [https://doi.org/10.1016/S0951-8320\(03\)00058-9](https://doi.org/10.1016/S0951-8320(03)00058-9), 2003.
- 760 Hickler, T., Prentice, I. C., Smith, B., Sykes, M. T., and Zaehle, S.: Implementing Plant Hydraulic Architecture within the LPJ Dynamic Global Vegetation Model, *Global Ecology and Biogeography*, 0, 060811081017001–???, <https://doi.org/10.1111/j.1466-822X.2006.00254.x>, 2006.
- Hickler, T., Vohland, K., Feehan, J., Miller, P. A., Smith, B., Costa, L., Giesecke, T., Fronzek, S., Carter, T. R., Cramer, W., 765 Kühn, I., and Sykes, M. T.: Projecting the Future Distribution of European Potential Natural Vegetation Zones with a Generalized, Tree Species-Based Dynamic Vegetation Model: Future Changes in European Vegetation Zones, *Global Ecology and Biogeography*, 21, 50–63, <https://doi.org/10.1111/j.1466-8238.2010.00613.x>, 2012.
- Jansen, M. J.: Analysis of Variance Designs for Model Output, *Computer Physics Communications*, 117, 35–43, [https://doi.org/10.1016/S0010-4655\(98\)00154-4](https://doi.org/10.1016/S0010-4655(98)00154-4), 1999.
- 770 Johnson, D., Woodruff, D., McCulloh, K., and Meinzer, F.: Leaf Hydraulic Conductance, Measured in Situ, Declines and Recovers Daily: Leaf Hydraulics, Water Potential and Stomatal Conductance in Four Temperate and Three Tropical Tree Species, *Tree Physiology*, 29, 879–887, <https://doi.org/10.1093/treephys/tpp031>, 2009.
- Johnson, D. M., Wortemann, R., McCulloh, K. A., Jordan-Meille, L., Ward, E., Warren, J. M., Palmroth, S., and Domec, J.-C.: A Test of the Hydraulic Vulnerability Segmentation Hypothesis in Angiosperm and Conifer Tree Species, *Tree Physiology*, 775 36, 983–993, <https://doi.org/10.1093/treephys/tpw031>, 2016.
- Jones, H. G. and Sutherland, R. A.: Stomatal Control of Xylem Embolism, *Plant, Cell & Environment*, 14, 607–612, <https://doi.org/10.1111/j.1365-3040.1991.tb01532.x>, 1991.
- Jönsson, A. M., Schroeder, L. M., Lagergren, F., Anderbrant, O., and Smith, B.: Guess the Impact of *Ips typographus*—An Ecosystem Modelling Approach for Simulating Spruce Bark Beetle Outbreaks, *Agricultural and Forest Meteorology*, 166–167, 780 188–200, <https://doi.org/10.1016/j.agrformet.2012.07.012>, 2012.
- Joshi, J., Stocker, B. D., Hofhansl, F., Zhou, S., Dieckmann, U., and Prentice, I. C.: Towards a Unified Theory of Plant Photosynthesis and Hydraulics, *Nature Plants*, 8, 1304–1316, <https://doi.org/10.1038/s41477-022-01244-5>, 2022.
- Kannenberg, S. A., Driscoll, A. W., Malesky, D., and Anderegg, W. R.: Rapid and Surprising Dieback of Utah Juniper in the Southwestern USA Due to Acute Drought Stress, *Forest Ecology and Management*, 480, 118 639, 785 <https://doi.org/10.1016/j.foreco.2020.118639>, 2021.
- Kennedy, D., Swenson, S., Oleson, K. W., Lawrence, D. M., Fisher, R., Lola da Costa, A. C., and Gentine, P.: Implementing Plant Hydraulics in the Community Land Model, Version 5, *Journal of Advances in Modeling Earth Systems*, 11, 485–513, <https://doi.org/10.1029/2018MS001500>, 2019.
- Kirschbaum, M. U. F. and Paul, K. I.: Modelling C and N Dynamics in Forest Soils with a Modified Version of the CENTURY 790 Model, *Soil Biology and Biochemistry*, 34, 341–354, [https://doi.org/10.1016/S0038-0717\(01\)00189-4](https://doi.org/10.1016/S0038-0717(01)00189-4), 2002.

- Klein, T.: The Variability of Stomatal Sensitivity to Leaf Water Potential across Tree Species Indicates a Continuum between Isohydic and Anisohydic Behaviours, *Functional Ecology*, 28, 1313–1320, <https://doi.org/10.1111/1365-2435.12289>, 2014.
- Körner, C.: Paradigm Shift in Plant Growth Control, *Current Opinion in Plant Biology*, 25, 107–114, <https://doi.org/10.1016/j.pbi.2015.05.003>, 2015.
- 795 Körner, C.: No Need for Pipes When the Well Is Dry—a Comment on Hydraulic Failure in Trees, *Tree Physiology*, 39, 695–700, <https://doi.org/10.1093/treephys/tpz030>, 2019.
- Köstner, B. M. M., Schulze, E. D., Kelliher, F. M., Hollinger, D. Y., Byers, J. N., Hunt, J. E., McSeveny, T. M., Meserth, R., and Weir, P. L.: Transpiration and Canopy Conductance in a Pristine Broad-Leaved Forest of *Nothofagus*: An Analysis of Xylem Sap Flow and Eddy Correlation Measurements, *Oecologia*, 91, 350–359, <https://doi.org/10.1007/BF00317623>, 1992.
- 800 Lagergren, F., Jönsson, A. M., Blennow, K., and Smith, B.: Implementing Storm Damage in a Dynamic Vegetation Model for Regional Applications in Sweden, *Ecological Modelling*, 247, 71–82, <https://doi.org/10.1016/j.ecolmodel.2012.08.011>, 2012.
- Lamarque, J.-F., Kyle, G. P., Meinshausen, M., Riahi, K., Smith, S. J., van Vuuren, D. P., Conley, A. J., and Vitt, F.: Global and Regional Evolution of Short-Lived Radiatively-Active Gases and Aerosols in the Representative Concentration Pathways, *Climatic Change*, 109, 191, <https://doi.org/10.1007/s10584-011-0155-0>, 2011.
- 805 Lambers, H. and Oliveira, R. S.: *Plant Water Relations*, pp. 187–263, Springer International Publishing, Cham, https://doi.org/10.1007/978-3-030-29639-1_5, 2019.
- Lan, X., Tans, P., Thoning, K., and NOAA Global Monitoring Laboratory: Trends in Globally-Averaged CO₂ Determined from NOAA Global Monitoring Laboratory Measurements., <https://doi.org/10.15138/9N0H-ZH07>, 2023.
- Lindeskog, M., Smith, B., Lagergren, F., Sycheva, E., Ficko, A., Pretzsch, H., and Rammig, A.: Accounting for Forest
- 810 Management in the Estimation of Forest Carbon Balance Using the Dynamic Vegetation Model LPJ-GUESS (v4.0, R9710): Implementation and Evaluation of Simulations for Europe, *Geoscientific Model Development*, 14, 6071–6112, <https://doi.org/10.5194/gmd-14-6071-2021>, 2021.
- Lloret, F., Siscart, D., and Dalmases, C.: Canopy Recovery after Drought Dieback in Holm-Oak Mediterranean Forests of Catalonia (NE Spain), *Global Change Biology*, 10, 2092–2099, <https://doi.org/10.1111/j.1365-2486.2004.00870.x>, 2004.
- 815 Martin-StPaul, N., Delzon, S., and Cochard, H.: Sureau Database : A Database Of Hydraulic And Stomatal Traits For Modelling Drought Resistance In Plants, <https://doi.org/10.5281/ZENODO.854700>, 2017.
- Martínez-Vilalta, J. and Garcia-Forner, N.: Water Potential Regulation, Stomatal Behaviour and Hydraulic Transport under Drought: Deconstructing the Iso/Anisohydic Concept, *Plant, Cell & Environment*, 40, 962–976, <https://doi.org/10.1111/pce.12846>, 2017.
- 820 Martínez-Vilalta, J., Poyatos, R., Aguadé, D., Retana, J., and Mencuccini, M.: A New Look at Water Transport Regulation in Plants, *New Phytologist*, 204, 105–115, <https://doi.org/10.1111/nph.12912>, 2014.
- Mcdowell, N., Pockman, W. T., Allen, C. D., Breshears, D. D., Cobb, N., Kolb, T., Plaut, J., Sperry, J., West, A., Williams, D. G., and Yezpe, E. A.: Mechanisms of Plant Survival and Mortality during Drought: Why Do Some Plants Survive While Others Succumb to Drought?, *New Phytologist*, 178, 719–739, <https://doi.org/10.1111/j.1469-8137.2008.02436.x>, 2008.
- 825 Mckay, M. D., Beckman, F. J., and Conover, W. J.: A Comparison of Three Methods for Selecting Values of Input Variables in the Analysis of Output From a Computer Code.

- Méndez-Alonzo, R., Ewers, F. W., Jacobsen, A. L., Pratt, R. B., Scoffoni, C., Bartlett, M. K., and Sack, L.: Covariation between Leaf Hydraulics and Biomechanics Is Driven by Leaf Density in Mediterranean Shrubs, *Trees*, 33, 507–519, <https://doi.org/10.1007/s00468-018-1796-7>, 2019.
- 830 Neycken, A., Scheggia, M., Bigler, C., and Lévesque, M.: Long-Term Growth Decline Precedes Sudden Crown Dieback of European Beech, *Agricultural and Forest Meteorology*, 324, 109–103, <https://doi.org/10.1016/j.agrformet.2022.109103>, 2022.
- Nolan, R. H., Gauthey, A., Losso, A., Medlyn, B. E., Smith, R., Chhajed, S. S., Fuller, K., Song, M., Li, X., Beaumont, L. J., Boer, M. M., Wright, I. J., and Choat, B.: Hydraulic Failure and Tree Size Linked with Canopy Die-back in Eucalypt Forest during Extreme Drought, *New Phytologist*, 230, 1354–1365, <https://doi.org/10.1111/nph.17298>, 2021.
- 835 Nolf, M., Creek, D., Duursma, R., Holtum, J., Mayr, S., and Choat, B.: Stem and Leaf Hydraulic Properties Are Finely Coordinated in Three Tropical Rain Forest Tree Species, *Plant, Cell & Environment*, 38, 2652–2661, <https://doi.org/10.1111/pce.12581>, 2015.
- Oberpriller, J., Herschlein, C., Anthoni, P., Arneth, A., Krause, A., Rammig, A., Lindeskog, M., Olin, S., and Hartig, F.: Climate and Parameter Sensitivity and Induced Uncertainties in Carbon Stock Projections for European Forests (Using LPJ-GUESS 4.0), *Geoscientific Model Development*, 15, 6495–6519, <https://doi.org/10.5194/gmd-15-6495-2022>, 2022.
- 840 Pammenter, N. W. and Van Der Willigen, C.: A Mathematical and Statistical Analysis of the Curves Illustrating Vulnerability of Xylem to Cavitation, *Tree Physiology*, 18, 589–593, <https://doi.org/10.1093/treephys/18.8.589>, 1998.
- Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., Phillips, O. L., Shvidenko, A., Lewis, S. L., Canadell, J. G., Ciais, P., Jackson, R. B., Pacala, S. W., McGuire, A. D., Piao, S., Rautiainen, A., Sitch, S., and Hayes, D.: A Large and Persistent Carbon Sink in the World’s Forests, *Science*, 333, 988–993, <https://doi.org/10.1126/science.1201609>, 2011.
- 845 Pan, Y., Birdsey, R. A., Phillips, O. L., Houghton, R. A., Fang, J., Kauppi, P. E., Keith, H., Kurz, W. A., Ito, A., Lewis, S. L., Nabuurs, G.-J., Shvidenko, A., Hashimoto, S., Lerink, B., Schepaschenko, D., Castanho, A., and Murdiyarso, D.: The Enduring World Forest Carbon Sink, *Nature*, 631, 563–569, <https://doi.org/10.1038/s41586-024-07602-x>, 2024.
- 850 Papastefanou, P., Zang, C. S., Pugh, T. A. M., Liu, D., Grams, T. E. E., Hickler, T., and Rammig, A.: A Dynamic Model for Strategies and Dynamics of Plant Water-Potential Regulation Under Drought Conditions, *Frontiers in Plant Science*, 11, 373, <https://doi.org/10.3389/fpls.2020.00373>, 2020.
- Papastefanou, P., Pugh, T. A. M., Buras, A., Fleischer, K., Grams, T., Hickler, T., Lapola, D., Liu, D., Zang, C., and Rammig, A.: Simulated Sensitivity of the Amazon Rainforest to Extreme Drought, *Environmental Research Letters*, 855 <https://doi.org/10.1088/1748-9326/ad8f48>, 2024.
- Pappas, C., Fatichi, S., Leuzinger, S., Wolf, A., and Burlando, P.: Sensitivity Analysis of a Process-Based Ecosystem Model: Pinpointing Parameterization and Structural Issues, *Journal of Geophysical Research: Biogeosciences*, 118, 505–528, <https://doi.org/10.1002/jgrg.20035>, 2013.
- Parton, W. J., Scurlock, J. M. O., Ojima, D. S., Gilmanov, T. G., Scholes, R. J., Schimel, D. S., Kirchner, T., Menaut, 860 J.-C., Seastedt, T., Garcia Moya, E., Kamnalrut, A., and Kinyamario, J. I.: Observations and Modeling of Biomass and Soil Organic Matter Dynamics for the Grassland Biome Worldwide, *Global Biogeochemical Cycles*, 7, 785–809, <https://doi.org/10.1029/93GB02042>, 1993.

Parton, W. J., Hanson, P. J., Swanston, C., Torn, M., Trumbore, S. E., Riley, W., and Kelly, R.: ForCent Model Development and Testing Using the Enriched Background Isotope Study Experiment, *Journal of Geophysical Research: Biogeosciences*, 115, <https://doi.org/10.1029/2009JG001193>, 2010.

Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W., Poindexter, C., Chen, J., Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Reichstein, M., Ribeca, A., Van Ingen, C., Vuichard, N., Zhang, L., Amiro, B., Ammann, C., Arain, M. A., Ardö, J., Arkebauer, T., Arndt, S. K., Arriga, N., Aubinet, M., Aurela, M., Baldocchi, D., Barr, A., Beamesderfer, E., Marchesini, L. B., Bergeron, O., Beringer, J., Bernhofer, C., Berveiller, D., Billesbach, D., Black, T. A., Blanken, P. D., Bohrer, G., Boike, J., Bolstad, P. V., Bonal, D., Bonnefond, J.-M., Bowling, D. R., Bracho, R., Brodeur, J., Brümmer, C., Buchmann, N., Burban, B., Burns, S. P., Buysse, P., Cale, P., Cavagna, M., Cellier, P., Chen, S., Chini, I., Christensen, T. R., Cleverly, J., Collalti, A., Consalvo, C., Cook, B. D., Cook, D., Coursolle, C., Cremonese, E., Curtis, P. S., D'Andrea, E., Da Rocha, H., Dai, X., Davis, K. J., Cinti, B. D., Grandcourt, A. D., Ligne, A. D., De Oliveira, R. C., Delpierre, N., Desai, A. R., Di Bella, C. M., Tommasi, P. D., Dolman, H., Domingo, F., Dong, G., Dore, S., Duce, P., Dufrêne, E., Dunn, A., Dušek, J., Eamus, D., Eichelmann, U., Elkhidir, H. A. M., Eugster, W., Ewenz, C. M., Ewers, B., Famulari, D., Fares, S., Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G., Fischer, M., Frank, J., Galvagno, M., Gharun, M., Gianelle, D., Gielen, B., Gioli, B., Gitelson, A., Goded, I., Goeckede, M., Goldstein, A. H., Gough, C. M., Goulden, M. L., Graf, A., Griebel, A., Gruening, C., Grünwald, T., Hammerle, A., Han, S., Han, X., Hansen, B. U., Hanson, C., Hatakka, J., He, Y., Hehn, M., Heinesch, B., Hinko-Najera, N., Hörtnagl, L., Hutley, L., Ibrom, A., Ikawa, H., Jackowicz-Korczynski, M., Janouš, D., Jans, W., Jassal, R., Jiang, S., Kato, T., Khomik, M., Klatt, J., Knohl, A., Knox, S., Kobayashi, H., Koerber, G., Kolle, O., Kosugi, Y., Kotani, A., Kowalski, A., Kruijt, B., Kurbatova, J., Kutsch, W. L., Kwon, H., Launiainen, S., Laurila, T., Law, B., Leuning, R., Li, Y., Liddell, M., Limousin, J.-M., Lion, M., Liska, A. J., Lohila, A., López-Ballesteros, A., López-Blanco, E., Loubet, B., Loustau, D., Lucas-Moffat, A., Lüers, J., Ma, S., Macfarlane, C., Magliulo, V., Maier, R., Mammarella, I., Manca, G., Marcolla, B., Margolis, H. A., Marras, S., Massman, W., Mastepanov, M., Matamala, R., Matthes, J. H., Mazzenga, F., Mccaughey, H., Mchugh, I., Mcmillan, A. M. S., Merbold, L., Meyer, W., Meyers, T., Miller, S. D., Minerbi, S., Moderow, U., Monson, R. K., Montagnani, L., Moore, C. E., Moors, E., Moreaux, V., Moureaux, C., Munger, J. W., Nakai, T., Neirynck, J., Nesic, Z., Nicolini, G., Noormets, A., Northwood, M., Nosetto, M., Nouvellon, Y., Novick, K., Oechel, W., Olesen, J. E., Ourcival, J.-M., Papuga, S. A., Parmentier, F.-J., Paul-Limoges, E., Pavelka, M., Peichl, M., Pendall, E., Phillips, R. P., Pilegaard, K., Pirk, N., Posse, G., Powell, T., Prasse, H., Prober, S. M., Rambal, S., Rannik, Ü., Raz-Yaseef, N., Rebmann, C., Reed, D., Dios, V. R. D., Restrepo-Coupe, N., Reverter, B. R., Roland, M., Sabbatini, S., Sachs, T., Saleska, S. R., Sánchez-Cañete, E. P., Sanchez-Mejia, Z. M., Schmid, H. P., Schmidt, M., Schneider, K., Schrader, F., Schroder, I., Scott, R. L., Sedláč, P., Serrano-Ortíz, P., Shao, C., Shi, P., Shironya, I., Siebicke, L., Šigut, L., Silberstein, R., Sirca, C., Spano, D., Steinbrecher, R., Stevens, R. M., Sturtevant, C., Suyker, A., Tagesson, T., Takanashi, S., Tang, Y., Tapper, N., Thom, J., Tomassucci, M., Tuovinen, J.-P., Urbanski, S., Valentini, R., Van Der Molen, M., Van Gorsel, E., Van Huissteden, K., Varlagin, A., Verfaillie, J., Vesala, T., Vincke, C., Vitale, D., Vygodskaya, N., Walker, J. P., Walter-Shea, E., Wang, H., Weber, R., Westermann, S., Wille, C., Wofsy, S., Wohlfahrt, G., Wolf, S., Woodgate, W., Li, Y., Zampedri, R., Zhang, J., Zhou, G., Zona, D., Agarwal, D., Biraud, S., Torn, M., and Papale, D.: The FLUXNET2015 Dataset and the ONEFlux Processing Pipeline for Eddy Covariance Data, *Scientific Data*, 7, <https://doi.org/10.1038/s41597-020-0534-3>, 2020.

- 900 Peters, R. L., Steppe, K., Cuny, H. E., De Pauw, D. J., Frank, D. C., Schaub, M., Rathgeber, C. B., Cabon, A., and Fonti, P.: Turgor – a Limiting Factor for Radial Growth in Mature Conifers along an Elevational Gradient, *New Phytologist*, 229, 213–229, <https://doi.org/10.1111/nph.16872>, 2021.
- Potkay, A., Hölttä, T., Trugman, A. T., and Fan, Y.: Turgor-Limited Predictions of Tree Growth, Height and Metabolic Scaling over Tree Lifespans, *Tree Physiology*, 42, 229–252, <https://doi.org/10.1093/treephys/tpab094>, 2022.
- 905 Powell, T. L., Galbraith, D. R., Christoffersen, B. O., Harper, A., Imbuzeiro, H. M. A., Rowland, L., Almeida, S., Brando, P. M., Da Costa, A. C. L., Costa, M. H., Levine, N. M., Malhi, Y., Saleska, S. R., Sotta, E., Williams, M., Meir, P., and Moorcroft, P. R.: Confronting Model Predictions of Carbon Fluxes with Measurements of Amazon Forests Subjected to Experimental Drought, *New Phytologist*, 200, 350–365, <https://doi.org/10.1111/nph.12390>, 2013.
- Pugh, T. A. M., Arneth, A., Kautz, M., Poulter, B., and Smith, B.: Important Role of Forest Disturbances in the Global Biomass Turnover and Carbon Sinks, *Nature Geoscience*, 12, 730–735, <https://doi.org/10.1038/s41561-019-0427-2>, 2019.
- 910 Puy, A., Piano, S. L., Saltelli, A., and Levin, S. A.: sensobol: An R Package to Compute Variance-Based Sensitivity Indices, *Journal of Statistical Software*, 102, 1–37, <https://doi.org/10.18637/jss.v102.i05>, 2022.
- Rouault, G., Candau, J.-N., Lieutier, F., Nageleisen, L.-M., Martin, J.-C., and Warzée, N.: Effects of Drought and Heat on Forest Insect Populations in Relation to the 2003 Drought in Western Europe, *Annals of Forest Science*, 63, 613–624, <https://doi.org/10.1051/forest:2006044>, 2006.
- 915 Ruffault, J., Pimont, F., Cochard, H., Dupuy, J.-L., and Martin-StPaul, N.: SurEau-Ecos v2.0: A Trait-Based Plant Hydraulics Model for Simulations of Plant Water Status and Drought-Induced Mortality at the Ecosystem Level, *Geoscientific Model Development*, 15, 5593–5626, <https://doi.org/10.5194/gmd-15-5593-2022>, 2022.
- Saltelli, A., ed.: *Global Sensitivity Analysis: The Primer*, Wiley, Chichester, West Sussex, 2008.
- 920 Saltelli, A., Annoni, P., Azzini, I., Campolongo, F., Ratto, M., and Tarantola, S.: Variance Based Sensitivity Analysis of Model Output. Design and Estimator for the Total Sensitivity Index, *Computer Physics Communications*, 181, 259–270, <https://doi.org/10.1016/j.cpc.2009.09.018>, 2010.
- Saxton, K. E., Rawls, W. J., Romberger, J. S., and Papendick, R. I.: Estimating Generalized Soil-water Characteristics from Texture, *Soil Science Society of America Journal*, 50, 1031–1036, <https://doi.org/10.2136/sssaj1986.03615995005000040039x>, 1986.
- 925 Schönbeck, L. C., Schuler, P., Lehmann, M. M., Mas, E., Mekarni, L., Pivovarov, A. L., Turberg, P., and Grossiord, C.: Increasing Temperature and Vapour Pressure Deficit Lead to Hydraulic Damages in the Absence of Soil Drought, *Plant, Cell & Environment*, 45, 3275–3289, <https://doi.org/10.1111/pce.14425>, 2022.
- Schuldt, B., Buras, A., Arend, M., Vitasse, Y., Beierkuhnlein, C., Damm, A., Gharun, M., Grams, T. E., Hauck, M., Hajek, P., Hartmann, H., Hiltbrunner, E., Hoch, G., Holloway-Phillips, M., Körner, C., Larysch, E., Lübke, T., Nelson, D. B., Rammig, A., Rigling, A., Rose, L., Ruehr, N. K., Schumann, K., Weiser, F., Werner, C., Wohlgemuth, T., Zang, C. S., and Kahmen, A.: A First Assessment of the Impact of the Extreme 2018 Summer Drought on Central European Forests, *Basic and Applied Ecology*, 45, 86–103, <https://doi.org/10.1016/j.baae.2020.04.003>, 2020.
- 930 Scoffoni, C., Rawls, M., McKown, A., Cochard, H., and Sack, L.: Decline of Leaf Hydraulic Conductance with Dehydration: Relationship to Leaf Size and Venation Architecture, *Plant Physiology*, 156, 832–843, <https://doi.org/10.1104/pp.111.173856>, 2011.

- Seiler, C., Melton, J. R., Arora, V. K., Sitch, S., Friedlingstein, P., Anthoni, P., Goll, D., Jain, A. K., Joetzjer, E., Lienert, S., Lombardozzi, D., Luyssaert, S., Nabel, J. E. M. S., Tian, H., Vuichard, N., Walker, A. P., Yuan, W., and Zaehle, S.: Are Terrestrial Biosphere Models Fit for Simulating the Global Land Carbon Sink?, *Journal of Advances in Modeling Earth Systems*, 14, <https://doi.org/10.1029/2021ms002946>, 2022.
- 940 Senf, C., Buras, A., Zang, C. S., Rammig, A., and Seidl, R.: Excess Forest Mortality Is Consistently Linked to Drought across Europe, *Nature Communications*, 11, 6200, <https://doi.org/10.1038/s41467-020-19924-1>, 2020.
- Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S., Lucht, W., Sykes, M. T., Thonicke, K., and Venevsky, S.: Evaluation of Ecosystem Dynamics, Plant Geography and Terrestrial Carbon Cycling in the LPJ Dynamic Global Vegetation Model, *Global Change Biology*, 9, 161–185, <https://doi.org/10.1046/j.1365-2486.2003.00569.x>, 2003.
- 945 Smith, B., Prentice, I. C., and Sykes, M. T.: Representation of Vegetation Dynamics in the Modelling of Terrestrial Ecosystems: Comparing Two Contrasting Approaches within European Climate Space: *Vegetation Dynamics in Ecosystem Models*, *Global Ecology and Biogeography*, 10, 621–637, <https://doi.org/10.1046/j.1466-822X.2001.t01-1-00256.x>, 2001.
- 950 Smith, B., Wårlind, D., Arneth, A., Hickler, T., Leadley, P., Siltberg, J., and Zaehle, S.: Implications of Incorporating N Cycling and N Limitations on Primary Production in an Individual-Based Dynamic Vegetation Model, *Biogeosciences*, 11, 2027–2054, <https://doi.org/10.5194/bg-11-2027-2014>, 2014.
- Sperry, J. S., Adler, F. R., Campbell, G. S., and Comstock, J. P.: Limitation of Plant Water Use by Rhizosphere and Xylem Conductance: Results from a Model, *Plant, Cell & Environment*, 21, 347–359, <https://doi.org/10.1046/j.1365-3040.1998.00287.x>, 1998.
- 955 Steppe, K., De Pauw, D. J. W., Lemeur, R., and Vanrolleghem, P. A.: A Mathematical Model Linking Tree Sap Flow Dynamics to Daily Stem Diameter Fluctuations and Radial Stem Growth, *Tree Physiology*, 26, 257–273, <https://doi.org/10.1093/treephys/26.3.257>, 2006.
- Sulman, B. N., Roman, D. T., Yi, K., Wang, L., Phillips, R. P., and Novick, K. A.: High Atmospheric Demand for Water Can Limit Forest Carbon Uptake and Transpiration as Severely as Dry Soil, *Geophysical Research Letters*, 43, 9686–9695, <https://doi.org/10.1002/2016GL069416>, 2016.
- 960 Sykes, Martint. and Prentice, I.: Climate Change, Tree Species Distributions and Forest Dynamics: A Case Study in the Mixed Conifer/Northern Hardwoods Zone of Northern Europe, *Climatic Change*, 34, <https://doi.org/10.1007/bf00224628>, 1996.
- Tardieu, F., Simonneau, T., and Parent, B.: Modelling the Coordination of the Controls of Stomatal Aperture, Transpiration, Leaf Growth, and Absciscic Acid: Update and Extension of the Tardieu–Davies Model, *Journal of Experimental Botany*, 66, 2227–2237, <https://doi.org/10.1093/jxb/erv039>, 2015.
- 965 Torres-Ruiz, J. M., Cochard, H., Delzon, S., Boivin, T., Burlett, R., Cailleret, M., Corso, D., Delmas, C. E. L., De Caceres, M., Diaz-Espejo, A., Fernández-Conradi, P., Guillemot, J., Lamarque, L. J., Limousin, J.-M., Mantova, M., Mencuccini, M., Morin, X., Pimont, F., De Dios, V. R., Ruffault, J., Trueba, S., and Martin-Stpaul, N. K.: Plant Hydraulics at the Heart of Plant, Crops and Ecosystem Functions in the Face of Climate Change, *New Phytologist*, 241, 984–999, <https://doi.org/10.1111/nph.19463>, 2024.
- 970 Tschumi, E., Lienert, S., Bastos, A., Ciais, P., Gregor, K., Joos, F., Knauer, J., Papastefanou, P., Rammig, A., van der Wiel, K., Williams, K., Xu, Y., Zaehle, S., and Zscheischler, J.: Large Variability in Simulated Response of Vegetation Composition

- and Carbon Dynamics to Variations in Drought-Heat Occurrence, *Journal of Geophysical Research: Biogeosciences*, 128, e2022JG007332, <https://doi.org/10.1029/2022JG007332>, 2023.
- 975 Tyree, M. T. and Sperry, J. S.: Vulnerability of Xylem to Cavitation and Embolism, *Annual Review of Plant Physiology and Plant Molecular Biology*, 40, 19–36, <https://doi.org/10.1146/annurev.pp.40.060189.000315>, 1989.
- Tyree, M. T., Davis, S. D., and Cochard, H.: Biophysical Perspectives of Xylem Evolution: Is There a Tradeoff of Hydraulic Efficiency for Vulnerability to Dysfunction?, *IAWA journal*, 15, 335–360, 1994.
- 980 van der Woude, A. M., Peters, W., Joetzjer, E., Lafont, S., Koren, G., Ciais, P., Ramonet, M., Xu, Y., Bastos, A., Botía, S., Sitch, S., de Kok, R., Kneuer, T., Kubistin, D., Jacotot, A., Loubet, B., Herig-Coimbra, P.-H., Loustau, D., and Luijkx, I. T.: Temperature Extremes of 2022 Reduced Carbon Uptake by Forests in Europe, *Nature Communications*, 14, 6218, <https://doi.org/10.1038/s41467-023-41851-0>, 2023.
- Wagner, Y., Volkov, M., Nadal-Sala, D., Ruehr, N. K., Hochberg, U., and Klein, T.: Relationships between Xylem Embolism and Tree Functioning during Drought, Recovery, and Recurring Drought in Aleppo Pine, *Physiologia Plantarum*, 175, e13995, <https://doi.org/10.1111/ppl.13995>, 2023.
- 985 Walthert, L., Ganthaler, A., Mayr, S., Saurer, M., Waldner, P., Walser, M., Zweifel, R., and von Arx, G.: From the Comfort Zone to Crown Dieback: Sequence of Physiological Stress Thresholds in Mature European Beech Trees across Progressive Drought, *Science of The Total Environment*, 753, 141792, <https://doi.org/10.1016/j.scitotenv.2020.141792>, 2021.
- 990 Warm Winter 2020 Team and ICOS Ecosystem Thematic Centre: Warm Winter 2020 Ecosystem Eddy Covariance Flux Product for 73 Stations in FLUXNET-Archive Format—Release 2022-1 (Version 1.0), <https://doi.org/10.18160/2G60-ZHAK>, 2022.
- Whitehead, D.: Regulation of Stomatal Conductance and Transpiration in Forest Canopies, *Tree Physiology*, 18, 633–644, <https://doi.org/10.1093/treephys/18.8-9.633>, 1998.
- Xu, C., Christoffersen, B., Robbins, Z., Knox, R., Fisher, R. A., Chitra-Tarak, R., Slot, M., Solander, K., Kueppers, L., Koven, C., and McDowell, N.: Quantification of Hydraulic Trait Control on Plant Hydrodynamics and Risk of Hydraulic Failure within a Demographic Structured Vegetation Model in a Tropical Forest (FATES-HYDRO V1.0), *Geoscientific Model Development*, 16, 6267–6283, <https://doi.org/10.5194/gmd-16-6267-2023>, 2023.
- 995 Xu, X., Medvigy, D., Powers, J. S., Becknell, J. M., and Guan, K.: Diversity in Plant Hydraulic Traits Explains Seasonal and Inter-Annual Variations of Vegetation Dynamics in Seasonally Dry Tropical Forests, *New Phytologist*, 212, 80–95, <https://doi.org/10.1111/nph.14009>, 2016.
- 1000 Yao, Y., Joetzjer, E., Ciais, P., Viovy, N., Cresto Aleina, F., Chave, J., Sack, L., Bartlett, M., Meir, P., Fisher, R., and Luyssaert, S.: Forest Fluxes and Mortality Response to Drought: Model Description (ORCHIDEE-CAN-NHA R7236) and Evaluation at the Caxiuanã Drought Experiment, *Geoscientific Model Development*, 15, 7809–7833, <https://doi.org/10.5194/gmd-15-7809-2022>, 2022.
- 1005 Zaehle, S., Sitch, S., Smith, B., and Hatterman, F.: Effects of Parameter Uncertainties on the Modeling of Terrestrial Biosphere Dynamics, *Global Biogeochemical Cycles*, 19, 2004GB002395, <https://doi.org/10.1029/2004GB002395>, 2005.
- Zhou, H., Tang, J., Olin, S., and Miller, P. A.: A Comprehensive Evaluation of Hydrological Processes in a Second-Generation Dynamic Vegetation Model, *Hydrological Processes*, 38, e15152, <https://doi.org/10.1002/hyp.15152>, 2024.
- Zhou, S., Duursma, R. A., Medlyn, B. E., Kelly, J. W. G., and Prentice, I. C.: How Should We Model Plant Responses to Drought? An Analysis of Stomatal and Non-Stomatal Responses to Water Stress, *Agricultural and Forest Meteorology*, 182–183, 204–214, <https://doi.org/10.1016/j.agrformet.2013.05.009>, 2013.
- 1010

Zweifel, R., Haeni, M., Buchmann, N., and Eugster, W.: Are Trees Able to Grow in Periods of Stem Shrinkage?, *New Phytologist*, 211, 839–849, <https://doi.org/10.1111/nph.13995>, 2016.

1015 A1: Species-specific comparison of modeled evapotranspiration and evapotranspiration from eddy-covariance flux towers




Figure A1. Of the 12 species analyzed in this study, monospecific eddy-covariance flux sites exist only for the four species shown here. In LPJ-GUESS-HYD (A) modeled ET better matches observed ET patterns for the relatively anisohydric species *Fagus sylvatica* and *Quercus robur* better than standard LPJ-GUESS (B) during three pan-European droughts. On the contrary, for the more isohydric species *Picea abies* and *Pinus sylvestris* both LPJ-GUESS-HYD (A) and LPJ-GUESS (B) underestimate observed ET. The dotted black line indicates perfect agreement between model and observations.




Figure A2. Species-specific daily evapotranspiration rates under differing levels of vapor pressure deficit from A) eddy-covariance flux towers, B) LPJ-GUESS-HYD, and C) standard LPJ-GUESS using ψ_{50} values from [Martin-StPaul et al. \(2017\)](#). The colors are ranked according to the λ of each species (see Fig. 2) from high λ (light) to low λ (dark). Daily VPD was binned into 6 equally-sized classes representing increasing levels of drought. Species-specific responses to drought remain constant in LPJ-GUESS while clear differences between more anisohydric and more isohydric species are seen in LPJ-GUESS-HYD.

A3: Influence of soil water retention curve on ψ_s

$$a = \frac{-e^{-4.396 - 0.0715 * \% \text{clay} - 4.88 * 10^{-4} * \% \text{sand}^2 - 4.285 * 10^{-5} * \% \text{sand}^2 * \% \text{clay}}}{10} \quad (\text{A1})$$

1020 and (see Saxton et al. (1986) Eq. 6):

$$b = -3.14 - 0.00222 * \% \text{clay}^2 - 3.484 * 10^{-5} * \% \text{sand}^2 * \% \text{clay} \quad (\text{A2})$$