

A shift in circadian stem increment patterns in a Pyrenean alpine treeline precedes spring growth after snow melting

5 Helen Flynn^{1,2}, J. Julio Camarero², Alba Sanmiguel-Vallelado³, Francisco Rojas Heredia², Pablo Domínguez Aguilar², Jesús Revuelto², Juan Ignacio López-Moreno²

¹Department of Geosciences- Colorado State University, Fort Collins, Colorado, 80521, USA

²Instituto Pirenaico de Ecología (IPE-CSIC), Zaragoza, 50059, Spain

³iuFOR, EiFAB, Universidad de Valladolid, Campus Duques de Soria, Soria, 42004, Spain.

10

Correspondence to: Helen Flynn (Helen.Flynn@colostate.edu) or Juan Ignacio López-Moreno (nlopez@ipe.csic.es)

Abstract. Changing snow regimes and warmer growing seasons are some climate factors influencing productivity and growth of high-elevation forests and alpine treelines. In low-latitude mountain regions with seasonal snow and drought regimes such as the Pyrenees, these climate factors could negatively impact forest productivity. To address this issue, we assessed the relationships between climate, snow, and inter- and intra-annual radial growth and stem increment data in an alpine *Pinus uncinata* treeline ecotone located in the central Spanish Pyrenees. First, we developed tree-ring width chronologies of the study site to quantify climate-growth relationships. Second, radial growth, tree water deficit, and shrinking/swelling cycles were quantified and identified at monthly to daily scales using fine-resolution dendrometer data. These variables were extracted for three climatically different years, including one of the hottest summers on record in Spain (2022), and were related to soil water content, soil and air temperature, and the dates of snow duration across the treeline ecotone. Warmer February and May temperatures enhanced tree radial growth, probably because of an earlier snow melt-out and start of the growing season and higher growth rates in spring, respectively. The characteristic circadian cycle of stem increment, defined by night swelling and day shrinking, was detected in summer and autumn. However, this pattern inverted during the snow season from November through April, suggesting a transition phase characterized by wet soils and swollen stems preceding the spring onset of growth. Air temperature, soil temperature and moisture, and the presence of snow are strong indicators of how much and for how long mountain trees can grow. Shifts in daily stem increment patterns reveal changes in early growth phenology linked to snow melting.

15
20
25

30

1 Introduction

In mountain areas, warming rates are much higher than in lowlands (Pepin et al., 2015) leading to changes in seasonal snow regimes and soil moisture available for tree growth (Harpold & Molotch, 2015). Forest productivity and tree growth in high-elevation, subalpine forests and alpine treelines are especially sensitive to the warming effects of climate change (Albrich et al., 2020). However, drier conditions could also negatively impact on low- to mid-latitude mountain forests making the late-winter soil moisture coming from snow melting critical for tree growth. This has been observed in mountain areas subjected to strong snow and drought seasonality, such as the southern Rockies, the central Andes, and often in Mediterranean mountain ranges like the Pyrenees (Andrus et al., 2018; Saavedra et al., 2018; Vicente-Serrano et al., 2021; Villalba et al., 1994). High elevation treeline forests are limited in productivity by climate variables like air and soil temperature (Peterson, 1998; Sanmiguel-Vallelado et al., 2021). Such forests are especially sensitive to the warming effects of climate change (Albrich et al., 2020) and can experience longer and, thus, more productive growing seasons (Yang et al., 2024).

Snow distribution and processes are clearly impacted by forest structures, but forests are also influenced by climate and snow as well. For instance, productivity was found to decrease with greater seasonal snow water equivalent (SWE) and later timing of melt out in the United States and southeast Canada (Yang et al., 2024). Similarly, greater snowpacks led to less radial growth in mountain pine (*Pinus uncinata* Ram.) high-elevation forests in the Spanish Pyrenees (Sanmiguel-Vallelado et al., 2019), whereas monthly growth rates were found to be enhanced by a higher soil temperature linked to an earlier snow melt (Sanmiguel-Vallelado et al., 2021). To make these different findings found at yearly to monthly scales compatible, we need closer, fine-resolution approaches to understand how climate factors, and particularly snow dynamics, drive tree radial growth, a proxy of carbon uptake in woody tissues.

Snow is an essential water resource in the Spanish Pyrenees, and is highly variable across the landscape, especially in the high-mountain treeline ecotone (Revuelto et al., 2017). Pyrenean forests have unique controls on snow distribution, redistribution, and sublimation which vary with forest density and structure (Hedstrom & Pomeroy, 1998; López-Moreno & Latron, 2008; Pomeroy et al., 2002; Revuelto et al., 2015; Storck et al., 2002). For this reason, it is important to gain a better understanding of how individual tree growing in alpine treelines respond to snowpack. Snow and vegetation dynamics are being studied more often (Dobbert et al., 2022; Huang et al., 2023; Pomeroy et al., 2006; Sanmiguel-Vallelado et al., 2019, 2021; Yang et al., 2024), specifically in treeline ecotones due to their relevance in forecasting future treeline shifts (Hagedorn et al., 2014; Huang et al., 2023).

The central Spanish Pyrenees, due to its strong seasonality in snow cover and soil water availability during the growing-season, provide an adequate setting to study snow-forest interactions. In this area, treelines are dominated by mountain pine, which shows a growing season from May through October (Camarero et al., 1998). Radial growth rates of this species peak in May

(Sanmiguel-Vallelado et al., 2021). In the Pyrenees, the areas above 1600 meters are snow covered at least 50% of the time
65 December through April with wet soil and low temperature conditions (Gascoin et al., 2015). However, climate change is
shifting the snow regimes in the Spanish Pyrenees leading to a shorter snowpack duration due to a lower snow accumulation
(López-Moreno, 2005; López-Moreno et al., 2020). Summer heatwaves and droughts are becoming common. This was
illustrated by the summer of 2022 which was the hottest in the last several hundred years (Serrano-Notivoli et al., 2023);
Izaguirre et al.). In addition to increasing summer temperatures, climate change is increasing vapor pressure deficit (VPD)
70 which could negatively impact forest productivity and growth by rising water evaporative demand (Noguera et al., 2023).

Trees swell and shrink throughout each day as a function of changes in soil moisture and atmospheric water demand usually
expressed as VPD. In late winter to early spring, usually from February to April, the snow has melted out and soils are wet,
but later in the growing season, higher temperatures dry soils, and increase the drought stress (Zweifel et al., 2016). During
75 times of high heat and low soil moisture, trees can be impacted by drought stress causing them to undergo reversible shrinking
which can be quantified through a metric called tree water deficit (TWD) derived from dendrometer measurements (Zweifel
et al., 2016). Trees predominantly exhibit drought stress (high TWD) during daylight hours which is why most radial growth
is occurring during the night during the growing season (Zweifel et al., 2021). Tree stems are shrinking during the high-stress,
TWD daylight hours, and they are expanding and contributing to irreversible radial growth during the cooler night hours. This
80 mechanism for growth is exhibited during the time of year when it is advantageous for the tree to do so. Circadian approaches
have been used to identify this trend in other biomes with varying results. Some researchers found that circadian clocks play
an important role in triggering the onset of growth processes (Lázaro-Gimeno et al., 2024; Mei-Jun et al., 2023) and are
influenced by environmental factors (Ziaco & Biondi, 2018), while others had variable results (Lüttge & Hertel, 2009).
Climatological conditions in the central Spanish Pyrenees vary greatly over the course of the year which could cause a shift in
85 daily circadian cycles.

If soil temperature and moisture, controlled by the presence of snow, drive the start of the growing season in high mountain
forests and alpine treelines (Sanmiguel-Vallelado et al., 2021), then soil temperature and the presence of snow dictate the
magnitude and timing of tree growth in the treeline ecotone with the shift being dictated by a change in the daily circadian
90 rhythm of shrinking and swelling dynamics. The purpose of this research is to test this hypothesis and to quantify the influence
of varying climate and snow conditions on radial growth in a Pyrenean treeline ecotone at several time scales (year, month,
day and hour). To do this, we: 1. compared tree growth and shrinking and swelling dynamics across three climatically
differentiated years, 2. analyzed daily stem radius fluctuations as related to changes in growth and shrinking/swelling dynamics
during warm and cold seasons, and 3. evaluate how climatic variables (air/soil temperature, snow, precipitation, soil water
95 content) affect growth and TWD.

2 Materials and Methods

2.1 Study Site


100 The study site is a relatively undisturbed treeline ecotone located in the Sierra de las Cutas (42.6371 °N, 0.0512° W), near the “Ordesa and Monte Perdido” National Park in northeastern Spain. The site sits at an average elevation of 2100 meters facing 186° south-southwest and is dominated by mountain pine. The average slope is 17°, reaching a maximum slope of 33° when descending towards the forest. The soils are rocky and basic. The understory is dominated by several shrub species such as *Juniperus communis* L. above the treeline and *Rhododendron ferrugineum* L. and *Calluna vulgaris* L. within the forest (Pardo et al., 2013).

2.2 Climate and dendrometer data

105 In an open area in the upper limit of the tree line, the automatic weather station (AWS) at Las Cutas is equipped to measure air temperature, relative humidity, wind velocity, and incoming solar radiation. The data from 2020 to 2024 were downloaded and processed, which revealed a gap in sensor data from December 6, 2021, to June 17, 2022, due to battery malfunction (Fig. 1). There is a meteorological station 2 km away at the Góriz refuge (42.6634 °N, 0.0148° E) with similar sensors, an additional precipitation gauge and a snow depth sensor. In addition to the AWS at Las Cutas, there were also four soil water content (SWC) sensors (ECH2O probe, model EC-5, Decagon Devices, Pullman, Washington, USA) installed at 10cm depths in the forested zone of the tree line ecotone. One of the four sensors malfunctioned, so SWC data was averaged across the three remaining sensors (Fig. 1). Other stand-scale climatological data were downloaded from the nearby Góriz refuge. The refuge lies on a similarly south-facing aspect at an elevation of 2200 m. The Góriz meteorological station provided additional daily air temperature, precipitation and snow depth data. After correlating the existing daily minimum and maximum air temperature data from Las Cutas and the Góriz refuge, it was determined that the Góriz data could supplement the data from Las Cutas (Fig. A1).

110

115



120 **Figure 1. Daily mean air (T_{air}) and soil (T_{soil}) temperatures, and mean soil water content (SWC) in the three study years. Values are means \pm SE. Snow cover dates are outlined in vertical blue boxes with inconsistent snow cover defined by light blue and consistent snow cover defined by dark blue.**

In September 2020, stainless steel band dendrometers (DR 26, EMS Brno, Czech Republic) were placed on 9 *P. uncinata* individuals in three zones of the tree line ecotone: the forest (TRA), transition to tree line (TRA-TRE), and treeline (TRE).
 125 The dendrometers recorded 15-minute perimeter measurements at a high resolution of 1 μm and air temperature (T_{air}) until October 2024. The diameter at breast height (DBH) and height were also measured for each individual tree (Table A1) when the dendrometers were installed in 2020. Radius values calculated using the perimeter measurements (assuming a spherical circumference) were then processed and cleaned using the program treenetproc in R (Knüsel et al., 2021). To examine the daily circadian rhythms of the trees, hourly stem radial increment values were calculated using the maximum hourly radius,

130 and then normalized across the 9 individuals. Hourly stem radius values were then averaged by month across all years to view the changes in daily shrinking and swelling patterns over the time period by month.

Soil temperature at each tree was recorded each hour using a combination of dataloggers (Tinytag-Plus-2, model TGP-4017, Gemini Dataloggers UK Ltd., Chichester, West Sussex, UK; EasyLog-USB, model EL-USB-1 PRO, Lascar Electronics Ltd.,
135 Whiteparish, Wiltshire, UK; Thermochron iButton, model DS-1922L, Dallas Semiconductors, Texas, USA). Data were downloaded and sensors replaced approximately every 6 months. Due to sensor malfunction, only 6 of the 9 trees had a complete T_{soil} time series for the period of record. Extraneous T_{soil} values (greater than 100°C away from 0°C) were removed.

2.3 Tree-ring width data and processing

140 In late October 2023, 20 mature trees were selected and 2 cores per tree were taken at 1.3 m using a Pressler increment borer. Cores were air dried in the laboratory, glued to wooden mounts, and sanded with progressively finer sandpapers for visualizing tree-ring boundaries (Fritz, Harold C, 1976). Then, they were scanned at 2400 dpi resolution and visually cross-dated under a stereoscope. Ring widths were measured with a 0.001 mm resolution using scanned images and the CooRecorder-CDendro software (Maxwell & Larsson, 2021). The quality of cross-dating was checked using the COFECHA software which calculates
145 moving correlations between individual series of ring-width values and the mean sites series (Holmes, 1983).

To calculate climate-growth relationships, first the individual ring-width series were detrended by fitting x -year cubic smoothing splines with a 50 % frequency response cut-off, where x was $2/3$ of the mean series length. The measured ring-width values were divided by fitted values. The resulting dimensionless ring-width indices were pre-whitened by fitting auto-
150 regressive models and were averaged using bi-weight robust means. This allowed building a mean series or chronology of ring-width indices preserving annual to decadal variability. Lastly, calculated over the common 1970–2023 period, an Expressed Population Signal (EPS) ≥ 0.85 indicated a high common signal of the chronology (Wigley et al., 1984). These are the statistics characterizing the chronology (values are means \pm SD): ring-width = 1.65 \pm 0.46 mm, first-order autocorrelation = 0.71 \pm 0.17 mm, mean sensitivity of standardized width indices = 0.24 \pm 0.05 mm, r_{bar} = 0.37. Lastly, Pearson correlations
155 were calculated from the prior September to the current September between climate data and the ring-width indices. Monthly climate data (T_x , mean maximum temperature; T_n , mean minimum temperature; Pr , precipitation) were obtained from the 0.1°-gridded E-OBS dataset v29.0e (Cornes et al., 2018) for the period 1970–2023. These procedures were done using the dplr (Bunn et al., 2023; Bunn, 2008, 2010) and treeclim (Zang & Biondi, 2015) R packages.

160 To verify that the patterns identified in the aforementioned methods were consistent with the Las Cutas experimental site, monthly correlations between climate variables (T_x , mean maximum temperature; T_n , mean minimum temperature; Pr , total precipitation) and the mean series of ring-width indices (period 1970-2023) were calculated using the Pearson method for the

September prior to the growing season of interest and up until the September after the growing season of interest. The results of that analysis were used to identify which climate variables during which months showed the highest correlation to growth. 165 Using that information, climate summaries from the months deemed influential were compared across years to identify possible interannual variability and test the hypothesis.

2.4 Relating climate and dendrometer data

Daily stem radial increment time series were calculated from the sub-hourly dendrometer data using the maximum daily radius following a daily approach (Deslauriers et al., 2007). The growing season was defined as day of year (DOY) 100 through 300 170 (Camarero et al., 1998; Sanmiguel-Valladolid et al., 2021). Using the daily radial increment data, the daily growth rate and TWD were extracted from the time series (Zweifel et al., 2016). The correlation between daily T_{air} and the growth rate was calculated during the growing period for values from the same day, as well as a lagged correlation between the growth rate and T_{air} from the previous day up to 10 days prior using the Pearson method. This correlation was also calculated between the SWC and TWD. In addition, 20-day moving correlations were calculated for daily T_{air} and the growth rate, daily T_{soil} and 175 growth rate, and SWC and TWD. This set of robust and simple statistics was useful to deal with non-parametric analyses and was selected due to the small sample size of trees.

With the complete T_{soil} data, daily temperature oscillation (DTO) was calculated by subtracting the daily minimum temperature from the daily maximum temperature. A novel approach was used to determine the time periods in which snow was present. 180 First, days with snow cover were identified using two different thresholds: a maximum DTO (DTOMax) of less than 2 degrees or less than 5 degrees. These two datasets for each of the 6 trees with T_{soil} data were used to define the snow cover duration periods. Snow cover was considered intermittent starting (IS) on the first date with snow presence and before the continuous (CS) period which was defined as the period with gaps less than 1 day. When gaps larger than 1 day began to occur again in the spring, the inconsistent period resumed until gaps with snow cover exceeded 15 days.

185 Because of the similar location, climatological conditions, and strong correlation between the T_{air} recorded at the Góriz refuge and the AWS at Las Cutas (Fig. A1), the calculated snow cover duration dates were compared with the Góriz refuge snow depth data. Although the refuge sits at an elevation approximately 100 meters higher than the Las Cutas average, the dates corresponded to accumulation and melting patterns identified in the Góriz snow depth data. The snow cover duration dates 190 with a DTOMax less than 2 were determined to be more accurate and were used for the rest of the analysis.


3 Results

Each year of the study period showed varying tree growth and shrinking and swelling dynamics. In 2021, the onset of growth and day of maximum growth rate were slightly delayed compared to the other two years in which the maximum growth rate

195 occurred 10 days earlier in 2022 and 15 days earlier in 2023, on average (Table 1). In addition, the length of the 2021 growing season was the shortest at an average of 83 days while the 2022 season was the longest with an average of 144 days (Table 1).

Table 1. Summary statistics of each growing season. All values are averages \pm SE. Different letters indicate significant ($p < 0.05$) differences between years according to t tests.

Variables	Year		
	2021	2022	2023
Growth start (DOY)	151 \pm 3a	143 \pm 2b	136 \pm 2c
Growth end (DOY)	249 \pm 2a	287 \pm 5b	244 \pm 2a
Growth length (days)	83 \pm 4a	144 \pm 5b	108 \pm 3c
Maximum growth rate ($\mu\text{m day}^{-1}$)	19.7 \pm 3.1	24.7 \pm 2.4	22.7 \pm 2.1
Date of maximum growth rate (DOY)	167 \pm 2a	157 \pm 3b	152 \pm 3b



200 **Figure 2. Daily values (means \pm SE) of radial stem variation (DMR) and minimum tree water deficit (TWD) in the three study years. DOY is the day of calendar year.**

The average DMR in 2022 and 2023 showed increased growth rates starting around DOY 130. The 2022 growing season was the most gainful (approximately 2300 μm) of the three examined, while the 2021 season showed the least total growth (approximately 1900 μm) (Fig. 2). The TWD also varied in timing between years. During the 2022 growing season, there were two large (greater than 60 μm) peaks before DOY 140 (Fig. 2). There was also one large peak in TWD in 2021 around DOY 140 and DOY 225 (Fig. 2).

TWD had high intra- and interannual variability throughout the time series (Fig. 2). There were small peaks in TWD (less than 100 μm) around DOY 145 and 130 in 2021 and 2022 respectively (Fig. 2). However, the largest peaks occurred in the late growing season between DOY 225 and 280 with 2023 demonstrating the largest peak (Fig. 2).




Figure 3. Climate-growth relationships (Pearson correlations) calculated by relating monthly climate variables (Tx, mean maximum temperature; Tn, mean minimum temperature; Pr, total precipitation) and the mean series of ring-width indices (period 1970–2023). Correlations were calculated from September of the previous year (months abbreviated by lowercase letters) to September of the current year (months abbreviated by uppercase letters). Dashed and dotted horizontal lines show the 0.05 and 0.01 significance levels, respectively.

Based on climate-growth correlations, mean maximum air temperature and mean minimum air temperature from the months of February and May showed the highest correlation with growth indices (Fig. 3). Thus, climate summaries were compared across years for these months and the overall growing season (Table 2).

Table 2. A summary of the climate variables at the study site and precipitation from the nearby Góriz station. Values are means \pm SE.

Climate variable	Time period	2021	2022	2023
Soil temperature (°C)	February	0.73 \pm 0.45	1.72 \pm 0.55	0.84 \pm 0.24
	May	5.88 \pm 0.27	8.77 \pm 0.79	7.12 \pm 0.57
	Growing season	9.89 \pm 0.52	10.54 \pm 0.86	11.45 \pm 0.66
Air temperature (°C)	February	1.14 \pm 0.47	2.5 \pm 0.57	0.06 \pm 0.82
	May	6.22 \pm 0.63	10.41 \pm 0.82	6.7 \pm 0.6
	Growing season	10.00 \pm 0.37	12.39 \pm 0.37	12.04 \pm 0.39
SWC (m ³ *m ⁻³)	February	0.20 \pm 0.01	—	0.25 \pm 0.01
	May	0.18 \pm 0.01	—	0.21 \pm 0.01
	Growing season	0.15 \pm 0.01	0.15 \pm 0	0.18 \pm 0.01
Precipitation (mm)	February	110.2	51.0	79.0
	May	104.4	21.6	118.0
	Growing season	684.4	944.7	1140.1

225

The average air temperatures in February, May and the fixed growing season (DOY 100-300) were highest in 2022, then 2023 (Table 2). For instance, in 2021, 2022 and 2023, growing-season T_{air} were 10.0, 12.4 and 12.0 °C, respectively. This was also true of the average soil temperatures except for the fixed growing season in 2023 which had average soil temperatures slightly less than 1 °C higher than in 2022 (Table 2). For instance, in 2021, 2022 and 2023, growing-season T_{soil} were 9.9, 10.5 and 11.4 °C, respectively. Due to SWC sensor malfunction in 2022, data are limited to the fixed growing season. The average SWC was slightly higher in February, May and the fixed growing season of 2023 than 2021 (Table 2). However, the SWC values were very similar despite the differences in T_{air} , T_{soil} and precipitation for those two years.


230

Table 3. Calculated snow season dates are included with IS meaning intermittent snow season and CS meaning continuous snow season.

Snow season index	2021	2022	2023
IS _{start} (DOY)	275 \pm 1	320 \pm 4	292 \pm 1
CS _{start} (DOY)	331 \pm 4	339 \pm 8	334 \pm 7
CS _{end} (DOY)	87 \pm 5	117 \pm 49	22 \pm 15
IS _{end} (DOY)	128 \pm 11	118 \pm 1	84 \pm 5
length of CS (days)	121 \pm 9	113 \pm 26	64 \pm 15
length of IS (days)	218 \pm 11	163 \pm 3	157 \pm 5
days w/ snow during IS (days)	156 \pm 13	119 \pm 13	106 \pm 11
% IS with snow	0.72 \pm 0.06	0.73 \pm 0.08	0.68 \pm 0

Precipitation from the Góriz refuge also varied highly between the first two years of this study. In 2021, the cumulative precipitation in February was 1102 mm (Table 2). This dropped to 51.0 mm in 2022, then rose in 2023 for a total of 79.0 mm (Table 2). Similar trends were observed in May as well (Table 2). However, the total precipitation during the entirety of each growing season was 684.4 mm, 944.7 mm, and 1140.1 mm for the years 2021, 2022, and 2023, respectively (Table 2).

240 Although snow presence was not directly recorded, days with snow were estimated using soil temperature. The longest consistent snow season occurred during the 2020-2021 winter, followed by 2021-2022, and, finally, 2022-2023 (Table 3). Unlike the two other years which appeared to have a longer period of inconsistent snow towards the end of the winter, the 2021-2022 winter snow cover ended abruptly around DOY 118 \pm 1, which was earlier than the 2020-2021 winter (DOY 128 \pm 11) and later than the 2022-2023 winter (DOY 84 \pm 5) (Table 3).



245

Figure 4. Moving correlations calculated by correlating air (T_{air}) or soil (T_{soil}) temperatures with daily growth rates along 20-day periods. Values are means \pm SE. Correlation values located outside the grey box are significant ($p < 0.05$).

The moving correlations between T_{air} or T_{soil} with the growth rate showed generally similar patterns for the fixed growing seasons during the years 2021 and 2022 (Fig. 4). Both years had a window with a peak in positive significance for both T_{air} and T_{soil} during May (Fig. 4). This pattern appears to have occurred earlier in 2023 with the peak in significance already beginning to decline by the start of May. The year 2023 also stands out as the year with the smallest difference between T_{air} and T_{soil} .




Figure 5. The hourly stem radius normalized between trees and averaged by month with each solid line representing an individual tree in the forest (TRA), transition to treeline (TRA-TRE) and treeline (TRE) zones.

The in-depth analysis of the seasonal changes in stem shrinking and swelling showed that the daily pattern shifts across the seasons (Fig. 5). The growing season pattern shows swelling during the evening and shrinking during the day beginning in July or August and ending in November (Fig. 5). However, the pattern that can be seen during the snow season (November or December through April) is inverted, meaning that it is often the reverse of the normal growing season daily pattern exhibiting expansion during the day and shrinking during the night (Fig. 5). The switch from the inverted pattern to the normal pattern occurred very rapidly during the months of May or June, as noted by the decoupling of the cycle between trees (Fig. 5).

4 Discussion and Conclusions


The onset of the growing season at this site is triggered by rising air and soil temperatures, and the disappearance of the winter snowpack which is consistent with previous studies conducted on the study species (Galván et al., 2014; Tardif et al., 2003). SWC was not identified as a strong indicator of length of the growing season or total growth. However, the data were limited due to the malfunctioning sensor during the late winter and onset of the growing season in 2022 which was a critical time

period in the study. While important, May air and soil temperatures were less significant when correlated to total growth than February air and soil temperatures. In winters like 2022 that are warmer in February, there is likely less precipitation falling as snow and accumulating which leads to an earlier melt out date, earlier warming of soil, and longer growing seasons resulting in more total growth. In a previous study, February was identified as an important month in the climate-growth relationships specifically between air temperature and radial growth (Sanmiguel-Vallelado et al., 2019). Additionally, researchers found that radial growth in *P. uncinata* was related to early and late growing season soil temperatures (Sanmiguel-Vallelado et al., 2021). They specifically found that May climatological conditions were more strongly correlated with growth than other months (Sanmiguel-Vallelado et al., 2021).

275

The high interannual variability of this region allows for the comparison of the impact of climatological differences on *P. uncinata* growth and TWD. Higher February air and soil temperatures in 2023 and especially in 2022 lead to an earlier growth onset, and a longer growing season which is consistent with the findings in (Sanmiguel-Vallelado et al., 2019)). In summer, the central Spanish Pyrenees become hotter and drier leading to high vapor pressure deficit (VPD) and increased TWD. Under this set of environmental conditions, trees tend to close its stomata to conserve water during the hottest hours of the day (Grossiord et al., 2020). For this reason, it is common for trees to grow at night and early morning during months when VPD and TWD during the day is too high (Tumajer et al., 2022; Zweifel et al., 2016). The normal growing season pattern was identified in previous research (Lázaro-Gimeno et al., 2024; Zweifel et al., 2016) and occurs during the warm months at Las Cutas, beginning in July or August and ending in October or November. This pattern could be attributed to normal responses to high VPD. During the cold months, when trees were not experiencing high VPD, and had much lower daily growth rates, radial increment swelling occurred during daylight hours. This switch in the swelling/shrinking cycle was attributed to the presence of snow which was influenced by winter air temperatures and led to cold and moist soil conditions.

285



290 **Figure 6. Graphic summarizing the patterns found in this study and showing the shifts in snow and climate conditions and swelling/shrinking stem cycles along the growing season. The rectangle around the trunk highlights that the tree is growing radially under the normal pattern and transitioning to a non-growing state during the transition phase.**

These findings could have implications for other high alpine treeline ecotones, especially in the Mediterranean region. Impacts of climate change like increasing winter and summer temperatures, and snow accumulation and meltout will likely alter the timing of the growing season and impact growth and productivity as clearly demonstrated by the year 2022 highlighted in this study (Table 1, Fig. 2). An overall increase in productivity could lead to greater carbon sequestration during the growing season, specifically in high elevation areas of the Pyrenees that were previously agricultural lands, now transitioning back to forests either naturally or through afforestation efforts (Khorchani et al., 2022). The growing season at Las Cutas is limited by cold temperatures and the presence of snow which causes the trees to remain in the inverted circadian pattern for longer (Fig. 300 6). However, in other more arid mountain regions, warmer and drier conditions may lead to a longer growing season but amplify drought stress which could limit carbon sequestration by reducing the rate of wood cell production (Ren et al., 2019).

It is important to acknowledge the limitations of this study. The sample size of the dendrometer data was small and the trees corresponded to a narrow range of ages, not allowing for widespread spatial and ontogenetic variability to be included. This is not ideal, especially in a treeline ecotone which experiences diverse and heterogeneous climate variability. Additionally, the presence of snow was determined using a novel method and a small sample size. However, overall snow accumulation patterns

were confirmed using data from the nearby Góriz refuge. There is likely some error associated with these data, but, for the purposes of this study, we believe the methods are useful and justified.

310 Future research could use in-situ and short-range remotely sensed snow observations to confirm that snow is the driver of the changes to the growing season and total growth that we saw in this study. Different locations of each tree (within the stand, outer edge, or krummholz) could also impact the timing of snow melt, the growth rate, total growth, TWD and circadian stem increment cycles. This is very relevant because the study treeline ecotone is rapidly encroaching which can be interpreted as a response to climate warming. In addition, dates of growth onset are inferred from dendrometer data, but they should be
315 confirmed using periodic wood monitoring and xylogenesis analyses (e.g. Sanmiguel-Vallelado et al., 2021). Unfortunately, we were limited by the number of sample trees in each zone to make robust conclusions, so a good starting point for another study would be to extend the tree-ring chronologies to discern if trees are turning more or less responsive to growing-season precipitation.

320 In agreement with prior studies, we observed that warmer February temperatures, specifically soil temperatures, enhanced overall growth and increased spring growth rates, and this positive effect was related to an earlier snow melt-out and, thus, a longer growing season. The characteristic circadian cycle of stem increment that leads to wood production, defined by night swelling and day shrinking, was detected in summer and autumn (approximately from July to November) during the growing season. However, this pattern was inverted during the snow season (approximately from November to April), prior to the onset
325 of growth (Fig. 6). Air temperature, soil temperature, and the presence of snow are strong indicators of how much and for how long trees can grow each year and cause the shift in daily stem increment patterns of tree radius.

Author contributions:

Conceptualization, J.J.C., A.S.V., H.F., J.I.L.M.; data curation and management, H.F., J.J.C., A.S.V., F.R.H.; formal analysis, H.F., J.J.C., A.S.V.; methodology, A.S.V., J.J.C, H.F., P.D.A.; resources, J.I.L.M., J.J.C., J.R.; writing - original draft
330 preparation, H.F.; writing - review and editing, H.F., A.S.V., J.J.C, J.I.L.M., J.R., F.R.H., P.D.A.; visualization, J.J.C., H.F., A.S.V.; project administration, J.I.L.M., J.J.C., J.R.

Funding:

This research was funded by a Fulbright Spain Predoctoral Fellowship to H.F. A.S.V. was supported by postdoctoral grant JDC2022-048316-I funded by MICIU/AEI/10.13039/501100011033 and by European Union NextGenerationEU/PRTR.

335 **Code Availability:**

The code used in the analysis of this research is available upon request from the first author.

Data Availability:

The data used in this research are available upon request from the first author.

Competing interests:

340 The authors declare that they have no conflicts of interest.

Acknowledgements:

The authors would like to thank the members of CryoPyr for carrying out the fieldwork, the Agencia Estatal de Meteorología of Spain for providing the Góriz refuge data, and the Pyrenees Institute of Ecology – Zaragoza for hosting H.F.

References

- 345 Albrich, K., Rammer, W., & Seidl, R. (2020). Climate change causes critical transitions and irreversible alterations of mountain forests. *Global Change Biology*, 26(7), 4013–4027.
<https://doi.org/10.1111/gcb.15118>
- Andrus, R. A., Harvey, B. J., Rodman, K. C., Hart, S. J., & Veblen, T. T. (2018). Moisture availability limits subalpine tree establishment. *Ecology*, 99(3), 567–575. <https://doi.org/10.1002/ecs.2134>
- 350 Bunn, A. G. (2008). A dendrochronology program library in R (dplR). *Dendrochronologia*, 26(2), 115–124. <https://doi.org/10.1016/j.dendro.2008.01.002>
- Bunn, A. G. (2010). Statistical and visual crossdating in R using the dplR library. *Dendrochronologia*, 28(4), 251–258. <https://doi.org/10.1016/j.dendro.2009.12.001>

- Bunn, A., Korpela, M., Biondi, F., Campelo, F., Mérian, P., Qeadan, F., & Zang, C. (2023). *dplR: dendrochronology program library in R* (Version R package version 1.7.6) [Computer software]. <https://CRAN.R-project.org/package=dplR>
- 355
- Camarero, J. J., Guerrero-Campo, J., & Gutiérrez, E. (1998). Tree-Ring Growth and Structure of *Pinus uncinata* and *Pinus sylvestris* in the Central Spanish Pyrenees. *Arctic and Alpine Research*, *30*(1), 1–10. <https://doi.org/10.2307/1551739>
- Cornes, R. C., van der Schrier, G., van den Besselaar, E. J. M., & Jones, P. D. (2018). An Ensemble Version of the E-OBS Temperature and Precipitation Data Sets. *Journal of Geophysical Research: Atmospheres*, *123*(17), 9391–9409. <https://doi.org/10.1029/2017JD028200>
- 360
- Deslauriers, A., Rossi, S., & Anfodillo, T. (2007). Dendrometer and intra-annual tree growth: What kind of information can be inferred? *Dendrochronologia*, *25*(2), 113–124. <https://doi.org/10.1016/j.dendro.2007.05.003>
- 365
- Dobbert, S., Pape, R., & Löffler, J. (2022). The application of dendrometers to alpine dwarf shrubs – a case study to investigate stem growth responses to environmental conditions. *Biogeosciences*, *19*(7), 1933–1958. <https://doi.org/10.5194/bg-19-1933-2022>
- Fritz, Harold C. (1976). *Tree rings and climate*. Academic Press Inc.
- 370
- Galván, J. D., Camarero, J. J., & Gutiérrez, E. (2014). Seeing the trees for the forest: Drivers of individual growth responses to climate in *Pinus uncinata* mountain forests. *Journal of Ecology*, *102*(5), 1244–1257. <https://doi.org/10.1111/1365-2745.12268>

- Gascoin, S., Hagolle, O., Huc, M., Jarlan, L., Dejoux, J.-F., Szczypta, C., Marti, R., & Sánchez, R. (2015). A snow cover climatology for the Pyrenees from MODIS snow products. *Hydrology and Earth System Sciences*, *19*(5), 2337–2351. <https://doi.org/10.5194/hess-19-2337-2015>
- 375
- Grossiord, C., Ulrich, D. E. M., & Vilagrosa, A. (2020). Controls of the hydraulic safety–efficiency trade-off. *Tree Physiology*, *40*(5), 573–576. <https://doi.org/10.1093/treephys/tpaa013>
- Hagedorn, F., Shiyatov, S. G., Mazepa, V. S., Devi, N. M., Grigor’ev, A. A., Bartysh, A. A., Fomin, V. V., Kapralov, D. S., Terent’ev, M., Bugman, H., Rigling, A., & Moiseev, P. A. (2014). Treeline advances along the Urals mountain range – driven by improved winter conditions? *Global Change Biology*, *20*(11), 3530–3543. <https://doi.org/10.1111/gcb.12613>
- 380
- Harpold, A. A., & Molotch, N. P. (2015). Sensitivity of soil water availability to changing snowmelt timing in the western U.S. *Geophysical Research Letters*, *42*(19), 8011–8020. <https://doi.org/10.1002/2015GL065855>
- 385
- Hedstrom, N. R., & Pomeroy, J. W. (1998). Measurements and modelling of snow interception in the boreal forest. *Hydrological Processes*, *12*(10–11), 1611–1625. [https://doi.org/10.1002/\(SICI\)1099-1085\(199808/09\)12:10/11<1611::AID-HYP684>3.0.CO;2-4](https://doi.org/10.1002/(SICI)1099-1085(199808/09)12:10/11<1611::AID-HYP684>3.0.CO;2-4)
- Holmes, R. L. (1983). *COMPUTER -ASSISTED QUALITY CONTROL IN TREE -RING DATING AND MEASUREMENT*.
- 390
- Huang, M., Wang, G., Bie, X., Jiang, Y., Huang, X., Li, J.-J., Shi, S., Zhang, T., & Peng, P.-H. (2023). Seasonal snow cover patterns explain alpine treeline elevation better than temperature at regional scale. *Forest Ecosystems*, *10*, 100106. <https://doi.org/10.1016/j.fecs.2023.100106>

- Khorchani, M., Nadal-Romero, E., Lasanta, T., & Tague, C. (2022). Carbon sequestration and water yield tradeoffs following restoration of abandoned agricultural lands in Mediterranean mountains. *Environmental Research*, 207, 112203. <https://doi.org/10.1016/j.envres.2021.112203>
- 395 Knüsel, S., Peters, R. L., Haeni, M., Wilhelm, M., & Zweifel, R. (2021). Processing and Extraction of Seasonal Tree Physiological Parameters from Stem Radius Time Series. *Forests*, 12(6), Article 6. <https://doi.org/10.3390/f12060765>
- Lázaro-Gimeno, D., Ferrari, C., Delhomme, N., Johansson, M., Sjölander, J., Singh, R. K., Mutwil, M., & Eriksson, M. E. (2024). The circadian clock participates in seasonal growth in Norway spruce (Picea abies). *Tree Physiology*, 44(11), tpae139. <https://doi.org/10.1093/treephys/tpae139>
- 400 López-Moreno, J. I. (2005). Recent Variations of Snowpack Depth in the Central Spanish Pyrenees. *Arctic, Antarctic, and Alpine Research*, 37(2), 253–260. [https://doi.org/10.1657/1523-0430\(2005\)037\[0253:RVOSDI\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2005)037[0253:RVOSDI]2.0.CO;2)
- 405 López-Moreno, J. I., & Latron, J. (2008). Influence of canopy density on snow distribution in a temperate mountain range. *Hydrological Processes*, 22(1), 117–126. <https://doi.org/10.1002/hyp.6572>
- López-Moreno, J. I., Soubeyroux, J. M., Gascoin, S., Alonso-Gonzalez, E., Durán-Gómez, N., Lafaysse, M., Vernay, M., Carmagnola, C., & Morin, S. (2020). Long-term trends (1958–2017) in snow cover duration and depth in the Pyrenees. *International Journal of Climatology*, 40(14), 6122–6136. <https://doi.org/10.1002/joc.6571>
- 410 Lüttge, U., & Hertel, B. (2009). Diurnal and annual rhythms in trees. *Trees*, 23(4), 683–700. <https://doi.org/10.1007/s00468-009-0324-1>

- Maxwell, R. S., & Larsson, L.-A. (2021). Measuring tree-ring widths using the CooRecorder software
415 application. *Dendrochronologia*, 67, 125841. <https://doi.org/10.1016/j.dendro.2021.125841>
- Mei-Jun, L. I. U., Qiu-Wen, C., Jin-Lin, L., Guo-Qing, L. I., & Sheng, D. U. (2023). Seasonal dynamics
of radial growth and micro-variation in stems of *Quercus mongolica* var. *Liaotungensis* and
Robinia pseudoacacia in loess hilly region. *Chinese Journal of Plant Ecology*, 47(2), 227.
<https://doi.org/10.17521/cjpe.2022.0100>
- 420 Noguera, I., Vicente-Serrano, S. M., Peña-Angulo, D., Domínguez-Castro, F., Juez, C., Tomás-
Burguera, M., Lorenzo-Lacruz, J., Azorin-Molina, C., Halifa-Marín, A., Fernández-Duque, B.,
& El Kenawy, A. (2023). Assessment of vapor pressure deficit variability and trends in Spain
and possible connections with soil moisture. *Atmospheric Research*, 285, 106666.
<https://doi.org/10.1016/j.atmosres.2023.106666>
- 425 Pardo, I., Camarero, J. J., Gutiérrez, E., & García, M. B. (2013). Uncoupled changes in tree cover and
field layer vegetation at two Pyrenean treeline ecotones over 11 years. *Plant Ecology &
Diversity*, 6(3–4), 355–364. <https://doi.org/10.1080/17550874.2013.811306>
- Pepin, N., Bradley, R. S., Diaz, H. F., Baraer, M., Caceres, E. B., Forsythe, N., Fowler, H., Greenwood,
G., Hashmi, M. Z., Liu, X. D., Miller, J. R., Ning, L., Ohmura, A., Palazzi, E., Rangwala, I.,
430 Schöner, W., Severskiy, I., Shahgedanova, M., Wang, M. B., ... Mountain Research Initiative
EDW Working Group. (2015). Elevation-dependent warming in mountain regions of the world.
Nature Climate Change, 5(5), 424–430. <https://doi.org/10.1038/nclimate2563>

- Peterson, D. L. (1998). Climate, limiting factors and environmental change in high-altitude forests of Western North America. In M. Beniston & J. L. Innes (Eds.), *The Impacts of Climate Variability on Forests* (pp. 191–208). Springer. <https://doi.org/10.1007/BFb0009773>
- 435
- Pomeroy, J. W., Bewley, D. S., Essery, R. L. H., Hedstrom, N. R., Link, T., Granger, R. J., Sicart, J. E., Ellis, C. R., & Janowicz, J. R. (2006). Shrub tundra snowmelt. *Hydrological Processes*, 20(4), 923–941. <https://doi.org/10.1002/hyp.6124>
- Pomeroy, J. W., Gray, D. M., Hedstrom, N. R., & Janowicz, J. R. (2002). Prediction of seasonal snow accumulation in cold climate forests. *Hydrological Processes*, 16(18), 3543–3558.
- 440 <https://doi.org/10.1002/hyp.1228>
- Ren, P., Ziaco, E., Rossi, S., Biondi, F., Prislán, P., & Liang, E. (2019). Growth rate rather than growing season length determines wood biomass in dry environments. *Agricultural and Forest Meteorology*, 271, 46–53. <https://doi.org/10.1016/j.agrformet.2019.02.031>
- 445 Revuelto, J., Azorin-Molina, C., Alonso-González, E., Sanmiguel-Vallelado, A., Navarro-Serrano, F., Rico, I., & López-Moreno, J. I. (2017). Meteorological and snow distribution data in the Izas Experimental Catchment (Spanish Pyrenees) from 2011 to 2017. *Earth System Science Data*, 9(2), 993–1005. <https://doi.org/10.5194/essd-9-993-2017>
- Revuelto, J., López-Moreno, J. I., Azorin-Molina, C., & Vicente-Serrano, S. M. (2015). Canopy influence on snow depth distribution in a pine stand determined from terrestrial laser data. *Water Resources Research*, 51(5), 3476–3489. <https://doi.org/10.1002/2014WR016496>
- 450

- Saavedra, F. A., Kampf, S. K., Fassnacht, S. R., & Sibold, J. S. (2018). Changes in Andes snow cover from MODIS data, 2000–2016. *The Cryosphere*, *12*(3), 1027–1046. <https://doi.org/10.5194/tc-12-1027-2018>
- 455 Sanmiguel-Valladolid, A., Camarero, J. J., Gazol, A., Morán-Tejeda, E., Sangüesa-Barreda, G., Alonso-González, E., Gutiérrez, E., Alla, A. Q., Galván, J. D., & López-Moreno, J. I. (2019). Detecting snow-related signals in radial growth of *Pinus uncinata* mountain forests. *Dendrochronologia*, *57*, 125622. <https://doi.org/10.1016/j.dendro.2019.125622>
- Sanmiguel-Valladolid, A., Camarero, J. J., Morán-Tejeda, E., Gazol, A., Colangelo, M., Alonso-
460 González, E., & López-Moreno, J. I. (2021). Snow dynamics influence tree growth by controlling soil temperature in mountain pine forests. *Agricultural and Forest Meteorology*, *296*, 108205. <https://doi.org/10.1016/j.agrformet.2020.108205>
- Serrano-Notivoli, R., Tejedor, E., Sarricolea, P., Meseguer-Ruiz, O., de Luis, M., Saz, M. Á., Longares, L. A., & Olcina, J. (2023). Unprecedented warmth: A look at Spain’s exceptional summer of
465 2022. *Atmospheric Research*, *293*, 106931. <https://doi.org/10.1016/j.atmosres.2023.106931>
- Storck, P., Lettenmaier, D. P., & Bolton, S. M. (2002). Measurement of snow interception and canopy effects on snow accumulation and melt in a mountainous maritime climate, Oregon, United States. *Water Resources Research*, *38*(11), 5-1-5–16. <https://doi.org/10.1029/2002WR001281>
- Tardif, J., Camarero, J. J., Ribas, M., & Gutiérrez, E. (2003). Spatiotemporal Variability in Tree Growth
470 in the Central Pyrenees: Climatic and Site Influences. *Ecological Monographs*, *73*(2), 241–257. [https://doi.org/10.1890/0012-9615\(2003\)073\[0241:SVITGI\]2.0.CO;2](https://doi.org/10.1890/0012-9615(2003)073[0241:SVITGI]2.0.CO;2)

- Tumajer, J., Scharnweber, T., Smiljanic, M., & Wilmking, M. (2022). Limitation by vapour pressure deficit shapes different intra-annual growth patterns of diffuse- and ring-porous temperate broadleaves. *New Phytologist*, 233(6), 2429–2441. <https://doi.org/10.1111/nph.17952>
- 475 Vicente-Serrano, S. M., Peña-Angulo, D., Murphy, C., López-Moreno, J. I., Tomas-Burguera, M., Domínguez-Castro, F., Tian, F., Eklundh, L., Cai, Z., Alvarez-Farizo, B., Noguera, I., Camarero, J. J., Sánchez-Salguero, R., Gazol, A., Grainger, S., Conradt, T., Boincean, B., & El Kenawy, A. (2021). The complex multi-sectoral impacts of drought: Evidence from a mountainous basin in the Central Spanish Pyrenees. *Science of The Total Environment*, 769, 144702. <https://doi.org/10.1016/j.scitotenv.2020.144702>
- 480
- Villalba, R., Veblen, T. T., & Ogden, J. (1994). Climatic Influences on the Growth of Subalpine Trees in the Colorado Front Range. *Ecology*, 75(5), 1450–1462. <https://doi.org/10.2307/1937468>
- Wigley, T. M. L., Briffa, K. R., & Jones, P. D. (1984). *On the Average Value of Correlated Time Series, with Applications in Dendroclimatology and Hydrometeorology*. https://journals.ametsoc.org/view/journals/apme/23/2/1520-0450_1984_023_0201_otavoc_2_0_co_2.xml
- 485
- Yang, J. C., Bowling, D. R., Smith, K. R., Kunik, L., Raczka, B., Anderegg, W. R. L., Bahn, M., Blanken, P. D., Richardson, A. D., Burns, S. P., Bohrer, G., Desai, A. R., Arain, M. A., Staebler, R. M., Ouimette, A. P., Munger, J. W., & Litvak, M. E. (2024). Forest carbon uptake as influenced by snowpack and length of photosynthesis season in seasonally snow-covered forests of North America. *Agricultural and Forest Meteorology*, 353, 110054. <https://doi.org/10.1016/j.agrformet.2024.110054>
- 490

- Zang, C., & Biondi, F. (2015). *treeclim: An R package for the numerical calibration of proxy-climate relationships* (38:431-436; Version 2.0.0) [Computer software]. *Ecography*.
- 495 Ziaco, E., & Biondi, F. (2018). Stem Circadian Phenology of Four Pine Species in Naturally Contrasting Climates from Sky-Island Forests of the Western USA. *Forests*, 9(7), Article 7. <https://doi.org/10.3390/f9070396>
- Zweifel, R., Haeni, M., Buchmann, N., & Eugster, W. (2016). Are trees able to grow in periods of stem shrinkage? *New Phytologist*, 211(3), 839–849. <https://doi.org/10.1111/nph.13995>
- 500 Zweifel, R., Sterck, F., Braun, S., Buchmann, N., Eugster, W., Gessler, A., Häni, M., Peters, R. L., Walthert, L., Wilhelm, M., Ziemińska, K., & Etzold, S. (2021). Why trees grow at night. *New Phytologist*, 231(6), 2174–2185. <https://doi.org/10.1111/nph.17552>

Appendix:

505 **Table A1. Tree features (DBH, height, location).**

Tree N°	DBH (cm)	Height (m)	Location	Location
311	18.9	14.6	TRA	Forest
326	19.0	15.0	TRA	Forest
306	14.0	13.2	TRA	Forest
316	18.4	14.0	TRA	Forest
314	14.7	11.0	TRA-TRE	Transition
304	12.2	8.5	TRA-TRE	Transition
798	12.7	9.0	TRA-TRE	Transition
799	20.0	6.0	TRE	Treeline

800

8.6

2.8

TRE

Treeline

Figure A1. a) Maximum and b) minimum daily temperatures at Las Cutas (treeline site) and the Góriz refuge (local station).

