

1 Assessing evapotranspiration dynamics across central Europe in the 2 context of land-atmosphere drivers

3 Anke Fluhrer^{1,2}, Martin J. Baur³, María Piles⁴, Bagher Bayat⁵, Mehdi Rahmati⁵, David Chaparro^{1,6},
4 Clémence Dubois^{7,8}, Florian M. Hellwig^{1,7}, Carsten Montzka⁵, Angelika Kübert⁹, Marlin M. Mueller^{7,8},
5 Isabel Augscheller¹, Francois Jonard¹⁰, Konstantin Schellenberg^{7,11}, Thomas Jagdhuber^{1,2}

6 1 Microwaves and Radar Institute (HR), German Aerospace Center (DLR), Wessling, Germany

7 2 Institute of Geography, University of Augsburg, Augsburg, Germany

8 3 Department of Geography, Cambridge University, Cambridge, UK

9 4 Image Processing Laboratory, University of Valencia, Valencia, Spain

10 5 Institute of Bio-and Geosciences: Agrosphere (IBG-3), Forschungszentrum Jülich, Jülich, Germany

11 6 Centre for Ecological Research and Forestry Applications (CREAF), Cerdanyola del Vallès, Spain.

12 7 Department for Earth Observation, Friedrich Schiller University Jena, Jena, Germany

13 8 Institute of Data Science (DW), German Aerospace Center (DLR), Jena, Germany

14 9 Institute for Atmospheric and Earth System Research (INAR)/Physics, University of Helsinki, Helsinki, Finland

15 10 Earth Observation and Ecosystem Modelling Laboratory, University of Liège, Liège, Belgium

16 11 Department of Biogeochemical Processes, Max-Planck Institute for Biogeochemistry, Jena, Germany

17 *Correspondence to:* Anke Fluhrer (Anke.Fluhrer@dlr.de)

18 *Keywords:* ICOS, Eddy-covariance, MODIS, SEVIRI, ERA5-land, GLDAS-2, GLEAM, soil moisture, vapor pressure deficit,
19 extended triple collocation, error cross-correlation, anomaly, binning

20 Abstract.

21 Evapotranspiration (ET) is an important variable for analysing ecosystems, biophysical processes, and drought-related changes
22 in the soil-plant-atmosphere system. In this study, we evaluated freely available ET products from satellite remote sensing
23 (i.e., MODIS, SEVIRI, and GLEAM) as well as modelling and reanalysis (i.e., ERA5-land and GLDAS-2) together with in-
24 situ observations at eight Integrated Carbon Observation System (ICOS) stations across central Europe between 2017 and
25 2020. The land cover at the selected ICOS stations ranged from deciduous broad-leaved, evergreen needle-leaved, and mixed
26 forests to agriculture. Trends in ET were analysed together with soil moisture (SM) and water vapor pressure deficit (VPD)
27 during four years including a severe summer drought in 2018, but contrasting wet conditions in 2017. The analyses revealed
28 the increased atmospheric aridity and decreased water supply for plant transpiration under drought conditions, showing that
29 ET was generally lower and VPD higher in 2018 compared to 2017. Across the study period, results indicate that during
30 moisture limited drought years, ET is strongly decreasing due to decreasing SM and increasing VPD. However, during normal
31 or rather wet years, when SM is not limited, ET is mainly controlled by VPD, and hence, the atmospheric demand.
32 The comparison of the different ET products based on time series, statistics, and extended triple collocation (ETC) shows in
33 general a good agreement with ETC correlations between 0.39 and 0.99 as well as root-mean-square errors lower than 1.07

34 mm/day. The greatest deviations are found at the agricultural-managed sites Selhausen (Germany) and Bilos (France), with
35 the former also showing the highest potential dependencies (error cross-correlation (ECC)) between the ET products (up to
36 7.6 and outside the acceptable range of $-0.5 < ECC < 0.5$). Hence, Our results indicate that ET products differ most at stations
37 with spatio-temporal varying land cover conditions (varying crops over growing periods and between seasons). This is because
38 complex heterogeneity in land cover complicates the estimation of ET, while ET products agree well at evergreen needle-
39 leaved stations with less temporal changes throughout the year and between years. The ET products from SEVIRI, ERA5-
40 land, and GLEAM performed best when compared to ICOS observations with either lowest errors or highest correlations.

41 1 Introduction

42 Land-atmosphere dynamics and interactions are of key importance for understanding exchange processes in the global water,
43 energy, and carbon cycles (Zhou et al., 2016). For a holistic and well-founded ecosystem survey, the uptake, consumption, and
44 release of matter and energy need to be monitored. Especially in times of climate change, availability of terrestrial water,
45 agricultural productivity assuring food security, as well as forest health guaranteeing, for instance, carbon uptake and
46 biodiversity preservation, are mainly monitored by soil moisture (SM) and water vapor pressure deficit (VPD; as measure for
47 atmospheric aridity) (Novick et al., 2016; Zhou et al., 2019; Liu et al., 2020). Many studies focus on these two variables when
48 analysing drought-related terrestrial ecosystem productivity and its spatio-temporal changes (Fu et al., 2022; Zhang et al.,
49 2021). Evapotranspiration (ET) is an important proxy for analysing water stress and its effects on ecosystems since
50 precipitation (P) and evaporation are the two key components of the global water cycle (Miralles et al., 2011). ~~Since~~
51 ~~precipitation (P) and evaporation are the two key components of the global water cycle~~ (Miralles et al., 2011), ~~another~~
52 ~~important proxy for analysing water stress and its effects on ecosystems is evapotranspiration (ET)~~. As the sum of evaporation
53 from land, vegetation, and water surfaces as well as transpiration from vegetation, ET directly links the terrestrial energy,
54 water, and carbon cycles (Zhang et al., 2016; Zhou et al., 2016), and integrates meteorological conditions along SM (Bayat et
55 al., 2022). Hence, ET is an important variable for quantifying biophysical processes, ecosystem functioning, land surface
56 energy and water budgets, as well as improving weather and climate model predictions (Bayat et al., 2024; Zhang et al., 2016;
57 Zhou et al., 2016). For example, Zhou et al., (2019) reported negative SM-VPD coupling, meaning low SM and high VPD,
58 due to land-atmosphere feedbacks, since high VPD stimulates ET, which reduces SM. Although there is a debate that ET alone
59 does not determine SM, and other factors such as precipitation should also be considered, as reduced P for constant ET can
60 lead to lower SM (Rahmati et al., 2023), ET should in any case be one of the essential variables to inform about ecosystem-
61 atmosphere dynamics and interactions along with SM and VPD (Bayat et al., 2021).

62 ET is controlled by biological (e.g., plant growth and plant stomatal regulation) and physical (e.g., temperature) processes. For
63 example, vegetation controls interannual changes and affects spatio-temporal patterns and trends in ET (Zhang et al., 2016).
64 ET can be theoretically linked to the independent physical control factors demand (humidity, temperature) and supply
65 (precipitation). Depending on environmental and meteorological conditions, ET is primarily influenced by one of these three

66 factors. For instance, across central Europe, ET is mainly driven by the available energy due to reduced solar radiation during
67 cloudy skies (Zhang et al., 2016). However, Seneviratne et al., (2010) stated that decreasing SM leads to decreasing ET due to
68 the less accessible SM for plant water uptake and increasing soil suction.

69 During summer 2018, Europe experienced an unprecedented drought event comparable to previous extreme droughts, such as
70 in 2003 and 2010, with a temperature anomaly of +2.8 K (Rakovec et al., 2022) and an abnormally reduced SM and increased
71 VPD (Fu et al., 2022). This extreme drought was characterized by a unique geographical distribution, focused on regions at
72 higher latitudes (central and northern Europe), a rapid change from a wet spring to a dry summer, and an intense heatwave in
73 the spring of 2018 (Bastos et al., 2020). As a result, it caused severe tree stress in central Europe, with low leaf water potential,
74 leaf discolouration, and premature shedding, leading to significant tree mortality and heavy drought-legacy effects in 2019,
75 leaving trees vulnerable to further damage from pests and pathogens (Schuldt et al., 2020).

76 The significance of ET can also be seen in relation to the precise parametrization of SM and its memory in Land Surface
77 Models (LSMs) (Rahmati et al., 2024). Due to its importance and influence on the entire soil-plant-atmosphere system (SPAS),
78 tracking ET in time and space, meaning ~~at-from~~ seasonal to multi-year scales and for wide areas, is necessary and calls for a
79 satellite remote sensing approach (complementary to current modelling and reanalysis approaches). Although it is not directly
80 measurable from remote sensing acquisitions, optical, thermal, infrared, or microwave observations are used to derive ET
81 based on surface energy balance, physical and empirical models (Zhang et al., 2016; Rahmati et al., 2020; Singh et al., 2020;
82 Bayat et al., 2021; Bhattacharya et al., 2022; Bayat et al., 2024). Still, research comparing the performance of remote sensing
83 with model and reanalysis data under drought conditions is lacking, and an analysis on how main ET drivers (SM and VPD)
84 impact these ET products is also needed. Bridging this gap is paramount to assess which products and in which conditions are
85 more suitable to track ET, especially under the increasingly frequency and severity of droughts due to climate change.

86 Several ~~sub-global~~regional studies exist for comparing various ET products, e.g., over China (Meng et al., 2024; Xu et al.,
87 2024), across the U.S. (Carter et al., 2018; Xu et al., 2019), over Africa (Trambauer et al., 2014), and across Europe (Ahmed
88 et al., 2020; Stisen et al., 2021). However, due to the complexity of ecosystems, findings from specific regions (e.g., China,
89 U.S., Africa) cannot be generalized for other regions (e.g., Europe). Further, European studies focused either only on spatial
90 product comparisons, evaluating the performance of hydrological models (e.g., Stisen et al., 2021), on former time periods
91 (e.g., 2003-2013) at basin scale (Liu et al., 2023), on analysing drought impacts on ET dynamics using solely a single ET
92 product (e.g., Sepulcre-Canto et al., 2014; Ahmed et al., 2020), and on evaluating new ET products (e.g., Hu et al., 2023). For
93 example, the focus in the study of Stisen et al., was the evaluation of the spatial pattern performance in different hydrological
94 models for ET estimation. For this, four remote sensing based ET products were inter-compared for years among each other
95 ~~between~~ 2002-2014, and they found high agreements in spatial patterns across continental Europe (Stisen et al., 2021). Further,
96 Ahmed et al., investigated the drought impact of 2018 on the MODerate Resolution Imaging Spectroradiometer (MODIS) ET
97 across European ecosystems and found that ET decreased up to 50% compared to a 10-year reference period, with agricultural
98 areas and mixed natural vegetation being most affected (Ahmed et al., 2020). However, there is a lack of studies comparing
99 various ET products among each other and with in-situ measurements across central Europe, especially during severe drought

100 years (e.g., 2018), as well as evaluating the potential of remote sensing for tracking seasonal ET dynamics. ~~But~~ The evaluation
101 of the varying employed retrieval techniques (e.g., eddy covariance, land surface schemes, Penman-Monteith equation) of
102 commonly used ET products ~~and how well these techniques perform~~ under drought conditions is paramount in order to assess
103 whether they capture ET dynamics correctly.
104 In this study, we first compare the most common ET products from field measurements, modelling, and remote sensing across
105 central Europe for the period 2017 to 2020. These selected six products (ICOS, MODIS, SEVIRI, ERA5-land, GLDAS-2, and
106 GLEAM) are well-known, commonly employed, and freely available. The focus hereby is on the evaluation and quality
107 assessment of the individual products regarding the estimation of absolute ET values and their time-dynamics. Second, we
108 compare ET products in the context of SM and VPD, disentangling the relative role of all three variables within the SPAS
109 under severe drought conditions in 2018 in comparison to the rather wet year 2017. This is to analyse how the ET products
110 catch drought conditions and to what extent they can be used as indicator for drought events.

111 2 Materials and Methods

112 2.1 Study area

113 The focus is on eight Integrated Carbon Observation System (ICOS) (Rebmann et al., 2018) stations within central Europe
114 between 2017 and 2020, where field-scale in-situ eddy-covariance (EC) ET measurements are available (see Fig. 1).





Figure 1: Location of the eight investigated Integrated Carbon Observation System (ICOS) stations in central Europe, and their classification according to the respective dominant land cover class. DBF = deciduous broad-leaved, ENF = evergreen needle-leaved, MF = mixed forest.

The study comprises two deciduous broad-leaved (DBF) — the German Hohes Holz (DE-HoH) and Hainich (DE-Hai), two evergreen needle-leaved (ENF) — the German Wuestebach (DE-Ruw) and Finnish Lettosuo (FI-Let), and two mixed forest (MF) stations — the Czech Landzhot (CZ-Lnz) and the Swiss Laegern (CH-Lae), as well as two agriculture stations — the German Selhausen (DE-Rus) and the French Bilos (FR-Bil). [Details regarding coordinates, altitude, and climate zone for every station are given in table S1.](#) At every station, a footprint of 3 km radius is analysed to account for differences in spatial resolutions among employed datasets (see Sec. 2.2 ~~and Tab. 1~~). As displayed in Figure 2 and Table [S1-S2](#) (supplement), the land cover types and their homogeneity within the 3 km × 3 km footprint around every station was analysed based on the Corine land cover (CLC) 2018 classification from the Copernicus Land Monitoring Service at 100 m spatial resolution (European Environment Agency, 2019).



128


129 Figure 2: Overview of land cover classes according to the Corine Land Cover (CLC) 2018 (European Environment Agency, 2019) within
130 the 3 km × 3 km footprint around every investigated ICOS station. Percentages inside the circles indicate the dominant land cover classes,
131 respectively. The percentages of all land cover classes at every station can be found in the supplement (see Tab. S4S2).

132 According to this classification, two stations can be considered as homogeneous with one dominant land cover class, i.e., 86.7
133 % of coniferous forest at DE-Ruw, and 82.4 % of broad-leaved forest at DE-Hai. Station DE-Rus is mainly (63.1 %) covered
134 by non-irrigated arable land. Further, two stations show a two-part split land cover with two almost equally dominant classes.
135 At DE-HoH, 45.6 % are covered by non-irrigated arable land and 45.5 % are covered by broad-leaved forest. At FR-Bil,
136 although it is officially labelled as ENF station, 44.4 % are covered by transitional woodland shrub, while 41.4 % are covered
137 by coniferous forest, a managed Pine forest plantation (Loustau et al., 2022). Hence, due to this heterogeneity and the fact that
138 14.2 % of non-irrigated arable land (see Tab. S4S2) are mostly directly located near the station (see Fig. 2), we ranked it as
139 agricultural station in order to account for the frequently changing land cover conditions and spatial heterogeneity. All other
140 stations are rather heterogeneous with a mix of more than two different land cover classes (see Tab. S4S2 and Fig. 2). However,
141 it is worth noting that the CLC 2018 classification is based on data from 2017 to 2018. Hence, changes in the land cover, e.g.,
142 such as differences between summer and winter months, deforestation, weather extremes (storms, floods), or varying
143 agricultural crop cultivation, at each station between 2017 to 2020 are not included here.

Figure 3 illustrates the meteorological conditions (precipitation P and air temperature T_{Air}) at every station during the investigation period. The mean annual P and T_{Air} values are given-summarized in table S1. Note that the in-situ P measurements contain missing values at stations DE-HoH, CZ-Lnz, and CH-Lae in 2020. The overall lowest T_{Air} is found at the northernmost ICOS station FI-Let, varying between -12.6 °C (absolute minimum) and 22.75 °C (absolute maximum) in the years 2017 to 2020, with an interannual average of 5.67 °C. In contrast, the highest average T_{Air} (between 2017 and 2020) of 14.1 °C is found at the southernmost ICOS station FR-Bil, which also has the highest average P value of 3.04 mm/day. The lowest P is found at DE-HoH with an average of 1.26 mm/day, which is similar to the other stations in the mid-latitudes (see Tab. S1). The overall highest T_{Air} and lowest P at every station are always found in 2018 with an average of 1.7°C higher T_{Air} and annual 0.76 mm higher P, compared to the second hottest and driest year in each case. Exceptions can be found at the station FR-Bil, where the highest T_{Air} are recorded in 2019 and lowest P in 2017, and DE-Ruw, as well as CH-Lae, where the lowest average annual P are recorded in 2020, respectively.

Based on the standardized precipitation-evapotranspiration index (SPEI) (Beguería et al., 2023) (see Fig. S1), which describes drought based on the amount and duration of water deficit (Yu et al., 2023), distinctly dry and wet years are identified for each ICOS station. While all stations show abnormally dry periods, especially for 2018, only stations FI-Let and FR-Bil show abnormally wet periods at the end of 2017 and 2019. These two are the northernmost and southernmost stations (see Fig. 1).

The choice of SPEI to identify drought conditions instead of the standardized precipitation index (SPI) or other indices (i.e., Palmer drought severity index) is due to the fact that the SPEI considers implicitly temporal changes in ET and hence, temperature, which is relevant for identifying abnormal (drought) conditions and for this study with focus on ET variations. Previous studies showed that not only the lack of precipitation defines drought events but also the level of temperature and consumption of rainfall by evaporation and/or transpiration (Vicente-Serrano et al., 2010).



164

165 Figure 3: Daily in-situ measured precipitation (P) [mm/day] and air temperature (T_{Air}) [°C] at investigated ICOS stations. T_{Air} was cleaned
 166 for daily and weekly dynamics using a Savitzky-Golay (Savitzky and Golay, 1964) filter with a window size of 31 days.

167 2.2 Data base

168 In the first part of this study, different ET products (see Tab. 1) are inter-compared in order to evaluate the potential of remote
 169 sensing for tracking seasonal ET dynamics. The in-situ ET data, recorded at the ICOS stations at field-scale, are mass balance-
 170 based measurements of sensible heat (H) and latent heat (LE) fluxes through the covariance of heat and moisture fluxes,
 171 respectively. The LE [W/m²] can then be converted to ET by dividing it by the latent heat of vaporization (2.434 [MJ/kg] at
 172 20 °C air temperature) (Allen et al., 1998). The ICOS network ~~has undertaken~~has undertaken significant a large efforts to
 173 ensure consistent high-quality LE measurements, ~~which are comparable among different~~across ICOS-stations (Rebmann et
 174 al., 2018). Besides in-situ ~~EC~~-ET measurements, we employ some of the most commonly employed optical/thermal remote
 175 sensing products from NASA's (National Aeronautics and Space Administration) Moderate-resolution Imaging
 176 Spectroradiometer (MODIS) sensor on Terra (Running et al., 2017), ESA's Spinning Enhanced Visible and Infrared Imager

(SEVIRI) sensor onboard of the Meteosat Second Generation (MSG) satellites, and the Global Land Evaporation Amsterdam Model (GLEAM) (Martens et al., 2017). Further, also well-known reanalysis and modelling products from the land component of the Earth system modelling product European Re-Analysis (ERA5-land) from the European Centre for Medium-Range Weather Forecasts (ECMWF) (Muñoz Sabater, 2019), and from NASA’s Global Land Data Assimilation System Version 2 (GLDAS-2) (Beaudoing et al., 2020) are used (see Tab. 1). It should be noted that the GLEAM product is based on various remote sensing observations and reanalysis datasets from, e.g., NASA’s SMOS (soil moisture and ocean salinity) mission, MODIS, GLDAS-Noah, and ERA-Interim (Martens et al., 2017). The MODIS product with nominal spatial resolution of 500 m is aggregated to the 3 km footprint, while the SEVIRI, ERA5-land, GLDAS-2, and GLEAM products are maintained at their original spatial resolutions of 3 km, 9 km and 25 km, respectively. ~~All datasets are temporally aggregated to daily time series.~~ Although several downscaling methods and data fusion techniques exist for improving the spatial resolution of remote sensing products (Ha et al., 2013; Mahour et al., 2017; Peng et al., 2017), we decided to keep ET products with a spatial resolution lower than 3 km at their original resolution (i.e., GLEAM at 25 km). ~~Since, for one, the intention of this study is a comparison of well-known and established ET products and not an optimization of rescaled comparisons. Second, we did not want to include additional uncertainties potentially originating from the employed downscaling method or auxiliary datasets. Especially, downscaling approaches intend to statistically correlate coarse-scale data and fine-scale auxiliaries, yielding to interpolation uncertainties and errors that cannot be quantified~~ (Peng et al., 2017). All datasets are, however, temporally aggregated to daily time series, in order to provide a temporal basis for comparison and analysis of the signal dynamics.

Table 1: Overview of investigated ET and auxiliary products presenting the data source, the original spatial and temporal resolution as well as the retrieval basis and method of each product.

PRODUCT (NAME)	SOURCE	ORIGINAL SPATIAL / TEMPORAL RESOLUTION	RETRIEVAL BASIS	RETRIEVAL METHOD
<u>ET PRODUCTS</u>				
ICOS (Level 2)	ICOS (ICOS RI et al., 2024)	Point scale / Half-hourly	In-situ measurements	Eddy covariance technique
MODIS (MOD16A2)	NASA (Running et al., 2017)	500m / 8-daily	Remote Sensing	Penman-Monteith
ERA5-land	ECMWF (Muñoz Sabater, 2019)	9 km / hourly	Reanalysis	ECMWF’s IFS, H-TESSEL land surface scheme

SEVIRI (METv3)	ESA (LSA SAF and EUMETSAT SAF On Land Surface Analysis, 2019)	3 km / half-hourly	Remote Sensing	SVAT, (H-) TESSEL land surface scheme
GLDAS-2 (GLDAS_NOAH_025_3H_2.1)	NASA (Beaudoin et al., 2020)	25 km / 3-hourly	Land Surface Model (NOAH) L4	Penman-Monteith
GLEAM (v3)	University of Amsterdam (Miralles et al., 2011; Martens et al., 2017)	25 km / daily	Remote Sensing	Priestley-Taylor

AUXILIARY PRODUCTS

<u>FLUXNET2015</u>	(Pastorello et al., 2020; Warm Winter 2020 Team et al., 2022)	<u>Point scale / half-hourly</u>	<u>In-situ measurements / Reanalysis</u>	<u>Downscaled and consolidated from ERA5-interim reanalysis data and gap filled</u>
<u>SMAP MT-DCA V5</u>	(Feldman et al., 2021)	<u>9 km / daily</u>	<u>Remote Sensing</u>	<u>Tau-Omega; Multi-temporal dual channel algorithm (MT-DCA)</u>
<u>SPEI V2.8</u>	(Beguería et al., 2023)	<u>0.5° / 3-monthly</u>	<u>Remote Sensing / Modelling</u>	<u>FAO-56 Penman-Monteith method</u>

197 In Table 1, the retrieval methods for each ET product are given. MODIS and GLDAS-2 are based on physically-based methods
198 employing the Penman-Monteith equation (Penman, 1948; Monteith, 1965), while GLEAM is based on the Priestley-Taylor
199 equation (Priestley and Taylor, 1972), and ERA5-land uses the ECMWF integrated forecasting system (IFS) and is derived
200 from the ERA5 product where the land surface model is based on the hydrology Tiled ECMWF Surface Scheme for Exchange
201 Processes over Land (H-TESSEL) (Hersbach et al., 2020). Further, SEVIRI employs a soil-vegetation-atmosphere-transfer
202 (SVAT) approach also based on the physics of the TESSEL and H-TESSEL land surface scheme (Balsamo et al., 2009; Bayat
203 et al., 2024; Ghilain et al., 2011). The officially reported ET biases after evaluation of each product (based on comparison with
204 multiple EC flux tower measurements) range from -0.11 mm/day (MODIS) (Running et al., 2019) and -0.12 mm/day (SEVIRI)
205 (The EUMETSAT Satellite Application Facility on Land Surface Analysis (LSA SAF), 2024), to -5% (GLEAM) (Miralles et
206 al., 2011). Meaning, all three products show in average slightly lower ET values compared to EC flux tower measurements.
207 All other products indicate no bias, but employ either bias corrected atmospheric reanalysis data for the forcings to avoid
208 discontinuity in ET retrievals (GLDAS-2) (Rui and Beaudoin, 2022), or found no significant difference in comparison to
209 other products (ERA-land) (Muñoz-Sabater et al., 2021). The Priestley-Taylor equation does not consider the impact of VPD

or canopy conductance (Wang and Dickinson, 2012), while within the Penman-Monteith equation VPD and relative humidity (RH) are used according to the function of Fisher et al., (2008) in order to account for soil water stress when calculating the actual soil evaporation. Further, the canopy conductance is retrieved from stomatal and cuticular conductance depending on LAI and the wet surface fraction, with the stomatal conductance constrained by VPD and minimum air temperature and the cuticular conductance fixed to a constant of 0.01 [mm/s] (Running et al., 2019; Wang and Dickinson, 2012). As stated by (He et al., 2022), the Penman-Monteith equation includes the most important modification by accounting for the physiological controls on ET, using stomatal resistance to explain water movement from leaves to the atmosphere and aerodynamic resistance to describe heat and water vapor transfer from the dry canopy surface to the air above (Running et al., 2019). Hence, the Penman-Monteith equation is, in theory, supposed to be more accurate than the Priestley-Taylor equation but, in turn, requires more ‘parameters that are difficult to characterize’ (Fisher et al., 2008). Within the TESSEL and H-TESSEL schemes, canopy conductance is formulated according to the modified Jarvis function and based on the stomatal conductance (retrieved from net assimilation and Kirchhoff’s resistance/conductance analogy) and cuticular conductance (fixed between 0 to 0.25 [mm/s] according to vegetation types), while SM at four layers, and therefore also deeper soil layers, are accounted when defining the soil water stress on soil evaporation (ECMWF, 2018). Lastly, for this study, it is interesting to note that GLEAM and ERA5-land employ the ECMWF atmospheric reanalysis data (Li et al., 2022), while GLDAS-2 is based on MODIS land surface parameters (Rui and Beaudoin, 2022). These product interdependencies should be kept in mind during interpretation of results.

In the second part of this study, the ET products are compared in relation to two dominant parameters of the SPAS, namely SM and VPD. While VPD comes from in-situ measurements of the Fluxnet network (point precise), SM comes from NASA’s Soil Moisture Active Passive (SMAP) mission, the multi-temporal dual channel algorithm (MT-DCA) L-band (1.4 GHz) dataset (9 km spatial resolution) (Konings et al., 2016; Feldman et al., 2021) (see Tab. 1). We employed the SMAP SM in this study instead of using available in-situ measurements of the Fluxnet network, since the latter were of poor quality at several stations and years, and we wanted to build our analyses on one continuous dataset. The SMAP MT-DCA dataset is quality controlled and filtered for, e.g., snow, frozen ground, and water bodies (Feldman et al., 2021).

2.3 Methods

2.3.1 Extended triple collocation

For the comparison of different ET products in sec. 3.1., the extended triple collocation (ETC) method (McColl et al., 2014) is employed. The ETC technique not only provides the root-mean-square-error σ_ϵ [mm/day] of the classical triple collocation (TC) method (Stoffelen, 1998) among three independent measurement systems, but also provides the correlation $\rho_{t,X}$ [-] among them, giving the sensitivity of the measuring systems. The most important advantage of the TC and ETC techniques is that one can calculate σ_ϵ and $\rho_{t,X}$ without considering any of the systems as the necessary reference. The product with the lowest σ_ϵ and highest $\rho_{t,X}$ identifies the one with the lowest uncertainty. As input to the ETC, the daily ET time series are filtered for

the growing season (April to October) of each year. With the aim of evaluating the performance of the remote sensing products (SEVIRI, MODIS, GLEAM), we compare them individually with ERA5-land and in-situ measurements (ICOS) on the one hand, and with GLDAS-2 and ICOS on the other hand. Sanity checks for Gaussian distributions and large sample sizes of ~853 values per product ensure precise and representative ETC analyses. Additionally, since one of the requirements for thorough ETC analyses is the independence among evaluated datasets (McColl et al., 2014), the error cross-correlation (ECC) values (Gruber et al., 2016) are calculated in order to evaluate product dependencies. In case the ECC lies between -0.5 and 0.5, the datasets can be regarded as independent from each other. The ECC for each product comparison (with ET product $\in [i,j,k,l]$) is calculated from the error cross covariance $\sigma_{\varepsilon_i \varepsilon_j}$ between two products as well as the random error variance $\sigma_{\varepsilon_i}^2$ of each dataset, respectively (Gruber et al., 2016):

$$ECC_{ij} = \frac{\sigma_{\varepsilon_i \varepsilon_j}}{\sigma_{\varepsilon_i}^2 \sigma_{\varepsilon_j}^2}, \quad (1)$$

with

$$\sigma_{\varepsilon_i \varepsilon_j} = \sigma_{ij} - \frac{\sigma_{ik} \sigma_{jl}}{\sigma_{kl}}, \quad (2)$$

and

$$\sigma_{\varepsilon_i}^2 = \sigma_i^2 - \frac{\sigma_{ij} \sigma_{ik}}{\sigma_{jk}}. \quad (3)$$

2.3.2 Anomalies

For the comparison of different SPAS parameters in sec. 3.2., the seasonal imprint is removed from the signals in order to focus on exceptional events in the time series. For that, we calculated the 30-day anomaly time series for each parameter. To do so, the daily average over all four years was calculated first. The resulting daily average was then smoothed using a Savitzky-Golay (Savitzky and Golay, 1964) filter with a window size of 61 days. Lastly, for every day between 2017 to 2020, the difference between the day of interest and the 30-day average of the filtered daily average before that day has been calculated.

2.3.3 Binning

To analyse the effects of water supply and demand on ET, we binned daily ET values into a grid of 30 by 30 SM and VPD conditions, with SM ranging between 0.0001 vol.% and 40 vol.%, and VPD ranging between 0.0001 hPa and 25 hPa, both in 31 steps (to create a grid of 30 by 30). While SM is indicative of the available water supply, VPD is an indicator of atmospheric water demand. The co-regulation of ET by SM and VPD is complex as it depends on stomatal and surface conductance, which in turn are dependent on SM and VPD, as well as vegetation and soil properties (Carminati and Javaux, 2020; Zhang et al., 2021; Vargas Zeppetello et al., 2023). To understand the main directionality of ET changes relative to SM, we calculated the

average slopes of ET relative to SM (equivalent to $\frac{\Delta ET}{\Delta SM}$). The same applies when we examine the directionality of the ET changes with respect to VPD ($\frac{\Delta ET}{\Delta VPD}$). These analyses are done in order to get an indication of the dominating control on ET.

3 Results

3.1 Differences in examined ET products

In Figure 4, times series of the employed ET products (see Tab. 1) are shown at all investigated ICOS stations (see Fig. 1) for the period 2017 to 2020. Apart from the seasonal dynamics of ET, with highest values in the summer months (June, July, August) and low values but with more frequent changes in the winter months (November, December, January), the overall good consistency between the different ET products can be noted.




Figure 4: Comparison of seasonal dynamics of ET [mm] products for the period 2017-2020 at investigated ICOS stations. All time series were cleaned for daily and weekly dynamics using a Savitzky-Golay (Savitzky and Golay, 1964) filter with a window size of 31 days.

281 The highest variability among products and ET dynamics can be observed during summer months, with greatest differences at
 282 stations DE-Hai and DE-Ruw when comparing all products to the ICOS measurements. Here, the ground-based ET shows
 283 always lower values across all years for DE-Hai, and in 2018 and 2019 for DE-Ruw. Additionally, for each year, the ICOS ET
 284 rises a few weeks later than the other products at both stations but decreases together with all other ET products. At station
 285 CZ-Lnz, ERA5-land shows the overall lowest ET values during the growing period (April to October). Further, the highest ET
 286 values are found at station FR-Bil for the GLDAS-2 product with most pronounced differences to all other products in 2018,
 287 while overall lowest values across all years and ET products are displayed at DE-Rus. At the latter, ET values never exceed 4
 288 mm/day. From this daily time series analyses, the largest differences among ET products can be seen at the DBF station DE-
 289 Hai, MF station CZ-Lnz, and agriculture station DE-Rus. At DE-Hai, the ICOS ET is overestimated by all other products, at
 290 CZ-Lnz, the ERA5-land product is lower compared to all other ET products, especially in the summer months, and at DE-Rus,
 291 the MODIS and often also the ICOS product are overestimated by the ERA5-land and SEVIRI products. Hence, no clear
 292 pattern at all stations and between different land cover classes can be found.

293 For more detailed analyses, daily time series of ET products are averaged to 8-daily sums in order to account for the coarse
 294 temporal resolution of the MODIS product (see Tab. 1). In Figure 5, the 8-daily ET products are compared with each other at
 295 the two agriculture stations. The same illustrations for the forest stations can be found in the supplement (see Figs. S2-S4).
 296 These figures show the scatter plots between ET products giving the probability density function (PDF) of points (by colour)
 297 below (left panels) and above (right panels) the matrix diagonal, as well as the PDF curves for each site and product in the
 298 diagonal of the matrix. They support the previously stated good consistency between ET products but outline the exact
 299 differences on 8-days scale in more detail. The highest density of values can be observed between 0 to 10 mm/8-days at all
 300 stations except at DE-Ruw and FR-Bil. This comes from the rather low ET values during the autumn, winter, and spring
 301 seasons due to the overall reduced solar radiation combined with decreased vegetation cover during cold months. However, at
 302 stations DE-Ruw (see Fig. S3, right panels) and FR-Bil (see Fig. 4, left panels), the density of values is shifted towards higher
 303 ET (0 to 20 mm/8-days). These are two out of the three stations covered by coniferous forest. While FR-Bil has a two-part
 304 split land cover in the footprint (shrub and coniferous forest), DE-Ruw is almost homogeneously covered by coniferous forest
 305 (see Fig. 2), and both stations show higher ET values during autumn and spring seasons compared to all other stations due to,
 306 e.g., the lack of leaf off conditions during that periods. The third station covered by coniferous forest (FI-Let), however, shows
 307 the density of values between 0 to 10 mm/8-days (see Fig. S3, left panels), similar to DBF and MF stations. This is the
 308 northernmost station, typically covered with snow between November and March.

309 Further, the over- or underestimation of values between two products can be seen, such as the overestimation of ICOS
 310 compared to all other ET products at DE-Hai for higher ET values, affirmed by the PDF for ICOS peaking at the highest
 311 density (see Fig. S2, left panels). There is also an overestimation of MODIS compared to all other products at DE-Rus (see
 312 Fig. 5, right panels) and CH-Lae (see Fig. S4, left panels) when ET values are higher. DE-Rus is the only homogeneously
 313 covered agricultural station with potentially most changes in land cover classes during the seasons and years, showing the
 314 greatest differences in ET products due to the overall higher complexity of agricultural plants and more frequent alterations.

While the PDF of MODIS at DE-Rus peaks at the highest density and gives the smallest range of ET values across all stations, a bimodal distribution of densities is displayed at CH-Lae. This bimodal distribution of densities is also noticeable at other products and stations but stronger always for MODIS.





Figure 5: Comparison of seasonal dynamics of ET [mm/8-days] products for the period 2017-2020 at investigated ICOS stations DE-Rus (right panels above the diagonal of the matrix) and FR-Bil (left panels below the diagonal of the matrix). All time series were averaged to 8-daily sums at MODIS dates, and cleaned for daily and weekly dynamics using a Savitzky-Golay (Savitzky and Golay, 1964) filter with a window size of 31 days. All statistics are included in supplement figures S5-S7.

This visual interpretation is also supported by statistics in supplement Figures S5-S7. In general, the highest coefficient of determination, R^2 [-], among all products can be found at station CH-Lae, while the overall lowest root-mean square errors, RMSE [mm/8-days], are retrieved at both ENF stations (DE-Ruw, FI-Let). DE-Ruw is also the station with, in general, lowest percentage bias, PBIAS [%], among all ET products. In detail, the highest R^2 of 0.94 is found between GLEAM and GLDAS-2 at CH-Lae, while the lowest RMSE of 2.3 mm/8-days and the lowest PBIAS of -0.05 % is found between GLEAM and

328 ERA5-land again at CH-Lae. The lowest R^2 of 0.62 and highest PBIAS of 91 % is found between ICOS and MODIS at the
 329 agricultural station DE-Rus, while the highest RMSE of 8.8 mm/8-days is found between MODIS and ERA5-land again at
 330 DE-Rus. In summary, the statistics indicate an overall worse consistency among products at the rather mixed agricultural
 331 station (DE-Rus) and better consistency at ENF stations.

332 In order to evaluate the performance of each ET product in more detail, the ETC method (McColl et al., 2014) is employed.
 333 Here, we use the ETC approach to compare the three remote sensing products individually first with ERA5-land and ICOS,
 334 and then with GLDAS-2 and ICOS. The preceding calculation of ECC values among all products (see Fig. S8) is conducted
 335 to ensure the independence of the examined products, which is required by ETC analysis (see Sec. 2.3.1). Overall, ECC values
 336 are always around zero or within the acceptable range of -0.5 to 0.5. Only at station DE-HoH between GLDAS-2 and GLEAM,
 337 at CZ-Lnz between ERA5-land and GLEAM, at CH-Lae between ERA5-land and MODIS as well as for all product
 338 comparisons at DE-Rus (except between ERA5-land and SEVIRI), ECC values outside the acceptable range can be found (see
 339 Fig. S8). The high ECC values at DE-HoH, CZ-Lnz, and DE-Rus between GLEAM and GLDAS-2 or ERA5-land is not
 340 surprising, since the GLEAM product is based on various remote sensing and reanalysis datasets, with among others GLDAS
 341 and ERA5 (see Sec. 2.2). Hence, at most stations ET products can be regarded as statistically independent from each other.
 342 Only some potential product dependencies, especially at the agricultural station DE-Rus, should be kept in mind during the
 343 interpretation of ETC results.

344 In Figure 6, the ETC statistics for the applied product combinations at all stations are shown. While the x- and y- axes represent
 345 the estimated root-mean-square-error σ_ϵ , the arcs give the correlation $\rho_{t,x}$. Hence, numbers (representing the eight stations)
 346 close to zero on the x- and y-axes and close to one on the arcs give the best ETC results, meaning lowest uncertainty of the ET
 347 product (represented by colours) compared to the other two products, respectively. It can be seen that all σ_ϵ values are below
 348 1.07 mm/day due to the overall high consistency among ET products, with correlations between $0.39 < \rho_{t,x} < 0.99$. However,
 349 products with highest $\rho_{t,x}$ necessarily do not have the lowest σ_ϵ . Hence, the discrepancy between products varies but does not
 350 dominate differences in the sensitivity among products. The highest σ_ϵ is found at station FR-Bil for GLDAS-2, when
 351 comparing GLDAS-2 with GLEAM and ICOS. The lowest $\rho_{t,x}$ of 0.33 is found at station DE-Ruw for ICOS as the results of
 352 the ETC among GLDAS, MODIS, and ICOS. Despite the high ECC values at DE-Rus (see Fig. S8) and hence, potential
 353 product dependencies, ETC results at this station are inconspicuous with comparable errors and correlations. Overall, ERA5-
 354 land, SEVIRI, and GLEAM perform better at all stations with either lowest errors or highest correlations within their ETC
 355 triplets. In summary, compared to ERA5-land and ICOS, the remote sensing products (SEVIRI, MODIS, GLEAM) show
 356 similar uncertainties as ERA5-land, but at most stations ERA5-land outperforms GLEAM and MODIS (see Fig. 6, upper row).
 357 Further, compared to GLDAS-2 and ICOS, the remote sensing products in most cases outperform GLDAS-2 and ICOS,
 358 showing the lowest uncertainties, i.e. lower errors and higher correlations (see Fig. 6, lower row). During all analyses, ICOS
 359 shows generally the highest uncertainties. Potential explanation is the discrepancy in spatial resolutions (see Tab. 1) as will be
 360 discussed in more detail in sec. 4.



361


362 Figure 6: Estimated root-mean-square-error (σ_e) [mm/day] (on the x- and y- axes) and correlation ($\rho_{t,x}$) [-] (on the arcs) among ET products
 363 at all stations based on the extended triple collocation (ETC) method from McColl et al., (2014). Numbers represent the eight stations and
 364 colours the different ET products. 1st row: ETC between SEVIRI, MODIS, and GLEAM datasets respectively with ERA5-land and ICOS.
 365 2nd row: ETC between SEVIRI, MODIS, and GLEAM datasets respectively with GLDAS-2 and ICOS.

366 3.2 Drought impacts on ET products

367 As shown in Figures 3 and S1, 2018 was an exceptional dry year across central Europe. In this section, the impact of the
 368 drought in 2018 on ET is investigated by comparing it to SM and VPD, the two main parameters that are used for monitoring
 369 drought-related terrestrial ecosystem productivity (see Sec. 1). For that, we will compare 2018 always to the rather wet year
 370 2017 to identify significant changes.

371 In Figure 7, the time series of ICOS ET, SMAP SM, and in-situ measured VPD for 2017 and 2018 are compared to their
 372 respective calculated anomalies (see Sec. 2.3.2) for DBF (DE-HoH, DE-Hai) and ENF (DE-Ruw, FI-Let) stations. While ET
 373 and VPD show a distinct seasonal pattern at all stations with highest values during summer months, SM shows a less clear
 374 seasonal pattern with more inter- and intra-annual variations. At both DBF stations and the ENF station DE-Ruw, the highest
 375 SM values are generally found during the winter months. In contrast, at ENF station FI-Let, an almost constantly increasing
 376 SM in 2017 can be observed with a distinct drop from in January 2018 and subsequent distinct increase in April 2018. The SM
 377 also stays at high values throughout the entire summer until mid of October in 2018, besides a smaller decrease from end of

378 May until August. However, these SM values may be an artefact of snow cover or frozen ground at the northernmost station
 379 and should be treated carefully, although the MT-DCA is quality controlled and filtered for that (see Sec. 2.2).




380 Figure 7: Time series of daily ICOS ET [mm/day], SMAP SM [vol.%], and in-situ VPD [hPa] for 2017 and 2018 at DBF (DE-HoH, DE-
 381 Hai), and ENF (DE-Ruw, FI-Let) stations compared to their respective anomalies (see Sec. 2.3.2). All time series were cleaned for daily and
 382 weekly dynamics using a Savitzky-Golay (Savitzky and Golay, 1964) filter with a window size of 31 days.

383 From these time series, in general lower ET and higher VPD values can be found in 2018 compared to 2017, reflecting the
 384 drought conditions with higher atmospheric aridity and decreased water supply for plant transpiration and soil evaporation in
 385 2018. At the MF (CZ-Lnz, CH-Lae) and agriculture (DE-Rus, FR-Bil) stations, the same trends can be observed but with minor
 386 differences in VPD maxima between 2017 and 2018, and sometimes higher ET peaks in 2018 at stations CZ-Lnz and FR-Bil
 387 (see Fig. S9). The overall lowest SM values can also be found in 2018, except at station FI-Let. At the DBF stations and station
 388 DE-Ruw, constantly low SM values over several months from mid of April to mid of October are shown without any significant
 389 increase during this time in 2018 (see Fig. 7). The same is true at MF station CH-Lae and the agricultural stations. At station
 390 CZ-Lnz, SM is varying monthly at low values between ~5 vol.% and 18.6 vol.% (see Fig. S9). When analysing the anomaly
 391 time series (seasonal detrending; see Sec. 2.3.2) of each parameter and station, in general higher ET and VPD anomalies and
 392 lower SM anomalies are found in 2018 compared to 2017, except at station FI-Let with higher SM anomalies in 2018 compared
 393 to 2017 (see Figs. 7 & S9).

394 These anomalies are subsequently used in Figure 8 to visualize the kernel densities of SM, VPD, and ET anomalies of all
 395 stations for 2017 and 2018. In Figure 8, only the vegetation periods from April to October within each year are analysed. It
 396 can be seen that in 2018 (drought year), the SM and ET anomalies peak at lower, negative values compared to 2017, where
 397 they peak around zero, while the VPD anomalies peak at higher, positive values compared to 2017. Also, the respective


398 anomaly medians are lower for SM and ET, and higher for VPD in 2018. The calculated p -values of always ≤ 0.045 prove the
 399 shift in yearly median values at the 5 % significance level.



400
 401 Figure 8: Kernel density estimates of daily SMAP SM, in-situ VPD, and ICOS ET anomalies (see Sec. 2.3.2) during April to October of
 402 2017 and 2018 across all investigated stations. The dashed lines represent the seasonal median of respective parameters and years. The p -
 403 values of a two-sided Wilcoxon rank-sum test indicate the acceptance (> 0.05) or rejection (< 0.05) of the null hypothesis regarding
 404 continuous distributions with equal medians at the 5 % significance level.

405 When comparing the anomalies for different ET products (see Fig. 9), a similar shift towards lower values for 2018 compared
 406 to 2017 can be found for MODIS and ERA5-land products. For SEVIRI, GLDAS-2, and GLEAM a shift towards higher
 407 anomalies in 2018 is found with medians at slightly higher values compared to 2017. However, while the ICOS p -value of
 408 0.045 being close to the 5 % significance level of equal medians, the ones of SEVIRI, GLDAS-2 and GLEAM are more
 409 significant around zero. GLEAM anomalies peak at the same value for both years but with higher positive anomalies for 2018
 410 at values greater than 0.6. In general, Gaussian distributions around zero are evident for both years at all anomalies of ET
 411 products. Only at MODIS, a clear bimodal distribution in ET anomalies of 2018 with a first peak around -0.4 and a subsequent
 412 second smaller peak at 0.55 can be found. This is also the ET product with the smallest anomaly range from -1.5 to 2.5. All
 413 other ET products vary at least between -3 and 3. For the ET products ERA5-land, GLDAS-2, and GLEAM, a non-linear
 414 decrease in 2018 can be found with almost stagnating anomalies around one. For the ICOS and SEVIRI data, this trend is first
 415 visible at values greater than one. In contrast, the density curves of ET anomalies for 2017 are smoother for all products,
 416 showing a clear Gaussian distribution. Again, the calculated p -values of ≤ 0.02 prove the shift in yearly median values at the
 417 5 % significance level, except for the MODIS product (p -value < 0.1). The MODIS product is also the ET product with the
 418 lowest temporal resolution of eight days (see Tab. 1). When analysing all other ET products at the same 8-daily resolution (see
 419 Fig. S10) similar bimodal distributions in 2018 can be found for ERA5-land, SEVIRI, and GLEAM. GLDAS-2 shows even a
 420 trimodal distribution with the highest density of ET anomalies around -4.5, a second peak around 1.4, and a third peak around
 421 6.3. Although no clear bimodal distribution can be seen for ICOS even at 8-daily resolution, the distribution smoothly increases
 422 from -15 to -4 and then non-linearly decreases with at least three smaller plateaus (see Fig. S10). And even for 2017, the
 423 Gaussian distributions are not that smooth as for the daily analyses. More detailed analyses revealed that there is a distinct
 424 drop in 8-daily anomaly time series, leading to this bimodal distribution. Between April and August almost only positive ET


425 anomalies are found, while during September and October almost only negative anomalies are found. The same trend is, of
 426 course, also visible for the daily time series but due to the preserved daily and intra-weekly dynamics, the difference between
 427 positive and negative anomalies during both periods (April-August, September-October) is not that distinct. These small-scale
 428 dynamics are excluded in the 8-daily analyses. However, the differences in ET anomalies between 2017 and 2018 are greater
 429 for the 8-daily anomaly analyses (see Fig. S10) compared to the daily anomaly analyses (see Fig. 9), indicating that drought
 430 impacts on ET are more pronounced at larger time scales (more than a week, monthly) than on smaller time scales (less than
 431 a week, daily). In summary, the reason for the bimodal distribution in ET anomalies within the MODIS products originates
 432 from the lower temporal resolution.



433
 434 Figure 9: Kernel density estimates of daily ET anomalies (see Sec. 2.3.2) for all investigated ET products during April to October of 2017
 435 and 2018 across all investigated stations. The dashed lines represent the seasonal median of respective parameters and years. The p -values
 436 of a two-sided Wilcoxon rank-sum test indicate the acceptance (> 0.05) or rejection (< 0.05) of the null hypothesis regarding continuous
 437 distributions with equal medians at the 5 % significance level.

438 For analysing the dependencies between ET, SM and VPD, respective ET products in SMAP SM and in-situ measured VPD
 439 bins (see Sec. 2.3.3) are visualized for the wet year 2017 (see Fig. 10) and the dry year 2018 (see Fig. 11) across all stations.
 440 ET for all stations and both years are similarly distributed across the SM and VPD phase space.
 441 For the rather wet year 2017, a general decreasing trend in ET values along increasing VPD and increasing SM can be found
 442 for all ET products except SEVIRI. Here, a decreasing trend along increasing VPD but decreasing SM is visible as indicated
 443 by the arrow within the inset plot (see Fig. 10). Overall, ET varies more with VPD than SM. Only ET from ICOS and to some

444 extend ERA5-land and GLEAM have highest values at intermediate VPD and SM, and lower ET at low SM. Especially ET
 445 products SEVIRI and GLDAS-2 do not display any reductions at low SM.



446
 447 Figure 10: ET [mm] relative to SMAP SM [vol.%] and in-situ VPD [hPa] for all investigated ET products and averaged over all investigated
 448 ICOS stations in 2017. The inset plots provide the corresponding median slope in SM and VPD changes.

449 For the dry year 2018, only MODIS and GLDAS-2 still show a decreasing trend along increasing VPD for increasing SM. All
 450 other products indicate decreasing ET for increasing VPD and decreasing SM (see. Fig. 11). At SEVIRI, the slope in SM
 451 direction is twice as low in 2018 compared to 2017 but almost the same for VPD, meaning greater decrease in ET along SM
 452 during the dry year. A similar trend is observable at MODIS with half of the slope along SM in 2018 compared to 2017,
 453 meaning half as strong increase in ET values with SM during the drought affected year 2018. Lastly, at GLDAS-2, the slope
 454 along SM bins is increased by a factor of almost seven in addition to a reduced slope in VPD of ~ 0.1 hPa in 2018, meaning
 455 stronger increase in ET values at increasing SM at simultaneously decreasing VPD during the drought year. Further, ET values
 456 are in general lower in 2018 compared to 2017, but in 2018, bins at higher VPD values with low ET can be found across the
 457 entire SM range (see Fig. 11).

458 In summary, for both years, ET is generally higher at high VPD, i.e., higher atmospheric water demand, and much lower below
 459 a VPD of 5 hPa. In figures 10 and 11, we do not really see very clear reductions of ET with decreasing SM. Hence, ET varies

more with VPD than SM. The influence of SM on ET is only noticeable when comparing the wet (2017) and dry (2018) years with each other, as the change along SM ($\frac{\Delta ET}{\Delta SM}$) is significantly higher during the drought affected year.




Figure 11: ET [mm] relative to SMAP SM [vol.%] and in-situ VPD [hPa] for all investigated ET products and averaged over all investigated ICOS stations in 2018. The inset plots provide the corresponding median slope in SM and VPD changes.

For more detailed analyses regarding the drought effect on ET products, we calculated the coefficient of variation (CV) [%] for 2017 and 2018 (see Fig. 12) for each ET bin relative to the SM and VPD ranges (see Sec. 2.3.3.). CV is defined as ratio between the standard deviation and the mean and provides the relative dispersion or amount of uncertainty of data. As can be seen, the differences in CV between 2017 and 2018 are highest for low SM. Here, the variability in ET values during the drought year of 2018 reach higher VPD values compared to 2017. Furthermore, overall higher CV are estimated for low VPD across the entire SM range in 2018 compared to 2017. In contrast, 2017 shows slightly higher CV values for intermediate values (SM between 10 and 30 vol.% and VPD between 4 and 8 hPa). When comparing the different ET products among each other, the CV show overall similar patterns. However, for both years the CV median for ICOS (49.23 %, 48.43 %) are always the highest compared to all other products, indicating a greater dispersion of data points within the ET time series. Lowest median CV are found within ERA5-land with 33.28 % in 2017 and 36.48 % in 2018 (see Fig. 12), indicating overall lowest amount of uncertainty in ET data.




Figure 12. Coefficient of variation (CV) [%] of ET values relative to SMAP SM [vol.%] and in-situ VPD [hPa] for all investigated ET products and averaged over all investigated ICOS stations for 2017 (upper two rows) and 2018 (lower two rows). Inlet plots give the histogram of displayed CV values with the red line indicating the location of the given median for each product.

477 4 Discussion

478 4.1 Differences in examined ET products

479 When evaluating the performance of all ET products from remote sensing, reanalysis, modelling and ground-based eddy
480 covariance measurements, analyses of their time series revealed that the ICOS ET almost always show a time lag of about few
481 weeks during spring ET rise compared to all other products (see Fig. 4). This could be explained by the discrepancy in spatial
482 resolutions, with the ICOS product providing local point-scale measurements compared to the other larger-scale remote sensing
483 and modelling ET products. This spatial mismatch alters the vegetation impact within the ET signal. Another reason is the
484 dependency of models on indicators for phenological changes in vegetation. For example, many models use the leaf-area index
485 (LAI) to track phenology dynamics, which influence ET simulations (Adeluyi et al., 2021). Further, the overall lowest ET
486 values were found for all products at the agricultural station DE-Rus, while highest values were found at the southernmost
487 station FR-Bil, where the highest average precipitation was recorded between 2017 to 2020 (see Fig. 3). Reason for that are
488 for one, reduced transpiration of agricultural sites throughout the year compared to forested sites, and second, the humid
489 Atlantic climate at the southernmost station at lowest altitude (see Tab. S1). The 8-day analyses showed that MODIS gives
490 higher values compared to all other ET products at two stations, while ICOS is higher than all other ET products at one station.
491 Further, the highest density of values was found between 0 to 10 mm/8-days due to the seasonal imprint with reduced ET
492 across Europe during months with reduced solar radiation and vegetation cover (November-March). Only at the two coniferous
493 forest stations (DE-Ruw, FR-Bil), the highest density of values is between 0 to 20 mm/8-days with lower ET values only
494 during winter months (December-February). However, this does not apply to the third coniferous station FI-Let, which is the
495 northernmost station with less dense forests and more snow fall between November and March, which influences the estimation
496 of ET. Hence, the lack of leave-off conditions and the reduced amount of days with snow cover influences the amount of ET.
497 Conducted statistics confirmed the noticeable differences among ET products and ICOS stations, which indicated an overall
498 lower agreement among products at the rather mixed agricultural station (DE-Rus) and better consistency at ENF stations (DE-
499 Ruw, FI-Let). Hence, products differ most at stations with complex land cover conditions, where varying crops and growing
500 seasons (changing phenology) make the estimation of ET more difficult, while evergreen needle-leaved stations with less
501 changes throughout the year and between years are easier to define (temporal homogeneity).

502 For more detailed product performance analyses, the extended triple collocation (ETC) method (McColl et al., 2014) and SM-
503 VPD binning revealed highest uncertainties (see Fig. 6) and coefficient of variation (see Fig. 12) for the ICOS product, and
504 lowest uncertainties for SEVIRI and GLEAM as well as ERA5-land. The highest ETC error was estimated for GLDAS-2,
505 when analysing with GLEAM and ICOS, while the lowest sensitivity (correlation) was found for ICOS, when analysing with
506 GLDAS-2 and MODIS (see Sec. 4.1). Hence, the remote sensing products (SEVIRI, GLEAM) and the reanalysis product
507 (ERA5-land) differed most from the in-situ field-scale (ICOS) and modelling (GLDAS-2) products. One reason for the
508 mismatch between the ICOS product and SEVIRI, GLEAM and ERA5-land is surely the spatial mismatch between the point-
509 scale ground-based EC tower measurements and the remote sensing (3 km) or reanalysis (9 km) products. However, in order

510 to capture vegetation stress, ecosystem health, and fine-scale variability in ET globally, adequate spatial (and temporal)
 511 resolutions are necessary. Here, detailed research regarding downscaling techniques, as reviewed in, e.g. (Mahour et al., 2017;
 512 Peng et al., 2017), that combine medium-scale ET data with fine-scale auxiliaries in order to improve the spatial resolution,
 513 are needed regarding its uncertainties and impact on product comparisons. Further, ET measurements based on the eddy
 514 covariance method tend to underestimate sensible heat (H) and latent heat (LE) fluxes (Petropoulos et al., 2015), are often
 515 temporally too short and spatially too sparse to sample drought conditions correctly (Zhao et al., 2022), and suffer from
 516 challenges to close the energy balance (Yu et al., 2023). Several studies (Twine et al., 2000; Petropoulos et al., 2015; Barrios
 517 et al., 2024) reported an error range of EC measurements of ~10-30 % due to, e.g., a ‘systematic closure problem in the surface
 518 energy budget’ (Twine et al., 2000). In order to identify potential product dependencies, which may impact the ETC results,
 519 the estimated error cross-correlations (ECC) were calculated, with high ECC between GLDAS-2 and GLEAM (at DE-HoH),
 520 between ERA5-land and GLEAM (at CZ-Lnz), and all products and GLEAM (at DE-Rus). These need to be accounted for
 521 when analysing the differences among ET products. Although in this study, we have analysed different land cover classes
 522 within a 3 km footprint around every ICOS station at daily resolution to account for the different resolutions, the SEVIRI
 523 product provides ET data every 30 minutes at moderate spatial resolution (3 km), and showed to capture ET dynamics on small
 524 as well larger temporal scales comparable or even better than other examined products, as also reported by previous studies,
 525 e.g., (Hu et al., 2015; Petropoulos et al., 2015; De Santis et al., 2022). None of the other examined products can provide similar
 526 spatio-temporal coverage, due to either lower temporal resolution (MODIS) or coarser spatial resolution (ERA5-land, GLDAS-
 527 2, GLEAM). Only the ICOS data provide similar temporal resolution to SEVIRI but at point-scale, which disqualifies it for
 528 global analyses. Although there exist other ET products from remote sensing and modelling, e.g., (Jiménez et al., 2011; Mueller
 529 et al., 2013; Fisher et al., 2020; De Santis et al., 2022; Yu et al., 2023), the examined ET products in this study are appropriate
 530 when addressing global analyses since other products have either a more coarse spatial or temporal resolution (Yu et al., 2023),
 531 ~~or~~ are limited to clear sky conditions (De Santis et al., 2022), which prohibits continuous time series of ET measurements, or
 532 are higher order derivatives from either field measured or merged remote sensing based products (Jung et al., 2019; Chen et al.,
 533 2021). We also analysed data from the ECOSystem Spaceborne Thermal Radiometer Experiment on Space Station
 534 (ECOSTRESS) launched by NASA in June 2018 (Fisher et al., 2020) at the beginning of our analyses. However, we found
 535 several problems with this product and worse performance compared to other ET products, meaning a clear overestimation
 536 using the ECO3ETPTJPL product, as reported also by previous studies, e.g., (Liu et al., 2021; De Santis et al., 2022; Wu et
 537 al., 2022). In our research with ECOSTRESS, data was unavailable at CZ-Lnz and FI-Let. Another ECOSTRESS ET product,
 538 the ECO3ETALEX (based on the DisALEXI model), has shown better performance, but it is more suited for agricultural
 539 applications, and it is limited to the United States (Cawse-Nicholson and Anderson, 2021). ECOSTRESS level 3 ET data come
 540 at the advantage of a high spatial resolution (70 m), but its temporal resolution is irregular due to the ISS orbit and the
 541 dependency on the product type and study region limited our preliminary analyses. For these reasons, we decided not to include
 542 it in our research.

543 4.2 Impact of droughts on ET products

544 Since remote sensing-based ET products are not purely observational, the performance of an ET product is highly dependent
545 on the employed retrieval model for ET estimation. This is in turn dependent on how the model deals with limitations in SM
546 or VPD and responses under drought conditions. Every retrieval method has its own strengths and weaknesses, but especially
547 under drought conditions, the ability of the employed algorithm to deal with water shortage and vegetation stress is essential
548 for valid ET estimation. Varying types of vegetation have different strategies how to deal with water stress, e.g., by closing
549 stomata to prevent water loss through leaves and increasing the water uptake from the soil or deeper soil depths by increasing
550 the water resistance (He et al., 2022). Many studies reported decreasing ET during droughts due to reduced SM supply and
551 hence, decreasing evaporation, but also decreasing transpiration since plants close their stomata to prevent water loss (Novick
552 et al., 2016; Zhao et al., 2022). However, during drought conditions with increasing air temperatures, ET can also increase due
553 to the higher atmospheric moisture demand (increasing VPD). Further, the generic statement that ET decreases due to
554 decreasing SM often ignores the fact that plants have access to SM from greater soil depths, which are not immediately affected
555 by meteorological droughts, or have different strategies for drought resistance (Gupta et al., 2020; He et al., 2022; Feldman et
556 al., 2024). Hence, the dynamics of ET to drought conditions remain highly variable (Zhao et al., 2022). Novick et al., (2016)
557 pointed out that SM and VPD may become more decoupled in the future and models need to resolve limitations in SM and
558 VPD independently from each other in order to capture the response of ecosystems to water stress correctly (Novick et al.,
559 2016; Zhao et al., 2022). How models react to limitations in SM and VPD varies significantly which impacts resulting ET.
560 Analyses performed in this study revealed that during the rather wet year 2017, ET varied more with VPD than with SM, with
561 almost no dependency of ET on SM in SEVIRI and GLDAS-2 products. Here, our results indicate that ET is more controlled
562 by atmospheric demand rather than ~~wateratmospheric~~ supply from atmosphere (precipitation) and soil (soil moisture) as
563 reported also by Zhou et al., (2019). However, it is suggested by previous work and the Budyko framework (Budyko and
564 Miller, 1974) that ET should exhibit some level of dependence on SM (Porporato et al., 2002; Zhang et al., 2021). One reason
565 could be that forests at selected ICOS stations might have substantial access to deeper SM (root zone) that exceeds the
566 measurement depths of the SMAP satellite (first 25 cm) (Feldman et al., 2022). When analysing the controls of SM and VPD
567 on ET during the dry year 2018 however, all ET products, except MODIS and GLDAS-2, showed that ET decreases with
568 increasing VPD and decreasing SM. For SEVIRI, even a twice as large decrease in ET along SM during the drought year could
569 be observed compared to the rather wet year. This declining trend of ET during dry years when ET is limited by moisture and
570 VPD is increasing due to increasing air temperatures is in line with previous studies (Jung et al., 2010; Seneviratne et al., 2010;
571 Zhou et al., 2019). Further, results show that VPD and SM are negatively coupled during extreme events as reported also by
572 (Zhou et al., 2019; De Santis et al., 2022). However, MODIS and GLDAS-2 products showed an increase of ET with increasing
573 SM and with decreasing VPD during 2018 (see Fig. 11). These are the two products that are based on the Penman-Monteith
574 equation (see Tab. 1), and that were outperformed by SEVIRI, ERA5-land and GLEAM in the ETC analyses (see Fig. 6). For
575 MODIS, one reason for the worse performance was found to be the coarse temporal resolution of 8-days, since at this time

576 scale the temporal variability of ET is significantly different lacking all diurnal and day-to-day ET dynamics. The
577 underperformance of MODIS compared to in-situ EC measurements was also reported by (De Santis et al., 2022), who found
578 that MODIS overestimated in-situ ET measurements at stations in Italy, as well as (Yu et al., 2023), who investigated several
579 stations with different land covers and varying climatic zones across the U.S. They concluded that daily or monthly ET products
580 performed best compared to EC tower measurements (Yu et al., 2023). Due to the temporal resolution, MODIS is the only
581 product showing a bimodal distribution of ET anomalies with a p -value above the 5 % significance level (see Fig. 9). In this
582 study, we could show that differences in ET anomalies between 2017 and 2018 are greater for the 8-daily anomaly analyses
583 (see Fig. S10) compared to the daily anomaly analyses (see Fig. 9), indicating that drought impacts on ET are more pronounced
584 at larger time scales (more than a week, monthly) than on smaller time scales (daily, less than a week). Hence, the temporal
585 scale for ET analyses is crucial in order to select which temporal component of the ET dynamics should be considered for a
586 respective application.

587 Further, although GLEAM is built on the less parameterized Priestly-Taylor equation compared to the Penman-Monteith
588 equation since it does not consider VPD or canopy conductance on soil water stress, the GLEAM ET product showed to deliver
589 better ETC results and statistics in this study. A comparable or even better performance of the Priestley-Taylor equation
590 compared to the Penman-Monteith was also reported in previous studies, e.g., (Akumaga and Alderman, 2019; Bottazzi et al.,
591 2021). Reasons could be the uncertainties of input variables within the Penman-Monteith equation, e.g., for stomatal-~~or~~
592 canopy, or aerodynamic resistances, which are often unknown, approximated (Widmoser, 2009), or parameterized based on
593 the wrong variable (Hu et al., 2015), or due to the overestimation of specific parameters, such as the net radiation, or other
594 aerodynamic factors as reported by (Hao et al., 2018). Similar, Hu et al., (2015) stated that MODIS tends to overestimate water
595 stress during thawing of frozen soil in Spring or over irrigated land, which leads to an underestimation of soil evaporation.
596 Moreover, several studies pointed out that the Penman-Monteith equation needs to be adapted for climate/weather extremes
597 and vegetation limited cases, e.g., (Widmoser, 2009; Hao et al., 2018; McColl, 2020).

598 The estimated coefficient of variation (CV) showed that during the drought year 2018 ET values display highest uncertainty
599 for low SM or low VPD, while during the rather wet year 2017, the highest variability was found for intermediate SM and
600 VPD values. Hence, our results show that during drought conditions the estimation of ET leads to highest uncertainties and is
601 most difficult for low SM and low VPD depending on the assumptions for controlling factors on ET. For example, within the
602 Penman-Monteith equation, aerodynamic as well as stomatal resistances are considered but since they can vary significantly
603 for drought and non-drought conditions erroneous assumptions for them can lead to significant errors. During normal or wet
604 conditions, as shown for 2017, CV results do not vary that much between investigated ET products but indicate for all highest
605 variability for intermediate values of SM and VPD, which originate most probably from different vegetation and ecosystem
606 types. Here, more research for individual ecosystem types is required to further address ET controls, as vegetation is the
607 controlling factor on ET estimates when SM and VPD are not limited (Brown et al., 2010; Jin et al., 2017).

608 In summary, it is important for ET retrieval algorithms to account for water droughts and vegetation stress as done with
609 adaptable stomatal closure and canopy resistance within the Penman-Monteith equation. However, analyses showed, that false

610 assumptions on these physiological stress indicators can decrease the performance and be exceeded by less parameterized and
611 simpler retrieval algorithms like the Priestley-Taylor equation.

612 **5 Conclusion and Outlook**

613 In this study, eight different ET products with varying temporal and spatial resolutions as well as varying ET retrieval methods
614 (see Tab. 1) are analysed across central Europe for the period of 2017 to 2020. Despite the spatial mismatch (in-situ vs. remote
615 sensing) and the spatial heterogeneity of the analysed landscapes (see Fig. 2), all products showed a concurrent seasonal pattern
616 and overall low uncertainties during ETC analyses. It was shown that ET varied from year to year for different forest and
617 agricultural stations due to changing seasonal weather and vegetation conditions over the years. Analyses revealed that
618 temporal and spatial homogeneity helps with the consistency and interpretability of the ET estimates. This is, since products
619 were most consistent with each other at stations with less complex land cover conditions and changes throughout the seasons
620 (the evergreen needle-leaved stations DE-Ruw and FI-Let). Despite the good match in seasonal patterns, differences in ET
621 products were noticeable. The remote sensing products, SEVIRI, MODIS, and GLEAM, performed equivalently well or even
622 better than the in-situ measured (ICOS), modelled (GLDAS-2) or reanalysis (ERA5-land) products for this specific study
623 concept (3 km footprint, daily analyses). Extended triple collocation (ETC) and SM-VPD binned ET analyses revealed that
624 SEVIRI and ERA5-land (the two products based on the (H-) Tessel land surface scheme) perform best. They provide low
625 uncertainties when compared with other products and reasonable SM and VPD controls on absolute ET. GLEAM also shows
626 a good performance, although this result should be taken with caution since potential product dependencies with ERA5-land
627 and GLDAS-2 may have affected the ETC results. When analysing the behaviour of ET in context of SM and VPD during the
628 rather wet year 2017 and dry year 2018, it was found that in 2017, ET is highly dependent on VPD and less on SM. Hence,
629 with sufficient moisture supply, ET is mainly controlled by atmospheric demand and the vegetation transpiration. In contrast,
630 in 2018, limited moisture supply because of decreasing SM and increasing VPD, which were in turn due to increasing air
631 temperatures, led to a decline in ET, in line with previous studies. Further, during the dry year 2018, SM and VPD were more
632 negatively coupled which could also had an impact on the ET decline. These behaviours were consistently found in all ET
633 products, except for GLDAS-2 and MODIS, the two products whose retrieval approaches-methods are based on the Penman-
634 Monteith equation. Hence, although GLEAM is based on the less parameterized Priestley-Taylor equation compared to the
635 Penman-Monteith equation, it is outperforming GLDAS-2 and MODIS within this study set-up, which supports the idea to
636 adapt the Penman-Monteith equation as reported by previous studies, e.g., (Widmoser, 2009; Hao et al., 2018; Akumaga and
637 Alderman, 2019; McColl, 2020; Bottazzi et al., 2021).

638 Further, the comparison between estimated coefficient of variation (CV) for 2017 and 2018 showed that the dispersion of data
639 is higher during the extreme drought year 2018 for extreme conditions, such as low SM or low VPD across all SM values. In
640 contrast, 2017 showed higher CV for intermediate conditions. However, the difference between investigated products is rather
641 minor, with median CV between 33.28 % (ERA-land) and 49.23 % (ICOS), and should be analysed in future studies for

individual stations and ecosystem types (requiring longer time series and more stations to have enough data points for binning) for determining the impact of varying vegetation types on ET controls. In summary, when considering all conducted analyses together (spatial and temporal resolutions, product dependencies, ETC results, SM and VPD controls on ET), the remote sensing products SEVIRI and GLEAM as well as reanalysis product ERA5-land seems to provide most reasonable results compared all other ET products, with SEVIRI providing a higher temporal and spatial resolution compared to GLEAM and ERA5-land. Hence, despite their coarse spatial resolution, GLEAM and ERA5-land are able to capture ET dynamics sufficiently even under drought conditions. Future research regarding data fusion techniques and downscaling approaches that combine coarse- or medium-scale ET data with fine-scale auxiliaries in order to improve the spatial resolution of certain ET products may help to decrease the spatial mismatch and optimize the comparison between point-scale field measurements and satellite remote sensing or modelling data.

This study served as a pathfinder to compare freely available and commonly employed ET products at highly monitored EC towers across central Europe. Whether these reported findings hold true across space and for other drought events has to be analysed further with focus on spatially larger regions and longer time series. Additionally, potential add-on studies could include the examination and comparison of ET dynamics from optical/thermal remote sensing observations with microwave remote sensing data, e.g. the Sentinel-1 backscatter, in order to evaluate the potential of active microwave remote sensing for drought monitoring, e.g., (Mueller et al., 2022; Jagdhuber et al., 2023). In order to identify relevant conditions and causal strengths with lagged and contemporaneous causal dependencies between different variables, like ET, the Sentinel-1 backscatter and other important SPAS parameters, like air temperature, relative humidity, and water potentials, the use of emerging powerful tools for causal discovery could prove useful (Runge et al., 2019; Díaz et al., 2022). Previous studies already outlined the potential of identifying causal relations between Earth system parameters (i.e., precipitation, ET, SM, air temperature) by using the wavelet coherency analysis (WCA) (Graf et al., 2014; Rahmati et al., 2020), or the PC algorithm Momentary Conditional Independence (PCMCI) method (Runge et al., 2019, 2023).

Data availability.

The SMAP MT-DCA V5 soil moisture dataset is available at <https://zenodo.org/records/5619583>, last access: 11 May 2022. The SPEI dataset is available at <https://spei.csic.es/database.html>, last access: 18 November 2023. The evapotranspiration products are available as follows: ICOS data are available at <https://www.icos-cp.eu/>, last access: 20 November 2023. SEVIRI data are available at <https://datasasaf.lasvcs.ipma.pt/PRODUCTS/MSG/MDMETv3/>, last access: 21 November 2023. MODIS data are available at <https://lpdaac.usgs.gov/products/mod16a2v061/>, last access: 20 November 2023. ERA5-land data are available at <https://cds.climate.copernicus.eu/datasets/reanalysis-era5-land?tab=overview>, last access: 20 November 2023. The GLDAS-2 data are at <https://ldas.gsfc.nasa.gov/gldas/model-output>, last access: 22 November 2023. The GLEAM data are available at <https://www.gleam.eu/>, last access: 23 August 2024. The Corine land cover classes are available at

675 <https://land.copernicus.eu/en/products/corine-land-cover/clc2018?hash=4ecde146e6ca8dd7a42f68a9f5370153d9731a95>, last
676 access: 14 March 2024.

677

678 *Author contributions.*

679 TJ designed the study concept and assembled the research team. AF, MB, MP, BB, MR, CM, and TJ were involved in the data
680 acquisition and in developing the methodology. AF led the data curation and visualization of results. The original draft was
681 written and prepared by AF and TJ. Draft editing and review were done by all authors.

682

683 *Competing interests.*

684 The contact author has declared that neither they nor their co-authors have any competing interests.

685

686 *Disclaimer.*

687

688 *Acknowledgements.*

689 The authors would like to thank the editor-in-chief, associate editor, and the two anonymous reviewers for their effort and help
690 during the review process.

691

692 *Financial support.*

693 David Chaparro was supported by ‘Fundació La Caixa’ project LCF/BQ/PI23/11970013, and by the H2020 Project
694 FORGENIUS (Improving access to FORest GENetic resources Information and services for end-USers).
695 María Piles thanks the support of *Conselleria de Innovación, Universidades, Ciencia y Sociedad Digital* through the project
696 AI4CS CIPROM/2021/56.

697

698 ~~*Review statement.*~~

699

700 **References**

- 701 Adeluyi, O., Harris, A., Verrelst, J., Foster, T., and Clay, G. D.: Estimating the phenological dynamics of irrigated rice leaf
702 area index using the combination of PROSAIL and Gaussian Process Regression, *International Journal of Applied Earth*
703 *Observation and Geoinformation*, 102, 102454, <https://doi.org/10.1016/j.jag.2021.102454>, 2021.
- 704 Ahmed, K. R., Paul-Limoges, E., Rascher, U., and Damm, A.: A First Assessment of the 2018 European Drought Impact on
705 Ecosystem Evapotranspiration, *Remote Sensing*, 13, 16, <https://doi.org/10.3390/rs13010016>, 2020.
- 706 Akumaga, U. and Alderman, P. D.: Comparison of Penman–Monteith and Priestley-Taylor Evapotranspiration Methods for
707 Crop Modeling in Oklahoma, *Agronomy Journal*, 111, 1171–1180, <https://doi.org/10.2134/agronj2018.10.0694>, 2019.

- 708 Allen, R. G., Pereira, L. S., Raes, D., Smith, M., and others: Crop evapotranspiration-Guidelines for computing crop water
709 requirements-FAO Irrigation and drainage paper 56, Fao, Rome, 300, D05109, 1998.
- 710 Balsamo, G., Beljaars, A., Scipal, K., Viterbo, P., van den Hurk, B., Hirschi, M., and Betts, A. K.: A Revised Hydrology for
711 the ECMWF Model: Verification from Field Site to Terrestrial Water Storage and Impact in the Integrated Forecast System,
712 *Journal of Hydrometeorology*, 10, 623–643, <https://doi.org/10.1175/2008JHM1068.1>, 2009.
- 713 Barrios, J. M., Arboleda, A., Dutra, E., Trigo, I., and Gellens-Meulenberghs, F.: Evapotranspiration and surface energy fluxes
714 across Europe, Africa and Eastern South America throughout the operational life of the Meteosat second generation satellite,
715 *Geoscience Data Journal*, gdj3.235, <https://doi.org/10.1002/gdj3.235>, 2024.
- 716 Bastos, A., Ciais, P., Friedlingstein, P., Sitch, S., Pongratz, J., Fan, L., Wigneron, J. P., Weber, U., Reichstein, M., Fu, Z.,
717 Anthoni, P., Arneth, A., Haverd, V., Jain, A. K., Joetzjer, E., Knauer, J., Lienert, S., Loughran, T., McGuire, P. C., Tian, H.,
718 Viovy, N., and Zaehle, S.: Direct and seasonal legacy effects of the 2018 heat wave and drought on European ecosystem
719 productivity, *Sci. Adv.*, 6, eaba2724, <https://doi.org/10.1126/sciadv.aba2724>, 2020.
- 720 Bayat, B., Camacho, F., Nickeson, J., Cosh, M., Bolten, J., Vereecken, H., and Montzka, C.: Toward operational validation
721 systems for global satellite-based terrestrial essential climate variables, *International Journal of Applied Earth Observation and*
722 *Geoinformation*, 95, 102240, <https://doi.org/10.1016/j.jag.2020.102240>, 2021.
- 723 Bayat, B., Montzka, C., Graf, A., Giuliani, G., Santoro, M., and Vereecken, H.: One decade (2011–2020) of European
724 agricultural water stress monitoring by MSG-SEVIRI: workflow implementation on the Virtual Earth Laboratory (VLab)
725 platform, *International Journal of Digital Earth*, 15, 730–747, <https://doi.org/10.1080/17538947.2022.2061617>, 2022.
- 726 Bayat, B., Raj, R., Graf, A., Vereecken, H., and Montzka, C.: Comprehensive accuracy assessment of long-term geostationary
727 SEVIRI-MSG evapotranspiration estimates across Europe, *Remote Sensing of Environment*, 301, 113875,
728 <https://doi.org/10.1016/j.rse.2023.113875>, 2024.
- 729 Beaudoin, H., Rodell, M., and NASA/GSFC/HSL: GLDAS Noah Land Surface Model L4 3 hourly 0.25 x 0.25 degree,
730 Version 2.1, Greenbelt, Maryland, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC),
731 Accessed: [02 Nov. 2022], <https://doi.org/10.5067/E7TYRXPJKWOQ>, 2020.
- 732 Beguería, S., Vicente Serrano, S. M., Reig-Gracia, F., and Latorre Garcés, B.: SPEIbase v.2.8 [Dataset]; DIGITAL.CSIC;
733 Version 2.8, <https://doi.org/10.20350/DIGITALCSIC/15121>, 2023.
- 734 Bhattacharya, B. K., Mallick, K., Desai, D., Bhat, G. S., Morrison, R., Clevery, J. R., Woodgate, W., Beringer, J., Cawse-
735 Nicholson, K., Ma, S., Verfaillie, J., and Baldocchi, D.: A coupled ground heat flux–surface energy balance model of
736 evaporation using thermal remote sensing observations, *Biogeosciences*, 19, 5521–5551, [https://doi.org/10.5194/bg-19-5521-](https://doi.org/10.5194/bg-19-5521-2022)
737 2022, 2022.
- 738 Bottazzi, M., Bancheri, M., Mobilia, M., Bertoldi, G., Longobardi, A., and Rigon, R.: Comparing Evapotranspiration Estimates
739 from the GEOframe-Prospero Model with Penman–Monteith and Priestley-Taylor Approaches under Different Climate
740 Conditions, *Water*, 13, 1221, <https://doi.org/10.3390/w13091221>, 2021.
- 741 Brown, S. M., Petrone, R. M., Mendoza, C., and Devito, K. J.: Surface vegetation controls on evapotranspiration from a sub-
742 humid Western Boreal Plain wetland, *Hydrological Processes*, 24, 1072–1085, <https://doi.org/10.1002/hyp.7569>, 2010.
- 743 Budyko, M. I. and Miller, D. H.: *Climate and life*, Academic Press, New York, 1974.

744 Carminati, A. and Javaux, M.: Soil Rather Than Xylem Vulnerability Controls Stomatal Response to Drought, *Trends in Plant*
745 *Science*, 25, 868–880, <https://doi.org/10.1016/j.tplants.2020.04.003>, 2020.

746 Carter, E., Hain, C., Anderson, M., and Steinschneider, S.: A Water Balance–Based, Spatiotemporal Evaluation of Terrestrial
747 Evapotranspiration Products across the Contiguous United States, *Journal of Hydrometeorology*, 19, 891–905,
748 <https://doi.org/10.1175/JHM-D-17-0186.1>, 2018.

749 Cawse-Nicholson, K. and Anderson, M.: ECOSTRESS Level-3 DisALEXI-JPL Evapotranspiration (ECO3ETALEXI) User
750 Guide., 2021.

751 Chen, X., Su, Z., Ma, Y., Trigo, I., and Gentile, P.: Remote Sensing of Global Daily Evapotranspiration based on a Surface
752 Energy Balance Method and Reanalysis Data, *JGR Atmospheres*, 126, e2020JD032873,
753 <https://doi.org/10.1029/2020JD032873>, 2021.

754 De Santis, D., D’Amato, C., Bartkowiak, P., Azimi, S., Castelli, M., Rigon, R., and Massari, C.: Evaluation of remotely-sensed
755 evapotranspiration datasets at different spatial and temporal scales at forest and grassland sites in Italy, in: 2022 IEEE
756 Workshop on Metrology for Agriculture and Forestry (MetroAgriFor), 2022 IEEE International Workshop on Metrology for
757 Agriculture and Forestry (MetroAgriFor), Perugia, Italy, 356–361,
758 <https://doi.org/10.1109/MetroAgriFor55389.2022.9964755>, 2022.

759 Díaz, E., Adsua, J. E., Martínez, Á. M., Piles, M., and Camps-Valls, G.: Inferring causal relations from observational long-
760 term carbon and water fluxes records, *Sci Rep*, 12, 1610, <https://doi.org/10.1038/s41598-022-05377-7>, 2022.

761 ECMWF: IFS Documentation CY45R1 - Part IV : Physical processes, <https://doi.org/10.21957/4WHWO8JW0>, 2018.

762 European Environment Agency: CORINE Land Cover 2018 (raster 100 m), Europe, 6-yearly - version 2020_20u1, May 2020
763 (20.01), <https://doi.org/10.2909/960998C1-1870-4E82-8051-6485205EBBAC>, 2019.

764 Feldman, A., Konings, A., Piles, M., and Entekhabi, D.: The Multi-Temporal Dual Channel Algorithm (MT-DCA) (5),
765 <https://doi.org/10.5281/ZENODO.5619583>, 2021.

766 Feldman, A., Gianotti, D., Dong, J., Akbar, R., Crow, W., McColl, K., Nippert, J., Tumber-Dávila, S. J., Holbrook, N. M.,
767 Rockwell, F., Scott, R., Reichle, R., Chatterjee, A., Joiner, J., Poulter, B., and Entekhabi, D.: Satellites capture soil moisture
768 dynamics deeper than a few centimeters and are relevant to plant water uptake, <https://doi.org/10.1002/essoar.10511280.1>, 6
769 May 2022.

770 Feldman, A. F., Feng, X., Felton, A. J., Konings, A. G., Knapp, A. K., Biederman, J. A., and Poulter, B.: Plant responses to
771 changing rainfall frequency and intensity, *Nat Rev Earth Environ*, 5, 276–294, <https://doi.org/10.1038/s43017-024-00534-0>,
772 2024.

773 Fisher, J. B., Tu, K. P., and Baldocchi, D. D.: Global estimates of the land–atmosphere water flux based on monthly AVHRR
774 and ISLSCP-II data, validated at 16 FLUXNET sites, *Remote Sensing of Environment*, 112, 901–919,
775 <https://doi.org/10.1016/j.rse.2007.06.025>, 2008.

776 Fisher, J. B., Lee, B., Purdy, A. J., Halverson, G. H., Dohlen, M. B., Cawse-Nicholson, K., Wang, A., Anderson, R. G., Aragon,
777 B., Arain, M. A., Baldocchi, D. D., Baker, J. M., Barral, H., Bernacchi, C. J., Bernhofer, C., Biraud, S. C., Bohrer, G., Brunzell,
778 N., Cappelare, B., Castro-Contreras, S., Chun, J., Conrad, B. J., Cremonese, E., Demarty, J., Desai, A. R., De Ligne, A.,
779 Foltýnová, L., Goulden, M. L., Griffis, T. J., Grünwald, T., Johnson, M. S., Kang, M., Kelbe, D., Kowalska, N., Lim, J.,
780 Maïnassara, I., McCabe, M. F., Missik, J. E. C., Mohanty, B. P., Moore, C. E., Morillas, L., Morrison, R., Munger, J. W.,
781 Posse, G., Richardson, A. D., Russell, E. S., Ryu, Y., Sanchez-Azofeifa, A., Schmidt, M., Schwartz, E., Sharp, I., Šigut, L.,

782 Tang, Y., Hulley, G., Anderson, M., Hain, C., French, A., Wood, E., and Hook, S.: ECOSTRESS: NASA's Next Generation
783 Mission to Measure Evapotranspiration From the International Space Station, *Water Resources Research*, 56,
784 e2019WR026058, <https://doi.org/10.1029/2019WR026058>, 2020.

785 Fu, Z., Ciais, P., Prentice, I. C., Gentine, P., Makowski, D., Bastos, A., Luo, X., Green, J. K., Stoy, P. C., Yang, H., and Hajima,
786 T.: Atmospheric dryness reduces photosynthesis along a large range of soil water deficits, *Nat Commun*, 13, 989,
787 <https://doi.org/10.1038/s41467-022-28652-7>, 2022.

788 Ghilain, N., Arboleda, A., and Gellens-Meulenberghs, F.: Evapotranspiration modelling at large scale using near-real time
789 MSG SEVIRI derived data, *Hydrol. Earth Syst. Sci.*, 15, 771–786, <https://doi.org/10.5194/hess-15-771-2011>, 2011.

790 Graf, A., Bogena, H. R., Drüe, C., Hardelauf, H., Pütz, T., Heinemann, G., and Vereecken, H.: Spatiotemporal relations
791 between water budget components and soil water content in a forested tributary catchment, *Water Resources Research*, 50,
792 4837–4857, <https://doi.org/10.1002/2013WR014516>, 2014.

793 Gruber, A., Su, C. -H., Crow, W. T., Zwieback, S., Dorigo, W. A., and Wagner, W.: Estimating error cross-correlations in soil
794 moisture data sets using extended collocation analysis, *JGR Atmospheres*, 121, 1208–1219,
795 <https://doi.org/10.1002/2015JD024027>, 2016.

796 Gupta, A., Rico-Medina, A., and Caño-Delgado, A. I.: The physiology of plant responses to drought, *Science*, 368, 266–269,
797 <https://doi.org/10.1126/science.aaz7614>, 2020.

798 Ha, W., Gowda, P. H., and Howell, T. A.: A review of downscaling methods for remote sensing-based irrigation management:
799 part I, *Irrig Sci*, 31, 831–850, <https://doi.org/10.1007/s00271-012-0331-7>, 2013.

800 Hao, X., Zhang, S., Li, W., Duan, W., Fang, G., Zhang, Y., and Guo, B.: The Uncertainty of Penman-Monteith Method and
801 the Energy Balance Closure Problem, *JGR Atmospheres*, 123, 7433–7443, <https://doi.org/10.1029/2018JD028371>, 2018.

802 He, Q.-L., Xiao, J.-L., and Shi, W.-Y.: Responses of Terrestrial Evapotranspiration to Extreme Drought: A Review, *Water*,
803 14, 3847, <https://doi.org/10.3390/w14233847>, 2022.

804 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers,
805 D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara,
806 G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L.,
807 Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P.,
808 Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.: The ERA5 global reanalysis, *Quart J Royal Meteor Soc*, 146, 1999–
809 2049, <https://doi.org/10.1002/qj.3803>, 2020.

810 Hu, G., Jia, L., and Menenti, M.: Comparison of MOD16 and LSA-SAF MSG evapotranspiration products over Europe for
811 2011, *Remote Sensing of Environment*, 156, 510–526, <https://doi.org/10.1016/j.rse.2014.10.017>, 2015.

812 Hu, T., Mallick, K., Hitzelberger, P., Didry, Y., Boulet, G., Szantoi, Z., Koetz, B., Alonso, I., Pascolini-Campbell, M.,
813 Halverson, G., Cawse-Nicholson, K., Hulley, G. C., Hook, S., Bhattarai, N., Oliso, A., Roujean, J., Gamet, P., and Su, B.:
814 Evaluating European ECOSTRESS Hub Evapotranspiration Products Across a Range of Soil-Atmospheric Aridity and Biomes
815 Over Europe, *Water Resources Research*, 59, e2022WR034132, <https://doi.org/10.1029/2022WR034132>, 2023.

816 ICOS RI, Aalto, J., Aalto, P., Aaltonen, H., Aigui, T., Akubia, J., Ala-Könni, J., Alivernini, A., Aluome, C., Andersson, T.,
817 Arca, A., Arriga, N., Aurela, M., BRECHET, L., Baab, F., Back, J., Baltes, U., Baneschi, I., Barten, S., Baur, T., Bauters, M.,
818 Bazot, S., Beauclair, P., Becker, N., Beletti Marchesini, L., Bergström, G., Bernhofer, C., Berveiller, D., Biermann, T.,
819 Bignotti, L., Biron, R., Bloor, J., Bodson, B., Boeckx, P., Bogaerts, G., Bonal, D., Boon, G., Bornet, F., Bortoli, M., Bosio, I.,

820 Brut, A., Brümmer, C., Buchmann, N., Bulonza, E., Burban, B., Buysse, P., Bâth, A., Calandrelli, D., Calvet, J.-C., Canut-
821 Rocafort, G., Carrara, A., Cavagna, M., Ceschia, E., Chabbi, A., Chan, T., Chebbi, W., Chianucci, F., Chipeaux, C., Chopin,
822 H., Christen, A., Chrysoulakis, N., Claverie, N., Cobbe, I., Cohard, J.-M., Colosse, D., Conte, A., Corsanici, R., Coulaud, C.,
823 Courtois, P., Coyle, M., Cremonese, E., Crill, P., Cuntz, M., Cuocolo, D., Czerný, R., DEPUYDT, J., Daelman, R., Darenová,
824 E., Darsonville, O., De Ligne, A., De Meulder, T., De Simon, G., Decau, M.-L., Dell'Acqua, A., Delorme, J.-P., Delpierre, N.,
825 Demoulin, L., Denou, J.-L., Di Tommasi, P., Dienstbach, L., Dignam, R., Dolfus, D., Domec, J.-C., Douxfils, B., Drösler, M.,
826 Drüe, C., Dufrêne, E., Dumont, B., Durand, B., et al.: Ecosystem final quality (L2) product in ETC-Archive format - release
827 2024-1, <https://doi.org/10.18160/G5KZ-ZD83>, 22 May 2024.

828 Jagdhuber, T., Fluhrer, A., Chaparro, D., Dubois, C., Hellwig, F. M., Bayat, B., Montzka, C., Baur, M. J., Ramati, M., Kübert,
829 A., Mueller, M. M., Schellenberg, K., Boehm, M., Jonard, F., Steele-Dunne, S., Piles, M., and Entekhabi, D.: On the Potential
830 of Active and Passive Microwave Remote Sensing for Tracking Seasonal Dynamics of Evapotranspiration, in: IGARSS 2023
831 - 2023 IEEE International Geoscience and Remote Sensing Symposium, IGARSS 2023 - 2023 IEEE International Geoscience
832 and Remote Sensing Symposium, Pasadena, CA, USA, 2610–2613, <https://doi.org/10.1109/IGARSS52108.2023.10283234>,
833 2023.

834 Jiménez, C., Prigent, C., Mueller, B., Seneviratne, S. I., McCabe, M. F., Wood, E. F., Rossow, W. B., Balsamo, G., Betts, A.
835 K., Dirmeyer, P. A., Fisher, J. B., Jung, M., Kanamitsu, M., Reichle, R. H., Reichstein, M., Rodell, M., Sheffield, J., Tu, K.,
836 and Wang, K.: Global intercomparison of 12 land surface heat flux estimates, *J. Geophys. Res.*, 116, D02102,
837 <https://doi.org/10.1029/2010JD014545>, 2011.

838 Jin, Z., Liang, W., Yang, Y., Zhang, W., Yan, J., Chen, X., Li, S., and Mo, X.: Separating Vegetation Greening and Climate
839 Change Controls on Evapotranspiration trend over the Loess Plateau, *Sci Rep*, 7, 8191, [https://doi.org/10.1038/s41598-017-](https://doi.org/10.1038/s41598-017-08477-x)
840 08477-x, 2017.

841 Jung, M., Reichstein, M., Ciais, P., Seneviratne, S. I., Sheffield, J., Goulden, M. L., Bonan, G., Cescatti, A., Chen, J., De Jeu,
842 R., Dolman, A. J., Eugster, W., Gerten, D., Gianelle, D., Gobron, N., Heinke, J., Kimball, J., Law, B. E., Montagnani, L., Mu,
843 Q., Mueller, B., Oleson, K., Papale, D., Richardson, A. D., Rouspard, O., Running, S., Tomelleri, E., Viovy, N., Weber, U.,
844 Williams, C., Wood, E., Zaehle, S., and Zhang, K.: Recent decline in the global land evapotranspiration trend due to limited
845 moisture supply, *Nature*, 467, 951–954, <https://doi.org/10.1038/nature09396>, 2010.

846 Jung, M., Koirala, S., Weber, U., Ichii, K., Gans, F., Camps-Valls, G., Papale, D., Schwalm, C., Tramontana, G., and
847 Reichstein, M.: The FLUXCOM ensemble of global land-atmosphere energy fluxes, *Sci Data*, 6, 74,
848 <https://doi.org/10.1038/s41597-019-0076-8>, 2019.

849 Konings, A., Piles, M., Rötzer, M., McColl, K., Chang, S. K., and Entekhabi, D.: Vegetation optical depth and scattering
850 albedo retrieval using time series of dual-polarized L-band radiometer observations, *Elsevier Remote Sensing of Environment*,
851 172, 178–189, <https://doi.org/10.1016/j.rse.2015.11.009>, 2016.

852 Li, C., Yang, H., Yang, W., Liu, Z., Jia, Y., Li, S., and Yang, D.: Error characterization of global land evapotranspiration
853 products: Collocation-based approach, *Journal of Hydrology*, 612, 128102, <https://doi.org/10.1016/j.jhydrol.2022.128102>,
854 2022.

855 Liu, H., Xin, X., Su, Z., Zeng, Y., Lian, T., Li, L., Yu, S., and Zhang, H.: Intercomparison and evaluation of ten global ET
856 products at site and basin scales, *Journal of Hydrology*, 617, <https://doi.org/10.1016/j.jhydrol.2022.128887>, 2023.

857 Liu, L., Gudmundsson, L., Hauser, M., Qin, D., Li, S., and Seneviratne, S. I.: Soil moisture dominates dryness stress on
858 ecosystem production globally, *Nat Commun*, 11, 4892, <https://doi.org/10.1038/s41467-020-18631-1>, 2020.

859 Liu, N., Oishi, A. C., Miniat, C. F., and Bolstad, P.: An evaluation of ECOSTRESS products of a temperate montane humid
860 forest in a complex terrain environment, *Remote Sensing of Environment*, 265, 112662,
861 <https://doi.org/10.1016/j.rse.2021.112662>, 2021.

862 Loustau, D., Chipeaux, C., and ICOS Ecosystem Thematic Centre: Warm winter 2020 ecosystem eddy covariance flux product
863 from Bilos (1.0), <https://doi.org/10.18160/MSRT-T1YA>, 2022.

864 LSA SAF and EUMETSAT SAF On Land Surface Analysis: MSG Evapotranspiration Version 3 (Metv3), available at
865 <https://data.saf.saf.gov/PRODUCTS/MSG/METv3/>, 2019.

866 Mahour, M., Tolpekin, V., Stein, A., and Sharifi, A.: A comparison of two downscaling procedures to increase the spatial
867 resolution of mapping actual evapotranspiration, *ISPRS Journal of Photogrammetry and Remote Sensing*, 126, 56–67,
868 <https://doi.org/10.1016/j.isprsjprs.2017.02.004>, 2017.

869 Martens, B., Miralles, D. G., Lievens, H., van der Schalie, R., de Jeu, R. A. M., Fernández-Prieto, D., Beck, H. E., Dorigo, W.
870 A., and Verhoest, N. E. C.: GLEAM v3: satellite-based land evaporation and root-zone soil moisture, *Geosci. Model Dev.*, 10,
871 1903–1925, <https://doi.org/10.5194/gmd-10-1903-2017>, 2017.

872 McColl, K. A.: Practical and Theoretical Benefits of an Alternative to the Penman-Monteith Evapotranspiration Equation,
873 *Water Resources Research*, 56, e2020WR027106, <https://doi.org/10.1029/2020WR027106>, 2020.

874 McColl, K. A., Vogelzang, J., Konings, A. G., Entekhabi, D., Piles, M., and Stoffelen, A.: Extended triple collocation:
875 Estimating errors and correlation coefficients with respect to an unknown target, *Geophysical Research Letters*, 41, 6229–
876 6236, <https://doi.org/10.1002/2014GL061322>, 2014.

877 Meng, X., Deng, M., Shu, L., Chen, H., Wang, S., Li, Z., Zhao, L., and Shang, L.: An evaluation of evapotranspiration products
878 over the Tibetan Plateau, *Journal of Hydrometeorology*, <https://doi.org/10.1175/JHM-D-23-0223.1>, 2024.

879 Miralles, D. G., Holmes, T. R. H., De Jeu, R. A. M., Gash, J. H., Meesters, A. G. C. A., and Dolman, A. J.: Global land-surface
880 evaporation estimated from satellite-based observations, *Hydrol. Earth Syst. Sci.*, 15, 453–469, [https://doi.org/10.5194/hess-](https://doi.org/10.5194/hess-15-453-2011)
881 15-453-2011, 2011.

882 Monteith, J. L.: Evaporation and environment, *Symposia of the society for experimental biology*, 19, 1965.

883 Mueller, B., Hirschi, M., Jimenez, C., Ciais, P., Dirmeyer, P. A., Dolman, A. J., Fisher, J. B., Jung, M., Ludwig, F., Maignan,
884 F., Miralles, D. G., McCabe, M. F., Reichstein, M., Sheffield, J., Wang, K., Wood, E. F., Zhang, Y., and Seneviratne, S. I.:
885 Benchmark products for land evapotranspiration: LandFlux-EVAL multi-data set synthesis, *Hydrol. Earth Syst. Sci.*, 17, 3707–
886 3720, <https://doi.org/10.5194/hess-17-3707-2013>, 2013.

887 Mueller, M. M., Dubois, C., Jagdhuber, T., Hellwig, F. M., Pathe, C., Schmullius, C., and Steele-Dunne, S.: Sentinel-1
888 Backscatter Time Series for Characterization of Evapotranspiration Dynamics over Temperate Coniferous Forests, *Remote*
889 *Sensing*, 14, 6384, <https://doi.org/10.3390/rs14246384>, 2022.

890 Muñoz Sabater, J.: ERA5-Land hourly data from 1981 to present. Copernicus Climate Change Service (C3S) Climate Data
891 Store (CDS). (Accessed on 10-08-2022), <https://doi.org/10.24381/CDS.E2161BAC>, 2019.

892 Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M.,
893 Harrigan, S., Hersbach, H., Martens, B., Miralles, D. G., Piles, M., Rodríguez-Fernández, N. J., Zsoter, E., Buontempo, C.,
894 and Thépaut, J.-N.: ERA5-Land: a state-of-the-art global reanalysis dataset for land applications, *Earth Syst. Sci. Data*, 13,
895 4349–4383, <https://doi.org/10.5194/essd-13-4349-2021>, 2021.

896 Novick, K. A., Ficklin, D. L., Stoy, P. C., Williams, C. A., Bohrer, G., Oishi, A. C., Papuga, S. A., Blanken, P. D., Noormets,
897 A., Sulman, B. N., Scott, R. L., Wang, L., and Phillips, R. P.: The increasing importance of atmospheric demand for ecosystem
898 water and carbon fluxes, *Nature Clim Change*, 6, 1023–1027, <https://doi.org/10.1038/nclimate3114>, 2016.

899 Pastorello, G., Poindexter, C., Chen, J., Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Reichstein, M., Ribeca, A., van
900 Ingen, C., Vuichard, N., Zhang, L., Amiro, B., Ammann, C., Arain, M. A., Ardö, J., Arkebauer, T., Arndt, S. K., Arriga, N.,
901 Aubinet, M., Aurela, M., Baldocchi, D., Barr, A., Beamesderfer, E., Marchesini, L. B., Bergeron, O., Beringer, J., Bernhofer,
902 C., Berveiller, D., Billesbach, D., Black, T. A., Blanken, P. D., Bohrer, G., Boike, J., Bolstad, P. V., Bonal, D., Bonnefond, J.-
903 M., Bowling, D. R., Bracho, R., Brodeur, J., Brümmer, C., Buchmann, N., Burban, B., Burns, S. P., Buysse, P., Cale, P.,
904 Cavagna, M., Cellier, P., Chen, S., Chini, I., Christensen, T. R., Cleverly, J., Collalti, A., Consalvo, C., Cook, B. D., Cook, D.,
905 Coursolle, C., Cremonese, E., Curtis, P. S., D’Andrea, E., da Rocha, H., Dai, X., Davis, K. J., Cinti, B. D., Grandcourt, A. de,
906 Ligne, A. D., De Oliveira, R. C., Delpierre, N., Desai, A. R., Di Bella, C. M., Tommasi, P. di, Dolman, H., Domingo, F., Dong,
907 G., Dore, S., Duce, P., Dufrêne, E., Dunn, A., Dušek, J., Eamus, D., Eichelmann, U., ElKhidir, H. A. M., Eugster, W., Ewenz,
908 C. M., Ewers, B., Famulari, D., Fares, S., Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G., Fischer, M., Frank, J., Galvagno,
909 M., Gharun, M., Gianelle, D., Gielen, B., Gioli, B., Gitelson, A., et al.: The FLUXNET2015 dataset and the ONEFlux
910 processing pipeline for eddy covariance data, *Scientific Data*, 7, 225, <https://doi.org/10.1038/s41597-020-0534-3>, 2020.

911 Peng, J., Loew, A., Merlin, O., and Verhoest, N. E. C.: A review of spatial downscaling of satellite remotely sensed soil
912 moisture, *Reviews of Geophysics*, 55, 341–366, <https://doi.org/10.1002/2016RG000543>, 2017.

913 Penman, H. L.: Natural evaporation from open water, bare soil and grass, *Proc. R. Soc. Lond. A*, 193, 120–145,
914 <https://doi.org/10.1098/rspa.1948.0037>, 1948.

915 Petropoulos, G. P., Ireland, G., Cass, A., and Srivastava, P. K.: Performance Assessment of the SEVIRI Evapotranspiration
916 Operational Product: Results Over Diverse Mediterranean Ecosystems, *IEEE Sensors J.*, 15, 3412–3423,
917 <https://doi.org/10.1109/JSEN.2015.2390031>, 2015.

918 Porporato, A., D’Odorico, P., Laio, F., Ridolfi, L., and Rodriguez-Iturbe, I.: Ecohydrology of water-controlled ecosystems,
919 *Advances in Water Resources*, 25, 1335–1348, [https://doi.org/10.1016/S0309-1708\(02\)00058-1](https://doi.org/10.1016/S0309-1708(02)00058-1), 2002.

920 Priestley, C. H. B. and Taylor, R. J.: On the Assessment of Surface Heat Flux and Evaporation Using Large-Scale Parameters,
921 *Mon. Wea. Rev.*, 100, 81–92, [https://doi.org/10.1175/1520-0493\(1972\)100<0081:OTAOSH>2.3.CO;2](https://doi.org/10.1175/1520-0493(1972)100<0081:OTAOSH>2.3.CO;2), 1972.

922 Rahmati, M., Groh, J., Graf, A., Pütz, T., Vanderborght, J., and Vereecken, H.: On the impact of increasing drought on the
923 relationship between soil water content and evapotranspiration of a grassland, *Vadose Zone Journal*, 19, e20029,
924 <https://doi.org/10.1002/vzj2.20029>, 2020.

925 Rahmati, M., Graf, A., Poppe Terán, C., Amelung, W., Dorigo, W., Franssen, H.-J. H., Montzka, C., Or, D., Sprenger, M.,
926 Vanderborght, J., Verhoest, N. E. C., and Vereecken, H.: Continuous increase in evaporative demand shortened the growing
927 season of European ecosystems in the last decade, *Commun Earth Environ*, 4, 236, [https://doi.org/10.1038/s43247-023-00890-](https://doi.org/10.1038/s43247-023-00890-7)
928 7, 2023.

929 Rahmati, M., Amelung, W., Brogi, C., Dari, J., Flammini, A., Bogen, H., Brocca, L., Chen, H., Groh, J., Koster, R. D.,
930 McColl, K. A., Montzka, C., Moradi, S., Rahi, A., Sharghi S., F., and Vereecken, H.: Soil Moisture Memory: State-Of-The-
931 Art and the Way Forward, *Reviews of Geophysics*, 62, e2023RG000828, <https://doi.org/10.1029/2023RG000828>, 2024.

932 Rakovec, O., Samaniego, L., Hari, V., Markonis, Y., Moravec, V., Thober, S., Hanel, M., and Kumar, R.: The 2018–2020
933 Multi-Year Drought Sets a New Benchmark in Europe, *Earth’s Future*, 10, e2021EF002394,
934 <https://doi.org/10.1029/2021EF002394>, 2022.

935 Rebmann, C., Aubinet, M., Schmid, H., Arriga, N., Aurela, M., Burba, G., Clement, R., De Ligne, A., Fratini, G., Gielen, B.,
936 Grace, J., Graf, A., Gross, P., Haapanala, S., Herbst, M., Hörtnagl, L., Ibrom, A., Joly, L., Kljun, N., Kolle, O., Kowalski, A.,
937 Lindroth, A., Loustau, D., Mammarella, I., Mauder, M., Merbold, L., Metzger, S., Mölder, M., Montagnani, L., Papale, D.,
938 Pavelka, M., Peichl, M., Roland, M., Serrano-Ortiz, P., Siebicke, L., Steinbrecher, R., Tuovinen, J.-P., Vesala, T., Wohlfahrt,
939 G., and Franz, D.: ICOS eddy covariance flux-station site setup: a review, *International Agrophysics*, 32, 471–494,
940 <https://doi.org/10.1515/intag-2017-0044>, 2018.

941 Rui, H. and Beaudoin, H.: README Document for NASA GLDAS Version 2 Data Products, NASA Goddard Earth Sciences
942 Data and Information Services Center (GES DISC), 2022.

943 Runge, J., Nowack, P., Kretschmer, M., Flaxman, S., and Sejdinovic, D.: Detecting and quantifying causal associations in
944 large nonlinear time series datasets, *Sci. Adv.*, 5, eaau4996, <https://doi.org/10.1126/sciadv.aau4996>, 2019.

945 Runge, J., Gerhardus, A., Varando, G., Eyring, V., and Camps-Valls, G.: Causal inference for time series, *Nat Rev Earth*
946 *Environ*, 4, 487–505, <https://doi.org/10.1038/s43017-023-00431-y>, 2023.

947 Running, S., Mu, Q., and Zhao, M.: MOD16A2 MODIS/Terra Net Evapotranspiration 8-Day L4 Global 500m SIN Grid V006,
948 <https://doi.org/10.5067/MODIS/MOD16A2.006>, 2017.

949 Running, S., Mu, Q., Zhao, M., and Moreno, A.: User’s guide MODIS global terrestrial evapotranspiration (ET) product
950 (MOD16A2/A3 and year-end gap-filled MOD16A2GF/A3GF), MODIS Land Team 40, 2019.

951 Savitzky, Abraham. and Golay, M. J. E.: Smoothing and Differentiation of Data by Simplified Least Squares Procedures.,
952 *Anal. Chem.*, 36, 1627–1639, <https://doi.org/10.1021/ac60214a047>, 1964.

953 Schuldt, B., Buras, A., Arend, M., Vitasse, Y., Beierkuhnlein, C., Damm, A., Gharun, M., Grams, T. E. E., Hauck, M., Hajek,
954 P., Hartmann, H., Hiltbrunner, E., Hoch, G., Holloway-Phillips, M., Körner, C., Larysch, E., Lübke, T., Nelson, D. B.,
955 Rammig, A., Rigling, A., Rose, L., Ruehr, N. K., Schumann, K., Weiser, F., Werner, C., Wohlgemuth, T., Zang, C. S., and
956 Kahmen, A.: A first assessment of the impact of the extreme 2018 summer drought on Central European forests, *Basic and*
957 *Applied Ecology*, 45, 86–103, <https://doi.org/10.1016/j.baee.2020.04.003>, 2020.

958 Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., Orlowsky, B., and Teuling, A. J.: Investigating
959 soil moisture–climate interactions in a changing climate: A review, *Earth-Science Reviews*, 99, 125–161,
960 <https://doi.org/10.1016/j.earscirev.2010.02.004>, 2010.

961 Sepulcre-Canto, G., Vogt, J., Arboleda, A., and Antofie, T.: Assessment of the EUMETSAT LSA-SAF evapotranspiration
962 product for drought monitoring in Europe, *International Journal of Applied Earth Observation and Geoinformation*, 30, 190–
963 202, <https://doi.org/10.1016/j.jag.2014.01.021>, 2014.

964 Singh, R. P., Paramanik, S., Bhattacharya, B. K., and Behera, M. D.: Modelling of evapotranspiration using land surface energy
965 balance and thermal infrared remote sensing, *Trop Ecol*, 61, 42–50, <https://doi.org/10.1007/s42965-020-00076-8>, 2020.

966 Stisen, S., Soltani, M., Mendiguren, G., Langkilde, H., Garcia, M., and Koch, J.: Spatial Patterns in Actual Evapotranspiration
967 *Climatologies for Europe*, *Remote Sensing*, 13, 2410, <https://doi.org/10.3390/rs13122410>, 2021.

968 Stoffelen, A.: Toward the true near-surface wind speed: Error modeling and calibration using triple collocation, *J. Geophys.*
969 *Res.*, 103, 7755–7766, <https://doi.org/10.1029/97JC03180>, 1998.

970 The EUMETSAT Satellite Application Facility on Land Surface Analysis (LSA SAF): Validation Report Evapotranspiration
971 & Turbulent Fluxes v3; SAF/LAND/RMI/VR/ETFv3/1.0; Issue: 2, 2024.

972 Trambauer, P., Dutra, E., Maskey, S., Werner, M., Pappenberger, F., Van Beek, L. P. H., and Uhlenbrook, S.: Comparison of
973 different evaporation estimates over the African continent, *Hydrol. Earth Syst. Sci.*, 18, 193–212, [https://doi.org/10.5194/hess-](https://doi.org/10.5194/hess-18-193-2014)
974 18-193-2014, 2014.

975 Twine, T. E., Kustas, W. P., Norman, J. M., Cook, D. R., Houser, P. R., Meyers, T. P., Prueger, J. H., Starks, P. J., and Wesely,
976 M. L.: Correcting eddy-covariance flux underestimates over a grassland, *Agricultural and Forest Meteorology*, 103, 279–300,
977 [https://doi.org/10.1016/S0168-1923\(00\)00123-4](https://doi.org/10.1016/S0168-1923(00)00123-4), 2000.

978 Vargas Zeppetello, L. R., McColl, K. A., Bernau, J. A., Bowen, B. B., Tang, L. I., Holbrook, N. M., Gentine, P., and Huybers,
979 P.: Apparent surface conductance sensitivity to vapour pressure deficit in the absence of plants, *Nat Water*, 1, 941–951,
980 <https://doi.org/10.1038/s44221-023-00147-9>, 2023.

981 Vicente-Serrano, S. M., Beguería, S., and López-Moreno, J. I.: A Multiscalar Drought Index Sensitive to Global Warming:
982 The Standardized Precipitation Evapotranspiration Index, *Journal of Climate*, 23, 1696–1718,
983 <https://doi.org/10.1175/2009JCLI2909.1>, 2010.

984 Wang, K. and Dickinson, R. E.: A review of global terrestrial evapotranspiration: Observation, modeling, climatology, and
985 climatic variability, *Reviews of Geophysics*, 50, 2011RG000373, <https://doi.org/10.1029/2011RG000373>, 2012.

986 Warm Winter 2020 Team, ICOS Ecosystem Thematic Centre, ICOS Ecosystem Thematic Centre, and Trotta, C.: Warm Winter
987 2020 ecosystem eddy covariance flux product for 73 stations in FLUXNET-Archive format—release 2022-1,
988 <https://doi.org/10.18160/2G60-ZHAK>, 1 February 2022.

989 Widmoser, P.: A discussion on and alternative to the Penman–Monteith equation, *Agricultural Water Management*, 96, 711–
990 721, <https://doi.org/10.1016/j.agwat.2008.10.003>, 2009.

991 Wu, J., Feng, Y., Liang, L., He, X., and Zeng, Z.: Assessing evapotranspiration observed from ECOSTRESS using flux
992 measurements in agroecosystems, *Agricultural Water Management*, 269, 107706,
993 <https://doi.org/10.1016/j.agwat.2022.107706>, 2022.

994 Xu, C., Wang, W., Hu, Y., and Liu, Y.: Evaluation of ERA5, ERA5-Land, GLDAS-2.1, and GLEAM potential
995 evapotranspiration data over mainland China, *Journal of Hydrology: Regional Studies*, 51, 101651,
996 <https://doi.org/10.1016/j.ejrh.2023.101651>, 2024.

997 Xu, T., Guo, Z., Xia, Y., Ferreira, V. G., Liu, S., Wang, K., Yao, Y., Zhang, X., and Zhao, C.: Evaluation of twelve
998 evapotranspiration products from machine learning, remote sensing and land surface models over conterminous United States,
999 *Journal of Hydrology*, 578, 124105, <https://doi.org/10.1016/j.jhydrol.2019.124105>, 2019.

1000 Yu, X., Qian, L., Wang, W., Hu, X., Dong, J., Pi, Y., and Fan, K.: Comprehensive evaluation of terrestrial evapotranspiration
1001 from different models under extreme condition over conterminous United States, *Agricultural Water Management*, 289,
1002 108555, <https://doi.org/10.1016/j.agwat.2023.108555>, 2023.

1003 Zhang, J., Guan, K., Peng, B., Pan, M., Zhou, W., Jiang, C., Kimm, H., Franz, T. E., Grant, R. F., Yang, Y., Rudnick, D. R.,
1004 Heeren, D. M., Suyker, A. E., Bauerle, W. L., and Miner, G. L.: Sustainable irrigation based on co-regulation of soil water
1005 supply and atmospheric evaporative demand, *Nat Commun*, 12, 5549, <https://doi.org/10.1038/s41467-021-25254-7>, 2021.

1006 Zhang, K., Kimball, J. S., and Running, S. W.: A review of remote sensing based actual evapotranspiration estimation, *WIREs*
1007 *Water*, 3, 834–853, <https://doi.org/10.1002/wat2.1168>, 2016.

1008 Zhao, M., A. G., Liu, Y., and Konings, A. G.: Evapotranspiration frequently increases during droughts, *Nat. Clim. Chang.*, 12,
1009 1024–1030, <https://doi.org/10.1038/s41558-022-01505-3>, 2022.

1010 Zhou, S., Yu, B., Zhang, Y., Huang, Y., and Wang, G.: Partitioning evapotranspiration based on the concept of underlying
1011 water use efficiency, *Water Resources Research*, 52, 1160–1175, <https://doi.org/10.1002/2015WR017766>, 2016.

1012 Zhou, S., Zhang, Y., Park Williams, A., and Gentile, P.: Projected increases in intensity, frequency, and terrestrial carbon
1013 costs of compound drought and aridity events, *Sci. Adv.*, 5, eaau5740, <https://doi.org/10.1126/sciadv.aau5740>, 2019.

1014