

Atmospheric moisture tracking with WAM2layers v3

Peter Kalverla¹, Imme Benedict², Chris Weijenborg², and Ruud J. van der Ent³

¹Netherlands eScience Center, Amsterdam, Netherlands

²Meteorology and Air Quality Group, Wageningen University and Research, Wageningen, Netherlands

³Department of Water Management, Faculty of Civil Engineering and Geosciences, Delft University of Technology, Delft, Netherlands

Correspondence: Peter Kalverla (p.kalverla@esciencecenter.nl) or Ruud J. van der Ent (r.j.vanderent@tudelft.nl)

Abstract. This manuscript documents the atmospheric moisture tracking model WAM2layers v3 (Water Accounting Model - 2 layers - version 3). WAM2layers may be used to gain understanding of atmospheric dynamics and to study rainfall patterns and extremes by mapping their sources or sinks, often in the context of climate and land-use changes. To this end, WAM2layers solves a prognostic equation for tagged moisture in gridded atmospheric datasets such as reanalysis data or climate model output. WAM2layers can be used in forward mode to determine where evaporated water eventually precipitates, or in backward mode to determine where precipitation originally evaporated.

WAM2layers v3 represents a complete rewrite of the WAM2layers model originally introduced in 2010 and subsequently used in more than sixty academic studies. This latest version incorporates performance optimisations to cope with the increased resolution of input data, and introduces various best practices aimed at improved user-friendliness and software sustainability. Since an increasing number of researchers is using the code, this manuscript is intended as an updated description and reference in the academic literature. After describing the history, model formulation, and numerical implementation, we present and evaluate two example cases to illustrate the use and skill of WAM2layers v3. We then discuss best practices, some important assumptions, and directions for future development.

1 Introduction

This paper documents the Water Accounting Model – 2 layers version 3, or WAM2layers v3 for short. WAM2layers is a moisture tracking model that can be used to study the transport of water in the atmosphere, from source (surface evaporation) to sink (precipitation) or vice versa. Since the previous version (see Van der Ent et al., 2014, Appendix B), the model has seen some substantial upgrades, and this manuscript is intended as an updated reference for those who have used, are using, or consider to use WAM2layers in their research. Before we delve into details, we briefly consider the scientific and historical context.

Moisture tracking in general has been applied in different fields (e.g., Gimeno et al., 2020) and research objectives range from obtaining better process understanding of specific events (e.g., Dirmeyer and Brubaker, 1999) and different modes of climate variability (e.g., Miralles et al., 2016) to general aspects of the global hydrological cycle (e.g., Theeuwens et al., 2023).




Figure 1. A: Example tracking domain (green outline) and tagging region (red outline and fill) for WAM2layers overlaid on a global input grid. B: Conceptual illustration of the inner structure of WAM2layers. Green arrows indicate horizontal transport, red the vertical flux, blue is precipitation and purple is evaporation. Horizontal transport in the upper layer (not drawn) is similar to that in the lower layer. Yellow square represents the surface.

Moreover, moisture tracking results have often been used in the context of studying the role of vegetation, potential effects of
 25 land-use changes and possible management options (e.g., Te Wierik et al., 2021).

Various types of moisture tracking models exist. A potential classification system was given in Dominguez et al. (2020, Fig. 1), who distinguished between analytical, offline numerical, and online numerical methods. Alternatively, numerical models can be distinguished based on the way they formulate the transport problem (Eulerian vs Lagrangian), such as done in the review paper by Gimeno et al. (2020). Other differences may arise due to the description of the land surface interaction,
 30 numerous assumptions and simplifications, and implementation details. The effect of the differences in modelling approach and the simplifications and assumptions has received some attention in literature (e.g. Cloux et al., 2021; Crespo-Otero et al., 2024; Goessling and Reick, 2013; Li et al., 2024; Van der Ent et al., 2013) and a larger intercomparison study is underway (Benedict et al., 2024).

WAM2layers is an offline, Eulerian atmospheric moisture tracking model. A conceptual illustration of WAM2layers is provided in Figure 1, which highlights the domain configuration and the transport between the grid cells and exchange with the surface. The model setup, including simplifications, assumptions and limitations are described in detail in Sections 2–4 and 6.

WAM2layers was originally developed as the single-layer Water Accounting Model (WAM v1; Van der Ent et al., 2010), driven by ERA-Interim data. The two-layer version was introduced because Van der Ent et al. (2013), who compared the capabilities of WAM to an online tracking method in a regional climate model, found that the one-layer version resulted in large errors for cases with strong directional wind shear. After further adjustments for use with global climate and reanalysis data, the version described in Van der Ent et al. (2014, appendix B) is referred to as WAM2layers v1. Originally written in MATLAB, the first published work with a full rewrite into Python was by Van der Ent and Tuinenburg (2017, WAM2layers v2). Hence, the update to the model described here is considered to be WAM2layers v3. This major rewrite was aimed at improving the code efficiency in order to keep up with the increased resolution of weather and climate model output. As of v3, we started using semantic versioning (<https://semver.org/>), which means each software release gets a three-number identifier using the format MAJOR.MINOR.PATCH. Within a major version, we strive to maintain backward compatibility, minimizing disruptions to existing workflows with WAM2layers as new features and fixes are introduced. This manuscript pertains to major version v3, including all its minor and patch releases. Each release gets its own DOI to enable referencing specific versions of the code. Additionally, a concept DOI is available for generic references to the software regardless of a specific version van der Ent et al. (2024a).

Since its introduction, WAM2layers has been used by many researchers and applied to different topics (Table 1). Fig. 2 shows a breakdown of use cases by time and data source. The use of WAM2layers has increased over time and the model has predominantly been used with reanalysis data, but also sometimes with climate model output. Studies that use the output data from WAM2layers for further analysis (Al Hasan et al., 2021; Berger et al., 2014, 2018; Cui et al., 2022; Link et al., 2021; Van der Ent and Savenije, 2011; Weng et al., 2019) are not included in this overview (Table 1, Figure 2).

Evidently, WAM2layers has developed into a mature piece of research software with a considerable user base. We envision that this reference manuscript will help to run it consciously with appropriate model settings, to interpret existing results, and that it will serve as a sound basis for future model improvements and sensitivity studies to understand the effects of certain modelling decisions in detail. The governing equations are derived in Section 2. Section 3 describes the pre-processing procedure, i.e., how to prepare data from various sources for a tracking experiment. Section 4 describes the actual tracking procedure. In Section 5 we briefly discuss two reference cases. Section 6 is dedicated to the best practices that have been incorporated in WAM2layers. Strengths and weaknesses, as well as opportunities for future development are discussed in Section 7.

2 Governing equations

Here, we describe the budget equations for total and tagged moisture as they are implemented in WAM2layers v3. To facilitate comparison with other (moisture tracking/tracer) models, we present a generic description that extends to more than two layers.

Table 1. Topics for which WAM2layers has been used, and associated references.

Topic	References
Moisture Recycling	Bedoya-Soto and Poveda (2024); Enciso et al. (2022); Keys et al. (2014, 2024); Posada-Marín et al. (2023); Shi et al. (2022); Van der Ent et al. (2010); Zemp et al. (2014); Zhang et al. (2020)
Climate Dynamics	Carr and Ummenhofer (2024); Chen et al. (2023); Cluett et al. (2021); Guo et al. (2019); Li et al. (2021, 2022); Link et al. (2020); Lobos-Roco et al. (2022); Mu et al. (2023); Van der Ent and Savenije (2013); Van der Ent and Tuinenburg (2017); Xia et al. (2022); Xiao and Cui (2021); Zhang (2020b); Zhao et al. (2016)
Water Resources Governance	Keys et al. (2017); Keys and Wang-Erlandsson (2018); Keys et al. (2018); Posada-Marín et al. (2024)
Vegetation Dynamics	Ampuero et al. (2020); Duerinck et al. (2016); Keys et al. (2016, 2022); Van der Ent et al. (2012, 2014)
Land-Use Change	De Hertog et al. (2024); Keys et al. (2012); Li et al. (2017); Wang-Erlandsson et al. (2018); Zemp et al. (2017)
Paleoclimate	Bosmans et al. (2020)
Climate Model Evaluation	Franco-Díaz et al. (2024); Guo et al. (2020)
Climate Change	Benedict et al. (2020); Findell et al. (2019); Liu et al. (2022, 2024); Yuan et al. (2023a); Zhang et al. (2019, 2023a, b)
Droughts	Benedict et al. (2021); Mu et al. (2021); Pranindita et al. (2022); Zhang (2020a); Zhou and Shi (2024)
Extreme Precipitation	Liu et al. (2021); Yuan et al. (2023b); Zhang et al. (2023b, 2024)
Method Differences	Li et al. (2024); Van der Ent et al. (2013)




Figure 2. Peer-reviewed scientific papers running WAM2layers (or its predecessor WAM) as of 30 September 2024. The data source only refers to the atmospheric wind and humidity fields as some studies used evaporation or precipitation from other data sources. The category ‘Climate Model’ groups different models. The underlying data is provided as supplement.

The two-layer concept will emerge more clearly in Sections 3 and 4, where we describe in detail how the input data is collapsed from an arbitrary number of model or pressure levels onto a 2-layer grid, and how the equations are implemented numerically.

2.1 Total moisture budget

- 70 As the name "Water Accounting Model" implies, WAM2layers is primarily concerned with bookkeeping the amount of water that is exchanged between grid cells. The total water vapour S per unit area $\mathcal{A}$ contained in a three-dimensional grid cell of volume $\mathcal{V} = \mathcal{A}\Delta z$ is

$$S = \frac{1}{\mathcal{A}} \int_{\mathcal{V}} q\rho \, d\mathcal{V} = q\rho\Delta z = \frac{q\Delta p}{g} \quad (1)$$

- where q is the specific humidity, ρ is the density of the air, g is the gravitational constant, and p is pressure. The change from
 75 height to pressure coordinates is accomplished through substitution of hydrostatic balance, which conveniently eliminates the density. Changes of grid cell area with height are neglected. Notice that WAM2layers uses specific humidity as a proxy for specific water content. As discussed in Section 3, it is possible to account for other phases of water to some extent by using the total column water, if this variable is available in the input data.

- The area-averaged moisture content is convenient to work with, as the corresponding unit of kg m^{-2} is commonly used and
 80 allows for straightforward interpretation of the values of the moisture fields. Older versions of WAM2layers have worked with $(S\mathcal{A})/\rho_{lw}$ instead, where ρ_{lw} represents the density of liquid water, so effectively working in units of m^3 . While the moisture balance is not affected, the input and output of WAM2layers are different between v3 and older versions.

Since the total moisture is a conserved quantity, the change of (area-averaged) moisture S in a grid cell is equal to the fluxes through the boundary of the grid cell plus the sources and sinks,

$$85 \quad \frac{\Delta_t S}{\Delta t} + \frac{1}{\mathcal{A}} (\Delta_x F_x + \Delta_y F_y + \Delta_p F_p) = E - P \quad (2)$$

where the terms on the left hand side represents the change of moisture in a grid cell and the horizontal (green arrows in figure 1B) and vertical (red arrow) fluxes respectively. The terms on the right side represent the rates of removal by precipitation (P , blue arrow) and addition through evaporation (E , purple arrow), both per unit area. Considering advection as the main mode of transport, the horizontal fluxes between grid cells are given by

$$90 \quad F_x = \frac{uq\Delta y\Delta p}{g} \quad , \quad F_y = \frac{vq\Delta x\Delta p}{g} \quad (3)$$

- where u and v are the zonal and meridional wind components, respectively. For our two-layer model, the vertical fluxes F_p at the top and bottom boundaries vanish, leaving only one vertical transport term at the interface between the two layers. Since the vertical flux is positive downwards, this term acts as a sink for the top layer, and a source for the bottom layer. Importantly, this vertical transport term is ill-defined. Advection is typically not the main process for vertical transport. Unresolved processes
 95 such as convection and transport associated with precipitation can also lead to a strong vertical redistribution of water. Another complicating factor is that the vertical extent of the grid cells may vary in time and space. Therefore, in solving for (2), all terms are estimated from the input data (see Sect. 3), except for the vertical transport, which is calculated as a closure term (see Sect. 4).

2.2 Tagged moisture budget

100 The core of WAM2layers consists of tracking a certain proportion of the total moisture, which we refer to as ‘tagged’ moisture. We define the tagged moisture contained in a grid cell as

$$S^* = cS \quad (4)$$

where c represents the tagged moisture concentration. Seeking a budget equation for S^* , we recast (2) in terms of tagged moisture,

$$105 \quad \frac{\Delta_t S^*}{\Delta t} + \frac{1}{\mathcal{A}} (\Delta_x F_x^* + \Delta_y F_y^* + \Delta_p F_p^*) = E^* - P^* \quad (5)$$

such that tagged moisture transport is described by same fluxes as total moisture (illustrated in Figure 1B), except that tagged moisture represents only a portion of the total.

As opposed to (2), which is complete in terms of the input data, (5) is incomplete and must be modelled. Our quest is thus to express the unknown fluxes of tagged moisture, denoted by asterisks, as a function of the known terms of the total moisture budget. For the sources and sinks we write

$$E^* = \delta_e E \quad (6)$$

$$P^* = \delta_p P \quad (7)$$

Here, δ_e and δ_p control where water is added or removed from the system. This is controlled by the tagging region, marked by the red outline and fill in Figure 1A. In case of forward tracking, δ_e is 1 inside the tagging (source) region and 0 elsewhere, and
 115 $\delta_p = c$, i.e., precipitation is removed proportionally to the total moisture in the upper and lower layer. For backward tracking, it is the exact opposite: δ_p is 1 inside the tagging (sink) region, and 0 elsewhere, and $\delta_e = c$, i.e., evaporation is removed proportionally from the lower layer. For the fluxes, WAM2layers uses

$$F_x^* = cF_x \quad F_y^* = cF_y \quad F_p^* = cF_p + \kappa \frac{\Delta_p c}{\Delta p} \quad (8)$$

that is, tagged moisture transport is proportional to the transport of total moisture in the horizontal direction. In the vertical, an
 120 additional dispersion term is added, since Van der Ent et al. (2014) noticed that there was insufficient vertical mixing of tagged moisture without it. Setting $\kappa = k_{vf}|F_p|\Delta p$ ensures that the dispersion rate is proportional to vertical advection rate F_p and the grid size. A similar form may be found in Cushman-Roisin and Beckers (2011, p. 138), who reason that κ is characterized by the largest unresolved eddies. Here, k_{vf} is a tunable parameter to control the dispersion rate. Van der Ent et al. (2014) suggested that a value of 3 gives realistic results, but this could be case-dependent.

125 2.3 Some notes on the formulation of WAM2layers

Though originally conceived as a simple bookkeeping exercise, the accounting method in WAM2layers has a formal basis in the finite volume approach. In the absence of sources and sinks, conservation of moisture reads

$$\frac{\partial(\rho q)}{\partial t} + \nabla \cdot \mathbf{f}(\rho q) = 0 \quad (9)$$

where $\mathbf{f}(x)$ represents a generic flux function (for advection only, $\mathbf{f}(x) = \mathbf{u}x$). Integrating all terms over a fixed control volume and applying Gauss' theorem to the divergence term yields an equivalent expression, stating that the rate of change of moisture in a grid cell is equivalent to the flux across its enclosing surface. Upon discretization, substitution of hydrostatic balance, and division by the grid cell area, we retrieve (2). Applying this procedure to grid cells with a variable vertical extent rather than a fixed control volume yields an additional term by virtue of Leibniz' rule. This term is effectively 'absorbed' by our loose definition of the vertical flux.

On a Cartesian grid with $\mathcal{A} = \Delta x \Delta y$, the horizontal transport in (2) can be written as $\nabla \cdot \mathbf{F}$ with

$$\mathbf{F} = (F_x, F_y) = \left(\frac{uq\Delta p}{g}, \frac{vq\Delta p}{g} \right) \quad (10)$$

This testifies to the equivalence of the continuous and integrated budget equations. However, since WAM2layers operates on a spherical latitude/longitude grid, we must stick with (2) to ensure meridional transport is dependent on latitude.

To facilitate numerical stability analysis and comparison to other moisture tracking methods and, more broadly, to tracer models in general, it is convenient to define an 'integrated moisture velocity' $\hat{\mathbf{u}} = (\hat{u}, \hat{v}, \hat{\omega})$ such that

$$F_x = \hat{u}S \quad F_y = \hat{v}S \quad F_p = \hat{\omega}S \quad (11)$$

With this, conservation of total moisture, tagged moisture, and tagged moisture concentration can all be expressed in the same form:

$$\frac{\partial S}{\partial t} + \nabla \cdot (\hat{\mathbf{u}}S) = 0 \quad (12)$$

$$\frac{\partial S^*}{\partial t} + \nabla \cdot (\hat{\mathbf{u}}S^*) = 0 \quad (13)$$

$$\frac{\partial c}{\partial t} + \nabla \cdot (\hat{\mathbf{u}}c) = 0 \quad (14)$$

In relation to existing literature, we point out that Trenberth and Guillemot (1995); Dominguez et al. (2006); Burde and Zangvil (2001) take a slightly different approach to calculate integrated vapour transport. They take the vertical integral of each term in (9) in isobaric coordinates and apply Leibniz' rule to each of the terms, upon which the vertical flux disappears from the budget equation. In their case, since they consider the budget equation vertically integrated over the full atmosphere, this makes perfect sense. In our case, as mentioned before, we need to retain a vertical transport term between the two layers of our model, due to unresolved and non-isobaric processes, and variability of the grid cells' vertical extent. Note that the aforementioned studies used the symbol w (sometimes referred to as 'precipitable water') for what we refer to as the total moisture S .

2.4 Summary

Recapitulating, the heart of WAM2layers consists of the equations (2) and (5) for total moisture S and tagged moisture S^* , respectively. These governing equations naturally lead to a two-step process for using WAM2layers. First, the input data are collapsed onto a two-layer grid, and the terms in (2) are reconstructed from the available input data. This is what we refer to

as the pre-processing step. Subsequently, (5) is integrated numerically, which we call the tracking step. In the next sections, we
 160 will discuss each of these steps in detail.

3 Pre-processing

The first step in any WAM2layers experiment is to collapse the input data onto a two-layer grid and to reconstruct the corresponding terms in (2) for each new grid cell, except for the vertical flux, which will be calculated as a closure term later on. This procedure is dataset-dependent, and therefore it is isolated from the rest of the code in a dedicated pre-processing module.
 165 This makes it possible to support multiple input datasets with minimal code duplication.

In developing WAM2layers v3, we have primarily focused on working with ERA5 reanalysis data (Hersbach et al., 2020). The comprehensiveness of this dataset enables scrutiny in the treatment of vertical levels, which also facilitates the presentation in this manuscript. We aim for a complete description such that a similar procedure can easily be derived for other data sources.

3.1 Data retrieval: required variables

170 ERA5 data can be downloaded at model levels or pressure levels. In either case, the forthcoming steps require as input the horizontal wind components u and v , as well as specific humidity q , at all or a subset of the available levels. Furthermore, precipitation (P) and evaporation (E), as well as surface pressure (p_s) are required at the surface, and we also include total column water (S_{tc_ERA5}). This last variable is not strictly required to run WAM2layers, but it enables us to estimate and mitigate (to some extent) the errors introduced by the negligence of liquid and ice water. Working with pressure level data additionally
 175 requires the surface values of wind and humidity, i.e., 10-meter wind components u_{10} and v_{10} and 2-meter dewpoint temperature $T_{d,2}$. While ERA5 data is available globally, WAM2layers can be configured to use only a subset of the data. This can be useful to speed up data downloads and simulations, or to reduce disk space requirements. The domain at which WAM2layers operates is referred to as the "tracking domain" (Fig. 1A), which can be provided as a bounding box or as a 2-dimensional raster to allow for arbitrary regions.

180 3.2 States and fluxes

We start by describing the ideal case where the three-dimensional fields of u , v , and q are available on all model levels. WAM2layers aggregates the input data onto a two-layer grid by partitioning the input grid in an upper ($k = 0$) and lower ($k = 1$) part, and taking the sum over each layer.

Summation of (1) over each partition k gives the total water in each grid cell in WAM2layers,

$$185 \quad S_k = \frac{1}{g} \sum_{i \in k} q_i \Delta p_i \quad (15)$$

Here the subscript i refers to indices in the ERA5 level definition, which can be found in the documentation of the underlying Integrated Forecasting System (IFS) (ECMWF, 2016). Horizontal advection terms are calculated according to (3), and

aggregated in a similar fashion

$$F_{x,k} = \frac{1}{g} \sum_{i \in k} u_i q_i \Delta p_i \quad , \quad F_{y,k} = \frac{1}{g} \sum_{i \in k} v_i q_i \Delta p_i \quad (16)$$

190 Following Van der Ent et al. (2013, 2014), we set the boundary to be at the interface between IFS levels 111 and 112 by default. This corresponds to around 812 hPa, for a standard atmospheric pressure of 1013.25 hPa. This boundary in WAM2layers is usually just above the atmospheric boundary layer with on average slightly more moisture below the boundary than above the boundary. Note that the boundary can be at a much lower pressure over mountainous terrain.

The vertical pressure difference over each input grid cell can be calculated with

$$195 \quad \Delta p_i = p_{i+1/2} - p_{i-1/2} \quad (17)$$

$$p_{i+1/2} = A_{i+1/2} + p_{s,i+1/2} B \quad (18)$$

where A and B are hybrid model level coefficients describing the vertical discretization of the model levels in IFS (ECMWF, 2016).

Accounting for liquid and solid water

200 Equation (15) represents the total mass of water *vapour* in each grid cell. By contrasting the calculated column water *vapour* with the *total* column water from ERA5 (S_{T_ERA5}), we can apply an adjustment to account for other phases of water:

$$S_k = S_k \frac{S_{T_ERA5}}{\sum_k S_k} \quad (19)$$

This adds missing cloud and rain/snow water to all levels proportionally to vertical distribution of water vapour. For the two example cases, this correction is typically within 1 % when averaged over the grid, but may be more than 10 % for specific individual grid cells and time steps. WAM2layers provides information about this in the onscreen logging and associated logfile. Note that this step is only possible when the total column water is available from the input data source, which is not always the case.

3.3 Sources and sinks

In the formulation of WAM2layers's equations we adopted the common convention in hydrology of having only positive values for both precipitation and evaporation. Precipitation represents all water transfer from the atmosphere to the surface, including condensation.

In ERA5, vertical fluxes are defined positive downwards. The evaporation variable can be either positive (condensation) or negative (evaporation). Therefore, during the pre-processing step for ERA5, the sign of evaporation is flipped and any remaining negative values are re-assigned to the precipitation variable instead.

215 In a forward-tracking experiment, a proportion of the total surface evaporation is tagged (δ_e). Similarly, for a backward-tracking experiment, we tag a proportion of the precipitation (δ_p). The tagging region is illustrated in Figure 1A. It can be




Figure 3. Illustration of how WAM2layers treats a subset of layers. Full black lines are full-levels, dashed grey lines are half-levels. In B, the top and bottom half-levels are retained, but the remaining half-levels are expired. New half-levels are obtained by interpolating between the selected full-levels.

supplied as a bounding box, or as a 2- or 3-dimensional raster. The three-dimensional option, which is still relatively experimental, would allow for a moving tagging region, e.g. for studying storms. In the two-dimensional case, a tagging start- and end date are given in the configuration file.

220 3.4 Dealing with fewer model levels

ERA5 data is available at all 137 model levels, which is quite unique in terms of resolution (Hersbach et al., 2020), but also quite demanding in terms of storage and processing. Since we collapse the data onto a two-layer grid anyway, it is not necessary to use all 137 levels. To use WAM2layers with only a subset of these 137 layers, the A and B in (18) are interpolated first to the full-levels of the layer subset, then to the interfaces between these full-levels. This procedure is illustrated in Figure 3.

225 Based on our experience with a few test cases, around 20 layers gives good results, provided that the sampling is dense enough near the surface where most of the moisture is located. The number of input levels is configurable though, so users of WAM2layers are encouraged to experiment with different subsets as they see fit.

3.5 Working with pressure-level data

230 Working with (a limited number of) pressure levels instead of model levels is sometimes the only option. This is, for example, the case for datasets from CMIP archives (Juckes et al., 2020, Table 4). ERA5 data is also available on pressure levels, and WAM2layers v3 provides code to work with pressure level data as well.

From a point of view of achieving accurate moisture tracking having wind and humidity data near the surface is crucial, but this could be absent in case of limited pressure levels. Therefore, in order to work with (ERA5) pressure-level data, we complement the data with additional values from the surface-level data (2-meter specific humidity and 10-meter wind) as these




Figure 4. Pressure-level data pre-processing. A: original pressure levels. B: The original levels are extended at the top, surface, and the interface is inserted. C: Wind and humidity are interpolated to the midpoints between the pressure coordinate, to become the new full levels (full black lines). Pressure is kept at the original levels, which thus become the new half-levels (dashed grey lines).

are generally more representative of the surface conditions than the closest available pressure level. In addition, at the top of the atmosphere, WAM2layers assumes humidity to be zero and wind equal to the pressure level with the lowest pressure. This procedure is illustrated in Figure 4.

The surface pressure is used to insert these values in between the right pressure levels. Since surface pressure is space and time dependent ($p_s = p_s(x, y, t)$), the same now applies to our vertical coordinate: $p = p(x, y, z, t)$. A further complication is that the pressure levels can intersect the surface, so we mask any data where the pressure is higher than the surface pressure.

The interface between the two layers in WAM2layers must also follow the terrain. For consistency with the treatment of model-level data, we define the interface according to (18). By default, we use the A and B values for level IFS 111, but since WAM2layers v3.1 this is configurable so users of WAM2layers can easily modify it as they see fit.

At this point, the variables of interest, wind speed and humidity, are collocated with the pressure coordinate. To calculate the grid cell moisture content and fluxes, we need to multiply the pressure difference over a volume with representative values for that same volume. Thus, we interpolate wind and humidity to the interfaces between the pressure coordinates. Finally, we can proceed with evaluating (15) and (16) as before.

3.6 Pre-processing other datasets

WAM2layers is set up in such a way that it is relatively easy to extend its pre-processing to any dataset. Most of the steps outlined above are implemented as generic routines that can be used for any dataset. To add a dataset, only a data loader script has to be implemented, which loads data into a standard internal format. As of version 3.2, built-in pre-processing is available for ERA5 on both model levels and pressure levels, and for CMIP data (on pressure levels). We also support ERA5 pressure level data from ARCO-ERA5 (Carver and Merose, 2023). We envision that more datasets will be included in the future, and we foster community contributions such that other researchers can also contribute their own custom pre-processing scripts.

The pre-processing step is designed in such a way that the intermediate data can be stored efficiently. As part of the tracking routine, the data is first interpolated to a finer time step (~ 10 minutes, see 4.3), after which we solve for the vertical transport term in (2). Finally, (5) is integrated numerically.

4.1 Solving for the vertical flux

260 In discretizing (2), we assign the full evaporation source to the lower layer, but we distribute the precipitation sink proportionally over both layers. Additionally, we introduce an error term to account for inaccuracies, numerical errors, and inconsistencies in the input data caused by both the raw (ERA5) data as well as the pre-processing in WAM2layers. This leads to the following, fully discretized system of equations

$$\begin{aligned}
 & \frac{S_{ijk}^{t+1} - S_{ijk}^t}{\Delta t} + (F_{x,i+1/2jk} - F_{x,i-1/2jk})/\mathcal{A}_j \\
 & \quad + (F_{y,ij+1/2k} - F_{y,ij-1/2k})/\mathcal{A}_j \\
 & \quad + (F_{p,ijk+1/2} - F_{p,ijk-1/2})/\mathcal{A}_j \\
 265 \quad & = \delta_k E_{ij} - \frac{S_{ijk}}{S_{ijT}} P_{ij} + \epsilon_{ijk}^{t+1}
 \end{aligned} \tag{20}$$

where all fluxes are evaluated at $t + 1/2$; the superscripts are omitted for clarity, S_{ijT} represents the calculated column total, whether or not corrected for cloud/rain/snow water as per (19), and δ_k is 1 for the lower layer ($k = 1$) and 0 otherwise. Note that in (20) the vertical flux is positive downward in accordance with the pressure coordinate. For convenience we can move all known terms to the left-hand side and all unknowns to the right. Using R to denote the known terms and leaving out the

270 subscripts i, j for clarity, we get

$$R_k = -(F_{p,k+1/2} - F_{p,k-1/2})/\mathcal{A} + \epsilon_k \tag{21}$$

After applying the boundary conditions $F_p = 0$ at the bottom and top, the only remaining net vertical flux in our two-layer model is at the interface $F_p = F_{p,1/2}$, such that

$$R_0 = -\frac{F_{p,1/2}}{\mathcal{A}} + \epsilon_0 \quad , \quad R_1 = +\frac{F_{p,1/2}}{\mathcal{A}} + \epsilon_1 \tag{22}$$

275 To solve this system, we introduce another constraint by requiring that the errors are proportionally distributed between the upper and lower layer:

$$\frac{\epsilon_k}{\epsilon_T} = \frac{S_k}{S_T} \tag{23}$$

Isolating ϵ_k and recognizing that the total error should be equal to the total residual, we have

$$\epsilon_k = \frac{\epsilon_T S_k}{S_T} = \frac{R_T S_k}{S_T} \tag{24}$$

280 Substituting this result in (22) we finally arrive at

$$\frac{F_{p,1/2}}{\mathcal{A}} = -R_0 + \frac{R_T S_0}{S_T} = R_1 - \frac{R_T S_1}{S_T} \quad (25)$$

In the software implementation, as well as in previous publication, this term $F_{p,1/2}/\mathcal{A}$ is typically simply referred to as ‘the vertical flux’, F_v .

4.2 Numerical integration

285 The formulation of WAM2layers naturally lends itself to a finite-volume implementation, where the incoming fluxes in one cell correspond to outgoing fluxes in its neighbours. Concretely, in case of forward tracking, we have for the upper and lower layers, respectively,

$$\begin{aligned} S_{ij0}^{t+1} = S_{ij0}^t + \Delta t [& \\ & - (F_{x,i+1/2,j0}^* - F_{x,i-1/2,j0}^*)/\mathcal{A} \\ & - (F_{y,i,j+1/2,0}^* - F_{y,i,j-1/2,0}^*)/\mathcal{A} \\ & - (F_{p,ij,1/2}^* + k_{vf}|F_{p,ij,1/2}|(c_{ij,1}^t - c_{ij0}^t))/\mathcal{A} \\ & - c_{ij0} \frac{S_{ij0}}{S_{ijT}} P_{ij}^{t+1/2}] \end{aligned} \quad (26)$$

290

$$\begin{aligned} S_{ij1}^{t+1} = S_{ij1}^t + \Delta t [& \\ & - (F_{x,i+1/2,j1}^* - F_{x,i-1/2,j1}^*)/\mathcal{A} \\ & - (F_{y,i,j+1/2,1}^* - F_{y,i,j-1/2,1}^*)/\mathcal{A} \\ & + (F_{p,ij,1/2}^* + k_{vf}|F_{p,ij,1/2}|(c_{ij,1}^t - c_{ij0}^t))/\mathcal{A} \\ & + \delta_e E_{ij}^{t+1/2} - c_{ij1} \frac{S_{ij1}}{S_{ijT}} P_{ij}^{t+1/2}] \end{aligned} \quad (27)$$

Note that the only remaining vertical transport term, which is shared by both equations, is directed from the upper to the lower layer. δ_e is the tagging region (see Section 3).

295 A crucial aspect of any finite volume scheme is how the fluxes at the interfaces are calculated. WAM2layers employs a simple donor cell scheme (see, e.g. Cushman-Roisin and Beckers, 2011). In this scheme, the flux of tagged moisture $F^* = cF$ is constructed by combining F , calculated at the interfaces of the grid cell and at $t + 1/2$, with c at t , taken from the *upstream* volume. Concretely,

$$(cF)_{x,i+1/2,jk} = \begin{cases} F_{x,i+1/2,jk}^{t+1/2} c_{ijk}^t, & \text{if } F_{x,i+1/2,jk}^{t+1/2} > 0 \\ F_{x,i+1/2,jk}^{t+1/2} c_{i+1,jk}^t, & \text{otherwise} \end{cases} \quad (28)$$

300 and similarly for the other directions.

In case of backward tracking, the equations are similar,

$$\begin{aligned}
S_{ij0}^{t-1} = & S_{ij0}^t - \Delta t [\\
& - (F_{x,i+1/2,j0}^* - F_{x,i-1/2,j0}^*) / \mathcal{A} \\
& - (F_{y,i,j+1/2,0}^* - F_{y,i,j-1/2,0}^*) / \mathcal{A} \\
& - (F_{p,i,j,1/2}^* + k_{vf} |F_p| (c_{ij,1}^t - c_{ij0}^t)) / \mathcal{A} \\
& - \delta_p \frac{S_{ij0}}{S_{ijT}} P_{ij}^{t-1/2}]
\end{aligned} \tag{29}$$

$$\begin{aligned}
305 \quad S_{ij1}^{t-1} = & S_{ij1}^t - \Delta t [\\
& - (F_{x,i+1/2,j1}^* - F_{x,i-1/2,j1}^*) / \mathcal{A} \\
& - (F_{y,i,j+1/2,1}^* - F_{y,i,j-1/2,1}^*) / \mathcal{A} \\
& + (F_{p,i,j,1/2}^* + k_{vf} |F_p| (c_{ij,1}^t - c_{ij0}^t)) / \mathcal{A} \\
& + c_{ij1} E_{ij}^{t-1/2} - \delta_p \frac{S_{ij1}}{S_{ijT}} P_{ij}^{t-1/2}]
\end{aligned} \tag{30}$$

Effectively, all fluxes change direction, which means that the condition in (28) should also be reversed. Also note that the tagged moisture mask is now applied to precipitation instead of evaporation.

4.3 Stability considerations

310 Here we briefly consider the timestep at which WAM2layers should operate. For an explicit solver, a common criterion is that the Courant number should not exceed unity. In other words, water should not be allowed to move more than one grid cell per time step. To explore this limit, we can use the moisture velocity defined in (11).

Figure 5 shows the corresponding maximum time step for a typical range of moisture velocity and the ERA5 grid. It shows that, due to the convergence of the meridians (see Figure 1A), the maximum time step rapidly decreases towards the poles.

315 For this reason, we strongly advise against using WAM2layers with input data on a latitude/longitude grid that extends beyond $\sim 75^\circ$ N/S. Even for tracking experiments with a tagging region located far away from the poles, it is wise to exclude the poles from the calculation domain to avoid spurious results. For other input datasets, a similar assessment should always be made. In section 7 we discuss this limitation in more detail.

In principle, the time step should be chosen sufficiently small so as to satisfy the CFL criterion. However, choosing a very small time step increases computation time and exacerbates numerical diffusion. So there is an incentive to choose a time step that is sufficiently small for, say, $\sim 95\%$ of the time, and accept that there may be some extreme circumstances where the CFL criterion would be violated. In this case, however, we do need to constrain the fluxes for these extremes. To this end, WAM2layers imposes some limits such that the combined moisture fluxes will at most be able to empty the upper or lower grid cell completely. The first limit is such that the meridional and zonal fluxes combined can at most empty the grid cell, but not create any numerical water gains. The second limit is such that the gross vertical flux can also at most empty the grid cell.




Figure 5. Timestep for which the Courant number resolves to unity, based on grid spacing of 0.25° as in the ERA5 input data and a typical range of moisture velocities seen in the example cases.

These limits are not entirely strict when both horizontal and vertical fluxes are large, which can theoretically lead to numerical tagged moisture gains. However, this is being monitored in the logging and should warrant a reconsideration of the time step.

At the end of each time step, WAM2layers checks against moisture surplus or deficit in any grid cell. Notably, tagged moisture cannot exceed the total moisture in a grid cell. Should there be any imbalance, WAM2layers attempts to redistribute the moisture between the upper and lower layer. If this is not sufficient, tagged moisture is lost from the system internally. In v3, we have added log messages to monitor the ongoing experiment, and the spatial fields are included in the output for later analysis. This provides confidence as long as everything is okay, and highlights problems early if they arise. Similarly, WAM2layers logs and report transport over the boundary of the domain, which is naturally expected. However if a research question requires, for instance, an identification of 80 % of the moisture sources, then the boundary transport and where that boundary transport occurred provide information for rerunning the model with a larger domain.

5 Example use cases

Here, we present two example cases to show some of the possibilities of WAM2layers v3. We selected one forward and one backward tracking example with different event duration. The input data as well as the configuration for these cases are available via the 4TU.ResearchData data repository (Benedict and Weijenborg, 2024; Gaasbeek and van der Ent, 2024), and WAM2layers implements a download utility to automatically retrieve these datasets by their DOI. As such, anyone can easily reproduce these example cases and get started with WAM2layers.

The backward tracking example case determines the moisture sources of the extreme precipitation event over the Eiffel region in western Europe (Belgium, France, Germany, Luxemburg and the Netherlands) on 13-14 July 2021. This was a catastrophic event where extreme precipitation over two days resulted in large floods in the tributaries of the Lower Rhine, such as the Ahr,




Figure 6. Outcome of the backward tracking example case ‘Eiffel’. The figure shows the accumulated tracked moisture sources of the extreme precipitation event of 13-14 July 2021. Total precipitation over the sink region during the event is displayed as inset. Backward tracking was performed until 1 July. By then, 42 % of moisture was tracked to its source; 3.4 % of was still in the domain’s atmosphere, and 54 % was associated with transport across the boundaries of the domain. Streamlines depict the average vertically integrated moisture flow during 1-14 July.

Erft and Wupper and the river Meuse, such as the Ourthe, Rur and Geul (Kreienkamp et al., 2021; Mohr et al., 2023). The moisture sources of this event have been previously quantified using different moisture tracking models as shown in Insua-Costa et al. (2022) and Staal and Koren (2023) and now by WAM2layers in Fig. 6. Results show that the evaporative sources of this heavy precipitation event were mostly located over Germany and France. Besides, there is a substantial contribution from the Atlantic ocean and a small contribution from the Mediterranean Sea. The spatial pattern of the sources corresponds to the sources determined by Insua-Costa et al. (2022) and Staal and Koren (2023). Please note that this configuration only considers a limited domain and time span, as it is intended as a lightweight example case. For a full comparison with previous studies, extra simulations are recommended. In this regard, it is worth noting that an intercomparison study of moisture tracking methods, including WAM2layers, is currently being conducted (Benedict et al., 2024).

The forward tracking example case tracks evaporation in the Lake Volta region (Ghana) forward in time for July 1998 to identify the precipitating sinks (Fig. 7). This case was also used in a comparison of moisture tracking models (Van der Ent et al., 2013), although it is not exactly the same. Input data for the previous study was from the regional climate model MM5 (Knoche and Kunstmann, 2013), the domain was smaller and the evaporation was tagged during August as well. Nonetheless, we can observe the similar moisture sink patterns. These patterns are caused by the southwesterly winds transporting moisture in the lower levels, until the African Easterly Jet picks up the tagged moisture and transport it back in westerly direction towards the




Figure 7. Outcome of the forward tracking example case ‘Volta’. The figure shows the tracked moisture sinks of the evaporation from the lake Volta region during July 1998 while the tracking continued until the end of August 1998. The inset shows the time-averaged evaporation in the source region. 86 % of the total tagged evaporated moisture was attributed to surface precipitation and 14 % of the tagged moisture was transported across the boundaries of the domain. Of the tagged evaporation 7.8 % precipitated within the source region. Streamlines depict the average vertically integrated moisture flow during July and August.

360 Atlantic. Note that the average vertically integrated moisture fluxes alone as shown by the streamlines in Fig. 7 cannot capture the complicated moisture transport in this system with strong wind shear and as such highlight the importance of having two layers in WAM2layers.

6 Best practices in research software development

During the development of WAM2layers v3, we have put a lot of emphasis on the incorporation of research software engineering best practices aimed at making WAM2layers more user-friendly, easier to maintain, and to enable a more open and transparent scientific process (Barker et al., 2022; Martinez-Ortiz et al., 2023).

One of the key innovations is that WAM2layers has been turned into an installable command-line program. Users no longer need to edit the source code in order to be able to configure their model. Instead, they simply edit a configuration file. This file contains all settings of the model, including the direction of tracking (forward or backward), the internal time step of the

370 model, the interval on which the output data is provided, and the dates to perform the tracking for. This makes it easier to share and reproduce experiments. To further enhance reproducibility, WAM2layers creates a dedicated output folder for each experiment, which includes a copy of the configuration file and a log file. Each release of WAM2layers has a corresponding DOI, so users can always refer to the exact version of WAM2layers used for their experiments, here version 3.1 (van der Ent et al., 2024b).

375 Importantly, we have added extensive user and developer documentation, which is available at <https://wam2layers.readthedocs.io>. The user documentation includes a quickstart guide, which lets the user run both example cases described in section 5 in a matter of minutes. To this end, we have made WAM2layers available as a Python package on PyPI, such that it can be installed with a single command. Additionally, we have made sample data available at the 4TU.ResearchData repository (Benedict and Weijenborg, 2024; Gaasbeek and van der Ent, 2024), and added functionality to automatically download these datasets. This
380 could easily be extended to include more reference cases, so we encourage users of WAM2layers to make their cases available in a similar manner. To wrap up the quickstart, WAM2layers includes a quick analysis module to quickly inspect the input or output of an experiment, but we recommend users to use more elaborate case-specific code to produce publication-quality figures.

Apart from technical contributions, we have invested in building a welcoming community on GitHub (<https://github.com/WAM2layers/WAM2layers/>). All development happens openly and we encourage users to engage and contribute through
385 opening issues, pull requests, or through the discussion forum where they can ask questions, showcase their results and share codes. The documentation includes community guidelines and instructions for developing the model, as well as an explanation of the collaborative development process. With respect to developer experience, we formulated guidelines for code formatting, and we have set up automated tests to ensure the continuity of the model code. It is verified that results for simple test cases do
390 not change upon making changes to the model code, unless this is expected based on the changes. This strategy is known as ‘regression testing’ (e.g., Pezze, 2008), and it can provide a lot of confidence for (new) developers.

7 Discussion

The Water Accounting Model has since its inception (Van der Ent et al., 2010) evolved from a loose collection of scripts to calculate moisture recycling characteristics into a widely used atmospheric moisture tracking software with a broad range of
395 applications (Fig. 2). This manuscript documents WAM2layers v3 (van der Ent et al., 2024a), which forms a solid basis for future experiments and further development. Here we discuss some key strengths, as well as aspects where we think the model could still be improved.

7.1 Important simplifications

In WAM2layers, the evolution of (tagged) moisture is completely governed by precipitation, evaporation, and redistribution
400 through advection, vertical mixing and, artificially, numerical diffusion. Horizontal mixing, microphysics and other subgrid processes are omitted. These simplifications, partly born of necessity, introduce inaccuracies that users of WAM2layers should

beware of. Below we discuss these simplifications in more detail and how future work may test the importance of these simplification and potentially improve on them.

Historically, WAM2layers has always operated on the same grid as the input data, i.e. reanalysis and climate model output that is typically retrieved at an equirectangular latitude-longitude grid. As shown in section 4, this limits its applicability over polar areas. While alternatives exist, assumptions about the current grid structure are deeply woven into the implementation and changing this is not trivial, especially since we foster continuity of the codebase to keep existing users onboard and avoid alienation. Nonetheless, WAM2layers v3 includes substantial efforts to improve the code structure, isolating the solver and related numerical aspects from other parts of the code that deal with things like I/O and orchestration. This enables experimentation with different solvers, which could eventually enable a different core design. For research questions that focus on moisture transport over polar areas (but not globally), a more feasible workaround could be to remap the input data to another equirectangular grid that is optimized for polar areas. This would keep the existing assumptions intact, though it would still require various code changes that are beyond the scope of this manuscript.

The donor cell scheme employed in WAM2layers (aka ‘upstream’ or ‘Godunov’ scheme) has some favourable properties: it is simple, intuitive, conservative, and numerically stable. Above all, it doesn’t permit overshoot or undershoot, which is a highly suitable property for a moisture tracking model considering the absolute minimum of 0 (tagged) moisture. On the downside, this scheme is known to be numerically diffusive (Hourdin and Armengaud, 1999), and this artificial diffusion may even exceed the true physical diffusion, which is undesirable (Cushman-Roisin and Beckers, 2011). It limits WAM2layers’ usability at high latitudes and may lead to an overestimation of far away sources. Ideally, one would want to suppress or eliminate numerical diffusion altogether, or at least be able to quantify it. One option could be to explore the use of alternative solvers, such as flux-limiters (Hourdin and Armengaud, 1999; Cushman-Roisin and Beckers, 2011) or MPDATA (Smolarkiewicz and Margolin, 1998). WAM2layers v3 isolates the numerical solver such that it becomes easier to experiment with such alternatives. However, we emphasize that any changes in this regard should be carefully evaluated to ensure that their benefits outweigh the existing advantages.

WAM2layers uses water vapour as a proxy for total water, and does not represent microphysical processes associated with phase changes or transport of ice and liquid water. This is a common limitation in offline moisture tracking models (e.g. Cheng and Lu, 2023; Dey and Döös, 2020; Dirmeyer and Brubaker, 1999; Holgate et al., 2020; Keune et al., 2022; Sodemann et al., 2008; Tuinenburg and Staal, 2020). As discussed in Section 3, we can sometimes partly correct the states and horizontal fluxes for the presence of ice and liquid water by application of (19). However, condensation and re-evaporation may be most relevant in the vicinity of precipitation, in which case they could bring about a strong vertical exchange of (tagged) moisture, e.g., condensating in the upper layer and re-evaporating in the lower layer. This is one of the reasons why it is important to realize that the vertical transport term, which is calculated as a closure term in WAM2layers, effectively acts as a generic redistribution term, incorporating the effects of unresolved processes such as microphysics and (turbulent) diffusion, and even compensating for discontinuities in the input data or errors in the horizontal fluxes.

Finally, a note on the reduction to two layers. Van der Ent et al. (2013) found that a two-layer model was both necessary and sufficient to represent large-scale atmospheric moisture transport. The two-layer model simplifies the computation of the

vertical moisture flux and the partitioning of precipitation between the layers of the model. However, it also leads to a two-step procedure where all three-dimensional fields first need to be aggregated onto the new grid. Generalizing to N layers (where N corresponds to the number of levels in the input data) could potentially simplify the code. However, it remains to be seen
440 whether this would actually improve the results, as a correct representation of all vertical exchange processes would remain challenging and should be thoroughly tested.

7.2 A note on performance

WAM2layers v3 is written in Python. This programming language is very popular among (atmospheric) scientists, which makes it easy to use. Moreover, Python’s simple syntax makes it easy to understand the models internal workings, which we deem
445 key from a transparency point-of-view. However, Python has somewhat of a poor reputation when it comes to computational performance. Evidently, the use of numerical computing libraries such as NumPy alleviates this to a certain extent (Harris et al., 2020; Langtangen and Cai, 2008). Further performance gains could potentially be achieved by using dedicated libraries for stencil computations, or by porting some of the performance critical code to more performant languages. A potential downside of using experimental third-party libraries or mixing languages is that it puts a bigger constraint on the maintainability of the
450 software. Treading this path therefore remains a balancing act, and we tend to favour usability and code transparency over performance.

7.3 Future developments

Software development is planned to continue also after the WAM2layers release associated with this manuscript (version 3.1.0 at the time of writing). Future developments notably aim to bring back some old features that have not yet been ported to version
455 3, such calculations for moisture recycling within a single grid cell for all grid cells of the domain at once (e.g., Van der Ent and Savenije, 2011; De Hertog et al., 2024), time tracking (e.g., Van der Ent et al., 2014; Van der Ent and Tuinenburg, 2017), distance tracking (e.g. Guo et al., 2019, 2020) and advanced spline interpolation of the humidity profile in case of limited vertical information (Benedict et al., 2020). We are planning to re-incorporate these features in a backwards-compatible manner. Similarly, we would like to port pre-processing codes for other reanalyses datasets (e.g., Keys et al., 2024; Li et al., 2022) as
460 well as climate model output with which WAM2layers has been used in the past (e.g., Bosmans et al., 2020; De Hertog et al., 2024; Findell et al., 2019; Guo et al., 2020) to future model releases as well.

In developing WAM2layers v3, we have made substantial efforts to generalize the derivation presented here, and to isolate the implementation of the numerical scheme in the code. We believe that this manuscript, together with the changes made for WAM2layers v3, form a good basis for modelling exercises such as implementing alternative solvers or further increasing the
465 performance.

Simultaneously with writing this manuscript, we are co-organizing a coordinated moisture tracking intercomparison study (Benedict et al., 2024), to gain understanding of the uncertainties in different tracking methods. We expect that these community activities will help to shed a better light on the strengths and weaknesses of WAM2layers and other models, and to gain confidence in moisture tracking results in general.

To conclude, this manuscript fills an important gap in the documentation of a widely used atmospheric moisture tracking model. We provided an updated description of the model that facilitates comparison with other models and literature, and pointed out some important assumptions, which facilitates interpretation of past and future results obtained with WAM2layers.

This manuscript, and the associated v3 release, mark a new milestone in the evolution of WAM2layers. While the core
475 tracking principle and assumptions remain unchanged, the code is now much more efficient and modular, making it ready for future developments. We have pointed out several directions for future work, including a more thorough investigation of the numerical aspects and opportunities for improved model validation in collaboration with the moisture tracking community.

With the incorporation of various best practices as outlined in the practical guide to software management plans (Martinez-Ortiz et al., 2023), and the engagement with the wider community, we have made important steps to carry out FAIR and open,
480 collaborative research software development, and we are committed to improving this further into the future. We hope this approach will inspire others to follow suit, as we believe that transparency and collaboration are crucial for the credibility of prosperity of the scientific field.

Code availability. WAM2layers is available on PyPI at <https://pypi.org/project/wam2layers/>. The source code is distributed under a permissive Apache 2.0 license and is available on GitHub at <https://github.com/WAM2layers/WAM2layers>. Documentation lives at <https://wam2layers.readthedocs.io>
485 which includes a description to reproduce the example cases. WAM2layers can be cited via the following DOI: <https://doi.org/10.5281/zenodo.7010594>

Data availability. The data for the example cases is hosted at the 4TU.ResearchData repository, see Benedict and Weijenborg (2024); Gaasbeek and van der Ent (2024). The data on WAM2layers literature underlying Fig. 2 is provided as supplement to this manuscript.

Author contributions. PK lead the development of WAM2layers v3 and writing of the manuscript. IB, CW and RvdE provided input and
490 feedback on the code and the manuscript. IB lead the project under which most of the development of WAM2layers v3 was achieved. RvdE developed the original version of WAM2layers and performed the literature analysis of WAM2layers papers.

Competing interests. The authors declare no competing interests.

Acknowledgements. The authors acknowledge funding from the Netherlands eScience Center (Grant ID: 027.021.S07), the Netherlands Organization for Scientific Research (NWO) under the Open Science Fund (Grant ID: OSF23.1.029) and the Mainstreaming OS Fund from

495 TU Delft. We thank Yang Liu, Bart Schilperoort and Vincent de Feiter for also contributing significantly to the code for WAM2layers v3.
Furthermore, we thank all other contributors to WAM2layers and its user community on GitHub.

References

- Al Hasan, F., Link, A., and van der Ent, R. J.: The Effect of Water Vapor Originating from Land on the 2018 Drought Development in Europe, *Water*, 13, 2856, <https://doi.org/10.3390/w13202856>, 2021.
- 500 Ampuero, A., Strfakis, N. M., Apaéstegui, J., Vuille, M., Novello, V. F., Espinoza, J. C., Cruz, F. W., Vonhof, H., Mayta, V. C., Martins, V. T. S., Cordeiro, R. C., Azevedo, V., and Sifeddine, A.: The Forest Effects on the Isotopic Composition of Rainfall in the Northwestern Amazon Basin, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031445, <https://doi.org/10.1029/2019JD031445>, 2020.
- Barker, M., Chue Hong, N. P., Katz, D. S., Lamprecht, A.-L., Martinez-Ortiz, C., Psomopoulos, F., Harrow, J., Castro, L. J., Gruenpeter, M., Martinez, P. A., et al.: Introducing the FAIR Principles for research software, *Scientific Data*, 9, 622, 2022.
- 505 Bedoya-Soto, J. M. and Poveda, G.: Moisture Recycling in the Colombian Andes, *Water Resources Research*, 60, e2022WR033601, <https://doi.org/10.1029/2022WR033601>, 2024.
- Benedict, I., Van Heerwaarden, C. C., Van Der Ent, R. J., Weerts, A. H., and Hazeleger, W.: Decline in terrestrial moisture sources of the mississippi river basin in a future climate, *Journal of Hydrometeorology*, 21, 299–316, <https://doi.org/10.1175/JHM-D-19-0094.1>, 2020.
- Benedict, I., van Heerwaarden, C., van der Linden, E., Weerts, A., and Hazeleger, W.: Anomalous moisture sources of the Rhine basin during
510 the extremely dry summers of 2003 and 2018, *Weather and Climate Extremes*, 31, <https://doi.org/10.1016/j.wace.2020.100302>, 2021.
- Benedict, I., Weijenborg, C., van der Ent, R., Keune, J., Koren, G., and Kalverla, P.: A moisture tracking intercomparison study-Addressing the uncertainty in modelling the origins of precipitation, in: EMS Annual Meeting, <https://doi.org/10.5194/ems2024-1040>, 2024.
- Benedict, I. B. and Weijenborg, C.: ERA5 data West Europe 2021 July for WAM2layers, <https://doi.org/10.4121/f9572240-f179-4338-9e1b-82c5598529e2.v1>, 2024.
- 515 Berger, M., Van Der Ent, R., Eisner, S., Bach, V., and Finkbeiner, M.: Water Accounting and Vulnerability Evaluation (WAVE): Considering Atmospheric Evaporation Recycling and the Risk of Freshwater Depletion in Water Footprinting, *Environmental Science & Technology*, 48, 4521–4528, <https://doi.org/10.1021/es404994t>, 2014.
- Berger, M., Eisner, S., Van Der Ent, R., Flörke, M., Link, A., Poligkeit, J., Bach, V., and Finkbeiner, M.: Enhancing the Water Accounting and Vulnerability Evaluation Model: WAVE+, *Environmental Science and Technology*, 52, 10757–10766,
520 <https://doi.org/10.1021/acs.est.7b05164>, 2018.
- Bosmans, J., Van der Ent, R., Haarsma, R., Drijfhout, S., and Hilgen, F.: Precession-and obliquity-induced changes in moisture sources for enhanced precipitation over the Mediterranean Sea, *Paleoceanography and Paleoclimatology*, 35, e2019PA003655, <https://doi.org/10.1029/2019PA003655>, 2020.
- Burde, G. and Zangvil, A.: The estimation of regional precipitation recycling. Part I: Review of recycling models, *Journal of climate*, 14,
525 2497–2508, 2001.
- Carr, T. and Ummenhofer, C. C.: Impact of Atmospheric Circulation Variability on U.S. Midwest Moisture Sources, *Journal of Climate*, 37, 59–75, <https://doi.org/10.1175/JCLI-D-23-0178.1>, 2024.
- Carver, R. W. and Merosé, A.: ARCO-ERA5: An Analysis-Ready Cloud-Optimized Reanalysis Dataset, in: 103rd AMS Annual Meeting, AMS, 2023.
- 530 Chen, J., Li, Y., Xiong, B., Wang, Y., Zhou, S., and Huang, Y.: Comparison of moisture sources of summer precipitation in 1998 and 2020 in the middle and lower reaches of Yangtze River basin, *International Journal of Climatology*, 43, 3493–3505, <https://doi.org/10.1002/joc.8040>, 2023.

- Cheng, T. F. and Lu, M.: Global Lagrangian Tracking of Continental Precipitation Recycling, Footprints, and Cascades, *Journal of Climate*, 36, 1923–1941, <https://doi.org/10.1175/JCLI-D-22-0185.1>, 2023.
- 535 Cloux, S., Garaboa-Paz, D., Insua-Costa, D., Miguez-Macho, G., and Pérez-Muñuzuri, V.: Extreme precipitation events in the Mediterranean area: Contrasting two different models for moisture source identification, *Hydrology and Earth System Sciences*, 25, 6465–6477, <https://doi.org/10.5194/hess-25-6465-2021>, 2021.
- Cluett, A. A., Thomas, E. K., Evans, S. M., and Keys, P. W.: Seasonal Variations in Moisture Origin Explain Spatial Contrast in Precipitation Isotope Seasonality on Coastal Western Greenland, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033 543, <https://doi.org/10.1029/2020JD033543>, 2021.
- 540 Crespo-Otero, A., Insua-Costa, D., Hernández-García, E., López, C., and Míguez-Macho, G.: Simple physics-based adjustments reconcile the results of Eulerian and Lagrangian techniques for moisture tracking, *Earth System Dynamics Discussions*, 2024, 1–25, 2024.
- Cui, J., Lian, X., Huntingford, C., Gimeno, L., Wang, T., Ding, J., He, M., Xu, H., Chen, A., Gentile, P., and Piao, S.: Global water availability boosted by vegetation-driven changes in atmospheric moisture transport, *Nature Geoscience*, 15, 982–988, [https://doi.org/10.1038/s41561-](https://doi.org/10.1038/s41561-022-01061-7)
- 545 022-01061-7, 2022.
- Cushman-Roisin, B. and Beckers, J.-M.: *Introduction to geophysical fluid dynamics: physical and numerical aspects*, Academic press, 2 edn., 2011.
- De Hertog, S. J., Lopez Fabara, C. E., van der Ent, R., Keune, J., Miralles, D. G., Portmann, R., Schemm, S., Havermann, F., Guo, S., Luo, F., and others: Effects of idealised land cover and land management changes on the atmospheric water cycle, *Earth System Dynamics*, 15, 265–291, <https://doi.org/10.5194/esd-15-265-2024>, 2024.
- 550 Dey, D. and Döös, K.: Atmospheric Freshwater Transport From the Atlantic to the Pacific Ocean: A Lagrangian Analysis, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2019GL086176>, 2020.
- Dirmeyer, P. A. and Brubaker, K. L.: Contrasting evaporative moisture sources during the drought of 1988 and the flood of 1993, *Journal of Geophysical Research*, 104, 19 383–19 397, <https://doi.org/10.1029/1999JD900222>, 1999.
- 555 Dominguez, F., Kumar, P., Liang, X.-Z., and Ting, M.: Impact of atmospheric moisture storage on precipitation recycling, *Journal of climate*, 19, 1513–1530, 2006.
- Dominguez, F., Hu, H., and Martinez, J.: Two-layer dynamic recycling model (2L-DRM): learning from moisture tracking models of different complexity, *Journal of Hydrometeorology*, 21, 3–16, 2020.
- Duerinck, H., van der Ent, R., van de Giesen, N., Schoups, G., Babovic, V., and Yeh, P.-F.: Observed soil moisture-precipitation feedback in Illinois: A systematic analysis over different scales, *Journal of Hydrometeorology*, 17, 1645–1660, [https://doi.org/10.1175/JHM-D-15-](https://doi.org/10.1175/JHM-D-15-0032.1)
- 560 0032.1, 2016.
- ECMWF: IFS Documentation CY41R2 - Part III: Dynamics and Numerical Procedures, pp. 1–31, 3, ECMWF, <https://doi.org/10.21957/83wouv80>, 2016.
- Enciso, A. M., Baquero, O. L., Escobar-Carbonari, D., Tapasco, J., and Cerón, W. L.: Variability of Precipitation Recycling and Moisture Sources over the Colombian Pacific Region: A Precipitationshed Approach, *Atmosphere*, 13, 1202, <https://doi.org/10.3390/atmos13081202>, 2022.
- 565 Findell, K. L., Keys, P. W., Van Der Ent, R. J., Lintner, B. R., Berg, A., and Krasting, J. P.: Rising temperatures increase importance of oceanic evaporation as a source for continental precipitation, *Journal of Climate*, 32, 7713–7726, <https://doi.org/10.1175/JCLI-D-19-0145.1>, 2019.

- Franco-Díaz, A., Klingaman, N. P., Turner, A. G., Dong, B., and Guo, L.: Effect of global and regional SST biases on the East Asian Summer Monsoon in the MetUM GA7 and GC3 configurations, *Climate Dynamics*, 62, 1535–1553, <https://doi.org/10.1007/s00382-023-06954-w>, 2024.
- Gaasbeek, T. and van der Ent, R. J.: ERA5 data West Africa 1998 July and August for WAM2layers, <https://doi.org/10.4121/bbe10a2a-39dc-4098-a69f-0f677d06ecdd.v2>, 2024.
- Gimeno, L., Vázquez, M., Eiras-Barca, J., Sorí, R., Stojanovic, M., Algarra, I., Nieto, R., Ramos, A. M., Durán-Quesada, A. M., and Dominguez, F.: Recent progress on the sources of continental precipitation as revealed by moisture transport analysis, *Earth-Science Reviews*, 201, 103 070, <https://doi.org/10.1016/j.earscirev.2019.103070>, 2020.
- Goessling, H. F. and Reick, C. H.: On the "well-mixed" assumption and numerical 2-D tracing of atmospheric moisture, *Atmospheric Chemistry and Physics*, 13, 5567–5585, <https://doi.org/10.5194/acp-13-5567-2013>, 2013.
- Guo, L., Van Der Ent, R., Klingaman, N., Demory, M.-E., Vidale, P., Turner, A., Stephan, C., and Chevuturi, A.: Moisture sources for East Asian precipitation: Mean seasonal cycle and interannual variability, *Journal of Hydrometeorology*, 20, 657–672, <https://doi.org/10.1175/JHM-D-18-0188.1>, 2019.
- Guo, L., Van Der Ent, R., Klingaman, N., Demory, M.-E., Vidale, P., Turner, A., Stephan, C., and Chevuturi, A.: Effects of horizontal resolution and air-sea coupling on simulated moisture source for East Asian precipitation in MetUM GA6/GC2, *Geoscientific Model Development*, 13, 6011–6028, <https://doi.org/10.5194/gmd-13-6011-2020>, 2020.
- Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., et al.: Array programming with NumPy, *Nature*, 585, 357–362, 2020.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, 2020.
- Holgate, C. M., Evans, J. P., van Dijk, A. I. J. M., Pitman, A. J., and Di Virgilio, G.: Australian Precipitation Recycling and Evaporative Source Regions, *Journal of Climate*, 33, 8721–8735, <https://doi.org/10.1175/JCLI-D-19-0926.1>, 2020.
- Hourdin, F. and Armengaud, A.: The use of finite-volume methods for atmospheric advection of trace species. Part I: Test of various formulations in a general circulation model, *Monthly weather review*, 127, 822–837, 1999.
- Insua-Costa, D., Senande-Rivera, M., Llasat, M. C., and Miguez-Macho, G.: The central role of forests in the 2021 European floods, *Environmental Research Letters*, 17, 064 053, 2022.
- Juckes, M., Taylor, K. E., Durack, P. J., Lawrence, B., Mizielinski, M. S., Pamment, A., Peterschmitt, J.-Y., Rixen, M., and Sényesi, S.: The CMIP6 data request (DREQ, version 01.00. 31), *Geoscientific Model Development*, 13, 201–224, 2020.
- Keune, J., Schumacher, D. L., and Miralles, D. G.: A unified framework to estimate the origins of atmospheric moisture and heat using Lagrangian models, *Geoscientific Model Development*, 15, 1875–1898, <https://doi.org/10.5194/gmd-15-1875-2022>, 2022.
- Keys, P. and Wang-Erlandsson, L.: On the social dynamics of moisture recycling, *Earth System Dynamics*, 9, 829–847, <https://doi.org/10.5194/esd-9-829-2018>, 2018.
- Keys, P., Barnes, E., Van Der Ent, R., and Gordon, L.: Variability of moisture recycling using a precipitationshed framework, *Hydrology and Earth System Sciences*, 18, 3937–3950, <https://doi.org/10.5194/hess-18-3937-2014>, 2014.
- Keys, P., Warrier, R., Van Der Ent, R., Galvin, K., and Boone, R.: Analysis of Kenya’s Atmospheric Moisture Sources and Sinks, *Earth Interactions*, 26, 139–150, <https://doi.org/10.1175/EI-D-21-0016.1>, 2022.
- Keys, P. W., Van der Ent, R., Gordon, L. J., Hoff, H., Nikoli, R., and Savenije, H.: Analyzing precipitationsheds to understand the vulnerability of rainfall dependent regions, *Biogeosciences*, 9, 733–746, <https://doi.org/10.5194/bg-9-733-2012>, 2012.

- Keys, P. W., Wang-Erlandsson, L., and Gordon, L. J.: Revealing Invisible Water: Moisture Recycling as an Ecosystem Service, *PLOS ONE*, 11, e0151 993, <https://doi.org/10.1371/journal.pone.0151993>, 2016.
- Keys, P. W., Wang-Erlandsson, L., Gordon, L. J., Galaz, V., and Ebbesson, J.: Approaching moisture recycling governance, *Global Environmental Change*, 45, 15–23, <https://doi.org/10.1016/j.gloenvcha.2017.04.007>, 2017.
- Keys, P. W., Wang-Erlandsson, L., and Gordon, L. J.: Megacity precipitation sheds reveal tele-connected water security challenges, *PLOS ONE*, 13, e0194 311, <https://doi.org/10.1371/journal.pone.0194311>, 2018.
- Keys, P. W., Collins, P. M., Chaplin-Kramer, R., and Wang-Erlandsson, L.: Atmospheric water recycling an essential feature of critical natural asset stewardship, *Global Sustainability*, 7, e2, <https://doi.org/10.1017/sus.2023.24>, 2024.
- 615 Knoche, H. R. and Kunstmann, H.: Tracking atmospheric water pathways by direct evaporation tagging: A case study for West Africa, *Journal of Geophysical Research Atmospheres*, 118, 12 345–12 358, <https://doi.org/10.1002/2013JD019976>, 2013.
- Kreienkamp, F., Philip, S. Y., Tradowsky, J. S., Kew, S. F., Lorenz, P., Arrighi, J., Belleflamme, A., Bettmann, T., Caluwaerts, S., Chan, S. C., Ciavarella, A., and De Cruz, Lesley, N.: Rapid attribution of heavy rainfall events leading to the severe flooding in Western Europe during July 2021, 2021.
- 620 Langtangen, H. P. and Cai, X.: On the efficiency of Python for high-performance computing: A case study involving stencil updates for partial differential equations, in: *Modeling, Simulation and Optimization of Complex Processes: Proceedings of the Third International Conference on High Performance Scientific Computing*, March 6–10, 2006, Hanoi, Vietnam, pp. 337–357, Springer, 2008.
- Li, J., Liu, D., Wang, T., Li, Y., Wang, S., Yang, Y., Wang, X., Guo, H., Peng, S., Ding, J., Shen, M., and Wang, L.: Grassland restoration reduces water yield in the headstream region of Yangtze River, *Scientific Reports*, 7, 2162, <https://doi.org/10.1038/s41598-017-02413-9>, 2017.
- 625 Li, Y., Wang, C., Peng, H., Xiao, S., and Yan, D.: Contribution of moisture sources to precipitation changes in the Three Gorges Reservoir Region, *Hydrology and Earth System Sciences*, 25, 4759–4772, <https://doi.org/10.5194/hess-25-4759-2021>, 2021.
- Li, Y., Wang, C., Huang, R., Yan, D., Peng, H., and Xiao, S.: Spatial distribution of oceanic moisture contributions to precipitation over the Tibetan Plateau, *Hydrology and Earth System Sciences*, 26, 6413–6426, <https://doi.org/10.5194/hess-26-6413-2022>, 2022.
- 630 Li, Y., Wang, C., Tang, Q., Yao, S., Sun, B., Peng, H., and Xiao, S.: Unraveling the discrepancies between Eulerian and Lagrangian moisture tracking models in monsoon- and westerly-dominated basins of the Tibetan Plateau, *Atmospheric Chemistry and Physics*, 24, 10 741–10 758, <https://doi.org/10.5194/acp-24-10741-2024>, 2024.
- Link, A., Van Der Ent, R., Berger, M., Eisner, S., and Finkbeiner, M.: The fate of land evaporation - A global dataset, *Earth System Science Data*, 12, 1897–1912, <https://doi.org/10.5194/essd-12-1897-2020>, 2020.
- 635 Link, A., Berger, M., Van Der Ent, R., Eisner, S., and Finkbeiner, M.: Considering the Fate of Evaporated Water Across Basin Boundaries—Implications for Water Footprinting, *Environmental Science & Technology*, 55, 10 231–10 242, <https://doi.org/10.1021/acs.est.0c04526>, 2021.
- Liu, X., Guo, C., Zhang, J., Liu, Y., Xiao, M., Wu, Y., Li, B., and Zhao, T.: Moisture sources of precipitation over the Pearl River Basin in South China, *International Journal of Climatology*, 44, 2160–2173, <https://doi.org/10.1002/joc.8447>, 2024.
- 640 Liu, Y., Zhang, C., Tang, Q., Hosseini-Moghari, S.-M., Haile, G. G., Li, L., Li, W., Yang, K., van der Ent, R. J., and Chen, D.: Moisture source variations for summer rainfall in different intensity classes over Huaihe River Valley, China, *Climate Dynamics*, 57, 1121–1133, <https://doi.org/10.1007/s00382-021-05762-4>, 2021.
- Liu, Y., Garcia, M., Zhang, C., and Tang, Q.: Recent decrease in summer precipitation over the Iberian Peninsula closely links to reduction in local moisture recycling, *Hydrology and Earth System Sciences*, 26, 1925–1936, <https://doi.org/10.5194/hess-26-1925-2022>, 2022.

- 645 Lobos-Roco, F., Hartogensis, O., Suárez, F., Huerta-Viso, A., Benedict, I., De La Fuente, A., and Vilà-Guerau De Arellano, J.: Multi-scale temporal analysis of evaporation on a saline lake in the Atacama Desert, *Hydrology and Earth System Sciences*, 26, 3709–3729, <https://doi.org/10.5194/hess-26-3709-2022>, 2022.
- Martinez-Ortiz, C., Martinez Lavanchy, P., Sesink, L., Olivier, B. G., Meakin, J., de Jong, M., and Cruz, M.: Practical guide to Software Management Plans, <https://doi.org/10.5281/zenodo.7589725>, 2023.
- 650 Miralles, D. G., Nieto, R., McDowell, N. G., Dorigo, W. A., Verhoest, N. E., Liu, Y. Y., Teuling, A. J., Dolman, A. J., Good, S. P., and Gimeno, L.: Contribution of water-limited ecoregions to their own supply of rainfall, *Environmental Research Letters*, 11, 124 007, <https://doi.org/10.1088/1748-9326/11/12/124007>, 2016.
- Mohr, S., Ehret, U., Kunz, M., Ludwig, P., Caldas-Alvarez, A., Daniell, J. E., Ehmele, F., Feldmann, H., Franca, M. J., Gattke, C., Hundhausen, M., Knippertz, P., Küpfer, K., Mühr, B., Pinto, J. G., Quinting, J., Schäfer, A. M., Scheibel, M., Seidel, F., and Wisotzky, C.: A multi-disciplinary analysis of the exceptional flood event of July 2021 in central Europe - Part 1: Event description and analysis, *Natural Hazards and Earth System Sciences*, 23, 525–551, <https://doi.org/10.5194/nhess-23-525-2023>, 2023.
- 655 Mu, Y., Biggs, T. W., and De Sales, F.: Forests Mitigate Drought in an Agricultural Region of the Brazilian Amazon: Atmospheric Moisture Tracking to Identify Critical Source Areas, *Geophysical Research Letters*, 48, e2020GL091 380, <https://doi.org/10.1029/2020GL091380>, 2021.
- 660 Mu, Y., Biggs, T. W., and Jones, C.: Importance in Shifting Circulation Patterns for Dry Season Moisture Sources in the Brazilian Amazon, *Geophysical Research Letters*, 50, e2023GL103 167, <https://doi.org/10.1029/2023GL103167>, 2023.
- Pezze, M.: Software testing and analysis: process, principles, and techniques, John Wiley & Sons, 2008.
- Posada-Marín, J., Salazar, J., Rulli, M. C., Wang-Erlandsson, L., and Jaramillo, F.: Upwind moisture supply increases risk to water security, *Nature Water*, 2, 875–888, <https://doi.org/10.1038/s44221-024-00291-w>, 2024.
- 665 Posada-Marín, J. A., Arias, P. A., Jaramillo, F., and Salazar, J. F.: Global Impacts of El Niño on Terrestrial Moisture Recycling, *Geophysical Research Letters*, 50, e2023GL103 147, <https://doi.org/10.1029/2023GL103147>, 2023.
- Pranindita, A., Wang-Erlandsson, L., Fetzer, I., and Teuling, A.: Moisture recycling and the potential role of forests as moisture source during European heatwaves, *Climate Dynamics*, 58, 609–624, <https://doi.org/10.1007/s00382-021-05921-7>, 2022.
- Shi, K., Li, T., Zhao, J., Su, Y., Gao, J., and Li, J.: Atmospheric recycling of agricultural evapotranspiration in the Tarim Basin, *Frontiers in Earth Science*, 10, 950 299, <https://doi.org/10.3389/feart.2022.950299>, 2022.
- 670 Smolarkiewicz, P. K. and Margolin, L. G.: MPDATA: A finite-difference solver for geophysical flows, *Journal of Computational Physics*, 140, 459–480, 1998.
- Sodemann, H., Schwierz, C., and Wernli, H.: Interannual variability of Greenland winter precipitation sources: Lagrangian moisture diagnostic and North Atlantic Oscillation influence, *Journal of Geophysical Research D: Atmospheres*, 113, D03 107, <https://doi.org/10.1029/2007jd008503>, 2008.
- 675 Staal, A. and Koren, G.: Comment on ‘The central role of forests in the 2021 European floods’, *Environmental Research Letters*, 18, 048 002, 2023.
- Te Wierik, S. A., Cammeraat, E. L. H., Gupta, J., and Artzy-Randrup, Y. A.: Reviewing the Impact of Land Use and Land-Use Change on Moisture Recycling and Precipitation Patterns, *Water Resources Research*, 57, e2020WR029 234, <https://doi.org/10.1029/2020WR029234>, 2021.
- 680 Theeuwens, J. J., Staal, A., Tuinenburg, O. A., Hamelers, B. V., and Dekker, S. C.: Local moisture recycling across the globe, *Hydrology and Earth System Sciences*, 27, 1457–1476, 2023.

- Trenberth, K. E. and Guillemot, C. J.: Evaluation of the global atmospheric moisture budget as seen from analyses, *Journal of Climate*, 8, 2255–2272, 1995.
- 685 Tuinenburg, O. A. and Staal, A.: Tracking the global flows of atmospheric moisture and associated uncertainties, *Hydrology and Earth System Sciences*, 24, 2419–2435, 2020.
- Van der Ent, R. and Savenije, H.: Length and time scales of atmospheric moisture recycling, *Atmospheric Chemistry and Physics*, 11, 1853–1863, <https://doi.org/10.5194/acp-11-1853-2011>, 2011.
- Van der Ent, R. and Tuinenburg, O.: The residence time of water in the atmosphere revisited, *Hydrology and Earth System Sciences*, 21, 779–790, <https://doi.org/10.5194/hess-21-779-2017>, 2017.
- 690 Van der Ent, R., Tuinenburg, O., Knoche, H.-R., Kunstmann, H., and Savenije, H.: Should we use a simple or complex model for moisture recycling and atmospheric moisture tracking?, *Hydrology and Earth System Sciences*, 17, 4869–4884, <https://doi.org/10.5194/hess-17-4869-2013>, 2013.
- Van der Ent, R., Wang-Erlandsson, L., Keys, P. W., and Savenije, H.: Contrasting roles of interception and transpiration in the hydrological cycle–Part 2: Moisture recycling, *Earth System Dynamics*, 5, 471–489, <https://doi.org/10.5194/esd-5-471-2014>, 2014.
- 695 Van der Ent, R. J. and Savenije, H. H.: Oceanic sources of continental precipitation and the correlation with sea surface temperature, *Water Resources Research*, 49, 3993–4004, <https://doi.org/10.1002/wrcr.20296>, 2013.
- Van der Ent, R. J., Savenije, H. H., Schaefli, B., and Steele-Dunne, S. C.: Origin and fate of atmospheric moisture over continents, *Water Resources Research*, 46, W09 525, <https://doi.org/10.1029/2010WR009127>, 2010.
- 700 Van der Ent, R. J., Coenders-Gerrits, A. M. J., Nikoli, R., and Savenije, H. H.: The importance of proper hydrology in the forest cover-water yield debate: commentary on Ellison et al.(2012) *Global Change Biology*, 18, 806–820., *Global Change Biology*, 18, <https://doi.org/10.1111/j.1365-2486.2012.02703.x>, 2012.
- van der Ent, R. J., Benedict, I. B., Weijenberg, C., Schilperoort, B., Liu, Y., Barnes, E., Cömert, T., Niek, v. d. K., Guo, L., de Feiter, V., and Kalverla, P.: WAM2layers, Zenodo, <https://doi.org/10.5281/zenodo.7010594>, 2024a.
- 705 van der Ent, R. J., Benedict, I. B., Weijenberg, C., Schilperoort, B., Liu, Y., Barnes, E., Cömert, T., van de Koppel, N., Guo, L., de Feiter, V., and Kalverla, P.: WAM2layers Version v3.1.0, Zenodo, <https://doi.org/10.5281/zenodo.12206708>, 2024b.
- Wang-Erlandsson, L., Fetzer, I., Keys, P. W., Van Der Ent, R. J., Savenije, H. H., and Gordon, L. J.: Remote land use impacts on river flows through atmospheric teleconnections, *Hydrology and Earth System Sciences*, 22, 4311–4328, <https://doi.org/10.5194/hess-22-4311-2018>, 2018.
- 710 Weng, W., Costa, L., Lüdeke, M. K., and Zemp, D. C.: Aerial river management by smart cross-border reforestation, *Land Use Policy*, 84, 105–113, <https://doi.org/10.1016/j.landusepol.2019.03.010>, 2019.
- Xia, Z., Welker, J. M., and Winnick, M. J.: The Seasonality of Deuterium Excess in Non-Polar Precipitation, *Global Biogeochemical Cycles*, 36, e2021GB007 245, <https://doi.org/10.1029/2021GB007245>, 2022.
- Xiao, M. and Cui, Y.: Source of Evaporation for the Seasonal Precipitation in the Pearl River Delta, China, *Water Resources Research*, 57, e2020WR028 564, <https://doi.org/10.1029/2020WR028564>, 2021.
- 715 Yuan, X., Yang, K., Lu, H., Sun, J., Wang, Y., Liu, Y., and Tang, Q.: Weakened Subtropical Westerlies Reduced Early Spring Precipitation in the Southeast Tibetan Plateau, *Journal of Climate*, 36, 4363–4373, <https://doi.org/10.1175/JCLI-D-22-0770.1>, 2023a.
- Yuan, X., Yang, K., Sun, J., Wang, Y., He, Y.-Y., Zou, M.-J., and Jiang, Y.-Z.: Why was Pakistan extreme precipitation stronger in 2022 than in 2010?, *Advances in Climate Change Research*, 14, 913–920, <https://doi.org/10.1016/j.accre.2023.11.016>, 2023b.

- 720 Zemp, D., Schleussner, C.-F., Barbosa, H., Van der Ent, R., Donges, J. F., Heinke, J., Sampaio, G., and Rammig, A.: On the importance of cascading moisture recycling in South America, *Atmospheric Chemistry and Physics*, 14, 13 337–13 359, <https://doi.org/10.5194/acp-14-13337-2014>, 2014.
- Zemp, D. C., Schleussner, C.-F., Barbosa, H. M. J., Hirota, M., Montade, V., Sampaio, G., Staal, A., Wang-Erlandsson, L., and Rammig, A.: Self-amplified Amazon forest loss due to vegetation-atmosphere feedbacks, *Nature Communications*, 8, 14 681, <https://doi.org/10.1038/ncomms14681>, 2017.
- 725 Zhang, C.: Moisture source assessment and the varying characteristics for the Tibetan Plateau precipitation using TRMM, *Environmental Research Letters*, 15, 104 003, <https://doi.org/10.1088/1748-9326/abac78>, 2020a.
- Zhang, C.: Moisture sources for precipitation in Southwest China in summer and the changes during the extreme droughts of 2006 and 2011, *Journal of Hydrology*, 591, <https://doi.org/10.1016/j.jhydrol.2020.125333>, 2020b.
- 730 Zhang, C., Tang, Q., Chen, D., Van der Ent, R. J., Liu, X., Li, W., and Haile, G. G.: Moisture source changes contributed to different precipitation changes over the northern and southern Tibetan Plateau, *Journal of Hydrometeorology*, 20, 217–229, <https://doi.org/10.1175/JHM-D-18-0094.1>, 2019.
- Zhang, C., Chen, D., Tang, Q., and Huang, J.: Fate and Changes in Moisture Evaporated From the Tibetan Plateau (2000–2020), *Water Resources Research*, 59, e2022WR034 165, <https://doi.org/10.1029/2022WR034165>, 2023a.
- 735 Zhang, C., Tang, Q., Zhao, Y., Chen, D., Huang, J., Liu, Y., and Zhang, X.: Moisture source differences between the 2020 and 1998 super Meiyu-flood events in the Yangtze River Valley, *Weather and Climate Extremes*, p. 100644, <https://doi.org/10.1016/j.wace.2024.100644>, 2024.
- Zhang, X., Liu, Z., Liu, Y., Jiang, L., Wang, R., Jiang, H., Li, J., Tang, Q., and Yao, Z.: Examining moisture contribution for precipitation in response to climate change and anthropogenic factors in Hengduan Mountain Region, China, *Journal of Hydrology*, 620, 129 562, <https://doi.org/10.1016/j.jhydrol.2023.129562>, 2023b.
- 740 Zhang, Y., Huang, W., Zhang, M., Tian, Y., Wang, G., and Zhong, D.: Atmospheric Basins: Identification of Quasi-Independent Spatial Patterns in the Global Atmospheric Hydrological Cycle Via a Complex Network Approach, *Journal of Geophysical Research: Atmospheres*, 125, <https://doi.org/10.1029/2020JD032796>, 2020.
- Zhao, T., Zhao, J., Hu, H., and Ni, G.: Source of atmospheric moisture and precipitation over China’s major river basins, *Frontiers of Earth Science*, 10, 159–170, <https://doi.org/10.1007/s11707-015-0497-4>, 2016.
- 745 Zhou, K. and Shi, X.: Understanding Precipitation Moisture Sources and Their Dominant Factors During Droughts in the Vietnamese Mekong Delta, *Water Resources Research*, 60, e2023WR035 920, <https://doi.org/10.1029/2023WR035920>, 2024.