

Accretionary prism deformation and fluid migration caused by slow earthquakes in the Nankai subduction zone

Takashi Tonegawa¹, Takeshi Akuhara², Yusuke Yamashita³, Hiroko Sugioka⁴, Masanao Shinohara², Shunsuke Takemura², Takeshi Tsuji⁵

5 ¹Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokohama, 236-0001 Japan

²Earthquake Research Institute, The University of Tokyo, Tokyo, 113-0032, Japan

³Disaster Prevention Research Institute, Kyoto University, Miyazaki, 889-2161, Japan

⁴Department of Planetology, Graduate School of Science, Kobe University, Kobe, 657-8501, Japan

⁵Department of Systems Innovation, Faculty of Engineering, The University of Tokyo, Tokyo, 113-8656, Japan

10 *Correspondence to:* Takashi Tonegawa (tonegawa@jamstec.go.jp)

Abstract. Slow earthquakes induce structural deformation around their source regions and can also be linked to fluid migrations. Both of these phenomena potentially induce temporal variations in the seismic structure; how these two factors behave for a slow earthquake episode remains unknown. In this study, we applied the ambient noise correlation technique to the continuous records acquired at the seafloor in the Nankai subduction zone, and investigate the changes in seismic velocity
15 (dv/v) and heterogeneous (correlation coefficient, CC) structures before and after the slow earthquake activity that occurred around the shallow plate interface from the end of 2020. As a result, temporal variations in dv/v and CC show different patterns for the slow earthquake episode. The dv/v variations show a step-like temporal reduction and the reduced velocity was not recovered to the original level until the end of the observation period, whereas the CC variations show transient reductions and were recovered to the original level after the episode. Thus, the dv/v reflects the variation in the aspect ratio of pre-existing
20 cracks and/or newly created cracks due to sediment deformation, where the extensional stresses normal to the trough were induced by the slips of the slow earthquakes, and the updated crack condition persisted even after the episode. We suggest that the CC variations correspond to transient fluid migrations from the source regions to shallow depths, activated by the fracturing of fluid caprocks that resulted from slow earthquakes. Our study indicates that monitoring these two quantities provides useful information to understand the variations in the subsurface structure due to slow earthquakes.

25

Short summary. This study demonstrates that for slow earthquakes in the shallow Nankai subduction zone, (1) the seismic velocity reductions in the accretionary prism reflect the stress field changes due to the slow earthquakes, and (2) the temporal variations in the heterogeneous structure correspond to upward fluid migrations from the source region of the slow earthquakes.

Slow earthquakes occur at shallow plate interfaces in the Nankai subduction zone in southwestern Japan, and their characteristics have been investigated from interdisciplinary viewpoints, such as seismic and geodetic observations, geology, experiments, and seismic structures (e.g., Takemura et al. 2023). The slow earthquakes in this region are mainly categorized into tremors, very low frequency earthquakes (VLFs), and slow slip events (SSEs), based on the dominant periods of their seismic signals (2–8 Hz for tremors and 0.01–0.1 Hz for VLFs) and durations of geodetic deformation (several days to several years for SSEs) (e.g., Fig. 1 in Obara and Kato, 2016). The occurrences of tremors and VLFs and their distributions have been determined by seismometers in both land and sea areas (e.g., Obara and Ito, 2005; Ito and Obara, 2006; Asano et al., 2008; Sugioka et al., 2012; Kaneko et al., 2018; Takemura et al., 2019; Yabe et al., 2019; Baba et al., 2020; Ogiso et al., 2022; Tamaribuchi et al., 2022; Yamamoto et al., 2022). SSEs have been observed by pore pressure measurements in the boreholes at the seafloor (Araki et al., 2017; Nakano et al., 2018; Ariyoshi et al., 2021) and seafloor geodetic observations using Global Navigation Satellite System (GNSS)–acoustic ranging combined with a seafloor positioning system (GNSS-A) (Yokota and Ishikawa, 2020). Compared to SSEs, the spatio-temporal variations of tremors and VLFs in this region have been elucidated using seismometer records. Tremors and VLFs are sporadically distributed along the strike direction of the Nankai Trough, and they occur at the plate interface at depths less than ~10 km (Obara and Kato, 2016). The recurrence intervals of these phenomena vary regionally along the Nankai Trough between 1 and 5 years (e.g., Baba et al., 2020).


Fluids are related to the generations of slow earthquakes. This linkage has been documented for slow earthquakes occurring at the both deep and shallow plate interfaces in various subduction zones. Seismological techniques, including tomography and receiver functions, have revealed that the subducting oceanic crust near the region where deep tremors occur shows specific seismic velocities, such as low P wave velocity (V_p) in the Nankai subduction zone (Shelly et al., 2007) and high V_p/V_s in the Cascadia subduction zone (Gosselin et al., 2020). A tomographic study suggested that low-frequency earthquakes occur under undrained conditions within the subducting oceanic crust (Nakajima and Hasegawa, 2016). Details on other subduction zones have been reviewed elsewhere (Audet and Kim, 2016). Pore pressure waves along the plate interface may influence the migration of tremor excitations (Cruz-Atienza et al., 2018). Such fluid migrations to different sites are also related to slow earthquake generations (e.g., Nakajima and Uchida, 2018; Ito and Nakajima, 2024). These seismological features indicate the prevalence of fluids with high pore pressure within the oceanic crust, which mechanically reduces the shear strength of the fault.

In the cases of shallow slow earthquakes, evidence of fluids and their migrations has been reported. Seismic exploration surveys have reported the presence of a low velocity layer within the accretionary prism in the central Nankai subduction zone (Park et al., 2010; Kamei et al., 2012), and this is linked to the generation of VLFs through the presence of fluids (Kitajima and Saffer, 2012; Tsuji et al., 2014; Tonegawa et al., 2017; Akuhara et al., 2020). Another exploration study in the Hikurangi subduction zone indicated that sediment lenses with low velocity left behind subducting seamounts are linked to SSEs (Bangs et al. 2023). In the Hyuga-nada region of the western Nankai subduction zone, low seismic velocity layers have been detected

within marine sediments above the regions where tremors occur (Akuhara et al., 2023a). Arai et al. (2023) has successfully imaged the vertical low velocity zones connecting between the plate boundary and seafloor, and found a spatial correlation between the vertical conduits and the distribution of slow earthquakes.

In order to unveil the relationship between fluids and slow earthquake generations, the temporal changes in the seismological structure around the region of slow earthquakes should be investigated. Using the continuous records of the Dense Oceanfloor Network system for Earthquakes and Tsunamis (DONET, Fig. 1) (Kaneda et al., 2015; Kawaguchi et al., 2015; Aoi et al., 2020) for the observation periods >10 years, which was acquired in the Nankai subduction zone. Tonegawa et al. (2022) found that the heterogeneous structure shows temporal changes associated with slow earthquakes, and interpreted that transient fluid migrations alter the fractional fluctuation of wave velocity in the heterogeneous structure within the sediments. In particular, the slow earthquake activity that started at the end of 2020 showed the temporal changes in seismic velocity and heterogeneous structure. However, the time after the slow earthquake activity has not been investigated in details because post-event data were not sufficiently available. Moreover, the mapping of temporal variations in the subsurface structure was limited to the DONET1 area (Tonegawa et al., 2022).

In this study, we include the temporary OBSs into the DONET stations to cover the gap between DONET1 and DONET2, and estimate temporal changes in the subsurface structure using ambient noise correlation techniques. Additionally, we investigate the connection between fluid and the occurrence of slow earthquakes, especially focusing on the activity that started from the end of 2020. Because the spatio-temporal variation of tremors in the activity has been investigated in detail and potentially reflects the slip area, we compare the obtained structural changes with the tremor activity. The tremor activity started on December 5, 2020, peaked on December 11 and 28 (Tamaribuchi et al., 2022; Ogiso and Tamaribuchi, 2022), and faded by early February 2021, which expanded from the eastern part of DONET2 to the eastern part of DONET1 (Fig. 1).



85 **Figure 1: (a) Frequency of tremors that occurred between December 1, 2020, and February 5, 2021, with the cumulative numbers**
by the solid line and counts per day by the histogram. The tremors are the same as those used in the panel (b). (b) Map showing the
locations of stations. Squares and circles represent DONET1 and DONET2 stations, respectively, and triangles show temporary
OBSs. The two lines correspond to the Nankai Trough and the contour lines of 10 km for the Philippine Sea Plate (Baba et al. 2002;
Hirose et al. 2008). Crosses indicate the locations of tremors that occurred between December 1st 2020 and February 5th, 2021
(Tamaribuchi et al. 2022), for which tremors with spatial errors $\leq 0.1^\circ$ are plotted and 16% of tremors are preserved from the original
catalog.


2 Data

We used the continuous vertical component records from 49 DONET stations (Fig. 1, and the stations codes used in this study
 95 are displayed in Fig. S1). However, those near the Nankai Trough were primarily considered because temporal changes due to
 the slow earthquakes were limited to this region (Tonegawa et al., 2022). DONET contains the eastern (DONET1) and western
 (DONET2) cabled networks, which are deployed at water depths of 1,000–4,400 m with a station spacing of 10–30 km and
 are installed at the end of 2010 and 2014, respectively. All stations include a broadband seismometer (Guralp CMG-3T, flat
 velocity response up to 360 s), which is buried 1 m below the seafloor (Nakano et al., 2013). The locations of the DONET1
 100 and DONET2 differ from each other by ~ 80 km. To fill the gap, 15 OBSs with short-period sensors (1Hz) were deployed
 between September 2019 and May 2021, among which 10 OBSs were used for two subarrays with a station spacing of ~ 2 km.
 In this study, we used 2 OBSs located at the center of the two subarrays. In total, 7 OBSs were used.

3 Methods

3.1 Calculation of cross correlation functions (CCFs)

105 We calculated cross-correlation functions (CCFs) from ambient noise records following established methods (e.g., Campillo and Paul, 2003; Shapiro et al., 2005; Brenguier et al., 2007). To suppress energetic signals, such as those of earthquakes, we applied lognormal-shaped functions (Tonegawa et al., 2020; 2022). A bandpass filter of 0.5–2.0 Hz combined with spectral whitening was used. In the chosen frequency band, the ambient seismic noise is dominated by acoustic-coupled Rayleigh (ACR) waves. These waves propagate through the ocean and the entire accretionary prism in this region with a propagation
110 velocity of 1.3–1.5 km/s (Tonegawa et al., 2015). CCFs were computed using 600-s time windows, and a 30-day moving average was obtained by stacking daily CCFs for each station pair. If the duration affected by the lognormal-shaped function was less than 70 % of 30-day period, the corresponding CCF was discarded. Additionally, we prepared 20-day CCFs, preserving the duration longer than 70% of 20 days, for estimating temporal variations in the subsurface structures. This is because 30-day CCFs are required for evaluating clock deviations that is introduced in the subsequent paragraph, but 20-day
115 CCFs can be used for assessing the subsurface structures. Reference CCFs for each station pair were generated by stacking CCFs over the entire observation period. In the CCFs, direct ACR wave propagation between station pairs appears at early lag times, while their scattered waves are observed in the coda (Fig. 2).



120 **Figure 2: Stacked CCFs at 0.5–2.0 Hz. (a) CCFs for station pairs stacked between –182 (October 2, 2019) and 183 (October 1, 2020) days from the reference date of April 1, 2020 (1 year). The two lines represent a propagation velocity of 1.5 km/s. (b) CCFs at the station pair of SHM6–KMD13 for the observation period (October 2, 2019–May 23, 2021).**

3.2 Clock error estimations

125 The clocks of the recorders are adjusted to the clocks of the GNSS at the timings before and after their observations, and are linearly interpolated during the observations. However, the deviation rate of recorder clocks from the accurate times may not be uniform through the observation period. These deviations can be measured using CCFs with ambient noise correlations (Takeo et al., 2014). Using the CCFs calculated in the previous section, we measure the clock deviations of the OBSs from the clocks of the DONET stations. The clocks of the DONET data are calibrated by the GNSS signal through the submarine cables to land (e.g., Tsuji et al., 2023). We used CCFs for the station pairs for which the DONET stations are located within a distance of 50 km from each OBS. When the CCFs are aligned as a function of the observation date, several coherent phases, including ACR waves and their scattered waves, are observed in the positive and negative lag times over the observation periods (Fig. 2b). However, if the clocks are deviated from the accurate times, the arrival times of the causal and acausal waves are simultaneously shifted in the same direction towards the positive or negative lag time during the observation periods. Here, in cases where temporal variations in the seismic velocity at the subsurface structure affect the arrival times of causal and acausal waves, the arrival times of causal waves in the positive lag time and acausal waves in the negative lag time are shifted symmetrically with respect to the lag time of 0 s (e.g., Brenguier et al. 2008), which is a different feature from the clock deviations to the CCFs.

We set a 2-s time window with an increment of 1 s between –20 and 20 s in the CCFs (Fig. 2b), and prepare the reference CCF for each station pair by stacking the CCFs between October 2019 and September 2020. In each 2-s time window, we calculate cross-correlation functions (referred to as CF, to distinguish this function from the CCFs using ambient noise records) between the reference CCF and individual CCFs. If, within a given time window, the cross-correlation coefficients of the CFs exceed 0.9 for more than 85% of the observation period, the time window is considered to contain coherent signals throughout the period. The delay times of these CFs are then interpreted as the clock deviation. The obtained temporal variation of the clock deviation from the starting time of the observation period may be slightly shifted from 0 s in lag time, because the coherent signals between the reference CCF and individual CCFs may have phase differences. Therefore, the time difference of a coherent signal averaged over the first 10 days is subtracted from the time differences of the subsequent days. This processing is repeated for different pairs between each OBS and the available DONET stations that satisfy the aforementioned criterion. On each day, the obtained delay times are counted at 0.01 s bins, and the counts are used for estimating the weighted average, S_j , for j -th day.

$$S_j = \frac{\sum_i w_{ij} T_{ij}}{\sum_i w_{ij}}, \quad (1)$$

$$w_{ij} = (C_{ij})^3, \quad (2)$$

where T_{ij} is the i -th time, which is assigned from -0.15 s to 0.15 s with an increment of 0.01 s, and C_{ij} is the counts at the bins. We chose the power of 3 in eq. (2) because, if the power is 1–2, the estimated average is slightly away from the large counts and is deviated to the center of the count distribution. Additionally, we took the moving average of S_j within 60 days for smoothing the obtained values (Fig. 3 and S2).




Figure 3: Clock deviations at SHM7c. (a) The locations of the pairs between SHM7c and DONET stations used for measuring the clock deviations, which are connected by solid lines. (b) The estimated clock deviations at SHM7c. The solid and dashed lines represent the weighted average and its standard deviations for the clock deviation. The background counts indicate the measurements of the clock deviations counted at bins of 0.01 s and 1 day.


3.3 Estimations of dv/v and CC

For each 30-day CCF, we applied the stretching technique for estimating the changes in seismic velocity ($dv/v = -dt/t$) and cross-correlation coefficients (CC) (Sens-Schönfelder and Wegler, 2006; Obermann et al., 2013), in which the relation of $dv/v = -dt/t$ has been widely used since an earlier work for coda waves from repetitive seismic sources (Poupinet et al. 1984). The searching range of the stretch (dt/t) is between -0.4 and 0.4 % with an increment of 0.01 %. CCs are estimated by cross-correlating 50-s segments between the reference and 30-day CCFs that are stretched by the obtained dv/v , and the CCs obtained for the positive and negative lag times are averaged. The starting times of the 50-s segments are the arrival times of the ACR waves in the positive and negative lag times, which were estimated by dividing the separation distance of two stations by a

propagation velocity of 1.5 km/s. After correcting the individual CCFs using the obtained dv/v , we calculated the CCs between the reference and the corrected CCFs.

The uncertainties in dv/v and CC due to the clock deviations are estimated by a bootstrapping technique. The dv/v and CC variations were estimated using the 30-day CCFs every day (between -14^{th} and $+15^{\text{th}}$ days from the reference days), and the clock deviations were also estimated for the OBSs every day. We randomly selected a value between the standard deviations, and time-shifted the CCFs using this value as the clock deviation. Using the time-shifted CCFs, we estimated dv/v and CC, and repeated this process 100 times to obtain dv/v and CC uncertainties. The results for the node KMC and OBSs are summarized in Figs. 4 and S3. Moreover, we compared temporal variations in dv/v and CC for several pairs of KMC stations with changing stacking periods for 10, 20, and 30 days (Fig. S4). dv/v for all stacking periods is consistent for three station pairs. The patterns of CC for stacking periods between 20 and 30 days are similar though there are small offsets. On the other hand, CC for a stacking period of 10 days disagrees with those with longer stacking periods. Considering the similarity of the CC, 20-day CCFs can be leveraged for detecting subsurface structure changes. However, 30-day CCFs are reliable for assessing clock deviations because of the small offsets of CC. Moreover, dv/v and CC could not be estimated at several pairs including MRE21 during the slow earthquake activity (Fig. S3d). This is because we discarded the 30-day CCFs for the reasons mentioned in Section 3.1, and we did not use the corresponding 20-day CCFs for dv/v and CC mappings, which will be explained in the subsequent section.

Because we use ACR waves, the depth sensitivities of the dv/v and CC obtained in this study are approximately 1 km from the seafloor (Tonegawa et al., 2022). Moreover, since the stretching technique measures the dt/t for scattered waves in the coda part of the CCFs, the obtained dv/v reflects relatively wide horizontal areas around two stations. In contrast, because the CC potentially varies with amplitude variations within a short time segment of the CCFs, the obtained CC reflects the temporal variations of the heterogeneous structure in relatively horizontally localized areas where the scattered waves within the short time segment sample. Indeed, an experimental study suggested that the CC variation was more sensitive than dv/v variations to fluid injection (Théry et al., 2020).



195 **Figure 4: (a) dv/v variations for node KMC. The shaded region represents the timing of the slow earthquake activity. The bold vertical line indicates the reference period, and the dotted lines show the dates for which the mappings of dv/v and Δg for non-slow-earthquakes are conducted (Figs. S5 and S6). (b) Same as panel (a), but for results for the temporary OBSs. Upper and lower lines indicate the uncertainties due to clock deviations. (c) Same as panel (a), but for CC variations. (d) Same as panel (b), but for CC variations.**

200

3.4 Spatial mapping of velocity change and seismic scattering coefficient

To estimate the region where the velocity and heterogeneous structure change, we map velocity changes and scattering coefficient changes (SCCs: Δg) using a method described previously (Obermann et al., 2013; 2014; 2015; Sánchez-Pastor et al., 2018; Hirose et al., 2019; 2020; Tonegawa et al., 2022). We limited the mapping of dv/v and Δg with available stations in nodes KMB, KMC, KMD, MRD, MRE, and temporary OBSs (Fig. 1), because the dv/v and CC variations are limited within the distribution of these stations. We investigate dv/v and CC changes during the tremor activities started on December 6, 2020, and finished on January 14, 2021. We used the reference values for the dv/v and CC averaged over November 21–30, 2020, and selected the values of dv/v and CC for Periods (1)–(6); (1) November 25, 2020, (2) December 10, 2020, (3) December 20,

205

2020, (4) December 30, 2020, (5) January 10, 2021, and (6) January 20, 2021. Because the used CCFs are stacked over 20
210 days, obtained results on dv/v and CC contain the effects between -9^{th} and $+10^{\text{th}}$ days from the reference dates. The methods
for estimating dv/v and are the same as those used in Tonegawa et al. (2022), and they are briefly summarized in Supplement.

4 Results

4.1 Clock deviation


215 Because stations of SHM1, SHM6c, and SHM7c are relatively close to DONET1 or DONET2, direct and scattered waves can
be retrieved in the CCFs. Consequently, as shown in Figs. 3 and S2, sufficient measurements of clock deviations can be
obtained for these OBSs. The clock deviations obtained at stations SHM4, and SHM6c are approximated by linear interpolation
before and after the observations, since the average values are close to 0 s over the entire observation period. On the other
hand, the average values of SHM1, SHM2, SHM3, SHM5 and SHM7c are slightly away from 0 s during the observation period,
220 which indicates non-linear clock deviations (Figs. 3 and S2). The average of the standard deviation of the clock deviations for
all the stations are <0.006 s (Table S1). Because the clocks of the recorders are adjusted to the clocks of the GNSS at the
timings before and after the observations, the estimated clock deviations at the starting and ending dates should be 0 s. However,
the clock deviation of SHM2 at the end of the observation is approximately -0.02 s. This is probably because the counts at
each day within 550–600 days are almost comparable and relatively high counts are not estimated, and hence the obtained
225 results are deviated from 0 s (Fig. S2b). Since the range of the high tremor activity is 445–485 days (December 5, 2020–
January 14, 2021; Fig. 1a), the clock estimations near the end of the observation period do not substantially influence the
results of dv/v and CC.

4.2 Spatio-temporal changes in dv/v and CC

230 Figures 4a and 4b show the temporal variations in dv/v at node KMC and OBSs, respectively. The dv/v changes at node KMC
shows an abrupt reduction of 0.1 % during the slow earthquake activity, while such reductions cannot be observed in the station
pairs of the OBSs. Substantial dv/v reductions are not observed at nodes MRD and MRE of DONET2 (Fig. S2c and S2d). This
fact indicates that the dv/v changes do not expand to the area of the OBSs. The reductions in dv/v at node KMC continued,
with a gradual recovery to the original dv/v values until the end of the observation period. Figures 4c and 4d display the
235 temporal variations in CC at node KMC and OBSs, respectively, and CC reductions are detected during the slow earthquake
activity in both the areas. In particular, the troughs of the CC variations in the OBS pairs are slightly shifted, which indicates
that they occur first in the eastern part of the survey area (SHM6c_SHM7c and SHM6c_SHM5) and later in the central part

(Fig. S4c). These reductions were almost recovered to the original levels of CC immediately after the slow earthquake activity. Another CC reduction can be observed in the middle of 2020 for 3 pairs of two stations at node KMC (Fig. 4c), and such features may be identified at other pairs, e.g., KMB06_KMB05 and KMB07_KMB08 with relatively weak levels (Fig. S4). The spatio-temporal variations in dv/v for Periods (1–6) are displayed in Fig 5. Substantial variations in dv/v are not observed in Period (1), whereas dv/v reduction can be observed in the eastern part of the survey area during Periods (2–6). This result is consistent with the results obtained at nodes KMC (Fig. 4), KMB and KMD (Fig. S2), where dv/v reduction is not recovered to the original level after the event. The spatiotemporal variations in Δg for Periods (1–6) are displayed in Fig 6. Large localized Δg is imaged at the eastern part of the survey area in Periods (2–5), and relatively weak Δg occurred at the western part in the same period. In addition, the large Δg region at the eastern part in Period (5) migrated to the central part in Period (6). Figure 7 shows the spatiotemporal relationship between Δg and tremor distributions (Tamaribuchi et al., 2022); here, tremors with estimated errors $\leq 0.1^\circ$ from the original catalog are preserved, and hence 84% of the tremors are excluded. These tremors occurred near the location of the maximum Δg . The large Δg in the eastern and western parts are merged on January 15–20, 2021, and substantial large Δg cannot be observed in February 2021.

We estimated dv/v and Δg for the dates of April 1, July 1, and October 1 in 2020 and March 1 and April 1 in 2021 using 20-day CCFs, and these dates in 2020 and 2021 correspond to non-slow-earthquake periods and post-slow-earthquake periods, respectively (Figs. S5 and S6). Δg does not change in these dates substantially. On the other hand, dv/v reductions relative to the reference period (November 21–30, 2020) are observed on April 1 and July 1 in 2020. This is attributed to the long-term increase of dv/v that has been observed in this area (Tonegawa et al. 2022). After the slow earthquake activity, although dv/v reductions are also observed on March 1 and April 1 in 2021, recoveries of the dv/v reductions are not clearly imaged. The dv/v changes between the activity and post activity appear to be small to detect in the mapping.



260 **Figure 5: Spatio-temporal dv/v variations in Periods (1–6).** Circles represent the tremor locations within ± 5 days from the reference dates, as determined by Tamaribuchi et al. (2022), from which tremors with spatial errors $\leq 0.1^\circ$ are plotted (16% of tremors from the original catalog). Squares and circles indicate the DONET1 and DONET2 stations, and triangles show temporary OBSs used in this analysis (KMB, KMC, KMD, MRD, MRE and temporary OBSs). Crosses represent the other stations.




Figure 6: Spatio-temporal scattering coefficient change (Δg) variations in Periods (1–6). Circles represent the tremor locations within ± 5 days from the reference dates, determined by Tamaribuchi et al. (2022), from which tremors with spatial errors $\leq 0.1^\circ$ are plotted (16% of tremors from the original catalog). Squares and circles indicate the DONET1 and DONET2 stations, and triangles show temporary OBSs used in this analysis (KMB, KMC, KMD, MRD, MRE and temporary OBSs). Crosses represent the other stations. The amplitudes in Period (1) are low, and the different color scale is used.

5 Discussion

5.1 Spatio-temporal relationship of dv/v and $SCC(\Delta g)$ with tremors

A localized area of the dv/v reduction is observed in the eastern part of the survey area (Fig. 5). The dv/v reduction and tremor activities started from Period (2). Although the location of the maximum dv/v reduction is only at a small distance from the high activity area of tremors during Periods (2–3), the dv/v reduction area cover the tremor distribution. The tremor activity area migrated westward in Periods (4–5) and reached the central and western parts of the survey area; however, there is no considerable change in the dv/v reduction area. These facts indicate that the areas of dv/v reduction and tremors are correlated only at the early stages of the episode.

The large Δg area is also localized in the eastern part of the survey area, and the extent of localization is higher than that of the dv/v reduction area. The large Δg was initiated at the timing of the tremor activity in Period (2). Although the location of the

maximum Δg slightly shifted westward from the tremor activity area, the relatively large Δg area surrounding the maximum Δg cover the area of the tremors as well as that of the dv/v reduction. The tremor area moved to the location of the maximum Δg in Periods (3–4). The areas of tremors moved westward in Period (5), but the CC reduction area did not change. However, in Period (6), the CC reduction area moved westward approximately 10 days after the westward tremor migrations observed in Period (5) (Figs. 6 and 7).




Figure 7: Spatiotemporal scattering coefficient change (Δg) variations with tremors. (a) Tremor distribution (Tamaribuchi et al., 2022) and the line X–X' at which the Δg and tremors are projected. Other symbols are the same as those used in Fig. 1. (b) Tremors and Δg projected onto the line X–X' in panel (a). Arrows show that large Δg in the western and eastern parts are merged.

5.2 Interpretation of temporal change difference in dv/v and $\text{SCC}(\Delta g)$

Our results indicate that the dv/v shows a step-like drop at the time of the slow earthquakes that began from the end of 2020, whereas the CC changes are recovered to the original level after the slow earthquake activity. Because CC reductions were observed during SSEs without or less tremor activities, the effects of the tremor signal contaminations on CC measurements appear minor (Tonegawa et al., 2022). Although a step-like dv/v reduction was observed in this area due to the strong seafloor motions of the 2016 off-Mie earthquake (Mw6.0) that occurred beneath the DONET1 (Ikeda and Tsuji, 2018; Tonegawa et al., 2022), such strong motions were not observed during the slow earthquake episode. Although slow earthquake activity with tremors and VLFs occurred in October 2015 (e.g., Annoura et al. 2017; Baba et al., 2020), dv/v reductions were not observed during this event (Ikeda and Tsuji, 2018; Tonegawa et al., 2022). Therefore, it appears that the dv/v and CC changes obtained in this study were a result of different factors that occurred within the accretionary prism.

A candidate for the observed dv/v reduction is the change in crack conditions due to the deformation within the accretionary prism (Fig. 8a). Prior to the slow earthquake activity, the compressional stress in the trough-normal direction is dominant in the accretionary prism, and cracks with a long axis oriented in the trough-normal direction are prevalent (e.g., Crampin 1977; 1981) (Fig. 8b). Slow earthquakes at the shallow plate interface generate extensional stress within the overlying plate, which

reduces the compressional stress and results in the extensional deformation of the accretionary prism along the trough-normal direction. Such deformations induce changes in the aspect ratio of the pre-existing cracks normal to the trough axis, and new cracks parallel to the axis are created. The updated conditions of the stress field and cracks are preserved with a small degree of recovery after the slow earthquakes, and such recoveries can be observed from the beginning of 2021 to the end of the observation period at nodes KMB, KMC, and KMD (Figs. 4 and S2). In our observation, because the obtained dv/v changes are spatially limited to the eastern part of the survey region owing to the concentrated occurrence of tremors here (Tamaribuchi et al. 2022; Ogiso and Tamaribuchi, 2022), the deformation of the sediment also appears to be concentrated in this region. Thus, the mechanism that crack states are altered by the sediment deformation supports the step-like dv/v reduction observed in the region where tremors prevalently occurred.

In contrast, the CC reductions are recovered to the original level after the slow earthquake activity. Tonegawa et al. (2022) interpreted that the CC reduction and its recovery are caused by transient upward fluid migrations at local scales (Fig. 8c). Based on the observations in this study and Tonegawa et al. (2022) that upward fluid migrations intermittently occur, we suggest a possible scenario for fluid storage and upward migration in this paragraph, and fluid trapping in the subsequent paragraph. Before slow earthquakes, the pore fluid pressure at the source region tends to increase with the fluid supply from the underlying oceanic crust through dehydration reactions (e.g., Kameda et al., 2011) and the accretionary prism itself by the tectonic compression (Saffer and Bekins, 1998). Although the uncertainties in the tremor depths are still high, the horizontal spatial coincidence between the VLFs and the low velocity layer within the accretionary prism has been observed (Tsuji et al. 2014; Tonegawa et al., 2022; Akuhara et al., 2020) (Fig. 8a). Several studies have also investigated whether tremor occurrences are related to the low velocity layer (Hendriyana and Tsuji, 2021; Fahrudin et al., 2022; Akuhara et al., 2023b). The low velocity layer has a thickness of ~ 1 km and a horizontal distance of ~ 15 km in the dip direction (Park et al., 2010). Because a high pore-pressure region with a width of several hundred meters was found by a seafloor drilling program (Hirose et al., 2021), it seems that the low velocity layer consists of many such high pore-pressure volumes. Thus, it appears that fluids are trapped by impermeable caprocks in localized areas with low seismic velocities. Numerical simulations with hydraulic models also indicate the importance of impermeable zones to locally increase pore pressures (Kaneki and Noda, 2023). A previous study has reported that temporal changes in seismic velocity and anisotropic structures occurred during the rupture of a slow-slip patch in the Hikurangi subduction zone (Zal et al., 2020). When slow earthquakes occurred at the end of 2020, the caprocks fractured, and the trapped fluids migrated upwards. Our results of CC reductions reflect such fluid migration passing through shallow depths. In particular, the CC reductions in Period (6) are observed approximately 10 days after the tremor activities in the central part (Figs. 7 and 8e), which may correspond to the fracture-induced fluid migrations, and this phenomenon was not documented in Tonegawa et al. (2022).

To increase pore fluid pressures at the source region, mechanisms that trap fluids within the accretionary prism are required, and the topographic relief of the top surface of the oceanic crust may be invoked for the fluid trapping. The topographic relief induces deformation at the downdip side of the relief (Fig. 8c). Such deformations intensively form horizontally sheeted clay minerals in and around the deformation area immediately above the oceanic crust, which can create impermeable caprocks.

As a result, fluids cannot escape laterally or vertically and are trapped below the impermeable sheets; eventually the pore pressure becomes higher than that in the surrounding regions. Such fluid-trapped areas would have been created locally in several parts of the Nankai subduction zone, which may be related to the sporadic occurrences of the VLFs. Indeed, VLFs occur on the downdip sides of the topographic relief (Shiraishi et al. 2020; Hashimoto et al. 2022). Moreover, several discontinuous reflections with high impedance contrasts are imaged within the low velocity zone (Tsuji et al., 2014), which may represent impermeable sheets due to relief-induced deformations.




Figure 8: Interpretation for our results with tremor and fluid distributions. (a) Illustration for crack creations due to the deformation causing step-like reduction in dV/V . (b) Crack conditions for trough-normal compressional stress, and newly created cracks and aspect ratio changes of pre-existing cracks during slow earthquakes. (c) Upward fluid migration during tremors (slow earthquakes) in the cross section along the dipping direction. (d) Fluids are trapped by the impermeable sheets that are created by the deformation by topographic reliefs. (e) Spatio-temporal distributions of tremors and upward fluid migrations in the cross sections along the strike direction. Crosses indicate the locations of caprock fractures.

6 Conclusion

We present the temporal variations in dv/v and CC in the Nankai subduction zone associated with the slow earthquake activity that began at the end of 2020. The dv/v variations exhibit the step-like reductions at several pairs of stations that are located in the eastern part of the survey area. These features are considered to be constructed by changes in the aspect ratios of pre-existing cracks and/or newly created cracks due to the sediment deformation induced by the slow earthquake episode. The CC variations show the transient reductions at the eastern and western parts of the survey area, which are spatially correlated with the tremor distribution. It is considered that the CC reductions can be linked to the transient fluid migration from the source region of the slow earthquakes. Thus, dv/v and CC are sensitive to different factors in the subseafloor structure, and we suggest that both the dv/v and CC variations be monitored for understanding the slow earthquake generations and their influence on the surrounding structure.

Data availability

DONET data can be downloaded from the website operated by National Research Institute for Earth Science and Disaster Resilience (NIED) (<https://doi.org/10.17598/NIED.0008>). dv/v and CC data used in this study can be downloaded from a repository Zenodo (<https://zenodo.org/records/15048965>).

Author contribution

T.T processed the data, and drafted the manuscript. T.A, S.T, T.Ts edited the manuscript and contributed to the interpretation. T.A, Y.Y, H.S, M.S aimed to acquire the data. All authors contributed to the final version of the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

The comments from Takashi Hirose and Köpfli, Manuela are constructive and greatly improved the manuscript. This work was supported by JSPS KAKENHI Grant No. JP21H05202 in Scientific Research on Transformative Research Areas “Science of Slow-to-Fast earthquakes” and Cooperative Research Program of Atmosphere and Ocean Research Institute, The University of Tokyo (Research Vessel Shinsei-maru, SH19-27 and S21-31).

References

- Akuhara, T., Tsuji, T., and Tonegawa, T.: Overpressured Underthrust Sediment in the Nankai Trough Forearc Inferred From Transdimensional Inversion of High-Frequency Teleseismic Waveforms, *Geophysical Research Letters*, 47, e2020GL088280, <https://doi.org/10.1029/2020GL088280>, 2020.
- Akuhara, T., Yamashita, Y., Ohyanagi, S., Sawaki, Y., Yamada, T., and Shinohara, M.: Shallow Low-Velocity Layer in the Hyuga-Nada Accretionary Prism and Its Hydrological Implications: Insights From a Passive Seismic Array, *JGR Solid Earth*, 128, e2022JB026298, <https://doi.org/10.1029/2022JB026298>, 2023a.
- Akuhara, T., Yamashita, Y., Sugioka, H., and Shinohara, M.: Locating tectonic tremors with uncertainty estimates: time- and amplitude-difference optimization, wave propagation-based quality control and Bayesian inversion, *Geophysical Journal International*, 235, 2727–2742, <https://doi.org/10.1093/gji/ggad387>, 2023b.
- Annoura, S., Hashimoto, T., Kamaya, N., and Katsumata, A.: Shallow episodic tremor near the Nankai Trough axis off southeast Mie prefecture, Japan, *Geophysical Research Letters*, 44, 3564–3571, <https://doi.org/10.1002/2017GL073006>, 2017.
- Aoi, S., Asano, Y., Kunugi, T., Kimura, T., Uehira, K., Takahashi, N., Ueda, H., Shiomi, K., Matsumoto, T., and Fujiwara, H.: MOWLAS: NIED observation network for earthquake, tsunami and volcano, *Earth Planets Space*, 72, 126, <https://doi.org/10.1186/s40623-020-01250-x>, 2020.
- Arai, R., Miura, S., Nakamura, Y., Fujie, G., Kodaira, S., Kaiho, Y., Mochizuki, K., Nakata, R., Kinoshita, M., Hashimoto, Y., Hamada, Y., and Okino, K.: Upper-plate conduits linked to plate boundary that hosts slow earthquakes, *Nat Commun*, 14, 5101, <https://doi.org/10.1038/s41467-023-40762-4>, 2023.
- Araki, E., Saffer, D. M., Kopf, A. J., Wallace, L. M., Kimura, T., Machida, Y., Ide, S., Davis, E., and IODP Expedition 365 shipboard scientists: Recurring and triggered slow-slip events near the trench at the Nankai Trough subduction megathrust, *Science*, 356, 1157–1160, <https://doi.org/10.1126/science.aan3120>, 2017.
- Ariyoshi, K., Iinuma, T., Nakano, M., Kimura, T., Araki, E., Machida, Y., Sueki, K., Yada, S., Nishiyama, T., Suzuki, K., Hori, T., Takahashi, N., and Kodaira, S.: Characteristics of Slow Slip Event in March 2020 Revealed From Borehole and DONET Observatories, *Front. Earth Sci.*, 8, 600793, <https://doi.org/10.3389/feart.2020.600793>, 2021.
- Asano, Y., Obara, K., and Ito, Y.: Spatiotemporal distribution of very-low frequency earthquakes in Tokachi-oki near the junction of the Kuril and Japan trenches revealed by using array signal processing, *Earth Planet Sp*, 60, 871–875, 2008.
- Audet, P. and Kim, Y.: Teleseismic constraints on the geological environment of deep episodic slow earthquakes in subduction zone forearcs: A review, *Tectonophysics*, 670, 1–15, <https://doi.org/10.1016/j.tecto.2016.01.005>, 2016.

- Baba, T., Tanioka, Y., Cummins, P. R., and Uhira, K.: The slip distribution of the 1946 Nankai earthquake estimated from tsunami inversion using a new plate model, *Physics of the Earth and Planetary Interiors*, 132, 59–73, [https://doi.org/10.1016/S0031-9201\(02\)00044-4](https://doi.org/10.1016/S0031-9201(02)00044-4), 2002.
- 415 Baba, S., Takemura, S., Obara, K., and Noda, A.: Slow Earthquakes Illuminating Interplate Coupling Heterogeneities in Subduction Zones, *Geophys. Res. Lett.*, 47, <https://doi.org/10.1029/2020GL088089>, 2020.
- Bangs, N. L., Morgan, J. K., Bell, R. E., Han, S., Arai, R., Kodaira, S., Gase, A. C., Wu, X., Davy, R., Frahm, L., Tilley, H. L., Barker, D. H. N., Edwards, J. H., Tobin, H. J., Reston, T. J., Henrys, S. A., Moore, G. F., Bassett, D., Kellett, R., Stucker, V., and Fry, B.: Slow slip along the Hikurangi margin linked to fluid-rich sediments trailing subducting seamounts, *Nat. Geosci.*, 16, 505–512, <https://doi.org/10.1038/s41561-023-01186-3>, 2023.
- 420 Brenguier, F., Shapiro, N. M., Campillo, M., Nercissian, A., and Ferrazzini, V.: 3-D surface wave tomography of the Piton de la Fournaise volcano using seismic noise correlations, *Geophys. Res. Lett.*, 34, 1351–5, <https://doi.org/10.1029/2006GL028586>, 2007.
- Brenguier, F., Shapiro, N. M., Campillo, M., Ferrazzini, V., Duputel, Z., Coutant, O., and Nercissian, A.: Towards forecasting volcanic eruptions using seismic noise, *Nature Geoscience*, 1, 126–130, 2008.
- 425 Campillo, M. and Paul, A.: Long-Range Correlations in the Diffuse Seismic Coda, *Science*, 299, 547–549, <https://doi.org/10.1126/science.1078551>, 2003.
- Crampin, S.: A review of the effects of anisotropic layering on the propagation of seismic waves, *Geophys J Int*, 49, 9–27, <https://doi.org/10.1111/j.1365-246X.1977.tb03698.x>, 1977.
- 430 Crampin, S.: A review of wave motion in anisotropic and cracked elastic-media, *Wave Motion*, 3, 343–391, [https://doi.org/10.1016/0165-2125\(81\)90026-3](https://doi.org/10.1016/0165-2125(81)90026-3), 1981.
- Cruz-Atienza, V. M., Villafuerte, C., and Bhat, H. S.: Rapid tremor migration and pore-pressure waves in subduction zones, *Nat Commun*, 9, 2900, <https://doi.org/10.1038/s41467-018-05150-3>, 2018.
- Fahrudin, Chhun, C., and Tsuji, T.: Influence of shear zone thickness and strike-slip faulting on tectonic tremor in the Nankai Trough, southwest Japan, *Tectonophysics*, 838, 229519, <https://doi.org/10.1016/j.tecto.2022.229519>, 2022.
- 435 Gosselin, J. M., Audet, P., Esteve, C., McLellan, M., Mosher, S. G., and Schaeffer, A. J.: Seismic evidence for megathrust fault-valve behavior during episodic tremor and slip, *Science Advances*, 6, eaay5174, 2020.
- Hashimoto, Y., Sato, S., Kimura, G., Kinoshita, M., Miyakawa, A., Moore, G. F., Nakano, M., Shiraishi, K., and Yamada, Y.: Décollement geometry controls on shallow very low frequency earthquakes, *Sci Rep*, 12, 2677, <https://doi.org/10.1038/s41598-022-06645-2>, 2022.
- 440 Hendriyana, A. and Tsuji, T.: Influence of structure and pore pressure of plate interface on tectonic tremor in the Nankai subduction zone, Japan, *Earth and Planetary Science Letters*, 558, 116742, <https://doi.org/10.1016/j.epsl.2021.116742>, 2021.
- Hirose, F., Nakajima, J., and Hasegawa, A.: Three-dimensional seismic velocity structure and configuration of the Philippine Sea slab in southwestern Japan estimated by double-difference tomography, *J. Geophys. Res.*, 113, 59–26, <https://doi.org/10.1029/2007JB005274>, 2008.
- 445

- Hirose, T., Nakahara, H., and Nishimura, T.: A Passive Estimation Method of Scattering and Intrinsic Absorption Parameters From Envelopes of Seismic Ambient Noise Cross-Correlation Functions, *Geophys. Res. Lett.*, 46, 3634–3642, <https://doi.org/10.1029/2018GL080553>, 2019.
- 450 Hirose, T., Nakahara, H., Nishimura, T., and Campillo, M.: Locating Spatial Changes of Seismic Scattering Property by Sparse Modeling of Seismic Ambient Noise Cross-Correlation Functions: Application to the 2008 Iwate-Miyagi Nairiku (M_w 6.9), Japan, Earthquake, *J. Geophys. Res. Solid Earth*, 125, <https://doi.org/10.1029/2019JB019307>, 2020.
- Hirose, T., Hamada, Y., Tanikawa, W., Kamiya, N., Yamamoto, Y., Tsuji, T., Kinoshita, M., Heuer, V. B., Inagaki, F., Morono, Y., and Kubo, Y.: High Fluid-Pressure Patches Beneath the Décollement: A Potential Source of Slow Earthquakes in the Nankai Trough off Cape Muroto, *JGR Solid Earth*, 126, e2021JB021831, <https://doi.org/10.1029/2021JB021831>, 2021.
- 455 Ikeda, T. and Tsuji, T.: Temporal change in seismic velocity associated with an offshore MW 5.9 Off-Mie earthquake in the Nankai subduction zone from ambient noise cross-correlation, *Prog Earth Planet Sci*, 5, 62, <https://doi.org/10.1186/s40645-018-0211-8>, 2018.
- Ito, Y. and Obara, K.: Very low frequency earthquakes within accretionary prisms are very low stress-drop earthquakes, *Geophys. Res. Lett.*, 33, L09302, <https://doi.org/10.1029/2006GL025883>, 2006.
- 460 Ito, Y. and Nakajima, J.: Temporal Variations in Frequency-Dependent Shear-Wave Anisotropy Above a Plate Interface Following Episodic Slow-Slip Events, *JGR Solid Earth*, 129, e2023JB028526, <https://doi.org/10.1029/2023JB028526>, 2024.
- Kameda, J., Yamaguchi, A., Saito, S., Sakuma, H., Kawamura, K., and Kimura, G.: A new source of water in seismogenic subduction zones, *Geophys. Res. Lett.*, 38, n/a-n/a, <https://doi.org/10.1029/2011GL048883>, 2011.
- Kamei, R., Pratt, R. G., and Tsuji, T.: Waveform tomography imaging of a megasplay fault system in the seismogenic Nankai subduction zone, *Earth and Planetary Science Letters*, 317–318, 343–353, <https://doi.org/10.1016/j.epsl.2011.10.042>, 2012.
- 465 Kaneda, Y., Kawaguchi, K., Araki, E., Matsumoto, H., Nakamura, T., Kamiya, S., Ariyoshi, K., Hori, T., Baba, T., and Takahashi, N.: Development and application of an advanced ocean floor network system for megathrust earthquakes and tsunamis, in: *SEAFLOOR OBSERVATORIES*, edited by: Favali, P., Beranzoli, L., and De Santis, A., Springer Berlin Heidelberg, Berlin, Heidelberg, 643–662, https://doi.org/10.1007/978-3-642-11374-1_25, 2015.
- 470 Kaneki, S. and Noda, H.: Steady-State Effective Normal Stress in Subduction Zones Based on Hydraulic Models and Implications for Shallow Slow Earthquakes, *JGR Solid Earth*, 128, e2022JB025995, <https://doi.org/10.1029/2022JB025995>, 2023.
- Kaneko, L., Ide, S., and Nakano, M.: Slow Earthquakes in the Microseism Frequency Band (0.1–1.0 Hz) off Kii Peninsula, Japan, *Geophysical Research Letters*, 45, 2618–2624, <https://doi.org/10.1002/2017GL076773>, 2018.
- 475 Kawaguchi, K., Kaneko, S., Nishida, T., and Komine, T.: Construction of the DONET real-time seafloor observatory for earthquakes and tsunami monitoring, in: *SEAFLOOR OBSERVATORIES*, edited by: Favali, P., Beranzoli, L., and De Santis, A., Springer Berlin Heidelberg, Berlin, Heidelberg, 211–228, https://doi.org/10.1007/978-3-642-11374-1_10, 2015.

- Kitajima, H. and Saffer, D. M.: Elevated pore pressure and anomalously low stress in regions of low frequency earthquakes along the Nankai Trough subduction megathrust, *Geophys. Res. Lett.*, 39, L23301, <https://doi.org/10.1029/2012GL053793>, 2012.
- Nakajima, J. and Hasegawa, A.: Tremor activity inhibited by well-drained conditions above a megathrust, *Nat Commun*, 7, 13863, <https://doi.org/10.1038/ncomms13863>, 2016.
- Nakajima, J. and Uchida, N.: Repeated drainage from megathrusts during episodic slow slip, *Nature Geosci*, 11, 351–356, <https://doi.org/10.1038/s41561-018-0090-z>, 2018.
- Nakano, M., Nakamura, T., Kamiya, S., Ohori, M., and Kaneda, Y.: Intensive seismic activity around the Nankai trough revealed by DONET ocean-floor seismic observations, *Earth Planet Sp*, 65, 5–15, <https://doi.org/10.5047/eps.2012.05.013>, 2013.
- Nakano, M., Hori, T., Araki, E., Kodaira, S., and Ide, S.: Shallow very-low-frequency earthquakes accompany slow slip events in the Nankai subduction zone, *Nature Communications*, 9, 984, <https://doi.org/10.1038/s41467-018-03431-5>, 2018.
- Obara, K. and Kato, A.: Connecting slow earthquakes to huge earthquakes, *Science*, 353, 253–257, <https://doi.org/10.1126/science.aaf1512>, 2016.
- Obara, K. and Ito, Y.: Very low frequency earthquakes excited by the 2004 off the Kii peninsula earthquakes: A dynamic deformation process in the large accretionary prism, *Earth Planet Sp*, 57, 321–326, 2005.
- Obermann, A., Planès, T., Larose, E., and Campillo, M.: Imaging preeruptive and coeruptive structural and mechanical changes of a volcano with ambient seismic noise, *J. Geophys. Res. Solid Earth*, 118, 6285–6294, <https://doi.org/10.1002/2013JB010399>, 2013.
- Obermann, A., Froment, B., Campillo, M., Larose, E., Planès, T., Valette, B., Chen, J. H., and Liu, Q. Y.: Seismic noise correlations to image structural and mechanical changes associated with the M_w 7.9 2008 Wenchuan earthquake, *J. Geophys. Res. Solid Earth*, 119, 3155–3168, <https://doi.org/10.1002/2013JB010932>, 2014.
- Obermann, A., Kraft, T., Larose, E., and Wiemer, S.: Potential of ambient seismic noise techniques to monitor the St. Gallen geothermal site (Switzerland):, *J. Geophys. Res. Solid Earth*, 120, 4301–4316, <https://doi.org/10.1002/2014JB011817>, 2015.
- Ogiso, M. and Tamaribuchi, K.: Spatiotemporal evolution of tremor activity near the Nankai Trough trench axis inferred from the spatial distribution of seismic amplitudes, *Earth Planets Space*, 74, 49, <https://doi.org/10.1186/s40623-022-01601-w>, 2022.
- Paasschens, J. C. J.: Solution of the time-dependent Boltzmann equation, *Phys. Rev. E.*, 56(1), 1135–1141, <https://doi.org/10.1103/PhysRevE.56.1135>, 1997.
- Park, J.-O., Fujie, G., Wijerathne, L., Hori, T., Kodaira, S., Fukao, Y., Moore, G. F., Bangs, N. L., Kuramoto, S., and Taira, A.: A low-velocity zone with weak reflectivity along the Nankai subduction zone, *Geology*, 38, 283–286, <https://doi.org/10.1130/G30205.1>, 2010.
- Saffer, D. M. and Bekins, B. A.: Episodic fluid flow in the Nankai accretionary complex: Timescale, geochemistry, flow rates, and fluid budget, 103, 30351–30370, 1998.

- Sánchez-Pastor, P., Obermann, A., and Schimmel, M.: Detecting and Locating Precursory Signals During the 2011 El Hierro, Canary Islands, Submarine Eruption, *Geophysical Research Letters*, 45, <https://doi.org/10.1029/2018GL079550>, 2018.
- Sato, H.: Energy transportation in one- and two-dimensional scattering media: Analytic solutions of the multiple isotropic scattering model, *Geophys. J. Int.*, 112, 141–146, <https://doi.org/10.1111/j.1365-246X.1993.tb01443.x>, 1993.
- Sen, M. and Stoffa, P.: *Global Optimization Methods in Geophysical Inversion*, Elsevier Sci., New York, 1995.
- Sens-Schonfelder, C. and Wegler, U.: Passive image interferometry and seasonal variations of seismic velocities at Merapi Volcano, Indonesia, *Geophys. Res. Lett.*, 33, 547–5, <https://doi.org/10.1029/2006GL027797>, 2006.
- Shapiro, N. M., Campillo, M., Stehly, L., and Ritzwoller, M. H.: High-Resolution Surface-Wave Tomography from Ambient Seismic Noise, *Science*, 307, 1615–1618, <https://doi.org/10.1126/science.1108339>, 2005.
- Shelly, D. R., Beroza, G. C., Ide, S., and Nakamura, S.: Low-frequency earthquakes in Shikoku, Japan, and their relationship to episodic tremor and slip, *Nature*, 442, 188–191, <https://doi.org/10.1038/nature04931>, 2006.
- Shiraishi, K., Yamada, Y., Nakano, M., Kinoshita, M., and Kimura, G.: Three-dimensional topographic relief of the oceanic crust may control the occurrence of shallow very-low-frequency earthquakes in the Nankai Trough off Kumano, *Earth Planets Space*, 72, 72, <https://doi.org/10.1186/s40623-020-01204-3>, 2020.
- Sugioka, H., Okamoto, T., Nakamura, T., Ishihara, Y., Ito, A., Obana, K., Kinoshita, M., Nakahigashi, K., Shinohara, M., and Fukao, Y.: Tsunamigenic potential of the shallow subduction plate boundary inferred from slow seismic slip, *Nature Geoscience*, 5, 414–418, <https://doi.org/10.1038/ngeo1466>, 2012.
- Takemura, S., Matsuzawa, T., Noda, A., Tonegawa, T., Asano, Y., Kimura, T., and Shiomi, K.: Structural Characteristics of the Nankai Trough Shallow Plate Boundary Inferred From Shallow Very Low Frequency Earthquakes, *Geophysical Research Letters*, 46, 4192–4201, <https://doi.org/10.1029/2019GL082448>, 2019.
- Takemura, S., Hamada, Y., Okuda, H., Okada, Y., Okubo, K., Akuhara, T., Noda, A., and Tonegawa, T.: A review of shallow slow earthquakes along the Nankai Trough, *Earth Planets Space*, 75, 164, <https://doi.org/10.1186/s40623-023-01920-6>, 2023.
- Takeo, A., Forsyth, D. W., Weeraratne, D. S., and Nishida, K.: Estimation of azimuthal anisotropy in the NW Pacific from seismic ambient noise in seafloor records, *Geophysical Journal International*, 199, 11–22, <https://doi.org/10.1093/gji/ggu240>, 2014.
- Tamaribuchi, K., Ogiso, M., and Noda, A.: Spatiotemporal Distribution of Shallow Tremors Along the Nankai Trough, Southwest Japan, as Determined From Waveform Amplitudes and Cross-Correlations, *JGR Solid Earth*, 127, e2022JB024403, <https://doi.org/10.1029/2022JB024403>, 2022.
- Thery, R., Guillemot, A., Abraham, O., and Larose, E.: Tracking fluids in multiple scattering and highly porous materials: Toward applications in non-destructive testing and seismic monitoring, *Ultrasonics*, 102, 106019, <https://doi.org/10.1016/j.ultras.2019.106019>, 2020.
- Tonegawa, T., Fukao, Y., Takahashi, T., Obana, K., Kodaira, S., and Kaneda, Y.: Ambient seafloor noise excited by earthquakes in the Nankai subduction zone, *Nature Communications*, 6, 6132, <https://doi.org/10.1038/ncomms7132>, 2015.

- 545 Tonegawa, T., Araki, E., Kimura, T., Nakamura, T., Nakano, M., and Suzuki, K.: Sporadic low-velocity volumes spatially correlate with shallow very low frequency earthquake clusters, *Nature Communications*, 8, 2048, <https://doi.org/10.1038/s41467-017-02276-8>, 2017.
- Tonegawa, T., Kimura, T., and Araki, E.: Near-Field Body-Wave Extraction From Ambient Seafloor Noise in the Nankai Subduction Zone, *Front. Earth Sci.*, 8, 610993, <https://doi.org/10.3389/feart.2020.610993>, 2021.
- 550 Tonegawa, T., Takemura, S., Yabe, S., and Yomogida, K.: Fluid Migration Before and During Slow Earthquakes in the Shallow Nankai Subduction Zone, *JGR Solid Earth*, 127, <https://doi.org/10.1029/2021JB023583>, 2022.
- Tsuji, T., Kamei, R., and Pratt, R. G.: Pore pressure distribution of a mega-splay fault system in the Nankai Trough subduction zone: Insight into up-dip extent of the seismogenic zone, *Earth and Planetary Science Letters*, 396, 165–178, <https://doi.org/10.1016/j.epsl.2014.04.011>, 2014.
- 555 Tsuji, S., Araki, E., Yokobiki, T., Nishida, S., Machida, Y., Zumberge, M., and Takahashi, K.: Precise tilt measurement by seafloor borehole tiltmeters at the Nankai Trough subduction zone, *Earth Planets Space*, 75, 188, <https://doi.org/10.1186/s40623-023-01946-w>, 2023.
- Yabe, S., Tonegawa, T., and Nakano, M.: Scaled Energy Estimation for Shallow Slow Earthquakes, *J. Geophys. Res. Solid Earth*, <https://doi.org/10.1029/2018JB016815>, 2019.
- 560 Yamamoto, Y., Ariyoshi, K., Yada, S., Nakano, M., and Hori, T.: Spatio-temporal distribution of shallow very-low-frequency earthquakes between December 2020 and January 2021 in Kumano-nada, Nankai subduction zone, detected by a permanent seafloor seismic network, *Earth Planets Space*, 74, 14, <https://doi.org/10.1186/s40623-022-01573-x>, 2022.
- Yokota, Y. and Ishikawa, T.: Shallow slow slip events along the Nankai Trough detected by GNSS-A, *Science Advances*, 6, eaay5786, 2020.
- 565 Zal, H. J., Jacobs, K., Savage, M. K., Yarce, J., Mroczek, S., Graham, K., Todd, E. K., Nakai, J., Iwasaki, Y., Sheehan, A., Mochizuki, K., Wallace, L., Schwartz, S., Webb, S., and Henrys, S.: Temporal and spatial variations in seismic anisotropy and V/V ratios in a region of slow slip, *Earth and Planetary Science Letters*, 532, 115970, <https://doi.org/10.1016/j.epsl.2019.115970>, 2020.