

Analysis of a saline dust storm from the Aralkum Desert - Part 1:  
Consistency **between** multisensor satellite aerosol products

Xin Xi<sup>1</sup>, Jun Wang<sup>2</sup>, Zhendong Lu<sup>3</sup>, Andrew M. Sayer<sup>4,5</sup>, Jaehwa Lee<sup>4,6</sup>, Robert Levy<sup>4</sup>, Yujie Wang<sup>4,5</sup>, Alexei Lyapustin<sup>4</sup>, Hongqing Liu<sup>7,8</sup>, Istvan Laszlo<sup>7,9</sup>, Changwoo Ahn<sup>4,10</sup>, Omar Torres<sup>4</sup>, Sabur Abdullaev<sup>11</sup>, James Limbacher<sup>12</sup>, and Ralph Kahn<sup>13</sup>

<sup>1</sup>Department of Geological and Mining Engineering and Sciences, Michigan Technological University, Houghton, MI, USA  
<sup>2</sup>Department of Chemical and Biochemical Engineering, The University of Iowa, Iowa City, IA, USA

<sup>3</sup>Interdisciplinary Graduate Program in Informatics, The University of Iowa, Iowa City, IA, USA

<sup>4</sup>NASA Goddard Space Flight Center, Greenbelt, MD, USA

<sup>5</sup>Goddard Earth Sciences Technology and Research (GESTAR) II, University of Maryland, Baltimore County, Baltimore, MD, USA

<sup>6</sup>Earth System Science Interdisciplinary Center, University of Maryland, College Park, MD, USA

<sup>7</sup>Center for Satellite Applications and Research, National Environmental Satellite, Data, and Information Service, National Oceanic and Atmospheric Administration, College Park, MD, USA

<sup>8</sup>I. M. Systems Group, Inc., College Park, MD, USA

<sup>9</sup>Department of Atmospheric and Oceanic Sciences, University of Maryland, College Park, MD, USA

<sup>10</sup>Science Systems and Applications Inc., Lanham, MD, USA

<sup>11</sup>Physical Technical Institute of the Academy of Sciences of Tajikistan, Dushanbe, Tajikistan

<sup>12</sup>Science and Technology Corporation, Hampton, VA, USA

<sup>13</sup>Laboratory for Atmospheric and Space Physics, The University of Colorado Boulder, Boulder, CO, USA

Correspondence: Xin Xi (xinxi@mtu.edu)

**Abstract,** The Aralkum Desert is a prominent source of saline dust and presents a challenging environment for satellite-based aerosol observations, due to lack of in situ observational constraints on aerosol microphysical properties and the region's highly heterogeneous, dynamic surfaces. Through a survey of satellite aerosol algorithms, we identify diverse retrieval techniques and differing assumptions about surface and aerosol microphysical properties, which may deviate from actual conditions and contribute to inconsistencies in characterizing the saline dust from Aralkum. The inconsistencies and potential biases of satellite products in detecting the dust presence, column burden, and vertical height are further assessed via detailed comparisons of multiple UVAI, midvisible and thermal-infrared AOD, and ALH products for a saline dust storm from Aralkum Desert. The key findings are: (1) UVAI products consistently delineate the dust plume extent, once differences in the dynamic range are considered. All products display large positive values over the Garabogazköl Gulf and northern Caspian Sea due to absorption by turbid saline waters, highlighting the need for caution to avoid misinterpreting surface effects as dust signals over dry or ephemeral lakes. (2) Midvisible aerosol products from MODIS and VIIRS show strong agreement in total and coarse-mode AOD retrievals over the Caspian Sea, despite using inconsistent aerosol optical models. Over desert surfaces, all operational products misclassify the fresh dust plume as clouds and exhibit strong non-linear relationships in AOD retrievals. The NOAA EPS product retrieves significantly lower AOD than other products, though its agreement improves after a dust optical model is selected for retrieval. The MISR research aerosol algorithm produces higher, more consistent AOD retrievals and improved

characterization of particle properties compared to the MISR operational product. (3) Among four IASI infrared aerosol products, the CLIP algorithm has the best performance in capturing dust plume features over both desert and water surfaces. Preprint Discussion started 2 December 2024

© ARKUM product overestimates aerosol heights at low aerosol loadings but exhibits good consistency with CALIOP. CALIOP-weighted ALH for dust-laden scenes. MISR height retrieval also aligns well with CALIOP and EPIC in detecting the elevated dust with well-defined upper boundary. On average, IASI-retrieved ALH over dust-affected areas is 0.4 km higher than EPIC. This study underscores the important insights from a synergistic, multisensor analysis into the complementary strengths and potential biases of satellite aerosol products in Central Asia and highlights the need for their proper use and careful interpretation when characterizing the saline dust from the Aralkum Desert.

## 1 Introduction

Emerging from the desiccated basin of the former Aral Sea, the Aralkum Desert has evolved into one of the most active sources of wind-blown dust in Central Asia, with adverse impacts on biodiversity, agriculture, and human well-being (Orlovsky and Orlovsky, 2001; Xi and Sokolik, 2016). Aralkum stands out among the extensive dust sources of Central Asia, not only because of its anthropogenic origin but also due to the distinct chemical and mineralogical compositions of its deflatable sediments. Figure 1 shows the spectral abundance of four mineral groups derived from the Earth Surface Mineral Dust Source Investigation (EMIT) instrument. EMIT reveals that Aralkum contains more abundant carbonate and sulfate minerals, but less iron oxides and clays than the nearby Karakum and Taklamakan deserts. Groll et al. (2019) confirmed that dust samples collected near Aralkum contained considerably higher sulfate and chloride content than those from sandy deserts. Given its distinct mineralogical compositions, Aralkum dust is expected to be more hygroscopic and less light-absorbing than typical desert dust (Sokolik and Toon, 1999). Indeed, ground lidar measurements in Tajikistan reported significantly lower extinction-to-backscatter or lidar ratios (23 sr at 532 nm) for salt dust compared to typical dust aerosols (44±9 sr) (Hofer et al., 2017).

Even within the desiccated Aral basin, there is substantial variability in the physiochemical properties of erodible sediments. As the Aral Sea continued shrinking, sediment grain size became progressively smaller towards the basin's lowest point, while evaporate minerals began to precipitate in a typical sequence of calcite, gypsum/anhydrite, halite, and finally, potassium and magnesium salts. Consequently, distinct spatial gradients in mineralogy, grain size, and soil texture have been observed in the exposed sediments. For instance, Jiang et al. (2021) reported increasing abundances of clay and evaporites minerals but decreasing abundance of carbonates and organic carbon, when moving from the older coastline towards the newly exposed seabed. Argaman et al. (2006) found that the highly erodible takyrs soils dominate the outer rim of Aralkum, whereas the newly formed solonchak soils are more likely protected by salt crusts, which created stable, coarse aggregates that are resistant to wind erosion.

Assessing the impacts of Aralkum-generated saline dust is greatly hindered by lack of in situ measurements of the physical and chemical properties of the deflatable sediments (e.g., soil texture, mineralogical composition, crusting), and airborne dust particles (e.g., particle size distribution, shape, nonsphericity, chemical composition, mixing state, solubility). In particular, the global network of ground-based sun/sky photometers, AERosol RObotic NETwork (AERONET), has no operating sites

Deleted: <object><object><object>

Deleted: potential of multi-sensor approaches for ... [57]

Formatted ... [56]

Formatted ... [58]

Deleted: biases and limitations

Deleted: -----Section Break (Next Page)----- ... [59]

Formatted ... [60]

Deleted: Following

Deleted: drying

Deleted: over the past several decades

Deleted: newly formed

Deleted: emerged as a major source

Formatted ... [61]

Formatted ... [62]

Formatted ... [63]

Formatted ... [64]

Formatted ... [65]

Deleted: saline

Deleted: aerosol, causing a range of

Deleted: the

Formatted ... [66]

Formatted ... [67]

Formatted ... [68]

Deleted: ¶ ... [69]

Deleted: vast drylands

Deleted: due to...ecause of its anthropogenic origin ... [72]

Formatted ... [70]

Formatted ... [71]

Deleted: composition

Deleted: the erodible

Formatted ... [73]

Formatted ... [74]

Formatted ... [75]

Deleted: The

Formatted ... [76]

Formatted ... [77]

Deleted: adjacent sandy deserts (such as Karakum ... [78]

Deleted: 25 Aralkum contained significantly higher ... [79]

Deleted: aerosol originating from Aralkum Desert is ... [81]

Deleted: desert

Deleted: (Rudich et al., 2002).

Formatted ... [80]

Formatted ... [82]

Formatted ... [83]

Deleted: Aralkum Desert

Deleted: intra-basin

Formatted ... [84]

Deleted: the dust-producing

Formatted ... [85]


Formatted ... [86]

Formatted ... [87]

Deleted: the

Deleted: of the basin

Formatted ... [88]



**Figure 1.** Aggregated spectral abundance of four mineral groups based on the Earth Surface Mineral Dust Source Investigation (EMIT) L3 product: (a) Clays (including chlorite, illite, muscovite, kaolinite, montmorillonite and vermiculite); (b) Carbonates (calcite and dolomite); (c) Iron Oxides (goethite and hematite); and (d) Gypsum. Boxed regions indicate the Aralkum Desert. Note that panel (d) uses a different color scale from others.

near Aralkum, presumably due to the region's remoteness. The two operational AERONET sites in Central Asia—Issyk-Kul and Dushanbe—are located too far away to provide representative measurements of the saline dust from Aralkum (Semenov et al., 2005; Rupakheti et al., 2020). AERONET serves two important purposes. First, it provides the climatological aerosol information needed to specify season- and region-dependent aerosol optical models used in many satellite retrieval algorithms (Dubovik et al., 2002). Second, AERONET data are used as benchmark for validating satellite retrievals. Consequently, the a priori assumptions about aerosol microphysical properties in satellite algorithms may not accurately capture the unique dust properties over Aralkum. The lack of ground truth also prevents an evaluation of satellite product performance for the region. In general, satellite algorithms are optimized for global performance, but may exhibit significant local biases when the local prevailing aerosol conditions deviate substantially from the algorithm assumptions on aerosol properties.

Large-scale dust monitoring has relied on satellite retrievals over ocean basins frequently affected by continental dust outflow. For dust sources located deep within continental interiors such as Central Asia, observing airborne dust from space is more challenging due to difficulty in separating the surface contribution from the top-of-the-atmosphere (TOA) radiance measurements. Central Asia comprises a range of dust sources with varying sediment abundance and erodibility, including sandy and hilly deserts, desert steppes, salt flats, and ephemeral or dry lakes (Xi and Sokolik, 2015a, b). These regions feature highly heterogeneous dynamic surface properties and conditions (e.g., reflectivity, emissivity, land/water boundary, water turbidity), posing a major challenge to isolate aerosol signals from surface contributions. Currently, a number of techniques have been

- Deleted: <object><object><object>
- Formatted ... [116]
- Formatted ... [117]
- Deleted: The boxed region indicates
- Formatted ... [119]
- Formatted ... [120]
- Formatted ... [118]
- Deleted: Characterization
- Deleted: large
- Deleted: variability...onitoring has primarily ...el/ ... [121]
- Deleted: the tropical Atlantic, North Pacific, and ... [123]
- Deleted: ),
- Deleted: a
- Formatted ... [122]
- Formatted ... [124]
- Formatted ... [125]
- Deleted: task
- Formatted ... [126]
- Formatted ... [127]
- Deleted: the
- Deleted: of
- Deleted: influence of land surfaces
- Deleted: measurements.
- Formatted ... [128]
- Formatted ... [129]
- Formatted ... [130]
- Deleted: ¶ ... [132]
- Deleted: diversity
- Formatted ... [131]
- Formatted ... [133]
- Deleted: water bodies (Xi and Sokolik, 2015a, b). The
- Formatted ... [134]
- Deleted: and bright surfaces pose
- Deleted: for retrieving the aerosol properties in th ... [136]
- Deleted: satellite
- Deleted: and products
- Formatted ... [135]
- Formatted ... [137]
- Formatted ... [138]
- Formatted ... [139]
- Deleted: <object>





### 2.1.1 OMPS


### 2.1.2 TROPOMI

There are two separate UVAI products from TROPOMI, one developed by [ESA](#) ([European Space Agency](#)), and the other developed by [NASA](#). The NASA version [uses](#) the TropOMAER algorithm [which](#) reports both LER- and Mie-based UVAI at 354/388 nm ([Torres, 2021](#)). The ESA version reports LER-based UVAI at three wavelength pairs: 354/388, 340/380, and 335/367 nm ([Stein Zweers, 2022](#)). The first two pairs were selected to continue the multidecadal heritage UVAI records, while the 335/367 nm pair was added to ensure compatibility with the future UVAI algorithm planned for the Sentinel-5 mission.

### 2.1.3 EPIC

EPIC is an imaging spectroradiometer, onboard the Deep Space Climate Observatory (DSCOVR) spacecraft, which operates in a Lissajous orbit about Lagrange-1 point in the Earth-Sun system. This allows EPIC to view the sunlit disk of Earth every 60–100 minutes. EPIC measures reflected radiances in 10 channels from UV to near-infrared (NIR), with a ground resolution of 8 km at 443 nm and 16 km in other bands. The EPIC EPICARUV algorithm reports both LER- and Mie-based UVAI at the 340/388 nm wavelengths [Torres, 2019b].

Deleted: <span>&lt;object&gt;&lt;object&gt;&lt;object&gt;</span>	
Formatted	[204]
Deleted: <span>¶</span>	[205]
Formatted	[206]
Moved down [2]: 130 <span>¶</span>	
Moved down [3]: 135 <span>¶</span>	
Moved down [4]: 140 <span>¶</span>	
Moved down [5]: 150 <span>¶</span>	[217]
Deleted: aerosol, and volcanic ash (Hsu, 1996; Herr	[213]
Deleted: <span>¶</span>	[214]
Deleted: <span>¶</span>	[215]
Deleted: <span>¶</span>	[216]
Deleted: .....Column Break.....	[218]
Deleted: spacecraft	
Formatted	[219]
Formatted	[220]
Deleted: every 0.42 nm	
Formatted	[221]
Deleted: size	
Formatted	[222]
Formatted	[207]
Formatted	[208]
Deleted: (Torres, 2019a).	
Formatted	[223]
Formatted	[209]
Deleted: deployed on	
Formatted	[224]
Formatted	[225]
Deleted: the ...NPP spacecraft	[226]
Formatted	[227]
Deleted: (Veeffkind et al., 2012).	
Deleted: in	
Deleted: -visible (270–500 nm), NIR (710–770 nm)	[230]
Formatted	[228]
Formatted	[229]
Formatted	[210]
Deleted: Royal Netherlands Meteorological Institut	[231]
Deleted: (ESA	
Formatted	[232]
Formatted	[233]
Deleted: the National Aeronautics and Space	[234]
Deleted: is based on	
Formatted	[235]
Formatted	[236]
Deleted: aerosol ...lgorithm, and	[237]
Formatted	[238]
Formatted	[211]
Deleted: 354/388...35/367 nm wavelengths (Torres	[239]
Formatted	[240]
Deleted: and measures the Earth’s reflected solar	[242]
Formatted	[241]
Formatted	[243]
Formatted	[212]
Deleted: which	



## 2.2 Midvisible AOD

<https://doi.org/10.5194/egusphere-2024-3416>

Inferring column-integrated aerosol properties from reflected visible-NIR observations is an ill-posed inverse problem due to limited information content to fully characterize the atmosphere-surface system. Midvisible AOD is generally the most readily-

parameter, and consequently algorithms must rely on a priori knowledge or assumptions of the aerosol size distribution, shape, and refractive index (known as aerosol optical models). A common strategy involves using radiative transfer codes to create look-up tables (LUTs) of TOA radiances for a set of predefined aerosol optical models, surface types, and solar/viewing geometries (e.g., Remer et al., 2013a). These LUTs are then used to match with measured TOA reflectances to derive optimal estimates of AOD and, in some cases, additional aerosol properties related to particle size and absorption. Aerosol retrieval using this strategy is highly sensitive to the aerosol optical models that are predefined in the algorithm and selected during the retrieval. Although these aerosol models are designed to represent season- and location-specific aerosol conditions as realistically as possible, satellite algorithms may still incorporate inconsistent or even contradictory models, especially for regions like Central Asia due to lack of observational constraints. In addition, satellite algorithms may fail to select the appropriate model during the retrieval, leading to mismatch between the assumed aerosol properties and the actual aerosol conditions. These factors can contribute to inconsistencies between retrieved AODs, including those from the same instrument.

We compare multiple AOD products from three polar-orbiting instruments onboard NASA's EOS and NOAA's JPSS satellites, including the Moderate Resolution Imaging Spectroradiometer (MODIS), Multi-angle Imaging Spectroradiometer (MISR), and Visible Infrared Imaging Radiometer Suite (VIIRS). MODIS and VIIRS employ three single-view aerosol algorithms—Dark Target (DT), Deep Blue (DB), and Enterprise Processing System (EPS)—and the time series-based Multi-Angle Implementation of Atmospheric Correction (MAIAC) algorithm. MISR utilizes two algorithms: an operational standard algorithm for generating routine global aerosol records, and a research algorithm designed to explore alternative retrieval methods and additional aerosol properties on a case-by-case basis.

Below, we briefly describe the MODIS, VIIRS, and MISR aerosol algorithms, focusing on the a priori assumptions about aerosol optical models and surface reflectances. Other algorithm components (e.g., screening of clouds/sun glint/snow, spectral fitting) are not discussed.

### 2.2.1 MODIS and VIIRS DT over-water algorithms

The DT algorithm includes different approaches for over-land and over-water retrievals. We focus on the over-water algorithm which is well suited for retrieving dust properties over the Caspian Sea. DT exploits the contrast of reflective aerosol layers against a dark background in the visible spectrum. Over water, the algorithm searches the LUT for the best fit of measured TOA reflectances at seven window bands from visible to SWIR, and obtains solutions for both AOD and fine mode fraction (FMF) (Tanré et al., 1997). The ambient aerosol scene is represented as a linear combination of one fine and one coarse mode weighted by FMF. DT considers nine aerosol optical models over water: four fine aerosol models, three sea salt models, and two coarse dust models, which are derived from long-term AERONET measurements near water bodies (Remer et al., 2005).

Deleted: <object><object><object>

Formatted ... [246]

Deleted: AOD...olumn-integrated aerosol properties ... [257]

Deleted: and

Formatted ... [258]

Moved (insertion) [2]

Formatted ... [247]

Formatted ... [259]

Deleted: on

Formatted ... [260]

Deleted: microphysical ... [261]

Moved up [7]: 160 ... [262]

Formatted ... [263]

Moved down [9]: 170 ... [264]

Deleted: ..... Column Break ..... [264]

Formatted ... [265]

Deleted: composition (or

Formatted ... [266]

Deleted: )

Deleted: general

Deleted: of AOD retrieval from passive, downward ... [269]

Deleted: a forward

Formatted ... [267]

Formatted ... [268]

Formatted ... [270]

Formatted ... [271]

Deleted: model...odes to compute the ...reate look- ... [272]

Deleted: expected to be encountered by the instrum ... [274]

Formatted ... [273]

Formatted ... [275]

Formatted ... [248]

Deleted: find the best fit ...rive optimal estimates ... [276]

Formatted ... [277]

Moved (insertion) [3]

Formatted ... [249]

Deleted: method

Deleted: predefined aerosol models, which aim...er ... [279]

Formatted ... [278]

Deleted: , sometimes conflicting, assumptions of th ... [281]

Formatted ... [280]

Formatted ... [250]

Moved (insertion) [4]

Formatted ... [282]

Deleted: Below, we briefly review the AOD algo ... [283]

Deleted: )

Formatted ... [284]

Formatted ... [251]

Deleted: ) sensors. The

Formatted ... [285]

Formatted ... [286]

Formatted ... [252]

Deleted: Multi-Angle Implementation of Atmo- sp ... [287]

Formatted ... [288]

Deleted: ) algorithms are suitable for observing du ... [289]

Due to lack of non-spherical dust models, the algorithm was known to yield biased retrievals over dusty oceanic scenes (Zhou et al., 2024). DT aerosol retrieval is performed at a nominal resolution of 10×10 km<sup>2</sup>. A higher-resolution product (DT3K) was later introduced at 3×3 km<sup>2</sup> to capture small-scale aerosol features (Remer et al., 2013b). DT3K employs the same technique as DT, but using different pixel aggregation and quality assurance (QA) rules. The DT algorithm has been ported to VIIRS, which retrieves AOD at a nominal resolution of 6×6 km<sup>2</sup> (Sawyer et al., 2020).

**2.2.2 MODIS and VIIRS DB algorithms**

Inspired by the aerosol detection capability in the near-UV, the DB algorithm employs the 0.41 μm or “deep blue” band, which has lower and more homogeneous surface reflectivity than the longer visible wavelengths (Hsu et al., 2004). The aerosol column is represented by a spheroidal dust model and a spherical fine-dominated anthropogenic model, which employ various SSA values depending on locations and seasons. The surface reflectances are determined based on a pre-calculated database over bright surfaces (e.g., deserts, urban areas), and empirical relationships between visible and SWIR bands over vegetated surfaces (Hsu et al., 2013).

DB was initially implemented to fill the data gap over bright surfaces in the MODIS aerosol product. In the VIIRS product suite, DB has been expanded to all cloud-, snow- and ice-free land surfaces, and also performs retrieval over water using the Satellite Ocean Aerosol Retrieval (SOAR) algorithm (Hsu et al., 2019; Lee et al., 2024). SOAR considers four aerosol optical models: maritime, dust, fine-dominated, and mixed, each represented by a bi-modal distribution consisting of one fine and one coarse mode (Sayer et al., 2018). The dust model consists of one spherical fine mode and one spheroidal coarse mode, derived from AERONET measurements at Cape Verde (Lee et al., 2017).

**2.2.3 VIIRS EPS algorithm**

The EPS algorithm was developed for NOAA’s next-generation polar-orbiting and geostationary environmental satellites. Here we focus on the VIIRS EPS aerosol product described in Laszlo (2018) and more recently Laszlo and Liu (2022). Over water, EPS is based on the MODIS heritage and represents the aerosol column as a linear combination of one fine and one coarse mode weighted by FMF, selected from four fine mode and five coarse mode aerosol models same as the MODIS DT algorithm (Remer et al., 2006). The algorithm searches for the AOD and FMF that give the best match between observed and pre-calculated TOA reflectances at seven VIIRS channels (Jackson et al., 2013; Laszlo and Liu, 2022).

Over land, EPS simultaneously retrieves AOD, aerosol optical model, and Lambertian surface reflectances in selected bands, by matching the observed and calculated TOA reflectance over both dark and bright (snow-free) surfaces. EPS considers four candidate aerosol optical models, of which three are spherical, fine-mode dominated aerosols (labeled as generic, urban, and smoke), and one is non-spherical, coarse-mode dominated aerosol (labeled as dust). The dust model is forcibly used for North Africa and the Arabian Peninsula to account for the dominant dust presence in these regions. The candidate aerosol models are



- Deleted: <object><object><object>
- Moved (insertion) [8]
- Formatted ... [312]
- Formatted ... [313]
- Formatted ... [314]
- Formatted ... [315]
- Moved down [14]: 225¶
- Deleted: ..... Column Break ..... [322]
- Formatted ... [323]
- Deleted: and thus provides sufficient contrast betwe ... [324]
- Moved (insertion) [9]
- Deleted: surface reflectance
- Deleted: the
- Deleted: (similar to the DT approach) as a function ... [327]
- Formatted ... [325]
- Formatted ... [326]
- Deleted: Over ocean, the
- Deleted: algorithm began
- Deleted: generate AOD retrieval from VIIRS based ... [331]
- Formatted ... [329]
- Formatted ... [330]
- Formatted ... [328]
- Formatted ... [332]
- Deleted: (Sayer et al., 2018).
- Formatted ... [333]
- Deleted: over water
- Formatted ... [334]
- Deleted: as
- Formatted ... [335]
- Deleted: modes.
- Formatted ... [336]
- Deleted: which currently is
- Formatted ... [337]
- Deleted: (Lee et al., 2017).
- Moved down [11]: ¶
- Moved down [12]: ¶
- Moved down [13]: 220¶
- Deleted: Over land, the DT algorithm uses three ba ... [316]
- Formatted ... [317]
- Deleted: ¶ ... [318]
- Formatted ... [319]
- Deleted: ¶ ... [320]
- Deleted: ¶ ... [321]
- Formatted ... [339]
- Formatted ... [340]
- Formatted ... [338]
- Moved (insertion) [10]
- Deleted: a suite of aerosol products from the new
- Deleted: (e.g., VIIRS)
- Deleted: (e.g., Advanced Baseline Imager) sensors ... [347]
- Formatted ... [344]
- Formatted ... [345]
- Formatted ... [346]
- Formatted ... [343]

195

200

205

210

215

220

adopted from the MODIS Collection 5 DT algorithm (Remer et al., 2006). Over bright desert surfaces, the surface reflectance is estimated from a state analysis of spectral reflectance ratios between VIIRS channels (Zhang et al., 2016).  
Preprint. Discussion started: 2 December 2024  
© Author(s) 2024. CC BY 4.0 License.


1.1.1 MODIS MAIAC algorithm

Unlike the single-view approach adopted by DT, DB and EPS algorithms, MAIAC uses a sliding window to accumulate up to 16 days of multi-angle observations from different orbits for the same location to retrieve bidirectional surface reflectance over land (including deserts) simultaneously with aerosol properties (Lyapustin et al., 2018). MAIAC uses a dynamic minimum reflectance method to define the surface reflectance, spectral ratios for each 1 km grid cell, which allows AOD retrieval over both dark and bright surfaces. The MODIS Collection 6 MAIAC algorithm uses nine aerosol optical models derived from AERONET climatology to represent the regional background aerosol conditions. For Central Asia, the background aerosol model ("Model 2") is derived from AERONET measurements over the western U.S., which represents a mixture of dust and fine mode aerosols. During the retrieval, a smoke/dust test is first applied to determine whether the background or dust model should be used. If dust is detected, the algorithm uses a spheroidal dust model ("Model 6") derived from AERONET measurements from the Solar Village site in Saudi Arabia (Dubovik et al., 2006).

2.2.5 MISR AOD and particle property algorithms

MISR measures reflected sunlight using nine push-broom cameras with view angles of  $\pm 70.5^\circ$ ,  $\pm 60.0^\circ$ ,  $\pm 45.6^\circ$ ,  $\pm 26.1^\circ$ , and  $0^\circ$  along-track, each in four spectral bands (446, 558, 672, and 866 nm) across a 380 km swath at pixel-resolution between 275 m and 1.1 km, depending on the channel (Diner et al., 1998). The MISR Standard Aerosol retrieval algorithm produces AOD and constraints on column-effective particle type operationally (Garay et al., 2020). The aerosol column is represented by 74 aerosol optical models as mixtures of single-composition components, including 50 mixtures of spherical components, 20 mixtures of spherical and dust components, and 4 mixtures of dust components. The algorithm accounts for the contribution of surface bidirectional reflectance factor (BRF) based on a principal component analysis of TOA radiances (Martonchik et al., 2009).

The MISR Research Aerosol (RA) retrieval algorithm is optimized to provide constraints on particle size, sphericity, and light-absorption under favorable observing conditions on a case-by-case basis (Limbacher et al., 2022). The algorithm considers a broader range of aerosol mixtures to allow more subtle particle property distinction. For the surface reflectance, two different approaches are considered: (a) surface BRF retrieved self-consistently with the atmosphere using only MISR data (similar to the standard algorithm), and (b) surface BRF contribution prescribed from the MODIS MAIAC product to separate the surface contribution. In addition, a XGBoost AI/ML approach has been developed using the retrieved and prescribed RA results, along with an aerosol microphysical property validation dataset developed by Anstett et al. (in prep). The XGBoost models employed here use MISR RA prescribed + retrieved geophysical output as input, training models separately against AERONET AOD, FMF, Ångström Exponent, non-sphericity, and SSA. The training dataset consists of ~50,000 global MISR/AERONET over-land coincidences, with each of the coincidences containing potentially >2,000 MISR pixels (retrievals). For the purposes of



Deleted: <object><object><object>

Deleted: MODIS

Deleted: (Remer et al., 2006; Levy et al., 2007).

Formatted ... [372]

Formatted ... [373]

Formatted ... [371]

Formatted ... [374]

Moved (insertion) [11]

Formatted ... [375]

Formatted ... [376]

Formatted ... [383]

Deleted: retrievals

Formatted ... [384]

Formatted ... [377]

Deleted: at a high spatial resolution (1×1 km<sup>2</sup>).

Deleted: also uses a sliding window technique to ac ... [386]

Formatted ... [385]

Formatted ... [378]

Deleted: USA, and

Formatted ... [387]

Formatted ... [388]

Deleted: in the retrieval. Once... If dust is detected ... [389]

Formatted ... [379]

Deleted: Section Break (Continuous) ... [380]

Moved down [17]: ¶

Deleted: ¶ ... [382]

Formatted ... [381]

Deleted: Algorithm implementation in MODIS

Deleted: VIIRS ¶ ... [392]

Deleted: perform retrieval at a nominal reso ... [394]

Formatted ... [391]

Formatted ... [393]

Formatted ... [390]

Moved (insertion) [12]

Formatted ... [395]

Formatted ... [396]

Formatted ... [397]

Formatted ... [398]

Formatted ... [399]

Moved (insertion) [13]


Formatted ... [400]

Deleted: <object>



225

developing optimal AI/ML coefficients, each of these quality-assessed pixels was treated as an independent data point, yielding <https://doi.org/10.5194/amt-2024-424> used for training. Preprint. Discussion started: 2 December 2024



EGU  
Open Access  
Preprint repository

230

Compared to visible and NIR-based techniques, aerosol retrieval in the thermal-infrared (TIR) offers several advantages, such as nighttime observation and enhanced contrast of dust signals over deserts. In addition, the infrared spectrum is primarily sensitive to coarse particles (diameter > 1 μm), providing better dust detection. Split-window techniques have been used to detect dust from passive radiometers by exploiting the distinct negative brightness temperature difference between two neighboring atmospheric window channels (e.g., 10.8 and 12 μm) (Legrand et al., 2001; Lensky and Rosenfeld, 2008). Based on this approach, the Dust RGB composite derived from the Spinning Enhanced Visible and InfraRed Imager (SEVIRI) is capable of tracking dust plumes every 15 minutes and mapping the source locations over Central Asia, as demonstrated in Xi (2023).

235

Hyperspectral infrared spectrometers have offered new potential to infer dust properties, including AOD<sub>10</sub> and dust layer height. A major challenge in TIR-based retrieval is isolating the aerosol signal from the infrared emissions of the atmosphere and the underlying surface. TIR-based algorithms rely on forward model simulations of TOA radiances or brightness temperatures for a range of atmospheric profiles (e.g., temperature, water vapor), surface properties (e.g., emissivity, temperature), and dust aerosol properties (e.g., AOD<sub>10</sub>, vertical distribution, size, and refractive index). Below, we describe four different algorithms used for retrieving AOD<sub>10</sub> from the Infrared Atmospheric Sounding Interferometer (IASI), a Fourier transform spectrometer onboard the European MetOp satellite series that observes the Earth's radiation spectra from 645 to 2760 cm<sup>-1</sup> (3.6–15.5 μm) at a spectral resolution of 0.5 cm<sup>-1</sup> and a nadir ground resolution of 12 km. The four algorithms differ in the retrieval methods, the atmosphere and surface input data, and the dust optical models assumed in forward model calculations, among other factors.

240

### 2.3.1 LMD algorithm

245

The Laboratoire de Météorologie Dynamique (LMD) algorithm employs a two-step, LUT-based approach (Capelle et al., 2018). In the first step, the atmospheric state is determined using 18 IASI channels that are insensitive to the surface or aerosols. In the second step, AOD<sub>10</sub>, mean altitude, and surface temperature are retrieved by fitting observed brightness temperatures against simulations at 8 aerosol-sensitive channels. These simulations incorporate a range of atmospheric profiles, surface properties (emissivity, temperature, and pressure), AOD<sub>10</sub>, altitudes, and dust microphysical properties. Dust is represented by a spherical, lognormal size distribution ( $R_{eff}=2.3\text{ }\mu\text{m}$ ,  $\sigma_g=0.65$ ) and two refractive indices from Balkanski et al. (2007) and Volz (1973), corresponding to strongly and weakly absorbing dust types, respectively. Surface emissivity is based on the monthly mean IASI retrievals by Capelle et al. (2012).

Deleted: <object><object><object>

Moved (insertion) [14]

Formatted: Space Before: 5.5 pt

Moved (insertion) [15]

Formatted: Font: Arial

Formatted: Space Before: 5.85 pt

Formatted: Space Before: 5.5 pt

Moved (insertion) [16]

Moved (insertion) [17]

Formatted: Font: Arial, Not Bold

Formatted: Space Before: 5.7 pt

Deleted: ~~~~~Section Break (Next Page)~~~~~

... [401]

Deleted: <object>

9

### 2.3.2 MAPIR algorithm

<https://doi.org/10.5194/egusphere-2024-3416>

Preprint Discussion started 2 December 2024

© Author(s) 2024. CC BY 4.0 License.

The Mineral Aerosol Retrieval Scheme (MAPIR) algorithm retrieves the vertical profiles of dust concentrations using the Rodgers optimal estimation method (Callewaert et al., 2019). The dust concentration profiles are converted to AOD<sub>10</sub> using Mie-calculated extinction cross section, based on an assumed size distribution and refractive index. A forward model (RTTOV) is used to simulate IASI radiances based on a priori of the state vector (i.e., dust concentration profiles at seven 1-km-thick layers centered at 0.5 to 6.5 km), along with temperature and water vapor profiles derived from IASI. The dust optical model assumes a lognormal size distribution ( $r_g=0.6\ \mu\text{m}$ ,  $\sigma_g=2$ ), and refractive index from Volz (1973). Surface emissivity is based on a monthly climatology derived from IASI clear sky spectra by Zhou et al. (2011).

### 2.3.3 ULB algorithm

The Université libre de Bruxelles (ULB) algorithm estimates dust AOD<sub>10</sub> using a neural network approach (Clarisse et al., 2019). The algorithm first performs dust detection by computing a dust index from a linear discrimination analysis of IASI-observed spectra. The dust index is then converted to dust AOD<sub>10</sub> via a neural network trained with synthetic spectra generated by a forward model. The forward model incorporates a representative set of atmospheric states from IASI Level-2 data, surface emissivity from Zhou et al. (2011), and a range of dust layer altitudes (from 0 to 7 km). Dust aerosol is represented by a lognormal size distribution ( $r_g=0.5\ \mu\text{m}$ ,  $\sigma_g=2$ ), and refractive index from Volz (1973).

### 2.3.4 IMARS algorithm

The Infrared Mineral Aerosol Retrieval Scheme (IMARS) algorithm performs probabilistic estimates of AOD<sub>10</sub>, composition, effective radius, and mean layer temperature (a proxy for dust layer height), based on forward model simulations of IASI-observed radiances under various dust and ice cloud conditions (Offenwanger et al., 2024). The algorithm considers 12 possible combinations of four dust mineralogical mixtures (China, Central Sahara, Niger, Iowa Loess) and three size distributions. Each dust mixture is associated with predefined mineral fractions (and hence refractive indices) and particle effective radius. Surface emissivity is based on the MODIS UCSB Emissivity Library.

### 2.4 ALH

Aerosol vertical distribution can be retrieved from both active and passive sensors. Active sensors, such as the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) lidar, provide vertically-resolved retrievals of aerosol volume extinction at high vertical resolutions. However, they are limited by narrow swaths and poor spatial coverage. In recent years, substantial progress has been made in retrieving ALH from passive sensors using various techniques, such as stereoscopic retrieval from polar-orbiting multiangle or geostationary imagers (Nelson et al., 2013; Carr et al., 2020), polarimetric observations in the near UV (Wu et al., 2016), differential optical absorption spectroscopy in oxygen absorption bands (Xu et al., 2017, 2019), and hyperspectral infrared measurements (Capelle et al., 2018; Callewaert et al., 2019; Clarisse et al., 2019; Offenwanger et al., 2024). Although passive techniques do not achieve the same level of accuracy as lidars and only estimate an effective height,

Deleted: <object><object><object>

Deleted: distribution affects the atmospheric lifetime ... [413]

Formatted ... [412]

Formatted ... [414]

Formatted ... [402]

Formatted ... [403]

Formatted ... [404]

Formatted ... [405]

Formatted ... [406]

Deleted: ..... Column Break ..... [407]

Deleted: 0.75×0.75 km<sup>2</sup> (Laszlo and Liu, 2022). [408]

Formatted ... [409]

Formatted ... [410]

Moved down [18]: 285 [409]

Deleted: [411]

Moved (insertion) [19]

Formatted ... [415]

Formatted ... [416]

Deleted: from AOD (Li et al., 2015; Sun et al., 2016). [418]

Formatted ... [417]

Formatted ... [419]

Deleted: spaceborne lidars onboard the ...loud-A ... [420]

Formatted ... [421]

Deleted: (Winker et al., 2009; Yorks et al., 2016). [422]

Deleted: swath

Deleted: there has been

Formatted ... [423]

Formatted ... [424]

Formatted ... [425]

Deleted: one piece information of aerosol vertical ... [428]

Deleted: based on

Deleted: including

Formatted ... [427]

Formatted ... [429]

Formatted ... [430]

Formatted ... [431]

Formatted ... [426]

Deleted:

Deleted: (Nelson et al., 2013; Carr et al., 2020),

Formatted ... [432]

Formatted ... [433]

Deleted: (Wu et al., 2016),

Deleted: (DOAS) in the NIR (e.g., ...n oxygen A ... [434]

Deleted: thermal ...nfrared (TIR) measurements ... [435]

Deleted: with limited information on the aerosol ... [437]

Formatted ... [436]

Deleted: <object>

with limited information on the aerosol layer thickness, they provide much better spatial coverage and revisit frequency (Lu et al. 2023). <https://doi.org/10.5194/egusphere-2024-3416>  
Preprint. Discussion started: 2 December 2024  
The ALH definition varies by retrieval techniques, and may refer to the top of aerosol layers such as from the ground or an effective central height corresponding to peak aerosol extinction (Xu et al., 2019). Like AOD, ALH is an optical quantity and varies with the retrieval wavelength. Except for stereoscopic techniques, passive ALH retrievals depend on assumptions about the aerosol vertical distribution. In this study, we compare four ALH products including the CALIOP aerosol extinction-weighted height, EPIC aerosol optical centroid height (AOCH) product, IASI mean dust layer altitude, and MISR plume height.

2.4.1 CALIOP aerosol extinction-weighted height

CALIOP was a nadir-viewing elastic backscatter lidar onboard the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO), and measures polarized backscatter at 532 nm and total attenuated backscatter at 532 nm and 1032 nm, with a vertical resolution of 30 m below 8.2 km and 60 m between 8.2 and 20.2 km (Winker et al., 2009). In the CALIOP data processing, calibrated attenuated backscatter coefficient profiles are used to detect the top and base altitudes of atmospheric features. A set of scene classification algorithms (SCA) then classifies these features as either aerosol or cloud, and determine the aerosol type and cloud phase. During this process, aerosol lidar ratios are selected to derive the aerosol extinction and backscatter coefficient profiles (Young et al., 2018). The selection of lidar ratios is based on the aerosol typing algorithm which considers six tropospheric aerosol types: clean marine, dusty marine, dust, polluted continental/smoke, polluted dust, and elevated smoke (Kim et al., 2018). Based on the aerosol extinction profile, the ALH with respect to the mean sea level can be calculated as 
$$\frac{\sum_{i=1}^n \beta_{ext,i} Z_i}{\sum_{i=1}^n \beta_{ext,i}}$$
 where  $\beta_{ext,i}$  is the 532 or 1064 nm aerosol extinction coefficient ( $\text{km}^{-1}$ ) at level  $i$ , and  $Z_i$  is the altitude (km) at level  $i$  (Koffi et al., 2012).

#### 2.4.1 CALIOP aerosol extinction-weighted height

2.4.2 EPIC AOCH product

The EPIC AOCH product is derived from the spectral contrast in TOA reflectances between the oxygen ( $\text{O}_2$ ) absorption and continuum bands. The physical principle is that ALH affects the path length of backscattered light and the amount of light absorbed by well-mixed  $\text{O}_2$  molecules. Consequently, the spectral contrast between the  $\text{O}_2$  absorption bands and the continuum bands depends on the ALH (Xu et al., 2019). The EPIC AOCH algorithm employs pre-computed TOA reflectances for a range of AOD<sub>0.68</sub>, AOCH, surface reflectivity, and surface pressure. Three aerosol optical models are considered: smoke, Saharan dust, and Asian dust. The dust models are derived from AERONET measurements at Cape Verde and over East Asia (Xu et al., 2017). The aerosol vertical distribution assumes a quasi-Gaussian profile characterized by a centroid altitude and a fixed half-width at half maxima of 1 km. Hence, the EPIC-retrieved AOCH represents the altitude of peak volume extinction.

#### 2.4.2 EPIC AOCH product

The EPIC AOCH product is derived from the spectral contrast in TOA reflectances between the oxygen ( $\text{O}_2$ ) absorption and continuum bands. The physical principle is that ALH affects the path length of backscattered light and the amount of light absorbed by well-mixed  $\text{O}_2$  molecules. Consequently, the spectral contrast between the  $\text{O}_2$  absorption bands and the continuum bands depends on the ALH (Xu et al., 2019). The EPIC AOCH algorithm employs pre-computed TOA reflectances for a range of AOD<sub>0.68</sub>, AOCH, surface reflectivity, and surface pressure. Three aerosol optical models are considered: smoke, Saharan dust, and Asian dust. The dust models are derived from AERONET measurements at Cape Verde and over East Asia (Xu et al., 2017). The aerosol vertical distribution assumes a quasi-Gaussian profile characterized by a centroid altitude and a fixed half-width at half maxima of 1 km. Hence, the EPIC-retrieved AOCH represents the altitude of peak volume extinction.

- Deleted: <object><object><object>
- Moved (insertion) [18]
- Formatted ... [438]
- Formatted ... [445]
- Deleted: generally capable of penetrating deeper in ... [446]
- Formatted ... [447]
- Deleted: layer thickness or profile. Below
- Deleted: describe two aerosol layer height
- Deleted: with coverage over Central Asia
- Formatted ... [448]
- Formatted ... [449]
- Formatted ... [450]
- Deleted: The definition of aerosol layer height vary ... [439]
- Deleted: derived from CALIPSO, and the EPIC
- Formatted ... [451]
- Formatted ... [440]
- Formatted ... [452]
- Deleted: <object>As part of the A-train constellation ... [453]
- Deleted: ,
- Deleted: with Orthogonal Polarization (CALIOP), which
- Formatted ... [454]
- Formatted ... [455]
- Formatted ... [441]
- Formatted ... [456]
- Deleted: an altitude of
- Formatted ... [457]
- Deleted: (Winker et al., 2009).
- Deleted: level-2
- Formatted ... [458]
- Formatted ... [459]
- Deleted: the
- Formatted ... [460]
- Deleted: using a selective iterated boundary location ... [461]
- Deleted: algo- rithms
- Deleted: are used to classify
- Deleted: and
- Formatted ... [462]
- Formatted ... [463]
- Formatted ... [464]
- Formatted ... [465]
- Deleted: the aerosol extinction-to-backscatter ratios ... [466]
- Formatted ... [467]
- Formatted ... [442]
- Deleted: (Young et al., 2018). The lidar ratio
- Formatted ... [468]
- Deleted: relies
- Formatted ... [469]
- Deleted: ¶ ... [443]
- Deleted: : ...clean marine, dusty marine, dust, pollu ... [470]
- Deleted: ¶ ... [444]
- Deleted: , each assigned a default lidar ratio (Omar ... [472]
- Deleted: CALIOP AOCH
- Deleted: ¶ ... [475]
- Deleted: computed

#### 2.4.3 IASI dust altitude from the LMD algorithm

<https://doi.org/10.5194/egusphere-2024-3416>

Preprint. Discussion started: 2 December 2024

© Author(s) 2024. CC BY 4.0 License.

The dust layer height employed in this study employs the sensitivity of infrared radiation to the temperature of the aerosol layer, which is intrinsically linked to its mean altitude. The IASI LMD algorithm estimates the dust AOD<sub>10</sub> and mean altitude by matching IASI observations to forward model simulations assuming eight mean layer altitudes from 750 to 5795 m (Capelle et al., 2018). Dust is assumed to be uniformly distributed within a single atmospheric layer, of which the thickness varies from 500 to 800 m. The retrieved mean altitude represents the height at which half of the AOD<sub>10</sub> is below and half of the AOD<sub>10</sub> is above, and considered as an infrared optical equivalent to the centroid of the aerosol vertical profiles.

#### 2.4.4 MISR plume height

In addition to the radiometrically derived AOD, ALH can be derived geometrically from the parallax in the MISR hyper- stereo imagery. The MISR Interactive Explorer (MINX) software performs this task interactively, on a case-by-case basis, and retrieves plume heights at 1.1 km horizontal resolution, and between 250 and 500 m vertical resolution (Nelson et al., 2013).

The MISR ALH retrievals identify the layer of maximum spatial contrast in the plume imagery, so the results are often skewed lower than CALIPSO, which is also sensitive to thin, less distinct aerosol layers above dense plume features (Flower and Kahn, 2017).

### 3 Evaluation of Cross-Sensor and Cross-Algorithm Consistency

Table J summarizes the satellite products and parameters used in this study. These products were chosen based on the data availability and similar overpass times over Central Asia, ensuring direct comparisons with minimal scene differences. While some products have duplicated parameters (e.g., AOD is also reported in the UVAI products), we focus on the primary and most widely used aerosol parameters from each product. The UVAI and midvisible AOD products also report Ångström Exponent, and single scattering albedo (SSA) over land; however, they are subject to larger uncertainties than AOD, and generally not recommended for scientific studies.

#### 3.1 Horizontal and Vertical Dust Distributions

This section describes the horizontal and vertical dust distributions based on coincident passive and active satellite observations. On 27 May 2018, MODIS Aqua observed an extensive whitish dust plume originating from the Aralkum Desert and moving southeastward towards Iran and Afghanistan (Fig. 2(a)). While CALIOP missed the Aralkum dust plume, it detected a shallow dust layer at an altitude of 0–1 km at the southeast coast of Caspian Sea (Fig. 2(d)). This dust likely originated from the dry channel of the ancient River Uzboy in western Turkmenistan, which has been identified as a highly active dust source in Central Asia (Nobakht et al., 2021).

On 28 May 2018, a high-pressure system developed just south of the Aralkum Desert. As shown in the nighttime SEVIRI Dust RGB composite (Fig. 2(b)), the anticyclonic flow carried the lofted dust across Ustyurt Plateau toward the Caspian Sea,

Deleted: <object><object><object>

Moved (insertion) [20]

Formatted ... [518]

Formatted ... [519]

Formatted ... [520]

Moved (insertion) [21]

Formatted ... [521]

Formatted ... [525]

Formatted ... [522]

Formatted ... [527]

Formatted ... [526]

Deleted: sensor...ensor and Cross-algorithm ... [528]

Formatted ... [529]

Deleted: 1

Formatted ... [530]

Deleted: aerosol

Deleted: considered

Deleted: The

Deleted: are selected

Formatted ... [531]

Formatted ... [532]

Formatted ... [533]

Formatted ... [534]

Deleted: are mostly from polar-orbiting sensors (ex ... [535]

Deleted: time over Aralkum Desert. For the AOD ... [537]

Formatted ... [536]

Formatted ... [538]

Moved (insertion) [22]

Formatted ... [523]

Deleted: of the considered

Deleted: available from

Deleted: NASA

Formatted ... [539]

Formatted ... [540]

Formatted ... [541]

Deleted: parameter

Deleted: parameters related to aerosol size (e.g., ... [543]

Deleted: )...and absorption (i.e. ...ingle scattering ... [545]

Formatted ... [542]

Formatted ... [544]

Formatted ... [524]

Formatted ... [546]

Deleted: ¶ ... [548]

Deleted: nearly

Deleted: observations from

Deleted: sensors. Fig. 2 shows the daytime MODIS ... [552]

Formatted ... [547]

Formatted ... [549]

Formatted ... [550]

Formatted ... [551]

Formatted ... [553]

Deleted: off...riginating from the Aralkum Desert ... [554]

Deleted: .

Formatted ... [555]





**Table 1.** List of satellite aerosol products considered in this study.

Sensor	Algorithm/Product	Resolution	Data Reference
<b>UV Aerosol Index (UVAI)</b>			
OMPS/SNPP	NMMIEAI v2	50×50 km <sup>2</sup>	Torres (2019a)
EPIC/DSCOVER	AER v3	12×12 km <sup>2</sup>	Torres (2019b)
TROPOMI/SSP	TROPOMAER v1	7.5×3 km <sup>2</sup>	Torres (2021)
	AER_AI v2	7.5×3 km <sup>2</sup>	ESA (2021)
<b>Midvisible Aerosol Optical Depth (AOD)</b>			
MODIS/Terra, Aqua	DT C6.1	10×10 km <sup>2</sup>	Levy and Hsu (2015a)
	DT3K C6.1	3×3 km <sup>2</sup>	Levy and Hsu (2015b)
	DB C6.1	10×10 km <sup>2</sup>	Levy and Hsu (2015a)
	MAIAC C6.1	1×1 km <sup>2</sup>	Lyapustin and Wang (2022)
VIIRS/SNPP, NOAA20	DT v2	6×6 km <sup>2</sup>	Levy et al. (2023)
	DB v2	6×6 km <sup>2</sup>	Hsu (2022)
	EPS v3r0	0.75×0.75 km <sup>2</sup>	Kondragunta et al. (2023)
MISR/Terra	v23 (operational)	4.4×3 km <sup>2</sup>	ASDC (1999)
	Research Algorithm	1.1×1.1 km <sup>2</sup>	Limbacher et al. (2022)
<b>Thermal Infrared AOD (AOD<sub>10</sub>)</b>			
IASI/METOP-A	LMD v2.2	1°×1°	C3S CDS (2019)
	MAPIR v5.1	1°×1°	C3S CDS (2019)
	ULB v9	1°×1°	C3S CDS (2019)
	IMARS v7	1°×1°	C3S CDS (2019)
<b>Aerosol Laver Height (ALH)</b>			
CALIOP/CALIPSO	05kmAPro v4.51	5 km	ASDC (2023)
EPIC/DSCOVER	AOCH v1	20×30 km <sup>2</sup>	ASDC (2018)
IASI/METOP-A	LMD v2.2	12 km	C3S CDS (2019)
MISR/Terra	Plume Height	1.1 km	This Study

Deleted: <object><object><object>

Formatted ... [568]

Formatted: Space Before: 2.35 pt

Formatted: Space Before: 0 pt

Deleted: <object> ... [569]

Formatted: Font: 9.5 pt, Not Bold

Deleted: UVAerosolIndex

Deleted Cells

Formatted ... [571]

Deleted: Torres (2019a)

Formatted Table

Formatted: Not Expanded by / Condensed by

Formatted ... [572]

Formatted: Indent: Left: 0.08", Line spacing: single

Formatted ... [570]

Formatted: Indent: Left: 0.09", Line spacing: single

Deleted: UVAerosolIndex

Deleted Cells

Formatted ... [574]

Deleted: Torres (2021)

Formatted: Indent: Left: 0.08"

Formatted ... [573]

Formatted: Indent: Left: 0.09"

Formatted Table

Deleted: AER

Deleted: UVAerosolIndex

Deleted: 12×12

Deleted: Torres (2019b)

Deleted Cells

Formatted: Font: Times New Roman

Formatted: Indent: Left: 0.08", Line spacing: single


Formatted ... [575]

Formatted: Indent: Left: 0.09", Line spacing: single

Formatted Table

Formatted: Line spacing: single

Deleted: <object>



On 29 May 2018, remnants of the Aralkum dust plume continued to affect western Uzbekistan and Turkmenistan. The suspended dust was primarily detected by CALIOP due to extensive cloud cover. The dust layer extended to an altitude of 3.5 km. Likely resulting from daytime convection and increased vertical motion ahead of a low-level jet.



## 3.2 Comparison of UVAI Products

Figure 3 compares multiple UVAI products in detecting the fresh dust plume on 27 May 2018, including the Mie-based UVAI from OMPS, TROPOMI, and EPIC, all developed by NASA, and the TROPOMI LER-based UVAI developed by ESA. The sensor scan times over Aralkum (45°N, 60°E), as shown in each panel, are less than 20 minutes apart. The UVAI products display similar spatial patterns, but with significant differences in the dynamic range and statistical distribution. Among the three NASA products (Fig. 3(b–d)), OMPS yields markedly lower UVAI (e.g., mean=0.5, maximum=3.2) compared to EPIC (1.1 and 11.4) and TROPOMI (0.6 and 5.2), likely due to its coarser spatial resolution and cloud interference. EPIC exhibits significantly higher values over dusty scenes, partly because the 340/388 nm pair is more sensitive to absorbing aerosols. Nonetheless, the cloudy scenes in EPIC appear more noisy, most likely due to EPIC L1 calibration issues (personal communications with Karin Blank). Statistically, both OMPS and TROPOMI (NASA) reveal a bimodal distribution with a prominent peak near zero and a secondary peak near 1. EPIC displays a unimodal distribution with a larger spread, as indicated by the standard deviation and interquartile range.

Comparing the TROPOMI 354/388 nm UVAI from NASA and ESA (Fig. 3(d, f)) reveals that the ESA LER-based method produces more negative values over cloudy, dust-free scenes (e.g., Caspian Sea) than the NASA Mie-based method. Consequently, the ESA product exhibits greater dispersion—indicated by a trimodal distribution—although their regional mean and median differ by less than 0.05. The choice of 354/388 nm versus 340/380 nm wavelength pair in the ESA product has a minor effect on the UVAI, with small differences in the regional averages, extremes, and dispersion. However, the 335/367 nm pair produces significantly lower UVAI and less dispersion, indicating reduced sensitivity to aerosol absorption.

Dust detection using UVAI products often used fixed thresholds to isolate the dust signal. For example, Prospero et al. (2002) used a UVAI threshold of 1 for North Africa and 0.7 elsewhere. Schepanski et al. (2012) used a threshold of 2 to detect major dust plumes over the Sahara. As shown in Fig. 3(g), EPIC yields more than 18% pixels exceeding a threshold of 2—over three times those observed in OMPS and TROPOMI. This disparity suggests that using a fixed UVAI threshold may lead to significant inconsistencies in dust detection. In contrast, adopting a percentile-based threshold can more effectively capture the dynamic range of each product, resulting in more coherent dust plume detection. As demonstrated in Fig. 3, using the 95th percentile produces a consistent delineation of the dust plume, and captured three clusters of elevated UVAI values separated by clouds (see Fig. 3(a)).

A persistent feature in all UVAI products is the occurrence of large positive values along the eastern coast of Caspian Sea, particularly over the northern Caspian Sea and Garabogazköl Gulf (i.e., above 2). To determine whether airborne dust was responsible for these anomalies, we conduct a comparison between UVAI and coincident CALIOP AOD<sub>0.53</sub> measurements. The CALIPSO ground track is displayed in Fig. 3. To account for the along-track scan time differences (<30 min apart) between

Deleted: <object><object><object>

Formatted ... [577]

Deleted: 7.5×3 km<sup>2</sup>→ ESA (2021)

Formatted ... [578]

Formatted ... [579]

Moved up [19]:

Deleted: Optical Centroid Height (AOCH)

Formatted ... [581]

Formatted ... [580]

Formatted ... [582]

Deleted: show ...n detecting the fresh dust plume or

Formatted ... [602]

Formatted ... [583]

Formatted ... [601]

Formatted ... [603]

Deleted: TROPOMI (NASA) yield near zero values

Formatted ... [604]

Formatted ... [605]

Deleted: display a bimodal distribution with a peak

Formatted ... [606]

Deleted: UVAI is generally smaller

Formatted ... [584]

Formatted ... [607]

Deleted: influence

Formatted ... [608]

Deleted: yields

Formatted ... [609]

Deleted: UVAI than OMPS and TROPOMI (NASA) for

Formatted ... [610]

Deleted: of

Formatted ... [611]

Deleted: higher sensitivity

Formatted ... [612]

Deleted: in the 340/380 nm wavelengths. The

Formatted ... [613]

Deleted: communication with Karin Blank).

Formatted ... [585]

Deleted: The LER-based UVAI

Moved (insertion) [27]

Formatted ... [586]

Formatted ... [587]

Formatted ... [588]

Deleted: ..... Column Break ..... [589]

Formatted ... [614]

Deleted: from TROPOMI (ESA) are lower than the

Formatted ... [615]

Formatted ... [616]

Deleted: for dust plume tracking and source mapping

Formatted ... [617]

Formatted ... [618]

Moved up [26]: Black lines indicate the ground

Deleted: reached an altitude of 3.5 km, likely due

Formatted ... [590]

Moved up [27]:

Formatted ... [591]

Deleted: 3.2 Comparison of UVAI products

Formatted ... [592]

Moved up [25]:

Formatted ... [593]

Deleted: 3. Comparison of five UVAI products or


Formatted ... [594]

Formatted ... [595]

Deleted: . Schepanski et al. (2012)

Formatted ... [596]

Deleted: <object><object><object>



**Figure 3.** Comparison of UVAI products on 27 May 2018: (a) MODIS/Aqua true color image; (b–f) Five UVAI products from OMPS, EPIC, and TROPOMI; (g) Violin plots and summary statistics of six UVAI products. The sensor scan time over the Aralkum and CALIPSO ground track are indicated on panel (a–f). Black contours in (b–f) indicate the 95% percentile of UVAI values. The summary statistics in panel (g) are pixel count (N), mean, median (Med), standard deviation (SD), median absolute deviation (MAD), minimum (Min), maximum (Max), 95% percentile (Q95), interquartile range (IQR), and proportion of UVAI $\geq 2$  pixels (F2).

Moved (insertion) [30]

Formatted: Font: 9 pt

Formatted: Indent: Left: 0"

Formatted: Space Before: 1.15 pt


Deleted: , we compare the CALIOP AOD with coincident UVAI products along the overpass shown in Fig. 3. We create the co-located data by first identifying the nearest

Formatted: Condensed by 0.55 pt

Formatted: Font: 9 pt

Deleted: <object>





**Figure 4.** Coincident measurements of CALIOP 532 nm AOD and UVAI products along the eastern coast of Caspian Sea on 27 May 2018. The CALIPSO ground track is shown in Fig. 3. The Garabogazköl Gulf and northern Caspian Sea are shaded.

the sensors, we shifted the UVAI products to match the UVAI and CALIOP AOD<sub>0.53</sub> maxima observed at the southeastern coast of Caspian Sea (see Fig. 2(d)). Next, we applied a 5-second moving averaging to smooth the co-located data.

Figure 4 shows that CALIOP detected only a very thin aerosol layer over the northern Caspian Sea and Garabogazköl Gulf. The large positive UVAI in these regions also remained stationary and persisted on dust-free days, indicating that the elevated UVAI was unlikely caused by dust and was most likely related to time-invariant surface features. Due to the influx of sediment-rich water from the Volga and Ural rivers (Modabberi et al., 2019; Moradi, 2022), the northern Caspian Sea displays persistent discoloration, as seen in Fig. 3(a). This shallow, sediment-rich water strongly absorbs in the near-UV, producing a spectral signature with enhanced absorption at shorter wavelengths that resembles the signature of airborne dust (He et al., 2012; Lee et al., 2013). Similarly, the shallow, saline waters of Garabogazköl Gulf (with a salinity of ~ 35‰) cause strong UV absorption and yield a dust-like UVAI signal, in contrast to the main body of the Caspian Sea, which has a much lower salinity (~ 1.3‰) and yields near zero UVAI. In addition, the Sor Kaydak salt marsh to the east of northern Caspian produces large positive UVAI due to enhanced UV absorption by salt minerals and organisms (Fig. 3). Therefore, enhanced UV absorption by turbid and saline waters and salt flats causes the water-leaving UV radiances to deviate from those for a pure scattering atmosphere, thereby generating large positive UVAI that resembles the effects of airborne dust. This surface-related UVAI signal is likely

Deleted: <object><object><object>

Moved (insertion) [31]

Formatted: Font: 9 pt

Formatted: Indent: Left: 0"

Formatted: Space Before: 4.65 pt

Formatted: Condensed by 0.15 pt

Deleted: the CALIOP footprints, and then shifting the UVAI pixels along track based on the sensor scan time differences. As shown in Fig. 3, CALIOP has an average time difference of 10, 6, and 23 minutes from OMPS, TROPOMI, and EPIC, respectively. The coincident AOD and UVAI retrievals are displayed in Fig. 4, after applying a moving average every 5 seconds to smooth the

Moved (insertion) [28]

Formatted: Space Before: 6.25 pt

Deleted: from all

Formatted: Indent: First line: 0", Space Before: 4.9 pt

Deleted: <object>

**Table 2.** Summary statistics of eight AOD products in observing the Aralkum dust aerosol during 27–29 May 2018, based on best-quality, over-land retrievals. SD, standard deviation; MAD, median absolute deviation; IQR, interquartile range; Sk, skewness; 90th, 90th percentile. Preprint. Discussion started: 2 December 2024

Q Author(s) 2024. CC BY 4.0 License.

Product	N	Mean	SD	Median	MAD	IQR	Sk	90th	Max
MODIS/Terra DB	21,338	0.49	0.73	0.25	0.17	0.38	3.0	1.0	3.5
MODIS/Aqua DB	21,659	0.46	0.55	0.31	0.18	0.37	3.4	0.9	3.5
MODIS/Terra MAIAC	5,223,130	0.35	0.54	0.18	0.11	0.26	4.7	0.7	6
MODIS/Aqua MAIAC	5,070,926	0.37	0.69	0.17	0.10	0.24	4.3	0.7	6
VIIRS/SNPP DB	110,726	0.44	0.76	0.15	0.10	0.39	3.6	1.0	5
VIIRS/NOAA20 DB	110,286	0.56	0.85	0.26	0.19	0.55	3.2	1.2	5
VIIRS/SNPP EPS	5,295,908	0.49	0.62	0.26	0.19	0.54	2.3	1.2	5
VIIRS/NOAA20 EPS	3,996,574	0.37	0.53	0.19	0.16	0.39	2.8	1.0	5

common over ephemeral and dried lakes, and constitutes a potential source of error in using UVAI products for dust plume detection and source mapping.

3.3 Comparison of Midvisible AOD Products

3.3.1 AOD retrieval over desert surfaces

Table 2 shows the summary statistics of eight AOD products, based on the best-quality, over-land retrievals of the Aralkum dust event between 27 and 29 May 2018. The number of pixels differs greatly among the products, indicating inconsistent sampling due to different spatial resolutions, screening of irretrievable scenes (e.g., clouds, snow/ice, sun glint, ocean color), and/or QA definitions. The inconsistent sampling, along with algorithm differences as discussed in Section 2.2, contribute to the disparities in the AOD statistics. In addition, the NOAA EPS product uses L1b data from the NOAA VIIRS Calibration/Validation group, whereas the NASA DT and DB products use the L1b data from the NASA VIIRS Calibration Support Team, which may further explain the inconsistent retrievals.

Based on the MODIS products, the AOD statistics appear more consistent between the Terra and Aqua platforms than between the DB and MAIAC algorithms. In particular, the mean AOD differs by less than 10% between Terra and Aqua, but exceeds 20% between DB and MAIAC. Compared to MAIAC, DB produces a larger upper tail, as indicated by a higher 90th percentile, and consequently a broader distribution (higher SD, MAD and IQR). Whereas, MAIAC produces a narrower distribution (lower MAD and IQR) despite more extreme high values (upper limit of 6 versus 3.5 for DB).

The VIIRS DB and EPS algorithms employ the same upper limit (5), but display more complex AOD statistics compared to MODIS. For VIIRS/SNPP, EPS yields a higher mean and median as well as more extreme values, as indicated by a slightly higher 90th percentile. In contrast, DB produces a larger standard deviation and skewness, indicating a broader upper tail. For VIIRS/NOAA20, DB yields a larger spread (higher SD and IQR), and a larger upper tail (higher 90th percentile), suggesting that DB captures more frequent high AOD values than EPS.

Deleted: <object><object><object>

Moved (insertion) [32]

Moved (insertion) [29]

Formatted: Font: Times New Roman, Bold

Formatted: Space Before: 0.6 pt

Deleted: CALIOP detected a dust layer at the southeastern coast of Caspian Sea (see Fig. 2d) with an AOD up to 1.8. Over the northern Caspian Sea and Garabogazköl gulf, CALIOP detected a very thin aerosol layer with negligible AOD, which may be transported dust from the adjacent deserts (e.g., Ryn Desert). This thin aerosol layer, however, is unlikely to explain the large positive UVAI (i.e. >2) found in these regions. Such high UVAI values are found to persist during dust-free days and not change with time based on the high-cadence EPIC observations. Both the northern Caspian Sea and Garabogazköl gulf have an average water depth of less than 10 m, and contain large bodies of turbid or salty water (Modabberi et al., 2019). The

... [632]

Formatted: Font: Arial

Formatted: Space Before: 5.5 pt


Formatted: Space Before: 0 pt

Formatted: Space Before: 0.05 pt

Deleted: ¶  
¶  
¶

... [633]

Deleted: <object>



**Figure 5.** Over-land AOD retrievals on 27–29 May 2018. (a–c) VIIRS/NOAA20 true color images, (d–f) MODIS/Aqua DB product, (g–i) MODIS/Aqua MAIAC product, (j–l) VIIRS/NOAA20 DB product, (m–o) VIIRS/NOAA20 EPS product, and (p–r) VIIRS/NOAA20 EPS product using the dust optical model as a test case.

Next, we examined the MODIS/Aqua DB and MAIAC products, and the VIIRS/NOAA20 DB and EPS products. By comparing AOD retrievals from the same sensor, we can attribute the disparity to the choice of algorithms. The instruments are hereafter referred to simply as MODIS and VIIRS unless otherwise noted. Due to the different spatial resolutions, all products are gridded onto a uniform  $0.2^{\circ} \times 0.2^{\circ}$  resolution by computing the mean value within each grid cell. Only the best-quality retrievals are used, based on the QA rules recommended in each product.

Figure 5 displays the daily gridded AOD products, together with VIIRS true color images to help attribute missing retrievals to either clouds or incorrect cloud screening. None of the algorithms captured the full extent of the fresh dust plume on 27 May 2018, while the retrieved AOD for the dust scene reached the upper limit of each algorithm. There are several possible reasons. First, the AOD retrieval from reflected sunlight relies on the scene brightening by aerosol scattering (primarily the forward scattering of surface reflection) against a relatively dark background. At high aerosol loadings, the TOA reflectance becomes less sensitive to additional AOD increases. Second, AOD retrieval is performed by matching observed TOA reflectances against pre-computed values within a specific AOD range, which may be a poor fit if the aerosol burden is high or exceeds the predefined AOD limit. These retrievals are often flagged as marginal quality, and consequently excluded from our screening for best quality data. Setting an upper AOD limit is necessary to minimize the impact of residual cloud contamination. Heavy dust scenes may be either retrieved but with low confidence in clear sky, or classified as clouds and not retrieved.

Deleted: <object><object><object>

Moved up [30]:

Moved down [34]:

Deleted: first evaluate

Deleted: consistency of AOD products in observing ... [643]

Deleted: . The MODIS regional mean (or median) A ... [645]

Deleted: 5. Multi-sensor AOD retrieval of the Arak ... [647]

Deleted: ..... Column Break ..... [648]

Deleted: from VIIRS/NOAA20

Formatted ... [641]

Formatted ... [642]

Formatted ... [644]

Formatted ... [646]

Formatted ... [649]

Formatted ... [650]

Deleted: specified. By com- paring two different pr ... [651]

Moved up [32]: Table 2.

Deleted: Summary statistics of eight AOD produc ... [634]

Formatted ... [635]

Deleted: grid, and calculate

Formatted ... [652]

Formatted ... [653]

Deleted: The

Formatted ... [636]

Deleted: AODs are displayed in Fig. 5

Deleted: the

Deleted: which can

Deleted: identify

Formatted ... [654]

Formatted ... [655]

Formatted ... [656]

Formatted ... [657]

Formatted ... [658]

Deleted: due

Deleted: erroneous

Deleted: ... [661]

Formatted ... [659]

Formatted ... [660]

Formatted ... [662]

Deleted: heavy

Formatted ... [663]

Deleted: . Among the four products, VIIRS EPS ha ... [664]

Deleted: of the scene

Formatted ... [665]

Formatted ... [666]

Deleted: As the

Formatted ... [667]

Deleted: burden increases beyond a certain magnitude


Formatted ... [668]

Formatted ... [637]


Deleted: start to flatten out and become insensitive ... [669]

Deleted: ... [638]

Deleted: ... [639]



**Figure 6.** Misclassification of the fresh dust plume as clouds on 27 May 2018. (a) Cloud fraction from MODIS/Aqua DB product; (b) Cloud mask from MODIS/Aqua MAIAC product; (c) Unsuitable pixel fraction from VIIRS/NOAA20 DB product; and (d) Cloud mask from VIIRS/NOAA20 EPS product. Red contours indicate the 95th percentile (2.1) of TROPOMI TropOMAER UVAI product.



**Figure 7.** Aerosol optical model used for over-land retrievals on 28 May 2018 in the (a) MODIS/Aqua MAIAC, (b) VIIRS/NOAA20 DB, and (c) VIIRS/NOAA20 EPS products.

To check whether the fresh dust plume was erroneously classified as clouds, Fig. 6 displays the cloud fraction and cloud masks reported in selected granules from each product, as well as the TROPOMI UVAI on 27 May 2018. Given the small overpass time differences between SSP and Aqua (5 minute) and between SSP and NOAA20 (1 hour), TROPOMI UVAI provides a nearly coincident view of the dust plume. Fig. 6 reveals that the thick dust plume is mostly flagged as clouds in all products. For MODIS, the DB algorithm detected parts of the plume fringes, where the AODs reached the algorithm limit (3.5). MAIAC classified the majority of dust scene as “cloudy”. For VIIRS, the DB algorithm does not report cloud masking or fraction; instead it reports an Unsuitable Pixel Fraction Land Ocean parameter, which indicates that the dust pixels near Arakum were not retrieved. Similarly, the dust scene was flagged as “confidently cloudy” in the EPS algorithm.

Deleted: <object><object><object>

Moved up [31]:

Figure

Deleted: on 27 May 2018 was erroneously categorized

Deleted: , Fig. 6 shows the cloud fraction or mask reported in selected L2 granules from each product. The TROPOMI TropOMAER UVAI is also shown to indicate the dust plume location. Given the slight difference in overpass time between SSP and Aqua (5 minute apart) and between SSP and NOAA20 (1 hour apart), TROPOMI UVAI provides a nearly coincident view of the dust plume as MODIS.

... [674]

Deleted: 6. Mis-classification of the heavy dust plume

Formatted ... [673]

Formatted: Font: 9 pt

Formatted ... [675]

Formatted ... [677]

Deleted: in the

Deleted: MAIC

Deleted: in the

Deleted: in the...from VIIRS/NOAA20 EPS product

Formatted ... [681]

Formatted ... [678]

Formatted ... [679]

Formatted ... [680]

Deleted: <object>

Formatted: Space Before: 6.3 pt

Deleted: The aerosol

Deleted: AOD retrieval

Formatted ... [682]

Formatted ... [683]

Formatted ... [684]

Formatted: Right: 0.65", Line spacing: Multiple 1.48 li

Moved (insertion) [33]

Formatted: Space Before: 6.2 pt

Moved (insertion) [35]

Deleted: <object>



On 28 May 2018, a high-pressure system produced cloud-free conditions over Central Asia, providing an optimal sky condi-  
https://doi.org/10.5194/amt-2024-341. All products successfully captured the extensive dust layer over western Turkmenistan;  
Preprint. Discussion started: 2 December 2024  
Q Author(s) 2024. CC BY 4.0 License.  
however, there are large differences in the AOD magnitudes. Notably, VIIRS EPS yields significantly lower AODs than  
MODIS DB than other products over dust-affected areas. As described in Section 2, all the algorithms use predefined aerosol  
optical models (including a coarse-mode dust models) to represent location- and season-dependent aerosol microphysical prop-  
erties. To examine whether the dust optical model was successfully selected, Fig. 7 displays the aerosol optical model reported  
in selected granules. MODIS DB does not report this information and thus is not shown. Both MODIS MAIAC and VIIRS DB  
successfully selected their respective dust models for the dust scene over western Turkmenistan. VIIRS EPS selected the urban  
aerosol model for the dust scene, presumably because the urban model produced a closer match between the observed and  
simulated TOA reflectances.  
According to Laszlo and Liu (2022), EPS first retrieves AOD in the blue channel (0.41  $\mu\text{m}$ ) and then scales it to longer  
wavelengths using the normalized spectral extinction coefficients from the aerosol optical model. The normalized extinction  
coefficient exhibits much stronger wavelength dependence in the urban model than the dust model (see Figure 3-4 in Laszlo  
and Liu (2022)). For the same  $\text{AOD}_{0.41}$  value, the dust model would yield more than 40% higher AOD compared to the urban  
model. We therefore hypothesize that had the dust model be selected as intended, EPS would have retrieved higher AOD that  
align more closely with DB and MAIAC. Indeed, when we forced EPS to use the dust model, as shown in Fig. 5(p-r), the EPS  
agreement with DB and MAIAC improved, although the retrieved AODs remained lower and produced more marginal  
retrievals close to the upper limit that were subsequently excluded.  
Using the gridded AOD products, we conducted a regression analysis to further investigate the inconsistencies among aerosol  
algorithms (Fig. 8). The choice of algorithms for both MODIS and VIIRS introduces a non-linear response in the retrieved  
AOD. For MODIS, MAIAC produces lower AOD than DB under low aerosol loadings (e.g., 36% lower for  $\text{AOD} < 1$ ), but  
higher AOD under heavy loadings (22% higher for  $2 < \text{AOD} < 3$ ). This nonlinearity may be partly explained by MAIAC's higher  
spatial resolution, which yields more extreme retrievals (either very clean or heavily polluted). The higher AOD limit in  
MAIAC (6 vs. 3.5 for DB) may further exacerbate the discrepancies at heavy aerosol loadings, as shown in Fig. 8(a). For  
VIIRS, EPS retrievals are generally lower than DB across all aerosol loadings, with the EPS-DB difference increasing from  
37% for  $\text{AOD} < 1$  to 51% for  $4 < \text{AOD} < 5$  (Fig. 8(b)). Indeed, Fig. 5(m-o) confirm that EPS yields approximately 50% lower  
AODs than other algorithms in areas affected by thick dust.  
Figure 9 displays the MISR AOD retrievals on 28 May 2018 (orbit 98092) based on the standard operational product and the  
MISR research algorithms. The standard algorithm exhibits several limitations. First, it uses an outdated land/water mask and  
marked the Aral Sea as "shallow water"; as a result, no retrievals were performed over the Aralkum Desert. Second, the thick  
dust over the Kopet-Dag foothills was misclassified as clouds. Lastly, the retrieved AOD is substantially lower than those from  
single-view sensors (Fig. 5). Kahn et al. (2010) noted that MISR tends to underestimate AOD under heavy aerosol loadings, as  
the weak surface reflection signal leads to poor surface-atmosphere separation and overestimation of the surface contribution  
to TOA radiances. Another contributor to the biased AOD retrieval is the lack of appropriate aerosol optical models for saline  
dust; Consequently, the particle property retrievals are of low confidence. We find that the standard algorithm selected spherical

Deleted: <object><object><object>

Moved (insertion) [34]

Formatted: Space Before: 9.65 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 5.5 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 5.5 pt

Formatted: Space Before: 5.45 pt

Formatted: Space Before: 0.05 pt

Deleted: (c) VIIRS/NOAA20 EPS products.

Formatted: Font: Arial

Formatted: Space Before: 5.45 pt

Deleted: ..... Column Break.....

Moved up [35]: . Fig.

Moved down [36]: bin-averaged AODs every 0.1 increments. RMSD, root mean square difference.

Deleted: 6 reveals that the majority of the heavy dust plume is classified as clouds in all products. For MODIS, the DB algorithm only detected the dust plume fringe area, where the retrieved AOD reached the algorithm limit (3.5). MAIAC classified the majority of dust pixels as "cloudy", but it was able to see the "clear" pixels through cloud gaps. Surprisingly, MAIAC successfully detected the sediment-rich water of the northern Caspian Sea and Garabogazköl gulf. For VIIRS, the DB algorithm does not report the cloud fraction or mask, but instead report the fraction of level-1 pixels not used in the retrieval, named

Deleted: .....Section Break (Next Page).....

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.1 pt


Formatted: Font: 10 pt

Formatted: Font: Arial

Formatted: Space Before: 5.5 pt

Deleted: <object>

21



**Figure 8.** Comparison of over-land AOD retrievals during 27–29 May 2018: (a) MODIS/Aqua MAIAC vs. DB; (b) VIIRS/NOAA20 EPS vs. DB. Black lines indicate the linear model fits. Red squares indicate bin-averaged AODs every 0.1 increments. RMSD, root mean square difference.

non-absorbing aerosol mixtures for the dust scene, resulting in underestimation of the nonspherical fraction and overestimation of FMF and SSA.

480 To explore the particle-property information content of MISR observations, we ran the MISR research algorithm in three modes: (a) with the surface retrieved self-consistently with the atmosphere using only MISR data, (b) with the surface re-

Deleted: <object><object><object>

Formatted: Justified, Right: 0.65", Space Before: 0.05 pt

Moved (insertion) [36]

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.15 pt

Formatted: Condensed by 0.1 pt

Formatted: Font: 10 pt

Formatted: Space Before: 2.25 pt

Deleted: ¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

480¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶

¶


¶

¶

Deleted: <object>

Deleted: <object><object><object>

<https://doi.org/10.5197>  
Preprint. Discussion  
Q Author(s) 2024. C



**Figure 9.** MISR aerosol retrievals on 28 May 2018 based on: (top row) Operational standard product; (2nd row) Research algorithm with retrieved surface reflectances from MISR data; (3rd row) Research algorithm with prescribed surface reflectances from MODIS MAIAC product; and (bottom row) Research algorithm using an AI/ML approach.

reflectance prescribed from the MODIS MAIAC product (Limbacher et al., 2022), and (c) with a modified algorithm based on an AI/ML approach. Initial statistics indicate that these models do well in the regional means. AOD is retrieved fairly consistently with all three approaches, except in the highest and lowest AOD regions. The prescribed surface results appear to perform bet-

Deleted: <object>

485

ter for extremely high AOD conditions and capture the increasing AOD gradient towards the Kopet-Dag Range. The retrieved  
https://doi.org/10.5194/amt-2024-1460  
Preprint. Discussion started: 2 December 2024  
in expectation in this case (i.e., NonSph = 0.8), whereas the retrieved- and prescribed-surface results may have been estimated  
Q Author(s) 2024. CC BY 4.0 License.

490

3.3.2 AOD retrieval over the Caspian Sea

In this section, we compare six AOD products in observing the dust outflow to the Caspian Sea on 28 May 2018, based on the  
best quality, over-water retrievals from MODIS/Aqua (DT, DT3K, and MAIAC) and VIIRS/NOAA20 (DT, DB, and EPS). All  
products are gridded to a 0.1°×0.1° resolution. Bilinear interpolation is applied to the MODIS DT product in order to fill the  
data gaps due to reduced pixel resolutions near sensor swath edges.

495

Figure 10 shows that all six products capture the enhanced aerosol burden associated with dust outflow to the Garabogazköl  
Gulf and Caspian Sea. Among the MODIS products, DT3K and MAIAC produce more extreme AOD over Garabogazköl Gulf  
than DT, resulting in higher means and greater spread (indicated by higher SD and MAD). These spurious high AODs create  
unnatural discontinuities along coastal regions, likely due to inaccurate surface reflectance characterization of the shallow,  
saline waters. In the VIIRS products, DT exhibits extremely high AOD over the shallow waters and salt flats of northern  
Caspian Sea, and to a lesser extent over the coastal region of Garabogazköl Gulf. The non-negligible surface signal from these  
500 areas deviated from the dark surface characteristics of deep water assumed in the algorithms, and was likely misinterpreted as  
aerosol signal, leading to spurious high AODs. As a result, VIIRS DT produced a higher mean, more extreme upper tail values  
(indicated by the 90th percentile and maximum), and a larger spread (higher SD and IQR) than the DB and EPS products.

505

We further examined the aerosol optical models selected during the over-water retrieval (Fig. 11). The MAIAC spheroidal  
dust model ("Model 6") was applied to only small parts of the dust scene, resulting in discontinuous AOD patterns. We  
computed the relative fractions of three aerosol groups in the VIIRS DT product: fine aerosol, sea salt, and coarse dust. The  
sea salt group dominated the retrieved AOD, indicating that the spherical dust model was not selected for retrieval. VIIRS  
DB successfully applied the spheroidal dust model for the full extent of the dust scene, and the maritime model for dust-  
free regions, while VIIRS EPS selected its oceanic aerosol model for the full scene, again failing to select the dust model as  
intended. In summary, the aerosol algorithms showed varied performance in selecting the dust optical model during retrievals.  
Figure 12 compares the gridded AOD products via linear regression to assess the cross-algorithm consistency. Overall, over-  
water retrievals exhibit stronger linear relationships and better agreement than over-land products (Fig. 8), with R<sup>2</sup> above 0.9  
and RMSD below 0.1 in all cases. In particular, MODIS DT and DT3K products show excellent agreement with a regression  
slope of 0.98 and an R<sup>2</sup> of 0.91, while other products show slightly weaker slopes of approximately 0.8. Notably, DT yields  
515 lower AOD under clean marine conditions (AOD<0.15), but higher AOD in dusty conditions than the MAIAC, DB and EPS  
algorithms.

Compared to aerosol retrieval over land, over-water retrieval provides greater information content to constrain particle prop-  
erties. As described in Section 2.2.1, over-water algorithms represent the aerosol column as a sum of a fine and a coarse mode,

Deleted: <object><object><object>

Formatted: Space Before: 5.5 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0.05 pt

Deleted: ..... Column Break .....  
Using the regridded AOD in Fig. 5, we further use regression analysis to examine the inter-algorithm AOD differences from MODIS and VIIRS. We find that the choice of algorithms for both MODIS or VIIRS causes a non-linear relationship between the AOD products. Hence a quadric model is used for fitting, as shown in Fig. 8. The model goodness of fit is measured by R<sup>2</sup> and root mean square difference (RMSD). Compared to DB, MAIAC produces lower AOD from MODIS under low aerosol burden, but higher AOD under heavy burdens. For example, the MAIAC product is 36% lower than the DB product when AOD<1, but 22% higher when 2<AOD<3. This nonlinear relationship may be partly explained by the higher resolution in

... [688]

Formatted: Font: Bold

Formatted: Space Before: 4.75 pt

Formatted: Space Before: 5.5 pt

Deleted: ¶

Deleted: ..... Column Break .....  
Dust outflow to Caspian Sea

... [689]


Formatted: Font: Arial

Formatted: Space Before: 5.5 pt

Deleted: ¶

... [690]

Deleted: <object>



**Figure 10.** Over-water AOD retrievals over the Caspian Sea on 28 May 2018. Top row panels are MODIS/Aqua (a) true color image, (b) DT AOD, (c) DT3K AOD, and (d) MAIAC AOD. Bottom row panels are VIIRS/NOAA20 (e) true color image, (f) DT AOD, (g) DB AOD, and (h) EPS AOD. All products are gridded to a  $0.1^\circ \times 0.1^\circ$  resolution.

and retrieves the total AOD and FMF. Here, we use the total AOD and FMF to compute the coarse-mode AOD from each product, and perform a similar comparison to that in Fig. 12. Figure 13 shows that the coarse-mode AODs exhibit strong linear relationships, with  $R^2$  above 0.8 and RMSD below 0.1, although the agreement is somewhat weaker than for the total AOD. In general, the DT algorithm tends to yield higher coarse-mode AOD under dust-laden conditions compared to DT3K, MAIAC, DB and EPS.

### 3.4 Comparison of Infrared AOD ( $AOD_{10}$ ) Products

In this section, we compare the IASI  $AOD_{10}$  products from four algorithms: LMD, MAPIR, ULB and IMARS, based on Level-3 daily gridded ( $1^\circ \times 1^\circ$ ) products from the Aerosol\_cci project. Table 3 summarizes the  $AOD_{10}$  statistics based on

Deleted: <object><object><object>

Formatted: Font: Times New Roman

Formatted ... [693]

Formatted: Space Before: 3.4 pt

Deleted: which is then regridded

Deleted: a  $0.1^\circ \times 0.1^\circ$  resolution. The regridded

Deleted: are compared via

Formatted ... [694]

Formatted ... [695]

Formatted ... [696]

Formatted: Space Before: 0 pt

Deleted: regression, as shown in Fig. 11. The coarse-mode AODs are strongly linearly correlated, but show larger inter-algorithm differences

Deleted: AODs

Formatted ... [697]

Formatted ... [698]

Deleted: produce

Deleted: ,

Deleted: other algorithms

Formatted ... [699]

Formatted ... [700]

Deleted: AOD products

Formatted ... [701]

Formatted: Left, Indent: Left: 0.07", Hanging: 0.35", Space Before: 9.7 pt, Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 3 + Alignment: Left + Aligned at: 0.41" + Indent at: 0.75", Tab stops: 0.42", Left + Not at 0.38"

Deleted: Figure 12 displays several parameters from the EPIC AOD product on 28 and 29 May 2018, including the 680 nm surface reflectivity, 680 nm AOD, AOD<sub>10</sub>, and AOD<sub>10</sub> after applying a filter of AOD>1.8. The heavy dust plume on 27 May 2018 was missed by EPIC and thus not shown. On 28 May 2018, the low surface reflectivity of Caspian Sea (i.e. below 0.03), combined with the elevated AOD (i.e. 0.5–1) associated with ... [702]

Formatted: Space Before: 0.05 pt

Moved down [37]:

Deleted: ... [691]


Deleted: ..... Column Break ..... [692]

Formatted: Font: Times New Roman


Deleted: <object>



Deleted: <object><object><object>



**Figure 11.** Aerosol optical models used for AOD retrieval over the Caspian Sea on 28 May 2018 in the (a) MODIS/Aqua MAIAC, (b) VIIRS/NOAA20 DT, (c) VIIRS/NOAA20 DB, and (d) VIIRS/NOAA20 EPS products.



**Figure 12.** Comparison of AOD retrievals over the Caspian Sea on 28 May 2018: (a) MODIS/Aqua DT3K vs. DT, (b) MODIS/Aqua MAIAC vs. DT, (c) VIIRS/NOAA20 DB vs. DT, (d) VIIRS/NOAA20 EPS vs. DT. Black lines indicate the linear regression fits. Red squares indicate bin-averaged AODs at 0.05 increments. RMSD, root mean square difference.

**Table 3.** Summary statistics of four IASI AOD<sub>10</sub> products in observing the Aralkum dust aerosol during 27–29 May 2018. SD, standard deviation. MAD, median absolute deviation. QR, interquartile range. Sk, skewness.

Algorithm	N	Mean	SD	Median	MAD	IQR	Sk	Max
LMD	1122	0.13	0.20	0.05	0.05	0.17	2.5	1.2
MAPIR	1380	0.11	0.11	0.08	0.03	0.07	6.0	1.7
ULB	1486	0.06	0.09	0.03	0.04	0.09	2.0	0.6
IMARS	365	0.08	0.07	0.07	0.05	0.10	1.4	0.4

Deleted: 10

Formatted: Not Expanded by / Condensed by

Deleted: products using different algorithms over

Formatted

Formatted

Formatted: Space Before: 0.05 pt

Deleted: the ...in-averaged AODs every

Formatted: Font: 9 pt

Formatted: Space Before: 0 pt

Deleted: By applying a filter of AOD>1.8, we find that the majority of the unrealistically high AOD pixels are removed, resulting in a spatially continuous AOD pattern from Turkmenistan to the Caspian Sea (Fig. 12g). The AOD-filtered AOD suggests that the lofted dust over Turkmenistan was located between 1–1.5 km on 28 May 2018, and between 2–3 km on 29 May 2018, or twice as high as the previous day. The temporal variation of dust height can be explained by the meteorological condition:

Deleted: <object>




Figure 13. Same as Fig. 12 but for coarse-mode AOD.




Figure 14. IASI AOD<sub>10</sub> retrievals from four algorithms on 27–29 May 2018. (a–c) LMD, (d–f) MAPIR, (g–i) ULB, and (j–l) IMARS.

separate daytime (descending node) and nighttime (ascending node) retrievals on 27–29 May 2018. Figure 14 displays the daily combined AOD<sub>10</sub> by merging the daytime and nighttime observations. The inclusion of both daytime and nighttime data substantially improved the data coverage compared to midvisible AOD products that relies solely on daytime observations

530 (Fig. 5).

Deleted: <object><object><object>

Moved down [38]: ¶  
540

Deleted: a prevailing high pressure system may have suppressed the vertical aerosol mixing on 28 May 2018, whereas the enhanced upward vertical motion ahead of a deepening cold front promoted the convective mixing of the lofted dust on 29 May 2018. ¶

... [706]

Deleted: 11

Deleted: 10

Deleted: the

Formatted: Font: Arial, 8.5 pt

Formatted: Condensed by 0.25 pt

Formatted: Condensed by 0.3 pt

Formatted: Condensed by 0.3 pt

Formatted: Condensed by 0.25 pt

Formatted: Condensed by 0.3 pt

Deleted: <object>

535

540

545

550

555

Among the four algorithms, LMD has the best performance in detecting the dust plume features both over deserts and the extensive plume stretching from the Aralkum Desert to the southern borders of Turkmenistan (Fig. 14(a)), outperforming midvisible techniques in detecting the fresh dust plume over deserts. The highest AOD<sub>10</sub> (1.2) was observed downwind over Uzbekistan rather than near the Aralkum Desert, due to difficulty in detecting the heavy dust near the source. Additionally, the retrieved AOD<sub>10</sub> reached the upper limit of the predefined range (0–1.4), indicating possible underestimation. Unlike midvisible AOD, which includes fine aerosol contributions, AOD<sub>10</sub> is sensitive to coarse particles only. The AOD<sub>10</sub>-to-AOD ratio depends on the particle size distribution, particularly the relative fraction of fine and coarse modes. Using a AOD<sub>10</sub>-to-AOD ratio of 0.6 for coarse-dominated dust would imply an AOD<sub>10</sub> of around 3—more than twice the upper limit set in the LMD algorithm—corresponding to the maximum observed VIIRS DB AOD on 27 May (Fig. 5(j)). On 28 May, LMD successfully retrieved the dust accumulated by the foothills of Kopet-Dag Range and dust outflow to the Caspian Sea (Fig. 14(b)), resulting in a spatially continuous AOD<sub>10</sub> pattern, in contrast to midvisible retrievals which exhibited unnatural discontinuities near coastal areas due to inconsistent treatment of water and land surface reflectances. The AOD<sub>10</sub> retrieval on 29 May captured the remnant dust from the previous emission event as well as the newly emitted dust from Aralkum, consistent with the midvisible AOD products.

Compared to LMD, MAPIR generally yields higher background AOD<sub>10</sub> over aerosol-free areas but lower AOD<sub>10</sub> over dust-laden scenes, resulting in less distinct plume features (Fig. 5(g–i)). The two products exhibit moderate correlations, with R<sup>2</sup> of 0.6 for daytime retrievals, and 0.4 for nighttime retrievals. MAPIR successfully retrieved the thick dust on 27 May, although with a smaller spatial extent than LMD. However, MAPIR struggled to detect dust over the complex terrains of the Ustyurt Plateau and Kopet-Dag foothills on 28–29 May, likely due to increased uncertainty in characterizing the heterogeneous surface properties (e.g., emissivity, temperature), to which infrared retrievals are particularly sensitive over elevated terrains (Capelle et al., 2014).

Overall, ULB and IMARS exhibit poorer performance compared to LMD and MAPIR. Specifically, ULB misclassified the entire dust plume as clouds on 27 May, and retrieves very low AOD<sub>10</sub> on 28–29 May. IMARS reported substantial missing data for dust-affected areas on all three days, likely due to overly restrictive cloud masking and/or low confidence associated with its probabilistic retrievals.

3.5 Comparison of ALH Products

In this section, we compare four ALH products: CALIOP aerosol extinction-weighted height, EPIC-retrieved AOC, IASI LMD mean dust layer altitude, and the MISR stereo-height retrieval. The CALIOP ALH is calculated from the aerosol extinction vertical profiles at 532 and 1064 nm, whereas the EPIC and IASI products retrieve an effective ALH using passive techniques that rely on a priori assumptions about aerosol vertical distributions (Section 2.4). Among the IASI products, we focus on the LMD algorithm due to its capability in detecting distinct dust features (Fig. 14). Below, we first examine the EPIC retrievals at around 09:00z on 28 and 29 May 2018, and use co-located CALIOP measurements for validation. The dust plume on 27 May is misclassified as clouds in the EPIC AOC product and thus not discussed. Next, we compare the coinci-

Deleted: <object><object><object>

Moved (insertion) [37]

Formatted: Font: Times New Roman

Formatted: Space Before: 4 pt

Formatted: Font: Arial, 8.5 pt


Formatted: Space Before: 0 pt

Moved (insertion) [38]

Formatted: Font: Arial, 8.5 pt

Formatted: Left, Indent: 0", First line: 0", Right: 0", Space Before: 5.45 pt, Line spacing: single

Deleted: Using the daytime CALIOP overpass on 29 May 2018, we evaluated the accuracy of EPIC AOC against the CALIOP aerosol extinction-weight height, and explored the effects of wavelength choice (532 vs 1064 nm) and AOD filter. The CALIPSO ground track is shown in Fig. 12f. Past studies suggested that the CALIOP 1064 nm measurement is more sensitive to coarse aerosols close to the ground, while the 532 nm channel is subject to sensitivity loss in the presence of carbonaceous aerosols (Torres et al., 2013). Fig. 13a shows that the choice of 532 or 1064 nm has a minor effect on the CALIOP extinction-weighted height, with a mean difference below 0.1 km or 5%. Both indicate that the dust layer was mostly located between 1–2 km.




Formatted: Condensed by 0.25 pt

Formatted: Font: Arial

Formatted: Space Before: 5.5 pt

Deleted: ¶

Deleted: <object>



**Figure 15.** EPIC AOC retrievals at 09:01z 28 May 2018 (top row) and 08:47z 29 May 2018 (bottom row): (a, b) 680 nm surface reflectivity; (c, d) 680 nm AOD; (e, f) AOC. The coincident CALIPSO ground track is shown in panel (f).

dent retrievals from EPIC and IASI to assess the consistency between these passive techniques for retrieving ALH over desert surfaces. Finally, we present results on MISR plume height retrieval on 28 May 2018.

Figure 15 displays the  $0.68 \mu\text{m}$  surface reflectivity,  $\text{AOD}_{0.68}$ , and AOC retrieved from EPIC. On 28 May, the low surface reflectivity and elevated dust loading (i.e.,  $\text{AOD}_{0.68} > 0.5$ ) over Caspian Sea provided optimal conditions for AOC retrieval. Figure 15(e) shows that the dust outflow to Garabogazköl Gulf and Caspian Sea was elevated to an altitude of 1.5–2.5 km, likely enhanced by orographic lifting as the dust was transported across the Ustyurt Plateau.

In contrast, the bright desert surfaces of Central Asia pose significant challenges in isolating aerosol scattering contributions to the EPIC measurements. Radiative transfer simulations indicate that the sensitivity to ALH in the  $\text{O}_2$  band spectral contrast diminishes over bright surfaces, requiring high aerosol loadings to generate sufficient signal-to-noise ratios for reliable AOC retrieval (Xu et al., 2019). As shown in Fig. 15, the EPIC-retrieved  $\text{AOD}_{0.68}$  and AOC exhibits inverse spatial patterns: AOC displays unrealistically high values over low  $\text{AOD}_{0.68}$  areas, and reasonably low values over high  $\text{AOD}_{0.68}$  (up to 6) areas that form a continuous pattern with the Caspian Sea retrievals. Thus,  $\text{AOD}_{0.68}$  provides a practical means for screening marginal-quality AOC retrievals. Focusing on dust scenes where  $\text{AOD}_{0.68}$  exceeds 2, the retrieved AOC over Turkmenistan's desert plains increased from 1–1.5 km on 28 May to 2–3 km on 29 May. This temporal evolution can be explained by the meteorological conditions: a prevailing high-pressure system suppressed vertical aerosol mixing on 28 May, whereas enhanced upward motion ahead of a deepening cold front promoted convective mixing on 29 May.

Deleted: <object><object><object>

Formatted: Font: Times New Roman

Formatted: Space Before: 3.4 pt

Formatted: Space Before: 0.05 pt

Deleted: ..... Column Break ..... [709]

Moved down [39]:  
Conclusions

Deleted: ¶  
Driven by the desiccation of the Aral Sea, the newly formed Aralkum Desert has emerged as a major source of wind-blown saline dust aerosol in Central Asia. There is generally a poor understanding of the performance and consistency of satellite aerosol remote sensing techniques in observing the airborne dust from Aralkum Desert. This paper addresses this knowledge gap in two aspects. ¶  
First, we provided a review of the physical principles and algorithms used to retrieve UVAI, AOD, and AOC over desert surfaces, focusing on the algorithms' treatment of regional aerosol optical properties and land surface reflectivity. As satellite remote sensing aerosol algorithms are generally optimized for global performance, the algorithm differences are expected to have a larger effect on the product performance and consistency over Central Asia, where the local aerosol and land surface ¶


[710]

Formatted: Font: Arial

Formatted: Space Before: 5.45 pt

Deleted: <object>

Deleted: <object><object><object>



**Figure 16.** (a) Coincident retrievals of CALIOP extinction-weighted ALH and EPIC AOC along the CALIPSO overpass on 29 May 2018 (ground track shown in Fig. 15f). Magenta triangles are EPIC-retrieved AOD<sub>0.68</sub>. Gray line indicates the ground surface. (b) Comparison between CALIOP 532 nm ALH and EPIC AOC. (c) Comparison between CALIOP 1064 nm ALH and EPIC AOC. N, number of coincident retrievals. MB, mean bias. RMSE, root mean square error. EE, percentage of co-located retrievals within expected errors of 0.5 and 1 km (shaded).

The EPIC AOC retrievals are evaluated against the CALIOP extinction-weighted ALH along the ground track shown in Fig. 16(f). Generally, the CALIOP 532 nm channel is more prone to signal attenuation in the presence of dense absorbing aerosols, resulting in lower ALH than the 1064 nm channel (Torres et al., 2013). Our analysis reveals that the wavelength choice has minor effects on the CALIOP-derived ALH, with a mean difference of 0.05 km between the two channels. EPIC AOC retrievals exhibit substantial positive biases in regions with low aerosol loadings. Using the CALIOP 532 nm ALH as

Moved (insertion) [40]

Formatted: Font: 10 pt


Formatted: Space Before: 4.05 pt

Deleted: ..... Column Break.....  
Second, we presented a case analysis of the cross-sensor and cross-algorithm consistency in observing a saline dust storm from the Aralkum Desert during 27–29 May 2018. We considered a wide range of aerosol products, including the UVA1 products from OMPS, TROPOMI, and EPIC, the AOD products from MODIS and VIIRS based on multiple algorithms (DT, DT3K, DB, MAIAC, and EPS), and the AOC derived from CALIOP aerosol extinction profile and EPIC oxygen absorption spectroscopic measurements

Formatted: Not Expanded by / Condensed by

Deleted: <object>





**Figure 17.** ALH retrievals in the (a, b) EPIC AOC and (c, d) IASI LMD products on 28–29 May 2018. Only the EPIC AOC retrievals with  $AOD_{0.68} > 1.5$  are shown. The sensor scan times over Aralkum (45°N, 60°E) are shown at the top right corner of each panel.

a benchmark, the EPIC AOC product has a mean bias of 3.6 km and an RMSE of 4.3 km, with only 17% (21%) of the co-located retrievals within 0.5 (1.0) km agreement. However, after applying a  $AOD_{0.68}$  threshold of 1.5 to isolate dust-dominated scenes, the AOC accuracy significantly improved, with a lower mean bias (0.4 km) and RMSE (0.8 km), as well as 61% (72%) of co-located retrievals within 0.5 (1.0) km agreement. In general, the higher the  $AOD_{0.68}$  threshold, the better the agreement between EPIC and CALIOP. The close agreement between EPIC and CALIOP, as well as between the two CALIOP channels, can be attributed to the well-defined upper boundary of the dust layer (as seen in Fig. 2(f)), such that the different techniques detected the strongest dust signal from the same altitude.

Deleted: <object><object><object>

Moved down [41]: . The main findings are as follows.

Moved up [40]:  
580

Deleted: .....Section Break (Next Page).....

Deleted:

Formatted: Font: 11 pt

Formatted: Not Expanded by / Condensed by

Formatted: Space Before: 5.5 pt

Formatted: Space Before: 0 pt

Deleted:

Deleted: <object>

595

600

605

610

615

To compare the EPIC AOC and IASI mean dust layer altitude retrievals, we selected the EPIC scans closest in time (06:50z and 09:01z on 28 May and 06:41z and 08:52z on 29 May) to IASI overpasses to minimize scene differences. We applied an AOD threshold of 1.5 to EPIC AOC data over land to exclude marginal-quality retrievals and enhance clarity. As illustrated in Fig. 17, the AOD threshold removes the majority of unrealistic AOC values over land. However, EPIC displays unrealistic retrievals over the northern Caspian Sea and coastal regions. This bias resembles the surface-related UVAI signal shown in Fig. 3, suggesting that the water-leaving radiances from turbid coastal waters may have been misinterpreted as aerosol signals in the EPIC AOC algorithm. Comparing EPIC retrievals on both days reveals no significant AOC changes between 06:50z and 09:01z on 28 May, likely due to a prevailing cold anticyclone that trapped dust within a shallow boundary layer. In contrast, the mean AOC increased by about 0.4 km between 06:41z and 08:52z on 29 May, likely due to the gradual lifting by an approaching cold front.

Figure 17 (c–d) suggests that IASI generally retrieves higher ALH than EPIC in dust-affected areas, both over deserts and the Caspian Sea. To further assess their consistency, we regridded the Level-2 data from both sensors onto a 1°×1° resolution and examined the ALH differences between co-located retrievals, shown in Fig. 18. EPIC and IASI have the poorest agreement at low aerosol loadings, exhibiting large dispersions. EPIC tends to produce much higher ALH under such conditions. As the aerosol loading increases, the EPIC–IASI difference decreases rapidly but stabilizes at AOD<sub>10</sub> ≥ 0.3 (Fig. 18(b)). Retrievals with AOD<sub>10</sub> ≥ 0.3 show a mean ALH difference of −0.4 km and an RMSD of 1.4 km, with 79% (85%) of co-located retrievals falling within 0.5 (1.0) km differences.

Finally, we used MINX to perform stereo height retrieval from MISR multi-angle images for two polygon regions located at the foothills of the Kopet-Dag Range (Fig. 19). The digitized pixels in both regions indicate an elevated dust layer extending from ~ 0.8 km above the ground to a sharp upper boundary at ~ 1.3 km, with a median plume height of 1.1 km. This estimate is ~ 1 km lower than the CALIOP-detected dust layer top (Fig. 2(f)), consistent with expected differences in the two techniques (Section 2.2.5). Overall, the MISR retrieval aligns with the CALIOP and EPIC products in detecting the dust layer height at the Kopet-Dag foothills (Fig. 16(a)).

#### 4 Conclusions

The Aralkum Desert has emerged as a prominent source of wind-blown saline dust in Central Asia. While satellite-based aerosol observations remain the most practical means for monitoring the region’s dust activity, they face considerable challenges due to the absence of in situ aerosol measurements near Aralkum and the area’s heterogeneous and highly variable surface characteristics. Although satellite algorithms are optimized for global performance, they may exhibit greater inconsistency and biases in Central Asia, where the region-specific aerosol and surface properties are inaccurately represented in the algorithms. To date, the performance and consistency of various observation techniques in characterizing the saline aerosol from Aralkum remain poorly understood. We address this knowledge gap through two key efforts.

First, we conducted a survey of current satellite techniques and algorithms capable of detecting the presence, column burden, and vertical height of airborne dust over Central Asia, focusing on four families of satellite aerosol products: UVAI, midvisible

Deleted: <object><object><object>

Formatted: Font: Times New Roman

Formatted: Space Before: 6.85 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 5.5 pt

Formatted: Space Before: 5.45 pt

Formatted: Space Before: 0.05 pt

Formatted: Space Before: 5.45 pt

Moved (insertion) [39]

Formatted: Space Before: 7.7 pt


Formatted: Indent: Left: 0.07", Hanging: 0.21", Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 3 + Alignment: Left + Aligned at: 0.07" + Indent at: 0.32"

Moved (insertion) [42]

Formatted: Space Before: 0 pt

Deleted: <object>

32




**Figure 18.** (a) Comparison of co-located ALH retrievals by EPIC and IASI. N, number of coincident retrievals. MD, mean difference. RMSD, root mean square difference. ED, percentage of co-located retrievals within expected differences of 0.5 and 1 km (shaded). (b) The EPIC-IASI difference as a function of IASI-retrieved AOD<sub>10</sub>.

625 AOD, thermal-infrared AOD, and ALH. The survey identifies the diverse methodologies employed by various algorithms and the differing a priori assumptions regarding surface properties and aerosol optical models (e.g., refractive index, particle shape, particle size distribution). In particular, the dust optical models predefined in these algorithms are primarily derived from AERONET measurements from West Africa and the Middle East. These models may deviate substantially from the

Deleted: <object><object><object>

Deleted: <object>



**Figure 19.** MINX-retrieved plume height profiles (middle and right panels) based on MISR multi-angle images on 28 May 2018 (orbit 98092), for two polygon areas denoted in the nadir-view RGB image (left panel). Surface elevation is represented in green, and retrieved plume heights in red.

microphysical properties of Aralkum-derived aerosols, which has been found to contain distinct chemical and mineralogical compositions (e.g., sulfate- and carbonate-rich minerals, low iron oxide content).

Currently, due to lack of in situ aerosol measurements (e.g., AERONET), the extent to which these dust optical models deviate from actual conditions in Central Asia has yet to be determined. The impact of inaccurate algorithm assumptions on the aerosol product performance is also not well understood. Addressing these knowledge gaps requires establishing in situ aerosol measurement capabilities near Aralkum, e.g., in the downwind Karakalpakstan region. Such measurements are essential not only for validating satellite products but also for developing new or refining existing aerosol optical models for Central Asia or similar environments.

Second, we performed a case study to assess the cross-sensor and cross-algorithm consistency in observing a saline dust storm originating from the Aralkum Desert during 27–29 May 2018. We analyzed a broad range of aerosol products, including UVAI from OMPS, TROPOMI and EPIC, midvisible AOD from MODIS, MISR, and VIIRS, thermal-infrared AOD from IASI using four different algorithms, and ALH from EPIC, CALIOP, IASI, and MISR. This synergistic, multisensor analysis allowed us to identify the complementary strengths of different techniques and uncover the potential limitations and biases of each product. The main findings are as follows.

The UVAI products show similar spatial patterns associated with the heavy dust plume on 27 May, but exhibit notable differences in magnitude and dynamic range. These discrepancies can be attributed, at least in part, to the choice of wavelength

Deleted: <object><object><object>

Moved (insertion) [43]

Formatted: Space Before: 0 pt

Moved (insertion) [44]

Formatted: Space Before: 0.05 pt

Formatted: Space Before: 0 pt

Moved (insertion) [45]

Moved (insertion) [41]

Deleted: ..... Column Break .....

Deleted: 2018

Deleted: reveal significant

Formatted: Not Expanded by / Condensed by

Formatted: Not Expanded by / Condensed by

Formatted: Not Expanded by / Condensed by

Deleted: The differences

Deleted: explained

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.55 pt

Deleted: by

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.5 pt

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.55 pt

Formatted: Condensed by 0.1 pt

Deleted: <object>

645 pair and the treatment of cloud scattering effect. Using the 95th percentile of each product as a threshold for dust detection, we find good agreement in delineating the dust plume extent. However, all UVAI products show large positive values over the northern Caspian Sea, Garabogazkol Gulf, and Sor Kaydak salt marsh. These dust-like signals are primarily caused by enhanced UV absorption by turbid and saline waters, which causes the water-leaving UV radiances to deviate from a pure Rayleigh scattering atmosphere, similar to the effect of absorbing aerosols. Consequently, this presents a significant limitation for using UVAI for dust detection and source mapping over Central Asia, due to interferences of turbid/salty waters, salt marshes, and saline deserts. Therefore, caution must be used to avoid misinterpreting surface effects as dust signals over ephemeral or dry lakes such as Aralkum.

650 Midvisible AOD retrievals over desert surfaces exhibit considerable inconsistencies across different sensors and algorithms. The choice of retrieval algorithms causes nonlinear relationships between the retrieved AOD from MODIS and VIIRS. Specifically, the MAIAC algorithm retrieves lower AOD than the DB algorithm under low aerosol loadings but higher AOD under high aerosol loadings. The EPS algorithm generally produces lower AOD than DB, with their divergence increasing as aerosol loading increases. In the operational EPS product, an urban aerosol optical model was selected for dust retrieval, leading to significantly lower AOD compared to other algorithms. A test retrieval using the dust optical model, however, shows significant improvement in the EPS agreement with other algorithms. The MISR operational product exhibits reduced data coverage and significant AOD underestimation compared to MODIS and VIIRS. In contrast, the MISR research algorithm produces higher, more consistent AOD retrievals as well as improved constraints on particle microphysical properties. Over the Caspian Sea, midvisible AOD retrievals show better overall agreement between algorithms, despite the DT and EPS algorithms failing to select the dust optical model during retrieval. Generally, the DT algorithm retrieves lower AOD under clean marine conditions but higher AOD under dust-laden conditions compared to MAIAC, DB, and EPS algorithms.

655 The high surface reflectivity and shallow or turbid coastal waters of Central Asia pose a great challenge for passive ALH retrieval from EPIC. The EPIC AOC product significantly overestimates aerosol heights under low aerosol loading or background conditions. However, over dust-laden scenes, EPIC shows good agreement with the CALIOP extinction-weighted height, with a mean bias of 0.4 km, an RMSE of 0.8 km, and 61% (72%) of co-located retrievals falling within 0.5 km (1.0 km) differences. At low aerosol loadings, EPIC AOC and IASI infrared-based mean dust height retrievals show considerable discrepancies, but their differences stabilize as aerosol loading increases. On average, IASI retrievals over dust-laden scenes are 0.4 km higher than those from EPIC. The MISR plume height retrieval, derived from parallax of multi-angle images, aligns well with EPIC and CALIOP in detecting dust height over the foothills of the Kopet-Dag Range.

665 The high surface reflectivity and shallow or turbid coastal waters of Central Asia pose a great challenge for passive ALH retrieval from EPIC. The EPIC AOC product significantly overestimates aerosol heights under low aerosol loading or background conditions. However, over dust-laden scenes, EPIC shows good agreement with the CALIOP extinction-weighted height, with a mean bias of 0.4 km, an RMSE of 0.8 km, and 61% (72%) of co-located retrievals falling within 0.5 km (1.0 km) differences. At low aerosol loadings, EPIC AOC and IASI infrared-based mean dust height retrievals show considerable discrepancies, but their differences stabilize as aerosol loading increases. On average, IASI retrievals over dust-laden scenes are 0.4 km higher than those from EPIC. The MISR plume height retrieval, derived from parallax of multi-angle images, aligns well with EPIC and CALIOP in detecting dust height over the foothills of the Kopet-Dag Range.

670 Data availability. Most of the data analyzed in the study are publicly available. The data repository links are provided in Table 1. The MISR research algorithm and plume height results will be available upon request. The regridded data products are available at <https://doi.org/10.5281/zenodo.13994593>.

Deleted: <object><object><object>

Deleted: the... threshold for dust detection, we find ... [722]

Moved (insertion) [46]

Deleted: Hence...consequently, this presents a signi ... [724]

Formatted ... [723]

Formatted ... [711]

Formatted ... [712]

Deleted: The

Deleted: products

Formatted ... [725]

Formatted ... [726]

Deleted: are subject to significant inconsistency between

Formatted ... [727]

Formatted ... [713]

Deleted: aerosol

Formatted ... [728]

Deleted: a

Formatted ... [729]

Deleted: response in

Formatted ... [730]

Deleted: Specifically

Formatted ... [731]

Deleted: yields

Formatted ... [732]

Deleted: burden,

Formatted ... [733]

Moved (insertion) [47]

Deleted: heavy

Deleted: burden

Deleted: yields

Deleted: under all aerosol burden conditions. The ... [737]

Formatted ... [734]

Formatted ... [735]

Formatted ... [736]

Formatted ... [738]

Deleted: aerosol burden. In addition, all AOD prod ... [739]

Formatted ... [740]

Deleted: in the VIIRS EPS product

Formatted ... [741]

Deleted: products. ... [742]

Formatted ... [714]

Moved (insertion) [48]

Formatted ... [715]

Moved (insertion) [49]

Formatted ... [716]

Formatted ... [717]

Deleted: Despite the

Deleted: which limits the

Deleted: AOC

Formatted ... [743]

Formatted ... [744]



Formatted ... [745]

Formatted ... [718]

Deleted: over Central Asia, we find that the EPIC A ... [746]



Author contributions. XX performed conceptualization, data curation, formal analysis, visualization, and writing of the initial draft. HL performed VIIRS EPS retrieval using dust model. JL and RK performed MISR research algorithm aerosols and climate health interview. All authors contributed to the result interpretation and edited the manuscript.  
Q Author(s) 2024. CC BY 4.0 License.



Competing interests. The authors declare no competing interests.

680 Acknowledgements. X, Xi acknowledges support from the NASA Land-Cover and Land-Use Change (LCLUC) Program and helpful discussions with Jason Tackett on CALIPSO data. The work of R. Kahn is supported in part by the NASA Terra-Aqua-NPP program, and the NASA Earth Observing System MISR project. The satellite data teams acknowledge funding support from multiple NASA and NOAA funding sources supporting the multi-decadal development and evaluation of these data records. The authors gratefully acknowledge the Copernicus Data Space Ecosystem and NASA ASDC, LAADS DAAC, and GES DISC for maintaining/distributing the data products used in this study.

685

Deleted: <object><object><object>

Deleted: designed the study and

Deleted: JW contributed to the writing of EPIC AOC algorithm. IL contributed to the writing of...L performed VIIRS EPS retrieval using dust model. JL and RK performed MISR research algorithm. CA and OT contributed to the UVAI analysis. AS, JL, RL, YW, AL, and IL contributed to AOD analysis. JW, ZL and OT contributed to the AOC analysis. [764]

Formatted [762]

Formatted [763]

Formatted: Indent: Left: 0.41", Space Before: 4.05 pt

Deleted: Section Break (Next Page)

Formatted: Justified, Space Before: 0.05 pt

Formatted: Font: 10 pt

Formatted: Space Before: 10.55 pt

Deleted: XX

Deleted: (grant no. 80NSSC20K1480).

Formatted [765]

Formatted: Not Expanded by / Condensed by

Moved (insertion) [52]

Formatted: Space Before: 5.35 pt

Formatted [766]

Deleted: devel- opment

Formatted: Space Before: 8.25 pt

Deleted:

Deleted: 625 LAADS DAAC, and GES DISC for maintaining/distributing the data products used in this study.

Deleted: <object>

36

## References

<https://doi.org/10.5194/egusphere-2024-3416>

Regman, D., Singer, A., and Tsao, H.: Erodibility of some crust forming soils/sediments from the Southern Aral Sea Basin as determined in a wind tunnel, Earth Surface Processes and Landforms, 31, 47–63, <https://doi.org/10.1002/esp.1230>, 2006.

Reuter, R.: Level 2 Aerosol parameters V003 [Dataset], [https://doi.org/10.5067/TERRA/MISR/MIL2ASAE\\_L2.003](https://doi.org/10.5067/TERRA/MISR/MIL2ASAE_L2.003), 1999.

ASDC: DSCOVR EPIC Aerosol Optical Centroid Height Version 1 [Dataset], [https://doi.org/10.5067/EPIC/DSCOVR/L2\\_AOCH.001](https://doi.org/10.5067/EPIC/DSCOVR/L2_AOCH.001), 2018.

ASDC: CALIPSO Lidar Level 2 Aerosol Profile, V4-51, [Dataset], [https://doi.org/10.5067/CALIP/CALIPSO/CAL\\_LID\\_L2\\_05kmAPro-Standard-V4-51](https://doi.org/10.5067/CALIP/CALIPSO/CAL_LID_L2_05kmAPro-Standard-V4-51), 2023.

Balkanski, Y., Schulz, M., Claquin, T., and Guibert, S.: Reevaluation of Mineral aerosol radiative forcings suggests a better agreement with satellite and AERONET data, Atmospheric Chemistry and Physics, 7, 81–95, <https://doi.org/10.5194/acp-7-81-2007>, 2007.

C3S CDS: Aerosol properties gridded data from 1995 to present derived from satellite observation [Dataset], <https://doi.org/10.24381/cds.239d815c>, 2019.

Callevaert, S., Vandenbussche, S., Kumps, N., Kylling, A., Shang, X., Komppula, M., Goloub, P., and De Mazière, M.: The mineral aerosol profiling from infrared radiances (MAPIR) algorithm: Version 4.1 description and evaluation, Atmospheric Measurement Techniques, 12, 3673–3698, <https://doi.org/10.5194/amt-12-3673-2019>, 2019.

Capelle, V., Chédin, A., Péquignot, E., Schlüssel, P., Newman, S. M., and Scott, N. A.: Infrared continental surface emissivity spectra and skin temperature retrieved from IASI observations over the tropics, Journal of Applied Meteorology and Climatology, 51, 1164–1179, <https://doi.org/10.1175/JAMC-D-11-0145.1>, 2012.

Capelle, V., Chédin, A., Siméon, M., Tsamalis, C., Pierangelo, C., Pondrom, M., Crevoisier, C., Crepeau, L., and Scott, N. A.: Evaluation of IASI-derived dust aerosol characteristics over the tropical belt, Atmospheric Chemistry and Physics, 14, 9343–9362, <https://doi.org/10.5194/acp-14-9343-2014>, 2014.

Capelle, V., Chédin, A., Pondrom, M., Crevoisier, C., Armante, R., Crepeau, L., and Scott, N. A.: Infrared dust aerosol optical depth retrieved daily from IASI and comparison with AERONET over the period 2007–2016, Remote Sensing of Environment, 206, 15–32, <https://doi.org/10.1016/j.rse.2017.12.008>, 2018.

Carr, J. L., Wu, D. L., Daniels, J., Friberg, M. D., Bresky, W., and Madani, H.: Geo-geo stereo-tracking of atmospheric motion vectors (AMVS) from the geostationary ring, Remote Sensing, 12, 1–47, <https://doi.org/10.3390/rs12223779>, 2020.

Clarisse, L., Clerbaux, C., Franco, B., Hadji-Lazaro, J., Whitburn, S., Kopp, A. K., Hurtmans, D., and Coheur, P. F.: A Decadal Data Set of Global Atmospheric Dust Retrieved From IASI Satellite Measurements, Journal of Geophysical Research: Atmospheres, 124, 1618–1647, <https://doi.org/10.1029/2018JD029701>, 2019.

de Graaf, M., Stammes, P., Torres, O., and Koelemeijer, R. B.: Absorbing Aerosol Index: Sensitivity analysis, application to GOME and comparison with TOMS, Journal of Geophysical Research D: Atmospheres, 110, 1–19, <https://doi.org/10.1029/2004JD005178>, 2005.

Diner, D. J., Beckert, J. C., Reilly, T. H., Bruegge, C. J., Conel, J. E., Kahn, R. A., Martonchik, I. V., Ackerman, T. P., Davies, R., Gerstel, S. A., Gordon, H. R., Muller, J. P., Myneni, R. B., Sellers, P. J., Pinty, B., and Verstraete, M. M.: Multi-angle imaging spectroradiometer (MISR) instrument description and experiment overview, IEEE Transactions on Geoscience and Remote Sensing, 36, 1072–1087, <https://doi.org/10.1109/36.700992>, 1998.

Dubovik, O., Holben, B., Eck, T. F., Smirnov, A., Kaufman, Y. J., King, M. D., Tanré, D., and Slutsker, I.: Variability of Absorption and Optical Properties of Key Aerosol Types Observed in Worldwide Locations, Journal of the Atmospheric Sciences, 59, 590–608, [https://doi.org/10.1175/1520-0469\(2002\)059<0590:vooaop>2.0.co;2](https://doi.org/10.1175/1520-0469(2002)059<0590:vooaop>2.0.co;2), 2002.

Deleted: <object><object><object>

Formatted ... [768]

Formatted ... [767]

Formatted ... [769]

Moved up [43]: 630

Moved up [46]:

Moved down [55]: G.

Deleted:

Deleted: Column Break

Deleted: Y.: Saline dust storms and their ecological

Formatted ... [777]

Formatted ... [770]

Deleted: org/10.5067/EPIC/DSCOVR/L2\_AOCH.001,

Formatted ... [779]

Moved (insertion) [53]

Deleted: Data set],

Formatted ... [780]

Moved (insertion) [56]

Formatted ... [782]

Moved (insertion) [57]

Formatted ... [784]

Moved (insertion) [54]

Formatted ... [771]

Moved (insertion) [59]

Deleted: Standard-V4-51, 2023.

Formatted ... [772]

Formatted ... [785]

Deleted: org/10.3390/rs12223779,

Moved (insertion) [60]

Deleted: Castellanos, P., Colarco, P., Espinosa, W.

Formatted ... [786]

Moved down [61]: H.,

Moved down [62]: J.,

Moved down [63]: R.,

Moved up [58]: M.,

Deleted: Nowotnick, E. P., Rocha-Lima, A., Smith

Deleted: Xu, X., Zhou, M., Zhang, H., Castro Garza

Deleted: Janz, S. J., Yorks, J., McGill, M., Reid, J.

Deleted: and Kondragunta, S.: First retrieval of

Formatted ... [788]

Formatted ... [790]

Formatted ... [792]

Formatted ... [794]

Moved (insertion) [64]

Moved (insertion) [62]

Moved (insertion) [65]

Moved (insertion) [66]

Formatted ... [797]

Formatted ... [798]


Formatted ... [799]

Formatted ... [800]

Moved (insertion) [67]

Deleted: Di Biagio, C., Formenti, P., Balkanski, Y.

Formatted ... [802]



	Dubovik, O., Sinyuk, A., Lapyonok, T., Holben, B. N., Mishchenko, M., Yang, P., Eck, T. F., Volten, H., Muñoz, O., Veihelmann, B., van der Zande, W. J., Leon, J. F., Sorokin, M., and Slutsker, I.: Application of spheroid models to account for aerosol particle nonsphericity in remote sensing of desert dust, <i>Journal of Geophysical Research Atmospheres</i> , 111, <a href="https://doi.org/10.1029/2006JD006618">https://doi.org/10.1029/2006JD006618</a> , 2006.	Deleted: <object><object><object>
725	ESA-TROPOMI Level 2 Ultraviolet Aerosol Index products Version 02 [Data set], <a href="https://doi.org/10.5270/SSP-3dgd66p">https://doi.org/10.5270/SSP-3dgd66p</a> , 2021.	Moved (insertion) [68]
	Lawson, T. J., and Kahn, R. A.: Assessing the altitude and dispersion of volcanic plumes using MISR multi-angle imaging from space: Sixteen years of volcanic activity in the Kamchatka Peninsula, Russia, <i>Journal of Volcanology and Geothermal Research</i> , 337, 1–15, <a href="https://doi.org/10.1016/j.jvolgeores.2017.03.010">https://doi.org/10.1016/j.jvolgeores.2017.03.010</a> , 2017.	Formatted ... [806]
730	Garay, M. J., Witek, M. L., Kahn, R. A., Seidel, F. C., Limbacher, J. A., Bull, M. A., Diner, D. J., Hansen, E. G. G., Kalashnikova, O. V., Lee, H., Nastan, A. M., and Yu, Y.: Introducing the 4.4km spatial resolution Multi-Angle Imaging SpectroRadiometer (MISR) aerosol product, <i>Atmospheric Measurement Techniques</i> , 13, 593–628, <a href="https://doi.org/10.5194/amt-13-593-2020">https://doi.org/10.5194/amt-13-593-2020</a> , 2020.	Formatted ... [807]
	Groll, M., Opp, C., Issanova, G., Vereshagina, N., and Semenov, O.: Physical and chemical characterization of dust deposited in the Turan Lowland (Central Asia), in: E3S Web of Conferences, vol. 99, ISSN 22671242, <a href="https://doi.org/10.1051/e3sconf/20199903005">https://doi.org/10.1051/e3sconf/20199903005</a> , 2019.	Deleted: org/10.1029/2005JD006619,
735	He, X., Bai, Y., Pan, D., Tang, J., and Wang, D.: Atmospheric correction of satellite ocean color imagery using the ultraviolet wavelength for highly turbid waters, <i>Optics Express</i> , 20, 20 754, <a href="https://doi.org/10.1364/oe.20.020754">https://doi.org/10.1364/oe.20.020754</a> , 2012.	Formatted ... [808]
	Herman, J. R., Bhartia, P. K., Torres, O., Hsu, C., Sefior, C., and Celarier, E.: Global distribution of UV-absorbing aerosols from Nimbus 7/TOMS data, <i>Journal of Geophysical Research Atmospheres</i> , 102, 16 911–16 922, <a href="https://doi.org/10.1029/96jd03680">https://doi.org/10.1029/96jd03680</a> , 1997.	Deleted: org/10.5270/SSP-3dgd66p,
740	Hofer, J., Althausen, D., Abdullaev, S. F., Makhmudov, A. N., Nazarov, B. I., Schettler, G., Engelmann, R., Baars, H., Fomba, K. W., Müller, K., Heinold, B., Kandler, K., and Ansmann, A.: Long-term profiling of mineral dust and pollution aerosol with multiwavelength polarization Raman lidar at the Central Asian site of Dushanbe, Tajikistan: Case studies, <i>Atmospheric Chemistry and Physics</i> , 17, 14 559–14 577, <a href="https://doi.org/10.5194/acp-17-14559-2017">https://doi.org/10.5194/acp-17-14559-2017</a> , 2017.	Formatted ... [819]
	Hsu, C.: VIIRS/NOAA20 Deep Blue Aerosol L2 6-Min Swath 6 km [Data set], <a href="https://doi.org/10.5067/VIIRS/AERDB_L2_VIIRS_NOAA20.002">https://doi.org/10.5067/VIIRS/AERDB_L2_VIIRS_NOAA20.002</a> , 2022.	Moved (insertion) [55]
745	Hsu, N. C., Tsay, S. C., King, M. D., and Herman, J. R.: Aerosol properties over bright-reflecting source regions, <i>IEEE Transactions on Geoscience and Remote Sensing</i> , 42, 557–569, <a href="https://doi.org/10.1109/TGRS.2004.824067">https://doi.org/10.1109/TGRS.2004.824067</a> , 2004.	Moved (insertion) [70]
	Hsu, N. C., Jeong, M. J., Bettenhausen, C., Sayer, A. M., Hansell, R., Sefior, C. S., Huang, J., and Tsay, S. C.: Enhanced Deep Blue aerosol retrieval algorithm: The second generation, <i>Journal of Geophysical Research Atmospheres</i> , 118, 9296–9315, <a href="https://doi.org/10.1002/jgrd.50712">https://doi.org/10.1002/jgrd.50712</a> , 2013.	Moved (insertion) [71]
750	Hsu, N. C., Lee, J., Sayer, A. M., Kim, W., Bettenhausen, C., and Tsay, S. C.: VIIRS Deep Blue Aerosol Products Over Land: Extending the EOS Long-Term Aerosol Data Records, <i>Journal of Geophysical Research: Atmospheres</i> , 124, 4026–4053, <a href="https://doi.org/10.1029/2018JD029688">https://doi.org/10.1029/2018JD029688</a> , 2019.	Formatted ... [820]
	Jackson, J. M., Liu, H., Laszlo, I., Kondragunta, S., Remer, L. A., Huang, J., and Huang, H. C.: Suomi-NPP VIIRS aerosol algorithms and data products, <i>Journal of Geophysical Research Atmospheres</i> , 118, 673–12, <a href="https://doi.org/10.1002/2013JD020449">https://doi.org/10.1002/2013JD020449</a> , 2013.	Formatted ... [821]
755	Jiang, H., Huang, J., Li, L., Huang, L., Manzoor, M., Yang, J., Wu, G., Sun, X., Wang, B., Egamberdieva, D., Panosyan, H., Birkeland, N. K., Zhu, Z., and Li, W.: Onshore soil microbes and endophytes respond differently to geochemical and mineralogical changes in the Aral Sea, <i>Science of the Total Environment</i> , 765, <a href="https://doi.org/10.1016/j.scitotenv.2020.142675">https://doi.org/10.1016/j.scitotenv.2020.142675</a> , 2021.	Formatted ... [822]
	Kahn, R. A., Gaitley, B. J., Garay, M. J., Diner, D. J., Eck, T. F., Smirnov, A., and Holben, B. N.: Multiangle Imaging SpectroRadiometer global aerosol product assessment by comparison with the Aerosol Robotic Network, <i>Journal of Geophysical Research Atmospheres</i> , 115, <a href="https://doi.org/10.1029/2010JD014601">https://doi.org/10.1029/2010JD014601</a> , 2010.	Formatted ... [810]
760		Moved (insertion) [72]
		Deleted: Groll, M., Opp, C., and Aslanov, I.: Spati...
		Formatted ... [823]
		Formatted ... [826]
		Formatted ... [825]
		Deleted: org/10.1051/e3sconf/20199903005,
		Formatted ... [811]
		Moved up [49]: ¶
		Moved up [50]: 670¶
		Deleted: -----Section Break (Next Page)-----
		Deleted: ¶
		Formatted ... [830]
		Formatted ... [828]
		Moved up [52]: 680¶
		Deleted: ¶
		Formatted ... [831]
		Moved up [53]: 690¶
		Deleted: -----Column Break-----
		Formatted ... [832]
		Moved (insertion) [69]
		Formatted ... [812]
		Deleted: Herman, J. R. and Celarier, E. A.: Earth
		Moved (insertion) [73]
		Deleted: Holben, B. N., Eck, T. F., Slutsker, I., Tar
		Formatted ... [836]
		Formatted ... [835]
		Formatted ... [813]
		Formatted ... [814]
		Moved up [65]: A.,
		Moved up [67]: . A.,
		Moved up [73]: F.,
		Moved down [75]: I., and
		Deleted: Vermote, E., Reagan, J
		Deleted: Kaufman, Y. J., Nakajima, T., Lavenu,
		Deleted: Jankowiak,
		Deleted: Smirnov, A.: AERONET - A federated
		Formatted ... [839]
		Formatted ... [837]
		Formatted ... [838]
		Formatted ... [815]
		Deleted: Hsu, N. C.: Detection of biomass burning
		Formatted ... [840]
		Deleted: org/10.1109/TGRS.2004.824067
		Formatted ... [841]

765

Kim, M. H., Omar, A. H., Tackett, J. L., Vaughan, M. A., Winker, D. M., Trepte, C. R., Hu, Y., Liu, Z., Poole, L. R., Pitts, M. C., Kar, J., and Magill, B. E. The CALIPSO version 4 automated aerosol classification and lidar ratio selection algorithm. *Atmospheric Measurement Techniques*, 11, 6107–6135, <https://doi.org/10.5194/amt-11-6107-2018>, 2018.

770

Laszlo, I.: Remote Sensing of Tropospheric Aerosol Optical Depth From Multispectral Monodirectional Space-Based Observations, in: Comprehensive Remote Sensing, edited by Liang, S. B. T. C. R. S., pp. 137–196, Elsevier, Oxford, ISBN 978-0-12-803221-3, <https://doi.org/https://doi.org/10.1016/B978-0-12-409548-9.10389-6>, 2018.

775

Laszlo, I. and Liu, H.: EPS Aerosol Optical Depth (AOD) Algorithm Theoretical Basis Document Version 3.4, Tech. rep., NOAA NESDIS CENTER for SATELLITE APPLICATIONS and RESEARCH, [https://www.star.nesdis.noaa.gov/jps/documents/ATBD/ATBD\\_EPS\\_Aerosol\\_AOD\\_v3.4.pdf](https://www.star.nesdis.noaa.gov/jps/documents/ATBD/ATBD_EPS_Aerosol_AOD_v3.4.pdf), 2022.

780

Lee, J., Hsu, N. C., Sayer, A. M., Bettenhausen, C., and Yang, P.: AERONET-Based Nonspherical Dust Optical Models and Effects on the VIIRS Deep Blue/SOAR Over Water Aerosol Product, *Journal of Geophysical Research: Atmospheres*, 122, 10 384–10 401, <https://doi.org/10.1002/2017JD027258>, 2017.

785

Lee, J., Hsu, N. C., Kim, W. V., Sayer, A. M., and Tsay, S.-C.: VIIRS Version 2 Deep Blue Aerosol Products, *Journal of Geophysical Research: Atmospheres*, 129, e2023JD040 082, <https://doi.org/https://doi.org/10.1029/2023JD040082>, 2024.

790

Lee, Z., Hu, C., Shang, S., Du, K., Lewis, M., Arnone, R., and Brewin, R.: Penetration of UV-visible solar radiation in the global oceans: Insights from ocean color remote sensing, *Journal of Geophysical Research: Oceans*, 118, 4241–4255, <https://doi.org/10.1002/jgrc.20308>, 2013.

795

Legrand, M., Plana-Fattori, A., and N'Doumé, C.: Satellite detection of dust using the IR imagery of Meteosat 1. Infrared difference dust index, *Journal of Geophysical Research Atmospheres*, 106, 18 251–18 274, <https://doi.org/10.1029/2000JD900749>, 2001.

800

Lensky, I. M. and Rosenfeld, D.: Clouds-Aerosols-Precipitation Satellite Analysis Tool (CAPSAT), *Atmospheric Chemistry and Physics*, 8, 6739–6753, <https://doi.org/10.5194/acp-8-6739-2008>, 2008.

805

Levy, R. C. and Hsu, N. C.: MODIS Atmosphere L2 Aerosol Product [Data set], [https://doi.org/10.5067/MODIS/MYD04\\_L2.006](https://doi.org/10.5067/MODIS/MYD04_L2.006), 2015a.

810

Levy, R. C. and Hsu, N. C.: MODIS Atmosphere L2 Aerosol Product [Dataset], [https://doi.org/10.5067/MODIS/MOD04\\_3K.061](https://doi.org/10.5067/MODIS/MOD04_3K.061), 2015b.

815

Levy, R. C., Mattoo, S., Sawyer, V., and Munchak, L.: VIIRS/NOAA20 Dark Target Aerosol 6-Min L2 Swath 6 km [Dataset], [https://doi.org/10.5067/VIIRS/AERDT\\_L2\\_VIIRS\\_NOAA20.002](https://doi.org/10.5067/VIIRS/AERDT_L2_VIIRS_NOAA20.002), 2023.

820

Limbacher, J. A., Kahn, R. A., and Lee, J.: The new MISR research aerosol retrieval algorithm: A multi-Angle, multi-spectral, bounded-variable least squares retrieval of aerosol particle properties over both land and water, *Atmospheric Measurement Techniques*, 15, 6865–6887, <https://doi.org/10.5194/amt-15-6865-2022>, 2022.

825

Lu, Z., Wang, J., Xu, X., Chen, X., Kondragunta, S., Torres, O., Wilcox, E. M., and Zeng, J.: Hourly Mapping of the Layer Height of Thick Smoke Plumes Over the Western U.S. in 2020 Severe Fire Season, *Frontiers in Remote Sensing*, 2, <https://doi.org/10.3389/frsen.2021.766628>, 2021.

Deleted: <object><object><object>

Moved up [77]: J.,

Moved (insertion) [80]

Deleted: Wald, A. E., Remer, L. A., Gao, B. C., Li ... [867]

Formatted ... [856]

Formatted ... [866]

Formatted ... [857]

Deleted: Section Break (Next Page) ... [868]

Moved up [54]: 700

Deleted: ... [869]

Moved up [68]:

Deleted: Column Break ... [871]

Formatted ... [870]

Formatted ... [858]

Formatted ... [859]

Moved up [66]: V.,

Moved up [78]: J.,

Deleted: Kokhanovsky, A. A., Davis, A. B., Cairn ... [872]

Deleted: Litvinov, P., Lapyonok, T., Kolomiets, I. ... [874]

Deleted: Xu, F., Natraj, V., Duan, M., Cheng, T., a ... [876]

Formatted ... [873]

Formatted ... [875]

Formatted ... [860]

Formatted ... [877]

Moved (insertion) [81]

Formatted ... [861]

Deleted: <https://www.star.nesdis.noaa.gov/jps/do> ... [878]

Formatted ... [862]

Deleted: Aerosol\_AOD\_v3.4.pdf,

Formatted ... [879]

Deleted: org/10.1002/2017JD027258,

Deleted: org/10.1002/jgrc.20308,

Deleted: org/10.5067/MODIS/MYD04\_L2.006,

Formatted ... [880]

Deleted: org/10.5067/MODIS/MOD04\_3K.061,

Deleted: Levy, R.

Moved up [64]: C.,

Deleted: Remer, L. A., and Dubovik, O.: Global aer ... [881]

Deleted: Li, J., Carlson, B. E., and Lacis, A. A.: H ... [882]

Formatted ... [863]

Formatted ... [864]

Deleted: Kaufman, Y.

Formatted ... [865]

Deleted: org/10.3389/frsen.2021.766628,

Deleted: <object>

39





835

Rupakheti, D., Rupakheti, M., Abdullaev, S. F., Yin, X., and Kang, S.: Columnar aerosol properties and radiative effects over Dushanbe, Tajikistan in Central Asia, *Environmental Pollution*, 265, <https://doi.org/10.1016/j.envpol.2020.114872>, 2020.

Sawyer, V., Levy, R. C., Mattoo, S., Cleton, G., Shi, Y., and Remer, L. A.: Continuing the MODIS dark target aerosol time series with VIIRS, *Remote Sensing*, 12, <https://doi.org/10.3390/rs12020308>, 2020.

Yuan, M., Hsu, N. C., Lee, J., Bettenhausen, C., Kim, W. V., and Smimov, A.: Satellite Ocean Aerosol Retrieval (SOAR) Algorithm Extension to S-NPP VIIRS as Part of the “Deep Blue” Aerosol Project, *Journal of Geophysical Research: Atmospheres*, 123, 380–400, <https://doi.org/10.1002/2017JD027412>, 2018.

Schepanski, K., Tegen, I., and Macke, A.: Comparison of satellite based observations of Saharan dust source areas, *Remote Sensing of Environment*, 123, 90–97, <https://doi.org/10.1016/j.rse.2012.03.019>, 2012.

Semenov, V. K., Smimov, A., Arefev, V. N., Sinyakov, V. P., Sorokina, L. I., and Ignatova, N. I.: Aerosol optical depth over the mountainous region in central Asia (Issyk-Kul Lake, Kyrgyzstan), *Geophysical Research Letters*, 32, 1–5, <https://doi.org/10.1029/2004GL021746>, 2005.

Sokolik, I. N. and Toon, O. B.: Incorporation of mineralogical composition into models of the radiative properties of mineral aerosol from UV to IR wavelengths, *Journal of Geophysical Research Atmospheres*, 104, 9423–9444, <https://doi.org/10.1029/1998JD200048>, 1999.

Stein Zweers, D. C.: TROPOMI ATBD of the UV aerosol index document number - SSP-KNMI-L2-0008-RP, Tech. Rep. 2.1.0, Royal Netherlands Meteorological Institute, De Bilt, 3731 GA, the Netherlands, <https://sentinel.esa.int/documents/247904/2476257/Sentinel-5P-TROPOMI-ATBD-UV-Aerosol-Index.pdf>, 2022.

Tanré, D., Kaufman, Y. J., Herman, M., and Mattoo, S.: Remote sensing of aerosol properties over oceans using the MODIS/EOS spectral radiances, *Journal of Geophysical Research Atmospheres*, 102, 16 971–16 988, <https://doi.org/10.1029/96jd03437>, 1997.

Torres, O.: OMPS-NPP L2 NM Aerosol Index swath orbital V2 [Data set], <https://doi.org/10.5067/40L92G8144IV>, 2019a.

Torres, O.: DSCOVR EPIC Level 2 UV Aerosol Version 3 [Data set], [https://doi.org/10.5067/EPIC/DSCOVR/L2\\_AER\\_03](https://doi.org/10.5067/EPIC/DSCOVR/L2_AER_03), 2019b.

Torres, O.: TROPOMI/Sentinel-5P Near UV Aerosol Optical Depth and Single Scattering Albedo L2 1-Orbit Snapshot 7.5 km x 3 km [Data set], <https://doi.org/10.5067/MEASURES/AER/DATA204>, 2021.

Torres, O., Bhartia, P. K., Herman, J. R., Ahmad, Z., and Gleason, J.: Derivation of aerosol properties from satellite measurements of backscattered ultraviolet radiation: Theoretical basis, *Journal of Geophysical Research Atmospheres*, 103, 17 099–17 110, <https://doi.org/10.1029/98JD00900>, 1998.

Torres, O., Ahn, C., and Chen, Z.: Improvements to the OMI near-UV aerosol algorithm using A-train CALIOP and AIRS observations, *Atmospheric Measurement Techniques*, 6, 3257–3270, <https://doi.org/10.5194/amt-6-3257-2013>, 2013.

Torres, O., Bhartia, P. K., Jethva, H., and Ahn, C.: Impact of the ozone monitoring instrument row anomaly on the long-term record of aerosol products, *Atmospheric Measurement Techniques*, 11, 2701–2715, <https://doi.org/10.5194/amt-11-2701-2018>, 2018.

Veeffkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de Haan, J. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sensing of Environment*, 120, 70–83, <https://doi.org/10.1016/j.rse.2011.09.027>, 2012.

Volz, F. E.: Infrared Optical Constants of Ammonium Sulfate, Sahara Dust, Volcanic Pumice, and Flyash, *Applied Optics*, 12, 564, <https://doi.org/10.1364/ao.12.000564>, 1973.

840

845

850

855

860

865

870

Deleted: <object><object><object>

Moved (insertion) [86]

Formatted: Space Before: 4.5 pt

Deleted: org/10.1016/j.envpol.2020.114872,

Moved up [56]: ., Schulz, M.,

Deleted: Samset, B. H., Stjern, C. W., Andrews, E., Kahn, R. A., Myhre, G

Deleted: and Schuster, G. L.: Aerosol Absorption: (... [915])

Formatted (... [914])

Deleted: org/10.1002/2017JD027412,

Deleted: org/10.1016/j.rse.2012.03.019,

Moved (insertion) [85]

Formatted (... [916])

Moved (insertion) [75]

Deleted: Shukurov, K. A., Simonenkov, D.

Formatted (... [917])

Moved up [84]: V.,

Moved up [83]: ¶

Moved up [86]: 835¶

Deleted: Nevzorov, A. V., Rashki, A., Hamzeh, N. (... [919])

Deleted: ¶ (... [920])

Deleted: .....Column Break..... (... [921])

Formatted (... [918])

Formatted: Font: Arial

Moved (insertion) [87]

Formatted: Font: Arial

Formatted: Space Before: 8.25 pt

Deleted: https://sentinel.esa.int/documents/247904/ (... [922])

Formatted: Space Before: 0.05 pt

Deleted: Sun, J., Veeffkind, P., Nanda, S., Van Veldt (... [923])

Deleted: Torres, O.: OMI/Aura Near UV Aerosol (... [924])

Formatted: Space Before: 8.25 pt

Deleted: org/10.5067/EPIC/DSCOVR/L2\_AER\_03,

Deleted: org/10.5067/MEASURES/AER/DATA204,

Deleted: org/10.1029/98JD00900,

Deleted: Torres, O., Jethva, H., and Bhartia, P. K.: (... [925])

Deleted: org/10.5194/amt-11-2701-2018,

Deleted: org/10.1016/j.rse.2011.09.027,

Deleted: <object>

41

Winker, D. M., Vaughan, M. A., Omar, A., Hu, Y., Powell, K. A., Liu, Z., Hunt, W. H., and Young, S. A.: Overview of the CALIPSO mission and CALIOP data processing algorithms, Journal of Atmospheric and Oceanic Technology, 26, 2310–2323, <https://doi.org/10.1175/2009JTECHA1281.1>, 2009.

Wink, T., Hasekamp, O., van Diedenhoven, B., Cairns, B., Yorks, J. E., and Chowdhary, J.: Passive remote sensing of aerosol layer height using multiangle polarization measurements, Geophysical Research Letters, 43, 8783–8790, <https://doi.org/10.1002/2016GL069848>, 2016.

Xi, X.: On the Geomorphic, Meteorological, and Hydroclimatic Drivers of the Unusual 2018 Early Summer Salt Dust Storms in Central Asia, Journal of Geophysical Research: Atmospheres, 128, <https://doi.org/10.1029/2022JD038089>, 2023.

Xi, X. and Sokolik, I. N.: Seasonal dynamics of threshold friction velocity and dust emission in Central Asia, Journal of Geophysical Research: Atmospheres, 120, 1536–1564, <https://doi.org/10.1002/2014JD022471>, 2015a.

Xi, X. and Sokolik, I. N.: Dust interannual variability and trend in Central Asia from 2000 to 2014 and their climatic linkages, Journal of Geophysical Research: Atmospheres, 120, 12 175–12 197, <https://doi.org/10.1002/2015JD024092>, 2015b.

Xi, X. and Sokolik, I. N.: Quantifying the anthropogenic dust emission from agricultural land use and desiccation of the Aral Sea in Central Asia, Journal of Geophysical Research: Atmospheres, 121, 270–12, <https://doi.org/10.1002/2016JD025556>, 2016.

Xu, X., Wang, J., Wang, Y., Zeng, J., Torres, O., Yang, Y., Marshak, A., Reid, J., and Miller, S.: Passive remote sensing of altitude and optical depth of dust plumes using the oxygen A and B bands: First results from EPIC/DSCOVER at Lagrange-1 point, Geophysical Research Letters, 44, 7544–7554, <https://doi.org/10.1002/2017GL073939>, 2017.

Xu, X., Wang, J., Wang, Y., Zeng, J., Torres, O., Reid, J. S., Miller, S. D., Vanderlei Martins, J., and Remer, L. A.: Detecting layer height of smoke aerosols over vegetated land and water surfaces via oxygen absorption bands: Hourly results from EPIC/DSCOVER in deep space, Atmospheric Measurement Techniques, 12, 3269–3288, <https://doi.org/10.5194/amt-12-3269-2019>, 2019.

Young, S. A., Vaughan, M. A., Garnier, A., Tackett, J. L., Lambeth, J. D., and Powell, K. A.: Extinction and optical depth retrievals for CALIPSO's Version 4 data release, Atmospheric Measurement Techniques, 11, 5701–5727, <https://doi.org/10.5194/amt-11-5701-2018>, 2018.

Zhang, H., Kondragunta, S., Laszlo, I., Liu, H., Remer, L. A., Huang, J., Superczynski, S., and Ciren, P.: An enhanced VIIRS aerosol optical thickness (AOT) retrieval algorithm over land using a global surface reflectance ratio database, Journal of Geophysical Research, 121, 717–10, <https://doi.org/10.1002/2016JD024859>, 2016.

Zhou, D. K., Larar, A. M., Liu, X., Smith, W. L., Strow, L. L., Yang, P., Schlüssel, P., and Calbet, X.: Global land surface emissivity retrieved from satellite ultraspectral IR measurements, IEEE Transactions on Geoscience and Remote Sensing, 49, 1277–1290, <https://doi.org/10.1109/TGRS.2010.2051036>, 2011.

Zhou, Y., Levy, R. C., Remer, L. A., Mattoo, S., and Espinosa, W. R.: Dust Aerosol Retrieval Over the Oceans With the MODIS/VIIRS Dark Target Algorithm: 2. Nonspherical Dust Model, Earth and Space Science, 7, <https://doi.org/10.1029/2020EA001222>, 2020a.

Zhou, Y., Levy, R. C., Remer, L. A., Mattoo, S., Shi, Y., and Wang, C.: Dust Aerosol Retrieval Over the Oceans With the MODIS/VIIRS Dark-Target Algorithm: 1. Dust Detection, Earth and Space Science, 7, <https://doi.org/10.1029/2020EA001221>, 2020b.

Deleted: <object><object><object>

Moved (insertion) [88]

Formatted: Font: Times New Roman

Formatted: Top: 0.35", Section start: New page, Height: 9.45", Header distance from edge: 0"

Deleted: org/10.1175/2009JTECHA1281.1,

Formatted: Space Before: 0.1 pt

Formatted: Space Before: 7.2 pt

Deleted: org/10.1002/2016GL069848,

Deleted: org/10.1029/2022JD038089,

Deleted: org/10.1002/2014JD022471,

Moved up [87]: ¶  
850¶  
¶  
¶  
¶  
¶  
855¶  
¶  
¶  
¶  
860¶  
¶

Moved up [88]: ¶

Deleted: .....Section Break (Next Page)..... ([926])

Deleted: .....Column Break.....

Formatted: Font: Arial

Formatted: Font: Times New Roman

Deleted: org/10.1002/2015JD024092,

Deleted: org/10.1002/2016JD025556,

Deleted: org/10.1002/2017GL073939,

Deleted: org/10.5194/amt-12-3269-2019,

Deleted: Yorks, J. E., McGill, M. J., Palm, S. P., HI (... [927])

Deleted: org/10.5194/amt-11-5701-2018,

Deleted: Yu, H., Yang, Y., Wang, H., Tan, Q., Chi (... [928])

Deleted: org/10.1002/2016JD024859,

Deleted: org/10.1029/2020EA001222,

Deleted: org/10.1029/2020EA001221, 2020b.

Deleted: <object>