

Effect of double diffusion processes in the deep ocean on the distribution and dynamics of particulate and dissolved matter: a case study in Tyrrhenian Sea

Xavier DURRIEU DE MADRON^{1*}, Paul BLIN¹, Mireille PUJO-PAY², Vincent TAILLANDIER³, Pascal CONAN^{2,4}

⁽¹⁾ CEFREM, CNRS-Université de Perpignan Via Domitia, Perpignan, France

⁽²⁾ LOMIC, CNRS-Sorbonne Université, Banyuls/Mer, France

⁽³⁾ LOV, CNRS-Sorbonne Université, Villefranche/Mer, France


⁽⁴⁾ OSU STAMAR, CNRS-Sorbonne Université, Paris, France

Correspondence to : Xavier Durrieu de Madron (demadron@univ-perp.fr)

Abstract: This study examines CTD, ADCP and optical data collected during the PERLE-3 cruise in March 2020 between the surface and 2000 m depth over an east-west section of the Tyrrhenian Sea in the Mediterranean. The focus is on the impact of double diffusion processes, in particular salt fingering, on the distribution and dynamics of particulate and dissolved matter. The staircases develop at the interface between the warm, saline Eastern Intermediate Water (EIW) and the colder, less saline Tyrrhenian Deep Water (TDW) in the centre of the basin with low hydrodynamic energy. The results show that thermohaline staircases formed by salt fingering significantly influence particle sedimentation and biogeochemical cycling in deep ocean environments by altering vertical flux patterns. These density steps create distinct vertical layers that act as physical barriers, slowing the descent of particles and facilitating their retention and aggregation. Retention of fine particles at density gradients promotes the formation of larger aggregates, thereby affecting particle size distribution. The staircases also affect dissolved matter by creating pronounced concentration gradients of oxygen and nutrients, which can influence microbial activity and nutrient fluxes.

Keywords: Mediterranean, Tyrrhenian Sea, Double diffusion, Salt fingering, Suspended particulate matter, Particle size distribution, Nitrate, Mineralization

Graphical abstract:



1. Introduction

Gravitational settling of biologically derived particles, produced by marine plankton in the surface layer, is the primary driver of vertical fluxes of material in the deep ocean (Newton and Liss, 1990). In general, vertical mass fluxes of particles in the ocean decrease exponentially with depth due to particle degradation and the increasing density of seawater (Omand et al., 2020). Marine particles have a wide range of sizes, from micrometres (clays) to centimetres (organic detritus). Particles generally tend to agglomerate to form aggregates of different sizes, shapes, densities, and characteristics (Kiko et al., 2022). The change in geometry of the aggregates as well as their excess density compared to seawater are the factors that determine their sedimentation rate.

The water column in the ocean is composed by the superposition of water masses with generally distinct thermohaline characteristics and often distant origins. Large density interfaces, associated with strong vertical gradients in temperature ($\sim 0.25\text{ }^{\circ}\text{C m}^{-1}$) at seasonal thermoclines, or salinity ($\sim 0.5\text{ PSU m}^{-1}$) at river plumes can occur in the upper ocean, resulting in the retention of the less dense particles. In the ocean temperature and salinity gradients are much weaker, and the effect of particle size becomes greater than the effect of density excess, so that settling velocity generally increases with particle size. Thus settling velocities in the open ocean are on the order of $5 \times 10^{-4}\text{ mm s}^{-1}$ to 1 mm s^{-1} for particles ranging from $2\text{ }\mu\text{m}$ to $300\text{ }\mu\text{m}$ (McCave, 1975).

However, in the deep ocean, in the transition zones between two water masses, areas of enhanced density gradients can occur due to double diffusion processes, in particular those associated with salt fingers. Salt fingering is a common process in the ocean (Kunze, 2003; Radko et al., 2014). It requires a stably stratified water column, where a layer of warm, saline

water overlies a layer of cooler, less saline water. Because molecular diffusion of temperature is about 100 times faster than that of salinity, the salinity interface initially remains essentially unchanged as temperature exchange occurs across it. As a result, the saltier water above this thin mixed layer of average temperature becomes denser than the less saline water below. This leads to salt finger instabilities at the interface: the saltier water sinks and the less saline water rises. The mixing process is then repeated at the new interfaces, resulting in a progressive thickening of the mixed layer and an increase in the vertical density gradients on either side of the interface. If favourable conditions persist, step structures can develop and reach thicknesses of several tens of meters. These staircases exhibit temperature and salinity steps ranging between 0.001 and 0.019 °C and 0.001 and 0.2 g kg⁻¹, respectively (Van Der Boog et al., 2021).

Various experimental and numerical studies have explored the impact of density interfaces on the sedimentation of particulate material, but to our knowledge very little research has been applied to the deep oceanic environment. Laboratory experiments conducted for a variety of individual particles geometries and densities (e.g. Prairie et al., 2013; Mrokowska, 2018; Doostmohammadi and Ardekani, 2014; Doostmohammadi and Ardekani, 2015; Verso et al., 2019) generally show a decrease in settling velocity, as well as a reorientation of the particles, during their passage through the transition layer formed by the density interface. The initial decrease is followed by an increase, once the particle reaches the base of the interface, but this settling velocity is always smaller in the lower layer due to its higher density. The initial decrease in velocity in the interface is likely due to the entrainment of less dense water by the particles and the drag of their wake as they cross the interface. Finally, Maggi (2013) shows that sedimentation rates are very well correlated with size for mineral particles, but are much less clear for biomineral or biological material. Thus, the sedimentation rate of a solid particle alone will depend almost exclusively on its size. This is not the case for aggregates, where composition, shape, excess density and porosity must also be taken into account. Kindler et al. (2010) demonstrated that slowly sinking particles, like highly porous aggregates, can be retained and therefore accumulate at density interfaces, increasing the likelihood of collisions and subsequent aggregation. They suggested that this increase in retention time may affect carbon transformation through increased microbial colonization and utilization of particles and release of dissolved organics.

The Mediterranean Sea is a prime location for the observation and study of thermohaline staircases due to its unique hydrographic conditions. The study presented here focuses on the behaviour of particulate matter and dissolved elements in the Tyrrhenian Sea (Fig. 1a), a region known to be favourable for salt fingers with the formation of large staircases at the transition between intermediate and deep waters (Durante et al., 2019). The Tyrrhenian Sea has a complex circulation pattern characterized by three main water masses: Atlantic Water (AW) at the surface, Eastern Intermediate Water (EIW) at intermediate depths and Tyrrhenian Deep Water (TDW) in the deeper layers (Millot and Taupier-Letage, 2005; Iacono et al., 2021). Note that the names of the water masses follow the nomenclature recommended by (Schroeder et al., 2024). The surface circulation shows strong seasonal variability, with a basin-wide cyclonic pattern in winter, which weakens and becomes more complex in summer. The EIW, formed in the eastern Mediterranean Basin, enters through the Strait of Sicily. It is found throughout the Tyrrhenian Sea at depths between 200 and 700 m. It participates in the cyclonic flow and exits through the Corsican and Sardinian channels. The deep circulation, involving the TDWs, is defined as resulting from the mixing of the deep waters of the western Mediterranean with the intermediate and deep waters of the eastern basin (Sparnocchia et al., 1999). Staircase

structures, characterized by remarkable spatial and temporal stability, extend over most of the central basin at depths between 600 and 2500 m, with individual layers spanning tens or even hundreds of meters in thickness. These persistent formations have been consistently observed and documented over several decades, demonstrating their long-term coherence in the region (Johannessen and Lee, 1974; Molcard and Tait, 1977; Zodiatis and Gasparini, 1996; Sparnocchia et al., 1999; Falco et al., 2016).

The dataset used in this work was obtained from the PERLE-3 cruise (Pujo-Pay et al., 2020) whose main objective is to study the formation of Levantine intermediate waters in the eastern basin and their fate and transformation along their course in the Mediterranean Sea. Here, we focused on a section between the Bay of Naples and southern Sardinia, which cuts across the Tyrrhenian Sea from east to west. The data collected included hydrological parameters (temperature, salinity, density), hydrodynamic parameters (current velocity and direction), and particulate (turbidity, large particle abundance) and dissolved (oxygen, nitrate) parameters in the upper 2000 m of the water column. The two questions addressed here are (1) what are the characteristics and development conditions of the notable staircase structures observed during this cruise, and (2) what is the impact of these staircases on the distribution of dissolved and particulate matter. The answers to these two questions will provide new insights into the ecological consequences of these small-scale structures.




Figure 1 (a) Map of the stations carried out during part of the PERLE-3 cruise (14–16 March 2020) in the Tyrrhenian Sea. The route followed by the ship is indicated by the dotted red line. The northward deviation in the centre of the basin is associated with the recovery of the profiling float. (b) Θ -S diagram for the different stations of the transect and associated water masses (AW, Atlantic Water; TIW, Tyrrhenian Intermediate Water; EIW, Eastern Intermediate Water; TDW, Tyrrhenian Deep Water), with depth in colour.

2. Material and Methods

2.1 Hydrographical and optical data

This hydrographic survey of the Tyrrhenian Sea was conducted in March 2020 as part of the PERLE-3 cruise, which was unexpectedly shortened due to the outbreak of the COVID-19 pandemic. It includes 16 stations for which hydrological profiles were obtained between the surface and 2000 m with a Seabird 911+ CTD-O2 probe mounted on a rosette carrying 22 twelve-litre Niskin bottles. Additional optical sensors (Wetlabs C-Star 0.25 m pathlength transmissometer at 650 nm wavelength, Seabird Suna V2 UV nitrate sensor, Hydroptic Underwater Vision Profiler UVP-5) were connected to the probe.

Data from the CTD sensors and the transmissometer were recorded at a frequency of 24 Hz. Data were therefore acquired every 2 cm vertically at a descent rate of 1 m s⁻¹. The temperature and conductivity sensors provided measurements with a resolution of 2.10⁻⁴ °C (4.10⁻⁵ S/m) and an accuracy of 2.10⁻³ °C (3.10⁻⁴ S m⁻¹), respectively. The dissolved oxygen sensor provided measurements with a resolution of 0.2 µmol L⁻¹. The transmissometer has a resolution of 1.25 mV over a range of 0 to 5 V (WET Labs, Inc., 2011), giving a beam attenuation coefficient [BAC= $-4 \times \ln(T\%)$, where T% is the transmittance in %] and resolution of 10⁻³ m⁻¹. The UVP-5 is a stand-alone camera mounted on the CTD/rosette frame to quantify the vertical distribution of large particles and zooplankton. Images were acquired at a frequency of up to 6 Hz, i.e. on average every 20 cm at a descent rate of 1 m/s. The small size limit of the UVP-5 is determined by the optical resolution (94.7 µm corresponding to 1 pixel), while the large size limit is determined by the volume of water illuminated (1.02 l).

The stand-alone Seabird Suna V2 UV nitrate sensor attached to the CTD collected data at 1 Hz from the surface down to 2000 m. It uses ultraviolet absorption spectroscopy to measure nitrate in situ. A good correlation was observed with bottle measurements over the entire water column collected during the cruise at station PERLE3-10 ($R^2 = 0.99$, $N = 27$ samples, $p < 10^{-5}$, $\text{NO}_3 \text{ SUNA} = 0.991 \times \text{NO}_3 \text{ btl} - 0.116$).

The CTD data processing was performed with the SBE Data Processing software, and derived variables (potential temperature, salinity, potential density anomaly, and density ratio) were estimated based on the TEOS-10 toolbox (IOC, SCOR and IAPSO, 2010; Picheral et al., 2010).

The transmissometer provides a beam attenuation coefficient (BAC), which is linearly correlated with particle concentration. This signal is influenced by the size and composition of the particles (Hill et al., 2011). Due to their preponderance and larger surface to volume ratio, this signal is essentially sensitive to the finest (sub-micron to micron) particles. As a result, turbidity spikes caused by the rare passage of large particles through the beam were eliminated by smoothing (Giering et al., 2020).

The UVP provides quantitative data on the abundance (in number of particles per litre, # L⁻¹) of large particles, estimated from raw images (Picheral et al., 2010), for different size classes between 80 µm and 2000 µm (equivalent circular diameter). This material is hereafter referred to in the text as large particulate matter (LPM). The UVP data were then processed to estimate the particle size distribution, which was modelled using a typical power law of the form $N(d) = C \times d^{-\alpha}$, where C represents a constant, d stands for particle diameter, and $-\alpha$ denotes the Junge index (Guidi et al., 2009), as a proxy of the particle size distribution. The Junge index is determined through linear regression involving log-transformed values of $N(d)$ and d. The calculation was performed on the size range between 80 and 400 µm, in order

to consider only the most abundant particles, which represent up to 99% of the total number observed and whose concentration is greater than 1 particle per litre (Table A1). The Junge index, α , varies between 2.5 and 4.5 for this data set. A slope of the differential particle size distribution of 4 indicates an equal amount of mass in logarithmically increasing size intervals. Higher values indicate a greater dominance of fine particles within the particle population, while lower values are associated with particle populations enriched in larger particles.

Another CTD data set was collected from a BGC-Argo profiling float (WMO 6,902,903). This float was equipped with SBE-41CP pumped CTD with a sampling rate of 0.5 Hz and an instrumental precision of 0.01 for salinity, 0.002 °C for temperature and 2.4 dbar for pressure. CTD profiles were collected during an ascent from the parking depth to the surface, which took between 3 and 6 hours, depending on the depth, at a nominal vertical speed of 0.1 m s⁻¹. The profiling float collected data for almost 2 years, between 23 June 2018 and 15 March 2020 (date of recovery during the PERLE-3 cruise). It remained in the centre of the Tyrrhenian Basin (between 39.1-39.7° N, 11.6-12.8° E) during this period. The resulting time series of CTD profiles includes 16 profiles between 0 and 1000 m from June 23 to July 6, 2018, and 118 profiles between 0 and 2000 m from July 13, 2018 to March 15, 2020, with a time resolution of 7 days until October 2019, then 3 days thereafter. The filtering step proved to be very important, given the scales on which the study focused, to remove the spikes and reduce the noise of the signal while maintaining the best vertical resolution. The 24 Hz raw data were subjected to outlier removal using two successive moving median filters with window lengths of 7 and 5 scans, respectively. The data were then binned at 1 m intervals and smoothed with a Loess regression filter. For temperature, salinity, and potential density anomaly signals, the loess regression smoothing window length was 10 scans (10 metres). For dissolved oxygen, nitrate, beam attenuation coefficient, UVP, and Junge index signals we applied a Loess regression smoothing with a window length of 50 scans (50 metres).

Thermohaline staircases are characterized by alternating mixed layers, which are thick regions of nearly constant temperature, salinity and density, and interfaces, which are thin layers with sharp changes in temperature, salinity, and density. Step structures were defined based on temperature and salinity profiles between 500 and 2000 m depth, following the procedure described in Durante et al. (2019). Relative maxima in the vertical gradient of salinity and potential temperature are used to identify interfaces that form well-marked steps and delimit a well-mixed layer. Gradient thresholds of 10⁻⁴°C m⁻¹ for potential temperature and 5 x10⁻⁴ PSU m⁻¹ for salinity were used.

2.2 Acoustical data

During the PERLE-3 survey, two shipboard Acoustic Doppler Current Profilers – S-ADCPs – collected continuous current data from 21 m to 1200 m depth. The first S-ADCP was an RDI OS150 with an acoustic frequency of 150 kHz, a sampling rate of 1 Hz, and a cell size of 8 m, allowing a total range of 220 m. The second S-ADCP was an RDI OS38 with an acoustic frequency of 38 kHz, a sampling rate of 1 Hz, and a cell size of 8 m, allowing a maximum range of 1200 m. Data averaged over a 2-min period were concatenated and processed using Cascade V7.2 processing software (Kermabon et al., 2018) to compute horizontal ocean current velocities with a spatial resolution of 2 km, corrected for navigation

and ship attitude parameters, and filtered according to various quality criteria (i.e., thresholds on vertical velocity error, vertical shear, correlation, minimum percentage of valid ensembles, Kermabon et al., 2018). Bathymetry (Etopo 1 with 1 arc-minute resolution) was incorporated in the processing to account for bottom detection. The profile data for the meridional and zonal components of the current for the two S-ADCPs were combined to obtain a complete profile between 21 and 1200 m depth with maximum resolution in the surface layer.

In addition to the S-ADCP current measurements, current data between the surface and 2000 m depth were also collected using a dual-head Lowered-Acoustic Doppler Current Profiler (L-ADCP) system. These measurements were collected with two RDI Workhorse 300 kHz current meters mounted on the CTD frame, one looking up and one looking down. The vertical profiling resolution was 8 m. The data were processed by the velocity inversion method using version IX of the LDEO software (Thurnherr, 2021). Qualified external data (CTD, S-ADCP, GPS) are used to process the L-ADCP data. The horizontal current profiles generated by the L-ADCPs were combined with the current profiles obtained by the S-ADCPs to obtain current sections between 21 m and 2000 m. Vertical ocean velocities were calculated using the LADCP_w_ocean utility from combined raw L-ADCP and CTD data (Thurnherr, 2022). The upward and downward looking data were processed separately and combined during post-processing to provide vertical velocity profiles for the downcast and upcast.

The vertical shear squared (S^2) was calculated from the vertical gradients of the velocity components measured by the L-ADCP using the formula: $S^2 = (\partial u / \partial z)^2 + (\partial v / \partial z)^2$, where u and v are the zonal and meridional horizontal velocity components, and z is depth. The gradient Richardson number (Ri), defined as $Ri = N^2 / S^2$, quantifies the balance between stratification and shear-driven turbulence, with N^2 representing the buoyancy frequency squared. To ensure consistency and reduce noise, the temperature, salinity (used to calculate density), and velocity components were smoothed using a Savitzky-Golay filter with an 48 m window length. The vertical gradients of buoyancy and shear frequencies were then computed over this smoothing interval.

The data from the 38 KHz S-ADCP were also used to derive the acoustic backscatter index (BI, Mullison, 2017), which is a proxy for the abundance of centimetre-scale reflectors (organic detritus, zooplankton, micronecton...) in the water column. This derivation takes into account the absorption and geometric dispersion of sound $BI = K_c * (RL - Er) + (TL_w + TL_g)$, where K_c is the conversion factor (*count to decibels*) of the ADCP used, RL the received signal, Er the signal noise, TL_w and TL_g are the absorption and geometric transmission losses of the acoustic signal in water respectively.

2.3 Selection of representative stations

A classification of the stations was made by threshold-based detection of buoyancy frequency anomalies between 700 and 1900 m relative to the background stratification. The ‘WStairs’ category includes stations with 2 to 4 sharp staircases characterized by N peaks (>0.3 cph), corresponding to strong density gradients, bounded above and below by well-mixed layers. The ‘NoStair’ category includes stations with smoother N profiles without multiple

strong stairs. The ‘WStairs’ category comprises stations 8, 9, 11, 12, 13 and 15, while the ‘NoStair’ category includes stations 7, 14, 16, 19 and 20. The vertical profiles of key thermohaline (temperature, potential, salinity, potential density anomaly), particulate (beam attenuation coefficient, large particle abundance, Junge index) and biogeochemical (dissolved oxygen, nitrate, apparent oxygen utilization) parameters are presented here for a representative station of both categories. Station 9 was chosen as a representative station ‘WStairs’ for the processes occurring at the density interfaces, while station 20 was chosen as representative of the ‘NoStair’ category.

3. Results

3.1 Hydrological and hydrodynamical features along the section

3.1.1 Temperature, salinity, oxygen, and nitrate

The temperature-salinity diagram (Fig. 1b) and the basin cross-section (Fig. 2) clearly identify the distribution of the water masses present in the study area. As in Falco et al. (2016), we used the isohaline 38.72 as the minimum salinity value to identify the shallowest and deepest levels of the EIW. The AW, which extends in the upper 200 m of the water column, has a higher temperature and lower salinity to the east. The colder Tyrrhenian Intermediate Water (TIW) can be distinguished between 150 and 220 m in a tongue-like feature trapped in an anticyclonic eddy in the middle of the section (see § 3.1.2). The warmer, saltier, and oxygen-depleted core of the EIW is found at depths between 300 and 600 m, as illustrated in Fig. 2 (sections a, b, and c). The core of the colder and less saline TDW is visible beyond 1200 m depth (Fig. 2a and b).

In March, the nitrate distribution in the Tyrrhenian Sea (Fig. 2d) shows a nutrient-poor surface layer and a nutrient-rich deep layer, typical of oligotrophic conditions. Near the surface, nitrate concentrations are low, about $1 \mu\text{mol kg}^{-1}$, due to biological uptake. With increasing depth, nitrate concentrations increase. At intermediate depths (250–650 m) they range from 4 to $7 \mu\text{mol kg}^{-1}$, indicating a transition zone with maximum vertical gradients. In deeper waters, concentrations reach 7 to $9 \mu\text{mol kg}^{-1}$ due to decomposing organic matter. An anticyclonic eddy around 12°E and the boundary current jet, especially over the eastern part of the section, modify the intermediate water depth (EIW) by a few hundred metres without significantly affecting the deep water depth (TDW). To the west, the signature of a cyclonic eddy is also visible and contributes to the vertical modulation of isopycnals. These mesoscale structures affect the distribution of oxygen and nitrate, which increase less rapidly with depth beneath them.





Figure 2: Cross-basin section of (a) potential temperature, (b) salinity, (c) dissolved oxygen, and (d) nitrate for the PERLE-3 cruise (March 2020). The upper x-axis indicates the station number along the section that runs from west (station 18) to east (station 4).

3.1.2 Currents

Dynamic topography and observed horizontal velocity in the water column during PERLE-3 reveal the presence of two large mesoscale eddies in the centre of the basin with maximum velocities of 10–15 cm s^{-1} (Fig. 3a, b). The southern edge of the cyclonic eddy at 10.5° E extends to a depth of about 200 m, while the anticyclonic eddy at 12° E extends to more than 500 m. The deep current running along the eastern edge of the basin defines the general along-slope cyclonic circulation. Below the EIW core, i.e. at 600 m depth, the currents are very weak, of the order of a few cm s^{-1} (Fig. 3c). The Richardson number (Ri, Fig. 3d) is high in the surface layers down to a depth of approximately 1000 metres, indicating high water column stability and low vertical mixing potential. Below this depth, Ri decreases significantly, with localised areas of very low values indicating regions where stability is lower and vertical mixing may be favoured.






Figure 3: (a) Absolute dynamic topography (in cm) and surface geostrophic currents derived from daily Copernicus Marine Environment Monitoring Service Gridded L4 data (interpolated to 3.75 arcmin), averaged over the period of the PERLE-3 cruise (14–16 March 2020). Black dots indicate station locations; see Figure 1 for station numbering. (b) Stick plot of vertically averaged S-ADCP currents between 21 m and 1200 m along the ship’s route across the basin, (c) Meridional component of combined S-ADCP and L-ADCP currents, and (d) Richardson number derived from the L-ADCP measurements. The upper x-axis indicates the station number along the section that runs from west (station 18) to east (station 4).

The estimated vertical velocities for the layer between the surface and 2000 m depth (Fig. 4) are between -15 and +15 mm s⁻¹ with a rms of 4 mm s⁻¹. There was no significant spatial variation in vertical velocities, apart from greater vertical shear in the first 300 m of the water column on either side of the eddy at around 12° E.




Figure 4: Histogram of the vertical velocities between 20 and 2000 m depth estimated from the L-ADCP measurements for all the stations during the PERLE 3 cruise.

3.1.3 Turbidity, large particle abundance and Junge index

The beam attenuation coefficient, an indicator of the abundance of small particles (Fig. 5a) is highest in the surface layer and along the continental slope on both sides of the basin. The tongue-like feature of turbid water that descends to 500–600 m in the western half of the basin (stations 19–20) is associated with the downward movement of water around the anticyclonic eddy.

Coarse particles observed with the UVP (Fig. 5b) are most abundant along the continental slope and between 400 and 900 m depth throughout the basin. The subduction effect of the anticyclonic eddy is also evident in the abundance of large particles.

a

b




Figure 5: Cross-basin section of (a) beam attenuation coefficient, (b) total large particle abundances, (c) backscatter index for the PERLE-3 cruise, and (d) Junge index. The upper x-axis indicates the station number along the section that runs from west (station 18) to east (station 4).

An accumulation of large reflectors between 200 and 500 m depth is clearly visible on the 38 kHz ADCP echo intensity (Fig. 5c). It corresponds to the deep scattering layer formed by the micronekton (small mesopelagic fishes, crustaceans, and cephalopods) (Kapelonis et al., 2023; Peña, 2024). Its fragmentation into elongated patches of lower intensity between 200-300 m is related to the diel vertical migrations (upward motion) of part of the organisms at night.

The Junge index (Fig. 5d), estimated from UVP data, shows an intermediate maximum – indicating the preponderance of smaller particles over coarser ones – between 400 and 1000 m, below the deep scattering layer. Below 1000 m, the index decreases by one unit to 2000 m depth, indicating a decrease in the relative abundance of smaller particles compared to coarser particles. This decrease is more pronounced on the continental slope.

3.2 Hydrological features of thermohaline staircases


3.2.1 Attributes of staircase station vs. “non-staircase” station

In this section we describe the vertical distribution of physical (potential temperature, salinity, potential density anomaly), dissolved and particulate parameters (dissolved oxygen, nitrate, apparent oxygen utilization, beam attenuation coefficient, large particle abundance, and Junge index) at station 20, which shows virtually no significant staircase (Fig. 6), and station 09 which shows marked staircases between 700 and 1700 m (Fig. 7).

At station 20, the vertical profiles of most variables exhibit relatively smooth characteristics between 800 m and 1070 m depth. At 1070 m, a homogeneous layer of approximately 70 m thickness is observed, immediately succeeded by a pronounced density interface. The water column structure below this interface demonstrates increased variability, characterized by minor step-like features in the measured parameters.

The profiles of physical variables at station 09 show a series of steps starting at 750 m depth. Density steps result in thin interfaces (about 9–73 m) with temperature steps of approximately 0.1–0.2°C and salinity steps of 0.01–0.03 PSU between adjacent mixed layers, resulting in density steps of a few thousandths of a kg m^{-3} . The thickness of the mixed layers varies between 10 and 230 m and increases significantly below 1000 m. The profiles of the biogeochemical variables also show step-like profiles, with larger gradients corresponding to the density steps and nearly homogeneous concentrations in the mixed layers between each step. This is clearly visible for dissolved elements (oxygen, nitrate) and small particle concentration (beam attenuation coefficient). The effect of the density steps on the abundance of coarser material is less obvious due to the variability of the measurements, but it still appears that the total abundance decreases significantly below each step. It is noteworthy that the decrease in the

379 Junge index between 700 and 1700 m is greater for station 9 with staircase steps, of the order
 380 of one unit, than for station 20, which is more irregular and of the order of 0.2 units.



381

382 *Figure 6: Profiles of potential temperature, salinity, potential density anomaly (top row)*
 383 *dissolved oxygen, nitrate, apparent oxygen utilization (AOU) (first middle row), beam*
 384 *attenuation coefficient, large particle abundance between 80 and 400 μm , Junge index (second*
 385 *middle row), buoyancy frequency, Richardson number, shear squared (S^2 , red line) and 4 times*
 386 *the buoyancy frequency squared, ($4 \times N^2$, black line), and the Richardson number (< 0.25 , blue*
 387 *dots) (bottom row) between 700 and 1600 m deep for station PERLE3-20. The grey dots are*
 388 *the data binned at 1-metre intervals and the solid red line indicates the smoothed profile.*

389




Figure 7: Profiles of potential temperature, salinity, potential density anomaly (top row) dissolved oxygen, nitrate, apparent oxygen utilization (AOU) (first middle row), beam attenuation coefficient, large particle abundance between 80 and 400 μm , Junge index (second middle row), buoyancy frequency, Richardson number, shear squared (S^2 , red line) and 4 times the buoyancy frequency squared ($4 \times N^2$, black line), and the Richardson number (< 0.25 , blue dots) (bottom row) between 700 and 1600 m deep for station PERLE3-09. The grey dots are the data binned at 1-metre intervals and the solid red line indicates the smoothed profile. See station position in Fig. 1.

Regarding the stratification and stability parameters (N , S^2 , Ri), at station 20, the buoyancy frequency (N) generally below 0.5 cph, and the Richardson number (Ri) remains high (mostly > 1). Only a few isolated depths below 1000 m have an S^2 value greater than $4 \times N^2$ ($Ri < 0.25$), suggesting that turbulent mixing is rare and that dynamic stability prevails throughout this depth range.

In contrast, station 09 displays sharp, repeated peaks in N often reaching or exceeding 0.75 cph. The S^2 profile frequently surpasses $4 \times N^2$, and the Ri drops below 0.25 at the mixed layer's level that encompasses the density interfaces. This pattern reflects a water column punctuated by dynamically unstable layers, where turbulent mixing is likely to be concentrated in the homogeneous layers of the staircases.

3.2.2 Positioning and persistence of main staircases

The transition zone between the EIW and the TDW thus provides the right conditions (warmer and saltier water mass overlying a colder and less saline water mass) for salt fingers. The thermohaline gradients from the profiles collected during the cruise and from the profiling float allowed us to identify the main stepped structures in the Tyrrhenian Basin (Fig. 8).

During the PERLE-3 cruise, distinct thermohaline staircases are observed between 600 and 2000 m depth, primarily in the central region of the basin (Fig. 8a). These staircases are notably absent in two areas: near the western slope of the basin and beneath the deep anticyclonic eddy located at about 12° E. The absence of staircases extends to 1000 m depth beneath this eddy.

The temporal evolution of the staircase in the centre of the basin, as seen by the profiling float between July 2018 and March 2020, underscores that these structures, particularly at depths greater than 1000 m, are relatively stable and have been maintained for several years (Fig. 8b).




Figure 8: (a) Depth of the main thermohaline staircases (dashed lines) across the basin during the PERLE-3 cruise and (b) temporal evolution of the staircases in the centre of the basin as seen by the profiling float between July 2018 and March 2020. The dots are the steps identified using the procedure referenced in the methods (§2.1). The blue lines show the isopycnals. The position of the stations and the trajectory of the profiling float are shown on the maps on the right.

4. Discussion

4.1 Circulation and thermohaline staircases

The hydrodynamic patterns observed during the PERLE-3 cruise are consistent with the established understanding of winter circulation in the southern Tyrrhenian Sea (Rinaldi et al., 2010; Iacono et al., 2013; Iacono et al., 2021). The winter circulation in this region is primarily driven by wind forcing, resulting in a vigorous cyclonic flow that encompasses the entire Tyrrhenian Basin and persistent mesoscale eddy structures. The observed anticyclonic gyre between 12°E-13°E and 39°N-40°N, and the cyclonic circulation between Sardinia and Sicily, corroborate the findings of previous studies.

The Tyrrhenian Sea is a region where overall turbulence levels remain low, as reported by Ferron et al. (2017), who estimated extremely low dissipation rates derived from microstructure. Our observations show well-defined thermohaline staircase structures associated with salt fingers developing between 700 m and 2000 m. The observed thermohaline staircases occur mainly in the centre of the Tyrrhenian Basin and are less defined on the continental slope at both ends of the section. These latter regions are the ones that contain the cyclonic boundary circulation that entrains the core of the EIWs originating from the Strait of Sicily. These results are consistent with those of Zodiatis and Gasparini (1996) from ship observations, of Buffett et al. (2017) from seismic observations, and Taillandier et al. (2020) from profiling float observations, who showed that the staircases with the most distinct step-like gradients appear in the centre of the basin, while they become less evident towards the boundaries and the bottom. They linked this change to increased vertical motions that prevent diffusive convection and staircase formation because the internal wave field and current shear are stronger near the boundaries. Similarly, Durante et al. (2021) found that internal gravity wave-induced mixing can modulate staircase structures in the basin interior.

Mesoscale eddies appear to be able to disrupt this balance. Seismic observations in the Caribbean (Yang et al., 2024) have shown that eddies generate turbulent mixing and vertical shear, especially at their peripheries, which can erode thermohaline staircases down to 750 m. In our study, staircases are notably absent up to 1000 m beneath the anticyclonic eddy at 12°E, suggesting a similar disruptive effect. This erosion is not observed near the neighbouring cyclonic structure to the west, possibly due to the focusing of near-inertial internal waves and enhanced mixing within anticyclones, as described by Fer et al. (2018) in the Lofoten Basin Eddy. While this co-occurrence does not establish causality, and in the absence of turbulence measurements, the potential disruptive effect of this eddy on staircase formation warrants consideration.

The contrasting conditions observed at the representative stations 20 and 09 suggest the role of background stratification (Fig. 8a). At the station 020 (Fig. 6), located under the

anticyclonic eddy, we observe relatively high Richardson numbers ($Ri > 2$) down to 1000 m, due to stronger stratification, probably caused by the deepening of the isopycnals under the eddy, creating a stable environment that prevents staircase formation. Conversely, station 09 exhibits localised Ri drops below 0.25 within mixed layers, creating discrete 'mixing hotspots' while maintaining the overall integrity of the staircase. In the density steps, the high buoyancy frequency, which overcomes the vertical shear, helps to prevent instability.

Below 1000 m, reduced stratification and shear create low Ri conditions, which appear to be important for both the formation and maintenance of thermohaline staircases, as evidenced by both float measurements and cruise data in the basin's central region. The observed persistence of thermohaline steps below 1000 m depth also corroborates previous observations that have highlighted the stability of these structures in the basin interior. These step-like features have been documented to maintain their integrity over temporal scales ranging from years (Falco et al., 2016; Taillandier et al., 2020; Zodiatis and Gasparini, 1996) to decades (Durante et al., 2019) and show strong lateral coherence, sometimes reaching several hundred kilometres (Buffett et al., 2017).

The subsequent sections provide a detailed examination of how these persistent, well-developed thermohaline staircases may influence particulate matter dynamics, specifically addressing their impact on sedimentation processes (§ 4.2) and the degradation of suspended particles (§ 4.3). This analysis aims to investigate the complex interactions between physical oceanographic features and biogeochemical cycles in the deep marine environment.

4.2 Impact of staircases on the distribution and settling of particulate matter

The Junge index has been shown to be a valid first-order description of the particle size distribution, particularly for the study of size-dependent processes such as particle sinking (Guidi et al., 2009). In our case, the observed abundance of coarse particles and the Junge index are greatest in the deep scattering layer between 400 and 800 m (Fig. 5). Deeper down to 2000 m, both the total particle abundance and the slope of the size distribution decrease, indicating that the contribution of coarser particles to the particle population increases with depth. Similar results regarding the size distribution of smaller particles in the Tyrrhenian Basin were obtained by (Chaikalis et al., 2021) during the trans-Mediterranean cruise in March 2018 using an in-situ laser scattering and transmissometry instrument (LISST-Deep). In their study, they considered the size range between 5.6 and 92.6 μm , complementary to that measured with the UVP. Their stations show similar particle distribution size with depth below the surface layer (>150 m), with an intermediate maximum at about 400–500 m depth and a steady decrease to 2000 m depth. Such an evolution of the abundance and particle size distribution is generally considered to be the result of aggregation, with smaller particles agglomerating to form larger ones, a common process in the ocean (McCave, 1984).




Figure 9: Vertical profiles of (a) mean Junge index and (b) mean Apparent Oxygen Utilization (AOU) in profiles for the groups of stations with staircases ("WStairs", red open circles) and without staircases ("NoStair", blue filled triangles). For each depth, the values represent the mean of all available stations within each group, and the error bars correspond to the standard deviation among stations of the same group. Linear regression equations and R^2 values are indicated in each panel (between 700 and 1900 m for a) and between 1000 and 1900 m for b)). In panel (b), AOU profiles have been shifted so both profiles start from the same value at 700 m, to facilitate comparison of their vertical trends.

To assess the influence of thermohaline staircases on particle dynamics, the vertical distribution of the Junge index was examined across depth under two distinct conditions: presence (“WStairs”) and absence (“NoStair”) of multiple staircases (see §3.2.2 for definitions). A multiple linear regression framework, incorporating an interaction term between depth and staircase condition (Aiken et al., 1991), was applied to evaluate whether the depth-dependent trends in the Junge index diverged significantly between conditions.

As illustrated in Fig. 9a, the regression model demonstrated a robust fit to the observational data ($R^2 = 0.986$, $p \ll 0.001$), with residuals conforming to a normal distribution (Shapiro-Wilk test: $p = 0.081$), validating the statistical assumptions. The interaction term between depth and staircase condition exhibited high statistical significance ($p < 0.001$), confirming distinct depth-dependent slopes for the Junge index under “WStairs” and “NoStair” conditions. In “NoStair” regions, the Junge index decreased markedly with depth (slope: -0.0007 m^{-1}), consistent with a gradual reduction in the relative abundance of smaller particles compared to larger particles. However, under “WStairs” conditions, this decrease was more pronounced (slope: -0.0008 m^{-1}), suggesting that stairs modulate grain size distribution, potentially retaining larger particle fractions during their sedimentation in the water column.

The overall diminution in suspended particle abundance, concurrent with an augmentation of the coarse particle fraction, is evident not only at large vertical scales throughout the water column beneath the Eastern Intermediate Waters (EIWs), but also at finer scales spanning several decametres within the thermohaline staircases. The influence of density steps on the vertical distribution of particle abundance and size is illustrated through an examination of three distinct interfaces, situated at different stations and depths ranging from 700 to 1400 m, i.e. station 08 (Fig. 10), station 11 (Fig. 11), and station 09 (Fig. 12). These interfaces, characterized by thicknesses varying between 20 and about 150 m, exhibit density variations of $3 \text{ to } 5 \times 10^{-3} \text{ kg m}^{-3}$.

These examples show a significant reduction in the abundance of fine particles, as evidenced by transmissometry (BAC) measurements, across each interface. For coarse particles detected by UVP, despite measurement variability, the change in abundance across the interface varies according to particle size. At the stations 08 (Fig. 10) and 11 (Fig. 11), a slight decrease in abundance is observed for the smallest size fraction (80.6-161 μm), while minimal variation is noted for intermediate sizes (161-256 μm), and an increase is evident for the largest particles ($> 256 \mu\text{m}$). At station 08, the Junge index exhibits a clear decrease across the density interface. The signal is less pronounced at station 11, where the index decreases at the interface's upper boundary but shows a localized increase at its base. For the interface at station 09 (Fig. 12), all parameters — turbidity, abundance across different size fractions, and the Junge index — decrease across the interface. These depth-dependent variations provide evidence that these density interfaces exert an influence on particle abundance and size distribution.




Figure 10: Profiles of potential density anomaly, beam attenuation coefficient, large particle abundances for the first five size classes and for all sizes $> 256 \mu m$ of the UVP, dissolved oxygen, nitrate, apparent oxygen utilization (AOU), and Junge Index between 1100 and 1300 m deep for station PERLE3-08. Grey points show individual in situ measurements; the red line represents the averaged vertical profile.




Figure 11: Profiles of potential density anomaly, beam attenuation coefficient, large particle abundances for the first five size classes and for all sizes $>256 \mu\text{m}$ of the UVP, dissolved oxygen, nitrate, apparent oxygen utilization (AOU), and Junge Index between 1100 and 1300 m deep for station PERLE3-11. Grey points show individual in situ measurements; the red line represents the averaged vertical profile.




Figure 12: Profiles of potential density anomaly, beam attenuation coefficient, large particle abundances for the first five size classes and for all sizes $> 256 \mu m$ of the UVP, dissolved oxygen, nitrate, apparent oxygen utilization (AOU), and Junge index between 700 and 900 m deep for station PERLE3-09. Grey points show individual in situ measurements; the red line represents the averaged vertical profile.

Based on laboratory and modelling experiments, strong density interfaces are known to cause particle retention and promote aggregation. According to (Doostmohammadi and Ardekani, 2015), density interfaces can induce preferential retention of fine, slow-sinking particles compared to larger particles, and can therefore promote the formation of clusters of fine particles, which can then aggregate. This retention phenomenon can also act on coarse particles, which are often formed by high porosity aggregates. According to Kindler et al. (2010), large porous particles whose effective density is close to that of ambient water are much more dependent on transition zones and large accumulations of porous particles can occur at interfaces due to their less dense water transport, which is also conducive to aggregation. In our case, the density difference at the interface between two staircases are really weak (a few thousandths of kg/m^3) and despite the measurement uncertainty of the particle abundances, the

results suggest that interfaces have an effect on the size distribution – through retention and aggregation – of the settling particles. The larger particles, which are generally denser and have higher settling velocities, are not affected by the change in water density.

In addition, convection phenomena in the mixed layers sandwiching the density interfaces are likely to even out the concentration of slowly sinking particles (especially the finest and most numerous). The estimated vertical velocities of the current, on the order of a few mm s⁻¹, are likely to alter the settling of particles, helping to homogenize their abundance with depth and increase their residence time in each layer.

4.3 Potential impact of staircases on the biogeochemical activity

The vertical export of organic particles to the deep ocean is a crucial component of the oceanic carbon cycle, with flux decreasing significantly (>70%) in the mesopelagic zone (100-1000 m depth). Several biogeochemical processes are responsible for the reduction of this flux and particle lability decrease with depth. Briggs et al. (2020) suggest that particle fragmentation could account for approximately half of the observed particle loss in the mesopelagic zone. Additionally, remineralization by heterotrophic respiration and enzymatic solubilization both by free-living and particle-attached microbes play a key role in particle degradation (Karl et al., 1988; Collins et al., 2015). It is also well known that marine snow aggregates act as hotspots for microbial activity, hosting diverse communities of bacteria, protozoans, and other microorganisms. Particle-attached microbial community evolves with depth, due to evolution of the initial attached community rather than de novo colonization, detachment or grazing as the marine snow sinks (Thiele et al., 2015). These communities are often orders of magnitude more concentrated and active than those in the surrounding water, leading to intense localized degradation and remineralization of organic matter (Kiørboe, 2001; Baumas and Bizic, 2024).

The presence of thermohaline staircases appears to affect the sedimentation patterns, promoting particle retention and aggregation, increasing the time particles spend in the water column. These structures also create distinct microenvironments with gradients of oxygen and nutrients, which are believed to be the result of the degradation of particulate organic. This effect is illustrated for three separate interfaces in figures 10 to 12. It can be then hypothesized that the increase in particle residence time favours the remineralization process of organic matter, which releases nutrients while consuming dissolved oxygen through heterotrophic respiration. Depending on the conditions and with a sufficiently long stability period, these activities can lead to localized oxygen depletion and an increase in AOU. In our data, this effect is observed in the shallower staircases, between 750 and 900 m, at station 09 (Fig. 12, bottom row). For deeper interfaces for stations 08 and 11, at around 1200 m (Figs. 10 and 11, bottom row), the increase in AOU and the potential accumulation of nitrate are masked by a stronger vertical oxygen gradient and higher nutrient concentrations compared to shallower depths. Moreover, biological activity within these deeper staircases is also reduced due to the decreased lability of organic matter with increasing depth (Ghiglione et al., 2009; Karl et al., 1988).

To assess the effect of thermohaline staircases on AOU profiles, we compared linear regressions of AOU versus depth between stations with ("WStairs") and without ("NoStair") multiple staircases (Fig. 9b), focusing on the 1000-1900 m depth range where the relationship

is approximately linear. The analysis shows a statistically significant difference in the slopes of the AOU between the “WStairs” and “NoStair” conditions ($p = 2.47 \times 10^{-22}$). Specifically, the AOU decreases less rapidly with depth in the “Wstairs” condition (slope = -0.00281) compared to the “NoStair” condition (slope = -0.00328). This difference suggests that oxygen demand decreases more slowly in the “Wstairs” condition. This could indicate increased in situ mineralization activity compared to the “NoStair” condition. A slower decline in AOU may reflect a more active microbial cycle and a greater contribution of organic matter degradation to the oxygen budget within the “Wstairs” setting. This pattern supports the idea that stairs may slow the decline of the respiration signal with depth, possibly due to retention at density interfaces, as previously proposed for particle dynamics.

The persistence of these staircase structures allows sufficient time for significant biogeochemical transformations to occur. Arístegui et al. (2009) and Nagata and Kirchman (1997) showed that in deep marine environments, the degradation kinetics of organic matter, which vary according to temperature, oxygen availability, and microbial composition, can range from a few days to several months. This can lead to the local accumulation of nitrates, creating pockets of higher concentration and intensifying vertical double diffusive nitrate fluxes toward upper layers. However, observing these accumulations with the present resolution and precision remains challenging. Taillandier et al. (2020) already showed that the thermohaline staircases in the Tyrrhenian Sea significantly influence the biogeochemical dynamics by contributing to the nitrate enrichment of the EIW through diffusion. They estimated upward nitrate fluxes (assuming weak turbulent diapycnal diffusivity) in the transition zone between 400 and 2000 m depth to be about $4 \mu\text{mol m}^{-2} \text{d}^{-1}$ using the salt diffusivity formulation of Radko and Smith (2012) based on the density ratio ($R_\rho = \alpha \times \partial\theta/\partial z / \beta \times \partial S/\partial z$), where α and β are the thermal expansion and haline contraction coefficients of seawater, respectively, and the molecular diffusivity of heat is $k_T = 1.4 \times 10^{-7} \text{ m}^2 \text{s}^{-1}$. We calculated nitrate diffusive fluxes, assuming weak turbulent diapycnal diffusivity) at each interface using the same formulation. The nitrate flux (F_{NO_3}) can be expressed as:

$$F_{\text{NO}_3} = K_{\text{sf}} \times \partial C_{\text{NO}_3} / \partial z = k_T \times R_\rho \times (135.7 / (R_\rho - 1)^{1/2} - 62.75) \times \partial C_{\text{NO}_3} / \partial z$$

where K_{sf} is the salt-fingering diffusivity of nitrate and $\partial C_{\text{NO}_3} / \partial z$ the vertical gradient in nitrate concentration. At station 09, the diffusive nitrate flux was assessed at the depth of each step between 600 and 1600 m (Fig. 13). Fluxes ranged from $3.8 \mu\text{mol m}^{-2} \text{d}^{-1}$ at the deepest depth (1254 m) to $19.1 \mu\text{mol m}^{-2} \text{d}^{-1}$ for the three shallower depths (728 to 870 m). The fluxes at the interfaces are therefore stronger than the overall flux estimated by Taillandier et al. (2020) for the 400-2000 m depth range. Fluxes at interfaces are the major contributor to the total nitrate flux. In agreement with our estimation of double diffusive upward nitrate flux, Durante et al. (2021) estimated that heat and salt fluxes are more pronounced in the upper part of the staircases zone (700 m – 1600 m) but decrease below 1800 m, reaching a minimum near the base of the staircases.




Figure 13: Estimated double diffusive nitrate fluxes across each interface (annotated in blue and expressed in $\mu\text{mol m}^{-2} \text{d}^{-1}$) from station 9. The blue arrows indicate the direction of the flux and the sizes are proportional to the intensity. In red: the vertical profile of the specific volume anomaly (σ_θ in kg m^{-3}) delimiting homogeneous and gradient zones. The grey points correspond to nitrate measurements by the SUNA, while the blue squares are the values averaged over 1 m for each point at the upper and lower interfaces, and used to calculate the nitrate gradient in the shaded areas.

5. Conclusion

This study examines the effect of observed weakly stepped density gradients induced by salt fingering on the distribution and fluxes of particulate and dissolved elements. While the effect of density steps on particle settling has been well studied experimentally and demonstrated for strong density gradients in natural environments (i.e. buoyant river plumes

and pycnocline), this is the first time, to our knowledge, that this effect has been identified for steps with the extremely low-density gradients that exist in the deep ocean.

In the Tyrrhenian Sea, the interface between the warm and saline intermediate waters (EIW) and the colder and less saline deep waters (TDW) presents thermohaline conditions favourable to the development of significant salt fingering staircases. These structures are predominantly observed in regions characterized by minimal hydrodynamic activity, situated in the interior of the basin, distanced from boundary influences, and beneath the depth of influence of mesoscale eddy dynamics. Furthermore, the investigation confirms that the thermohaline staircases in the central region of the Tyrrhenian basin exhibit remarkable temporal stability, persisting for several years.

Overall, this study demonstrates that the interplay between physical oceanographic processes and biogeochemical dynamics in the deep Tyrrhenian Sea is strongly mediated by the presence of thermohaline staircases. The presence of thermohaline staircases appears to have a significant influence on the distribution of particulate and dissolved matter. Profiles of particle abundance resolved down to a few hundred microns, or profiles of dissolved substances such as oxygen and nitrate, tend to mirror the step-like structure observed in the density profiles.

Thermohaline staircases create distinct density interfaces that act as physical barriers, slowing the descent of particles. The retention of near-floating particles at density interfaces is thought to increase their residence time, promote particle aggregation and, incidentally, allow the larger particles thus formed to cross the density interface. At the same time, the staircases influence the concentration of nutrients (such as nitrate) and dissolved oxygen levels, thereby potentially affecting microbial activity and nutrient cycling within these interfaces. The retention of part of the particulate material at the density interfaces allows the mineralization of organic matter, contributing to nitrate enrichment and increasing the upward diapycnal diffusive fluxes of oxygen and nitrate. Ecologically, it is suggested that these staircases can play a crucial role in promoting diverse habitats and influencing the lability of organic matter and the nutrient distribution and fluxes. Such findings underscore the importance of fine-scale physical structures in shaping the biogeochemical landscape of the ocean interior and emphasize the need for further research to fully understand their implications for marine ecosystems.

References

- Aiken, L.S., West, S.G., Reno, R.R., 1991. Multiple Regression : Testing and Interpreting Interactions. Sage publications, Newbury Park (Calif.).
- Arístegui, J., Gasol, J.M., Duarte, C.M., Herndl, G.J., 2009. Microbial oceanography of the dark ocean's pelagic realm. *Limnology & Oceanography* 54, 1501–1529. <https://doi.org/10.4319/lo.2009.54.5.1501>
- Baumas, C., Bizic, M., 2024. A focus on different types of organic matter particles and their significance in the open ocean carbon cycle. *Progress in Oceanography* 224, 103233. <https://doi.org/10.1016/j.pocean.2024.103233>

720 Briggs, N., Dall’Olmo, G., Claustre, H., 2020. Major role of particle fragmentation in regulating
721 biological sequestration of CO₂ by the oceans. *Science* 367, 791–793.
722 <https://doi.org/10.1126/science.aay1790>

723 Buffett, G.G., Krahmann, G., Klaeschen, D., Schroeder, K., Sallarès, V., Papenberg, C.,
724 Ranero, C.R., Zitellini, N., 2017. Seismic Oceanography in the Tyrrhenian Sea: Thermohaline
725 Staircases, Eddies, and Internal Waves. *JGR Oceans* 122, 8503–8523.
726 <https://doi.org/10.1002/2017JC012726>

727 Chaikalis, S., Parinos, C., Möbius, J., Gogou, A., Velaoras, D., Hainbucher, D., Sofianos, S.,
728 Tanhua, T., Cardin, V., Proestakis, E., Amiridis, V., Androni, A., Karageorgis, A., 2021.
729 Optical Properties and Biochemical Indices of Marine Particles in the Open Mediterranean Sea:
730 The R/V Maria S. Merian Cruise, March 2018. *Front. Earth Sci.* 9, 614703.
731 <https://doi.org/10.3389/feart.2021.614703>

732 Collins, J.R., Edwards, B.R., Thamtrakoln, K., Ossolinski, J.E., DiTullio, G.R., Bidle, K.D.,
733 Doney, S.C., Van Mooy, B.A.S., 2015. The multiple fates of sinking particles in the North
734 Atlantic Ocean. *Global Biogeochemical Cycles* 29, 1471–1494.
735 <https://doi.org/10.1002/2014GB005037>

736 Doostmohammadi, A., Ardekani, A.M., 2015. Suspension of solid particles in a density
737 stratified fluid. *Physics of Fluids* 27, 023302. <https://doi.org/10.1063/1.4907875>

738 Doostmohammadi, A., Ardekani, A.M., 2014. Reorientation of elongated particles at density
739 interfaces. *Phys. Rev. E* 90, 033013. <https://doi.org/10.1103/PhysRevE.90.033013>

740 Durante, S., Oliveri, P., Nair, R., Sparnocchia, S., 2021. Mixing in the Tyrrhenian Interior Due
741 to Thermohaline Staircases. *Front. Mar. Sci.* 8, 672437.
742 <https://doi.org/10.3389/fmars.2021.672437>

743 Durante, S., Schroeder, K., Mazzei, L., Pierini, S., Borghini, M., Sparnocchia, S., 2019.
744 Permanent Thermohaline Staircases in the Tyrrhenian Sea. *Geophysical Research Letters* 46,
745 1562–1570. <https://doi.org/10.1029/2018GL081747>

746 Falco, P., Trani, M., Zambianchi, E., 2016. Water mass structure and deep mixing processes in
747 the Tyrrhenian Sea: Results from the VECTOR project. *Deep Sea Research Part I: Oceanographic Research Papers* 113, 7–21. <https://doi.org/10.1016/j.dsr.2016.04.002>

749 Fer, I., Bosse, A., Ferron, B., Bouruet-Aubertot, P., 2018. The Dissipation of Kinetic Energy in
750 the Lofoten Basin Eddy. *Journal of Physical Oceanography* 48, 1299–1316.
751 <https://doi.org/10.1175/JPO-D-17-0244.1>

752 Ferron, B., Bouruet Aubertot, P., Cuypers, Y., Schroeder, K., Borghini, M., 2017. How
753 important are diapycnal mixing and geothermal heating for the deep circulation of the Western
754 Mediterranean? *Geophysical Research Letters* 44, 7845–7854.
755 <https://doi.org/10.1002/2017GL074169>

756 Ghiglione, J.-F., Conan, P., Pujo-Pay, M., 2009. Diversity of total and active free-living vs.
757 particle-attached bacteria in the euphotic zone of the NW Mediterranean Sea. *FEMS Microbiology Letters* 299, 9–21. <https://doi.org/10.1111/j.1574-6968.2009.01694.x>

759 Giering, S.L.C., Cavan, E.L., Basedow, S.L., Briggs, N., Burd, A.B., Darroch, L.J., Guidi, L.,
760 Irisson, J.-O., Iversen, M.H., Kiko, R., Lindsay, D., Marcolin, C.R., McDonnell, A.M.P.,
761 Möller, K.O., Passow, U., Thomalla, S., Trull, T.W., Waite, A.M., 2020. Sinking Organic

762 Particles in the Ocean—Flux Estimates From in situ Optical Devices. *Front. Mar. Sci.* 6, 834.
763 <https://doi.org/10.3389/fmars.2019.00834>

764 Guidi, L., Stemmann, L., Jackson, G.A., Ibanez, F., Claustre, H., Legendre, L., Picheral, M.,
765 Gorsky, G., 2009. Effects of phytoplankton community on production, size, and export of large
766 aggregates: A world-ocean analysis. *Limnology & Oceanography* 54, 1951–1963.
767 <https://doi.org/10.4319/lo.2009.54.6.1951>

768 Hill, P.S., Boss, E., Newgard, J.P., Law, B.A., Milligan, T.G., 2011. Observations of the
769 sensitivity of beam attenuation to particle size in a coastal bottom boundary layer. *J. Geophys.*
770 *Res.* 116, C02023. <https://doi.org/10.1029/2010JC006539>

771 Iacono, R., Napolitano, E., Marullo, S., Artale, V., Vetrano, A., 2013. Seasonal Variability of
772 the Tyrrhenian Sea Surface Geostrophic Circulation as Assessed by Altimeter Data. *Journal of*
773 *Physical Oceanography* 43, 1710–1732. <https://doi.org/10.1175/JPO-D-12-0112.1>

774 Iacono, R., Napolitano, E., Palma, M., Sannino, G., 2021. The Tyrrhenian Sea Circulation: A
775 Review of Recent Work. *Sustainability* 13, 6371. <https://doi.org/10.3390/su13116371>

776 IOC, SCOR and IAPSO, 2010. The international thermodynamic equation of seawater – 2010:
777 Calculation and use of thermodynamic properties.

778 Johannessen, O.M., Lee, O.S., 1974. A deep stepped thermo-haline structure in the
779 Mediterranean. *Deep Sea Research and Oceanographic Abstracts* 21, 629–639.
780 [https://doi.org/10.1016/0011-7471\(74\)90047-3](https://doi.org/10.1016/0011-7471(74)90047-3)

781 Kapelonis, Z., Siapatis, A., Machias, A., Somarakis, S., Markakis, K., Giannoulaki, M.,
782 Badouvas, N., Tsagarakis, K., 2023. Seasonal patterns in the mesopelagic fish community and
783 associated deep scattering layers of an enclosed deep basin. *Sci Rep* 13, 17890.
784 <https://doi.org/10.1038/s41598-023-44765-5>

785 Karl, D.M., Knauer, G.A., Martin, J.H., 1988. Downward flux of particulate organic matter in
786 the ocean: a particle decomposition paradox. *Nature* 332, 438–441.
787 <https://doi.org/10.1038/332438a0>

788 Kermabon, C., Lherminier, P., Le Bot, P., Gaillard, F., 2018. Chaîne Automatisée de Suivi des
789 Courantomètres Acoustiques Doppler Embarqués - Cascade V7.2: Logiciel de validation et de
790 visualisation des mesures ADCP de coque.

791 Kiko, R., Picheral, M., Antoine, D., Babin, M., Berline, L., Biard, T., Boss, E., Brandt, P.,
792 Carlotti, F., Christiansen, S., Coppola, L., De La Cruz, L., Diamond-Riquier, E., Durrieu De
793 Madron, X., Elineau, A., Gorsky, G., Guidi, L., Hauss, H., Irisson, J.-O., Karp-Boss, L.,
794 Karstensen, J., Kim, D., Lekanoff, R.M., Lombard, F., Lopes, R.M., Marec, C., McDonnell,
795 A.M.P., Niemeyer, D., Noyon, M., O'Daly, S.H., Ohman, M.D., Pretty, J.L., Rogge, A.,
796 Searson, S., Shibata, M., Tanaka, Y., Tanhua, T., Taucher, J., Trudnowska, E., Turner, J.S.,
797 Waite, A., Stemmann, L., 2022. A global marine particle size distribution dataset obtained with
798 the Underwater Vision Profiler 5. *Earth Syst. Sci. Data* 14, 4315–4337.
799 <https://doi.org/10.5194/essd-14-4315-2022>

800 Kindler, K., Khalili, A., Stocker, R., 2010. Diffusion-limited retention of porous particles at
801 density interfaces. *Proc. Natl. Acad. Sci. U.S.A.* 107, 22163–22168.
802 <https://doi.org/10.1073/pnas.1012319108>

803 Kiørboe, T., 2001. Formation and fate of marine snow: small-scale processes with large- scale

804 implications. *Sci. mar.* 65, 57–71. <https://doi.org/10.3989/scimar.2001.65s257>

805 Kunze, E., 2003. A review of oceanic salt-fingering theory. *Progress in Oceanography* 56, 399–
806 417. [https://doi.org/10.1016/S0079-6611\(03\)00027-2](https://doi.org/10.1016/S0079-6611(03)00027-2)

807 Maggi, F., 2013. The settling velocity of mineral, biomineral, and biological particles and
808 aggregates in water: SPM SETTLING VELOCITY IN WATER. *J. Geophys. Res. Oceans* 118,
809 2118–2132. <https://doi.org/10.1002/jgrc.20086>

810 McCave, I.N., 1984. Size spectra and aggregation of suspended particles in the deep ocean.
811 *Deep Sea Research Part A. Oceanographic Research Papers* 31, 329–352.
812 [https://doi.org/10.1016/0198-0149\(84\)90088-8](https://doi.org/10.1016/0198-0149(84)90088-8)

813 McCave, I.N., 1975. Vertical flux of particles in the ocean. *Deep Sea Research and*
814 *Oceanographic Abstracts* 22, 491–502. [https://doi.org/10.1016/0011-7471\(75\)90022-4](https://doi.org/10.1016/0011-7471(75)90022-4)

815 Millot, C., Taupier-Letage, I., 2005. Circulation in the Mediterranean Sea, in: Salot, A. (Ed.),
816 *The Mediterranean Sea, Handbook of Environmental Chemistry*. Springer Berlin Heidelberg,
817 Berlin, Heidelberg, pp. 29–66. <https://doi.org/10.1007/b107143>

818 Molcard, R., Tait, R.I., 1977. The steady state of the step structure in the Tyrrhenian Sea, in: A
819 *Voyage of Discovery, George Deacon 70th Anniversary*. Oxford, pp. 221–233.

820 Mrokowska, M.M., 2018. Stratification-induced reorientation of disk settling through ambient
821 density transition. *Sci Rep* 8, 412. <https://doi.org/10.1038/s41598-017-18654-7>

822 Mullison, J., 2017. Backscatter Estimation Using Broadband Acoustic Doppler Current
823 Profilers-Updated.

824 Nagata, T., Kirchman, D.L., 1997. Roles of Submicron Particles and Colloids in Microbial Food
825 Webs and Biogeochemical Cycles within Marine Environments, in: *Advances in Microbial*
826 *Ecology*. Springer, Boston, MA.

827 Newton, P.P., Liss, P.S., 1990. Particles in the oceans (and other natural waters). *Science*
828 *Progress* (1933-) 74, 91–114.

829 Omand, M.M., Govindarajan, R., He, J., Mahadevan, A., 2020. Sinking flux of particulate
830 organic matter in the oceans: Sensitivity to particle characteristics. *Sci Rep* 10, 5582.
831 <https://doi.org/10.1038/s41598-020-60424-5>

832 Peña, M., 2024. Atlantic versus Mediterranean deep scattering layers around the Iberian
833 peninsula. *Progress in Oceanography* 221, 103211.
834 <https://doi.org/10.1016/j.pocean.2024.103211>

835 Picheral, M., Guidi, L., Stemann, L., Karl, D.M., Iddaoud, G., Gorsky, G., 2010. The
836 Underwater Vision Profiler 5: An advanced instrument for high spatial resolution studies of
837 particle size spectra and zooplankton. *Limnology & Ocean Methods* 8, 462–473.
838 <https://doi.org/10.4319/lom.2010.8.462>

839 Prairie, J., Ziervogel, K., Arnosti, C., Camassa, R., Falcon, C., Khatri, S., McLaughlin, R.,
840 White, B., Yu, S., 2013. Delayed settling of marine snow at sharp density transitions driven by
841 fluid entrainment and diffusion-limited retention. *Mar. Ecol. Prog. Ser.* 487, 185–200.
842 <https://doi.org/10.3354/meps10387>

843 Pujo-Pay, M., Durrieu de Madron, X., Conan, C., 2020. PERLE3 cruise, Pourquoi pas ? R/V.
844 <https://doi.org/10.17600/18001342>

845 Radko, T., 2013. Double-diffusive convection. Cambridge university press, Cambridge.

846 Radko, T., Bulters, A., Flanagan, J.D., Campin, J.-M., 2014. Double-Diffusive Recipes. Part I:
847 Large-Scale Dynamics of Thermohaline Staircases. *Journal of Physical Oceanography* 44,
848 1269–1284. <https://doi.org/10.1175/JPO-D-13-0155.1>

849 Radko, T., Smith, D.P., 2012. Equilibrium transport in double-diffusive convection. *J. Fluid*
850 *Mech.* 692, 5–27. <https://doi.org/10.1017/jfm.2011.343>

851 Rinaldi, E., Buongiorno Nardelli, B., Zambianchi, E., Santoleri, R., Poulain, P. -M., 2010.
852 Lagrangian and Eulerian observations of the surface circulation in the Tyrrhenian Sea. *J.*
853 *Geophys. Res.* 115, 2009JC005535. <https://doi.org/10.1029/2009JC005535>

854 Schroeder, K., Ben Ismail, S., Bensì, M., Bosse, A., Chiggiato, J., Civitarese, G., Falcieri M,
855 F., Fusco, G., Gačić, M., Gertman, I., Kubin, E., Malanotte-Rizzoli, P., Martellucci, R., Menna,
856 M., Ozer, T., Taupier-Letage, I., Vargas-Yáñez, M., Velaoras, D., Vilibić, I., 2024. A
857 consensus-based, revised and comprehensive catalogue for Mediterranean water masses
858 acronyms. *Mediterr. Mar. Sci.* 25, 783–791. <https://doi.org/10.12681/mms.38736>

859 Sparnocchia, S., Gasparini, G.P., Astraldi, M., Borghini, M., Pistek, P., 1999. Dynamics and
860 mixing of the Eastern Mediterranean outflow in the Tyrrhenian basin. *Journal of Marine*
861 *Systems* 20, 301–317. [https://doi.org/10.1016/S0924-7963\(98\)00088-8](https://doi.org/10.1016/S0924-7963(98)00088-8)

862 Taillandier, V., Prieur, L., D’Ortenzio, F., Ribera d’Alcalà, M., Pulido-Villena, E., 2020.
863 Profiling float observation of thermohaline staircases in the western Mediterranean Sea and
864 impact on nutrient fluxes. *Biogeosciences* 17, 3343–3366. [https://doi.org/10.5194/bg-17-3343-](https://doi.org/10.5194/bg-17-3343-2020)
865 2020

866 Thiele, S., Fuchs, B.M., Amann, R., Iversen, M.H., 2015. Colonization in the Photic Zone and
867 Subsequent Changes during Sinking Determine Bacterial Community Composition in Marine
868 Snow. *Appl Environ Microbiol* 81, 1463–1471. <https://doi.org/10.1128/AEM.02570-14>

869 Thurnherr, A.M., 2022. How to Process LADCP Data For Vertical Velocity (w) and Derive
870 Parameterized Estimates for Turbulent Kinetic Energy Dissipation (ϵ) using LADCP w
871 Software V2.2.

872 Thurnherr, A.M., 2021. How To Process LADCP Data With the LDEO Software (Version
873 IX.14).

874 Van Der Boog, C.G., Koetsier, J.O., Dijkstra, H.A., Pietrzak, J.D., Katsman, C.A., 2021. Global
875 dataset of thermohaline staircases obtained from Argo floats and Ice-Tethered Profilers. *Earth*
876 *Syst. Sci. Data* 13, 43–61. <https://doi.org/10.5194/essd-13-43-2021>

877 Verso, L., Reeuwijk, M.V., Liberzon, A., 2019. Transient stratification force on particles
878 crossing a density interface. *International Journal of Multiphase Flow* 121, 103109.
879 <https://doi.org/10.1016/j.ijmultiphaseflow.2019.103109>

880 WET Labs, Inc., 2011. Transmissometer C-Star User’s Guide.

881 Yang, S., Zhang, K., Song, H., Ruddick, B., Liu, M., Meng, L., 2024. Disruptions in
882 thermohaline staircases caused by subsurface mesoscale eddies in the eastern Caribbean Sea.
883 *Commun Earth Environ* 5, 408. <https://doi.org/10.1038/s43247-024-01577-3>

884 Zodiatis, G., Gasparini, G.P., 1996. Thermohaline staircase formations in the Tyrrhenian Sea.
885 *Deep Sea Research Part I: Oceanographic Research Papers* 43, 655–678.
886 [https://doi.org/10.1016/0967-0637\(96\)00032-5](https://doi.org/10.1016/0967-0637(96)00032-5)

887

888 **Data Availability Statement**

889 The CTD and ADCP data used in this study are available at
890 <https://doi.org/10.17600/18001342>. The UVP data are available at <https://ecopart.obs-vlfr.fr>.

891

892 **Author Contributions**

893 Conceptualization and Methodology, XDDM, PB and MPP; Writing – Original Draft
894 Preparation, XDDM and PB; Writing – Review & Editing, All.

895

896 **Acknowledgements**

897 This study benefited from the data obtained during the PERLE-3 cruise
898 (<https://doi.org/10.17600/18001342>) with the R/V Pourquoi-Pas? and was supported by the
899 MISTRALS-MERMEX project.

900

901 **Competing interests**

902 The authors declare that they have no conflict of interest.

903

Appendix A

Table A1 Mean abundance (#/L) percentage, and cumulative percentage by size class of UVP between 80 and 2000 mm for all stations between the surface and 2000 m depth

Size class	Abundance (#/L)	Percentage	Cumulative percentage
80.6-102 μm	26.272	45.617	45.62
102-128 μm	10.465	18.171	63.79
128-161 μm	5.547	9.631	73.42
161-203 μm	8.542	14.832	88.25
203-256 μm	3.13	5.435	93.68
256-323 μm	1.944	3.375	97.06
323-406 μm	0.985	1.710	98.77
406-512 μm	0.391	0.679	99.45
512-645 μm	0.169	0.293	99.74
645-813 μm	0.078	0.135	99.88
0.813-1020 μm	0.038	0.066	99.94
1.02-1290 μm	0.018	0.031	99.98
1.29-1630 μm	0.009	0.016	99.99
1630-2050 μm	0.005	0.009	100.00