

1 **Short ~~lived~~^{times} ~~of~~ organic nitrates in a sub-urban temperate**
2 **forest: An indication of efficient assimilation of reactive nitrogen**
3 **by the biosphere?**
4

5 Simone T. Andersen¹, Rolf Sander¹, Patrick Dewald¹, Laura Wüst¹, Tobias Seubert¹, Gunther N.
6 T. E. Türk¹, Jan Schuladen¹, Max R. McGillen², Chaoyang Xue², Abdelwahid Mellouki^{2,3},
7 Alexandre Kukui⁴, Vincent Michoud⁵, Manuela Cirtog⁵, Mathieu Cazaunau⁵, Astrid Bauville⁵,
8 Hichem Bouzidi⁵, Paola Formenti⁵, Cyrielle Denjean⁶, Jean-Claude Etienne⁶, Olivier Garrouste⁶,
9 Christopher Cantrell⁷, Jos Lelieveld¹, John N. Crowley¹

10 ¹Atmospheric Chemistry Department, Max-Planck-Institute for Chemistry, 55128-Mainz,
11 Germany

12 ²Institut de Combustion, Aérothermique, Réactivité Environnement (ICARE), CNRS, 1C Avenue
13 de la Recherche Scientifique, CEDEX 2, 45071 Orléans, France

14 ³University Mohammed VI Polytechnic (UM6P), Lot 660, Hay Moulay Rachid Ben Guerir, 43150,
15 Morocco

16 ⁴Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), CNRS Orléans,
17 France

18 ⁵Université Paris Cité and Univ Paris Est Creteil, CNRS, LISA, F-75013 Paris, France

19 ⁶CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France

20 ⁷Univ Paris Est Creteil and Université de Paris Cité, CNRS, LISA, F-94010 Créteil, France

21 *Correspondence to:* Simone T. Andersen (simone.andersen@mpic.de) and John N. Crowley
22 (john.crowley@mpic.de)
23

1 Abstract

Alkyl nitrates (ANs) and peroxy-carboxylic nitric anhydrides (PANs) are important reservoirs of reactive nitrogen that contribute significantly to the rate of formation and growth of secondary organic aerosols and support the transport of reactive nitrogen from polluted areas to remote areas. It is therefore critical to understand their sources and sinks in different environments. In this study we use measurements of OH, O₃, NO₃ reactivity, VOCs, ΣANs and ΣPANs during the ACROSS (Atmospheric ChemistRy Of the Suburban foreSt) campaign to investigate different production and loss processes of ANs and PANs in a temperate forest. At daytime OH-initiated processes were the dominant source of ANs (69-72 %) followed by NO₃ (18-20 %) and O₃ (8-12 %). At nighttime the contribution from OH decreased to 43-53 %, and NO₃ increased to 26-40 % with that of O₃ largely unchanged. Of the measured ΣPANs, 48-78 % was modelled to be peroxyacetic nitric anhydride (PAN, commonly known as peroxyacetyl nitrate). Physical loss (e.g. deposition) was an important sink for both ANs and PANs and contributed significantly to the very short lifetimes of 1 - 41.5 ± 1 h for ANs and 0.08 - 1.5 h for PANs observed during the campaign.

2 Introduction

Approximately 1000 Tg of biogenic volatile organic compounds (BVOCs) are emitted into the atmosphere annually, whereof ~50% is isoprene and ~15% are monoterpenes (Guenther et al., 2012). The oxidation of BVOCs is initiated by ozone (O₃) and the hydroxyl (OH) and nitrate (NO₃) radicals. When BVOCs are oxidized in the presence of (largely anthropogenic) NO_x they lead to the formation of alkyl nitrates (RONO₂, ANs) and peroxy-carboxylic nitric anhydrides (RC(O)O₂NO₂, PANs) as illustrated in Figure 1. Note that the term PANs includes the most common carboxylic nitrate (CH₃C(O)O₂NO₂, commonly referred to as PAN). ANs are produced in the daytime in a minor branch of the reaction between organic peroxy radicals (RO₂) which do not have an α-carbonyl group and nitrogen oxide (NO, Lightfoot et al. (1992)) and through the reaction of unsaturated BVOCs with NO₃ at nighttime. Recent studies suggest that the NO₃-initiated oxidation of BVOCs during the day could also be an important pathway for the formation of alkyl nitrates (Liebmann et al., 2019; Ayres et al., 2015; Liebmann et al., 2018a; Liebmann et al., 2018b; Dewald et al., 2024). PANs are formed when α-carbonyl peroxy radicals (RC(O)O₂) react with nitrogen dioxide (NO₂). The stability of PANs is highly temperature dependent, resulting in boundary layer lifetimes of the order of hours at temperate mid-latitudes with respect to dissociation to RC(O)O₂ and NO₂ (IUPACupae, 2024).

The yield of alkyl nitrates in the RO₂ + NO reaction strongly depends on the composition of BVOCs in the atmosphere, the oxidant that initiates the degradation of the BVOC (OH, O₃ or NO₃), and the ratio of NO to other reactants with which the RO₂ can react such as NO₂, HO₂ and other RO₂ (IUPACupae, 2024; Perring et al., 2013; Wennberg et al., 2018; Hallquist et al., 1999; Fry et al., 2014). The yield of the PANs formed from the corresponding α-carbonyl peroxy radical depends only on the fraction of RC(O)O₂ that reacts with NO₂ as opposed to reacting with HO₂ or NO. Production and loss processes of ANs and PANs are described in detail through reaction mechanisms and equations in section 4.1-4.3 and 4.6, respectively.

The formation of ANs and PANs serves to sequester reactive nitrogen (NO_x = NO + NO₂) into reservoir species, which can release NO_x following transport to regions remote from NO_x sources; both can also be removed from the lowermost atmosphere through deposition, which thus represents a sink of NO_x in the atmosphere. As NO₂ is formed from primary emitted NO

and its photolysis drives the formation of O₃, understanding the fate of NO_x is critical for predicting O₃ levels in the troposphere. It has additionally been shown that ANs contribute significantly to the rate of formation and growth of secondary organic aerosols (SOA), and thereby impacting human health and air quality (Hallquist et al., 2009; Shiraiwa et al., 2017; Kanakidou et al., 2005).

This study investigates the production and loss of PANs and ANs in an anthropogenically impacted temperate forest using field measurements of BVOCs, oxidants (OH, O₃, and NO₃), products (PANs and ANs), and meteorological data. The lifetimes of PANs and ANs are derived from their production rates and measured total mixing ratios.

3 Methodology

3.1 Site Description

The measurements used in this study were conducted at the Rambouillet forest site located approximately 50 km southwest of Paris, France, (48.687, 1.704) during the ACROSS (Atmospheric Chemistry Of the Suburban forest) campaign between June 13th 2022 and July 25th 2022 (Cantrell and Michoud, 2022). The forest consists of approximately 70% oak, 20% pine, and small contributions from beech and chestnut. The top of the forest canopy around the site was approximately 20-25 m. A 40 m measurement tower and multiple containers with a large variety of instruments were located in a roughly quadratic clearing (26.5 m × 26.3 m, ~697 m²) as shown in Figure S1. 48h HYSPLIT back trajectories showed that the airmasses sampled during the campaign passed either over the Atlantic Ocean or Continental regions in Europe before reaching the site (Draxler and Rolph, 2011; Andersen et al., 2024). All the instruments used in this study are described briefly below.

3.2 Measurements

3.2.1 Reactive Nitrogen and O₃

NO₂, total peroxy-carboxylic nitric anhydrides (ΣPANs), total alkyl nitrates (ΣANs), NO₃ reactivity (k^{NO_3}), and O₃ were measured by instruments inside the MPIC (Max Planck Institute for Chemistry) container with co-located inlets sampling from a high-volume-flow stainless steel tube (10 m³ min⁻¹; 15 cm diameter, 0.2 s residence time) taking air from a height of 5.4 m above ground.

A 5-channel thermal dissociation cavity-ringdown spectrometer (5CH-TD-CRDS, Sobanski et al. (2016)) was used to measure NO₂, ΣPANs, and ΣANs. NO₂ was measured directly at 408 nm with a limit of detection (LOD) of 9.7 parts per trillion by volume (pptv) for 1 min averaging (3σ) and a total uncertainty of 7% + 9.7 pptv + (20 pptv * RH/100). ΣPANs and ΣANs were thermally dissociated to NO₂ by heating their separate inlets to 448 and 648 K, respectively, followed by detection of NO₂ at 408 nm. Numerical simulations were run for both ΣPANs and ΣANs to correct for NO₂ loss via recombination with RO₂, the reactions of peroxy radicals with ambient NO, NO oxidation to NO₂ by O₃, and pyrolysis of O₃ (Thieser et al., 2016; Sobanski et al., 2016). The majority of the correction factors for both ΣPANs and ΣANs were between 0.9 and 1.2 as shown in Figure S4₂, which is consistent with low NO_x levels. The LOD was determined to be 6.3 pptv and 8.6 pptv for ΣPANs and ΣANs for 10 min averaging (3σ of the noise on the zero measurements), respectively, and the total uncertainty

was 21% + 6.3 pptv and 28% + 8.6 pptv. A timeseries of NO₂, ΣPANs, and ΣANs is shown in Figure 2. Two additional cavities, operated at 662 nm, measured NO₃ and (via thermal dissociation to NO₃, 373 K) N₂O₅ (Sobanski et al., 2016). NO₃ was throughout the campaign below the LOD of 0.25 pptv for 1 min averaging and N₂O₅ was only detected above the LOD of 0.9 pptv on 5 nights and only one of those nights had consistently high N₂O₅ for more than an hour. NO₃ and N₂O₅ data are therefore not presented or used in this study. ~~A timeseries of NO₂, ΣPANs, and ΣANs is shown in Figure 2.~~

A second CRDS-instrument was used ~~to~~ primarily to measure the NO₃ reactivity towards VOCs (~~k^{NO3}~~) in the forest, but it also has a cavity operated at 405 nm for the measurement of NO₂ (Liebmann et al., 2018b). The ambient NO₃ reactivity was quantified by a CRDS-measurement (at 662 nm) of in-situ-generated NO₃ after its residence in a flow-tube reactor when mixed with either synthetic or ambient air. A numerical simulation procedure was used to correct the measurements for competing reactions taking place inside the flow-tube in order to extract the VOC contribution to the measured NO₃ consumption. A detailed analysis of the NO₃ reactivity measurements is presented in Dewald et al. (2024). During the ACROSS campaign, the reactivity of NO₃ towards organics was dominated by those of biogenic origin, so henceforth we refer to this as k^{BVOC} .

O₃ was measured with a commercial instrument (2B Technologies model 205) using UV absorption at ~254 nm. The LOD is 2 ppbv for 10 s averaging time. A timeseries of O₃ can be observed in Figure 2.

NO was measured using a commercial chemiluminescence instrument (Ecophysics CLD 780 TR, henceforth CLD) with an LOD of 10 pptv for 1 min averaging time. The sampling height for NO measurements was about 3.2 m above the ground surface and the inlet was approximately 17 m from the MPIC container at the ICARE-LPC2E container in Figure S1. The NO measurements were corrected for a change in the CLD sensitivity during the campaign caused by an interruption in the instrument's oxygen supply as described in Andersen et al. (2024).

3.2.2 OH and XO₂ (HO₂+RO₂)

The OH radical was measured by its conversion (via reaction with isotopically labelled SO₂) to H₂SO₄ which was subsequently detected using nitrate chemical ionization mass spectrometry (Eisele and Tanner, 1991). The lower limit of detection for OH radicals at signal-to-noise-ratio (S/N)=3 and a 15-minute integration time was 5×10⁴ molecule cm⁻³. The sum of peroxy radicals, XO₂=HO₂+RO₂, was measured by their conversion to OH in the presence of NO. The OH calibration coefficient was determined using N₂O actinometry and OH generation in a turbulent flow reactor by photolysis of N₂O or H₂O at 184.9 nm (Kukui et al., 2008). The calibration of HO₂, CH₃O₂ and other RO₂ was performed by adding into the calibration reactor CO, CH₄ (or other RO₂ precursors) converting OH to RO₂. The overall estimated calibration accuracy (2σ) for OH is about 25% and about 30% for calibrated XO₂, although the uncertainty of the XO₂ measurements is typically higher due to variable detection efficiency (i.e. yields of OH) of different XO₂. Peroxy radicals derived from terpene oxidation will be detected with a lower efficiency than e.g. short chained RO₂ so that the XO₂ measurements during ACROSS should be regarded as lower limits. The lower limit of detection for XO₂ radicals at S/N=3 and a 4-minute integration time is 2×10⁶ molecule cm⁻³.

The background signal is determined by adding OH and XO₂ scavenger, NO₂ in this case, before their conversion to H₂SO₄. As discussed in Kukui et al. (2021), formation of H₂SO₄ in the reaction of stabilized Criegee intermediates (SCI) with SO₂ in the conversion reactor may lead to some positive interference in radical measurements. However, this interference estimated to correspond to several 10⁴ cm⁻³ of radicals concentrations is negligible compared to measured OH concentrations. A detailed description of the instrument and calibration system have been presented elsewhere (Kukui et al., 2008; Kukui et al., 2021). During the ACROSS field campaign the instrument was installed in the ICARE-LPC2Ea shipping container with the chemical conversion reactor fixed to the roof of the container via an interface cap covered with a polytetrafluorethylene (PTFE) sheet. The sampling aperture of the chemical conversion reactor (3 mm diameter) was positioned 50 cm above the roof and about 3 m above the ground surface. A detailed analysis of the OH measurements will be presented in a forthcoming publication.

3.2.3 Photolysis frequencies and meteorology

Spectral radiometers (Mettcon GmbH) were installed near the high-volume-flow stainless-steel tube on top of the MPIC container and on top of the tower to measure actinic fluxes, which were converted to photolysis frequencies using recommended absorption spectra and quantum yields (IUPAC, 2024; Burkholder et al., 2020) as described in Meusel et al. (2016). Note that upwelling radiation is not accounted for resulting in a potential underestimation of the photolysis frequencies of 5-10 %. A comparison of the two measurement heights is shown in Figure S3. It is clear that the measurements below the canopy are significantly impacted by shading from trees in the morning and afternoon as well as by the tower around midday. Differences between the two measurements when both instruments are in direct sunlight is caused by integration of solar flux over one complete hemisphere (basically above the height of the integrating dome), which is influenced by trees at angles close to horizontal.

Ambient temperature was measured at four different heights on the tower; 5 m, 13 m, 21 m, and 41 m using temperature sensors from Atexis (PT1000) and Thermoest (PT100). Relative humidity was measured at 5 m using a Vaisala humidity sensor (HMP45A).

3.2.4 Biogenic Volatile Organic Compounds (BVOCs)

The LISA (Laboratoire Interuniversitaire des Systèmes Atmosphériques) Proton Transfer Reaction Time of Flight Mass Spectrometer (PTR-ToF-MS, hereafter called PTRMS), manufactured by Kore Technology Ltd., was used for monitoring concentrations of VOCs. Air samples were drawn at a flow rate of approximately 300 mL min⁻¹ through a 3 m long Silcosteel® coated stainless steel tube (2.1 mm inner diameter) from an inlet height of 4.6 m in the LISA observation container in Figure S1. Calibration was performed approximately every 3 days using VOC standards (5 – 20 ppb) from a certified National Physical Laboratory (NPL) calibration mixture with nominally 1 ppmv ±5% of several trace gases including acetaldehyde, methanol, ethanol, isoprene, acetone, dimethyl sulphide, acetonitrile, and 3-carene. Humidity corrections were applied for each trace-gas. Mixing ratios of non-calibrated trace gases were retrieved from reactor conditions, rate constants, fragmentations, and ion

transmissions determined using the same NPL standard cylinder. The time series of acetaldehyde, isoprene and total monoterpenes (Σ MT) is shown in Figure 2.

As there were no measurements of speciated monoterpenes, due to instrumental issues with the gas chromatography instrument deployed during the campaign, different potential monoterpene mixtures were determined using ~~the total mixing ratios measured by the PTRMS together with~~ the measured reactivity of NO_3 towards BVOCs (k^{BVOC}) after subtracting the reactivity due to isoprene ($k_{\text{NO}_3+\text{isoprene}}[\text{Isoprene}]$) as described in Eq. (1). Here, $k_{\text{effective}}$ is an effective rate coefficient for the reaction of NO_3 with an assumed monoterpene mixture. $k_{\text{effective}}$ was determined by adding all the fractional contributions (a_i) from different monoterpenes as described in equation (2), where ~~a_i and k_{NO_3+i} are the fractional contribution and the rate coefficient with NO_3 for monoterpene i , respectively,~~ which are listed in Table 1. The estimated Σ MT from equation (1) using different $k_{\text{effective}}$ can then be compared to the total mixing ratios measured by the PTRMS.

$$[\Sigma \text{Monoterpenes}] = \frac{k^{\text{BVOC}} - k_{\text{NO}_3+\text{isoprene}}[\text{Isoprene}]}{k_{\text{effective}}} \quad (1)$$

$$k_{\text{effective}} = \sum_i (a_i \times k_{\text{NO}_3+i}) \quad (2)$$

Only limonene, α -pinene, and β -pinene were used to determine ~~the~~ potential mixtures since the box-model used (see below) contained schemes for their degradation only. Four examples of potential mixtures constrained by the measured NO_3 -reactivity and consistent (for parts of the day) with the measured sum of monoterpenes (Σ MT) from the PTRMS were determined ~~Bby~~ varying the fractional contribution (a_i) of limonene between 10 and 30% of the sum of monoterpenes (Σ MT) ~~three potential mixtures which were consistent with the measured NO_3 -reactivity were determined.~~ Mixture 1: 30% limonene, 10% β -pinene, and 60% α -pinene. Mixture 2: 20% limonene, 15% β -pinene, and 65% α -pinene. Mixture 3: 10% limonene, 5% β -pinene, and 85% α -pinene. Mixture 4: 57% β -pinene and 43% α -pinene. All ~~four~~three scenarios are plotted together with the measured Σ MT over a 48-hour period in Figure 3B. Here the three scenarios, which include limonene, can be observed to agree well with the measurements except when a temperature inversion occurred (Figure 3A). The same kind of discrepancy is observed every night with a temperature inversion, which is shown in Figure 3C, where mixture 2 is plotted against the measured Σ MT for the entire campaign and coloured by the difference in temperature (ΔT) between the top of the tower (41 m) and measurement height (5m). A significant temperature inversion can result in the formation of a shallow nocturnal surface layer with weak vertical mixing. In contrast, the fourth mixture, which does not include limonene, does agree well with the measured Σ MT when a temperature inversion is observed, but it would require a much larger (than measured by the PTRMS) total mixing ratio of monoterpenes when no temperature inversion is observed.

As BVOCs continue to be emitted at night, weak vertical mixing leads to a strong gradient in monoterpene mixing ratios with higher values at low heights above ground level. As the gradient (both horizontally and vertically) will depend on the lifetime of each monoterpene, the mixture of monoterpenes measured by the NO_3 -reactivity instrument (at 5.4 m) might not be the same as that emitted by the vegetation at different heights. In Figure S42, the average derived diel profiles of the lifetime of α -pinene, β -pinene, and limonene when taking reactions with OH, O_3 , and NO_3 into account using the rate coefficients in Table 1 are plotted. During the daytime all three monoterpenes have short lifetimes of around 0.5-1.5 hours; however, at

nighttime, limonene clearly has the shortest lifetime, whereas β -pinene has a slightly longer lifetime than α -pinene. On nights with $\Delta T > 1^\circ\text{C}$ (26% of the total measurements between June 17th 2022 and July 22nd 2022), limonene is therefore assumed not to be sampled by the NO_3 -reactivity instrument due to the slow vertical and horizontal mixing and ~~the~~ monoterpene mixture 1-3 ~~are~~^{is} changed to 57% β -pinene and 43% α -pinene (Mixture 4), which can be seen in Figure 3D for ~~Mixture scenario~~ 2. This correction aligns the calculated and measured ΣMT throughout the entire campaign (Figure 3E). It should also be noted that the NO_3 -reactivity and PTRMS measurements were not co-located and there could therefore also be a gradient between the measurements, which we do not take into account as the magnitude of this gradient is unknown.


3.3 Box Model

To simulate PANs numerically, we have used the atmospheric chemistry box model CAABA/MECCA (Chemistry As A Boxmodel Application/Module Efficiently Calculating the Chemistry of the Atmosphere) by Sander et al. (2019). The code is based on model version 4.7.0, and it has been adapted to simulate the ACROSS campaign. To allow for a detailed calculation of monoterpenes and PANs, reactions were exported from the Master Chemical Mechanism (MCM, <https://mcm.york.ac.uk>), including the MCM species APINENE, BPINENE, C_3H_8 , C_5H_8 , CH_3CHO , CH_4 , LIMONENE, and NC_4H_{10} in the marklist. This resulted in a gas-phase chemical mechanism with 1536 species and 4550 reactions. The setup of the individual model runs will be described in Sect. 4.7.

4 Results and Discussion

4.1 AN production from NO_3 reactions with BVOCs

NO_3 radicals are produced from the reaction between NO_2 and O_3 (R1) and are usually lost rapidly during daytime to photolysis and reactions with unsaturated BVOCs and NO (R2-R4). Reactions between NO_3 radicals and unsaturated BVOCs lead to the formation of alkyl nitrates (ANs, R3a) as well as other products (R3b). At nighttime, ground-level NO_3 mixing ratios can vary greatly from < 1 pptv to > 100 pptv depending on atmospheric composition (Ng et al., 2017; Brown and Stutz, 2012), ~~i.e. their production and loss terms~~. During ACROSS, the NO_3 reactivity within the canopy was generally high at nighttime due to the emission of biogenic volatile organic compounds (BVOCs, by vegetation) and NO (from soil) into a shallow, poorly mixed nocturnal surface layer (Dewald et al., 2024; Andersen et al., 2024). This led to NO_3 mixing ratios, at the ground, of < 0.5 pptv, which were generally lower than the limit of detection of instrumentation at the site.



The total production rate of ANs from the NO₃-initiated oxidation of unsaturated BVOCs can be calculated using equation (3), where [NO₃]_{ss} is the NO₃ concentration at steady state and $\alpha_i^{\text{NO}_3}$, $k_i^{\text{NO}_3}$ and [C_i] are the ANs yield, the rate coefficient and BVOC concentration for compound *i*, respectively. [NO₃]_{ss} is determined from the production and loss terms described by reactions (R1-R4) as described in equation (4), where [NO], [NO₂], and [O₃] are the concentrations of NO, NO₂, and O₃, respectively, k_1 and k_4 are the rate coefficients of reaction (R1) and (R4), respectively, k^{BVOC} is the first-order loss frequency for NO₃ towards BVOCs, and J_{NO_3} is the photolysis frequency of NO₃ radicals (Liebmann et al., 2019). This calculation ignores physical losses of NO₃ (e.g. deposition) which will not compete with its reactive losses in this environment.

$$\sum P_{\text{ANs}}^{\text{NO}_3+\text{OH}} = [\text{NO}_3]_{\text{ss}} \sum_i \alpha_i^{\text{NO}_3} k_i^{\text{NO}_3} [\text{C}_i] \quad (3)$$

$$[\text{NO}_3]_{\text{ss}} = \frac{k_1[\text{NO}_2][\text{O}_3]}{k^{\text{BVOC}} + J_{\text{NO}_3} + k_4[\text{NO}]} \quad (4)$$

4.2 AN production from OH reactions with BVOCs

At daytime, primary OH radicals are produced e.g. from the photolysis of O₃ followed by the reaction between O(¹D) and water vapour (R5-R6) with secondary production through reaction of HO₂ (formed in peroxy-radical (RO₂) reactions) with NO. In the absence of photochemistry, OH concentrations are generally lower at nighttime than at daytime, with average hourly concentrations of around $3.5\text{--}5 \times 10^5$ molecules cm⁻³ observed across the campaign. The most important nighttime source of OH radicals is generally believed to be the reaction between unsaturated VOCs and ozone e.g. (R11-R12). As will be shown in a forthcoming publication, the nighttime OH levels are broadly consistent with measured OH-reactivity, HO₂-recycling via reaction of HO₂ with NO and production via the ozonolysis of terpenoids. The nighttime OH results from the ozonolysis of unsaturated BVOCs (see below) in the nocturnal boundary layer, which was very shallow and stable due to temperature inversions (Andersen et al., 2024). When OH reacts with BVOCs in the presence of O₂, peroxy radicals are produced (R7), which can then react with NO to give alkyl nitrates (R8a), as well as alkoxy radicals and NO₂ (R8b). Additional competing processes that lower the yield of ANs from RO₂ are reactions with itself and other RO₂ (R9) and reaction with HO₂ (R10).



The total production rate of ANs from OH-initiated oxidation of BVOCs is described in equation (5), where [OH] is the OH concentration, $\alpha_i^{\text{RO}_2+\text{NO}}$ is the fraction of the organic

peroxy radicals from BVOCs which (via R8a) forms an alkyl nitrate when reacting with NO, k_i^{OH} and $[C_i]$ are the rate coefficient and BVOC concentration for compound i , respectively, and β is the fraction of peroxy radicals that reacts with NO (rather than RO₂ or HO₂) as described in equation (6) (Liebmann et al., 2019). β was calculated using the measured XO₂ (HO₂+RO₂) and NO together with a generic rate coefficient for reaction (R8), $8 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, and a generic rate coefficient for the combination of reactions (R9) and (R10), $k_{9/10} = 1 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (IUPACupae, 2024; Lightfoot et al., 1992). k_8 was set to this value because the vast majority of organic peroxy radicals react with NO with a rate coefficient of $8 \pm 1 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (IUPAC, 2024; Lightfoot et al., 1992). Since we do not have separate measurements of RO₂ and HO₂ an effective rate coefficient of $1 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ was chosen. This was derived by considering the MCM rate coefficients ($2.3 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ at 298 K) for reaction (R10) and $2\text{--}3 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ for reaction between isoprene derived peroxy radicals and other RO₂ and calculating the geometric mean. As we do not know the contributions of HO₂ and RO₂ to XO₂ this is clearly a coarse approximation. We therefore undertook a sensitivity test in which the effective rate coefficient was varied by a factor of two in both directions to establish the uncertainty in β using median mixing ratios across the entire campaign of NO (41 pptv) and XO₂ ($6 \times 10^8 \text{ molecule cm}^{-3}$). When halving the effective rate coefficient β is increased by 30% and thereby increasing the production rate and when doubling the effective rate coefficient β is decreased by 30% and thereby decreasing the production rate.

$$\sum \text{P}_{\text{ANS}}^{\text{OH}} = [\text{OH}] \beta \sum_i \alpha_i^{\text{RO}_2+\text{NO}} k_i^{\text{OH}} [C_i] \sum_i \alpha_i^{\text{RO}_2} / k_i^{\text{OH}} [C_i] \quad \text{---}$$

(5)

$$\beta = \frac{k_8[\text{NO}]}{k_8[\text{NO}] + k_9[\text{RO}_2] + k_{10}[\text{HO}_2]} = \frac{k_8[\text{NO}]}{k_8[\text{NO}] + k_{9/10}[\text{XO}_2]} \quad \text{---}$$

(6)

4.3 AN production from O₃ reactions with BVOCs

Ozone (O₃) addition to an unsaturated BVOC forms a primary ozonide (POZ, R11), which in the presence of O₂ can rapidly decompose via Criegee intermediates to OH and organic peroxy radicals (RO₂) (R12a). The POZ can also react through other processes that do not result in organic peroxy radicals (R12b). The RO₂ formed in (R12a) further reacts through reactions (R8-R10) as described above.



Equation (7) describes the total production rate of ANs from the O₃-initiated oxidation of unsaturated BVOCs, where [O₃] is the O₃ concentration, $\alpha_i^{\text{O}_3}$ is the yield of RO₂ from reactions (R11-R12a), $\alpha_i^{\text{RO}_2+\text{NO}}$ is the fraction of the organic peroxy radicals (from BVOCs) formed in (R12a) which when reacting with NO form an alkyl nitrate, $k_i^{\text{O}_3}$ and $[C_i]$ are the rate coefficient and BVOC concentration for compound i , respectively, and β is the fraction of peroxy radicals that react with NO as described in equation (6) (Liebmann et al., 2019), calculated as described

in section 4.2. Here we assume that the yield of ANs from RO₂ + NO is independent of whether RO₂ is formed by OH or O₃-initiated oxidation.

$$\sum P_{\text{ANs}}^{\text{O}_3} = [\text{O}_3] \beta \sum_i \alpha_i^{\text{O}_3} \alpha_i^{\text{RO}_2+\text{NO}} k_i^{\text{O}_3} [\text{C}_i] \quad (7)$$

4.4 Relative importance of OH, O₃ and NO₃ oxidation for the production of ANs

To analyse the production and loss processes of alkyl nitrates during the ACROSS campaign, the measurements have been separated into two phases: phase 1 (lower photochemical activity) from June 28th 2022 to July 7th 2022, and phase 2 (higher photochemical activity) from July 8th 2022 to July 20th 2022. Average diel profiles of O₃, NO₂, NO, temperature, OH radicals, and isoprene for the two phases are plotted in Figure 4. Phase 1 is characterised by low levels of oxidants and organics and a maximum average daytime temperature around 25 degrees Celsius, whereas phase 2 is characterised by higher levels of oxidants and organics due to generally higher temperatures (maximum average daytime temperature around 30 degrees Celsius). The average monoterpene mixtures for the two phases are displayed in Figure 5 for ~~the mixture 2~~ with 20% limonene, 15% β-pinene, and 65% α-pinene when no temperature inversion is observed and 57% β-pinene and 43% α-pinene when temperature inversions higher than 1°C is observed. The monoterpene mixtures derived for Mixture 1 and 3 using 57% β-pinene and 43% α-pinene when temperature inversions higher than 1°C is observed scenario 1 and 3 are plotted in Figure S53. At daytime the average mixture reflects the percentages used for each scenario, whereas at nighttime, the mixture is dominated by β-pinene and α-pinene during both phases, associated with the temperature inversions observed on many nights.

The diel profile of the \sum ANs production rates from NO₃-, OH-, and O₃-initiated oxidation ~~for mixture 2~~ using the rate coefficients and yields in Table 1 and the monoterpene mixture in Figure 5 is shown in Figure 6A and B for phases 1 and 2, respectively. The total ANs production during phase 1 is fairly constant at 35-75 pptv h⁻¹ throughout the diel profile. In contrast, phase 2 shows large variation with 80-100 pptv h⁻¹ in the early morning between 01:00-05:00 UTC (03:00-07:00 LT) and 220-280 pptv h⁻¹ around late morning to midday between 07:00-12:00 UTC (09:00-14:00 LT). For both phases, the variability in the \sum ANs production rate from O₃ and NO₃-initiated oxidation of BVOCs is small compared to that of OH-initiated oxidation. The calculated ~~averaged~~ O₃-initiated \sum ANs production rate ~~varied from 4-15 to 9 ± 3 and 12-2518 ± 4 (±1σ)~~ pptv h⁻¹ for phase 1 and 2, respectively, and the derived ~~averaged~~ NO₃-initiated \sum ANs production rate ~~varied from 4-2212 ± 5 and 22-6241 ± 11 (±1σ)~~ pptv h⁻¹ for phase 1 and 2, respectively. The differences between the two phases are therefore caused by the OH-initiated oxidation, which is relatively stable during phase 1 at ~~13-6134 ± 11 (±1σ)~~ pptv h⁻¹ across the diel profile on average due to the lower daytime levels of OH and BVOCs (see Figure 4), but varies between 30-70 pptv h⁻¹ at nighttime and 100-230 pptv h⁻¹ at daytime during phase 2. The production rates are associated with uncertainties, which have been evaluated using the average midday measurements and derived monoterpene mixing ratios in table S1, the uncertainties in rate coefficients (see table 1 and S2), measurement uncertainties and yield uncertainties (see table S2) for the three monoterpene mixtures containing limonene. The results are summarized in table S3 and show that the uncertainty varies only a little within the different monoterpene mixtures. At midday the propagated uncertainty of the NO₃-, O₃-, OH-initiated production rate, and the total production rate is around 35%, 40%, 30%, and 26%, respectively.

Figure 6C and D show the fractional contributions of NO₃-, OH-, and O₃-initiated oxidation (~~mixture 2~~) to the ΣANs production rate for phase 1 and 2, respectively. OH clearly dominates at daytime (06:00-18:00 UTC) with, on average, 69-72 % for both phases, followed by NO₃ with 18-20 % and O₃ with 8-12 %. At nighttime (18:00-06:00 UTC), the picture is not as clear: During phase 1 OH-initiated oxidation is still the dominant ANs production pathway with, on average, 53 % and the remainder is close to evenly split at 21 and 26 % between O₃- and NO₃-initiated oxidation, respectively. At nighttime during phase 2, OH- and NO₃-initiated oxidation contributed similarly with 43 and 40 % of the total ANs production rate, respectively, leaving only 17 % for O₃-initiated processes. All the fractional contributions are associated with uncertainties around 40-50% when propagating the uncertainties in the individual and total production rates at midday. The differences between the two phases at nighttime can be explained by the availability of the precursors, where there is approximately double the amount of O₃ and NO₂ during phase 2, leading to a higher production rate of NO₃ radicals and thereby an increased ANs production rate from NO₃-initiated oxidation.

Both phases give significantly different fractions at both daytime and nighttime ~~to~~ from those observed by Liebmann et al. (2019) in a boreal forest, where, in the absence of measurements, OH was calculated from the actinic flux, which thus resulted in zero OH at nighttime. However, both studies agree on NO₃ oxidation being a significant source of ANs, both at daytime and nighttime. If [NO₃]_{ss} was calculated using photolysis frequencies measured above the clearing instead of inside the clearing, the contribution from NO₃-initiated oxidation would be reduced in the morning and evening (see Figure S3 for comparison of J_{NO_3} above (tower) and inside (ground) the clearing. Further as discussed by (Dewald et al., 2024) the relative contributions of e.g. OH and NO₃ would be significantly modified to favour NO₃ if we consider the greatly reduced photolysis frequencies of NO₃ and OH precursors in non-cleared parts of the forest.

4.5 ANs loss and lifetime

Neglecting the role of transport, we now combine the diel profile of the total production rate of ANs (described above) with the average measured diel profile of ANs to evaluate the loss processes and lifetime of the ANs using equation (8). P_{ANs} is the production rate of ANs, $[\text{ANs}]_0$ is the average ANs mixing ratio at 00:00 UTC, and $k_L(\text{ANs})$ is the loss rate of the ANs, which is defined as the inverse of the lifetime of the ANs ($(\tau_{\text{ANs}})^{-1}$).

$$\frac{d[\text{ANs}]}{dt} = P_{\text{ANs}} - k_L(\text{ANs})[\text{ANs}]_0 \quad (8)$$

The ANs mixing ratio at any subsequent time to $[\text{ANs}]_0$ can then be calculated as described in Eq. (9) with the variation of $k_L(\text{ANs})$ to match the observed ANs mixing ratio.

$$[\text{ANs}]_t = \int_0^t \frac{d[\text{ANs}]}{dt} + [\text{ANs}]_0 \quad (9)$$

Figure 6E and F show the average diel profiles of the measured ANs in black for phase 1 and 2, respectively. The orange lines show how the ANs mixing ratios would have increased if there was no chemical or transport-induced loss throughout the day and the blue lines show how the diel profiles would look when applying lifetimes of 1-10 hours for the ANs. The best fit to the measured diel profile of ANs results from using an effective lifetime in the clearing of 1.5 ± 1 h throughout the diel cycle for both phases despite the very different production rates described above. Table 2 gives an overview of the average daytime and nighttime loss rate

frequencies and the resulting effective lifetimes. No difference was observed between daytime and nighttime during phase 1 and only a small difference within the uncertainties was observed during phase 2.

The total ANs production rate resulting from the three different monoterpene mixtures derived ~~above~~ in section 3.2.4 (–plotted in Figure S4A–S6A for the two phases), clearly shows that the choice of mixture does not have a significant impact on the total production rate. This means that the determined lifetime for the two phases of 1.5 ± 1 h is consistent across all three mixtures, as can be observed in Figure S4C–S6C. This lifetime is similar to the 2 ± 3 h estimated in a boreal forest (Liebmann et al., 2019), and that determined for isoprene nitrates using flux measurements over the Ozark Mountains (Wolfe et al., 2015).




In section 4.7, it is shown that the box model described ~~above~~ in Section 3.3 predicts the mixing ratio of XO_2 to be up to 4 times higher than the measured XO_2 depending on the day, which is related in part to the variable detection efficiency of different RO_2 by the XO_2 instrument. Strictly speaking the XO_2 measurements represent a lower limit in regions with high biogenic contribution to RO_2 . Owing to the large uncertainty associated with the measurements ~~As the origin of this model-observation discrepancy is not known~~ we have also calculated the total ANs production rate and the lifetime of ANs for the three monoterpene mixtures when applying 4 times the measured XO_2 . The results (Figures S4B–S6B and S4D–S6D) show that β and thus P_{ANs} decrease when increasing XO_2 , which results in an increase in the calculated lifetime from 1.5 ± 1 h to 2.5 ± 1.5 h, which remains consistent with the previous ~~measurements conducted in a boreal forest~~ observations described above.

4.6 PANs major production and loss processes


A dominant fraction of the measured PANs is expected to be in the form of peroxyacetic nitric anhydride ($\text{CH}_3\text{C}(\text{O})\text{O}_2\text{NO}_2$, also known as peroxyacetyl nitrate, PAN), which is formed in the reaction between the peroxyacetyl radical ($\text{CH}_3\text{C}(\text{O})\text{O}_2$) and NO_2 (R13). In the boundary layer, $\text{CH}_3\text{C}(\text{O})\text{O}_2$ is produced directly from the OH-initiated oxidation of acetaldehyde (CH_3CHO , R14) and the photolysis of dicarbonyls such as methylglyoxal ($\text{CH}_3\text{C}(\text{O})\text{CHO}$, R15, Crowley et al. (2018)) and from the oxidation of BVOCs such as isoprene (via methacrolein ($\text{CH}_2\text{C}(\text{CH}_3)\text{CHO}$, MACR) and methyl vinyl ketone ($\text{CH}_2\text{CHC}(\text{O})\text{CH}_3$, MVK)) and α -pinene after multiple reaction steps.



In ~~the summertime~~, forested environments in the summer, where isoprene emissions are high, peroxyacetic nitric anhydride ($\text{CH}_2\text{C}(\text{CH}_3)\text{C}(\text{O})\text{O}_2\text{NO}_2$, MPAN), an OH-initiated oxidation product of MACR (R16–R17), is expected to make a significant contribution to the total peroxy nitrates. Note that $\text{CH}_2\text{C}(\text{CH}_3)\text{C}(\text{O})\text{O}_2$ can also isomerize (Crounse et al., 2012), which is not included in the MCM. With an annual average propane (C_3H_8) measurement around 507 pptv at La Tardiere, France, in 2018 (Ge et al., 2024), peroxypropionic nitric anhydride ($\text{CH}_3\text{CH}_2\text{C}(\text{O})\text{O}_2\text{NO}_2$, PPN), an OH-initiated oxidation product of propanal ($\text{CH}_3\text{CH}_2\text{CHO}$, R18–R19), is expected to be present as well.



For both PAN, MPAN, PPN and any other PANs, the production rate strongly depends on the ~~concentration~~~~presence~~ of NO, hydroperoxyl radicals (HO₂) and other peroxy radicals (RO₂) that can lead to competing reactions (R20-R21) to e.g. (R13), (R17) and (R19). XO₂ ~~represents~~~~describes~~ the sum of HO₂ + RO₂.



PANs are permanently removed through deposition (R22) and can be lost through thermal decomposition (R23) reforming RC(O)O₂, which can subsequently react as described above (R20-R21). Larger and/or unsaturated PANs, such as MPAN, can also be lost through oxidation (R24).



The thermal decomposition of PANs is strongly temperature dependent resulting in lifetimes with respect to (R23) ~~from~~~~of~~ 7.5 hours at 283 K and 40 minutes at 298 K (IUPAC^{upae}, 2024). The effective lifetime increases from that calculated from the thermal decomposition rate coefficient when regeneration of PANs through e.g. reaction (R13), (R17), and (R19) occurs. Thermal decomposition is thus expected to be the dominant loss process of PANs at high temperatures in the presence of NO and/or XO₂, however, at nighttime, when the temperature is lower and the mixing ratios of NO and XO₂ also are lower, deposition can play an important role, depending on boundary layer height, humidity and surfaces.

4.7 Measured and modelled PANs

Figure 7A plots the ~~measured~~ mixing ratio for ΣPANs (~~=PAN + MPAN + PPN + other PANs~~) for which maximum daytime mixing ratios are between 100 pptv and 1600 pptv. The large variability is presumably caused by the observed variability in temperature, concentrations of oxidants, and BVOCs. The temperature fluctuations measured during the campaign results in a thermal lifetime of the PANs that spans two orders of magnitude, from 15 hours at 279 K ~~and to~~ 3 minutes at 314 K (~~the extremes measured during ACROSS~~) as shown in Figure 7B.

Due to the many different production pathways of PANs, we used a detailed chemical box-model (see section 3.3 for details) to assess the contributions of various precursors and compare to measured ΣPANs. As the calculated thermal loss rate varies significantly from day to day, two single days (marked in grey in Figure 7), where measurements of OH, O₃, NO, NO₂, and BVOCs are available, have been modelled instead of using average diel profiles ~~aslike~~ in the ANs analysis. One day (July 4th 2022) is in phase 1 from the ANs analysis, where the

temperature reaches around 27°C (300 K) resulting in a thermal lifetime of 25 minutes (without considering recombination) and during which the mixing ratios of precursors (oxidants and BVOCs) were low. The second day (July 13th 2022) is in phase 2 from the ANs analysis, where the temperature has a maximum of around 39°C (312 K) resulting in a thermal lifetime of around 5 minutes (without considering recombination) and higher precursor levels than during the first day.

The box model was constrained by measurements (240-minute running averages) of temperature, humidity, JNO₂, J(O¹D), OH, O₃, NO, NO₂, isoprene, and acetaldehyde (CH₃CHO), and the prescribed 20% limonene monoterpene mixture described above ~~estimated monoterpene mixture of α -pinene, β -pinene, and limonene as described above for the case with 20% limonene~~. Note that varying the fraction of limonene did not change the modelled Σ PANs significantly. The model was additionally initiated using average CO and pressure for the modelled day, ~~and~~ 507 pptv propane (C₃H₈), 193 pptv n-butane (n-C₄H₁₀), and 107 pptv isobutane (i-C₄H₁₀) (Ge et al., 2024). To ensure that all the trace gases not constrained in the model were in steady-state, a spin-up time of nine days was used. The thermal losses of PANs were included in the model as described above and an additional first-order loss term (physical losses from e.g. deposition) was varied in two separate model runs to get satisfactory model-measurement agreement for Σ PANs during daytime (06:00-18:00 UTC) and nighttime (18:00-06:00 UTC) for the individual days. The PANs identified in the MCM are listed in table S4 and the fractional contributions of PAN, MPAN, and PPN to the total are described in the supplementary information.

Modelled loss rates optimized to match nighttime observations of PANs: The modelled PANs, separated into PAN and other PANs when optimized for nighttime are plotted for the low and high precursor day in Figure 8A and B together with the measured mixing ratios. Reasonable agreement between the measured and modelled Σ PANs can be observed for both the high and low precursor day at nighttime, but overestimates part of the day. The daytime was therefore modelled separately (see below).

The physical loss frequencies required to align the modelled Σ PANs at nighttime with the observations were $1.8 \times 10^{-4} \text{ s}^{-1}$ and $1.1 \times 10^{-3} \text{ s}^{-1}$ for the low and high precursor day, respectively. By comparison the nighttime loss frequency attributable to thermal decomposition when taking recombination into account (at temperatures of 10-20 °C, see Figure 8C and D) is negligible. The resulting average nighttime lifetime (1.5 and 0.24 h, see Table 2) is therefore determined almost solely by physical losses.

Modelled loss rates optimized to match daytime observations of PANs: When optimizing for daytime agreement in the model (see Figure S75A and B) reasonable agreement can be observed between 06:00-18:00 for the high precursor day. For the low precursor day, the model overestimates the measurement between 06:00 and 12:00 UTC and underestimates between 12:00 and 18:00 UTC. The physical loss frequencies required to align the modelled Σ PANs at daytime with the observations were thus derived were at daytime $5.5 \times 10^{-4} \text{ s}^{-1}$ and $2.6 \times 10^{-3} \text{ s}^{-1}$ for the low and high precursor day, respectively, and ~~$1.8 \times 10^{-4} \text{ s}^{-1}$ and $1.1 \times 10^{-3} \text{ s}^{-1}$ at nighttime.~~

The results shown above indicate that daytime physical losses are significantly higher than at nighttime, for which ~~there are several potential explanations, for the enhanced physical loss of PANs at daytime compared to nighttime.~~ The high daytime values increase could be

explained by a reduction in the surface resistance to foliar uptake when plant-stomata are open, (similar to O₃, Shepson et al. (1992)) or a reduction in the transport resistance to uptake due to turbulent mixing. Rapid vertical mixing (venting) out of the canopy would also contribute to the net physical losses during daytime (Bohn, 2006) if significant concentration gradients exist. However, as effects of venting were not observed for alkyl nitrates (i.e. no significant difference between physical loss frequency during day and night was observed), we do not consider venting to be significant.

~~According to the model, at daytime (06:00–18:00 UTC, 08:00–20:00 LT), PAN contributes 61–78 % and 58–72 % of Σ PANs for the low and high precursor day, respectively. The nighttime PAN contribution to the Σ PANs is lower at 48–70 % and 48–60 %, respectively. The lower contribution of PAN to Σ PANs at nighttime is related to the high nighttime mixing ratios of monoterpenes (caused by the vegetation emitting BVOCs into a shallow boundary layer), which degrade to form larger PANs (see Table S1) compared to PAN. During the BEARPEX-2007 campaign in the Sierra Nevada Mountains, PAN was determined at local noon time to contribute 70–90 % of Σ PANs (Wolfe et al., 2011), which is in reasonable agreement with our daytime results. A list of the modelled PANs is given in Table S1.~~

~~The model generated high amounts of HPAN (HOCH₂C(O)O₂NO₂) from glycolaldehyde, despite HPAN having a very short thermal lifetime (Zheng et al., 2011) and never having been detected in ambient air samples. In order to reduce the modelled contribution of HPAN to Σ PANs, an arbitrary loss term (10 times that of the other PANs) was included in the model.~~

~~The modelled PANs, separated into PAN and other PANs when optimized for nighttime are plotted for the low and high precursor day in Figure 8A and B together with the measured mixing ratios. Reasonable agreement between the measured and modelled Σ PANs can be observed for both the high and low precursor day at nighttime and from 12:00–18:00 UTC when optimizing for nighttime agreement. However, the model overestimates the measurement between 06:00 and 12:00 UTC on both days by up to a factor of 4. When optimizing for daytime agreement in the model (see Figure S5A and B) reasonable agreement can be observed between 06:00–18:00 for the high precursor day. For the low precursor day, the model overestimates the measurement between 06:00 and 12:00 UTC and underestimates between 12:00 and 18:00 UTC.~~

~~For both days optimizing for daytime agreement by adjusting the Σ PANs physical loss term using a constant value for the whole day results in a significant underestimation of modelled Σ PANs compared to the measured at nighttime. The lower nighttime physical loss results from temperature inversions and weak vertical mixing.~~

The overall loss of PAN is caused by a combination of physical losses such as deposition, and (at daytime) transport and thermal decomposition, which are plotted together in Figure S79C and D. ~~In this case, the physical losses (modelled as a single loss frequency) were adjusted to optimize measurement-model agreement at nighttime. The thermal lifetime derived in Figure 7B from the thermal decomposition rate coefficient need to be modified to account for reformation via reaction between RC(O)O₂ and NO₂. To calculate an effective thermal loss rate coefficient, a correction factor which takes recombination through for example reaction (R13) into account was applied.~~ The correction factor (f_{NO_2}) is described in equation (10) and represents the fraction of RC(O)O₂ formed from the thermal decomposition that does not lead to reformation of PANs.

$$f_{\text{NO}_2} = 1 - \left(\frac{k_{\text{RC(O)O}_2+\text{NO}_2}[\text{NO}_2]}{k_{\text{RC(O)O}_2+\text{NO}_2}[\text{NO}_2] + k_{20}[\text{NO}] + k_{21}[\text{XO}_2]} \right) \quad (10)$$

where k_i is the rate coefficient for reaction (Ri), and $[\text{NO}_2]$, $[\text{NO}]$, and $[\text{XO}_2]$ are the concentrations of NO_2 , NO , and XO_2 ($\text{HO}_2 + \text{RO}_2$). As seen in equation (10), this correction factor requires information not only of NO and NO_2 , but also XO_2 concentrations. In the box model, NO and NO_2 are constrained to the measurements whereas XO_2 is calculated. The modelled XO_2 is compared to the measured value in Figure S86 for the two chosen days. On the low precursor day, the model and measurements agree well, but on the high precursor day, the model predicts that XO_2 should be 4 times higher than the measurements at midday. As the measured XO_2 is strictly a lower limit, the effective thermal decomposition rate coefficient for PAN (considering recombination of $\text{CH}_3\text{C(O)O}_2$ with NO_2 , see above) ~~was~~ therefore calculated~~shown~~ using both the measured and the modelled XO_2 to evaluate the uncertainty in this parameter. If the measured XO_2 is correct (and the modelled values are too high on the day during phase 2), the modelled derived physical loss frequency would be too low and would have to be incremented by the difference in effective thermal decomposition frequencies (approximately $5 \times 10^{-4} \text{ s}^{-1}$).

Thermal decomposition accounts for approximately 40 % of the total loss when using modelled XO_2 and optimizing for daytime agreement (see Figure S74C and D). The 40% contribution of thermal loss is inconsistent with the observations of Wolfe et al. (2009) who showed through flux measurements that thermal losses of PAN were dominant. Combining physical and thermal loss rates results in daytime lifetimes of $0.42 \pm 0.05 \text{ h}$ and $0.08 \pm 0.01 \text{ h}$ for the low and high precursor day, respectively. These are significantly shorter than those determined at nighttime (see above or Table 2).

~~Table 2 shows an overview of the average modelled thermal, physical, and total loss frequencies of PANs at daytime and nighttime and the resulting effective lifetimes. The nighttime effective lifetime for Σ PANs during the low photochemical activity day was similar to that determined for ANs with $1.5 \pm 0.1 \text{ h}$, but the daytime effective lifetime was significantly shorter with $0.42 \pm 0.05 \text{ h}$. On the day with high photochemical activity, the average effective lifetime was even shorter at 0.08 ± 0.01 and $0.24 \pm 0.02 \text{ h}$ for daytime and nighttime, respectively. The total loss frequency peaked at 0.0038 s^{-1} and 0.00085 s^{-1} for the high and low photochemical activity days resulting in effective lifetimes of around 4.4 min (0.07 h) and 20 min (0.32 h), respectively, when adjusting the physical loss frequency for measurement-model agreement at daytime.~~

5 Conclusion:

Measurements of NO_x , O_3 , BVOCs, the sum of alkyl nitrates (Σ ANs), and the sum of peroxydicarboxylic nitric anhydrides (Σ PANs) have been used to analyse the sources, sinks and lifetime of ANs and PANs in a temperate forest influenced by anthropogenic emissions.

The ANs analysis has been performed for two phases. The first phase is characterised by relatively low temperatures, oxidants, NO and ~~reactant~~ BVOCs compared to the entire campaign, and the second phase was characterised by higher temperatures, oxidants, NO and BVOCs~~reactants~~. This led to significantly different production rates, but very similar lifetimes. The production was dominated by OH-initiated reactions at midday for both phases, but large

differences were estimated at nighttime. NO₃-initiated reactions play a similarly important role as OH at nighttime for the second phase, however, for the first phase OH still dominates at nighttime. NO₃-initiated reactions have also been shown to be important at daytime despite the rapid photolysis. The lifetime for both phases was short at 1-4 hours, which agrees with a previous study in a forest environments.

For the PANs analysis, a box model was used to simulate two individual days; one in the first phase of the ANs analysis and one in the second phase. Two constant physical loss terms are applied for each of the two days optimized to match the average daytime and nighttime mixing ratios. For the low precursor day (July 4th 2022), loss frequencies of $5.5 \times 10^{-4} \text{ s}^{-1}$ and $1.8 \times 10^{-4} \text{ s}^{-1}$ were used to align measurement and model for daytime and nighttime, respectively, while for the high precursor day (July 13th 2022) $2.6 \times 10^{-3} \text{ s}^{-1}$ and $1.1 \times 10^{-3} \text{ s}^{-1}$ were used. This resulted in lifetimes of around 20 min and 4 min at midday for the low and high precursor day, respectively, where thermal decomposition contributed approximately 40 %. Peroxyacetic nitric anhydride (PAN) represents 48-78% of Σ PANs according to the box model, with the highest fractions predicted at daytime.

Lifetimes of organic nitrates in the forested environment are very short, with a potential reason for the short lifetime is dry deposition to e.g., presumably to soil and foliar surfaces playing an important role in removing them from the gas phase. If deposition is the cause of the short lifetimes this would be consistent with a picture of Through emission of reactive organic trace gases, the forest ecosystem thus capturing essential nitrogen-containing nutrients originating from anthropogenic sources and transferring them to the biosphere.

6 Data Availability:

All measurements from the ACROSS campaign including NO_x (Andersen and Crowley, 2023a; Xue et al., 2023), O₃ (Crowley, 2023), organic nitrates (Andersen and Crowley, 2023b), NO₃ reactivity (Dewald and Crowley, 2023), BVOCs (Michoud et al., 2024), meteorological quantities (Denjean, 2023), photolysis frequencies (Dusanter and Jamar, 2023), OH (Kukui, 2023a), and peroxy radicals (Kukui, 2023b) can be found at <https://across.aeris-data.fr/catalogue/> (last access: 31 August 2024).

7 Author contribution:

All authors contributed with measurements. Data analysis was conducted by STA with contributions from JNC. RS did the box modelling. CC and VM organized the field campaign with contributions from the individual group leads. STA and JNC developed the manuscript with contributions from all authors.

8 Competing Interests:

The authors declare that they have no conflict of interest.

9 Acknowledgements:

STA is thankful to the Alexander von Humboldt foundation for funding her stay at MPIC.

PD gratefully acknowledges the Deutsche Forschungsgemeinschaft (project “MONOTONS”, project number: 522970430).

We thank Sebastien Dusanter and Marina Jamar from IMT Nord Europe, France, for supplying photolysis frequency measurements on the tower.

The ACROSS project has received funding from the French National Research Agency (ANR) under the investment program integrated into France 2030, with the reference ANR-17-MPGA-0002, and it was supported by the French National program LEFE (Les Enveloppes Fluides et l'Environnement) of the CNRS/INSU (Centre National de la Recherche Scientifique/Institut National des Sciences de l'Univers). CNRS-INSU provides support to the PEGASUS platform as a national facility. Data from the ACROSS campaign and the PEGASUS facility are hosted by the French national center for Atmospheric data and services AERIS.

10 References:

Andersen, S. T. and Crowley, J. N.: ACROSS_MPIC_RambForest_5ch-NO2_1min_L2 [dataset], <https://doi.org/10.25326/705>, 2023a.

Andersen, S. T. and Crowley, J. N.: ACROSS_MPIC_RambForest_5ch-PNs-ANs_10min_L2 [dataset], <https://doi.org/10.25326/706>, 2023b.

Andersen, S. T., McGillen, M. R., Xue, C., Seubert, T., Dewald, P., Türk, G. N. T. E., Schuladen, J., Denjean, C., Etienne, J. C., Garrouste, O., Jamar, M., Harb, S., Cirtog, M., Michoud, V., Cazaunau, M., Bergé, A., Cantrell, C., Dusanter, S., Picquet-Varrault, B., Kukui, A., Mellouki, A., Carpenter, L. J., Lelieveld, J., and Crowley, J. N.: Measurement report: Sources, sinks, and lifetime of NO_x in a suburban temperate forest at night, *Atmos. Chem. Phys.*, 24, 11603-11618, 10.5194/acp-24-11603-2024, 2024.

Ayres, B. R., Allen, H. M., Draper, D. C., Brown, S. S., Wild, R. J., Jimenez, J. L., Day, D. A., Campuzano-Jost, P., Hu, W., de Gouw, J., Koss, A., Cohen, R. C., Duffey, K. C., Romer, P., Baumann, K., Edgerton, E., Takahama, S., Thornton, J. A., Lee, B. H., Lopez-Hilfiker, F. D., Mohr, C., Wennberg, P. O., Nguyen, T. B., Teng, A., Goldstein, A. H., Olson, K., and Fry, J. L.: Organic nitrate aerosol formation via NO₃ + biogenic volatile organic compounds in the southeastern United States, *Atmos. Chem. Phys.*, 15, 13377-13392, 10.5194/acp-15-13377-2015, 2015.

Barnes, I., Bastian, V., Becker, K. H., and Tong, Z.: KINETICS AND PRODUCTS OF THE REACTIONS OF NO₃ WITH MONOALKENES, DIALKENES, AND MONOTERPENES, *Journal of Physical Chemistry*, 94, 2413-2419, 10.1021/j100369a041, 1990.

Berndt, T. and Boge, O.: Gas-phase reaction of NO₃ radicals with isoprene: A kinetic and mechanistic study, *International Journal of Chemical Kinetics*, 29, 755-765, 10.1002/(sici)1097-4601(1997)29:10<755::aid-kin4>3.0.co;2-l, 1997.

Berndt, T. and Böge, O.: Products and mechanism of the gas-phase reaction of NO₃ radicals with α -pinene, *Journal of the Chemical Society, Faraday Transactions*, 93, 3021-3027, 10.1039/A702364B, 1997.

Bohn, B.: Solar spectral actinic flux and photolysis frequency measurements in a deciduous forest, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006902>, 2006.

Brown, S. S. and Stutz, J.: Nighttime radical observations and chemistry, *Chem. Soc. Rev.*, 41, 6405–6447, 2012.

745 Burkholder, J., Sander, S., Abbatt, J., Barker, J., Cappa, C., Crounse, J., Dibble, T., Huie, R.,
Kolb, C., and Kurylo, M.: Chemical kinetics and photochemical data for use in atmospheric
studies; evaluation number 19, Jet Propulsion Laboratory, National Aeronautics and Space
Administration, 2020.

750 Cantrell, C. and Michoud, V.: An Experiment to Study Atmospheric Oxidation Chemistry and
Physics of Mixed Anthropogenic–Biogenic Air Masses in the Greater Paris Area, *Bulletin
of the American Meteorological Society*, 103, 599-603, <https://doi.org/10.1175/BAMS-D-21-0115.1>, 2022.

755 Claflin, M. S. and Ziemann, P. J.: Identification and Quantitation of Aerosol Products of the
Reaction of β -Pinene with NO_3 Radicals and Implications for Gas- and Particle-Phase
Reaction Mechanisms, *Journal of Physical Chemistry A*, 122, 3640-3652,
10.1021/acs.jpca.8b00692, 2018.

Cox, R. A., Ammann, M., Crowley, J. N., Herrmann, H., Jenkin, M. E., McNeill, V. F.,
Mellouki, A., Troe, J., and Wallington, T. J.: Evaluated kinetic and photochemical data for
atmospheric chemistry: Volume VII – Criegee intermediates, *Atmos. Chem. Phys.*, 20,
13497-13519, 10.5194/acp-20-13497-2020, 2020.

760 Crounse, J. D., Knap, H. C., Ørnsø, K. B., Jørgensen, S., Paulot, F., Kjaergaard, H. G., and
Wennberg, P. O.: Atmospheric Fate of Methacrolein. 1. Peroxy Radical Isomerization
Following Addition of OH and O_2 , *The Journal of Physical Chemistry A*, 116, 5756-5762,
10.1021/jp211560u, 2012.

765 Crowley, J.: ACROSS_MPIC_RambForest_O3_10min_L1 [dataset],
<https://doi.org/10.25326/707>, 2023.

Crowley, J. N., Pouvesle, N., Phillips, G. J., Axinte, R., Fischer, H., Petäjä, T., Nölscher, A.,
Williams, J., Hens, K., Harder, H., Martinez-Harder, M., Novelli, A., Kubistin, D., Bohn,
B., and Lelieveld, J.: Insights into HO_x and RO_x chemistry in the boreal forest via
measurement of peroxyacetic acid, peroxyacetic nitric anhydride (PAN) and hydrogen
770 peroxide, *Atmospheric Chemistry and Physics*, 18, 13457-13479, 10.5194/acp-18-13457-2018, 2018.

Denjean, C.: ACROSS_CNRM_RambForest_MTO-1MIN_L2 [dataset],
<https://doi.org/10.25326/437>, 2023.

775 DeVault, M. P., Ziola, A. C., and Ziemann, P. J.: Products and Mechanisms of Secondary
Organic Aerosol Formation from the NO_3 Radical-Initiated Oxidation of Cyclic and
Acyclic Monoterpenes, *ACS Earth and Space Chemistry*, 6, 2076-2092,
10.1021/acsearthspacechem.2c00130, 2022.

Dewald, P. and Crowley, J. N.: ACROSS_MPIC_RambForest_KNO3_10min_L2 [dataset],
<https://doi.org/10.25326/545>, 2023.

780 Dewald, P., Seubert, T., Andersen, S. T., Türk, G. N. T. E., Schuladen, J., McGillen, M. R.,
Denjean, C., Etienne, J. C., Garrouste, O., Jamar, M., Harb, S., Cirtog, M., Michoud, V.,
Cazaunau, M., Bergé, A., Cantrell, C., Dusanter, S., Picquet-Varrault, B., Kukui, A., Xue,
C., Mellouki, A., Lelieveld, J., and Crowley, J. N.: NO_3 reactivity during a summer period
in a temperate forest below and above the canopy, *Atmos. Chem. Phys.*, 24, 8983-8997,
785 10.5194/acp-24-8983-2024, 2024.

Draxler, R. R. and Rolph, G. D.: HYSPLIT (HYbrid Single-Particle Lagrangian Integrated
Trajectory) Model access via NOAA ARL READY Website
(<http://ready.arl.noaa.gov/HYSPLIT.php>). NOAA Air Resources Laboratory, Silver
Spring, MD., 2011.

790 Dusanter, S. and Jamar, M.: ACROSS_IMTNE_RambForest_Jvalues_AboveCanopy_L2
[dataset], <https://doi.org/10.25326/774>, 2023.

Eisele, F. L. and Tanner, D. J.: Ion-Assisted Tropospheric Oh Measurements, *Journal of
Geophysical Research-Atmospheres*, 96, 9295-9308, 1991.

795 Fry, J. L., Draper, D. C., Barsanti, K. C., Smith, J. N., Ortega, J., Winkle, P. M., Lawler, M. J.,
Brown, S. S., Edwards, P. M., Cohen, R. C., and Lee, L.: Secondary Organic Aerosol
Formation and Organic Nitrate Yield from NO₃ Oxidation of Biogenic Hydrocarbons,
Environmental Science & Technology, 48, 11944-11953, 10.1021/es502204x, 2014.

800 Fry, J. L., Kiendler-Scharr, A., Rollins, A. W., Wooldridge, P. J., Brown, S. S., Fuchs, H.,
Dube, W., Mensah, A., dal Maso, M., Tillmann, R., Dorn, H. P., Brauers, T., and Cohen,
R. C.: Organic nitrate and secondary organic aerosol yield from NO₃ oxidation of beta-
pinene evaluated using a gas-phase kinetics/aerosol partitioning model, Atmospheric
Chemistry and Physics, 9, 1431-1449, 2009.

805 Fry, J. L., Kiendler-Scharr, A., Rollins, A. W., Brauers, T., Brown, S. S., Dorn, H. P., Dube,
W. P., Fuchs, H., Mensah, A., Rohrer, F., Tillmann, R., Wahner, A., Wooldridge, P. J., and
Cohen, R. C.: SOA from limonene: role of NO₃ in its generation and degradation,
Atmospheric Chemistry and Physics, 11, 3879-3894, 10.5194/acp-11-3879-2011, 2011.

810 Ge, Y., Solberg, S., Heal, M. R., Reimann, S., van Caspel, W., Hellack, B., Salameh, T., and
Simpson, D.: Evaluation of modelled versus observed non-methane volatile organic
compounds at European Monitoring and Evaluation Programme sites in Europe, Atmos.
Chem. Phys., 24, 7699-7729, 10.5194/acp-24-7699-2024, 2024.

Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and
Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1
(MEGAN2.1): an extended and updated framework for modeling biogenic emissions,
Geoscientific Model Development, 5, 1471-1492, 10.5194/gmd-5-1471-2012, 2012.

815 Hallquist, M., Wangberg, I., Ljungstrom, E., Barnes, I., and Becker, K. H.: Aerosol and product
yields from NO₃ radical-initiated oxidation of selected monoterpenes, Environmental
Science & Technology, 33, 553-559, 10.1021/es980292s, 1999.

820 Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M.,
Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H.,
Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A.,
Maenhaut, W., McFiggans, G., Mentel, T. F., Monod, A., Prevot, A. S. H., Seinfeld, J. H.,
Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of
secondary organic aerosol: current and emerging issues, Atmospheric Chemistry and
Physics, 9, 5155-5236, 10.5194/acp-9-5155-2009, 2009.

825 IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, (Ammann, M., Cox,
R.A., Crowley, J.N., Herrmann, H., Jenkin, M.E., McNeill, V.F., Mellouki, A., Rossi, M.
J., Troe, J. and Wallington, T. J.). Last access April. 2024: <https://iupac.aeris-data.fr/>, last
Kanakidou, M., Seinfeld, J., Pandis, S., Barnes, I., Dentener, F., Facchini, M., Dingenen, R.
V., Ervens, B., Nenes, A., and Nielsen, C.: Organic aerosol and global climate modelling:
a review, Atmospheric Chemistry and Physics, 5, 1053-1123, 10.5194/acp-5-1053-2005,
830 2005.

Kukui, A.: ACROSS_LPC2E_Rambforest_OH_L2 [dataset], <https://doi.org/10.25326/510>,
2023a.

835 Kukui, A.: ACROSS_LPC2E_Rambforest_RO2_L2 [dataset], <https://doi.org/10.25326/509>,
2023b.

Kukui, A., Ancellet, G., and Le Bras, G.: Chemical ionisation mass spectrometer for
measurements of OH and Peroxy radical concentrations in moderately polluted
atmospheres, Journal of Atmospheric Chemistry, 61, 133-154, 10.1007/s10874-009-9130-
9, 2008.

840 Kukui, A., Chartier, M., Wang, J., Chen, H., Dusanter, S., Sauvage, S., Michoud, V., Locoge,
N., Gros, V., Bourrianne, T., Sellegri, K., and Pichon, J. M.: Role of Criegee intermediates
in the formation of sulfuric acid at a Mediterranean (Cape Corsica) site under influence of

- biogenic emissions, *Atmos. Chem. Phys.*, 21, 13333-13351, 10.5194/acp-21-13333-2021, 2021.
- 845 Kwan, A. J., Chan, A. W. H., Ng, N. L., Kjaergaard, H. G., Seinfeld, J. H., and Wennberg, P. O.: Peroxy radical chemistry and OH radical production during the NO₃-initiated oxidation of isoprene, *Atmospheric Chemistry and Physics*, 12, 7499-7515, 10.5194/acp-12-7499-2012, 2012.
- 850 Liebmann, J., Karu, E., Sobanski, N., Schuladen, J., Ehn, M., Schallhart, S., Quéléver, L., Hellen, H., Hakola, H., Hoffmann, T., Williams, J., Fischer, H., Lelieveld, J., and Crowley, J. N.: Direct measurement of NO₃ radical reactivity in a boreal forest, *Atmospheric Chemistry and Physics* 2018, 3799-3815, 10.5194/acp-18-3799-2018, 2018a.
- 855 Liebmann, J., Sobanski, N., Schuladen, J., Karu, E., Hellén, H., Hakola, H., Zha, Q., Ehn, M., Riva, M., Heikkinen, L., Williams, J., Fischer, H., Lelieveld, J., and Crowley, J. N.: Alkyl nitrates in the boreal forest: formation via the NO₃-, OH- and O₃-induced oxidation of biogenic volatile organic compounds and ambient lifetimes, *Atmos. Chem. Phys.*, 19, 10391-10403, 10.5194/acp-19-10391-2019, 2019.
- 860 Liebmann, J. M., Muller, J. B. A., Kubistin, D., Claude, A., Holla, R., Plass-Dülmer, C., Lelieveld, J., and Crowley, J. N.: Direct measurements of NO₃ reactivity in and above the boundary layer of a mountaintop site: identification of reactive trace gases and comparison with OH reactivity, *Atmospheric Chemistry and Physics*, 18, 12045-12059, 10.5194/acp-18-12045-2018, 2018b.
- 865 Lightfoot, P. D., Cox, R. A., Crowley, J. N., Destriau, M., Hayman, G. D., Jenkin, M. E., Moortgat, G. K., and Zabel, F.: Organic peroxy radicals - kinetics, spectroscopy and tropospheric chemistry, *Atmospheric Environment, Part A: General Topics*, 26, 1805-1961, 1992.
- 870 Mellouki, A., Ammann, M., Cox, R. A., Crowley, J. N., Herrmann, H., Jenkin, M. E., McNeill, V. F., Troe, J., and Wallington, T. J.: Evaluated kinetic and photochemical data for atmospheric chemistry: volume VIII – gas-phase reactions of organic species with four, or more, carbon atoms., *Atmos. Chem. Phys.*, 21, 4797-4808, 10.5194/acp-21-4797-2021, 2021.
- 875 Meusel, H., Kuhn, U., Reiffs, A., Mallik, C., Harder, H., Martinez, M., Schuladen, J., Bohn, B., Parchatka, U., Crowley, J. N., Fischer, H., Tomsche, L., Novelli, A., Hoffmann, T., Janssen, R. H. H., Hartogensis, O., Pikridas, M., Vrekoussis, M., Bourtsoukidis, E., Weber, B., Lelieveld, J., Williams, J., Pöschl, U., Cheng, Y., and Su, H.: Daytime formation of nitrous acid at a coastal remote site in Cyprus indicating a common ground source of atmospheric HONO and NO, *Atmospheric Chemistry and Physics*, 16, 14475-14493, 10.5194/acp-16-14475-2016, 2016.
- 880 Michoud, V., Bouzidi, H., Formenti, P., Bauville, A., Cazaunau, M., Alage, S., Cirtog, M., De Haan, D., and Rafla, M.: ACROSS_2022_RambForest_LISA_PTRMS_VOCs_Belowcanopy_10min_20220617 - 20220723 [dataset], <https://doi.org/10.25326/685>, 2024.
- 885 Ng, N. L., Brown, S. S., Archibald, A. T., Atlas, E., Cohen, R. C., Crowley, J. N., Day, D. A., Donahue, N. M., Fry, J. L., Fuchs, H., Griffin, R. J., Guzman, M. I., Herrmann, H., Hodzic, A., Iinuma, Y., Jimenez, J. L., Kiendler-Scharr, A., Lee, B. H., Luecken, D. J., Mao, J., McLaren, R., Mutzel, A., Osthoff, H. D., Ouyang, B., Picquet-Varrault, B., Platt, U., Pye, H. O. T., Rudich, Y., Schwantes, R. H., Shiraiwa, M., Stutz, J., Thornton, J. A., Tilgner, A., Williams, B. J., and Zaveri, R. A.: Nitrate radicals and biogenic volatile organic compounds: oxidation, mechanisms, and organic aerosol, *Atmospheric Chemistry and Physics*, 17, 2103-2162, 10.5194/acp-17-2103-2017, 2017.
- 890

- Nozière, B., Barnes, I., and Becker, K.-H.: Product study and mechanisms of the reactions of α -pinene and of pinonaldehyde with OH radicals, *Journal of Geophysical Research: Atmospheres*, 104, 23645-23656, <https://doi.org/10.1029/1999JD900778>, 1999.
- 895 Perring, A. E., Pusede, S. E., and Cohen, R. C.: An observational perspective on the atmospheric impacts of alkyl and multifunctional nitrates on ozone and secondary organic aerosol, *Chemical Reviews*, 113, 5848-5870, doi:10.1021/cr300520x, 2013.
- Perring, A. E., Wisthaler, A., Graus, M., Wooldridge, P. J., Lockwood, A. L., Mielke, L. H., Shepson, P. B., Hansel, A., and Cohen, R. C.: A product study of the isoprene+NO₃ reaction, *Atmospheric Chemistry and Physics*, 9, 4945-4956, 2009.
- 900 Rindelaub, J. D., McAvey, K. M., and Shepson, P. B.: The photochemical production of organic nitrates from alpha-pinene and loss via acid-dependent particle phase hydrolysis, *Atmospheric Environment*, 100, 193-201, 10.1016/j.atmosenv.2014.10.010, 2015.
- 905 Rollins, A. W., Kiendler-Scharr, A., Fry, J. L., Brauers, T., Brown, S. S., Dorn, H. P., Dubé, W. P., Fuchs, H., Mensah, A., Mentel, T. F., Rohrer, F., Tillmann, R., Wegener, R., Wooldridge, P. J., and Cohen, R. C.: Isoprene oxidation by nitrate radical: alkyl nitrate and secondary organic aerosol yields, *Atmospheric Chemistry and Physics*, 9, 6685-6703, 10.5194/acp-9-6685-2009, 2009.
- Sander, R., Baumgaertner, A., Cabrera-Perez, D., Frank, F., Gromov, S., Grooss, J. U., Harder, H., Huijnen, V., Jockel, P., Karydis, V. A., Niemeyer, K. E., Pozzer, A., Hella, R. B., 910 Schultz, M. G., Taraborrelli, D., and Tauer, S.: The community atmospheric chemistry box model CAABA/MECCA-4.0, *Geoscientific Model Development*, 12, 1365-1385, 10.5194/gmd-12-1365-2019, 2019.
- Schwantes, R. H., Teng, A. P., Nguyen, T. B., Coggon, M. M., Crounse, J. D., St Clair, J. M., Zhang, X., Schilling, K. A., Seinfeld, J. H., and Wennberg, P. O.: Isoprene NO₃ Oxidation 915 Products from the RO₂ + HO₂ Pathway, *Journal of Physical Chemistry A*, 119, 10158-10171, 10.1021/acs.jpca.5b06355, 2015.
- Shepson, P. B., Bottenheim, J. W., Hastie, D. R., and Venkatram, A.: Determination of the relative ozone and PAN deposition velocities at night, *Geophysical Research Letters*, 19, 1121-1124, 10.1029/92gl01118, 1992.
- 920 Shiraiwa, M., Ueda, K., Pozzer, A., Lammel, G., Kampf, C. J., Fushimi, A., Enami, S., Arangio, A. M., Fröhlich-Nowoisky, J., Fujitani, Y., Furuyama, A., Lakey, P. S. J., Lelieveld, J., Lucas, K., Morino, Y., Pöschl, U., Takahama, S., Takami, A., Tong, H., Weber, B., Yoshino, A., and Sato, K.: Aerosol Health Effects from Molecular to Global Scales, *Environmental Science & Technology*, 51, 13545-13567, 10.1021/acs.est.7b04417, 925 2017.
- Sobanski, N., Schuladen, J., Schuster, G., Lelieveld, J., and Crowley, J. N.: A five-channel cavity ring-down spectrometer for the detection of NO₂, NO₃, N₂O₅, total peroxy nitrates and total alkyl nitrates, *Atmospheric Measurement Techniques*, 9, 5103-5118, 10.5194/amt-9-5103-2016, 2016.
- 930 Spittler, M., Barnes, I., Bejan, I., Brockmann, K. J., Benter, T., and Wirtz, K.: Reactions of NO₃ radicals with limonene and alpha-pinene: Product and SOA formation, *Atmospheric Environment*, 40, S116-S127, 10.1016/j.atmosenv.2005.09.093, 2006.
- Thieser, J., Schuster, G., Phillips, G. J., Reiffs, A., Parchatka, U., Pöhler, D., Lelieveld, J., and Crowley, J. N.: A two-channel, thermal dissociation cavity-ringdown spectrometer for the 935 detection of ambient NO₂, RO₂NO₂ and RONO₂, *Atmos. Meas. Tech.*, 9, 553-576, 10.5194/amt-9-553-2016, 2016.
- Wängberg, I., Barnes, I., and Becker, K. H.: Product and Mechanistic Study of the Reaction of NO₃ Radicals with α -Pinene, *Environmental Science & Technology*, 31, 2130-2135, 1997.

- 940 Wennberg, P. O., Bates, K. H., Crounse, J. D., Dodson, L. G., McVay, R. C., Mertens, L. A.,
 Nguyen, T. B., Praske, E., Schwantes, R. H., Smarte, M. D., St Clair, J. M., Teng, A. P.,
 Zhang, X., and Seinfeld, J. H.: Gas-Phase Reactions of Isoprene and Its Major Oxidation
 Products, *Chemical Reviews*, 118, 3337-3390, 10.1021/acs.chemrev.7b00439, 2018.
- 945 Wolfe, G. M., Thornton, J. A., Yatavelli, R. L. N., McKay, M., Goldstein, A. H., LaFranchi,
 B., Min, K. E., and Cohen, R. C.: Eddy covariance fluxes of acyl peroxy nitrates (PAN,
 PPN and MPAN) above a Ponderosa pine forest, *Atmospheric Chemistry and Physics*, 9,
 615-634, 2009.
- 950 Wolfe, G. M., Thornton, J. A., Bouvier-Brown, N. C., Goldstein, A. H., Park, J. H., McKay,
 M., Matross, D. M., Mao, J., Brune, W. H., LaFranchi, B. W., Browne, E. C., Min, K. E.,
 Wooldridge, P. J., Cohen, R. C., Crounse, J. D., Faloona, I. C., Gilman, J. B., Kuster, W.
 C., de Gouw, J. A., Huisman, A., and Keutsch, F. N.: The Chemistry of Atmosphere-Forest
 Exchange (CAFE) Model – Part 2: Application to BEARPEX-2007 observations, *Atmos.*
Chem. Phys., 11, 1269-1294, 10.5194/acp-11-1269-2011, 2011.
- 955 Wolfe, G. M., Hanisco, T. F., Arkinson, H. L., Bui, T. P., Crounse, J. D., Dean-Day, J.,
 Goldstein, A., Guenther, A., Hall, S. R., Huey, G., Jacob, D. J., Karl, T., Kim, P. S., Liu,
 X., Marvin, M. R., Mikoviny, T., Misztal, P. K., Nguyen, T. B., Peischl, J., Pollack, I.,
 Ryerson, T., St. Clair, J. M., Teng, A., Travis, K. R., Ullmann, K., Wennberg, P. O., and
 Wisthaler, A.: Quantifying sources and sinks of reactive gases in the lower atmosphere
 using airborne flux observations, *Geophysical Research Letters*, 42, 8231-8240,
 960 <https://doi.org/10.1002/2015GL065839>, 2015.
- Xue, C., McGillen, M., and Mellouki, A.: ACROSS_ICARE_RambForest_NO_L2 [dataset],
<https://doi.org/10.25326/512>, 2023.
- 965 Zheng, W., Flocke, F. M., Tyndall, G. S., Swanson, A., Orlando, J. J., Roberts, J. M., Huey, L.
 G., and Tanner, D. J.: Characterization of a thermal decomposition chemical ionization
 mass spectrometer for the measurement of peroxy acyl nitrates (PANs) in the atmosphere,
Atmospheric Chemistry and Physics, 11, 6529-6547, 2011.

11 Figures:





Figure 1: Schematic diagram showing the formation of PANs ($\text{RC(O)O}_2\text{NO}_2$) and ANs (RONO_2) from the oxidation of VOCs by OH, O_3 , and NO_3 . Reactions of RO_2 not relevant to the formation of $\text{RC(O)O}_2\text{NO}_2$ and RONO_2 have been left out. Note that the scheme does not attempt to capture

974 all formation routes of the primary oxidants, especially those of the OH-radical, which may
975 additionally be formed in e.g. reactions of HO₂ with ~~NOOH~~ and photolysis of HONO.






Figure 2: Time series of 10-minute averages of Σ PANs, Σ ANs, acetaldehyde, isoprene, and sum of monoterpenes (Σ MT), 12-minute averages of OH radicals, and 1-minute averages of NO₂, O₃, and temperature at 5 and 41 m during the ACROSS campaign.




Figure 3: NO_3 -reactivity based estimation of monoterpene mixtures during ACROSS. Panel A shows the difference in temperature between 41 m and 5 m ($\Delta T = T_{41\text{m}} - T_{5\text{m}}$) for 48 hours during

985 ACROSS. Panel B shows the measured sum of monoterpenes (Σ MT) and that calculated using eq.
986 (1) with ~~four~~three different monoterpene mixtures. Panel C plots calculated against measured
987 Σ MT for the scenario with 20% limonene coloured by ΔT . Panel D shows the scenario with 20%
988 limonene from panel C using ~~the~~a mixture of 57% β -pinene and 43% α -pinene when $\Delta T > 1^\circ\text{C}$ and
989 panel E shows the scatter plot after performing the correction for temperature inversions.






Figure 4: Diel profiles of O₃, NO₂, NO, temperature, OH, and isoprene for the two phases used to analyse organic nitrates during the ACROSS campaign; phase 1 from June 28th 2022 to July 7th 2022 and phase 2 from July 8th 2022 to July 20th 2022. The shaded area is $\pm 1\sigma$.

997



998

999 Figure 5: Average diel profiles of the PTRMS measurements of the sum of monoterpenes (black)
1000 and the calculated, average mixture of limonene (dark blue), α -pinene (light blue), and β -pinene
1001 (orange) for the scenario with 20% limonene.






Figure 6: Panel A and B show the mean diel profiles of the Σ ANs production rates from NO₃-, OH-, and O₃-initiated oxidation of a monoterpene mixture consisting of 20% limonene, 15% β -pinene, and 65% α -pinene for phase 1 and 2, respectively. The fractional contribution to the Σ ANs for each oxidant is plotted in panel C and D. Panel E and F show the average diel profile across the two phases for the measured alkyl nitrates (black), expected alkyl nitrates with (blue) and without (orange) any losses, where the different blue lines were calculated using different effective lifetimes of the alkyl nitrates.




Figure 7: Time series of ΣPANs (A) and the thermal lifetime of PAN (B) during ACROSS, where k_{dec} is the temperature dependent rate coefficient for the thermal decomposition of PAN using the expression preferred by the IUPAC panel ([IUPACupac, 2024](#)). Days marked in grey are used for analysis in Figure 8.




Figure 8: The measured and modelled when optimizing for nighttime agreement Σ PANs is plotted for two individual days; one with low precursors (A) and one with high precursors (B). The optimised physical loss for each day is shown in panel C and D together with the thermal decomposition when taking recombination into account using both the measured and modelled mixing ratio of XO_2 .

12 Tables:

Table 1: Rate coefficients and relevant yields for the calculation of Σ PANs

VOC	$k(\text{NO}_3)$ at 298 K ($\text{cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$)	α^{NO_3}	$k(\text{OH})$ at 298 K ($\text{cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$)	$\alpha^{\text{RO}_2+\text{NO}}$	$k(\text{O}_3)$ at 298 K ($\text{cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$)	α^{O_3} ⁱ
α -pinene	6.2×10^{-12} ($\pm 26\%$) ^a	0.18 ^b	5.3×10^{-11} ($\pm 20\%$) ^a	0.22 ^f	9.6×10^{-17} ($\pm 41\%$) ^a	0.80 ^a
β -pinene	2.5×10^{-12} ($\pm 32\%$) ^a	0.49 ^c	7.6×10^{-11} ($\pm 12\%$) ^a	0.24 ^g	1.9×10^{-17} ($\pm 78\%$) ^a	0.30 ^a
<i>d</i> -limonene	1.2×10^{-11} ($\pm 32\%$) ^a	0.50 ^d	1.7×10^{-10} ($\pm 12\%$) ^a	0.23 ^g	2.2×10^{-16} ($\pm 26\%$) ^a	0.66 ^a
Isoprene	6.5×10^{-13} ($\pm 41\%$) ^a	0.77 ^e	1.0×10^{-10} ($\pm 15\%$) ^a	0.13 ^h	1.28×10^{-17} ($\pm 20\%$) ^a	0.26 ^a

α^{NO_3} : Yield of ANs from NO_3 +BVOC in air.

$\alpha^{\text{RO}_2+\text{NO}}$: Yield of ANs from RO_2 +NO for the specific BVOC when the RO_2 is formed from BVOC+OH.

α^{O_3} : Yield of RO_2 from the ozonolysis of BVOC in air.

^aRate coefficients and yields recommended by IUPAC (IUPAC ^{аpaе}, 2024; Mellouki et al., 2021; Cox et al., 2020).

^bAverage of Wängberg et al. (1997), Berndt and Böge (1997), Hallquist et al. (1999), Spittler et al. (2006), Fry et al. (2014), and Devault et al. (2022).

^cAverage of Hallquist et al. (1999), Fry et al. (2009), Fry et al. (2014), Clafflin and Ziemann (2018), and Devault et al. (2022).

^dAverage of Hallquist et al. (1999), Spittler et al. (2006), Fry et al. (2011), Fry et al. (2014), and Devault et al. (2022).

^eAverage of Barnes et al. (1990), Berndt and Boge (1997), Perring et al. (2009), Kwan et al. (2012), Rollins et al. (2009), and Schwantes et al. (2015).

^fAverage of the yields given by Nozière et al. (1999) and Rindelaub et al. (2015).

^gPerring et al. (2013).

^hRecommended by Wennberg et al. (2018) based on multiple studies.

ⁱSet equal to the OH yield of the ozonolysis since a RO_2 is formed with each OH.

Table 2: Overview of the thermal, physical and total loss frequencies (and lifetimes) of ANs and PANs needed to explain the measured ANs and PANs during ACROSS.

		Phase 1 (Low photochemical activity)		Phase 2 (High photochemical activity)	
		Day	Night	Day	Night
ANs	Thermal loss frequency (s^{-1}) ^a	-	-	-	-
	Physical loss frequency (10^{-4} s^{-1})	1.7 ± 0.3	1.7 ± 0.4	3.3 ± 0.8	2.5 ± 0.9
	Total loss frequency (10^{-4} s^{-1})	1.7 ± 0.3	1.7 ± 0.4	3.3 ± 0.8	2.5 ± 0.9
	Total lifetime (h)	1.7 ± 0.2	1.7 ± 0.5	0.9 ± 0.2	1.3 ± 0.5
PANs	Thermal loss frequency (10^{-4} s^{-1}) ^a	1.2 ± 0.8	0.1 ± 0.1	8.8 ± 5.9	0.8 ± 1.3
	Physical loss frequency (10^{-4} s^{-1}) ^b	5.5	1.8	26	11
	Total loss frequency (10^{-4} s^{-1})	6.7 ± 0.8	1.9 ± 0.1	34.8 ± 5.9	11.8 ± 1.3
	Total lifetime (h)	0.42 ± 0.05	1.5 ± 0.1	0.08 ± 0.01	0.24 ± 0.02

^aThe thermal loss frequency has been corrected for the fractional recombination of $\text{CH}_3\text{C}(\text{O})\text{O}_2$ with NO_2 .

Day is defined as 06:00-18:00 UTC (08:00-20:00 LT). Night is defined as 18:00-06:00 UTC (20:00-08:00 LT).

All values except the modelled physical loss frequencies for PANs are given as the mean $\pm 1\sigma$. The ANs are based on hourly averages for each phase and the PANs are based on modelled loss frequencies for one day during each phase.

^aThe thermal loss frequency has been corrected for the fractional recombination of $\text{CH}_3\text{C}(\text{O})\text{O}_2$ with NO_2 .

^bIn total four model runs were conducted to examine the two phases and to differentiate between day and night.