

Estimating breakpoints ~~between climate states~~ in the Cenozoic Era using econometric methods

Mikkel Bennedsen^{1,2,3}, Eric Hillebrand^{1,3}, Siem Jan Koopman^{4,5}, Kathrine By Larsen^{1,3}, and
Rachel Lupien⁶

Authors are listed in alphabetical order

¹Department of Economics and Business Economics, Aarhus University, Denmark.

²Aarhus Center for Econometrics, Aarhus University, Denmark.

³Center for Research in Energy: Economics and Markets (CoRE), Aarhus University, Denmark.

⁴Department of Econometrics and Data Science, Vrije Universiteit Amsterdam, The Netherlands.

⁵Tinbergen Institute, Amsterdam, The Netherlands.

⁶[Department of Geoscience, Aarhus University, Denmark.](#)

Corresponding author: Kathrine By Larsen (kblarsen@econ.au.dk)

Abstract

This study presents a statistical time-domain approach for identifying transitions between climate states, referred to as breakpoints, using well-established econometric tools. ~~We analyze a~~ Our approach offers the advantage of constructing time-domain confidence intervals for the breakpoints, and it includes procedures to determine how many breakpoints are present in the time series. We apply these tools to the 67.1 ~~million-year record of the oxygen isotope ratio~~ million-year-long compilation of benthic foraminiferal oxygen isotopes ($\delta^{18}\text{O}$ ~~derived from benthic foraminifera. The~~), which signify global temperature and ice volume throughout the Cenozoic. This foundational dataset is presented in Westerhold et al. (2020), where the authors use recurrence analysis to identify five breakpoints that define six climate states. Fixing the number of breakpoints to five, our procedure results in breakpoint estimates that closely align with those identified by Westerhold et al. (2020). ~~By treating~~ However, by allowing the number of breakpoints ~~as a parameter to be estimated~~ to vary, we provide the statistical justification for more than five breakpoints in the time series. ~~Further, our approach offers the advantage of constructing confidence intervals for the breakpoints, and it allows for testing the number of breakpoints present in the~~, which is used to characterize Cenozoic paleoclimate and as a reference for many paleoclimate studies. This adds to our understanding of Cenozoic climate history, in terms of the timing and rate of transitions between climate states, and provides a tool to assess many other paleoclimate time series.

1 Introduction

~~Westerhold et al. (2020) present a time series dataset spanning from 67.1~~ Understanding the transitions between climate states in Earth's past is crucial for constraining nonlinear and feedback dynamics of our climate system, and anticipating potential climate system responses to anthropogenic warming. The Cenozoic Era,

spanning from 66 million years ago (Ma) to the present time, covering the Cenozoic Era. Using recurrence analysis, the authors identify four major climate states today, is particularly informative in this regard, as it is well-studied and it includes major shifts from hothouse climates with temperatures 10°C warmer than today to the onset of permanent glaciations at both poles (Zachos et al., 2001; Hansen et al., 2013). These transitions, or *breakpoints*, reflect large-scale changes in the climate system, involving shifts in the carbon cycle, ocean circulation, ice volume, and more (Zachos et al., 2008). As emphasized by Tierney et al. (2020), paleoclimate records are essential for assessing climate sensitivity and evaluating climate models under warmer-than-present conditions. Evidence suggests that the sensitivity of the climate system to external forcings may depend on the climate state (Caballero and Huber, 2013), and that projected future climates may increasingly resemble early Cenozoic conditions under continued emissions (Burke et al., 2018; Steffen et al., 2018). These insights underscore the importance of identifying when past climate state transitions occurred, how many there were, and how certain we are about their timing. Addressing these questions is crucial for understanding the dynamics of long-term climate variability, and recent work has increasingly emphasized transition detection as a key task in climate data analysis (e.g., Marwan et al., 2021; Trauth, 2025).

A widely used approach to identify breakpoints in paleoclimate records is recurrence analysis, which identifies when a system returns to similar states over time, helping to detect changes in the underlying dynamics of time series (Marwan et al., 2007; Marwan, 2023). Westerhold et al. (2020) apply this technique to a record of $\delta^{18}\text{O}$ from benthic foraminifera spanning from 67.1 Ma to the present, covering the Cenozoic Era. Based on the recurrence structure of the record, the authors identify four major climate states — ~~Hothouse, Warmhouse, Coolhouse, and Icehouse~~ Hothouse, Warmhouse, Coolhouse, and Icehouse — which are further divided into six states. We refer to these as *Warmhouse I* (66–56 Ma), *Hothouse* (56–47 Ma), *Warmhouse II* (47–34 Ma), *Coolhouse I* (34–13.9 Ma), *Coolhouse II* (13.9–3.3 Ma), and *Icehouse* (3.3 Ma — present). The climate states in the Cenozoic Era range from very warm climates to the glaciation of Earth’s polar regions (Zachos et al., 2001). The climatic transitions contain important information about variations in Earth’s climate system; see Tierney et al. (2020) for a review. Our study presents a statistical approach for identifying the transitions between climate states, referred to as *breakpoints*, using econometric time-domain tools proposed by Bai and Perron (1998, 2003). This approach offers the advantages of constructing confidence intervals for the dates of the breakpoints, providing a measure of estimation uncertainty, as well as testing for the number of breakpoints in the time series which are further divided into six states through time.

The methodology used by Westerhold et al. (2020) to identify breakpoints is recurrence analysis, as described by Marwan et al. (2007). To conduct this analysis, Westerhold et al. (2020) resampled their data at a frequency they resampled the data at an interval of 5 thousand years (kyr) and used both un-detrended and detrended versions of the data. This analysis leads to the identification of the six climate states. Recurrence analysis provides valuable insights into the recurrence structure and shifts in a time series, and recurrence quantification analysis offers complementary summary measures, such as recurrence rate and determinism. However, the identification of transitions remains largely based on visual interpretation of recurrence plots, and the method lacks formal procedures for selecting the number and statistical certainty

of the transitions.

The use of recurrence analysis to identify climate states is common in the study of paleoclimate time series, see the review in Marwan et al. (2021). Many extensions of this methodology have been suggested to address different issues. Several methodological extensions have sought to address these limitations. For instance, Goswami et al. (2018) propose a breakpoint detection method using a probability density function sequence representation of the time series, which allows for uncertainties in the time stamping of the time series accounts for timestamping uncertainty. Bagniewski et al. (2021) combine recurrence analysis with Kolmogorov–Smirnov tests to detect abrupt transitions in a time series. Rousseau et al. (2023) applies this method on statistically assess abrupt shifts in recurrence distributions. Rousseau et al. (2023) apply this method to the Westerhold et al. (2020) data and find similar climate states as the ones reported in Westerhold et al. (2020), identifying a similar set of transitions along with several additional ones. As discussed by Marwan et al. (2021), there are several other approaches to identify transitions in paleoclimate time series. Among these, Livina et al. (2010) developed a new statistical method of potential analysis and applied it to detect the number of states in a geophysical time series—an ice core record. In a Bayesian framework, Schütz and Holschneider (2011) develop a method for detecting changes in trend, and Ruggieri (2013) introduce a Bayesian algorithm for identifying multiple breakpoints. Reviews of breakpoint detection techniques in more general climate time series are provided by Reeves et al. (2007) and Lund and Shi (2023).

Our approach contributes to the existing breakpoint detection methods in paleoclimate research by applying well-established econometric tools in the time domain, developed in Bai and Perron (1998, 2003). Recently, Trauth et al. (2024) explored a suite of methods, including recurrence analysis, changepoint detection, and nonlinear curve fitting (e.g., sigmoid and ramp functions), to identify climate states in the paleo record. It enables the estimation of multiple breakpoints along with confidence intervals and provides procedures to estimate the number of breakpoints.

The estimation methodology of Bai and Perron (1998, 2003) necessitates a constant observation frequency and a predetermined model specification. To obtain a constant observation frequency, we use mean binning, which entails dividing the data into intervals of fixed length and calculating the mean in each bin. We explore three different model specifications and implement these using the R package by Nguyen et al. (2023). The first model is a state-dependent mean model, which posits an abrupt break in the mean of $\delta^{18}\text{O}$ for each climate state. The second model generalizes this by including a state-independent autoregressive term, which makes the transitions between states more gradual. The final model extends the second model by letting the autoregressive term be state-dependent as well, allowing for state-specific transition dynamics. All models incorporate an error term with state-dependent variance. Given that the time series appears state-wise non-stationary, meaning that the mean and/or the variance of the time series vary over time within a state, we conduct a simulation study to demonstrate the applicability of the approach by Bai and Perron (1998, 2003) in this non-stationary setting transitions in a paleoclimate record. They apply a changepoint detection algorithm by Killick et al. (2012), which efficiently detects multiple changepoints by minimizing a cost function that balances goodness-of-fit with a penalty for additional changes. This approach enables the detection of shifts in the mean, variance, and trend. The

sigmoid functions are characterized by their S-shaped curves and allow for modeling gradual transitions (Crowley and Hyde, 2008; Trauth et al., 2021). In contrast, the ramp functions consist of two horizontal segments connected by a linear trend and represent gradual transitions bounded by abrupt changes in slope, which can be fitted using regression techniques. This method was proposed by Mudelsee (2000) and has been applied to various paleoclimate records (e.g., Fleitmann et al., 2003; Mudelsee and Raymo, 2005). Furthermore, Mudelsee et al. (2014) apply this ramp-function method, among others, to detect major climate transitions in the Cenozoic. For further details, we refer to the textbook treatments in Mudelsee (2014) and Trauth (2025).

~~The number of breakpoints is a parameter in the statistical approach of Bai and Perron (1998, 2003).~~
~~—Fixing—~~We employ a statistical approach based on least-squares that allows the researcher to assess the uncertainty in dating breakpoints through confidence intervals. Information criteria give guidance on the number of breakpoints~~to five, the resulting breakpoint estimates align closely with those identified by Westerhold et al. (2020) across various binning frequencies and model specifications. This demonstrates the robustness of the approach and corroborates the dating of the climate states of Westerhold et al. (2020) with statistical analysis in the time domain. However, when we allow the number of breakpoints to vary and treat this as a parameter to be estimated using information criteria, we find strong statistical evidence for the presence of more than five breakpoints. Specifically, our analysis suggests that Warmhouse II and Coolhouse II can each be split into two separate substates.~~

~~The remainder of the paper is structured as follows: In Section 2.1, we present the . The approach is an econometric time-domain framework (Bai and Perron, 1998, 2003), which was originally applied to detect shifts in real interest rates data in economics (Garcia and Perron, 1996). We henceforth refer to this as the Bai-Perron framework. This framework offers the advantages of constructing confidence intervals for the timestamps of the breakpoints, providing a measure of estimation uncertainty, as well as procedures for selecting the number of breakpoints in the time series that provides the best fit. These additional measures are crucial for understanding the certainty, significance, and timing of climate transition periods in the past. The Bai-Perron framework offers flexibility in modeling both abrupt and gradual transitions. We demonstrate its application and benefits by using the benthic $\delta^{18}\text{O}$ -dataset and climate states by Westerhold et al. (2020). In Section 2, the statistical breakpoint detection methodology applied in this paper is outlined. In Section 3, we conduct the analysis and discuss the results. Section 5 concludes. The finite sample performance of the methodology under state-wise non-stationarity is investigated in a simulation study in Appendix C record from Westerhold et al. (2020), though the framework is broadly applicable to a wide range of paleoclimate time series.~~

2 DataMethodology

2.1 Data

~~The paleoclimate variable $\delta^{18}\text{O}$ measures the ratio of ^{18}O to ^{16}O in the shells of benthic foraminifera obtained from ocean sediment cores, relative to a standard sample. The weight difference between the oxygen isotopes leads to an inverse relationship between $\delta^{18}\text{O}$ and ocean temperatures; see for instance~~

~~Epstein et al. (1951) and Shackleton (1967).~~

~~In this paper, we~~ We use the dataset provided by Westerhold et al. (2020), which compiles measurements of oxygen ~~and carbon~~-isotope ratios from benthic foraminifera across 34 different studies and 14 ocean drilling locations into a single ~~data file~~stack covering the Cenozoic. Our study focuses on the benthic $\delta^{18}\text{O}$ record, specifically the correlation-corrected ~~observations of values of benthic~~ $\delta^{18}\text{O}$ (column “benthic d18O VPDB Corr” from the data file). Westerhold et al. (2020) provide an estimated chronology of the data, which has accuracy ranging from ± 100 kyr in the older part of the sample period to ± 10 kyr in the younger part. We ignore the uncertainty of.¹

Benthic $\delta^{18}\text{O}$ measures the deviations in the ratio of the stable oxygen isotopes ^{18}O to ^{16}O in the shells of benthic foraminifera relative to the Vienna Pee Dee Belemnite (VPDB) standard. The weight difference between the stable oxygen isotopes is a function of deep ocean temperatures (Epstein et al., 1951; Shackleton, 1967; Lisiecki and Raymo, 2005) and of the ~~time stamps in this study~~. The data cover the period 67.10113 Ma to 0.000564 Ma, and we order the observations from oldest to most recent. We remove the 74 missing values in the record, leaving us with 24,259 data points. The top panel of Fig. 1 shows the $\delta^{18}\text{O}$ data with the breakpoints between the climate states as identified by Westerhold et al. (2020). Summary statistics of the dataset for the full sample length and for each climate state are presented in Appendix B.1. of the seawater, which in turn is a function of ice volume and salinity (e.g., Waelbroeck et al., 2002; Oerlemans, 2004). Thus, the benthic stack is an important reference record for global climate history across the Cenozoic. Hereafter, we refer to benthic $\delta^{18}\text{O}$ simply as $\delta^{18}\text{O}$.

~~Top panel: $\delta^{18}\text{O}$ data from Westerhold et al. (2020). The order of the vertical axis is reversed, following standard practice. Bottom panel: The time between data points of $\delta^{18}\text{O}$ data measured in kyr. The vertical dashed lines show transitions between the climate states by Westerhold et al. (2020). The horizontal axis represents time, measured in millions of years before present.~~

The $\delta^{18}\text{O}$ data present unique challenges due to its irregular nature and intermittent gaps. The bottom panel of Fig.1 shows the time series of increments between consecutive time stamps in the record. This graph reveals that the time series is relatively sparse in the older part of the record and relatively dense in the younger part. The average time between two adjacent data points is compilation by Westerhold et al. (2020) spans 67.10113 Ma to 564 years before present (Fig. 1). It contains 24,333 entries, of which 74 are missing in the published version. After excluding these, we retain 24,259 data points, ordered from oldest to most recent. The $\delta^{18}\text{O}$ record is irregularly spaced in time, as is typical for paleoclimate proxy data, and sampling density increases through time, with an average resolution of approximately 2.8 kyr, ~~and the longest gap between data points is approximately~~. The longest gap spans about 115.4 kyr. ~~There are and 533 occurrences of gaps between two data points lasting longer than gaps exceed 10 kyr. Moreover, there are~~ Additionally, 591 instances of multiple observations at the same time stamp time stamps contain multiple $\delta^{18}\text{O}$ values, with up to four simultaneous observations.

Methodology

¹These are the values in column “benthic d18O VPDB Corr”, found in Sheet 33 of the file aba6853_tables_s8_s34.xlsx provided in the Supplementary Materials of Westerhold et al. (2020).

In this section, we present and discuss the methodology developed by Bai and Perron (1998, 2003) for estimating breakpoints in linear regression models. Section 2.2 provides an outline of the framework for detecting multiple structural changes in linear regression coefficients. Section 2.3 discusses the model specifications employed—observations recorded at the same time. Westerhold et al. (2020) provide an age model, which has an accuracy ranging from ± 100 kyr in the early Cenozoic to ± 10 kyr in the latest Cenozoic. We do not account for age model uncertainty in this study for breakpoint estimation. Section 2.4 outlines the implementation of the model specifications, but we return to this issue in the results section. Using recurrence analysis, Westerhold et al. (2020) identify six climate states, and we refer to these as *Warmhouse I* (66–56 Ma), *Hothouse* (56–47 Ma), *Warmhouse II* (47–34 Ma), *Coolhouse I* (34–13.9 Ma), *Coolhouse II* (13.9–3.3 Ma), and *Icehouse* (3.3 Ma–present). Summary statistics for the full record and for each climate state identified by Westerhold et al. (2020) are reported in Appendix B.1.




Figure 1: $\delta^{18}\text{O}$ data from Westerhold et al. (2020). The order of the vertical axis is reversed, following standard practice. The vertical dashed lines show transitions between the climate states by Westerhold et al. (2020). The horizontal axis represents time, measured in millions of years before present.

2.2 General The Bai-Perron framework

The method-Bai-Perron framework is based on minimizing the sum of squared residuals while treating the breakpoints as unknown parameters to be estimated (Bai and Perron, 1998, 2003). Consider a linear regression framework for the dependent variable y_t , for $t = 1, \dots, T$, and with m breakpoints, corresponding to $m + 1$ distinct states in the sample. The general model equation is

$$y_t = x_t' \beta + z_t' \delta_j + u_t, \quad t = T_{j-1} + 1, \dots, T_j, \quad (1)$$

with $j = 1, \dots, m + 1$. The m break dates are denoted by $(T_1, \dots, T_m)$, with the convention that $T_0 = 0$ and $T_{m+1} = T$, and u_t is a disturbance term with mean zero and variance σ_j^2 . The $(p \times 1)$ -vector x_t and the $(q \times 1)$ -vector z_t comprise two sets of covariate vectors, for which β is the state-independent vector of coefficients and δ_j is the state-dependent vector of coefficients. Since only specific coefficients are subject to structural breaks, this model is referred to as a partial structural change model. Moreover, we consider breaks in the variance of u_t at the break dates $T_1, \dots, T_m$, such that $\sigma_i^2 \neq \sigma_j^2$ for $i \neq j$. The parameters β and δ_j are estimated alongside the breakpoints but are not of primary interest here.

We initially treat the number of breakpoints, m , as known and estimate the coefficients and the break-

points using a sample of T observations of $\{y_t, x_t, z_t\}$. The estimation method is based on least squares for both the coefficients and the breakpoints. For each possible set of m breakpoints $(T_1, \dots, T_m)$ denoted as $\{T_i\}_{i=1}^m$, we obtain estimates of β and δ_j by minimizing the sum of squared residuals (SSR), that is,

$$SSR = \sum_{j=1}^{m+1} \sum_{t=T_{j-1}+1}^{T_j} (y_t - x'_t \beta - z'_t \delta_j)^2, \quad (2)$$

where β is common to all states, while δ_j is specific for the state j , which is the period between $T_{j-1} + 1$ and T_j . The resulting estimated coefficients are denoted as $\hat{\beta}(\{T_i\}_{i=1}^m)$ and $\hat{\delta}(\{T_i\}_{i=1}^m) \hat{\delta}_j(\{T_i\}_{i=1}^m)$. These coefficients are then used to determine the SSR associated with each set of breakpoints,

$$SSR_T(\{T_i\}_{i=1}^m) \equiv \sum_{j=1}^{m+1} \sum_{t=T_{j-1}+1}^{T_j} \left(y_t - x'_t \hat{\beta}(\{T_i\}_{i=1}^m) - z'_t \hat{\delta}_j(\{T_i\}_{i=1}^m) \right)^2. \quad (3)$$

The estimated breakpoints are then given by

$$(\hat{T}_1, \dots, \hat{T}_m) = \underset{T_1, \dots, T_m}{\operatorname{argmin}} SSR_T(\{T_i\}_{i=1}^m). \quad (4)$$

The minimization is conducted over all partitions $(T_1, \dots, T_m)$ such that $T_j - T_{j-1} \geq \dim(z_t)$ to ensure that there are enough data points to estimate the parameters δ_j in each partition. This procedure leads to estimated parameters for the m breakpoints, i.e., $\{\hat{T}_i\}_{i=1}^m$, $\hat{\beta} = \hat{\beta}(\{\hat{T}_i\}_{i=1}^m)$, and $\hat{\delta} = \hat{\delta}(\{\hat{T}_i\}_{i=1}^m) \hat{\delta}_j = \hat{\delta}_j(\{\hat{T}_i\}_{i=1}^m)$. Since the possible combinations of the placement of the breakpoints is finite, this optimization can be conducted using a grid search, which can be computationally heavy, especially for many breakpoints. Bai and Perron (2003) introduce an efficient method for determining the global minimizers.

An essential advantage of ~~this~~ the Bai-Perron framework is that it allows for constructing confidence intervals for the breakpoints, something that is not available for the recurrence analysis approach implemented in Westerhold et al. (2020). The construction of confidence intervals is based on the asymptotic distribution of the ~~estimated~~ break dates. The convergence results for the construction of confidence intervals rely on a number of assumptions (see Bai and Perron, 2003), ~~which may possibly be violated for the $\delta^{18}\text{O}$ data. To examine whether the framework is adequate for the type of data studied here, we have conducted a large simulation study, of which the details are reported in Appendix C. We discuss the results from the simulation study and how they relate to the analysis of the $\delta^{18}\text{O}$ data in Section 3.~~

2.3 Model specifications

~~In this section, we introduce the model specifications employed in this paper for estimating breakpoints.~~ Three distinct specifications are considered ~~within the Bai-Perron framework~~, referred to as the ‘‘Mean’’, ‘‘Fixed AR’’, and ‘‘AR’’ models, where AR refers to the autoregressive model of order one with intercept. These are all special cases of the framework outlined in Eq. (1). The simplest among them, the Mean model, is specified as follows,

$$y_t = c_j + u_t, \quad t = T_{j-1} + 1, \dots, T_j, \quad (5)$$

for $j = 1, \dots, m + 1$, where c_j is the state-dependent intercept and u_t is an error term. This model is equivalent to setting $x_t = 0$, $z_t = 1$, and $\delta_j = c_j$ in Eq. (1). A breakpoint in this model specification leads to an abrupt change in the mean of the dependent variable y_t .

The Fixed AR model extends the Mean model by incorporating an autoregressive term. We obtain the model

$$y_t = c_j + \varphi y_{t-1} + u_t, \quad t = T_{j-1} + 1, \dots, T_j, \quad (6)$$

for $j = 1, \dots, m + 1$, where y_{t-1} is the dependent variable lagged by one period, and φ is the autoregressive coefficient that is constant over the whole sample. In this model, the effect of a change in the coefficient c_j is more gradual, since it depends on the autoregressive dynamics. The Fixed AR model is obtained from Eq. (1) by specifying $x_t = y_{t-1}$, $\beta = \varphi$, $z_t = 1$, and $\delta_j = c_j$.

The general AR specification also allows the autoregressive term to be state-dependent, resulting in the AR model,

$$y_t = c_j + \varphi_j y_{t-1} + u_t, \quad t = T_{j-1} + 1, \dots, T_j, \quad (7)$$

for $j = 1, \dots, m + 1$, where the autoregressive coefficient φ in Eq. (6) is now state-dependent and is denoted by φ_j . This model is obtained from Eq. (1) by setting $x_t = 0$, $z_t = (1, y_{t-1})$, and $\delta_j = (c_j, \varphi_j)$. Here, both the intercept and the autoregressive coefficient are state-dependent. Thus, the three specifications are nested: The AR model is the most general, the Fixed AR model is nested in the AR model by setting $\varphi_1 = \varphi_2 = \dots = \varphi_{m+1} = \varphi$, and the Mean model is nested in the Fixed AR model by setting $\varphi = 0$. Figure 2 illustrates how the models capture breakpoints. The Mean model is designed to detect abrupt breaks in the mean of a time series, while the Fixed AR model is for smoother breaks. The AR model is more flexible, allowing for both relatively gradual (e.g., T_1) and abrupt (e.g., T_2) breakpoints compared to the Fixed AR model.




Figure 2: Simulated time series using the three model specifications, each with breakpoints $T_1 = 25$ and $T_2 = 75$, and total sample size $T = 100$. For the Mean model, we set $c_1 = 1.0$, $c_2 = 1.2$, and $c_3 = 0.8$. In the Fixed AR model, the parameters are $\varphi = 0.7$, $c_1 = 0.30$, $c_2 = 0.36$, and $c_3 = 0.24$, chosen to yield comparable state-wise means. Likewise in the AR model, we set $\varphi_1 = 0.7$, $\varphi_2 = 0.9$, $\varphi_3 = 0.4$, $c_1 = 0.30$, $c_2 = 0.12$, and $c_3 = 0.48$. In all specifications, we set $u_t = 0$ for all t .

2.4 Implementation

The ~~models are implemented using the~~ Bai-Perron framework is implemented using *mbreaks* ~~R-package~~ (Nguyen et al., 2023) ~~based on the methodology of Bai and Perron (1998, 2003)~~, an R package specifically

designed for this purpose (Nguyen et al., 2023). For all model specifications, we set the minimum length of a state, h , to 2.5 million years (Myr), facilitating the estimation of shorter climate states. Also, we let the variance of the error term, denoted as σ_j^2 , be state-dependent.

As outlined by Bai and Perron (2003), no serial correlation is permitted in the ~~errors of the regressions~~ regression residuals. However, the time series of $\delta^{18}\text{O}$ is likely subject to both autocorrelation and heteroscedasticity. ~~The assumption of no serial correlation in the errors may be violated~~, as documented in ice core records (Davidson et al., 2015; Keyes et al., 2023). Autocorrelation occurs when current values correlate with past values, which is common in paleoclimate data due to long-term persistence in climate dynamics (Mudelsee et al., 2014). Since only up to one lag is included in the covariates in the model specifications ~~considered~~ in this paper, ~~since the incorporation of only up to one lag in the covariates is unlikely to remove serial correlation in the errors~~ residual serial correlation is likely to remain. Heteroscedasticity, or time-varying error variance, is already partially addressed in the model specifications through state-dependent variance. However, additional heteroscedasticity may arise within the estimated states due to factors such as orbital forcing and changes in ice sheet extent. Addressing both autocorrelation and heteroscedasticity is essential to ensure unbiased parameter estimates and valid confidence intervals for the estimated breakpoints.

To ~~address account for~~ these issues, we use the autocorrelation and heteroscedasticity consistent (HAC) covariance matrix estimator with prewhitening in ~~our implementation~~ the Bai-Perron framework. The prewhitening procedure, proposed by Andrews and Monahan (1992), entails applying an autoregressive model with one lag to $z_t \hat{u}_t$, where $\hat{u}_t$ denotes the residuals. The HAC covariance matrix estimator by Andrews (1991) is then constructed based on the filtered series using the quadratic spectral kernel with bandwidth selected by an AR of order one approximation. This approach is used for ~~both the Mean, Fixed AR, and AR models~~ all model specifications and is straightforward to implement using the R package (Nguyen et al., 2023).

Analysis and results

~~This section presents the results of the breakpoint analysis of the $\delta^{18}\text{O}$ record. The irregular sampling is addressed by data binning in Section 2.5, where the stationarity properties of binned data are assessed. In Section 3.1, we fix the number of breakpoints to five as in Westerhold et al. (2020). In Section 3.2, we treat the number of breakpoints as a parameter to be estimated.~~

2.5 Constant data frequency

To conduct breakpoint estimation using the ~~methodology of Bai and Perron (1998, 2003)~~ Bai-Perron framework, we need ~~an equidistant~~ a regularly sampled time series. We use a binning approach to construct a dataset with ~~evenly spaced~~ evenly spaced observations, which is common practice in the analysis of paleoclimate data; see for instance Boettner et al. (2021) ~~or Reikard (2021)~~. We divide the dataset into bins of fixed time intervals and compute the mean of the observations within each bin. In the case of gaps in the binned data, we use the values immediately preceding and succeeding the section with missing data to perform linear interpolation. We consider six different bin sizes, namely 5, 10, 25, 50, 75, and 100 kyr -

(Fig. 3). Summary statistics for the full sample length and for each climate state identified by Westerhold et al. (2020) for all binning frequencies are provided in Appendix B.1. ~~Figure 3 illustrates the binning approach, with the top panel showing unaltered data and binned data at 5 kyr and 100 kyr frequencies, and the bottom panels zooming in on two sub-samples.~~

~~Top panel: The original data and the 5 and 100 kyr binned data. Bottom left panel: The period 36-35 Ma. Bottom right panel: The period 3-2 Ma.~~

~~The top panel of Fig. 3 shows that data~~ Data binned at higher frequencies follow the variations in the dataset more closely, whereas data binned at lower frequencies tend to be smoother ~~. The longest gap in the dataset spans approximately 115 kyr and occurs between 36 Ma and 35 Ma. The bottom panels in Fig. 3 zoom in on the periods 36 to 35 Ma (left) and 3 to 2 Ma (right). These plots illustrate that in (Fig. 3).~~ In case of large gaps (left), a high binning frequency results in linear interpolation between observations ~~(Fig. 3 bottom left). This effect does not occur for periods with many observations (right), where low binning frequencies capture only a small part of the variation in the original data (Fig. 3 bottom right). Binning offers a simple approach to handle the uneven frequency of the dataset. However, it leads to data loss at lower binning frequencies and to the introduction of artificial data points resulting from linear interpolation at higher binning frequencies. The selection of binning frequencies can therefore alter the properties of the time series, potentially misrepresenting the dynamics of the original data.~~

The ~~theoretical framework of Bai and Perron (1998, 2003)~~ Bai-Perron framework is developed for estimating and testing for multiple breakpoints in linear regression models where the regressors are non-trending or state-wise stationary ~~. However, the (Bai and Perron, 2003). A time series is considered stationary if its statistical properties, such as mean and variance, do not change over time. The $\delta^{18}\text{O}$ data appears state-wise non-stationary over most of the record, even within climate states found by Westerhold et al. (2020). As pointed out by Kejriwal et al. (2013), if the time series maintains its stationarity properties over the respective states, the methods developed for stationary data are still applicable for these cases. However, if the process alternates between stationary and non-stationary states, the theoretical properties of the methodology are unknown.~~

To investigate whether the time series is non-stationary, we apply the Augmented Dickey-Fuller (ADF) test (Dickey and Fuller, 1979), with the null hypothesis of non-stationarity. For the entire 25 kyr binned data sample, the ADF test does not reject the null hypothesis at the 1% significance level, indicating non-stationarity. However, when examining the binned data for each climate state identified by Westerhold et al. (2020) separately, the ADF test rejects the null hypothesis at the 1% significance level for the Warmhouse II, Coolhouse I, and Icehouse states. These tests indicate the presence of state-wise non-stationarity, and we therefore need to examine whether the ~~methodology of Bai and Perron (1998, 2003)~~ Bai-Perron framework is applicable to data-generating processes that are state-wise non-stationary or alternating between stationary and non-stationary states.




Figure 3: Top panel: The original data and the 5 and 100 kyr binned data. Bottom left panel: The period 36-35 Ma. Bottom right panel: The period 3-2 Ma.

We conduct a simulation study to examine potential challenges in conducting breakpoint estimation on For this purpose, we conduct a large simulation study designed to verify that the Bai-Perron framework works as intended when applied to these types of data-generating processes using the three model specifications. The study is conducted for both independent and identically distributed (~~i.i.d.~~) error terms and serially correlated error terms in Appendices C.1 and C.2, respectively. The results show that the procedure works well with non-stationarity and is robust to processes with one stationary and one non-stationary state for Fixed AR and AR models. However, the Mean model performs poorly ~~with highly persistent when the~~ data-generating ~~processes~~process exhibits high persistence. In the case of serial correlation, the results are less conclusive, but if the states are sufficiently different, the methodology still appears effective. The study also reveals that the coverage rates ~~of the estimated for~~ confidence intervals are generally adequate for the Fixed AR model~~specification in cases of large breaks. In contrast, , while the confidence intervals for of~~ the AR model are too narrow in many ~~of the cases~~. Overall, the Fixed AR model performs best across the data-generating processes considered.

3 Results

3.1 Setting the number of breakpoints to five ~~Fixing the number of breakpoints to five~~

As an initial step, we fix the number of breakpoints to five, which is the number ~~found in~~ used in the

recurrence analysis presented in Westerhold et al. (2020). We estimate the breakpoints and corresponding 95% confidence intervals for each of the binning frequencies, 5, 10, 25, 50, 75, and 100 kyr. ~~In each estimation, we use a minimum state length of 2.5 Myr, allowing us to estimate relatively short climate states. The estimated breakpoints are tabulated in Appendix B.2 and are shown in Fig. 4, with subfigures for the~~, using Mean, Fixed AR, and AR models ~~for each (App. B.2; Fig. 4).~~ The estimated confidence intervals around the breakpoints are often asymmetrical. Bai and Perron (2003) advocate the use of asymmetric confidence intervals, as these provide better coverage rates when the data are non-stationary.

For the Mean model ~~presented in Fig. 4(a), it is evident that~~, the estimated breakpoints generally remain at the same dates throughout as the binned data frequency decreases step-by-step from 5 kyr to 100 kyr ~~(Fig. 4.a)~~. The width of the 95% confidence intervals increases as the frequency decreases, which can be attributed to the resultant decrease in the number of binned observations available for estimation at the lower frequencies. All the breakpoints align with those identified by recurrence analysis in Westerhold et al. (2020). A similar pattern of alignment is observed in the Fixed AR model, albeit with tighter confidence intervals ~~as depicted in Fig. 4(b)~~ (Fig. 4.b).


~~Figure 4(c) presents the findings for the AR model, which~~ The AR model exhibits more sensitivity to the frequency of the binned data ~~(Fig. 4.c)~~. At higher frequencies, the breakpoints tend to appear in the more recent parts of the sample. However, as the frequency decreases further, the breakpoints are estimated to be in the older parts of the sample period.

For the results using 25 kyr, we find that the estimated breakpoints from the three model specifications align closely ~~The estimated breakpoints align almost with each other and nearly perfectly with those identified by Westerhold et al. (2020) and hence strongly corroborate their findings. The parameters of the~~. The three model specifications ~~estimated using the 25 kyr binned data with the number of breakpoints fixed at five, are provided in Appendix B.3~~ yield parameter estimates that differ across states, reflecting differences in mean and autoregressive dynamics (App. B.3).


(a) Mean model (b) Fixed AR model (c) AR model

~~A comparison of estimated breakpoints using binned data with frequencies of 5, 10, 25, 50, 75, and 100 kyr from top to bottom, fixing the number of breakpoints to five for each model specification. The black dots represent estimated breakpoints, while colored shaded rectangles indicate 95% confidence intervals. The results overlay the $\delta^{18}\text{O}$ data from Westerhold et al. (2020) and their climate states.~~


As a robustness check, we re-estimate the model specifications for five breakpoints using the 25 kyr binned data reversed with respect to the time dimension, ~~i.e., letting time run backwards. The results are shown in Appendix A. Iso~~ that the time series is ordered from present to past rather than past to present (App. A.1). We find that the results of the Mean and Fixed AR models are robust to ~~reversing the time frame~~ the ordering of the time axis, with almost unchanged estimated breakpoints. Conversely, the AR model leads to estimated breakpoints in the more recent part of the sample, resulting in breakpoints at 16.9 Ma and 9.7 Ma, which differ from those estimated using the same model and binning frequency with time running forward (Fig. 4).



(a) Mean model



(b) Fixed AR model



(c) AR model

Figure 4: A comparison of estimated breakpoints using binned data with frequencies of 5, 10, 25, 50, 75, and 100 kyr from top to bottom, fixing the number of breakpoints to five for each model specification. The black dots represent estimated breakpoints, while colored shaded rectangles indicate 95% confidence intervals. The results overlay the $\delta^{18}\text{O}$ data from Westerhold et al. (2020) and their transitions.

In summary, ~~the results of the~~ changing the binning frequency mainly affects the width of the confidence intervals, while the estimated breakpoint timing remains largely unchanged for both the Mean and Fixed AR models ~~exhibit robustness across different binning frequencies, whereas~~. In contrast, the AR model ~~appears~~ is more sensitive to ~~variations in binning frequency and the reversal resolution and the direction of the time dimension~~ frame. As detailed in the simulation study in ~~Appendix C~~ (App. C), the Mean model fails to

accurately detect breakpoints in highly persistent data-generating processes. Consequently, in what follows, we focus on the Fixed AR model for the estimation of breakpoints in the $\delta^{18}\text{O}$ time series. ~~Furthermore, we recommend using binning frequencies~~ Among the binning frequencies, we proceed with 10 kyr and 25 kyr ~~as they result in~~, as these yield the most consistent ~~outcomes~~ results across model specifications and strike a good balance between temporal resolution and signal quality.

3.2 ~~Estimating the number of breakpoints~~ Selecting the number of breakpoints

We use information criteria to ~~estimate the~~ guide the choice of the number of breakpoints. These criteria are model selection tools that balance goodness of fit with model complexity, helping to avoid overfitting. We initially consider the following three criteria: the Bayesian Information Criterion (BIC) by Yao (1988), the modified Schwarz Information Criterion (LWZ) by Liu et al. (1997), and the modified BIC (KT) by Kurozumi and Tuvaandorj (2011). For all criteria, the ~~estimated~~ preferred number of breakpoints is determined as the number of breakpoints that minimizes the information criterion in question. A few paleoclimate studies use information criteria for model selection, for example, Valler et al. (2024) show it can be beneficial.

Bai and Perron (2006) note that the BIC and LWZ criteria perform well in the absence of serial correlation, but both ~~of them~~ lead to overestimation of the number of breakpoints in case of serial correlation in the error term. In simulation studies ~~, reported in~~ (Appendices C.1 and C.2), we find that the KT information criterion performs poorly, and hence, we exclude it from the subsequent analysis. We also find that the number of breakpoints ~~estimated~~ determined using the Mean model specification is generally ~~overestimated too large~~ when employing the information criteria. For the Fixed AR and AR models, the BIC and LWZ criteria typically perform well, especially in data-generating processes with a large break. With serial correlation in the error term, the BIC criterion tends to overestimate the number of breakpoints, whereas the LWZ criterion generally performs well in the Fixed AR and AR model specifications.

Bin size	Mean		Fixed AR		AR	
	BIC	LWZ	BIC	LWZ	BIC	LWZ
5	19	17	17	7	15	5
10	17	17	14	7	14	3
25	17	14	12	6	8	3
50	17	14	10	0	7	0
75	17	14	6	0	5	0
100	17	12	6	0	5	0

Table 1: The number of breakpoints estimated using BIC and LWZ criteria for all models and binning frequencies considered. The minimum state length is set to $h = 2.5$ Myr and the maximum number of breakpoints is 26.

We use the BIC and LWZ information criteria for each model specification and binning frequency to determine the number of breakpoints, and set the minimum state length to $h = 2.5$ Myr. ~~Table B.4 shows the estimated number of breakpoints. There is a tendency towards higher numbers with increasing binning frequency, and for the BIC to indicate higher numbers than the LWZ criterion.~~ (App. B.4). For our preferred specification, the Fixed AR model with 25 kyr binning frequency, the LWZ and BIC criteria suggests six and

~~twelve~~ 12 breakpoints, respectively. For a 10 kyr binning frequency, the estimated number of breakpoints are seven and ~~fourteen~~ 14, respectively. Thus, the information criteria indicate that the number of distinct climate states in the $\delta^{18}\text{O}$ record is larger than the five suggested in Westerhold et al. (2020).




Figure 5: A comparison of estimated breakpoints using the Fixed AR model for one to 15 breakpoints on 25 kyr binned data. The minimum state length is set to $h = 1$ Myr. The black dots represent estimated breakpoints, while colored shaded rectangles indicate 95% confidence intervals. The results overlay the $\delta^{18}\text{O}$ data from Westerhold et al. (2020) and their transitions.

To further investigate the potential for a higher number of breakpoints, we consider the estimation of up to ~~fifteen~~ breakpoints. ~~As previously discussed, we advocate for using the Fixed AR model and mid-range binning frequency. Therefore, we estimate one to fifteen breakpoints using 15~~ with the minimum length of a state set of $h = 1$ Myr. This analysis is conducted with the Fixed AR model and 25 kyr binned data. ~~The results are shown in Fig. 5 (Fig. 5).~~ The same analysis conducted using the 10 kyr binned data led to nearly identical breakpoint estimates ~~and is therefore omitted for conciseness~~ (App. A.2). Comparable findings are presented in Appendices A.3 and A.4, which detail the results of estimating one to 15 breakpoints using the Mean and AR models, respectively, with 25 kyr binned data.

The final estimated breakpoint is placed at 1.425 Ma for the Fixed AR model, just below the upper boundary of the detection window at 1 Ma, imposed by the minimum state length of 1 Myr. Additionally, the estimated breakpoint is located near the midpoint of a linear trend in the time series from approximately 3.3 Ma to the present, suggesting it may be driven by the trend rather than representing a break in the time series (cf. Fig. 5). To investigate this further, we re-estimate the breakpoints for the Fixed AR model, focusing solely on the Icehouse period, with the minimum length of a state set to 250 kyr and 5 kyr binning, leveraging the denser sampling in this part of the record. For the Fixed AR model, the LWZ criterion suggests one breakpoint, while the BIC indicates two. With one breakpoint, the estimate is 1.355 Ma, and with two, the estimated breakpoints are 2.54 Ma and 0.95 Ma (Fig. 6). Estimating more than two breakpoints leads to overlap between the confidence intervals, reducing the interpretability, and these models are therefore excluded. The results are comparable for the Mean and AR models (Appendices A.5 and A.6).




Figure 6: A comparison of estimated breakpoints using the Fixed AR model for one and two breakpoints on 5 kyr binned data for the Icehouse period. The minimum state length is set to $h = 250$ kyr. The black dots represent estimated breakpoints, while colored shaded rectangles indicate 95% confidence intervals. The results overlay the $\delta^{18}\text{O}$ data from Westerhold et al. (2020).

3.3 Limitations of the Bai–Perron framework

Although the Bai–Perron framework provides a flexible and well-established method for detecting breaks, it does have some limitations. The approach assumes piecewise linearity and white noise residuals (Bai and Perron, 2003). However, in the estimations conducted in this study, the residuals are not white noise, indicating that some dynamics are left unexplained. The simulation results show that the Bai–Perron framework nevertheless performs well even when residuals exhibit complex dynamics (App. C). Confidence intervals should still be interpreted with caution. The method is also computationally demanding for high-resolution data, although it remains possible to run on personal computers.

Another limitation is that the method does not account for age model uncertainty, which is important for interpreting the timing and significance of time series analytical output (Marwan et al., 2021). In the Westerhold et al. (2020) data, dating uncertainty ranges from about ± 10 kyr in the younger parts to ± 100 kyr in the older parts. This can affect the timing of transitions and lead to differences when comparing across records (Franke and Donner, 2019). Previous work has shown that age-depth models often underestimate the true uncertainty in the chronology, which would amplify these effects (Telford et al., 2004). While some progress has been made in including age uncertainty into recurrence analyses (Goswami et al., 2018), incorporating it into the Bai–Perron framework remains a challenge. One could however consider the use of age ensembles which are multiple plausible realizations of the time axis to assess robustness of the estimated breakpoints. Fully integrating age uncertainty into the estimation process, for example by modeling timestamps as random variables, would require further methodological development. However, since the age model uncertainties reported by Westerhold et al. (2020) are small compared to the duration of the estimated climate states, we expect our main findings to be robust.

In addition to age uncertainty, another direction for methodological advancement is developing a breakpoint detection framework for irregularly spaced time series. This would obviate the need for aggregating the data to fixed time intervals, preserving more of the original record. Steps in this direction have already been made in concurrent research (Bennedsen et al., 2024), where the full $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ stacks

(Westerhold et al., 2020) are analyzed while taking the climate state transitions as given and addressing measurement errors.

4 Discussion

Our results demonstrate that the Bai-Perron time-domain framework is a flexible and effective tool for detecting breakpoints in paleoclimate time series. When fixing the number of breakpoints to five and binning the data at 25 kyr intervals, all model specifications lead to breakpoint estimates that closely match those identified by Westerhold et al. (2020), providing strong statistical support for their climate-state classification. This not only corroborates their results obtained through recurrence analysis, but also lends credibility to the use of this econometric tool for analyzing paleoclimate records.

Moreover, information criteria point to a higher number of transitions than previously reported, suggesting the potential for a more detailed classification of Cenozoic climate variability. To explore this, we estimate between one and 15 breakpoints using the Fixed AR model (Fig. 5). Allowing for six breakpoints, we get an additional breakpoint in Warmhouse II at 40.075 Ma that remains for higher numbers of breakpoints. This breakpoint aligns with the cooling following the Middle Eocene Climatic Optimum ~~, a known climatic event~~ (MECO) described by Bohaty and Zachos (2003). Allowing for seven, we ~~get-identify~~ another one in Coolhouse II ~~; allowing for eight, another in Icehouse; allowing for nine, another in Coolhouse I; and so on~~ around 9.975 Ma in the Miocene, which saw the expansion of C4 grasslands, altering the global carbon cycle and land surface with potential effects on climate (Polissar et al., 2019; Strömberg, 2011). Using the same $\delta^{18}\text{O}$ dataset, Rousseau et al. (2023) apply recurrence plots and a Kolmogorov–Smirnov test and identify similar transitions near 40 Ma and 9.7 Ma. Allowing for eight in the Bai-Perron framework, the additional breakpoint occurs in the Icehouse. Some of ~~these breakpoints~~ the additional breakpoints up to 15 coincide with other known climatic events, ~~for instance, the Latest Danian Event at 62.2~~ like the onset of the Mid-Miocene Climatic Optimum (MMCO) with an estimated age of 16.95 Ma (Flower and Kennett, 1994; Zachos et al., 2001). Particularly noteworthy is the lack of breakpoints, even with 15 detections, between the Eocene–Oligocene Transition (EOT) at 34 Ma and the onset of the ~~Miocene Climatic Optimum at 16.9 Ma~~ (Westerhold et al., 2020).

~~The estimation results based on information criteria justify dividing the climate states Warmhouse II and Coolhouse II into two substates each at approximately 39.7 Ma and 10 Ma, respectively~~ MMCO around 17 Ma. This is supported by the presence of breakpoints estimated approximately at these time stamps in the estimations with seven or more breakpoints. ~~Comparable findings are presented in Appendices A.3 and A.4, which detail the results of estimating one to fifteen breakpoints using the Mean and AR models, respectively, with 25-kyr binned data.~~ This is consistent with this being a relatively stable period in the Cenozoic Era following the establishment of the Antarctic ice sheet (Zachos et al., 2001; Mudelsee et al., 2014).

To explore transitions within the relatively higher-resolution Icehouse period more closely, we have re-estimated breakpoints using a finer 5 kyr binning and a reduced minimum regime length of $h = 250$ kyr. This setup yields a single breakpoint at 1.355 Ma, which may reflect a midpoint in the record rather than a distinct climatic shift. When allowing for two breakpoints as also suggested by the BIC, they are estimated at 2.54 Ma and 0.95 Ma, corresponding well to the onset of Northern Hemisphere Glaciation

(Lisiecki and Raymo, 2005) and the Mid-Pleistocene Transition (MPT) (Pisias and Moore, 1981), respectively. The MPT marks a change in the rhythm of glacial cycles, with its timing still debated. For instance, Clark et al. (2006) describe it as a gradual transition between 1.25 and 0.7 Ma. This uncertainty is also reflected in the estimated confidence interval of the latter breakpoint, which spans from 1.545 Ma to 0.66 Ma. These results underscore the capability of the Bai-Perron framework to detect key transitions in Earth’s recent climate history and emphasize the importance of climate system knowledge when interpreting breakpoint estimates. The same period is examined by Rousseau et al. (2023), who apply recurrence plots and Kolmogorov–Smirnov tests to a benthic $\delta^{18}\text{O}$ record (Hodell and Channell, 2016) and identify six transitions at approximately 2.93 Ma, 2.52 Ma, 1.51 Ma, 1.25 Ma, 0.61 Ma, and 0.35 Ma. Their transitions at 2.52, 1.51, and 0.61 Ma broadly align with the first estimated breakpoint and the confidence interval bounds of the second. In contrast to the Bai-Perron framework used here, their approach neither determines the number of breakpoints nor estimates confidence intervals for their placements.

Based on our findings, we offer several general recommendations for the application of this breakpoint detection method in paleoclimate research and related fields. First, careful consideration should be given to the choice of binning frequency. While finer binning enhances temporal resolution, it may also preserve measurement errors and introduce artifacts by linear interpolations, particularly in unevenly sampled records. Also, coarser binning can lead to loss of information. In our application, we find that the bin width 10 and 25 kyr provide a good balance between detail and signal quality. For the 25 kyr bin width, the mean number of observations per bin is approximately 9, and 3.6 for 10 kyr. However, these numbers vary across the sample, being only 3.5 and 1.4, respectively, in the Warmhouse II and increasing to 28.3 and 11.3, respectively, in the Icehouse period. This highlights the importance of accounting for variable sampling resolution when selecting bin widths. For other records, we recommend seeking a similar balance. If the data are already evenly spaced in time, retaining the original resolution is preferable. Second, the model specification should reflect the statistical features of the data, such as trends and autocorrelation. Although the Fixed AR model has performed well in our study, the flexibility of the Bai–Perron framework allows users to adapt the model specification to suit different datasets. Third, the number of breakpoints can be selected based on information criteria.

5 Conclusion

This study ~~presented~~ presents a statistical time-domain approach to ~~identify breakpoints between climate states~~ estimate breakpoints in the Cenozoic Era, using the econometric tools developed by Bai and Perron (1998, 2003). We ~~analyzed~~ analyze the time series of benthic $\delta^{18}\text{O}$ provided by Westerhold et al. (2020), which ~~we made equidistant through mean binning~~ is a widely cited foundational record for many corners of the field of paleoclimatology. Westerhold et al. (2020) identified five breakpoints using recurrence analysis, and our analysis strongly ~~corroborated~~ corroborates the placements of these breakpoints across various model specifications and binning frequencies. Our approach offered the advantage of constructing confidence intervals for the dates of the breakpoints, providing a measure of estimation uncertainty. Based on the results of our simulation study, we advocate using the model specification with a state-independent autoregressive term and state-dependent intercept.

By ~~treating selecting~~ the number of breakpoints ~~as a parameter to be estimated~~ using information criteria, we ~~provided~~ provide statistical justification for more than five breakpoints in the time series. ~~We found evidence that~~ For instance, in specifications with seven or more breakpoints, the five transitions identified by Westerhold et al. (2020) are preserved, while additional breakpoints suggest that both the Warmhouse II and Coolhouse II ~~could each be split into two substates, which were preserved along with the five breakpoints also found by Westerhold et al. (2020) when considering seven or more breakpoints.~~ states could be further divided into substates. This points to the potential for a more detailed classification of Cenozoic climate states, enhancing our understanding of Earth system dynamics. The results also show that the Bai-Perron framework can robustly capture important transitions in Earth’s climate history. Many of the estimated breakpoints align with known climatic events, such as the cooling after the Middle Eocene Climatic Optimum (MECO), the onset of the Mid-Miocene Climatic Optimum (MMCO), and the Mid-Pleistocene Transition (MPT), supporting its use in broader paleoclimate studies.

~~In concurrent research (Bennedsen et al., 2024), we address some other key challenges specific to paleo time series, including uneven time stamps, multiple observations at the same time point, and measurement errors, while taking the breakpoints identified by Westerhold et al. (2020) as given. Future research can then focus on conducting breakpoint detection using the~~ The applicability of the Bai-Perron framework extends well beyond the benthic $\delta^{18}\text{O}$ ~~time series without applying data aggregation techniques.~~ O stack by Westerhold et al. (2020). It can be employed across a wide range of different paleoclimate archives, allowing for investigations of regional differences in the estimated breakpoints, for instance. It can also be applied to other proxies, such as $\delta^{13}\text{C}$ and greenhouse gas concentrations. Furthermore, the method is suitable for detecting both gradual and abrupt transitions, including climatic events such as Dansgaard-Oeschger events (Dansgaard et al., 1993; Livina et al., 2010).

The framework allows for the inclusion of covariates, opening up many possibilities for future applications. For example, incorporating orbital parameters (e.g., eccentricity, obliquity, and precession; Laskar et al., 2004) provides the potential for detecting transitions while controlling for these external effects. Alternatively, one could investigate breaks in the relationship between orbital forcings and paleoclimate variables, reflecting changes in how strongly these external factors influence climate dynamics. A key example is the MPT, marked by a shift in the dominant glacial cycle from 41 kyr to 100 kyr (Berends et al., 2021), the timing of which could be estimated using the Bai-Perron framework.

These examples highlight the broader potential of the framework as a flexible tool for paleoclimate data analysis. Understanding when and how breakpoints in the climate system occurred is essential for interpreting past climate variability and, ultimately, for informing future projections. The Bai-Perron framework provides a statistically rigorous way of estimating these breakpoints, offering new opportunities to deepen our understanding of long-term climate dynamics.

Code and data availability. The data used in this study are available as the supplementary material of Westerhold et al. (2020). The code used to conduct the analysis is based on the R-package *mbreaks* by Nguyen et al. (2023) and the implementation is available upon request.

Author contributions. All authors contributed equally and are listed alphabetically.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. For helpful comments and suggestions, we thank participants at the conferences on Econometric Models of Climate Change (EMCC-VII and VIII) in Amsterdam in 2023 and in Cambridge, UK, in 2024, at the General Assembly of the EGU in Vienna in 2024, and seminar participants at Aarhus University. MB acknowledges funding from the International Research Fund Denmark under grant 7015-00018B.

References

- Andrews, D. W. K. (1991). Heteroskedasticity and Autocorrelation Consistent Covariance Matrix Estimation. *Econometrica*, 59(3):817–858.
- Andrews, D. W. K. and Monahan, J. C. (1992). An Improved Heteroskedasticity and Autocorrelation Consistent Covariance Matrix Estimator. *Econometrica*, 60(4):953–966.
- Bagniewski, W., Ghil, M., and Rousseau, D. D. (2021). Automatic detection of abrupt transitions in paleoclimate records. *Chaos*, 31(11):113129.
- Bai, J. and Perron, P. (1998). Estimating and Testing Linear Models with Multiple Structural Changes. *Econometrica*, 66(1):47–78.
- Bai, J. and Perron, P. (2003). Computation and analysis of multiple structural change models. *Journal of Applied Econometrics*, 18(1):1–22.
- Bai, J. and Perron, P. (2006). *Multiple Structural Change Models: A Simulation Analysis*, page 212–238. Cambridge University Press.
- Bennedsen, M., Hillebrand, E., Koopman, S. J., and Larsen, K. B. (2024). Continuous-time state-space methods for delta-O-18 and delta-C-13. arXiv:2404.05401.
- Berends, C. J., Köhler, P., Lourens, L. J., and van de Wal, R. S. W. (2021). On the Cause of the Mid-Pleistocene Transition. *Reviews of Geophysics*, 59(2):e2020RG000727.
- Boettner, C., Klinghammer, G., Boers, N., Westerhold, T., and Marwan, N. (2021). Early-warning signals for Cenozoic climate transitions. *Quaternary Science Reviews*, 270:107177.
- Bohaty, S. M. and Zachos, J. C. (2003). Significant Southern Ocean warming event in the late middle Eocene. *Geology*, 31(11):1017–1020.
- Burke, K. D., Williams, J. W., Chandler, M. A., Haywood, A. M., Lunt, D. J., and Otto-Bliesner, B. L. (2018). Pliocene and Eocene provide best analogs for near-future climates. *Proceedings of the National Academy of Sciences*, 115(52):13288–13293.

- Caballero, R. and Huber, M. (2013). State-dependent climate sensitivity in past warm climates and its implications for future climate projections. *Proceedings of the National Academy of Sciences*, 110(35):14162–14167.
- Clark, P. U., Archer, D., Pollard, D., Blum, J. D., Rial, J. A., Brovkin, V., Mix, A. C., Pisias, N. G., and Roy, M. (2006). The middle Pleistocene transition: characteristics, mechanisms, and implications for long-term changes in atmospheric pCO₂. *Quaternary Science Reviews*, 25(23):3150–3184. Critical Quaternary Stratigraphy.
- Crowley, T. J. and Hyde, W. T. (2008). Transient nature of late Pleistocene climate variability. *Nature*, 456:226–230.
- Dansgaard, W., Johnsen, S., Clausen, H., Dahl-Jensen, D., Gundestrup, N., Hammer, C., Hvidberg, C., Steffensen, J., Sveinbjörnsdóttir, A., Jouzel, J., and GC, B. (1993). Evidence of general instability of past climate from a 250-kyr ice-core record. *Nature*, 364:218–220.
- Davidson, J., Stephenson, D., and Turasie, A. (2015). Time series modeling of paleoclimate data. *Environmetrics*, 27.
- Dickey, D. and Fuller, W. (1979). Distribution of the Estimators for Autoregressive Time Series With a Unit Root. *Journal of the American Statistical Association*, 74.
- Epstein, S., Buchsbaum, R., Lowenstam, H., and Urey, H. C. (1951). Carbonate-Water Isotopic Temperature Scale. *GSA Bulletin*, 62(4):417–426.
- Fleitmann, D., Burns, S. J., Mudelsee, M., Neff, U., Kramers, J., Mangini, A., and Matter, A. (2003). Holocene Forcing of the Indian Monsoon Recorded in a Stalagmite from Southern Oman. *Science*, 300(5626):1737–1739.
- Flower, B. P. and Kennett, J. P. (1994). The middle miocene climatic transition: East antarctic ice sheet development, deep ocean circulation and global carbon cycling. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 108:537–555.
- Franke, J. G. and Donner, R. V. (2019). Correlating paleoclimate time series: Sources of uncertainty and potential pitfalls. *Quaternary Science Reviews*, 212:69–79.
- Garcia, R. and Perron, P. (1996). An Analysis of the Real Interest Rate under Regime Shifts. *The Review of Economics and Statistics*, 78(1):111–125.
- Goswami, B., Boers, N., Rheinwalt, A., Marwan, N., Heitzig, J., Breitenbach, S., and Kurths, J. (2018). Abrupt transitions in time series with uncertainties. *Nature Communications*, 9(48).
- Hansen, J., Sato, M., Russell, G., and Kharecha, P. (2013). Climate sensitivity, sea level and atmospheric carbon dioxide. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 371(2001):20120294.

- Hodell, D. A. and Channell, J. E. T. (2016). Mode transitions in northern hemisphere glaciation: co-evolution of millennial and orbital variability in quaternary climate. *Climate of the Past*, 12(9):1805–1828.
- Kejriwal, M., Perron, P., and Zhou, J. (2013). Wald tests for detecting multiple structural changes in persistence. *Econometric Theory*, 29(2):289–323.
- Keyes, N. D. B., Giorgini, L. T., and Wettlaufer, J. S. (2023). Stochastic paleoclimatology: Modeling the EPICA ice core climate records. *Chaos*, 33(9):093132. Special Collection: Theory-informed and Data-driven Approaches to Advance Climate Sciences.
- Killick, R., Fearnhead, P., and Eckley, I. A. (2012). Optimal detection of changepoints with a linear computational cost. *Journal of the American Statistical Association*, 107(500):1590–1598.
- Kurozumi, E. and Tuvaandorj, P. (2011). Model selection criteria in multivariate models with multiple structural changes. *Journal of Econometrics*, 164(2):218–238.
- Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A. C. M., and Levrard, B. (2004). A long-term numerical solution for the insolation quantities of the earth. *Astronomy & Astrophysics*, 428(1):261–285. Published online: 23 November 2004.
- Lisiecki, L. E. and Raymo, M. E. (2005). A Pliocene-Pleistocene stack of 57 globally distributed benthic ^{18}O records. *Paleoceanography*, 20(1):PA1003.
- Liu, J., Wu, S., and Zidek, J. V. (1997). On Segmented Multivariate Regressions. *Statistica Sinica*, 7:497–525.
- Livina, V. N., Kwasniok, F., and Lenton, T. M. (2010). Potential analysis reveals changing number of climate states during the last 60 kyr. *Climate of the Past*, 6(1):77–82.
- Lund, R. and Shi, X. (2023). Changepoint Methods in Climatology. *CHANCE*, 36(2):4. ISSN: 1867-2280.
- Marwan, N. (2023). Challenges and perspectives in recurrence analyses of event time series. *Frontiers in Applied Mathematics and Statistics*, 9.
- Marwan, N., Carmen Romano, M., Thiel, M., and Kurths, J. (2007). Recurrence plots for the analysis of complex systems. *Physics Reports*, 438(5):237–329.
- Marwan, N., Donges, J. F., Donner, R. V., and Eroglu, D. (2021). Nonlinear time series analysis of palaeoclimate proxy records. *Quaternary Science Reviews*, 274:107245.
- Mudelsee, M. (2000). Ramp function regression: a tool for quantifying climate transitions. *Computers Geosciences*, 26(3):293–307.
- Mudelsee, M. (2014). *Climate Time Series Analysis: Classical Statistical and Bootstrap Methods*, volume 51 of *Atmospheric and Oceanographic Sciences Library*. Springer, Cham, Heidelberg, New York, Dordrecht, London, 2nd edition.

- Mudelsee, M., Bickert, T., Lear, C. H., and Lohmann, G. (2014). Cenozoic climate changes: A review based on time series analysis of marine benthic $\delta^{18}\text{O}$ records. *Reviews of Geophysics*, 52(3):333–374.
- Mudelsee, M. and Raymo, M. E. (2005). Slow dynamics of the northern hemisphere glaciation. *Paleoceanography*, 20(4):PA4022.
- Nguyen, L., Yamamoto, Y., and Perron, P. (2023). *mbreaks: Estimation and Inference for Structural Breaks in Linear Regression Models*. R package version 1.0.0.
- Oerlemans, J. (2004). Correcting the Cenozoic $\delta^{18}\text{O}$ deep-sea temperature record for Antarctic ice volume. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 208(3):195–205.
- Pisias, N. G. and Moore, T. C. (1981). The evolution of the pleistocene climate: A time series approach. *Earth and Planetary Science Letters*, 52(2):450–458.
- Polissar, P. J., Rose, C., Uno, K. T., Phelps, S. R., and deMenocal, P. (2019). Synchronous rise of African C4 ecosystems 10 million years ago in the absence of aridification. *Nature Geoscience*, 12(8):657–660.
- Reeves, J., Chen, J., Wang, X. L., Lund, R., and Lu, Q. (2007). A Review and Comparison of Changepoint Detection Techniques for Climate Data. *Journal of Applied Meteorology and Climatology*, 46(6):900–915.
- Reikard, G. (2021). Forecasting paleoclimatic data with time series models. *Results in Geophysical Sciences*, 6:100015.
- Rousseau, D.-D., Bagniewski, W., and Lucarini, V. (2023). A Punctuated Equilibrium Analysis of the Climate Evolution of Cenozoic Exhibits a Hierarchy of Abrupt Transitions. *Scientific Reports*, 13:11290.
- Ruggieri, E. (2013). A bayesian approach to detecting change points in climatic records. *International Journal of Climatology*, 33(2):520–528.
- Schütz, N. and Holschneider, M. (2011). Detection of trend changes in time series using bayesian inference. *Physical Review E*, 83(4):041131.
- Shackleton, N. (1967). Oxygen Isotope Analyses and Pleistocene Temperatures Re-assessed. *Nature*, 215:15–17.
- Steffen, W., Rockström, J., Richardson, K., Lenton, T., Folke, C., Liverman, D., Summerhayes, C., Barnosky, A., Cornell, S., Crucifix, M., Donges, J., Fetzer, I., Lade, S., Scheffer, M., and Schellnhuber, H. (2018). Trajectories of the earth system in the anthropocene. *Proceedings of the National Academy of Sciences*, 115:201810141.
- Strömberg, C. A. E. (2011). Evolution of Grasses and Grassland Ecosystems. *Annual Review of Earth and Planetary Sciences*, 39:517–544.
- Telford, R., Heegaard, E., and Birks, H. (2004). All age-depth models are wrong: but how badly? *Quaternary Science Reviews*, 23(1-2):1–5.

- Tierney, J. E., Poulsen, C. J., Montañez, I. P., Bhattacharya, T., Feng, R., Ford, H. L., Hönisch, B., Inglis, G. N., Petersen, S. V., Sagoo, N., Tabor, C. R., Thirumalai, K., Zhu, J., Burls, N. J., Foster, G. L., Godd  ris, Y., Huber, B. T., Ivany, L. C., Turner, S. K., Lunt, D. J., McElwain, J. C., Mills, B. J. W., Otto-Bliesner, B. L., Ridgwell, A., and Zhang, Y. G. (2020). Past climates inform our future. *Science*, 370(6517):eaay3701.
- Trauth, M. H. (2025). *MATLAB   Recipes for Earth Sciences*. Springer Textbooks in Earth Sciences, Geography and Environment. Springer. Access provided by Vrije Universiteit Amsterdam.
- Trauth, M. H., Asrat, A., Berner, N., Bibie, F., Foerster, V., Grove, M., Kaboth-Bahr, S., Maslin, M. A., Mudelsee, M., and Sch  bitz, F. (2021). Northern hemisphere glaciation, african climate and human evolution. *Quaternary Science Reviews*, 268:107099.
- Trauth, M. H., Asrat, A., Fischer, M. L., Hopcroft, P. O., Foerster, V., Kaboth-Bahr, S., Kindermann, K., Lamb, H. F., Marwan, N., Maslin, M. A., Sch  bitz, F., and Valdes, P. J. (2024). Early warning signals of the termination of the african humid period(s). *Nature Communications*, 15(1):3697.
- Valler, V., Franke, J., Brugnara, Y., Samakinwa, E., Hand, R., Lundstad, E., Burgdorf, A.-M., Lipfert, L., Friedman, A. R., and Br  nnimann, S. (2024). Mode-ra: a global monthly paleo-reanalysis of the modern era 1421 to 2008. *Scientific Data*, 11:36.
- Waelbroeck, C., Labeyrie, L., Michel, E., Duplessy, J. C., McManus, J. F., Lambeck, K., Balbon, E., and Labracherie, M. (2002). Sea-level and deep water temperature changes derived from benthic foraminifera isotopic records. *Quaternary Science Reviews*, 21(1–3):295–305.
- Westerhold, T., Marwan, N., Drury, A. J., Liebrand, D., Agnini, C., Anagnostou, E., Barnet, J. S. K., Bohaty, S. M., Vleeschouwer, D. D., Florindo, F., Frederichs, T., Hodell, D. A., Holbourn, A. E., Kroon, D., Lauretano, V., Littler, K., Lourens, L. J., Lyle, M., P  like, H., R  hl, U., Tian, J., Wilkens, R. H., Wilson, P. A., and Zachos, J. C. (2020). An astronomically dated record of Earth’s climate and its predictability over the last 66 million years. *Science*, 369(6509):1383–1387.
- Yao, Y.-C. (1988). Estimating the number of change-points via Schwarz’ criterion. *Statistics & Probability Letters*, 6(3):181–189.
- Zachos, J., MO, P., Sloan, L., Thomas, E., and Billups, K. (2001). Trends, Rhythms, and Aberrations in Global Climate 65 Ma to Present. *Science (New York, N.Y.)*, 292:686–93.
- Zachos, J. C., Dickens, G. R., and Zeebe, R. E. (2008). An early Cenozoic perspective on greenhouse warming and carbon-cycle dynamics. *Nature*, 451:279–283.

A Graphs

A.1 Reversed time




Figure 7: A comparison of estimated breakpoints using the Mean, Fixed AR, and AR model specifications for five breakpoints on 25 kyr binned data where the time frame is reversed. The black dots represent estimated breakpoints, while colored shaded rectangles indicate 95% confidence intervals. The results overlay the $\delta^{18}\text{O}$ data from Westerhold et al. (2020) and their transitions.

A.2 One to 15 breakpoints: Fixed AR model 10 kyr




Figure 8: A comparison of estimated breakpoints using the Fixed AR model for one to 15 breakpoints on 10 kyr binned data. The minimum state length is set to $h = 1$ Myr. The black dots represent estimated breakpoints, while colored shaded rectangles indicate 95% confidence intervals. The results overlay the $\delta^{18}\text{O}$ data from Westerhold et al. (2020) and their transitions.

A.3 One to 15 breakpoints: Mean model




Figure 9: A comparison of estimated breakpoints using the Mean model for one to 15 breakpoints on 25 kyr binned data. The minimum state length is set to $h = 1$ Myr. The black dots represent estimated breakpoints, while colored shaded rectangles indicate 95% confidence intervals. The results overlay the $\delta^{18}\text{O}$ data from Westerhold et al. (2020) and their transitions.

A.4 One to 15 breakpoints: AR model




Figure 10: A comparison of estimated breakpoints using the AR model for one to 15 breakpoints on 25 kyr binned data. The minimum state length is set to $h = 1$ Myr. The black dots represent estimated breakpoints, while colored shaded rectangles indicate 95% confidence intervals. The results overlay the $\delta^{18}\text{O}$ data from Westerhold et al. (2020) and their transitions.

A.5 One and two breakpoints in the Icehouse: Mean model 5 kyr




Figure 11: A comparison of estimated breakpoints using the Mean model for one and two breakpoints on 5 kyr binned data for the Icehouse period. The minimum state length is set to $h = 250$ kyr. The black dots represent estimated breakpoints, while colored shaded rectangles indicate 95% confidence intervals. The results overlay the $\delta^{18}\text{O}$ data from Westerhold et al. (2020).

A.6 One and two breakpoints in the Icehouse: AR model 5 kyr




Figure 12: A comparison of estimated breakpoints using the AR model for one and two breakpoints on 5 kyr binned data for the Icehouse period. The minimum state length is set to $h = 250$ kyr. The black dots represent estimated breakpoints, while colored shaded rectangles indicate 95% confidence intervals. The results overlay the $\delta^{18}\text{O}$ data from Westerhold et al. (2020).

B Tables

B.1 Summary statistics: State-wise and full sample

Bin size	State	Mean	Sd.	Max.	Min.	Data points
5	Warmhouse I	0.417	0.249	1.07	-0.215	2221
5	Hothouse	-0.269	0.261	0.391	-2.014	1800
5	Warmhouse II	0.897	0.366	1.894	-0.254	2600
5	Coolhouse I	2.239	0.233	2.991	1.266	4020
5	Coolhouse II	3.072	0.237	4.172	1.885	2120
5	Icehouse	4.037	0.463	5.405	3.05	660
5	Full sample period	1.561	1.277	5.405	-2.014	13421
10	Warmhouse I	0.417	0.245	0.977	-0.12	1111
10	Hothouse	-0.269	0.256	0.308	-2.014	900
10	Warmhouse II	0.897	0.366	1.777	-0.254	1300
10	Coolhouse I	2.239	0.221	2.877	1.324	2010
10	Coolhouse II	3.072	0.228	4.122	1.975	1060
10	Icehouse	4.034	0.447	5.33	3.181	330
10	Full sample period	1.561	1.276	5.33	-2.014	6711
25	Warmhouse I	0.418	0.237	0.912	-0.065	445
25	Hothouse	-0.269	0.245	0.218	-1.871	360
25	Warmhouse II	0.898	0.358	1.688	0.01	520
25	Coolhouse I	2.239	0.202	2.749	1.391	804
25	Coolhouse II	3.073	0.213	3.793	2.087	424
25	Icehouse	4.033	0.401	5.158	3.258	132
25	Full sample period	1.561	1.273	5.158	-1.871	2685
50	Warmhouse I	0.419	0.233	0.867	-0.042	223
50	Hothouse	-0.268	0.233	0.197	-1.871	180
50	Warmhouse II	0.898	0.354	1.656	0.182	260
50	Coolhouse I	2.24	0.188	2.713	1.567	402
50	Coolhouse II	3.072	0.206	3.72	2.156	212
50	Icehouse	4.042	0.359	4.757	3.264	66
50	Full sample period	1.562	1.271	4.757	-1.871	1343
75	Warmhouse I	0.42	0.229	0.837	0.006	148
75	Hothouse	-0.26	0.203	0.167	-0.985	120
75	Warmhouse II	0.894	0.351	1.553	0.156	173
75	Coolhouse I	2.239	0.181	2.717	1.691	268
75	Coolhouse II	3.068	0.214	3.652	2.072	142
75	Icehouse	4.041	0.351	4.753	3.283	44
75	Full sample period	1.563	1.268	4.753	-0.985	895
100	Warmhouse I	0.42	0.229	0.832	0.007	112
100	Hothouse	-0.263	0.203	0.155	-0.985	90
100	Warmhouse II	0.898	0.349	1.601	0.228	130
100	Coolhouse I	2.241	0.175	2.685	1.739	201
100	Coolhouse II	3.073	0.201	3.625	2.353	106
100	Icehouse	4.047	0.344	4.673	3.4	33
100	Full sample period	1.562	1.269	4.673	-0.985	672
Without binning	Warmhouse I	0.428	0.25	1.07	-0.215	2761
Without binning	Hothouse	-0.279	0.255	0.391	-2.46	3030
Without binning	Warmhouse II	0.916	0.357	1.894	-0.254	1786
Without binning	Coolhouse I	2.251	0.242	3.263	1.026	6669
Without binning	Coolhouse II	3.102	0.254	4.49	1.84	6282
Without binning	Icehouse	4.064	0.533	5.53	2.66	3731
Without binning	Full sample period	2.128	1.445	5.53	-2.46	24259

Table 2: Summary statistics of the binned data with bin sizes (5, 10, 25, 50, 75, and 100 kyr) and the $\delta^{18}\text{O}$ data without binning for each of the states identified by Westerhold et al. (2020) and the full sample period.

B.2 Estimated breakpoints: 5 breakpoints

Bin size	BP index	Mean		Fixed AR		AR	
		Estimate	95% CI	Estimate	95% CI	Estimate	95% CI
5	1	55.965	(56.085, 55.885)	55.995	(56.085, 55.92)	33.745	(33.745, 33.72)
5	2	46.725	(46.845, 46.675)	46.73	(46.76, 46.68)	16.96	(17.365, 16.78)
5	3	34.02	(34.025, 33.915)	34.05	(34.075, 34.015)	13.825	(13.84, 13.775)
5	4	13.36	(13.395, 13.325)	13.41	(13.465, 13.34)	9.555	(9.585, 9.505)
5	5	2.735	(2.845, 2.715)	2.74	(3.1, 2.715)	3.36	(3.815, 3.355)
10	1	55.97	(56.15, 55.79)	55.99	(56.15, 55.88)	33.77	(33.77, 33.72)
10	2	46.73	(46.84, 46.64)	46.73	(46.77, 46.64)	17.88	(18.32, 17.64)
10	3	34.02	(34.03, 33.9)	34.15	(34.18, 34.09)	13.82	(13.84, 13.75)
10	4	13.36	(13.4, 13.3)	13.82	(13.89, 13.72)	9.59	(9.72, 9.45)
10	5	2.73	(2.81, 2.7)	2.74	(3.18, 2.71)	2.74	(2.88, 2.72)
25	1	55.975	(56.3, 55.1)	56.025	(56.575, 55.7)	55.825	(55.85, 55.675)
25	2	46.725	(47.3, 46.55)	46.725	(46.825, 46.45)	48.35	(48.625, 47.85)
25	3	34.025	(34.05, 33.5)	34.15	(34.225, 34.0)	33.75	(33.75, 33.675)
25	4	13.4	(13.525, 13.275)	13.875	(13.975, 13.65)	13.875	(14.05, 13.55)
25	5	2.725	(2.8, 2.625)	2.775	(3.075, 2.7)	2.575	(2.6, 2.55)
50	1	55.95	(56.2, 54.6)	56	(57.1, 55.35)	56	(56.65, 55.7)
50	2	46.7	(48.15, 46.45)	47.1	(47.25, 46.55)	48.8	(49.1, 40.45)
50	3	34.05	(34.05, 32.8)	34.2	(34.3, 33.9)	33.75	(33.75, 33.6)
50	4	13.8	(14.15, 13.6)	13.85	(14.0, 13.45)	16.95	(17.35, 16.7)
50	5	2.75	(2.9, 2.5)	3.15	(3.4, 3.0)	14.3	(14.55, 12.8)
75	1	55.95	(56.325, 53.775)	56.25	(57.45, 54.75)	55.95	(56.325, 55.5)
75	2	46.725	(50.625, 46.425)	47.1	(47.475, 46.425)	53.325	(53.625, 50.1)
75	3	34.05	(34.05, 30.9)	34.2	(34.425, 33.675)	34.05	(34.05, 33.825)
75	4	13.35	(13.8, 12.975)	13.875	(14.1, 13.125)	16.95	(17.325, 16.5)
75	5	2.775	(3.375, 2.4)	3.15	(3.525, 2.925)	14.475	(15.075, 14.25)
100	1	56	(56.4, 54.0)	56.2	(57.7, 54.5)	56	(56.3, 55.5)
100	2	46.7	(52.5, 46.3)	47.1	(47.7, 46.3)	53.4	(53.8, 52.1)
100	3	34.1	(34.1, 29.4)	34.2	(34.5, 33.4)	49.1	(50.8, 48.8)
100	4	13.8	(14.7, 13.4)	13.9	(14.1, 12.9)	34.1	(34.1, 33.8)
100	5	2.9	(4.2, 2.3)	3.4	(3.8, 3.2)	13.8	(15.7, 12.9)

Table 3: Estimated breakpoints and their 95% confidence intervals (in Ma) where the number of breakpoints is fixed to 5, and all values are rounded to three decimals. The table shows estimates for each method across bin sizes 5, 10, 25, 50, 75, and 100 kyr.

B.3 Estimated parameters: 5 breakpoints and 25 kyr binned data

Parameter	Mean		Fixed AR		AR	
	Estimate	SE	Estimate	SE	Estimate	SE
c_1	0.418	0.051	0.069	0.008	-0.001	0.026
c_2	-0.256	0.040	-0.043	0.007	-0.108	0.015
c_3	0.911	0.072	0.153	0.013	0.028	0.007
c_4	2.247	0.017	0.373	0.031	0.660	0.061
c_5	3.119	0.027	0.519	0.043	0.421	0.073
c_6	4.140	0.051	0.698	0.057	2.423	0.326
φ	×	×	0.833	0.014	×	×
φ_1	×	×	×	×	0.990	0.054
φ_2	×	×	×	×	0.631	0.037
φ_3	×	×	×	×	0.970	0.008
φ_4	×	×	×	×	0.706	0.027
φ_5	×	×	×	×	0.865	0.024
φ_6	×	×	×	×	0.419	0.081
σ_1^2	0.237	×	0.095	×	0.106	×
σ_2^2	0.255	×	0.154	×	0.140	×
σ_3^2	0.347	×	0.112	×	0.107	×
σ_4^2	0.210	×	0.141	×	0.140	×
σ_5^2	0.208	×	0.111	×	0.116	×
σ_6^2	0.351	×	0.340	×	0.315	×

Table 4: Estimated parameters and their corresponding standard errors (SE) for each model specification. Parameters absent in a given model specification are denoted by ×. The number of breakpoints is set to 5, and the parameters are estimated with a binning frequency of 25 kyr and $h = 2.5$ Myr. All values are rounded to three decimals.

B.4 The number of breakpoints selected by information criteria

Bin size	Mean		Fixed AR		AR	
	BIC	LWZ	BIC	LWZ	BIC	LWZ
5	19	17	17	7	15	5
10	17	17	14	7	14	3
25	17	14	12	6	8	3
50	17	14	10	0	7	0
75	17	14	6	0	5	0
100	17	12	6	0	5	0

Table 5: The number of breakpoints selected using BIC and LWZ criterion for all models and binning frequencies considered. The minimum state length is set to $h = 2.5$ Myr and the maximum number of breakpoints is 26.

C Simulation study

C.1 Serially uncorrelated error term

In this appendix, we assess whether the methodology by Bai and Perron (1998, 2003) can be used to accurately estimate the number and timing of breakpoints in a state-wise non-stationary time series. We conduct 1000 simulations for each data-generating process (DGP) with a sample size of 500. All the DGPs considered have the following form,

$$\begin{aligned} y_t &= c_1 + \varphi_1 y_{t-1} + \varepsilon_t, & \varepsilon_t &\stackrel{i.i.d.}{\sim} \mathcal{N}(0, \sigma^2) & \text{for } t \leq T/2 \\ y_t &= c_2 + \varphi_2 y_{t-1} + \varepsilon_t, & \varepsilon_t &\stackrel{i.i.d.}{\sim} \mathcal{N}(0, \sigma^2) & \text{for } t > T/2. \end{aligned} \quad (8)$$

Hence, we consider a single breakpoint in the middle of the sample interval, namely at $t = 250$. We examine eight DGPs, each specified and described in Table 6.

DGP	σ	c_1	c_2	φ_1	φ_2	Description
1	1	0.1	0.2	1	1	Small break in the drift term of a RW
2	1	0.1	1	1	1	Large break in the drift term of a RW
3	1	0.1	1	0.95	0.95	Large break in the intercept and a fixed AR-coefficient
4	1	0.1	1	0.95	1	Break in the intercept and small break in the AR-coefficient
5	1	0.1	1	0.5	1	Break in the intercept and large break in the AR-coefficient
6	1	1	1	1	1	RW with a drift <i>without</i> a breakpoint
7	0.5	0.1	1	1	1	Large break in the drift of a RW with low variance
8	1	0.1	1	0.5	0.5	Large break in the intercept and a low fixed AR-coefficient

Table 6: Data-generating processes for the simulation study and short descriptions. RW: random walk.

The DGPs range from random walk models with a break in the drift term to models with breaks in both the intercept and the AR coefficient. For comparison, we include a random walk without breakpoints as the sixth model. For each of the DGPs, we are interested in the performance of the methodology by Bai and Perron (1998, 2003) in estimating the breakpoint and confidence intervals. The model specifications from Section 2.3 are estimated on the data generated by the DGPs, and we use the implementation outlined in Section 2.4. We use the R-package *mbreaks* by Nguyen et al. (2023), and we impose a single breakpoint in the estimation. The left and right panels of Figs. 13 through 20 display realizations of the DGP and density plots of the estimated breakpoints for each of the models, respectively. The results are summarized in Table 7, which provides the mean of the estimated breakpoints, and medians of the lower and upper boundaries of the estimated 95% CIs are tabulated along with their coverage rates for each model and DGP.

DGP	Mean				Fixed AR				AR			
	BP est.	Lower	Upper	Coverage	BP est.	Lower	Upper	Coverage	BP est.	Lower	Upper	Coverage
1	301	174	655	57.1%	251	216	336	43.4%	290	240	316	22.7%
2	333	-386	332	95.4%	249	237	262	93%	249	236	256	77.2%
3	263	253	284	41.4%	256	239	260	89.9%	251	241	260	85.9%
4	340	-190	340	97.5%	249	239	260	95.8%	249	238	250	65.8%
5	340	-114	340	97.1%	250	239	258	97%	250	241	250	72.9%
6	249	-3325	3976	×	253	142	371	×	254	202	312	×
7	333	-282	330	92%	249	246	253	97.8%	249	246	253	96%
8	249	237	264	95.1%	248	236	263	95.2%	248	236	263	94.5%

Table 7: Mean of the estimated breakpoints and medians of the lower and upper boundary of the estimated confidence intervals, along with the coverage rates for each model specification and DGP. DGP 6 is simulated without a breakpoint, so the coverage rate is irrelevant and indicated by $\times$.

In the first DGP, a random walk with a small drift term break, we observe that the mean of the estimated breakpoints is later than the true breakpoint in all model specifications. Additionally, the density plots exhibit asymmetry around the true breakpoint. This is expected due to the low magnitude of the break in the drift term, which creates a subtle change in the overall stochastic trend, making accurate breakpoint detection difficult. In the second DGP with a larger drift term break, the estimated breakpoints exhibit a narrower and more bell-shaped density. The mean estimated breakpoints for the Fixed AR and AR models slightly precede the true breakpoint. However, the Mean model performs poorly, with the mean of the estimated breakpoints far from the true breakpoint.

In the third DGP, both the Fixed AR and AR models produce mean estimated breakpoints slightly later than the true breakpoint. The Mean model exhibits better performance in this DGP than in the second DGP. The fourth DGP has a break in the intercept and the AR-coefficient from 0.95 to 1, resulting in a state-wise non-stationary model. This change leads to breakpoint estimates very close to the true breakpoint, except in the Mean model. A similar outcome is observed in the fifth DGP, which features a larger increase in the AR-coefficient. In the sixth DGP, which is defined without any breakpoints, the Mean model estimates breakpoints near the midpoint of the sample period, while the other two specifications yield inconclusive results. In the seventh DGP, the AR and Fixed AR models produce estimates close to the true breakpoint. However, the Mean model continues to produce breakpoint estimates far from the true value. Examining the eighth DGP, the three models perform almost equally well.

Overall, the Fixed AR and AR models tend to perform well in non-stationary scenarios, estimating breakpoints close to the true breakpoints. The methodology, however, appears to struggle with accurately estimating the true breakpoint in cases of minor changes between states and large error term variance. In contrast, the Mean model does not perform well in DGPs featuring gradual changes, aligning with theoretical expectations as detailed in Bai and Perron (2003).

The coverage rate of a CI is the proportion of times the CI covers the true breakpoint, here at $t = 250$. We find that the CIs of the Mean model are generally very wide and have varying coverage. In the Fixed AR and AR models, the CIs are typically narrower. The coverage rates are best in the DGPs with large differences between the states as seen in DGPs 4, 5, 7 and 8 using the Fixed AR model specification, which is in line with the findings of Bai and Perron (2003). For the AR model, the coverage rates are only close

to the desired 95% in the seventh and eighth DGP, indicating that the CIs are inadequate in most of the DGPs considered.

DGP	Mean			Fixed AR			AR		
	BIC	LWZ	KT	BIC	LWZ	KT	BIC	LWZ	KT
1	3.0 (0%)	3.0 (0%)	3.0 (0%)	0.2 (15%)	0.0 (0%)	3.0 (0%)	0.1 (6%)	0.0 (0%)	0.0 (3%)
2	3.0 (0%)	3.0 (0%)	3.0 (0%)	1.0 (97%)	0.8 (82%)	3.0 (0%)	1.0 (94%)	0.5 (46%)	1.0 (93%)
3	2.9 (0%)	2.7 (4%)	3.0 (0%)	1.0 (94%)	0.2 (16%)	2.9 (0%)	0.9 (85%)	0.0 (0%)	0.7 (70%)
4	3.0 (0%)	3.0 (0%)	3.0 (0%)	1.0 (98%)	1.0 (98%)	2.8 (0%)	1.0 (99%)	0.9 (92%)	1.0 (99%)
5	3.0 (0%)	3.0 (0%)	3.0 (0%)	1.0 (99%)	1.0 (97%)	2.7 (0%)	1.0 (99%)	1.0 (100%)	1.0 (99%)
6	3.0 (0%)	3.0 (0%)	3.0 (0%)	0.0 (98%)	0.0 (100%)	3.0 (0%)	0.0 (100%)	0.0 (100%)	0.0 (100%)
7	3.0 (0%)	3.0 (0%)	3.0 (0%)	1.0 (99%)	1.0 (100%)	3.0 (0%)	1.0 (98%)	1.0 (100%)	1.0 (98%)
8	1.5 (63%)	1.0 (98%)	1.3 (72%)	1.0 (99%)	1.0 (100%)	1.3 (73%)	1.0 (100%)	1.0 (98%)	1.0 (100%)

Table 8: Means of the estimated number of breakpoints for each model specification across different DGPs, rounded to one decimal. Percentages indicate the proportion of estimates equal to the true number of breakpoints.

Table 8 shows the mean number of breakpoints estimated for each DGP and method, along with the proportion of correctly estimated breakpoints. The difficulty in accurately estimating gradual changes using the Mean model is also evident when estimating the number of breakpoints. This model specification leads to overestimating the number of breakpoints in all DGPs considered except DGP 8, where it performs well. The BIC criterion in the Fixed AR specification performs very well, with an estimated number of breakpoints equal to the true number in most simulations in DGP 2-8. The LWZ criterion performs almost equally well except in the third DGP, while the KT criterion vastly overestimates the number of breakpoints in DGP 1-7. In the AR model, the information criteria all perform well in DGPs 2-8 except for the third DGP where the LWZ criterion underestimates the number of breakpoints.




Figure 13: DGP 1: Left: Five process realizations. Right: The densities of the estimated breakpoints for each specification.




Figure 14: DGP 2: Left: Five process realizations. Right: The densities of the estimated breakpoints for each specification.




Figure 15: DGP 3: Left: Five process realizations. Right: The densities of the estimated breakpoints for each specification.




Figure 16: DGP 4: Left: Five process realizations. Right: The densities of the estimated breakpoints for each specification.




Figure 17: DGP 5: Left: Five process realizations. Right: The densities of the estimated breakpoints for each specification.




Figure 18: DGP 6: Left: Five process realizations. Right: The densities of the estimated breakpoints for each specification.




Figure 19: DGP 7: Left: Five process realizations. Right: The densities of the estimated breakpoints for each specification.




Figure 20: DGP 8: Left: Five process realizations. Right: The densities of the estimated breakpoints for each specification.

C.2 Serially correlated error term

A possible extension of the simulation study outlined in Eq. (8) is allowing the error term to exhibit serial correlation. We use the same DGPs as before, but generate $\{\varepsilon_t\}_{t=1}^T$ as follows,

$$\varepsilon_t = \psi\varepsilon_{t-1} + \theta\eta_{t-1} + \eta_t, \quad \eta_t \stackrel{i.i.d.}{\sim} \mathcal{N}(0, \sigma_\eta^2) \quad \forall t. \quad (9)$$

We conduct 1000 simulations for each, with a sample size of 500. Here, we consider DGPs 2, 3, 4, 5, 7, and 8 as outlined in Table 6 and refer to these DGPs in the serially correlated cases as models 2_s , 3_s , 4_s , 5_s , 7_s , and 8_s . We set $\psi = \theta = 0.5$ and the standard deviation σ_η , such that the standard deviation of ε_t corresponds to the σ in Table 6. This is accomplished as follows,

$$\begin{aligned} \text{Var}(\varepsilon_t) &= \text{Var}(\psi\varepsilon_{t-1} + \theta\eta_{t-1} + \eta_t) \\ &= \psi^2 \text{Var}(\varepsilon_{t-1}) + \theta^2 \text{Var}(\eta_{t-1}) + 2\psi\theta \text{Cov}(\varepsilon_{t-1}, \eta_{t-1}) + \text{Var}(\eta_t) \\ &= \psi^2 \text{Var}(\varepsilon_{t-1}) + \theta^2 \sigma_\eta^2 + 2\psi\theta \sigma_\eta^2 + \sigma_\eta^2, \end{aligned}$$

since ε_{t-1} and η_{t-1} have zero means and $\mathbb{E}[\varepsilon_t \eta_t] = \phi \mathbb{E}[\varepsilon_{t-1} \eta_t] + \theta \mathbb{E}[\eta_t \eta_{t-1}] + \mathbb{E}[\eta_t^2] = \sigma_\eta^2$. Given stationarity of the process, which implies $\sigma^2 = \text{Var}(\varepsilon_t)$ for all t , we derive,

$$\sigma_\eta^2 = \sigma^2 \frac{1 - \psi^2}{1 + \theta^2 + 2\psi\theta}.$$

This adjustment ensures the comparability of the results between the two error term types.

In Figs. 21 through 26, we plot examples of realizations and frequency plots of the estimated breakpoints using each of the models while imposing a single breakpoint in the estimation. The results are summarized

in Table 9, which provides means of the estimated breakpoints and medians of the lower and upper boundary of the estimated confidence intervals, along with the coverage rates for each model specification and DGP. Generally speaking, the mean of the estimated breakpoints are further from the true breakpoint and the CIs become wider compared to the results from the corresponding DGPs without serial correlation. It is evident that serial correlation in the error term makes it more difficult to estimate the dating of breaks. We find that the Fixed AR and AR models perform well for DGP 7_s , which has a large difference between the states and low variance. This is in line with the theoretical framework by Bai and Perron (2003), who note that the estimated break dates are consistent even in the presence of serial correlation. The Fixed AR model performs well in DGPs 2_s , 4_s and 5_s where the mean of the estimated breakpoints is close to the true breakpoint, and confidence intervals are reasonably wide with acceptable coverage rates. The results of the AR model are less conclusive.

For the Mean and Fixed AR models, the coverage rates are generally close to the desired 95% and even higher in some DGPs. However, the CIs are also extremely wide, reaching outside the sample window in many DGPs. The CIs seem reasonable in the Fixed AR model for DGPs 2_s , 4_s , 5_s , and 7_s , where the coverage rates are close to 95% and the medians of the lower and upper bounds of the CIs are not too extreme. The CIs for the AR model are generally wider than in the version without serial correlation in the error term. In the AR model, the coverage rates are lower than the desired 95%, but it seems that DGPs with large breaks have higher coverage rates. The relatively poor performance is in line with the theoretical framework by Bai and Perron (2003). The authors note that the construction of the CIs rely on having no serial correlation in the error term if a lagged dependent variable is included as a regressor that has coefficients that are subject to breakpoints.

DGP	Mean				Fixed AR				AR			
	BP est.	Lower	Upper	Coverage	BP est.	Lower	Upper	Coverage	BP est.	Lower	Upper	Coverage
2_s	332	-1400	335	95.9%	247	188	312	95.7%	261	190	299	79.9%
3_s	266	60	787	90.6%	285	-112	656	97.2%	276	156	421	77.1%
4_s	340	-776	339	94.9%	252	197	301	96.9%	264	195	277	84.9%
5_s	342	-329	340	96.2%	256	196	266	96.4%	259	192	250	70.8%
7_s	333	-1708	329	92.3%	249	230	270	97.6%	251	230	267	92.8%
8_s	250	122	370	98.3%	245	-5	492	99.8%	247	23	490	97.4%

Table 9: Mean of the estimated breakpoints and medians of the lower and upper boundary of the estimated confidence intervals, along with the coverage rates for each model specification and DGP.

Table 10 shows the mean number of breakpoints estimated for each DGP and method, along with the proportion of correctly estimated number. In the Mean model, all information criteria overestimate the number of breakpoints. An important exception is the eighth DGP, where the performance is better, as in the case without serial correlation. In the Fixed AR and AR model specifications, the LWZ criterion generally performs well, while both the BIC and the KT criteria generally overestimate. However, the LWZ criterion leads to underestimating the number of breakpoints in DGPs 3_s and 8_s . These two DGPs are characterized by fixed AR-coefficients that are lower than one. This implies that these two processes do not exhibit an autoregressive unit root. Hence, it seems that the LWZ criterion performs well in cases of state-wise non-stationarity or switching between stationary and non-stationary states.

Compared to the findings in the DGPs without serial correlation, it is clear that the proportion of correct estimates are lower for most DGPs and model specifications. Overall, the best performing criterion seems to be the LWZ criterion in the Fixed AR and AR models, while the Mean model typically leads to overestimating the number of breakpoints.

DGP	Mean			Fixed AR			AR		
	BIC	LWZ	KT	BIC	LWZ	KT	BIC	LWZ	KT
2_s	3.0 (0%)	3.0 (0%)	3.0 (0%)	1.9 (32%)	0.9 (70%)	2.9 (0%)	1.8 (37%)	0.7 (61%)	1.9 (33%)
3_s	3.0 (0%)	2.8 (2%)	3.0 (0%)	0.7 (33%)	0.0 (0%)	2.7 (3%)	0.3 (19%)	0.0 (0%)	0.4 (17%)
4_s	3.0 (0%)	3.0 (0%)	3.0 (0%)	1.7 (45%)	1.0 (85%)	2.8 (1%)	1.6 (51%)	0.8 (79%)	1.6 (47%)
5_s	3.0 (0%)	3.0 (0%)	3.0 (0%)	1.8 (5%)	1.1 (85%)	2.8 (0%)	1.7 (40%)	1.0 (92%)	1.6 (49%)
7_s	3.0 (0%)	3.0 (0%)	3.0 (0%)	1.9 (34%)	1.1 (89%)	3.0 (0%)	1.9 (34%)	1.0 (96%)	1.9 (32%)
8_s	2.2 (21%)	1.2 (78%)	2.2 (23%)	0.4 (35%)	0.0 (0%)	1.9 (36%)	0.0 (4%)	0.0 (0%)	0.0 (3%)

Table 10: Means of the estimated number of breakpoints for each model specification across different DGPs, rounded to one decimal. Percentages indicate the proportion of estimates equal to the true number of breakpoints.




Figure 21: DGP 2_s : Left: Five process realizations. Right: The densities of the estimated breakpoints for each specification.




Figure 22: DGP 3_s : Left: Five process realizations. Right: The densities of the estimated breakpoints for each specification.




Figure 23: DGP 4_s: Left: Five process realizations. Right: The densities of the estimated breakpoints for each specification.




Figure 24: DGP 5_s: Left: Five process realizations. Right: The densities of the estimated breakpoints for each specification.




Figure 25: DGP 7_s: Left: Five process realizations. Right: The densities of the estimated breakpoints for each specification.




Figure 26: DGP 8_s: Left: Five process realizations. Right: The densities of the estimated breakpoints for each specification.