



The importance of an informed choice of CO₂-equivalence metrics for contrail avoidance

Audran Borella¹, Olivier Boucher¹, Keith P. Shine², Marc Stettler³, Katsumasa Tanaka^{4,5}, Roger Teoh³ and Nicolas Bellouin^{1,2}

5 ¹Institut Pierre-Simon Laplace, Sorbonne Université/CNRS, Paris, France

²Department of Meteorology, University of Reading, Reading, UK

³Centre for Transport Studies, Department of Civil and Environmental Engineering, Imperial College London, London, UK

⁴Laboratoire des Sciences du Climat et de l'Environnement (LSCE), IPSL, CEA/CNRS/UVSQ, Université Paris-Saclay, Gif-sur-Yvette, France

10 ⁵Earth System Division, National Institute for Environmental Studies (NIES), Tsukuba, Japan

Correspondence to: Audran Borella (audran.borella@ipsl.fr)

Abstract. One of the proposed ways to reduce the climate impact of civil aviation is rerouting aircraft to minimise the formation of warming contrails. As this strategy may increase fuel consumption, it would only be beneficial if the climate impact reduction from the avoided contrails exceeds the negative impact of any additional carbon dioxide (CO₂) emitted by the rerouted flight. In this study, we calculate the surface temperature response of almost half-a-million flights that crossed the North Atlantic sector in 2019 and compare to the response of hypothetical rerouted flights. The climate impacts of contrails and CO₂ are assessed through the perspective of CO₂-equivalence metrics, defined here as nine combinations of different definitions and time horizons. We estimate that the total emitted CO₂ and the persistent contrails formed will have warmed the climate by 16.9 μK in 2039, 13.5 μK in 2069, and 14.0 μK in 2119. Under a scenario where 1% additional carbon dioxide is enough to reroute all contrail-forming flights and avoid contrail formation completely, total warming would decrease by 4.6 (−27%), 2.4 (−18%), and 1.8 (−13%) μK in 2039, 2069, and 2119, respectively. In most rerouting cases, the results based on the nine different CO₂-equivalence metrics agree that rerouting leads to a climate benefit, assuming that contrails are avoided as predicted. But the size of that benefit is very dependent on the choice of CO₂-equivalence metrics, contrail efficacy and CO₂ penalty. Sources of uncertainty not considered here could also heavily influence the perceived benefit. In about 10% of rerouting cases, the climate damage resulting from contrail avoidance indicated by CO₂-equivalence metrics integrated over a 100-year time horizon is not predicted by metrics integrated over a 20-year time horizon. This study highlights, using North Atlantic flights as a case study, the implications of the choice of CO₂-equivalence metrics for contrail avoidance, but the choice is ultimately political.

1 Introduction

30 Civil aviation mostly contributes to climate change through the in-flight emissions of commercial aircraft. These emissions include carbon dioxide (CO₂): the global aviation fleet represented about 2.4% of the global annual fossil-fuel CO₂ emissions

in 2018 (ICCT, 2018). However, kerosene combustion also emits non-CO₂ species such as nitrogen oxides (NO_x), water vapour, and soot particles. These emissions exert the so-called non-CO₂ effects of aviation (EASA, 2020). According to current knowledge, non-CO₂ effects could exert two thirds of the total effective radiative forcing due to civil aviation since its inception in 1940 (Lee et al., 2021). Uncertainties in the estimates of non-CO₂ effects are large and subject of active research.

Contrails (condensation trails) form when atmospheric conditions allow water vapour to condense onto aerosols emitted by the incomplete combustion of kerosene impurities in the aircraft engines (Schumann et al., 2012; Kärcher, 2018). Contrails that form in ice supersaturated regions (ISSRs) of the upper atmosphere persist to form contrail cirrus clouds. These clouds reflect a fraction of incoming sunlight back to space, cooling the climate system, but they also absorb and re-emit infrared radiation at colder temperatures than the underlying surface and atmosphere, leading to a warming. The overall impact depends on the contrail properties, its ambient environment, and the time of day. Some persistent contrails cool the Earth, but most have a warming effect (Wilhelm et al., 2021; Wang et al., 2023). Taken together, contrail cirrus are currently thought to be the largest contributor of climate forcing by aviation (Figure 6.16 of Szopa et al. 2021) and hence are the focus of this study. A more comprehensive discussion would need to assess the impact of other non-CO₂ effects, such as NO_x and aerosols. There is an interdependency between them: for example, aerosols play a role in contrail formation but also exert their own radiative forcing, directly by interacting with radiation or via their interactions with clouds, but the magnitude and size of this forcing is highly uncertain (Lee et al., 2021).

On average, 13% of flight time occurs in ISSRs (Gierens et al., 1999), and studies suggest that only a minority of flights in a given airspace are responsible for most of the higher-impact persistent contrails (Teoh et al., 2020, 2022, 2023). It is therefore tempting to avoid flight trajectories or altitudes that encounter ISSRs to avoid the formation of persistent contrails. This strategy assumes that the location and properties of ISSRs can be accurately predicted on both the original and rerouted trajectories, and it also assumes that other considerations, including impacts on air traffic management, safety and flight duration, make a reroute practical. In addition, the rerouted aircraft could deviate from its optimal trajectory in terms of fuel consumption and therefore CO₂ emissions, thus requiring methods to quantify the trade-off between CO₂ and non-CO₂ climate impacts.

The radiative and climate impacts of CO₂ and non-CO₂ effects are usually compared using a CO₂-equivalence metric as a multiplicative factor that operates on the CO₂-only effect (Fuglestedt et al., 2010). For example, a factor of 2 indicates a net CO₂ and non-CO₂ effect that is twice as large as that of the CO₂ effect alone, while a factor of 1 indicates that the net non-CO₂ effects are zero. The multiplicative factor can thus translate emissions of aviation into CO₂-equivalent emissions.

It is now well accepted that the choice of CO₂-equivalency metric depends on the perspective and objectives of climate policies (Fuglestedt et al., 2010; Tanaka et al., 2010; Forster et al., 2021). The UNFCCC (United Nations Framework Convention on Climate Change) decided in 1995 for its Kyoto protocol to compute CO₂-equivalent emissions of greenhouse gas emissions using a metric definition based on the integration of radiative forcing, called the Global Warming Potential (GWP), over a time horizon of 100 years. That decision was reconfirmed in 2018 (UNFCCC, 1995, 2019). The initial choice

was informed by the First Assessment Report of the Intergovernmental Panel on Climate Change (IPCC), which assessed GWP for 20-, 100-, and 500-year time horizons in its Table 2.8 (Shine et al., 1990). The report noted that these “three different time horizons are presented as candidates for discussion and should not be considered as having any special significance” (Shine, 2009). UNFCCC therefore had a limited choice of CO₂-equivalence metric, since GWP was the only
70 metric assessed by the IPCC at the time. It is believed that the 100-year time horizon may have been chosen as a middle-ground between shorter and longer options. The EU Emission Trading System followed in 2003 the UNFCCC choice of GWP (European Commission, 2003, Directive 2003/87/EC). This decision does not mandate a particular time horizon, but 100 years is likely to remain the most widely used. In its most recent assessment (AR6), the IPCC refrained from recommending an emission metric (Box 7.3 of Forster et al. 2021), noting that the choice “depends on the purposes for
75 which [...] forcing agents are being compared”. Although the UNFCCC choice of GWP100 was based on the limited research available in the mid-1990s, it is becoming clearer that GWP100 is relatively close to an economically optimal metric at least currently for achieving the long-term temperature goals of the Paris Agreement (see Cross Chapter 2 of Dhakal et al., 2022; Tanaka et al., 2021).

In the case of aviation, Grewe et al. (2014) studied the impact of metric choice on estimating the short, medium, and long-
80 term climate impacts of re-routing strategies, and found that some combinations of metrics and horizon put a greater value on reducing the radiative forcing of short-lived components like contrails, while other combinations put a greater value on reducing the forcing due to CO₂. Aviation is not the only sector where non-CO₂ emissions are a significant fraction of the CO₂-equivalent emissions. Most notable are methane emissions from the agricultural sector for which the CO₂-equivalence is also heavily dependent on the choice of metric and time horizon, and the debate and lack of scientific consensus on that
85 choice continues, as noted in the latest report of the Food and Agriculture Organization (FAO) on the topic (FAO, 2023).

The present study revisits the question of the choice of a CO₂-equivalence metric definition and time horizon in the context of contrail avoidance, using the actual 2019 North Atlantic air traffic to quantify the climate outcome of that choice. It includes sensitivity analysis of the results to the amount of additional CO₂ needed to reroute a flight, to the capability of the rerouting to fully avoid the formation of a persistent contrail, and to the contrail efficacy. Section 2 details the metrics used
90 to quantify the impact of contrails, and the CO₂-equivalence metric definition that allows comparison of CO₂ and contrails climate impacts. Section 3 describes the air traffic dataset used in this study and the contrail and Earth system models used to calculate the surface temperature change exerted by a flight based on its CO₂ emissions and contrail radiative forcing. Section 4 discusses how the different time scales of the climate perturbations of CO₂ and contrails affect the results based on different CO₂-equivalence metrics, focusing on nine metrics, which are the combinations of three different metric definitions
95 (global warming potential, global temperature change potential, and average temperature response) with three alternative time horizons (20, 50, and 100 years). Section 5 uses case studies to illustrate how the climate outcome of re-routing decisions depends on the combination of metric definition and time horizon. Section 6 assesses the climate implications of re-routing strategies applied to contrail-forming aircraft traffic over the North Atlantic for the year 2019, with simplified assumptions on the additional CO₂ emitted to avoid contrail formation. Sections 7 and 8 investigate the sensitivity of the



100 results to more realistic rerouting and to contrail efficacy, respectively. Section 9 concludes the study, highlighting the implications of radiative forcing uncertainties.

2. CO₂-equivalence metrics

It is impractical to run climate models to quantify the different contributions to temperature change for individual activities, or individual flights in our case, so different CO₂-equivalence metrics (also called “climate metrics”, “emission metrics” or
105 simply “metrics” in the literature) have been developed to compare CO₂ and non-CO₂ effects (Fuglestvedt et al., 2010). Such a comparison can be done in terms of the time evolution of the radiative forcing ΔF exerted by the constituent, or of the temperature change ΔT that follows from ΔF .

The Absolute Global Warming Potential (AGWP) measures the cumulative radiative forcing exerted by a pulse emission of the constituent of interest over a given time horizon H (Fuglestvedt et al., 2010):

$$110 \quad AGWP_H = \int_0^H \Delta F(t) dt \quad (1)$$

Relating the AGWP_H of contrails to a pulse emission is not straightforward, since contrails are not emitted. Instead, contrails form owing to the release of water vapour and aerosols from fuel combustion, and one estimates the AGWP_H of a contrail formed from the energy it adds to or removes from the Earth system during its lifetime. AGWP_H is a time-integrated metric, and because it is based on radiative forcing, it is not an explicit measure of the climate response.

115 To consider the climate response, the Absolute Global Temperature Change Potential (AGTP; Shine et al., 2005) measures the change in global surface temperature at time horizon H after a pulse emission:

$$AGTP_H = \Delta T(H) \quad (2)$$

Unlike AGWP_H, AGTP_H is an endpoint metric. Corresponding integrated metrics have been proposed that measure global surface temperature change averaged over H : the integrated GTP (Peters et al., 2011), mean GTP (Gillett and Matthews,
120 2010), or Average Temperature Response (ATR, Grewe et al. 2021). The latter has been used regularly in aviation-climate studies following the publication of Grewe et al. (2014), so will be considered here. ATR_H is the average change in temperature between the time of an emission at $t = 0$ and the end of the time horizon H :

$$ATR_H = \frac{1}{H} \int_0^H \Delta T(t) dt \quad (3)$$

Combined pulse/sustained metrics, like the GWP* (Allen et al. 2018) and CGTP (Collins et al. 2020), have been proposed to
125 ensure equivalency in the temperature response of short-lived and long-lived climate forcers (Allen et al., 2022). Those metrics are not considered here because they are most suited for quantifying CO₂/non-CO₂ trade-off for future emission scenarios rather than for quantifying potential reductions in climate impact for individual flights, which is the topic of this study.



Note that, as detailed in the next section, we calculate the time-dependent temperature response, then apply the perspective of CO₂-equivalence metrics to interpret the results, without relying on previously published values of CO₂-equivalence metrics (see Cross Chapter 2 of Dhakal et al., 2022).

3. Methods

This study considers flight trajectories, their associated fuel consumption and CO₂ emissions, and, when relevant, the persistent contrails formed. Each specific persistent contrail is described using its energy forcing EF_{contrail} (Teoh et al., 2020), in joules, which quantifies the total energy added to or lost by the Earth system by the contrail during its lifetime while accounting for the length of the contrail L , its width W , and lifetime τ :

$$EF_{\text{contrail}} = \int_0^{\tau} \Delta F(t) \times W(t) \times L(t) \times dt, \quad (4)$$

where ΔF is the instantaneous contrail RF. EF_{contrail} is almost equal to $AGWP_H$ (but in a different unit) for time horizons greater than 1 year (see Supplementary Data of Irvine et al., 2014). Such time horizons are much longer than the lifetime of a contrail, therefore $AGWP_H$ integrates the whole contrail radiative forcing. However, the equality is not exact because of the existence of slow carbon-cycle feedbacks within the Earth system, and $AGWP_H$ is about 7, 12 and 21% greater than EF_{contrail} for a time horizon of 20, 50 and 100 years, respectively.

This study considers all segments of flights that operated over the North Atlantic in the Shanwick and Gander Oceanic Control Areas, which are approximatively bounded by longitudes 50°W and 10°W and latitude 40°N and 75°N. We use the data from the year 2019, before the disruption caused to intercontinental routes by the Covid-19 pandemic. That represents a total of 477,923 flights.

The dataset provides fuel consumption and EF_{contrail} , which were calculated by Teoh et al. (2022) using: (i) actual flight trajectories provided by the UK National Air Traffic Services (NATS); (ii) the European Centre for Medium Range Weather Forecasts (ECMWF) ERA5 high-resolution realization (HRES) reanalysis, which, for the purposes of this study, is presumed to be completely accurate; and (iii) the contrail cirrus prediction model CoCiP (Schumann et al., 2012). CoCiP tests the meteorological conditions over flight segments and if they are favourable to contrail formation, a contrail segment is formed. CoCiP then advects the contrail in a Lagrangian framework to simulate coverage and radiative properties over the contrail lifetime. In this study, EF_{contrail} is expressed in terms of energy forcing per flown distance, dividing total energy forcing by total flown distance for each flight.

According to those calculations, 260,854 flights (55%) formed contrails and 208,965 flights (80% of contrail-forming flights) formed contrails that exerted a positive contrail energy forcing. Contrail energy forcing per flown distance ranges from -8.7×10^{11} to 5.0×10^{12} J km⁻¹, with a mean of 9.6×10^{10} J km⁻¹ and a median of 2.9×10^{10} J km⁻¹. The 10th and 90th percentiles are 6.0×10^9 and 2.8×10^{11} J km⁻¹, respectively. It should be noted that the uncertainties associated with those energy forcing estimates are large. Uncertainties were estimated by Teoh et al. (2022) but are left out of the present analysis.



160 Contrail energy forcing and emitted CO₂ are enough to calculate the radiative forcing of each flight but are not directly
comparable quantities. So this study uses the OSCAR reduced-complexity Earth system model (Gasser et al., 2017a; 2020)
to compute for each individual flight, with a 1-year timestep, the time evolution of the globally averaged radiative forcing
and the globally averaged surface temperature change that occurs in response to that forcing. For CO₂, OSCAR computes the
time evolution of radiative forcing from the provided flight CO₂ emissions using a multi-region box model. For contrails,
165 energy forcing, in joules, is added to the climate system uniformly over 1 year, which is the time step of the model. That is
done without geographical considerations. OSCAR calculates the climate response to radiative forcing by using an impulse
response function. The carbon cycle response to a pulse CO₂ emission depends on the background concentration of CO₂.
That background concentration is simulated using the emission scenario SSP4-3.4 in our experiment.
Uncertainties in radiative forcing and surface temperature change calculated by OSCAR are assessed using a Monte Carlo
170 approach based on 2000 simulations with different model parameters. Then, the 1726 simulations with simulated historical
CO₂ concentrations closest to observations provide the best estimate and standard deviation.
CoCiP and OSCAR do not simulate the tropospheric and stratospheric adjustments triggered by the contrail climate forcing.
A contrail efficacy factor is used here for contrails and contrail cirrus to account for those adjustments: the contrail radiative
forcing calculated by Teoh et al. (2022) is thus multiplied by contrail efficacy before being used in OSCAR. Contrail
175 efficacy is divided in two parts: the ERF-to-RF ratio and the “Ponater” efficacy (Hansen et al., 2005, Ponater et al, 2021).
The ERF-to-RF ratio quantifies the atmospheric adjustments that follow contrail formation, i.e. changes in stratospheric and
tropospheric temperature, humidity, or cloudiness that happen without large-scale changes in surface temperature (Sherwood
et al. 2015). The Ponater efficacy quantifies the ability of a unit of contrail ERF to change surface temperatures compared to
a unit of CO₂ ERF. It can differ from 1 because different ERF patterns trigger different surface temperature responses (e.g.
180 Shindell and Faluvegi, 2009). Total contrail efficacy is the product of these two efficacies. It is set to 0.35 in this study,
which is the best estimate of ERF-to-RF obtained in the contrail-climate model of Bickel et al. (2020), multiplied by 1 for
the Ponater efficacy, for lack of a better best estimate, and the same factor is assumed to apply to all contrails. Contrail
efficacy is taken into account in all the results in this study, in particular when using the AGWP. The sensitivity of the
climate outcome of contrail avoidance to the uncertainty in contrail efficacy is investigated in Section 5.3.

185 **4. CO₂ and contrail time scales**

The climate perturbations of CO₂ and contrails are associated with very different time scales. These time scales are
illustrated by Figure 1, which shows the radiative forcing and surface temperature response caused by a typical contrail-
forming flight flying 2450 km in the Shanwick and Gander sectors of the North Atlantic. This typical segment of flight burns
17 tons of kerosene and emits a total of 53 tons of CO₂ into the atmosphere. We assume in addition that the flight is
190 responsible for forming a contrail cirrus with an energy per flown distance of 2.9×10^{10} J km⁻¹, which is the median value for
contrail-forming flights in the dataset.



The emission of CO₂ molecules from kerosene combustion is instantaneous, while the formation of the ice crystals that form persistent contrails and contrail cirrus takes a few seconds (Kärcher, 2018). Both the CO₂ molecules or ice crystals will exert a radiative forcing for as long as their respective atmospheric concentrations remain perturbed. The radiative forcing exerted by the contrail ice crystals, which is assumed in Figure 1 to be positive, lasts up to 10-15 hours (Kärcher, 2018), except for very small carbon cycle adjustments that develop over a few decades (Gasser et al., 2017b). In stark contrast, the emission of fossil CO₂ causes an initial increase in the CO₂ concentration in the atmosphere, and about 30% of this increase persists after 100 years, and even about 20% persists after 1000 years, thus exerting a radiative forcing long in the future (Archer et al., 2009; Joos et al., 2013). Note however that although the radiative forcing of contrail operates on a time scale orders of magnitude shorter than that of CO₂, it is also orders of magnitude stronger.

The stark differences in CO₂ and contrail radiative forcing time scales do not fully propagate to the temperature response (bottom row of Figure 1) because the ocean absorbs the energy perturbation resulting from the radiative forcing, increasing its heat content, before returning that heat to the atmosphere over several decades (Stjern et al., 2023). CO₂ is associated with both short and long timescales because its radiative forcing is exerted over many years, effectively providing a slowly decreasing source of energy for the ocean to absorb and then release. The warming of contrails is felt by the climate system through the ocean response over several decades despite contrail cirrus only lasting for several hours. This long-term response is further amplified by so-called carbon cycle feedback, which causes the small increase in contrail radiative forcing that can be seen in Figure 1 a few years after the contrail has dissipated. That feedback occurs because warming leads to a release of CO₂, primarily through increased decomposition of soil organic matter, which in turn leads to more warming.





Figure 1: Time evolution of (top row) globally averaged effective radiative forcing, in pW m^{-2} ($10^{-12} \text{ W m}^{-2}$), and (bottom row) global surface temperature change, in pK (10^{-12} K), for a flight that emits 53 tons of CO₂ (grey) in the atmosphere and forms a



215 **contrail (red) with an energy forcing of $2.9 \times 10^{10} \text{ J km}^{-1}$ over the whole 2450 km of its journey. Contrail efficacy is set to 0.35. Background CO_2 concentrations follow the SSP4-3.4 scenario.**

This discussion of time scales already implies that the behaviour of CO_2 -equivalence metrics will depend on whether they are based on radiative forcing or temperature response, and whether they are integrated or endpoint metrics. Figure 2
220 compares the multiplicative factors associated with the AGWP, AGTP, and ATR using three time horizons, 20, 50, and 100 years, for the same flight presented in Figure 1. Multiplicative factors are here presented relative to the emitted CO_2 , which is therefore given a value of 1. The values shown in Figure 2 are about half of those in Table 5 of Lee et al. (2021) because this study assumes a contrail efficacy of 0.35 compared to 0.42 in Lee et al. (2021), but mostly because Lee et al. (2021) computes the mean over all flights which gives an energy forcing of $3.6 \times 10^{10} \text{ J km}^{-1}$, while this study uses the median over
225 contrail-forming flights which is equal to $2.9 \times 10^{10} \text{ J km}^{-1}$. The distribution is strongly skewed by a few flights that form very strongly warming contrails, therefore the mean radiative forcing is shifted toward higher values than the median. Figure 2 includes the uncertainty in the climate response obtained using the Monte Carlo approach described above, which does not account for the uncertainties in contrail ERF and its efficacy, which are both large. According to the 5-95% confidence level in contrail cirrus ERF given in Lee et al. (2021), the contrail multiplicative factors could be between one-third and double the
230 length shown in Figure 2.

The largest CO_2 -equivalence metrics are obtained for a 20-year time horizon, regardless of the metric definition, and CO_2 -equivalence tends toward 1 when time horizon increases. From a radiative forcing point of view, this is because the AGWP of contrails only slightly increases once the contrail has dissipated due to the carbon cycle feedback, while the AGWP of CO_2 keeps increasing with increasing time horizon. From a temperature response point of view, this is because the AGTP of
235 contrails decays by two orders of magnitude after about 50 years. The AGWP20 and ATR20 of contrail cirrus are therefore large, ranging from 1.3 to 1.7 times the respective metric for CO_2 . If these larger values were used in a practical setting, they would indicate a larger perceived benefit of contrail avoidance. Overall, AGWP and ATR yield greater multiplicative factor than AGTP, which places emphasis on the longer-term warming (Figure 1).



240 **Figure 2: CO₂-equivalence of a median North Atlantic contrail with a contrail efficacy of 0.35 (red) compared to the emitted CO₂ during the median flight (grey) when using Absolute Global Warming Potential (AGWP), Absolute Global Temperature Change Potential (AGTP), and Average Temperature Response (ATR) with time horizons of 20, 50, and 100 years. Error bars quantify the one-sigma uncertainty arising from the physical climate and carbon cycle of OSCAR, rather than uncertainties in the contrail radiative forcing and efficacy. The values at the top of each bar are the ratio of non-CO₂ to CO₂ effects for each metric choice.**

245


5. Contrail avoidance in the North Atlantic

The OSCAR model is now run for each of the 477,923 flights in 2019 considered by Teoh et al. (2022), including 260,854 contrail-forming flights. Overall, we estimate that the CO₂ emitted and the contrails formed by those flights will have warmed the climate by a total of 16.9 μK in 2039, 13.5 μK in 2069, and 14.0 μK in 2119 (third panel of Figure 3a. Figure 3b shows that flights that form persistent warming contrails, which represent 44% of the database, are responsible for 70%, 72% and 54% of warming from both CO₂ and contrails according to AGWP100, ATR100 and AGTP100. For AGWP20, ATR20,

250



and AGTP20, contributions increase to 86%, 90% and 63%, respectively. Total cooling from flights where contrail cooling exceeds CO₂ warming offsets at most 2% of total warming.



255 **Figure 3: Cumulative histogram of the warming or cooling by emitted CO₂ and persistent contrails formed by the 477,923 flights that crossed the North Atlantic sector in 2019 (Teoh et al., 2022) for nine CO₂-equivalence metrics. The first row (a) shows the decomposition into forcers: warming contrails (red), CO₂ (grey), and cooling contrails (blue). The second row (b) shows the decomposition into different categories of flights: warming flights that formed cooling contrails (green), warming flights that formed warming contrails (dark red), warming flights that did not form persistent contrails (yellow), and cooling flights (blue), including the effects of CO₂ for those flights. The corresponding numbers are presented in Tables S1 and S2 of the Supplementary Materials.**


260

OSCAR is used again to simulate the hypothetical rerouted flights. For each contrail-forming flight, we assume that it is possible to reroute the flight to avoid forming the persistent contrail. In practice, the increase in flight time and distance
 265 would depend on the size of the ISSR in which the contrail is formed and whether it is avoided horizontally or vertically. But what matters here is the additional CO₂ emitted by the rerouted flight. We set that additional CO₂ to 1% of the original flight emissions, which is in the range of increased fuel burn found in the case studies of Zengerling et al. (2023). In addition, fuel consumption and flight delays are both associated with large costs to airlines, so they might be unlikely to commit to larger increases. The additional CO₂ due to the rerouting would in practice depend also on the initial trajectory flown, ambient
 270 wind fields, and aircraft characteristics (Irvine et al., 2014). Rerouting is assumed to have been successful with persistent contrail formation being completely avoided, although it should be noted that this assumption is overly optimistic and cannot be currently guaranteed in any realistic setting.

Figure 4 shows the climate outcome of the rerouting, calculated here as the difference in AGTP100 between the rerouted and original flight, as a function of the contrail energy forcing of the original flight. AGTP100 is taken here as the reference



275 because it is by definition equal to the temperature change after 100 years, so is easy to interpret physically. Rerouting, done
within the assumptions above, leads to climate benefit for 68% of the contrail-forming flights (bottom right quadrant), for a
total decrease in warming of $1.9 \mu\text{K}$ (-14% compared to total warming from all original flights). Rerouting the remaining
32% of the flights would damage climate, for a total increase in warming of $0.1 \mu\text{K}$ ($+0.7\%$ compared to total warming from
all original flights). For those flights, the temperature change contribution by the rerouted flight is 5% larger than that of the
280 original flight, either because the original flight was cooling (top left quadrant) or because the contrail formed had too weak
positive radiative forcing to be worth the additional CO_2 under the 1% additional CO_2 assumption (top right quadrant).
Illustrative case studies to clarify the results shown in Figure 4 are presented in Supplementary Materials.
It could be of interest to focus on flights in the bottom right corner of Figure 4 because their energy forcing might be
sufficiently large to indicate with high likelihood that they do indeed warm the climate despite uncertainties in contrail EF.
285 Such flights might be good candidates for “lower risk” contrail avoidance provided ISSR forecasts are reliable. That subset
of flights can be selected in various ways. For example, selecting flights with an energy forcing per contrail length larger
than $10^{11} \text{ J km}^{-1}$ and whose climate benefit from contrail avoidance is 100 times larger than the climate damage from
emitting 1% more CO_2 leaves 8.5% of contrail-forming flights. They contribute $1.34 \mu\text{K}$ to the initial AGTP100, which is
reduced to $0.49 \mu\text{K}$ after rerouting. In other words, those 8.5% of flights represent 55% of the potential climate benefit of
290 contrail avoidance. However, Figure 5 shows that the number of flights that match the selection criteria is a strong function
of the choice of the CO_2 -equivalence metric. There are many other options, which set the level of ambition and complexity
of the rerouting, but also the climate benefit, which will be dependent on the CO_2 -equivalence metric used.



295 **Figure 4: Difference in AGTP100 between rerouted and original flights, in pK (10^{-12} K), as a function of the energy forcing of the original flight, in J km^{-1} . Flights are all the contrail-forming flights that flew over the North Atlantic in 2019. Colours indicate the number of flights for each combination, for a total of 260,854 contrail-forming flights. The rerouted flights emit 1% more CO_2 but do not form a contrail. Contrail efficacy is set to 0.35. The same figure expressed in other CO_2 -equivalence metrics are in the Supplementary Materials.**

300

5.1 Sensitivity to additional CO_2

So far, the rerouting scenario assumed that contrails could be fully avoided by consuming 1% more fuel. Additional fuel consumption is now set to 0, 2 and 5%, to cover the range of estimated additional fuel burn from the case studies of Zengerling et al. (2023). Results are shown in Figure 5 for the nine CO_2 -equivalence metrics. Emitting more CO_2 forces more flights into the climate damage category. For AGTP100, the effect is sizeable in terms of number of flights: at 5% additional CO_2 , rerouting damages climate for 43% of contrail-forming flights, 11 percentage points more than the assumption with 1% additional CO_2 (32%). In the scenario where no additional CO_2 is emitted, the number of reroutings that damage climate is decreased by 12 percentage points compared to the 1% additional CO_2 scenario, falling to 20% of contrail-forming flights. These flights damage the climate because a cooling contrail is avoided. The number of “lower risk” reroutings is 550% larger. Choosing a shorter time horizon enhances the perceived climate benefit but underestimates the

305


310



long-term climate damage. Using AGTP20 and the 1% additional CO₂ scenario, 11% fewer flights fall in the climate damage category than with AGTP100.

For a given CO₂ scenario and for most contrail-forming flights, all CO₂-equivalence metrics agree that rerouting would benefit climate. Disagreements between CO₂-equivalence metrics happen for about 10% of flights, which form low energy 315 contrails that do not contribute much to climate damage anyway. The range of contrail EF where at least one CO₂-equivalence metric gives a different climate outcome of rerouting than the other metrics depends on the flight and CO₂ scenario. For the median flight with 1% additional CO₂, that range spans 7% of the contrail-forming flights, from the 24th to the 31st percentiles. This range represents very low-energy contrails, with EF per flown kilometre from 1.7×10⁸ to 3.1×10⁹ J km⁻¹.

320



325 **Figure 5: (First row) Number of flights flown over the North Atlantic in 2019, for a total of 260,854 contrail-forming flights, for which rerouting leads to climate benefit (colours) or climate damage (white) according to nine CO₂-equivalence metrics and depending on the additional CO₂ emitted to reroute. Colours indicate the amount of additional CO₂ emitted to reroute: +0% (blue), +1% (orange), +2% (green) or +5% (red). Rerouting is assumed to be successful, so the rerouted flight does not form a contrail, and a contrail would have formed on the original route. Lower risk reroutings (see text) are marked with hatches. (Second row) As first row but showing the climate benefit from rerouting all contrail-forming flights. For ATR, the unit is denoted μK to distinguish an average over the time horizon from an endpoint temperature change. The corresponding numbers are presented in Table S3 of the Supplementary Materials.**

330

5.2 Imperfect contrail avoidance

The results presented in Figure 5 are idealised because the additional CO₂ is independent of the distance over which a contrail is formed. It is plausible that 1% additional CO₂, and indeed even 5%, would not allow the complete avoidance of



335 some contrails. Determining the best avoidance strategy for a given flight requires accurate forecasts of meteorological conditions for that flight, which is out of the scope of this study. There is no simple relationship between additional CO₂ and contrail EF: ISSRs could be orthogonal to the plane trajectory so difficult to avoid, or conversely vertical avoidance might be very easy, irrespective of contrail EF. There might also be cases where two different routes have very different ISSR occurrence but identical fuel burn.

340 But to illustrate the impact of imperfect contrail avoidance, we assume here that the longer the contrail, the more difficult it is to completely avoid. A rerouting efficiency factor is therefore calculated as the ratio of the contrail length to total distance flown for the original flight. So, rerouting efficiency is set to 0 for a contrail that is as long as the flight length, meaning that the rerouting fails to reduce the energy forcing of the contrail. Rerouting efficiency will tend toward 1 for contrails that are very small compared to the length of the flight, meaning that the contrail is fully avoided. For the contrail-forming flights flown over the North Atlantic in 2019, the average rerouting efficiency is 0.71, and the 5th and 95th percentiles are 0.31 and 345 0.95, respectively.

Table 1 shows the number of reroutings beneficial to climate and the absolute benefit of all reroutings (both beneficial and detrimental to climate) as a function of the CO₂-equivalence metric. The total decrease in warming in 100 years (i.e., using AGTP100) is 0.97 μK, a reduction of 46% compared with assuming a rerouting efficiency of 1 for all flights, where total warming was decreased by 1.79 μK. That reduction is larger than the reduction in average rerouting efficiency, which is only 350 29% (from 1 to 0.71), because high-energy contrails are often longer than low-energy ones. Therefore, their rerouting efficiency is lower, although they contribute more to the absolute benefit of the rerouting. In contrast, the number of reroutings leading to climate benefit in terms of temperature rise in 100 years decreases by only 3%. That is because reducing the climate impact of strongly warming contrails leads to a climate benefit even when that reduction is incomplete. The relative decrease in climate benefit depends only slightly on the CO₂-equivalence metric. The number of reroutings that 355 benefit climate does not change much in this imperfect contrail avoidance calculation, ranging from -1% to -3%. This indicates that the previous finding that qualitative decision making is almost independent on the choice of CO₂-equivalence metric remains valid. Moreover, the absolute benefit is reduced by around 45% for all CO₂-equivalence metrics. As we showed earlier in this Section, this is because a large part of the absolute benefit comes from flights for which the impact of the additional emitted CO₂ is much greater than the impact of the avoided contrail. For these flights, CO₂ is of secondary 360 importance, therefore the CO₂-equivalence metrics mostly quantify the climate outcome of contrail avoidance, which is reduced in an imperfect avoidance scenario.

Table 1: Number of flights flown over the North Atlantic in 2019, out of a total of 260,854 contrail-forming flights, whose rerouting is beneficial to climate, and absolute benefit of all reroutings (beneficial and detrimental to climate), assuming that contrail formation is fully or imperfectly avoided (see text), as a function of CO₂-equivalence metric. The unit of absolute benefit depends on the CO₂-equivalence metric and is denoted <μK> for ATR to highlight that it is an average temperature change over the time horizon.



| CO ₂ -equivalence metric | Number of flight reroutings beneficial to climate | | Absolute climate benefit of all reroutings | |
|-------------------------------------|---|---------------------|--|-----------------------------------|
| | Full avoidance | Imperfect avoidance | Full avoidance | Imperfect avoidance |
| AGWP100 | 191,119 | 188,372 (-1%) | 1.52 mW m ⁻² yr | 0.83 mW m ⁻² yr (-45%) |
| AGWP50 | 193,989 | 191,580 (-1%) | 1.49 mW m ⁻² yr | 0.82 mW m ⁻² yr (-45%) |
| AGWP20 | 196,703 | 194,608 (-1%) | 1.46 mW m ⁻² yr | 0.80 mW m ⁻² yr (-45%) |
| AGTP100 | 178,502 | 173,623 (-3%) | 1.79 μK | 0.97 μK (-46%) |
| AGTP50 | 182,755 | 178,558 (-2%) | 2.41 μK | 1.30 μK (-46%) |
| AGTP20 | 187,445 | 184,130 (-2%) | 4.61 μK | 2.51 μK (-46%) |
| ATR100 | 191,936 | 189,309 (-1%) | 8.76 <μK> | 4.80 <μK> (-45%) |
| ATR50 | 194,780 | 192,395 (-1%) | 15.5 <μK> | 8.49 <μK> (-45%) |
| ATR20 | 197,694 | 195,631 (-1%) | 34.2 <μK> | 18.8 <μK> (-45%) |

370 5.3 Sensitivity to contrail efficacy

Contrail efficacy has been set so far to the best estimate of 0.35 from Bickel et al. (2020). To study the sensitivity of our results to that choice, contrail efficacy is varied from 1 to 0.05 by considering five values:

- 0.23 and 0.51, which are the limits of 5-95% confidence interval of Bickel et al. (2020).
- 1.0, which is not supported by climate modelling but corresponds to a focus on contrail radiative forcing, rather than effective radiative forcing.
- 0.05 and 0.1, which are very low efficacies that might be possible if the lower end of the Bickel et al. (2020) confidence interval was multiplied by a Ponater efficacy of 0.38 (Bickel, 2023).

Figure 6a shows that the number of reroutings beneficial to climate does not change drastically between different choices of contrail efficacy. For contrail efficacies between 0.05 to 1, the maximum difference is 25% for AGTP100 for a contrail efficacy of 0.05. This implies that contrail efficacy plays a minor role in the decision making of a rerouting: in the worst case, 75% of the reroutings that benefit climate remain beneficial. This again comes from the fact that a warming contrail is, most of the time, a few orders of magnitudes more warming than the additional CO₂ emitted to avoid it.

In contrast, the perceived climate benefit associated to a rerouting depends highly on contrail efficacy. Figure 6b shows that the impact of contrail efficacy on the quantification of the climate benefit of reroutings is much larger. Climate benefit is 3 times larger for a contrail efficacy of 1 than for an efficacy of 0.35, and 10 times smaller for a contrail efficacy of 0.05. It should be noted that this impact does not depend much on the CO₂-equivalence metric, because the climate benefit of very high-energy contrails is much greater than the climate damage by the additional emitted CO₂. Therefore, changing the contrail efficacy by a given factor changes the absolute climate benefit by about the same factor, for all the considered metrics. However, this is only true when considering rerouted flight together, not on a flight-to-flight basis.



390




Figure 6: Relative number of reroutings that benefit climate (a) and relative climate benefit of all reroutings (b), for the nine CO₂-equivalence metrics considered in this study (colours and symbols) and as a function of contrail efficacy. Additional CO₂ emitted is 1%, and contrail avoidance is assumed to be complete. Values are normalised to the case where contrail efficacy is 0.35.

395

6. Conclusion

The different time scales of CO₂ and contrail cirrus radiative forcing and temperature response cause different behaviour of CO₂-equivalent metrics. Time-integrated metrics defined on a short time horizon, like AGWP20 or ATR20, put more weight on contrail cirrus, while endpoint metrics on a long time horizon, like AGTP100, put more weight on CO₂. This behaviour, which had already been noted by Grewe et al. (2014) and others, means that the choice of metric definition and time horizon can have implications for decision making related to contrail avoidance. Using an analysis of potential contrail avoidance strategies in the North Atlantic flight corridor, this study shows that the decision whether to reroute a flight or not is generally independent of the choice of CO₂-equivalent metric. This is because when a persistent warming contrail is formed,

400



it is often orders of magnitude more warming than the potential additional emission of CO₂ to avoid its formation. In many
405 of these situations, all the investigated metrics indicate that the flight should be rerouted, although the weight put on contrails
is not the same.

The lack of consensus on what is a suitable or the correct CO₂-equivalence metric is therefore not an obstacle to
implementing contrail avoidance policies. However, the quantification of how much additional CO₂ can be emitted on a
rerouted flight and the resulting climate outcome of a rerouting both depend on that choice. For a flight for which all metrics
410 agree that a rerouting would benefit climate, time-integrated metrics with short time horizons, like AGWP20 or ATR20, give
a much greater climate benefit than endpoint metrics with long time horizons like AGTP100, which in many situations
indicates that the benefit is moderate. Assuming that additional CO₂ emissions are fixed percentages of original emissions, or
an increasing function of contrail length does not change these findings. The same conclusion is also drawn when
considering sensitivity to contrail efficacy: the number of reroutings that benefit climate is a weak function of contrail
415 efficacy, but the size of the climate benefit is very dependent on contrail efficacy.

This study used the contrail energy forcing distribution from Teoh et al. (2022) computed with CoCiP (Schumann et al.,
2012) as radiative forcing input to the OSCAR compact Earth system model (Gasser et al., 2020). Rerouting was represented
in a simplified way because the knowledge of flight trajectories and meteorology has been lost into the contrail energy
forcing distribution. Future studies should investigate more realistic cases, considering the meteorology and particularly the
420 horizontal and vertical sizes and humidity profiles of ISSRs. However, our work shows that even a partially avoided contrail
can be beneficial for the climate if the contrail is sufficiently warming (e.g., with a contrail energy forcing per contrail length
greater than 10¹¹ J km⁻¹), and that many flights have formed such contrails over the North Atlantic in 2019 (30% of all flights
for the latter condition). The contrail climate impact is often orders of magnitude greater than the impact of the additional
emitted CO₂, so a rerouting can still be effective if it leads to the formation of much weaker contrail. Moreover, this study
425 has assumed that contrail avoidance translates into unchanged or increased fuel burn and CO₂ emissions. In fact, current
flight trajectories are predominantly optimised to minimise total cost (including, but not limited to, fuel burn, airspace
charges, and staff costs), so there is a possibility that re-routing flights to minimise in addition the climate impact by
avoiding contrails could lead to a reduction in overall fuel burn for some flights.

This study focuses on the North Atlantic, so its results do not necessarily carry over to other regions which may have
430 different characteristics, for example in the diurnal variation of air traffic, the probability of encountering ISSRs and the
average altitude of flights, as well as the practicality and operational constraints of re-routing due to the volume of air traffic.
However, the large difference between contrail warming and that of the additional emitted CO₂ seen for the most warming
flights is likely a defining feature of such flights in any region, so the fundamental results are expected to remain valid.

Implementing a contrail avoidance scheme is likely to be complex because of the difficulties in predicting ISSRs (Gierens et
435 al., 2020), or the large uncertainties in calculating the radiative forcing of contrails (Wilhelm et al., 2021), or operational
challenges with air traffic management. Sausen et al. (2023) found in a case study of contrail avoidance over the Netherlands
that persistent contrails were not observed in about half the cases when they were forecast, which indicates the risk of



unnecessary diversions. Lee et al. (2021) found a 5-95% confidence level for contrail cirrus radiative forcing of 33 to 189 mW m⁻². By contrast, the CO₂ forcing is uncertain by only ±10% (31 – 38 mW m⁻²). This study considered uncertainties originating from the OSCAR climate model, which are comprised within a 10% range, but did not account for uncertainties in contrail radiative forcing or ISSR prediction in the ERA5 reanalysis. On a flight-by-flight basis, rerouting would need to use operational weather forecasts rather than a reanalysis. However, such forecasts are known to have a greater uncertainty. So, the jury is still out as to whether contrail avoidance has in practice the potential to substantially reduce the total radiative forcing of aviation, and whether that reduction can be validated. One way to avoid dealing directly with uncertainty could be to target the subset of strongly warming contrails whose energy forcing is sufficiently strong to remain positive despite large contrail uncertainties, assuming these can be adequately forecast. But our results suggest that the number of flights selected in that way strongly depends on the choice of CO₂-equivalence metrics. A cautious approach to select such flights could be to use AGTP100 to maximise the probability that the rerouting does indeed benefit climate. The next step is therefore to link the present analysis to estimates of contrail radiative forcing uncertainty and to use real cases of contrail avoidance reroutings to quantify the potential contrail energy forcing reduction with the corresponding additional emitted CO₂.

Which CO₂-equivalence metric works best for contrail avoidance? This study offers some insights. In terms of choice of time horizon, rerouted flights can be perceived to be a climate benefit using short time horizons. However, an increasing proportion of these become climate damaging as the time horizon is increased, as shown in Figure 7. Indeed, it has been noted that the use of GWP20 is equivalent to use a discount rate of more than 10% (Sarofim and Giordano, 2018; Mallapragada and Mignone, 2020), which is much higher than the discount rate of 4 to 5% typically assumed for designing cost-effective mitigation scenarios using Integrated Assessment Models (Emmerling et al. 2019). Furthermore, it can be argued that aviation would have to be at an advanced stage of decarbonisation, at least in some regions, to justify using a shorter time horizon than the 100 years used for surface emissions from other economic sectors. In terms of choice of metric definition, AGWP and ATR, which both involve a time integration over the time horizon, behave very similarly, both qualitatively and quantitatively. AGWP has the advantage over ATR of being less inflicted with uncertainties, whereas AGWP is less comprehensive than ATR in terms of the consideration of climate impacts. AGTP behaves differently from the other two metrics, and generally predicts less positive climate outcomes of contrail avoidance. But it has the advantage of measuring surface temperature change, which is directly relevant to the warming target of the Paris Agreement.

Until such time that a political decision has been made, any proposed contrail avoidance schemes should, in addition to considering the many scientific uncertainties inherent in such schemes, consider a range of metrics and time horizons, such as those used in this study, to assess the robustness of rerouting decisions and to quantify the actual climate benefit. Such a recommendation has already been made (Levasseur et al., 2016; Cherubini et al., 2016; Jolliet et al., 2018) and applied (e.g., Tanaka et al., 2019; Reisinger et al., 2017; Tibrewal et al., 2020) by the Life Cycle Impact Assessment community, which uses by consensus GWP100, GWP20, GTP100, and GTP20 in their assessments.

470

Data availability

CO₂ emission and contrail energy forcing data were made available from Roger Teoh (roger.teoh15@imperial.ac.uk; Teoh et al., 2020). The source code of OSCAR is available at <https://github.com/tgasser/OSCAR>. Additional scripts and data are available upon request from the corresponding author.

475

Author contribution

AB and NB conceptualised and conducted the study, and prepared the manuscript. All the authors discussed and commented on the manuscript.

480 Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

485 AB, OB, and NB acknowledge the support of the French Ministère de la Transition écologique et solidaire (N° DGAC 382 N2021-39), with support from France's Plan National de Relance et de Résilience (PNRR) and the European Union's NextGenerationEU. K.T. benefited from State assistance managed by the National Research Agency in France under the Programme d'Investissements d'Avenir under the reference ANR-19-MPGA-0008.

References

- 490 Allen, M., Fuglestedt, J., Shine, K. et al. New use of global warming potentials to compare cumulative and short-lived climate pollutants. *Nature Clim Change*, 6, 773–776. <https://doi.org/10.1038/nclimate2998>, 2016.
- Allen, M.R., Shine, K.P., Fuglestedt, J.S. et al. A solution to the misrepresentations of CO₂-equivalent emissions of short-lived climate pollutants under ambitious mitigation. *npj Clim Atmos Sci* 1, 16. <https://doi.org/10.1038/s41612-018-0026-8>, 2018.
- 495 Allen, M.R., Peters, G.P., Shine, K.P. et al. Indicate separate contributions of long-lived and short-lived greenhouse gases in emission targets. *npj Clim Atmos Sci* 5, 5. <https://doi.org/10.1038/s41612-021-00226-2>, 2022.
- Archer, David, Michael Eby, Victor Brovkin, Andy Ridgwell, Long Cao, Uwe Mikolajewicz, Ken Caldeira, Katsumi Matsumoto, Guy Munhoven, Alvaro Montenegro, Kathy Tokos. Atmospheric Lifetime of Fossil Fuel Carbon Dioxide. *Annual Review of Earth and Planetary Sciences* 37:1, 117-134, <https://doi.org/10.1146/annurev.earth.031208.100206>, 2009.
- 500 Bickel, M., M. Ponater, L. Bock, U. Burkhardt, and S. Reineke: Estimating the Effective Radiative Forcing of Contrail Cirrus. *J. Climate*, 33, 1991–2005, <https://doi.org/10.1175/JCLI-D-19-0467.1>, 2020.
- Bickel, Marius. Climate Impact of Contrail Cirrus. DLR-Forschungsbericht. DLR-FB-2023-14. Dissertation. Ludwig-Maximilians-Universität München. 133 S. <https://doi.org/10.57676/mzmg-r403>, 2023.



- Cherubini F., J. Fuglestedt, T. Gasser, A. Reisinger, O. Cavalett, M. A.J. Huijbregts, D. J.A. Johansson, S. V. Jørgensen, M. 505
Raugei, G. Schivley, A. H. Strømman, K. Tanaka, A. Levasseur Bridging the gap between impact assessment methods and
climate science, *Environmental Science & Policy*, Volume 64, 129-140, <https://doi.org/10.1016/j.envsci.2016.06.019>, 2016.
- Collins, W. J. (B.), Frame, D. J., Fuglestedt, J. and Shine, K. P. Stable climate metrics for emissions of short and long-lived
species – combining steps and pulses. *Environmental Research Letters*, 15 (2). 024018. <https://doi.org/10.1088/1748-9326/ab6039>, 2020.
- 510 Dhakal, S., J.C. Minx, F.L. Toth, A. Abdel-Aziz, M.J. Figueroa Meza, K. Hubacek, I.G.C. Jonckheere, Yong-Gun Kim,
G.F.Nemet, S. Pachauri, X.C. Tan, T. Wiedmann, 2022: Emissions Trends and Drivers. In IPCC, 2022: Climate Change
2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the
Intergovernmental Panel on Climate Change [P.R. Shukla, J. Skea, R. Slade, A. Al Khourdajie, R. van Diemen, D.
McCollum, M. Pathak, S. Some, P. Vyas, R. Fradera, M. Belkacemi, A. Hasija, G. Lisboa, S. Luz, J. Malley, (eds.)].
515 Cambridge University Press, Cambridge, UK and New York, NY, USA. <https://doi.org/10.1017/9781009157926.004>, 2022.
- EASA European Union Aviation Safety Agency, Updated analysis of the non-CO₂ climate impacts of aviation and potential
policy measures pursuant to the EU Emissions Trading System Directive Article 30(4), available at
<https://www.easa.europa.eu/en/document-library/research-reports/report-commission-european-parliament-and-council>,
accessed 17/10/2023, 192 pp, 2020.
- 520 Emmerling, Johannes, Laurent Drouet, Kaj-Ivar van der Wijst, Detlef van Vuuren, Valentina Bosetti and Massimo Tavoni.
The role of the discount rate for emission pathways and negative emissions. *Environmental Research Letters*, 14, 104008,
<https://doi.org/10.1088/1748-9326/ab3cc9>, 2019.
- European Commission, Directive 2003/87/EC of the European Parliament and of the Council of 13 October 2003
establishing a scheme for greenhouse gas emission allowance trading within the Community and amending Council
525 Directive 96/61/EC, available at <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex%3A32003L0087>, accessed
17/10/2023, 2003.
- FAO. *Methane emissions in livestock and rice systems – Sources, quantification, mitigation and metrics*.
Rome. <https://doi.org/10.4060/cc7607en>, 2023.
- Forster, P., T. Storelvmo, K. Armour, W. Collins, J.-L. Dufresne, D. Frame, D.J. Lunt, T. Mauritsen, M.D. Palmer, M.
530 Watanabe, M. Wild, and H. Zhang, 2021: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity. In
Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the
Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N.
Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield,
O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY,
535 USA, pp. 923–1054, <https://doi.org/10.1017/9781009157896.009>, 2021.
- Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J., Landschützer, P., Le Quéré, C.,
Luijckx, I. T., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B.,
Alin, S. R., Anthoni, P., Barbero, L., Bates, N. R., Becker, M., Bellouin, N., Decharme, B., Bopp, L., Brasika, I. B. M.,
Cadule, P., Chamberlain, M. A., Chandra, N., Chau, T.-T.-T., Chevallier, F., Chini, L. P., Cronin, M., Dou, X., Enyo, K.,



- 540 Evans, W., Falk, S., Feely, R. A., Feng, L., Ford, D. J., Gasser, T., Ghattas, J., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Hefner, M., Heinke, J., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Joos, F., Kato, E., Keeling, R. F., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Lan, X., Lefèvre, N., Li, H., Liu, J., Liu, Z., Ma, L., Marland, G., Mayot, N., McGuire, P. C., McKinley, G. A., Meyer, G., Morgan, E. J., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O'Brien, K. M., Olsen, A., Omar, A.
- 545 M., Ono, T., Paulsen, M., Pierrot, D., Pocock, K., Poulter, B., Powis, C. M., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Séférian, R., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Sun, Q., Sutton, A. J., Sweeney, C., Takao, S., Tans, P. P., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., van Ooijen, E., Wanninkhof, R., Watanabe, M., Wimart-Rousseau, C., Yang, D., Yang, X., Yuan, W., Yue, X., Zaehle, S., Zeng, J., and Zheng, B.: Global Carbon Budget 2023, *Earth Syst. Sci. Data*, 15, 5301–5369, [https://doi.org/10.5194/essd-15-5301-](https://doi.org/10.5194/essd-15-5301-2023)
- 550 2023, 2023.
- Fuglestedt, J.S., Berntsen, T.K., Godal, O., Sausen, R., Shine, K.P., Skodvin, T. Metrics of Climate Change: Assessing Radiative Forcing and Emission Indices. *Climatic Change* 58, 267–33, <https://doi.org/10.1023/A:1023905326842>, 2003.
- Fuglestedt, J.S., K.P. Shine, T. Berntsen, J. Cook, D.S. Lee, A. Stenke, R.B. Skeie, G.J.M.M. Velders, I.A. Waitz. Transport impacts on atmosphere and climate: metrics. *Atmos. Environ.*, 44, pp. 4648-4677, <https://doi.org/10.1016/j.atmosenv.2009.04.044>, 2010.
- Gasser, T., Ciais, P., Boucher, O., Quilcaille, Y., Tortora, M., Bopp, L., and Hauglustaine, D.: The compact Earth system model OSCAR v2.2: description and first results, *Geosci. Model Dev.*, 10, 271–319, [https://doi.org/10.5194/gmd-10-271-](https://doi.org/10.5194/gmd-10-271-2017)
- 555 [2017](https://doi.org/10.5194/gmd-10-271-2017), 2017a.
- Gasser, T., Peters, G. P., Fuglestedt, J. S., Collins, W. J., Shindell, D. T., and Ciais, P.: Accounting for the climate–carbon
- 560 feedback in emission metrics, *Earth Syst. Dynam.*, 8, 235–253, <https://doi.org/10.5194/esd-8-235-2017>, 2017b.
- Gasser, T., Crepin, L., Quilcaille, Y., Houghton, R. A., Ciais, P., and Obersteiner, M.: Historical CO₂ emissions from land use and land cover change and their uncertainty, *Biogeosciences*, 17, 4075–4101, <https://doi.org/10.5194/bg-17-4075-2020>, 2020.
- Gierens, K., Schumann, U., Helten, M., Smit, H., and Marenco, A.: A distribution law for relative humidity in the upper
- 565 troposphere and lower stratosphere derived from three years of MOZAIC measurements, *Ann. Geophys.*, 17, 1218–1226, <https://doi.org/10.1007/s00585-999-1218-7>, 1999.
- Gierens, K.; Matthes, S.; Rohs, S. How well can persistent contrails be predicted? *Aerospace*, 7, 169. <https://doi.org/10.3390/aerospace7120169>, 2020.
- Gillett, Nathan P, and H Damon Matthews. Accounting for carbon cycle feedbacks in a comparison of the global warming
- 570 effects of greenhouse gases. *Environmental Research Letters*, 5, 034011, <https://doi.org/10.1088/1748-9326/5/3/034011>, 2010.
- Grewe, V., Frömming, C., Matthes, S., Brinkop, S., Ponater, M., Dietmüller, S., Jöckel, P., Garny, H., Tsati, E., Dahlmann, K., Søvde, O. A., Fuglestedt, J., Berntsen, T. K., Shine, K. P., Irvine, E. A., Champougny, T., and Hullah, P.: Aircraft routing with minimal climate impact: the REACT4C climate cost function modelling approach (V1.0), *Geosci. Model Dev.*,
- 575 7, 175–201, <https://doi.org/10.5194/gmd-7-175-2014>, 2014.



- Grewe, V., Gangoli Rao, A., Grönstedt, T. et al. Evaluating the climate impact of aviation emission scenarios towards the Paris agreement including COVID-19 effects. *Nat Commun* 12, 3841, <https://doi.org/10.1038/s41467-021-24091-y>, 2021.
- Hansen, J., M. Sato, R. Ruedy, L. Nazarenko, A. Lacis, G. A. Schmidt, G. Russell, I. Aleinov, M. Bauer, S. Bauer, N. Bell, B. Cairns, V. Canuto, M. Chandler, Y. Cheng, A. Del Genio, G. Faluvegi, E. Fleming, A. Friend, T. Hall, C. Jackman, M. Kelley, N. Kiang, D. Koch, J. Lean, J. Lerner, K. Lo, S. Menon, R. Miller, P. Minnis, T. Novakov, V. Oinas, Ja. Perlwitz, Ju. Perlwitz, D. Rind, A. Romanou, D. Shindell, P. Stone, S. Sun, N. Tausnev, D. Thresher, B. Wielicki, T. Wong, M. Yao, S. Zhang. Efficacy of climate forcings, *J. Geophys. Res.*, 110, D18104, <https://doi.org/10.1029/2005JD005776>, 2005.
- ICCT International Council on Clean Transportation (Brandon Graver, Kevin Zhang, and Dan Rutherford), CO₂ emissions from commercial aviation 2018, available at <https://theicct.org/publication/co2-emissions-from-commercial-aviation-2018/>, accessed 17/10/2023, 2018.
- Irvine, E A, B J Hoskins, and K P Shine. A simple framework for assessing the trade-off between the climate impact of aviation carbon dioxide emissions and contrails for a single flight. *Environmental Research Letters*, 9, 064021, <https://doi.org/10.1088/1748-9326/9/6/064021>, 2014.
- Jolliet, O., Antón, A., Boulay, AM. *et al.* Global guidance on environmental life cycle impact assessment indicators: impacts of climate change, fine particulate matter formation, water consumption and land use. *Int J Life Cycle Assess* 23, 2189–2207, <https://doi.org/10.1007/s11367-018-1443-y>, 2018.
- Joos, F., Roth, R., Fuglestedt, J. S., Peters, G. P., Enting, I. G., von Bloh, W., Brovkin, V., Burke, E. J., Eby, M., Edwards, N. R., Friedrich, T., Frölicher, T. L., Halloran, P. R., Holden, P. B., Jones, C., Kleinen, T., Mackenzie, F. T., Matsumoto, K., Meinshausen, M., Plattner, G.-K., Reisinger, A., Segschneider, J., Shaffer, G., Steinacher, M., Strassmann, K., Tanaka, K., Timmermann, A., and Weaver, A. J.: Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis, *Atmos. Chem. Phys.*, 13, 2793–2825, <https://doi.org/10.5194/acp-13-2793-2013>, 2013.
- Lee, D. S., Fahey, D. W., Skowron, A., Allen, M. R., Burkhardt, U., Chen, Q., Doherty, S. J., Freeman, S., Forster, P. M., Fuglestedt, J., Gettelman, A., De León, R. R., Lim, L. L., Lund, M. T., Millar, R. J., Owen, B., Penner, J. E., Pitari, G., Prather, M. J., Sausen, R., and Wilcox, L. J.: The contribution of global aviation to anthropogenic climate forcing for 2000 to 2018, *Atmos. Environ.*, 244, 117 834, <https://doi.org/10.1016/j.atmosenv.2020.117834>, 2021.
- Levasseur, A., et al. Enhancing life cycle impact assessment from climate science: Review of recent findings and recommendations for application to LCA. *Ecological Indicators*, 71, 163-174, <https://doi.org/10.1016/j.ecolind.2016.06.049>, 2016.
- Mallapragada, D.S., Mignone, B.K. A theoretical basis for the equivalence between physical and economic climate metrics and implications for the choice of Global Warming Potential time horizon. *Climatic Change* 158, 107–124. <https://doi.org/10.1007/s10584-019-02486-7>, 2020.



- Peters, Glen P, Borgar Aamaas, Terje Berntsen and Jan S Fuglestedt. The integrated global temperature change potential (iGTP) and relationships between emission metrics. *Environmental Research Letters*, 6 044021, 610 <https://doi.org/10.1088/1748-9326/6/4/044021>, 2011.
- Ponater, M., Bickel, M., Bock, L., Burkhardt, U. Towards Determining the Contrail Cirrus Efficacy. *Aerospace*, 8, 42, <https://doi.org/10.3390/aerospace8020042>, 2021.
- Reisinger A., S. F. Ledgard, S. J. Falconer Sensitivity of the carbon footprint of New Zealand milk to greenhouse gas metrics, *Ecological Indicators*, Volume 81, 74-82, <https://doi.org/10.1016/j.ecolind.2017.04.026>, 2017.
- 615 Sarofim, M. C. and Giordano, M. R.: A quantitative approach to evaluating the GWP timescale through implicit discount rates, *Earth Syst. Dynam.*, 9, 1013–1024, <https://doi.org/10.5194/esd-9-1013-2018>, 2018.
- Sausen, R. and Hofer, S. M. and Gierens, K. M. and Bugliaro Goggia, L. and Ehrmanntraut, R. and Sitova, I. and Walczak, K. and Burridge-Diesing, A. and Bowman, M. and Miller, N. Can we successfully avoid persistent contrails by small altitude adjustments of flights in the real world? *Meteorologische Zeitschrift.* <https://doi.org/10.1127/metz/2023/1157>, 2023.
- 620 Shine, K.P., R.G. Derwent, D.J. Wuebbles, J-J. Morcrette. Radiative Forcing of Climate. Chapter 2 of the First Assessment Report of the Intergovernmental Panel on Climate Change, 1990.
- Shine, K. P., J. S. Fuglestedt, K. Hailemariam and N. Stuber. Alternatives to the Global Warming Potential for Comparing Climate Impacts of Emissions of Greenhouse Gases. *Climatic Change* volume 68, pages 281–302, <https://doi.org/10.1007/s10584-005-1146-9>, 2005.
- 625 Shine, K.P. The global warming potential—the need for an interdisciplinary retrieval. *Climatic Change* 96, 467–472. <https://doi.org/10.1007/s10584-009-9647-6>, 2009.
- Stjern, C. W., et al. The time scales of climate responses to carbon dioxide and aerosols. *J. Climate*, 36, 3537–3551, <https://doi.org/10.1175/JCLI-D-22-0513.1>, 2023.
- Szopa, S., V. Naik, B. Adhikary, P. Artaxo, T. Berntsen, W.D. Collins, S. Fuzzi, L. Gallardo, A. Kiendler-Scharr, Z. Klimont, H. Liao, N. Unger, and P. Zanis Short-Lived Climate Forcers. In *Climate Change 2021: The Physical Science Basis*. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 817–922, 630 <https://doi.org/10.1017/9781009157896.008>, 2021.
- Tanaka, K., Peters, G.P. and Fuglestedt, J.S. Policy update: Multicomponent climate policy: why do emission metrics matter?, *Carbon Management*, 1:2, 191-197. <https://doi.org/10.4155/cmt.10.28>, 2010.
- Tanaka, K., Cavalett, O., Collins, W.J. et al. Asserting the climate benefits of the coal-to-gas shift across temporal and spatial scales. *Nat. Clim. Chang.* 9, 389–396. <https://doi.org/10.1038/s41558-019-0457-1>, 2019.
- 640 Tanaka, K. et al., Cost-effective implementation of the Paris Agreement using flexible greenhouse gas metrics. *Sci. Adv.* 7, eabf9020, <https://doi.org/10.1126/sciadv.abf9020>, 2021.



- Teoh, Roger, Ulrich Schumann, Arnab Majumdar, and Marc E. J. Stettler. Mitigating the Climate Forcing of Aircraft Contrails by Small-Scale Diversions and Technology Adoption. *Environmental Science & Technology*, 54 (5), 2941–2950, <https://doi.org/10.1021/acs.est.9b05608>, 2020.
- 645 Teoh, R., Schumann, U., Gryspeerdt, E., Shapiro, M., Molloy, J., Koudis, G., Voigt, C., and Stettler, M. E. J.: Aviation contrail climate effects in the North Atlantic from 2016 to 2021, *Atmos. Chem. Phys.*, 22, 10919–10935, <https://doi.org/10.5194/acp-22-10919-2022>, 2022.
- Teoh, R., Engberg, Z., Shapiro, M., Dray, L., and Stettler, M. A high-resolution Global Aviation emissions Inventory based on ADS-B (GAIA) for 2019–2021, *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2023-724>, 2023.
- 650 Tibrewal, K., Venkataraman, C. Climate co-benefits of air quality and clean energy policy in India. *Nat Sustain* 4, 305–313, <https://doi.org/10.1038/s41893-020-00666-3>, 2021.
- UNFCCC United Nations Framework Convention on Climate Change, Decision 4/CP.1 Methodological issues, Available at <https://unfccc.int/decisions?f%5B0%5D=session%3A3851>, Accessed 17/10/2023, 63 pp, 1995.
- UNFCCC United Nations Framework Convention on Climate Change, “Report of the Conference of the Parties serving as the meeting of the Parties to the Paris Agreement on the third part of its first session, held in Katowice from 2 to 15 December 2018. Addendum 2. Part two: Action taken by the Conference of the Parties serving as the meeting of the Parties to the Paris Agreement” (FCCC/PA/CMA/2018/3/Add.2 2019). Available at https://unfccc.int/sites/default/files/resource/cma2018_3_add2_new_advance.pdf, Accessed 12/12/2023, 65 pp, 2019.
- 655 Wang, Z., Bugliaro, L., Jurkat-Witschas, T., Heller, R., Burkhardt, U., Ziereis, H., Dekoutsidis, G., Wirth, M., Groß, S., Kirschler, S., Kaufmann, S., and Voigt, C.: Observations of microphysical properties and radiative effects of a contrail cirrus outbreak over the North Atlantic, *Atmos. Chem. Phys.*, 23, 1941–1961, <https://doi.org/10.5194/acp-23-1941-2023>, 2023.
- Wilhelm, L.; Gierens, K.; Rohs, S. Weather Variability Induced Uncertainty of Contrail Radiative Forcing. *Aerospace*, 8, 332. <https://doi.org/10.3390/aerospace8110332>, 2021.
- 665 Zengerling, Z.L.; Linke, F.; Weder, C. M.; Dietmüller, S.; Matthes, S.; Peter, P. Flying low and slow: Application of algorithmic climate change functions to assess the climate mitigation potential of reduced cruise altitudes and speeds on different days. *Meteorologische Zeitschrift*, <https://doi.org/10.1127/metz/2023/1194>, 2023.