

Cold pools mediate mesoscale adjustments of trade-cumulus fields to changes in cloud-droplet number concentration

Pouriya Alinaghi¹, Fredrik Jansson¹, Daniel A. Blázquez¹, and Franziska Glassmeier¹

¹Department of Geoscience & Remote Sensing, Delft University of Technology, Delft, The Netherlands

Correspondence: Pouriya Alinaghi (p.alinaghi@tudelft.nl)

Abstract. The mesoscale self-organization of trade-cumulus cloud fields is a major cloud-climate uncertainty. Cold pools, i.e. pockets of cold, dense air resulting from rain evaporation, are a key mechanism in shaping these dynamics and are controlled by the large-scale forcing. We study the microphysical sensitivity of cloud-field self-organization through cold pools by varying cloud-droplet number concentration N_c from 20 to 1000 /cm³ in large-eddy simulations on large 154×154 km²-domains.

5 We find that cold pools exhibit two distinct regimes of mesoscale self-organization. In very low N_c -conditions, cold pools transition from a stage where they are small and randomly distributed to forming large, long-lived structures that perpetuate due to the collisions of cold pools at their fronts. Under high- N_c conditions, cold pools display strongly intermittent behaviour and interact with clouds through small, short-lived structures. While N_c thus influences the number of cold pools and, in turn, mesoscale organization, cloud depth, and cloud albedo, we find its effect on cloud cover to be minimal. Comparing the
10 microphysical sensitivity of cold-pool-mediated mesoscale dynamics to the external, large-scale forcing shows that N_c is as important as horizontal wind and large-scale subsidence for trade-cumulus albedo. Our results highlight that cold pools mediate adjustments of trade-cumulus cloud fields to changes in N_c . Such mesoscale adjustments need to be considered if we are to better constrain the effective aerosol forcing and cloud feedback in the trade-wind regime.

1 Introduction

15 Clouds play a crucial role in the climate system by modulating the Earth's energy budget through their interactions with radiation. Their net effect is to cool the planet by reflecting incoming solar radiation back into space (Stephens et al., 2012). Clouds are one of the most important sources of uncertainty in climate projections. Firstly, the cloud feedback is the most uncertain feedback to the anthropogenic forcing of the climate system, which is mainly due to the uncertain response of shallow clouds to climate change (Schneider et al., 2017; Nuijens and Siebesma, 2019; IPCC AR6, 2023). Secondly, the
20 complex interactions between clouds and aerosols lead to the process uncertainty that makes the effective radiative forcing due to aerosol-cloud interactions the most uncertain forcing in the climate system (IPCC AR6, 2023; Bellouin et al., 2020).

Aerosol perturbations change the concentration of cloud-condensation nuclei and in turn cloud-droplet number concentrations N_c . Assuming a fixed cloud-liquid-water path, increased N_c results in a larger number of smaller cloud droplets, leading to a larger surface area to interact with radiation, and in turn increased cloud-optical depth and cloud albedo, known as [the](#)
25 *Twomey* effect (Twomey, 1977). In addition to this quasi-immediate effect that can be considered to occur on the spatiotempo-

ral scales of individual cloud parcels, changes in N_c can also propagate to larger scales. On the single-cloud scale, increased N_c reduces the efficiency of collision-coalescence processes through smaller radii, decreasing the rain-formation efficiency, thereby delaying precipitation formation, known as [the Albrecht or lifetime effect](#) (Albrecht, 1989). This delay in precipitation formation allows clouds to live longer and in turn get deeper, which in the end precipitate more intensely. Such effects can
30 lead to an internal re-organization on the cloud field, or mesoscale, [which ranges from tens to hundreds of kilometers and evolves on timescales of hours to days](#). They can be considered a form of self-organization because they are not prescribed by a large-scale forcing. In addition to delayed precipitation formation, increased N_c has also been described to affect entrainment rates with effects on meso-timescales (Glassmeier et al., 2021).

For trade-cumulus cloud fields, large-eddy simulations (LESs) were first employed on small domains (6.4×6.4 - 12.8×12.8
35 km^2) to investigate the response of shallow cumuli to aerosol perturbations (Xue et al., 2008; Zuidema et al., 2008). A decade later, LESs on larger domains (Seifert et al., 2015; Yamaguchi et al., 2019) showed that the compensating internal adjusting processes as proposed by Stevens and Feingold (2009) occur on the scales of $50 \times 50 \text{ km}^2$ cumulus cloud fields, which is far beyond the scale of an individual cloud. While not their focus, these studies give a clear indication of cold-pool activity. Cold pools are pockets of cold, dense air resulting from downdrafts associated with rain evaporation. When these downdrafts
40 reach the surface, cold pools spread outward in a circular pattern. Studies across various regimes—stratocumulus, shallow, and deep convection—indicate that cold-pool boundaries feature strong moist updrafts, which trigger cloud formation along their edges, forming cloud rings often visible in satellite imagery (Xue et al., 2008; Savic-Jovicic and Stevens, 2008; Zuidema et al., 2012; Böing et al., 2012; Jeevanjee and Roms, 2013; Schlemmer and Hohenegger, 2014; Langhans and Roms, 2015; Torri et al., 2015; Drager and van den Heever, 2017; Zuidema et al., 2017; Haerter and Schlemmer, 2018; Helfer and Nuijens, 2021;
45 Lochbihler et al., 2021; Vogel et al., 2021; Touzé-Peiffer et al., 2022). When cold pools form in close proximity, their boundaries can collide (Torri and Kuang, 2019), which intensifies the next convective event. Such cold-pool interactions through collisions implement self-organization, as has been conceptually modeled for both open-cell stratocumulus (Glassmeier and Feingold, 2017) and deep convective regimes (Haerter et al., 2019; Nissen and Haerter, 2021).

Shallow cumuli in the trades are frequently precipitating (Nuijens et al., 2009; Snodgrass et al., 2009; Radtke et al., 2022),
50 leading to the frequent presence of cold pools in the trade-wind regime (Zuidema et al., 2012; Vogel et al., 2021; Touzé-Peiffer et al., 2022). The size and frequency of occurrence of cold pools covary with the mesoscale organization of trade-cumulus clouds (Vogel et al., 2021). Even under large-scale conditions that are invariant in time and space, LES studies show that trade-cumulus cold pools self-organize and in turn pattern trade-cumulus fields into arc-shape structures (Seifert and Heus, 2013; Vogel et al., 2016). By generating strong moist updrafts at their fronts, cold pools affect and interact with clouds (Zuidema
55 et al., 2012; Li et al., 2014; Vogel et al., 2021; Alinaghi et al., 2025). On large-domain LESs of the trade-wind regime, Alinaghi et al. (2025) recently showed that the cold-pool-cloud interaction expresses itself in the form of structures resembling shallow squall lines. Thus, cold pools are coupled to clouds through shallow circulations at the mesoscales which were frequently observed in the trades (George et al., 2023) and affect cloudiness (Vogel et al., 2022; Janssens et al., 2023; Alinaghi et al., 2025).




Figure 1. Conceptual picture of the study. The diagram summarizes how (link 1) cloud fields respond to changes in N_c , and how (link 2) cold pools feed back to clouds (self-organization). The diagram further shows that the trade-cumulus system is forced by (link 3) the large-scale cloud-controlling factors (CCFs) whose relative importance for (link 4) the cloud-radiative effect will be quantified compared to N_c .

60 The mesoscale dynamics of trade cumulus are typically discussed in the context of cloud feedback. The trade-cumulus feedback has long been a large source of uncertainty in climate projections (Cesana and Del Genio, 2021; Myers et al., 2021). Trade-cumulus fields pattern into structures at the mesoscales that influence cloud-radiative effect (Bony et al., 2020; Alinaghi et al., 2024a; Denby, 2023). Therefore, it is crucial to explore processes through which these clouds organize and how these processes respond to the variations in large-scale cloud-controlling factors (CCFs). By employing a large ensemble of LESs, 65 the *Cloud Botany* ensemble (Jansson et al., 2023), Alinaghi et al. (2025) illustrated that cold pools in the trades are strongly controlled by the variations in the large-scale external CCFs. They particularly quantified the relative importance of CCFs with respect to each other. Additionally, diurnality in insolation, which acts as a time-varying CCF, was shown to strongly control the temporal evolution of cold pools throughout the entire *Cloud Botany* ensemble (Alinaghi et al., 2025).

Given the direct impact of N_c on precipitation formation, it is expected that trade-cumulus cold pools respond to variations in 70 N_c (Fig. 1, link 1), and thereby to feed back to cloud fields at the mesoscales (Fig. 1, link 2). Despite the previously investigated sensitivity of shallow cold pools to microphysics schemes (Li et al., 2015), the response of shallow cold pools to N_c has not been directly explored and quantified. Furthermore, it is unknown how such N_c variations change the interplay between cold pools and clouds in the trade-wind regime. This study explores this response by performing large-domain, large-eddy simulations in which we only vary cloud-droplet number concentrations N_c from 20 to 1000 /cm³. The newly added dimension of variability 75 in N_c here also enables us to systematically investigate the relative importance of N_c compared to the other CCFs (Fig. 1, link 3) for cold pools and the radiative effect of clouds in the trade-wind regime (Fig. 1, link 4). Hence, our work serves as a step towards understanding the significance of the aerosol forcing in comparison to the trade cumulus feedback and exploring the corresponding role of mesoscale dynamics.

This paper is structured as follows. Based on simulations discussed in section 2, we first investigate how trade-cumulus cold
80 pools respond to N_c variability (Fig. 1, link 1; sections 3.1, 3.2) and how this response shapes the mesoscale organization of
clouds (Fig. 1, link 2; section 3.2). Second, we explore how the diurnal cycle in insolation, as a time-varying CCF (Fig. 1, link
3), controls the evolution and response of cold pools to N_c (section 3.3). Next, we investigate the implications of our results for
the cloud-field adjustments to N_c (Fig. 1, links 1, 2; section 3.4). Finally, we compare the effect N_c on cloud-field properties
and radiative effects to that of large-scale external CCFs (Fig. 1, links 3, 4; section 3.5). Conclusions are presented in section
85 4.

2 Data and Methods

We perform large-eddy simulations (LESs) with the Dutch Atmospheric LES (DALES) model over domains of 153.6×153.6
 km^2 , featuring a horizontal resolution of 100 m and a vertical resolution of about 20 m. All simulations are forced by the
same large-scale CCFs. These follow the central reference simulation of the *Cloud Botany* ensemble (Jansson et al., 2023),
90 which corresponds to the mean large-scale conditions of the winter trades as derived from the ERA5 reanalysis data (Hersbach
et al., 2020). The corresponding profiles, which are also used for initialization are shown in Fig. 2. Moreover, all simulations
feature the same horizontal tendencies of cooling and drying through advection as shown in the *Cloud Botany* paper (Jansson
et al., 2023, their Fig. 3). Most of the simulations feature diurnality in the solar incoming radiation, while all other CCFs are
fixed in time. Thus, the variability and evolution in the simulations are driven by the interaction between the components of
95 the system, allowing the study of processes via which the system self-organizes. For more details on the design of the *Cloud*
Botany simulations, including the selection of parameters for the large-scale forcing, refer to Jansson et al. (2023).

Simulations utilize the two-moment cloud-microphysics scheme of Seifert and Beheng (2001) with a constant cloud-droplet
number concentration N_c . We conduct six, 72-hour simulations with varying N_c values in the set $\{20, 50, 70, 100, 200, 1000\}$
 $/\text{cm}^3$. The selected range of N_c variability, from 20 to 100, is similar to that used by Seifert et al. (2015), which is based on
100 observations of the trades (Colón-Robles et al., 2006; Gerber et al., 2008; Hudson and Noble, 2014). We also included N_c
values of 200 and 1000, as recent observations from the EUREC⁴A field campaign (Bony et al., 2017; Stevens et al., 2021)
report N_c values as high as $1000/\text{cm}^3$, primarily due to the presence of dust (see Fig. 9 in Quinn et al., 2021, and Figs. 9 and 10
in Bony et al., 2022). Note that since N_c is fixed in time and space and does not evolve in our simulations, our study excludes
microphysical adjustments. To gauge this limitation, we compare our results to cases from the literature with more complex
105 microphysics.

Figure 3 visualizes that all these simulations start from a homogeneous non-cloudy state and develop into randomly dis-
tributed cumulus clouds. Afterward, clouds self-aggregate due to the presence of self-reinforcing shallow mesoscale over-
turning circulations (Bretherton and Blossey, 2017; Narenpitak et al., 2021; Janssens et al., 2023). As clouds aggregate, they
deepen and eventually start to precipitate, leading to the presence of mesoscale arc-like structures, indicating the presence of
110 cold pools in the field.




Figure 2. Large-scale and initial conditions of the simulation for potential temperature θ_l , total moisture q_t , horizontal wind u , and updraft w following the assumptions of the *Cloud Botany* ensemble (Jansson et al., 2023) with parameter values of sea-surface (potential) temperature $\theta_{l0} = 299$ K, near-surface wind speed $u_0 = -10.6$ m/s, moisture scale-height $h_{q_t} = 1810$ m, temperature lapse-rate $\Gamma = 5$ K/km, large-scale vertical velocity variability $w_1 = 0.0393$ cm/s, and the horizontal wind shear $u_z = 0.0022$ (m/s)/m.

Alinaghi et al. (2025) showed that the evolution of cold pools across the entire *Botany* ensemble is strongly controlled by the diurnality of insolation, mirroring observations of the trades (Vial et al., 2021; Vogel et al., 2021). To investigate how the evolution of cold pools is affected by variations in N_c independently of the diurnal cycle, we switch off the diurnality in insolation. To this end, we re-run simulations with N_c values of 20, 70, and 1000 /cm³, while keeping the solar zenith angle
115 time-invariant, ensuring that the total incoming solar radiation over the entire 24-hour period is equal to that of the simulations with the diurnal cycle (Alinaghi et al., 2025, their section 3.2).

To diagnose cold pools, we use the 2D outputs of the mixed-layer height h_{mix} , as h_{mix} has been shown to be a reliable indicator for trade-cumulus cold pools in both models (Rochetin et al., 2021) and observations (Touzé-Peiffer et al., 2022). We identify cold pools following Alinaghi et al. (2025): for each cloud field, we find the mode and the upper boundary (99th
120 percentile) of an assumed symmetric probability density function (PDF) of h_{mix} that would have been observed in the absence of cold pools. ~~The lower boundary of h_{mix} is determined by subtracting the difference between the upper boundary and the mode from the mode~~ The lower boundary of h_{mix} is calculated by first determining the difference between the upper boundary and the mode, and then subtracting this difference from the mode. Cold pools are identified where h_{mix} is smaller than the lower boundary of h_{mix} (see Fig. 3 in Alinaghi et al. (2025)). In essence, this method identifies cold pools where h_{mix} is relatively shallower compared
125 to other parts of the field. Alinaghi et al. (2025) showed that the response of cold pools to CCFs is not sensitive to the details of the cold-pool diagnosis performed with this method.

Using the identified cold-pool mask and a clustering method, we define cold-pool objects as 2D contiguous structures within the simulation domain at each model time step. We then quantify the number of cold pools, n_{cp} , within the domain. Additionally, we compute the domain-mean cold-pool size as $s_{cp} = \sum_{i=1}^{n_{cp}} \sqrt{A_i} / n_{cp}$, where A_i denotes the area of each cold-pool object
130 i within the domain.




Figure 3. Cloud-field albedo examples. The figures show how cloud fields develop in our LES simulations featuring the diurnal cycle of insolation with N_c of 20 (1st row), 50 (2nd row), 70 (3rd row), 100 (4th row), 200 (5th row), and 1000 /cm³ (6th row), at hours 15 (1st column), 30 (2nd column), 40 (3rd column), and 50 (4th column) after the start of simulations.

3 Results and Discussion

3.1 Cloud-droplet number concentration affects the spatial and temporal properties of trade-cumulus cold pools

In this section, we investigate how the spatial and temporal properties of cold pools are influenced by N_c (Fig. 1, link 1, 2). As cold pools result from rain-evaporation (Fig. 1, link 2), we first examine how clouds and rain respond to N_c in simulations without diurnal cycle, which feature N_c values of 20, 70, and 1000 /cm³. According to theory (Albrecht, 1989) and previous
135 LES studies (Seifert et al., 2015; Yamaguchi et al., 2019), increased N_c reduces the efficiency of auto-conversion, delaying rain formation and allowing clouds to deepen and persist longer. Consistent with this, increased N_c leads to the accumulation of liquid water $\mathcal{L}$ (Fig. 4a), eventually resulting in the production of more intense rain $\mathcal{R}$ (Fig. 4b). Furthermore, the amplitude of fluctuations of $\mathcal{L}$ and $\mathcal{R}$ increases with increasing N_c , while their frequency decreases (Figs. 4a,b).

We quantitatively compare our results to those of Yamaguchi et al. (2019) and Seifert et al. (2015) (Figs. 4e,f). Results of Yamaguchi et al. (2019) were obtained on ten times smaller domains but using a microphysics scheme with prescribed aerosol and prognostic cloud-droplet number concentrations. The latter converges to a certain value after approximately 20 hours (Yamaguchi et al., 2019, their Fig. 5e). We therefore compare our fixed- N_c results to averages of the last 20 hours of their simulations. The systematic difference between the values of cloud-field properties is expected due to differences in the large-
145 scale cloud-controlling factors; notably, their geostrophic wind speed is 60% smaller than that of our simulations (Yamaguchi et al., 2019, their Table 1). Similarly, we present the results from Seifert et al. (2015), which feature a domain size similar to that of Yamaguchi et al. (2019) but fixed N_c as in our simulations. We selected their simulations with interactive radiation and prescribed large-scale advective cooling to match our setup as closely as possible. It is worth noting that the cloud fields in Seifert et al. (2015) have larger cloud- and rain-water paths (Figs. 4e,f), which we mainly attribute to their larger meridional
150 geostrophic wind.

Consistent with Yamaguchi et al. (2019) and Seifert et al. (2015), increased N_c leads to an increased domain-mean liquid-water path, $\mathcal{L}$ (Fig. 4e). As shown by the error bars, the temporal variance in $\mathcal{L}$ increases in response to increased N_c in our simulations. In addition, Fig. 4f illustrates that the domain-mean rain-water path, $\mathcal{R}$, decreases with increasing N_c in all studies. Similar to $\mathcal{L}$, the temporal variations in $\mathcal{R}$ increase with increasing N_c in our simulations. This is in contrast to the
155 small-domain LESs of Yamaguchi et al. (2019, 50×50 km²) and Dagan et al. (2018, 12×12–50×50 km²), where increased N_c was found to reduce the amplitude of fluctuations in the time series of $\mathcal{R}$. The temporal variance of liquid- and rain-water path in simulations of Seifert et al. (2015) do not show a systematic response to N_c . The response of the temporal variations of cloud-field properties to N_c thus appears markedly different in our large domains.

The temporal variations of cold-pool characteristics follow those of the rain-water path, $\mathcal{R}$ (Figs. 4b-d). Notably, in the
160 simulation with $N_c = 20/\text{cm}^3$, once cold pools form around hour 30, they grow until around hour 36 and persist until the end of the simulation. In contrast, cold pools in the simulation with $N_c = 1000/\text{cm}^3$ form, develop, reach a maximum, decay, and completely vanish. Therefore, increased N_c enhances the intermittency in the evolution of cold pools. Averaged over the last day, increased N_c leads to a smaller number of cold pools, n_{cp} (Fig. 4g), while the domain-mean size of cold pools, s_{cp} , shows a muted response to N_c variations (Fig. 4h).




Figure 4. Time series of clouds, rain, and spatial properties of cold pools in simulations without the diurnal cycle. Panels (a-d) show time series of $\mathcal{L}$, $\mathcal{R}$, n_{cp} , and s_{cp} for simulations without the diurnal cycle of solar incoming radiation, where solid lines shown for cold-pool number and size provide a guide to the eye obtained through Gaussian filtering of the original data points. Note that there is significant precipitation in the $N_c = 20 / \text{cm}^3$ case before cold pools with diameters larger than the 5-km threshold appear.. Panels (e-h) show the response of $\mathcal{L}$, $\mathcal{R}$, n_{cp} , and s_{cp} to N_c . The results from Yamaguchi et al. (2019) averaged over the last 20 hours of their simulations are shown in orange. The results from Seifert et al. (2015) for their near-equilibrium state are shown in blue. Dashed lines are added as a visual guide where the mean values show a trend. Note that the reason for the cold-pools in simulation $N_c = 70 / \text{cm}^3$ forming earlier than $N_c = 20 / \text{cm}^3$ is that cold pools featuring a diameter of < 5 km is not considered to be part of the cold-pool mask. Time series of cold-pool number and size are slightly smoothed using Gaussian filtering.

165 3.2 Cold-pool evolution shows two distinct mesoscale behaviours for low and high cloud-droplet number concentrations

To understand the difference in cold-pool dynamics for different cloud-droplet number concentrations, Fig. 5 contrasts the extreme N_c cases. Consistent with observations (Zuidema et al., 2012; Vogel et al., 2021), both have in common that cold




Figure 5. Effects of cold pools on the organization of the (sub-)cloud layer properties for the simulations without diurnal cycle and with $N_c = 20/\text{cm}^3$ and $N_c = 1000/\text{cm}^3$. For each condition, columns 1-4 indicate the 2D top-views of cloud albedo, total moisture anomalies (q'_t) at the 200-m level, the horizontal wind speed anomalies (U') at the first level of the model, and the vertical velocity (w) at the 200-m height. The green dashed contour line marks the cold-pool boundaries quantified from the mixed-layer height fields. In the U' fields, red indicates that cold pools accelerate the wind, while blue indicates they decelerate it.

pools alter the spatial pattern of moisture in the sub-cloud layer. Cold pools are characterized by relatively dry air inside but relatively moist air at their fronts. Cold pools further modify the spatial pattern of horizontal velocity in the sub-cloud layer, thereby altering the spatial pattern of convergence and, in turn, vertical velocity. Cold pools feature large values of downward vertical velocity inside but strong updrafts at their fronts. Therefore, cold-pool fronts actively contribute to moist convergence and cloud formation. In contrast, the downward motions inside cold pools can potentially suppress convection and lead to cloud-free regions.

Despite the similar evolution of individual cold pools independent of N_c , there are notable differences in the collective, i.e. self-organization, behavior. In the low- N_c simulation (i.e., $N_c = 20/\text{cm}^3$), cold pools tend to form in close proximity. As their fronts converge, they trigger the formation of new cold pools at their collision points. This triggering mechanism occurs due to (i) the collision of anomalously moist cold-pool fronts, which mechanically forces the moist air upward due to mass conservation, and (ii) the high efficiency of rain formation, where a small amount of cloud water quickly turns into rainwater, leading to the formation of a new cold pool at the collision point. Consequently, the small, space-filling cold pools at hours

30-36 collide and transition into a stage where they become organized into a large front, sustained by the interaction (collision) of its cold pools at its fronts (see Fig. 5, $N_c = 20/\text{cm}^3$, hour 48). This is consistent with the time series of cold-pool number n_{cp} and size s_{cp} shown in Figs. 4c,d, which show that after around hour 38, the metrics of cold pools stabilize: s_{cp} remains around 15 ± 5 km, and n_{cp} around 6 ± 4 . This behaviour of cold pools resembles the mathematical toy model of colliding circle
185 cold pools presented by Nissen and Haerter (2021, their Fig. 5), which demonstrates (i) similar transitions from randomly distributed cold pools to a band-like structure, and (ii) that cold-pool collision is the key mechanism for the self-organization of the system. Notably, their model was motivated by cold pools in the regime of deep convection.

In contrast, cold pools do not interact as readily in the high- N_c simulation (i.e., $N_c = 1000/\text{cm}^3$). This is attributable to two factors. First, the efficiency of rain formation and, consequently, cold-pool formation, is significantly lower in high- N_c than in
190 low- N_c simulations. Second, cold pools form at greater distances from each other. We speculate that this distance is determined by the horizontal length scale of the self-reinforcing shallow circulations that lead to moisture aggregating in the absence of rain (Bretherton and Blossey, 2017; Narenpitak et al., 2021; Janssens et al., 2023; George et al., 2023). As Janssens et al. (2023) showed, the aggregation of non-precipitating cumuli develops atop such anomalous mesoscale moist regions, whose size expands in tandem with the growth of anomalous mesoscale dry regions. The scale growth of dry regions in the descending
195 branch of circulations effectively separates moist regions—and consequently, their associated aggregated clouds—preventing interactions between their subsequent cold pools, which form at greater distances from each other. AsConsistently, suggested by Fig. 5 ($N_c = 1000/\text{cm}^3$, hour 38) suggests that the downward branches of these circulations are so large that they effectively separate two cloud clusters and cause their cold pools to form at a distance from each other, thus preventing their interactions.

The intermediate case of $N_c = 70/\text{cm}^3$ exhibits a cold-pool evolution similar to that of the high- N_c case with $N_c = 1000/\text{cm}^3$:
200 (i) cold pools in the $N_c = 70/\text{cm}^3$ case also form far apart, resulting in very infrequent cold-pool interactions, and (ii) its cold-pool evolution displays strong intermittency, as shown by Figs. 4c,d. The high- N_c behaviour corresponds to the behaviorThe self-organization dynamics of this case were recently discussed by Alinaghi et al. (2025) ~~for simulation with $N_c = 70/\text{cm}^3$~~ , where cold pools are characterized in analogy to squall lines in deep mesoscale convective systems (Rotunno et al., 1988; Weisman and Rotunno, 2004; Stensrud et al., 2005). In this regime, cold pools reinforce and sustain their parent clouds due to the convergence of moist
205 air at their fronts. Once cold pools get mature, their moist updrafts at their fronts become so strong such that they impinge on the inversion, leading to the formation of stratiform anvils with stratiform precipitation. This weakens the cold-pool-induced updraft, ultimately causing parent clouds to detach from their cold-pool children (Alinaghi et al., 2025, their Figs. 7, 8, 11). This self-organizing behaviour of cold pools and clouds in simulations where they cannot interact is also evident in the cold-pool time series (Fig. 4cd), which exhibit an intermittent behaviour with large amplitudes and low frequencies for both $N_c = 70$
210 and $1000/\text{cm}^3$.

An interesting observation is that rain and cold pools tend to develop more rapidly once the initial event has occurred. For instance, the time series shown in Fig. 4b-d demonstrates that it takes approximately 36 hours for the $N_c = 1000/\text{cm}^3$ simulation to produce rain and cold pools. However, subsequent cold-pool events occur within about 8 hours, indicating a faster formation of rain and cold pools. In this simulation, the next convective event takes place precisely where the fronts of the previous
215 cold-pool event had accumulated moisture, thereby expediting the development of subsequent convection. We hypothesize that

cold pools in the high- N_c simulation act as a “moisture memory” similar to deep convection (Colin et al., 2019), facilitating aggregation over shorter timescales compared to when cold pools are absent. This suggests that the moisture variance induced by cold pools ~~compensates~~decreases for the induced delay in rain formation associated with increased N_c .

As the final point in this section, we investigate how the two N_c -induced regimes of cold-pool self-organization dynamics relate to the mesoscale organization of trade-cumulus cloud fields (Fig. 1, link 2). To address this, we quantify several metrics based on the geometry of clouds that effectively capture the variability in the mesoscale organization of cloud fields in the trades (Janssens et al., 2021). These include the domain-averaged size of cloud objects, the mean fraction of open-sky areas, the domain-mean depth of clouds, and the degree of organization I_{org} . The details of these metrics and their calculations are explained by Janssens et al. (2021). In addition to geometry-based cloud metrics, we further compute more organization metrics that are based on spatial distribution of cloud-liquid-water and total moisture fields. First, we quantify the spatial standard deviation of the liquid-water path $\sigma\mathcal{L}$. Second, following Radtke et al. (2023), we consider the metric ΔQ , which quantifies the moisture aggregation at the mesoscales. This metric is calculated as the difference between the 5th and 95th percentiles of the mesoscale total moisture anomaly fields, derived by coarse-graining the total moisture anomaly fields as outlined by Janssens et al. (2023, see their Fig. 3).

Interestingly, the geometry-based organization metrics quantifying the mean size of cloud objects and the open-sky areas, which were shown to explain most of the variability within the mesoscale organization of trade cumuli (Janssens et al., 2021), do not capture the two distinct behaviours of cold pools at low- and high- N_c regimes discussed in the context of Fig. 5 (Fig. A1). However, Fig. 6a,b shows that the spatial variance in cloud-liquid-water path $\sigma\mathcal{L}$ is strongly affected by N_c . First, increased N_c translates into a strong intermittent behaviour in $\sigma\mathcal{L}$ evolution (Fig. 6a). Second, Fig. 6b indicates that, when averaged over 24-48 and 48-72 hour intervals, increased N_c leads to greater spatial heterogeneity in liquid-water content: shallow cumuli become more aggregated in response to increased N_c , as visually evident in the snapshots of cloud fields shown in Figs. 3 and 5. The $\sigma\mathcal{L}$ - N_c relationship suggests that a reduced number of cold pools, in response to increased N_c , enhances cloud aggregation. This echoes Radtke et al. (2023)’s findings, which also show that rain (auto-conversion) is less efficient in more aggregated fields of trade cumuli (Radtke et al., 2023, their Fig. 2).

With the delay in precipitation formation due to increased N_c , moisture is expected to continue aggregating through shallow circulations driven by latent heating from condensation in the non-precipitating cumulus layer (Bretherton and Blossey, 2017; Janssens et al., 2023). Fig. 6c shows that this is exactly what happens in simulations with high N_c . In simulations with N_c of 70 and 1000 /cm³, the moisture aggregation metric ΔQ keeps growing until cold pools start to form, after which ΔQ stabilizes and shows an intermittent behaviour. In contrast, in the lowest case of N_c , the moisture does not aggregate at the mesoscales until hour 30, where cold pools start to form, after which the moisture aggregation metric ΔQ starts developing and keeps increasing until the end of the simulation. This implies that this metric of moisture aggregation at the mesoscales is intriguingly able to encapsulate the contrasts between these distinct behaviours of cold pools at low- and high- N_c . Averaged over hours 24-48, ΔQ associated with the low- N_c is by 50% smaller than of that of the high- N_c cases (Fig. 6d). However, as the simulations progress into hours 48-72, cold pools in low- N_c simulations develop into large squall lines, increasing their moisture aggregation and reducing the sensitivity of ΔQ to N_c (Fig. 6d).




Figure 6. Dependence of mesoscale organization of cloud fields on N_c . The panels (a,c) show the time series of the spatial standard deviation of the liquid-water path σ_L and the difference between the 5th and 95th percentiles of the mesoscale total moisture anomaly fields ΔQ for the simulations without the diurnal cycle of insolation. Panels (b,d) show their responses to N_c for hours 24-48 in purple and hours 48-72 in green. Dashed lines are added as a visual guide where the mean values have a trend.

In summary, the evolution of cold pools and trade-cumulus fields is significantly influenced by N_c , exhibiting two distinct behaviours at low and high N_c (Fig. 1, links 1, 2). In simulations with $N_c = 20/\text{cm}^3$, cold pools form in close proximity, leading to interactions and collisions that trigger the formation of clouds and due to high rain efficiency formation of new cold pools at the collision points. This results in persistent long-lived structures resembling squall lines. Conversely, in simulations with higher N_c values (70 and 1000/ cm^3), cold pools form at greater distances, preventing interactions and resulting in intermittent behaviour. Cold pools in such cases are short-lived structures resembling squall lines, hypothetically facilitating convection by providing moisture anomalies at their fronts, thereby compensating/decreasing for N_c -induced delays in subsequent cold-pool formation. This N_c -driven contrast in cold-pool dynamics affects moisture and cloud-water variance, with higher N_c leading to more aggregated trade-cumulus cloud fields.

260 3.3 Diurnal cycle synchronizes the phases of cold-pool evolution across simulations with perturbed N_c

We have shown that the temporal evolution of cold pools, and the trade-cumulus system in general, is controlled by N_c (Fig. 1, links 1, 2). Observations have shown that the evolution of trade-cumulus fields, their mesoscale organization, precipitation, and cold pools in the trades feature diurnality (Nuijens et al., 2009; Vogel et al., 2021; Vial et al., 2021; Radtke et al., 2022). This raises the question: to what extent does the diurnal cycle of insolation, as a time-varying external forcing, control or affect the influence of N_c on the evolution of cold pools (Fig. 1, link 3)?

To answer this question, we plot the time series of the domain-mean rain-water path $\mathcal{R}$, the number n_{cp} , and the size s_{cp} of cold pools in our simulations with the diurnal cycle, featuring N_c values of 20, 50, 70, 100, 200, and 1000 / cm^3 . As expected,




Figure 7. Effects of the diurnal cycle of insolation on the evolution and N_c -response of rain and cold pools. Panels (a-c) show time series of rain-water path $\mathcal{R}$, cold-pool number n_{cp} , and cold-pool size s_{cp} for simulations with the diurnal cycle of solar incoming radiation. Nighttime is shown by the grey color. Solid lines shown for cold-pool number and size provide a guide to the eye obtained through Gaussian filtering of the original data points. Note that there is significant precipitation in the $N_c = 20$ /cm³ case before cold pools with diameters larger than the 5-km threshold appear. Panels (d-e) show the response of $\mathcal{R}$, n_{cp} , and s_{cp} to N_c during the transient (purple), and near-equilibrium (green) phases, which are marked in the time series plots (a-c). Dashed lines are added as a visual guide where there is a trend. Note that the reason for the cold pools in simulation $N_c = 70$ /cm³ forming earlier than $N_c = 20$ /cm³ is that cold pools featuring a diameter of < 5 km are not considered to be part of the cold-pool mask. Time-series of cold-pool number and size are slightly smoothed using Gaussian filtering.

Figs. 7a-c show that, across all simulations, the evolution of rain and cold pools follows the diurnal cycle of radiation, peaking around sunrise and reaching a minimum around sunset. This pattern is due to the absence of solar radiation combined with longwave radiative cooling during the night. This strong nighttime radiative cooling destabilizes the atmosphere, stimulating convection and leading to the formation of deeper clouds that precipitate more intensely. In contrast, during the daytime, radiative heating from solar radiation stabilizes the atmosphere, suppressing convection and causing a notable decrease in rain, and in the number and size of cold pools in almost all simulations. Thus, the diurnal cycle serves as an external forcing (Fig. 1, link 3) that synchronizes the periodicity and amplitude of cold-pool variability that were discussed in the context of Fig. 4 across simulations with different N_c . The simulated diurnal cycle in precipitation and cold pools is consistent with observations of the trades (Nuijens et al., 2009; Vogel et al., 2021; Vial et al., 2021; Radtke et al., 2022).

Although all simulations show evolution synchronized with the diurnal cycle, the cold pools in the simulation with the lowest N_c continue to persist even during the daytime when convection suppression due to reduced net radiative cooling is at its peak. This behaviour is notable as it indicates that the mesoscale self-organization of cold pools through collisions, as discussed in Section 3.2, outweighs the externally imposed suppressive effect of the diurnal cycle during the day. Consequently, cold pools in the $N_c = 20/\text{cm}^3$ simulation remain active throughout the day. However, the duration of cold-pool's persistence during the day decreases with increasing N_c . Specifically, during daytime hours around 34-46, increased N_c leads to an earlier disappearance of cold pools and a more delayed formation of the next generation of cold pools, as shown in Fig. 7c. To summarize, we find that the diurnal cycle externally synchronizes the mesoscale self-organization dynamics of cold pools, which is in turn modulated by the details of microphysics and rain formation.

The synchronization of cold-pool events by the diurnal cycle enables us to compare the responses of rain-water path $\mathcal{R}$ and cold pools to N_c during the same time window across all simulations. In all simulations, except for $N_c = 20/\text{cm}^3$, rain and subsequently cold pools begin to form after hour 24 (Figs. 7a-c), with the evolution of cold pools following the diurnal cycle of net radiative cooling. We refer to the day starting at hour 22 as the "transient" phase and the subsequent day beginning at hour 46 as the "near-equilibrium" phase. We selected these hours based on the development of the total-water (cloud and water vapor) path in our simulations, which consistently increases until hour 48 across all simulations, after which it stabilizes and becomes time-invariant (Fig. A2).

All simulations consistently show a higher daily mean rain-water path $\mathcal{R}$ during the near-equilibrium phase compared to the transient phase (Fig. 7d). This is because, in the near-equilibrium phase, our simulations are more developed and feature deeper boundary layers with larger total water that can develop more rain. Consistent with our results based on the last 24 hours of simulations without the diurnal cycle (Figs. 4f-h), Figs. 7d,e illustrate that during the both transient and near-equilibrium phases, rain-water path $\mathcal{R}$ and the number of cold pools n_{cp} decrease with increasing N_c . Also, the size of cold pools s_{cp} do not show a notable change in response to N_c (Fig. 7f).

3.4 Twomey effect primarily controls the dependence of cloud-radiative effect on N_c

In this section, we investigate the sensitivity of the relative cloud-radiative effect (Xie and Liu, 2013), $\text{rCRE} = f_c \cdot \mathcal{A}_c$, to N_c (Fig. 1, link 4), with f_c and $\mathcal{A}_c$ as cloud fraction and albedo. Assuming the plane-parallel approximation (Lacis and Hansen, 1974), cloud albedo is given by $\mathcal{A}_c = \frac{\tau}{\tau + 7.7}$ with the cloud-optical thickness $\tau \approx N_c^{1/3} \mathcal{L}^{5/6}$ following Zhang et al. (2005). Therefore, we explore the response of cloud fraction f_c , domain-mean liquid-water path $\mathcal{L}$, mean cloud albedo $\mathcal{A}_c$ over the cloudy columns where the cloud-liquid-water path is larger than zero, and the relative cloud-radiative effect rCRE to N_c . These sensitivities are explored during different phases of the simulations with the diurnal cycle: non-precipitating (hours 5-15), transient (hours 22-46), and near-equilibrium (hours 46-72). For comparison, we also include results from Seifert et al. (2015) and Yamaguchi et al. (2019) , where the latter features more comprehensive microphysics than our study with its fixed cloud-droplet number concentration.

During the non- (or weakly) precipitating phase, the response of the cloud fraction f_c and liquid-water path $\mathcal{L}$ to N_c is negligible (Figs. 8a,b). Thus, the relative cloud-radiative effect (rCRE) is influenced by N_c primarily through the cloud albedo




Figure 8. Sensitivity of relative cloud-radiative effect to cloud-droplet number. Panels (a-d) show the response of cloud fraction f_c , domain-mean liquid-water path $\mathcal{L}$, cloud albedo $\mathcal{A}_c$, and relative cloud-radiative effect rCRE to N_c during the non-precipitating (dark grey), transient (purple), and near-equilibrium (green) phases. The results from Yamaguchi et al. (2019) averaged over the last 20 hours of their simulations are shown in orange. The results from Seifert et al. (2015) for their near-equilibrium state are shown in blue. Note that Yamaguchi et al. (2019); Seifert et al. (2015) do not report on the cloud albedo and rCRE in their studies. Dashed lines are added as a visual guide where [there is a trend](#).

response, or the Twomey effect (Figs. 8c,d). During both the transient and near-equilibrium phases, the cloud fraction f_c decreases very slightly with increasing N_c , though this response is much smaller than the temporal variance of f_c within each simulation (Fig. 8a). Additionally, the liquid-water path $\mathcal{L}$ shows a small increase with increased N_c (Fig. 8b). However, similar to the non-precipitating phase, the Twomey effect continues to strongly control the response of rCRE to N_c during both the transient and near-equilibrium phases (Figs. 8c,d).

The response of the liquid-water path in the near-equilibrium phase is consistent with the results of Yamaguchi et al. (2019) and Seifert et al. (2015). This small positive sensitivity of the liquid-water path in our simulations does not significantly affect the impact of N_c on rCRE compared to the Twomey effect. Cloud fraction f_c decreases with increasing N_c in the simulations of Yamaguchi et al. (2019). Similarly, our simulations and those of Seifert et al. (2015) both show a decrease in f_c in response to increased N_c . However, this decrease seems to be smaller compared with the f_c response in Yamaguchi et al. (2019)’s simulations. It would be interesting to revisit this difference with a cloud microphysics scheme that does not fix cloud-droplet number as in the current study but allows for full microphysical adjustments.

It is worth noting that although variations in N_c modulate the number of cold pools and in turn, mesoscale self-organization of trade-cumulus fields, cloud fraction f_c is very weakly affected. Our results here resonate with Janssens et al. (2025) who hypothesize that circulations associated with self-organization symmetrically distribute cloudiness at the mesoscales such that the increased cloudiness at their ascending branch is compensated by the decreased cloudiness at their descending branch. This implies that although decreased N_c increases the number of cold pools, the increased cloudiness at their fronts, where convection is triggered, appears to be buffered by the decreased cloudiness at their interiors, where convection is suppressed. Future studies are encouraged to explicitly investigate this.

3.5 N_c induces comparable variations in cloud-radiative effect to the large-scale cloud-controlling factors

In this section, we quantify the relative importance of N_c (Fig. 1, link 1) compared with the large-scale cloud-controlling factors (CCFs; Fig. 1, link 3) for driving changes in cloud-field properties and radiative effects (Fig. 1, links 2, 4). Using the data from the *Botany* dataset (Jansson et al., 2023) as well as our new simulations of this study, we employ a multivariate regression model

$$\overline{C} \approx \sum_{i=1}^7 \beta_i \times \widetilde{CCF}_i \quad \text{with} \quad \widetilde{CCF}_i := \frac{CCF_i - \overline{CCF}_i}{\sigma(CCF_i)},$$

where, the vector $\overline{C}$ represents the mean of the metric $C \in \{\mathcal{A}_c, \text{rCRE}\}$, averaged over the last 2 days (hours 12–60) for each member of the *Botany* ensemble. Each regressor $\widetilde{CCF}_i$ is a vector containing the associated CCF_i values for the simulation members of the *Botany* ensemble. The regressors (CCFs) include sea-surface (potential) temperature (θ_{l0}), near-surface wind speed (u_0), moisture scale height (h_{qt}), temperature lapse rate (Γ), large-scale vertical velocity variability (w_1), and horizontal wind shear (u_z). In addition, we consider N_c as a 7th regressor. For the regressor N_c , we use the simulations with the diurnal cycle over hours 12-60 to be consistent with the simulations of the *Botany* ensemble. All regressors are standardized by subtracting their mean $\overline{CCF}$ and dividing by their standard deviation $\sigma(CCF)$ across the ensemble, which allows the comparison of CCFs and N_c with an equal weighting. In our regression analysis, we only include simulations that develop clouds and run for at least 48 hours. This leaves us with 80 simulations out of the initial 103 in the *Botany* ensemble. Including the simulations with the diurnal cycle from this study (six in total) and noting that the simulation with $N_c = 70/\text{cm}^3$ being already part of the ensemble as the central reference simulation, our regression analysis features 85 data points in total. This means that the target value $\overline{C}$ and regressors $\widetilde{CCF}_i$ of the regression model are vectors of size 85×1 .




Figure 9. Cloud-field response to large-scale cloud-controlling factors and cloud-droplet number concentration. The standardized β coefficients of the multiple regression analysis for (a) cloud albedo $\mathcal{A}_c$ and (b) relative cloud-radiative effect $rCRE$, all averaged over the last two days of the LESs of the *Botany* ensemble. The error bars show the 95% confidence interval for each regressor. The p -values of the F -statistic test of all models are smaller than 10^{-22} .

Figure 9 shows the results of the multivariate regression for cloud albedo $\mathcal{A}_c$ and $rCRE$. Note that the response of cloud albedo $\mathcal{A}_c$ and $rCRE$ to CCFs of the *Botany* ensemble has already been addressed and discussed by Janssens (2023a); Janssens et al. (2025), to which we refer for details. Figure 9a illustrates that the effect of N_c on cloud albedo $\mathcal{A}_c$, known as the Twomey effect, is comparable to that of large-scale subsidence as quantified by w_1 . Additionally, the effect of N_c on trade-cumulus brightening is about 75% of the effect of horizontal wind speed, 150% of the effect of stability, and 300% of the effects of free-tropospheric humidity and vertical wind shear. Eventually, the response of $rCRE$ to N_c is statistically significant at the 95% level and accounts for about 66%, 28%, and 25% of the response of $rCRE$ to free-tropospheric humidity, wind speed, and large-scale subsidence, respectively (Fig. 9b). Note that the sensitivity of $rCRE$ to N_c is smaller than that of cloud albedo to N_c in our regression analysis, which is due to the very weak impact of N_c on cloud fraction f_c (Fig. 8a). Similar regression results for other cloud-field and cold-pool properties are presented in Fig. A3.

4 Conclusions & Outlook

Cold pools, resulting from rain-evaporation, affect the mesoscale organization of trade-cumulus fields (Zuidema et al., 2012; Seifert and Heus, 2013; Vogel et al., 2016, 2021; Alinaghi et al., 2025). We have used an ensemble of large-domain LESs to investigate the sensitivity of mesoscale organization to cloud-droplet number concentration N_c and the role of cold pools therein as conceptualized in Fig. 1.

Investigating the sensitivity of mesoscale cold-pool dynamics to microphysics (Fig. 1, link 1), we find that cold pools show two distinct behaviours at low and high N_c . In low- N_c cases, there are many cold pools within the simulation domain, which form in close proximity (Figs. 4c,d,g,h, 5). This allows them to interact with each other through collisions. Efficient rain formation then leads to the swift triggering of new cold pools at collision points. Consequently, cold pools organize into a large long-lived front with a resemblance to a squall line that perpetuates through collisions of cold pools at its leading edge. In contrast, high- N_c cases feature sparsely distributed cold pools, which prevents their interaction. In this regime, cold pools exhibit an intermittent behaviour, manifesting as small, short-lived fronts resembling squall lines that form, develop, decay, and vanish (Figs. 4c,d,g,h, 5).

For the effect of N_c on the interaction between clouds and cold pools (Fig. 1, link 2), our analysis shows that increased N_c suppresses the formation of cold pools (Figs. 4g,h, 5), while enhancing the self-aggregation of cloud fields (Figs. 6). In other words, by delaying precipitation formation, increased N_c allows non-precipitating cumulus fields to aggregate moisture (Fig. 6c,d) through self-reinforcing mesoscale overturning circulations (Bretherton and Blossey, 2017; Narenpitak et al., 2021; Janssens et al., 2023). We quantified this effect, which clearly influences the mesoscale organization of cloud fields, particularly by increasing the spatial variance in the cloud-liquid-water path (Fig. 6a,b). Interestingly, despite the suppression of cold pools and the boost in aggregation due to increased N_c (Figs. 6a-d), the overall daily mean responses of cloud fraction and liquid-water path are notably small (Figs. 4e, 8a,b). This echoes recent findings of Janssens et al. (2025) that shallow mesoscale circulations appear to symmetrically modulate cloudiness within trade-cumulus fields such that the increased cloudiness at their ascending branch is compensated by the decreased cloudiness at their descending branch. This suggests that although by increasing N_c , mesoscale organization is affected by the variations in cold pools, the increased cloudiness at the edges of cold pools appears to be buffered by the decreased cloudiness at their interiors.

For the effect of the diurnal cycle as an external forcing on the microphysical sensitivity (Fig. 1, link 3), we find that the diurnal cycle synchronizes the self-organization dynamics of cold pools on the mesoscale across simulations with varying N_c (Fig. 7). Cold pool activity is thus controlled by both, the external forcing as well as the self-organization dynamics. The contribution of self-organization dynamics increases with decreasing N_c as showcased by the fact that cold-pool activity survives the day-time suppression for the lowest N_c case (Figs. 7b,c).

To compare the importance of microphysical and large-scale controls on the cold-pool dynamics (Fig. 1, links 1 and 3) occurring on the intermediate mesoscale, we have made use of the *Cloud Botany* ensemble (Jansson et al., 2023). We demonstrate that the Twomey effect is as significant as the primary cloud-controlling factors for the brightening of trade-cumulus fields (Fig. 9a). Despite the very small response of cloud fraction to N_c (Fig. 8a), the response of the relative cloud-radiative effect to N_c (Fig. 1, link 4) is about 25% as significant as the response of rCRE to horizontal wind speed and large-scale subsidence (Fig. 9b).

We have obtained these results using a prescribed cloud-droplet number as a proxy for microphysical influences. [This assumption excludes microphysical adjustments. Despite a broad agreement between our results and those of](#) Yamaguchi et al. (2019) [that are based on](#) more detailed microphysics scheme [with prognostic \$N_c\$](#) , Li et al. (2015) showed that cold pools are sensitive to choices of microphysics schemes. We therefore encourage future research, such as the cold-pool model

intercomparison project (Kazil et al., 2025, CP-MIP), to focus on the sensitivity of mesoscale cold-pool dynamics to such microphysical choices if we truly want to understand rain-evaporation, cold pools, and their relevance for trade-cumulus fields.

400 Irrespective of its idealizations, this study clearly highlights that aerosol-cloud interactions are affected by processes happening at multiple spatial and temporal scales, ranging from the microscale via the mesoscale to the large scales. For shallow cloud fields in the trades, we demonstrate that variations in the microscale can manifest themselves at the mesoscale to a degree that is comparable to the influence of large-scale controls. We consider these findings a valuable step towards understanding the mesoscales as a pre-requisite for constraining trade-cumulus climate feedbacks as well as trade-cumulus mediated aerosol

405 forcings.

Data availability. The *Cloud Botany* dataset is publicly accessible through the EUREC⁴A intake catalog (https://howto.eurec4a.eu/botany_dailes.html). The simulation outputs of Yamaguchi et al. (2019) are publicly available from the NOAA dataset platform (https://csl.noaa.gov/groups/csl9/datasets/data/cloud_phys/2019-Yamaguchi-Feingold-Kazil/). The data is analyzed using Python (libraries: Numpy (Harris et al., 2020), Xarray (Hoyer and Joseph, 2017), Pandas (Wes McKinney, 2010), Scipy (Virtanen et al., 2020), Statsmodel (Seabold and Perktold, 2010), Matplotlib (Hunter, 2007), and Seaborn (Waskom, 2021)). The basic profiles and time series associated with cloud-field properties, cloud organization metrics, and cold-pool properties as well as a movie of simulations are publicly available by Alinaghi et al. (2024b) and from the link <https://zenodo.org/records/13868738>. The coarse-graining of the total-water path anomaly fields is done using the code (SMOCs.ipynb) by Janssens (2023b) which is publicly available (<https://doi.org/10.5281/zenodo.8089287>).

410

Author contributions. The concept for this study was developed by PA and FG. LES simulations were performed by FJ. PA analyzed and visualized the data, with DB assisting during the analysis. PA and FG interpreted the results with contributions from FJ and DB. PA drafted the initial manuscript, and all authors contributed to its revision and finalization.

415

Competing interests. At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics.

Acknowledgements. FG and PA acknowledge support from The Branco Weiss Fellowship - Society in Science, administered by ETH Zurich A. FG also acknowledges support by the European Union (ERC, MesoClou, 101117462). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them. FJ acknowledges support from the European Union's Horizon 2020 research and innovation program under grant agreement no. 820829 (CONSTRAIN project). PA and FJ thank SURF (www.surf.nl) for making the National Supercomputer Snellius accessible for running and analyzing the droplet number simulations.

420

Appendix A: Supplementary Figures




Figure A1. Dependence of several mesoscale cloud organization metrics on N_c . The panels (a,c,e,g) show the time series of the domain-mean size of cloud objects L_c , the mean fraction of the open-sky areas L_o , the domain-mean of cloud-top height z_t , and the degree of organization I_{org} for the simulations without the diurnal cycle of insolation. Panels (b,d,f,h) show their mean responses to N_c for hours 24-48 in purple and 48-72 in green. The dashed line is shown as a visual guide where there is a trend.




Figure A2. Total-water path time series for several N_c . The figure shows the development of the domain-mean total-water path, which is the sum of both cloud-water and water-vapor paths. The transient and near-equilibrium phases are marked by purple and green.

References

Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, <https://doi.org/10.1126/science.245.4923.1227>, 1989.

Alinaghi, P., Janssens, M., Choudhury, G., Goren, T., Siebesma, A. P., and Glassmeier, F.: Shallow cumulus cloud fields are optically thicker when they are more clustered, *Quarterly Journal of the Royal Meteorological Society*, 150, 3566–3577, <https://doi.org/https://doi.org/10.1002/qj.4783>, 2024a.

Alinaghi, P., Jansson, F., A. Blázquez, D., and Glassmeier, F.: Datasets and the movie for the manuscript "Cold pools mediate mesoscale adjustments of trade- cumulus fields to changes in cloud-droplet number concentration", <https://doi.org/10.5281/zenodo.13868738>, 2024b.

Alinaghi, P., Siebesma, A. P., Jansson, F., Janssens, M., and Glassmeier, F.: External Drivers and Mesoscale Self-Organization of Shallow Cold Pools in the Trade-Wind Regime, *Journal of Advances in Modeling Earth Systems*, 17, e2024MS004540, <https://doi.org/https://doi.org/10.1029/2024MS004540>, e2024MS004540 2024MS004540, 2025.

Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K. S., Christensen, M., Daniau, A.-L., et al.: Bounding global aerosol radiative forcing of climate change, *Reviews of Geophysics*, 58, e2019RG000660, 2020.

Böing, S. J., Jonker, H. J., Siebesma, A. P., and Grabowski, W. W.: Influence of the subcloud layer on the development of a deep convective ensemble, *Journal of the Atmospheric Sciences*, 69, 2682–2698, 2012.

Bony, S., Stevens, B., Ament, F., Bigorre, S., Chazette, P., Crewell, S., Delanoë, J., Emanuel, K., Farrell, D., Flamant, C., et al.: EUREC 4 A: A field campaign to elucidate the couplings between clouds, convection and circulation, *Surveys in Geophysics*, 38, 1529–1568, <https://doi.org/https://doi.org/10.1007/s10712-017-9428-0>, 2017.

Bony, S., Schulz, H., Vial, J., and Stevens, B.: Sugar, gravel, fish, and flowers: Dependence of mesoscale patterns of trade-wind clouds on environmental conditions, *Geophysical research letters*, 47, e2019GL085988, 2020.

Bony, S., Lathon, M., Delanoë, J., Coutiris, P., Etienne, J.-C., Aemisegger, F., Albright, A. L., André, T., Bellec, H., Baron, A., Bourdinot, J.-F., Brilouet, P.-E., Bourdon, A., Canonici, J.-C., Caudoux, C., Chazette, P., Cluzeau, M., Cornet, C., Desbios, J.-P., Duchanoy, D.,




Figure A3. Cloud-field response to large-scale cloud-controlling factors and cloud-droplet number perturbations. The standardized β coefficients of the multiple regression analysis for (a) rain-water path $\bar{\mathcal{R}}$, (b) cold-pool fraction $\bar{f}_{cp}$, (c) cold-pool number $\bar{n}_{cp}$, (d) cold-pool size $\bar{s}_{cp}$ (e) liquid-water path $\bar{\mathcal{L}}$, and (f) cloud fraction $\bar{f}_c$, all averaged over the last two days of the LESs of the *Botany* ensemble. The error bars show the 95% confidence interval for each regressor. The larger the distance of the confidence interval with zero, the more significant the corresponding regressor. The p -values of the F -statistic test of all models are smaller than 10^{-15} .

- Flamant, C., Fildier, B., Goubeyre, C., Guiraud, L., Jiang, T., Lainard, C., Le Gac, C., Lendroit, C., Lernoold, J., Perrin, T., Pouvesle, F., Richard, P., Rochetin, N., Salaün, K., Schwarzenboeck, A., Seurat, G., Stevens, B., Totems, J., Touzé-Peiffer, L., Vergez, G., Vial, J., Villiger, L., and Vogel, R.: EUREC⁴A observations from the SAFIRE ATR42 aircraft, *Earth System Science Data*, 14, 2021–2064, <https://doi.org/10.5194/essd-14-2021-2022>, 2022.
- Bretherton, C. and Blossey, P.: Understanding mesoscale aggregation of shallow cumulus convection using large-eddy simulation, *Journal of Advances in Modeling Earth Systems*, 9, 2798–2821, 2017.
- Cesana, G. V. and Del Genio, A. D.: Observational constraint on cloud feedbacks suggests moderate climate sensitivity, *Nature Climate Change*, 11, 213–218, <https://doi.org/10.1038/s41558-020-00970-y>, 2021.
- Colin, M., Sherwood, S., Geoffroy, O., Bony, S., and Fuchs, D.: Identifying the Sources of Convective Memory in Cloud-Resolving Simulations, *Journal of the Atmospheric Sciences*, 76, 947 – 962, <https://doi.org/10.1175/JAS-D-18-0036.1>, 2019.
- Colón-Robles, M., Rauber, R. M., and Jensen, J. B.: Influence of low-level wind speed on droplet spectra near cloud base in trade wind cumulus, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2006GL027487>, 2006.
- Dagan, G., Koren, I., Kostinski, A., and Altaratz, O.: Organization and oscillations in simulated shallow convective clouds, *Journal of Advances in Modeling Earth Systems*, 10, 2287–2299, 2018.

- Denby, L.: Charting the Realms of Mesoscale Cloud Organisation using Unsupervised Learning, arXiv preprint arXiv:2309.08567, 2023.
- Drager, A. J. and van den Heever, S. C.: Characterizing convective cold pools, *Journal of Advances in Modeling Earth Systems*, 9, 1091–1115, <https://doi.org/https://doi.org/10.1002/2016MS000788>, 2017.
- George, G., Stevens, B., Bony, S., Vogel, R., and Naumann, A. K.: Widespread shallow mesoscale circulations observed in the trades, *Nature Geoscience*, 16, 584–589, 2023.
- Gerber, H. E., FRICK, G. M., JENSEN, J. B., and HUDSON, J. G.: Entrainment, Mixing, and Microphysics in Trade-Wind Cumulus, *Journal of the Meteorological Society of Japan. Ser. II*, 86A, 87–106, <https://doi.org/10.2151/jmsj.86A.87>, 2008.
- Glassmeier, F. and Feingold, G.: Network approach to patterns in stratocumulus clouds, *Proceedings of the National Academy of Sciences*, 114, 10 578–10 583, <https://doi.org/https://doi.org/10.1073/pnas.1706495114>, 2017.
- 470 Glassmeier, F., Hoffmann, F., Johnson, J. S., Yamaguchi, T., Carslaw, K. S., and Feingold, G.: Aerosol-cloud-climate cooling overestimated by ship-track data, *Science*, 371, 485–489, <https://doi.org/10.1126/science.abd3980>, 2021.
- Haerter, J. O. and Schlemmer, L.: Intensified cold pool dynamics under stronger surface heating, *Geophysical Research Letters*, 45, 6299–6310, <https://doi.org/https://doi.org/10.1029/2017GL076874>, 2018.
- Haerter, J. O., Böing, S. J., Henneberg, O., and Nissen, S. B.: Circling in on convective organization, *Geophysical Research Letters*, 46, 475 7024–7034, 2019.
- Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., van Kerkwijk, M. H., Brett, M., Haldane, A., del Río, J. F., Wiebe, M., Peterson, P., Gérard-Marchant, P., Sheppard, K., Reddy, T., Weckesser, W., Abbasi, H., Gohlke, C., and Oliphant, T. E.: Array programming with NumPy, *Nature*, 585, 357–362, <https://doi.org/10.1038/s41586-020-2649-2>, 2020.
- 480 Helfer, K. C. and Nuijens, L.: The Morphology of Simulated Trade-Wind Convection and Cold Pools Under Wind Shear, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD035 148, 2021.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Hoyer, S. and Joseph, H.: xarray: N-D labeled Arrays and Datasets in Python, *Journal of Open Research Software*, 5, 485 <https://doi.org/10.5334/jors.148>, 2017.
- Hudson, J. G. and Noble, S.: Low-altitude summer/winter microphysics, dynamics, and CCN spectra of northeastern Caribbean small cumuli, and comparisons with stratus, *Journal of Geophysical Research: Atmospheres*, 119, 5445–5463, <https://doi.org/https://doi.org/10.1002/2013JD021442>, 2014.
- Hunter, J. D.: Matplotlib: A 2D graphics environment, *Computing in Science & Engineering*, 9, 90–95, 490 <https://doi.org/10.1109/MCSE.2007.55>, 2007.
- IPCC AR6, C. .: The Earth’s Energy Budget, Climate Feedbacks and Climate Sensitivity, p. 923–1054, Cambridge University Press, 2023.
- Janssens, M.: Mesoscale Cloud Patterns in the Trade-Wind Boundary Layer, Ph.D. thesis, Wageningen University, ISBN 978-94-6447-829-7, <https://doi.org/https://doi.org/10.18174/635857>, 2023a.
- Janssens, M.: Supporting data for Ch. 7 of "Mesoscale Cloud Patterns in the Trade-Wind Boundary Layer", Zenodo, 495 <https://doi.org/10.5281/zenodo.8089287>, 2023b.
- Janssens, M., Vilà-Guerau de Arellano, J., Scheffer, M., Antonissen, C., Siebesma, A. P., and Glassmeier, F.: Cloud patterns in the trades have four interpretable dimensions, *Geophysical Research Letters*, 48, e2020GL091 001, 2021.

- Janssens, M., De Arellano, J. V.-G., Van Heerwaarden, C. C., De Roode, S. R., Siebesma, A. P., and Glassmeier, F.: Nonprecipitating Shallow Cumulus Convection Is Intrinsically Unstable to Length Scale Growth, *Journal of the Atmospheric Sciences*, 80, 849–870, 2023.
- 500 Janssens, M., Jansson, F., Alinaghi, P., Glassmeier, F., and Siebesma, A. P.: Symmetry in Mesoscale Circulations Explains Weak Impact of Trade Cumulus Self-Organization on the Radiation Budget in Large-Eddy Simulations, *Geophysical Research Letters*, 52, e2024GL112288, <https://doi.org/https://doi.org/10.1029/2024GL112288>, 2025.
- Jansson, F., Janssens, M., Grönqvist, J. H., Siebesma, A. P., Glassmeier, F., Attema, J., Azizi, V., Satoh, M., Sato, Y., Schulz, H., and Kölling, T.: Cloud Botany: Shallow Cumulus Clouds in an Ensemble of Idealized Large-Domain Large-Eddy Simulations of the Trades, *Journal of Advances in Modeling Earth Systems*, 15, e2023MS003796, <https://doi.org/https://doi.org/10.1029/2023MS003796>, e2023MS003796 2023MS003796, 2023.
- 505 Jeevanjee, N. and Romps, D. M.: Convective self-aggregation, cold pools, and domain size, *Geophysical Research Letters*, 40, 994–998, <https://doi.org/https://doi.org/10.1002/grl.50204>, 2013.
- Kazil, J., Vogel, R., Alinaghi, P., Bariteau, L., Bayley, C., Blossey, P., Boeing, S., Chandrakar, K. K., Denby, L. C., Falk, N. M., et al.: Cold Pool Analysis from The Cold Pool Model Intercomparison Project (CP-MIP), in: 105th AMS Annual Meeting, AMS, 2025.
- 510 Lacis, A. A. and Hansen, J.: A parameterization for the absorption of solar radiation in the earth’s atmosphere, *Journal of Atmospheric Sciences*, 31, 118–133, 1974.
- Langhans, W. and Romps, D. M.: The origin of water vapor rings in tropical oceanic cold pools, *Geophysical Research Letters*, 42, 7825–7834, <https://doi.org/https://doi.org/10.1002/2015GL065623>, 2015.
- 515 Li, Z., Zuidema, P., and Zhu, P.: Simulated convective invigoration processes at trade wind cumulus cold pool boundaries, *Journal of the Atmospheric Sciences*, 71, 2823–2841, <https://doi.org/https://doi.org/10.1175/JAS-D-13-0184.1>, 2014.
- Li, Z., Zuidema, P., Zhu, P., and Morrison, H.: The sensitivity of simulated shallow cumulus convection and cold pools to microphysics, *Journal of the Atmospheric Sciences*, 72, 3340–3355, 2015.
- Lochbihler, K., Lenderink, G., and Siebesma, A. P.: Cold pool dynamics shape the response of extreme rainfall events to climate change, *Journal of Advances in Modeling Earth Systems*, 13, e2020MS002306, <https://doi.org/https://doi.org/10.1029/2020MS002306>, 2021.
- 520 Myers, T. A., Scott, R. C., Zelinka, M. D., Klein, S. A., Norris, J. R., and Caldwell, P. M.: Observational constraints on low cloud feedback reduce uncertainty of climate sensitivity, *Nature Climate Change*, 11, 501–507, <https://doi.org/10.1038/s41558-021-01039-0>, 2021.
- Narenpitak, P., Kazil, J., Yamaguchi, T., Quinn, P., and Feingold, G.: From sugar to flowers: A transition of shallow cumulus organization during ATOMIC, *Journal of Advances in Modeling Earth Systems*, 13, e2021MS002619, <https://doi.org/https://doi.org/10.1029/2021MS002619>, 2021.
- 525 Nissen, S. B. and Haerter, J. O.: Circling in on convective self-aggregation, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD035331, 2021.
- Nuijens, L. and Siebesma, A. P.: Boundary layer clouds and convection over subtropical oceans in our current and in a warmer climate, *Current Climate Change Reports*, 5, 80–94, 2019.
- 530 Nuijens, L., Stevens, B., and Siebesma, A. P.: The environment of precipitating shallow cumulus convection, *Journal of the Atmospheric Sciences*, 66, 1962–1979, 2009.
- Quinn, P. K., Thompson, E. J., Coffman, D. J., Baidar, S., Bariteau, L., Bates, T. S., Bigorre, S., Brewer, A., de Boer, G., de Szoeke, S. P., Drushka, K., Foltz, G. R., Intrieri, J., Iyer, S., Fairall, C. W., Gaston, C. J., Jansen, F., Johnson, J. E., Krüger, O. O., Marchbanks, R. D., Moran, K. P., Noone, D., Pezoa, S., Pincus, R., Plueddemann, A. J., Pöhlker, M. L., Pöschl, U., Quinones Melendez, E., Royer, H. M.,
- 535 Szczodrak, M., Thomson, J., Upchurch, L. M., Zhang, C., Zhang, D., and Zuidema, P.: Measurements from the RV *Ronald H. Brown*

- and related platforms as part of the Atlantic Tradewind Ocean-Atmosphere Mesoscale Interaction Campaign (ATOMIC), *Earth System Science Data*, 13, 1759–1790, <https://doi.org/10.5194/essd-13-1759-2021>, 2021.
- Radtke, J., Naumann, A. K., Hagen, M., and Ament, F.: The relationship between precipitation and its spatial pattern in the trades observed during EUREC4A, *Quarterly Journal of the Royal Meteorological Society*, <https://doi.org/10.1002/qj.4284>, 2022.
- 540 Radtke, J., Vogel, R., Ament, F., and Naumann, A. K.: Spatial organisation affects the pathway to precipitation in simulated trade-wind convection, *Geophysical Research Letters*, 50, e2023GL103 579, <https://doi.org/https://doi.org/10.1029/2023GL103579>, 2023.
- Rochetin, N., Hohenegger, C., Touzé-Peiffer, L., and Villefranque, N.: A Physically Based Definition of Convectively Generated Density Currents: Detection and Characterization in Convection-Permitting Simulations, *Journal of Advances in Modeling Earth Systems*, 13, e2020MS002 402, 2021.
- 545 Rotunno, R., Klemp, J. B., and Weisman, M. L.: A theory for strong, long-lived squall lines, *Journal of Atmospheric Sciences*, 45, 463–485, [https://doi.org/https://doi.org/10.1175/1520-0469\(1988\)045<0463:ATFSL>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0469(1988)045<0463:ATFSL>2.0.CO;2), 1988.
- Savic-Jovicic, V. and Stevens, B.: The Structure and Mesoscale Organization of Precipitating Stratocumulus, *Journal of the Atmospheric Sciences*, 65, 1587 – 1605, <https://doi.org/10.1175/2007JAS2456.1>, 2008.
- Schlemmer, L. and Hohenegger, C.: The formation of wider and deeper clouds as a result of cold-pool dynamics, *Journal of the Atmospheric*
- 550 *Sciences*, 71, 2842–2858, <https://doi.org/https://doi.org/10.1175/JAS-D-13-0170.1>, 2014.
- Schneider, T., Teixeira, J., Bretherton, C. S., Brient, F., Pressel, K. G., Schär, C., and Siebesma, A. P.: Climate goals and computing the future of clouds, *Nature Climate Change*, 7, 3–5, 2017.
- Seabold, S. and Perktold, J.: statsmodels: Econometric and statistical modeling with python, in: 9th Python in Science Conference, 2010.
- Seifert, A. and Beheng, K. D.: A double-moment parameterization for simulating autoconversion, accretion and selfcollection, *Atmospheric*
- 555 *research*, 59, 265–281, 2001.
- Seifert, A. and Heus, T.: Large-eddy simulation of organized precipitating trade wind cumulus clouds, *Atmospheric Chemistry and Physics*, 13, 5631–5645, <https://doi.org/https://doi.org/10.5194/acp-13-5631-2013>, 2013.
- Seifert, A., Heus, T., Pincus, R., and Stevens, B.: Large-eddy simulation of the transient and near-equilibrium behavior of precipitating shallow convection, *Journal of Advances in Modeling Earth Systems*, 7, 1918–1937, 2015.
- 560 Snodgrass, E. R., Girolamo, L. D., and Rauber, R. M.: Precipitation Characteristics of Trade Wind Clouds during RICO Derived from Radar, Satellite, and Aircraft Measurements, *Journal of Applied Meteorology and Climatology*, 48, 464 – 483, <https://doi.org/10.1175/2008JAMC1946.1>, 2009.
- Stensrud, D. J., Coniglio, M. C., Davies-Jones, R. P., and Evans, J. S.: Comments on “‘A theory for strong long-lived squall lines’ revisited”, *Journal of the Atmospheric Sciences*, 62, 2989–2996, <https://doi.org/https://doi.org/10.1175/JAS3514.1>, 2005.
- 565 Stephens, G. L., Li, J., Wild, M., Clayson, C. A., Loeb, N., Kato, S., L’ecuyer, T., Stackhouse Jr, P. W., Lebsock, M., and Andrews, T.: An update on Earth’s energy balance in light of the latest global observations, *Nature Geoscience*, 5, 691–696, 2012.
- Stevens, B. and Feingold, G.: Untangling aerosol effects on clouds and precipitation in a buffered system, *Nature*, 461, 607–613, 2009.
- Stevens, B., Bony, S., Farrell, D., Ament, F., Blyth, A., Fairall, C., Karstensen, J., Quinn, P. K., Speich, S., Acquistapace, C., et al.: EUREC 4 a, *Earth System Science Data Discussions*, 2021, 1–78, <https://doi.org/https://doi.org/10.5194/essd-13-4067-2021>, 2021.
- 570 Torri, G. and Kuang, Z.: On cold pool collisions in tropical boundary layers, *Geophysical Research Letters*, 46, 399–407, <https://doi.org/https://doi.org/10.1029/2018GL080501>, 2019.
- Torri, G., Kuang, Z., and Tian, Y.: Mechanisms for convection triggering by cold pools, *Geophysical Research Letters*, 42, 1943–1950, <https://doi.org/https://doi.org/10.1002/2015GL063227>, 2015.

Touzé-Peiffer, L., Vogel, R., and Rochetin, N.: Cold pools observed during EUREC 4 A: Detection and characterization from atmospheric soundings, *Journal of Applied Meteorology and Climatology*, 61, 593–610, 2022.

Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *Journal of Atmospheric Sciences*, 34, 1149 – 1152, [https://doi.org/https://doi.org/10.1175/1520-0469\(1977\)034<1149:TIOPOT>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2), 1977.

Vial, J., Vogel, R., and Schulz, H.: On the daily cycle of mesoscale cloud organization in the winter trades, *Quarterly Journal of the Royal Meteorological Society*, 147, 2850–2873, <https://doi.org/https://doi.org/10.1002/qj.4103>, 2021.

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., Carey, C. J., Polat, İ., Feng, Y., Moore, E. W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E. A., Harris, C. R., Archibald, A. M., Ribeiro, A. H., Pedregosa, F., van Mulbregt, P., and SciPy 1.0 Contributors: SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python, *Nature Methods*, 17, 261–272, <https://doi.org/10.1038/s41592-019-0686-2>, 2020.

Vogel, R., Nuijens, L., and Stevens, B.: The role of precipitation and spatial organization in the response of trade-wind clouds to warming, *Journal of Advances in Modeling Earth Systems*, 8, 843–862, <https://doi.org/https://doi.org/10.1002/2015MS000568>, 2016.

Vogel, R., Konow, H., Schulz, H., and Zuidema, P.: A climatology of trade-wind cumulus cold pools and their link to mesoscale cloud organization, *Atmospheric Chemistry and Physics*, 21, 16 609–16 630, 2021.

Vogel, R., Albright, A. L., Vial, J., George, G., Stevens, B., and Bony, S.: Strong cloud–circulation coupling explains weak trade cumulus feedback, *Nature*, 612, 696–700, 2022.

Waskom, M. L.: seaborn: statistical data visualization, *Journal of Open Source Software*, 6, 3021, <https://doi.org/10.21105/joss.03021>, 2021.

Weisman, M. L. and Rotunno, R.: “A theory for strong long-lived squall lines” revisited, *Journal of the Atmospheric Sciences*, 61, 361–382, [https://doi.org/https://doi.org/10.1175/1520-0469\(2004\)061<0361:ATFSLS>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0469(2004)061<0361:ATFSLS>2.0.CO;2), 2004.

Wes McKinney: Data Structures for Statistical Computing in Python, in: *Proceedings of the 9th Python in Science Conference*, edited by Stéfan van der Walt and Jarrod Millman, pp. 56 – 61, <https://doi.org/10.25080/Majora-92bf1922-00a>, 2010.

Xie, Y. and Liu, Y.: A new approach for simultaneously retrieving cloud albedo and cloud fraction from surface-based shortwave radiation measurements, *Environmental Research Letters*, 8, 044 023, <https://doi.org/10.1088/1748-9326/8/4/044023>, 2013.

Xue, H., Feingold, G., and Stevens, B.: Aerosol effects on clouds, precipitation, and the organization of shallow cumulus convection, *Journal of the Atmospheric Sciences*, 65, 392–406, <https://doi.org/https://doi.org/10.1175/2007JAS2428.1>, 2008.

Yamaguchi, T., Feingold, G., and Kazil, J.: Aerosol-cloud interactions in trade wind cumulus clouds and the role of vertical wind shear, *Journal of Geophysical Research: Atmospheres*, 124, 12 244–12 261, 2019.

Zhang, Y., Stevens, B., and Ghil, M.: On the diurnal cycle and susceptibility to aerosol concentration in a stratocumulus-topped mixed layer, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 131, 1567–1583, <https://doi.org/https://doi.org/10.1256/qj.04.103>, 2005.

Zuidema, P., Xue, H., and Feingold, G.: Shortwave radiative impacts from aerosol effects on marine shallow cumuli, *Journal of the atmospheric sciences*, 65, 1979–1990, <https://doi.org/https://doi.org/10.1175/2007JAS2447.1>, 2008.

Zuidema, P., Li, Z., Hill, R. J., Bariteau, L., Rilling, B., Fairall, C., Brewer, W. A., Albrecht, B., and Hare, J.: On trade wind cumulus cold pools, *Journal of the Atmospheric Sciences*, 69, 258–280, <https://doi.org/https://doi.org/10.1175/JAS-D-11-0143.1>, 2012.

Zuidema, P., Torri, G., Muller, C., and Chandra, A.: A survey of precipitation-induced atmospheric cold pools over oceans and their interactions with the larger-scale environment, *Surveys in Geophysics*, 38, 1283–1305, <https://doi.org/https://doi.org/10.1007/s10712-017-9447-x>, 2017.