



# Nitrogen Fixation in Arctic Coastal Waters (Qeqertarsuaq, West Greenland): Influence of Glacial Melt on Diazotrophs, Nutrient Availability, and Seasonal Blooms

Schlangen Isabell<sup>1</sup>, Leon-Palmero Elizabeth<sup>1,2</sup>, Moser Annabell<sup>1</sup>, Xu Peihang<sup>1</sup>, Laursen Erik<sup>1</sup>, and Löscher Carolin R.<sup>1,3</sup>

<sup>1</sup>Nordcee, Department of Biology, University of Southern Denmark, Campusvej 55, 5230 Odense M, Denmark

<sup>2</sup>Department of Geosciences, Princeton University, Princeton, New Jersey

<sup>3</sup>DIAS, University of Southern Denmark, Odense, Denmark

**Correspondence:** Carolin R. Löscher (cloescher@biology.sdu.dk)

**Abstract.** The Arctic Ocean is undergoing rapid transformation due to climate change, with decreasing sea ice contributing to a predicted increase in primary productivity. A critical factor determining future productivity in this region is the availability of nitrogen, a key nutrient that often limits biological growth in Arctic waters. The fixation of dinitrogen (N<sub>2</sub>) gas, a biological process mediated by diazotrophs, not only supplies new nitrogen to the ecosystem but also plays a central role in shaping the biological productivity of the Arctic. Historically it was believed to be limited to oligotrophic tropical and subtropical oceans, Arctic N<sub>2</sub> fixation has only garnered significant attention over the past decade, leaving a gap in our understanding of its magnitude, the diazotrophic community, and potential environmental drivers. In this study, we investigated N<sub>2</sub> fixation rates and the diazotrophic community in Arctic coastal waters, using a combination of isotope labeling, genetic analyses and biogeochemical profiling, in order to explore its response to glacial meltwater, nutrient availability and its impact on primary productivity. Here we show N<sub>2</sub> fixation rates ranging from 0.16 to 2.71 nmol N L<sup>-1</sup> d<sup>-1</sup>, to be notably higher than those observed in many other oceanic regions, suggesting a previously unrecognized significance of N<sub>2</sub> fixation in these high-latitude waters. The diazotrophic community is predominantly composed of UCYN-A. We found highest N<sub>2</sub> fixation rates co-occurring with maximum chlorophyll *a* concentrations and primary production rates at a station in the Vaigat Strait close impacted by glacier meltwater inflow, possibly providing otherwise limiting nutrients. Our findings illustrate the importance of N<sub>2</sub> fixation in an environment previously not considered important for this process and provide insights into its response to the projected melting of the polar ice cover.

## 1 Introduction

Nitrogen is a key element for life and often acts as a growth-limiting factor for primary productivity (Gruber and Sarmiento, 1997; Gruber, 2004; Gruber and Galloway, 2008). Despite nitrogen gas (N<sub>2</sub>) making up approximately 78% of the atmosphere, it remains inaccessible to most marine life forms. Diazotrophs, which are specialized bacteria and archaea, have the ability to convert N<sub>2</sub> into biologically available nitrogen, facilitated by the nitrogenase enzyme complex carrying out the process of



biological nitrogen fixation ( $N_2$  fixation) (Capone and Carpenter (1982)). Despite the fact that these organisms are highly specialized and  $N_2$  fixation is energetically demanding, the ability to carry out this process is widespread amongst prokaryotes. However, it is controlled by several factors such as temperature, light, nutrients and trace metals such as iron and molybdenum (Sohm et al., 2011; Tang et al., 2019). Oceanic  $N_2$  fixation is the major source of nitrogen to the marine system (Karl et al., 2002; Gruber and Sarmiento, 1997), thus, diazotrophs determine the biological productivity of our planet (Falkowski et al. (2008), impact the global carbon cycle and the formation of organic matter (Galloway et al., 2004; Zehr and Capone, 2020). Traditionally it has been believed that the distribution of diazotrophs was limited to warm and oligotrophic waters (Buchanan et al., 2019; Sohм et al., 2011; Luo et al., 2012) until putative diazotrophs were identified in the central Arctic Ocean and Baffin Bay (Farnelid et al., 2011; Damm et al., 2010). First rate measurements have been reported for the Canadian Arctic by Blais et al. (2012) and recent studies have reported rate measurements in adjacent seas (Harding et al., 2018; Sipler et al., 2017; Shiozaki et al., 2017, 2018), drawing attention to cold and temperate waters as significant contributors to the global nitrogen budget through diverse organisms.

$N_2$  fixation is performed by diverse group of cyanobacteria as well as by non-cyanobacteria diazotrophs (NCDs). UCYN-A has been described as the dominant active  $N_2$  fixing cyanobacterial diazotroph in those waters (Harding et al. (2018)), while other cyanobacteria have only occasionally been reported (Díez et al., 2012; Fernández-Méndez et al., 2016; Blais et al., 2012). Recent studies found that the majority of the arctic marine diazotrophs are NCDs and those may contribute significantly to  $N_2$  fixation in the Arctic Ocean (Shiozaki et al., 2018; Fernández-Méndez et al., 2016; Harding et al., 2018; Von Friesen and Riemann, 2020). Still, studies on the Arctic diazotroph community remain scarce, leaving Arctic environments poorly understood regarding  $N_2$  fixation. Shao et al. (2023) note the impossibility of estimating Arctic  $N_2$  fixation rates due to the sparse spatial coverage, which currently represents only approximately 1 % of the Arctic Ocean. Increasing data coverage in future studies will aid in better constraining the contribution of  $N_2$  fixation to the global oceanic nitrogen budget (Tang et al. (2019)).

Additionally, the Arctic ecosystem is undergoing significant changes driven by rising temperatures and the consequent reduction in sea ice extent due to accelerated melting, which is predicted to increase in the future (Arrigo et al., 2008; Hanna et al., 2008; Haine et al., 2015). These severe climate shifts have stimulated primary productivity in the Arctic by 57 % from 1998 to 2018, thereby elevating nutrient demands in the Arctic Ocean (Ardyna and Arrigo, 2020; Arrigo and van Dijken, 2015; Lewis et al., 2020). This increase can be attributed to the extension of the phytoplankton growing seasons and the expansion of ice-free areas available for phytoplankton growth (Arrigo et al. (2008)). The Greenland Ice Sheet is strongly affected by climate change and the waters of Baffin Bay have experienced a substantial sea surface temperature (SST) increase of 47.4 % along with a significant increase in chlorophyll *a* (Chl *a*) concentration of 26.4 % over the last two decades (1998-2018) (Lewis et al. (2020)). Coastal sites are particularly impacted by melting, receiving glacial runoff enriched with nutrients and trace elements triggering phytoplankton blooms and altering near-shore biogeochemical cycling (Ardyna and Arrigo, 2020; Arrigo et al., 2017; Hendry et al., 2019; Bhatia et al., 2013). Given the changes, there is an urgency to explore the role of  $N_2$  fixation in shaping the response of the Arctic ecosystem to these environmental changes. While the general magnitude of  $N_2$  fixation is suspected to have a substantial impact (Sipler et al. (2017)), the complexity of Arctic biogeochemical processes necessitates further studies and broader spatial and temporal investigations to facilitate robust predictions. The question of whether primary

production in the Arctic will be limited by nitrogen availability and the extent to which species will adapt to these conditions remains unknown and needs to be addressed. This study aims to contribute to the understanding of N<sub>2</sub> fixation dynamics and its implications for ecosystem productivity with the rapidly evolving Arctic Ocean.

60 We explored the diazotroph diversity in combination with N<sub>2</sub> fixation rate measurements, to elucidate the importance of this process in the Arctic ecosystem. We hope that understanding the dynamics of N<sub>2</sub> fixation and its impact on the ecosystem productivity can inform predictions and help managing the consequences of ongoing environmental changes in the Arctic Ocean. Our study has been carried out in Disko Bay (Qeqertarsuaq), which can serve as a model for Arctic coastal systems influenced by large meltwater runoff and thus potentially an addition of high levels of iron and nutrients, both of which have the ability to  
65 affect N<sub>2</sub> fixation (Lewis et al., 2020; Bhatia et al., 2013).

## 2 Material and methods

### 2.1 Seawater sampling

The research expedition was conducted from August 16 to 26 in 2022 aboard the Danish military vessel P540 within the waters of Qeqertarsuaq, located in the western region of Greenland (Kalaallit Nunaat). A comprehensive sampling strategy was employed at 10 stations (Fig. 1), covering the surface to a depth of 100 m. The sampled parameters included water characteristics,  
70 such as nutrient concentrations, chl *a*, particulate organic carbon (POC) and nitrogen (PON), molecular samples for nucleic acid extractions (DNA), dissolved inorganic carbon (DIC) as well as CTD sensor data. At three selected stations (3,7,10) N<sub>2</sub> fixation and primary production rates were quantified through concurrent incubation experiments.

Samples for nutrient analysis, nitrate (NO<sub>3</sub><sup>-</sup>), nitrite (NO<sub>2</sub><sup>-</sup>) and phosphate (PO<sub>4</sub><sup>3-</sup>) were taken in triplicates, filtered through  
75 a 0.22 μm syringe filter (Avantor VWR® Radnor, Pa, USA) and stored at -20 °C until further analysis. Concentrations were spectrophotometrically determined (Thermo Scientific, Genesys 10S UV-VIS spectrophotometer) following the established protocols of Murphy and Riley (1962) for PO<sub>4</sub><sup>3-</sup>; García-Robledo et al. (2014) for NO<sub>3</sub><sup>-</sup> & NO<sub>2</sub><sup>-</sup>. Chl *a* samples were filtered onto 47 mm ø GF/F filters (GE Healthcare Life Sciences, Whatman, USA), placed into darkened 15 mL LightSafe centrifuge tubes (Merck, Rahway, NJ, USA) and were subsequently stored at -20 °C until further analysis. To determine the Chl *a* concentration, the samples were immersed in 8 mL of 90 % acetone overnight at 5 °C. Subsequently, 1 mL of the resulting solution  
80 was transferred to a 1.5 mL glass vial (Mikrolab Aarhus A/S, Aarhus, Denmark) the following day and subjected to analysis using the Triology® Fluorometer (Model #7200-00) equipped with a Chl *a* in vivo blue module (Model #7200-043, both Turner Designs, San Jose, CA, USA). Measurements of serial dilutions from a 4 mg L<sup>-1</sup> stock standard and 90 % acetone (serving as blank) were performed to calibrate the instrument. In addition, measurements of a solid-state secondary standard  
85 were performed every 10 samples. Water (1 L) water from each depth was filtered for the determination of POC and PON concentrations, as well as natural isotope abundance (δ <sup>13</sup>C POC / δ <sup>15</sup>N PON) using 47 mm ø, 0.7 μm nominal pore size precombusted GF/F filter (GE Healthcare Life Sciences, Whatman, USA), which were subsequently stored at -20 °C until further analysis. Seawater samples for DNA were filtered through 47 mm ø, 0.22 μm MCE membrane filter (Merck, Millipore Ltd., Ireland) for a maximum of 20 minutes, employing a gentle vacuum (200 mbar). The filtered volumes varied depending



90 on the amount of material captured on the filter, ranging from 1.3 L to 2 L, with precise measurements recorded. The filters were promptly stored at -20 °C on the ship and moved to -80 °C upon arrival to the lab until further analysis. Seawater (40 ml) was filtered through a 0.22  $\mu\text{m}$  syringe filter (Avantor VWR® Radnor, Pa, USA) and stored at 4 °C in an amber glass vial, sealed with closed caps, affixed with a PTFE-faced silicon liner (Thermo Fisher Scientific, Waltham, MA, USA) for subsequent DIC measurements in the laboratory using an AS-C5 DIC analyzer (ApolloSciTech, Newark, Delaware, USA) equipped with a laser-based CO<sub>2</sub> detector. Sample analysis was carried out following the manufacturer's guidelines and the use of a certified seawater reference (Batch 187, Scripps Institution of Oceanography, University of California, San Diego, USA).  
95 To achieve detailed vertical profiles, a conductivity-temperature-depth-profiler (CTD, Seabird X) equipped with supplementary sensors for dissolved oxygen (DO), photosynthetic active radiation (PAR), and fluorescence (Fluorometer) was manually deployed. In the same manner, discrete water samples were obtained using a 10 L Niskin bottle, manually lowered with a hand  
100 winch to five distinct depths (Surface, 5, 25, 50, 100 m). These systematic and multifaceted sampling methodologies provide a robust dataset for a comprehensive analysis of the hydrographic conditions in Qeqertarsuaq.

## 2.2 Nitrogen fixation and primary production

Water samples were collected at three distinct depths (0, 25 and 50 m) for the investigation of N<sub>2</sub> fixation rates and primary production rates, encompassing the euphotic zone, chlorophyll maximum, and a light-absent zone. Three incubation stations  
105 (Fig. 2: station 3, 7, 10) were chosen, in a way to cover the variability of the study area. This strategic sampling aimed to capture a gradient of the water column with varying environmental conditions, relevant to the aim of the study. N<sub>2</sub> fixation rates were assessed through triplicate incubations employing the modified <sup>15</sup>N-N<sub>2</sub> dissolution technique after Großkopf et al. (2012) and Mohr et al. (2010).

To ensure minimal contamination, 2.3 L glass bottles (Schott-Duran, Wertheim, Germany) underwent pre-cleaning and acid  
110 washing before being filled with seawater samples. Oxygen contamination during sample collection was mitigated by gently and bubble-free filling the bottles from the bottom, allowing the water to overflow. Each incubation bottle received a 100 mL amendment of <sup>15</sup>N-N<sub>2</sub> enriched seawater (98 %, Cambridge Isotope Laboratories, Inc., USA) achieving an average dissolved N<sub>2</sub> isotope abundance (<sup>15</sup>N atom %) of  $3.90 \pm 0.02$  atom % (mean  $\pm$  SD). Additionally, 1 mL of H<sup>13</sup>CO<sub>3</sub> (1g/50 mL) (Sigma-Aldrich, Saint Louis Missouri US) was added to each incubation bottle, roughly corresponding to 10 atom % enrichment and  
115 thus measurements of primary production and N<sub>2</sub> fixation were conducted in the same bottle. Following the addition of both isotopic components, the bottles were closed airtight with septa-fitted caps and incubated for 24 hours on-deck incubators with a continuous surface seawater flow. These incubators, partially shaded to simulate in situ photosynthetically active radiation (PAR) conditions, aimed to replicate environmental parameters experienced at the sampled depths. Control incubations utilizing atmospheric air served as controls to monitor any natural changes in  $\delta^{15}\text{N}$  not attributable to <sup>15</sup>N-N<sub>2</sub> addition. These control  
120 incubations were conducted using the dissolution method, like the <sup>15</sup>N-N<sub>2</sub> enrichment experiments, but with the substitution of atmospheric air instead of isotopic tracer.

After the incubation period, subsamples for nutrient analysis were taken from each incubation sample, and the remaining content was subjected to the filtration process and were gently filtered (200 mbar) onto precombusted GF/F filters (Advantec,



47 mm  $\phi$ , 0.7  $\mu\text{m}$  nominal pore size). This step ensured a comprehensive examination of both nutrient dynamics and the  
125 isotopic composition of the particulate pool in the incubated samples. Samples were stored at  $-20\text{ }^{\circ}\text{C}$  until further analysis.  
Upon arrival in the lab, the filters were dried at  $60\text{ }^{\circ}\text{C}$  and to eliminate particulate inorganic carbon, subsequently subject to  
acid fuming during which they were exposed to concentrated hydrochloric acid (HCL) vapors overnight in a desiccator. After  
undergoing acid treatment, the filters were carefully dried, then placed into tin capsules and pelletized for subsequent analysis.  
The determination of POC and PON, as well as isotopic composition ( $\delta^{13}\text{C}$  POC /  $\delta^{15}\text{N}$  PON), was carried out using an  
130 elemental analyzer (Flash EA, ThermoFisher, USA) connected to a mass spectrometer (Delta V Advantage Isotope Ratio MS,  
ThermoFisher, USA) with the ConFlo IV interface. This analytical setup was applied to all filters. These values, derived from  
triplicate incubation measurements, exhibited no omission of data points or identification of outliers. Final rate calculations for  
 $\text{N}_2$  fixation rates were performed after Mohr et al. (2010) and primary production rates after Slawyk et al. (1977).

### 2.3 Molecular methods

135 The filters were flash-frozen in liquid nitrogen, crushed and DNA was extracted using the Qiagen DNA/RNA AllPrep Kit (Qi-  
agen, Hildesheim, DE), following the procedure outlined by the manufacturer. The concentration and quality of the extracted  
DNA was assessed spectrophotometrically using a MySpec spectrofluorometer (VWR, Darmstadt, Germany). The prepara-  
tion of the metagenome library and sequencing were performed by BGI (China). Sequencing libraries were generated using  
MGIEasy Fast FS DNA Library Prep Set following the manufacturer's protocol. Sequencing was conducted with 2x150bp  
140 on a DNBSEQ-G400 platform (MGI). SOAPnuke1.5.5 (Chen et al. (2018)) was used to filter and trim low quality reads and  
adaptor contaminants from the raw sequence reads, as clean reads. In total, fifteen metagenomic datasets were produced with  
an average of 9.6G bp per sample.

#### 2.3.1 Metagenomic De Novo assembly, gene prediction, and annotation

Megahit v1.2.9 (Li et al. (2015)) was used to assemble clean reads for each dataset with its minimum contig length as 500.  
145 Prodigal v2.6.3 (Hyatt et al. (2010)) with the setting of “-p meta” was then used to predict the open reading frames (ORFs) of  
the assembled contigs. ORFs from all the available datasets were filtered ( $>100\text{bp}$ ), dereplicated and merged into a catalog of  
non-redundant genes using cd-hit-est ( $>95\%$  sequence identity) (Fu et al. (2012)). Salmon v1.10.0 (Patro et al. (2017)) with  
the “- meta” option was employed to map clean reads of each dataset to the catalog of non-redundant genes and generate the  
GPM (genes per million reads) abundance. Eggnog mapper v2.1.12 (Cantalapiedra et al. (2021)) was then performed to assign  
150 KEGG Orthology (KO) and identify specific functional annotation for the catalog of non-redundant genes. The marker genes,  
*nifDK* (K02586, K02591 nitrogenase molybdenum-iron protein alpha/beta chain), *nifH* (K02588, nitrogenase iron protein),  
were used for the evaluation of microbial potential of  $\text{N}_2$  fixation. *RbcL* (K01601, ribulose-bisphosphate carboxylase large  
chain) and *psbA* (K02703, photosystem II P680 reaction center D1 protein) were selected to evaluate the microbial potential  
of carbon fixation and photosynthesis, respectively.




## 155 3 Results and discussion

### 3.1 Hydrographic conditions in Qeqertarsuaq (Disco Bay) and Sullorsuaq (Vaigat) Strait

Disko Bay (Qeqertarsuaq) is located along the west coast of Greenland (Kalaallit Nunaat) at approximately 69 °N (Figure 1), and is strongly influenced by the West Greenland Current (WGC) which is associated with the broader Baffin Bay Polar Waters (BBPW) (Mortensen et al., 2022; Hansen et al., 2012). The WGC does not only significantly shape the hydrographic conditions within the bay but also plays an important role in the larger context of Greenland Ice Sheet melting (Mortensen et al. (2022)).  
160 Central to the hydrographic system of the Qeqertarsuaq area is the Jakobshavn Isbræ, which is the most productive glacier in the northern hemisphere and believed to drain about 7 % of the Greenland Ice Sheet and thus contributes substantially to the water influx into the Qeqertarsuaq (Holland et al. (2008)). A predicted increased inflow of warm subsurface water, originating from North Atlantic waters, has been suggested to further affect the melting of the Jakobshavn Isbræ and thus adds another  
165 layer of complexity to this dynamic system (Holland et al., 2008; Hansen et al., 2012).

The hydrographic conditions in Qeqertarsuaq have a significant influence on biological processes, nutrient availability, and the




**Figure 1.** Map of Greenland (Kalaallit Nunaat) with indication of study area (red box), on the left. Interpolated distribution of Sea Surface Salinity (SSS) values with corresponding isosurface lines and indication of 10 sampled stations (normal stations in black, incubation stations in blue), in the middle. Scatterplot of the conservative temperature and salinity for all station data. The plot is used for the identification of the main water masses within the study area. Isopycnals ( $\text{kg m}^{-3}$ ) are depicted in grey lines, on the right. Figures were created in Ocean Data View (ODV) (Schlitzer (2022)).

broader marine ecosystem (Munk et al., 2015; Hendry et al., 2019; Schiøtt, 2023).

During our survey, we found very heterogenous hydrographic conditions at the different stations across Qeqertarsuaq (Fig. 1 & Fig. 2). The three selected stations for  $\text{N}_2$  fixation analysis (stations 3, 7, and 10) were strategically chosen to capture the spatial



170 variability of the area. Surface salinity and temperature measurements at these stations indicate the influence of freshwater  
input. The surface temperature exhibit a range of 4.5 to 8 °C, while surface salinity varies between 31 and 34, as illustrated  
in Fig. 1. The profiles sampled during our survey extend to a maximum depth of 100 m. Comparison of temperature/salinity  
(T/S) plots with recent studies suggests the presence of previously described water masses, including Warm Fjord Water  
(WFjW) and Cold Fjord Water (CFjW) with an overlaying surface glacial meltwater runoff. Those water masses are defined  
175 with a density range of  $27.20 \leq \sigma_\theta \leq 27.31$  but different temperature profiles. Thus water masses can be differentiated by  
their temperature within the same density range (Gladish et al. (2015)). Other water masses like upper subpolar mode water  
(uSPMW), deep subpolar mode water (dSPMW) and Baffin Bay polar Water (BBPW) which has been identified in the Disko  
Bay (Qeqertarsuaq) before, cannot be identified from this data and may be present in deeper layers (Mortensen et al., 2022;  
Sherwood et al., 2021; Myers and Ribergaard, 2013; Rysgaard et al., 2020). The temperature and salinity profiles across the 10



**Figure 2.** Profiles of temperature (°C), salinity and Chl *a* ( $\text{mg m}^{-3}$ ) across stations 1 to 10 with depth (m). Stations 3, 7, and 10 are highlighted in red, blue, and yellow, respectively, to emphasize incubation stations. Figures were created in Ocean Data View and R-Studio (Schlitzer (2022)).

180 stations in the study area show distinct stratification and variability, which is represented through the three incubation stations  
(highlighted stations 3, 7, and 10 in Fig. 2). They display varying degrees of stratification and mixing, with notable differences  
in the salinity and temperature profiles. Station 3 and station 7 exhibit clear stratification in both temperature and salinity  
marked by the presence of thermoclines and haloclines. These features suggest significant freshwater input influenced by local  
weather conditions and climate dynamics, like surface heat absorption. In contrast station 10 shows more homogeneous salinity  
and temperature throughout the water column, indicative of well-mixed conditions. This uniformity is likely influenced by the  
185 regional circulation pattern and partial upwelling (Hansen et al., 2012; Krawczyk et al., 2022). The distinct characteristics  
observed at station 10, as illustrated in the surface plot (Fig. 1), show an elevated salinity and colder temperatures compared



to the other stations. This feature suggests upwelling of deeper waters along the shallower shelf, likely influenced by the seafloor shallowing off the coast of station 10, which acts as a barrier and disrupts typical circulation. The presence of water masses forced to the surface due to this topographical feature may explain the observed properties at station 10. Furthermore, the variability in temperature and salinity conditions between stations, particularly in relation to topography, aligns with the findings of Krawczyk et al. (2022). Their description of the bathymetry in Qeqertarsuaq, featuring depths ranging from ca. 50 to 900 m, suggests its impact on turbulent circulation patterns, leading to the mixing of different water masses. Evident variability in oceanographic conditions that can be observed throughout the study area occurs particularly along characteristic topographical features like steep slopes, canyons, and shallower areas. The summer melting of sea ice and glaciers introduces freshwater influxes that create distinct vertical and horizontal gradients in salinity and temperature in the Qeqertarsuaq area Hansen et al. (2012). Additionally, the accelerated melting of the Jakobshavn Isbrae, influenced by the warmer inflow from the West Greenland Intermediate Current (WGIC), further alters the hydrographic conditions. Recent observations indicate significant warming and shoaling of the WGIC, potentially enabling it to overcome the sill separating the Illulissat Fjord from the Qeqertarsuaq area (Hansen et al., 2012; Holland et al., 2008; Myers and Ribergaard, 2013). This shift intensifies glacier melting, driving substantial changes in the local ecological dynamics (Ardyna et al., 2014; Arrigo et al., 2008; Bhatia et al., 2013).

### 3.2 Elevated N<sub>2</sub> fixation rates might play a role in nutrient dynamics and bloom development

We quantified N<sub>2</sub> fixation rates within the waters of Qeqertarsuaq, spanning from the surface to a depth of 50 m (Table 1). The rates ranged from 0.16 to 2.71 nmol N L<sup>-1</sup> d<sup>-1</sup> with all rates surpassing the detection limit. Our findings represent rates at the upper range of those observed in the Arctic Ocean. Previous measurements in the region have been limited, with only one study in Baffin Bay by Blais et al. (2012), reporting rates of 0.02 nmol N L<sup>-1</sup> d<sup>-1</sup>, which are 1-2 orders of magnitude lower than our observations. Compared to other European Arctic waters, our rates at the surface and at 25 m water depth fall within the reported range for Arctic estuarine stations (1.04 to 1.87 nmol N L<sup>-1</sup> d<sup>-1</sup>, (SD ± 0.76 to 1.19) and marine stations (0.11 to 0.12 nmol N L<sup>-1</sup> d<sup>-1</sup>, (SD ± 0.09 to 0.09) (Blais et al. (2012)). However, we observed some of the highest rates reported so far, particularly at the surface. Simultaneous primary production rate measurements ranged from 0.07 to 3.79 μmol N L<sup>-1</sup> d<sup>-1</sup>, with the highest rates observed at station 7 and generally higher values in the surface layers. Employing Redfield stoichiometry, the measured N<sub>2</sub> fixation rates accounted for 0.47 to 2.6 % (averaging 1.57 %) of primary production at our stations. The relatively modest contribution to primary production suggests that N<sub>2</sub> fixation may not exert a substantial influence on the productivity of these waters during the time of the sampling. Rather, our N<sub>2</sub> fixation rates suggest primary production to depend mostly on additional nitrogen sources including regenerated, meltwater or land based sources.

The N:P ratio, calculated as DIN to DIP, indicates a deficit in N for primary production based on Redfield stoichiometry (Fig. 3). This aligns with findings presented by Jensen et al. (1999) and Tremblay and Gagnon (2009), who observed a similar nitrogen limitation in this region. Such biogeochemical conditions would be expected to generate a niche for N<sub>2</sub> fixing organisms (Sohm et al. (2011)). While N<sub>2</sub> fixation did not chiefly sustain primary production during our sampling campaign, we hypothesize that the relatively high N<sub>2</sub> fixation rates observed may play a role in bloom dynamics. As nitrogen availability decreases





during a bloom, it may provide a niche for  $N_2$  fixation, potentially extending the productive period of the bloom (Reeder et al. (2021)). Satellite data indicates that a fall bloom began in early August, following the annual spring bloom, as described by Ardyna et al. (2014). This double bloom situation may be driven by increased melting and the subsequent input of bioavailable nutrients and iron (Fe) from meltwater runoff (Arrigo et al., 2017; Hopwood et al., 2016; Bhatia et al., 2013). The meltwater from the Greenland Ice Sheet is a significant source of Fe (Bhatia et al., 2013; Hawkings et al., 2015, 2014), which is a limiting factor especially for diazotrophs (Sohm et al. (2011)). Consequently, it is possible that nutrients and Fe from the Isbræ glacier introduced into the Qeqertarsuaq are promoting a bloom and further provide a niche for diazotrophs to thrive (Arrigo et al. (2017)).

230

**Table 1.**  $N_2$  fixation ( $\text{nmol N L}^{-1} \text{d}^{-1}$ ), standard deviation (SD), primary productivity (PP;  $\mu\text{mol C L}^{-1} \text{d}^{-1}$ ), SD, percentage of estimated new primary productivity (% New PP) sustained by  $N_2$  fixation, dissolved inorganic nitrogen compounds ( $\text{NO}_x$ ), phosphorus ( $\text{PO}_4$ ), and the molar nitrogen-to-phosphorus ratio (N:P) at stations 3, 7, and 10.


Station (no.)	Depth (m)	$N_2$ fixation ( $\text{nmol N L}^{-1} \text{d}^{-1}$ )	SD ( $\pm$ )	Primary Productivity ( $\mu\text{mol C L}^{-1} \text{d}^{-1}$ )	SD ( $\pm$ )	% New PP (%)	$\text{NO}_x$ ( $\mu\text{mol L}^{-1} \text{d}^{-1}$ )	$\text{PO}_4$ ( $\mu\text{mol L}^{-1} \text{d}^{-1}$ )	N:P (mol:mol)
3	0	1.20	0.21	0.466	0.08	1.71	0	0	0
3	25	1.88	0.11	0.588	0.04	2.11	0	0.70	0
3	50	0.29	0.01	0.209	0.00	0.91	0.33	1.48	0.22
7	0	2.49	0.44	0.63	0.20	2.60	0	0	0
7	25	2.71	0.22	3.79	2.45	0.47	0	0.45	0
7	50	0.53	0.24	0.33	0.36	1.08	0	0.97	0
10	0	1.48	0.12	0.74	0.15	1.33	0	0	0
10	25	0.31	0.01	0.29	0.07	0.73	0	0	0
10	50	0.16	0	0.07	0.07	1.40	0	0	0

A near-Redfield stoichiometry in POC:PON 3 indicates that the particulate organic matter (POM) is freshly derived from an ongoing bloom. However, the absence of  $\text{NO}_x$  (with the exception of one station) and the observed low N:P ratios suggest that any available nitrogen from earlier phases of the bloom has likely been depleted. This could create a niche for  $N_2$  fixation as a supplementary nitrogen source, potentially supporting continued production during this stage of the bloom. The onset and development of the bloom would be expected to lead to high nitrogen demands and intense competition for nitrogen sources. Notably, despite the apparent balance in the POM pool, the N ratio indicates strong nitrogen depletion and nutrient exhaustion within the ecosystem. This deficiency can be partly alleviated by  $N_2$  fixation, providing possibly increasing amounts of nitrogen over the course of the bloom. Moreover, DIP is generally limited in the environment (Table 1); however, some organisms may still access it through luxury phosphorus uptake, storing excess phosphate when it is sporadically available. A recent study

235



240 by Laso Perez et al. (2024) documented changes in microbial community composition during an Arctic bloom, focusing on nitrogen cycling. They observed a shift from chemolithotrophic to heterotrophic organisms throughout the summer bloom and noted increased activity to compete for various nitrogen sources. However, no *nifH* gene copies, indicative of nitrogen-fixing organisms, were found in their dataset based on metagenome-assembled genomes (MAGs). This is not unexpected due to the classically low abundance of diazotrophs in marine microbial communities which has often been described (Turk-Kubo et al.,  
245 2015; Farnelid et al., 2019). Given the high productivity and metabolic activity observed in Qeqertarsuaq during a similar bloom period, the detected diazotrophs (Section 3.3) may play a more significant role than previously thought. Across the 10



**Figure 3.** The POC/PON and DIN/DIP ratios at all 10 stations. The red line represents the Redfield ratios of POC/PON (106:16) and DIN/DIP (16:1).

stations there is considerable variability in POC and PON concentrations (Fig. 3). PON concentrations range from 0.5 µmol N L<sup>-1</sup> to 4.0 µmol N L<sup>-1</sup> (n=124), while POC concentrations range from 2.5 µmol C L<sup>-1</sup> to 32.6 µmol C L<sup>-1</sup> (n=144). The highest concentrations for both PON and POC were observed at station 7 at a depth of 25 m and coincide with the highest reported  
250 N<sub>2</sub> fixation rate (Figure Appendix A2 & A3). Generally, POC and PON concentrations decrease with depth, peaking at the deep chl *a* maximum (DCM), identified between 15 to 30 m across all stations. The variability in chl *a* concentrations indicates differences in phytoplankton abundance among the stations, with concentrations ranging between 0 to 0.42 mg m<sup>-3</sup>. Excluding station 7, which exhibited the highest chl *a* concentration at the DCM (1.51 mg m<sup>-3</sup>). Tang et al. (2019) have found that N<sub>2</sub> fixation measurements strongly correlated to satellite estimates of chl *a* concentrations and thus may be an explanation for the  
255 presented N<sub>2</sub> fixation rates. The elevated concentration of chl *a* likely result from a local phytoplankton bloom induced by meltwater outflow from the Isbræ glacier and sea ice melting (Arrigo et al., 2017; Wang et al., 2014). This can also be seen



from satellite images (Appendix:A1). This study's findings are in agreement with prior reports of analogous blooms occurring in the region (Fox and Walker, 2022; Jensen et al., 1999).

### 3.3 UCYN-A might contribute to N<sub>2</sub> fixation during a diatom bloom


260 In our metagenomic analysis, we filtered the *nifH*, *nifD*, *nifK* genes, which code for the nitrogenase enzyme responsible for catalyzing N<sub>2</sub> fixation. We could identify sequences related to UCYN-A, which dominated the sequence pool of diazotrophs, particularly in the upper water masses (0 to 5 m) (Fig. 4). UCYN-A, a unicellular cyanobacterial symbiont, has a cosmopolitan distribution and is thought to substantially contribute to global N<sub>2</sub> fixation, as documented by (Martínez-Pérez et al., 2016; Tang et al., 2019). Due to the lack of genes such as those encoding Photosystem II and Rubisco, UCYN-A plays a significant  
265 role within the host cell and participates in fundamental cellular processes. Consequently it has evolved to become a closely integrated component of the host cell. Very recent findings demonstrate that UCYN-A imports proteins encoded by the host genome and has been described as an early form of N<sub>2</sub> fixing organelle termed a "Nitroplast" (Coale et al. (2024)).

Previous investigations document that they are critical for primary production, supplying up to 85% of the fixed nitrogen to their haptophyte host (Martínez-Pérez et al. (2016)). In addition to its high contribution to primary production, studies have shown  
270 that UCYN-A in high latitude waters fix similar amounts of N<sub>2</sub> per cell as in the tropical Atlantic Ocean, even in nitrogen-replete waters (Harding et al., 2018; Shiozaki et al., 2020; Martínez-Pérez et al., 2016; Krupke et al., 2015; Mills et al., 2020). However, estimating their contribution to N<sub>2</sub> fixation in our study is challenging, particularly since we detected cyanobacteria only at the surface but observe significant N<sub>2</sub> fixation rates below 5 m. The diazotrophic community is often underrepresented in metagenomic datasets due to the low abundance of nitrogenase gene copies, implying our data may not present a complete  
275 picture. We suspect a more diverse diazotrophic community exists, with UCYN-A being a significant contributor to N<sub>2</sub> fixation in Arctic waters. However, the exact proportion of its contribution requires further investigation.

The contribution of N<sub>2</sub> fixation to carbon fixation (as percent of PP) is relatively low, but may increase with a further onset of bloom periods. We identified genes such as *rbcL*, which encodes Rubisco, a key enzyme in the carbon fixation pathway and *psbA*, a gene encoding Photosystem II, involved in light-driven electron transfer in photosynthesis, in our metagenomic dataset.  
280 The gene *rbcL* (for the carbon fixation pathway) and the gene *psbA* (for primary producers) were used to track the community of photosynthetic primary producers in our metagenomic dataset. At station 7, elevated carbon fixation rates are correlated with high diatom (*Bacillariophyta*) abundance and increased chl *a* concentration (Fig. 4), suggesting the onset of a bloom, which is also observable via satellite images (Appendix A1). We hypothesize that meltwater, carrying elevated nutrient and trace metal concentrations, was rapidly transported away from the glacier through the Vaigat Strait by strong winds, leading  
285 to increased productivity, as previously described by Fox and Walker (2022) & Jensen et al. (1999). The elevated diatom abundance and primary production rates at station 7 coincide with the highest N<sub>2</sub> fixation rates, which could possibly point towards a possible diatom-diazotroph symbiosis (Foster et al., 2022, 2011; Schvarcz et al., 2022). However, we did not detect any relevant diazotrophic group associated with the observed diatoms in our metagenomic dataset, which might be due to their absence or due to the general underrepresentation of diazotrophs in metagenomes. Therefore, we can only assume that such



290 a symbiosis explains the elevated rates. Additional molecular approaches would be necessary to enhance our understanding show a more detailed picture of the diazotrophic community.



**Figure 4.** Upper left image: *psbA* with correlation plot. Lower left image: *rbcL* with correlation plot. Right image: *nifH*, *nifD*, *nifK* genes per million reads in the metagenomic datasets. All figures display molecular data from metagenomic dataset for all sampled depth of station 3,7,10


There is evidence that UCYN-A have a higher Fe demand, with input through meltwater or river runoff potentially being advantageous to those organisms (Shiozaki et al., 2017, 2018; Cheung et al., 2022). Consequently, UCYN-A might play a more critical role in the future with increased Fe-rich meltwater runoff. UCYN-A can potentially fuel primary productivity by supplying nitrogen, especially with increased melting, nutrient inputs, and more light availability due to rising temperatures associated with climate change. This predicted enhancement of primary productivity may contribute to the biological drawdown of CO<sub>2</sub>, acting as a negative feedback mechanism. These projections are based on studies forecasting increased temperatures, melting, and resulting biogeochemical changes leading to higher primary productivity. However large uncertainties make predictions very difficult and should be handled with care. Thus we can only hypothesize that UCYN-A might be coupled to these dynamics by providing essential nitrogen.

### 3.4 $\delta^{15}\text{N}$ Signatures in particulate organic nitrogen show no clear evidence of nitrogen fixation

Stable isotopic composition, expressed using the  $\delta^{15}\text{N}$  notation, serve as indicators for understanding nitrogen dynamics because different biogeochemical processes fractionate nitrogen isotopes in distinct ways (Montoya (2008)). However, it is important to keep in mind that the final isotopic signal is a combination of all processes and an accurate distinction between processes cannot be made. N<sub>2</sub> fixation tends to enrich nitrogenous compounds with lighter isotopes, producing OM with isotopic values ranging approximately from -2 to +2 ‰ (Dähnke and Thamdrup (2013)). Upon complete remineralization and oxidation, organic matter contributes to a reduction in the average  $\delta$ -values in the open ocean (e.g. Montoya et al. (2002);



Emeis et al. (2010)). Whereas processes like denitrification and anammox preferentially remove lighter isotopes, leading to enrichment in heavier isotopes and delta values up to  $-25\text{‰}$ .



**Figure 5.** Vertical profiles of  $\delta^{15}\text{N}$  natural abundance signatures in PON across 10 stations in the study area. Incubation stations 3, 7, and 10 are highlighted in red, blue, and yellow, respectively. The figure shows variations in  $\delta^{15}\text{N}$  signatures with depth at each station, providing insight into nitrogen cycling in the study area.

310 Thus,  $\delta^{15}\text{N}$  values help to identify different processes of the nitrogen cycle generally present in a system (Dähnke and Thamdrupe (2013)). In our study, the  $\delta^{15}\text{N}$  values of PON from all 10 stations, range between  $2.45\text{‰}$  and  $8.30\text{‰}$  within the 0 to 100 m depth range, thus do not exhibit a clear signal indicative of  $\text{N}_2$  fixation. This suggests that  $\text{N}_2$  fixation likely contributes only a certain fraction to export production or that it only started to contribute to isotope fractionation in the bloom dynamic. The composition of OM in the surface ocean is influenced by the nitrogen substrate and the fractionation factor during photosynthesis. When nitrate is depleted in the surface ocean, the isotopic signature of OM produced during photosynthesis will mirror that of the nitrogen substrate. This substrate can originate from either nitrate in the subsurface or  $\text{N}_2$  fixation. Notably, nitrate, the primary form of dissolved nitrogen in the open ocean, typically exhibits an average stable isotope value of around

315



5 ‰. No fractionation occurs during photosynthesis because the nitrogen source is entirely taken up in the surface waters (Sigman et al. (2009)). In Qeqertarsuaq, where similar conditions prevail, this suggests that factors other than N<sub>2</sub> fixation may be influencing the observed  $\delta$ -values and POM is sustained by nitrogen sources from deeper subsurface waters, as observed in earlier studies (Fox and Walker (2022)).

In the eastern Baffin Bay waters, Atlantic water masses serve as an important source of nitrate for sustaining primary productivity, which is also reflected in the nitrogen isotopic signature in this study (Sherwood et al. (2021)). The influx of Atlantic waters, characterized by NO<sub>3</sub><sup>-</sup> values of approximately 5 ‰, closely matches the  $\delta^{15}\text{N}$  values of observed PON concentrations in our study. This suggests that Atlantic-derived NO<sub>3</sub><sup>-</sup> serves as a primary source of new nitrogen to the initial stages of bloom development (Fox and Walker, 2022; Knies, 2022). As the bloom progresses and nitrogen from Atlantic waters is depleted, N<sub>2</sub> fixation may provide an additional nitrogen source, supporting continued primary productivity. The mechanisms through which subsurface nitrate reaches the euphotic layer are not well understood. However, potential pathways include vertical migration of phytoplankton and physical mixing. Subsequently, nitrogen undergoes rapid recycling and remineralization processes to meet the system's nitrogen demands (Jensen et al. (1999)).

#### 4 Conclusion

Our study highlights the occurrence of elevated rates of N<sub>2</sub> fixation in Arctic coastal waters, particularly prominent at station 7, where they coincide with high chl *a* values, indicative of heightened productivity. Satellite observations tracing the origin of a bloom near the Isbræ Glacier, subsequently moving through the Vaigat strait, suggest a recurring phenomenon likely triggered by increased nutrient-rich meltwater originating from the glacier. This aligns with previous reports by Jensen et al. (1999) & Fox and Walker (2022), underlining the significance of such events in driving primary productivity in the region. The contribution of N<sub>2</sub> fixation to primary production was low (average 1.57 %) across the stations. Since the demand was high relative to the new nitrogen provided by N<sub>2</sub> fixation, the observed primary production must be sustained by the already present or adequate amount of subsurface supply of NO<sub>x</sub> nutrients in the seawater. This is also visible in the isotopic signature of the POM (Fox and Walker, 2022; Sherwood et al., 2021). However, the detected N<sub>2</sub> fixation rates are likely linked to the development of the fresh secondary summer bloom, which could be sustained by high nutrient and Fe availability from melting, potentially leading the system into a nutrient-limited state. The ongoing high demand for nitrogen compounds may suggest an onset to further sustain the bloom, but it remains speculative whether Fe availability definitively contributes to this process. The occurrence of such double blooms has increased by 10 % in the Qeqertarsuaq and even 33 % in the Baffin Bay, with further projected increases moving north from Greenland (Kalaallit Nunaat) waters (Ardyna et al. (2014)). Thus, nutrient demands are likely to increase, and the role of N<sub>2</sub> fixation may become more significant. The diazotrophic community in this study is dominated by UCYN-A in surface waters and may be linked to diatom abundance in deeper layers. This co-occurrence of diatoms and N<sub>2</sub> fixers in the same location is probably due to the co-limitation of similar nutrients, rather than a symbiotic relationship. Thus, this highlights the significant presence of diazotrophs despite their limited representation in datasets. It also highlights the potential for further discoveries, as existing datasets likely underestimate the full extent of the diazotrophic community (Laso Perez et al., 2024;




Shao et al., 2023; Shiozaki et al., 2017, 2023). The reported  $N_2$  fixation rates in the Vaigat strait within the Arctic Ocean are notably higher than those observed in many other oceanic regions, emphasizing that  $N_2$  fixation is an active and significant process in these high-latitude waters. When compared to measured rates across various ocean systems using the  $^{15}N$  approach, the significance of these findings becomes clear. For instance,  $N_2$  fixation rates are sometimes below the detection limit and often relatively low ranging from 0.8 to 4.4  $nmol\ N\ L^{-1}\ d^{-1}$  (Löscher et al., 2020, 2016; ?; Turk et al., 2011). In contrast, higher rates reach up to 20  $nmol\ N\ L^{-1}\ d^{-1}$  (Rees et al. (2009)) and sometime exceptional high rates range from 38 to 610  $nmol\ N\ L^{-1}\ d^{-1}$  (Bonnet et al. (2009)). The Arctic Ocean rates are thus significant in the global context, underscoring the region's role in the global nitrogen cycle and the importance of  $N_2$  fixation in supporting primary productivity in these waters.

These findings highlight the urgent need to understand the interplay between seasonal variations, sea-ice dynamics, and hydrographic conditions in Qeqertarsuaq. As climate change accelerates the melting of the Greenland Ice Sheet at Jakobshavn Isbræ, shifts in hydrodynamic patterns and hydrographic conditions in Qeqertarsuaq are anticipated. The resulting influx of warmer waters could significantly reshape the bay's hydrography, making it crucial to comprehend the coupling of climate-driven changes and oceanic processes in this vital Arctic region. Our study provides key insights into these dynamics and underscores the importance of continued investigation to predict Qeqertarsuaq's future hydrographic state. By detailing the environmental and hydrographic changes, we contribute valuable knowledge to the broader context of  $N_2$  fixation in the Arctic Ocean. Given nitrogen's pivotal role in Arctic ecosystem productivity, it is essential to explore diazotrophs, quantify  $N_2$  fixation, and assess their impact on ecosystem services as climate change progresses.







## Appendix A



**Figure A1.** Chlorophyll *a* concentration  $\text{mg m}^{-3}$  at four time points before, during, and after sea water sampling in August 2022, obtained from MODIS-Aqua; <https://giovanni.gsfc.nasa.gov> (Aqua MODIS Global Mapped Chl *a* Data, version R2022.0, DOI:10.5067/AQUA/MODIS/L3M/CHL/2022), 4 km resolution, last access 03 June 2024



**Figure A2.** Correlation matrix of environmental and biological variables. The plot shows the correlation coefficients between the following parameters: N<sub>2</sub> fixation rates, PON, POC, PP rates, the contribution N<sub>2</sub> fixation to PP (PP contribution), Chl *a*, temperature (Temp), salinity (Sal), and Oxygen. The scale ranges from -1 to 1, where values close to 1 or -1 indicate strong positive or negative correlations, respectively, and values near 0 indicate weak or no correlation. The color intensity represents the strength and direction of the correlations, facilitating the identification of relationships among the variables



**Figure A3.** This figure displays a ggpairs plot, showing pairwise relationships and correlations between biological and environmental variables. Pearson correlation coefficients displayed in the upper triangular panel, indicating the strength and significance of linear relationships. Statistical significance levels are indicated by stars (\*), where \* indicates  $p < 0.05$ , \*\* indicates  $p < 0.01$  and \*\*\* indicates  $p < 0.001$



370 *Data availability.* The presented data collected during the cruise will be made accessible on PANGEA. The molecular datasets have been deposited with the accession number: Bioproject PRJNA1133027/Bioproject PRJNA1133027.

375 *Author contributions.* IS carried out fieldwork and laboratory work at the University of Southern Denmark, and wrote the majority of the manuscript. ELP, AM, and EL conducted fieldwork and laboratory work at the University of Southern Denmark. PX performed metagenomic analysis and created the corresponding graphs. CRL designed the study, provided supervision and guidance throughout the project, and contributed to the writing and revision of the manuscript. All authors contributed to the conception of the study and participated in the writing and revision of the manuscript.

*Competing interests.* The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. One of the authors, CRL, serves as an Associate Editor for Biogeosciences.

380 *Acknowledgements.* This work was supported by the Velux Foundation (grant no.29411 to Carolin R. Löscher) and through the DFF grant from the the Independent Research Fund Denmark (grant no. 0217-00089B to Lasse Riemann, Carolin R. Löscher and Stiig Markager). ELP was supported by a postdoctoral contract from Danmarks Frie Forskningsfond (DFF, 1026-00428B) at SDU, and by a Marie Skłodowska-Curie postdoctoral fellowship (HORIZON291 MSCA-2021-PF-01, project number: 101066750) by the European Commission at Princeton University. We sincerely thank the captain and crew of the P540 during the cruise on the Danish military vessel for their invaluable support and cooperation at sea. Our gratitude extends to Isaaffik Arctic Gateway for providing the infrastructure and opportunities that made this project possible. We also acknowledge Zarah Kofoed for her technical support in the laboratory and thank all the Nordcee laboratory technicians for  
385 their general assistance.



## References

- Ardyna, M. and Arrigo, K. R.: Phytoplankton dynamics in a changing Arctic Ocean, *Nature Climate Change*, 10, 892–903, 2020.
- Ardyna, M., Babin, M., Gosselin, M., Devred, E., Rainville, L., and Tremblay, J.-É.: Recent Arctic Ocean sea ice loss triggers novel fall phytoplankton blooms, *Geophysical Research Letters*, 41, 6207–6212, 2014.
- 390 Arrigo, K. R. and van Dijken, G. L.: Continued increases in Arctic Ocean primary production, *Progress in oceanography*, 136, 60–70, 2015.
- Arrigo, K. R., van Dijken, G., and Pabi, S.: Impact of a shrinking Arctic ice cover on marine primary production, *Geophysical Research Letters*, 35, 2008.
- Arrigo, K. R., van Dijken, G. L., Castelao, R. M., Luo, H., Rennermalm, Å. K., Tedesco, M., Mote, T. L., Oliver, H., and Yager, P. L.: Melting glaciers stimulate large summer phytoplankton blooms in southwest Greenland waters, *Geophysical Research Letters*, 44, 6278–  
395 6285, 2017.
- Bhatia, M. P., Kujawinski, E. B., Das, S. B., Breier, C. F., Henderson, P. B., and Charette, M. A.: Greenland meltwater as a significant and potentially bioavailable source of iron to the ocean, *Nature Geoscience*, 6, 274–278, 2013.
- Blais, M., Tremblay, J.-É., Jungblut, A. D., Gagnon, J., Martin, J., Thaler, M., and Lovejoy, C.: Nitrogen fixation and identification of potential diazotrophs in the Canadian Arctic, *Global Biogeochemical Cycles*, 26, 2012.
- 400 Bonnet, S., Biegala, I. C., Dutrieux, P., Slemmons, L. O., and Capone, D. G.: Nitrogen fixation in the western equatorial Pacific: Rates, diazotrophic cyanobacterial size class distribution, and biogeochemical significance, *Global Biogeochemical Cycles*, 23, 2009.
- Buchanan, P. J., Chase, Z., Matear, R. J., Phipps, S. J., and Bindoff, N. L.: Marine nitrogen fixers mediate a low latitude pathway for atmospheric CO<sub>2</sub> drawdown, *Nature Communications*, 10, 4611, 2019.
- Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P., and Huerta-Cepas, J.: eggNOG-mapper v2: functional annotation, orthology  
405 assignments, and domain prediction at the metagenomic scale, *Molecular biology and evolution*, 38, 5825–5829, 2021.
- Capone, D. G. and Carpenter, E. J.: Nitrogen fixation in the marine environment, *Science*, 217, 1140–1142, 1982.
- Chen, Y., Chen, Y., Shi, C., Huang, Z., Zhang, Y., Li, S., Li, Y., Ye, J., Yu, C., Li, Z., et al.: SOAPnuke: a MapReduce acceleration-supported software for integrated quality control and preprocessing of high-throughput sequencing data, *Gigascience*, 7, gix120, 2018.
- Cheung, S., Liu, K., Turk-Kubo, K. A., Nishioka, J., Suzuki, K., Landry, M. R., Zehr, J. P., Leung, S., Deng, L., and Liu, H.: High biomass  
410 turnover rates of endosymbiotic nitrogen-fixing cyanobacteria in the western Bering Sea, *Limnology and Oceanography Letters*, 7, 501–509, 2022.
- Coale, T. H., Loconte, V., Turk-Kubo, K. A., Vanslebrouck, B., Mak, W. K. E., Cheung, S., Ekman, A., Chen, J.-H., Hagino, K., Takano, Y., et al.: Nitrogen-fixing organelle in a marine alga, *Science*, 384, 217–222, 2024.
- Dähnke, K. and Thamdrup, B.: Nitrogen isotope dynamics and fractionation during sedimentary denitrification in Boknis Eck, Baltic Sea,  
415 *Biogeosciences*, 10, 3079–3088, 2013.
- Damm, E., Helmke, E., Thoms, S., Schauer, U., Nöthig, E., Bakker, K., and Kiene, R.: Methane production in aerobic oligotrophic surface water in the central Arctic Ocean, *Biogeosciences*, 7, 1099–1108, 2010.
- Díez, B., Bergman, B., Pedrós-Alió, C., Antó, M., and Snoeijis, P.: High cyanobacterial nifH gene diversity in Arctic seawater and sea ice brine, *Environmental microbiology reports*, 4, 360–366, 2012.
- 420 Emeis, K.-C., Mara, P., Schlarbaum, T., Möbius, J., Dähnke, K., Struck, U., Mihalopoulos, N., and Krom, M.: External N inputs and internal N cycling traced by isotope ratios of nitrate, dissolved reduced nitrogen, and particulate nitrogen in the eastern Mediterranean Sea, *Journal of Geophysical Research: Biogeosciences*, 115, 2010.



- Falkowski, P. G., Fenchel, T., and Delong, E. F.: The microbial engines that drive Earth's biogeochemical cycles, *science*, 320, 1034–1039, 2008.
- 425 Farnelid, H., Andersson, A. F., Bertilsson, S., Al-Soud, W. A., Hansen, L. H., Sørensen, S., Steward, G. F., Hagström, Å., and Riemann, L.: Nitrogenase gene amplicons from global marine surface waters are dominated by genes of non-cyanobacteria, *PloS one*, 6, e19223, 2011.
- Farnelid, H., Turk-Kubo, K., Ploug, H., Ossolinski, J. E., Collins, J. R., Van Mooy, B. A., and Zehr, J. P.: Diverse diazotrophs are present on sinking particles in the North Pacific Subtropical Gyre, *The ISME journal*, 13, 170–182, 2019.
- Fernández-Méndez, M., Turk-Kubo, K. A., Buttigieg, P. L., Rapp, J. Z., Krumpfen, T., and Zehr, J. P.: Diazotroph diversity in the sea ice, melt  
430 ponds, and surface waters of the Eurasian Basin of the Central Arctic Ocean, *Frontiers in microbiology*, 7, 217140, 2016.
- Foster, R. A., Kuypers, M. M., Vagner, T., Paerl, R. W., Musat, N., and Zehr, J. P.: Nitrogen fixation and transfer in open ocean diatom–cyanobacterial symbioses, *The ISME journal*, 5, 1484–1493, 2011.
- Foster, R. A., Tienken, D., Littmann, S., Whitehouse, M. J., Kuypers, M. M., and White, A. E.: The rate and fate of N<sub>2</sub> and C fixation by marine diatom-diazotroph symbioses, *The ISME journal*, 16, 477–487, 2022.
- 435 Fox, A. and Walker, B. D.: Sources and Cycling of Particulate Organic Matter in Baffin Bay: A Multi-Isotope  $\delta^{13}\text{C}$ ,  $\delta^{15}\text{N}$ , and  $\Delta^{14}\text{C}$  Approach, *Frontiers in Marine Science*, 9, 846025, 2022.
- Fu, L., Niu, B., Zhu, Z., Wu, S., and Cd-hit, W. L.: Accelerated for clustering the next-generation sequencing data, *Bioinformatics*, 28, 3150–3152, 2012.
- Galloway, J., Dentener, F., Capone, D., Boyer, E., Howarth, R., Seitzinger, S., Asner, G., Cleveland, C., Green, P., Holland, E., et al.: Nitrogen  
440 cycles: past, present, and future. *Biogeochemistry* 70, 153e226, 2004.
- García-Robledo, E., Corzo, A., and Pappaspyrou, S.: A fast and direct spectrophotometric method for the sequential determination of nitrate and nitrite at low concentrations in small volumes, *Marine Chemistry*, 162, 30–36, 2014.
- Gladish, C. V., Holland, D. M., and Lee, C. M.: Oceanic boundary conditions for Jakobshavn Glacier. Part II: Provenance and sources of variability of Disko Bay and Ilulissat icefjord waters, 1990–2011, *Journal of Physical Oceanography*, 45, 33–63, 2015.
- 445 Großkopf, T., Mohr, W., Baustian, T., Schunck, H., Gill, D., Kuypers, M. M., Lavik, G., Schmitz, R. A., Wallace, D. W., and LaRoche, J.: Doubling of marine dinitrogen-fixation rates based on direct measurements, *Nature*, 488, 361–364, 2012.
- Gruber, N.: The dynamics of the marine nitrogen cycle and its influence on atmospheric CO<sub>2</sub> variations, in: *The ocean carbon cycle and climate*, pp. 97–148, Springer, 2004.
- Gruber, N. and Galloway, J. N.: An Earth-system perspective of the global nitrogen cycle, *Nature*, 451, 293–296, 2008.
- 450 Gruber, N. and Sarmiento, J. L.: Global patterns of marine nitrogen fixation and denitrification, *Global biogeochemical cycles*, 11, 235–266, 1997.
- Haine, T. W., Curry, B., Gerdes, R., Hansen, E., Karcher, M., Lee, C., Rudels, B., Spreen, G., de Steur, L., Stewart, K. D., et al.: Arctic freshwater export: Status, mechanisms, and prospects, *Global and Planetary Change*, 125, 13–35, 2015.
- Hanna, E., Huybrechts, P., Steffen, K., Cappelen, J., Huff, R., Shuman, C., Irvine-Fynn, T., Wise, S., and Griffiths, M.: Increased runoff from  
455 melt from the Greenland Ice Sheet: a response to global warming, *Journal of Climate*, 21, 331–341, 2008.
- Hansen, M. O., Nielsen, T. G., Stedmon, C. A., and Munk, P.: Oceanographic regime shift during 1997 in Disko Bay, western Greenland, *Limnology and Oceanography*, 57, 634–644, 2012.
- Harding, K., Turk-Kubo, K. A., Sipler, R. E., Mills, M. M., Bronk, D. A., and Zehr, J. P.: Symbiotic unicellular cyanobacteria fix nitrogen in the Arctic Ocean, *Proceedings of the National Academy of Sciences*, 115, 13371–13375, 2018.



- 460 Hawkings, J., Wadham, J., Tranter, M., Lawson, E., Sole, A., Cowton, T., Tedstone, A., Bartholomew, I., Nienow, P., Chandler, D., et al.:  
The effect of warming climate on nutrient and solute export from the Greenland Ice Sheet, *Geochemical Perspectives Letters*, pp. 94–104,  
2015.
- Hawkings, J. R., Wadham, J. L., Tranter, M., Raiswell, R., Benning, L. G., Statham, P. J., Tedstone, A., Nienow, P., Lee, K., and Telling, J.:  
Ice sheets as a significant source of highly reactive nanoparticulate iron to the oceans, *Nature communications*, 5, 1–8, 2014.
- 465 Hendry, K. R., Huvenne, V. A., Robinson, L. F., Annett, A., Badger, M., Jacobel, A. W., Ng, H. C., Opher, J., Pickering, R. A., Taylor, M. L.,  
et al.: The biogeochemical impact of glacial meltwater from Southwest Greenland, *Progress in Oceanography*, 176, 102–126, 2019.
- Holland, D. M., Thomas, R. H., De Young, B., Ribergaard, M. H., and Lyberth, B.: Acceleration of Jakobshavn Isbræ triggered by warm  
subsurface ocean waters, *Nature geoscience*, 1, 659–664, 2008.
- Hopwood, M. J., Connelly, D. P., Arendt, K. E., Juul-Pedersen, T., Stinchcombe, M. C., Meire, L., Esposito, M., and Krishna, R.: Seasonal  
470 changes in Fe along a glaciated Greenlandic fjord, *Frontiers in Earth Science*, 4, 15, 2016.
- Hyatt, D., Chen, G.-L., LoCascio, P. F., Land, M. L., Larimer, F. W., and Hauser, L. J.: Prodigal: prokaryotic gene recognition and translation  
initiation site identification, *BMC bioinformatics*, 11, 1–11, 2010.
- Jensen, H. M., Pedersen, L., Burmeister, A., and Winding Hansen, B.: Pelagic primary production during summer along 65 to 72 N off West  
Greenland, *Polar Biology*, 21, 269–278, 1999.
- 475 Karl, D., Michaels, A., Bergman, B., Capone, D., Carpenter, E., Letelier, R., Lipschultz, F., Paerl, H., Sigman, D., and Stal, L.: Dinitrogen  
fixation in the world's oceans, *The nitrogen cycle at regional to global scales*, pp. 47–98, 2002.
- Knies, J.: Nitrogen isotope evidence for changing Arctic Ocean ventilation regimes during the Cenozoic, *Geophysical Research Letters*, 49,  
e2022GL099512, 2022.
- Krawczyk, D. W., Yesson, C., Knutz, P., Arboe, N. H., Blicher, M. E., Zinglensen, K. B., and Wagnholt, J. N.: Seafloor habitats across  
480 geological boundaries in Disko Bay, central West Greenland, *Estuarine, Coastal and Shelf Science*, 278, 108–117, 2022.
- Krupke, A., Mohr, W., LaRoche, J., Fuchs, B. M., Amann, R. I., and Kuypers, M. M.: The effect of nutrients on carbon and nitrogen fixation  
by the UCYN-A–haptophyte symbiosis, *The ISME journal*, 9, 1635–1647, 2015.
- Laso Perez, R., Rivas Santisteban, J., Fernandez-Gonzalez, N., Mundy, C. J., Tamames, J., and Pedros-Alio, C.: Nitrogen cycling during an  
Arctic bloom: from chemolithotrophy to nitrogen assimilation, *bioRxiv*, pp. 2024–02, 2024.
- 485 Lewis, K., Van Dijken, G., and Arrigo, K. R.: Changes in phytoplankton concentration now drive increased Arctic Ocean primary production,  
*Science*, 369, 198–202, 2020.
- Li, D., Liu, C.-M., Luo, R., Sadakane, K., and Lam, T.-W.: MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics  
assembly via succinct de Bruijn graph, *Bioinformatics*, 31, 1674–1676, 2015.
- Löscher, C. R., Bourbonnais, A., Dekaezacker, J., Charoenpong, C. N., Altabet, M. A., Bange, H. W., Czeschel, R., Hoffmann, C., and  
490 Schmitz, R.: N<sub>2</sub> fixation in eddies of the eastern tropical South Pacific Ocean, *Biogeosciences*, 13, 2889–2899, 2016.
- Löscher, C. R., Mohr, W., Bange, H. W., and Canfield, D. E.: No nitrogen fixation in the Bay of Bengal?, *Biogeosciences*, 17, 851–864, 2020.
- Luo, Y.-W., Doney, S., Anderson, L., Benavides, M., Berman-Frank, I., Bode, A., Bonnet, S., Boström, K. H., Böttjer, D., Capone, D., et al.:  
Database of diazotrophs in global ocean: abundance, biomass and nitrogen fixation rates, *Earth System Science Data*, 4, 47–73, 2012.
- Martínez-Pérez, C., Mohr, W., Löscher, C. R., Dekaezacker, J., Littmann, S., Yilmaz, P., Lehnen, N., Fuchs, B. M., Lavik, G., Schmitz,  
495 R. A., et al.: The small unicellular diazotrophic symbiont, UCYN-A, is a key player in the marine nitrogen cycle, *Nature Microbiology*, 1,  
1–7, 2016.



- Mills, M. M., Turk-Kubo, K. A., van Dijken, G. L., Henke, B. A., Harding, K., Wilson, S. T., Arrigo, K. R., and Zehr, J. P.: Unusual marine cyanobacteria/haptophyte symbiosis relies on N<sub>2</sub> fixation even in N-rich environments, *The ISME Journal*, 14, 2395–2406, 2020.
- Mohr, W., Grosskopf, T., Wallace, D. W., and LaRoche, J.: Methodological underestimation of oceanic nitrogen fixation rates, *PloS one*, 5, e12583, 2010.
- 500
- Montoya, J. P.: Nitrogen stable isotopes in marine environments, *Nitrogen in the marine environment*, 2, 1277–1302, 2008.
- Montoya, J. P., Carpenter, E. J., and Capone, D. G.: Nitrogen fixation and nitrogen isotope abundances in zooplankton of the oligotrophic North Atlantic, *Limnology and Oceanography*, 47, 1617–1628, 2002.
- Mortensen, J., Rysgaard, S., Winding, M., Juul-Pedersen, T., Arendt, K., Lund, H., Stuart-Lee, A., and Meire, L.: Multidecadal water mass dynamics on the West Greenland Shelf, *Journal of Geophysical Research: Oceans*, 127, e2022JC018724, 2022.
- 505
- Munk, P., Nielsen, T. G., and Hansen, B. W.: Horizontal and vertical dynamics of zooplankton and larval fish communities during mid-summer in Disko Bay, West Greenland, *Journal of Plankton Research*, 37, 554–570, 2015.
- Murphy, J. and Riley, J. P.: A modified single solution method for the determination of phosphate in natural waters, *Analytica chimica acta*, 27, 31–36, 1962.
- 510
- Myers, P. G. and Ribergaard, M. H.: Warming of the polar water layer in Disko Bay and potential impact on Jakobshavn Isbrae, *Journal of Physical Oceanography*, 43, 2629–2640, 2013.
- Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., and Kingsford, C.: Salmon provides fast and bias-aware quantification of transcript expression, *Nature methods*, 14, 417–419, 2017.
- Reeder, C. F., Stoltenberg, I., Javidpour, J., and Löscher, C. R.: Salinity as a key control on the diazotrophic community composition in the Baltic Sea, *Ocean Science Discussions*, 2021, 1–30, 2021.
- 515
- Rees, A. P., Gilbert, J. A., and Kelly-Gerrey, B. A.: Nitrogen fixation in the western English Channel (NE Atlantic ocean), *Marine Ecology Progress Series*, 374, 7–12, 2009.
- Rysgaard, S., Boone, W., Carlson, D., Sejr, M., Bendtsen, J., Juul-Pedersen, T., Lund, H., Meire, L., and Mortensen, J.: An updated view on water masses on the pan-west Greenland continental shelf and their link to proglacial fjords, *Journal of Geophysical Research: Oceans*, 125, e2019JC015564, 2020.
- 520
- Schiøtt, S.: The Marine Ecosystem of Ilulissat Icefjord, Greenland, Ph.D. thesis, Department of Biology, Aarhus University, Denmark, 2023.
- Schlitzer, R.: Ocean data view, 2022.
- Szvarcz, C. R., Wilson, S. T., Caffin, M., Stancheva, R., Li, Q., Turk-Kubo, K. A., White, A. E., Karl, D. M., Zehr, J. P., and Steward, G. F.: Overlooked and widespread pennate diatom-diazotroph symbioses in the sea, *Nature communications*, 13, 799, 2022.
- 525
- Shao, Z., Xu, Y., Wang, H., Luo, W., Wang, L., Huang, Y., Agawin, N. S. R., Ahmed, A., Benavides, M., Bentzon-Tilia, M., et al.: Global oceanic diazotroph database version 2 and elevated estimate of global N<sub>2</sub> fixation, *Earth System Science Data*, 15, 2023.
- Sherwood, O. A., Davin, S. H., Lehmann, N., Buchwald, C., Edinger, E. N., Lehmann, M. F., and Kienast, M.: Stable isotope ratios in seawater nitrate reflect the influence of Pacific water along the northwest Atlantic margin, *Biogeosciences*, 18, 4491–4510, 2021.
- Shiozaki, T., Bombar, D., Riemann, L., Hashihama, F., Takeda, S., Yamaguchi, T., Ehama, M., Hamasaki, K., and Furuya, K.: Basin scale variability of active diazotrophs and nitrogen fixation in the North Pacific, from the tropics to the subarctic Bering Sea, *Global Biogeochemical Cycles*, 31, 996–1009, 2017.
- 530
- Shiozaki, T., Fujiwara, A., Ijichi, M., Harada, N., Nishino, S., Nishi, S., Nagata, T., and Hamasaki, K.: Diazotroph community structure and the role of nitrogen fixation in the nitrogen cycle in the Chukchi Sea (western Arctic Ocean), *Limnology and Oceanography*, 63, 2191–2205, 2018.





- 535 Shiozaki, T., Fujiwara, A., Inomura, K., Hirose, Y., Hashihama, F., and Harada, N.: Biological nitrogen fixation detected under Antarctic sea ice, *Nature geoscience*, 13, 729–732, 2020.
- Shiozaki, T., Nishimura, Y., Yoshizawa, S., Takami, H., Hamasaki, K., Fujiwara, A., Nishino, S., and Harada, N.: Distribution and survival strategies of endemic and cosmopolitan diazotrophs in the Arctic Ocean, *The ISME journal*, 17, 1340–1350, 2023.
- 540 Sigman, D. M., DiFiore, P. J., Hain, M. P., Deutsch, C., Wang, Y., Karl, D. M., Knapp, A. N., Lehmann, M. F., and Pantoja, S.: The dual isotopes of deep nitrate as a constraint on the cycle and budget of oceanic fixed nitrogen, *Deep Sea Research Part I: Oceanographic Research Papers*, 56, 1419–1439, 2009.
- Sipler, R. E., Gong, D., Baer, S. E., Sanderson, M. P., Roberts, Q. N., Mulholland, M. R., and Bronk, D. A.: Preliminary estimates of the contribution of Arctic nitrogen fixation to the global nitrogen budget, *Limnology and Oceanography Letters*, 2, 159–166, 2017.
- 545 Slawyk, G., Collos, Y., and Auclair, J.-C.: The use of the  $^{13}\text{C}$  and  $^{15}\text{N}$  isotopes for the simultaneous measurement of carbon and nitrogen turnover rates in marine phytoplankton I, *Limnology and Oceanography*, 22, 925–932, 1977.
- Sohm, J. A., Webb, E. A., and Capone, D. G.: Emerging patterns of marine nitrogen fixation, *Nature Reviews Microbiology*, 9, 499–508, 2011.
- 550 Tang, W., Wang, S., Fonseca-Batista, D., Dehairs, F., Gifford, S., Gonzalez, A. G., Gallinari, M., Planquette, H., Sarthou, G., and Cassar, N.: Revisiting the distribution of oceanic  $\text{N}_2$  fixation and estimating diazotrophic contribution to marine production, *Nature communications*, 10, 831, 2019.
- Tremblay, J.-É. and Gagnon, J.: The effects of irradiance and nutrient supply on the productivity of Arctic waters: a perspective on climate change, in: *Influence of climate change on the changing arctic and sub-arctic conditions*, pp. 73–93, Springer, 2009.
- Turk, K. A., Rees, A. P., Zehr, J. P., Pereira, N., Swift, P., Shelley, R., Lohan, M., Woodward, E. M. S., and Gilbert, J.: Nitrogen fixation and nitrogenase (*nifH*) expression in tropical waters of the eastern North Atlantic, *The ISME journal*, 5, 1201–1212, 2011.
- 555 Turk-Kubo, K. A., Frank, I. E., Hogan, M. E., Desnues, A., Bonnet, S., and Zehr, J. P.: Diazotroph community succession during the VAHINE mesocosm experiment (New Caledonia lagoon), *Biogeosciences*, 12, 7435–7452, 2015.
- Von Friesen, L. W. and Riemann, L.: Nitrogen fixation in a changing Arctic Ocean: an overlooked source of nitrogen?, *Frontiers in Microbiology*, 11, 596426, 2020.
- 560 Wang, S., Bailey, D., Lindsay, K., Moore, J., and Holland, M.: Impact of sea ice on the marine iron cycle and phytoplankton productivity, *Biogeosciences*, 11, 4713–4731, 2014.
- Zehr, J. P. and Capone, D. G.: Changing perspectives in marine nitrogen fixation, *Science*, 368, eaay9514, 2020.