

# Marine heatwaves deeply alter marine food web structure and function

Vianney Guibourd de Luzinai<sup>1,2\*</sup>, William W. L. Cheung<sup>2</sup>, Didier Gascuel<sup>1</sup>

<sup>1</sup>UMR Dynamics and sustainability of ecosystems: from source to sea (DECOD), Institut Agro, Ifremer, INRAE, Rennes, France

<sup>2</sup>Institute for the Oceans and Fisheries, the University of British Columbia, Vancouver, British Columbia, Canada

*Correspondence to:* Vianney Guibourd de Luzinai (vianney.gdl@orange.fr)

## 1. Abstract

Marine heatwaves (MHWs) have become longer, more frequent and more intense in recent decades. MHWs have caused large-scale ecological impacts, such as coral bleaching, mass mortality of seagrass, fishes and invertebrates, and shifts in abundance and distribution of marine species. However, the implications of these MHW-induced impacts on marine species for the structure and functioning of marine food webs are not clearly understood. In this study, we use the EcoTroph-Dyn ecosystem modelling approach to examine the impacts of MHWs occurring during the year's warmest month on the trophodynamics of marine ecosystems. EcoTroph-Dyn represents marine ecosystem dynamics at a spatial resolution of 1° longitude by 1° latitude and a temporal resolution of 14 days. We applied the model to simulate changes in trophodynamic processes, energy transfer and ecosystem biomass using daily temperature and monthly net primary production (NPP) data that were derived from satellite observations from 1998 to 2021. We compared and contrasted the simulated changes in biomass by trophic level with results generated from temperature and NPP time series that had been filtered to remove MHWs. Our results show a significant decline in biomass between 1998 and 2021 specifically caused by MHWs. For example, in the Northeast Pacific Ocean, our model simulated a MHW-induced decline in biomass of  $8.7\% \pm 1.0$  (standard error) from 2013 to 2016. Overall, MHW-induced biomass declines are more pronounced in the northern hemisphere and Pacific Ocean. Moreover, the MHW-induced declines in high trophic level biomass were larger than in lower trophic levels and lasted longer post-MHW. Finally, this study highlights the need to integrate MHWs into modelling the effects of climate change on marine ecosystems. It shows that the EcoTroph approach, and especially its new dynamic version, provides a framework to understand more comprehensively the implications of climate change for marine ecosystem structure and functioning.

## 2. Introduction

Over the last century, marine heatwaves (MHWs) - defined as more than 5 days period of anomalously warm sea surface temperatures (SSTs) exceeding a specific threshold, typically determined by natural climatological variations - have increased in frequency, duration, and intensity (Frölicher and Laufkötter, 2018; Oliver et al., 2018). Between the 1920s and the 2010s, their duration and frequency increased by 17% and 34%, respectively, resulting in more than a doubling of the number of MHW days at a global scale (Oliver et al., 2018). Since the 2000s, MHWs that are spatially and temporally extensive have been recorded in the world oceans, such as the North Pacific MHW in 2013-2015 (Bond et al., 2015; Cavole et al., 2016), the Mediterranean Sea 2003 and 2022 MHW (Garrahou et al., 2009, 2022), the Western Australia MHW in 2011 (Pearce et al., 2011), and the Tasman Sea MHW in 2015-2016 (Oliver et al., 2017).

Marine ectotherms' physiological functions are directly affected by ocean temperature changes that are closely related to their body temperature (Guibourd de Luzinais et al., 2024; Pörtner and Farrell, 2008). These species are adapted to perform optimally at a range of body temperature, with certain upper and lower temperature limits within which they can survive (Pörtner and Farrell, 2008). When environmental temperatures exceed this temperature range, e.g., during MHWs, the organism is stressed, leading to functional constraints and declines in performance (Pörtner and Farrell, 2008). Particularly, abnormally high temperatures during MHWs often exceed organisms' thermal limits, impacting their distribution, growth and survival (Guibourd de Luzinais et al., 2024; Smale et al., 2019; Smith et al., 2023). Moreover, the impacts of MHWs at the population level have cascading effects at the community and ecosystem levels. For example, MHW-induced declines in phytoplankton biomass and diversity have led to significant changes in zooplankton and other marine invertebrate diversity and biomass (Cavole et al., 2016). MHWs cause coral bleaching that also impacts coral reef ecosystems (Garrahou et al., 2009, 2022; Pearce et al., 2011). Range shifts driven by MHWs result in "tropicalization" of fish communities (Wernberg et al., 2016). Ultimately, MHWs imply mass mortality of fish and invertebrates modifying ecosystem functioning (Cannell et al., 2019; Cavole et al., 2016; Collins et al., 2019). However, these ecological impacts of MHWs are not ubiquitous and vary largely between MHW events, species and ecosystems (Fredston et al., 2023; Oliver et al., 2021; Pershing et al., 2018; Smale et al., 2019; Smith et al., 2023).

The ecological impacts of MHWs have predominantly been studied using laboratory experimentations, analyses of ecological time series, and numerical modelling (Joyce et al., 2024). For example, Carneiro et al. (2020) assessed the evolution of physiological and biochemical parameters and survival rates of the clam *Anomalocardia flexuosa* in response to simulated MHWs. Under laboratory conditions, *Anomalocardia flexuosa* was allowed to adapt to a stable control condition

before being exposed to simulated conditions of MHWs occurrence lasting up to 21 days by warming the tank water by 3°C above the control temperature. Fredston et al. (2023) studied the effects of MHWs on marine species biomass by analysing scientific trawl survey data (FishGlob data) and historical temperature records. Previous studies assessed MHWs impacts through numerical modelling approaches. For example Cheung et al. (2021) and Cheung & Frölicher (2020) employed employed climate-fish-fisheries models to investigate MHW implications for biomass and potential catches of exploited marine species and their implication for fisheries. They found that MHWs may cause biomass decreases and shifts in the biogeography of fish stocks that are faster and bigger in magnitude than the effects of decadal-scale mean changes. They projected a doubling of impact levels by 2050 amongst the most important fisheries species over previous assessments that focus only on long-term climate change. Gomes et al. (2024) use the Ecopath with Ecosim (EwE) modelling approach to assess the ecological impacts of 'the Blob' MHW. They highlighted the alteration of trophic interactions and energy flux following the MHWs which might have profound consequences for the specific ecosystem structure and function. However, there is a gap in applying ecosystem modelling framework to study global impacts of MHWs on ecosystem structure and functions.

Trophic dynamics of marine ecosystems are affected by ocean temperature (Eddy et al., 2021; du Pontavice et al., 2020). In particular, ocean warming affects biomass transfer efficiency and the speed of energy transfer through the food web, i.e., flow kinetic (du Pontavice et al., 2020). Faster flow kinetic under ocean warming represents the increasing dominance of short-lived species so that each unit of biomass spends less time at a given TL and, on average, across all TLs, ultimately leading to a decrease in total biomass (Gascuel et al., 2008). Simultaneously, ocean warming is expected to induce a decrease in biomass transfer efficiency, altering both consumer production and biomass due to larger energy losses between each TL (du Pontavice et al., 2020). Therefore, ocean warming alters both the amount and speed of matter and energy transfer within the food web, potentially leading to a decline in consumer biomass through independent and cumulative effects (du Pontavice et al., 2021; Guibourd de Luzinais et al., 2023). We expect that temperature changes during MHWs will also impact trophodynamics and marine animal biomass. However, the effects of MHWs on ecosystem structure and function have not yet been clearly understood on a global scale.

This study aimed to disentangle the additional or synergistic consequences of MHWs occurring during the year's warmest month, when species are undergoing a thermal stress, with the effects of the slow-onset climate change in marine ecosystems. We used the EcoTroph-Dyn ecosystem model that was developed and applied to study the responses of marine ecosystems to MHWs (see Guibourd de Luzinais et al., 2024). We explored the effects of different scenarios of MHW-induced community mortality that were dependent on climatic conditions and species' resistance capacities. We undertook a global-scale hindcast analysis over the 1998 to 2021 period and analysed the added impacts of MHWs

on marine ecosystem biomass and trophodynamics under climate change. Subsequently, we delved into distinct geographical characteristics and identified marine ecosystems that are particularly sensitive to MHW-induced biological impacts. Last but not least, as a case study, we examined a recognised MHW ('the Blob', which occurred in the Northeast Pacific Ocean from 2013 to 2016) and highlighted how MHWs can trigger long-term changes in the ecosystem.

### 3. Material and methods

#### 3.1 The EcoTroph dynamic model

We used EcoTroph-Dyn, a dynamic version derived from the steady-state EcoTroph trophodynamic modelling approach first proposed by Gascuel, (2005) and further developed by Gascuel & Pauly, (2009). In EcoTroph, biomass is produced by primary producers (trophic level TL = 1) and consumed by heterotrophic organisms (TL>1). Thus, the food webs functioning is represented as a continuous flow of biomass surging up the food webs, from primary producers (low TLs) to top predators (high TLs). The resulting ecosystem structure is a continuous distribution of biomass along a gradient of TLs, i.e., biomass trophic spectra (Gascuel et al., 2005, Figure 1). Practically, each biomass trophic spectrum with TL above 1 is split into small trophic classes bounded by pre-defined lower and upper trophic levels (with conventionally TL width = 0.1 - primarily because of computational efficiency while maintaining a good representation of food web structure and functions - in the steady-state version of EcoTroph). The biomass spectrum in EcoTroph generally refers to the total biomass of all consumers, including organisms living in pelagic, mid-water, and benthic habitats. The time needed for the biomass to flow from one to the next trophic class varies along the food chain, with biomass transfers generally being faster in lower TLs (as species generally have short life-expectancy) than in the higher ones.

EcoTroph has been applied to study the long-term effects of fishing, e.g., (du Pontavice et al., 2023; Gasche et al., 2012; Gasche & Gascuel, 2013; Halouani et al., 2015; Tremblay-Boyer et al., 2011) and climate change (du Pontavice et al., 2021) on ecosystem biomass and production. EcoTroph-Dyn, the dynamic version of EcoTroph, simulates changes in biomass flows in each trophic biomass spectrum at a 15 days time step. This time step was used as it represents the average duration of most naturally occurring or experimentally simulated MHWs (Smale et al., 2015). EcoTroph-Dyn includes algorithms that model changes in flow kinetics, boundaries of trophic classes and the resulting biomass states and flow between trophic classes, and an algorithm that models the biomass loss associated with MHW occurrence. EcoTroph-Dyn's algorithm details are described in Guibourd de Luzinais et al., 2024. Here, we provided a summary of these algorithms.

The quantity of biomass flowing from a trophic class to the next, due to predation or ontogeny, is not conservative and can be calculated at each time step as:

$$\Phi_{\tau+\Delta\tau,t+1} = \Phi_{\tau,t} \cdot e^{-(\mu_t+\eta_{\tau,t})\cdot\Delta\tau_t}, \quad (1)$$

where  $\mu_t$  (expressed in  $TL^{-1}$ ) represents the mean natural losses (due to non-predation mortalities, excretion and respiration). The  $\eta_{\tau,t}$  parameter (expressed in  $TL^{-1}$ ) represents the additional loss rate specifically due to mortalities induced by MHW events (section 3.2.2). Equation (1) additionally describes the transfer efficiency (TE) between continuous trophic levels (TLs), representing the estimated proportion of biomass flow passed from one TL to the next. TE can be formulated as an empirical equation based on SST, as presented by du Pontavice et al. (2021):

$$TE = e^{-\mu_\tau} = e^{(-2.162+b+(-0.025+a)\cdot SST)} \cdot 1.038013, \quad (2)$$

where  $a$  and  $b$  are specific parameters for each biome type (du Pontavice et al., 2021) and SST the sea surface temperature of the time simulated.

The flow kinetic  $K_\tau$  (expressed in  $TL\cdot year^{-1}$ ), representing the speed of biomass flows (trophic transfers) from low to high TLs, is inversely proportional to biomass residence time (the time each biomass unit stays at a given TL).  $K_\tau$  is expressed as a function of trophic level, using the empirical equation described in Gascuel et al., (2008):

$$K_\tau = 20.19 \cdot \tau_j^{-3.258} \cdot e^{0.041 \cdot SST_y}, \quad (3)$$

where  $SST_y$  corresponds to the annual moving average sea surface temperature in year  $y$  and  $\tau_j$  corresponds to the trophic level of the  $j$  TL class.

MHWs cause marine organism mortality, impacting their life expectancy (Smith et al., 2023). In EcoTroph-Dyn, these changes in life expectancy are reflected in the loss rate within the biomass spectrum, representing the proportion of biomass that neither persists nor progresses through the food web (Gascuel et al., 2008; Guibourd de Luzinais et al., 2024; du Pontavice et al., 2021). Therefore, according to equations used in the steady-state version of EcoTroph, the flow kinetics during MHWs is increased as follows:

$$K_\tau = 20.19 \cdot \tau_j^{-3.258} \cdot e^{0.041 \cdot SST_y} \cdot (1+\eta_\tau), \quad (4)$$

where  $\eta_\tau$  is the MHW-associated additional loss rate.

Biomass spectra in EcoTroph-Dyn are split into trophic classes of variable width. The width of trophic classes  $[\tau_j; \tau_{j+1}[$  was determined based on the estimated mean flow kinetics so that biomass could transfer up a trophic class in each time step ( $\Delta t = 1/24$  year). Thus:

$$\Delta\tau = \tau_{j+1} - \tau_j = K_\tau \cdot \Delta t = 20.19 \cdot \tau_j^{-3.258} \cdot e^{0.041 \cdot SST_y} \cdot (1+\eta_\tau) \cdot \Delta t, \quad (5)$$

where  $\eta_\tau$ , the MHW-associated additional loss rate, is only considered during MHWs.

According to fluid dynamic equations, the EcoTroph-Dyn state variable  $B_{\tau,t}$ , represents the biomass present at time step  $t$  within the TL class  $[\tau, \tau+\Delta\tau[$ . It is expressed as:


$$B_{\tau,t} = \frac{1}{K_{\tau,t}} \cdot \Phi_{\tau,t} \cdot \Delta\tau, \quad (6)$$

where  $\Phi_{\tau,t}$  is the mean quantity of biomass flowing within the trophic class  $[\tau, \tau + \Delta\tau[$  at time step  $t$ , and  $K_{\tau,t}$  is the mean flow kinetic within the trophic class  $[\tau, \tau + \Delta\tau[$ .

170 Finally, the production  $P_{\tau,t}$  of the trophic class  $[\tau, \tau + \Delta\tau[$  at time step  $t$  is:

$$P_{\tau,t} = \int_{s=\tau}^{s=\tau+\Delta\tau} \Phi(s,t) \cdot ds = \Phi_{\tau,t} \cdot \Delta\tau, \quad (7).$$

Hence, according to equations (6) and (7), EcoTroph highlights that biomass stems from the ratio of the production to the flow kinetic. Production is expressed in  $t \cdot TL \cdot year^{-1}$ , that is, biomass in weight moving up the food web by 1 TL on average during one year.



175 **Figure 1: Schematic diagram illustrating the ecological framework represented by EcoTroph-Dyn (adapted from du Pontavice et al., 2021).** The trophic functioning of marine food webs is represented by a biomass flow, starting with biomass entering the system at trophic level 1 through net primary production (NPP). Biomass then moves through each trophic level according to trophic transfer efficiency. The flow kinetic, temperature and MHW dependant, is crucial in defining trophic class boundaries and estimating biomass for each trophic level over time. When MHWs occur, there is a loss of biomass flow at each trophic level. The empirical models used for each parameter (MHW losses ( $\eta$ ), transfer Efficiency (TE), and flow kinetic (K)) and where satellite data (SST and NPP) are incorporated into the model to account for marine heatwaves are noted in the figure.

### 3.2 MHW loss rate algorithm

185 We compute the loss rate in the biomass flow associated with MHWs based on the percentage of species undergoing thermal stress, i.e., species exposed to temperatures exceeding their thresholds. A detailed description of the MHW loss rate algorithm is described in Appendix A1. Here, we provided the main steps for the incorporation of the effects of marine heatwaves into EcoTroph-Dyn:

1. Detection and characterisation of MHWs with climatology (period 1982-2011)
- 190 2. Estimation of species distribution and associated thermal niche
3. Matching historical MHW distributions and characteristics with species distribution

4. Calculation of the percentage of species undergoing thermal stress in each ocean spatial cell
5. Based on this percentage estimation of an additional loss rate associated with MHWs

Finally, we assumed in the mathematical expression of loss rate ( $\eta_i$ ) that species are continuously challenged by increased MHW intensity, which is expressed as:

$$\eta_i = e^{-e^{b_{tl_i} \cdot \alpha \left( MHW_{cat,i} - \frac{lt_{50\_tl_i}}{\alpha} \right)}} \cdot \beta, (7)$$

where  $b_{tl_i}$  and  $lt_{50\_tl_i}$  correspond to the slope of the function and the index of marine heatwave intensity ( $MHW_{cat,i}$ ) at which 50% of the species are undergoing thermal stress in cell<sub>i</sub>,  $\beta$  corresponds to the MHW duration, with  $\beta$  ranging from  $\beta=0$  (no MHW) to  $\beta=1$  (MHW lasting 15 days (see section 3.3.1)).  $MHW_{cat,i}$  corresponds to an MHW intensity index (defined in Appendix A1) and  $\alpha$  is a coefficient assumed to represent the species' resistance capacity to MHW conditions, thus reducing the mortality rate due to species' exposure to thermal stress.

We used an  $\alpha = 0.2$  in our simulations. However, we also explored the sensitivity of the results to community resistance capacity by testing four alternative values of  $\alpha$ . The  $\alpha$  values that we tested were full resistance ( $\alpha = 0$ , i.e., we fixed  $\eta_i=0$ ; no mortality due to thermal stress), partial resistance ( $\alpha = 0.2, 0.5$ ; 20%, and 50% of the species die because of thermal stress, respectively) and no resistance ( $\alpha = 1$ ; all species die when they are under thermal stress). We performed a three-way ANOVA, that is a statistical test we used to analyse the effects of trophic levels, biomes, and  $\alpha$  value on biomass change. We also related the loss rate to the MHW duration over the 15 days by assuming that the duration increased the mortality rate.

### 3.3 EcoTroph-Dyn simulations

The EcoTroph-Dyn model was applied to simulate consumer biomass and production (between TLs 2 and 5.5) from 1998 to 2021 for each  $1^\circ \times 1^\circ$  spatial cell of the global ocean (Fig. S1). Model outputs were summarised spatially by biomes: tropical, temperate, or upwelling biomes, based on Reygondeau et al., (2013). We excluded polar biomes because net primary production (NPP) data, one of the forcing variables of the EcoTroph-Dyn model, were not available in high-latitude regions during the winter period.

#### 3.3.1 Environmental forcing data

We simulated and compared consumer biomass and production under scenarios with and without MHWs. The simulations were driven by daily SST observations from the National Oceanic and Atmospheric Administration - Advanced Very High-Resolution Radiometer (NOAA \_AVHRR) data (<https://www.ncei.noaa.gov/access/metadata/landing-page/bin/iso?id=gov.noaa.ncdc:C00680>) and NPP data predicted from the EPPLEY-VGPM algorithm and satellite remote sensing data ([http://orca.science.oregonstate.edu/npp\\_products.php](http://orca.science.oregonstate.edu/npp_products.php)).

From the daily SST time series in each ocean cell, we identified every MHW day. We defined an MHW as when the daily SSTs exceed an extreme temperature threshold value for at least five consecutive days (Hobday et al., 2016). The extreme temperature threshold value was calculated for each 1° latitude x 1° longitude spatial cell as the 90th percentile of daily SST from the 30-year historical time series from January 1982 to December 2011. We did not calculate threshold values by season; thus, MHW events were identified by a single threshold across the year. As a result, we detected MHWs mostly occurring during the year's warmest months. We used the R package *heatwaveR* described at <https://robwschlegel.github.io/heatwaveR/> (Schlegel and Smit, 2018) to detect MHWs in each spatial cell from January 1998 to December 2021.

For the scenarios 'with MHWs', we used the SST time series from NOAA \_AVHRR data.

To simulate scenarios without MHW in the SST time series, we decomposed the daily SST time series ( $Y_t$ ) of each ocean spatial cell using a Census X-11 procedure (Pezzulli et al., 2005; Shiskin, 1967; Vantrepotte and Mélin, 2011). With this method, the time series can be decomposed as:

$$Y_t = T_t + S_t + H_t ,$$

where  $Y_t$  represents the daily SST at day  $t$ ,  $T_t$  represents the long-term mean annual changes in temperature,  $S_t$  represents the seasonal component (repetitive pattern over time), and  $H_t$  represents the additional temperature variability that is not attributed to the annual trend or seasonality.

The  $T_t$  underlying long-term direction is obtained from the 365-day running average of the initial series  $Y_t$ . The seasonal component ( $S_t$ ) is then computed by applying a centered moving average to the trend-adjusted series ( $Y_t - T_t$ ). The estimation of  $S_t$  on the trend-adjusted series avoids any confusion with the inter-annual (trend) signal. After the revised estimation of these two components (see the method in Pezzulli et al., 2005; Vantrepotte & Mélin, 2011), The additional temperature variability component was computed as  $H_t = Y_t - S_t - T_t$ .

We applied the following procedure to create a daily SST time series for the scenarios without MHW. When the daily  $Y_t$  value was identified as MHW and  $Y_t$  was above the value ( $T_t + S_t$ ), we replaced the daily SST value ( $Y_t$ ) with the expected temperature without the additional variability ( $H_t$ ) component, i.e.,  $T_t + S_t$ . For MHW days with  $Y_t$  below  $T_t + S_t$  or non-MHW-days, we keep the daily SST value  $Y_t$  (see Fig. S2, where the process of creating this time series is illustrated). In our study, we can have an MHW day declared even though  $Y_t < T_t + S_t$ . This specific situation is rare (less than 0.5% of time series) and occurs because of the use of an annual threshold value to detect MHW events that mainly occurred during the year's warmest months. (See Fig. S2 just before April month for schematic visualisation). The time series created using this algorithm is referred to here as 'without MHW'.

Finally, to adapt the daily SST time series to EcoTroph-Dyn time-step resolution (1/24 year), we aggregated the initial temperature time series  $Y_t$  and the 'without MHW' time series at a 15 days scale. Specifically, we averaged the daily SST for every 15 days of a year. We then computed the



proportion of MHW days ( $\beta$ ) within each 15 days. Thus,  $\beta = 0$  when no MHW day is identified in a 15 days time step and  $\beta=1$  when an MHW last the 15 days of the time step.

Gridded monthly NPP data from 1998 to 2021 were obtained from satellite-derived estimation. The NPP data were estimated using the EPPLEY-Vertically Generalized Production Model (VGPM) computation method (Behrenfeld and Falkowski, 1997). The EPPLEY-VGPM method is a hybrid model that employs the basic model structure and parameterisation of the standard VGPM (Vertically Generalized Production Model) computation. This model estimates net primary production (NPP) based on chlorophyll concentration, incorporating the vertical distribution of primary production. The specificity of the EPPLEY-VGPM method is that the polynomial description of the maximum daily net primary production found within a given water column ( $Pb_{opt}$ , expressed in units of mg carbon fixed per mg chlorophyll per hour) is replaced by the exponential relationship described by Morel(1991), based on the curvature of the temperature-dependent growth function described by Eppley (1972).

After excluding spatial cells from the polar biome, we had a total of 34,643 cells, 13,340 of which contained incomplete time series of NPP. For cells with incomplete time series, we utilised the spline function from the R package "stats" to interpolate the missing Net Primary Production (NPP) values. In each ocean cell, the interpolation was constrained by the minimum and maximum satellite data values of the NPP observed over their respective time series. This approach ensured reliable interpolation and reduced potential bias.

Finally, to match the NPP time series with the resolution of the EcoTroph-Dyn time step (1/24 of a year), we duplicated the monthly NPP values to cover the two 15 days of each month.

### **3.3.2 Biomass simulations**

We simulated the changes in biomass spectra in each spatial cell from 1998 to 2021 for the scenarios with and without MHW. We calculated the differences in consumer biomass change between scenarios with and without MHW. Biomass change was computed using the reference period from 1998 to 2009 under the 'without MHW' scenario, allowing us to examine the projected declines in biomass specifically attributable to MHWs. We explored the sensitivity of the results to species' resistance capacity to MHW conditions using the four resistance capacity settings represented by the values of the coefficient  $\alpha$ .

We initialised the model by applying a 'burn-in' period of 12 years without any environmental forcing. Indeed, simulation testing indicates that a 'burn-in' period of at least 10 years is needed for biomass spectra to reach equilibrium (Fig. S3). Increasing the 'burn-in' period beyond 12 years does not significantly change the equilibrium biomass spectra (t test p-value=0.6118). We thus used the average of the time series (1998 to 2021) as the 'burn-in' period in each ocean spatial cell.


### 3.3.3 Northeast Pacific Ocean MHW case study

An analysis focusing on the MHW that occurred in the Northeast Pacific Ocean from 2013 to 2016 (commonly known as ‘the Blob’, Bond et al., 2015) was undertaken to assess the capability of EcoTroph-Dyn to transcribe past MHW events. First, we extracted the EcoTroph-Dyn simulation outputs (with and without MHW) in the region, delineated by the boundary of nine biogeochemical provinces that had been identified by Reygondeau et al., (2013) and adapted from Longhurst, (2007). These biogeochemical provinces include: Central American coast (CAMR), California Ocean and Coastal current (OCAL and CCAL), Alaska coastal downwelling (ALSK), North Pacific Tropical gyre (NPTG), Northeast Pacific subtropical (NPSE), North Pacific polar front (NPPF), Eastern Pacific subarctic gyres (PSAE), and the Western Pacific subarctic gyres (PSAW). Second, we computed biomass change using the reference period from 1998 to 2009 under the ‘without MHW’ scenario, allowing us to examine the projected declines in biomass specifically attributable to ‘the Blob’. Finally, we discussed and compared our results with the literature.

## 4. Results


### 4.1 Environmental forcing with and without MHWs

In response to climate change, the NPP and SST are already perturbed. Global total NPP decreased by 1% in the period 2015-2021 compared to the 1998-2009 period. However, large spatial variability was observed in NPP changes (Figure 2a). In particular, NPP decreased by 20% in the Northeast Pacific, while an increase of 20% was estimated for the Southern Ocean. Under the ‘without MHW’ scenario (Figure 2b), global average SST increased by 0.28°C in the period 2015-2021 compared to the 1998-2009 period, with some areas warming up to 0.5°C. However, under the ‘with MHWs’ scenario (Figure 2c), SST increased by 0.32°C in 2015-2021 compared to the period 1998-2009, with some areas, such as the Northeast Pacific, warming up by 1°C over the same period. Globally, the estimated increase in average SST from 2015 to 2021 was 4% higher in the ‘with MHWs’ scenario compared to the ‘without MHW’ scenario.



**Figure 2: Changes in SST and NPP between the average of 2015-2021 relative to the average between 1998-2009. (a) NPP, (b) SST under the ‘without MHW’ scenario, and (c) SST under the ‘with MHWs’ scenario.**

Under the ‘with MHWs’ scenario, MHWs occurring during the year's warmest month increased in intensity, duration, and surface extent from 1998 to 2021 (Figures 3a, b) with large spatial variability (Figures 3c, d). MHWs with intensity lower than 3°C above the climatology were identified on average in 28.5 % of the ocean surface (Figures 3a). These MHWs lasted, on average, more than 40 days (Figures 3b). In contrast, MHWs characterised as higher intensity ( $\geq 3^{\circ}\text{C}$  above climatology) were identified in <20% of the ocean surface area (Figures 3a). These relatively more intensive MHWs lasted, on average, 32 days (Figures 3b). Furthermore, more MHW days of lower intensity were identified for low latitude regions (23°N - 6°S) (Figure 3c, 3d) compared to MHW days identified in higher latitude regions (> 23°N and 25°S).




**Figure 3: Temporal and spatial characteristics of MHWs identified for the period 1998 to 2021.** (a) Changes in the percentage of the oceans' total surface area with MHW in each year categorised by their intensity, (b) Evolution of MHWs averaged duration categorised by their intensity, and (c) Average duration of each MHW event in days that occurred over the period 2015-2021. (d) Average intensity of each MHW event over the period 2015-2021.

## 4.2 Biological impacts of MHWs at the global scale

### 4.2.1 Impact on total consumer biomass

The projected changes in total consumer biomass were higher under the 'with MHWs' scenario relative to the 'without MHW' scenario. Globally, total consumer biomass was projected to decrease, on average by  $0.07 \pm 0.02\%$  per year (standard error) relative to the baseline period of 1998-2009 (Figure 4a) (GLS, Generalized Least Squares,  $p$ -value  $< 0.05$ ). In contrast, simulations under the 'with MHWs' scenario with an MHW resistance capacity setting of  $\alpha=0.2$  projected an average decrease in total consumer biomass of  $0.12 \pm 0.02\%$  per year relative to the baseline period of 1998-2009 ( $p$ -value  $< 0.05$ ). Thus, simulations that focused on annual mean changes in temperature only masked the effects of MHWs on the long-term changes in consumer biomass in the ecosystem (Fig. S4).



**Figure 4: Simulated total consumer biomass change in the ocean from 1998 to 2021 under the ‘without MHW’ (black lines) and with MHWs (light orange lines) scenarios, with a resistance capacity setting of  $\alpha=0.2$ .** (a) the world ocean (excluding polar biome), (b) temperate biomes, (c) tropical biomes, and (d) upwelling biomes, respectively (S1 for biome spatial definition). The horizontal grey line separates positive and negative total consumer biomass changes.


While total consumer biomass was projected to decrease slightly across the three biomes (temperate, tropical, and upwelling biomes represented in Figures 4b, c, and d, respectively), in the scenario ‘with MHWs’ the declines in tropical biomes were larger than the global average. Under the ‘without MHW’ scenario, total consumer biomass in temperate biomes was projected to increase by  $0.01\% \pm 0.01$  (standard error), while in tropical and upwelling biomes, it was projected to decrease by  $0.18\% \pm 0.01$ , and  $0.02\% \pm 0.01$  per year, respectively, relative to the baseline period. Under the ‘with MHWs’ scenario with  $\alpha=0.2$ , total consumer biomass in temperate, tropical, and upwelling biomes was projected to decrease more strongly by  $0.03\% \pm 0.01\%$ ,  $0.23\% \pm 0.02\%$ , and  $0.1\% \pm 0.02$ , respectively, relative to the baseline.

#### 4.2.2 Impacts by trophic level

MHWs exacerbated the projected climate change impacts on biomass, particularly for higher trophic level groups. Indeed, our model projected a similar level of biomass loss across trophic levels in the ‘without MHW’ scenario, with a projected decrease in global total consumer biomass by  $1 \pm 0.1\%$  SE in the period 2015-2021 relative to 1998-2009 (Figure 5a). In contrast, under the ‘with MHWs’ scenario and with a resistance capacity setting of  $\alpha=0.2$ , total consumer biomass was projected to decrease by  $4.4 \pm 0.1\%$  for high trophic level classes ( $TL \in [4;5.5]$ ), while the decrease was smaller for

low trophic level ( $TL \in [2;3[$ ,  $3.4 \pm 0.1\%$ ) and mid trophic level ( $TL \in [3;4[$ ,  $4.1 \pm 0.1\%$ ) classes (Figure 5e).

The impact on high trophic levels differed between biomes, with the tropical and upwelling biomes being notably more impacted. Under the ‘without MHW’ scenario, our model projected a decrease in high trophic level consumer biomass of  $0.5 \pm 0.1\%SE$ ,  $2.4 \pm 0.1\%$ , and  $2.2 \pm 0.1\%$  in temperate, tropical, and upwelling biomes, respectively, in the period 2015-2021 relative to 1998-2009 (Figures 5b, c, and d). In contrast, under the ‘with MHWs’ scenario and with a resistance capacity setting of  $\alpha=0.2$ , high trophic level consumer biomass was projected to decrease by  $3.3 \pm 0.1\%$ ,  $6.6 \pm 0.1\%$ , and  $5.9 \pm 0.1\%$  in temperate, tropical, and upwelling biomes, respectively, while a smaller biomass decrease for low and mid-trophic level was expected (Figures 5e, g, and h).




**Figure 5: Projected changes in consumer biomass by trophic level and biome under the ‘without MHW’ and ‘with MHWs’ scenarios relative to the 1998-2009 average.** Low trophic level ( $TL \in [2;3[$  in black), medium TL ( $TL \in [3;4[$  in blue), and high TL ( $TL \in [4;5.5]$  in red). (a, e) correspond to global scale, (b, f) to the temperate biome, (c, g) to the tropical biome, and (d, h) to the upwelling biomes. (a, b, c, and d) correspond to ‘without MHW’ scenarios while (e, f, g, and h) correspond to ‘with MHWs’ scenarios. The simulation results under the ‘with MHWs’ scenario used a resistance capacity setting of  $\alpha=0.2$ .

#### 4.2.2 Various responses of ecosystems to MHWs

Over the 2015 to 2021 period, MHWs impacted total consumer biomass with large variations in both the direction and magnitude of the impacts between biomes (Figure 6). Under the ‘without MHW’ scenario, the model projected an increase in total consumer biomass in 33.5% of the ocean area, especially in temperate biomes (Figure 6a). Under the ‘with MHWs’ scenario, this proportion



decreased and a biomass increase was projected to occur in only 24% of the global ocean, with a projected global biomass decrease of 4.8% in the period 2015-2021 relative to 1998-2009 compared to an only 2.4% biomass loss under 'without MHW' scenario. In high latitudes ( $> 25^{\circ}\text{N}$  and  $25^{\circ}\text{S}$ ), MHWs exhibiting high intensity but short duration (Figures 3c and d) resulted in moderate additional biomass losses, up to 8% in 2015-2021 (Figure 6c). Conversely, in low latitudes ( $25^{\circ}\text{N} - 25^{\circ}\text{S}$ ), MHWs led to substantial additional biomass losses, exceeding 10% on average (Figure 6c) and up to 20% in biodiversity hotspots such as Indonesia, off the coast of Papua New Guinea, and Central America. Thus, high latitude and upwelling areas appeared to be refuge zones from MHWs compared to low latitudes. Looking at the MHW's additional impact within food webs (Figs. S5, S6, and S7), similar spatial patterns and responses are projected with a higher impact (biomass decreases) on high trophic levels compared to lower trophic levels.




**Figure 6: Changes in total consumer biomass in 2015-2021 compared to 1998-2009.** (a) total consumer biomass under the “without MHW” scenario, (b) total consumer biomass under the ‘with MHWs’ scenario with  $\alpha=0.2$ , and (c) differences in consumer biomass between scenarios.

### 4.3 Sensitivity to resistance capacity setting ( $\alpha$ )

Ecosystem total consumer biomass response to MHWs depended on community resistance capacity. Assuming species to have full resistance ( $\alpha=0$ ) or no resistance ( $\alpha=1$ ) to MHWs, total consumer biomass was projected to decrease by 2.3% and 15.3% in the period 2015-2021 relative to 1998-2009, respectively (Figure 7a). Simulations using different  $\alpha$  values projected changes in total consumer biomass that were consistent in direction but differed significantly in magnitude. A three-way ANOVA was performed to analyse the effect of trophic levels, biomes, and  $\alpha$  value on biomass change. This ANOVA revealed a statistically significant effect on biomass change of trophic levels, biomes, and  $\alpha$  values ( $p < 2e-16$ ), individually. Additionally, the interaction between biomes, trophic levels, and  $\alpha$  value also had a significant effect on biomass change ( $F(27, 16737) = 442.4$ ,  $p\text{-value} < 2e-16$ ). The lower the resistance capacity of species (higher  $\alpha$  value), the greater the loss of total consumer biomass. On a biome scale, these variations in biomass loss were greater for the tropics, with values ranging from 3.1% ( $\alpha=0$ ) to 20.2% ( $\alpha=1$ ) (Figure 7c). In comparison, sensitivity was lower in the

temperate and upwelling biomes with a decrease between 1.4% ( $\alpha=0$ ) and 11.2% ( $\alpha=1$ ) and 3.8% ( $\alpha=0$ ) and 12.3% ( $\alpha=1$ ), respectively (Figures 7b and d).



**Figure 7: Sensitivity of changes in total consumer biomass to different resistance capacity settings. Changes are aggregated for the whole biomass spectrum (TCB) and by trophic level (TL) (low TL, medium TL and high TL). Colours represent different resistance capacity settings ( $\alpha$  values). (a) Global ocean, (b) temperate biomes, (c) tropical biomes, and (d) upwelling biomes. The different horizontal lines of the box plot refer to the median, 25<sup>th</sup>, and 75<sup>th</sup> quantiles.**

The sensitivity of changes in total consumer biomass to the resistance setting increased with trophic level. At a global scale, the response of low TLs ranged from a consumer biomass decrease of 1.4% ( $\alpha=0$ ) to 8.0% ( $\alpha=1$ ), while high TLs exhibited values ranging from 0% ( $\alpha=0$ ) to 11% ( $\alpha=1$ ) (Figure 7a). Furthermore, the sensitivity of trophic levels to the  $\alpha$  value was greater in the tropical biome and lower in the temperate and upwelling biomes relative to that of the global ocean.

#### 4.4 A case study of a Northeast Pacific MHW


Our model projected that the MHW in the Northeast Pacific from 2013 - 2016 ('the Blob') resulted in long-term changes in the biomass spectrum in the region (Figure 8a). Temperature anomalies were on average  $\geq 4$  °C (between 2013 to 2016) and up to 8 °C in 2015 relative to 1982-2011. This temperature anomaly had some ecological repercussions. On average, without accounting for MHWs, total consumer biomass in our simulation was hindcast to decrease by  $1.5\% \pm 0.3$ ,  $3.1\% \pm 0.4$ ,  $6.6\% \pm 0.2$ , and  $5.2\% \pm 0.3$  in 2013, 2014, 2015, and 2016, respectively, compared to the reference period of 1998 to 2009 (black line in Figure 8b). However, when accounting for MHWs with  $\alpha=0.2$ , the total consumer biomass loss decreased on average by an additional 3.1% in 2013, 2014, 2015, and



2016. Furthermore, the difference in linear trend (slope and intercept) between the biomass time series 1998 to 2013 and 2017 to 2021 indicated that the MHW ('the Blob') had a significant effect on long-term changes in biomass (GLS, Generalized least squares,  $p\_value < 0.05$ ).

The magnitude of the hindcast total consumer biomass differed significantly between biogeochemical provinces (ANOVA,  $F(1, 2906) = 7.854$ ,  $p\_value < 0.05$ , Figure 8c). Comparing the consumer biomass before (1998 to 2012) and during (2013 to 2016) the occurrence of 'the Blob', all biogeochemical provinces in the Northeast Pacific, except Alaska coastal downwelling (ALSK) and Central American coast (CAMR), exhibited a significant total consumer biomass decrease (ANOVA,  $F(1, 151) = 155.443$ ,  $p\_value < 2e-16$ ) under the scenario with and without MHWs. Under the 'without MHW' scenario, total consumer biomass in the North Pacific Tropical gyre (NPTG) biogeochemical provinces, followed by the California current (CCAL+OCAL biogeochemical provinces), was hindcast to decline the most amongst the provinces when comparing before, during, and after 'the Blob' (Figure 8c). Under the 'with MHWs' scenario and with a resistance capacity setting of  $\alpha=0.2$ , all biogeochemical provinces were hindcast to have an additional biomass decrease during and after 'the Blob'. This additional biomass decrease ranged from 0.9% to 5% and 1.4% to 5.1% during and after 'the Blob', with an average decrease of  $2.7\% \pm 0.4\%$  and  $3.1\% \pm 0.3\%$ , respectively (Figure 8c). Particularly, the California current and the Alaska coastal downwelling provinces were most affected by the MHW, with additional biomass decreases of 5% and 3.8% relative to the 'without MHW' scenario, during and after 'the Blob'.

The impact of 'the Blob' on various trophic levels was similar to the global pattern, with a rapid reaction (in terms of biomass loss) observed at lower TLs and higher TLs exhibiting a delayed response (S8). Considering the influence of MHWs from 2013 to 2016 using the 'with MHWs' scenario and  $\alpha=0.2$  resistance capacity, there was a hindcast biomass reduction of  $6.8\% \pm 0.7$ ,  $6.3\% \pm 0.6$ , and  $4.3\% \pm 0.3$  for low, medium, and high TLs, respectively, when compared to the reference period of 1998 to 2009. Compared to pre-event conditions, by 2021, aside from low TLs, medium and high TLs gave no sign of recovery from 'the Blob' MHW event.



**Figure 8: Hindcast of temporal and spatial biological impacts of the MHW from 2013 to 2016 in the Northeast Pacific ('the Blob').** (a) Temperature anomalies between 2013 and 2016, (b) biogeochemical provinces definition, with Central American coast (CAMR), California Ocean and Coastal current (OCAL and CCAL), Alaska coastal downwelling (ALSK), North Pacific Tropical gyre (NPTG), Northeast Pacific subtropical (NPSE), North Pacific polar front (NPPF), Eastern Pacific subarctic gyres (PSAE), and the Western Pacific subarctic gyres (PSAW). (c) Generalised least squares models (lines) and total consumer biomass change in the 'without MHW' and 'with MHWs' scenarios in the black and light orange colour, respectively. The vertical grey shaded area indicates the duration of 'the Blob'. (d) Average total consumer biomass changes before (1998 to 2012, pink), during (2013 to 2016, green), and after (2017 to 2021, blue) 'the Blob' for each biogeochemical province. The error bars correspond to standard errors.

485 Trophic dynamics of the biomass spectra in the Northeast Pacific were hindcast to be impacted  
by ‘the Blob’ (S9). The trophodynamic indicators flow kinetic and trophic efficiency increased and  
decreased, respectively, in the model. Under the scenario ‘without MHW’, the average hindcast flow  
kinetic of the biomass spectra in the Northeast Pacific increased by  $0.6\% \pm 0.2$ , while the transfer  
efficiency decreased by  $0.7\% \pm 0.2$  in the period 2013-2016 relative to 1998-2009. Consideration of the  
490 MHW under the ‘with MHWs’ scenario exacerbated the increase in flow kinetic and decrease in trophic  
efficiency significantly by 0.7% and 0.4%, respectively (t test p-value < 0.05).

## 5. Discussion

In this study, we accounted for MHWs in the last four decades using hindcast simulations and  
showed the potential of synergic impacts of MHWs (pulses) and long-term climate change (presses)  
495 on biomass and trophodynamics of ecosystems (Bender et al., 1984; Harris et al., 2018). These  
reconstructed MHW impacts varied spatially because of MHW spatial dynamics, regional differences  
in ocean biogeochemical and physical conditions and ecosystem trophodynamic characteristics.  
Furthermore, studying ‘the Blob’, we highlighted that MHWs could exacerbate the long-term impacts  
of climate change that vary between biogeochemical provinces.

### 5.1 Impacts of MHWs on global marine consumer biomass

Over the past two decades, MHWs have led to increased mortality rates and significant  
alterations in the functioning and structure of ecosystems (Smith et al., 2023; Wernberg et al., 2013,  
2016). Our modelling analysis suggests that, without accounting for MHWs and their ecological effects,  
the decline in global-scale biomass associated with climate change may have been significantly  
505 underestimated. We show that MHWs exacerbated the impacts of long-term climate change,  
impacting trophodynamic parameters such as the flow kinetic and transfer efficiency within  
ecosystems, which is congruent with studies by Arimitsu et al. 2021, Gomes et al. 2024, Smith et al.  
2023 . These perturbations of ecosystem functioning result in biomass loss through food webs.  
Although MHW temperature anomalies have mostly lasted for weeks to months only, we emphasise  
510 that the resulting ecosystem perturbations last for a longer time (decades) and influence long-term  
biomass change (Figure 8, Babcock et al., 2019; Cheung & Frölicher, 2020; Guibourd de Luzinais et al.,  
2024). Finally, the intensity and duration of MHWs influenced the magnitude of the perturbation in  
ecosystem functioning (Oliver et al., 2021; Smith et al., 2023). We highlighted that the intensity and  
duration of MHWs have continuously increased since the beginning of the 21st century (Figure 3),  
515 leading to a sharp biomass decrease over the hindcast period (Figure 4). Such short-term biomass  
decreases strengthened the impacts of long-term climate change (Cheung and Frölicher, 2020; Collins  
et al., 2019).

At a global scale, over the whole time series and different MHW resistance capacity scenarios, high TL biomass experienced greater impacts from MHWs, and was not able to recover to pre-perturbation levels as effectively as the low and medium TL biomass. The gap in biomass recovery was most apparent in tropical and upwelling biomes, where the hindcast biomass of high TLs consistently decreased over time. The time needed for the biomass to return to pre-perturbed levels was related to the rate of biomass turnover, which was also dependent on the speed of biomass flows. This turnover decreased with trophic level and was low for organisms at the top of the food chain (Gascuel et al., 2008; du Pontavice et al., 2020; Schoener, 1983). Thus, the high frequency of MHWs in recent years may be greater than the time that high TLs need to recover from the impacts of individual MHWs and could contribute to the continued decline in high-TL biomass.

## **5.2 Trophodynamics approach as a framework for deciphering ecosystem responses to MHWs**

In our modelling approach, trophodynamic changes were assessed through transfer efficiency (TE) and flow kinetics. TE (dimensionless) is defined as the fraction of energy transferred from one trophic level (TL) to the next and summarises all the losses in the trophic network (Jennings et al., 2002; Libralato et al., 2008; Lindeman, 1942; Niquil et al., 2014; Pauly and Christensen, 1995; Schramski et al., 2015; Stock et al., 2017). It is an emergent property of marine ecosystems and an essential parameter in many applications of marine ecology, such as estimating biomass flux in production models (e.g., Carozza et al., 2016; du Pontavice et al., 2021; Gascuel et al., 2011; Jennings et al., 2008; Tremblay-Boyer et al., 2011). Unlike previous applications of EcoTroph at an annual timescale (du Pontavice et al., 2021, 2023), where changes in transfer efficiency reflected long-term changes in species assemblages, having a transfer efficiency that evolved on a biweekly basis (14 days) allowed us to take into account metabolic fluctuations over a shorter period. The temperature during MHWs increases the basal metabolism (catabolism) of marine organisms (Grimmelpont et al., 2023; Minuti et al., 2021), thus reducing the energy available for anabolic processes that can only occur once catabolic needs are met (Eddy et al., 2021). The estimation of transfer efficiency only considers anabolic processes (Eddy et al., 2021); hence, this balance between catabolism and anabolism is how MHWs could disrupt transfer efficiency on a biweekly scale. The flow kinetics expressed in  $\text{TL} \cdot \text{year}^{-1}$  quantifies the speed of the trophic flux, i.e., the rate of biomass transfers from lower trophic levels to higher ones, due to predation and/or ontogenesis (Gascuel et al., 2008; du Pontavice et al., 2020). This rate is inversely proportional to the biomass residence time (BRT, du Pontavice et al., 2020), which is the average time a unit of biomass spends moving from one TL to the next higher one through predation (Gascuel et al., 2008; Schramski et al., 2015). This flow kinetics can be measured by the

production/biomass ratio (P/B, Gascuel et al., 2008), which is used in many ecosystem models, particularly EwE (Ecopath with Ecosim, Christensen & Pauly, 1992).

MHWs disrupt the metabolism of individuals and, on a larger scale, impact the entire ecosystem with trophodynamic changes throughout the trophic network (S9, Arimitsu et al., 2021; Collins et al., 2019; Gomes et al., 2024; Smith et al., 2023). However, due to their different physical characteristics depending on the ecosystems and the varying functioning of these ecosystems, they do not induce the same structural and functional changes, leading to different biomass losses. Tropical ecosystems are composed of species with low transfer efficiency (TE, low ratio between stored energy and ingested energy), with significant energy losses that increase with temperature (Brown et al., 2004; du Pontavice et al., 2020; Schramski et al., 2015). To compensate for this low efficiency, predation activity is high, resulting in rapid biomass transfers between prey and predators (biomass flow, du Pontavice et al., 2020). In these ecosystems, communities (which may be dominated by short-lived and fast-growing species) are generally living at the upper end of their thermal preferences (Begon and Townsend, 2021; Pinsky et al., 2019; Vinagre et al., 2016). They experience thermal stress associated with MHWs, leading to substantial mortality across the trophic network, with a higher proportion occurring in lower trophic levels regardless of the intensity of the MHWs as low TLs tend to have lower thermal limit than high TLs (Guibourd de Luzinai et al., 2024). MHWs scarcely affect transfer efficiency (TE), with an estimated average decrease of 0.05% between 1998 and 2021 because the metabolism of species is already very high (du Pontavice et al., 2020). However, MHWs exacerbate prey mortality rates, resulting in proportionally higher predation rates; i.e., biomass remains for a shorter time at each trophic level, corresponding to an acceleration of biomass flow (estimated at 1 % on average between 1998 and 2021). Given the relatively high number of MHW days (Figure 3, Hobday et al., 2018; Marin et al., 2021; Oliver et al., 2018), these modifications to biomass flow are "persistent" and involve significant biomass losses.

In temperate ecosystems, biomass is transferred more slowly between trophic levels with less loss (higher trophic efficiency, du Pontavice et al., 2020; Eddy et al., 2021). In EcoTroph-Dyn, temperate communities experience thermal stress and mortality only during high-intensity MHWs (Guibourd de Luzinai et al., 2024). MHWs affect the metabolic efficiency of species by increasing basal metabolism and respiration-related losses, which decreases TE across the trophic spectrum, estimated in our simulations at an average of 0.2% between 1998 and 2021. These higher metabolic demands lead to an increase in predation activity, which, combined with lower mortality (compared to tropical waters), reduces the biomass residence time at each trophic level in the food chain, i.e., a slight acceleration of the speed of biomass flow between each trophic level (estimated in our simulations at an average of 0.3% between 1998 and 2021). Since MHWs in these ecosystems are characterised by significant

temperature anomalies but over relatively short durations (Figure 3, Frölicher et al., 2018; Oliver et al., 2018), these modifications to biomass flow are "temporary" and involve lower biomass losses.

Finally, in upwelling ecosystems, communities are characterised by low transfer efficiencies with high biomass residence times, indicating slow biomass transfers between prey and predators (du Pontavice et al., 2020). MHWs cause thermal stress and moderate mortality (Guibourd de Luzinais et al., 2024). They lead to a decrease in the metabolic efficiency of species (estimated in our simulations at an average of 0.03% between 1998 and 2021), along with an increase in predation activity, moderately impacting the rate of biomass flows within the ecosystem (estimated in our simulations at an average of 0.3% between 1998 and 2021). The MHWs occurring in these ecosystems are of high intensity and last relatively long (an average of 23 days per year between 1998 and 2021). One might expect more significant impacts than in tropical environments; however, this was not the case. This could potentially be explained by (i) the high productivity of these ecosystems, which supports a highly biodiverse food web (Largier, 2020; Pauly and Christensen, 1995; Rykaczewski and Checkley, 2008; Ryther, 1969), leading to better resilience to environmental changes and extreme temperature events (Bernhardt and Leslie, 2013), and (ii) the specific functioning of these ecosystems with cool water rising from depth to the surface, tends to reduce the number of MHW days compared to their adjacent open ocean (Varela et al., 2021). More generally, it has been highlighted that ocean warming does not affect coastal regions with upwelling in the same way as the open ocean (Varela et al., 2021).

Generally, in our simulations, higher temperatures and frequent mortality events promoted the emergence of species and the growth of populations adapted to warm waters, both characterised by rapid growth and short lifespans (Beukhof et al., 2019; du Pontavice et al., 2020). This phenomenon was observed, for example, during 'the Blob' event in the California Ocean and Coastal Current (OCAL and CCAL) provinces, with an increase of tropical species that usually live much further south, such as tuna, sailfish, and marlin (Cavole et al., 2016).

The specific examination of 'the Blob' allowed us to elucidate how this MHW differentially affected the biogeochemical provinces (Longhurst, 2007; Reygondeau et al., 2013) and influenced their long-term responses. According to our simulations (Figure 8), the biomass in the affected oceanic regions had not returned to their reference levels (1998-2009) by 2021. Following the MHW, all biogeochemical regions experienced significant biomass losses (ANOVA,  $p\_value < 0.05$ ), though with varying magnitudes. Differences in exposure to the intensity and duration of temperature anomalies can potentially explain these differences in responses. For example, the coastal part of the California Current and the subarctic gyres of the North Eastern Pacific were subject to lower anomalies over different durations (Varela et al., 2021). In addition, differences in the structure and functioning of trophic networks may also have contributed to the variation in responses to MHWs between provinces (Morgan et al., 2019; Peterson et al., 2017; Ruzicka et al., 2012). The biogeochemical zones have

different compositions of pelagic communities that may respond differently to MHWs (Peterson et al., 2017). For instance, the Gulf of Alaska saw its planktonic community shift towards smaller plankton and zooplankton, resulting in a loss in the nutritional quality and quantity of the forage portion of the pelagic community for their predators, highlighting the disruption of biomass flow (Arimitsu et al., 2021; Piatt et al., 2020; Suryan et al., 2021). In contrast, in the California Current biogeochemical region, the MHW was associated with a substantial increase in the abundance of pyrosomes which implied a limitation of energy flow moving toward higher trophic levels (Gomes et al., 2024). This was not necessarily the case for the oceanic part of the California Current. These different changes in the composition and abundance of the lower trophic levels are represented in EcoTroph-Dyn by changes in the amount of energy flowing through the food web. EcoTroph-Dyn also takes into account the differences in trophodynamic characteristics (e.g., transfer efficiency, flow kinetics) between biogeochemical zones. For example, communities in the Gulf of Alaska are more efficient than those in the Californian Current (du Pontavice et al., 2020), and the energy entering the food web was less disrupted than in the California Current, which may explain the greater impact of the MHW on the California Current.


### **5.3 Model validation and sources of uncertainties**

We have developed an innovative approach to modelling marine ecosystems, linking trophic ecology with MHW hindcast to assess their ecological impacts on a seasonal time scale and at a global scale. Despite its apparent simplicity and the reduced number of parameters, EcoTroph-Dyn is part of the family of "complete ecosystem models and dynamic system models" (Plagányi, 2007) as it represents all trophic levels, from primary producers to top predators. It merges individual "species" into categories defined solely by their trophic level and describes ecosystems through a continuous distribution of biomass (trophic biomass spectrum), from primary producers to top predators. EcoTroph-Dyn does not account for specific climate effects on individual species and populations. The model assumes that variations in environmental conditions will lead to new biomass transfer dynamics in theoretical ecosystems at equilibrium. EcoTroph-Dyn meets the criteria described by Link, (2010) to be considered a plausible representation of marine ecosystems. These criteria include: i) the biomass values of all functional groups must cover 5 to 7 orders of magnitude, ii) there must be a 5 to 10% decrease in biomass density (on a logarithmic scale) for each unit increase in trophic level, iii) specific biomass production values (P/B) must never exceed specific biomass consumption values (C/B), and iv) the ecotrophic efficiency (EE) for each group must be less than 1. Additionally, the model relies on empirically obtained equations (Gascuel et al., 2008; du Pontavice et al., 2020) and has demonstrated its ability to replicate the effects of fishing and climate change on marine ecosystems (e.g., du Pontavice et al., 2021, 2023; Tremblay-Boyer et al., 2011).



To discuss the validity of our hindcasts, we focussed on the most studied MHW ('the Blob') in the literature with available quantitative data (e.g., Arimitsu et al., 2021; Bond et al., 2015; Cavole et al., 2016; Cheung & Frölicher, 2020; Gomes et al., 2024; Piatt et al., 2020; Suryan et al., 2021). The historical biomass changes simulation from our 'with MHWs' simulations and the  $\alpha=0.2$  scenarios showed the same biomass evolution per trophic group as the observational data based on species biomass surveys (Suryan et al., 2021). However, our historical simulations estimated lower biomass and also showed a smaller biomass decrease compared to observed data (Suryan et al., 2021). This would indicate a possible underestimation of biomass and ecosystem response to MHWs by EcoTroph-Dyn. The underestimation of ecological responses to MHWs is likely caused by the choice of a lower  $\alpha$  value that lowers the sensitivity of the ecosystem to MHWs. To reduce the uncertainty over the  $\alpha$  value, future studies could calibrate it for specific region using observational data of MHWs impacts on marine ecosystems' biomass.

Our simulations of biomass changes, according to the 'with MHWs' and  $\alpha=0.2$  scenarios, also fell within the range of biomass historical simulations reported by studies using other approaches to represent ecosystem functioning (e.g., species-based Dynamic Bioclimate Envelope Model (DBEM) by Cheung & Frölicher, 2020, and Ecotran, an extension of the popular Ecopath modelling framework by Gomes et al., 2024, see Figure 9). Additionally, we projected that the trophodynamic kinetic parameter of biomass flow increased by 3% in the 'with MHWs' and  $\alpha=0.2$  scenarios, which is consistent with the estimates highlighted by Gomes et al., 2024 (3.7%). Finally, the decreases in trophodynamic transfer efficiency parameters highlighted in EcoTroph-Dyn with 'the Blob' case study correspond to previous estimates (e.g., Arimitsu et al., 2021) showing a reduction in the availability/quality of food in the food web. It is worth noting that historical simulations obtained using a smaller (larger)  $\alpha$  led to an underestimation (overestimation) of biomass losses and changes in biomass flow parameters relative to the estimates of Cheung & Frölicher (2020) and Gomes et al. (2024) estimates.



**Figure 9: Biomass change associated with the Northeast Pacific MHW (2013-2016) per biogeochemical province.** Grey violin plots correspond to results from Cheung & Frölicher (2020), while the red ones correspond to our hindcast EcoTroph-Dyn simulation with  $\alpha=0.2$ . (Figure adapted from Cheung & Frölicher (2020)).

One of the main uncertainties in modelling MHWs in EcoTroph-Dyn is the assumption of the resilience of the biomass spectrum to MHWs, i.e., parameter  $\alpha$ . It is important to note that the  $\alpha$  values applied in this study were chosen arbitrarily, albeit reasonably, and were able to capture a broad range of potential responses to MHWs. “It would therefore have been valuable to test EcoTroph-Dyn against other MHWs in the world ocean in order to acquire better estimates of the  $\alpha$  parameter and more reliable results regarding the consequences of MHWs. However, to date, few MHWs are as well studied as ‘the Blob’, limiting these analyses. As discussed in Guibourd de Luzinais et al., 2024, given the diversity of ecosystem functioning, there is no reason why communities in different ecosystems, biogeochemical regions or trophic levels should have the same capacity to resist MHWs. Thus, we encourage future studies to use various observational data on the impacts of MHWs across the ocean (if available) to better estimate the  $\alpha$  parameter as a function of different ecosystems, biogeochemical regions or trophic levels in order to reduce the uncertainties in projections.

Here, we focused solely on the direct impacts of MHWs occurring during the year's warmest month via thermal stress, resulting in species mortality (Oliver et al., 2021; Smith et al., 2023). However, MHWs in other seasons can also have consequences on populations by affecting a specific stage of the life cycle of certain species (Crickenberger and Wetthey, 2018; Oliver et al., 2021; Smith and Thatje, 2013). For example, MHWs that stress adult breeders can lead to a decrease in reproductive investment and, consequently, fewer, smaller, and lower quality gametes (e.g., Shanks et al., 2020), resulting in a loss of abundance and biomass of some species (Johansen et al., 2021). While taking seasonality into account will increase the number of detected extreme events, some may

not have ecological consequences (Oliver et al., 2021; Smith et al., 2023). Thus, our approach can be seen as conservative and may underestimate the impact of MHWs. Nevertheless, we detected MHWs with potentially significant ecological impacts. Studying MHWs occurring in seasons other than summer would involve considering the phenological effects of MHWs. However, since the EcoTroph-Dyn model does not directly represent this phenological aspect of marine organisms, future studies can apply other approaches that explicitly represent seasonal processes such as spawning and migration to elucidate the effects of phenology.

Furthermore, uncertainties in our results arise from the environmental drivers of EcoTroph-Dyn. In our current investigation, EcoTroph-Dyn was driven by satellite data. For the NPP forcing variable, we implemented the EPPLEY-VGMP algorithm (Behrenfeld and Falkowski, 1997; Morel, 1991), which, like other algorithms such as CbPM and CAFE, derives NPP from satellite-derived estimates of Chl-a and SST. Firstly, in this study, we did not consider the 'with' and 'without' MHWs scenarios for NPP. Acknowledging that MHWs generally increase NPP at high latitudes while decreasing it at low latitudes (Arteaga and Rousseaux, 2023; Bouchard et al., 2017; Le Grix et al., 2022; LeBlanc et al., 2020), we may have overestimated MHWs impacts on ecosystem functioning at high latitudes and underestimated their impact at low latitudes. Furthermore, the variability and uncertainty in the estimation of the NPP by satellites directly affects the reliability of our results. In EcoTroph-Dyn, as well as in other Marine Ecosystem Models, ocean primary production (and its related phytoplankton biomass) plays a crucial role in both sustaining and constraining the biomass of higher trophic levels (e.g., Blanchard et al., 2012; Carozza et al., 2016; Cheung et al., 2011; Jennings & Collingridge, 2015). However, there exists significant variability in NPP estimation among satellite NPP algorithms (Milutinović and Bertino, 2011; Westberry et al., 2023). These discrepancies are particularly pronounced over continental shelves and oligotrophic gyres, primarily due to variations in model parametrisation and growth rate representation (Milutinović and Bertino, 2011; Westberry et al., 2023). Secondly, in this study, in order to propose a suitable representation of the world ocean, we use an interpolation method to reconstruct an incomplete NPP time series. The interpolation was constrained by the minimum and maximum satellite data values of the NPP observed over their respective time series to ensure reliable interpolation and reduce potential bias. Thirdly and lastly, in this study, we duplicated NPP monthly value to able the EcoTroph-Dyn to run at a 15 days basis. This duplication may have smoothed marine ecosystem response to the historical changes in marine environment; However, it has not changed trends and conclusions of our results. Consequently, elucidating the sources of the current uncertainty associated with satellite-derived NPP and refining these estimates pose significant challenges in comprehending the responses of marine food webs to MHWs.

Moreover, EcoTroph-Dyn does not account explicitly for species; thus, we could only assess aggregated food web responses to MHWs. The model projected that the aggregated response across species is expected to be negative, although it is known that some species would exhibit positive effects from MHWs occurrence (Cavole et al., 2016; Smith et al., 2023; Suryan et al., 2021; Wernberg et al., 2016). To be cautious, we considered various loss rate scenarios to obtain a complete range of responses from marine ecosystems. Running the aforementioned five MHW-induced loss rate-scenarios, we found that the aggregated resistance capacities  $\alpha$  had a significant (ANOVA,  $p\_value < 2e-16$ ) and large effect. All scenarios are consistent with each other, with an increasing biomass loss over marine ecosystems with decreasing resistance capacities (increasing  $\alpha$ ). Even though the global impact of MHWs is negative, species explicit modelling could improve our understanding of how various impacts of climate change and species-level responses will affect trophodynamics and ecosystem structure and function.

## 5.4 Implications and future research

In our study, we highlighted the specific impacts of MHWs on ecosystem structure and function, particularly through the case study of the MHW known as ‘the Blob’ (the longest and strongest period of abnormal temperature ever recorded (Collins et al., 2019; Oliver et al., 2021)). The anomalously low wind during the 2013-2014 winter induced anomalously weak Ekman transport of colder water from the north and, coupled with anomalously low air-sea heat exchange, triggered ‘the Blob’ (Bond et al., 2015). Furthermore, these processes as well as the El Niño Southern Oscillation (ENSO) have already contributed to the increased average duration and intensity of MHWs in the North-eastern Pacific Ocean. Given that projections for the 21st century indicate that ENSO events will increase in intensity and frequency (Holbrook, Gupta, et al., 2020; Oliver et al., 2021), events such as ‘the Blob’ are expected to occur more frequently (Holbrook, Gupta, et al., 2020; Oliver et al., 2021). This underscores the need for ongoing research to better understand how MHWs disrupt ecosystems. Finally, throughout the 21st century, ecosystem responses will depend on the ability of communities to adapt to the long-term increase in ocean temperature and their ability to withstand short-term extreme temperatures (Johansen et al., 2021). Therefore, to enhance our understanding of how marine ecosystems will respond to climate change, future studies should focus on potential scenarios of adaptive responses to future climate and associated MHWs. The EcoTroph Dyn model is a tool to understand the ecological consequences of MHWs at global and local scales, and to project their impacts under future scenarios. However, the model focuses on aggregated energy flows between trophic groups while ecological responses to MHWs between species within each group may vary substantially. Some species may acclimatise or adapt to MHWs. Consideration of the potential

acclimatization/adaptation in the model requires the development of specific adaptation scenarios and model settings in addition to the model settings presented here.

## 6. Conclusion

Utilising the EcoTroph-Dyn trophodynamic framework for MHWs, we highlighted substantial and latent repercussions of MHWs, notably biomass loss and biomass flow alteration, which are particularly consequential for higher TLs. As a result, the recovery/restoration time can extend over several years, if not decades. EcoTroph-Dyn model demonstrates its capacity to characterize the impacts of MHWs on ecosystem structure and functions, with a slight underestimation of the magnitude of the impacts when the model is applied to examine 'the Blob' MHW. However, considering the dynamics and characteristics of current and future MHWs, it can be anticipated that ecosystems might not be afforded the necessary temporal window to recover between successive MHW events, which can significantly disrupt long-term trends associated with climate change.


## 7. Appendices

### Appendix A

- MHW characterisation and detection

To characterise MHW in each ocean spatial cell, we analysed daily SST observations from the NOAA's AVHRR data (Reynolds et al., 2007; <https://www.ncei.noaa.gov/access/metadata/landing-page/bin/iso?id=gov.noaa.ncdc:C00680>). We defined MHW as a discrete prolonged anomalously warm water event when the daily SSTs exceed an extreme temperature threshold value for at least five consecutive days (A. J. Hobday et al., 2016). The extreme temperature threshold value was calculated for each 1° latitude x 1° longitude spatial cell as the 90th percentile of daily SST from the 30-year historical time series from January 1982 to December 2011. We did not calculate threshold values by season; thus, MHW events were identified by a single threshold across the year. As a result, we detected MHWs mostly occurring during the year's warmest months (see figure A1(a) for schematic explanation). This approach to identifying the MHW threshold represents biological extreme temperature in the local (spatial cell) context. It is appropriate to assess the direct mortality associated with MHWs (Oliver et al., 2021). We determined a reference average sea surface temperature (SST average) for each spatial cell by analysing data from the 1<sup>st</sup> of January 1982 to the 31<sup>st</sup> of December 2011. Utilising this reference average SST, we classified each spatial cell into 'thermal classes,' with

each class representing a 1°C increment of the reference SST average ranging from -1°C to 29°C (refer to Figure A1(b)). We used the R package `heatwaveR` described at <https://robwschlegel.github.io/heatwaveR/> to compute MHWs characteristics in each spatial cell from January 1982 to December 2021. Thus, we finally obtained MHW characteristics for each 1° per 1° of longitude and latitude ocean cell up to December 2021. The considered MHW characteristics are the threshold value defining MHWs, MHW's duration (in days), category and intensity (mean SST anomaly) and declaration of MHW days over the SST time series.



**Figure A1: Marine heatwave detection method and reference average SST of each ocean cell.**

(a) Schematic explanation of MHWs detection for a spatial cell. The solid horizontal green and black lines represented the extreme threshold value and the reference temperature, respectively. (b) A map of thermal classes of 10°C intervals from -1 to 29°C categorised based

on the reference temperature at each spatial cell over the period from 1st January 1982 to 31st December 2011.

- Estimation of species distribution and associated thermal niche

We developed an algorithm depending on the MHWs' characteristics to express the loss rate of trophic transfers associated with MHW. First, we identified a list of marine species with their occurrence records (3242 bivalves, 500 cephalopods species, 3116 crabs species, and 12782 fish species) gathered from the publicly accessible databases: OBIS ([www.iobis.org](http://www.iobis.org)), the Intergovernmental Oceanographic Commission ([ioc-unesco.org](http://ioc-unesco.org)), GBIF ([www.gbif.org](http://www.gbif.org)), Fishbase ([www.fishbase.org](http://www.fishbase.org)), and the International Union for the Conservation of Nature (<http://www.iucnredlist.org/technical-documents/spatial-data>). We then cleaned the data by removing duplicate entries, terrestrial occurrences, and occurrences outside the known species habitat from the aggregated species occurrence dataset (Froese & Pauly, 2018). Additionally, we excluded zooplankton from the algorithm development due to limited evidence of direct mortality induced by Marine Heatwaves (MHWs), with observed responses mainly manifesting as range shifts and alterations in community structure (Arimitsu et al., 2021; Suryan et al., 2021; Winans et al., 2023). Marine mammals and seabirds were also omitted from algorithm development, as their mortality linked to MHWs primarily stems from secondary effects such as diminished quality and quantity of food supply (Cavole et al., 2016; Piatt et al., 2020) rather than direct heat stress impact. Subsequently, the data were rasterised into a grid covering the global oceans (1° longitude by 1° latitude), denoting the historical presence of each species. Species with occurrence records in fewer than 30 cells were excluded from further analysis (Hernandez et al., 2006).

In a second step, we utilised an ensemble species distribution modelling approach (Asch et al., 2018; Reygondeau, 2019) at a 1° grid scale. Four environmental niche models (ENMs) were applied: Bioclim, Boosted Regression Trees models (Thuiller et al., 2009), Maxent (Phillips et al., 2006), and the Non-Parametric Probabilistic Ecological Niche model (Beaugrand et al., 2011), using global climatology satellite data (AVHRR). Model accuracy was assessed using the area under the curve (AUC) analysis of the receiver operating characteristic (ROC), discarding models with AUC below 0.5 (Sing et al., 2005). The evaluation employed the pROC package in R (Robin et al., 2011). We then calculated the average Habitat Suitability Index (HSI) weighted by the AUC values of each ENM for each spatial cell and species. An HSI threshold for each species was estimated using their prevalence. Spatial cells with HSI below the threshold were deemed non-viable habitats.

Species' predicted thermal niches were quantified from spatial distributions using averaged satellite sea surface temperature (SST) data (AVHRR) from 1982 to 2011. The average SST from 1982 to 2011 was recorded for each spatial cell above the HSI threshold. We then characterised the

predicted thermal niche (histogram of all the average values of SST), and more specifically, the upper-  
 850 temperature threshold, for each species from the 95th percentile of the SST records where they were  
 predicted to occur.

- Additional loss rate associated with MHW

For each spatial cell belonging to each thermal class, we calculated the percentage of species  
 in each trophic class exposed to thermal stress induced by MHWs intensity above their estimated  
 855 temperature threshold (95th percentile of their thermal niche). Matching MHW intensity in each  
 spatial cell from 1981 to 2021 with species' temperature thresholds, we determined the percentage  
 of species exposed to temperatures exceeding their thresholds. Thermal stress of a species was  
 assumed dependent on MHW category (1 to 4, based on SST anomaly) and species' trophic level (<2.5,  
 2.5-3.0, 3.0-3.5, 3.5-4.0, 4.0-4.5, 4.5-5.0, >5.0), estimated from FishBase and SeaLifeBase.

860 To obtain a continuous representation of the percentage of species undergoing a thermal stress  
 as the intensity of MHW increases, we decided to transform the discrete MHW categorisation (Hobday  
 et al., 2018) to a continuous MHW intensity index as follows:

$$MHW_{cat,i} = \frac{MHW \text{ mean anomaly},i}{cat1 \text{ associated anomaly},i} \quad (\text{Eq. A1})$$

MHW mean anomaly was calculated as the difference between the MHW mean SST anomaly  
 865 and the reference temperature of each thermal class (i), and “cat1 associated anomaly” is the mean  
 threshold value used to identify category 1 MHWs in each spatial cell.

We fit the estimated percentage of species undergoing thermal stress with the MHW intensity  
 index and species' trophic class to a nonlinear function. A Gompertz function was selected after  
 preliminary tests because it is better fitted to data than logistic or other mathematical functions with  
 870 similar shapes. The Gompertz function is expressed as:

$$\text{Percentage of species undergoing a thermal stress} = \exp^{-\exp^{b\_tl_i \cdot (MHW_{cat,i} - lt50\_tl_i)}} \quad (\text{Eq. A2});$$

We estimated the parameters  $b\_tl_i$ ,  $lt50\_tl_i$ , and  $MHW_{cat,i}$  for each thermal class i. The  
 parameters  $b\_tl_i$  and  $lt50\_tl_i$  correspond to the slope of the function and the index of marine  
 heatwave intensity ( $MHW_{cat,i}$ ) at which 50% of the species are undergoing thermal stress,  
 875 respectively.


For each thermal class i, parameters  $b\_tl_i$  and  $lt50\_tl_i$  were expressed as (figure A2):

$$b\_tl_i = -1.4511 - e^{0.4223 \cdot (i - 22.4926)} \quad (\text{Eq. A3}) \text{ and}$$

$$lt50\_tl_i = 3.29 - 0.485 \cdot i + 0.0306 \cdot i^2 - 0.000608 \cdot i^3 \text{ when trophic level } < 2.5 \text{ and}$$

$$lt50\_tl_i = 3.55 - 0.271 \cdot i + 0.014 \cdot i^2 - 0.000304 \cdot i^3 \text{ when trophic level } \geq 2.5. \quad (\text{Eq. A4})$$






**Figure A2: coefficient of each reference temperature individual thermal stress algorithm model.** With (1) corresponding to the Gompertz slope at the inflexion point coefficient (b) and (2) the MHW category that causes 50% of species to undergo thermal stress (lt50) with blue and other colours corresponding to the trophic levels below 2.5 and above 2.5, respectively.

Finally, to move from the thermally stressed stage to the loss rate ( $\eta_i$ ), we assumed that species were continuously challenged by MHW increased intensity (Figure A3 for schematic differences) expressed as.

$$\eta_i = \exp^{-\exp^{b_i \cdot \alpha \cdot \left( MHW_{cat,i} - \frac{lt50_i}{\alpha} \right)}} \cdot \beta \quad (\text{Eq. A5})$$



**Figure A3: loss rate associated with MHWs occurrences.** With the ecological hypothesis that MHW increases in intensity/category continuously challenge the aggregated response across species. On the plots,  $\beta$  is equal to 1. The top bottom and bottom rows correspond to the loss rate of trophic level  $<2.5$  and trophic level  $\geq 2.5$ , respectively.

We explored the sensitivity of the results to species' acclimation capacity to MHW conditions by assuming that acclimation reduces the mortality rate due to species' exposure to thermal stress. We tested four acclimation capacity settings represented by the values of the coefficient  $\alpha$ . These settings are full acclimation ( $\alpha = 0$ ; no mortality due to thermal stress), partial acclimation ( $\alpha = 0.2, 0.5$ ; 20%, and 50% of the species die because of thermal stress, respectively) and no acclimation ( $\alpha = 1$ ; all species die when they are under thermal stress). We also related the loss rate to the MHW duration over the fortnight by assuming that the duration increases the mortality rate. The duration of MHW is represented by  $\beta$  and ranges from  $\beta=0$ ; no MHW to  $\beta=1$ ; MHW lasting 15 days of the fortnight (see section 2.3.2 for  $\beta$  computation).

## 8. Code availability

The code for the EcoTroph\_Dyn model that supports the findings of this study is openly available at <https://doi.org/10.57745/NHVPCCR>.

## 9. Data availability

910 Daily SST observations from the NOAA \_ AVHRR data are publicly available on the link  
<https://www.ncei.noaa.gov/access/metadata/landing-page/bin/iso?id=gov.noaa.ncdc:C00680>.

Species occurrence data and associated trophic levels that support the findings of this study are openly available at <https://doi.org/10.57745/PION92>.

## 915 10. Authors contributions

V.G.D.L, W.W.L.C, and D.G undertook conceptualisation, methodology, validation, and writing – review & editing. W.W.L.C and D.G undertook funding acquisition and supervision. V.G.D.L handled the data curation, formal analysis, investigation, project administration, software, visualisation, and writing – original draft.

## 920 11. Competing interests

The authors declare that they have no conflict of interest

## 12. Acknowledgement

We thank the collaboration between L'Institut Agro, Rennes and the University of British Columbia, Vancouver. We further thank Jerome Guitton for his technical support.

## 925 13. Financial support

V.G.D.L and D.G acknowledges funding support from the Region Bretagne. W.W.L.C and V.G.D.L acknowledge funding support from the NSERC Discovery Grant and the SSHRC through the Solving-FCB partnership.

## 14. References

930 Arimitsu, M. L., Piatt, J. F., Hatch, S., Suryan, R. M., Batten, S., Bishop, M. A., Campbell, R. W., Coletti, H., Cushing, D., Gorman, K., Hopcroft, R. R., Kuletz, K. J., Marsteller, C., McKinstry, C., McGowan, D., Moran, J., Pegau, S., Schaefer, A., Schoen, S., Straley, J., and Biela, V. R.: Heatwave-induced synchrony within forage fish portfolio disrupts energy flow to top pelagic predators, *Glob. Change Biol.*, 27, 1859–1878, <https://doi.org/10.1111/gcb.15556>, 2021.

- 935 Arteaga, L. A. and Rousseaux, C. S.: Impact of Pacific Ocean heatwaves on phytoplankton community composition, *Commun Biol*, 6, 263, <https://doi.org/10.1038/s42003-023-04645-0>, 2023.
- Babcock, R. C., Bustamante, R. H., Fulton, E. A., Fulton, D. J., Haywood, M. D. E., Hobday, A. J., Kenyon, R., Matear, R. J., Plagányi, E. E., Richardson, A. J., and Vanderklift, M. A.: Severe Continental-Scale Impacts of Climate Change Are Happening Now: Extreme Climate Events Impact Marine Habitat Forming Communities Along 45% of Australia's Coast, *Front. Mar. Sci.*, 6, 411, <https://doi.org/10.3389/fmars.2019.00411>, 2019.
- 940 Begon, M. and Townsend, C. R.: *Ecology: From Individuals to Ecosystems*, John Wiley & Sons, 868 pp., 2021.
- Behrenfeld, M. J. and Falkowski, P. G.: Photosynthetic rates derived from satellite-based chlorophyll concentration, *Limnology and Oceanography*, 42, 1–20, <https://doi.org/10.4319/lo.1997.42.1.0001>, 1997.
- 945 Bender, E. A., Case, T. J., and Gilpin, M. E.: Perturbation Experiments in Community Ecology: Theory and Practice, *Ecology*, 65, 1–13, 1984.
- Bernhardt, J. R. and Leslie, H. M.: Resilience to Climate Change in Coastal Marine Ecosystems, *Annu. Rev. Mar. Sci.*, 5, 371–392, <https://doi.org/10.1146/annurev-marine-121211-172411>, 2013.
- 950 Beukhof, E., Dencker, T., Pecuchet, L., and Lindegren, M.: Spatio-temporal variation in marine fish traits reveals community-wide responses to environmental change, *Mar. Ecol. Prog. Ser.*, 610, 205–222, <https://doi.org/10.3354/meps12826>, 2019.
- Blanchard, J. L., Jennings, S., Holmes, R., Harle, J., Merino, G., Allen, J. I., Holt, J., Dulvy, N. K., and Barange, M.: Potential consequences of climate change for primary production and fish production in large marine ecosystems, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367, 2979–2989, <https://doi.org/10.1098/rstb.2012.0231>, 2012.
- 955 Bond, N. A., Cronin, M. F., Freeland, H., and Mantua, N.: Causes and impacts of the 2014 warm anomaly in the NE Pacific: 2014 WARM ANOMALY IN THE NE PACIFIC, *Geophys. Res. Lett.*, 42, 3414–3420, <https://doi.org/10.1002/2015GL063306>, 2015.
- 960 Bouchard, C., Geoffroy, M., LeBlanc, M., Majewski, A., Gauthier, S., Walkusz, W., Reist, J. D., and Fortier, L.: Climate warming enhances polar cod recruitment, at least transiently, *Progress in Oceanography*, 156, 121–129, <https://doi.org/10.1016/j.pocean.2017.06.008>, 2017.
- Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., and West, G. B.: TOWARD A METABOLIC THEORY OF ECOLOGY, *Ecology*, 85, 1771–1789, <https://doi.org/10.1890/03-9000>, 2004.
- 965 Cannell, B., Thompson, P., and Schoepf, V.: *Marine extremes: ocean safety, marine health and the blue economy*, Routledge, Taylor & Francis Group, London ; New York, 1 pp., 2019.
- Carneiro, A. P., Soares, C. H. L., Manso, P. R. J., and Pagliosa, P. R.: Impact of marine heat waves and cold spell events on the bivalve *Anomalocardia flexuosa*: A seasonal comparison, *Marine Environmental Research*, 156, 104898, <https://doi.org/10.1016/j.marenvres.2020.104898>, 2020.
- 970 Carozza, D. A., Bianchi, D., and Galbraith, E. D.: The ecological module of BOATS-1.0: a bioenergetically constrained model of marine upper trophic levels suitable for studies of fisheries and ocean biogeochemistry, *Geosci. Model Dev.*, 9, 1545–1565, <https://doi.org/10.5194/gmd-9-1545-2016>, 2016.

- 975 Cavole, L., Demko, A., Diner, R., Giddings, A., Koester, I., Pagniello, C., Paulsen, M.-L., Ramirez-Valdez, A., Schwenck, S., Yen, N., Zill, M., and Franks, P.: Biological Impacts of the 2013–2015 Warm-Water Anomaly in the Northeast Pacific: Winners, Losers, and the Future, *Oceanog*, 29, <https://doi.org/10.5670/oceanog.2016.32>, 2016.
- Cheung, W. W. L. and Frölicher, T. L.: Marine heatwaves exacerbate climate change impacts for  
980 fisheries in the northeast Pacific, *Sci Rep*, 10, 6678, <https://doi.org/10.1038/s41598-020-63650-z>, 2020.
- Cheung, W. W. L., Dunne, J., Sarmiento, J. L., and Pauly, D.: Integrating ecophysiology and plankton dynamics into projected maximum fisheries catch potential under climate change in the Northeast Atlantic, *ICES Journal of Marine Science*, 68, 1008–1018, <https://doi.org/10.1093/icesjms/fsr012>,  
985 2011.
- Cheung, W. W. L., Frölicher, T. L., Lam, V. W. Y., Oyinlola, M. A., Reygondeau, G., Sumaila, U. R., Tai, T. C., Teh, L. C. L., and Wabnitz, C. C. C.: Marine high temperature extremes amplify the impacts of climate change on fish and fisheries, *Sci. Adv.*, 7, eabh0895, <https://doi.org/10.1126/sciadv.abh0895>, 2021.
- Christensen, V. and Pauly, D.: ECOPATH II — a software for balancing steady-state ecosystem models and calculating network characteristics, *Ecological Modelling*, 61, 169–185, [https://doi.org/10.1016/0304-3800\(92\)90016-8](https://doi.org/10.1016/0304-3800(92)90016-8), 1992.  
990
- Collins, M., Sutherland, M., Bouwer, L., Cheong, S.-M., Combes, H. J. D., Roxy, M. K., Losada, I., McInnes, K., Ratter, B., Rivera-Arriaga, E., Susanto, R. D., Swingedouw, D., Tibig, L., Bakker, P., Eakin, C. M., Emanuel, K., Grose, M., Hemer, M., Jackson, L., Käb, A., Kajtar, J., Knutson, T., Laufkötter, C., Noy, I.,  
995 Payne, M., Ranasinghe, R., Sgubin, G., Timmermans, M.-L., Abdulla, A., González, M. H., and Turley, C.: SPM6 Extremes, Abrupt Changes and Managing Risks, 68, <https://doi.org/10.1017/9781009157964.008>, 2019.
- Crickenberger, S. and Wetthey, D. S.: Reproductive physiology, temperature and biogeography: the role of fertilization in determining the distribution of the barnacle *Semibalanus balanoides*, *J. Mar. Biol. Ass.*, 98, 1411–1424, <https://doi.org/10.1017/S0025315417000364>, 2018.  
1000
- Eddy, T. D., Bernhardt, J. R., Blanchard, J. L., Cheung, W. W. L., Colléter, M., du Pontavice, H., Fulton, E. A., Gascuel, D., Kearney, K. A., Petrik, C. M., Roy, T., Rykaczewski, R. R., Selden, R., Stock, C. A., Wabnitz, C. C. C., and Watson, R. A.: Energy Flow Through Marine Ecosystems: Confronting Transfer Efficiency, *Trends in Ecology & Evolution*, 36, 76–86, <https://doi.org/10.1016/j.tree.2020.09.006>, 2021.
- 1005 EPPLEY, R. W.: TEMPERATURE AND PHYTOPLANKTON GROWTH IN THE SEA, 1972.
- Fredston, A. L., Cheung, W. W. L., Frölicher, T. L., Kitchel, Z. J., Maureaud, A. A., Thorson, J. T., Auber, A., Mérigot, B., Palacios-Abrantes, J., Palomares, M. L. D., Pecuchet, L., Shackell, N. L., and Pinsky, M. L.: Marine heatwaves are not a dominant driver of change in demersal fishes, *Nature*, 621, 324–329, <https://doi.org/10.1038/s41586-023-06449-y>, 2023.
- 1010 Frölicher, T. L. and Laufkötter, C.: Emerging risks from marine heat waves, *Nat Commun*, 9, 650, <https://doi.org/10.1038/s41467-018-03163-6>, 2018.
- Frölicher, T. L., Fischer, E. M., and Gruber, N.: Marine heatwaves under global warming, *Nature*, 560, 360–364, <https://doi.org/10.1038/s41586-018-0383-9>, 2018.
- 1015 Garrabou, J., Coma, R., Bensoussan, N., Bally, M., Chevaldonné, P., Cigliano, M., Diaz, D., Harmelin, J. G., Gambi, M. C., Kersting, D. K., Ledoux, J. B., Lejeusne, C., Linares, C., Marschal, C., Pérez, T., Ribes,

- M., Romano, J. C., Serrano, E., Teixido, N., Torrents, O., Zabala, M., Zuberer, F., and Cerrano, C.: Mass mortality in Northwestern Mediterranean rocky benthic communities: effects of the 2003 heat wave, *Global Change Biology*, 15, 1090–1103, <https://doi.org/10.1111/j.1365-2486.2008.01823.x>, 2009.
- 1020 Garrabou, J., Gómez-Gras, D., Medrano, A., Cerrano, C., Ponti, M., Schlegel, R., Bensoussan, N., Turicchia, E., Sini, M., Gerovasileiou, V., Teixido, N., Mirasole, A., Tamburello, L., Cebrian, E., Rilov, G., Ledoux, J., Souissi, J. B., Khamassi, F., Ghanem, R., Benabdi, M., Grimes, S., Ocaña, O., Bazairi, H., Hereu, B., Linares, C., Kersting, D. K., la Rovira, G., Ortega, J., Casals, D., Pagès-Escalà, M., Margarit, N., Capdevila, P., Verdura, J., Ramos, A., Izquierdo, A., Barbera, C., Rubio-Portillo, E., Anton, I., López-Sendino, P., Díaz, D., Vázquez-Luis, M., Duarte, C., Marbà, N., Aspillaga, E., Espinosa, F., Grech, D.,
- 1025 Guala, I., Azzurro, E., Farina, S., Cristina Gambi, M., Chimienti, G., Montefalcone, M., Azzola, A., Mantas, T. P., Frascchetti, S., Ceccherelli, G., Kipson, S., Bakran-Petricioli, T., Petricioli, D., Jimenez, C., Katsanevakis, S., Kizilkaya, I. T., Kizilkaya, Z., Sartoretto, S., Elodie, R., Ruitton, S., Comeau, S., Gattuso, J., and Harmelin, J.: Marine heatwaves drive recurrent mass mortalities in the Mediterranean Sea, *Global Change Biology*, 28, 5708–5725, <https://doi.org/10.1111/gcb.16301>, 2022.
- 1030 Gasche, L. and Gascuel, D.: EcoTroph: a simple model to assess fishery interactions and their impacts on ecosystems, *ICES Journal of Marine Science*, 70, 498–510, <https://doi.org/10.1093/icesjms/fst016>, 2013.
- Gasche, L., Gascuel, D., Shannon, L., and Shin, Y.-J.: Global assessment of the fishing impacts on the Southern Benguela ecosystem using an EcoTroph modelling approach, *Journal of Marine Systems*, 90,
- 1035 1–12, <https://doi.org/10.1016/j.jmarsys.2011.07.012>, 2012.
- Gascuel, D.: The trophic-level based model: A theoretical approach of fishing effects on marine ecosystems, *Ecological Modelling*, 189, 315–332, <https://doi.org/10.1016/j.ecolmodel.2005.03.019>, 2005.
- Gascuel, D. and Pauly, D.: EcoTroph: Modelling marine ecosystem functioning and impact of fishing, *Ecological Modelling*, 220, 2885–2898, <https://doi.org/10.1016/j.ecolmodel.2009.07.031>, 2009.
- 1040 Gascuel, D., Morissette, L., Palomares, M. L. D., and Christensen, V.: Trophic flow kinetics in marine ecosystems: Toward a theoretical approach to ecosystem functioning, *Ecological Modelling*, 217, 33–47, <https://doi.org/10.1016/j.ecolmodel.2008.05.012>, 2008.
- Gascuel, D., Guénette, S., and Pauly, D.: The trophic-level-based ecosystem modelling approach: theoretical overview and practical uses, *ICES Journal of Marine Science*, 68, 1403–1416, <https://doi.org/10.1093/icesjms/fsr062>, 2011.
- 1045 Gomes, D. G. E., Ruzicka, J. J., Crozier, L. G., Huff, D. D., Brodeur, R. D., and Stewart, J. D.: Marine heatwaves disrupt ecosystem structure and function via altered food webs and energy flux, *Nat Commun*, 15, 1988, <https://doi.org/10.1038/s41467-024-46263-2>, 2024.
- 1050 Grimmelpont, M., Milinkovitch, T., Dubillot, E., and Lefrançois, C.: Individual aerobic performance and anaerobic compensation in a temperate fish during a simulated marine heatwave, *Science of The Total Environment*, 863, 160844, <https://doi.org/10.1016/j.scitotenv.2022.160844>, 2023.
- 1055 Guibourd de Luzinai, Du Pontavice, H., Reygondeau, G., Barrier, N., Blanchard, J. L., Bornarel, V., Büchner, M., Cheung, W. W. L., Eddy, T. D., Everett, J. D., Guiet, J., Harrison, C. S., Maury, O., Novaglio, C., Petrik, C. M., Steenbeek, J., Tittensor, D. P., and Gascuel, D.: Trophic amplification: A model intercomparison of climate driven changes in marine food webs, *PLoS ONE*, 18, e0287570, <https://doi.org/10.1371/journal.pone.0287570>, 2023.

- Guibourd de Luzinais, Gascuel, D., Reygondeau, G., and Cheung, W. W. L.: Large potential impacts of marine heatwaves on ecosystem functioning, *Global Change Biology*, 30, e17437, <https://doi.org/10.1111/gcb.17437>, 2024.
- Halouani, G., Gascuel, D., Hattab, T., Lasram, F. B. R., Coll, M., Tsagarakis, K., Piroddi, C., Romdhane, M. S., and Le Loc'h, F.: Fishing impact in Mediterranean ecosystems: an EcoTroph modeling approach, *Journal of Marine Systems*, 150, 22–33, <https://doi.org/10.1016/j.jmarsys.2015.05.007>, 2015.
- Harris, R. M. B., Beaumont, L. J., Vance, T. R., Tozer, C. R., Remenyi, T. A., Perkins-Kirkpatrick, S. E., Mitchell, P. J., Nicotra, A. B., McGregor, S., Andrew, N. R., Letnic, M., Kearney, M. R., Wernberg, T., Hutley, L. B., Chambers, L. E., Fletcher, M.-S., Keatley, M. R., Woodward, C. A., Williamson, G., Duke, N. C., and Bowman, D. M. J. S.: Biological responses to the press and pulse of climate trends and extreme events, *Nature Clim Change*, 8, 579–587, <https://doi.org/10.1038/s41558-018-0187-9>, 2018.
- Hobday, A., Alexander, L. V., Perkins, S. E., Smale, D. A., Straub, S. C., Oliver, E. C. J., Benthuyssen, J. A., Burrows, M. T., Donat, M. G., Feng, M., Holbrook, N. J., Moore, P. J., Scannell, H. A., Sen Gupta, A., and Wernberg, T.: A hierarchical approach to defining marine heatwaves, *Progress in Oceanography*, 141, 227–238, <https://doi.org/10.1016/j.pocean.2015.12.014>, 2016.
- Hobday, A., Oliver, E., Sen Gupta, A., Benthuyssen, J., Burrows, M., Donat, M., Holbrook, N., Moore, P., Thomsen, M., Wernberg, T., and Smale, D.: Categorizing and Naming Marine Heatwaves, *Oceanog*, 31, <https://doi.org/10.5670/oceanog.2018.205>, 2018.
- Holbrook, N. J., Sen Gupta, A., Oliver, E. C. J., Hobday, A. J., Benthuyssen, J. A., Scannell, H. A., Smale, D. A., and Wernberg, T.: Keeping pace with marine heatwaves, *Nat Rev Earth Environ*, 1, 482–493, <https://doi.org/10.1038/s43017-020-0068-4>, 2020.
- Jennings, S. and Collingridge, K.: Predicting Consumer Biomass, Size-Structure, Production, Catch Potential, Responses to Fishing and Associated Uncertainties in the World's Marine Ecosystems, *PLoS ONE*, 10, e0133794, <https://doi.org/10.1371/journal.pone.0133794>, 2015.
- Jennings, S., Warr, K., and Mackinson, S.: Use of size-based production and stable isotope analyses to predict trophic transfer efficiencies and predator-prey body mass ratios in food webs, *Mar. Ecol. Prog. Ser.*, 240, 11–20, <https://doi.org/10.3354/meps240011>, 2002.
- Jennings, S., Mélin, F., Blanchard, J. L., Forster, R. M., Dulvy, N. K., and Wilson, R. W.: Global-scale predictions of community and ecosystem properties from simple ecological theory, *Proc. R. Soc. B.*, 275, 1375–1383, <https://doi.org/10.1098/rspb.2008.0192>, 2008.
- Johansen, J. L., Nadler, L. E., Habary, A., Bowden, A. J., and Rummer, J.: Thermal acclimation of tropical coral reef fishes to global heat waves, *eLife*, 10, e59162, <https://doi.org/10.7554/eLife.59162>, 2021.
- Joyce, P. W. S., Tong, C. B., Yip, Y. L., and Falkenberg, L. J.: Marine heatwaves as drivers of biological and ecological change: implications of current research patterns and future opportunities, *Mar Biol*, 171, 20, <https://doi.org/10.1007/s00227-023-04340-y>, 2024.
- Largier, J. L.: Upwelling Bays: How Coastal Upwelling Controls Circulation, Habitat, and Productivity in Bays, *Annu. Rev. Mar. Sci.*, 12, 415–447, <https://doi.org/10.1146/annurev-marine-010419-011020>, 2020.
- Le Grix, N., Zscheischler, J., Rodgers, K., Yamaguchi, R., and Frölicher, T. L.: Hotspots and drivers of compound marine heatwave and low net primary production extremes, *Biogeochemistry: Open Ocean*, <https://doi.org/10.5194/egusphere-2022-451>, 2022.

- 1100 LeBlanc, M., Geoffroy, M., Bouchard, C., Gauthier, S., Majewski, A., Reist, J. D., and Fortier, L.: Pelagic production and the recruitment of juvenile polar cod *Boreogadus saida* in Canadian Arctic seas, *Polar Biol*, 43, 1043–1054, <https://doi.org/10.1007/s00300-019-02565-6>, 2020.
- Libralato, S., Coll, M., Tudela, S., Palomera, I., and Pranovi, F.: Novel index for quantification of ecosystem effects of fishing as removal of secondary production, *Mar. Ecol. Prog. Ser.*, 355, 107–129, <https://doi.org/10.3354/meps07224>, 2008.
- 1105 Lindeman, R. L.: The Trophic-Dynamic Aspect of Ecology, *Ecology*, 23, 399–417, <https://doi.org/10.2307/1930126>, 1942.
- Link, J. S.: Adding rigor to ecological network models by evaluating a set of pre-balance diagnostics: A plea for PREBAL, *Ecological Modelling*, 221, 1580–1591, <https://doi.org/10.1016/j.ecolmodel.2010.03.012>, 2010.
- 1110 Longhurst, A. R.: *Ecological geography of the sea*, 2. ed., Elsevier, Amsterdam Heidelberg, 542 pp., 2007.
- Marin, M., Bindoff, N. L., Feng, M., and Phillips, H. E.: Slower Long-Term Coastal Warming Drives Dampened Trends in Coastal Marine Heatwave Exposure, *JGR Oceans*, 126, e2021JC017930, <https://doi.org/10.1029/2021JC017930>, 2021.
- 1115 Milutinović, S. and Bertino, L.: Assessment and propagation of uncertainties in input terms through an ocean-color-based model of primary productivity, *Remote Sensing of Environment*, 115, 1906–1917, <https://doi.org/10.1016/j.rse.2011.03.013>, 2011.
- 1120 Minuti, J. J., Byrne, M., Hemraj, D. A., and Russell, B. D.: Capacity of an ecologically key urchin to recover from extreme events: Physiological impacts of heatwaves and the road to recovery, *Science of The Total Environment*, 785, 147281, <https://doi.org/10.1016/j.scitotenv.2021.147281>, 2021.
- Morel, A.: Light and marine photosynthesis: a spectral model with geochemical and climatological implications, *Progress in Oceanography*, 26, 263–306, [https://doi.org/10.1016/0079-6611\(91\)90004-6](https://doi.org/10.1016/0079-6611(91)90004-6), 1991.
- 1125 Morgan, C. A., Beckman, B. R., Weitkamp, L. A., and Fresh, K. L.: Recent Ecosystem Disturbance in the Northern California Current, *Fisheries*, 44, 465–474, <https://doi.org/10.1002/fsh.10273>, 2019.
- Niquil, N., Baeta, A., Marques, J., Chaalali, A., Lobry, J., and Patrício, J.: Reaction of an estuarine food web to disturbance: Lindeman’s perspective, *Mar. Ecol. Prog. Ser.*, 512, 141–154, <https://doi.org/10.3354/meps10885>, 2014.
- 1130 Oliver, E. C. J., Benthuisen, J. A., Bindoff, N. L., Hobday, A. J., Holbrook, N. J., Mundy, C. N., and Perkins-Kirkpatrick, S. E.: The unprecedented 2015/16 Tasman Sea marine heatwave, *Nat Commun*, 8, 16101, <https://doi.org/10.1038/ncomms16101>, 2017.
- 1135 Oliver, E. C. J., Donat, M. G., Burrows, M. T., Moore, P. J., Smale, D. A., Alexander, L. V., Benthuisen, J. A., Feng, M., Sen Gupta, A., Hobday, A. J., Holbrook, N. J., Perkins-Kirkpatrick, S. E., Scannell, H. A., Straub, S. C., and Wernberg, T.: Longer and more frequent marine heatwaves over the past century, *Nat Commun*, 9, 1324, <https://doi.org/10.1038/s41467-018-03732-9>, 2018.
- Oliver, E. C. J., Benthuisen, J. A., Darmaraki, S., Donat, M. G., Hobday, A. J., Holbrook, N. J., Schlegel, R. W., and Sen Gupta, A.: Annual Review of Marine Science Marine Heatwaves, *Annu. Rev. Mar. Sci.*, 13, 313–342, <https://doi.org/10.1146/annurev-marine-032720-095144>, 2021.



- 1140 Pauly, D. and Christensen, V.: Primary production required to sustain global fisheries, *Nature*, 374, 255–257, <https://doi.org/10.1038/374255a0>, 1995.
- Pearce, A., Lenanton, R., Jackson, G., Moore, J., Feng, M., and Gaughan, D.: The “marine heat wave” off Western Australia during the summer of 2010/1, 2011.
- 1145 Pershing, A., Mills, K., Dayton, A., Franklin, B., and Kennedy, B.: Evidence for Adaptation from the 2016 Marine Heatwave in the Northwest Atlantic Ocean, *Oceanog*, 31, <https://doi.org/10.5670/oceanog.2018.213>, 2018.
- Peterson, W. T., Fisher, J. L., Strub, P. T., Du, X., Risien, C., Peterson, J., and Shaw, C. T.: The pelagic ecosystem in the Northern California Current off Oregon during the 2014–2016 warm anomalies within the context of the past 20 years, *JGR Oceans*, 122, 7267–7290, <https://doi.org/10.1002/2017JC012952>, 2017.
- 1150 Pezzulli, S., Stephenson, D. B., and Hannachi, A.: The Variability of Seasonality, *Journal of Climate*, 18, 71–88, <https://doi.org/10.1175/JCLI-3256.1>, 2005.
- 1155 Piatt, J. F., Parrish, J. K., Renner, H. M., Schoen, S. K., Jones, T. T., Arimitsu, M. L., Kuletz, K. J., Bodenstein, B., García-Reyes, M., Duerr, R. S., Corcoran, R. M., Kaler, R. S. A., McChesney, G. J., Golightly, R. T., Coletti, H. A., Suryan, R. M., Burgess, H. K., Lindsey, J., Lindquist, K., Warzybok, P. M., Jahncke, J., Roletto, J., and Sydeman, W. J.: Extreme mortality and reproductive failure of common murrelets resulting from the northeast Pacific marine heatwave of 2014–2016, *PLoS ONE*, 15, e0226087, <https://doi.org/10.1371/journal.pone.0226087>, 2020.
- 1160 Pinsky, M. L., Eikeset, A. M., McCauley, D. J., Payne, J. L., and Sunday, J. M.: Greater vulnerability to warming of marine versus terrestrial ectotherms, *Nature*, 569, 108–111, <https://doi.org/10.1038/s41586-019-1132-4>, 2019.
- Plagányi, É. E.: Models for an ecosystem approach to fisheries, Food and Agriculture Organization of the United Nations, Rome, 108 pp., 2007.
- 1165 du Pontavice, H., Gascuel, D., Reygondeau, G., Maureaud, A., and Cheung, W. W. L.: Climate change undermines the global functioning of marine food webs, *Glob Change Biol*, 26, 1306–1318, <https://doi.org/10.1111/gcb.14944>, 2020.
- du Pontavice, H., Gascuel, D., Reygondeau, G., Stock, C., and Cheung, W. W. L.: Climate-induced decrease in biomass flow in marine food webs may severely affect predators and ecosystem production, *Glob Change Biol*, 27, 2608–2622, <https://doi.org/10.1111/gcb.15576>, 2021.
- 1170 du Pontavice, H., Gascuel, D., Kay, S., and Cheung, W.: Climate-induced changes in ocean productivity and food-web functioning are projected to markedly affect European fisheries catch, *Mar. Ecol. Prog. Ser.*, 713, 21–37, <https://doi.org/10.3354/meps14328>, 2023.
- Pörtner and Farrell, A. P.: Physiology and Climate Change, *Science*, 322, 690–692, <https://doi.org/10.1126/science.1163156>, 2008.
- 1175 Reygondeau, G., Longhurst, A., Martinez, E., Beaugrand, G., Antoine, D., and Maury, O.: Dynamic biogeochemical provinces in the global ocean: DYNAMIC BIOGEOCHEMICAL PROVINCES, *Global Biogeochem. Cycles*, 27, 1046–1058, <https://doi.org/10.1002/gbc.20089>, 2013.
- Ruzicka, J. J., Brodeur, R. D., Emmett, R. L., Steele, J. H., Zamon, J. E., Morgan, C. A., Thomas, A. C., and Wainwright, T. C.: Interannual variability in the Northern California Current food web structure:

- Changes in energy flow pathways and the role of forage fish, euphausiids, and jellyfish, *Progress in Oceanography*, 102, 19–41, <https://doi.org/10.1016/j.pocean.2012.02.002>, 2012.
- Rykaczewski, R. R. and Checkley, D. M.: Influence of ocean winds on the pelagic ecosystem in upwelling regions, *Proc. Natl. Acad. Sci. U.S.A.*, 105, 1965–1970, <https://doi.org/10.1073/pnas.0711777105>, 2008.
- Ryther, J. H.: Photosynthesis and Fish Production in the Sea, *Science*, 166, 72–76, <https://doi.org/10.1126/science.166.3901.72>, 1969.
- Schlegel, R. and Smit, A.: heatwaveR: A central algorithm for the detection of heatwaves and cold-spells, *JOSS*, 3, 821, <https://doi.org/10.21105/joss.00821>, 2018.
- Schoener, T. W.: Rate of Species Turnover Decreases from Lower to Higher Organisms: A Review of the Data, *Oikos*, 41, 372, <https://doi.org/10.2307/3544095>, 1983.
- Schramski, J. R., Dell, A. I., Grady, J. M., Sibly, R. M., and Brown, J. H.: Metabolic theory predicts whole-ecosystem properties, *Proc. Natl. Acad. Sci. U.S.A.*, 112, 2617–2622, <https://doi.org/10.1073/pnas.1423502112>, 2015.
- Shanks, A. L., Rasmuson, L. K., Valley, J. R., Jarvis, M. A., Salant, C., Sutherland, D. A., Lamont, E. I., Hainey, M. A. H., and Emlet, R. B.: Marine heat waves, climate change, and failed spawning by coastal invertebrates, *Limnology & Oceanography*, 65, 627–636, <https://doi.org/10.1002/lno.11331>, 2020.
- Shiskin, J.: The X-11 Variant of the Census Method II Seasonal Adjustment Program, U.S. Department of Commerce, Bureau of the Census, 74 pp., 1967.
- Smale, D. A., Yunnice, A. L. E., Vance, T., and Widdicombe, S.: Disentangling the impacts of heat wave magnitude, duration and timing on the structure and diversity of sessile marine assemblages, *PeerJ*, 3, e863, <https://doi.org/10.7717/peerj.863>, 2015.
- Smale, D. A., Wernberg, T., Oliver, E. C. J., Thomsen, M., Harvey, B. P., Straub, S. C., Burrows, M. T., Alexander, L. V., Benthuyssen, J. A., Donat, M. G., Feng, M., Hobday, A. J., Holbrook, N. J., Perkins-Kirkpatrick, S. E., Scannell, H. A., Sen Gupta, A., Payne, B. L., and Moore, P. J.: Marine heatwaves threaten global biodiversity and the provision of ecosystem services, *Nat. Clim. Chang.*, 9, 306–312, <https://doi.org/10.1038/s41558-019-0412-1>, 2019.
- Smith, Burrows, M. T., Hobday, A. J., King, N. G., Moore, P. J., Sen Gupta, A., Thomsen, M. S., Wernberg, T., and Smale, D. A.: Biological Impacts of Marine Heatwaves, *Annu. Rev. Mar. Sci.*, 15, 119–145, <https://doi.org/10.1146/annurev-marine-032122-121437>, 2023.
- Smith, K. E. and Thatje, S.: Nurse egg consumption and intracapsular development in the common whelk *Buccinum undatum* (Linnaeus 1758), *Helgol Mar Res*, 67, 109–120, <https://doi.org/10.1007/s10152-012-0308-1>, 2013.
- Stock, C. A., John, J. G., Rykaczewski, R. R., Asch, R. G., Cheung, W. W. L., Dunne, J. P., Friedland, K. D., Lam, V. W. Y., Sarmiento, J. L., and Watson, R. A.: Reconciling fisheries catch and ocean productivity, *Proc Natl Acad Sci USA*, 114, E1441–E1449, <https://doi.org/10.1073/pnas.1610238114>, 2017.
- Suryan, R. M., Arimitsu, M. L., Coletti, H. A., Hopcroft, R. R., Lindeberg, M. R., Barbeaux, S. J., Batten, S. D., Burt, W. J., Bishop, M. A., Bodkin, J. L., Brenner, R., Campbell, R. W., Cushing, D. A., Danielson, S. L., Dorn, M. W., Drummond, B., Esler, D., Gelatt, T., Hanselman, D. H., Hatch, S. A., Haught, S., Holderied, K., Iken, K., Irons, D. B., Kettle, A. B., Kimmel, D. G., Konar, B., Kuletz, K. J., Laurel, B. J., Maniscalco, J.

- 1220 M., Matkin, C., McKinstry, C. A. E., Monson, D. H., Moran, J. R., Olsen, D., Palsson, W. A., Pegau, W. S., Piatt, J. F., Rogers, L. A., Rojek, N. A., Schaefer, A., Spies, I. B., Straley, J. M., Strom, S. L., Sweeney, K. L., Szymkowiak, M., Weitzman, B. P., Yasumiishi, E. M., and Zador, S. G.: Ecosystem response persists after a prolonged marine heatwave, *Sci Rep*, 11, 6235, <https://doi.org/10.1038/s41598-021-83818-5>, 2021.
- 1225 Tremblay-Boyer, L., Gascuel, D., Watson, R., Christensen, V., and Pauly, D.: Modelling the effects of fishing on the biomass of the world's oceans from 1950 to 2006, *Mar. Ecol. Prog. Ser.*, 442, 169–185, <https://doi.org/10.3354/meps09375>, 2011.
- Vantrepotte, V. and Mélin, F.: Inter-annual variations in the SeaWiFS global chlorophyll a concentration (1997–2007), *Deep Sea Research Part I: Oceanographic Research Papers*, 58, 429–441, <https://doi.org/10.1016/j.dsr.2011.02.003>, 2011.
- 1230 Varela, R., Rodríguez-Díaz, L., de Castro, M., and Gómez-Gesteira, M.: Influence of Eastern Upwelling systems on marine heatwaves occurrence, *Global and Planetary Change*, 196, 103379, <https://doi.org/10.1016/j.gloplacha.2020.103379>, 2021.
- Vinagre, C., Leal, I., Mendonça, V., Madeira, D., Narciso, L., Diniz, M. S., and Flores, A. A. V.: Vulnerability to climate warming and acclimation capacity of tropical and temperate coastal organisms, *Ecological Indicators*, 62, 317–327, <https://doi.org/10.1016/j.ecolind.2015.11.010>, 2016.
- 1235 Wernberg, T., Smale, D. A., Tuya, F., Thomsen, M. S., Langlois, T. J., de Bettignies, T., Bennett, S., and Rousseaux, C. S.: An extreme climatic event alters marine ecosystem structure in a global biodiversity hotspot, *Nature Clim Change*, 3, 78–82, <https://doi.org/10.1038/nclimate1627>, 2013.
- 1240 Wernberg, T., Bennett, S., Babcock, R. C., de Bettignies, T., Cure, K., Depczynski, M., Dufois, F., Fromont, J., Fulton, C. J., Hovey, R. K., Harvey, E. S., Holmes, T. H., Kendrick, G. A., Radford, B., Santana-Garcon, J., Saunders, B. J., Smale, D. A., Thomsen, M. S., Tuckett, C. A., Tuya, F., Vanderklift, M. A., and Wilson, S.: Climate-driven regime shift of a temperate marine ecosystem, *Science*, 353, 169–172, <https://doi.org/10.1126/science.aad8745>, 2016.
- 1245 Westberry, T. K., Silsbe, G. M., and Behrenfeld, M. J.: Gross and net primary production in the global ocean: An ocean color remote sensing perspective, *Earth-Science Reviews*, 237, 104322, <https://doi.org/10.1016/j.earscirev.2023.104322>, 2023.