

# Ocean circulation, sea ice, and productivity simulated in Jones Sound, Canadian Arctic Archipelago, between 2003-2016

Tyler Pelle<sup>1</sup>, Paul G. Myers<sup>2</sup>, Andrew Hamilton<sup>2</sup>, Matthew Mazloff<sup>1</sup>, Krista M. Soderlund<sup>3</sup>, Lucas Beem<sup>3</sup>, Donald D. Blankenship<sup>3</sup>, Cyril Grima<sup>3</sup>, Feras Habbal<sup>3</sup>, Mark Skidmore<sup>4</sup>, and Jamin S. Greenbaum<sup>1</sup>

<sup>1</sup>Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA, USA

<sup>2</sup>Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, Canada

<sup>3</sup>Institute for Geophysics, Jackson School of Geosciences, University of Texas at Austin, Austin, TX, USA

<sup>4</sup>Department of Earth Sciences, Montana State University, Bozeman, Montana, USA

**Correspondence:** Tyler Pelle (tpelle@ucsd.edu)

**Abstract.** Jones Sound is one of three critical waterways in the Canadian Arctic Archipelago that regulate liquid exchange between the Arctic Ocean and northern Atlantic Ocean. However, to date, no high-resolution ocean circulation model exists to study the recent evolution of Jones Sound, meaning that our understanding of circulation within the sound is based either on temporally and spatially sparse oceanographic observations or on extrapolating conditions within Baffin Bay, which has a more 5 dense observational record. To address this, we develop a high-resolution (1/120°, 0.9 km) Jones Sound configuration of the Massachusetts Institute of Technology general circulation model and perform coupled ocean-sea ice-biological productivity simulations between 2003-2016. We find that circulation through Lady Ann Strait, Fram Sound, and Glacier Strait comprises 71%, 14%, and 15% of the volumetric transport into and out of Jones Sound, with tidal flushing enhancing the magnitude of 10 volumetric transport through Fram Sound. Warming Atlantic Water within western Baffin Bay flows into Jones Sound through Lady Ann Strait, becomes well-mixed, and circulates counter-clockwise, encroaching on the terminus of most tidewater glaciers that line the eastern periphery of the sound. Furthermore, we find that sustained atmospheric and oceanic warming drive an 11% reduction in the 2003-2016 mean summertime sea ice area, decreased wintertime sea ice thickness, and delayed onset of sea ice 15 refreeze in the fall (thus lengthening the amount of time in which Jones Sound is ice free). Tidal flushing through Cardigan Strait is critical in triggering meltback of sea ice across northern Jones Sound. Lastly, this decline in sea ice increases light availability and, when coupled with warming of the subsurface waters in Jones Sound, facilitates enhanced primary productivity down to ~21 meters depth. While we note that the modeled warming signal in Baffin Bay is overestimated relative to observations, the results presented here improve our general understanding of how this critical waterway might change under continued polar 20 amplified global warming and underscores the need for sustained oceanographic observations in this region.


## 1 Introduction

The Canadian Arctic Archipelago (CAA) is a tangle of shallow basins and narrow straits that connect the Arctic Ocean to the northern Atlantic Ocean. Flow through the CAA has been found to be primarily controlled by the baroclinic gradient that exists

between the Beaufort Gyre and northern Baffin Bay (Kliem and Greenberg, 2003; Prinsenberg and Bennett, 1987; Wang et al., 2017; Wekerle et al., 2013), with high frequency variability driven by the wind (Peterson et al., 2012). The Queen Elizabeth Islands (QEI) in the north is an area with relatively small islands surrounded by the larger Ellesmere, Devon, Cornwallis, 25 Bathurst, Melville and Prince Patrick islands. Arctic Ocean waters flow through the QEI and are transported into northern Baffin Bay (and eventually the north Atlantic Ocean) through three main passageways: Lancaster Sound, Nares Strait, and Jones Sound. Within these waterways, moorings reveal a mean transport of 0.46 Sv (1 Sv is equal to  $10^6 \text{ m}^3/\text{s}$ ) in western Lancaster Sound between 1998-2010 (Peterson et al., 2012; Prinsenberg and Hamilton, 2005),  $0.71 \pm 0.09$  and  $1.03 \pm 0.11$  Sv 30 between 2003 to 2006 and 2007 to 2009, respectively, along Nares Strait (Munchow and Humfrey, 2008; Münchow, 2016), and 0.3 Sv between 1998-2002 flowing through Jones Sound (Melling et al., 2008). Thus, the rough balance of transport through these three main passages is 46% Nares Strait, 34% Lancaster Sound and 20% Jones Sound (Melling et al., 2008; Grivault et al., 2018).

Jones Sound is home to the Inuit hamlet of Ausuiktuq (Grise Fiord) and is a marine region surrounded by glaciers draining large ice fields and ice caps on both the Ellesmere and Devon Islands. It is the third largest export pathway in the CAA (Grivault 35 et al., 2018; Melling et al., 2008; Zhang et al., 2016), connecting directly to the Arctic Ocean at the narrow (15 km) and shallow (150 m) western gateways of Hellsgate/Cardigan Strait (which merge into Fram Sound) and exchanging with Baffin Bay on the east side of the sound at a depth of 450 m (figure 1). The western half of Jones Sound is shallow, being  $\sim 200$  m depth, while the eastern basin is deeper, having a maximum depth of  $\sim 840$  m and greater exchange with external waterways. Aside 40 from facilitating liquid exchange between the Arctic and northern Atlantic Oceans, Jones Sound also hosts a diverse biological ecosystem that is sustained in part by ice-ocean interactions of tidewater glacier termini as well as seasonally-varying ocean and sea ice conditions (Bhatia et al., 2021). However, as part of the broader Arctic, Jones Sound is vulnerable to changing climate conditions that threaten these natural resources. For instance, Gardner et al. (2012) found that ice mass loss across the QEI has tripled from  $31 \pm 8$  Gt/yr in 2004-2006 to  $92 \pm 12$  Gt/yr in 2007-2009, largely driven by Arctic amplified atmospheric warming that outpaces the global average by three-times. Below the halocline within the Baffin Bay, mid-depth Atlantic Water 45 (AW) that penetrates into the CAA and Arctic Ocean has been warming steadily since at least the early 2000s (Polyakov et al., 2013; Wang et al., 2020; Ballinger et al., 2022). This atmospheric and oceanic warming has also driven recent declines in sea ice extent, with the summer sea ice area in northern Canadian waters and Baffin Bay declining at a rate of 7.1% per decade and 11.6% per decade, respectively (Tivy et al., 2011). Given that sea ice serves as hunting platforms for polar bears, resting 50 grounds and nursery areas for walruses and seals, and hosts for algae that grow on the ice base, these reductions in sea ice have cascading impacts on marine ecosystems.

Due to the maze of islands, narrow straits, complex coastlines, and shallow bathymetry, modeling ocean circulation, sea ice dynamics, biological productivity, and their joint interactions within the CAA is challenging but remains one of the best ways to begin understanding recent change of CAA waterways. Recent ocean modeling studies are largely capable of resolving mean transport through Lancaster Sound and Nares Strait (McGeehan and Maslowski, 2012; Shroyer et al., 2015; Wang et al., 55 2017; Wekerle et al., 2013; Zhang et al., 2016); however, coarse horizontal meshes cause issues with resolving baroclinic flow in narrower channels, such as Fram Sound, leading to large uncertainty in volume estimates into and general circulation within



**Figure 1.** Domain featuring ocean bathymetry (m) from the SRTM15+ dataset (Tozer et al., 2019), where blue shading denotes bathymetry below sea level. Gray shading denotes the present-day extent of land ice, the red box denotes the domain of the high-resolution ocean circulation model, and the black box denotes where new bathymetry observations were implemented in the ocean model in Brae Bay. Geographic features mentioned in the text are labeled, including glaciers, water ways, and islands.

Jones Sound. This, in turn, limits the fidelity of sea ice and productivity models in this region. To address this, we develop a high-resolution, Jones Sound configuration of the Massachusetts Institute of Technology general circulation model and perform coupled ocean-sea ice-biological productivity simulations between 2003-2016 to investigate the following: (1) The magnitude and spatial/temporal distribution of volumetric transport into and out of Jones Sound; (2) Fine-scale circulation features within Jones Sound and their impact on watercolumn structure on seasonal timescales; (3) Seasonal and decadal variations in sea ice dynamics; (4) The impact of simulated changes in ocean and sea ice conditions on productivity within Jones Sound. By investigating these four topics, we seek to improve our understanding of circulation, sea ice dynamics, and productivity within

Jones Sound and how it fits within the broader context of the Canadian Arctic Archipelago. Below, we provide information on  
65 the setup of the numerical ocean, sea ice, and biogeochemical model as well as an overview of the simulations.

## 2 Methods

Here, we provide information on the setup of the ocean, sea ice, and biogeochemical model that we use to simulate circulation  
in Jones Sound between 2003-2016. We note that we first developed a low resolution ocean-sea ice model to simulate the region  
surrounding Jones Sound between 2002-2016 (hereon referred to as the “low-resolution simulation”), from which we extract  
70 initial and boundary conditions to force our high-resolution ocean-sea ice-productivity simulation of Jones Sound between  
2003-2016 (hereon referred to as the “high-resolution simulation”). Most of the model setup between the two simulations is  
identical aside from the source of the boundary and initial conditions as well as the domain extent and resolution.

### 2.1 Ocean model setup

We model ocean circulation using a regional Canadian Arctic configuration of the ocean component of the Massachusetts  
75 Institute of Technology general circulation model (MITgcm; Marshall et al., 1997). This includes use of the hydrostatic approximation, a dynamic/thermodynamic model to simulate sea ice dynamics (Losch et al., 2010), and the Biogeochemistry with Light, Iron, Nutrients, and Gases with Nitrogen (N-BLING) productivity module to simulate photosynthetic biological productivity (only used in the high-resolution simulation and described in section 2.2; Galbraith et al., 2010). The low-resolution simulation domain extends from 73.25 - 76.75°N, 74 - 91.5°E, has a nominal horizontal resolution of 1/24 degree (~5 km)  
80 and contains 70 vertical levels (with vertical resolution of 7 m through 266 m depth, then decreasing to a minimum resolution of 62 m at the lowest ocean level; 1090 m). The high-resolution simulation domain extends from 75.40 – 76.70°N, -77.45 – -91.45°E (red outline in figure 1), has a nominal horizontal resolution of 1/120 degree (~900 m) and contains 70 vertical levels (with vertical resolution of 7 m through 266 m depth, then decreasing to a minimum resolution of 56 m at the lowest depth of 963 m). Model bathymetry is based on the SRTM15+ digital elevation model (Tozer et al., 2019) with corrections applied  
85 in Brae Bay (black box near Sverdrup Glacier in figure 1) and the oceanic regions near Grise Fiord following multibeam and point measurements. Ocean and sea ice parameter values that differ from Nakayama et al. (2018) are provided in Table 1.

Initial and boundary conditions for the low-resolution ocean simulation are extracted from the 1/12 degree Arctic and Northern Hemisphere Atlantic (ANHA12) configuration of the NEMO model that covers the entire North Atlantic Ocean down to 20°S and was run from 2002-2018 (Hu et al., 2019; Gillard et al., 2020). Fields used as initial and boundary conditions in our  
90 ocean simulations include temperature, salinity, the zonal (u) and meridional (v) velocity components, and sea surface height. In addition, fields used to initialize and drive the sea ice model include sea ice area, thickness, snow content, salt content, and the u and v sea ice velocity components. Initial conditions were extracted at model date Jan. 1, 2002 and boundary conditions are extracted monthly through Dec. 31, 2016. Initial and bi-monthly boundary conditions in the high-resolution simulation are extracted from the low-resolution simulation, with an initialization date of Jan. 1, 2003.

**Table 1.** MITgcm ocean and sea ice model parameters and values used in this study. Only parameters that are different from Nakayama et al. (2018) are shown.

| Parameter (unit)                          | Value   |
|---|---------|
| Ocean/air drag coefficient scaling factor | 0.00125 |
| Air/sea ice drag coefficient              | 0.00125 |
| Lead closing (m)                          | 1       |
| Sea ice dry albedo                        | 0.72    |
| Sea ice wet albedo                        | 0.63    |
| Snow dry albedo                           | 0.78    |
| Snow wet albedo                           | 0.65    |
| Ocean emissivity                          | 0.97    |
| Ice emissivity                            | 0.95    |
| Snow emissivity                           | 0.95    |

95 Atmospheric forcing is taken in three-hour intervals from the Arctic System Reanalysis version 2 (ASRv2; Bromwich et al., 2018) and interpolated onto the model grids. We use the following variables: 2 m air temperature, 2 m specific humidity, precipitation, 10 m u and v wind components, short/long wave radiation, atmospheric pressure, evaporation, and river/glacial runoff. The time series of all atmospheric forcing fields are shown in Figure 2. In addition, tidal forcing (amplitude and phase) is prescribed in the high-resolution simulation using the Arctic 2 kilometer Tide Model (Arc2kmTM; Howard and Padman, 100 2021) and includes the following constituents: M<sub>2</sub>, S<sub>2</sub>, N<sub>2</sub>, K<sub>2</sub>, K<sub>1</sub>, O<sub>1</sub>, P<sub>1</sub>, and Q<sub>1</sub>. In MITgcm, tidal forcing is applied along the ocean model boundaries as a surface elevation perturbation and as barotropic flow that propagates throughout the domain.

## 2.2 N-BLING setup

The Nitrogen version of the Biogeochemistry with Light, Iron, Nutrients, and Gases (N-BLING) productivity module simulates biogeochemical cycling of key elements/micronutrients as well as photosynthetic productivity and has been implemented in 105 MITgcm as a module (Galbraith et al., 2010; Verdy and Mazloff, 2017). N-BLING is driven by the physical ocean model as well as atmospheric carbon dioxide concentrations, which are taken monthly between 2003-2016 from a meteorological station in Alert, Canada and assumed to be spatially uniform across our high-resolution domain. Incoming solar radiation, taken from ASRv2, also drives N-BLING. Initial and monthly boundary conditions for N-BLING include the following: dissolved inorganic carbon (DIC) and alkalinity taken from the Global Ocean Data Analysis Project version 2 (GLODAPv2; Olsen et al., 110 2016); O<sub>2</sub>, NO<sub>3</sub>, PO<sub>4</sub>, and Silica taken from the World Ocean Atlas (WOA; Garcia et al., 2018); Fe, dissolved organic nitrogen, dissolved organic phosphorus, and initial small/large/diazotroph phytoplankton from a global run of BLINGv2 (Pers. Comm. Eric Galbraith); and iron dust deposition from Mahowald et al. (2005). N-BLING runs on the same computational grid and

115 timestep as the ocean model and coupling is only one-way, meaning that N-BLING-simulated productivity does not influence the radiative fluxes, and thus the ocean circulation and sea ice growth.

### 115 2.3 Model runs


We first run the low-resolution ocean-sea ice simulation between Jan. 1, 2002 to Dec. 31, 2016 using a baroclinic timestep of 120 seconds. From the low-resolution simulation, we extract initial and boundary conditions to force the high-resolution ocean-sea ice-biological productivity simulation, which is run between January 1, 2003 to Dec. 31, 2016 using a baroclinic timestep of 70 seconds. We performed simulations that do and do not resolve tidal forcing to explore the impact of tides on 120 ocean circulation and sea ice productivity in Jones Sound (see Appendix-a). All results discussed below are taken from the high-resolution simulation. We note that N-BLING crashed in May 2015, so we only report productivity results up to then. We describe the cause of this crash in the discussion section.

## 3 Results

### 3.1 Model Forcing

125 Between 2002-2016, there exist large trends in atmospheric forcing variables in ASRV2 over Jones Sound. The time series of all atmospheric forcing variables are shown in figure 2, where domain-wide means are taken except for evaporation, precipitation, and runoff, which are integrated across the domain. In panels a-i, we overlay the overall and summer linear trends as colored solid and dotted lines, respectively, and report the total trend in the title of each panel. In panel-j, we show trend lines on variables that have been normalized. Overall, we find strong increasing trends in the 2 m air temperature, which is increasing 130 at an overall rate of 0.87°C/decade and a summer rate of 1.19°C/decade (1.37 times higher than the overall trend). This strong surface warming trend drives strong positive trends in summer relative humidity (3.76e-4 kg/kg/decade; 2.81 times stronger than the overall relative humidity trend), summer and overall evaporation, and summer runoff (6.583e-4 m/s/decade). We see minimal changes in mean atmospheric pressure, mean short and long wave radiation, mean 2 m wind speed, and total integrated precipitation.


135 Along the ocean model boundaries, we find large changes in the ocean state between 2002 and 2016 as simulated by the ANHA12 configuration of NEMO, from which we extract our ocean boundary conditions. Figure A1 shows vertical profiles of ocean potential temperature applied to the Eastern boundary (-74°W) of the low-resolution ocean simulation in Jan. of 2002, 2007, 2012, and 2016. A strong warming signal of the winter mid-depth Atlantic Water (between 100-300 m depth) is evident after 2007, where waters warm by over 2°C in ~10 years in this simulation. We note that this simulated warming signal of 140 the Atlantic Water is overestimated, as oceanic observations taken on the Arcticnet Amundsen Ice Breaker report 1-1.5°C of warming of this water mass in northern Baffin Bay over this same time period. We discuss the implications of this overestimated ocean warming in the discussion section.



**Figure 2.** Time series of ASRv2 (a) 2 m air temperature, (b) surface atmospheric pressure, (c) relative humidity, (d) downward longwave radiation, (e) downward shortwave radiation, (f) 2 m wind speed, (g) evaporation, (h) precipitation and (i) runoff. Panels a-f show domain-wide averages while panels g-i (names appended with “\*”) show sums integrated across the domain. In each panel, the best fit lines for all-data (solid colored line) and for summer-data (dotted colored line) are overlaid and the slope of the all-data line (per decade) is included in the title. Panel-j plots all best-fit lines computed on data that have been normalized. The lines are then vertically shifted so that the initial value is 0 to facilitate trend comparison.

### 3.2 Model Evaluation


We evaluate our coupled ocean-sea ice model of Jones Sound against repeat Summertime casts of Conductivity, Temperature, 145 Depth (CTD) ocean sensors made aboard the Amundsen Ice Breaker between 2005 and 2021 (Amu, 2003-2021). We selected five sample sites based on data availability within our model domain: two locations spanning longitudinally across northern Baffin Bay, one site in the center of the eastern entrance of Lancaster Sound, one site close to Belcher Glacier, and one site near the center of Jones Sound. Observed and modeled temperature profiles are plotted in figure 3a/c/e/g and figure 3b/d/f/h, respectively (solid and dotted lines, respectively). The color of the profile is associated with the year it was taken. Note that 150 gray solid profiles indicate observed profiles taken after 2016 and dotted gray profiles were extracted in summer 2016. We included these profiles because the only currently publicly available CTD casts within Jones Sound in this dataset were taken



**Figure 3.** Comparison of observed (panels a/c/e/g, solid lines) and modeled (panels b/d/f/h, dotted lines) ocean temperature vertical profiles taken at western Baffin Bay (panels a-b), central Baffin Bay (panels c-d), the eastern entrance of Lancaster Sound (panels e-f), and near Belcher Glacier (panels g-h). The color of the line indicates the year of the profile and gray denotes any profile taken after 2016. Observations are summertime CTD-rosettes taken aboard the Amundsen Ice Breaker (Amu, 2003-2021) and the brown line in panel-b is a profile from Bhatia et al. (2021) that was taken in summer 2019 (B-19'). (i-j) Temperature-salinity plots taken in (i) 2019 in Jones Sound and (j) 2015 in the eastern entrance of Lancaster Sound. The color of the marker corresponds to the depth. The observed and modeled profiles are labeled.

in 2019 and 2021. The brown line in panel-g was measured in the summer of 2019 by Bhatia et al. (2021). Note that we also present evaluation of salinity profiles in Appendix A1 (figure A2).


In the profiles taken across northern Baffin Bay and in Lancaster Sound (figure 3a-f), we find that our model is generally able to replicate the summertime thermal properties of the ocean in these sample locations. In particular, we properly model the depth of the thermocline (located between 100-300 m depth) and mid-depth Atlantic Water temperatures at all sample sites. One notable exception to this is a warm bias in the modeled seabed ocean temperatures in the sample site in Northern Baffin Bay (figure 3c-d), where observed ocean temperatures increase from  $-0.5^{\circ}\text{C}$  to  $-1^{\circ}\text{C}$  between 2005-2016 but modeled ocean temperatures increase from  $0^{\circ}\text{C}$  to  $1.5^{\circ}\text{C}$  during this same time period. That is, we overestimate warming of the bottom ocean water in Baffin Bay but reasonably match warming of the mid-depth Atlantic Water in the summer. We expect that this overestimated bottom ocean warming stems from the overestimated warming in our NEMO-derived boundary conditions



**Figure 4.** Monthly snapshots of sea ice concentration observations in 2015 from the Canadian Ice Service, which are created through the manual analysis of in situ, satellite, and aerial reconnaissance data (Service, 2009). Gray shading denotes land.

(figure A1). Near Belcher Glacier, we properly simulate ocean thermal conditions between 2005-2007; however, we find that ocean temperatures below the thermocline are  $\sim 1^{\circ}\text{C}$  too warm at the end of our simulation, meaning that the warm bias simulated in Baffin Bay extends into Jones Sound as well (figure A4g-i). Furthermore, we note that simulated vertical profiles 165 of salinity generally show good agreement with observed profiles (figure A2 and figure 3i-j). However, we note that modeled salinity gradients with depth are weaker than observed. For instance, in Lancaster Sound, ocean waters below  $\sim 300$  m are too fresh by  $\sim 0.5$  PSU (figure A4j). These reduced salinity gradients could impact the magnitude of current velocities in our model. However, as we correctly simulate the depth of the thermocline, thermal properties of water above the thermocline, and 170 salinity gradients in mixed-layer, we believe the modeled circulation, the extent to which warmer waters can spatially extend, and surface properties of the ocean model are well resolved.

We further evaluate the sea ice model by comparing monthly observed (figure 4) and modeled (figure 5) timestamps of 2015 sea ice concentration. While we only show the 2015 sea ice cycle, we note that the other yearly cycles feature similar spatiotemporal patterns of annual change. Observed 2015 sea ice concentration fields are taken from the Canadian Ice Service and are created through manual analysis of in situ, satellite, and aerial reconnaissance data (Service, 2009). We find generally 175 good agreement between the observed and modeled sea ice concentration states within Jones Sound for the 2015 sea ice cycle,




**Figure 5.** Monthly timestamps of simulated surface ocean grid cell sea ice concentration and contoured sea ice thickness (contoured every 0.5 m). Gray shading indicates land.

with complete sea ice coverage simulated across the domain between November-April and near complete meltback of sea ice in August and September. Importantly, we also find that our model is able to replicate the spatial pattern of sea ice meltback, with flow through Fram Sound initiating sea ice decline in northwestern Jones Sound and partial clearing of sea ice across southeast Jones Sound. We note that modeled sea ice thickness east of Coburg Island is overestimated and sea ice does not clear out of this region during the summer months, disagreeing with observations. As such, in section 3.4, we only integrate sea ice quantities west of Coburg Island.


### 3.3 Ocean circulation through Jones Sound

Circulation through Jones Sound is dictated by flow through three waterways: Fram Sound (the confluence of Cardigan Strait and Hell Gate) in the northwest, Lady Ann Strait in the southeast, and Glacier Strait in the northeast. In figure 6a-d, we show 185 timeseries (a-b) and time-mean bar plots (c-d) of volumetric transport integrated through these three gateways into Jones Sound (lines in figure 8a). Of the temporal-mean 1.31 Sv that flows into and out of Jones Sound, flow through Lady Ann Strait, Fram Sound, and Glacier Strait comprise 71.14% (0.93 Sv), 14.26% (0.19 Sv), and 14.40% (0.19 Sv) of this volumetric transport, respectively (figure 6). Flow through each of these waterways is a mix of inflow and outflow into the sound; Lady Ann Strait



**Figure 6.** Time series of net (solid lines) and the magnitude (dotted lines) of volumetric transport through Fram Sound (blue), Glacier Strait (green), and Lady Ann Strait (red) with (a) tidal forcing included and (b) tidal forcing not included. Positive and negative values indicate transport into and out of Jones Sound and note the different y-axis limits between the two panels. (c-d) Bar plots of the time-mean transport into (blue) out of (orange) Jones Sound, with the mean net transport indicated by the yellow bar. Panel-c includes tides and panel-d does not include tides. Error bars indicate one standard deviation from the mean. (e-f) Vertical cross sections of time-mean ocean velocity through (e) Fram Sound, (f) Lady Ann Strait, and (g) Glacier Strait.

is comprised of  $0.37 \pm 0.12$  Sv inflow and  $0.56 \pm 0.15$  Sv outflow, Fram Sound is comprised of  $0.12 \pm 0.04$  Sv inflow and 190  $0.06 \pm 0.02$  Sv outflow, and Glacier Strait is comprised of  $0.16 \pm 0.09$  Sv inflow and  $0.03 \pm 0.03$  Sv outflow (taken from figure 6c-d). Of particular importance is the influence of tidal flushing on transport through Fram Sound, enhancing its total volumetric transport by 47%. For the other two waterways, tidal flushing had minimal impact on total volumetric transport.




**Figure 7.** (a-f) Time and depth (down to 150 m) mean ocean velocity (m/s) fields with flow vectors overlaid. Panels a-c and d-f show March and September mean velocity fields, respectively. (g-l) Vertical profiles of ocean temperature ( $^{\circ}$ C) taken along the red line in panel-a. Note that the first, second, and third columns show 2014, 2015, and 2016 mean fields, respectively.

In figure 6e-g, we plot vertical cross sections of the 2003-2016 mean velocity fields through transects across Fram Sound (e), Lady Ann Strait (f), and Glacier Strait (g; yellow, red, and green lines in figure 8a, respectively) to visualize regions of flow into (red, positive) and out of (blue, negative) Jones Sound. Time-mean ocean velocities through Fram Sound are the highest across all three waterways (maximum velocity of  $\sim 0.7$  m/s), with regions of inflow occupying the top 65 m of the water column and outflow located below (figure 6e). That is, while Fram Sound is the smallest waterway leading to Jones Sound by cross-sectional area, it has the second greatest integrated volumetric transport passing through it because of these high flow velocities. For Lady Ann Strait, the primary region of inflow is located near the south end of Coburg Island and persists through 195 depth (figure 6f). We model a secondary weak region of inflow on the southern end of the strait below 200 m depth. Otherwise, outflow dominates circulation through the strait, with the strongest outflow velocities comprised of a persistent strong current that wraps around eastern Devon Island in the upper 100 m of the water column. Flow through Glacier Strait is dominated by inflow into Jones Sound, with a small region of outflow modeled on the extreme southern end of this waterway (figure 6g). 200 Overall, these results highlight the complex spatial and temporal regimes of transport into and out of Jones Sound.

205 In figure 7a-f, we show March (a-c) and September (d-f) mean ocean velocity fields that are averaged through 150 m depth at year 2014 (a/d), 2015 (b/e), and 2016 (c/f). We selected March and September to capture circulation in which Jones Sound is completely covered with sea ice and completely sea ice free. In March under full sea ice coverage, the flow field is quiescent, with strong inflow (up to 0.5 m/s) through Fram Sound in the west and strong outflow (>0.5 m/s) through Lady Ann Strait in the east. The strongest currents from northern Baffin Bay tend to flow east of Coburg Island; however, we do simulate weak flow 210 of up to 0.2 m/s through Glacier Strait into Jones Sound. In September, however, reduced sea ice coverage allows atmosphere-ocean interactions, which drives the formation of persistent eddies that dominate circulation within Jones Sound (figure 7d-f). These eddies do not always form in the same place and rotate in the same direction. For instance, in 2014 (figure 7d), we model three sustained cyclonic eddies (rotating counterclockwise), where the easternmost eddy does not interact with strong inflow through Glacier Strait. In 2015 (figure 7e), we observe one anticyclonic eddy in the center of Jones Sound that is flanked by 215 two cyclonic eddies, where circulation through Glacier Strait interacts with the easternmost eddy. Lastly, in 2016 (figure 7f), we model a similar eddy configuration as in 2015. The diversity of summertime circulation patterns across 2014-2016 in Jones Sound highlights the complex dependence of current-patterns on inflow/outflow regimes and ocean-atmosphere interactions.

220 These circulation patterns influence the thermodynamic structure of the water column within Jones Sound. In figure 7g-l, we plot vertical temperature profiles across Jones Sound at 76°N (red line in figure 7a). As in the velocity fields, the third and 225 fourth row of the figure correspond to March and September mean temperature fields and the first, second, and third columns correspond to 2014, 2015, and 2016 fields, respectively. In all temperature fields, we note that the modeled thermal structure is homogeneous beneath the thermocline, as has been observed in Jones Sound (figure 3i). Under complete sea ice coverage, the modeled thermocline is nearly uniform at a depth of ~100 m across Jones Sound (figure 7g-i). In September, the development 230 of eddies drives spatial variation in the modeled depth of the thermocline, with the location of the eddy-centers corresponding to locations of thermocline upwelling and downwelling. In September 2015, we see that the thermocline is nearly uniform at ~200 m depth across Jones Sound, which is ~100 m deeper than the March thermocline depth.

235 The simulated spatial reach of warm waters below the thermocline in Jones Sound is topographically constrained. In figure 8, we plot the modeled ocean bottom temperature and velocity vectors in March and September of 2010, 2013, and 2016. As noted in the validation above, our model overestimates warming of waters below the thermocline in Jones Sound; however, as we correctly model the depth of the thermocline, we intend for the temperature fields shown here to be viewed as tracers of “warm” ocean water and its general circulation throughout the sound. It is evident that warm ocean waters flow into the model domain from the eastern model boundary, highlighting that the source of warm water inflow into Jones Sound is through northern Baffin Bay. This warm water then circulates into Jones Sound from the southeast through deep bathymetry that underlies Lady Ann Strait (>600 m; figure 1). Inflow of deep warm ocean water is blocked by shallow bathymetry along Glacier Strait, which sits between 100-300 m depth. Once the warm bottom water intrudes into Jones Sound after 2010, it circulates counter-clockwise and although topographically constrained, warm bottom water breaches the entrance of most major fjords along the Eastern periphery of the Sound by 2012 (where bathymetry is generally deeper than that of the western sound). In particular, warm bottom water reaches the termini of Sverdrup, Jakeman, and Belcher Glaciers (figure A3). Depression of the thermocline in September relative to March constrains the reach of this warm bottom water. This can be seen in Brae Bay and western Jones




**Figure 8.** Modeled ocean bottom potential temperature (shaded) and velocity (vectors) in Jones Sound, Canadian Arctic Archipelago. The top row shows fields in March of (a) 2010, (b) 2013, and (c) 2016 while the bottom row shows fields in September (d) 2010, (e) 2013, and (f) 2016. In panel-a, the yellow, red, and green lines denote the location of the vertical cross sections through Fram Sound, Lady Ann Strait, and Glacier Strait that are shown in figure 6e-g, respectively.

240 Sound when comparing figure 8b and figure 8e. In regions where this warm water cannot circulate (regions of bathymetry above the thermocline), September water temperatures are  $\sim 0.5^{\circ}\text{C}$  warmer than March water temperatures due to enhanced mixing.

### 3.4 Sea ice decline

245 Sea ice is a prominent feature of ocean circulation and dynamics within Jones Sound, regulating the exchange of heat, moisture, and momentum between the ocean and atmosphere, as well as blocking the penetration of sunlight that fuels photosynthetic biological productivity. In figure 5, we plot monthly timestamps of grid cell sea ice concentration and contoured sea ice thickness during model year 2015. Sea ice undergoes a yearly cycle in which the sound becomes largely ice-free in September and completely ice-full in the winter months. Oceanic flow through Fram Sound initiates sea ice decline in northwestern Jones Sound typically beginning in April. Sea ice then retreats south across Jones Sound, clearing within the northern fiords first and 250 then melting back across the rest of the sound by September. While most of the sound remains ice free in September, Coburg Island acts as a sea ice bottle neck in our model (and in observations), trapping ice that circulates between the open waters of Baffin Bay and the confined waters of Jones Sound. During the winter, sea ice thickness generally fluctuates between 1-2 m across most of the sound. However, there are isolated pockets of thicknesses  $>4$  m, which are primarily constrained to within



**Figure 9.** Integrated sea ice (a-b) area and (c-d) volume. In panels a/c, marker colors correspond to the area of integration: black (all of Jones Sound), purple (western Jones Sound), and green (eastern Jones Sound). Circles and triangles denote yearly sea ice maxima and minima. Best fit lines are overlaid on the East Jones Sound area and volume minima (green dashed lines) and the full Jones Sound volume maxima (black dashed line) to show long-term trends. In panels b/d, the 2003-2009 and 2010-2016 mean yearly cycles and standard deviations are plotted as the blue and red lines, respectively. Each yearly cycle is overlaid as transparent blue or red lines.

narrow fiords, along the periphery of Coburg Island, and in the open waters of Baffin Bay. This thick sea ice along Coburg  
255 Island and within Baffin Bay is not supported by observations and represents a model bias. We note that while we only show  
the 2015 sea ice cycle, these general patterns persist in other modeled years.

To investigate how the annual sea ice cycle within Jones Sound changes between 2003-2016, we present integrated sea  
ice area (summed areas of grid cells that have a sea ice fraction greater than 0) and volume (area multiplied by the sea ice  
thickness) in figure 9 across the full extent of Jones Sound (using the western end of Coburg Island as our eastern limit; black  
260 lines), the western half of Jones Sound (purple lines), and the eastern half of Jones Sound (green lines). In figure 9, over these  
regions of integration, we plot the integrated sea ice maxima (circles) and minima (triangles) and overlay linear trendlines  
as dashed lines where trends are evident. In panels b and d, we plot the average 2003-2009 and 2010-2016 yearly cycles of  
the aforementioned variables and their associated standard deviation, respectively. Beginning with the integrated sea ice area

(figure 9a), we observe a temporal decline in minimum sea ice area of  $316 \text{ km}^2/\text{yr}$  in Eastern Jones Sound, mainly associated  
265 with enhanced melt of sea ice pinned against Coburg Island and within Eastern Jones Sound (figure 5). When the yearly total  
Jones Sound sea ice area cycles are binned to 2003-2009 and 2010-2016 and averaged (blue and red lines in figure 9b), we  
further observe that the initiation of sea ice decline is occurring earlier and the onset of sea ice refreezing is occurring later  
270 in the 2010-2016 profiles. Furthermore, although we find that winter sea ice area remains stable in our model (Jones Sound  
becomes completely ice-covered in winter), we observe that the thickness of this winter-ice is declining in time (linear volume  
275 trend of  $-0.384 \text{ km}^3/\text{yr}$ ; black-circle line in figure 9c). In fact, the binned-yearly cycles of sea ice volume reveal that sea ice is  
generally thinner throughout the entire yearly cycle aside from in June and July (figure 9d).

To investigate the impact of tides on sea ice formation and melt within Jones Sound, we perform another simulation that  
does not include tides and find that tidal flushing through Fram Sound can enhance mean flow velocities by up to  $0.75 \text{ m/s}$   
280 over that of the non-tidal run in these regions (figure A6). These tidally-enhanced flow velocities through Fram Sound trigger  
accelerated sea ice meltback in northern Jones Sound between May and July (blue shading in figure A7e-g) while leading to  
generally thicker sea ice in the southwest sector of Jones Sound (due to ice advection into the southwestern fiords). Similar to  
figure 9, we integrate sea ice area and volume across Jones Sound and provide the associated time series in figure A8. Zooming  
into the 2014-2016 cycles, we find that tidal forcing drives a longer ice-free season within Jones Sound, decreasing the summer  
integrated sea area by up to  $6000 \text{ km}^2$  (figure A8b-c). Furthermore, tidal forcing also decreases sea ice thickness year-round  
due to both enhanced circulation velocities and mixing that entrains heat at depth to the surface (figure A8e-f).

### 3.5 Productivity enhancement

To begin deducing how these temporal changes in the state of Jones Sound sea ice and ocean circulation feedback on photosynthetic biological productivity, we couple our ocean-sea ice model to the MITgcm N-BLING biological productivity module and  
285 plot yearly- and depth-integrated profiles of Net Primary Production (NPP; figure 10a), yearly- and depth-integrated profiles  
of chlorophyll mass (CHL; figure 10b), yearly integrated profiles of Net Community Production (NCP; figure 10c/d), yearly-  
integrated profiles of light limitation (figure 10e), and yearly-integrated profiles of iron limitation (figure 10f). Light and iron  
limitation are computed as the percentage of ocean grid cells per vertical level that experience light or iron limitation, so a  
value of 100 indicates 100% of ocean grid cells are light or iron limited. NCP represents the difference between gross primary  
290 production and total community respiration; that is, when NCP is positive, photosynthetic primary productivity is greater than  
community respiration and vice versa when negative. In figure 10c/d, we observe that NCP increases in the first and second  
vertical ocean levels (corresponding to depths of  $3.5 \text{ m}$  and  $10.5 \text{ m}$ , respectively, as the tracer point within MITgcm is in the  
mid-depth of the grid cell). When we zoom into the transition zone between positive and negative NCP, we further observe  
that the third ocean level (corresponding to a depth of  $17.5 \text{ m}$ ) transitions from pure respiration before 2010 to a source of  
295 community production in the fall and summer beyond 2010. To investigate this further, we observe a strong increase in surface  
and subsurface NPP after 2010. Prior to 2010, total time integrated surface ( $3.5 \text{ m}$  depth level) NPP averaged  $\sim 0.135 \text{ GtC/yr}$   
and after 2010, averaged  $\sim 0.170 \text{ GtC/yr}$ . These productivity enhancements are also elevated at  $10.5 \text{ m}$  depth (the second ocean  
depth level), where NPP before and after 2010 averaged  $0.0051 \text{ GtC/yr}$  and  $0.0075 \text{ GtC/yr}$ , respectively. We observe that the


light limitation time series follows suit, as the mean percentage of summer light limitation of the surface level before and after 2010 is modeled as 48.08% and 43.73%, respectively. Similarly, at 10.5 m depth, the 2005-2010 and 2010-2015 mean summer 300 light limitation is 64.24% and 56.48%, respectively. Interestingly, for the first two ocean depth levels (through 10.5 m depth), we observe that the percentage of ocean grid cells that are iron limited in the summer increases after 2010; however, for these same levels, winter iron limitation steadily decreases throughout the simulation at a rate of  $\sim 0.615\%/\text{decade}$ , implying more mixing. For depth levels below 10.5 m, the yearly cycle of iron limitation decreases linearly at approximately this same rate 305 (yellow-to-red lines in figure 10f). We note that there are no temporal trends evident in the time series of phosphorus and nitrogen limitation and that both fields displayed decreasing percentages of limitation with depth.

## 4 Discussion

### 4.1 Broader implications of Jones Sound circulation

The modeling results presented here highlight the dynamic ocean state and circulation patterns within Jones Sound as well as 310 its joint impact on sea ice dynamics and biological productivity between 2003-2016. In our model, we find that the primary response of Jones Sound broad-scale atmospheric and oceanic changes is warming of the waters below 200 m depth. While 315 the magnitude of this warming signal is overestimated in our model due to the Atlantic Water at our Eastern model boundary warming too quickly in winter, we correctly model the depth of the thermocline as well as currents across our model domain, giving us confidence that we correctly simulate the spatial distribution of where these warmer mid-depth waters can circulate 320 as well as volumetric transport throughout the domain. In particular, we find that circulation through Lady Ann Strait dominates volumetric transport into and out of Jones Sound, followed equally by Glacier Strait and Fram Sound. In the latter, the magnitude of transport through Fram Sound reaches  $\sim 0.25 \text{ Sv}$  in summer months, agreeing with mooring data collected by Melling et al. (2008). These regions of inflow/outflow drive complex spatial patterns of circulation within Jones Sound, with 325 the formation of multiple eddies driving variability in the thermocline depth in the summer. In the winter, under full sea ice coverage, the thermocline shoals under reduced atmospheric forcing. We further find that warm water within Jones Sound is topographically constrained, flowing through Lady Ann Strait and circulating counter-clockwise within the sound, reaching 330 many of the tidewater glaciers that line the sound's eastern coast (figures 8 and A3). In combination with continued observed atmospheric warming (figure 2a), these results underscore the increasing vulnerability of the ice masses that populate the Devon and Ellesmere Islands.

Furthermore, our modeling results highlight the importance of circulation through Fram Sound in triggering sea ice decline 325 in northern Jones Sound during the summer. Such relationships between tidal forcing and sea ice decline have been studied in other sectors of the Arctic and CAA (e.g., Rotermund et al., 2021; Armitage et al., 2020) but have yet to include Jones Sound due to the relatively small size of the channels that feed this waterway. The complex dependence of Jones Sound sea ice dynamics/thermodynamics on small-scale tidally-forced ocean circulation features (as well as the importance of properly 330 modeling sea ice change for summertime transportation through these waters) highlights the need for ocean models of this



**Figure 10.** Bar plots of yearly and depth binned (a) net primary productivity, and (b) chlorophyll. Panels c-d show time series of net community productivity integrated on the top 20 vertical ocean levels (through 140 m depth), where the bold black line in panel-d denotes values for the third vertical ocean level (17.5 m depth). Panels e and f show the percent of ocean grid cells per each vertical level that are light and iron limited, respectively.

330 region to both explicitly include tidal forcing and be run at sufficient resolution to resolve the eddies that disperse subsurface heat.

In addition to these sub-annual patterns in simulated sea ice dynamics, we observe long-term declines in the Jones Sound integrated summer sea ice area as well as both the summer and winter sea ice volume. The summertime sea ice area declines on average  $\sim 11\%/\text{decade}$ , which is inline with the observed losses over Baffin Bay of  $11.7\%/\text{decade}$  between 1968-2022 and 335 slightly higher than the  $7.1\%/\text{decade}$  losses observed across all of northern Canada's waters over this same time period (Tivy et al., 2011). These losses mimic broader patterns of summer Arctic sea ice decline, which are cited to be driven by both natural

climate variability (Kinnard et al., 2011; Ding et al., 2017, 2019) and human-induced global warming (Kay et al., 2011; Notz and Stroeve, 2016; Stroeve and Notz, 2018). Here, however, we also show that winter sea ice in Jones Sound is also thinning (figure 9; winter sea ice area does not change in time but winter volume decreases in time), which is likely driven by winter warming of subsurface ocean temperatures (figure 2a and figure A5). Lastly, we simulate earlier onset of sea ice decline in the summer and later onset of sea ice refreeze in the fall between 2003-2016. That is, the period of time in which Jones Sound is not completely filled with sea ice is extending in time in our model, which impacts the timing and integrated magnitude of photosynthetic oceanic primary productivity.

We observe in our biological modeling results that as the time in which Jones Sound is sea ice free becomes longer, the total integrated oceanic productivity and the depth at which productivity takes place increases. We note that aside from the atmospheric carbon dioxide forcing time series (which largely increases linearly), the boundary conditions in the biological productivity module do not include significant temporal trends and also do not account for nutrient release from enhanced glacial meltwater/discharge. That is, the response of ocean primary productivity in our model is driven primarily by changes in local oceanic and atmospheric conditions. Local enhancements to primary productivity have been reported across the Arctic, with the mean annual (March-September) trend of primary productivity increasing  $\sim$ 50-75 g C/m<sup>2</sup>/year/decade within the central portion of Jones Sound between 2003-2022 (Frey et al., 2022). While this observed trend includes enhanced productivity due to increased nutrient availability from glacial runoff, the results presented here indicate that changing sea ice and ocean conditions are also partly responsible for driving these local enhancements to Jones Sound primary productivity. Plausible explanations that could either partially or wholly drive the simulated increase in Jones Sound productivity include: (1) increased availability of light resulting from sea ice decline; (2) increased overturning of the mixed layer from enhanced wind stress as sea ice declines, resulting in greater nutrient upwelling; (3) increased temperature of sub-surface ocean waters can drive enhanced productivity since the carbon-specific photosynthesis rate in N-BLING is temperature-dependent (Galbraith et al., 2010; Noh et al., 2024). It is expected that a combination of these factors will drive enhanced productivity in our model, and this is evidenced in the light limitation and ocean temperature time series (figure 10e and figure A5, respectively). Specifically, we observe that the pattern of light limitation and ocean temperature through  $\sim$ 31.5 m depth mirrors that of the yearly-integrated chlorophyll mass. Beyond 2010, surface and subsurface light limitation and ocean temperature decrease and increase, respectively, driving enhanced productivity. In terms of nutrient limitation, we observe that iron limitation down to  $\sim$ 10.5 m depth increases in the summer and decreases in the winter beyond 2010, possibly denoting increased vertical advection of nutrient-rich waters in the winter that drive productivity blooms once light becomes available. In all, these results highlight that complex interplay between the atmosphere, ocean, and sea ice will likely continue to drive enhanced productivity in the future in Jones Sound under increasing polar-amplified global climate change.

## 4.2 Study limitations and uncertainties

The results presented here are subject to a high degree of uncertainty that stems from model limitations, the sparsity of input data used to drive and validate our model, as well as processes that are currently unaccounted for. As noted in the methods section, N-BLING crashed in May 2015. Upon investigating the cause of this, it was determined that the pH of a coastal grid

cell along the northern coast of Jones Sound reached infinity. In N-BLING, computing carbon chemistry requires carbonate alkalinity; however, we only model total alkalinity. As such, we must estimate carbonate alkalinity as the difference between total alkalinity and the contributions from borate, silicate, and phosphate. However, we do not model silicate and instead prescribe it. If simulated water properties in a grid cell become exceedingly fresh, the total alkalinity can become zero; however, 375 silicate can never reach zero because we are prescribing it, causing the model to reach a threshold where total alkalinity is equal to silicate alkalinity (causing carbonate alkalinity to become zero). Thus, when N-BLING solves for pH, which has carbonate alkalinity in the denominator, it divides by zero, causing the model to crash. Future modeling studies can avoid this issue by directly modeling silicate or improving the N-BLING code to detect when silicate should be manually set to zero.

Furthermore, we previously noted that the ANHA12 ocean model output that was used to derive our ocean and sea ice 380 model initial and boundary conditions features large warm biases in winter Atlantic Water (100-300 m), which then propagates throughout our model solution. We selected the ANHA12 model because it has sufficient resolution to resolve key circulation features in Baffin Bay and was also run through the time period of interest. However, this warm bias limits our model's ability to realistically simulate change in the thermal properties of mid-depth Atlantic Water, a key measure of the impact of global 385 climate change on Arctic waters. In addition, errors in the bathymetry we use in our model, especially near coastal outlet glaciers, will lead to erroneous paths of warm water circulation. We corrected for bathymetry that was too deep near Sverdrup Glacier and too shallow near Grise Fiord; however, it is likely that there are other locations in Jones Sound in which the ocean bathymetry is incorrect and we do not yet have bathymetric observations to apply corrections. For the biological productivity 390 model, recent studies have highlighted the importance of ocean-glacier interactions in driving near-glacier spatiotemporal patterns of productivity within Jones Sound. In particular, subglacial discharge plumes that originate beneath the nutricline can promote vertical advection of nutrients into the euphotic zone while nutrient-rich glacial runoff can feed the upper-ocean; both of these processes have been observed to drive coastal productivity blooms (Achterberg et al., 2018; Bhatia et al., 2021). While we do prescribe runoff as an atmospheric boundary condition in our ocean model, we assume it includes no nutrients and further do not resolve ocean-glacier interactions. As such, we do not capture the full extent to which atmospheric and oceanic 395 warming drives change in Jones Sound productivity between 2003-2016 and we flag this as an important next step in this work.

395 In addition, Jones Sound remains understudied from both a modeling and observational perspective, which limits the amount of publicly available data that can be used as model inputs and validation. On the observational side, it is only since 2019 that recurring observational campaigns have targeted Jones Sound, so repeat oceanographic measurements only exist beyond this date. As such, ocean models of Jones Sound prior to 2019 must be validated based on how well they represent the time-evolving circulation within nearby Baffin Bay and Lancaster Sound, where annual-repeat observations are available since the 400 early 2000's. This is not sufficient, as the modeling results presented here demonstrate that while circulation within Jones Sound is driven by inflow from Baffin Bay and Fram Sound, water masses undergo transformation within Jones Sound and circulation around the sound is sensitive to small-scale bathymetric features. Furthermore, to our knowledge, this is the first publicly available ocean model that was developed and validated with the purpose of studying ocean circulation within Jones Sound, meaning that there exists no other models of sufficient resolution by which we can compare modeling results. As Jones 405 Sound is an important passageway of transport between the Arctic and Atlantic Oceans and is critical in supporting local

communities, we emphasize that future observational campaigns (especially within the inflow regions of Lady Ann Strait and Fram Sound) and modeling studies of Jones Sound should be prioritized so that we can gain a better understanding of the long-term impact of global climate change on this region and improve the fidelity of numerical ocean models.

## 5 Conclusions


410 In this study, we modeled ocean circulation, sea ice dynamics, and biological productivity within Jones Sound, Canadian Arctic Archipelago, between 2003-2016 with a high-resolution regional configuration of the Massachusetts Institute of Technology general circulation model. Atmospheric forcing was taken from the Arctic System Reanalysis version 2 and ocean boundary conditions were derived from a North Atlantic configuration of the NEMO ocean model (Hu et al., 2019; Gillard et al., 2020). We find that volumetric transport through the three waterways that connect Jones Sound to the Arctic and Atlantic oceans  
415 is partitioned as 71%, 14%, and 15% via Lady Ann Strait, Fram Sound, and Glacier Strait, respectively. Surface circulation in the Summer within Jones Sound is dominated by eddies, whereas winter circulation is quiescent due to sea ice cover. The spatial distribution of summertime eddies varies considerably year-to-year and drives variability in the depth of the thermocline across the sound, impacting the spatial-reach of warm Atlantic Water that circulates at depth. This warm water, although topographically constrained, circulates counterclockwise around Jones Sound and expands its spatial reach in the winter when  
420 the thermocline shoals. Sea ice dynamics within Jones Sound are sensitive to small-scale circulation features that are generally not resolved within broad-scale CAA ocean models, such as tidal flushing through Fram Sound which triggers sea ice meltback in the Spring. In addition, we find that wintertime sea ice thickness decreases and the onset of sea ice refreeze in the fall is delayed due to oceanic and atmospheric warming that is simulated in our model. These changes have the impact of lengthening the time and spatial extent in which Jones Sound is sea ice free, thus leading to enhanced productivity at all ocean depth levels  
425 through 17.5 meters. While we note that the modeled warming signal in Baffin Bay and Jones Sound is overstated compared to observations, the results presented here improve our general understanding of circulation into, out of, and within Jones Sound as well as how it impacts sea ice and biological productivity dynamics. These results also emphasize the utility of high-resolution models in simulating complex waterways and underscore the need for sustained oceanographic observations in this region.

430 *Code and data availability.* All MITgcm parameter files, boundary conditions (including bathymetry), and initial conditions associated with the “high-resolution” Jones Sound ocean-sea ice-biological productivity model, as well as validation data shown in figures, have been archived in the Dryad Digital Repository (Pelle et al., 2024). These data are available during the peer review process at the following link: <http://datadryad.org/stash/share/DHOe1caYiqpIvQm6-fcBuD-r4vL5yoLFztLOvJ2mN3M>. MITgcm is also open source and is available for download from [mitgcm.org](http://mitgcm.org) (checkpoint66j). The sea ice and biological productivity (N-BLING) modules are built into MITgcm and are  
435 included in its download. Atmospheric forcing used in this study is publicly available via the Arctic System Reanalysis version 2 (ASRv2; Bromwich et al., 2018). Canadian sea ice charts covering the Eastern Arctic are publicly available through the National Snow and Ice Data Center (Service, 2009). Oceanographic profiles collected aboard the Amundsen Ice Breaker are publicly available through the Polar Data


Catalogue (Amu, 2003-2021). Lastly, due to the large amount of model output produced, output is available by request to the corresponding author.

#### 440 Appendix A: Additional Figures


##### A1 Additional ocean model figures




**Figure A1.** Vertical ocean temperature profiles ( $^{\circ}\text{C}$ ) of the Eastern model boundary (in Western Baffin Bay), taken from the ANHA12 NEMO ocean model in January of (a) 2002, (b) 2007, (c) 2012, and (d) 2016. Black shading denotes bedrock.




**Figure A2.** Same as in figure 3, but comparing modeled and observed vertical profiles of salinity (units on the Practical Salinity Scale; PSU).



**Figure A3.** Vertical profiles of ocean potential temperature in April of (a-c) 2004, (d-f) 2008, (g-i) 2012, and (j-l) 2016 taken through (a/d/g/j) Sverdrup Bay, (b/e/h/k) Grise Fiord, and (c/f/i/l) the oceanic region adjacent to Belcher Glacier (white, yellow, and red lines in figure 8a). Areas of black denote land and white contours correspond to ocean temperature levels that start at  $-2^{\circ}\text{C}$  and increase in  $0.5^{\circ}\text{C}$  increments.




**Figure A4.** Depth-time Hovmöller diagram of (a-b) ocean temperature ( $^{\circ}\text{C}$ ) and (c-d) salinity (units on the practical salinity scale; PSU) taken at (a/c) the center of Jones Sound (green triangle in figure 8a) and (b/d) the center of Lady Ann Strait (orange circle in figure 8a). In panels a/b, the white-black dashed line is the vertical-integrated ocean heat content ( $10^{17}$  J) and corresponds to the right y-axis. In panels c/d, the black dotted line is the 33.80 PSU salinity contour.




**Figure A5.** Ocean temperature time series ( $^{\circ}\text{C}$ ) of the first five ocean depth layers (0-35 m depth) taken in the center of Jones Sound. The black and green dotted lines mark the 2003-2010 mean maximum winter and summer ocean temperatures, respectively.


## A2 Impact of tides on sea ice thickness



**Figure A6.** Magnitude (b/d) and difference (a/c) in 50 m ocean velocity in (a-b) January 2015 and (c-d) April 2015 between model runs with and without tides (red shading denotes where tidal forcing results in faster ocean velocities). Black shading denotes bathymetry above 50 m and land is shaded in gray. Absolute velocities in panels b/d are taken from the model run with tides.



**Figure A7.** Difference in sea ice thickness between model runs with and without tides averaged at each month of the year in 2015 (blue shading denotes where sea ice is thinner in the tidal run).



**Figure A8.** Jones Sound integrated sea ice area (a-c) and volume (d-f) timeseries, where red and blue lines represent results from the tidal and non-tidal simulation, respectively. Panels b and e are zoomed into the 2014-2016 sea ice cycle in panels a and d, respectively. In panels c and f, the black line denotes the difference of the tidal and non-tidal time series.

*Author contributions.* TP, JSG, MS, DDB, and KS conceptualized the study with input from all co-authors. PGM and AH provided NEMO ocean model output, updated bathymetry datasets, and domain expertise. MM provided files needed to run N-BLING. TP and KS performed 445 all simulations. TP led manuscript preparation with input from all co-authors. JSG provided TP with supervision and funding.

*Competing interests.* The authors declare no competing interests.

*Acknowledgements.* This project is a result of the international collaborative Exploration of Saline Cryospheric Habitats with Europa Relevance (ESCHER) project with support from NASA grant 80NSSC20K1134. M R Mazloff acknowledges support from NASA Grant 80NSSC24K0243 and NSF Grants OPP-2149501 and OPP-1936222. Oceanographic profiles within the model domain that were presented 450 herein were collected by the Canadian research icebreaker CCGS Amundsen and made available by the Amundsen Science program, which is supported through Université Laval by the Canada Foundation for Innovation.

## References

Amundsen Science Data Collection (2005-2021). CTD-Rosette data collected by the CCGS Amundsen in the Canadian Arctic, Processed data. version 1., Arcticnet Inc., Québec, Canada, Canadian Cryospheric Information Network (CCIN), Waterloo, Canada., 455 <https://doi.org/10.5884/12713>, 2003-2021.

Achterberg, E. P., Steigenberger, S., Marsay, C. M., LeMoigne, F. A. C., Painter, S. C., Baker, A. R., Connelly, D. P., Moore, C. M., Tagliabue, A., and Tanhua, T.: Iron Biogeochemistry in the High Latitude North Atlantic Ocean, *Scientific Reports*, 8, 1283, <https://doi.org/10.1038/s41598-018-19472-1>, 2018.

Armitage, T. W. K., Manucharyan, G. E., Petty, A. A., Kwok, R., and Thompson, A. F.: Enhanced eddy activity in the Beaufort Gyre in 460 response to sea ice loss, *Nature Communications*, 11, 761, <https://doi.org/10.1038/s41467-020-14449-z>, 2020.

Ballinger, T. J., Moore, G. W. K., Garcia-Quintana, Y., Myers, P. G., Imrit, A. A., Topál, D., and Meier, W. N.: Abrupt Northern Baffin Bay Autumn Warming and Sea-Ice Loss Since the Turn of the Twenty-First Century, *Geophysical Research Letters*, 49, e2022GL101472, <https://doi.org/https://doi.org/10.1029/2022GL101472>, 2022.

Bhatia, M. P., Waterman, S., Burgess, D. O., Williams, P. L., Bundy, R. M., Mellett, T., Roberts, M., and Bertrand, E. M.: 465 Glaciers and Nutrients in the Canadian Arctic Archipelago Marine System, *Global Biogeochemical Cycles*, 35, e2021GB006976, <https://doi.org/https://doi.org/10.1029/2021GB006976>, 2021.

Bromwich, D. H., Wilson, A. B., Bai, L., Liu, Z., Barlage, M., Shih, C. F., Maldonado, S., Hines, K. M., Wang, S. H., Woollen, J., Kuo, B., Lin, H. C., Wee, T. K., Serreze, M. C., and Walsh, J. E.: The Arctic System Reanalysis, Version 2, *Bulletin of the American Meteorological Society*, 99, 805–828, <https://doi.org/https://doi.org/10.1175/BAMS-D-16-0215.1>, 2018.

Ding, Q., Schweiger, A., L'Heureux, M., Battisti, D. S., Po-Chedley, S., Johnson, N. C., Blanchard-Wrigglesworth, E., Harnos, K., Zhang, Q., Eastman, R., and Steig, E. J.: Influence of high-latitude atmospheric circulation changes on summertime Arctic sea ice, *Nature Climate Change*, 7, 289–295, <https://doi.org/10.1038/nclimate3241>, 2017.

Ding, Q., Schweiger, A., L'Heureux, M., Steig, E. J., Battisti, D. S., Johnson, N. C., Blanchard-Wrigglesworth, E., Po-Chedley, S., Zhang, Q., Harnos, K., Bushuk, M., Markle, B., and Baxter, I.: Fingerprints of internal drivers of Arctic sea ice loss in observations and model 475 simulations, *Nature Geoscience*, 12, 28–33, <https://doi.org/10.1038/s41561-018-0256-8>, 2019.

Frey, K. E., Comiso, J. C., Cooper, L. W., Garcia-Eidell, C., Grebmeier, J. M., and Stock, L. V.: Arctic Ocean Primary Productivity: The Response of Marine Algae to Climate Warming and Sea Ice Decline, *Noaa technical report oar arc* ; 22-08, National Oceanic and Atmospheric Administration, <https://doi.org/10.25923/0je1-te61>, 2022.

Galbraith, E. D., Gnanadesikan, A., Dunne, J. P., and Hiscock, M. R.: Regional impacts of iron-light colimitation in a global biogeochemical 480 model, *Biogeosciences*, 7, 1043–1064, <https://doi.org/10.5194/bg-7-1043-2010>, 2010.

Garcia, H. E., Weathers, K. W., Paver, C. R., Smolyar, I., Boyer, T. P., Locarnini, R. A., Zweng, M. M., Mishonov, A. V., Baranova, O. K., Seidov, D., and Reagan, J. R.: World Ocean Atlas 2018, Volume 4: Dissolved Inorganic Nutrients (phosphate, nitrate and nitrate+nitrite, silicate), Tech. Rep. 84, A. Mishonov Technical Ed.; NOAA Atlas NESDIS, 2018.

Gardner, A., Moholdt, G., Arendt, A., and Wouters, B.: Accelerated contributions of Canada's Baffin and Bylot Island glaciers to sea level 485 rise over the past half century, *The Cryosphere*, 6, 1103–1125, <https://doi.org/10.5194/tc-6-1103-2012>, 2012.

Gillard, L. C., Hu, X., Myers, P. G., Ribergaard, M. H., and Lee, C. M.: Drivers for Atlantic-origin waters abutting Greenland, *The Cryosphere*, 14, 2729–2753, <https://doi.org/10.5194/tc-14-2729-2020>, 2020.

Grivault, N., Hu, X., and Myers, P. G.: Impact of the Surface Stress on the Volume and Freshwater Transport Through the Canadian Arctic Archipelago From a High-Resolution Numerical Simulation, *Journal of Geophysical Research: Oceans*, 123, 9038–9060, 490 [https://doi.org/https://doi.org/10.1029/2018JC013984](https://doi.org/10.1029/2018JC013984), 2018.

Howard, S. and Padman, L.: Arc2kmTM: Arctic 2 kilometer Tide Model., Arctic Data Center, <https://doi.org/10.18739/A2PV6B79W>, 2021.

Hu, X., Myers, P. G., and Lu, Y.: Pacific Water Pathway in the Arctic Ocean and Beaufort Gyre in Two Simulations With Different Horizontal Resolutions, *Journal of Geophysical Research: Oceans*, 124, 6414–6432, [https://doi.org/https://doi.org/10.1029/2019JC015111](https://doi.org/10.1029/2019JC015111), 2019.

Kay, J. E., Holland, M. M., and Jahn, A.: Inter-annual to multi-decadal Arctic sea ice extent trends in a warming world, *Geophysical Research Letters*, 38, <https://doi.org/https://doi.org/10.1029/2011GL048008>, 2011.

Kinnard, C., Zdanowicz, C. M., Fisher, D. A., Isaksson, E., de Vernal, A., and Thompson, L. G.: Reconstructed changes in Arctic sea ice over the past 1,450 years, *Nature*, 479, 509–512, <https://doi.org/10.1038/nature10581>, 2011.

Kliem, N. and Greenberg, D. A.: Diagnostic simulations of the summer circulation in the Canadian arctic archipelago, *Atmosphere-Ocean*, 41, 273–289, <https://doi.org/10.3137/ao.410402>, 2003.

500 Losch, M., Menemenlis, D., Campin, J.-M., Heimbach, P., and Hill, C.: On the formulation of sea-ice models. Part 1: Effects of different solver implementations and parameterizations, *Ocean Model.*, 33, 129–144, <https://doi.org/10.1016/j.ocemod.2009.12.008>, 2010.

Mahowald, N. M., Baker, A. R., Bergametti, G., Brooks, N., Duce, R. A., Jickells, T. D., Kubilay, N., Prospero, J. M., and Tegen, I.: Atmospheric global dust cycle and iron inputs to the ocean, *Global Biogeochemical Cycles*, 19, <https://doi.org/https://doi.org/10.1029/2004GB002402>, 2005.

505 Marshall, J., Adcroft, A., Hill, C., Perelman, L., and Heisey, C.: A finite-volume, incompressible Navier Stokes model for studies of the ocean on parallel computers, *J. Geophys. Res.*, 102, 5753–5766, <https://doi.org/10.1029/96JC02775>, 1997.

McGeehan, T. and Maslowski, W.: Evaluation and control mechanisms of volume and freshwater export through the Canadian Arctic Archipelago in a high-resolution pan-Arctic ice-ocean model, *Journal of Geophysical Research: Oceans*, 117, <https://doi.org/https://doi.org/10.1029/2011JC007261>, 2012.

510 Melling, H., Agnew, T. A., Falkner, K. K., Greenberg, D. A., Lee, C. M., Münchow, A., Petrie, B., Prinsenberg, S. J., Samelson, R. M., and Woodgate, R. A.: Fresh-Water Fluxes via Pacific and Arctic Outflows Across the Canadian Polar Shelf, pp. 193–247, Springer Netherlands, Dordrecht, ISBN 978-1-4020-6774-7, [https://doi.org/10.1007/978-1-4020-6774-7\\_10](https://doi.org/10.1007/978-1-4020-6774-7_10), 2008.

Münchow, A.: Volume and Freshwater Flux Observations from Nares Strait to the West of Greenland at Daily Time Scales from 2003 to 2009, *Journal of Physical Oceanography*, 46, 141–157, <https://doi.org/https://doi.org/10.1175/JPO-D-15-0093.1>, 2016.

515 Munchow, A. and Humfrey, M.: Ocean current observations from Nares Strait to the west of Greenland: Interannual to tidal variability and forcing, *Journal of Marine Research*, 66, 2008.

Nakayama, Y., Menemenlis, D., Zhang, H., Schodlok, M., and Rignot, E.: Origin of Circumpolar Deep Water intruding onto the Amundsen and Bellingshausen Sea continental shelves, *Nat. Commun.*, 2018.

Noh, K.-M., Oh, J.-H., Lim, H.-G., Song, H., and Kug, J.-S.: Role of Atlantification in Enhanced Primary Productivity in the Barents Sea, 520 *Earth's Future*, 12, e2023EF003709, <https://doi.org/https://doi.org/10.1029/2023EF003709>, 2024.

Notz, D. and Stroeve, J.: Observed Arctic sea-ice loss directly follows anthropogenic CO<sub>2</sub> emission, *Science*, 354, 747–750, <https://doi.org/10.1126/science.aag2345>, 2016.

Olsen, A., Key, R. M., van Heuven, S., Lauvset, S. K., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T., Hoppe, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Pérez, F. F., and Suzuki, T.: The Global Ocean Data Analysis Project version 2 (GLODAPv2) – an

525 internally consistent data product for the world ocean, *Earth System Science Data*, 8, 297–323, <https://doi.org/10.5194/essd-8-297-2016>, 2016.

Pelle, T., Myers, P. G., Hamilton, A., Mazloff, M. R., Soderlund, K. M., Beem, L. H., Blankenship, D. D., Grima, C., Habbal, F. A., Skidmore, M., and Greenbaum, J. S.: Data From: Ocean circulation, sea ice, and productivity simulated in Jones Sound, Canadian Arctic Archipelago, between 2003–2016, Dryad, <https://doi.org/10.5061/dryad.1vhmgro0b>, 2024.

530 Peterson, I., Hamilton, J., Prinsenberg, S., and Pettipas, R.: Wind-forcing of volume transport through Lancaster Sound, *Journal of Geophysical Research: Oceans*, 117, <https://doi.org/https://doi.org/10.1029/2012JC008140>, 2012.

Polyakov, I. V., Bhatt, U. S., Walsh, J. E., Abrahamsen, E. P., Pnyushkov, A. V., and Wassmann, P. F.: Recent oceanic changes in the Arctic in the context of long-term observations, *Ecological Applications*, 23, 1745–1764, <https://doi.org/https://doi.org/10.1890/11-0902.1>, 2013.

535 Prinsenberg, S. J. and Bennett, E. B.: Mixing and transports in Barrow Strait, the central part of the Northwest passage, *Continental Shelf Research*, 7, 913–935, [https://doi.org/https://doi.org/10.1016/0278-4343\(87\)90006-9](https://doi.org/https://doi.org/10.1016/0278-4343(87)90006-9), 1987.

Prinsenberg, S. J. and Hamilton, J.: Monitoring the volume, freshwater and heat fluxes passing through Lancaster sound in the Canadian arctic archipelago, *Atmosphere-Ocean*, 43, 1–22, <https://doi.org/10.3137/ao.430101>, 2005.

540 Rotermund, L. M., Williams, W. J., Klymak, J. M., Wu, Y., Scharien, R. K., and Haas, C.: The Effect of Sea Ice on Tidal Propagation in the Kitikmeot Sea, Canadian Arctic Archipelago, *Journal of Geophysical Research: Oceans*, 126, e2020JC016786, <https://doi.org/https://doi.org/10.1029/2020JC016786>, 2021.

Service, C. I.: Canadian Ice Service Arctic Regional Sea Ice Charts in SIGRID-3 Format, Version 1. [Eastern Arctic], Tech. rep., <https://doi.org/10.7265/N51V5BW9>, 2009.

Shroyer, E. L., Samelson, R. M., Padman, L., and Münchow, A.: Modeled ocean circulation in Nares Strait and its dependence on landfast-ice cover, *Journal of Geophysical Research: Oceans*, 120, 7934–7959, <https://doi.org/https://doi.org/10.1002/2015JC011091>, 2015.

545 Stroeve, J. and Notz, D.: Changing state of Arctic sea ice across all seasons, *Environ. Res. Lett.*, 13, 103 001, <https://doi.org/10.1088/1748-9326/aade56>, 2018.

Tivy, A., Howell, S. E. L., Alt, B., McCourt, S., Chagnon, R., Crocker, G., Carrieres, T., and Yackel, J. J.: Trends and variability in summer sea ice cover in the Canadian Arctic based on the Canadian Ice Service Digital Archive, 1960–2008 and 1968–2008, *Journal of Geophysical Research: Oceans*, 116, <https://doi.org/https://doi.org/10.1029/2009JC005855>, 2011.

550 Tozer, B., Sandwell, D. T., Smith, W. H. F., Olson, C., Beale, J. R., and Wessel, P.: Global Bathymetry and Topography at 15 Arc Sec: SRTM15+, *Earth and Space Science*, 6, 1847–1864, <https://doi.org/https://doi.org/10.1029/2019EA000658>, 2019.

Verdy, A. and Mazloff, M. R.: A data assimilating model for estimating Southern Ocean biogeochemistry, *Journal of Geophysical Research: Oceans*, 122, 6968–6988, <https://doi.org/https://doi.org/10.1002/2016JC012650>, 2017.

555 Wang, Q., Wekerle, C., Wang, X., Danilov, S., Koldunov, N., Sein, D., Sidorenko, D., von Appen, W.-J., and Jung, T.: Intensification of the Atlantic Water Supply to the Arctic Ocean Through Fram Strait Induced by Arctic Sea Ice Decline, *Geophysical Research Letters*, 47, e2019GL086682, <https://doi.org/https://doi.org/10.1029/2019GL086682>, 2020.

Wang, Z., Hamilton, J., and Su, J.: Variations in freshwater pathways from the Arctic Ocean into the North Atlantic Ocean, *Progress in Oceanography*, 155, 54–73, <https://doi.org/https://doi.org/10.1016/j.pocean.2017.05.012>, 2017.

560 Wekerle, C., Wang, Q., Danilov, S., Jung, T., and Schröter, J.: The Canadian Arctic Archipelago throughflow in a multiresolution global model: Model assessment and the driving mechanism of interannual variability, *Journal of Geophysical Research: Oceans*, 118, 4525–4541, <https://doi.org/https://doi.org/10.1002/jgrc.20330>, 2013.

Zhang, Y., Chen, C., Beardsley, R. C., Gao, G., Lai, Z., Curry, B., Lee, C. M., Lin, H., Qi, J., and Xu, Q.: Studies of the Canadian Arctic Archipelago water transport and its relationship to basin-local forcings: Results from AO-FVCOM, *Journal of Geophysical Research: Oceans*, 121, 4392–4415, <https://doi.org/https://doi.org/10.1002/2016JC011634>, 2016.