

An ERA5 Climatology of Synoptic-Scale Negative Potential Vorticity-Jet Interactions over the Western North Atlantic

Alexander Lojko¹, Andrew C. Winters², Annika Oertel³, Christiane Jablonowski¹, and Ashley Payne^{1,4}

¹Department of Climate and Space Sciences and Engineering, University of Michigan, Ann Arbor, MI 48109, USA

²Department of Atmospheric and Oceanic Sciences, University of Colorado: Boulder, Boulder, CO 80309, USA

³Institute of Meteorology and Climate Research Troposphere Research (IMKTRO), Karlsruhe Institute of Technology, Karlsruhe, Germany

⁴Tomorrow.io, 9 Channel Center St, 7th Floor, Boston, MA 02210, USA

Correspondence: Alexander Lojko (alojko@umich.edu)

Abstract. Recent numerical modelling and theoretical work deduce that potential vorticity (PV) can turn negative in the Northern Hemisphere as a result of localized, convective heating embedded in vertical wind shear. It is further postulated that negative potential vorticity (NPV) may be relevant for the large-scale circulation as it has been observed to grow upscale into elongated, mesoscale bands when in close proximity to the jet stream, accelerating jet stream winds and degrading numerical weather prediction skill. However, these findings are largely confined to case studies. Here, we use a climatological and composite perspective to evaluate the occurrence of elongated bands of NPV over the North-West Atlantic and its implications for jet stream dynamics. This research focuses on synoptic-scale bands (>1650 km) of NPV that are in close proximity (<100 km) to the jet stream (termed NPV-jet interactions) using ERA5 data from January 2000 – December 2021. Climatological characteristics show that NPV-jet interactions occur most frequently over the coastal West Atlantic during Boreal Winter along 40°N. This latitude band has also seen an 11% increase (relative-change) in NPV-jet interactions over the 22-year time-period. Separating NPV-jet interactions into three distinct, large-scale flow patterns using K-means clustering conceptually illustrates the evolution of NPV features from their initial formation along the westward flank of the ridge to the eastern flank of the ridge. The large-scale environment of NPV-jet interactions is characterized by a trough-ridge couplet adjacent to positive integrated vapor transport (IVT) anomalies, conducive to warm-conveyor belts and mesoscale convective systems. Even when NPV is positioned in a more adiabatic environment (far away from regions of strong IVT anomalies), robust, positive PV gradient and wind speed anomalies exist along the jet stream. Inspecting three detailed case-studies that serve as archetypes to the three clusters, it is shown that the presence of NPV near the jet stream adiabatically enhances wave activity flux due to NPV mutually strengthening momentum transport and the ageostrophic flux of geopotential. The results show that the close proximity of synoptic-scale NPV to the jet stream is conducive to the occurrence of wind speed maxima, and could be dynamically relevant in enhancing downstream development, despite NPV's theorized origin from sub-mesoscales.

1 Introduction

The mid-latitude jet stream is a fast flowing current of westerly air in the upper-levels of the mid-latitude troposphere. Daily variations of the jet stream wind speeds are influenced by transient eddies (Lorenz and Hartmann, 2003; Barnes et al., 2010). These eddies can take on the form of cloud diabatic heating processes, which act to perturb the jet stream's large-scale circulation away from its base state (Woollings et al., 2016). Notably, the mechanisms involved in the interactions between cloud processes and the large-scale circulation remains a significant source of theoretical uncertainty (Bony et al., 2015; Ceppi et al., 2017; Baumgart and Riemer, 2019). This uncertainty is reflected in practical applications, particularly in numerical weather prediction, where the quality and reliability of weather forecasting models can be rapidly degraded by cloud microphysical processes interacting with the large-scale flow (Rodwell et al., 2013; Gray et al., 2014; Grams and Archambault, 2016; Grams et al., 2018; Spreitzer et al., 2019). Accordingly, it is crucial to further understand how cloud processes impact jet stream dynamics, both for theoretical advancement and practical applications.

Large-scale cloud systems (i.e., Warm Conveyor Belts, Mesoscale Convective Systems) that frequently develop equatorward of the jet stream are associated with vigorous vertical heating gradients that drive divergent outflow at the tropopause (Wernli and Davies, 1997; Baumgart and Riemer, 2019; Steinfeld and Pfahl, 2019). The irrotational wind field established by large-scale cloud systems causes the air arising from the outflow to move polewards and enhance wind speeds (Grams and Archambault, 2016; Steinfeld and Pfahl, 2019). This is a key mechanism by which forecast errors along the jet stream manifest (Baumgart and Riemer, 2019; Berman and Torn, 2019). Using a potential vorticity (PV) perspective, the warm and moist air that is brought to the tropopause in cloud systems associated with vigorous diabatic heating can be visualized as an enclosed area of low, but positive PV (in the Northern Hemisphere). Several studies have shown that diabatic processes along the equatorward side of the jet stream substantially contribute to strengthening the PV gradient (Bukemberger et al., 2023; Wernli and Davies, 1997). Sharpening of the PV gradient can serve to kinematically strengthen the jet stream, resulting in faster wind speeds, and the enhanced westerly propagation of Rossby waves (Harvey et al., 2016).

The PV perspective has also been applied to smaller scales to study mesoscale and convective scale weather systems (Hertenstein and Schubert, 1991; Braun and Houze Jr, 1996; Conzemius and Montgomery, 2009; Chagnon and Gray, 2009). At scales where the Rossby number is large, the PV distribution within a diabatic weather system is significantly influenced by horizontal heating gradients, which can lead to the generation of quasi-horizontal PV dipoles (Chagnon and Gray, 2009; Weijenborg et al., 2015). Their generation has been analogously compared to the tilting of horizontal vorticity onto the vertical axis, resulting in a cyclonic and anticyclonic relative vorticity pair (Davies-Jones, 1984; Müller et al., 2020). Composite studies show that quasi-horizontal PV dipoles are coherent features that form around convective updrafts in a vertically sheared environment (Weijenborg et al., 2017; Oertel et al., 2020). Notably, the diabatically reduced PV pole can turn negative (in the Northern hemisphere). PV impermeability theory states that vertical heating gradients cannot lead to the generation of NPV (Haynes and McIntyre, 1990). Hence, NPV can only arise from localized, horizontal heating gradients embedded in a vertically sheared

55 environment (Harvey et al., 2020). Oertel et al. (2021) illustrate that a strong upper-level jet and wind shear are key ingredients for elongating convectively generated NPV onto larger scales. When a collection of convective-scale NPV features are present, they will preferentially coalesce together (Gray, 1999). Horizontal shear near the jet stream acts to stretch NPV onto the mesoscales but dilutes the elongated NPV towards a near-zero but still negative value (Oertel et al., 2020; Prince and Evans, 2022).

60

Elongated NPV features have been observed in observations (Harvey et al., 2020) and in a number of non-hydrostatic numerical modelling simulations (Braun and Houze Jr, 1996; Rowe and Hitchman, 2016; Oertel et al., 2020). NPV is unique from other large-scale regions of diabatically reduced PV such as large-scale negative PV anomalies (Hoskins, 1997) as NPV has been linked to the occurrence of frontal rainbands (Bennetts and Hoskins, 1979; Schultz et al., 2000), sting jets (Volonté et al., 2018) and enhanced stratosphere-troposphere exchange (Rowe and Hitchman, 2015). Additionally, since NPV has a negative PVU value, this implies that NPV is associated with hydrostatic, inertial or symmetric instability (Schultz et al., 2000). Oertel et al. (2020) theorized that since mesoscale bands of NPV exhibit negative absolute vorticity (relative vorticity magnitude exceeding the Coriolis parameter) and a temporal persistence of several hours, elongated bands of NPV may likely be analogous to an inertial instability.

70

When elongated NPV is embedded in large-scale negative PV anomalies such as ridges, the anticyclonic relative vorticity associated with NPV appears to be an order of magnitude larger than the large-scale negative PV anomaly it is embedded within (Rowe and Hitchman, 2016; Lojko et al., 2022). The vigorous anticyclonic relative vorticity associated with NPV features is noted to be an important contributor to the enhancement of momentum transport along the jet stream (Rowe and Hitchman, 2016) and may be linked to the occurrence of wind speed maximum (Harvey et al., 2020; Oertel et al., 2020). Misrepresenting anticyclonic vorticity associated with mesoscale NPV in global numerical weather models can lead to the rapid introduction of non-divergent wind errors along the jet stream (Lojko et al., 2022). Additionally, the strong horizontal shear associated with NPV can trigger the occurrence of clear-air-turbulence along the equatorward side of the jet stream (Trier and Sharman, 2016; Thompson and Schultz, 2021).

80

While the interaction of elongated bands of NPV with the jet stream can be dynamically relevant and have implications for aviation and weather prediction, the aforementioned studies examining NPV interactions with the jet stream employ case-study perspectives. These cases provide detailed mechanistic insights into NPV dynamics. However, a climatological and composite analysis of NPV and its interactions with the jet stream is lacking. Such an analysis will provide insight into common mechanisms and synoptic situations associated with elongated bands of NPV. The obtained results will serve to climatologically identify regions that may be relevant to forecast errors and turbulence associated with NPV features. Two key questions are proposed via the climatology and composite analysis:

- What are the climatological characteristics of elongated bands of NPV when they interact with the jet stream?

- What are the typical circulation patterns and dynamical mechanisms involved when NPV interacts with the jet stream?

90 The proposed analysis focuses on the Western North Atlantic, a region with numerous well-documented case studies evaluating NPV interactions with the jet stream (Rowe and Hitchman, 2016; Harvey et al., 2020; Lojko et al., 2022). These studies provide a basis for comparison against the proposed composite work. The region of Eastern North America and the West Atlantic is frequently associated with convective storms (Li et al., 2020) and warm conveyor belts (Madonna et al., 2014) which develop in close proximity to the jet stream, and are candidate weather events for the generation of elongated NPV features
95 (Clarke et al., 2019; Oertel et al., 2020). Another rationale for focusing on this specific region is that previous composite studies have identified the Western North Atlantic as a mid-latitude climatological hotspot for regions of inertial instability along the tropopause (Thompson et al., 2018).

The paper is structured such that the data and methodology is presented in Section 2. The climatological characteristics of
100 NPV are presented in Section 3.1. The large-scale circulation patterns during NPV-jet interactions are evaluated in Section 3.2. Three detailed case-studies of synoptic-scale NPV that focus on the dynamics involved in NPV-jet interactions are presented in Section 3.3. A discussion of synoptic-scale NPV is had in Section 4, with particular focus on its relevance for the large-scale atmospheric circulation. The work is concluded in Section 5.

105 2 Data & Methods

2.1 Data

The ECMWF reanalysis version 5 (ERA5; Hersbach et al., 2020) is downloaded at a grid spacing of 0.25° (~ 31 km) for the years of 2000 – 2021. The fine grid spacing and temporal resolution of ERA5 compared to other global reanalysis datasets provides an opportunity to study elongated NPV features in a climatological framework. PV, geopotential height (Z), horizontal winds and integrated vapor transport (IVT) with 6-hourly resolution are downloaded directly from the ERA5 archive.
110 Data is predominantly obtained at 250 hPa to focus on how NPV interacts with the mid-latitude tropopause. While it is more common for PV analysis to be performed on isentropic levels, the isentropic level associated with the tropopause can vary notably with respect to season (Röthlisberger et al., 2018). Using only one particular isentropic level can miss NPV features depending on the season examined. Hence, for simplicity, a single isobaric level that is representative of the tropopause during
115 all seasons is selected as isobaric surfaces near the tropopause tend to vary less with height across seasons. It is also noted that elongated bands of NPV are maximized in frequency at specific isobaric levels near the tropopause (Fig. A1). Prior to any further climatological analysis, the latitude and longitude data are filtered to only include the Western North Atlantic (25°N , 100°W)–(65°N , 50°W). An additional 10° buffer is kept on each side of the domain to minimize any boundary effects when identifying NPV features.

120

2.2 Methods

2.2.1 NPV-Jet Interaction Algorithm

An algorithm is designed to search for time-steps where elongated NPV features are in close proximity to the jet stream, specifically < 100 km to the jet stream. Henceforth, these time-steps are termed NPV-jet interactions. The algorithm is split
125 into three parts. First, the identification of elongated NPV features. Second, the identification of jet stream features and, last, identification of instances when elongated NPV is within close proximity to the jet stream. A schematic is provided below (Fig. 1) to give a general overview of the methodology. PV data is bilinearly interpolated to 0.5° to improve computational efficiency. This interpolation has no impact on the final results relating to the frequency of NPV-jet interactions.

130 The first step in the algorithm is to identify and label all connected regions of NPV, specifically where PV is ≤ -0.01 PVU. The major axis length-scale is then calculated for each label by using the two latitude, longitude coordinate pairs that are located furthest away from each other within the NPV label. Only the longest 2% of features are kept. This threshold corresponds to NPV length-scales longer than 1650 km (Fig. 2a). These features are henceforth referred to as a synoptic-scale NPV features. As a sanity check, the 97th and 99th percentile are also tested; however, these thresholds do not impact conclusions made re-
135 garding NPV-jet interaction frequency. The final step involves obtaining and saving the coordinates of the synoptic-scale NPV features.

For jet stream identification, contours where the PV field has a value of 2 PVU are identified (Barnes et al., 2010). Prior to jet stream identification, time-steps in which no synoptic-scale NPV features were identified are filtered out. Next, the PV field
140 is smoothed by a 10 point Gaussian smoother to improve contour identification by smoothing out mesoscale PV filaments. Further filtering of contours is applied by only keeping circumpolar contours. Circumpolar contours are defined as continuous contours of 2 PVU that extend across the longitudinal extent of the NPV-jet interaction domain. Non-continuous contours are likely associated with cut-off features (Hoskins, 1997) and are chosen to be removed. A small subset of times (4 time-steps) were not able to identify continuous 2 PVU lines. These times were also filtered out from the analysis. The remaining 2 PVU
145 contour coordinates are saved for evaluation with respect to the identified synoptic-scale NPV features.

In the final step of the algorithm, we use the coordinates from the synoptic-scale NPV feature and 2 PVU contour to find the minimum distance between the two features using the Haversine formula. If an NPV feature is within 100 km to the jet stream, the event is retained. If the minimum distance between NPV and 2 PVU exceeds 100 km for a particular event, the event is
150 filtered out. An interaction point is also defined, which is the coordinate of the 2 PVU contour that is in closest proximity to the synoptic-scale NPV feature. The interaction point is used to perform the centered composites described in the latter section of the methods. Sometimes, multiple synoptic-scale NPV features are detected within the 100 km threshold for a particular time-step. In these instances, all NPV features are retained for the climatological analysis. It is also worth noting that no temporal filtering is applied in this study. Hence, instances of consecutive time-steps with NPV-jet interactions are retained as we are not




Figure 1. Schematic overview of the NPV - Jet Interaction Identification Procedure outlining three key steps in the algorithm design. The images at the bottom of the schematic denote how the algorithm works on a single case. Grey shading shows individual NPV labels. Dark dashed contours denote the perimeter of synoptic-scale NPV features. Dark, continuous contours illustrate the 2 PVU contour (jet stream). The orange circle denotes the 'interaction-point', which is defined as the 2 PVU contour coordinate that is in closest proximity to the synoptic-scale NPV feature. The grey dashed box shows the domain over which NPV-jet interactions are searched for. Identified synoptic-scale NPV features can lie outside of the domain as long as the interaction point itself lies within the domain.

155 focused on evaluating the life-time of the NPV features in this study.

To provide analogue cases for NPV-jet interactions, the algorithm is also modified to search for synoptic-scale NPV features within 100 - 300 km to the 2 PVU contour. These events are specifically referred to as NPV-jet (100-300 km) interactions. Note that when referring to synoptic-scale NPV features that are within 100 km to the jet stream, these instances are always referred to as NPV-jet interactions. However, when simultaneously comparing both of these interaction distance thresholds, we will use the terminology: NPV-jet (< 100 km) and NPV-jet (100-300km) interactions to prevent confusion.

In total, 21341 synoptic-scale NPV features are detected within the domain during the 22-year time-period. From those, 4983 (23%) synoptic-scale NPV features are detected within 100 km of the 2 PVU contour. Some statistics of the NPV features are detailed in Fig. 2. Figure 2a shows a histogram of the major axis length of all NPV features detected. The vast majority of identified NPV features in ERA5 have rather small length-scales. Over 80% (more than 10^6) of identified NPV features in ERA5 are characterized by a major axis length shorter than 200 km. We also note that the synoptic-scale NPV features retained in this study are of much larger length-scale than the mesoscale filaments observed in high-resolution numerical modelling studies (Oertel et al., 2020; Blanchard et al., 2021). Deep convection resolving simulations result in a much noisier PV field compared to the smoother and coarser ERA5 dataset.

For additional reference, the area sizes of all NPV features (no matter their size) are plotted in Fig. 2b. The area sizes of NPV features largely follow the distribution of their major axis length scale (Fig. 2a). The majority (over 90%) of NPV features detected in ERA5 are smaller than $1 \times 10^5 \text{ km}^2$ in area size. Figure 2c shows that the frequency of synoptic-scale NPV features decreases quasi-exponentially with distance away from the jet stream. Most synoptic-scale NPV features are identified in very close proximity to the jet stream (approximately 25% of synoptic-scale NPV features are detected within 100 km to the jet stream). Of the 32144 time-steps that comprise the period of study, 14580 have a synoptic-scale NPV feature within the domain (45% of the time). 3845 time-steps have a synoptic-scale NPV feature that is within 100 km to the jet stream (12% of the time-steps detect an NPV-jet interaction). For reference, 3835 time-steps have a synoptic-scale NPV feature that is within 100 - 300 km to the jet stream, (approximately 25% of synoptic-scale NPV features are detected between 100-300 km to the jet stream).

2.2.2 Composite Approach

The purpose of the centered composite approach is to identify the typical circulation patterns and kinematic processes that occur when synoptic-scale NPV interacts with the jet stream. When computing the centered composite, the mean interaction point is computed from all events. Subsequently, all NPV-jet interaction events are shifted towards the mean using a latitude weighting following Winters (2021). Only unique time-steps are included in the centered composite approach. Hence, in the occasional circumstance that more than one NPV-jet interaction occurs at the same time, only the NPV feature that is in closest




Figure 2. Statistics of the identified NPV objects. (a) The total number of NPV objects detected in ERA5 on the y-axis (logged) and the major axis length scale (km) of each NPV object binned into intervals of 200 km. The red dashed lines show where the 98th percentile lies for the major axis length scale. (b) is the same as (a) but with the area size ($\times 10^6$ km^2) shown on the x-axis. Binning intervals are set to 0.1×10^6 km^2 . (c) shows the distance between each synoptic-scale NPV feature identified and its closest proximity to the jet stream. The red dashed line is used to show where the 100 km threshold lies for NPV-jet interactions (horizontal line) and how many NPV-jet interactions are identified within the 100 km threshold (vertical line).

proximity to the jet stream is used for centering on the selected time-step.

190

A problem that arises when computing a mean composite is that important circulation pattern features become smoothed. For example, when computing a principal component analysis on the PV field of all retained events, the dominant modes of variability are characterized by ridging environments (not shown). In contrast, computing a single mean composite of all events illustrates zonal flow. This is due to the synoptic-scale NPV features occurring in different locations along the ridge, hence a single composite centered on the location of NPV-jet interaction leads to too much smoothing of the attendant ridge. To mitigate this effect, K-means clustering is applied to separate events into distinct groups of circulation patterns. The K-means clustering is informed by the latitude weighted PV field. The size of the domain used in the clustering algorithm was a $10^\circ \times 10^\circ$ box centered on the interaction coordinate. Enlarging the domain ($15^\circ \times 15^\circ$ and $20^\circ \times 20^\circ$) had no discernible impact on the K-means clustering results. We also tested the robustness of the clusters by using other fields to organize NPV-jet interaction time-steps such as the use of a binary PV field (1 for stratosphere, 0 for troposphere) in which we obtained similar clustering results.

200

The number of clusters selected is informed both objectively and subjectively. The Silhouette score is used as an objective metric to determine the optimal number of clusters to use in this study. The Silhouette score metric evaluates how closely grouped together events are to each cluster centroid (Grazzini et al., 2020). The metric ranges from a score of -1 (poor clustering) to 1 (fully separated clusters). A value of 0 indicates that events tend to equally resemble other cluster centroids (i.e., equal distance to more than one cluster). The Silhouette score is computed for 2-8 clusters. Two clusters provides the highest

205

Silhouette score (0.24). However, it was noted that the interaction point was located in the same region (along the western flank of the ridge) for both of the clusters, so there was not a lot of variability in the location of NPV-jet interactions. Three, four and five clusters provided the next best Silhouette scores (0.18, 0.16 and 0.17 respectively). While the use of more clusters reduced the Silhouette score (the large-scale circulation patterns became more similar between each cluster), there was more variance in the location of the NPV-jet interaction coordinate within the broader large-scale ridging environment. Increasing the number of clusters to three led to an interaction point being located along the eastern flank of the ridge in one of the clusters. Increasing the number of clusters to four and five did not improve the variability of the interaction point location, hence three clusters was subjectively deemed appropriate for the analysis.

As an analogue comparison against NPV-jet (< 100 km) interaction time-steps, NPV-jet (100-300 km) interactions are also evaluated through the centered composite and clustering framework. However, a modification is made to the clustering approach. Each time-step associated with an NPV-jet (100-300 km) interaction is grouped to one of the three previously identified clusters. These time-steps are grouped separately from the NPV-jet (< 100 km) interaction time-steps (i.e., two separate groups of three clusters). NPV-jet (100-300 km) interaction time-steps are assigned to one of the three clusters depending on the similarity of the PV field during NPV-jet (100-300 km) interactions to the mean of the PV field of one of the clusters during NPV-jet (< 100 km) interactions (i.e., how similar is the field of a particular time-step to the cluster centroid?). Similarity is measured using a simple Euclidean distance metric as in Pohorsky et al. (2019). This provides clusters of similar synoptic-scale conditions (i.e., flow analogues) that differ with respect to the distance of elongated NPV to the jet.

2.2.3 Kinematic Analysis of NPV

Several methods are used to interpret the impact of NPV on the large-scale flow. Amplification of the jet stream by different components of the wind is examined by calculating PV advection by the irrotational and non-divergent wind (Archambault et al., 2013) at 250 hPa. The partitioning of the winds into these two components is completed via Helmholtz partitioning using the Python Package *Windspharm* (Dawson, 2016), which utilizes spherical harmonics on the global domain. The PV advection fields are computed for each event prior to applying the composite approach.

The wave activity flux (WAF; Takaya and Nakamura, 2001) is computed for each NPV-jet time-step to explain the relevance of how NPV enhances kinetic energy along the jet stream. The horizontal form of WAF can quantify the propagation and energy transport associated with horizontally propagating atmospheric waves along a single level of the atmosphere. WAF assumes quasi-geostrophy and is often used in the evaluation of large-scale flow patterns, particularly in the study of atmospheric Rossby waves and their downstream propagation. The WAF equation is shown below:

$$\vec{W} = \frac{1}{2|\vec{U}|} \begin{pmatrix} U[(\psi'_x)^2 - \psi'_x \psi'_{xx}] + V[\psi'_x \psi'_y - \psi'_x \psi'_{xy}] \\ U[\psi'_x \psi'_y - \psi'_x \psi'_{xy}] + V[(\psi'_y)^2 - \psi'_y \psi'_{yy}] \end{pmatrix} \quad (1)$$

U and V denote the base state wind in the zonal and meridional direction, where the base state is determined from the seasonal climatology: Boreal Winter, Spring, Summer and Autumn (DJF, MAM, JJA, SON). $|\vec{U}|$, which is the combination of the $|(U, V)|$ components, is the magnitude of the base state wind. ψ' is the streamfunction anomaly. ψ' is computed from ERA5 wind data using the Windspharm package. The x and y derivatives relate to latitude and longitude. xx , xy and yy denote second derivative terms. Derivatives are computed using spherical harmonics.

As mentioned above, 2D WAF is a quasi-geostrophic and dry kinematic metric. While the generation of NPV implies diabatic activity, once NPV grows to mesoscales, it has been observed to persist quasi-adiabatically (Oertel et al., 2020; Lojko et al., 2022). Hence, a dry kinematic approach employed in the WAF equation is assumed to be an appropriate framework to study how synoptic-scale NPV interacts with the jet stream. Individual components of the WAF equation can be assessed to understand the dry, dynamical mechanism by which NPV influences the magnitude of 2D WAF. Note that the phase velocity terms of the WAF equation are not used (Takaya and Nakamura, 2001). This is because only instantaneous time-steps are evaluated, hence Equation 1 assumes the analysis of standing waves.

The first terms inside the square bracket refer to momentum transport, providing information on Rossby wave propagation by geostrophic motion. This term represents the square of the non-divergent wind terms. The second terms in the square brackets refer to the ageostrophic flux of geopotential (Takaya and Nakamura, 2001), and serves as the source or sink for wave activity (Orlanski and Katzfey, 1991). ψ'_{xx} and ψ'_{yy} are of particular interest when focusing on the ageostrophic flux of geopotential as these terms can be easily related to the NPV feature. ψ'_{xx} and ψ'_{yy} represent wind shear anomalies that relate to the v_x and u_y terms of the relative vorticity equation. NPV is uniquely associated with anticyclonic vorticity maxima (Rowe and Hitchman, 2016; Lojko et al., 2022), hence it is expected that these terms will be magnified in regions of NPV. A partitioning of the WAF equation is performed in Section 3.3 to mechanistically illustrate how synoptic-scale NPV contributes to WAF.

One additional technique that is used involves relative vorticity inversion (Oertel and Schemm, 2021) to illustrate the circulation pattern associated with a spatially confined relative vorticity field. NPV can not be inverted via PV inversion as it is dynamically unstable (Davis and Emanuel, 1991; Davis et al., 1993; Oertel and Schemm, 2021). In contrast, relative vorticity inversion is independent of the sign of the relative vorticity field. The relative vorticity is equal to the Laplacian of the streamfunction on a horizontal surface. Solving the Poisson equation, and using the streamfunction's relation to vorticity, the non-divergent winds can be obtained along a horizontal, 2D surface (i.e., at 250 hPa) (Oertel and Schemm, 2021). This is particularly useful in examining synoptic-scale NPV, as it appears to largely be a shallow-layer feature that resides within the upper troposphere (Fig. A1).

3 Results

3.1 Climatology of NPV over the West Atlantic

The climatological frequency of NPV features are presented for the North American-West Atlantic region in Fig. 3. Figure 3a shows the frequency of all NPV identified using ERA5 regardless of size or distance to the jet stream. A meridional gradient of NPV frequency is observed with a maximum in the sub-tropics ($>12\%$) that decreases towards higher latitudes, where the frequency drops below 4% northward of 50°N . The higher frequency of NPV at lower latitudes is also consistent with maxima in inertial instability frequency (Thompson et al., 2018) where anticyclonic relative vorticity is of greater magnitude than the Coriolis parameter.

Other spatial patterns, such as effects of topography, are also captured in the NPV frequency distribution. A low percentage of NPV frequency downstream of the Rocky Mountains (100°W) sharply transitions into a region of higher NPV frequency farther east until reaching a maximum over the coastal Western Atlantic (NPV frequency $>9\%$). The comparatively higher frequency of NPV over the coastal West Atlantic and Eastern North America aligns with regions of strong latent heating attributed to mesoscale convective systems (Liu et al., 2021), and warm Gulf Stream waters that can drive vigorous diabatic weather systems (Minobe et al., 2008). This region also coincides with climatological hotspots of warm conveyor belts (Madonna et al., 2014)

The spatial distribution and percentage frequency of NPV changes when focusing on synoptic-scale NPV features that meet the NPV-jet interaction criteria (Fig. 3b). NPV-jet interaction features are most frequent about the mid-latitudes, with a maximum ($>1.2\%$) located over the coastal Western Atlantic near 41°N , 65°W . Generally, NPV-jet interaction features are frequent across the coastal Western Atlantic along a latitude band of 40°N . The location of maximum NPV-jet interaction features coincides with warm-conveyor belt ascent climatologies, notably 24 hours after ascent has begun (Madonna et al., 2014; Joos et al., 2023). For comparison, the NPV-jet algorithm is modified to only search for synoptic-scale NPV features within 100 - 300 km of the jet stream (Fig. 3c). The frequency of synoptic-scale NPV features during NPV-jet (100-300 km) interactions still resembles the pattern in Fig. 3b, with NPV-jet interactions maximized over the West Atlantic. However, the general frequency pattern is shifted south and the maxima frequency is located about 5° south of the maxima in Fig. 3b.

To initially motivate the link between NPV-jet interaction events and jet stream dynamics, the frequency of jet streaks is shown in Fig. 3. In this figure, jet streaks are defined as wind speed anomalies in excess of 40 m s^{-1} . Figure 3d shows the frequency of jet streaks when NPV-jet interactions are not occurring. Jet streak frequency is maximized east of 50°W , with maximum frequency values reaching 2% at particular grid-points. Figure 3e illustrates that when NPV-jet interactions are present, jet streaks become much more frequent over the West Atlantic, with maxima values reaching 6%. In other words, jet streaks become 5 times more likely to occur over the Coastal West Atlantic compared to time-steps with no NPV-jet interactions detected. The maximum frequency of jet streaks is located slightly polewards (about 5°) of the maximum frequency of




Figure 3. Climatological frequency of NPV between 2000-2021 is shown in (a) and (b). (a) shows the percentage of time NPV is observed at a grid-point irrespective of its size. (b) shows the frequency of synoptic-scale NPV features during NPV-jet interactions. (c) shows the frequency of synoptic-scale NPV features during NPV-jet (100-300 km) interactions. The frequency of wind speed anomalies that exceed 40 m s^{-1} , termed jet streaks (where anomalies are computed with respect to season) is shown in (d-f). (d) shows the frequency of jet-streaks during time-steps that do not have an NPV-jet interaction over the West Atlantic domain. A random sample of cases are selected such that the number of cases match the number and seasonal distribution of NPV-jet interaction cases. Bootstrapping is then performed 100 times with replacement and a mean is computed. (e) shows the frequency of jet streaks during NPV-jet interactions used to form the climatology in (b). (f) shows the frequency of jet streaks during NPV-jet (100-300 km) interactions used to perform the climatology in (c). Note that time-steps detected in (d), (e) and (f) do not overlap.

305 synoptic-scale NPV features (Fig. 3b). Figure 3f shows jet streak frequency by using the 100 - 300 km threshold for NPV-jet interactions. The overall frequency of jet streaks is displaced equatorwards, with the region of maxima being displaced by approximately 2° . The overall frequency of jet streaks also slightly decreases, with maxima values reaching 4.2%. A two sided student t-test is also performed (not shown) at the 2% significance level where it is found that the reduced frequency and equatorward displacement of jet streaks in NPV-jet (100 - 300 km) compared to NPV-jet (<100 km) exhibits statistical significance
310 over much of the West Atlantic. More on the implication of these results is shown in Section 3.2.

NPV-jet interaction frequencies exhibit a pronounced seasonal cycle. DJF is associated with the most frequent NPV-jet interactions (Fig. 4a) with NPV-jet interaction frequencies exceeding 2.5% over the Western Atlantic. MAM NPV-jet interaction frequency (Fig. 4b) maxima reduce to 1.25% and shift westward towards the Eastern US coastline. JJA frequency maxima
315 (Fig. 4c) of 0.5% are predominantly located over continental North America. SON (Fig. 4d) NPV-jet interaction frequency maxima exceeding 1.5% are predominantly located over the Western Atlantic. The spatial location of the JJA frequency pattern suggests that JJA NPV is predominantly induced by continental convection. In contrast, the location of wintertime NPV features over the West Atlantic is consistent with the aforementioned warm conveyor belt climatology. It is also noted that the spatial seasonal variations of NPV-jet interaction maxima agree with the locations of air masses that contribute to downstream
320 blocking (Pfahl et al., 2015; Steinfeld and Pfahl, 2019), which are postulated to arise from rapidly ascending air-streams in regions of strong latent heating. The seasonal co-location of NPV with the Steinfeld and Pfahl (2019) climatology is not necessarily surprising, as NPV is a byproduct of latent heating within convective weather systems (Harvey et al., 2020; Oertel et al., 2020).

325 Additional detail on the seasonal and spatial climatology of all NPV and NPV-jet interactions is provided in Fig. 4e-f. For all NPV features in the domain (Fig. 4e), the total monthly area coverage of NPV demonstrates a seasonal cycle. NPV area coverage is maximized in October and minimized in February. JJA has higher NPV area coverage in the domain compared to DJF, although, the interannual variability in NPV coverage during DJF is larger and can exceed JJA area coverage for particular years (not shown). It is also noted that the total area coverage for all NPV and NPV-jet interactions is sensitive to the location
330 of the domain. When experimenting with shifting the domain westward, the summer (winter) month frequencies increase (decrease). This relationship reverses when moving the domain eastward.

In Fig. 4f, the seasonal cycle for the area coverage of NPV-jet interactions is shown. In contrast to the area coverage of all NPV occurrences, the area coverage of NPV is now maximized during DJF alongside maxima in interannual variability. JJA
335 has a minimum in area coverage, and an order of magnitude lower area coverage and interannual variability in area coverage relative to DJF. The rapid decrease in NPV-jet interaction area from MAM to JJA is consistent with the climatological, rapid decrease of jet stream wind speeds over the US (Iqbal et al., 2018). The much lower frequency of JJA NPV-jet interaction area suggests synoptic-scale NPV is much less frequent during summer months, with the absence of strong jet stream winds




Figure 4. As in Fig. 3b but showing the seasonal frequency of NPV-jet interactions between 2000-2021 (a) for DJF, (b) MAM, (c) JJA, and (d) SON. (e) and (f) show boxplots of the monthly area covered by all NPV and NPV-jet interactions, respectively. Area coverage is computed within the dashed box domain shown in (a) (25°N, 100°W)–(65°N, 50°W). The whiskers of the box plot denote the upper and lower extrema. The top and bottom of the box denote the upper and lower quartile. The horizontal lines within the box illustrate the median area coverage of NPV.




Figure 5. Interannual trend of NPV frequency from 2000-2021 expressed as a relative percentage change. (a) shows the total change in NPV frequency per year at each grid-point. (b) shows the same as (a) but for NPV-jet interactions. The false-discovery rate (FDR) is applied to provide a more conservative estimate of statistical significance regarding observed trends (Wilks, 2016). Grid-points that exhibit statistical significance at the 2% level ($\alpha = 0.02$) are shaded. Grid-points where the p value is small enough to satisfy the FDR criterion ($\alpha FDR = 0.1$) are stippled. (c) and (d) shows the total area covered by all NPV (c) and NPV-jet interactions (d) per year computed within the domain (25°N, 100°W)–(65°N, 50°W).

potentially limiting the upscale growth of NPV (Oertel et al., 2021).

340

Linear trend analysis for NPV frequency from 2000 to 2021 shows an increase in NPV frequency (Fig. 5a), whereby much of the increase is attributed to a narrow latitude band between 35°N-45°N, coincident with the maximum in NPV-jet interaction frequency (Fig. 3b). An increasing, relative trend of 1% per year extends from 100°W-50°W, and farther downstream into the Atlantic. Localized maxima of 3% per year in the linear trend can be observed over the eastern US and coastal Western Atlantic, with an additional increasing trend area that encompasses the Gulf of Mexico and southern US (Over a 22-year time-period, this equates to a relative increase of NPV by 66% in some localized regions).

345

The relative trend in NPV-jet interactions is weaker but also positive (Fig. 5b), with maximum values exceeding 0.5% per year (a relative increase of 11% at these localized locations over the 22-year time-period). Interestingly, these values coincide with maxima in the trend for all NPV events, predominantly over the eastern US and coastal Western Atlantic. The increasing

350

trend is also generally contained within a narrow latitude band between 35°N-45°N, suggesting that the overall trend is driven, in part, by synoptic-scale NPV features that develop adjacent to, or are advected and interact with, the jet stream. Unlike Fig. 5a, however, the observed increasing trend over the 22-year time-period does not satisfy the false discovery rate.

355 Addressing the interannual area coverage of all NPV events in the domain (Fig. 5c), much of the increasing NPV frequency occurs after 2010. Prior to 2010, interannual variability of NPV area coverage was considerably less compared to post 2010. In contrast, the interannual variability is much more pronounced for NPV-jet interactions across the full period of study (Fig. 5b). 2010, 2016 and 2019 denote years with maxima in the area coverage of NPV-jet interactions within the domain. Since just 22-years of data are used, the observed trends could be predominantly influenced by decadal variability rather than longer term
360 climatic trends. However, recent work published by Lee et al. (2023) indicates that NPV at 250 hPa during the winter months over the Western Atlantic has experienced a statistically significant (using a Student's t-test) increasing trend over a 40-year time-period starting from 1979. Additionally, Prosser et al. (2023) use a variety of metrics related to turbulence (including NPV) over the same 40-year time-period to illustrate a statistically significant increasing trend in turbulence metrics over the continental United States and Western Atlantic. However, our work implies that NPV-jet interactions have approximately in-
365 creased by a relative amount of 11% during 2000-2021 near the climatological location of the jet stream, and that this pattern is consistently seen even when not filtering for synoptic-scale NPV features near the jet stream.

3.2 Circulation Characteristics

3.2.1 NPV-Jet Centered Composite

370 To link NPV-jet interaction events over the West Atlantic with distinct circulation pattern, a composite analysis leveraging the use of K-means clustering is performed. NPV-jet interaction events are separated into 3 clusters based on the pattern of the PV field within a 10°x10° box centered on the NPV-jet interaction point. The three clusters reveal ridging environments with different amplitudes and phases (Fig. 6). Cluster 1 shows an amplified ridge with a pronounced trough-ridge couplet. The interaction point is located on the westward flank of the ridge. Cluster 2 has a broader ridge pattern, with the interaction point
375 also located along the westward flank of the ridge. Lastly, Cluster 3 illustrates another pronounced trough-ridge couplet but with the interaction point located along the eastern flank of the ridge.

In Fig. 6a-c, composites of the PV anomalies, PV gradient anomalies and the frequency of NPV-jet interactions are shown. The maximum frequency of NPV-jet events lie adjacent to the interaction point (orange dot) along the equatorward side of
380 the jet (per definition of the composite). The interaction point is straddled by two large-scale PV anomalies of opposing sign. Note that, to a first order, the anomalies result from the large-scale flow pattern due to the presence of the trough-ridge couplet (Teubler and Riemer, 2021). However, NPV features are embedded within the large-scale negative PV anomalies and thus contribute to the large-scale negative PV anomaly signal. Directly adjacent to the interaction point lies a region of positive PV

gradient anomaly, which reaches maximum values in excess of 2.5 PVU per 100 km. The strengthened gradient lies on the
385 polar side of the 2 PVU contour, where the PV gradient rapidly sharpens towards much higher PVU values (and stratification
is particularly large).

In the second set of cluster composites (Fig. 6d-f), wind speed anomalies, upper-level geopotential height anomalies (Z) and
IVT anomalies are shown. In each cluster, the Z anomalies align with the location of the aforementioned PV anomalies. Cluster
390 1 has the most amplified trough-ridge couplet with both negative and positive Z anomalies reaching magnitudes in excess of
150 m. Cluster 1 also coincides with the strongest IVT anomalies that reach values in excess of $300 \text{ kg m}^{-1} \text{ s}^{-1}$. The close
proximity of the IVT anomaly adjacent to the trough indicates a favorable environment for large-scale ascent conducive to
squall line and warm conveyor belt development (Dacre et al., 2019).

395 Cluster 3 features NPV-jet interaction events in a comparatively drier region of the ridge (Fig. 6f). Hence, moist processes
are likely to be less important for NPV-jet interactions within this cluster. The IVT anomaly in this case is weakest but still re-
mains positive and in excess of $100 \text{ kg m}^{-1} \text{ s}^{-1}$ near the western flank of the ridge. The weakened IVT anomaly in this cluster
may partly result from the region of IVT being further away from the interaction point, and thus more radially smoothed out
by the compositing approach. Nevertheless, this cluster suggests that the IVT anomaly tends to be weaker when synoptic-scale
400 NPV interactions are located within the eastern flank of the ridge. Conceptually, the three clusters resemble the evolution of
synoptic-scale NPV features from their initial formation along the westward flank of the ridge, where strong diabatic processes
(i.e., latent heating) dominate, through their subsequent advection downstream along the apex and eastern flank of the ridge
(Oertel et al., 2020).

405 The composite of wind speed anomalies shows that all NPV-jet interactions are associated with wind speeds exceeding 40
 m s^{-1} (Fig. 6d-f) immediately poleward to the regions of high synoptic-scale NPV feature frequency (Fig. 6a-c). This result
holds regardless if synoptic-scale NPV is located on the upstream (Fig. 6a-b) or downstream (Fig. 6c) flank of the ridge. The
weaker IVT signal in Fig. 6f suggests that a highly amplified PV gradient and spatially coincident positive wind speed anoma-
lies may not require strong, positive IVT anomalies within the near-jet environment, as in Fig. 6d-e. Cluster 3 suggests that
410 synoptic-scale NPV is associated with enhanced jet wind speeds without in-situ influence from moist processes that accom-
pany the positive IVT anomaly. Of course, other larger-scale mechanisms may be co-occurring with the NPV feature, such as
supergeostrophic winds around the apex of the ridge due to flow curvature effects (Martin, 2014), which may also contribute
to the positive wind speed anomalies.

415 Figure 6g-i display the ageostrophic wind speed anomalies and PV advection by the irrotational wind (not anomaly). In
each cluster, the positive ageostrophic wind anomaly exceeds 15 m s^{-1} and is centered on the interaction point, denoting that
NPV-jet interactions are associated with highly ageostrophic environments. The ageostrophic wind speed magnitudes are of
equatable magnitude for each cluster. Attributing the ageostrophic wind anomalies to the NPV feature itself is complicated as




Figure 6. The k-means clusters ($k = 3$) derived from the latitude-weighted PV field at 250 hPa. Cluster 1 = 1136 cases, Cluster 2 = 1458 cases, Cluster 3 = 1251 cases. Fields are plotted at 250 hPa. The x and y axis represent a pseudo latitude and longitude centered on the interaction point (orange dot). (a-c) shows the PV anomaly computed from seasonal climatology (shaded, PVU) and the magnitude of the PV gradient anomaly (contoured in black at 1.5 and 2.5 PVU per 100 km). Regions where synoptic-scale NPV frequency exceeds 50% is contoured in blue. (d-f) shows the IVT anomaly (shaded, $\text{kg m}^{-1} \text{s}^{-1}$), Z anomaly contours (red: +50, 100, 150 and blue: -50, 100, 150, m) and positive wind speed anomalies (black contour: 30, 40, m s^{-1}). (g-i) illustrates the absolute PV advection by the irrotational wind (shaded, PVU day^{-1}), vectors plotted show irrotational wind anomalies $> 3 \text{ m s}^{-1}$. The black contour shows where the magnitude of the ageostrophic wind anomaly exceeds 15 m s^{-1} . j-l shows the absolute PV advection by the non-divergent wind (shading, PVU day^{-1}). Vectors show non-divergent wind anomalies $> 30 \text{ m s}^{-1}$. m-o shows the time-lagged WAF. Black vectors show WAF on the day of interaction. Blue (red) arrows show WAF 24 hours before (after) interaction. The time-lagged 2 PVU contours are shown as solid lines with the same respective colors as the WAF vectors.

the curvature of the flow pattern alongside strong latent heating can also contribute to ageostrophic winds within the near-jet
420 environment (Winters, 2021). The particular amplified flow and much stronger IVT signal in Cluster 1 could provide mechanisms that contribute to the distinct positive ageostrophic wind speed anomaly pattern that cross the westward flank of the ridge (Fig. 6g).

PV advection in Cluster 1 highlights a strong contribution from the irrotational wind field to sharpening the PV gradient
425 along the 2 PVU contour (Fig. 6g). NPV-jet interactions along the westward side of a ridge co-occur with regions of strong divergent wind field anomalies (strong upper-level outflow; Fig. 6g-i). Coupled with positive IVT anomalies, the divergent outflow is likely influenced by strong latent heating in this region (Steinfeld and Pfahl, 2019). Cluster 3 (Fig. 6i) differs with a weaker PV advection signal from the irrotational wind. The irrotational wind vectors do not directly lie over the 2 PVU line, suggesting a weak influence from the irrotational winds on perturbing the jet stream. Hence, NPV-jet interactions along the
430 downstream flank of the ridge appear to be predominantly associated with PV advection by the non-divergent wind (Fig. 6l). To a first-order, the advection signal from the non-divergent wind is associated with the eastward advection of PV by the mean flow in all clusters, and consistent with other studies assessing PV advection signals within amplified ridges (Steinfeld and Pfahl, 2019; Winters, 2021). The strongest signal in PV advection by the non-divergent wind arises in Cluster 1, which also coincides with a stronger contribution from the irrotational wind to the total PV advection (Fig. 6g) and is associated with the
435 most amplified flow pattern.

Lastly, the lagged WAF for each of the three clusters is shown in Fig. 6m-o. 24 hours before NPV-jet interaction, the large-scale circulation pattern consists of either a weakly amplified ridge (Fig. 6m-n) or zonal flow pattern (Fig. 6o). WAF is relatively small, and does not exceed the WAF threshold shown in Figs. 6m and 6o. At the time of NPV-jet interaction,
440 the ridge becomes more pronounced and amplified in all clusters. WAF vectors also emerge about the interaction point. The WAF packet emerges predominantly on the equatorward side of the jet stream in Fig. 6n-o. Additionally, the WAF packet is displaced slightly upstream of the interaction point, closer to the base of the trough in Fig. 6n. Examining the WAF 24 hours later, the WAF packet persists following its emission on the day of the NPV-jet interaction event. Furthermore, the WAF packet coherently propagates downstream in all three of the clusters, maintaining similar magnitudes as on the day of the NPV-jet
445 interaction. This maintenance of the amplified WAF packet coincides with the persistence of a more amplified ridge that is of comparable magnitude to the day of NPV-jet interaction. Remember that only the stationary component of WAF (Equation 1) is plotted here, and that there is likely additional contribution from the transient component (Takaya and Nakamura, 2001).

3.2.2 NPV-Jet (100 - 300 km) Centered Composite

The NPV-jet (100-300 km) interaction events serve as an analogue to compare against the circulation characteristics during
450 NPV-jet (< 100 km) interactions. In Fig. 7a-c, the large-scale environments largely resemble those illustrated in Fig. 6a-c, with two large-scale PV anomaly dipoles straddling a trough-ridge couplet. The maximum frequency of synoptic-scale NPV features is displaced slightly further equatorward from the jet stream compared to Fig. 6a-c. Some notable differences during




Figure 7. As in Figure 6 (first two rows) but for NPV-jet interactions using the 100-300 km threshold. Cluster 1 = 642 cases, Cluster 2 = 1969 cases, Cluster 3 = 1224 cases. Each NPV-jet (100-300 km) event is categorized into one of these clusters based on the event's similarity to the cluster centroids from Figure 6. A modification is made to the NPV frequency (dark blue contour) to show regions where synoptic-scale NPV is more frequent than 45% of the time. Stippling shows statistical significance at the 2% level using a student t-test. (a-c) shows statistically significant differences with respect to Figure 6 and where the PV gradient is greater than 1.25 PVU per 100 km. (d-f) shows statistically significant differences with respect to Figure 6 and where the wind speed differences are greater than 5 m s^{-1} .

NPV-jet (100-300 km) interactions include the occurrence of a much weaker negative PV anomaly and a slightly stronger positive PV anomaly compared to Fig. 6a-c. For reference, the average difference in the negative PV anomaly minima for the three clusters in Fig. 7a-c and Fig. 6a-c is approximately 0.8 PVU. In contrast, the positive PV anomaly maxima difference is about 0.3 PVU.

One reason for observing stronger positive PV anomalies and weaker negative PV anomalies arises from NPV-jet interactions (100 - 300 km) occurring on average at slightly lower latitudes compared to NPV-jet ($< 100 \text{ km}$) interactions (Fig. 3b-c). The computation of anomalies is sensitive to latitude. However, the comparatively weaker negative PV anomaly in Fig. 7 implies that processes that lead to deeper, large-scale negative PV anomalies are more pronounced during NPV-jet ($< 100 \text{ km}$) interactions compared to NPV-jet (100-300 km) interactions. This coincides with the PV gradient about the interaction point being over 1.25 PVU per 100 km weaker in Fig. 7a-c compared to Fig. 6a-c.

Figure 7d-f reaffirms the results of Fig. 6d-f, indicating that positive Z anomalies are weaker on the equatorward side of the jet stream in Fig. 7d-f relative to Fig. 6d-f, but stronger on the poleward side of the jet stream. Similarly, wind speed anomalies in Fig. 7d-f are comparatively weaker compared to Fig. 6d-f. In particular, wind speed anomaly maxima are weaker by 8 m s^{-1}

in Fig. 7d-e and over 10 m s^{-1} in Fig. 7f. It is difficult to ascertain whether the weaker wind speeds result from synoptic-scale NPV being positioned further away or whether it results from differences in the large-scale fields. The results do confirm that
 470 NPV-jet ($< 100 \text{ km}$) interactions occur in more amplified flow patterns compared to NPV-jet ($100\text{-}300 \text{ km}$) interaction events, and that the combination of more amplified flow patterns and closer proximity of synoptic-scale NPV to the jet stream coincide with statistically significant enhancements of PV gradient and wind speeds about the interaction point.

The changes in jet stream kinematics associated with diabatic heating are likely to play a lesser role in Cluster 3 (Fig. 7f)
 475 due to the distance of the interaction point from the positive IVT anomaly. In this drier region of the ridge, it is suggested that changes in wind speeds and PV gradient are predominantly driven by adiabatic processes (Buknenberger et al., 2023). Thus, the differences in PV gradient and wind speeds between NPV-jet ($< 100 \text{ km}$) compared to NPV-jet ($100\text{-}300 \text{ km}$) imply that large-scale regions of diabatic heating attributed to the IVT anomaly may not explain the observed kinematic differences, and that their may be significant adiabatic differences between the two distance thresholds. To more explicitly demonstrate the
 480 importance of synoptic-scale NPV within the near-jet environment and to better separate its impact from the large-scale flow, the WAF equation is partitioned in the following section.

3.3 Archetype Case-Study Analysis

The composite approach is not well suited for evaluating the evolution of temporally fast, mesoscale features. Consequently,
 485 a case-study approach is favored to further illustrate the kinematic impacts of NPV on the jet stream for each cluster. Three archetype cases are selected that are most representative of their corresponding cluster. Archetype cases are identified using a Euclidean distance metric (lower Euclidean distance denotes greater similarity to the cluster's mean PV field). From 10 cases for each cluster that best resemble the mean PV field, one case for each cluster is subjectively selected. The subjectively chosen cases are deemed to be the most illustrative for highlighting the influence of NPV features on the jet stream. To summarize,
 490 three cases that best represent their respective clusters are chosen to evaluate NPV-jet interactions through a detailed, wave activity flux perspective.

3.3.1 Synoptic Overview

In Fig. 8a-c, a synoptic overview is provided for each archetype case. The large-scale circulation for each case illustrates a ridge with strong IVT along its westward flank. Fig. 8a-b depict a synoptic-scale NPV feature predominantly along the western flank
 495 of the ridge, while the synoptic-scale NPV feature in Fig. 8c is mainly located along the apex and eastern flank of the ridge. For the second and third cluster, the NPV feature does not overlap with the region of strong IVT, implying the NPV feature is located away from the immediate influence of broad regions of latent heat release. For each case, the point where NPV is in closest proximity to the jet stream coincides with wind speed maxima. Wind speeds in excess of 70 m s^{-1} are observed in Fig. 8b-c, and surpassing 90 m s^{-1} in Fig. 8a).

500

Figure 8d-f analyzes relative vorticity and its inversion to assess the impact of relative vorticity within NPV on the large-scale flow. Regions of anticyclonic vorticity with a magnitude exceeding $1 \times 10^{-4} \text{ s}^{-1}$ are predominantly situated within the NPV features, indicating that the features are inertially unstable. On the polar side of the 2 PVU line, strips of cyclonic relative vorticity are observed adjacent to the NPV feature for each case. An inversion of the relative vorticity within the boxed domain predominantly results in an anticyclonic non-divergent wind field with wind speeds maximized about the 2 PVU contour reaching values of 45 m s^{-1} in Fig. 8d and Fig. 8f. Figure 8e involves an NPV feature adjacent to a strip of cyclonic vorticity exceeding $2 \times 10^{-4} \text{ s}^{-1}$ and a non-divergent wind maximum of 55 m s^{-1} . The vorticity inversion illustrates the influence of the strong cyclonic vorticity in this case (Fig. 8e). Namely, two dipoles rotating in opposing directions are illustrated, resulting from the interaction between anticyclonic vorticity (associated with NPV) and cyclonic vorticity on the polar side of the 2 PVU contour (Cunningham and Keyser, 2004; Pyle et al., 2004).

The wind speed magnitude derived from the inversion can account for over 50% of the maximum wind speeds in Fig. 8a, 75% in Fig. 8b, and 65% in Fig. 8c. The relative vorticity inversion implies that the interaction between NPV and strips of cyclonic vorticity on the polar side of the jet stream account for the majority of the total wind field signal. As a final test (not shown), the relative vorticity inversion was exclusively performed over the region of NPV (without incorporating the influence of the cyclonic vorticity). Maximum non-divergent wind speed values of 20 m s^{-1} were obtained for Fig. 8d and 25 m s^{-1} for Fig. 8e-f. In other words, the anticyclonic vorticity associated with NPV appeared to contribute to approximately 50% of the total non-divergent wind field in the three cases. The maxima in the magnitude of the non-divergent winds attributed to NPV are larger compared to the $5 - 10 \text{ m s}^{-1}$ attributed to a mesoscale NPV feature studied in Oertel et al. (2020). Although a different methodology was used in their study to quantify the influence of NPV on jet stream wind speeds based on 2-hourly wind circulation anomalies. Either way, NPV appears to lead to additional positive contributions to the non-divergent wind speed.

3.3.2 Individual Wave Activity Flux Terms

In this section, the terms that contribute to the total WAF equation are evaluated with respect to the synoptic-scale NPV feature in each archetypal case. As the WAF equation is constructed from multiple different components (Eq. 1), the analysis is narrowed to focus on the first part of the WAF equation, $U(\psi_x'^2 - \psi' \psi_{xx}')$ (although other terms will also be discussed when appropriate). As will be shown, these terms contribute the most to the WAF equation for each case.

In Fig. 9a-c, the non-divergent wind anomaly magnitude maxima is observed to lie adjacent to the NPV feature along the 2 PVU contour. Drawing insights from the relative vorticity inversion discussed in the preceding sub-section, it is suspected that the maxima in non-divergent wind anomalies stem from the dynamic interplay between the NPV feature and strong cyclonic relative vorticity on the polar side of the jet stream. First, focusing on $\psi_x'^2$, which is found in $U(\psi_x'^2 - \psi' \psi_{xx}')$, and the $\psi_y'^2$ momentum term, it follows that the maximum in momentum transport overlaps with regions of strong non-divergent wind




Figure 8. Synoptic overview of three cluster archetypes. The Cluster 1 Case (left column) occurs on: 2004-03-24 00 UTC, Cluster 2 Case (middle column): 2014-12-09 18 UTC, Cluster 3 Case (right column): 2021-05-24 18 UTC. In each panel, the 2 PVU contour (black line) and NPV (blue line) are shown. The orange dot illustrates the interaction point for each case (the coordinate at which the NPV feature is in closest proximity to the 2 PVU contour). (a-c) shows the IVT, $\text{kg m}^{-1} \text{s}^{-1}$ (shaded), and wind speed contours in red at 70 m s^{-1} (the wind speed is shown at 80 m s^{-1} in a). (d-f) shows relative vorticity (shaded), s^{-1} , and its subsequent inversion. The vectors show the non-divergent winds obtained from the inversion (computed within the grey, dashed box)

anomalies. For each case, $\psi_y'^2$ (zonal momentum transport) is maximized along the zonal extent of the jet stream. The $\psi_x'^2$ term (meridional momentum transport) is most prominent along the meridional extent of the jet stream (i.e., the western and eastern flanks of the large-scale ridge). For each case, these terms are maximized adjacent to the NPV features. In the three cases presented, $\psi_x'^2$ exceeds $2000 \text{ m}^2 \text{s}^{-2}$, reaching its maxima precisely where the NPV feature is closest to the jet stream. The $\psi_x'^2$ term reaches $4000 \text{ m}^2 \text{s}^{-2}$ where NPV is in closest proximity to the jet stream in Fig. 9a. The larger magnitude of the $\psi_x'^2$ term in this case corresponds to the much stronger non-divergent wind anomalies observed along the entirety of the jet stream. It can be inferred that while the large-scale environment varies in terms of its magnitude of momentum transport across the different cases, the presence of NPV features appears to consistently coincide with a maximum in momentum transport in all three archetype cases.

In Fig. 9d-f, the ageostrophic geopotential flux is evaluated. Focus is placed on the ψ' and ψ_{xx}' terms of $U(\psi_x'^2 - \psi' \psi_{xx}')$. The ridge is characterized by positive ψ' (positive streamfunction anomaly). The ψ_{xx}' term is related to relative vorticity as it equates to a v_x anomaly. In each case, negative values of this term largely overlap with the NPV feature observed in Fig. 9a-c. This is because NPV is associated with a maximum in anticyclonic vorticity (Fig. 8d-f). It can also be seen in Fig. 9d-f that negative ψ_{xx}' overlaps with positive ψ' . These two terms must be multiplied together to obtain $\psi' \psi_{xx}'$. Given that $\psi' \psi_{xx}'$ is negative when anticyclonic shear overlaps with a positive streamfunction anomaly, subtraction of this term following the sign

convention in $U(\psi_x'^2 - \psi' \psi_{xx}')$ means that $\psi' \psi_{xx}'$ becomes a positive contribution to WAF.

In Fig. 9e-f, $-\psi' \psi_{xx}'$ is maximized within the ridge where overlap between ψ_{xx}' and ψ' occurs. In Fig. 9d, ψ_{xx}' does not completely overlap with ψ' , thus reducing the contribution of $-\psi' \psi_{xx}'$ to WAF. In other words, NPV must be optimally embedded within a ridge environment such that its anticyclonic shear can enhance the ageostrophic flux of geopotential.

In Fig. 9g-i, $\psi_x'^2$ and $-\psi' \psi_{xx}'$ are combined to obtain $U(\psi_x'^2 - \psi' \psi_{xx}')$. In other words, the ageostrophic flux associated with anticyclonic shear is additive with the momentum flux term, $\psi_x'^2$. This means that $U(\psi_x'^2 - \psi' \psi_{xx}')$ positively contributes to the WAF equation (assuming the base state wind, U , is positive). For each case shown in Fig. 9g-i, the region where NPV interacts with the jet stream is characterized by a maximum in $U(\psi_x'^2 - \psi' \psi_{xx}')$. It is proposed that this maximum arises due to the NPV feature enhancing both the momentum transport and the ageostrophic geopotential flux. In Fig. 9g, the contribution to the $U(\psi_x'^2 - \psi' \psi_{xx}')$ component of the WAF equation predominantly arises from momentum transport, with ageostrophic geopotential flux becoming more dominant further away from the 2 PVU contour where momentum transport is minimized. In Fig. 9h and Fig. 9i, the ageostrophic flux term contributes more to $U(\psi_x'^2 - \psi' \psi_{xx}')$ as a result of the weaker momentum transport in the latter two cases. This indicates case-to-case dependence as to which terms contribute more to enhancing wave activity.

3.3.3 Full Wave Activity Flux

Following the evaluation of the individual terms in $U(\psi_x'^2 - \psi' \psi_{xx}')$, the full WAF equation is now evaluated with respect to NPV-jet interactions. Figure 10a-c illustrates spatial agreement between regions where large magnitudes of the full WAF are observed and large magnitudes of $U(\psi_x'^2 - \psi' \psi_{xx}')$ are observed (Fig. 9g-i). WAF is maximized about the 2 PVU contour in Fig. 10a where momentum transport dominates the contribution to the WAF equation for the case-study for Cluster 1 (fig, 9g). In contrast, WAF is maximized over the NPV feature in Fig. 10c, where the ageostrophic geopotential flux plays a more dominant role over momentum transport (Fig. 9i).

The $U(\psi_x'^2 - \psi' \psi_{xx}')$ term appears to dominate the WAF signal for each of the three cases. This observation is made due to the similarity of the spatial patterns shown in Figs. 9a-c with respect to Fig. 9g-i. To confirm this, all four terms of the WAF equation are shown as a bar graph in Fig. 10d-f. When focusing on the first two parts of the equation, which contribute to the zonal component of the WAF equation, $U(\psi_x'^2 - \psi' \psi_{xx}')$ dominates for each of the three cases. In contrast, the $U(\psi_x' \psi_y' - \psi' \psi_{xy}')$ part tends to be weaker. One reason for this occurrence is that in the second part of the WAF equation, ψ_{xy}' is associated with flow divergence. In the cases selected, relative vorticity associated with the NPV feature tends to dominate the WAF signal over divergence, indicating that the NPV region is predominantly defined by its relative vorticity.




Figure 9. Component analysis focusing on the first part of the wave activity flux equation, $U(\psi'_x{}^2 - \psi'\psi'_{xx})$. (a-c) focuses on momentum: The non-divergent wind anomaly magnitude, m s^{-1} is shaded in red, the magenta contours show the momentum transport anomaly terms at a threshold of $2000 \text{ m}^2 \text{ s}^{-2}$. Solid contours denote meridional momentum, ψ'^2_x while dashed contours show zonal momentum, ψ'^2_y . The solid dark blue contour shows the synoptic-scale NPV feature. (d-f) examines the ageostrophic flux of geopotential. The solid (dashed) grey contours denote positive (negative) streamfunction anomaly, ψ' at intervals of $\pm 1, 3 \times 10^7 \text{ m}^2 \text{ s}^{-1}$. The dark blue dashed contour denotes a negative shear anomaly ψ'_{xx} at $-0.75 \times 10^{-4} \text{ s}^{-1}$. The red shading illustrates the $-\psi'\psi'_{xx}$ term, relating to the ageostrophic flux of geopotential. The term is computed by multiplying the streamfunction anomaly and shear anomaly. Note that following sign convention in the WAF equation, a negative sign is placed at the front of $-\psi'\psi'_{xx}$. In (g-i), the entirety of $U(\psi'^2_x - \psi'\psi'_{xx})$ is shown in shading. Regions where $\psi'\psi'_{xx}$ is of larger magnitude than ψ'^2_x are stippled. The solid dark blue contour is as in (a-c). The orange dot in all panels denotes the interaction point.




Figure 10. Evaluation of all terms in the WAF equation (Section 2.2.3, Equation 1). (a-c) shows the magnitude of WAF, $\text{m}^2 \text{s}^{-2}$, shaded in red and the direction of WAF propagation shown by the vectors. All other features plotted are as in Fig. 9 (g-i). (d-f) shows a bar-chart in which the WAF equation is split into its four different parts as shown by the legend with units of $\text{m}^3 \text{s}^{-3}$. The units are this way because the $\frac{1}{|\bar{U}|}$ term is excluded. Exclusion of the base state wind term is chosen to more explicitly focus on the momentum transport and ageostrophic geopotential flux terms. Part 1 and Part 2 refer to the zonal component of the WAF equation. Part 3 and Part 4 are associated with meridional component of the WAF equation. The bar charts represent the mean of each WAF part, computed within the dashed box domain from grid-points that exceed $1200 \text{ m}^2 \text{s}^{-2}$.

The meridional component of WAF is weaker than the zonal component for each case, which likely results from the merid-
585 ional base-state wind (V) being weaker than the zonal base-state wind (U) when computing WAF. When evaluating the merid-
ional WAF terms further, the $V(\psi_x'^2 - \psi_y'\psi_{yy})$ part also dominates $V(\psi_x'\psi_y' - \psi_y'\psi_{xy})$ in Fig. 10e-f, implying once more that
relative vorticity may be dominating divergence in the boxed regions. However, Fig. 10d highlights the importance of case-to-
case variability. Strong shear linked to divergence nearby the NPV feature could also modulate the WAF signal during NPV-jet
interactions by potentially enhancing the $\psi'\psi_{xy}$ parts. Hence, while the WAF equation can be used to clearly illustrate the
590 impact of synoptic-scale NPV features on the jet stream, a partitioning approach is recommended to carefully determine how
relevant each term is in the equation for explaining how NPV modifies wave activity.

To summarize, the three archetype cases suggest that the presence of synoptic-scale NPV within a ridge (on the equatorward
side of the jet stream) can be dynamically relevant for enhancing the instantaneous WAF. The enhancement of WAF arises
595 from the increase in momentum transport and the anticyclonic shear in the vicinity of the jet across all cases resulting from
the presence of strong anticyclonic-vorticity embedded in the synoptic-scale NPV. Given that these cases examine a single, in-
stantaneous time-step of WAF, it is suspected that synoptic-scale NPV may be particularly relevant in enhancing wave activity

during instances when multiple, synoptic-scale NPV features are present that exhibit temporal persistence.

600 4 Discussion

This section contextualizes the results with respect to relevant literature and addresses some limitations of the study. Our climatological analysis demonstrates that NPV occurs more frequently at lower latitudes and decreases towards higher latitudes. These patterns are consistent with other climatological studies of NPV using ERA5 (Lee et al., 2023) and of inertial instability using ERA-interim (Thompson et al., 2018) at 250 hPa. These studies also agree in the observed poleward extension
605 of enhanced NPV (and inertial instability) frequency over the western North Atlantic (Fig. 3a). Lee et al. (2023) illustrates that the more common occurrence of turbulence indices, such as NPV, over the western North Atlantic coincides with more frequent instances of negative Brunt–Väisälä frequency, particularly during winter months. We suspect that enhanced negative Brunt–Väisälä frequency over the western North Atlantic may result from the common occurrence of trough-ridge couplets characterized by positive IVT anomalies (Fig. 6d,e,f), which prime the environment to be more conducive to severe convection
610 (O’Brien et al., 2024). Since NPV is generated in environments characterized by pronounced convective instability (Weijenborg et al., 2017), winter season trough-ridge couplets may provide a suitable genesis region for NPV-jet interactions.

The increasing trend of NPV-jet interactions from 2000 to 2022 (Fig. 5b) could imply that the synoptic setup favoring NPV-jet interactions has become more common over the last two decades. However, due to the limited time frame of our study,
615 we cannot infer whether the source of this trend is anthropogenic or natural variability. Studies examining longer time-periods suggest that planetary-scale anthropogenic influences could explain positive trends in wind shear (Lee et al., 2019; Prosser et al., 2023) and jet streaks (Shaw and Miyawaki, 2024) due to the strengthening of the meridional temperature gradient through lower stratospheric cooling in polar regions and upper tropospheric warming in the tropics. Given that NPV is closely associated with jet streaks and strong horizontal and vertical shear (Oertel et al., 2021), we recommend that future research
620 on kinematic jet stream trends also evaluate the significance of more localized dynamical features, such as regional trends in NPV. Further climatological evaluation of NPV is also important in the context of warm-conveyor belts over the West Atlantic, which are projected to become more intense in terms of their diabatic heating and precipitation (Joos et al., 2023), which could facilitate the increased occurrence of NPV features.

625 Some additional potential biases to consider in our climatological analysis relate to the use of reanalysis data. ERA5 is an optimal dataset due to the dense observational network over the North Atlantic (Tenenbaum et al., 2022). However, improvements in the quality and quantity of observations from 2000 to 2022 (Hersbach et al., 2020) introduce an unquantified bias that could influence trend analysis. More details on this are discussed in Tenenbaum et al. (2022). Another caveat is the resolution of the dynamical model used to construct the dataset. We advise against solely using dynamical models that do not explicitly re-
630 solve deep convection for detailed evaluations of NPV features, as coarse models struggle with resolving NPV features (Clarke

et al., 2019; Lojko et al., 2022). While ERA5 uses a dynamical model that parameterizes deep convection, we suspect that the data assimilation process is crucial for correcting for the location of NPV. New regional reanalysis datasets like CONUS404 (Rasmussen et al., 2023) use horizontal resolutions that explicitly resolve deep convection, but higher-resolution simulations will have noisier PV fields (Oertel and Schemm, 2021). Consequently, a climatological analysis of NPV using datasets such as
635 CONUS404 may identify fewer (more) synoptic-scale (convective-scale) NPV features compared to coarser reanalysis products. Thus, the results presented here are unique to ERA5 and likely to reanalysis datasets of comparable resolution.

The composite perspective indicates that NPV-jet interactions occur within ridge environments in which positive IVT anomalies are present on the westward flank of the ridge and are most frequent during boreal winter months over the western North
640 Atlantic. Oertel et al. (2021) postulate that a pre-existing, strong jet stream is a conducive environment that favors the development of elongated NPV features. Winter months tend to be associated with the climatologically fastest jet stream winds (Iqbal et al., 2018), hence alongside increased convective instability during winter (Lee et al., 2023), the winter-time maximum in NPV-jet interactions over the Coastal West-Atlantic may arise from the more favorable jet stream environment for elongating NPV features. Further evidence for this hypothesis is presented in Fig. 4e-f, in which, despite the summer months having a 1.5
645 times greater area coverage in NPV of all sizes, synoptic-scale NPV was 10 times less likely to be observed compared to winter.

The interaction of synoptic-scale NPV with the jet stream in the composite approach highlights the interaction between two opposing large-scale potential vorticity vorticity dipoles. The archetype case-studies provide additional detail, noting that maximum values in jet stream wind speeds are observed to be situated between more localized relative vorticity dipoles, with
650 anticyclonic relative vorticity being linked to the NPV features. Cunningham and Keyser (2004) use a barotropic framework to illustrate that the interaction of two-dimensional relative vorticity dipoles can explain the development of jet streaks, which has also been observed in real-cases within trough-ridge couplet environments (Pyle et al., 2004). In our research, the interaction of anticyclonic vorticity (NPV) with cyclonic vorticity on the poleward side of the jet stream appears to coincide with strong, positive wind speed anomalies along the jet stream. Hence, while our analysis is predominantly focused on the NPV feature
655 itself, it is likely important to also acknowledge the importance of cyclonic vorticity on the poleward side of the jet stream reinforcing with the anticyclonic vorticity associated with NPV.

Computing an inversion of the anticyclonic relative vorticity associated with the NPV features indicates that it substantially contributes to the non-divergent wind speeds along the jet stream. This may explain why wind speed maxima is commonly
660 observed adjacent to synoptic-scale NPV features in the case-study archetypes (Fig. 9), as opposed to elsewhere in the ridge where the anticyclonic circulation is benign. Acknowledging that the archetypes are representative of the composites in Fig. 6, it should also be expected that the circulation associated with NPV contributes to their distinct positive wind speed and PV gradient anomalies about the NPV-jet interaction point. During NPV-jet (100-300 km) interactions, wind speeds and PV gradient maxima are comparatively weaker (Fig. 7). While an explanation could be that the reinforcement between anticyclonic
665 relative vorticity associated with NPV and cyclonic relative vorticity on the poleward side of the jet stream is weakened, we also

note that the large-scale flow pattern tends to be less amplified. Hence, the weaker jet stream kinematics in Fig. 7 may likely result from the concomitant influence of NPV being positioned further away from the jet and the large-scale flow being less amplified. However, determining which factor is more influential for jet stream kinematics is not accomplished here.

670 As noted in a review paper by Keller et al. (2019), two frameworks by which to understand the development of jet streaks is through PV advection by the irrotational wind or through the Cunningham and Keyser (2004) approach to define jet streaks via the interaction between two opposing vorticity dipoles. In our composite analysis, strong irrotational wind fields were present in cluster one and two when synoptic-scale NPV was located near the westward flank of the ridge. This suggests that NPV can be embedded within synoptic-scale divergent outflow and could contribute to the strengthening the magnitude of the overall PV
675 advection signal, and thus favor the occurrence of jet streaks. A caveat of this circulation pattern is that the synoptic-scale NPV features are embedded within the diabatically reduced, large-scale negative PV anomaly that is also experiencing advection by the irrotational outflow. This flow set-up makes it difficult to quantify how much the NPV feature additionally contributes to enhancing wind speeds and PV gradient compared to the surrounding, diabatically influenced PV field.

680 We do note that one of the NPV-jet interaction clusters identifies that synoptic-scale NPV can also frequently interact with the jet stream even when strong irrotational outflow is weak, such as along the eastern flank of a ridge (Fig. 6i). This location is observed to have the same wind speed anomaly values as the other clusters which have comparatively stronger irrotational wind fields and much more pronounced IVT anomalies. The result suggests that broad regions of strong latent heating in combination with a strong irrotational wind field may not always be necessary to obtain jet streaks when NPV is present. Instead,
685 this cluster suggests that synoptic-scale NPV features may induce adiabatic enhancement of jet stream wind speeds, despite its initial diabatic origin (Buknenberger et al., 2023).

As an additional point of clarification, care must also be taken in treating jet streaks associated with NPV-jet interactions as 2D, geostrophic features. While our analysis is performed on a single level and appropriate comparisons are made to the 2D
690 idealized jet streaks discussed in Cunningham and Keyser (2004), the composites here identify NPV-jet interactions to be frequently associated with highly ageostrophic environments (Fig. 6g-i). Ageostrophy can significantly modify the intensification of the PV gradient along the tropopause (Winters, 2021) and influence the three dimensional wind speed profile. Hence, a 3D kinematic analysis of how NPV influences jet stream wind speeds would certainly be warranted.

695 A WAF approach was recently used to show that synoptic-scale NPV can degrade jet stream forecast skill within global numerical weather prediction models (Lojko et al., 2022). Specifically, it was found that the magnitude of the synoptic-scale NPV feature's anticyclonic relative vorticity was under-represented, and coincident with the manifestation of WAF errors. This finding is particularly interesting given that NPV-jet interactions appear to be geographically focused over the western North Atlantic. Grazzini and Vitart (2015) show that Rossby wave packets initiated over the western North Atlantic tend to
700 be associated with poorer medium-range predictability over Europe. Our study shows that WAF packets (analogous to Rossby

waves) tend to be emitted during NPV-jet interactions (Fig. 6m-o) and that synoptic-scale NPV can be dynamically relevant in enhancing the magnitude of WAF (Fig. 10a-f). It would thus be interesting to further explore the relevance of synoptic-scale NPV for predictability within a composite perspective.

705 5 Conclusions

The study presents a composite overview on the climatology and dynamical impact of synoptic-scale bands of negative potential vorticity (NPV) on the Western North Atlantic jet stream. The study is conducted using ERA5, and the obtained results should be contextualized with respect to the characteristics of the dataset. NPV features are identified using 6-hourly ERA5 data at 250 hPa over the period 2000 - 2021. Using the PV field, PV values ≤ -0.01 PVU and > 1650 km (> 98 th percentile) are used to identify synoptic-scale NPV. 'Interactions' of NPV with the jet stream are identified when NPV features are located within 100 km of a circumpolar 2 PVU contour. The 2 PVU contour is used to represent the jet stream and these interactions are referred to as NPV-jet interactions.

The results are split into three parts: A climatological quantification of the frequency of NPV-jet interactions, a composite analysis of the dynamics during NPV-jet interactions, and three case studies involving a mechanistic evaluation of the impact of synoptic-scale NPV on the jet stream through the wave activity flux perspective. The climatological analysis shows that NPV-jet interactions have a probability of occurring up to 1.2% of the time at particular grid-points over the Western Atlantic, maximized at a latitude of 40°N . Interactions are most frequent during winter ($> 2.5\%$) and least frequent during the summer months ($< 0.5\%$). The seasonal frequencies reaffirm previous case-study work (Harvey et al., 2020; Oertel et al., 2020) hypothesizing that a pre-existing, strong jet stream (which is climatologically more likely to occur in winter) is an ideal environment in which synoptic-scale NPV features can occur.

An investigation of NPV trends in the study region illustrates an increasing trend over a narrow latitude band centered at 40°N encompassing Eastern North America and the Western Atlantic. For (all NPV) NPV-jet interactions, there has been a relative frequency increase of over (45%) 11% in some localized regions of the Western Atlantic over the 22-year time-period based on a linear trend analysis. This result bares resemblance to Lee et al. (2023) who report increasing NPV frequency during Boreal Winter over the past four decades across localized regions of the North-Western Atlantic. Albeit, the spatial extent and magnitude of the trend in Lee et al. (2023) tends to be weaker than the percentages obtained in our work which uses a shorter (more recent) time-period.

Generally, NPV-jet interactions are characterized by strong, positive PV gradient anomalies (2.5 PVU per 100 km) which coincide with enhanced (ageostrophic) wind speed anomalies exceeding 40 m s^{-1} (15 m s^{-1}). Even when NPV features are located in regions far away from regions of diabatic heating that originally produced NPV, these positive anomalies persist,

highlighting that NPV can be associated with strengthened wind speeds without co-occurring alongside strong moisture anomalies. The results suggest that synoptic-scale NPV features could predominantly resemble a dry dynamics interaction with the jet stream (Cunningham and Keyser, 2004) in some circumstances. This finding is complemented by previous case-studies that have observed elongated bands of NPV to advect quasi-adiabatically along the jet stream (Oertel et al., 2020; Lojko et al., 2022).

To gain additional insight on how NPV amplifies the jet stream, the 2D WAF equation at 250 hPa is applied. Composite analysis of the WAF equation denotes that a packet of WAF manifests at the time of NPV-jet interaction and subsequently propagates downstream. This co-occurs alongside the amplification of a trough ridge couplet, indicating that synoptic-scale NPV features tend to develop as the jet stream becomes increasingly amplified. Partitioning the WAF equation into its momentum and ageostrophic geopotential flux components within three archetype case-studies illustrates that the anticyclonic relative vorticity associated with NPV can locally enhance the magnitude of wave activity flux. Synoptic-scale NPV features appear to coincide with localized regions of wave activity maxima when NPV features are optimally embedded within a ridge environment (i.e., in close proximity to the jet stream but situated within an amplified, positive stream function anomaly).

Overall, our results reaffirm previous case-study examinations of elongated NPV features. Elongated NPV features tend to exist along the equatorward side of an amplified jet stream pattern (i.e., trough-ridge couplets). In these flow patterns, the close proximity of synoptic-scale NPV features to the jet stream coincides with PV gradient and wind speed maxima. The vigorous anticyclonic relative vorticity within NPV features appears to play an important role in the kinematic strengthening of the jet stream, both in-situ and potentially downstream by serving to additionally enhance wave activity within pre-existing amplified flow patterns.

Designing a more elegant framework for quantifying the role of NPV on jet stream amplification would be a suitable next-step in constraining kinematic uncertainties associated with NPV-jet interactions. While PV inversion might not suit dynamically unstable NPV regions, the application of a 2D relative vorticity inversion (Oertel and Schemm, 2021) and the 2D WAF equation (Takaya and Nakamura, 2001) shows a pathway to quantitatively assess the effect of elongated NPV on the jet stream. A 2D approach may also be appropriate given that synoptic-scale NPV features appear to predominantly occur along isobaric levels related to the tropopause.

More generally, there still remains plenty of mystery regarding the concept and properties of NPV. For example, it remains unknown why synoptic-scale NPV can persist for long periods of time and by which processes NPV dissipates. Further, it would be interesting to synthesize the full life-cycle of NPV from its suspected generation within convective-scale PV dipoles to its upscale growth into an elongated filament of synoptic-scale NPV. Such research will progress our understanding of the dynamics of NPV, and its implications for weather prediction and aviation applications.

Code and data availability. The Python based algorithm for identification of negative PV features and their interaction with the jet stream can be downloaded from: https://github.com/AlexLojko/NPV_Algorithm. The Python script for computing a relative vorticity inversion can be downloaded from: <https://github.com/evans36/miscellany>. The ERA5 dataset is available from <https://cds.climate.copernicus.eu/> which is downloaded using era5cli: <https://github.com/eWaterCycle/era5cli>

Appendix A: NPV-jet Interactions: Sensitivity Tests

This section illustrates the motivation for selecting the 250 hPa level and the 100 km distance threshold when defining NPV-jet interactions. Figure A1 illustrates the identification of synoptic-scale NPV features for a selected year. NPV features are most frequent at 250 and 300 hPa for all length-scales of NPV. While the amount of NPV features is of the same order at each isobaric level when examining features smaller than 1000 km, the number of NPV features identified at 250 and 300 hPa are an order of magnitude more frequent when identifying NPV features greater than 1000 km. These results indicate that synoptic-scale NPV is predominantly an upper-tropospheric feature, maximized at isobaric levels that are in close proximity to the tropopause.

Author contributions. AL designed the project, downloaded the data, performed the data analysis and wrote the manuscript. AW and AO contributed to editing the manuscript. AW, AO and CJ worked on discussing the methodology with AL. All authors contributed to discussing the results.

Competing interests. The authors have no competing interests.

Acknowledgements. AL conducted the research with grant support from the Momental Foundation: Mistletoe Research Fellowship and the National Center for Atmospheric Research Graduate Visitor Program Fellowship. AW was supported by the National Science Foundation under AGS Grant No. 2114011. AO was supported by the Transregional Collaborative Research Center SFB / TRR 165 “Waves to Weather” (www.wavestoweather.de) funded by the German Research Foundation (DFG). The contribution of AO was also partly carried out within the Italia – Deutschland science-4-services network in weather and climate (IDEA-S4S; INVACODA, 4823IDEAP6). This Italian-German research network of universities, research institutes and Deutscher Wetterdienst is funded by the BMDV (Federal Ministry of Digital and Transport). Research visits by AL to scientists at the Karlsruhe Institute of Technology and the National Center for Atmospheric Research provided fruitful discussions on the project and the concept of negative potential vorticity. We acknowledge help from Dr. Kevin Prince and Dr. Clark Evans regarding discussions on the relative vorticity inversion script. We also thank the editor and two anonymous reviewers for their comments which have served to improve the quality of the manuscript.




Figure A1. Frequency of NPV labels at different isobaric levels in 2016. The year 2016 is selected as it is associated with the most NPV-jet interactions at 250 hPa Fig. 5d. The x-axis illustrates the major axis length scale of NPV labels, binned at intervals of 1000 km. The y-axis shows the logged, total count of NPV labels.

References

- 795 Archambault, H. M., Bosart, L. F., Keyser, D., and Cordeira, J. M.: A climatological analysis of the extratropical flow response to recurving western North Pacific tropical cyclones, *Monthly Weather Review*, 141, 2325–2346, 2013.
- Barnes, E. A., Hartmann, D. L., Frierson, D. M., and Kidston, J.: Effect of latitude on the persistence of eddy-driven jets, *Geophysical research letters*, 37, 2010.
- Baumgart, M. and Riemer, M.: Processes governing the amplification of ensemble spread in a medium-range forecast with large forecast
800 uncertainty, *Quarterly Journal of the Royal Meteorological Society*, 145, 3252–3270, 2019.
- Bennetts, D. A. and Hoskins, B.: Conditional symmetric instability-a possible explanation for frontal rainbands, *Quarterly Journal of the Royal Meteorological Society*, 105, 945–962, 1979.
- Berman, J. D. and Torn, R. D.: The impact of initial condition and warm conveyor belt forecast uncertainty on variability in the downstream waveguide in an ECWMF case study, *Monthly Weather Review*, 147, 4071–4089, 2019.
- 805 Blanchard, N., Pantillon, F., Chaboureau, J.-P., and Delanoë, J.: Mid-level convection in a warm conveyor belt accelerates the jet stream, *Weather and Climate Dynamics*, 2, 37–53, 2021.
- Bony, S., Stevens, B., Frierson, D. M., Jakob, C., Kageyama, M., Pincus, R., Shepherd, T. G., Sherwood, S. C., Siebesma, A. P., Sobel, A. H., et al.: Clouds, circulation and climate sensitivity, *Nature Geoscience*, 8, 261–268, 2015.
- Braun, S. A. and Houze Jr, R. A.: The heat budget of a midlatitude squall line and implications for potential vorticity production, *Journal of*
810 *Atmospheric Sciences*, 53, 1217–1240, 1996.
- Bukenberger, M., Rüdüsühli, S., and Schemm, S.: Jet stream dynamics from a potential vorticity gradient perspective: The method and its application to a kilometre-scale simulation, *Quarterly Journal of the Royal Meteorological Society*, 149, 2409–2432, 2023.
- Cepi, P., Brient, F., Zelinka, M. D., and Hartmann, D. L.: Cloud feedback mechanisms and their representation in global climate models, *Wiley Interdisciplinary Reviews: Climate Change*, 8, e465, 2017.
- 815 Chagnon, J. M. and Gray, S. L.: Horizontal potential vorticity dipoles on the convective storm scale, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 135, 1392–1408, 2009.
- Clarke, S. J., Gray, S. L., and Roberts, N. M.: Downstream influence of mesoscale convective systems. Part 1: influence on forecast evolution, *Quarterly Journal of the Royal Meteorological Society*, 145, 2933–2952, 2019.
- Conzemius, R. J. and Montgomery, M. T.: Clarification on the generation of absolute and potential vorticity in mesoscale convective vortices,
820 *Atmospheric Chemistry and Physics*, 9, 7591–7605, 2009.
- Cunningham, P. and Keyser, D.: Dynamics of jet streaks in a stratified quasi-geostrophic atmosphere: Steady-state representations, *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 130, 1579–1609, 2004.
- Dacre, H. F., Martinez-Alvarado, O., and Mbengue, C. O.: Linking atmospheric rivers and warm conveyor belt airflows, *Journal of Hydrometeorology*, 20, 1183–1196, 2019.
- 825 Davies-Jones, R.: Streamwise vorticity: The origin of updraft rotation in supercell storms, *Journal of Atmospheric Sciences*, 41, 2991–3006, 1984.
- Davis, C. A. and Emanuel, K. A.: Potential vorticity diagnostics of cyclogenesis, *Monthly weather review*, 119, 1929–1953, 1991.
- Davis, C. A., Stoelinga, M. T., and Kuo, Y.-H.: The integrated effect of condensation in numerical simulations of extratropical cyclogenesis,
830 *Monthly Weather Review*, 121, 2309–2330, 1993.

- Dawson, A.: Windspharm: A high-level library for global wind field computations using spherical harmonics, *Journal of Open Research Software*, 4, 2016.
- Grams, C. M. and Archambault, H. M.: The key role of diabatic outflow in amplifying the midlatitude flow: A representative case study of weather systems surrounding western North Pacific extratropical transition, *Monthly Weather Review*, 144, 3847–3869, 2016.
- 835 Grams, C. M., Magnusson, L., and Madonna, E.: An atmospheric dynamics perspective on the amplification and propagation of forecast error in numerical weather prediction models: A case study, *Quarterly Journal of the Royal Meteorological Society*, 144, 2577–2591, 2018.
- Gray, M.: An investigation into convectively generated potential-vorticity anomalies using a mass-forcing model, *Quarterly Journal of the Royal Meteorological Society*, 125, 1589–1605, 1999.
- Gray, S. L., Dunning, C., Methven, J., Masato, G., and Chagnon, J. M.: Systematic model forecast error in Rossby wave structure, *Geophysical Research Letters*, 41, 2979–2987, 2014.
- 840 Grazzini, F. and Vitart, F.: Atmospheric predictability and Rossby wave packets, *Quarterly Journal of the Royal Meteorological Society*, 141, 2793–2802, 2015.
- Grazzini, F., Craig, G. C., Keil, C., Antolini, G., and Pavan, V.: Extreme precipitation events over northern Italy. Part I: A systematic classification with machine-learning techniques, *Quarterly Journal of the Royal Meteorological Society*, 146, 69–85, 2020.
- 845 Harvey, B., Methven, J., Sanchez, C., and Schäfler, A.: Diabatic generation of negative potential vorticity and its impact on the North Atlantic jet stream, *Quarterly Journal of the Royal Meteorological Society*, 146, 1477–1497, 2020.
- Harvey, B. J., Methven, J., and Ambaum, M. H.: Rossby wave propagation on potential vorticity fronts with finite width, *Journal of Fluid Mechanics*, 794, 775–797, 2016.
- Haynes, P. H. and McIntyre, M.: On the conservation and impermeability theorems for potential vorticity, *Journal of Atmospheric Sciences*, 47, 2021–2031, 1990.
- 850 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, 2020.
- Hertenstein, R. F. and Schubert, W. H.: Potential vorticity anomalies associated with squall lines, *Monthly weather review*, 119, 1663–1672, 1991.
- 855 Hoskins, B.: A potential vorticity view of synoptic development, *Meteorological Applications*, 4, 325–334, 1997.
- Iqbal, W., Leung, W.-N., and Hannachi, A.: Analysis of the variability of the North Atlantic eddy-driven jet stream in CMIP5, *Climate Dynamics*, 51, 235–247, 2018.
- Joos, H., Sprenger, M., Binder, H., Beyerle, U., and Wernli, H.: Warm conveyor belts in present-day and future climate simulations–Part 1: Climatology and impacts, *Weather and Climate Dynamics*, 4, 133–155, 2023.
- 860 Keller, J. H., Grams, C. M., Riemer, M., Archambault, H. M., Bosart, L., Doyle, J. D., Evans, J. L., Galarneau, T. J., Griffin, K., Harr, P. A., et al.: The extratropical transition of tropical cyclones. Part II: Interaction with the midlatitude flow, downstream impacts, and implications for predictability, *Monthly Weather Review*, 147, 1077–1106, 2019.
- Lee, J. H., Kim, J.-H., Sharman, R. D., Kim, J., and Son, S.-W.: Climatology of clear-air turbulence in upper troposphere and lower stratosphere in the Northern Hemisphere using ERA5 reanalysis data, *Journal of Geophysical Research: Atmospheres*, 128, e2022JD037 679, 2023.
- 865 Lee, S. H., Williams, P. D., and Frame, T. H.: Increased shear in the North Atlantic upper-level jet stream over the past four decades, *Nature*, 572, 639–642, 2019.

- Li, J., Feng, Z., Qian, Y., and Leung, L. R.: A high-resolution unified observational data product of mesoscale convective systems and isolated deep convection in the United States for 2004–2017, *Earth System Science Data Discussions*, 2020, 1–48, 2020.
- 870 Liu, N., Leung, L. R., and Feng, Z.: Global mesoscale convective system latent heating characteristics from GPM retrievals and an MCS tracking dataset, *Journal of Climate*, 34, 8599–8613, 2021.
- Lojko, A., Payne, A., and Jablonowski, C.: The Remote Role of North-American Mesoscale Convective Systems on the Forecast of a Rossby Wave Packet: A Multi-Model Ensemble Case-Study, *Journal of Geophysical Research: Atmospheres*, 127, e2022JD037 171, 2022.
- Lorenz, D. J. and Hartmann, D. L.: Eddy–zonal flow feedback in the Northern Hemisphere winter, *Journal of climate*, 16, 1212–1227, 2003.
- 875 Madonna, E., Wernli, H., Joos, H., and Martius, O.: Warm conveyor belts in the ERA-Interim dataset (1979–2010). Part I: Climatology and potential vorticity evolution, *Journal of climate*, 27, 3–26, 2014.
- Martin, J. E.: Quasi-geostrophic diagnosis of the influence of vorticity advection on the development of upper level jet-front systems, *Quarterly Journal of the Royal Meteorological Society*, 140, 2658–2671, 2014.
- Minobe, S., Kuwano-Yoshida, A., Komori, N., Xie, S.-P., and Small, R. J.: Influence of the Gulf Stream on the troposphere, *Nature*, 452, 880 206–209, 2008.
- Müller, A., Niedrich, B., and N  vir, P.: Three-dimensional potential vorticity structures for extreme precipitation events on the convective scale, *Tellus A: Dynamic Meteorology and Oceanography*, 72, 1–20, 2020.
- Oertel, A. and Schemm, S.: Quantifying the circulation induced by convective clouds in kilometer-scale simulations, *Quarterly Journal of the Royal Meteorological Society*, 147, 1752–1766, 2021.
- 885 Oertel, A., Boettcher, M., Joos, H., Sprenger, M., and Wernli, H.: Potential vorticity structure of embedded convection in a warm conveyor belt and its relevance for large-scale dynamics, *Weather and Climate Dynamics*, 1, 127–153, 2020.
- Oertel, A., Sprenger, M., Joos, H., Boettcher, M., Konow, H., Hagen, M., and Wernli, H.: Observations and simulation of intense convection embedded in a warm conveyor belt–how ambient vertical wind shear determines the dynamical impact, *Weather and Climate Dynamics*, 2, 89–110, 2021.
- 890 Orlanski, I. and Katzfey, J.: The life cycle of a cyclone wave in the Southern Hemisphere. Part I: Eddy energy budget, *Journal of Atmospheric Sciences*, 48, 1972–1998, 1991.
- O’Brien, T. A., Loring, B., Dufek, A. S., Islam, M. R., Kamnani, D., Quagraine, K. T., and Kirkpatrick, C.: Atmospheric rivers in the eastern and midwestern United States associated with baroclinic waves, *Geophysical Research Letters*, 51, e2023GL107 236, 2024.
- Pfahl, S., Schwierz, C., Croci-Maspoli, M., Grams, C. M., and Wernli, H.: Importance of latent heat release in ascending air streams for 895 atmospheric blocking, *Nature Geoscience*, 8, 610–614, 2015.
- Pohorsky, R., R  thlisberger, M., Grams, C. M., Riboldi, J., and Martius, O.: The climatological impact of recurving North Atlantic tropical cyclones on downstream extreme precipitation events, *Monthly Weather Review*, 147, 1513–1532, 2019.
- Prince, K. C. and Evans, C.: Convectively Generated Negative Potential Vorticity Enhancing the Jet Stream through an Inverse Energy Cascade during the Extratropical Transition of Hurricane Irma, *Journal of the Atmospheric Sciences*, 79, 2901–2918, 2022.
- 900 Prosser, M. C., Williams, P. D., Marlton, G. J., and Harrison, R. G.: Evidence for Large Increases in Clear-Air Turbulence Over the Past Four Decades, *Geophysical Research Letters*, 50, e2023GL103 814, 2023.
- Pyle, M. E., Keyser, D., and Bosart, L. F.: A diagnostic study of jet streaks: Kinematic signatures and relationship to coherent tropopause disturbances, *Monthly weather review*, 132, 297–319, 2004.
- Rasmussen, R., Chen, F., Liu, C., Ikeda, K., Prein, A., Kim, J., Schneider, T., Dai, A., Gochis, D., Dugger, A., et al.: CONUS404: The NCAR- 905 USGS 4-km long-term regional hydroclimate reanalysis over the CONUS, *Bulletin of the American Meteorological Society*, 2023.

- Rodwell, M. J., Magnusson, L., Bauer, P., Bechtold, P., Bonavita, M., Cardinali, C., Diamantakis, M., Earnshaw, P., Garcia-Mendez, A., Isaksen, L., et al.: Characteristics of occasional poor medium-range weather forecasts for Europe, *Bulletin of the American Meteorological Society*, 94, 1393–1405, 2013.
- Röthlisberger, M., Martius, O., and Wernli, H.: Northern Hemisphere Rossby wave initiation events on the extratropical jet—A climatological analysis, *Journal of Climate*, 31, 743–760, 2018.
- Rowe, S. M. and Hitchman, M. H.: On the role of inertial instability in stratosphere–troposphere exchange near midlatitude cyclones, *Journal of the Atmospheric Sciences*, 72, 2131–2151, 2015.
- Rowe, S. M. and Hitchman, M. H.: On the relationship between inertial instability, poleward momentum surges, and jet intensifications near midlatitude cyclones, *Journal of the Atmospheric Sciences*, 73, 2299–2315, 2016.
- Schultz, D. M., Schumacher, P. N., and Doswell III, C. A.: The intricacies of instabilities, *Monthly weather review*, 128, 4143–4148, 2000.
- Shaw, T. A. and Miyawaki, O.: Fast upper-level jet stream winds get faster under climate change, *Nature Climate Change*, 14, 61–67, 2024.
- Spreitzer, E., Attinger, R., Boettcher, M., Forbes, R., Wernli, H., and Joos, H.: Modification of potential vorticity near the tropopause by nonconservative processes in the ECMWF model, *Journal of the atmospheric sciences*, 76, 1709–1726, 2019.
- Steinfeld, D. and Pfahl, S.: The role of latent heating in atmospheric blocking dynamics: a global climatology, *Climate Dynamics*, 53, 6159–6180, 2019.
- Takaya, K. and Nakamura, H.: A formulation of a phase-independent wave-activity flux for stationary and migratory quasigeostrophic eddies on a zonally varying basic flow, *Journal of the Atmospheric Sciences*, 58, 608–627, 2001.
- Tenenbaum, J., Williams, P. D., Turp, D., Buchanan, P., Coulson, R., Gill, P. G., Lunnon, R. W., Oztunali, M. G., Rankin, J., and Rukhovets, L.: Aircraft observations and reanalysis depictions of trends in the North Atlantic winter jet stream wind speeds and turbulence, *Quarterly Journal of the Royal Meteorological Society*, 148, 2927–2941, 2022.
- Teubler, F. and Riemer, M.: Potential-vorticity dynamics of troughs and ridges within Rossby wave packets during a 40-year reanalysis period, *Weather and Climate Dynamics*, 2, 535–559, 2021.
- Thompson, C. F. and Schultz, D. M.: The release of inertial instability near an idealized zonal jet, *Geophysical Research Letters*, 48, e2021GL092649, 2021.
- Thompson, C. F., Schultz, D. M., and Vaughan, G.: A global climatology of tropospheric inertial instability, *Journal of the Atmospheric Sciences*, 75, 805–825, 2018.
- Trier, S. B. and Sharman, R. D.: Mechanisms influencing cirrus banding and aviation turbulence near a convectively enhanced upper-level jet stream, *Monthly Weather Review*, 144, 3003–3027, 2016.
- Volonté, A., Clark, P. A., and Gray, S. L.: The role of mesoscale instabilities in the sting-jet dynamics of windstorm Tini, *Quarterly Journal of the Royal Meteorological Society*, 144, 877–899, 2018.
- Weijenborg, C., Friederichs, P., and Hense, A.: Organisation of potential vorticity on the mesoscale during deep moist convection, *Tellus A: Dynamic Meteorology and Oceanography*, 67, 25705, 2015.
- Weijenborg, C., Chagnon, J., Friederichs, P., Gray, S., and Hense, A.: Coherent evolution of potential vorticity anomalies associated with deep moist convection, *Quarterly Journal of the Royal Meteorological Society*, 143, 1254–1267, 2017.
- Wernli, H. and Davies, H. C.: A Lagrangian-based analysis of extratropical cyclones. I: The method and some applications, *Quarterly Journal of the Royal Meteorological Society*, 123, 467–489, 1997.
- Wilks, D.: “The stippling shows statistically significant grid points”: How research results are routinely overstated and overinterpreted, and what to do about it, *Bulletin of the American Meteorological Society*, 97, 2263–2273, 2016.

- Winters, A. C.: Kinematic processes contributing to the intensification of anomalously strong North Atlantic jets, *Quarterly Journal of the Royal Meteorological Society*, 147, 2506–2532, 2021.
- 945 Woollings, T., Papritz, L., Mbengue, C., and Spengler, T.: Diabatic heating and jet stream shifts: A case study of the 2010 negative North Atlantic Oscillation winter, *Geophysical Research Letters*, 43, 9994–10, 2016.