

Global scenarios of anthropogenic mercury emissions

Flora Maria Brocza^{1,2}, Peter Rafaj¹, Robert Sander¹, Fabian Wagner¹, Jenny Marie Jones²


¹International Institute for Applied Systems Sciences, Schlossplatz 1, 2361 Laxenburg, Austria

²School of Chemical and Process Engineering, University of Leeds, LS2 9J, United Kingdom

5 *Correspondence to:* Flora M. Brocza (brocza@iiasa.ac.at)

Abstract. Anthropogenic mercury (Hg) emissions to the atmosphere are a long-lived hazard to human and environmental health. The UN Minamata Convention on Mercury is seeking to lower anthropogenic mercury emissions through a mix of policies from banning ~~certain Hg uses and trade~~, to reducing unintentional Hg releases from different activities. In addition to independent Hg policy, ~~strategies to mitigate greenhouse gases~~, particulate matter (PM) and SO₂ ~~reduction policies~~ may also
10 lower Hg emissions as a co-benefit. This study uses the Greenhouse Gas – Air Pollution Interactions and Synergies (GAINS) model to examine the effect of different clean air and climate policy on future global Hg emissions. The Baseline scenario assumes current ~~trends for~~ energy use and Hg emissions, as well as current legislation for clean air, mercury and climate policy. In addition, we explore the impact of the Minamata Convention, co-benefits of climate policies and of stringent air pollution policies, as well as a maximum feasible reduction ~~measures scenario~~ for Hg (~~Hg-MFR~~). Hg emission projections until 2050
15 show noticeable reductions in combustion sectors for all scenarios, due to a decrease in global fossil fuels and traditional biomass use, leading to emission reductions of 33% ~~in the~~ (Baseline) up to 90% when combining stringent climate and ~~most efficient Hg-controls~~MFR. Cement and non-ferrous metal emissions increase in all ~~activity~~-scenarios with current air pollution policy, but can be reduced by up to 72% and 46% respectively in 2050 with stringent Hg-specific measures. Other emissions (including waste) are a ~~significant~~large source of uncertainty in this study, and ~~their~~ projections range between a 22% increase
20 and 54% decrease in 2050 depending on both climate and clean air policy. The largest absolute reduction potential for Hg abatement, but also the largest uncertainties of absolute emissions lie in ~~the in~~ small-scale and artisanal gold production, where ~~Hg-specific~~ abatement measures could eliminate annual Hg emissions in the range of 601-1371 t (95% confidence interval), ~~but which uncertainties in the estimate are so high that they might eclipse reduction efforts in all other sectors~~. 90% of the Hg emissions ~~in GAINS are~~ covered by ~~provisions of the~~ the Minamata Convention. Overall, the findings emphasize the necessity
25 of implementing targeted Hg control policies in addition to stringent climate, PM and SO₂ policies to achieve significant reductions in Hg emissions.

Graphical Abstract.



30

Abbreviations

- ASGM Artisanal and small-scale gold mining
- BAS Baseline scenario
- BAT Best available technology
- 35 BEP Best environmental practice
- CLE Current legislation
- CLIM1 Climate policy scenario
- CLIM2 Net-zero scenario
- co-MFR co-benefits for Hg from PM, SO₂ MFR
- 40 EC Emission control
- EF Emission factor
- EU-IED Industrial Emissions Directive of the European Union
- GDP Gross domestic product

	GMA'18	Global Mercury Assessment ((AMAP/UNEP 2019))
45	Hg-MFR	Maximum feasible reduction for Hg
	MEX	Market exchange rate
	MFR	Maximum feasible reduction
	MCM	Minamata Convention on Mercury
	Mina	Minamata policy scenario
50	NAP	National Action Plan
	NFME	Non-ferrous metals
	NOC	No Control Scenario
	PM	Particulate matter
	PPP	Purchasing power parity
55	POP	Population
	SRES	IPCC Special Report on Emission Scenarios
	UEF	Unabated emission factors
	VCM	Vinyl chloride monomer production
	WEO	World Energy Outlook
60	WHO	World Health Organization

1 Introduction

Mercury (Hg) is one of the top ten chemicals of major public health concern designated by the World Health Organization (WHO) (WHO, 2021) (ref). Hg has a high (redox-)reactivity at ambient conditions, facilitating frequent species changes. Elemental mercury (Hg⁰) exhibits high volatility and vapor pressure that are unique for a metal and the metal's unique volatility and (redox-)reactivity at ambient conditions facilitate frequent Hg species changes, leading to long-range atmospheric transport, subsequent deposition and re-emission of the metal and its derivative compounds, as well as methylation and subsequent bioaccumulation of the most toxic Hg species, methyl mercury, in the aquatic food chain as methyl mercury (Selin, 2009). Mercury has been known to be highly toxic to humans since the Minamata Disease tragedy in the 1950s. The extent of the pollution and health problems caused by atmospheric Hg emissions has been known for two decades (e.g. UNEP Chemicals 2002) and increasing awareness of the global dimensions of the Hg problem led to its explicit inclusion in the Aarhus protocol on heavy metals in 1998, as part of the CLRTAP convention (Selin and Selin, 2006; CLRTAP, 1979; Aarhus Protocol, 1998). In 2015, Cumulative anthropogenic emissions have increased the Hg content in the atmosphere by 450% above natural levels (AMAP/UNEP, 2019). The time for mercury to return to a permanent sink such as deep ocean sediments has been estimated as up to 3000 years (Selin, 2009), demonstrating that Hg pollution will continue to pose a serious environmental threat for years to come, but also highlighting that today's action will have a long-lasting effect to reduce levels of environmental Hg

Formatted: Superscript

Formatted: Not Superscript/ Subscript

Formatted: Not Highlight

[pollution](#) (Angot et al., 2018). From a health perspective, it has been estimated that accumulated health effects of Hg pollution will cost \$19 trillion globally between 2010 and 2050 (2020 dollars; Zhang et al., 2021), further demonstrating the importance of fast action.

80 To break the cycle of ~~Hg emissions to air, releases into the environment, and subsequent emissions, re-emissions and build-up of Hg pollution in the environment heightening pollution~~, the Minamata Convention on Mercury (MCM) has been adopted in 2013. It entered into force in 2017 and is ~~presently~~ ratified by 147 countries ~~as of January 2024~~ (UNEP, 2013). The first international health and environment treaty on hazardous substances in almost a decade, it recognizes that Hg emissions must be tackled urgently at the global level. The MCM aims to ~~“protect the human health and the environment from anthropogenic~~
85 ~~emissions and releases of mercury and mercury compounds”~~ ~~reduce releases of “mercury and mercury compounds”~~ by targeting ~~those emissions to air and releases to the environment them at different entry-points/levels of the release cycle~~, such as trade, use in production, use in products, emission sources, and wastes. Mercury ~~pollution releases to the atmosphere and environment is/are~~ on one hand addressed by technical solutions, such as ~~limiting emissions and releases through~~ best available technology / best environmental practice (BAT/BEP) recommendations for Hg handling, industrial emissions or waste storage.
90 On the other hand, ~~they require political and there are provisions for~~ regulatory action ~~on other domains~~, such as ~~severely limiting primary mercury mining and bans on~~ mercury trade, ~~bans on~~ specific products, and small scale or traditional (artisanal) gold mining practices, demonstrating a “life-cycle approach” to limiting Hg emissions (e.g. Selin, 2014; ~~UNEP, 2013; Giang et al., 2015~~). Despite these efforts, global anthropogenic emissions of Hg were estimated to have risen by 20% by 2015 compared to “pre-Minamata” 2010 levels. Small emission decreases in North America and the EU were offset by a mix of
95 increased economic activity, as well as the production, use and disposal of mercury-containing products (AMAP/UNEP, 2019; Pacyna et al., 2016). For a better understanding of future Hg levels in the atmosphere, scenarios of future anthropogenic Hg releases are needed. Such scenarios need to consider the wide range of Hg emission sources, their emission intensity, ~~abatement potential~~, as well as their drivers.

Where Hg is emitted in energy-intensive sectors, such as from the combustion of fossil fuels or different industries, future
100 emissions strongly depend on the assumptions on future energy demand and the decarbonization of those sectors. Emission trends from other sectors, such as waste generation, ~~can be/are~~ derived from macroeconomic factors and population growth. Other activities, such as artisanal and small-scale gold production, are specific to mercury pollution and ~~not influenced/unaffected~~ by other air quality and climate policy. Emission intensity is always specific to the emission source, its geographic location and the application of control measures which lower the amount of Hg released into the atmosphere. Such
105 measures include policies or technologies targeted directly at Hg. In addition, stringent clean air policy targeted at reducing particulate matter (PM), SO₂ and NO_x is well known to lower Hg emissions, as the applied pollution control technologies interact with the mercury present in the flue gas streams and are able to retain it (e.g. Granite et al. 2000, Pavlish et al. 2003). Scenario analysis is a powerful tool to quantify future pathways of anthropogenic mercury emissions and to understand interdependencies of various mitigation factors.

110 Only a small number of studies have produced global scenarios of speciated future mercury emissions; Streets et al. 2009
created a Hg inventory spanning different combustion and industry sectors as well as artisanal and small-scale gold mining
(ASGM)¹. These emissions were projected to 2050 based on four climate scenarios from the IPCC's Special Report on
Emission Scenarios (SRES) (Nakicenovic et al., 2000ref). The SRES is also used as a source of different energy scenarios for
Hg projections by Lei et al. 2014. Rafaj et al. (2013) present a set of global Hg projections for one baseline and one climate
115 scenario based on the POLES energy modelWorld Energy Outlook 2012 and cover somekey Hg-specific sectors includingsueh
as gold and caustic soda production; Pacyna et al. (2010); project Hg emissions up to 2020 based on different scenarios focusing
on Hg-specific policies. Pacyna et al. (2016); projected Hg emissions up to 2035 based on the GMA'13 inventory and own
projections. They; lookeding at a mix of scenarios including current legislation, new policies and a maximum feasible reduction
scenario with and a 450 ppm CO₂-climate scenario, suggesting that emissions would stay constant at the 2010 level of 1960t
120 by 2035 under current legislation, decrease to 1020t in the 'New Policies' scenario and project 300t Hg emissions in 2035 in
the MFR case. Additionally, several regional studies are available for China (Giang et al., 2015; Zhao et al., 2015; Ancora
et al., 2016; Wu et al., 2018a,b; Mulvaney et al., 2020), India (Chakraborty et al., 2013; Giang et al., 2015) and Europe (Pacyna
et al., 2006; Glodek et al., 2010 (Poland); Rafaj et al., 2014). The base years for the global scenario studies lie between 2000
and 2010 and the projection years are between 2020 and 2050. The most recent Global Mercury Assessment 2018 (GMA'18)
125 was published for 2015 and includes significant data quality improvements compared to the 2010 inventory, also including
quantification of more emission sources. Similarly, our outlook on energy and climate scenarios has significantly changed
since the COVID-19 pandemic and recent geopolitical developments. Up-to-date scenarios – as reported in this paper – arewith
be needed to understand future mercury emissions and the effectiveness of clean air and climate policy on curbing them.

130 IIASA's Greenhouse Gas and Air Pollution Interactions and Synergies (GAINS) model is uniquely suited for the creation of
global mercury emission scenarios (Amann et al. 2011). Originally built to inform policy questions regarding acid rain and
particulate matter, its database was extended to Hg in 2013 (Rafaj et al., 2013; Rafaj et al., 2014). In GAINS, sector- and
region-specific control strategies represent the pollution control measures and policies which are in place. Timing and
stringency ofDevelopments and policy infor air pollution control, greenhouse gas reduction strategies, as well as co-benefits
135 from PM and SO₂ abatement and changes in the energy and industrial sectors are represented for each of the 182 GAINS
regions. This study presents a recent update on the methodology of accounting for co-benefit control of Hg from PM and SO₂
in the GAINS model. Furthermore, Ddedicated mercury control options and their combinations were also updated in the
modelling framework. We presentdemonstrate the results of scenario analysis using three different energy/climate pathways
combined with four different scenarios of mercury control measures, including mercury control options consistent with current
140 legislation and Minamata commitments, as well as a maximum feasible reduction scenario. Scenarios were designed to identify

¹ The Minamata Convention further defines ASGM as 'gold mining conducted by individual miners or small enterprises with limited capital investment and production' (Minamata Convention on Mercury, 2013)

the impact of climate policy, co-benefits from air pollution control policy, and dedicated Hg measures and are presented on the level of 7 world regions and 8 exemplary sub-regions.

2 Modelling Framework

2.1 The GAINS model

145 The GAINS model quantifies emissions to the atmosphere, costs, [ecosystems-](#) and ~~(health)-~~impacts of different strategies to reduce different air pollutants and greenhouse gases (Amann et al., 2011). GAINS computes Hg emissions on a global level up to 2050 in 5-year time steps with a resolution of 182 regions. A region represents either a country, a group of neighboring countries or sub-national regions. Current and future emissions of mercury (E^{Hg}) are computed via equation (1) for each mercury species (f) from activity data (A) of different [combinations of](#) pollution sources (s) and ~~activity combinations-(p,s)~~,
150 and uncontrolled emission factors (e), which are lowered by taking into account the removal efficiencies (r) of different emission control technologies and other measures (t) and their application rates (x) in a specific sector [over time](#):

$$E^{Hg} = \sum_f E_{p,s,t} = \sum_{f,p,s,t} A_{p,s} e_{p,s} (1 - r_{f,t}) x_{p,s,t} \quad (1)$$

An illustration of this is provided by Figure 1. In total, 13 fuel types used in 52 combustion sectors and 22 non-combustion
155 emission sources are covered in GAINS, as summarized in the appendix (Tables S1, S2)². For the purpose of this study, GAINS sector-activity combinations have been grouped into the main emission sources for Hg, described in Table 1. Similarly, the 182 GAINS regions are grouped into 7 main regions (Africa, Asia Pacific, Central and South America, Europe, Eurasia, Middle East, North America) [consistent in accordance](#) with the IEA World Energy Outlook (2022). Additionally, [emission projections for](#) Brazil, China, the European Union, India, Japan, Russia, Southeast Asia and the USA are computed separately as sub-
160 regional case studies (see Table S10 in the appendix).

² The full list of activities, sectors and technologies in GAINS can be found in the GAINS glossary: <https://gains.iiasa.ac.at/gains/GOD/abbreviations.info>




Figure 1: Schematic of multi-pollutant control technology application in GAINS.

2.2 Emission Factors

Uncontrolled emission factors (UEFs) are derived from literature sources and are specific to each sector-activity combination (see Table 1). Emission factors for hard coal and brown coal combustion (power sector, industries and buildings) remain unchanged from previous versions of GAINS (Rafaj et al., 2013; Rafaj et al., 2014). Factors for the production of cement, non-ferrous metals, aluminum, iron and steel, gold, and caustic soda were updated in accordance with the Global Mercury Assessment 2018 (‘GMA’18; AMAP/UNEP, 2019). The non-ferrous metals (NFME) sector in GAINS includes emissions from copper, lead, nickel and zinc smelters. Metal-specific emission factors (either country-specific or generic, depending on data availability) were weighted with the share of the relevant metals of the total activity for a particular GAINS region, based on USGS Mineral Yearbook production data (Klochko, 2021; Flanagan, 2022; Tolcin, 2022). The shares were calculated for 2015 and assumed to be static, so this composite emission factor was subsequently applied to all years for a particular region. Total gold production was similarly based on USGS Mineral Yearbook data. This data was split into country-level shares of large-scale (sector ‘GOLD’) and artisanal/small-scale gold mining (sector ‘ASGM’), based on data from the World Gold Council as presented in GMA’18. Hg mining and vinyl chloride monomer production are not specified explicitly in the model

and no control measures ~~are~~ applied to them directly, but their emissions are included in the aggregate category “Other Hg emissions” on a region-by-region basis and their activities reflect projected sectoral emissions (George, 2021). In the waste sector, Hg emissions were derived from the GMA’18 and attributed evenly to industrial, rural and urban waste categories.

Information on the average speciation of emissions from each source ~~is also considered~~ ~~are also implemented~~, dividing total unabated emissions into shares of Hg⁰, Hg^{II} and particulate Hg (Hg_p). Due to a lack of regional data, ~~identical~~ speciation data was implemented for each sector-activity combination on a global level. The values represent the best available literature data and modelled speciation from the [Interactive Process Optimization Guidance \(iPOG™\)](#) ~~iPOG-tool~~ (Niksa Energy Associates LLC, 2011) at the time of writing, but care has to be taken in their interpretation, as they are associated with large uncertainty.

Table 1 summarizes the ranges of unabated emission factors used in the GAINS model for aggregated Hg-relevant sectors, as well as the Hg⁰ / Hg^{II} / Hg_p of uncontrolled stack emissions.

Table 1: Ranges of unabated emission factors (UEF) and speciation in GAINS on the global scale. UEFs vary on a regional scale, as well as due to different, aggregated ~~s~~ for sector-fuel combinations. Bold categories represent the sector aggregation level that are plotted in the results figures (Figs. 3, 4, 5, S1).

Sector	Abbreviation (as in Fig. 7)	Emission factor min - max (unit)	Speciation inlet Hg ⁰ / Hg ^{II} / Hg _p	Sources
<i>COMBUSTION - by sector</i>				
Combustion in power plants	COMB_POWER	0.0001 - 0.0477 (t/PJ)	50-60 / 30-40 / 10	Rafaj et al. (2013)
Industrial combustion	COMB_IND	0.0001 - 0.063 (t/PJ)	20-60 / 30-60 / 10-20	Rafaj et al. (2013)
Other combustion (Residential, service, conversion sectors)	COMB_OTHER	0.0001 - 0.0477 (t/PJ)	20-60 / 30-60 / 10-20	Rafaj et al. (2013)
<i>COMBUSTION - by fuel</i>				
All coal grades		0.0005 - 0.0477 (t/PJ)	50-60 / 40-60 / 10-20	Rafaj et al. (2013)
Gasoline		0.0001 (t/PJ)	50 / 40 / 10	Rafaj et al. 2013
Liquid fuels		0.0001 - 0.0005 (t/PJ)	50 / 40 / 10	Rafaj et al. (2013); GMA'18
Biomass		0.001 (t/PJ)	50 / 40 / 10	Rafaj et al. (2013); GMA'18
Waste		0.063 (t/PJ)	20 / 60 / 20	Own estimate, derived from GMA'18

INDUSTRY

Non-ferrous metals (Cu, Zn, Pb, Al)	NFME	0.0002 - 117.84 (g/t)	80 / 15 / 5	GMA'18
Large-scale gold	GOLD	12000 – 55000 (g/t)	80 / 15 / 5	GMA'18
Artisanal and small-scale gold	ASGM	975000 – 1500000 (g/t)	100 / 0 / 0	GMA'18
Cement production	CEM	0.022 - 0.124 (g/t)	80 / 15 / 5	GMA'18
Other industrial processes	IND_PROC	0.00025 – 20 (g/t)	70-80 / 15-30 / 0-5	
Iron and steel production		0.0061 - 0.41475 (g/t)	80 / 15 / 5	Wang et al. (2016) ; GMA'18
Oil Refining		0.0003 – 0.0166 (g/t)	80 / 15 / 5	GMA'18
Caustic Soda Production		2.5 – 20 (g/t)	70 / 30 / 0	GMA'18
OTHERS				
Cremation	OTHER	2 – 2.5 (g/Million)	80 / 15 / 5	Rafaj et al. (2013)
Waste	OTHER	0.0315 (g/t)	96 / 4 / 0	GMA'18
VCM production, Hg mining	OTHER	1 (t Hg/year)	100 / 0 / 0	GMA'18
Transport	OTHER	0.0001 – 0.063 (t/PJ)	20-60 / 30-60 / 10-20	Own estimate, derived from GMA'18

Notes: GMA'18 ... AMAP/UNEP (2019)

190 2.3 Control Technologies

2.3.1 Mercury control measures

195 A review of Hg control technologies and measures has been conducted. Relevant technologies have been implemented into the GAINS model in addition to previously available co-benefit controls from PM and SO₂ abatement. Such new controls include: the option of low-mercury or halogen-treated coal (LHGCO); sorbent injection (such as activated carbon) with or without an additional baghouse filter (SINJ); acid plants for the non-ferrous metal and gold sectors (PR_AP); and stationary sorbent modules (SPC), which represent the possibility of removing Hg not only from the atmosphere but bringing it into a permanent sink such as a controlled hazardous waste landfill, rather than re-directing emissions into other environmental releases. Removal efficiencies and Hg speciation of the control technologies operating on Hg are summarized for each emission source category in the appendix (Tables S3-S6).

200 2.3.2 Quantification of co-benefits for mercury from particulate matter and SO₂ control

GAINS has been used extensively to inform policies on the reduction of [air pollutants such as](#) particulate matter (PM), SO₂ and NO_x, whose abatement is known to strongly influence Hg emissions and their speciation. To compute impacts of traditional air pollution control devices on total reduction of mercury in the GAINS model, current and projected control strategies of PM and SO₂ are considered in addition to Hg-specific control measures³. The concept of ‘overlapping control measures’ has already
205 been introduced in an earlier publication (Rafaj et al., 2013). Where Hg-relevant PM and/or SO₂ [controls](#) co-exist in a region, sector and year, their compounded impact [on](#) Hg emissions is considered, increasing Hg removal efficiency for the relevant portion of installations. Where appropriate, combinations of Hg-specific measures and PM/SO₂ measures also lead to increased Hg removal efficiency. The relevant technology combinations are listed in Tables S4-S6 [together with their](#)~~and have~~ sector- and activity- specific removal efficiencies. Relevant technology combinations are [defined](#) in the power and industry sectors
210 ~~between-as~~ different [combinations of](#) particle filters and flue gas desulphurization ~~/-and-acid plants-in-industrial-processes~~. Lastly, there has been a significant update of the representation [of](#) emissions from waste in GAINS in the past years (Gómez-Sanabria et al., 2022). Control measures in the GAINS waste sector are multi-pollutant controls representing different types of landfill and other waste management options, not all of which can be [directly](#) linked to reduced atmospheric Hg emissions. In this modelling work, the following three measures [found in literature review](#) have been associated with Hg removal
215 ~~efficiencies, as Hg reduction can be expected from literature review~~: waste incineration with energy recovery and pollution controls (emissions accounted for in the power sector), landfill compression, [and](#) landfill covering. Details can be found in the appendix, Table S6.

2.3.3 Effect of control measures on Hg speciation


220 Mercury speciation is also altered through the application of control measures. The speciated removal efficiency of the controls operate on the speciated no-control emissions. The resulting total Hg emissions are then split again into Hg⁰ / Hg^{II} / Hg_p, based on reported stack speciated emissions of the technology from literature review. Tables of all applied control technologies, their speciated removal efficiencies and resulting emission speciation can be found in the appendix (Tables S3-S6).

3 Activity Projections

225 GAINS uses exogenous projections of anthropogenic activities and energy use to estimate future Hg emissions. For this study, three scenarios of energy and industrial production until 2050 were implemented in GAINS, based on trends reported by the

³ Co-benefits ~~from~~ NO_x ~~controls~~ were not [considered in this exercise because](#)~~implemented as NO_x control~~, when combined with PM and SO₂ ~~abatement techniques~~~~control~~, ~~has~~~~ve~~ been reported to only bring Hg removal efficiency improvements of a few percent lower than the standard deviation of the ~~constructed~~-technology categories used in this study (see e.g., Li et al. 2020 (SI, Table S10), or the iPOG tool at reference conditions (Niksa Energy Associates LLC, 2011).

World Energy Outlook 2022 (IEA, 2022). The three scenarios share assumptions on macroeconomic drivers (GDP growth, GDP per capita and annual population growth). They differ in their assumption on the stringency of climate policy and already include first effects of the Russian war in Ukraine. While the total, global energy demand either rises or stays similar to 2015 levels until 2050 in all three scenarios, there are differences in the energy sources which meet this demand (see Fig. 2). The Baseline (BAS) scenario represents developments in the energy sector leading to a 2.5C average global temperature rise by 2100 and is characterized by plateauing emissions in at 37 Gt and a reduction to 32 Gt energy-related CO₂ emissions in 2050. The demand growth is mostly met by renewable sources and the share of fossil fuels falls to 60% in 2050. The Climate Policy (CLIM1) scenario is consistent with a global average temperature rise of 1.7 C by 2100. Demand for all fossil fuels already declines by 2030, leading to a decrease in CO₂ emissions to 12 Gt by 2050. The Net Zero (CLIM2) scenario is the only scenario leading to <1.5 C temperature rise until 2100 (IEA, 2022). Although both climate scenarios are very if not unrealistically ambitious, they provide a scope to quantify a Hg-reduction potential induced through a rapid decarbonization of the global energy system. The scenarios are summarized in Table 2.



240 **Figure 2: Key activity data for the BASELINE, CLIM1 and CLIM2 scenarios. COMB_POWER ... Power plants, COMB_IND ... Combustion in industry, COMB_OTHER ... Other combustion (residential, commercial, conversion losses), IND_PROC ... Industrial Processes, CEM ... Cement, NFME ... Non-ferrous metal production, ASGM ... Artisanal and Small-scale gold mining,**

POP ... Population, GDP ... Gross domestic product, MEX ... Market exchange rate, PPP ... Purchasing power parity. Projections of fuel use (COMB and different fuels), transport, IND_PROC, CEM, NFME, POP & GDP from the World Energy Outlook 2022 (IEA 2022). Waste data from Gomez-Sanabria et al. 2022. PGOLD and ASGM ... this study (see Section 3.)

In addition, activities of Hg-specific sectors⁴ have been modelled in GAINS as follows. Gold production volumes have been cross-checked between the USGS Mineral Yearbook data (Sheaffer, 2022) and reports on ASGM from the World Gold Council via GMA'18, based on which shares between small-scale and large-scale gold production were calculated for 2020. Where the ASGM amounts in a specific country were larger than reported total gold production by the USGS, this higher number was taken and the share of ASGM was assumed to be 1. Shares between ASGM and large-scale gold production were then taken as fixed for all past and projection years. Gold production and ASGM activity have been projected into the future following the conservative assumption that levels will stay the same to 2020 levels, only to be modified by changing Hg policy through the control strategy (e.g., a ban on ASGM).

Caustic soda production from chlor-alkali plants using mercury cells has been adopted from the Rafaj et al. (2013) implementation of Hg in GAINS; phase-out of this technology is imminent, mandated through the Minamata convention, and no updates were necessary. Similarly, the methodology on estimating cremation emissions has been previously described in Rafaj et al. 2013. Activity projections for vinyl chloride monomer (VCM) production and Hg mining are only represented implicitly in the "OTHER_HG" sector. The MCM mandates that the Hg intensity of VCM production needs to be reduced by 50% in 2020, relative to production in 2010. For this study, 2015 VCM production values were assumed to be constant based on the data reported in the GMA'18, but the Hg emission intensity was adjusted phase-outs of the activity as mandated by the Minamata convention were applied to diminish activity projections.

The following mercury emission sources are not included in any GAINS sector for the current study: open savannah and forest burning, coal bed fires, and intentional mercury use in batteries, lamps, or other devices.

4 Scenarios

4.1 Control Strategies

Combined with the three energy and activity pathways listed in Section 3, scenarios of mercury control measures were devised. They span the full range of possible anthropogenic mercury emissions to the atmosphere – from the complete absence of control measures – to current legislation, all the way to maximizing either co-benefit controls and applying stringent Hg-specific controls where this is feasible (see Table 2).

⁴ 'Hg-specific sector' means: a sector which, in the GAINS model, is only associated with Hg emissions (e.g. no PM, SO₂ or other emissions)

Table 2: List of scenarios in this paper. The presented seven scenarios vary in two elements: Activity data incorporates assumptions on energy and climate policy, pollutant control strategies incorporate different [assumptions on scenarios](#) of clean air policy.

Activity data (Energy/Climate policy)	Pollutant control strategy (Clean Air policy)	Scenario ID
<p>Baseline scenario (BAS): Adapted from the WEO 2022 Stated Policies (STEPS) scenario (IEA, 2022). Global energy demand growth met mostly by renewables; share of fossil fuels in global energy mix falls to < 75% by 2030, 60% by 2050. Global energy-related CO₂ emissions plateau at 37 Gt and fall to 32 Gt in 2050, leading to 2.5° C global average temperature rise by 2100.</p>	<p>No control (NOC): Hypothetical baseline of unabated emissions. No PM, SO₂ or Hg controls implemented.</p>	00_BAS_NOC
	<p>Current legislation (CLE): Current legislation for Hg, PM and SO₂</p>	01_BAS_CLE
	<p>CLE + Minamata scenario (MINA): CLE for PM and SO₂; full implementation of Minamata BAT/BEP technologies and process phase-outs, as well as National Action Plans (NAPs) for ASGM</p>	02_BAS_MINA
	<p>Co-benefit control for Hg, maximum feasible reduction for PM and SO₂ (co-MFR): Maximum co-benefits from PM and SO₂ controls for Hg emissions in all sectors; no additional Hg-specific controls.</p>	03_BAS_coMFR
	<p>Maximum Feasible Reduction for Hg (Hg-MFR): Application of the most efficient Hg control implemented in the model for each GAINS sector.</p>	04_BAS_HgMFR
<p>Climate Policy scenario (CLIM1): Adapted from WEO 2022 Announced Pledges (AP) scenario (IEA, 2022). Demand for all fossil fuels declines by 2030. CO₂ emissions fall to 12 Gt in 2050, leading to 1.7 ° C global average temperature rise by 2100.</p>	<p>Current legislation (CLE): Current legislation for Hg, PM and SO₂</p>	05_CLIM1_CLE
<p>Net Zero scenario (CLIM2): Adapted from WEO 2022 Net Zero Emissions scenario (IEA, 2022). CO₂ emissions fall to 23 Gt in 2030 before reaching 0 Gt in 2050, leading to < 1.5 ° C in 2100.</p>	<p>Current legislation (CLE): Current legislation for Hg, PM and SO₂</p>	06_CLIM2_CLE
	<p>Maximum Feasible Reduction for Hg (Hg-MFR): Application of the most efficient Hg</p>	07_CLIM2_HgMFR

	control implemented in the model for each GAINS sector.	
--	---	--

275 The **No Control scenario (NOC, as in 00_BAS_NOC)** represents the complete absence of control measures for any pollutant – unabated emission factors are ~~applied~~ for all emission sources and all years. It displays higher than actual emissions, serving as a hypothetical end point that shows the efficacy of current legislation.

The **Current Legislation (CLE, as in 01_BAS_CLE, 05_CLIM1_CLE, 06_CLIM2_CLE)** control strategy represents existing and planned air pollution control policy for all pollutants implemented in GAINS (e.g. Rafaj et al. (2018), Amann et al. (2020) for global control strategies; Li et al. (2019) for China). Of these, the PM and SO₂ control strategies directly influence Hg emissions. In addition to co-benefit controls, Hg-specific controls were added in the extended modeling framework for this study; Hg-specific control measures were added for the cremation and waste incineration sectors in Europe and control measures for non-ferrous metal production were adjusted to acid plants in line with existing legislation (e.g. the European Union Industrial Emissions Directive (Directive 2010/75/EU), Indian emissions regulations (CPCB, 1998)). Table S7 in the appendix summarizes all Hg-specific control strategy changes.

For the **Minamata scenario (MINA, as in 02_BAS_MINA_CLE)**, the CLE control strategy was extended by information from the available Minamata National Action Plans (NAP) for ASGM. Targets for good practice or elimination of mercury use in this sector were collected on the country level, then aggregated into the GAINS regional levels and then WEO regional level (Table S7)⁵. It is important to note that this scenario does not include any assumptions on the implementation of emissions reductions pursuant to MCM Article 8 beyond co-benefits from current legislation.

Scenarios combined with Maximum Feasible Reduction (MFR) ~~strategies~~ scenarios, assume the implementation of currently available emission reduction technologies that achieves the lowest air pollution emission factors. Such scenarios have been computed for pollutants including PM and SO₂ in GAINS using optimization procedures (e.g. Amann et al. (2011) and Wagner et al. (2013)). An MFR scenario with maximized PM and SO₂ controls, but CLE Hg controls ~~generated~~. ~~This scenario~~ is called the **co-benefit MFR scenario (co-MFR, as in 03_BAS_coMFR)** and it simulates the maximum Hg reduction that can be achieved without Hg-specific measures, solely through co-benefits from air pollution abatement policy.

Lastly, to demonstrate the end point of the control measures represented in GAINS, the Hg-specific **maximum feasible reduction scenario (Hg-MFR, as in 04_BAS_HgMFR, 07_CLIM2_HgMFR)** was generated. It represents the full application of the APCDs and Hg-specific control measures (or their combinations) with the highest removal efficiency for each sector and each GAINS region in 2050. For combustion of coal, ~~oil products~~ ~~heavy fuel oil~~ ~~diesel~~ and waste, as well as ~~for~~ most industrial processes including gold production, ~~these are~~ Hg-specific control measures are applied. In sectors with

⁵ It is important to note that at the time of writing, not all ASGM-producing countries have published NAPs, meaning that ~~likely~~, not all ASGM reduction targets ~~might~~ can be fully represented in this scenario as of yet and that Hg reductions in this sector will likely be larger than estimated in this work once all NAPs are ~~enforced~~ published.

low emission factors where no Hg-specific controls are currently ~~commercially applicable~~ such as ~~road transport~~ or biomass combustion domestic/residential fuel combustion, the co-benefit control with the greatest removal efficiency for Hg was applied. Activities where a ban is a viable policy option, such as ASGM, are banned. The only exceptions are Hg mining and VCM production, as they are implemented indirectly in GAINS (see the discussion in section 3 of 'HG_OTHER') and do not have control measures applied to them, but ~~the~~ activities ~~rather reflect~~~~represent an adoption of~~ current Minamata policies. For the waste sector, ~~the~~ multi-pollutant waste management controls were applied, the most Hg-efficient of which is incineration, coupled with sorbent injection before an additional fabric filter (FFSINJ). Table S8 in the SI ~~list~~~~summarizes~~ the MFR control strategy in the year 2050.

4.2 Uncertainty of scenario results

Uncertainties for the aggregated sectors were estimated using a Monte Carlo Simulation approach ~~(ref?)~~. Uncertainties for uncontrolled emissions (scenario 00_BAS_NOC) were modelled by varying unabated emission factors and activity based on uncertainty estimates, ~~given in Table S11~~. The Monte Carlo Simulation was conducted by varying UEFs and activity data on the most granular GAINS resolution (182 regions, all sectors as listed in Tables S1, S10). The results were then aggregated to the regional and sectoral level used in this study ~~and are given in Table S12~~. Assuming that the abated emission factors, which take into account Hg emission reductions through abatement measures, have the same levels of uncertainty as the UEFs, the calculated relative uncertainties are applied to 01_BAS_CLE (see Figure S1).

5 Results and discussion

5.1 The Baseline and current legislation scenario (01_BAS_CLE)

The global emissions trajectory for the baseline scenario of this study (01_BAS_CLE, as displayed in Table 3 and in Fig. 3, compared with other control strategies), sees a slight increase in Hg emissions until 2050 to 109.6% of 2015 levels. Decreases from power generation, residential combustion and a small decrease in non-ferrous metal production are offset by emissions from waste and industrial emissions during both combustion and production processes. However, much of the increase is due to the increase in reported gold production between 2015 and 2020, ~~thereafter~~~~since~~ it is assumed that gold production is constant from 2020 to 2050. ~~As a result, so~~ 75% of the increase ~~in Hg emissions by 2050~~ is due to ~~the near-term~~~~near-term~~ ~~growth in~~ ASGM. Similarly, emissions of Hg from waste sector increase as they are driven by projected population increases (see Fig. 3). If these highly uncertain estimates are discounted, the combustion, metallurgy and other processes sectors reduce their emissions slightly, by 80 t/year until 2050. On a regional level, these trends are largely confirmed, but depend strongly on the dominating emission sectors and assumed controls in each region as plotted in Figure 4.

In **Europe**, emissions decrease by 18%, largely due to a reduction in Hg from combustion and industrial processes, followed by ~~other sources (mainly waste)transport and waste (OTHER)~~. There is a significant relative shift in the dominant emission

sector as well: the main emission source becomes non-ferrous metals (63% of 2050 emissions) and cement (10%). For the NFME sector, there is not much potential of emission reduction left in the GAINS model as acid plants, the most efficient control technology currently implemented in e.g. copper smelters, are already mandated⁶. A very similar trend can be observed for North America.

African, Central and South American as well as Southeast Asian Hg emissions are dominated by ASGM and the uncertainty intrinsic to these estimates (-36.8% up to +44.5%, see Table S12), eclipses trends in all other sectors such as emissions, which remain constant at around three quarters of the total. Small increases in cement and waste sectors point at a growing trend for population and resulting building activity, while combustion driven emissions remain at the same level as currently. Central and South American emissions show a very similar trend picture with ASGM emissions being 84-85% of total emissions. For all three both regions, it is important to note that ASGM estimates for 2015 are subject to large uncertainty (Keane et al., 2023) and the purpose of projections in all scenarios can only be to show the influence of Hg policy such as the Minamata convention, as the activity is kept constant, not reflecting the current forecast for production numbers of this sector.

Table 3: Mercury emissions in the Baseline + Current Legislation (01_BAS_CLE) scenario by world regions and by sectors (tons year⁻¹).

⁶ To further reduce Hg in NFME, the removal efficiency of acid plants would need to be increased. The GAINS removal efficiency of acid plants is significantly lower than their assumed removal efficiency in the GMA'18. This is due to the speciated emission accounting approach. The removal efficiency of PR_AP for Hg⁰ is 91% (see Table A6 in the appendix and sources therein). As 80% of the emissions are assumed to be Hg⁰, this leads to an overall removal efficiency of 92.7% as opposed to 99.98% in the GMA'18. Better data on emission speciation would be needed for a more exact estimate for this sector, and there might be an overestimation of European NFME emissions in GAINS.

Formatted: Heading 3

		ASGM	CEM	COMB_ IND	COMB_ OTHER	COMB_ POWER	GOLD	IND_ PROC	NFME	OTHER	Total
2015	Global	952.61	153.69	85.20	99.27	263.21	43.60	110.09	349.69	263.24	2320.61
	Africa	280.24	11.18	1.94	15.86	8.98	11.80	2.89	12.12	14.00	359.01
	Asia Pacific	301.03	104.85	73.47	69.91	179.56	3.46	53.41	163.40	142.32	1091.41
	China	48.27	61.76	46.87	47.97	68.19	0.69	33.07	85.66	93.02	485.51
	India	6.37	21.91	18.82	14.88	92.04	0.00	9.82	8.12	25.46	197.43
	Japan	0.00	1.89	0.69	0.22	2.04	0.03	3.19	7.62	4.63	20.33
	Southeast Asia	240.91	12.08	2.99	3.16	6.38	0.44	3.01	14.00	11.17	294.13
	Central & South America	352.80	5.46	2.25	1.84	2.36	7.32	7.31	18.60	12.99	410.93
	Brazil	50.27	0.81	1.46	0.65	0.74	0.27	3.92	6.20	6.42	70.73
	Eurasia	14.31	3.90	1.42	1.84	10.09	14.60	11.90	55.51	17.17	130.73
	Russia	7.57	1.43	0.85	0.96	8.68	8.55	10.09	35.81	15.19	89.12
	Europe	0.00	13.14	2.46	7.67	35.33	1.31	19.89	81.90	20.18	181.89
	EU 27	0.00	6.97	1.54	4.88	26.43	0.42	15.17	78.21	11.40	145.02
	Middle East	0.23	12.39	0.69	0.30	1.43	0.29	6.28	4.07	27.50	53.18
	North America	4.00	2.78	2.96	1.84	25.47	4.82	8.42	14.09	29.08	93.45
	USA	0.00	1.78	2.41	1.13	24.25	0.85	5.52	1.44	20.08	57.45
2050	Global	1130.62	223.51	131.57	42.27	127.90	25.84	115.38	337.23	319.12	2598.27
	Africa	333.84	25.55	4.14	16.79	2.30	1.53	4.81	1.83	29.74	440.24
	Asia Pacific	369.21	150.76	115.59	14.85	108.09	3.52	73.04	202.59	139.68	1251.32
	China	44.21	63.73	34.45	6.88	49.64	0.63	31.76	91.05	55.53	377.87
	India	8.72	51.63	62.09	3.80	30.58	0.00	26.52	19.18	43.37	245.90
	Japan	0.00	1.79	1.14	0.18	0.55	0.03	2.87	7.14	4.47	18.17
	Southeast Asia	310.17	18.33	6.88	1.18	13.61	0.29	7.27	21.83	23.98	403.56
	Central & South America	401.99	10.35	3.65	2.66	1.19	6.05	6.63	15.78	24.27	478.83
	Brazil	55.99	1.09	2.50	1.84	0.64	0.30	3.46	5.37	13.30	84.49
	Eurasia	22.00	4.50	2.45	0.82	7.56	7.72	7.16	2.14	18.81	94.55
	Russia	9.14	1.65	2.04	0.28	4.80	1.18	6.15	2.14	16.79	44.16
	Europe	0.00	14.53	2.98	2.69	6.91	1.84	11.03	95.11	13.17	148.26
	EU 27	0.00	8.06	1.82	2.32	0.77	0.52	5.31	89.19	6.04	114.04
	Middle East	0.29	13.34	0.57	0.26	1.41	0.01	5.74	2.08	61.74	85.44
	North America	3.29	4.48	2.21	4.21	0.44	5.17	6.97	17.70	31.70	76.17
	USA	0.00	2.44	1.72	3.31	0.31	0.79	4.32	1.90	17.84	32.63




Figure 3: Global mercury emissions for the BAS scenario [combined with a set of all](#) different control strategies – by sector.




Figure 4: Regional data: Hg emissions for the Baseline, Minamata, co-MFR and BAS_HgMFR scenarios (01_BAS_CLE, 02_BAS_MINA, 03_BAS_coMFR, 04_BAS_HgMFR). (a) Africa (b) Central and South America, (c) Middle East, (d) Europe, (e) Eurasia, (f) North America, (g) Asia Pacific and Asia-Pacific sub-regions; (h) China, (i) India, (j) Southeast Asia.

365 The Asia-Pacific region spans the majority of the world population and produces the majority of the world's Hg emissions. Emissions represent 46% of global emissions in 2015 and 48% in 2050. Globally, most emissions from power generation and industrial combustion come from this region. Emission reductions are projected in the Baseline (01_BAS_CLE) only in combustion in power plants, as well as in residential combustion. Emissions related to manufacturing and building activities, such as cement production, industrial processes, NFME smelting and also population-related releases such as waste are projected to increase by 2050. ASGM emissions are regionally highly variable. Within the Asia-Pacific region, large

differences pervade on the country and subregion level (Fig. 4 (h) – (j)). Emissions in China are projected to decrease in the Baseline due to both decarbonization and co-benefits from rapid and stringent application of PM and SO₂ controls. Indian Hg emissions from fossil fuel combustion (in COMB_POWER) are projected to increase, and ASGM emissions dominate in Southeast Asia. ~~Japan is a typical example of an industrialized country—there is a decrease in combustion emissions to 2050, but a stable trend in NFME emissions, as there is little scope for reduction left within the air pollution control policies, except for lowering production volumes.~~

Emissions from the **Middle East** are dominated by cement production and unmanaged landfill waste emissions.

5.1.1 Uncertainty

Table S12 presents the percent ranges for the aggregated emission sectors on a global level. ASGM emission variations are largest on a relative level, but due to the large uncertainty in emission factors and the high unabated emission factors in the NFME sector, the absolute range of emissions was largest in this sector, followed by OTHER emissions and CEM. Combustion sectors show a small spread, reflecting the good data quality. While large-scale gold emissions give a large relative spread in the upper and lower bounds, their low total emissions mean that their contribution to the overall uncertainty of the results is small. ~~Assuming that the abated emission factors, which take into account Hg emission reductions through abatement measures, have the same levels of uncertainty as the UEFs, the calculated relative uncertainties are applied to 01_BAS_CLE (see Figure S1).~~ The result is that the main uncertainties of this model lie within the ASGM sector; ASGM emissions make up between a third and 47% of total emissions, looking at the lower and upper range estimates, respectively, varying between 602 and 1373 t in 2015. After this, Waste emissions have the most variability in absolute terms, followed by emissions from the non-ferrous metal and power sector. To conclude, improved data quality in ASGM and non-ferrous metals and waste, as well as cement will significantly reduce/improve the overall uncertainty/quality of the baseline mercury emissions estimates.

5.2 Mercury emission reductions as co-benefits from PM and SO₂ controls/benefits

Comparing the No Control and Baseline (~~BAS_CLE~~) scenarios, the full extent of emissions reductions from current clean air policy becomes apparent, as NOC emissions are more than double those of the Baseline in 2015, illustrating that already, more than 50% of potential unabated Hg emissions are avoided through existing clean air policy. An extended discussion can be found in ~~S~~section S4 and ~~Fig. S1~~ of the appendix.

The further potential for Hg emission reductions through co-benefits from PM and SO₂ policy are assessed by comparing the Baseline with a co-MFR case. ~~It shows the total Hg emission reduction potential from PM and SO₂ abatement,~~ without considering any Hg-specific measures beyond those already implemented in the CLE case. In 2050, the co-MFR scenario projects 2023 t emissions compared to 2455 t in the Baseline – a reduction of 18%.

In most of the world's regions, the comparison of co-MFR compared to CLE scenarios reveals that a large fraction of power sector capacities and industrial installations are already controlled by at least PM controls and some form of SO₂ control policy is already put in place by 2050, reflected in an leading to implementation of the Hg control measure 'PM_FGD' in the power sector in GAINS—the most efficient co-benefit control technology. An extended discussion of the technology shares can be found in the SI, section S3.2. In China alone, the retrofitting measures for coal-fired power plants in the 12th Five-Year-Plan from 2010 to 2015 have reduced Hg emissions by 23.5 tons, explaining why further reductions in the power sector are limited as current policy already mandates the strictest levels of APCD deployment, closure of inefficient small plants (e.g. Li et al. 2020) and coal phase-out policies (e.g. Li et al., 2020). On the other hand, there is still scope for improvement in the industry sectors such as NFME, cement, gold and other production: in 2050, emissions in the co-MFR for China are only 6.2% (22.5t) lower than in the Baseline. The difference is mainly found in the OTHER (17.4 t) and IND_PROC (4.6 t) sectors, indicating that co-benefits from PM and SO₂ control measures are already planned to be maximized in the currently active policy. The results for India imply that co-benefits from air pollution abatement policy can still make significant contributions to lower Hg emissions, totaling 12.4% reduction in the co-MFR case compared to the Baseline. The largest reductions are 18.0 t Hg reduction in cement production in 2050 between the Baseline and the co-MFR case and 10.1 t Hg reduction in OTHER, with contributions mostly from waste management (see Fig. S3). In the Middle East, the potential for emission reductions through better waste management becomes especially clear (Figs. 3, 4, S3), as waste emissions in OTHER dominate the picture. In the Baseline, unmanaged waste allows Hg to be emitted into the atmosphere and waste generation is expected to increase with population growth. However, in the co-MFR scenario, these emissions are minimized by 2050 thanks to an application of waste incineration with efficient Hg capture, the OTHER sector alone causing a 69.2% reduction in total Hg emissions in 2050 in the co-MFR compared to the Baseline. In North America, Europe including the EU 27 and the Asia Pacific region, the most significant co-benefit reduction potential lies in the NFME sector, owing to its exceptionally high emission factors, where even small improvements in the co-benefit controls are able to cut tons of Hg emissions. For example, in the EU27, over 90% of the emission reduction is reached only in the NFME sector, which could see a further 54.1 t Hg reductions EU-wide if the strictest Acid Plant controls are employed at all plants.

5.3 The Maximum Feasible Reduction scenario



Comparison with a maximum feasible reduction (MFR) scenarios can serve to quantify the maximum potential of targeted Hg abatement (04_BAS_HgMFR, 07_BAS_HgMFR, short Hg-MFR) relative to other approaches. Here, the results of BAS_HgMFR are discussed relative to the BAS co-MFR scenario, which uses only air pollution co-benefit measures for Hg reduction (co-MFR, discussed in section 5.2.2). Both scenarios represent end points of possible policy developments, so this comparison is only made for the year 2050. Co-benefits from PM and SO₂ control impact mainly the power, industry and waste sectors, however, there are more efficient technologies available targeting Hg emissions applicable to a large range of sectors. For industry and combustion sectors where Hg-specific measures are expected, the Hg-MFR scenario assumes adoption of the

most efficient pollution control measures available, which in most sectors is sorbent injection in front of a fabric filter (FFSINJ). The Hg-MFR scenario implements a complete, ~~global~~ ban on ASGM and is thus the only scenario where the 1130 t of ASGM Hg emissions ~~are projected to disappear~~, reducing total Hg emissions in 2050 drastically by 79% compared to the Baseline, ~~from 2455 t to 521 t~~. As seen in Figs. 3 and S3, the difference in 2050 emissions is also large between the Hg-MFR and the co-MFR: the co-MFR projects 2050 emissions to be 2023 t, 3.9 times higher compared to the 521 t in Hg-MFR. Even when discounting the fact that ASGM emissions are completely removed from the Hg-MFR, and when comparing all sectors except ASGM, the co-MFR would still produce 371 t annual Hg emissions more than the Hg-MFR case. ~~Besides Apart from ASGM,~~ —the largest absolute reduction of 135 t can be achieved in the cement sector, where emissions in Hg-MFR are just 24% of those in the co-MFR scenario. This is followed by emissions from industrial processes (mainly iron and steel production metal production (?)), which, in 2050, are halved to 43 t in the Hg-MFR compared to ~~86 t in~~ co-MFR. Little to no further emission reductions are estimated in the COMB_OTHER, NFME and OTHER sectors. Waste-related emissions (~~from OTHER~~) are captured by multi-pollutant controls. This is especially visible in the Middle East region, where the majority of emissions is modeled to come from waste (in OTHER), as shown in the CLE and MINA scenarios (e.g. Fig. 4). This emission source is addressed/problem is solved in the co-benefit MFR scenario, where better landfill practices are adopted and the loosely managed waste is re-directed into waste incineration with mercury capture. Similarly, in NFME, the most advanced SO₂ controls, sulfuric acid plants, are expected to already comply with strict Hg legislations and are expected to contain Hg-specific sorbents, as sulfuric acid is sold on as a product and therefore needs to comply with the Hg-content/quality standards applicable for limit values of this product.

Combined with the Net Zero CO₂ policy, (scenario-07_CLIM2_HgMFR) presents the lowest-possible primary Hg emissions in 2050: Co-benefits from climate policy, clean air policy and Hg control policy are maximized and taken into account at the same time, leading to only 446.7 t Hg emissions in 2050, a further reduction of 56.9 t ~~are reduced in the combustion sectors compared relative~~ to BAS_HgMFR (Figs. 3, 5).

5.4 Mercury emission changes with climate policy

To compare climate policy impacts/outcomes on Hg pollution, the three energy scenarios (BAS, CLIM1 and CLIM2) with CLE controls (BAS_CLE, CLIM1_CLE, CLIM2_CLE) have been analyzed. (see Fig. 5). Considering the large decline in CO₂ emissions by 2050 driven by climate mitigation goals simulated in the Stated Policies, Announced Pledges and Net-Zero Emissions scenarios (IEA, 2022), our analysis suggests that the reductions in Hg emissions, while apparent in Fig. 5, occur at a significantly lower rate than for CO₂ and the differences between the three scenarios (277.7 t Hg between BAS_CLE and CLIM2_CLE) is smaller than the uncertainty attached to the ASGM estimates. Nevertheless, we believe that a discussion of trends is warranted as this helps to quantify co-benefits from climate action. ~~As ASGM emissions are not affected by climate policy in our model, they have been excluded from Fig. 5 for easier comparison of the remaining sectors.~~



465

Figure 5: Global mercury emissions excluding ASGM for the three energy scenarios BAS, CLIM1 and CLIM 2 under the CLE control strategy—by sector. The CLIM2 with HgMFR controls is included as the scenario with lowest-possible Hg emissions.

470 Noticeable emission reductions relative to 2015 are projected in the combustion sectors, caused by a reduction in global consumption of coal, oil, natural gas and traditional biomass use in the three scenarios. In contrast, emissions from cement production (CEM) and ‘OTHER’ (transport and mostly waste) increase in all three scenarios between 2015 and 2050. While this doesn’t offset the emission reductions in CLIM1 and CLIM2, it leads to a slight increase in global emissions in the 01_BAS_CLE scenario. A shift of emissions from the combustion sector towards industrial processes, gold production and waste treatment can be seen due to changes in the energy and industrial systems and is most pronounced in scenario compliant with the most stringent climate policy (Fig. 5, Panel b).

475 The decarbonization of global economy and transition towards renewable energy sources and associated infrastructure induces an increased demand for critical minerals for electrification in all three scenarios. In GAINS, this is reflected as increased demand for non-ferrous metals in 2050 (see Fig. 3, NFME activity). However, improved pollution control measures lead to 3.6% lower emissions in 2050 than 2015. A projected increase in activities and resulting emissions in all three scenarios relative to 2015 are also shown in other industrial processes sector (IND_PROC), largely due to iron and steel production. In the 480 CLIM1 scenario, IND_PROC emissions dip to 93t in 2030, then increase again to 101t until 2050. Overall, the baseline (BAS) scenario projects slightly rising emissions from IND_PROC, while both CLIM1 and CLIM2 show slight declines from 110 t/a in 2015 to 101 t/a Hg (CLIM1) and 91 t/a (CLIM2) until 2050, mostly due to declining iron and steel emissions.

485 The largest differentiation between the different energy scenarios and their resulting Hg emissions is apparent within the combustion sectors. Globally, emissions from the power sector (COMB_POWER) roughly halve from 2015 to 2050 in the Baseline and drop further by 82% and 97% in CLIM1 and CLIM2 respectively, virtually removing the combustion sources of Hg from the emissions profile. It is noted that the largest reductions in Hg emissions occur in regions where coal power is a significant contributor to the energy mix. Globally, the largest relative share of combustion-related Hg emissions in 2050 in all scenarios is estimated within the industrial combustion sector. In the Baseline, emission levels from industries are projected 490 to rise by 50% between 2015 to 2050, however, in comparison these are reduced by a quarter in CLIM1 and by 56% in CLIM2. While fossil fuel combustion in the power sector is projected to decline on a global level, in some countries such as India, fossil fuel combustion in industries (COMB_IND) is projected to grow beyond 2015 values in all scenarios, with Hg emissions projected to more than triple from 18.8 t/a in the Baseline to 62.1 t/a by 2050. In CLIM1, COMB_IND emissions rise by 7.7 t/a, and only in the net-zero CLIM2 scenario, Indian industrial combustion emissions decrease to 6.7 t/a by 2050.

495 Emissions of Hg from activities which are scaled by population growth, such as waste generation, rise equally in all scenarios. In the category ‘OTHER’, Hg emissions from transport are combined with emissions from waste and some other Hg emission activities. Consequently, waste emission increases are offset by emissions savings from fossil fuel-based road and off-road transport. Nevertheless, an overall increase in these emissions is seen across all three scenarios, although slightly higher in the Baseline than in the climate scenarios.

500 **5.5 Implications of Minamata convention on future Hg emission trends**

For the purpose of direct comparison between the modelling results and [impacts of these the](#) Minamata Convention on Mercury (MCM) concerning Hg emissions to the atmosphere, a sector mapping between the MCM provisions and GAINS was conducted, as detailed in Table 4: GAINS sectors which are also included in the MCM ([UNEP, 2013ref](#)) are grouped into Annex B, C and D sources. Smaller emission sources in GAINS including some industrial processes, combustion of biomass, liquid and gaseous fossil fuels, as well as transport emissions are not covered by the MCM⁷. Figure 6 shows the results of the BAS scenarios (CLE, MINA, coMFR, HgMFR), mapped to the Minamata sectors.

Table 4. Source and process categories covered by Minamata Convention provisions, with their corresponding GAINS sector representation in the ‘Minamata’ sector aggregation.

Source and process categories	Convention provisions	GAINS representation
Extraction and use of fuels/energy sources		
Coal combustion in power plants	Article 8, Annex D	COMB_POWER
Coal combustion in coal fired industrial boilers	Article 8, Annex D	COMB_IND
Primary (virgin) metal production		
Gold (and silver) extraction with mercury amalgamation processes	Article 7, Annex C	ASGM
Gold extraction and initial processing by other methods	Article 8, Annex D	GOLD
Zinc extraction and initial processing	Article 8, Annex D	NFME
Copper extraction and initial processing		
Lead extraction and initial processing		
Cement production		
Cement and clinker production	Article 8, Annex D	CEM
Intentional use of mercury in industrial processes		
Chlor-alkali production with mercury-technology	Article 5, Annex B	OTHER
Vinyl chloride monomer production with mercury catalyst	Article 5, Annex B	OTHER
Acetaldehyde production with mercury catalyst	Article 5, Annex B	not in GAINS

⁷ It is noted that there are also emission sources which the MCM covers, that are not explicitly represented in GAINS. This concerns emissions from acetaldehyde production, other chemicals and polymers production, incineration of medical waste, hazardous waste and sewage sludge.

	Other production of chemicals and polymers with mercury	Article 5, Annex B	not in GAINS
Waste incineration			
	Incineration of municipal/general waste	Article 8, Annex D	OTHER
	Incineration of hazardous waste	Article 8, Annex D	not in GAINS
	Incineration of medical waste	Article 8, Annex D	not in GAINS
	Sewage sludge incineration	Article 8, Annex D	not in GAINS
Other emission sources			
	E.g. transport emissions, various industrial processes, domestic coal burning	Not covered	NOT_COVERED

510




Figure 6: Emissions from sectors covered by the Minamata convention.

5.5.1. Annex D sources

515 Article 8 of the MCM addresses atmospheric emissions directly by 'controlling and, where feasible, reducing emissions of mercury and mercury compounds, often expressed as "total mercury", to the atmosphere through measures to control emissions from the point sources falling within the source categories listed in Annex D' (Minamata Convention on Mercury, 2013, art. 8.1). Annex D sources are represented well in GAINS (Table 4). Article 8 does not mandate a phase-out of these activities, but rather mandates use of BAT/BEP for new sources and it is up to the parties of the MCM to formulate appropriate steps to reduce Hg emissions from existing sources – which may be co-benefit reduction of Hg from other air pollution control or Hg-

520 specific measures. The different control strategies in the CLE, co-MFR and Hg-MFR scenarios (01_BAS_CLE, 03_BAS_coMFR, 04_BAS_HgMFR) represent different narratives of how the art. 8 objectives could be achieved, as different Hg reduction measures may be considered BAT/BEP. For example, the EU's Industrial Emissions Directive (EU-IED) states for large coal and lignite combustion plants that PM, SO₂ or NO_x as well as Hg-specific reduction technologies are considered BAT (see BAT 27 in Commission Implementing Decision (EU) 2021/2326 (2021)). Fuel choice is also listed as a Hg-specific control measure but might be considered a co-benefit from energy and climate policy in the case of Hg emissions from coal. Thus, each scenario (Baseline, co-MFR, Hg-MFR, CLIM1_CLE and CLIM2_CLE, and combinations thereof) would ~~be represent a variation of possible~~ compliance with Art. 8 for coal combustion. For non-ferrous metals, BAT is to consider raw materials with low Hg contents, as well as using Hg-specific sorbents in the EU (BAT 11 in Commission Implementing Decision (EU) 2016/1032 (2016)), the end point of which are represented in the Hg-MFR scenario. ~~For cement, the EU BAT conclusions are a combination of limiting Hg content in the feed materials, as well as utilizing dust control co-benefits (BAT 43 and 54, Commission Implementing Decision (EU) 2013/163/EU (2013)), represented in the CLE and co-MFR scenarios. Dedicated Hg controls are required in waste incineration (BAT 31, Commission Implementing Decision (EU) 2019/2010 (2019)) which would represent a Hg-MFR scenario if implemented globally.~~ Annex D sources are globally dominated by NFME and coal power emissions in 2015, but power sector emissions are expected to fall in importance until 2050, as discussed earlier. Industrial emissions (NFME, CEM) as well as waste emissions are projected to become the largest contributors to total emissions in Annex D in all scenarios. The ~~present scenario~~ results suggest that targeted, Hg-specific ~~measures~~ BAT recommendations result in the most ~~efficient~~ significant Hg reduction for Annex D sources, as they are generally associated with lowest emission factors. ~~However, after taking into account region-specific and economical factors, this may not translate into BAT/BEP for many countries and represents and represents an end point of lowest possible emissions from these sources.~~

540 5.5.2. Annex C sources

Article 7 does not exclusively address air emissions but concerns the phase-out of ASGM activities: Each party with ASGM activities on their territory 'shall take steps to reduce, and where feasible eliminate, the use of mercury and mercury compounds in, and the emissions and releases to the environment of mercury from, such mining and processing.' (Minamata Convention on Mercury, 2013, art. 7.3). The National Action Plans (NAPs) required by Art. 7 lay out the planned reduction / phase-out for each relevant party. ~~Where NAPs are already published, the reduction targets have been implemented in the BAS_MINA scenario.~~ With already currently published targets fully implemented, ASGM emissions in BAS_ MINA are projected to sink to 738.3 t in 2030 and 700.0 t in 2050 and ASGM is bound to make up roughly a third of global Hg emissions in 2050, the largest absolute reduction stemming from Southeast Asia (272.0 t) and Africa (116.3 t). ~~However,~~ There are large uncertainties connected to activity levels as well as emission factors of ASGM (determined to be -37% to +44% ~~uncertainty error~~ in this study's Baseline on a global level).~~;~~

Formatted: Normal

5.5.3 Sources not covered by the Minamata Convention

Further, it is important to note that the MCM does not cover all potential emission sources and not all mercury uses, instead focusing on intentional use of Hg and Hg compounds, as well as the largest emitters. In this study, only roughly 10% (of the emissions computed by GAINS fall into the 'NOT_COVERED' category: 234.6 t Hg in 2015, increasing slightly to 259.8t by 2050 in BAS_CLE. The number includes emissions from residential and domestic coal combustion, biomass combustion, transport emissions, emissions related to oil refining, production of iron and steel, paper, fertilizer, glass and aluminium. This number is higher than the GMA'18 estimate (171.5 t Hg, not including paper, fertilizer and glass). It, highlighting that the MCM has the potential to impact 90% of the future emissions examined shown in this paper.

5.6 Speciation

The speciation of Hg emissions released to the atmosphere strongly influences their fate and spatial distribution (e.g. Selin 2009). In this study, Hg⁰ dominate the speciation profile in all scenarios, making up between 76% and 90% of total emissions in the year 2050, as can be seen in Fig. 7. The scenarios with Hg-specific control measures (BAS_HgMFR, CLIM2_HgMFR) project the lowest proportions of Hg⁰ emissions with 76% and 77%, respectively, while the scenario BAS_coMFR, which leans most heavily on co-benefit control of Hg displays the highest share of 90% Hg⁰ emissions. This is due to the fact that the PM and SO₂ controls implemented in GAINS tend to have higher removal efficiencies for Hg^{II} and Hg_p, while the harder-to-abate Hg⁰ requires targeted approaches as implemented in the Hg-MFR scenarios. As ASGM emissions are assumed to be only elemental Hg⁰, scenarios with higher ASGM abatement such as BAS_MINA has a lower share of Hg⁰ emissions than BAS_CLE (84% and 87%, respectively).




Figure 7: Share of mercury species in different scenarios.

5.7 Further work

575 While the future scenarios are indicative of the abatement potential of different types of policy, there are some simplifications
 580 built into the current implementation of Hg within the GAINS model that would warrant further attention and present areas of
 future work.

(1) When considering future projections of **non-ferrous metal smelting**, the share between Cu, Pb and Zn and the proportions
 of primary vs. secondary [production/mining](#) within the NFME category is fixed to the 2015 level, not dynamic. This is a
 580 simplification that can lead to an overestimation of Hg emissions from this sector, as copper production generates relatively
 lower Hg emissions compared with lead and zinc, but its share in the NFME category is projected to increase as it is a critical
 metal in the decarbonized economy.

(2) Similarly, assumptions on Hg levels in the **waste sectors** have one fixed, global unabated emission factor, derived from
 total estimates of emissions from waste in 2015. These numbers are projected into the future using shares of different waste
 585 types, as well as population and macroeconomic projections. However, again, the fixed emission factor might change, as Hg
 policy reduces Hg levels in consumer products. Furthermore, the emission [factors?](#) are likely to be heterogeneous on a regional
 scale. Better projections of [Hg-content](#) in wastes will increase the accuracy of waste emissions estimates as well can better
 simulate shifts in waste composition under assumptions of circular economy.

(3) The Hg removal efficiency of **NOx controls** could be studied further and [impacts of](#) NOx control policy and their interplay
 590 with Hg, PM and SO₂ control technologies could be included into the GAINS algorithm, thus making the calculation of removal

efficiencies more detailed. This is especially relevant for the consideration of Hg speciation, as selective catalytic reduction technology for NOx removal from flue gas systems can be optimized to oxidize Hg⁰ to Hg^{II}, thus changing the share of species as well as increasing the efficiency of particle filters and flue gas desulfurization units in keeping mercury from the atmosphere (Usberti et al., 2016).

595 (4) Projections of **artisanal and small-scale gold mining** into the future need to be improved as additional data and a better understanding of the ASGM drivers are emerging from the Minamata Convention process and the scientific community. Similarly, policy and technical measures for ASGM emission reduction might be refined to better and added in GAINS reflect the complexity of this social phenomenon.

Formatted: Not Highlight

600 (5) As is done for traditional pollutants, optimization of mitigation options would allow the calculation of cost-optimal achievement meeting of reduction targets. Further, GAINS model output could be used as input to dispersion modelling of Hg transport and deposition, and the subsequent calculation of health and environmental impacts. The authors are also working on making some of the Hg scenarios accessible to the public via the GAINS online tool (include the link).

6 Conclusions

605 This study explores future anthropogenic mercury emissions through seven scenarios combining different energy and climate strategies (BAS, CLIM1, CLIM2) with policies to abate mercury as well as traditional pollutants such as PM and SO₂ (CLE, MINA, co-MFR, Hg-MFR).

Of all studied sectors, ASGM offers the largest absolute Hg reduction potential, but also the largest uncertainty in the emission estimates, and in world regions with significant ASGM contribution, all other emission sources fall within the uncertainty of ASGM emissions, making the studied scenarios (with exception of Hg-MFR) virtually undistinguishable. While caution should be exercised when interpreting absolute emission values, valuable information can be gained from considering trends across years and between scenarios, and sectors.

Formatted: Not Highlight

615 The Baseline (BAS_CLE) projects a slight increase of 5.7% in global emissions in 2050 compared to 2015, despite a 32.6% reduction of emissions from combustion, due to increased cement and other emissions including waste (+45.4% and +21.4% resp.). The comparison of three climate scenarios under current legislation for clean air policy shows that the Hg emission reduction from the fossil fuel combustion sectors depends on the level of climate policy ambition, which prompts a transformation in the energy system towards non-coal sources, enabling a range of 30% (BAS_CLE) - 86% (CLIM2_CLE) emission reduction from combustion sources and causing. However, there are trade-offs such as a 45% increase in cement emissions, a shift of emissions towards industrial processes, gold production and waste treatment.

620 In all studied sectors except ASGM, emission increases can be dampened or reversed by conventional air pollution control measures. High levels of co-benefits from PM and SO₂ control are already induced by present in the current legislation, as comparison to the No Control Scenario demonstrates. Compared to the Baseline, a 17.6% reduction in emissions can be

achieved in 2050 solely through maximizing PM and SO₂ control (BAS_coMFR). This strategy is especially effective strategy for the waste sector (represented in OTHER), where emissions fall by 58.9% compared to the Baseline in the 2050. Drastic reduction of ASGM as set out in the Minamata Convention is only possible with a Hg-focused approach, as the MINA and Hg-MFR scenarios show. ASGM offers the largest absolute Hg reduction potential of any sector, but also the largest uncertainty in the baseline emission estimates. OTHER emissions (including waste) are also a large source of uncertainty in this study, and projections range between a 22% increase and 54% decrease in 2050 depending on both climate and clean air policy.

The Minamata convention covers roughly 90% of the Hg emissions computed by the GAINS model. Annex D sources of the convention are may be regulated differently by each convention party and the CLE, co-MFR and Hg-MFR scenarios combined with different climate policy delineate the option space of possible Hg reductions from these sources, through BAT/BEP, which oftentimes means co-benefits from pollution control, as set out in the co-MFR scenario, so implying the CLE, co-MFR and Hg-MFR scenarios represent three different narratives that could represent pathways of different countries to reduce Hg from Annex D sources.

Overall, the findings emphasize the importance of the ASGM sector for total Hg emission reduction. Further, implementing targeted Hg control policies in addition to stringent climate and air pollution abatement, PM and SO₂ policies is vital to achieve significant reductions in Hg emissions.

Code and data availability.

The aggregated data discussed in this paper is in the electronic supplement. The GAINS model can be accessed upon registration through the online interface (http://gains.iiasa.ac.at/models/gains_models4.html~~http://gains.iiasa.ac.at/gains/GOD/index.login~~).

Author contributions

Conception of and study design ... PR, FB. Concept and methodology ... FB, PR, FB, FW. Implementation into the GAINS model ... FB, RS. Data collection, uncertainty calculation, preparation of manuscript ... FB. Manuscript review and supervision ... PR, MJM, FW.

Competing interests.

The authors declare no competing interests.

Acknowledgements

650 This work has been supported by the IIASA Young Scientists Summer Programme 2021, as well as by the RC-UK Centre for Doctoral Training in Bioenergy [Grant number EP/L014912/1]. FB thanks Zbigniew Klimont for welcoming and supporting the project, as well as comments on the manuscript. [Feedback from Francesco De Simone and an anonymous reviewer during review has been gratefully received and taken on board, sharpening the overall discussion.](#)

655 *Supplementary Information.*

The following files were submitted separately:

2023-ACP_SI_ScenarioData.xlsx

2023-ACP_SI_Tables.docx

References

660 [Amann, Markus, Imrich Bertok, Jens Borken-Kleefeld, Janusz Cofala, Chris Heyes, Lena Höglund-Isaksson, Zbigniew Klimont, et al. 2011. "Cost-Effective Control of Air Quality and Greenhouse Gases in Europe: Modeling and Policy Applications." *Environmental Modelling & Software* 26 \(12\): 1489–1501. doi:10.1016/j.envsoft.2011.07.012.](#)

665 [Amann, M., Bertok, I., Borken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., Klimont, Z., Nguyen, B., Posch, M., Rafaj, P., Sandler, R., Schöpp, W., Wagner, F., and Winiwarter, W.: Cost-effective control of air quality and greenhouse gases in Europe: Modeling and policy applications, *Environmental Modelling & Software*, 26, 1489–1501, <https://doi.org/10.1016/j.envsoft.2011.07.012>, 2011.](#)

670 Amann, M., Kiesewetter, G., Schoepp, W., Klimont, Z., Winiwarter, W., Cofala, J., Rafaj, P., Hoglund-Isaksson, L., Gomez Sanabria, A., Heyes, C., Purohit, P., Borken-Kleefeld, J., Wagner, F., Sander, R., Fagerli, H., Nyiri, A., Cozzi, L., and Pavarini, C. 2020. "Reducing global air pollution: The scope for further policy interventions." *Philosophical Transactions of the Royal Society A*. doi:10.1098/rsta.2019.0331.

AMAP/UNEP. 2013. "Technical Background Report for the Global Mercury Assessment 2013." Arctic Monitoring and Assessment Programme. Arctic Monitoring; Assessment Programme, Oslo, Norway / UNEP Chemicals Branch, Geneva, Switzerland.

675 AMAP/UNEP. 2019. Technical Background Report to the Global Mercury Assessment 2018. Troms: Arctic Monitoring; Assessment Programme, Oslo, Norway / UN Environment Programme, Chemicals; Health Branch. https://www.mercuryconvention.org/sites/default/files/documents/forms_and_guidance_document/BAT_BEP_E_interractif.pdf.

Formatted: Font: 10 pt

Formatted: Bibliography

Formatted: Font: 10 pt

Formatted: Font: 10 pt

680 [Angot, H., Hoffman, N., Giang, A., Thackray, C. P., Hendricks, A. N., Urban, N. R., and Selin, N. E.: Global and Local Impacts of Delayed Mercury Mitigation Efforts, Environ. Sci. Technol., 52, 12968–12977, <https://doi.org/10.1021/acs.est.8b04542>, 2018.](#)

Formatted: Font: 10 pt

Formatted: Bibliography

685 [Angot, Helene, Nicholas Hoffman, Amanda Giang, Colin P. Thackray, Ashley N. Hendricks, Noel R. Urban, Noelle E. Selin. 2018. "Global and Local Impacts of Delayed Mercury Mitigation Efforts." Environmental Science & Technology 52 \(22\). ACS Publications: 12968–12977. doi: 10.1021/acs.est.8b04542.](#)

690 CPCB. 1998. "Emission Regulations. Part – Two." Comprehensive Industry Document Series: COINDS/18/1984-85. Central Pollution Control Board (Ministry of Environment & Forests, Government of India), Parivesh Bhawan, East Arjun Nagar, Delhi.
<https://cpcb.nic.in/openpdf.php?id=UmVwb3J0RmlsZXMvTmV3SXRlbV8xNjRFRU1JU1NJT05FukVHVUxBVEIPTINfUEFSVF8yLnBkZg==> (Last accessed 7.9.2023).

Field Code Changed

695 [Cheng, Chin-Min, Pauline Hack, Paul Chu, Yung-Nan Chang, Ting-Yu Lin, Chih-Sheng Ko, Po-Han Chiang, Cheng-Chun He, Yuan-Min Lai, and Wei-Ping Pan. 2009. "Partitioning of Mercury, Arsenic, Selenium, Boron, and Chloride in a Full-Scale Coal Combustion Process Equipped with Selective Catalytic Reduction, Electrostatic Precipitation, and Flue Gas Desulfurization Systems" 23 \(10\). American Chemical Society \(ACS\): 4805–16. doi:10.1021/ef900293u. Cheng, C.-M., Hack, P., Chu, P., Chang, Y.-N., Lin, T.-Y., Ko, C.-S., Chiang, P.-H., He, C.-C., Lai, Y.-M., and Pan, W.-P.: Partitioning of Mercury, Arsenic, Selenium, Boron, and Chloride in a Full-Scale Coal Combustion Process Equipped with Selective Catalytic Reduction, Electrostatic Precipitation, and Flue Gas Desulfurization Systems†, 23, 4805–4816, <https://doi.org/10.1021/ef900293u>, 2009.](#)

Formatted: Font: 10 pt

700 [Convention on long-range transboundary air pollution, CHAPTER XXVII. - ENVIRONMENT, Geneva, 13 November 1979. \[hereinafter CLRTAP\] \[https://treaties.un.org/Pages/ViewDetails.aspx?src=IND&mtdsg_no=XXVII-1&chapter=27&clang=en\]\(https://treaties.un.org/Pages/ViewDetails.aspx?src=IND&mtdsg_no=XXVII-1&chapter=27&clang=en\) \(retrieved April 5, 2024.\)](#)

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Hyperlink, Font: 12 pt

Field Code Changed

Commission Implementing Decision (EU) of 26 March 2013 establishing the best available techniques (BAT) conclusions under Directive 2010/75/EU of the European Parliament and of the Council on industrial emissions for the production of cement, lime and magnesium oxide (notified under document C(2013) 1728). http://data.europa.eu/eli/dec_impl/2013/163/oj

Formatted: English (United States)

Formatted: English (United States)

705 Commission Implementing Decision (EU) 2016/1032 of 13 June 2016 establishing best available techniques (BAT) conclusions, under Directive 2010/75/EU of the European Parliament and of the Council, for the non-ferrous metals industries. <https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX:32016D1032>.

710 Commission Implementing Decision (EU) 2019/2010 of 12 November 2019 establishing the best available techniques (BAT) conclusions, under Directive 2010/75/EU of the European Parliament and of the Council, for waste incineration (notified under document C(2019) 7987) (Text with EEA relevance). http://data.europa.eu/eli/dec_impl/2019/2010/oj/eng.

Commission Implementing Decision (EU) 2021/2326 of 30 November 2021, establishing best available techniques (BAT) conclusions, under Directive 2010/75/EU of the European Parliament and of the Council, for large combustion plants. <https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX:32021D2326>.

715 [Cui, Jian, Lunbo Duan, Ying Jiang, Changsui Zhao, and Edward John Anthony. 2018. "Migration and Emission of Mercury from Circulating Fluidized Bed Boilers Co-Firing Petroleum Coke and Coal." Fuel 215 \(March\). Elsevier BV: 638–46. doi:10.1016/j.fuel.2017.11.062. Cui, J., Duan, L., Jiang, Y., Zhao, C., and Anthony, E. J.: Migration and emission of mercury from circulating fluidized bed boilers co-firing petroleum coke and coal, Fuel, 215, 638–646, <https://doi.org/10.1016/j.fuel.2017.11.062>, 2018.](#)

Formatted: English (United States)

720 Dehoust, G., ~~inter, Peter~~-Gebhardt P., ~~Christian~~-Tebert C., and ~~Heinz~~-Köser, H.: ~~2021~~. Quecksilberemissionen Aus Industriellen Quellen – Status Quo Und Perspektiven; Abschlussbericht - Teil 2: Quecksilberminderungstechniken Und Überführung von Quecksilber in Senken. Edited by Katja Kraus. Vol. TEXTE 68/2021. Dessau-Rosslau: German Environment Agency. 2021.

725 Directive 2010/75/EU of the European Parliament and of the Council of 24 November 2010 on industrial emissions (integrated pollution prevention and control) (recast) (Text with EEA relevance), OJ L, 2010.

730 ~~Feeley, T. J., Jones, A. P., Brickett, L. A., O'Palko, B. A., Miller, C. E., and Murphy, J. T.: An update on DOE's Phase II and Phase III mercury control technology R&D program. Fuel Processing Technology, 90, 1388–1391, <https://doi.org/10.1016/j.fuproc.2009.05.012>, 2009.~~

735 ~~Feeley, Thomas J., Andrew P. Jones, Lynn A. Brickett, B. Andrew O'Palko, Charles E. Miller, and James T. Murphy. 2009. "An Update on DOE's Phase II and Phase III Mercury Control Technology R&D Program." Fuel Processing Technology 90 (11). Elsevier BV: 1388–91. doi:10.1016/j.fuproc.2009.05.012.~~

735 Flanagan, Daniel M.: ~~2022~~. "2018 Minerals Yearbook - Copper (Advance Release)." U.S. Geological Survey. <https://prd-wret.s3-us-west-2.amazonaws.com/assets/palladium/production/atoms/files/myb1-2015-manga.pdf>. 2022.

George, Michael, W. ~~2021~~. "2018 Minerals Yearbook - Mercury (Advance Release)." U.S. Geological Survey. <https://prd-wret.s3-us-west-2.amazonaws.com/assets/palladium/production/atoms/files/myb1-2015-manga.pdf>. 2021.

740 Giang, A., Stokes, L. C., Streets, D. G., Corbitt, E. S., and Selin, N. E.: Impacts of the Minamata Convention on Mercury Emissions and Global Deposition from Coal-Fired Power Generation in Asia. *Environmental Science & Technology*, 49, 5326–5335. <https://doi.org/10.1021/acs.est.5b00074>, 2015.

745 ~~Gómez-Sanabria, A., Kiesewetter, G., Klimont, Z., Schoepp, W., and Haberl, H.: Potential for future reductions of global GHG and air pollutants from circular waste management systems. Nat Commun, 13, 106, <https://doi.org/10.1038/s41467-021-27624-7>, 2022.~~

~~Granite, E. J., Pennline, H. W., and Hargis, R. A.: Novel Sorbents for Mercury Removal from Flue Gas. *Industrial & Engineering Chemistry Research*, 39, 1020–1029. <https://doi.org/10.1021/ie990758v>, 2000.~~

750 ~~Han, Y.-J., Kim, P.-R., Lee, G.-S., Lee, J.-I., Noh, S., Yu, S.-M., Park, K.-S., Seok, K.-S., Kim, H., and Kim, Y.-H.: Mercury concentrations in environmental media at a hazardous solid waste landfill site and mercury emissions from the site. *Environmental Earth Sciences*, 76. <https://doi.org/10.1007/s12665-017-6700-z>, 2017.~~

~~Giang, Amanda, Leah C. Stokes, David G. Streets, Elizabeth S. Corbitt, and Noelle E. Selin. 2015. "Impacts of the Minamata Convention on Mercury Emissions and Global Deposition from Coal-Fired Power Generation in Asia." *Environmental Science & Technology* 49 (9). American Chemical Society (ACS): 5326–35. doi:10.1021/aes.est.5b00074.~~

755 ~~Gómez-Sanabria, Adriana, Gregor Kiesewetter, Zbigniew Klimont, Wolfgang Schoepp, and Helmut Haberl. 2022. "Potential for Future Reductions of Global GHG and Air Pollutants from Circular Waste Management Systems." *Nature Communications* 13 (1). Springer Science+Business Media LLC. doi:10.1038/s41467-021-27624-7.~~

~~Granite, E. J., Pennline, H. W., and Hargis, R. A. 2000. "Novel Sorbents for Mercury Removal from Flue Gas." *Industrial & Engineering Chemistry Research*, 39, 1020–1029. doi: 10.1021/ie990758v.~~

Formatted: Normal

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Bibliography

Formatted: Font: 10 pt

Formatted: Font: 10 pt

760 Han, Young Ji, Pyung Rae Kim, Gang San Lee, Jae In Lee, Seam Noh, Seok Min Yu, Kwang Su Park, Kwang Seol Seok, Hyuk Kim, and Young Hee Kim. 2017. "Mercury Concentrations in Environmental Media at a Hazardous Solid Waste Landfill Site and Mercury Emissions from the Site." *Environmental Earth Sciences* 76 (10). Springer Science+Business Media LLC. doi:10.1007/s12665-017-6700-z.

IEA. 2022. *World Energy Outlook 2022*. Paris.

765 Keane, S., Bernaudat, L., Davis, K. J., Stylo, M., Mutemeri, N., Singo, P., Twala, P., Mutemeri, I., Nakafeero, A., and Etui, I. D.: Mercury and artisanal and small-scale gold mining: Review of global use estimates and considerations for promoting mercury-free alternatives, *Ambio*, <https://doi.org/10.1007/s13280-023-01843-2>, 2023.

770 Keane, Susan, Ludovic Bernaudat, Kenneth J. Davis, Malgorzata Stylo, Nellia Mutemeri, Patience Singo, Pontsho Twala, Hai Mutemeri, Anne Nakafeero, and Imelda Dossou Etui. 2023. "Mercury and Artisanal and Small-Scale Gold Mining: Review of Global Use Estimates and Considerations for Promoting Mercury-Free Alternatives." *Ambio* 52 (5). Springer Science+Business Media LLC: 833–52. doi:10.1007/s13280-023-01843-2.

Klochko, K. 2021. "2017 Minerals Yearbook - Lead (Advance Release)." U.S. Geological Survey. <https://prd-wret.s3-us-west-2.amazonaws.com/assets/palladium/production/atoms/files/myb1-2015-manga.pdf>. 2021.

775 Lee, S.-W., Lowry, G. V., and Hsu-Kim, H.: Biogeochemical transformations of mercury in solid waste landfills and pathways for release, *Environ. Sci.: Processes Impacts*, 18, 176–189, <https://doi.org/10.1039/C5EM00561B>, 2016.

Li, J., Zhou, S., Wei, W., Qi, J., Li, Y., Chen, B., Zhang, N., Guan, D., Qian, H., Wu, X., Miao, J., Chen, L., Feng, K., and Liang, S.: China's retrofitting measures in coal-fired power plants bring significant mercury-related health benefits, *One Earth*, 3, 777–787, <https://doi.org/10.1016/j.oneear.2020.11.012>, 2020.

780 Li, N., Chen, W., Rafaj, P., Kiesewetter, G., Schöpp, W., Wang, H., Zhang, H., Krey, V., and Riahi, K.: Air Quality Improvement Co-benefits of Low-Carbon Pathways toward Well Below the 2 °C Climate Target in China, *Environ. Sci. Technol.*, 53, 5576–5584, <https://doi.org/10.1021/acs.est.8b06948>, 2019.

785 Lindberg, S. E., Southworth, G. R., Bogle, M. A., Blasing, T. J., Owens, J., Roy, K., Zhang, H., Kuiken, T., Price, J., Reinhart, D., and Sfeir, H.: Airborne Emissions of Mercury from Municipal Solid Waste. I: New Measurements from Six Operating Landfills in Florida, *Journal of the Air & Waste Management Association*, 55, 859–869, <https://doi.org/10.1080/10473289.2005.10464684>, 2005.

Liu, K., Wang, S., Wu, Q., Wang, L., Ma, Q., Zhang, L., Li, G., Tian, H., Duan, L., and Hao, J.: A Highly Resolved Mercury Emission Inventory of Chinese Coal-Fired Power Plants, *Environmental Science & Technology*, 52, 2400–2408, <https://doi.org/10.1021/acs.est.7b06209>, 2018.

790 Lee, Sung Woo, Gregory V. Lowry, and Heileen Hsu Kim. 2016. "Biogeochemical Transformations of Mercury in Solid Waste Landfills and Pathways for Release." *Environmental Science: Processes & Impacts* 18 (2). Royal Society of Chemistry (RSC): 176–89. doi:10.1039/e5em00561b.

Li, N., Chen, W., Rafaj, P., Kiesewetter, G., Schöpp, W., Wang, H., Zhang, H., Krey, V., and Riahi, K.: Air Quality Improvement Co-benefits of Low-Carbon Pathways toward Well Below the 2 °C Climate Target in China, *Environ. Sci. Technol.*, 53, 5576–5584, <https://doi.org/10.1021/acs.est.8b06948>, 2019.

795 Li, Jiashuo, Sili Zhou, Wendong Wei, Jianchuan Qi, Yumeng Li, Bin Chen, Ning Zhang, et al. 2020. "Chinas Retrofitting Measures in Coal Fired Power Plants Bring Significant Mercury Related Health Benefits." *One Earth* 3 (6). Elsevier BV: 777–87. doi:10.1016/j.oneear.2020.11.012.

Formatted: Font: 10 pt

Formatted: Bibliography

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Field Code Changed

800 Lindberg, Steve E., George R. Southworth, Mary Anna Bogle, T.J. Blasing, Jim Owens, Kelly Roy, Hong Zhang, et al. 2005. "Airborne Emissions of Mercury from Municipal Solid Waste. I: New Measurements from Six Operating Landfills in Florida." *Journal of the Air & Waste Management Association* 55 (7). Informa UK Limited: 859–69. doi:10.1080/10473289.2005.10464684.

Liu, Kaiyun, Shuxiao Wang, Qingru Wu, Long Wang, Qiao Ma, Lei Zhang, Guoliang Li, Hezhong Tian, Lei Duan, and Jiming Hao. 2018. "A Highly Resolved Mercury Emission Inventory of Chinese Coal-Fired Power Plants." *Environmental Science & Technology* 52 (4). American Chemical Society (ACS): 2400–2408. doi:10.1021/acs.est.7b06209.

805 Minamata Convention on Mercury, CHAPTER XXVII, - ENVIRONMENT, Kumamoto, 10 October 2013. [hereinafter MCM] https://treaties.un.org/pages/ViewDetails.aspx?src=IND&mtdsg_no=XXVII-17&chapter=27&clang=en (retrieved 5 April, 2024).

810 Nakicenovic, N., Swart, R., Alcamo, J., Davis, G., Vries, B. de, Fenhann, J., Gaffin, S., Gregory, K., and Gruebler, A.: Special Report on Emissions Scenarios, Working Group III of the Intergovernmental Panel on Climate Change (IPCC), Cambridge University Press, Cambridge, UK, 2000.

Niksa Energy Associates LLC. 2011. *Interactive Process Optimization Guidance. User Guide and Tutorial*. Niksa Energy Associates LLC. 2011.

815 Pacyna, J. M., Sundseth, K., Pacyna, E. G., Jozewicz, W., Munthe, J., Belhaj, M., and Aström, S.: An Assessment of Costs and Benefits Associated with Mercury Emission Reductions from Major Anthropogenic Sources, *Journal of the Air & Waste Management Association*, 60, 302–315, <https://doi.org/10.3155/1047-3289.60.3.302>, 2010.

Pacyna, J. M., Travnikov, O., Simone, F. D., Hedgecock, I. M., Sundseth, K., Pacyna, E. G., Steenhuisen, F., Pirrone, N., Munthe, J., and Kindbom, K.: Current and future levels of mercury atmospheric pollution on a global scale, *Atmospheric Chemistry and Physics*, 16, 12495–12511, <https://doi.org/10.5194/acp-16-12495-2016>, 2016.

820 Pavlish, J. H., Sondreal, E. A., Mann, M. D., Olson, E. S., Galbreath, K. C., Laudal, D. L., and Benson, S. A.: Status review of mercury control options for coal-fired power plants, *Fuel Processing Technology*, 82, 89–165, [https://doi.org/10.1016/s0378-3820\(03\)00059-6](https://doi.org/10.1016/s0378-3820(03)00059-6), 2003.

825 Pacyna, Jozef M., Kyrre Sundseth, Elisabeth G. Pacyna, Wojciech Jozewicz, John Munthe, Mohammed Belhaj, and Stefan Aström. 2010. "An Assessment of Costs and Benefits Associated with Mercury Emission Reductions from Major Anthropogenic Sources." *Journal of the Air & Waste Management Association* 60 (3). Informa UK Limited: 302–15. doi:10.3155/1047-3289.60.3.302.

830 Pacyna, Jozef M., Oleg Travnikov, Francesco De Simone, Ian M. Hedgecock, Kyrre Sundseth, Elisabeth G. Pacyna, Frits Steenhuisen, Nicola Pirrone, John Munthe, and Karin Kindbom. 2016. "Current and Future Levels of Mercury Atmospheric Pollution on a Global Scale." *Atmospheric Chemistry and Physics* 16 (19). Copernicus GmbH: 12495–12511. doi:10.5194/acp-16-12495-2016.

835 Pavlish, J. H., Sondreal, E. A., Mann, M. D., Olson, E. S., Galbreath, K. C., Laudal, D. L., and Benson, S. A. 2003. "Status review of mercury control options for coal-fired power plants." *Fuel Processing Technology*. 82. 89–165, doi: 10.1016/s0378-3820(03)00059-6, Protocol to the 1979 Convention on Long-Range Transboundary Air Pollution on Heavy Metals, CHAPTER XXVII, - ENVIRONMENT, Aarhus, 24 June 1998, [hereinafter Aarhus Protocol] https://treaties.un.org/pages/ViewDetails.aspx?src=TREATY&mtdsg_no=XXVII-1-f&chapter=27&clang=en (retrieved April 5, 2024)

Formatted: Font: 10 pt

Formatted: Bibliography

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Bibliography

Formatted: Font: 10 pt

Formatted: Font: 10 pt, Not Highlight

Formatted: Font: 10 pt

Formatted: Font: 10 pt, German (Germany)

Formatted: Font: 10 pt

Rafaj, P., Bertok, I., Cofala, J., and Schöpp, W.: Scenarios of global mercury emissions from anthropogenic sources, *Atmospheric Environment*, 79, 472–479, <https://doi.org/10.1016/j.atmosenv.2013.06.042>, 2013.

Formatted: Font: 10 pt

Formatted: Bibliography

Rafaj, P., Cofala, J., Kuenen, J., Wyrwa, A., and Zysk, J.: Benefits of European Climate Policies for Mercury Air Pollution, *Atmosphere*, 5, 45–59, <https://doi.org/10.3390/atmos5010045>, 2014.

Formatted: Font: 10 pt

Rafaj, P., Kieseewetter, G., Gül, T., Schöpp, W., Cofala, J., Klimont, Z., Purohit, P., Heyes, C., Amann, M., Borken-Kleefeld, J., and Cozzi, L.: Outlook for clean air in the context of sustainable development goals, *Global Environmental Change*, 53, 1–11, <https://doi.org/10.1016/j.gloenvcha.2018.08.008>, 2018.

Formatted: Font: 10 pt

Rafaj, P., I. Bertok, J. Cofala, and W. Schöpp. 2013. “Scenarios of Global Mercury Emissions from Anthropogenic Sources.” *Atmospheric Environment* 79 (November). Elsevier BV: 472–79. doi:10.1016/j.atmosenv.2013.06.042.

Rafaj, Peter, Janusz Cofala, Jeroen Kuenen, Artur Wyrwa, and Janusz Zysk. 2014. “Benefits of European Climate Policies for Mercury Air Pollution.” *Atmosphere* 5 (1). MDPI AG: 45–59. doi:10.3390/atmos5010045.

Rafaj, P., Kieseewetter, G., Gül, T., Schöpp, W., Cofala, J., Klimont, Z., Purohit, P., Heyes, C., Amann, M., Borken-Kleefeld, J., and Cozzi, L.: Outlook for clean air in the context of sustainable development goals, *Global Environmental Change*, 53, 1–11, <https://doi.org/10.1016/j.gloenvcha.2018.08.008>, 2018.

Selin, N. E. and Selin, H.: Global Politics of Mercury Pollution: The Need for Multi-Scale Governance, *Review of European Community & International Environmental Law*, 15, 258–269, <https://doi.org/10.1111/j.1467-9388.2006.00529.x>, 2006.

Formatted: Normal

Selin, N. E.: Global Biogeochemical Cycling of Mercury: A Review, *Annual Review of Environment and Resources*, 34, 43–63, <https://doi.org/10.1146/annurev.environ.051308.084314>, 2009.

Selin, Noelle E. 2009. “Global Biogeochemical Cycling of Mercury: A Review.” *Annual Review of Environment and Resources* 34 (1). *Annual Reviews*: 43–63. doi:10.1146/annurev.environ.051308.084314.

Sheaffer, Kristin N.: 2022. “2018 Minerals Yearbook - Gold (Advance Release).” U.S. Geological Survey. <https://prd-wret.s3-us-west-2.amazonaws.com/assets/palladium/production/atoms/files/myb1-2015-manga.pdf>. 2022.

Tolcin, Amy C.: 2022. “2018 Minerals Yearbook - Zinc (Advance Release).” U.S. Geological Survey. <https://prd-wret.s3-us-west-2.amazonaws.com/assets/palladium/production/atoms/files/myb1-2015-manga.pdf>. 2022.

UNEP. 2013. “Minamata Convention on Mercury—Text and Annexes,” 69p.

UNEP Chemicals. 2002. “Global Mercury Assessment.” United Nations Environment Programme. <https://wedocs.unep.org/bitstream/handle/20.500.11822/12297/final-assessment-report-25nov02.pdf?sequence=1&isAllowed=y>.

Usberti, N., Alcove Clave, S., Nash, M., and A. Beretta, 2016. “Kinetics of Hg⁰ oxidation over a V2O5/MoO3/TiO2 catalyst: Experimental and modelling study under DeNOX inactive conditions.” *Applied Catalysis B: Environmental* 193, 121–132, <https://doi.org/10.1016/j.apcatb.2016.03.071>, 2016. Usberti, N., Alcove Clave, S., Nash, M., and Beretta, A.: Kinetics of Hg⁰ oxidation over a V2O5/MoO3/TiO2 catalyst: Experimental and modelling study under DeNOX inactive conditions, *Applied Catalysis B: Environmental*, 193, 121–132, <https://doi.org/10.1016/j.apcatb.2016.03.071>, 2016.

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Wagner, F., C. Heyes, C. Z. Klimont, Z., and W. Schoepp W.: 2013. “The Gains Optimization Module: Identifying Cost-Effective Measures for Improving Air Quality and Short-Term Climate Forcing.” IIASA Interim Report. IIASA, Laxenburg, Austria: IR-13-001, <https://pure.iiasa.ac.at/id/eprint/10755/>, 2013.

Formatted: Default Paragraph Font, Font: 12 pt

Field Code Changed

875 Wang, F., Wang, S., Zhang, L., Yang, H., Gao, W., Wu, Q., and Hao, J.: Mercury mass flow in iron and steel production process and its implications for mercury emission control. Journal of Environmental Sciences, 43, 293–301, <https://doi.org/10.1016/j.jes.2015.07.019>, 2016.

Formatted: Font: 10 pt

Formatted: Bibliography

880 Wang, Z., Zhang, Y., Wang, L., Li, X., Zhou, X., Li, X., Yan, M., Lu, Q., Tang, Z., Zhang, G., and Wang, D.: Characteristics and Risk Assessments of Mercury Pollution Levels at Domestic Garbage Collection Points Distributed within the Main Urban Areas of Changchun City, Toxics, 9, 309, <https://doi.org/10.3390/toxics9110309>, 2021.

Formatted: Default Paragraph Font, Font: 10 pt

Formatted: Normal

885 Wang, Fengyang, Shuxiao Wang, Lei Zhang, Hai Yang, Wei Gao, Qingru Wu, and Jiming Hao. 2016. "Mercury Mass Flow in Iron and Steel Production Process and Its Implications for Mercury Emission Control." Journal of Environmental Sciences 43 (May). Elsevier BV: 293–301. doi:10.1016/j.jes.2015.07.019.

890 Wang, Zhaojun, Yangjie Zhang, Lei Wang, Xu Li, Xuhang Zhou, Xiangyun Li, Mengping Yan, et al. 2021. "Characteristics and Risk Assessments of Mercury Pollution Levels at Domestic Garbage Collection Points Distributed Within the Main Urban Areas of Changchun City." Toxics 9 (11). MDPI AG: 309. doi:10.3390/toxics9110309.

World Health Organization 2021. Chemicals. In: Compendium of WHO and other UN guidance on health and environment. (WHO/HEP/ECH/EHD/21.02). Geneva: World Health Organization. 2021

895 Wu, Qingru, Guoliang Li, Shuxiao Wang, Kaiyun Liu, and Jiming Hao. 2018. "Mitigation Options of Atmospheric Hg Emissions in China." Environmental Science & Technology 52 (21). American Chemical Society (ACS): 12368–75. doi:10.1021/acs.est.8b03702. Wu, Q., Li, G., Wang, S., Liu, K., and Hao, J.: Mitigation Options of Atmospheric Hg Emissions in China, Environmental Science & Technology, 52, 12368–12375, <https://doi.org/10.1021/acs.est.8b03702>, 2018.

Formatted: Bibliography

Formatted: Font: 10 pt

900 Zhang, S., Hui, L., Wang, S.: 2016. "Potential of Co-Benefit Mercury Control for Coal-Fired Power Plants and Industrial Boilers in China." Natural Resource Defence Council. <https://www.nrdc.org/sites/default/files/co-benefit-mercury-control-report.pdf>, 2016.

Formatted: English (United States)

Formatted: Font: 10 pt

Formatted: English (United States)

905 Zhang, Yanxu, Zhengcheng Song, Shaojian Huang, Peng Zhang, Yiming Peng, Peipei Wu, Jing Gu, et al. 2021. "Global Health Effects of Future Atmospheric Mercury Emissions." Nature Communications 12 (1). Springer Science+Business Media LLC. doi:10.1038/s41467-021-23391-7.

910 Zhang, Y., Song, Z., Huang, S., Zhang, P., Peng, Y., Wu, P., Gu, J., Dutkiewicz, S., Zhang, H., Wu, S., Wang, F., Chen, L., Wang, S., and Li, P.: Global health effects of future atmospheric mercury emissions, Nature Communications, 12, <https://doi.org/10.1038/s41467-021-23391-7>, 2021.

Formatted: Font: 10 pt

Formatted: Bibliography