

Comprehensive increase in CO₂ release by drying-rewetting cycles among Japanese forests and pastureland soils and exploring predictors of increasing magnitude

Yuri Suzuki^{1,2}, Syuntaro Hiradate³, Jun Koarashi⁴, Mariko Atarashi-Andoh⁴, Takumi Yomogida⁴, Yuki Kanda¹, and Hirohiko Nagano⁵

¹Faculty of Agriculture, Niigata University, Niigata 950-2181, Japan

²Graduate School of Environmental Science, Niigata University, Niigata 950-2181, Japan

³Graduate school of Agriculture, Kyushu University, Fukuoka 819-0395, Japan

⁴Nuclear Science and Engineering Center, Japan Atomic Energy Agency, Tokai, Ibaraki 319-1195, Japan

⁵Institute of Science and Technology, Niigata University, Niigata 950-2181, Japan

Correspondence: Hirohiko Nagano (hnagano@agr.niigata-u.ac.jp)

Abstract. It is still difficult to precisely quantify and predict the effects of drying-rewetting cycles (DWCs) on soil carbon dioxide (CO₂) release due to the paucity of studies using constant moisture conditions equivalent to the mean water content during DWC incubation. The present study was performed to evaluate overall trends in the effects of DWCs on CO₂ release and to explore environmental and soil predictors for variations in the effect size in 10 Japanese forests and pastureland soils variously affected by volcanic ash during their pedogenesis. Over an 84-day incubation period including three DWCs, CO₂ release was 1.3- to 3.7-fold greater than under continuous constant moisture conditions ($p < 0.05$) with the same mean water content as in the DWC incubations. Analysis of the relations between this increasing magnitude of CO₂ release by DWCs (IF_{CO₂}) and various environmental and soil properties revealed significant positive correlations between IF_{CO₂} and soil organo-metal complex contents ($p < 0.05$), especially pyrophosphate extractable aluminum (Alp) content ($r = 0.74$). Molar ratios of soil total carbon (C) and pyrophosphate-extractable C (Cp) to Alp contents and soil carbon content-specific CO₂ release rate under continuous constant moisture conditions (qCO₂_soc) were also correlated with IF_{CO₂} ($p < 0.05$). The covariations among Alp, total C, and Cp to Alp molar ratios and qCO₂_soc suggested Alp as the primary predictor of IF_{CO₂}. Additionally, soil microbial biomass C and nitrogen (N) levels were significantly lower in DWCs than under continuous constant moisture conditions, whereas there was no significant relation between the microbial biomass decrease and IF_{CO₂}. The present study showed a comprehensive increase in soil CO₂ release by DWC in Japanese forests and pastureland soils, suggesting that Alp is a predictor of the effect size likely due to vulnerability of organo-Al complexes to DWC.

Copyright statement. TEXT

1 Introduction

There is accumulating evidence of climate change-induced alterations in global water cycling (IPCC, 2021; Allan et al., 2020; 20 Dai, 2012; Donat et al., 2016; Pfliegerer et al., 2019). Of the consequent water regime changes, decreasing precipitation frequency (e.g., number of rainy days) and increasing intensity (e.g., number of heavy rainy days) are becoming more frequent (Dai, 2012; Donat et al., 2016), although decadal trends in annual precipitation levels are not significant over global scales (IPCC, 2021). For example, the annual number of precipitation days in Japan has decreased by 15% during the past 120 years, whereas the annual number of heavy precipitation days (more than 100 mm in a day) has increased by 26% (Ministry of 25 Education, Science, and Technology (MEXT) and Japanese Meteorological Agency (JMA), 2020). There is a non-significant trend in annual precipitation level during the same period (MEXT and JAM, 2020). This changing pattern in precipitation is often observed in the temperate region of the northern hemisphere (IPCC, 2021) and is related to increased fluctuation of soil water environments, especially DWCs, and consequent alterations in ecosystem functions (Borken and Matzner, 2009; Jin et al., 2023; Zhang et al., 2020, 2023).

30 Carbon dioxide (CO₂) release from soil is an ecosystem process that is sensitive to DWCs (Birch, 1958; Borken and Matzner, 2009; Lee et al., 2002; Nagano et al., 2019; Unger et al., 2010, 2012; Zhang et al., 2020, 2023) and has substantial feedback potential to the ongoing climate change due to its magnitude reaching as much as seven times greater than anthropogenic CO₂ emission on a global scale (Bond-Lamberty and Thomson, 2010; Friedlingstein et al., 2020). The effects of DWCs on soil CO₂ release were first shown by Birch (1958) as the marked increase in soil organic matter (SOM) decomposition and CO₂ release 35 after the rapid rewetting of dried soil, and has since been the subject of intensive investigation (Borken and Matzner, 2009; Kpemoua et al., 2023; Lee et al., 2002, 2004; Miller et al., 2005; Nagano et al., 2019; Unger et al., 2010, 2012; Xiang et al., 2008), including meta-analyses (Kim et al., 2012; Jin et al., 2023; Zhang et al., 2020). However, it is still difficult to precisely quantify and predict the effects of DWCs on soil CO₂ release.

The significant uncertainties in the effects of DWCs on soil CO₂ release include the inconsistent trends and sizes of effects 40 likely due to the paucity of studies using constant moisture conditions equivalent to the mean water content during DWC incubation (Kpemoua et al., 2023; Zhang et al., 2020, 2023). According to a meta-analysis by Zhang et al. (2020) using 208 data from 34 sites in 29 reports, the effects of DWCs vary according to soil and water contents in continuous constant moisture conditions. Especially, changes in CO₂ release rate associated with DWCs ranged from -4% to +19% with an average of +4% in comparison with the medium level of constant moisture content, which should be the same with the mean water content during 45 DWC incubation, whereas only 9 of 38 data representing CO₂ release rates were measured for such medium level of constant moisture conditions (Zhang et al., 2020). Another 29 data were calculated as the average of two CO₂ release rates at the wettest and driest water contents of constant moisture conditions, which should be the same with the maximum and minimum water contents in DWC treatment, respectively (Zhang et al., 2020). In the experiment using three Alfisols from Chinese long-term experimental field studies, Zhang et al. (2023) showed similar or somewhat lower CO₂ release in the DWC compared with 50 the constant moisture conditions with the same mean water content for the DWC incubation. Using two Luvisols from French long-term field experiment sites, Kpemoua et al. (2023) also showed similar features of changes in CO₂ release associated with

DWCs, indicating the need for further comparison of CO₂ release between DWCs and constant moisture conditions with the same mean water content. In contrast to these studies, Nagano et al. (2019) found a 49% increase in CO₂ release rate associated with DWCs in an Andisol collected from a Japanese forest. This increase was more than double that of another non-volcanic ash soil from the same forest. Thus, there are substantial variations in trends of effects of DWCs in comparison with constant moisture conditions having the same mean water content during incubation, remaining knowledge gaps about environmental and soil predictors for variations in effect sizes. There are proposed roughly three mechanisms for CO₂ release increase by DWCs (Schimel, 2018; Barnard et al., 2020): (i) increase in available carbon source via the releases of cellular metabolites from microbial cells destroyed by rewetting after the strong drought, (ii) increase in available carbon source by the releases of carbon from macroaggregates destroyed by repeated DWC, and (iii) changes in the microbial communities in response to transient moisture conditions. Nevertheless, there are still substantial knowledge gaps for critical mechanisms or the relative importance of those mechanisms among multiple soils.

We perform the present study to evaluate overall trends in the effect of DWCs on soil CO₂ release and to explore the predictors of variations in its effect size among 10 Japanese forests and pastureland soils. These soils are variously affected by volcanic ash during their pedogenesis and, therefore, include several Andisols, which are known to have a high SOM storage capacity (Morisada et al., 2004), likely due to the protection of SOM from microbial decomposition by abundant reactive minerals and metals in these soils (Asano and Wagai, 2014; Imaya et al., 2007; Shirato et al., 2004). Reactive minerals and metals that contribute to the protection of SOM are iron (Fe) and aluminum (Al), constituting short-range-order minerals and organo-metal complexes (Asano and Wagai, 2014; Rasmussen et al., 2018; Shirato et al., 2004; Wagai et al., 2018). Although global coverage of Andisols is about 1% (FAO/IIASA/ISRIC/ISS-CAS/JRC, 2009), determination of the responses of carbon cycling in Andisols to DWCs will help in understanding the responses of non-volcanic ash soils, because reactive minerals and metals are also essential in high carbon stocks of those soils (Rasmussen et al., 2018; Hall and Thompson, 2022) and may be sensitive to climate and land use management, including water regime (Georgiou et al., 2022; Kramer and Chadwick, 2018).

2 Materials and methods

2.1 Site description and soil sampling

We collected 10 soil samples from depths of 0–5 cm or 0–10 cm in six forests and a pastureland located in Niigata (six soils from four forests), Ibaraki (two soils from a forest) (Nagano et al., 2019), and Oita (two soils from a forest and a pastureland) (Wijesinghe et al., 2021) prefectures in Japan. Figure 1 and Table 1 present the locations and site characteristics, i.e., elevation, mean annual temperature (MAT), mean annual precipitation (MAP), potential evapotranspiration (PET), and net primary production (NPP). Briefly, all of the investigated sites have a humid temperate climate with MAT of 9.1–10.8°C and MAP of 1474–2930 mm. All of the forests are dominated by beech (*Fagus crenata* and *Fagus japonica*) and oak (*Quercus serrata*), except for Oita forest which is a deciduous/evergreen mixed forest dominated by Siebolds maple (*Acer sieboldianum*), Japanese snowbell (*Styrax japonicus*), and Japanese holly (*Ilex crenata*). The pastureland in Oita is dominated by Japanese lawn grass




Figure 1. Locations of Japanese forests and pastureland where investigated soils were collected.

Table 1. Environmental properties of Japanese forests and pastureland where investigated soils were collected^{a)}

Prefecture	Soil	Latitude °N	Longitude °E	Elevation m	MAT °C	MAP mm	Annual PET mm	MAP-PET mm	Annual NPP g C m ⁻²
Niigata	Aga1, Aga2	37.55	139.51	474	10.6	1714	747	967	633
	Tsunan1, Tsunan2	37.04	138.60	716	10.8	1899	795	1104	628
	Asahi1	38.38	139.70	655	9.2	1887	697	1190	605
	Oshirakawa1	37.35	139.16	627	9.1	1963	668	1295	563
Ibaraki	Ogawa13, Ogawa 14	36.93	140.59	643	10.7	1474	764	710	785
Oita	Kuju_pasture, Kuju_forest	33.06	131.23	841	10.8	2930	786	2144	819

^{a)} MAT, MAP, and PET were obtained as averages for 1981–2020 in a global data set of climate and climatic water balance (i.e., TerraClimate) by Abatzoglou et al. (2018). NPP data were obtained as averages for 2001–2020 in the global distribution of NPP estimated from MODIS observation products (i.e., MOD17A3HGF) by Running and Zhao. (2021). MAT, mean annual temperature; MAP, mean annual precipitation; PET, potential evapotranspiration; NPP, net primary production.

(*Zoysia japonica*), dwarf fountain grass (*Pennisetum alopecuroides*), cranesbill (*Geranium thunbergii*), white clover (*Trifolium repens*), and Indian strawberry (*Potentilla indica*).

Soil sampling was conducted in the snow-free season (April to October) of 2021. We also collected soil samples from layers below the target depth (i.e., 0–5 cm or 0–10 cm depth) down to 50 cm as the maximum depth to examine whether the soil could be classified as Andisol. According to the USDA (United States Department of Agriculture) Soil Taxonomy criteria (USDA Soil Survey Staff, 2022), soils with 60% or more of the thickness containing more than 20 mg of acid oxalate-extractable Al (Alo) plus 1/2 Fe (Feo) per 1 g soil within a depth of 0–60 cm are classified as Andisols. According to these criteria, 4 of the 10 soils were Andisols (one each from Niigata and Ibaraki, and two from Oita) with non-allophanic properties determined by a

high ratio of pyrophosphate-extractable Al (Alp) content to Alo content (Alp/Alo > 0.5; USDA Soil Survey Staff (2022); Fig. S1). Collected soil samples were transferred to the laboratory and stored at 4°C before further analysis. Before analysis, soils were gently passed through a 4-mm sieve to remove gravel and plant tissue. Fine roots in the sieved samples were removed with tweezers. In our study, we considered that the soil water content at the soil sampling reflected the ability of soil to hold the water and thus the usual water contents in the field because the soil water content showed significantly positive correlations with water holding capacity (WHC) ($r = 0.87$, $p < 0.01$). Therefore, CO₂ release rate for constant moisture conditions in the present study should represent the release rate under the usual field moisture conditions of each soil.

2.2 Soil analysis

The soil properties analyzed were pH(H₂O), electrical conductivity, water content, water-holding capacity (WHC), total carbon (C) and nitrogen (N) contents, particle size distributions as relative compositions of clay, silt, and sand-sized particles, and selectively dissolved Al and Fe contents (Tables 2 – 4). The pH(H₂O) was measured in soil and water mixtures consisting of 1 g of soil and 2.5 mL of water. For measurement of electrical conductivity, 5 mL of water was added to 1 g of soil. Water content was measured by determining the difference in soil weight before and after drying at 105°C for 24 h. WHC was measured as the difference in soil weight before and after water saturation referring to the Hilgard method (Mabuhay et al., 2003; Ahn et al., 2008). Here, water contents when soil is completely saturated in the Hilgard method should equal zero pF value (0 kPa) as soil water potential. Soil total C and N contents were measured for air-dried and well-ground soil samples using an elemental analyzer (vario PYRO cube; Elementar, Manchester, UK). Particle size distributions were determined using the Stokes' law-based sedimentation method (Miller et al., 1988; Nakai, 1997) using soil mineral particles after removing organic matter by hydrogen peroxide solution digestion. Selectively dissolved metals such as reactive Al and Fe were measured according to the procedure described previously in Nagano et al. (2023). Briefly, the contents of Al and Fe extractable with 2.0 M acid ammonium oxalate (i.e., Alo and Feo, respectively) were measured as contents of organo-metal complexes and short-range-order minerals, while Al and Fe extractable with 0.1 M sodium pyrophosphate solution (i.e., Alp and Fep, respectively) were measured as contents of organo-metal complexes (Asano and Wagai, 2014; Takahashi and Dahlgren, 2016; Wagai et al., 2018). The difference between acid oxalate- and pyrophosphate-extractable metals (i.e., Alo-p and Feo-p) represented the contents of short-range-order minerals (Courchesne and Turmel, 2008). Contents of Al and Fe in the solution were measured with an inductively coupled plasmaoptical emission spectrometer (ICP-OES) (5110; Agilent Technologies, Santa Clara, CA, USA). Carbon concentrations in pyrophosphate-extracted solution (Cp) were also measured with a total organic carbon (TOC) analyzer (TOC-L; Shimadzu, Kyoto, Japan). For soils from Kuju forest and grassland and two Ogawa forests, we also measured C concentration in free light density fraction (fLF) (Leuthold et al., 2023) obtained by the density fractionation method using sodium polytungstate solution having 2.0 g cm⁻³ (Koarashi et al., 2012), in order to evaluate possible effects of labile C abundance in fLF on CO₂ release increase by DWCs.

For the soils after incubation (see Section 2.3), soil microbial biomass C and N were measured by the chloroform fumigation-extraction method (Vance et al., 1987). Organic C and total N concentrations in 0.5 M potassium sulfate solution used for

Table 2. Basic properties of 10 Japanese forests and pastureland soils collected from depths of 0–5 or 0–10 cm

Soil	Soil type ^{a)}	pH(H ₂ O)	EC	Water content	WHC	Particle size-distribution		
						Sand	Silt	Clay
			$\mu\text{S cm}^{-1}$	$\text{g water g}^{-1} \text{ soil}$	$\text{g water g}^{-1} \text{ soil}$	%		
Aga1	Inceptisols or Entisols	4.36	23	0.77	0.80	27	30	43
Aga2	Inceptisols or Entisols	4.39	27	1.07	1.12	17	59	23
Tsunan1	Inceptisols (Brown forest soils)	4.72	17	0.78	1.06	47	44	9
Tsunan2	Inceptisols (Brown forest soils)	4.08	39	1.06	1.32	34	48	18
Asahi1	Inceptisols (Brown forest soils)	4.07	39	1.70	2.14	29	41	30
Oshirakawa1	Andisols	4.66	24	1.07	1.26	38	45	17
Ogawa13	Inceptisols (Brown forest soils)	5.29	18	0.46	1.00	66	25	8
Ogawa14	Andisols	5.30	27	1.14	1.68	68	23	9
Kuju_pasture	Andisols	5.35	133	1.32	1.43	72	22	5
Kuju_forest	Andisols	3.80	168	1.49	1.58	70	25	4

^{a)} Andisols were determined according to the USDA Soil Taxonomy criteria (USDA Soil Survey Staff, 2022) based on acid oxalate-extractable Al plus 1/2 Fe contents (see text for details). All Andisols were non-allophanic. Other soil types were determined using a Japanese soil digital map, i.e., Japan Soil Inventory NARO (2023).

125 extraction of fumigated and nonfumigated soils were measured using another TOC analyzer (TOC-L; Shimadzu) equipped with a total nitrogen (TN) unit (TNM-L, Shimadzu).

2.3 Incubation experiment and soil CO₂ release rate measurement

130 Soils were incubated aerobically at 20°C for 84 days including three DWCs (i.e., 28 days per cycle). Simultaneously, soils were incubated in the same manner but without DWCs, during which water content of the soils was maintained at a constant level equivalent to the mean water content for the DWC treatment (Fig. 2). A pre-incubation was conducted at the constant water content for 7 days prior to the 84-day incubation. A post-incubation was also conducted at the constant water content for 28 days after the 84-day incubation to evaluate the remaining effect of DWCs on soil CO₂ release. We consider that the incubated soils have been aerobic even after the rewetting to increase the water content by twice the WHC, because the CO₂ concentrations in our experiment never overwhelmed 1%, thus the oxygen concentrations in the incubation jar have likely never

Table 3. Soil carbon (C) and nitrogen (N) properties

Soil	Total C %	Total N %	Total C/N	C in free light density fraction (fLF) % to total C
Aga1	8.8	0.56	15.7	No data
Aga2	14.3	0.79	18.2	No data
Tsunan1	12.4	0.63	19.8	No data
Tsunan2	15.3	0.98	15.6	No data
Asahi1	23.7	1.41	16.9	No data
Oshirakawa1	15.1	0.76	19.8	No data
Ogawa13	6.8	0.31	22.2	10.0
Ogawa14	16.1	0.86	18.6	10.0
Kuju_pasture	21.8	1.37	15.9	14.0
Kuju_forest	22.3	1.18	18.9	5.9

Table 4. Selectively dissolved minerals and associated carbon contents in soils

Soil	Acid oxalate- extractable metals			Pyrophosphate- extractable metals			Acid oxalate – Pyrophosphate			Alp/Alo	Pyrophosphate- extractable C mg g ⁻¹ dry soil
	Al	Fe	Al+0.5Fe	Al	Fe	Al+0.5Fe	Al	Fe	Al+0.5Fe		
	mg g ⁻¹ dry soil			mg g ⁻¹ dry soil			mg g ⁻¹ dry soil				
Aga1	3.3	7.9	7.2	2.8	5.0	5.3	0.5	2.8	2.0	0.84	21.0
Aga2	6.9	10.6	12.2	6.8	10.5	12.0	0.1	0.2	0.2	0.98	42.9
Tsunan1	8.9	13.8	15.8	8.5	11.3	14.2	0.4	2.5	1.6	0.96	39.9
Tsunan2	6.9	8.5	11.1	7.1	7.2	10.6	-0.2	1.4	0.5	1.03	47.4
Asahi1	4.8	10.2	9.9	4.7	9.2	9.3	0.1	1.0	0.6	0.98	45.4
Oshirakawa1	9.5	12.4	15.7	9.1	11.0	14.6	0.4	1.4	1.2	0.95	43.0
Ogawa13	9.2	8.7	13.5	6.9	6.5	10.2	2.3	2.2	3.4	0.75	19.4
Ogawa14	24.9	16.6	33.2	23.2	12.7	29.6	1.6	3.9	3.6	0.93	59.1
Kuju_pasture	22.0	20.3	32.2	19.0	14.1	26.0	3.1	6.3	6.2	0.86	96.7
Kuju_forest	19.5	19.1	29.0	19.0	16.2	27.1	0.4	2.9	2.0	0.98	109.8

135 decreased below 19% or lower. Also, a sufficiently large volume of our incubation jar (1.0L) against contained soil amounts (i.e., 5.31-10.63 g) and added water contents in the rewetting (i.e., ca. 6 to 7 mL) support the state of aerobic condition during the incubation.




Figure 2. Schematic time courses of soil moisture during incubation with constant moisture content (a) and DWCs (b) with measurement period of CO₂ release rate.

Mason jars (1.0 L volume; Ball, Buffalo, NY, USA) with lids equipped with tube fitting systems for gas sample collection (Koarashi et al., 2012; Nagano et al., 2019) were used as incubation jars. Small vials (300 mL, SM sample glass vial; Sansyo, Tokyo, Japan) containing 5.31–10.63 g of soil sample depending on water content were placed in Mason jars. Under the constant water treatment, we surrounded the small vial with 20 mL of water within the incubation jar to prevent the soil from drying. For each DWC, Day 1 to Day 7 and Day 18 to Day 24 were drying stages (Fig. 2), during which the soils were incubated with silica gel (20 g jar⁻¹), which lowered the water content to < 5% WHC by Day 7. Day 8 to Day 12, Day 13 to Day 17, and Day 25 to Day 28 were the driest, wettest, and moderately wet stages, respectively, and CO₂ release rates in these three stages were measured using a gas chromatograph equipped with a thermal conductivity detector (GC-14B; Shimadzu). The CO₂ release rates were also measured in the pre- and post-incubation periods. At the beginning of the wettest stage, soils were rapidly rewetted with distilled water to double the soil water content from the initial status of DWC incubation. For CO₂ release rate measurements, at the start of each stage, the headspace of the incubation jar was flushed with CO₂-free air for 15 min at a rate of 0.5 L min⁻¹ and the jar was closed. At the end of each stage, 15 mL of gas sample was collected from the jar using a 20-mL plastic syringe (Terumo, Kyoto, Japan) and stored in a pre-evacuated 5-mL glass vial (SVG-5; Nichiden Rika, Osaka, Japan). Then, the CO₂ release rate was determined from the increase in CO₂ concentration during this period. After gas sampling, the jars were flushed with CO₂-free air and closed for the next incubation stage. For 84 days in the incubations

with the constant moisture condition as controls, the CO₂ release rates were measured for Day 1 to Day 12, Day 13 to Day 28, Day 29 to Day 40, Day 41 to Day 56, Day 57 to Day 68, and Day 69 to Day 84, in addition to the pre- and post-incubation periods. Soil water contents during the incubation were measured periodically and maintained by adding water to ensure the same mean water content between the two treatments. Even in the drying stage under the DWC treatment for Day 1 to Day 7 and Day 18 to Day 24, we conducted measurements of soil water content once to twice. The measurements were performed by weighing those soils. Based on these data, we confirmed that the mean soil water content during DWC incubation was equal to that during constant moisture incubation. All incubations were conducted with three replicates for each treatment and soil.

2.4 Data processing and statistical analysis

The CO₂ release rates were compared between the DWC and constant water content treatments. For the DWC treatment, the CO₂ release rates in the drying stages (i.e., Day 1 to Day 7 and Day 18 to Day 24) could not be measured and therefore had to be estimated to evaluate the mean CO₂ release rates for the individual cycle and total of three cycles. The rates in the drying stages were estimated as the mean values of the CO₂ release rates measured before and after the period of interest, in the same manner as described previously (Nagano et al., 2019). Then, the effect size of the DWCs on CO₂ release (defined here as the increase factor, IF_{CO₂}) was quantified as the ratio of CO₂ release rate under the DWC condition to that under the constant water content condition (Nagano et al., 2019; Zhang et al., 2020). Here, doubling of CO₂ release by DWC resulted in IF_{CO₂} of 2, while halving resulted in IF_{CO₂} of 0.5.

The pairwise *t* test was applied to examine the statistical significance of differences in CO₂ release rates between the DWC and constant water content treatments with adjustment of site-by-site variations in the metrics. Differences in soil microbial biomass C and N between the two treatments were also evaluated with the pairwise *t* test. To explore predictors explaining the variation in IF_{CO₂} among soils, relations between IF_{CO₂} and environmental and soil properties were visualized on scatter plots and evaluated by linear correlation analysis. All statistical analyses were conducted with R 4.1.1 (R Core Team, Vienna, Austria), and *p* < 0.05 was taken to indicate statistical significance.

3 Results

3.1 Quantifying the effect of DWCs on soil CO₂ release

The CO₂ release rates under DWC conditions showed large fluctuations for all soils along with fluctuations in soil water content (Fig. 3). The CO₂ release rates in the driest stages (3.5%–18.2% of WHC depending on the soil) were 3.0–41.5 μg C g⁻¹ dry soil day⁻¹. In contrast, CO₂ release rates in the wettest stages (90.8%–201.1% of WHC) reached 18.8–194.1 μg C g⁻¹ dry soil day⁻¹. In the moderately wet stages, CO₂ release rates were 8.4–63.9 μg C g⁻¹ dry soil day⁻¹. For the constant water content treatment, CO₂ release rates observed for each soil showed little fluctuation during incubation although the rates varied from 7.3 to 69.9 μg C g⁻¹ dry soil day⁻¹ depending on the soil.




Figure 3. Time courses of soil CO₂ release rates (a and b) and water contents (c and d) during 84-day incubation under constant water content (a and c) and three DWC conditions (b and d).

For all soils, the observed large fluctuations in CO₂ release under DWC treatment resulted in a greater CO₂ release rate than under constant water content treatment, although the IF_{CO₂} values varied among the soils (Fig. 4). At the wettest stage in the first DWC, the CO₂ release rates increased by 47.2–127.7 μg C g⁻¹ dry soil day⁻¹ compared with those under the constant water content conditions, resulting in IF_{CO₂} values of 2.9–12.2. For the whole of the first cycle (i.e., the first 28 days), the CO₂ release rates increased by 15.4–43.4 μg C g⁻¹ dry soil day⁻¹ under the DWCs compared with the constant water content conditions, resulting in IF_{CO₂} values of 1.6–5.2. For the whole incubation period (84 days) including three DWCs, IF_{CO₂} values were 1.3–3.7, with an increase in CO₂ release rate by 7.4–23.8 μg C g⁻¹ dry soil day⁻¹ by DWCs. These increases in CO₂ release by DWCs were observed in all cycles during the 84-day incubation period, whereas no increase was observed in the 28-day post-incubation period after the three DWCs (Fig. S2). Considering these results, we focused on the IF_{CO₂} variations obtained for the whole incubation period, including three DWCs.

3.2 Exploring predictors of the effect sizes on the increase in CO₂ release

Among the environmental and soil physiochemical properties, reactive mineral and metal contents (especially, Alo+0.5Feo, Alo, Feo, Alp+0.5Fep, and Alp) in soils showed significant positive correlations with IF_{CO₂} ($p < 0.05$; Table 5). In particular, Alp content appeared to be a key predictor of the variation in IF_{CO₂} among the soils, given that Alp accounted for most (73%–99%) of Alo (Table 4) and showed a higher correlation coefficient than others (Table 5). Scatterplots for Alp content and IF_{CO₂} values are presented in Fig. 5. In addition, the molar ratios of soil total C and Cp contents to Alp contents showed significant negative correlations to IF_{CO₂} ($p < 0.05$; Table 5, Fig. 5). Here, it should be noted that IF_{CO₂} had no significant relationship with soil water content at soil sampling and WHC, suggesting the variation in IF_{CO₂} among soils resulting from




Figure 4. Comparisons of mean CO_2 release rates after rewetting in the first cycle, the whole of the first cycle, and for the total of three cycles between DWc and constant water content conditions (a-c), and the factor of increase in CO_2 release by DWCs (IF_{CO_2}) for individual periods (d-f). Statistically significant differences ($p < 0.05$, pairwise t test) in CO_2 release rate between the two treatments are presented.

other than such hydrogenic properties of soils. The SOM quality, such as C/N ratios of total bulk soils and K_2SO_4 extractable fractions, also had no significant relationship with IF_{CO_2} . The insignificant relations of SOM quality to IF_{CO_2} were also supported by the facts that Kuju forest and grassland soils having more than doubling differences in fLF abundance showed only 12% differences in IF_{CO_2} , and two Ogawa forest soils having almost identical fLF abundances showed more than doubling differences in IF_{CO_2} . The amounts of clay and sand-sized particles showed significant correlations with IF_{CO_2} after rewetting in the first cycle ($p < 0.05$, $r = -0.66$ for clay and 0.71 for sand particles). However, those correlations between the particle contents and IF_{CO_2} were insignificant for IF_{CO_2} for a total of three cycles. Thus, in the present study, reactive mineral and metal content, especially Alp content in soils, was the significant predictor for IF_{CO_2} variation among soils, rather than soil water environments, SOM quality, and particle size distributions.

Among the additional soil properties related to soil microbial activity and abundance, the soil C content-specific CO_2 release rate under constant moisture conditions (defined as the CO_2 release rate per unit of C in soil, $\text{qCO}_2\text{-soc}$, in $\mu\text{g CO}_2\text{-C mg}^{-1}$ soil-C day^{-1}) showed significant negative correlations with IF_{CO_2} values in all incubation stages ($p < 0.01$; Table 6, Fig. 6). Here $\text{qCO}_2\text{-soc}$ should be an index for microbial availability of carbon substrate normalized by total C contents in soils. Thus, using $\text{qCO}_2\text{-soc}$, we can consider whether the microbially available carbon substrate in interested soil should be much more than in other soils. Microbial biomass N also showed significantly negative correlations with IF_{CO_2} values after rewetting in the first cycle and in the whole of the first cycle ($p < 0.05$) but not in the whole incubation period including three cycles ($p = 0.12$).

Table 5. Pearsons correlation coefficients between IF_{CO_2} and environmental and soil properties

Environmental or soil property	After rewetting in first cycle	Total in first cycle	Total in three cycles
Elevation	0.36	0.37	0.22
MAT	0.54	0.55	0.46
MAP	0.05	0.08	-0.11
MAP-PET	0.01	0.04	-0.14
Water content at sampling	-0.11	-0.07	-0.15
WHC	0.06	0.08	0.04
pH(H ₂ O)	0.54	0.51	0.57
Electronic conductivity	0.19	0.21	-0.01
Total C	-0.03	0.01	-0.11
Total N	-0.07	-0.03	-0.12
C/N ratio	0.27	0.23	0.22
Sand	0.72*	0.70*	0.56
Silt	-0.50	-0.48	-0.35
Clay	-0.69*	-0.68*	-0.57
Alo+0.5Feo	0.81**	0.82**	0.70*
Alo	0.84**	0.85**	0.73*
Feo	0.64*	0.66*	0.53
Alp+0.5Fep	0.81**	0.82**	0.70*
Alp	0.84**	0.85**	0.74*
Fep	0.55	0.58	0.54
Alo-p+0.5Feo-p	0.46	0.43	0.37
Alo-p	0.56	0.57	0.48
Feo-p	0.55	0.55	0.46
Total C/Alp molar ratio	-0.79**	-0.78**	-0.72*
Cp	0.39	0.42	0.23
Cp/Alp molar ratio	-0.79**	-0.75*	-0.74*

There were also considerable relations among Alp contents, total C/Alp molar ratio, and qCO_{2_soc} (Fig. 7). Soil Alp content and total C/Alp molar ratio showed a negative correlation at $p < 0.05$. The qCO_{2_soc} showed a positive correlation with




Figure 5. Relations between IF_{CO_2} and soil pyrophosphate-extractable Al (Alp) content (upper panels), total C to Alp molar ratio (middle panels), and pyrophosphate-extractable C (Cp) to Alp molar ratio (bottom panels). Significant correlation coefficients at $p < 0.01$ and $p < 0.05$ are indicated with single (*) and double asterisks (**), respectively.




Figure 6. Relations between IF_{CO_2} and total C content-specific CO_2 release rate under constant moisture content (qCO_{2_soc}). Significant correlation coefficients at $p < 0.01$ and $p < 0.05$ are indicated with single (*) and double asterisks (**), respectively.

220 total C/Alp molar ratio and negative correlation with Alp content at $p < 0.01$. The Cp/Alp molar ratio also showed similar covariations with these variables, showing a strongly positive correlation with total C/Alp molar ratio ($r = 0.90$, $p < 0.01$; Fig. S3).

Table 6. Pearsons correlation coefficients between IF_{CO_2} and soil microbial properties

Soil microbial property	After rewetting in first cycle	Total in first cycle	Total in three cycles
Total C content-specific CO_2 release rate under no water fluctuation, qCO_{2_soc}	-0.87*	-0.86**	-0.77*
K_2SO_4 extractable C	-0.25	-0.21	-0.33
K_2SO_4 extractable N	-0.63	-0.60	-0.55
K_2SO_4 extractable C/N	0.15	0.18	-0.01
Microbial biomass C	-0.62	-0.61	-0.48
Microbial biomass N	-0.67*	-0.66*	-0.52
Microbial biomass C/N	0.47	0.48	-0.31




Figure 7. Relations among soil Alp contents, total C/Alp molar ratio, and qCO_{2_soc} . Significant correlation coefficients at $p < 0.01$ and $p < 0.05$ are indicated with single (*) and double asterisks (**), respectively.

It should be noted that both microbial biomass C and N were significantly lower in soils incubated under DWC conditions than constant water content conditions ($p < 0.05$; Fig. 8). Microbial biomass C under DWC conditions was lower by 20.2% than those under constant water content conditions. Microbial biomass N under DWC conditions was 12.6% lower than under constant water content conditions. Nevertheless, there were no significant correlations between the differences in microbial biomass and the value of IF_{CO_2} ($p = 0.15$; Fig. 9).

4 Discussion

An increase in CO_2 release due to DWCs was consistently observed across 10 forest and pastureland soils in Japan (Figs. 3 and 4). The comprehensive increases in CO_2 release by DWCs were different from the findings of a recent meta-analysis of studies showing no significant increases in CO_2 release under DWCs compared with constant water content with an equivalent mean water content during the period of interest (Zhang et al., 2020). Furthermore, our observations quantified the increase




Figure 8. Comparisons of microbial biomass C (upper panel) and N (bottom panel) after incubation between DWC and constant water content conditions. Significant differences ($p < 0.05$ by pairwise t test) in microbial biomass between two treatments are presented.

in CO_2 release due to three DWCs as IF_{CO_2} values of 1.3–3.7. The observed effect size of DWCs on the CO_2 release from soils was large given that even a 20% increase in CO_2 release from the worlds soils can exceed the annual CO_2 emission from anthropogenic processes (Friedlingstein et al., 2020).

Furthermore, analyzing the relations between IF_{CO_2} and fundamental soil properties showed a significantly positive correlation between IF_{CO_2} and soil Alp content (Table 5, Fig. 5). Of the organo-metal complexes measured as pyrophosphate-extractable metals, Fep is known to be sensitive to DWCs, especially in seasonally flooded forests and wetlands, likely due to their vulnerability to redox potential changes caused by alterations to the water regime (Lacroix et al., 2019; Chen et al., 2017, 2018; Chen and Thompson, 2018). Less is known about the vulnerability of Alp to DWCs. However, soil Alp content may be affected by DWCs through changes in soil acidity. A previous field survey conducted in Japanese forest and arable soils by Takahashi et al. (2006) showed that liming of non-allophanic Andisols increased soil pH and decreased Alp content. Miyazawa et al. (2013) verified this behavior by a laboratory incubation experiment for Andisols. Although we did not monitor pH during incubation in the present study, increases in pH after DWCs have been widely observed in upland agricultural (Wang et al., 2020; Meng et al., 2020), seasonally submerged paddy, and wetland soils (Majumdar et al., 2023). Therefore, the acidity mitigation by DWCs would destroy the organo-Al complexes, and increase microbially available C through the release of C protected by the organo-Al complexes or other soil elements, such as macro- and microaggregates, which are tightly bonded by




Figure 9. Relations between IF_{CO_2} and microbial biomass C or N decrease by DWCs. There were no statistically significant correlations ($p > 0.05$).

organo-Al complexes and thus physically protect organic C from microbial decomposition (Asano and Wagai, 2014; Takahashi and Dahlgren, 2016; Wagai et al., 2018).

250 Covariations among soil Al_p contents, total C/Al_p molar ratio, and qCO_{2_soc} (Fig. 7) also support the state of soil Al_p content as the primary predictor of variations in IF_{CO_2} . Negative correlations of soil Al_p contents with both total C/Al_p molar ratio and qCO_{2_soc} suggested that more Al_p to total C (i.e., high total C/Al_p values) strengthens binding between Al_p and organic matter, resulting in resistance of organic matter to microbial decomposition under constant water conditions (i.e., low qCO_{2_soc} values). These features among reactive Al and SOM dynamics would result in positive correlations of qCO_{2_soc} and total C/Al_p molar ratio with IF_{CO_2} (Table 6, Figs. 5 and 6). However, such ability of Al_p-rich soils to protect SOM from microbial decomposition likely does not persist under conditions of increased water fluctuations associated with DWCs, as suggested above. Given that amounts of pyrophosphate extractable C (C_p) in investigated soils (>19,000 $\mu\text{g/g}$ dry soil, Table 4), which represented C associated with the organo-metal complex, were substantially greater than observed CO_2 release increase by DWCs ($620\text{--}1999 \mu\text{g C g}^{-1}$ dry soil 84 days^{-1}), limited and specific C substrates in the organo-metal complex likely
 260 associated with CO_2 increase by DWCs.

We also found substantially lower microbial biomass in soils subjected to DWCs than constant water content conditions (Fig. 8), suggesting a decrease in microbial biomass through the destruction of microbial cells by DWCs (Kaiser et al., 2015; Marumoto et al., 1977, 1982; Marumoto, 1984; Nagano et al., 2023; Unger et al., 2010, 2012). The destruction of microbial cells is expected to release soluble organic matter available for microbes that have survived the DWC and to cause a marked
 265 increase in CO_2 release after rewetting (Marumoto et al., 1977, 1982; Marumoto, 1984; Nagano et al., 2019; Unger et al., 2010, 2012). Nevertheless, the contribution of microbially derived substances to the increase in CO_2 release remained unclear

in the present study because of the lack of a significant correlation between IF_{CO_2} and the decrease in microbial biomass (Fig. 9). Whereas there was also no significant correlation between the decrease in microbial biomass and the increase in CO_2 release due to DWCs (Fig. S4), the decrease in microbial biomass C ($246\text{--}1134 \mu\text{g C g}^{-1}$ dry soil) was within the amount of increase
270 in CO_2 release ($620\text{--}1999 \mu\text{g C g}^{-1}$ dry soil 84 days^{-1}). Therefore, the strict mechanisms of these carbon sources to CO_2 release increase, including the persistence and timing of their contribution, still require further works (Schimel, 2018; Barnard et al., 2020), considering the significant contribution of more than two carbon pools to the CO_2 release increase (Slessarev and Schimel, 2020; Warren and Manzoni, 2023), as well as carbon likely released by destructions of organo-metal complexes and microbial biomass by DWC as suggested in the present study.

275 It should be noted that shortcomings from unmeasured CO_2 release during the drying periods for DWCs treatments (i.e., Day 1 to Day 7 and Day 18 to Day 24 in each cycle) should be minor, while the linear changes in CO_2 release rate during the drying also assumed in other studies (e.g., Nagano et al. (2019); Zhang et al. (2020); Jin et al. (2023)). This is because we observed significant relationships between IF_{CO_2} and organo-Al complex also in IF_{CO_2} after the rewetting in addition to the IF_{CO_2} for total 1st cycle and three cycles (Table 5).

280 5 Conclusions

Through the present study, a comprehensive increase in CO_2 release by DWCs (i.e., 1.3–3.7-fold greater than CO_2 release under constant water conditions) was observed in Japanese forest and pastureland soils. These magnitudes of increase in CO_2 release were strongly correlated with soil Al_p content, total C/ Al_p molar ratio, and total C content-specific CO_2 release rate under
285 constant water conditions (i.e., qCO_{2_soc}), suggesting the possible vulnerability of SOM protection by organo-Al complexes against DWCs. A decrease in microbial biomass by DWCs was also suggested, whereas their relations with the increase in CO_2 release remains to be determined in future studies.

Data availability. The data that support the findings of this study are available from the corresponding author upon reasonable request.

Author contributions. YS and HN established the basic research design, conducted all data analyses, including software preparation, validation, and visualization, and wrote the original manuscript. SH, MA-A, JK, and HN conducted soil sampling. YS, SH, MA-A, JK, and HN
290 contributed to the detailed research design, soil analysis, data validation, interpretation of the results, and editing of the manuscript. TY and YK provided essential support in analysis of soil properties and data. All authors contributed to editing the article and approved the submitted version.

Competing interests. The authors declare that they have no known competing financial interests or personal relations that could have appeared to influence the work reported in this paper.

295 *Acknowledgements.* This work was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI (grant numbers 21H02231,
21H05313, 22H05717). The authors thank Prof. Kouki Hikosaka and Dr. Hirofumi Kajino of Tohoku University for providing the vegetation
information on the Oita forest site. Ms. Ayako Tamaki and Mr. Masahiro Otaki at Niigata University supported sample preparation and anal-
ysis during the experiment and text editing in preparing the manuscript. Dr. Rei Shibata at Niigata University and Dr. Masami Tsukahara,
Mr. Kosuke Ito, and Mr. Tatsuki Tanaka at Niigata Prefectural Forest Research Institute helped in site selection and soil sampling in Niigata
300 Prefecture, Japan. Prof. Naoki Harada, Dr. Kazuki Suzuki, and Dr. Asiloglu Rasit of Niigata University and Dr. Hiroyuki Sase and Dr. Rieko
Urakawa of Asia Center for Air Pollution Research (ACAP) provided valuable preliminary discussion in writing the manuscript. The authors
also thank Ms. Misuzu Kaminaga, Ms. Kikuko Yoshigaki, Ms. Makiko Ishihara, and Ms. Kazumi Matsumura at JAEA for support with
laboratory work.

References

- 305 Abatzoglou, J. T., Dobrowski, S. Z., Parks, S. A., and Hegewisch, K. C.: TerraClimate, a high-resolution global dataset of monthly climate and climatic water balance from 19582015, *Scientific Data*, 5, 170 191, <https://doi.org/10.1038/sdata.2017.191>, 2018.
- Ahn, H. K., Richard, T. L., and Glanville, T. D.: Optimum moisture levels for biodegradation of mortality composting envelope materials, *Waste Management*, 28, 1411–1416, <https://doi.org/10.1016/J.WASMAN.2007.05.022>, 2008.
- Allan, R. P., Barlow, M., Byrne, M. P., Cherchi, A., Douville, H., Fowler, H. J., Gan, T. Y., Pendergrass, A. G., Rosenfeld, D., Swann, A. L.,
310 Wilcox, L. J., and Zolina, O.: Advances in understanding large-scale responses of the water cycle to climate change, *Annals of the New York Academy of Sciences*, 1472, 49–75, <https://doi.org/10.1111/NYAS.14337>, 2020.
- Asano, M. and Wagai, R.: Evidence of aggregate hierarchy at micro- to submicron scales in an allophanic Andisol, *Geoderma*, 216, 62–74, <https://doi.org/10.1016/j.geoderma.2013.10.005>, 2014.
- Barnard, R. L., Blazewicz, S. J., and Firestone, M. K.: Rewetting of soil: Revisiting the origin of soil CO₂ emissions, *Soil Biology and
315 Biochemistry*, 147, 107 819, <https://doi.org/10.1016/j.soilbio.2020.107819>, 2020.
- Birch, H. F.: The effect of soil drying on humus decomposition and nitrogen availability, *Plant and Soil*, 10, 9–31, <https://doi.org/10.1007/BF01343734>, 1958.
- Bond-Lamberty, B. and Thomson, A.: Temperature-associated increases in the global soil respiration record, *Nature*, 464, 579–582, <https://doi.org/10.1038/nature08930>, 2010.
- 320 Borken, W. and Matzner, E.: Reappraisal of drying and wetting effects on C and N mineralization and fluxes in soils, *Global Change Biology*, 15, 808–824, <https://doi.org/10.1111/j.1365-2486.2008.01681.x>, 2009.
- Chen, C. and Thompson, A.: Ferrous Iron Oxidation under Varying pO₂ Levels: The Effect of Fe(III)/Al(III) Oxide Minerals and Organic Matter, *Environmental Science and Technology*, 52, 597–606, <https://doi.org/10.1021/acs.est.7b05102>, 2018.
- Chen, C., Kukkadapu, R. K., Lazareva, O., and Sparks, D. L.: Solid-Phase Fe Speciation along the Vertical Redox Gradients in Flood-
325 plains using XAS and Mössbauer Spectroscopies, 51, 7903–7912, <https://pubs.acs.org/doi/10.1021/acs.est.7b00700><https://pubs.acs.org/sharingguidelines>, 2017.
- Chen, C., Meile, C., Wilmoth, J., Barcellos, D., and Thompson, A.: Influence of pO₂ on Iron Redox Cycling and Anaerobic Organic Carbon Mineralization in a Humid Tropical Forest Soil, *Environmental Science and Technology*, 52, 7709–7719, <https://doi.org/10.1021/acs.est.8b01368>, 2018.
- 330 Courchesne, F. and Turmel, M.-C.: Extractable Al, Fe, Mn, and Si, in: *Soil Sampling and Methods of Analysis*, edited by Carter, M. and Gregorich, E., chap. 26, pp. 307–316, CRC Press, second edi edn., ISBN 978-0-8493-3586-0, 2008.
- Dai, A.: Increasing drought under global warming in observations and models, *Nature Climate Change*, 3, 52–58, <https://doi.org/10.1038/nclimate1633>, 2012.
- Donat, M. G., Lowry, A. L., Alexander, L. V., O’Gorman, P. A., and Maher, N.: More extreme precipitation in the world’s dry and wet
335 regions, *Nature Climate Change*, 6, 508–513, <https://doi.org/10.1038/nclimate2941>, 2016.
- FAO/IIASA/ISRIC/ISS-CAS/JRC: Harmonized World Soil Database (version 1.1), FAO, Italy and IIASA, Laxenburg, Austria, 2009.
- Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le
Quéré, C., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragão, L. E. O. C., Arneth, A., Arora, V., Bates, N. R., Becker, M.,
Benoit-Cattin, A., Bittig, H. C., Bopp, L., Bultan, S., Chandra, N., Chevallier, F., Chini, L. P., Evans, W., Florentie, L., Forster, P. M.,
340 Gasser, T., Gehlen, M., Gilfillan, D., Gkritzalis, T., Gregor, L., Gruber, N., Harris, I., Hartung, K., Haverd, V., Houghton, R. A., Ilyina,

- T., Jain, A. K., Joetzjer, E., Kadono, K., Kato, E., Kitidis, V., Korsbakken, J. I., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Liu, Z., Lombardozzi, D., Marland, G., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Niwa, Y., O'Brien, K., Ono, T., Palmer, P. I., Pierrot, D., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Smith, A. J. P., Sutton, A. J., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., van der Werf, G., Vuichard, N., Walker, A. P., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, X., and Zaehle, S.: Global Carbon Budget 2020, *Earth System Science Data*, 12, 3269–3340, <https://doi.org/10.5194/essd-12-3269-2020>, 2020.
- Georgiou, K., Jackson, R. B., Vindušková, O., Abramoff, R. Z., Ahlström, A., Feng, W., Harden, J. W., Pellegrini, A. F., Polley, H. W., Soong, J. L., Riley, W. J., and Torn, M. S.: Global stocks and capacity of mineral-associated soil organic carbon, *Nature Communications*, 13, 1–12, <https://doi.org/10.1038/s41467-022-31540-9>, 2022.
- 345 Hall, S. J. and Thompson, A.: What do relationships between extractable metals and soil organic carbon concentrations mean?, *Soil Science Society of America Journal*, 86, 195–208, <https://doi.org/10.1002/saj2.20343>, 2022.
- Imaya, A., Inagaki, Y., Tanaka, N., and Ohta, S.: Free oxides and short-range ordered mineral properties of brown forest soils developed from different parent materials in the submontane zone of the Kanto and Chubu districts, Japan, *Soil Science and Plant Nutrition*, 53, 621–633, <https://doi.org/10.1111/j.1747-0765.2007.00175.x>, 2007.
- 355 IPCC: Climate Change 2021: The Physical Science Basis. Working Group I Contribution to the IPCC Sixth Assessment Report, Cambridge University Press, United Kingdom and New York, <https://doi.org/10.1017/9781009157896>, 2021.
- Jin, X., Wu, F., Wu, Q., Heděnc, P., Peng, Y., Wang, Z., and Yue, K.: Effects of drying-rewetting cycles on the fluxes of soil greenhouse gases, *Heliyon*, 9, e12984, <https://doi.org/10.1016/j.heliyon.2023.e12984>, 2023.
- Kaiser, M., Kleber, M., and Berhe, A. A.: How air-drying and rewetting modify soil organic matter characteristics: An assessment to improve data interpretation and inference, *Soil Biology and Biochemistry*, 80, 324–340, <https://doi.org/10.1016/j.soilbio.2014.10.018>, 2015.
- 360 Kim, D.-G., Vargas, R., Bond-Lamberty, B., and Turetsky, M. R.: Effects of soil rewetting and thawing on soil gas fluxes: a review of current literature and suggestions for future research, *Biogeosciences*, 9, 2459–2483, <https://doi.org/10.5194/bg-9-2459-2012>, 2012.
- Koarashi, J., Hockaday, W. C., Masiello, C. A., and Trumbore, S. E.: Dynamics of decadal cycling carbon in subsurface soils, *Journal of Geophysical Research: Biogeosciences*, 117, 1–13, <https://doi.org/10.1029/2012JG002034>, 2012.
- 365 Kpemoua, T. P., Barré, P., Houot, S., and Claire, C.: Accurate evaluation of the Birch effect requires continuous CO₂ measurements and relevant controls, *Soil Biology and Biochemistry*, 180, 109007, <https://doi.org/10.1016/J.SOILBIO.2023.109007>, 2023.
- Kramer, M. G. and Chadwick, O. A.: Climate-driven thresholds in reactive mineral retention of soil carbon at the global scale, *Nature Climate Change*, 8, 1104–1108, <https://doi.org/10.1038/s41558-018-0341-4>, 2018.
- Lacroix, R. E., Tfaily, M. M., McCreight, M., Jones, M. E., Spokas, L., and Keiluweit, M.: Shifting mineral and redox controls on carbon cycling in seasonally flooded mineral soils, *Biogeosciences*, 16, 2573–2589, <https://doi.org/10.5194/BG-16-2573-2019>, 2019.
- 370 Lee, M.-s., Nakane, K., Nakatsubo, T., Mo, W.-h., and Koizumi, H.: Effects of rainfall events on soil CO₂ flux in a cool temperate deciduous broad-leaved forest, *Ecological Research*, 17, 401–409, <https://doi.org/10.1046/j.1440-1703.2002.00498.x>, 2002.
- Lee, X., Wu, H. J., Sigler, J., Oishi, C., and Siccama, T.: Rapid and transient response of soil respiration to rain, *Global Change Biology*, 10, 1017–1026, <https://doi.org/10.1111/j.1365-2486.2004.00787.x>, 2004.
- 375 Leuthold, S. J., Haddix, M. L., Lavalley, J., and Cotrufo, M. F.: Physical fractionation techniques, *Encyclopedia of Soils in the Environment*, Second Edition, pp. 68–80, <https://doi.org/10.1016/B978-0-12-822974-3.00067-7>, 2023.
- Mabuhay, J. A., Nakagoshi, N., and Horikoshi, T.: Microbial biomass and abundance after forest fire in pine forests in Japan, *Ecological Research*, 18, 431–441, <https://doi.org/10.1046/J.1440-1703.2003.00567.X/METRICS>, 2003.

- Majumdar, A., Dubey, P. K., Giri, B., Moulick, D., Srivastava, A. K., Roychowdhury, T., Bose, S., and Jaiswal, M. K.: Combined effects of dry-wet irrigation, redox changes and microbial diversity on soil nutrient bioavailability in the rice field, *Soil and Tillage Research*, 232, 105 752, <https://doi.org/10.1016/J.STILL.2023.105752>, 2023.
- Marumoto, T.: Mineralization of C and N from microbial biomass in paddy soil, *Plant and Soil*, 76, 165–173, <https://doi.org/10.1007/BF02205577>, 1984.
- Marumoto, T., Kai, H., Yoshida, T., and Harada, T.: Drying effect on mineralizations of microbial cells and their cell walls in soil and contribution of microbial cell walls as a source of decomposable soil organic matter due to drying, *Soil Science and Plant Nutrition*, 23, 9–19, <https://doi.org/10.1080/00380768.1977.10433017>, 1977.
- Marumoto, T., Anderson, J. P., and Domsch, K. H.: Decomposition of ¹⁴C- and ¹⁵N-labelled microbial cells in soil, *Soil Biology and Biochemistry*, 14, 461–467, [https://doi.org/10.1016/0038-0717\(82\)90105-5](https://doi.org/10.1016/0038-0717(82)90105-5), 1982.
- Meng, Z., Huang, S., Xu, T., Deng, Y., Lin, Z., and Wang, X.: Transport and transformation of Cd between biochar and soil under combined dry-wet and freeze-thaw aging, *Environmental Pollution*, 263, 114 449, <https://doi.org/10.1016/J.ENVPOL.2020.114449>, 2020.
- MEXT and JAM: Climate Change in Japan: Report on Assessment of Observed/Projected Climate Changes Relating to the Atmosphere, Land and Oceans., Tech. rep., Ministry of Education, Culture, Sports, Science and Technology; Japan Meteorological Agency, <https://www.data.jma.go.jp/cpdinfo/ccj/index.html>, 2020.
- Miller, A. E., Schimel, J. P., Meixner, T., Sickman, J. O., and Melack, J. M.: Episodic rewetting enhances carbon and nitrogen release from chaparral soils, *Soil Biology and Biochemistry*, 37, 2195–2204, <https://doi.org/10.1016/j.soilbio.2005.03.021>, 2005.
- Miller, W. P., Radcliffe, D. E., and Miller, D. M.: An historical perspective on the theory and practice of soil mechanical analysis, *Journal of Agronomic Education*, 17, 24–28, <https://eric.ed.gov/?id=EJ374021>, 1988.
- Miyazawa, M., Takahashi, T., Sato, T., Kanno, H., and Nanzyo, M.: Factors controlling accumulation and decomposition of organic carbon in humus horizons of Andosols, *Biology and Fertility of Soils*, 49, 929–938, <https://doi.org/10.1007/s00374-013-0792-8>, 2013.
- Morisada, K., Ono, K., and Kanomata, H.: Organic carbon stock in forest soils in Japan, *Geoderma*, 119, 21–32, [https://doi.org/10.1016/S0016-7061\(03\)00220-9](https://doi.org/10.1016/S0016-7061(03)00220-9), 2004.
- Nagano, H., Atarashi-Andoh, M., and Koarashi, J.: Effect of dry-wet cycles on carbon dioxide release from two different volcanic ash soils in a Japanese temperate forest, *Soil Science and Plant Nutrition*, 65, 525–533, <https://doi.org/10.1080/00380768.2019.1649976>, 2019.
- Nagano, H., Atarashi-Andoh, M., Tanaka, S., Yomogida, T., Kozai, N., and Koarashi, J.: Stable C and N isotope abundances in water-extractable organic matter from air-dried soils as potential indices of microbially utilized organic matter, *Frontiers in Forests and Global Change*, 6, 1228 053, <https://doi.org/10.3389/ffgc.2023.1228053>, 2023.
- Nakai, M.: Soil particle distribution [The pipette method], in: *Dojo-Kankyo-Bunseki-Ho (Analytical Methods for Soil Environment)*, edited by Editorial Committee for Analytical Methods for Soil Environment, pp. 24–29, Hakuyusha, Tokyo, ISBN 4-8268-0168-8, 1997.
- NARO: Japan Soil Inventory, <https://soil-inventory.rad.naro.go.jp/index.html>, 2023.
- Pfleiderer, P., Schleussner, C.-F., Kornhuber, K., and Coumou, D.: Summer weather becomes more persistent in a 2°C world, *Nature Climate Change*, pp. 1–6, <https://doi.org/10.1038/s41558-019-0555-0>, 2019.
- Rasmussen, C., Heckman, K., Wieder, W. R., Keiluweit, M., Lawrence, C. R., Berhe, A. A., Blankinship, J. C., Crow, S. E., Druhan, J. L., Hicks Pries, C. E., Marin-Spiotta, E., Plante, A. F., Schädel, C., Schimel, J. P., Sierra, C. A., Thompson, A., and Wagai, R.: Beyond clay: towards an improved set of variables for predicting soil organic matter content, *Biogeochemistry*, <https://doi.org/10.1007/s10533-018-0424-3>, 2018.

- Running, S. and Zhao., M.: MODIS/Terra Net Primary Production Gap-Filled Yearly L4 Global 500m SIN Grid V061., <https://doi.org/10.5067/MODIS/MOD17A3HGF.061>, 2021.
- Schimel, J. P.: Life in Dry Soils: Effects of Drought on Soil Microbial Communities and Processes, *Annual Review of Ecology, Evolution, and Systematics*, 49, 409–432, <https://doi.org/10.1146/annurev-ecolsys-110617-062614>, 2018.
- 420 Shirato, Y., Hakamata, T., and Taniyama, I.: Modified rothamsted carbon model for andosols and its validation: changing humus decomposition rate constant with pyrophosphate-extractable Al, *Soil Science and Plant Nutrition*, 50, 149–158, <https://doi.org/10.1080/00380768.2004.10408463>, 2004.
- Slessarev, E. W. and Schimel, J. P.: Partitioning sources of CO₂ emission after soil wetting using high-resolution observations and minimal models, *Soil Biology and Biochemistry*, 143, 107–115, <https://doi.org/10.1016/j.soilbio.2020.107753>, 2020.
- 425 Takahashi, T. and Dahlgren, R. A.: Nature, properties and function of aluminum-humus complexes in volcanic soils, *Geoderma*, 263, 110–121, <https://doi.org/10.1016/j.geoderma.2015.08.032>, 2016.
- Takahashi, T., Ikeda, Y., Fujita, K., and Nanzyo, M.: Effect of liming on organically complexed aluminum of nonallophanic Andosols from northeastern Japan, *Geoderma*, 130, 26–34, <https://linkinghub.elsevier.com/retrieve/pii/S0016706105000108>, 2006.
- Unger, S., Máguas, C., Pereira, J. S., David, T. S., and Werner, C.: The influence of precipitation pulses on soil respiration - Assessing the
430 " Birch effect" by stable carbon isotopes, *Soil Biology and Biochemistry*, 42, 1800–1810, <https://doi.org/10.1016/j.soilbio.2010.06.019>, 2010.
- Unger, S., Máguas, C., Pereira, J. S., David, T. S., and Werner, C.: Interpreting post-drought rewetting effects on soil and ecosystem carbon dynamics in a Mediterranean oak savannah, *Agricultural and Forest Meteorology*, 154–155, 9–18, <https://doi.org/10.1016/j.agrformet.2011.10.007>, 2012.
- 435 USDA Soil Survey Staff: Keys to Soil Taxonomy, USDA-Natural Resources Conservation Service, 13th editi edn., <https://www.nrcs.usda.gov/resources/guides-and-instructions/keys-to-soil-taxonomy#keys>, 2022.
- Vance, E. D., Brookes, P. C., and Jenkinson, D. S.: An extraction method for measuring soil microbial biomass C, *Soil Biology and Biochemistry*, 19, 703–707, [https://doi.org/10.1016/0038-0717\(87\)90052-6](https://doi.org/10.1016/0038-0717(87)90052-6), 1987.
- Wagai, R., Kajiura, M., Uchida, M., and Asano, M.: Distinctive Roles of Two Aggregate Binding Agents in Allophanic Andisols: Young
440 Carbon and Poorly-Crystalline Metal Phases with Old Carbon, *Soil Systems*, 2, 29, <https://doi.org/10.3390/soilsystems2020029>, 2018.
- Wang, D., Felice, M. L., and Scow, K. M.: Impacts and interactions of biochar and biosolids on agricultural soil microbial communities during dry and wet-dry cycles, *Applied Soil Ecology*, 152, 103–110, <https://doi.org/10.1016/J.APSOIL.2020.103570>, 2020.
- Warren, C. R. and Manzoni, S.: When dry soil is re-wet, trehalose is respired instead of supporting microbial growth, *Soil Biology and Biochemistry*, 184, 109–115, <https://doi.org/10.1016/J.SOILBIO.2023.109121>, 2023.
- 445 Wijesinghe, J. N., Asaoka, H., Mori, Y., and Hiradate, S.: Depth profile of plant nutrients and acidity in a non-allophanic Andosol as affected by a half-century of fertilization, *Soil Science and Plant Nutrition*, pp. 1–12, <https://doi.org/10.1080/00380768.2021.1887711>, 2021.
- Xiang, S. R., Doyle, A., Holden, P. A., and Schimel, J. P.: Drying and rewetting effects on C and N mineralization and microbial activity in surface and subsurface California grassland soils, *Soil Biology and Biochemistry*, 40, 2281–2289, <https://doi.org/10.1016/j.soilbio.2008.05.004>, 2008.
- 450 Zhang, S., Yu, Z., Lin, J., and Zhu, B.: Responses of soil carbon decomposition to drying-rewetting cycles: A meta-analysis, *Geoderma*, 361, 114–121, <https://doi.org/10.1016/j.geoderma.2019.114069>, 2020.
- Zhang, S., Lin, J., Wang, P., and Zhu, B.: The direct and legacy effects of dryingrewetting cycles on active and relatively resistant soil carbon decomposition, *Land Degradation and Development*, 34, 2124–2135, <https://doi.org/10.1002/ldr.4594>, 2023.