

Regional and sectoral contributions of NO_x and reactive carbon emission sources to global trends in tropospheric ozone during the 2000-2018 period

Aditya Nalam ^{1,2}, Aura Lupaşcu ^{2,*}, Tabish Ansari ², Tim Butler ^{1,2}

¹ Institut für Meteorologie, Freie Universität Berlin, Berlin, Germany

² Research Institute for Sustainability, Helmholtz Centre Potsdam, Germany

* Now at European Center for Medium Range Weather Forecast, Bonn, Germany

Correspondence to: Aditya Nalam (adinalam@zedat.fu-berlin.de)

Abstract. Over the past few decades, the tropospheric ozone precursor anthropogenic emissions: nitrogen oxides (NO_x) and reactive carbon (RC) from northern mid/high-latitude regions (North America (NAM), Europe (EUR) etc.), have been decreasing, and those from Asia and (sub)-tropical regions (South Asia (SAS), Middle-East (MDE), etc.) have been increasing, leading to an equatorward emission redistribution. In this study, we quantify the contributions of various sources of NO_x and RC emissions to tropospheric ozone using a source attribution technique during the 2000-2018 period in a global chemistry transport model: CAM4-Chem. We tag the ozone molecules with the source of their NO_x or RC precursor emission in two separate simulations, one for each of: NO_x-tagged and RC-tagged. These tags include various natural (biogenic, biomass burning, lightning NO_x and RC from methane oxidation), and regional anthropogenic (North American, European, East Asian, South Asian etc.) precursor emission sources, and influx from stratosphere. We simulate ~336 Tg O₃ of tropospheric ozone burden (TOB) with an increasing trend of 0.91 Tg O₃/yr (0.28 %/yr), largely contributed (and trend driven) by anthropogenic NO_x emissions and methane oxidation. The ozone production efficiency of regional anthropogenic NO_x emissions increases significantly when emissions decrease (Europe, North American and Russia-Belarus-Ukraine region's emissions EUR, NAM etc.) and decreases significantly when emissions increase (South Asian, Middle Eastern SAS, MDE, International Shipping etc.). Tropical regions, despite smaller emissions, contribute more to tropospheric ozone burden TOB compared to emissions from higher latitudes, consistent with previous work, predominantly due to large convection (combined with intense sunlight and larger reaction rates) at the tropics thereby lifting O₃ and its precursor NO_x molecules into the free troposphere where ozone's lifetime is longer. We contrast the contribution to tropospheric ozone burden with that of the contribution to the global surface ozone. We simulate a smaller relative contribution from tropical regions to the global mean surface ozone (GMS-O3) compared to their contribution to the tropospheric ozone burden TOB. The global population-weighted mean ozone (GPW-O3 related to ozone exposure) is much larger compared to GMS-O3 surface mean, mainly due to large anthropogenic emissions from densely populated regions: East Asia, South Asia, and other tropical regions, and a substantial contribution from international ship NO_x emissions. The increasing trends in anthropogenic precursor emissions from these regions are the main drivers of increasing GPW-O3 global population-weighted mean ozone.

Short Summary (500 Characters).

Tropospheric O₃ molecules are labelled with the identity of their precursor source ~~in CAM-Chem to quantify-simulate the contributions~~ from various emission sources to the global tropospheric O₃ burden (TOB) and its trends. With an equatorward shift, anthropogenic NO_x emissions become significantly more efficient at producing O₃ and play a major role in driving TOB trends. ~~This is, mainly~~ due to larger convection at the tropics effectively lifting O₃ and its precursors to the free troposphere where O₃ lifetime is longer.

1 Introduction

Tropospheric ozone is a major greenhouse gas (Myhre et al. 2013), after carbon dioxide and methane. It also has an indirect influence on the lifetime of methane through its impact on the hydroxyl radical, OH (Monks et al., 2015). At the surface, ozone is a major air pollutant causing a large number of premature deaths (~423100 Ozone attributable deaths [95 % confidence interval: 223200, 659400] in 2019; Malashock et al. 2022) and heavy yield losses in the production of staple food crops worldwide (losses of up to 79–121 Tg annually; Avnery et al. 2011, Mills et al. 2018). Tropospheric ozone initially thought to be mostly of stratospheric origin (Junge, 1962), is also a secondary pollutant produced from the reaction between ozone precursor gases: nitrogen oxides (NO_x: NO and NO₂) and reactive carbon (RC: CH₄, CO and Volatile Organic Compounds (VOCs)); Levy, 1972; Chameides and Walker, 1973; Crutzen, 1974). A multi-model assessment study showed that the tropospheric ozone burden has increased by ~44 % since the pre-industrial period, mainly driven by increasing anthropogenic emissions of ozone precursor gases (Griffiths et al., 2021). Gaudel et al., (2020) utilize the data from In-service Aircraft for a Global Observing System database (IAGOS) and identify remarkable increase in tropospheric ozone since 1994 over several regions in the Northern Hemisphere. Addressing the effects of tropospheric ozone both as a pollutant and a greenhouse gas requires mitigation of the anthropogenic emissions of ozone precursors.

Since 1980s, anthropogenic emissions of tropospheric ozone precursors from North America and Western Europe have decreased in response to the implementation of control measures while the emissions from Asia, Central America, and Eastern Europe have increased due to economic and population growth (Granier et al., 2011, Cooper et al., 2014) leading to an equatorward shift in global emission pattern. A recent study (Li et al., 2024) confirms an equatorward shift in Asian NO_x emissions between 2010 and 2017 according to bottom-up estimates. Zhang et al., (2016) found that this equatorward shift in precursor emissions is the dominant factor, compared to the change in magnitude of emissions and methane concentration, that has led to an increase in the tropospheric ozone burden between 1980 and 2010. This is mainly due to larger convection of polluted air masses from the boundary layer into the free troposphere, in addition to larger reaction rates and NO_x sensitivity over the tropical regions (Wild et al., 2001) compared to extra-tropical regions. Further, Wang et al., (2022) show that increasing aircraft emissions play a major role in the increasing trend in tropospheric ozone burden over the 1995-2017 period. Similarly, the increase in methane concentration during the industrial period also had a substantial impact on tropospheric ozone burden (Stevenson et al., 2013).

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Source apportionment methods are used in chemical transport modelling studies to quantify the influence of specific emission sources on ozone concentration at any given location. Typically, global modelling studies have used two different methods for source apportionment: perturbation and tagging, which will be described in more detail below.

Several previous studies have used the perturbation method of source attribution to study the impact of emission changes from a particular region/sector on tropospheric ozone (e.g. West et al., 2007, Fiore et al., 2009, Jonson et al., 2018 etc). Zhang et al. (2021) use this method to study the influence of changing emissions from various world regions between 1980 and 2010 on tropospheric ozone burden. In their study, multiple model simulations were performed: one base simulation with all emissions at 2010 levels and other perturbed simulations with 1980 emissions from the region/sector of interest. [Zhang et al., \(2021\) The study](#) found that the change in tropospheric ozone burden responded the most to perturbation in prescribed methane concentration. Of the geographical source regions, tropospheric ozone [burden](#) responded the most to perturbations in emissions of anthropogenic ozone precursor emissions at tropical regions: South Asian and Southeast Asian regions.

The second method of source apportionment is called tagging, which involves labelling the modelled ozone molecules with the name of its precursor emission sources. As opposed to the previously discussed perturbation method that delivers the sensitivity of simulated ozone to emission changes, this tagging method delivers the exact contribution to the simulated ozone concentration from each of the tagged emission sources (see Table 1 in Mertens et al., 2020 for a detailed comparison between the two methods). Several previous studies have implemented the tagging method in box (e.g., Butler et al., 2011, Coates and Butler, 2015), regional (Kwok et al., 2015, Lupascu and Butler 2019, Zhao et al., 2022), and global models (Grewe et al., (2010, 2017), Emmons et al. 2012, Butler et al., 2018) to study the contribution of various emission sources on ozone concentrations (see Butler et al., (2018) for a review of various tagging approaches). The formation of one tropospheric ozone molecule requires two precursors (NO_x and RC), a major challenge in its emission source attribution is to know the label of which of the precursors should the ozone molecule inherit.

Among the previous global modelling approaches that implement tagging, Grewe et al., (2010, 2017) calculate the production rate of the tagged ozone molecules, by giving equal weight to the proportion of tagged NO_x and RC molecules emitted from a given sector (“combinatorial” tagging). While this approach can deliver information on the role of each emission source in its contribution to the tropospheric ozone concentration, it does not communicate the contrasting roles of NO_x and RC from those sources. For example, Mertens et al., (2018) using the combinatorial approach, attribute ~8 % of tropospheric ozone burden to ozone precursor emissions from land transportation but cannot determine which emitted precursor from land transportation: NO_x or RC contributes predominantly. Butler et al., 2018 formulated the tropospheric ozone source attribution system with tagging (TOAST) within the CAM4-Chem ([Community Atmosphere Model version 4 with chemistry](#)) model that requires running two separate simulations: NO_x -tagged and RC-tagged, to separately attribute ozone to NO_x and [reactive-carbonRC](#) sources respectively, by labelling them with the name of their originating source region/sector. With this approach, the

105 previously mentioned problems related to the contrasting roles of NO_x and RC, are avoided as the contribution from each precursor source is separately attributed. To illustrate: O₃ attributed to biogenic emissions in NO_x-tagged simulation would clearly mean that the contribution is from emitted biogenic NO_x only, irrespective of where the RC comes from to produce the O₃ molecule. Contribution from biogenic RC emissions can be simulated in the RC-tagged simulation. In this way, the roles of NO_x and RC emissions from a given sector are exclusively simulated in separate simulations.

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

110 Butler et al., (2020) use the TOAST tagging technique to quantify the contribution of ozone precursor emissions for the year 2010 from several regions/sectors. They demonstrated that the ozone production efficiency can directly be calculated using this method, as the ratio of tropospheric ozone attributed to a tagged emission source to the amount of precursor emission from that source. NO_x emissions from tropical regions such as South Asia, Southeast Asia and Central America were found to be the most efficient at producing tropospheric ozone compared to the emissions from other regions, consistent with the earlier
115 work of Zhang et al., (2016). They further showed ~~that~~-using a methane perturbation ~~simulation~~-introduced to NO_x-tagged simulation, that the contribution to tropospheric ozone burden by NO_x emitted from international shipping increases especially strongly in response to changes in methane concentration.

Formatted: Subscript

While Butler et al., (2020) discuss the ozone precursor contributions to tropospheric ozone burden for one year: 2010, we use
120 the TOAST tagging approach (further explained in section 2) in this study over the 2000-2018 period to answer the following questions: (i) In Sections 3.1 and 3.2: What is the contribution of precursor emissions from various regions/sectors to the global tropospheric O₃ burden-? How do changes in these emissions during 2000-2018 affect their contribution to the trend in tropospheric ozone burden? and its trend? (ii) Section 3.3: How does the ozone production efficiency (OPE) of ozone precursor emissions respond to the changes in these emissions during the 2000-2018 period? And (iii) in section 3.4: How do
125 contributions of different ozone precursors to the tropospheric O₃ burden contrast with their contribution to global mean surface O₃ and population weighted O₃? We conclude our article with the summary, limitations, and future scope in section 4.

2 Methods

2.1 Simulation setup

We use the ozone source attribution methods described in Butler et al., (2018) and Butler et al., (2020). We perform simulations
130 with CAM4-chem-~~(Community Atmosphere Model version 4 with chemistry)~~, which is a component of the CESM (Community Earth System Model) version 1.2.2 (Lamarque et al., 2012). The model is run at a horizontal resolution of 1.9° ×2.5°, with 56 vertical levels for the 2000-2018 period with specified dynamics from MERRA2 reanalysis (Molod et al., 2015). The temperature, horizontal winds, and ~~surface-sensible and latent heat~~ fluxes from MERRA2 reanalysis dataset are nudged every time step (30 min) by 10 % towards analysis fields (i.e., a 5 h Newtonian relaxation timescale for nudging).

135

140

145

150

155

160

165

Anthropogenic emissions of NO_x, ~~and non-methane reactive carbon (NMRC)~~, CO, and ~~non-methane~~-volatile organic compounds (~~NM~~VOCs) collectively including land-based emissions, international shipping emissions, and aircraft emissions are taken from Hemispheric Transport of Air Pollution version 3 (HTAPv3; Crippa et al., 2023) emissions inventory. We specify aircraft emissions at ~~three-different~~various altitudes effectively representing ~~three~~ different flight phases (landing/take-off, ascent/descent, and cruising). Biomass burning emissions are taken from GFED-v4 inventory (van der Werf et al., 2010). The biogenic ~~NM~~RVOC emissions are taken from CAMS-GLOB-BIO-v3.0 (Sindelarova et al., 2021), and biogenic NO_x (from soil) is prescribed as in Tilmes et al., (2015). While we interpolate the emissions from HTAPv3's high-resolution dataset to our coarser model resolution, we also correct the interpolated land-based emissions over ocean grid cells by moving them to the nearest land grid cell (vice-versa for ocean-based emissions) to make sure that the emissions are allocated to the correct region for the source attribution.

We impose methane concentration as a surface boundary condition. The methane concentration is taken from the ~~2010-2000~~-2018 average mole fraction fields from the CAMS CH₄ flux inversion product v18r1 (<https://ads.atmosphere.copernicus.eu/cdsapp#!/dataset/cams-global-greenhouse-gas-inversion?tab=overview>) and is specified as a zonally and monthly varying transient boundary condition. As in Butler et al., (2018) and Butler et al., (2020), we use the MOZART-4 chemical mechanism (Emmons et al., 2012) further modified to include tagged ozone tracers. This tagging system allows attribution of tropospheric ozone to reactive nitrogen (NO_x) and reactive carbon (CH₄ ~~and~~, ~~CO~~ ~~and~~ ~~NM~~RVOC~~NMRC~~) precursors in two separate simulations. Additionally, stratospheric influx as well as other minor production pathways of tropospheric ozone are also tagged in our model. For a complete attribution, we perform two separate simulations: (i) NO_x-tagged; and (ii) RC-tagged (reactive carbon tagged) with their respective tagged emission sources.

We specify separate tag identities for biogenic, biomass burning, aircraft and surface-based anthropogenic emission sources of ozone precursors, and for ozone from production in the stratosphere. We tag anthropogenic emissions according to the HTAP Tier 1 regions (Galmarini et al., 2017) from which the respective NO_x or ~~NM~~VOCs~~NMRCs~~ are emitted. We focus our study on the northern hemispheric anthropogenic emissions by individually tagging the major northern hemispheric regions (East Asia, South Asia, North America, Europe, Middle East, and Russia-Belarus-Ukraine), and combining the ~~southern hemispheric~~other regions under the "Rest of the World" tag. Within our "Rest of the World" tag, some regions (e.g. Central Asia, Southeast Asia etc.) have been explicitly tagged in our NO_x-tagged simulation but not in our RC-tagged simulation. We also specify additional tags for emissions from international shipping and aircraft emissions (see Table 1 and Fig. S1 for a summary of tags used in this study).

We specify an additional tag for NO_x emission from lightning in our NO_x-tagged simulation, and for RC from methane oxidation in our RC-tagged simulation. In both NO_x and RC tagged simulations, the sum of tagged ozone tracers is equal to

the total ozone simulated by the model. The maximum difference between the total ozone simulated by the NO_x and RC-tagged simulations is less than 1%, and these small differences are mostly related to the truncation of simulated values beyond a few decimal places.

2.2 Model Evaluation

CAM4-Chem has been evaluated in previous studies for simulating tropospheric ozone and precursors (Lamarque et al., 2012, Tilmes et al., 2015), and was also evaluated in its ozone-tagged configuration by Butler et al., (2018) and Butler et al., (2020).

Here we evaluate the configuration of CAM4 Chem used in this study, which primarily differs from the earlier work through its use of the HTAPv3 global emissions. In Fig. 1 we compare our simulated surface ozone against gridded observation dataset provided by Tropospheric ozone assessment report until year 2014 (TOAR; Schultz et al., 2017) until year 2014. We also make the comparison using simulation output from “CESM2-WACCM6” model which is part of the CMIP6 ensemble (Emmons et al., 2020), using this as a standard reference model. Results are shown as monthly averages over various HTAP Tier 2 regions (Galmarini et al., 2017) at the grid cells where the TOAR data is available. We overestimate the surface ozone mixing ratio by up to ~4-12 ppbv over most regions where TOAR data is available, largely during the summer months. This overestimation is also simulated by CESM2-WACCM6 model (shown in Fig. 1) and is consistent with high model bias of ~7 ppbv simulated by most models in various ensembles, as discussed in Young et al., (2018). We also simulate an underestimation of up to 2-6 ppbv over northern and eastern United States during winter months, which is not simulated by CESM2-WACCM6. For the year 2010, we simulate slightly smaller surface ozone at most regions over grid cells where TOAR data is available compared to that simulated by Butler et al., (2020), and is closer to the observations from TOAR dataset.

We evaluate free tropospheric ozone against the ozone sonde-based climatology compiled by Tilmes et al., (2012). Figure 2 shows the comparison our simulated climatology (2000-2010) to the ozone sonde climatology (1995–2010) in Taylor-like diagrams (Taylor, 2001; Tilmes et al., 2012) at 900, 500, and 250 hPa pressure levels, grouped by latitude ranges. The correlation coefficient between the observed and simulated monthly regional O₃ average is usually more than 0.8, and the fractional mean difference is usually within 25 % at most regions in the troposphere. The model captures the vertical distribution of ozone derived from ozone sonde climatology very well, although slightly biased high particularly in the upper troposphere at most sites (Fig. S2). Our free troposphere evaluation results are therefore largely consistent with previously evaluated versions of CAM4-Chem (Tilmes et al., 2012, Zhang et al., 2016, Emmons et al., 2020).

2.3 Trend Analysis

We express the trend for the 2000-2018 period in various quantities discussed in our study: Precursor emissions, tropospheric ozone burden, global surface mean and population-weighted mean ozone concentration as a slope of the timeseries in respective units/yr, and in %/yr relative to the first absolute value (for year 2000). This slope is calculated using the Theil-Sen

estimator method available as a python module
(<https://docs.scipy.org/doc/scipy/reference/generated/scipy.stats.theilslopes.html>), which also calculates the 95 % confidence interval of the slope. We further use the Mann-Kendall test to determine the extent to which the detected trend is monotonic (Hussain and Mahmud, 2019), expressed by the p-value. In the TOAR Guidelines for best statistical practices (Chang et al., 2023), it is recommended not to use a dichotomized expression such as significant/insignificant trend. Hence, we also provide the p-value and the 95 % confidence interval in the supplementary information for the interested readers to assess the meaningfulness of the estimated trend in our study. For the sake of discussion in sections 3 and 4, we categorize the trends with p-values less than 0.05 to be of high certainly, between 0.05 and 0.1 to be of medium certainty and larger than 0.1 to be of low certainty (Table 3 in Chang et al., 2023).

3 Results

We discuss our results related to the ozone precursor emissions, contribution to tropospheric ozone burden, ozone production efficiency, and contribution to surface mean and population-weighted mean ozone from each of the tags used in our study. Each subsection is further divided into two parts to discuss the role of NO_x and RC tagged contributions separately. All our results are discussed as timeseries of annual means for the 2000-2018 period, unless otherwise specified. When referring to anthropogenic emissions and their contributions to tropospheric ozone in our discussion, we only refer to surface-based anthropogenic emissions, and exclude aircraft emissions.

3.1 Ozone precursor emission trends

3.1.1 NO_x emissions

Table 2 shows the global NO_x emissions (averaged over the 2000-2018 period) from various regions and sectors, and their relative contribution (in %) to the total NO_x emissions. Fig. 3 (a and c) shows the time-series of NO_x emissions from all sources: natural/global sources (a), and regional anthropogenic sources (c).

Over the period 2000-2018, the average annual NO_x ~~being~~ emissions are ~~~67~~49 TgN/yr, which includes both surface and aloft emissions. There is a significantly increasing trend in the total NO_x emissions over the 2000-2018 period, with 63.3 TgN/yr emitted in 2000, peaking at 70 TgN/yr in 2011 and decreasing to 66 TgN/yr emitted in 2018 (Fig. 3(a); Table 3). The trend in total NO_x emissions is mainly driven by the anthropogenic NO_x emissions which are the largest source of NO_x (~52.21 TgN/yr (78 %); Table 2), much larger than natural sources such as lightning, biomass burning and biogenic emissions which contribute by ~~~5~~6~~7~~8.5 and 10 % respectively (Table 2). We also notice a slight decrease in the global anthropogenic NO_x emissions in 2008-2009, mostly related to the global financial crisis (Schneider and van der A 2012).

230 Among the anthropogenic NO_x emissions, East Asian emissions are the largest (~20 %; Table 2), with ~~significantly an~~
increasing trend ~~with high certainty~~ (~~~0.25 TgN/yr²~~; Table 3 ~~and Fig. 3~~) over the ~~period~~ 2000-2018 period. These emissions
~~increase at 0.49 TgN/yr²~~, peak in 2011 and start declining after that ~~at -0.44 TgN/yr²~~ (Zheng et al., 2018), as shown in Fig. 3c
~~and Table S2~~, which largely explain the timeseries of anthropogenic and total NO_x emissions timeseries shown in Fig. 3a. The
NO_x emissions increase ~~in-over~~ South Asia (~~2.350.13~~ TgN/yr²), Middle East (~~1.760.06~~ TgN/yr²) and “Rest of the world”
235 (~~5.780.15~~ TgN/yr²), whereas they decreased ~~in-theover~~ North America (4.16 TgN/yr²), Russia-Belarus-Ukraine region (~~1.42-~~
~~0.02~~ TgN/yr), and Europe (~~2.88-0.11~~ TgN/yr²) ~~regions~~. The magnitude of international ship NO_x emissions (~~5.924.04~~ TgN/yr)
is comparable to that of NO_x emissions from continental regions such as North America, Europe, and South Asia (~~Table 2~~),
and have an ~~significantly~~-increasing trend (~~~0.08 TgN/yr²~~; Table 3) ~~with high certainty~~. These trends effectively also indicate
an overall equatorward shift in anthropogenic emissions as discussed in several previous studies (e.g., Zhang et al., 2016,
240 Gaudel et al., 2020, Zhang et al., 2021). ~~We further illustrate this equatorward shift using zonal sum profiles of deviation in~~
~~anthropogenic NO_x emissions from year 2000 (Fig. S4a). The zonal NO_x emissions relative to year 2000 show a positive~~
~~(negative) deviation south (north) of ~35°N latitude, indicating an equatorward shift in the overall global NO_x emissions.~~

Formatted: Superscript

Emissions from the aircraft sector contribute only ~~1.40.93~~ TgN (~~~2.141.9~~ %) to the total NO_x emissions, with an increasing
245 trend of ~2.5 %/yr. This is comparable to the emissions taken from CEDS inventory reported by Wang et al., (2022): 0.88 TgN
(contributing ~3.3 % to total NO_x emissions) increasing at ~3.46 %/yr between 1995 and 2017.

Formatted: Subscript

3.1.2 Reactive Carbon emissions

The total amount of tropospheric reactive carbon emissions averaged over the 2000-2018 period in our simulations is ~~~1407~~
~~1342~~ TgC/yr (Table 4) with small increasing trend (~~0.258~~ %/yr²~~ear~~; Table 5) (Fig. 3(b)). Methane and biogenic ~~NM₂VO₂C₂s~~
250 ~~NM₂RC₂s~~ are the largest sources of reactive carbon (each about ~430-~~500~~-TgC (~~30-3532~~ %); Table 4; Fig. 3b; Heald and Kroll
2020). Total anthropogenic ~~NM₂VO₂C₂-NM₂RC₂~~ emissions contribute ~~only~~ ~300 TgC/yr (~~~224~~%; Table 4)-) ~~and~~ Biomass burning
~~emissions contribute ~171 TgC/yr (~12.77 %) and~~. Aircraft emissions contribute very small amounts (Table 4).

Formatted: Superscript

Since the concentration of methane is fixed at the surface in our simulations, we consider the methane oxidation rate derived
255 from methane’s oxidation reaction with atmospheric hydroxyl radical as the effective source of methane for subsequent ozone
production through rapid chemical reactions. We simulate ~~~492-437~~ TgC/yr of methane oxidation with an ~~significantly~~
increasing trend ~~of high certainty~~ over the 2000-2018 period. The prescribed CH₄ concentrations are such that there is a plateau
until the year 2006 (Lan et al., 2024) followed by a steep increase (Fig. 4a: ~~black-red~~ line). ~~Our simulated methane oxidation~~
~~rate also increases with high certainty but does not strictly follow this plateau followed by steep increase pattern and has large~~
260 ~~inter-annual variability (Fig. 4a). This could possibly be due to variability in the prescribed meteorology and OH concentration~~

(Fig. 4b) resulting in variations in the oxidation rate of methane. Our simulated methane oxidation rate does not strictly follow this pattern due to some inter-annual variability in the model. We simulate slightly larger methane oxidation for the year 2010 compared to that of simulated by Butler et al., (2020), possibly due to the usage of a different anthropogenic emissions dataset, prescribed CH₄ concentration-mixing ratio and prescribed meteorology dataset. A detailed comparison of methane related variables such as prescribed concentration, burden, lifetime, oxidation rate etc., between our study and Butler et al., 2020 for the year 2010 is provided in Table S4. While the absolute contributions from methane oxidation, and anthropogenic and biogenic NMVOC-NMRC emissions to total reactive carbon emissions increases significantly with medium-high certainty (Fig. 3b), its relative contributions doesn't show any significant trends with low certainty (Supplementary Figure S3b)

The tropospheric methane lifetime (~8.54 years), calculated as the total atmospheric methane burden divided by tropospheric methane oxidation rate, decreases significantly with high certainty (-0.01 years/yr; Table S5) in our simulations (Fig. 4a; Magenta line). This result is contrary to the expectation that methane lifetime might increase due to increasing methane concentrations leading to a smaller availability of OH radicals to oxidize methane (Prather et al., 1996). CH₄ oxidation with OH radical being the major loss pathway for CH₄, we show the time-series of OH in Fig. 4b. Here we simulate an increasing trend in air-mass-weighted tropospheric OH concentration (a prominent indicator for tropospheric oxidizing capacity; e.g., Voulgarakis et al., 2013, Chua et al., 2023) but with low certainty. Nevertheless, the CH₄-reaction-weighted tropospheric OH concentration (Lawrence et al., 2001) shows an increasing trend with high sensitivity. This increasing trend in tropospheric OH concentration is consistent with the trend discussed in Chua et al., (2023) and explains the decreasing lifetime of CH₄ as these quantities are inversely proportional. The increasing OH availability despite increasing CH₄ concentration could be due to increasing NO_x emissions over the simulated period that recycle HO₂ to OH (e.g. Lelieveld et al., 2008; Chua et al., 2023). The mean magnitude of our air-mass-weighted tropospheric OH concentration (~12.63 * 10⁵ molec/cm³) is slightly larger than previous studies (~10.75 * 10⁵ molec/cm³ in Chua et al., 2023 or ~11.7 * 10⁵ molec/cm³ in Voulgarakis et al., 2013) which is likely why we simulate a mean CH₄ lifetime slightly smaller than the aforementioned studies (~9 years), but within the multi-model range (7.1-10.6 years) reported in Voulgarakis et al., (2013). In our simulation, however, there is a significant increase in total NO_x emissions, which is consistent with more HO₂ to OH recycling (Lelieveld et al., 2008) leading to more methane oxidation and a decreasing lifetime.

Among the anthropogenic NMVOC emissions, East Asian emissions are the largest (~95.36 TgC/yr (6.78 %); Table 4), increasing at 3.54 TgC/yr² peaking in 2011 and decreasing after that at -2.82 TgC/yr² (Fig. 3(d); Table S6) as for the NO_x-tagging simulation emissions. Second largest emitter is the “Rest of the World” region (~90.77 TgC/yr (6.45 %); Table 4), with an significantly increasing trend of high certainty (Fig. 3 (d)). Remaining regions each contribute to less than 5 % of reactive carbon emissions. As for the anthropogenic NO_x emissions: South Asian, Middle Eastern, and ship NMVOC emissions show an increasing trend, and North American, European, Russia-Belarus-Ukraine region's NMVOC emissions

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Superscript

Formatted: Superscript

Formatted: Subscript

show a decreasing trend (Fig. 3 (d)). We also see an equatorward shift in global anthropogenic NMRC emissions, similar to the anthropogenic NO_x emissions in our zonal sum profiles of deviation in anthropogenic NMRC from year 2000 (Fig. S4b).

Formatted: Subscript

3.2 Tropospheric ozone burden

The tropospheric ozone burden is calculated as the mass of ozone in the model grid cells below the ozone-tropopause, defined as the where ozone concentration is highest layer in the upper troposphere with an ozone mixing ratio less than 150 ppb (E.g., Bak et al., 2022, Liu et al., 2022). We simulate a climatological average tropospheric ozone burden of ~ 336.418 Tg O_3 , with an significantly-increasing trend of $0.94-72$ Tg O_3/yr over the 2000-2018 period. Our simulated tropospheric ozone burden is within the range towards the lower end of values simulated by several multi-model studies for the 2005-2014 period (values from Griffiths et al., 2021): the CMIP 6 ensemble 356 ± 31 Tg O_3 ; ACCENT: 336 ± 27 Tg O_3 ; Atmospheric Chemistry and Climate Model Intercomparison Project, ACCMIP: 337 ± 23 Tg O_3 ; Tropospheric Ozone Assessment Report, TOAR: 340 ± 34 Tg O_3 ; and Intergovernmental Panel on Climate Change, IPCC: 347 ± 28 Tg O_3 (Szopa et al., 2021). The trend in tropospheric ozone burden is also within the range of trends simulated by CMIP 6 model ensemble: 0.4 to 1.3 Tg O_3/yr , as reported by Wang et al. 2022. When integrated from 60°S to 60°N , we simulate a climatological mean tropospheric ozone burden of $\sim 302.28287.38$ Tg O_3 (within lower end of the range of 287-311 Tg O_3 estimated by satellite products for 2014-2016 period; Gaudel et al. 2018), with an significantly-increasing trend of $0.84-68$ Tg O_3/yr (close to the trend of 0.82 ± 0.13 Tg O_3/yr simulated by CMIP 6 ensemble members; Griffiths et al. 2021).

We simulate a substantial amount of tropospheric ozone attributed to stratospheric influx in the NO_x -tagged ($\sim 87.691.5$ Tg O_3 (~ 296 %); Table 2) and RC-tagged simulation (~ 764 Tg O_3 (~ 242 %); Table 4). The larger amount of ozone attributed to stratospheric influx in the NO_x -tagged simulation compared to the RC-tagged simulation is because of the production of tropospheric O_3 attributed to NO_x from the stratosphere formed by the oxidation of N_2O , which is then subsequently transported into the troposphere (Butler et al., 2018). As described in Section 3.1.2 of Butler et al. (2018), ozone production in the stratosphere is handled directly in the chemical mechanism with the addition of new reactions producing the relevant odd oxygen species. Similarly, production of NO in the stratosphere from the oxidation of N_2O by O1D is also handled directly with an additional chemical reaction.

Formatted: Subscript

We attribute minor ozone production pathways with the “Extra Production” tag. In our NoxNO_x -tagged simulation this category consists of O_x production from the self-reaction of OH radicals, and from reactions between HO_2 and organic peroxy-radicals. In our RC-tagged simulation this category consists of O_x production from the self-reaction of OH radicals, and production of HO_2 from the reaction of OH with H_2O_2 . The contribution of these minor production pathways to the total tropospheric ozone burden is not very substantial (up to 2 %; Table 2 and Table 4)

325 **3.2.1 Tropospheric ozone burden attributed to ~~NOx~~NO_x emission sources.**

Formatted: Subscript

Our simulation shows that ~713 % of tropospheric ozone is attributed to NO_x emissions (both surface-based and aloft). The remaining ~297 % of tropospheric ozone burden is attributed to the stratospheric influx, and a small contribution from minor production pathways, as discussed above. We note that the tropospheric ozone attributable to stratospheric influx in our ~~NOx~~NO_x-tagged simulation is larger than the stratospheric contribution in the reactive carbon tagged simulation (243%, described below) due to the inclusion of tropospheric ozone production from stratospheric NO_x (as described above).

Anthropogenic ~~NOx~~NO_x emissions are the largest contributors to tropospheric ozone (11634 Tg O₃ (~ 3740 %)), followed by lightning NO_x (7165 Tg O₃ (229 %)), biogenic NO_x emissions (230 Tg O₃(7.46.5 %)) and aircraft ~~NOx~~NO_x emissions (13.47.88 Tg O₃ (~4.235.32 %); Table 2). Lightning NO_x has a large contribution despite a small amount of NO_x emissions because of large Ozone Production Efficiency (OPE) of NO_x when emitted aloft (Table 2). Further discussion about OPE is provided in section 3.3. Biomass burning NO_x emissions contribute less-than-5~4 % to the total tropospheric ozone burden.

Formatted: Subscript

Formatted: Subscript

Among regional anthropogenic NO_x sources, the contribution from the “Rest of the World” used in this study (Table 1) to the tropospheric ozone burden is largest among the anthropogenic NO_x emissions, followed by ship NO_x emissions. This “Rest of the World” contribution is even larger than that of East Asian contribution which is the region with the largest anthropogenic NO_x emission among continental regions considered in this study (Fig. 3, Table 2). This is due to more efficient production of ozone and convection into the free troposphere at the tropical regions (Zhang et al., 2016, Zhang et al., 2021, See further discussion in section 3.3.1) that are included within the “Rest of the World” tag. The disaggregated contribution of explicitly tagged regions within the “Rest of the World” tag in our ~~NOx~~NO_x-tagged simulation: Southeast Asia, Central Asia, North Africa, and Mexico-Central America (Table 1; Fig. S1b) are provided in the Supplement (Tables S52 and S63). We notice that the largest contributors among these regions are the NO_x emissions from the tropical regions: Mexico and Central America and Southeast Asia (Table S52). Similarly, the tropically situated South Asian contribution to the tropospheric ozone burden is larger than that of European contribution despite having a smaller amount of NO_x emissions (Table 2), due to the larger convection into the free troposphere as stated above.

To illustrate the predominant role of convection in transporting ozone and it’s precursors at the tropics into the free troposphere, we show the zonal mean vertical profiles of annual mean O₃ and NO_x (sum of NO and NO₂) mixing ratios for the 2000-2018 climatological mean attributed to NO_x emitted from a typical tropical region: Southeast Asia and a typical northern mid-latitude region: Europe (Fig. 6; see Fig. S1 for the location of defined regions). The vertical gradient in the ozone attributed to European NO_x is consistent with summertime production of ozone primarily within the boundary layer with subsequent vertical transport into the free troposphere. While there is indeed some ozone attributed to Southeast Asian NO_x emissions present in the

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Font color: Auto

Formatted: Subscript

360 boundary layer, the mixing ratio of this attributed ozone is much higher in the free troposphere, which is consistent with ozone
production aloft. This is due to emitted NO_x directly being transported aloft, eventually leading to free tropospheric ozone
production. Further, we also note the increasing trend in NO_x burden attributed to anthropogenic NO_x emissions in the free
troposphere (above 700 hPa; Figure S6)

365 The contribution from ship NO_x is also large due being emitted at remote ocean regions where ship NO_x is the only source of
NO_x emissions. This results in less competition among tagged NO_x sources in producing ozone leading to very efficient ozone
production from ship NO_x (Butler et al., 2020). Further discussion about ozone production efficiency of ship NO_x is provided
in section 3.3.1.

370 The sign of the trend in the contribution to the tropospheric ozone burden from each of the tags (Fig. 5 (a,c)) is consistent with
the sign of the NO_x emissions trend (Fig. 43 (a,c), Table 3). The percentage slope (ignoring the sign) of the trend in the
contribution to tropospheric ozone burden by regional anthropogenic NO_x sources, however, is generally smaller than that of
NO_x emission trend (Table 3). Similarly, the magnitude of the percentage slope in the trend in tropospheric ozone burden
attributed to East Asian NO_x emissions for both the pre-2011 (2000-2011) and post-2011 (2011-2018) periods is smaller
compared to the corresponding percentage slope in East Asian NO_x emissions trend (Table S2). The reason for this is due to
changing ozone production efficiency with changing NO_x emissions, explained in detail in section 3.3.1.

375 We simulate a trend of ~0.2835 Tg O₃/yr (2.4129 %/yr; Table 3) in ozone burden attributed to aircraft NO_x emissions, which
is comparable to 0.3 TgO₃/yr estimated by Wang et al. (2022) using sensitivity simulations, where the difference between a
simulation with transient aircraft emissions and that with fixed aircraft emissions delivers the exclusive impact of changing
aircraft emissions over the 1995-2017 period. While the trend derived from our tagged simulation only delivers the trend in
contribution from aircraft NO_x emissions, a sensitivity simulation would also cover the changes in simulated ozone contributed
380 from various sectors when their ozone production efficiency changes as a result of emissions being held constant. A
combination of sensitivity simulations with tagging included would help us track these non-linearities and compensating
feedbacks that would otherwise not be seen from sensitivity-only or tagging-only simulations.

3.2.2 Tropospheric ozone burden attributed to reactive carbon emission sources.

385 In our VORC-tagged simulation ~768 % of the total amount of tropospheric ozone burden is attributed to tropospheric reactive
carbon, while the remaining ~243 % is attributed to stratospheric influx and minor production pathways as discussed above.
Of the reactive carbon species sources, methane oxidation is the largest contributor to the tropospheric ozone burden
(~132.4548.6 Tg O₃ (42.4%); Table 4) and is consistent with previous studies (Young et al., 2013, Butler et al., 2018). We
simulate an increasing significant trend with high certainty in the part of the tropospheric ozone burden attributed to RC from

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Justified

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

methane oxidation (Fig. 5b), consistent with the significantly-increasing methane oxidation rate during the 2000-2018 period as shown in Fig. 4a.

The contribution to tropospheric ozone burden from biogenic reactive-carbonNMRC emissions (534 Tg O₃ (~17 %); Table 4) is much smaller than that of methane, despite having a comparable mass of carbon-RC emissions from both speciesources. This is consistent with the smaller number of oxidisable bonds per carbon atom in biogenic emissions, mainly isoprene (2.8) compared with methane (4) and the subsequently lower number of NO to NO₂ conversions possible (per carbon atom) during isoprene oxidation compared with methane oxidation (Edwards and Evans, 2017). We simulate much smaller contributions of NMVOG-NMRC emissions from biomass burning, and aircraft NMVOGs-NMRCs compared to biogenic and anthropogenic sources (consistent with Butler et al., 2020; Table 4).

Contribution to tropospheric ozone from-attributed to anthropogenic reactive-carbonNMRC emissions is relatively low at ~460 Tg O₃ (142 %; Table 4).

Regional anthropogenic tags each contribute less than 5 % to the total tropospheric ozone burden. The sign of the trend in tropospheric ozone attributed to most of the tags, including regional anthropogenic NMRC tags, in our RC-tagged simulation is consistent with that of the trend in the corresponding reactive carbon emissions (Table 5). We also note that the percentage trend in contribution to tropospheric ozone burden from anthropogenic NMRC (1.03 %/yr) is more than that of the trend in anthropogenic NMRC emissions (0.61 %/yr), whereas the trend in contribution from methane oxidation (0.33 %/yr) is less than that of trend in methane oxidation rate (0.42 %/yr). While the biogenic NMRC emissions show an increasing trend with medium certainty, the trend in its absolute contribution to tropospheric ozone burden shows a small trend with low certainty.

3.3 Ozone production efficiency (OPE)

We estimate ozone production efficiency of emissions of any given sector as the ratio of annual mean tropospheric ozone burden (in mol O₃) attributed to that sector to the annual-meanamount of ozone precursors emissions-emitted in a given year (either NO_x (in mol N) or reactive carbon (in mol C)). Several previous studies report OPE as the ratio of response in tropospheric ozone burden to change in emissions introduced in the model, or as a ratio of the production rate of ozone to the loss rate of NO_x (e.g. Liu et al., 1988, Kim et al., 2016, Miyazaki et al., 2020, Archibald et al., 2020, etc.). Since our model directly calculates the ozone attributed to emissions from tagged emission sources, the ratio of the attributed tropospheric ozone burden to the emitted amount of ozone precursor is calculated as OPE of that tagged precursor source. This can only be made possible when the NO_xNO_x and reactive carbon precursors are tagged in two separate simulations, as explained in Butler et al., (2020). A similar metric: Ozone Burden Efficiency (OBE) has been defined by Mertens et al., (2024), as the ratio of

Formatted: Subscript

ozone attributed of precursor emissions (both NO_x and RC) from a given source using the combinatorial tagging approach (in Tg O3) to the emitted NO_x (Tg NO) for various tagged sectors.

3.3.1 OPE of NO_x emissions.

Lightning NO_x (5.9346 mol $\text{O}_3/\text{mol N}$) and aircraft NO_x (4.193.64 mol $\text{O}_3/\text{mol N}$) are most efficient at producing ozone (Fig. 6(a), Table 2) directly into the free troposphere, due to being emitted aloft (Hoor et al., 2009, Dahlmann et al., 2011). NO_x emissions aloft are highly efficient at producing ozone due to the relatively low quantities of NO_x at higher altitudes. Low NO_x concentrations increase the ozone production efficiency of NO_x due to a lower frequency of radical termination reactions compared with higher NO_x concentrations, so that each molecule of NO_x can produce more molecules of ozone before being lost by reaction with OH (Seinfeld and Pandis, 2016). In contrast, biogenic (1.423 mol $\text{O}_3/\text{mol N}$), biomass burning (0.882 mol $\text{O}_3/\text{mol N}$), and anthropogenic NO_x (0.9674 mol $\text{O}_3/\text{mol N}$) are comparatively less efficient at producing ozone as they are emitted at the surface into regions with comparatively large NO_x concentrations.

Among regional anthropogenic NO_x emissions, Rest of the World is the most efficient (1.844), as it mainly consists of tropical regions where the convection into the free troposphere is large. We further elaborate the explicitly tagged regions in our NO_x -tagged simulation within the Rest of the World tag in Table S52. The tropical regions: Mexico and Central America (1.816 mol $\text{O}_3/\text{mol N}$), and Southeast Asia (1.9873 mol $\text{O}_3/\text{mol N}$) are the most efficient among all the explicitly tagged regions. The larger sensitivity of tropospheric ozone to emission changes in these tropical regions has also been noted in Zhang et al., (2021). Similarly, OPE of South Asian NO_x (0.7866 mol $\text{O}_3/\text{mol N}$) and to some extent Middle Eastern NO_x (0.654 mol $\text{O}_3/\text{mol N}$) is much larger compared to that of North American (0.544 mol $\text{O}_3/\text{mol N}$), European (0.36 mol $\text{O}_3/\text{mol N}$), Russia-Belarus-Ukraine region's (0.433 mol $\text{O}_3/\text{mol N}$) and East Asian (0.5419 mol $\text{O}_3/\text{mol N}$) NO_x , for being more tropically situated (e.g., Wild et al., 2001, Butler et al., 2020, Zhang et al., 2021 etc.). International shipping is the second most efficient (1.6733 mol $\text{O}_3/\text{mol N}$) source of anthropogenic NO_x emissions in producing ozone, due to their presence at remote pristine regions where there is a much smaller availability of other sources of NO_x (Lawrence and Crutzen 1999, Butler et al., 2020). We note that the chemistry occurring within the expanding plume emitted by ships is not considered in our coarse resolution model, leading to an instantaneous dilution of emitted NO_x into the large grid cells (Vinken et al. 2011) thereby making ozone production from Ship NO_x more efficient than observed (Kim et al. 2016).

The increasing trend in anthropogenic NO_x emissions from South Asia, Middle East, and international shipping (Fig. 3c and Table 3) results in less efficient production of tropospheric ozone from these sources over time (Fig. 76c and Table 3). Similarly, there is an increasing trend in OPE of anthropogenic NO_x emissions from regions with decreasing emissions (North American, European, and Russia-Belarus-Ukraine regions; Fig. 76c). The decreasing (increasing) NO_x emissions from regional anthropogenic sources becoming more (less) efficient at producing ozone, and this leads to a dampening effect where there is

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

a smaller percentage slope (ignoring the sign) in tropospheric ozone burden compared to the slope in NO_x emissions (Table 3). This result is also consistent with Mertens et al., (2024) where an increase in NO_x emission from various tagged sectors is associated with a decrease in OBE in both present-day and future scenarios.

Fig. 7c shows that the OPE of East Asian NO_x has a trend of low certainty over the 2000-2018 period. However, when shorter periods are considered, it decreases with high certainty during the pre-2011 period when emissions increase (2000-2011 period) and increases with high certainty for the post-2011 period when emissions decrease (2011-2018 period; Table S2).

Based on the results on trends in OPE of various anthropogenic NO_x emissions from regions discussed above, we could expect a significant increase in OPE of East Asian anthropogenic NO_x emissions. However, we do not simulate a significant trend in OPE (Fig. 6c, Table 3), likely due to variability within the 2000-2018 period. This could be introduced by the variability in emissions, other boundary conditions during this period such as prescribed meteorology, methane concentration etc., or the thresholds in the trend significance estimation tool used in our study (Hussain and Mahmud 2019).

In case of total anthropogenic NO_x emissions, we do not simulate a significant decrease in their OPE despite their significantly increasing emission trend. We rather simulate an significant increase in OPE of anthropogenic NO_x with high certainty. This is due to an equatorward shift in emissions over the 2000-2018 period (Fig. S4), where the emitted NO_x from the tropics and the subsequently formed ozone and NO_y (NO_x and reservoir species for NO_x; e.g., NO₃, N₂O₅ etc.) molecules are rapidly lifted into the free troposphere as discussed previously (Zhang et al., 2016). We also notice that the increase in anthropogenic emissions leads to a decreasing trend in OPE of other natural sources of NO_x such biogenic, biomass burning and lightning, as these natural sources must compete with more NO_x emissions from anthropogenic sources in order to produce ozone.

3.3.2 OPE of reactive carbon emissions.

Consistent with Butler et al., (2020), we simulate methane oxidation as the most efficient (0.073-08 mol O₃/mol C; Table 4) among the tagged reactive carbon sources in our VOCRC-tagged simulation. Similarly, we simulate smaller OPEs for biogenic (0.0347 mol O₃/mol C), biomass burning (0.0298 mol O₃/mol C), and anthropogenic (0.0326-04 mol O₃/mol C) reactive carbon sources. As noted in section 3.2.2, RC from methane oxidation is expected to be more efficient at producing ozone per unit of carbon due to the higher number of oxidisable bonds per carbon atom (Edwards and Evans., 2017). Anthropogenic reactive carbon emissions might be expected to be more efficient than biogenic reactive carbon emissions at producing ozone due to their proximity to anthropogenic NO_x emissions, but we see no large differences in OPE in our study (Table 4).

We simulate no significant trend a trend with low certainty in the OPE RC from natural sources such as of biogenic, biomass burning and methane as reactive carbon sources oxidation (Fig. 7; Table 5). We do simulate an significantly increasing trend with high certainty in the OPE of global anthropogenic reactive carbon emissions. Among the regional anthropogenic sources,

Formatted: Subscript

Formatted: Subscript

Formatted: Not Superscript/ Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

485 we ~~do not~~ simulate ~~any significant~~ trends ~~with low-medium certainty~~ for European, South Asian and Rest of the World ~~RC~~
emissions. We simulate increasing trend for North American and East Asian emissions, and a decreasing trend for the OPE of
the remaining tagged regional anthropogenic reactive carbon emissions. ~~OPE of East Asian anthropogenic RC emissions have~~
~~a small trend with low certainty over the 2000-2011 period and then increases with high certainty during the 2011-2018 period~~
(Table S6)

490

We would expect that the ozone production by reactive carbon emissions should mainly depend on the availability of NO_x,
especially in pristine environments. The trends in the OPE of ~~regional~~ anthropogenic ~~reactive carbon~~ NMRC emissions largely,
but ~~do not~~ always, follow the trends in the emissions of NO_x in their vicinity (Table 3 and 5). However, the trends in the
emission of the ~~reactive carbon~~ RC from a given sector itself could influence the OPE of its own emissions, and the OPE of
495 the NMRC emissions ~~of reactive carbon~~ from other sources. For example, Butler et al., (2020) illustrate how an increase in
prescribed methane concentration leads to smaller OPE of not just ~~RC from~~ methane ~~oxidation~~ but also of other tagged
~~NM VOC-RC~~ emission sectors. To better understand the behaviour of changes in OPE of reactive carbon emissions, further
studies could investigate the OPE changes from perturbation simulations, ~~with by enabling~~ ozone source attribution with
tagging ~~enabled~~ in the simulations. Mertens et al., (2018) demonstrate that the combination of tagging and perturbation can be
500 used to explain the compensating feedbacks induced by perturbation, from various tags used in the simulations.

3.4 Contributions from tagged precursor emission sources to surface ozone.

In this subsection, we discuss the contribution of various NO_x and reactive carbon sources to both the global surface mean and
population weighted mean ozone. We further contrast this contribution with that of their contribution to the tropospheric ozone
burden (discussed in section 3.2).

505

We select the population count for the year 2020 Gridded Population of the World (Center for International Earth Science
Information Network – CIESIN – Columbia University, 2018) and scale the surface ozone according to this distribution to
calculate the ~~global~~ population-weighted ~~mean~~ ozone, which is related to the ozone exposure. Choosing only one year for
population weighting gives the exclusive effect of changing emissions and not the effect of changing population. Future studies
510 may consider scaling according to a transient population over the analysis period rather than fixing it at one year (year 2020
in this study). Table S94 provides the population, and their fraction of world population (in %), for the regions considered in
this study.

We simulate ~ ~~27.13~~ ~~25.39~~ ppbv of global mean surface ozone with ~~an~~ ~~significantly~~-increasing trend ~~of high certainty~~ (0.08
515 ~~ppbv/yr~~ (0.37 %/yr); Fig. 7 and Table 3). Similarly, we simulate ~33.6-01 ppbv of global population-weighted mean ozone
with ~~an~~ ~~significantly~~-increasing trend ~~with and~~ slightly larger slope (0.17 ppbv/yr (0.48 %/yr); Fig. 8 and Table 3) ~~compared~~

to global surface mean ozone. We will explain the main contributors responsible for this increasing trend and the larger slope in global mean population-weighted ozone in the subsequent discussion. DeLang et al., 2021 showed that the increasing trend in global ozone exposure is mainly driven by increasing ozone exposure at highly populated regions such as Asia and Africa. In our study, we quantify the contribution of various ozone precursor emission sources to the global population-weighted surface ozone and its trend.

The share of the stratospheric contribution to the global mean surface ozone (4-5 ppbv (~15-20+8%); Table 2 and Table 4), is smaller than its contribution to tropospheric ozone burden (22-27 %; Table 2 and Table 4). We simulate an significantly increasing trend with high certainty (~ 0.3-0.4 %/yr; Table 3 and Table 5) in this contribution mainly over the southern hemisphere (not shown). The stratospheric contribution to the population weighted surface ozone is much smaller (~3 ppbv (9-11 %); Table 2 and Table 4) compared to the quantities discussed above, as this is mainly contributing over remote regions such as polar and oceanic regions (not shown).

3.4.1 Surface ozone attributed to ~~NO_x~~NO_x emissions.

The percentage contribution to global mean surface ozone from sources aloft (lightning, aircraft, stratosphere) is smaller compared to the contribution to tropospheric ozone burden (Table 2). Consequently, the surface-based ~~NO_x~~NO_x sources (biogenic, biomass burning and anthropogenic) contribute a larger share to the global mean surface ozone compared to their share of the tropospheric ozone burden.

The share of the anthropogenic NO_x contribution has the largest difference in its contribution from 379 % (to tropospheric ozone burden) to 548 % (to global mean surface ozone; Table 2). ~~Most-All the tagged~~ regional anthropogenic NO_x sources contribute a larger percentage share to global mean surface ozone than to the tropospheric ozone burden (Table 2). ~~However, in the case of South Asian anthropogenic NO_x, there is a smaller contribution to global mean surface ozone (2.7 %) than to tropospheric ozone burden (2.81 %; Table 2) due of being situated in a tropical region with large convection to the free troposphere (Zhang et al., 2016; Zhang et al., 2021). Similarly~~However, the tropically situated explicitly tagged regions within the “Rest of the Word” tag in our NO_x-tagged simulation: Mexico and Central America, and Southeast Asia have smaller share in their contribution to global mean surface ozone compared to that of tropospheric ozone burden (Table S72), largely due to being situated in a tropical region with large convection to the free troposphere (Fig 6; Zhang et al., 2016; Zhang et al., 2021). We simulate a small increasing trend (0.08 ppbv/yr (0.307 %/yr); Table 3) in global mean surface ozone, mainly driven by increasing stratospheric contribution (0.02 ppbv/yr (~0.32 %/yr)) and increasing trend in contribution from anthropogenic NO_x sources (0.08 ppbv/yr (0.7 %/yr); Table 3 and Fig 7c).

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

The contribution from anthropogenic NO_x to the population weighted surface ozone (~20 ppbv (627 %); Table 2) is larger than its contribution to global mean surface ozone. This larger contribution is compensated by smaller remote contributions such as stratosphere, lightning, aircraft, and biomass burning. Among the regional anthropogenic sources, the NO_x emissions from highly populated regions (see Table S94 for population of world regions): South Asia (4.6 ppbv (~145.5 %)), East Asia (~4.3 ppbv (132.34 %)) and Rest of the World (~4.3 ppbv (135.4 %)) are the largest contributors to global population-weighted mean ozone. Ship NO_x emissions also have a substantial contribution (~2.5 ppbv (7.5 %8.38 %)) to the global population-weighted surface mean ozone, larger than contributions from ~~major anthropogenic NO_x emissions by relatively sparsely~~ populated world regions such as North America, Europe, Russia-Belarus-Ukraine region, or Middle East (Table 2). The increasing trend in global population-weighted surface ozone (0.17 ppbv/yr (0.5448%/yr); Table 3) which is slightly larger compared to the trend in global mean surface ozone, is mainly driven by increasing anthropogenic NO_x emissions especially from highly populated regions (South Asia, East Asia and “Rest of the World”) and from international shipping. Despite NO_x emissions from international shipping happening only over ocean grid cells, the ozone attributed to ship NO_x spreads across land areas and contributes to the total surface ozone by ~3-6 ppb (Fig. 10a). Although there is no population over the regions where international ship NO_x is emitted, the global population weighted mean ozone attributed to ship NO_x is comparable to that of global mean surface ozone (Table 2) as large populations at several coastal areas and densely populated inland regions in East Asia and South Asia are exposed to ozone attributed to ship NO_x emissions (Fig. 10b). Although there is no population over the regions where international ship NO_x is emitted, there is a substantial exposure to ozone attributed to ship NO_x. We notice is that while there is a steep decreasing slope after year 2013 in global mean surface ozone (Fig. 7e) attributed to East Asian NO_x emissions (Fig. 8c; due to emission reduction; shown in Fig. 3c), we do not find a similar feature for ozone exposure global population-weighted mean ozone attributed to East Asian NO_x emissions (Fig. 89c and Table S2). The reason for this is unclear at this point and could be an interesting topic for future studies.

3.4.2 Surface ozone attributed to reactive carbon emissions.

RC from Methane oxidation contributes ~12 ppbv (4750 %) to global mean surface ozone, biogenic and anthropogenic reactive-carbonNMRC emissions contribute ~4 ppbv (16 %) and 3.72 ppbv (~14.6742 %; (Table 4) respectively. Regional anthropogenic tags each contribute less than 5 % to the global mean surface ozone. Since the percentage contribution from stratospheric influx on global mean surface ozone is smaller than it's contribution to tropospheric ozone burden (as explained above), there is larger contribution from methane oxidation, anthropogenic and biomass burning RC emissions. We do not simulate any significant small trend with low certainty in the contribution to global mean surface ozone from total anthropogenic reactive-carbonNMRC emissions (Table 5). The increasing trend in global mean surface ozone is mainly driven by increasing contribution from stratosphere, RC from methane oxidation, and biogenic reactive-carbonNMRC emissions. Although there is no trend with high certainty in emissions of biogenic reactive-carbonNMRC, its increasing contribution to

Formatted: Subscript

~~both tropospheric ozone burden and~~ global mean surface ozone (Table 5) could be explained by the ~~significantly~~ increasing anthropogenic NO_x emissions (Butler et al., 2018, Lupascu et al., 2022).

~~There is a smaller contribution from biogenic reactive carbon emissions to global mean surface ozone (15.61 %) compared to its contribution to tropospheric ozone burden (16.25 %; Table 4). This could be because biogenic reactive carbon is mainly emitted from tropical regions where there is a strong convection leading to ozone molecules being lifted into the free troposphere.~~

The largest contribution to global population-weighted mean surface ozone is from ozone attributed to ~~RC from~~ methane oxidation (~12 ppbv (37.40 %); Table 4). ~~Methane~~ This Contribution to global population-weighted mean ozone is smaller compared to its contribution to global mean surface ozone (~47 %), ~~as because~~ a substantial part of this contribution to global mean surface ozone is at remote oceanic regions (not shown). Consequently, there is a larger relative contribution from biogenic (22 %) and anthropogenic (24%) ~~reactive carbon~~ NMRC sources to global population weighted mean ozone compared to their contribution to global mean surface ozone. Among regional anthropogenic NMRC emissions, there is a larger share of contributions from highly populated regions such as South Asian, East Asian and Rest of the World, and a smaller share from North American, European, and Russia-Belarus-Ukraine region's anthropogenic NMVOC-NMRC emissions to ~~ozone exposure~~ global population-weighted mean ozone compared to contribution to global mean surface ozone (Table 4).

The increasing trend ~~in~~ population-weighted surface ozone is mainly ~~contributed-driven~~ by increasing trends in the contribution of ~~RC from~~ methane oxidation (0.05 ppbv/yr (0.3841 %/yr)), anthropogenic (0.09 ppbv/yr (1.243 %/yr)) and biogenic (0.03 ppbv/yr (0.442 %/yr)) ~~reactive carbon~~ NMRC emissions (Table 5). McDuffie et al. (2023) show that with a ~100 ppb of methane pulse, the population-weighted ozone response is larger than the response in the global mean surface ozone. They further explain that this larger response was due to the larger availability of NO_x precursor emissions at populated regions leading to larger ozone production in populated regions. In contrast, we simulate a larger trend in ~~methane contribution~~ contribution from methane oxidation to global mean (0.425 %/year) than to population-weighted mean surface ozone (0.3841 %/year; Table 5), despite having ~~nearly the same~~ comparable change in prescribed methane concentration (~135 ppb; Fig. 4) over the 2000-2018 period. This is due to a dominant trend in contribution from anthropogenic NMVOC-NMRC emissions from populated regions in our simulation, which is lacking in the methane pulse simulated by McDuffie et al. (2023). This dominant trend in anthropogenic NMRC contribution to population-weighted surface ozone can also be seen in its significantly increasing relative contribution (Figure S84), which is not seen in other major contributors such as methane and biogenic NMVOCs-NMRCs.

The total NO_x emissions show a positive trend leading to increasing contribution from methane oxidation. Similarly, the increasing anthropogenic NO_x emitted at densely populated regions reacts with ~~reactive carbon~~ NMRC emitted from

anthropogenic and biogenic sources (Fig. 8b; Lupascu et al., 2022). Among regional anthropogenic ~~reactive carbon~~NMRC emissions, the ~~contribution from highly populated regions (Table S9); South Asia (~0.07 ppbv/yr (3.61 %/yr)) and East Asia (~0.06 ppbv/yr (2.85 %/yr); Table 4)~~ ~~contribution~~ to global population-weighted ozone shows the largest increasing trend (~~3.34 %/yr~~). ~~As for the NO_x-tagged case discussed in the previous subsection, the global population-weighted ozone attributed to East Asian anthropogenic NMRC emissions shows a small trend with low certainty during the post-2011 period in response to the reduction in emissions (Table S6).~~

4 Summary and Conclusions

In this study we quantify the contributions of ozone precursor emissions from various sources to tropospheric ozone during the 2000-2018 period. For this, we use the tagging approach using CAM4-Chem introduced in Butler et al., 2018. We perform separate simulations for NO_x and RC-tagged contributors to tropospheric ozone which allows us to quantify the absolute contribution from the tagged emission sources to the tropospheric ozone burden and calculate the ozone production efficiency per tagged sector/region.

We note a decreasing trend in both anthropogenic NO_x and reactive carbon ozone precursor emissions from North America, Europe, and Russia-Belarus-Ukraine region, and increasing trend from East Asia, South Asia, Middle East, international shipping, and Rest of the World. We simulate the largest contribution to tropospheric ozone burden from anthropogenic NO_x emissions (in our ~~NO_x~~NO_x-tagged simulation) and reactive carbon from methane oxidation (in our RC-tagged simulation), ~~both with significantly increasing trend, followed by contribution of ozone influx from stratosphere but with no significant trend in its contribution.~~ We simulate a relatively larger contribution to the tropospheric ozone burden from emissions at tropical regions compared to other regions, as previously discussed in Zhang et al., (2021). For example, the anthropogenic NO_x emissions from “rest of the world” regional tag that mainly consists of tropical and southern hemisphere regions, contributes more to the tropospheric ozone burden compared with East Asia despite smaller NO_x emissions. ~~We show using vertical profiles of tagged O₃ and NO_x mixing ratio fields (Fig. 6) that it is the emitted NO_x in the tropics which predominantly gets lofted into the free troposphere and eventually form ozone. Our tagging method only conveys information on the precursor source of the simulated ozone and associated gas molecules. It, however, does not convey information about where exactly the ozone molecule was formed: in the free troposphere, or within the boundary layer. While addressing this is beyond the scope of our study, further studies could employ ozone tagging method used in Sudo and Akimoto (2007) and Derwent et al., (2015) that tag/label ozone molecules based on the location at which the ozone molecules are formed. Our study highlights the dominant effect of equatorward shifting of O₃ precursor emissions which contributes to the tropospheric O₃ burden trend. Nevertheless, there remain other climatic factors that may play an amplifying or offsetting role to these contributions. These~~

Formatted: Subscript
Formatted: Subscript
Formatted: Subscript

Formatted: Subscript
Formatted: Subscript

include, for example, changes in the general circulation of the atmosphere and monsoonal changes under a warming planet, as well as the natural variability of climate.

The trend in each tagged sectors' contribution to the total tropospheric ozone burden is consistent with the sign of trend in the respective precursor emission from that tagged sector. We simulate an ~~significantly~~-increasing trend in OPE of NO_x emissions from regions where the emissions are decreasing (European, North American, and Russia-Belarus-Ukraine region's emissions) and vice versa (South Asian and Middle East). The anthropogenic ~~reactive-carbon~~NMRC becomes more efficient at producing ozone, following the increasing trend in anthropogenic NO_x emissions in their vicinity. We, however, do not simulate any ~~significant~~-trends with medium-high certainty in the OPE of other major reactive carbon sources such as methane oxidation, biogenic and biomass burning. We therefore recommend that further studies perform simulations combining perturbation and tagging to investigate how perturbation in the emissions of one sector induces changes in OPE of itself and other tagged sectors.

We contrast the contribution of emissions from tagged sectors to the global tropospheric ozone burden with their contribution to the global mean surface ozone and the population weighted mean surface ozone. Except for tropical source regions such as South Asia, we simulate a larger share of contributions from regional anthropogenic NO_x emissions to the global mean surface ozone compared to the tropospheric ozone burden. We also simulate an increasing trend in contribution ~~from by RC from~~ methane oxidation, and biogenic NMRC and anthropogenic reactive-carbon to global mean surface ozone and population-weighted mean ozone, leading to an overall global increasing trend in both these quantities. The increasing trend in population weighted ozone is mainly ~~contributed-driven~~ by increasing trend in ~~contribution from~~ anthropogenic NO_x and (to a lesser extent) ~~reactive-carbon~~NMRC emissions over highly populated regions: South Asia and East Asia. We also simulate a substantial contribution of NO_x emissions from international shipping to the population weighted mean ozone. While the focus of the present study has been the quantification of the influences on global ozone metrics, future work using the methods employed in this study could focus on quantification of the intra- and extra-regional contributions to surface ozone in the regions studied here.

In our study, we discuss the results related to the 2000-2018 trends in global annual ozone metrics and its precursor emissions. Further studies could investigate the tagged contributions to the trends in the spatial and seasonal distribution of ozone (e.g., Wang et al. 2022, Fiore et al., 2022), which may help us enhance our understanding of various processes responsible for these trends. While previous studies used the TOAST tagging to understand the trends in surface ozone over United States (e.g., Li et al., 2023a), free tropospheric ozone over Southeast Asia (e.g., Li et al., 2023b) and over East Asia in future scenarios (e.g., Hou et al., 2023), we recommend such studies be extended to other regions of interest. Comparison of smaller time periods (2000-2019, and 2019-2018) could also be considered to understand the contrasting effect of increasing and decreasing East Asian precursor emissions respectively. Finally, we recommend other global models implement the tropospheric ozone source

attribution with NO_x and RC tagging separately, which would enable a more rigorous model intercomparison and a better understanding of the underlying processes leading to the inter-model spread in their estimation of tropospheric ozone.

Acknowledgements

680 The authors would like to thank Mark Lawrence, Louisa Emmons, Claire Granier, Jukka-Pekka Jalkanen, Jan Eiof Jonson and Mariano Mertens for their helpful discussions during the performance of our simulations and preparation of this paper. Simulations were performed on the high-performance supercomputing cluster GLIC at the GFZ, Potsdam.

Financial support

This research has been supported by the Bundesministerium für Bildung und Forschung (grant no. 01US1701).

685 Data Availability

The values in the timeseries plotted in this study (Figures 3, 5, ~~6, 7, and 8~~, 7, 8, and 9) are provided in the Supplementary material. Please contact tim.butler@rifs-potsdam.de for availing the model output of our simulations.

Supplementary materials

Attached documents are: noxtagged_annual.csv, rctagged_annual.csv, Supplementary-figures-tables.docx

690 Author Contributions

AN designed the study and model simulations with support from AL and TB. TA has performed the simulations with support from AN, AL and TB. AN has written the paper with inputs from AL, TB and TA. TA supported with population-weighted ozone analysis.

695 **Competing Interests**

At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics

References

700 [Archibald, A., Neu, J., Elshorbany, Y., Cooper, O., Young, P., Akiyoshi, H., Cox, R., Coyle, M., Derwent, R., Deushi, M., et al.: Tropospheric Ozone Assessment Report: A critical review of changes in the tropospheric ozone burden and budget from 1850 to 2100, Elem Sci Anth, 8, 034, 2020.](#)

[Avnery, S., Mauzerall, D. L., Liu, J., and Horowitz, L. W.: Global crop yield reductions due to surface ozone exposure: 1. Year 2000 crop production losses and economic damage, Atmospheric Environment, 45, 2284–2296, 2011.](#)

705 [Bak, J., Song, E.-J., Lee, H.-J., Liu, X., Koo, J.-H., Kim, J., Jeon, W., Kim, J.-H., and Kim, C.-H.: Temporal variability of tropospheric ozone and ozone profiles in the Korean Peninsula during the East Asian summer monsoon: insights from multiple measurements and reanalysis datasets, Atmospheric Chemistry and Physics, 22, 14 177–14 187, 2022.](#)

[Butler, T. J., Vermeylen, F. M., Rury, M., Likens, G. E., Lee, B., Bowker, G. E., and McCluney, L.: Response of ozone and nitrate to stationary source NOx emission reductions in the eastern USA, Atmospheric Environment, 45, 1084–1094, 2011.](#)

710 [Butler, T., Lupascu, A., and Nalam, A.: Attribution of ground-level ozone to anthropogenic and natural sources of nitrogen oxides and reactive carbon in a global chemical transport model, Atmospheric Chemistry and Physics, 20, 10 707–10 731, 2020.](#)

[Butler, T., Lupascu, A., Coates, J., and Zhu, S.: TOAST 1.0: Tropospheric ozone attribution of sources with tagging for CESM 1.2. 2, Geoscientific Model Development, 11, 2825–2840, 2018.](#)

715 [Center for International Earth Science Information Network - CIESIN - Columbia University. 2018. Gridded Population of the World, Version 4 \(GPWv4\): Population Count. Palisades, NY: NASA Socioeconomic Data and Applications Center \(SEDAC\). <http://dx.doi.org/10.7927/H4X63JVC>](#)

[Chameides, W. and Walker, J. C.: A photochemical theory of tropospheric ozone, Journal of Geophysical Research, 78, 8751–8760, 1973.](#)

720 [Chang, K.-L., Schultz, M. G., Koren, G., and Selke, N.: Guidance note on best statistical practices for TOAR analyses, arXiv preprint, arXiv:2304.14236, 2023.](#)

- Coates, J. and Butler, T. M.: A comparison of chemical mechanisms using tagged ozone production potential (TOPP) analysis, *Atmospheric Chemistry and Physics*, 15, 8795–8808, 2015.
- Cooper, O. R., Parrish, D., Ziemke, J., Balashov, N., Cupeiro, M., Galbally, I., Gilge, S., Horowitz, L., Jensen, N., Lamarque, J.-F., et al.: Global distribution and trends of tropospheric ozone: An observation-based review, *Elementa*, 2, 000 029, 2014.
- 725 Crippa, M., Guizzardi, D., Butler, T., Keating, T., Wu, R., Kaminski, J., Kuenen, J., Kurokawa, J., Chatani, S., Morikawa, T., et al.: HTAP v3 emission mosaic: a global effort to tackle air quality issues by quantifying global anthropogenic air pollutant sources, *Earth System Science Data Discussions*, 2023, 1–34, 2023.
- Crutzen, P. J.: Photochemical reactions initiated by and influencing ozone in unpolluted tropospheric air, *Tellus*, 26, 47–57, 1974.
- 730 Dahlmann, K., Grewe, V., Ponater, M., and Matthes, S.: Quantifying the contributions of individual NO_x sources to the trend in ozone radiative forcing, *Atmospheric Environment*, 45, 2860–2868, 2011.
- DeLang, M. N., Becker, J. S., Chang, K.-L., Serre, M. L., Cooper, O. R., Schultz, M. G., Schroder, S., Lu, X., Zhang, L., Deushi, M., et al.: Mapping yearly fine resolution global surface ozone through the Bayesian maximum entropy data fusion of observations and model output for 1990–2017, *Environmental science & technology*, 55, 4389–4398, 2021.
- 735 Derwent, R. G., Utembe, S. R., Jenkin, M. E., and Shallcross, D. E.: Tropospheric ozone production regions and the intercontinental origins of surface ozone over Europe, *Atmospheric Environment*, 112, 216–224, 2015.
- Edwards, P. M. and Evans, M. J.: A new diagnostic for tropospheric ozone production, *Atmospheric Chemistry and Physics*, 17, 13 669–13 680, 2017.
- 740 Emmons, L. K., Schwantes, R. H., Orlando, J. J., Tyndall, G., Kinnison, D., Lamarque, J.-F., Marsh, D., Mills, M. J., Tilmes, S., Bardeen, C., et al.: The chemistry mechanism in the community earth system model version 2 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001 882, 2020.
- Emmons, L., Hess, P., Lamarque, J.-F., and Pfister, G.: Tagged ozone mechanism for MOZART-4, CAM-chem and other chemical transport models, *Geoscientific Model Development*, 5, 1531–1542, 2012.
- 745 Fiore, A. M., Dentener, F., Wild, O., Cuvelier, C., Schultz, M., Hess, P., Textor, C., Schulz, M., Doherty, R., Horowitz, L., et al.: Multimodel estimates of intercontinental source-receptor relationships for ozone pollution, *Journal of Geophysical Research: Atmospheres*, 114, 2009.

- 750 [Fiore, A. M., Hancock, S. E., Lamarque, J.-F., Correa, G. P., Chang, K.-L., Ru, M., Cooper, O., Gaudel, A., Polvani, L. M., Sauvage, B., et al.: Understanding recent tropospheric ozone trends in the context of large internal variability: a new perspective from chemistry-climate model ensembles, *Environmental Research: Climate*, 1, 025 008, 2022.](#)
- [Galmarini, S., Koffi, B., Solazzo, E., Keating, T., Hogrefe, C., Schulz, M., Benedictow, A., Griesfeller, J. J., Janssens-Maenhout, G., Carmichael, G., et al.: Coordination and harmonization of the multi-scale, multi-model activities HTAP2, AQMEII3, and MICS-Asia3: simulations, emission inventories, boundary conditions, and model output formats, *Atmospheric Chemistry and Physics*, 17, 1543–1555, 2017.](#)
- 755 [Gaudel, A., Cooper, O. R., Ancellet, G., Barret, B., Boynard, A., Burrows, J. P., Clerbaux, C., Coheur, P.-F., Cuesta, J., Cuevas, E., et al.: Tropospheric Ozone Assessment Report: Present-day distribution and trends of tropospheric ozone relevant to climate and global atmospheric chemistry model evaluation, *Elem Sci Anth*, 6, 39, 2018.](#)
- [Gaudel, A., Cooper, O. R., Chang, K.-L., Bourgeois, I., Ziemke, J. R., Strode, S. A., Oman, L. D., Sellitto, P., Nédélec, P., Blot, R., et al.: Aircraft observations since the 1990s reveal increases of tropospheric ozone at multiple locations across the Northern Hemisphere, *Science Advances*, 6, eaba8272, 2020.](#)
- 760 [Granier, C., Bessagnet, B., Bond, T., D'Angiola, A., Denier van der Gon, H., Frost, G. J., Heil, A., Kaiser, J. W., Kinne, S., Klimont, Z., et al.: Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980–2010 period, *Climatic change*, 109, 163–190, 2011.](#)
- [Grewe, V., Tsati, E., and Hoor, P.: On the attribution of contributions of atmospheric trace gases to emissions in atmospheric model applications, *Geoscientific Model Development*, 3, 487–499, 2010.](#)
- 765 [Grewe, V., Tsati, E., Mertens, M., Frömming, C., and Jöckel, P.: Contribution of emissions to concentrations: The TAGGING 1.0 submodel based on the Modular Earth Submodel System \(MESSy 2.52\), *Geoscientific Model Development*, 10, 2615–2633, 2017.](#)
- [Griffiths, P. T., Murray, L. T., Zeng, G., Shin, Y. M., Abraham, N. L., Archibald, A. T., Deushi, M., Emmons, L. K., Galbally, I. E., Hassler, B., et al.: Tropospheric ozone in CMIP6 simulations, *Atmospheric Chemistry and Physics*, 21, 4187–4218, 2021.](#)
- 770 [Heald, C. L. and Kroll, J.: The fuel of atmospheric chemistry: Toward a complete description of reactive organic carbon, *Science Advances*, 6, eaay8967, 2020.](#)

Hoor, P., Borken-Kleefeld, J., Caro, D., Dessens, O., Endresen, O., Gauss, M., Grewe, V., Hauglustaine, D., Isaksen, I. S., Jöckel, P., et al.: The impact of traffic emissions on atmospheric ozone and OH: results from QUANTIFY, *Atmospheric Chemistry and Physics*, 9, 3113–3136, 2009.

Hou, X., Wild, O., Zhu, B., and Lee, J.: Future tropospheric ozone budget and distribution over east Asia under a net-zero scenario, *Atmospheric Chemistry and Physics*, 23, 15 395–15 411, 2023.

Hussain, M. and Mahmud, I.: pyMannKendall: a python package for non-parametric Mann Kendall family of trend tests., *Journal of Open Source Software*, 4, 1556, <https://doi.org/10.21105/joss.01556>, 2019.

Jonson, J. E., Schulz, M., Emmons, L., Flemming, J., Henze, D., Sudo, K., Tronstad Lund, M., Lin, M., Benedictow, A., Koffi, B., et al.: The effects of intercontinental emission sources on European air pollution levels, *Atmospheric Chemistry and Physics*, 18, 13 655–13 672, 2018.

Junge, C. E.: Global ozone budget and exchange between stratosphere and troposphere, *Tellus*, 14, 363–377, 1962.

Kim, H. S., Kim, Y. H., Han, K. M., Kim, J., and Song, C. H.: Ozone production efficiency of a ship-plume: ITCT 2K2 case study, *Chemosphere*, 143, 17–23, 2016.

Kwok, R., Baker, K., Napelenok, S., and Tonnesen, G.: Photochemical grid model implementation and application of VOC, NO_x, and O₃ source apportionment, *Geoscientific Model Development*, 8, 99–114, 2015.

Lamarque, J.-F., Emmons, L., Hess, P., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C., Holland, E. A., Lauritzen, P., Neu, J., et al.: CAM-chem: Description and evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geoscientific Model Development*, 5, 369–411, 2012.

Lan, X., K.W. Thoning, and E.J. Dlugokencky: Trends in globally-averaged CH₄, N₂O, and SF₆ determined from NOAA Global Monitoring Laboratory measurements. Version 2024-01, <https://doi.org/10.15138/P8XG-AA10>, 2024.

Lawrence, M. G. and Crutzen, P. J.: Influence of NO_x emissions from ships on tropospheric photochemistry and climate, *Nature*, 402, 167–170, 1999.

Lawrence, M., Jöckel, P., and Von Kuhlmann, R.: What does the global mean OH concentration tell us?, *Atmospheric Chemistry and Physics*, 1, 37–49, 2001.

Lelieveld, J. a., Butler, T., Crowley, J., Dillon, T., Fischer, H., Ganzeveld, L., Harder, H., Lawrence, M., Martinez, M., Taraborrelli, D., et al.: Atmospheric oxidation capacity sustained by a tropical forest, *Nature*, 452, 737–740, 2008.

800 [Levy II, H.: Photochemistry of the lower troposphere, Planetary and Space Science, 20, 919–935, 1972.](#)

[Li, P., Yang, Y., Wang, H., Li, S., Li, K., Wang, P., Li, B., and Liao, H.: Source attribution of near-surface ozone trends in the United States during 1995–2019, Atmospheric Chemistry and Physics, 23, 5403–5417, 2023a.](#)

[Li, S., Yang, Y., Wang, H., Li, P., Li, K., Ren, L., Wang, P., Li, B., Mao, Y., and Liao, H.: Rapid increase in tropospheric ozone over Southeast Asia attributed to changes in precursor emission source regions and sectors, Atmospheric Environment, 304, 119 776, 2023b.](#)

805 [Lin, X., Trainer, M., and Liu, S.: On the nonlinearity of the tropospheric ozone production, Journal of Geophysical Research: Atmospheres, 93, 15 879–15 888, 1988.](#)

[Liu, Z., Doherty, R. M., Wild, O., O’connor, F. M., and Turnock, S. T.: Tropospheric ozone changes and ozone sensitivity from the present day to the future under shared socio-economic pathways, Atmospheric Chemistry and Physics, 22, 1209–1227, 2022.](#)

810 [Lupascu, A. and Butler, T.: Source attribution of European surface O₃ using a tagged O₃ mechanism, Atmospheric Chemistry and Physics, 19, 14 535–14 558, 2019.](#)

[Lupascu, A., Otero, N., Minkos, A., and Butler, T.: Attribution of surface ozone to NO_x and volatile organic compound sources during two different high ozone events, Atmospheric chemistry and physics, 22, 11 675–11 699, 2022.](#)

815 [Malashock, D. A., DeLang, M. N., Becker, J. S., Serre, M. L., West, J. J., Chang, K.-L., Cooper, O. R., and Anenberg, S. C.: Estimates of ozone concentrations and attributable mortality in urban, peri-urban and rural areas worldwide in 2019, Environmental Research Letters, 17, 054 023, 2022.](#)

[McDuffie, E. E., Sarofim, M. C., Raich, W., Jackson, M., Roman, H., Seltzer, K., Henderson, B. H., Shindell, D. T., Collins, M., Anderton, J., et al.: The Social Cost of Ozone-Related Mortality Impacts From Methane Emissions, Earth’s future, 11, e2023EF003 853, 2023.](#)

820 [Mertens, M., Grewe, V., Rieger, V. S., and Jöckel, P.: Revisiting the contribution of land transport and shipping emissions to tropospheric ozone, Atmospheric Chemistry and Physics, 18, 5567–5588, 2018.](#)

[Mertens, M., Kerkweg, A., Grewe, V., Jöckel, P., and Sausen, R.: Attributing ozone and its precursors to land transport emissions in Europe and Germany, Atmospheric Chemistry and Physics \(ACP\), pp. 7843–7873, 2020.](#)

825 [Mills, G., Pleijel, H., Malley, C. S., Sinha, B., Cooper, O. R., Schultz, M. G., Neufeld, H. S., Simpson, D., Sharps, K., Feng, Z., et al.: Tropospheric Ozone Assessment Report: Present-day tropospheric ozone distribution and trends relevant to vegetation, *Elem Sci Anth*, 6, 47, 2018.](#)

[Miyazaki, K., Bowman, K., Sekiya, T., Takigawa, M., Neu, J. L., Sudo, K., Osterman, G., and Eskes, H.: Global tropospheric ozone responses to reduced NO_x emissions linked to the COVID-19 worldwide lockdowns, *Science Advances*, 7, eabf7460, 2021.](#)

830 [Molod, A., Takacs, L., Suarez, M., and Bacmeister, J.: Development of the GEOS-5 atmospheric general circulation model: Evolution from MERRA to MERRA2, *Geoscientific Model Development*, 8, 1339–1356, 2015.](#)

[Monks, P. S., Archibald, A., Colette, A., Cooper, O., Coyle, M., Derwent, R., Fowler, D., Granier, C., Law, K. S., Mills, G., et al.: Tropospheric ozone and its precursors from the urban to the global scale from air quality to short-lived climate forcer, *Atmospheric chemistry and physics*, 15, 8889–8973, 2015.](#)

835 [Myhre, G., D. Shindell, F.-M. Bréon, W. Collins, J. Fuglestad, J. Huang, D. Koch, J.-F. Lamarque, D. Lee, B. Mendoza, T. Nakajima, A. Robock, G. Stephens, T. Takemura and H. Zhang: Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* \[Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley \(eds.\)\]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2021.](#)

[Prather, M. J.: Time scales in atmospheric chemistry: Theory, GWPs for CH₄ and CO, and runaway growth, *Geophysical Research Letters*, 23, 2597–2600, 1996.](#)

[Schneider, P. and Van Der A, R.: A global single-sensor analysis of 2002–2011 tropospheric nitrogen dioxide trends observed from space, *Journal of Geophysical Research: Atmospheres*, 117, 2012.](#)

845 [Schultz, M. G., Schröder, S., Lyapina, O., Cooper, O. R., Galbally, I., Petropavlovskikh, I., Von Schneidmesser, E., Tanimoto, H., Elshorbany, Y., Naja, M., et al.: Tropospheric Ozone Assessment Report: Database and metrics data of global surface ozone observations, *Elem Sci Anth*, 5, 58, 2017.](#)

[Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John Wiley & Sons, 2016.](#)

850

Sindelarova, K., Markova, J., Simpson, D., Huszar, P., Karlicky, J., Darras, S., Granier, C.: Copernicus Atmosphere Monitoring Service Global Biogenic VOC emissions version 3.0 (CAMS-GLOB-BIO v3.0), ECCAD, <https://doi.org/10.24380/xs64-gj42>, 2021.

855 Sudo, K. and Akimoto, H.: Global source attribution of tropospheric ozone: Long-range transport from various source regions, *Journal of Geophysical Research: Atmospheres*, 112, 2007.

Szopa, S., V. Naik, B. Adhikary, P. Artaxo, T. Bernsten, W.D. Collins, S. Fuzzi, L. Gallardo, A. Kiendler-Scharr, Z. Klimont, H. Liao, N. Unger, and P. Zanis, 2021: Short-Lived Climate Forcers. In *Climate Change: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 817–922, doi:10.1017/9781009157896.008, 2021.*

865 Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *Journal of geophysical research: atmospheres*, 106, 7183–7192, 2001.

Tilmes, S., Lamarque, J.-F., Emmons, L., Conley, A., Schultz, M., Saunio, M., Thouret, V., Thompson, A., Oltmans, S., Johnson, B., et al.: Ozonesonde climatology between 1995 and 2011: description, evaluation and applications, *Atmospheric Chemistry and Physics*, 12, 7475–7497, 2012.

870 Tilmes, S., Lamarque, J.-F., Emmons, L., Kinnison, D., Ma, P.-L., Liu, X., Ghan, S., Bardeen, C., Arnold, S., Deeter, M., et al.: Description and evaluation of tropospheric chemistry and aerosols in the Community Earth System Model (CESM1. 2), *Geoscientific Model Development*, 8, 1395–1426, 2015.

Van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R., Jin, Y. v., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmospheric chemistry and physics*, 10, 11 707–11 735, 2010.

875 Vinken, G. C., Boersma, K. F., Jacob, D. J., and Meijer, E. W.: Accounting for non-linear chemistry of ship plumes in the GEOS-Chem global chemistry transport model, *Atmospheric Chemistry and Physics*, 11, 11 707–11 722, 2011.

880 Voulgarakis, A., Naik, V., Lamarque, J.-F., Shindell, D. T., Young, P., Prather, M. J., Wild, O., Field, R., Bergmann, D., Cameron-Smith, P., et al.: Analysis of present day and future OH and methane lifetime in the ACCMIP simulations, Atmospheric Chemistry and Physics, 13, 2563–2587, 2013.

Wang, W., Parrish, D. D., Wang, S., Bao, F., Ni, R., Li, X., Yang, S., Wang, H., Cheng, Y., and Su, H.: Long-term trend of ozone pollution in China during 2014–2020: Distinct seasonal and spatial characteristics and ozone sensitivity, Atmospheric Chemistry and Physics, 22, 8935–8949, 2022.

885 West, J. J., Fiore, A. M., Naik, V., Horowitz, L. W., Schwarzkopf, M. D., and Mauzerall, D. L.: Ozone air quality and radiative forcing consequences of changes in ozone precursor emissions, Geophysical Research Letters, 34, 2007.

Young, P. J., Naik, V., Fiore, A. M., Gaudel, A., Guo, J., Lin, M., Neu, J., Parrish, D., Rieder, H., Schnell, J., et al.: Tropospheric Ozone Assessment Report: Assessment of global-scale model performance for global and regional ozone distributions, variability, and trends, Elem Sci Anth, 6, 10, 2018.

890 Young, P., Archibald, A., Bowman, K., Lamarque, J.-F., Naik, V., Stevenson, D., Tilmes, S., Voulgarakis, A., Wild, O., Bergmann, D., et al.: Pre-industrial to end 21st century projections of tropospheric ozone from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), Atmospheric Chemistry and Physics, 13, 2063–2090, 2013.

Zhang, Y., Cooper, O. R., Gaudel, A., Thompson, A. M., Nédélec, P., Ogino, S.-Y., and West, J. J.: Tropospheric ozone change from 1980 to 2010 dominated by equatorward redistribution of emissions, Nature Geoscience, 9, 875–879, 2016.

895 Zhang, Y., West, J. J., Emmons, L. K., Flemming, J., Jonson, J. E., Lund, M. T., Sekiya, T., Sudo, K., Gaudel, A., Chang, K.-L., et al.: Contributions of world regions to the global tropospheric ozone burden change from 1980 to 2010, Geophysical Research Letters, 48, e2020GL089 184, 2021.

Zhao, Y., Li, Y., Kumar, A., Ying, Q., Vandenberghe, F., and Kleeman, M. J.: Separately resolving NOx and VOC contributions to ozone formation, Atmospheric Environment, 285, 119 224, 2022.

900 Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., et al.: Trends in China’s anthropogenic emissions since 2010 as the consequence of clean air actions, Atmospheric Chemistry and Physics, 18, 14 095–14 111, 2018.
Archibald, A., Neu, J., Elshorbany, Y., Cooper, O., Young, P., Akiyoshi, H., Cox, R., Coyle, M., Derwent, R., Deushi, M., et al.: Tropospheric Ozone Assessment Report: A critical review of changes in the tropospheric ozone burden and budget from 1850 to 2100, Elem Sci Anth, 8, 034, 2020.

Formatted: Justified, Space Before: 12 pt, After: 0 pt, Line spacing: 1.5 lines, Pattern: Clear (White)

Avnery, S., Mauzerall, D. L., Liu, J., and Horowitz, L. W.: Global crop yield reductions due to surface ozone exposure: 1-
905 Year 2000 crop production losses and economic damage, *Atmospheric Environment*, 45, 2284–2296, 2011.

Butler, T., Lupaseu, A., Coates, J., and Zhu, S.: TOAST 1.0: Tropospheric ozone attribution of sources with tagging for CESM
1.2. 2, *Geoscientific Model Development*, 11, 2825–2840, 2018.

Butler, T., Lupaseu, A., and Nalam, A.: Attribution of ground-level ozone to anthropogenic and natural sources of nitrogen
oxides and reactive carbon in a global chemical transport model, *Atmospheric Chemistry and Physics*, 20, 10 707–10 731,
910 2020.

Butler, T. J., Vermeylen, F. M., Rury, M., Likens, G. E., Lee, B., Bowker, G. E., and McCluney, L.: Response of ozone and
nitrate to stationary source NO_x emission reductions in the eastern USA, *Atmospheric Environment*, 45, 1084–1094, 2011.

Chameides, W. and Walker, J. C.: A photochemical theory of tropospheric ozone, *Journal of Geophysical Research*, 78, 8751–
8760, 1973.

915 Coates, J. and Butler, T. M.: A comparison of chemical mechanisms using tagged ozone production potential (TOPP) analysis,
Atmospheric Chemistry and Physics, 15, 8795–8808, 2015.

Cooper, O. R., Parrish, D., Ziemke, J., Balashov, N., Cupeiro, M., Galbally, I., Gilge, S., Horowitz, L., Jensen, N., Lamarque,
J. F., et al.: Global distribution and trends of tropospheric ozone: An observation-based review, *Elementa*, 2, 000 029, 2014.

Crippa, M., Guizzardi, D., Butler, T., Keating, T., Wu, R., Kaminski, J., Kuenen, J., Kurokawa, J., Chatani, S., Morikawa, T.,
920 et al.: HTAP_v3 emission mosaic: a global effort to tackle air quality issues by quantifying global anthropogenic air pollutant
sources, *Earth System Science Data Discussions*, 2023, 1–34, 2023.

Center for International Earth Science Information Network – CIESIN – Columbia University. 2018. Gridded Population of
the World, Version 4 (GPWv4): Population Count. Palisades, NY: NASA Socioeconomic Data and Applications Center
(SEDAC). <http://dx.doi.org/10.7927/H4X63JVC>

925 Crutzen, P. J.: Photochemical reactions initiated by and influencing ozone in unpolluted tropospheric air, *Tellus*, 26, 47–57,
1974.

Dahlmann, K., Grewe, V., Ponater, M., and Matthes, S.: Quantifying the contributions of individual NO_x sources to the trend
in ozone radiative forcing, *Atmospheric Environment*, 45, 2860–2868, 2011.

- DeLang, M. N., Becker, J. S., Chang, K. L., Serre, M. L., Cooper, O. R., Schultz, M. G., Schröder, S., Lu, X., Zhang, L.,
930 Deushi, M., et al.: Mapping yearly fine resolution global surface ozone through the Bayesian maximum entropy data fusion of
observations and model output for 1990–2017, *Environmental science & technology*, 55, 4389–4398, 2021.
- Edwards, P. M. and Evans, M. J.: A new diagnostic for tropospheric ozone production, *Atmospheric Chemistry and Physics*,
17, 13 669–13 680, 2017.
- Emmons, L., Hess, P., Lamarque, J.-F., and Pfister, G.: Tagged ozone mechanism for MOZART-4, CAM-chem and other
935 chemical transport models, *Geoscientific Model Development*, 5, 1531–1542, 2012.
- Emmons, L. K., Schwantes, R. H., Orlando, J. J., Tyndall, G., Kinnison, D., Lamarque, J.-F., Marsh, D., Mills, M. J., Tilmes,
S., Bardeen, C., et al.: The chemistry mechanism in the community earth system model version 2 (CESM2), *Journal of
Advances in Modeling Earth Systems*, 12, e2019MS001 882, 2020.
- Fiore, A. M., Dentener, F., Wild, O., Cuvelier, C., Schultz, M., Hess, P., Textor, C., Schulz, M., Doherty, R., Horowitz, L., et
940 al.: Multimodel estimates of intercontinental source-receptor relationships for ozone pollution, *Journal of Geophysical
Research: Atmospheres*, 114, 2009.
- Fiore, A. M., Hancock, S. E., Lamarque, J.-F., Correa, G. P., Chang, K.-L., Ru, M., Cooper, O., Gaudel, A., Polvani, L. M.,
Sauvage, B., et al.: Understanding recent tropospheric ozone trends in the context of large internal variability: a new
perspective from chemistry-climate model ensembles, *Environmental Research: Climate*, 1, 025–008, 2022.
- 945 Galmarini, S., Koffi, B., Solazzo, E., Keating, T., Hogrefe, C., Schulz, M., Benedictow, A., Griesfeller, J. J., Janssens-
Maenhout, G., Carmichael, G., et al.: Coordination and harmonization of the multi-scale, multi-model activities HTAP2,
AQMEII3, and MICS-Asia3: simulations, emission-inventories, boundary-conditions, and model-output-formats, *Atmospheric
Chemistry and Physics*, 17, 1543–1555, 2017.
- Gaudel, A., Cooper, O. R., Ancellet, G., Barret, B., Boynard, A., Burrows, J. P., Clerbaux, C., Coheur, P. F., Cuesta, J., Cuevas,
950 E., et al.: Tropospheric Ozone Assessment Report: Present-day distribution and trends of tropospheric ozone relevant to climate
and global atmospheric chemistry model evaluation, *Elem Sci Anth*, 6, 39, 2018.
- Gaudel, A., Cooper, O. R., Chang, K. L., Bourgeois, I., Ziemke, J. R., Strode, S. A., Oman, L. D., Sellitto, P., Nédélec, P.,
Blot, R., et al.: Aircraft observations since the 1990s reveal increases of tropospheric ozone at multiple locations across the
Northern Hemisphere, *Science Advances*, 6, eaba8272, 2020.

- 955 Granier, C., Bessagnet, B., Bond, T., D'Angiola, A., Denier van der Gon, H., Frost, G. J., Heil, A., Kaiser, J. W., Kinne, S., Klimont, Z., et al.: Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980–2010 period, *Climatic change*, 109, 163–190, 2011.
- Grewe, V., Tsati, E., and Hoor, P.: On the attribution of contributions of atmospheric trace gases to emissions in atmospheric model applications, *Geoscientific Model Development*, 3, 487–499, 2010.
- 960 Grewe, V., Tsati, E., Mertens, M., Frömming, C., and Jöckel, P.: Contribution of emissions to concentrations: The TAGGING 1.0 submodel based on the Modular Earth Submodel System (MESSy 2.52), *Geoscientific Model Development*, 10, 2615–2633, 2017.
- Griffiths, P. T., Murray, L. T., Zeng, G., Shin, Y. M., Abraham, N. L., Archibald, A. T., Deushi, M., Emmons, L. K., Galbally, I. E., Hassler, B., et al.: Tropospheric ozone in CMIP6 simulations, *Atmospheric Chemistry and Physics*, 21, 4187–4218, 2021.
- 965 Heald, C. L. and Kroll, J.: The fuel of atmospheric chemistry: Toward a complete description of reactive organic carbon, *Science Advances*, 6, eaay8967, 2020.
- Hoor, P., Borken-Kleefeld, J., Caro, D., Dessens, O., Endresen, O., Gauss, M., Grewe, V., Hauglustaine, D., Isaksen, I. S., Jöckel, P., et al.: The impact of traffic emissions on atmospheric ozone and OH: results from QUANTIFY, *Atmospheric Chemistry and Physics*, 9, 3113–3136, 2009.
- 970 Hou, X., Wild, O., Zhu, B., and Lee, J.: Future tropospheric ozone budget and distribution over east Asia under a net-zero scenario, *Atmospheric Chemistry and Physics*, 23, 15 395–15 411, 2023.
- Hussain, M. and Mahmud, I.: pyMannKendall: a python package for non-parametric Mann-Kendall family of trend tests., *Journal of Open Source Software*, 4, 1556, <https://doi.org/10.21105/joss.01556>, 2019.
- Jonson, J. E., Schulz, M., Emmons, L., Flemming, J., Henze, D., Sudo, K., Tronstad Lund, M., Lin, M., Benedictow, A., Koffi, B., et al.: The effects of intercontinental emission sources on European air pollution levels, *Atmospheric Chemistry and Physics*, 18, 13 655–13 672, 2018.
- Junge, C. E.: Global ozone budget and exchange between stratosphere and troposphere, *Tellus*, 14, 363–377, 1962.
- Kim, H. S., Kim, Y. H., Han, K. M., Kim, J., and Song, C. H.: Ozone production efficiency of a ship plume: ITCT 2K2 case study, *Chemosphere*, 143, 17–23, 2016.
- 980 Kwok, R., Baker, K., Napelenok, S., and Tonnesen, G.: Photochemical grid model implementation and application of VOC, NO_x, and O₃ source apportionment, *Geoscientific Model Development*, 8, 99–114, 2015.

Lamarque, J. F., Emmons, L., Hess, P., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C., Holland, E. A., Lauritzen, P., Neu, J., et al.: CAM-chem: Description and evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geoscientific Model Development*, 5, 369–411, 2012.

985 Lan, X., K.W. Thoning, and E.J. Dlugokencky: Trends in globally-averaged CH₄, N₂O, and SF₆ determined from NOAA Global Monitoring Laboratory measurements. Version 2024-01, https://doi.org/10.15138/P8XG-AA10_2024.

Lawrence, M. G. and Crutzen, P. J.: Influence of NO_x emissions from ships on tropospheric photochemistry and climate, *Nature*, 402, 167–170, 1999.

990 Lelieveld, J. a., Butler, T., Crowley, J., Dillon, T., Fischer, H., Ganzeveld, L., Harder, H., Lawrence, M., Martinez, M., Taraborrelli, D., et al.: Atmospheric oxidation capacity sustained by a tropical forest, *Nature*, 452, 737–740, 2008.

Levy II, H.: Photochemistry of the lower troposphere, *Planetary and Space Science*, 20, 919–935, 1972.

Li, P., Yang, Y., Wang, H., Li, S., Li, K., Wang, P., Li, B., and Liao, H.: Source attribution of near-surface ozone trends in the United States during 1995–2019, *Atmospheric Chemistry and Physics*, 23, 5403–5417, 2023a.

995 Li, S., Yang, Y., Wang, H., Li, P., Li, K., Ren, L., Wang, P., Li, B., Mao, Y., and Liao, H.: Rapid increase in tropospheric ozone over Southeast Asia attributed to changes in precursor emission source regions and sectors, *Atmospheric Environment*, 304, 119–776, 2023b.

Lin, X., Trainer, M., and Liu, S.: On the nonlinearity of the tropospheric ozone production, *Journal of Geophysical Research: Atmospheres*, 93, 15 879–15 888, 1988.

1000 Lupascu, A. and Butler, T.: Source attribution of European surface O₃ using a tagged O₃ mechanism, *Atmospheric Chemistry and Physics*, 19, 14 535–14 558, 2019.

Lupascu, A., Otero, N., Minkos, A., and Butler, T.: Attribution of surface ozone to NO_x and volatile organic compound sources during two different high ozone events, *Atmospheric chemistry and physics*, 22, 11 675–11 699, 2022.

1005 Malashock, D. A., DeLang, M. N., Becker, J. S., Serre, M. L., West, J. J., Chang, K. L., Cooper, O. R., and Anenberg, S. C.: Estimates of ozone concentrations and attributable mortality in urban, peri-urban and rural areas worldwide in 2019, *Environmental Research Letters*, 17, 054 023, 2022.

McDuffie, E. E., Sarofim, M. C., Raich, W., Jackson, M., Roman, H., Seltzer, K., Henderson, B. H., Shindell, D. T., Collins, M., Anderton, J., et al.: The Social Cost of Ozone-Related Mortality Impacts From Methane Emissions, *Earth's future*, 11, e2023EF003853, 2023.

Mertens, M., Grewe, V., Rieger, V. S., and Jöckel, P.: Revisiting the contribution of land transport and shipping emissions to tropospheric ozone, *Atmospheric Chemistry and Physics*, 18, 5567–5588, 2018.

Mertens, M., Kerkweg, A., Grewe, V., Jöckel, P., and Sausen, R.: Attributing ozone and its precursors to land transport emissions in Europe and Germany, *Atmospheric Chemistry and Physics (ACP)*, pp. 7843–7873, 2020.

Mills, G., Pleijel, H., Malley, C. S., Sinha, B., Cooper, O. R., Schultz, M. G., Neufeld, H. S., Simpson, D., Sharps, K., Feng, Z., et al.: Tropospheric Ozone Assessment Report: Present day tropospheric ozone distribution and trends relevant to vegetation, *Elem Sci Anth*, 6, 47, 2018.

Miyazaki, K., Bowman, K., Sekiya, T., Takigawa, M., Neu, J. L., Sudo, K., Osterman, G., and Eskes, H.: Global tropospheric ozone responses to reduced NO_x emissions linked to the COVID-19 worldwide lockdowns, *Science Advances*, 7, eabf7460, 2021.

Molod, A., Takaacs, L., Suarez, M., and Baemeister, J.: Development of the GEOS-5 atmospheric general circulation model: Evolution from MERRA to MERRA2, *Geoscientific Model Development*, 8, 1339–1356, 2015.

Monks, P. S., Archibald, A., Colette, A., Cooper, O., Coyle, M., Derwent, R., Fowler, D., Granier, C., Law, K. S., Mills, G., et al.: Tropospheric ozone and its precursors from the urban to the global scale from air quality to short-lived climate forcer, *Atmospheric chemistry and physics*, 15, 8889–8973, 2015.

Myhre, G., D. Shindell, F.-M. Bréon, W. Collins, J. Fuglestad, J. Huang, D. Koch, J.-F. Lamarque, D. Lee, B. Mendoza, T. Nakajima, A. Robock, G. Stephens, T. Takemura and H. Zhang: Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G. K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2021.

Prather, M. J.: Time scales in atmospheric chemistry: Theory, GWPs for CH₄ and CO, and runaway growth, *Geophysical Research Letters*, 23, 2597–2600, 1996.

Schneider, P. and Van Der A, R.: A global single-sensor analysis of 2002–2011 tropospheric nitrogen-dioxide trends observed from space, *Journal of Geophysical Research: Atmospheres*, 117, 2012.

1035 Schultz, M. G., Schröder, S., Lyapina, O., Cooper, O. R., Galbally, I., Petropavlovskikh, I., Von Schneidmesser, E., Tanimoto, H., Elshorbany, Y., Naja, M., et al.: Tropospheric Ozone Assessment Report: Database and metrics data of global surface ozone observations, *Elem Sci Anth*, 5, 58, 2017.

Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John Wiley & Sons, 2016.

1040 Sindelarova, K., Markova, J., Simpson, D., Huszar, P., Karlicky, J., Darras, S., Granier, C.: Copernicus Atmosphere Monitoring Service Global Biogenic VOC emissions version 3.0 (CAMS GLOB-BIO-v3.0), ECCAD, <https://doi.org/10.24380/xs64-gj42>, 2021.

Szopa, S., V. Naik, B. Adhikary, P. Artaxo, T. Berntsen, W.D. Collins, S. Fuzzi, L. Gallardo, A. Kiendler-Scharr, Z. Klimont, H. Liao, N. Unger, and P. Zanis, 2021: Short Lived Climate Forcers. In *Climate Change: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 817–922, doi:10.1017/9781009157896.008, 2021.

1050 Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *Journal of geophysical research: atmospheres*, 106, 7183–7192, 2001.

Tilmes, S., Lamarque, J.-F., Emmons, L., Conley, A., Schultz, M., Saunio, M., Thouret, V., Thompson, A., Oltmans, S., Johnson, B., et al.: Ozone-sonde climatology between 1995 and 2011: description, evaluation and applications, *Atmospheric Chemistry and Physics*, 12, 7475–7497, 2012.

1055 Tilmes, S., Lamarque, J.-F., Emmons, L., Kinnison, D., Ma, P.-L., Liu, X., Ghan, S., Bardeen, C., Arnold, S., Deeter, M., et al.: Description and evaluation of tropospheric chemistry and aerosols in the Community Earth System Model (CESM1.2), *Geoscientific Model Development*, 8, 1395–1426, 2015.

Van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R., Jin, Y., v., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmospheric chemistry and physics*, 10, 11 707–11 735, 2010.

- 1060 Vinken, G. C., Boersma, K. F., Jacob, D. J., and Meijer, E. W.: Accounting for non-linear chemistry of ship plumes in the GEOS-Chem global chemistry transport model, *Atmospheric Chemistry and Physics*, 11, 11 707–11 722, 2011.
- Wang, W., Parrish, D. D., Wang, S., Bao, F., Ni, R., Li, X., Yang, S., Wang, H., Cheng, Y., and Su, H.: Long-term trend of ozone pollution in China during 2014–2020: Distinct seasonal and spatial characteristics and ozone sensitivity, *Atmospheric Chemistry and Physics*, 22, 8935–8949, 2022.
- 1065 West, J. J., Fiore, A. M., Naik, V., Horowitz, L. W., Schwarzkopf, M. D., and Mauzerall, D. L.: Ozone air quality and radiative forcing consequences of changes in ozone-precursor emissions, *Geophysical Research Letters*, 34, 2007.
- Young, P., Archibald, A., Bowman, K., Lamarque, J.-F., Naik, V., Stevenson, D., Tilmes, S., Voulgarakis, A., Wild, O., Bergmann, D., et al.: Pre-industrial to end-21st century projections of tropospheric ozone from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmospheric Chemistry and Physics*, 13, 2063–2090, 2013.
- 1070 Young, P. J., Naik, V., Fiore, A. M., Gaudel, A., Guo, J., Lin, M., Neu, J., Parrish, D., Rieder, H., Schnell, J., et al.: Tropospheric Ozone Assessment Report: Assessment of global-scale model performance for global and regional ozone distributions, variability, and trends, *Elem-Sci Anth*, 6, 10, 2018.
- Zhang, Y., Cooper, O. R., Gaudel, A., Thompson, A. M., Nédélec, P., Ogino, S.-Y., and West, J. J.: Tropospheric ozone change from 1980 to 2010 dominated by equatorward redistribution of emissions, *Nature Geoscience*, 9, 875–879, 2016.
- 1075 Zhang, Y., West, J. J., Emmons, L. K., Flemming, J., Jonson, J. E., Lund, M. T., Sekiya, T., Sudo, K., Gaudel, A., Chang, K.-L., et al.: Contributions of world regions to the global tropospheric ozone burden change from 1980 to 2010, *Geophysical Research Letters*, 48, e2020GL089 184, 2021.
- Zhao, Y., Li, Y., Kumar, A., Ying, Q., Vandenbergh, F., and Kleeman, M. J.: Separately resolving NO_x and VOC contributions to ozone formation, *Atmospheric Environment*, 285, 119 224, 2022.
- 1080 Zheng, B., Tong, D., Li, M., Liu, F., Hong, C., Geng, G., Li, H., Li, X., Peng, L., Qi, J., et al.: Trends in China's anthropogenic emissions since 2010 as the consequence of clean air actions, *Atmospheric Chemistry and Physics*, 18, 14 095–14 111, 2018.




Fig. 1: Comparing the time-series of monthly mean surface ozone (in ppb) simulated over the 2000-2018 period in this study with the gridded observation dataset from TOAR available until 2014 (<https://toar-data.fz-juelich.de/>), at various regions defined by HTAP2 (Galmarini et al. 2017). The comparison is performed for simulated surface ozone values only over those grid cells where the TOAR data is available. Also shown is the comparison with the CESM2-WACCM6 model which is a CMIP6 ensemble member (also until 2014).




Fig. 2: Taylor diagram of comparisons between modelled monthly mean ozone climatology (2000-2018) and ozone sonde climatology (1995-2010) data from Tilmes et al. 2012 in the high Tropics (bottom), mid-latitudes (middle), and high latitudes (top) for three different altitude levels (900 hPa, 500 hPa and 250 hPa) in the troposphere. The x-axis shows the relative ozone normalized bias of the simulations compared to the observations, whereas the radial in the y-axis describes the correlation coefficient of seasonal averaged ozone values between simulated and observed values. Numbers indicate different regions as difference in previous studies^{26,40}. Bottom panels: 1-NH Subtropics; 2-W-Pacific/E-India Ocean; 3-equat. America; 4-Atlantic/Africa; Middle panels: 1-Western Europe; 2-Eastern US; 3-Japan; 4-SH Mid Latitudes; Top panels: 1-NH Polar West; 2-NH Polar East; 3-Canada; 4-SH Polar.




Fig. 3: Global annual emissions of ozone precursors from various sources used in this study. Left panels: NO_x emissions (in TgN/yr), right panels: reactive carbon emissions (TgC/yr). Bottom panels are the regional contributions to the Total Anthropogenic emission (dark pink line) shown in the top panels. The symbols indicate the sign and certainty of trend in the plotted quantity. The timeseries of relative contributions (in %) from each of the emission sectors to the total emissions is provided in Supplementary Figure S3.




Fig. 4: a) Red line: Annually varying Methane mixing ratio (in ppm) prescribed in our model. Green line: Methane oxidation rate (in TgC/yr) simulated by our model. Magenta line: Lifetime of Methane (in years). b) Airmass weighted and CH₄-reaction weighted tropospheric OH concentration (in 10^5 molec/cm³). The symbols on the plotted timeseries indicate the sign of trend in the plotted quantity. The mean values, slope of trend, 95 % confidence interval and p-value for the 2000-2018 period are provided in Table S5.




Fig. 5: Simulated global annual mean tropospheric ozone burden (in Tg O₃) time-series over the 2000-2018 period. Shown are the total simulated ozone burden (black line) and the contributions from the NOx-tagged (left panels) and RC-tagged (right panels) simulations. Bottom panels are the regional contributions to the Total Anthropogenic component (dark pink line) shown in the top panels. The symbols indicate the sign and certainty of trend in the plotted quantity. The timeseries of relative contributions (in %) from each of the emission sectors to the total emissions is provided in Supplementary Figure S5.




Fig. 6: Vertical profiles of zonal climatological (2000-2018) mean mixing ratio within the troposphere: O₃ (top panels) and NO_x (bottom panels) attributed to Southeast Asian (left panels) and European (right panels) anthropogenic NO_x emissions.




Fig. 7: Time-series (2000-2018) of the Ozone production efficiency (OPE; in mol O₃/mol N: NOx-tagged left panels, mol O₃/mol C: RC-tagged right panels) of various ozone precursor emission regions/sectors tagged in this study. Bottom panels are the OPE values of regional anthropogenic emissions tagged in this study. The symbols indicate the sign and certainty of trend in the plotted quantity.




Fig. 8: Same as Figure 5, but for global annual area-weighted mean surface ozone (in ppb). The timeseries of relative contributions (in %) from each of the tagged components to the global annual area-weighted mean surface ozone is provided in Supplementary Figure S7.




Fig. 9: Same as Figure 5, but for global annual population-weighted mean surface ozone (in ppb). The timeseries of relative contributions (in %) from each of the tagged components to the global annual population-weighted mean surface ozone is provided in Supplementary Figure S8.




Fig. 10: Spatial distribution of climatological (2000-2018) annual mean O_3 attributed to NO_x emissions from international shipping (in ppbv): a) Surface ozone and b) Population weighted surface ozone.

<u>Tag name</u>	<u>NOx-tagged</u>	<u>RC-tagged</u>
<u>HTAP Tier 1 regions</u>		
<u>International Shipping</u>	<u>Explicit</u>	<u>Explicit</u>
<u>North America</u>	<u>Explicit</u>	<u>Explicit</u>
<u>Europe</u>	<u>Explicit</u>	<u>Explicit</u>
<u>East Asia</u>	<u>Explicit</u>	<u>Explicit</u>
<u>South Asia</u>	<u>Explicit</u>	<u>Explicit</u>
<u>Russia, Belarus, Ukraine</u>	<u>Explicit</u>	<u>Explicit</u>
<u>Middle East</u>	<u>Explicit</u>	<u>Explicit</u>
<u>Southeast Asia</u>	<u>Explicit^S->RoW</u>	<u>RoW</u>
<u>Northern Africa</u>	<u>Explicit^S->RoW</u>	<u>RoW</u>
<u>Mexico and Central America</u>	<u>Explicit^S->RoW</u>	<u>RoW</u>
<u>Central Asia</u>	<u>Explicit^S->RoW</u>	<u>RoW</u>
<u>Pacific, Australia, and New Zealand</u>	<u>Southern Hemisphere Regions^S ->RoW</u>	<u>RoW</u>
<u>Southern Africa</u>	<u>Southern Hemisphere Regions^S ->RoW</u>	<u>RoW</u>
<u>South America</u>	<u>Southern Hemisphere Regions^S ->RoW</u>	<u>RoW</u>
<u>Antarctica</u>	<u>Southern Hemisphere Regions^S ->RoW</u>	<u>RoW</u>
<u>Arctic</u>	<u>Shipping</u>	<u>Shipping</u>
<u>Other tags</u>		
<u>Stratosphere[*]</u>	<u>Global</u>	<u>Global</u>
<u>Aircraft</u>	<u>Global</u>	<u>Global</u>
<u>Biogenic</u>	<u>Global</u>	<u>Global</u>
<u>Biomass Burning</u>	<u>Global</u>	<u>Global</u>
<u>NOx from Lightning</u>	<u>Global</u>	<u>N/A</u>
<u>RC from Methane Oxidation</u>	<u>N/A</u>	<u>Global</u>
<u>Extra Production</u>	<u>Global</u>	<u>Global</u>

^SAlthough explicitly tagged in the NOx-tagged simulation, these regions have been lumped into “Rest of the world” tag (RoW) used in this paper.

^{*}NOx-tagged simulation attributes ozone to influx from the stratosphere (same as RC-tagged), and also to NO produced from oxidation of N₂O.

Table 1: List of tags used in this study. See Fig. S1 for a map of tagged regions considered in this study.

<u>Source</u>	<u>Emissions</u> <u>(TgN/yr)</u>	<u>Tropospheric</u> <u>O₃ burden</u> <u>(TgO₃)</u>	<u>OPE</u> <u>(mol O₃</u> <u>/mol N)</u>	<u>Surface</u> <u>mean</u> <u>(ppbv)</u>	<u>Population</u> <u>weighted mean</u> <u>(ppbv)</u>
Total	49.12	317.93	N/A	25.39	33.01
Stratosphere	N/A	91.50 (28.78)	N/A	4.99 (19.65)	3.45 (10.44)
Aircraft	0.93 (1.90)	13.44 (4.23)	4.19	0.99 (3.91)	1.18 (3.59)
Biogenic	4.94 (10.05)	23.67 (7.44)	1.40	2.29 (9.01)	3.43 (10.40)
Biomass Burning	4.18 (8.52)	12.68 (3.99)	0.88	1.32 (5.20)	1.59 (4.82)
Lightning	3.47 (7.06)	70.51 (22.18)	5.93	3.51 (13.82)	2.86 (8.68)
Extra Production	N/A	0.97 (0.31)	N/A	0.08 (0.33)	0.11 (0.33)
Anthropogenic	35.60 (72.47)	116.94 (36.78)	0.96	12.20 (48.07)	20.38 (61.74)
<u>Regional contribution to Anthropogenic component</u>					
International Shipping	4.04 (8.22)	23.02 (7.24)	1.67	2.97 (11.69)	2.47 (7.47)
North America	5.51 (11.22)	9.79 (3.08)	0.54	1.43 (5.64)	1.61 (4.87)
Europe	3.54 (7.20)	4.36 (1.37)	0.36	0.84 (3.32)	1.47 (4.46)
East Asia	9.05 (18.43)	16.62 (5.23)	0.54	1.58 (6.23)	4.27 (12.92)
South Asia	2.91 (5.92)	7.64 (2.40)	0.78	0.62 (2.44)	4.60 (13.93)
Russia, Belarus, Ukraine	1.67 (3.39)	2.30 (0.72)	0.40	0.51 (2.01)	0.53 (1.62)
Middle East	1.68 (3.42)	3.64 (1.14)	0.64	0.49 (1.95)	1.15 (3.50)
Rest of the World	7.20 (14.66)	44.20 (13.90)	1.80	3.76 (14.80)	4.28 (12.97)

Table 2: Attribution of O₃ to NOx-tagged precursor emissions. 2000-2018 Mean contribution from NOx-tagged components to various metrics: Precursor emissions, tropospheric ozone burden, Ozone production efficiency, grid cell area weighted, and population weighted global mean surface ozone. In brackets is the percentage contribution to the total, wherever applicable.

<u>Source</u>	<u>Emissions</u> <u>(TgN/yr⁻²)</u>	<u>Tropospheric O₃</u> <u>burden (TgO₃/vr)</u>	<u>OPE</u> <u>((mol O₃</u> <u>/mol N)/yr)</u>	<u>Surface mean</u> <u>(ppbv/yr)</u>	<u>Population</u> <u>weighted mean</u> <u>(ppbv/yr)</u>
Total	*0.13(0.29)	0.72(0.23)	N/A	0.08(0.30)	0.17(0.54)
Stratosphere	N/A	*1.83e-03 (2.06e-03)	N/A	0.02(0.32)	*-8.59e-03(-0.25)
Aircraft	0.02(2.52)	0.28(2.41)	*2.74e-03	0.02(2.18)	0.02(1.70)
Biogenic ¹	**0 (0)	-0.06(-0.26)	-3.74E-03	-3.86e-03(-0.16)	-0.02(-0.66)
Biomass Burning	-0.04(-0.91)	-0.20(-1.38)	-5.47E-03	-0.02(-1.14)	-0.02(-1.12)
Lightning	-0.01(-0.35)	-0.43(-0.56)	-0.01	-0.02(-0.44)	-0.03(-0.82)
Extra Production	N/A	6.61e-03(0.74)	N/A	3.62e-04(0.46)	7.16e-04(0.71)
Anthropogenic	0.19(0.59)	1.35(1.29)	7.35E-03	0.08(0.69)	0.22(1.19)
<u>Regional contribution to Anthropogenic component</u>					
International Shipping	0.08(2.47)	0.31(1.48)	-9.80E-03	0.05(1.92)	0.04(1.60)
North America	-0.33(-3.82)	-0.34(-2.59)	0.01	-0.04(-2.35)	-0.05(-2.24)
Europe	-0.11(-2.52)	-0.11(-2.12)	2.62E-03	-0.02(-1.97)	-0.03(-1.59)
East Asia	0.25(4.04)	0.40(3.29)	*2.27e-04	0.04(2.87)	0.06(1.74)
South Asia	0.13(7.41)	0.24(4.58)	-0.01	0.02(3.97)	0.11(3.09)
Russia, Belarus, Ukraine	-0.02(-0.87)	-0.02(-0.62)	1.34E-03	-2.64e-03 (-0.50)	-3.59e-03(-0.63)
Middle East	0.06(4.79)	0.07(2.26)	-0.01	6.97e-03(1.62)	0.01(1.12)
Rest of the World	0.15(2.58)	0.73(1.87)	-8.69E-03	0.05(1.35)	0.08(2.09)

*0.5<p<0.1 (medium certainty)

**p>0.1 (low certainty)

¹Since we prescribe an annually repeating seasonal cycle of biogenic/soil NO_x emissions, the annual mean value is held constant leading to zero slope and p-value=1

Table 3: Theil-Sen estimator/slope of trends in contributions of NO_x-tagged metrics, wherever applicable. Trend slope and p-value estimation has been summarized in section 2.3. The 95 % confidence interval and p-value for each of these estimated trends is provided in Table S1. In brackets is the trend in %/year. Note that this trend in %/year is the trend slope relative to the first value of the absolute contribution (for year 2000), and not to be confused with the relative contribution to the total as shown in Supplementary Figures S3, S5, S7 and S8.

<u>Source</u>	<u>Emissions</u> <u>(TgC/yr)</u>	<u>Tropospheric</u> <u>O₃ burden</u> <u>(TgO₃)</u>	<u>OPE</u> <u>(mol O₃</u> <u>/mol C)</u>	<u>Surface</u> <u>mean (ppbv)</u>	<u>Population</u> <u>weighted mean</u> <u>(ppbv)</u>
Total	1342.06	317.92	N/A	25.39	33.01
Stratosphere	N/A	76.04 (23.92)	N/A	3.99 (15.73)	3.05 (9.25)
Aircraft	0.24 (0.02)	0.03 (8.58e-03)	0.03	2.33e-03 (9.19e-03)	2.97e-03 (8.98e-03)
Biogenic	430.93 (32.11)	52.95 (16.66)	0.03	4.07 (16.04)	7.38 (22.37)
Biomass Burning	171.39 (12.77)	13.85 (4.36)	0.02	1.17 (4.62)	1.34 (4.06)
Methane oxidation	437.98 (32.63)	132.45 (41.66)	0.08	11.99 (47.22)	12.36 (37.46)
Extra Production	N/A	4.52 (1.42)	N/A	0.42 (1.67)	0.62 (1.88)
Anthropogenic	301.52 (22.47)	45.95 (14.45)	0.04	3.72 (14.67)	8.24 (24.96)
<u>Regional contribution to Anthropogenic component</u>					
International Shipping	0.76 (0.06)	0.17 (0.05)	0.06	0.02 (0.07)	0.02 (0.07)
North America	36.47 (2.72)	4.03 (1.27)	0.03	0.57 (2.24)	0.68 (2.07)
Europe	21.19 (1.58)	2.40 (0.75)	0.03	0.43 (1.69)	0.66 (1.99)
East Asia	95.36 (7.11)	10.87 (3.42)	0.03	1.03 (4.06)	2.52 (7.63)
South Asia	41.89 (3.12)	5.55 (1.74)	0.03	0.40 (1.59)	2.36 (7.15)
Russia, Belarus, Ukraine	7.97 (0.59)	0.92 (0.29)	0.03	0.17 (0.67)	0.21 (0.63)
Middle East	7.12 (0.53)	1.22 (0.38)	0.04	0.17 (0.68)	0.36 (1.10)
Rest of the World	90.77 (6.76)	15.65 (4.92)	0.04	0.93 (3.66)	1.43 (4.33)

165 **Table 4:** Same as Table 2 but for RC-tagged components

<u>Source</u>	<u>Emissions (TgC/yr²)</u>	<u>Tropospheric O₃ burden (TgO₃/yr)</u>	<u>OPE ((mol O₃ /mol C)/yr)</u>	<u>Surface mean (ppbv/yr)</u>	<u>Population weighted mean (ppbv/yr)</u>
Total	3.18(0.25)	0.71(0.23)	N/A	0.08(0.30)	0.17(0.54)
Stratosphere	N/A	**0.03(0.04)	N/A	0.02(0.40)	** -4.2e-03(-0.14)
Aircraft	3.9e-03(1.74)	4.66e-04(1.81)	**4.05e-05	3.60e-05(1.62)	4.65e-05(1.64)
Biogenic	*1.26(0.31)	**0.06(0.11)	** -1.38e-05	9.53e-03(0.24)	0.03(0.44)
Biomass Burning	** -0.77(-0.51)	** -0.08(-0.63)	** -2.89e-05	** -1.58e-03(-0.15)	** -1.72e-03(-0.14)
Methane oxidation	1.81(0.42)	0.44(0.33)	** -7.00e-05	0.05(0.42)	0.05(0.38)
Extra Production	N/A	0.02(0.54)	N/A	1.92e-03(0.48)	4.33e-03(0.76)
Anthropogenic	1.74(0.61)	0.45(1.03)	2.17e-04	**4.6e-03(0.12)	0.09(1.23)
Regional contribution to Anthropogenic component					
International Shipping	0.01(1.79)	-2.73e-03(-1.03)	-1.38e-03	-1.91e-04(-0.73)	** -7.06e-05(-0.22)
North America	-1.80(-3.16)	-0.17(-2.83)	1.58e-04	-0.02(-2.71)	-0.03(-2.58)
Europe	-0.72(-2.53)	-0.08(-2.53)	* -4.09e-05	-0.01(-2.46)	-0.02(-2.26)
East Asia	1.58(2.20)	0.23(2.77)	2.30e-04	0.02(2.91)	0.06(2.85)
South Asia	1.19(3.76)	0.15(3.51)	** -3.11e-05	0.01(3.54)	0.07(3.61)
Russia, Belarus, Ukraine	-0.06(-0.72)	-0.01(-0.98)	-1.03E-04	-1.3e-03(-0.71)	-1.87e-03(-0.82)
Middle East	* -0.05(-0.60)	-0.01(-1.01)	-1.42e-04	* -1.3e-03(-0.70)	* -2.62e-03(-0.65)
Rest of the World	1.50(1.95)	0.26(1.87)	**2.59e-05	0.01(1.62)	0.03(2.31)

*0.5<p<0.1 (medium certainty)

**p>0.1 (low certainty)

Table 5: Same as Table 3 but for RC-tagged metrics. The 95 % confidence interval and p-value for each of these estimated trends is provided in Table S3.




Figure 1: Comparing the time-series of monthly mean surface ozone (in ppb) simulated over the 2000–2018 period in this study with the gridded observation dataset from TOAR available until 2014 (<https://toar-data.fz-juelich.de/>), at various regions defined by HTAP2 (Galmarini et al. 2017). The comparison is performed for simulated surface-ozone values only over those grid-cells where the TOAR data is available. Also shown is the comparison with the CESM2-WACCM6 model which is a CMIP6 ensemble member (also until 2014).




Figure 2: Taylor diagram of comparisons between modelled monthly mean ozone climatology (2000-2018) and ozone sonde climatology (1995-2010) data from Tilmes et al. 2012 in the high Tropics (left), mid-latitudes (middle), and high latitudes (right) for three different altitude levels (900 hPa, 500 hPa and 250 hPa) in the troposphere. The x-axis shows the relative ozone normalized bias of the simulations compared to the observations, whereas the radius in the y-axis describes the correlation coefficient of seasonal averaged ozone values between simulated and observed values. Numbers indicate different regions as difference in previous studies^{26,40}. Left panels: 1-NH Subtropics; 2-W-Pacific/E-India Ocean; 3-equat. America; 4-Atlantic/Africa; Middle panels: 1-Western Europe; 2-Eastern US; 3-Japan; 4-SH Mid Latitudes; Right panels: 1-NH Polar West; 2-NH Polar East; 3-Canada; 4-SH Polar.




Figure 3: Global annual emissions of ozone precursors from various sources used in this study. Left panels: NO_x emissions (in TgN/yr), right panels: reactive carbon emissions (TgC/yr). Bottom panels are the regional contributions to the Total Anthropogenic emission (dark pink line) shown in the top panels. The symbols show the sign of trend in the plotted quantity. Trend significance has been estimated by an original Mann-Kendall test at 5 % significance level using the pymannkendall python module described in Hussain and Mahmud (2019). The timeseries of relative contributions (in %) from each of the emission sectors to the total emissions is provided in Supplementary Figure S1.




Figure 4: Black line: Annually varying Methane concentration (in ppm) prescribed in our model. Green line: Methane oxidation rate (in TgC/yr) simulated by our model. Magenta line: Lifetime of Methane (in years). The symbols show the sign of trend in the plotted quantity. Trend significance has been estimated by an original Mann-Kendall test at 5 % significance level using the pymannkendall python module described in Hussain and Mahmud (2019).




Figure 5: Simulated global Annual-mean tropospheric-ozone burden (in Tg O₃) time-series over the 2000-2018 period. Shown are the total simulated ozone burden (black line) and the contributions from tagged components from the NOx-tagged (left panels) and RC-tagged (right panels) simulations. Bottom panels are the regional contributions to the Total Anthropogenic component (dark pink line) shown in the top panels. The symbols show the sign of trend in the plotted quantity. Trend significance has been estimated by an original Mann-Kendall test at 5% significance level using the pymannkendall python module described in Hussain and Mahmud (2019). The timeseries of relative contributions (in %) from each of the tagged components to the total tropospheric ozone burden is provided in Supplementary-Figure S2.




Figure 6: Time-series (2000-2018) of annual-mean Ozone-production efficiency (OPE; in mol O₃/mol N: NOx-tagged left panels, mol O₃/mol C: RC-tagged right panels) of various ozone-precursor emission regions/sectors tagged in this study. Bottom panels are the OPE values of regional anthropogenic emissions tagged in this study. The symbols show the sign of trend in the plotted quantity. Trend significance has been estimated by an original Mann-Kendall test at 5 % significance level using the pymannkendall python module described in Hussain and Mahmud (2019).




Figure 7: Same as Figure 5, but for global annual area-weighted mean surface ozone (in ppbv). The timeseries of relative contributions (in %) from each of the tagged components to the global annual area-weighted mean surface ozone is provided in Supplementary Figure S3.




Figure 8: Same as Figure 5, but for global annual population-weighted mean surface ozone (ozone exposure; in ppb). The timeseries of relative contributions (in %) from each of the tagged components to the global annual population-weighted mean surface ozone is provided in Supplementary Figure S4.

Tag name	NOx-tagged	VOC-tagged
HTAP-Tier 1 regions		
International Shipping	Explicit	Explicit
North America	Explicit	Explicit
Europe	Explicit	Explicit
East Asia	Explicit	Explicit
South Asia	Explicit	Explicit
Russia, Belarus, Ukraine	Explicit	Explicit
Middle East	Explicit	Explicit
Southeast Asia	Explicit ^b ->RoW	RoW
Northern Africa	Explicit ^b ->RoW	RoW
Mexico and Central America	Explicit ^b ->RoW	RoW
Central Asia	Explicit ^b ->RoW	RoW
Pacific, Australia, and New Zealand	RoW	RoW
Southern Africa	RoW	RoW
South America	RoW	RoW
Antarctica	RoW	RoW
Arctic ^a	Ocean-based->Shipping Land-Based->RoW	Ocean-based->Shipping Land-Based->RoW
Other tags		
Stratosphere	Global	Global
Aircraft	Global	Global
Biogenic	Global	Global
Biomass Burning	Global	Global
Lightning	Global	N/A
Methane	N/A	Global
Extra-Production	Global	Global

^aAlthough explicitly tagged in the NOx-tagged simulation, these regions have been lumped into “Rest of the world” tag (RoW) used in this paper. Refer to the Supplementary material Tables S2 and S3 for explicit results related to these tags.

^bShip emissions are included in the Shipping tag, and land-based anthropogenic emissions included in “Rest of the World” tag.

Formatted Table

Formatted Table

Formatted Table

Anthropogenic emissions here only refer to surface-based anthropogenic emissions.

Table 1: List of tags used in this study.

230

Source	Emissions (TgN/yr)	Tropospheric O ₃ burden (TgO ₃)	OPE (mol O ₃ /mol N)	Surface mean (ppbv)	Population weighted mean (ppbv)
Total	67.0	336.4	N/A	27.13	33.6
Stratosphere	N/A	87.58(26.03)	N/A	4.7(17.32)	3.0(8.93)
Aircraft	1.43(2.14)	17.88(5.32)	3.64	1.27(4.68)	1.42(4.22)
Biogenic	4.94(7.37)	20.85(6.2)	1.23	1.98(7.29)	2.76(8.21)
Biomass Burning	4.18(6.25)	11.76(3.5)	0.82	1.21(4.47)	1.43(4.25)
Lightning	3.47(5.18)	64.94(19.31)	5.46	3.15(11.61)	2.41(7.18)
Extra Production	N/A	0.91(0.27)	N/A	0.08(0.29)	0.1(0.28)
Anthropogenic	52.21(77.93)	132.47(39.38)	0.74	14.74(54.34)	22.49(66.92)
Regional contribution to Anthropogenic component					
International Shipping	5.92(8.83)	26.82(7.97)	1.33	3.75(13.81)	2.82(8.38)
North America	8.09(12.07)	11.67(3.47)	0.44	1.62(5.99)	1.7(5.06)
Europe	5.19(7.75)	5.09(1.51)	0.29	0.94(3.45)	1.56(4.64)
East Asia	13.27(19.81)	20.21(6.01)	0.19	1.8(6.63)	4.15(12.34)
South Asia	4.27(6.37)	9.45(2.81)	0.66	0.73(2.7)	5.21(15.5)
Russia, Belarus, Ukraine	2.44(3.65)	2.72(0.81)	0.33	0.59(2.16)	0.58(1.71)
Middle East	2.47(3.68)	4.5(1.34)	0.54	0.57(2.11)	1.3(3.88)
Rest of the World	10.56(15.76)	52.01(15.46)	1.44	4.74(17.49)	5.18(15.4)

Table 2: Attribution of O₃ to NOx-tagged precursor emissions. 2000–2018 Mean contribution from NOx-tagged components to various metrics: Precursor emissions, tropospheric ozone burden, Ozone production efficiency, Area-weighted, and Population-weighted mean surface ozone. In brackets is the percentage contribution to the total wherever applicable.

Source	Emissions (TgN/yr ²)	Tropospheric O ₃ burden (TgO ₃ /yr)	OPE (mol-O ₃ /mol N)/yr)	Surface mean (ppbv/yr)	Population weighted mean (ppbv/yr)
Total	0.24(0.38)	0.91(0.28)	N/A	0.1(0.37)	0.16(0.48)
-Stratosphere	N/A	Insig	N/A	0.01(0.32)	Insig
Aircraft	0.03(2.52)	0.35(2.29)	Insig	0.02(2.22)	0.02(1.5)
Biogenic	Insig	-0.08(-0.35)	-0.004	-0.008(-0.23)	-0.03(-0.82)
Biomass Burning	-0.04(-0.91)	-0.2(-1.48)	-0.006	-0.02(-1.21)	-0.02(-1.22)
Lightning	-0.01(-0.35)	-0.47(-0.65)	-0.018	-0.02(-0.52)	-0.03(-0.97)
Extra Production	N/A	0.01(0.66)	N/A	0.0003(0.43)	0.0005(0.55)
Anthropogenic	0.29(0.59)	1.31(1.09)	0.004	0.1(0.7)	0.21(0.99)
Regional contribution to Anthropogenic component					
International Shipping	0.12(2.47)	0.33(1.37)	-0.01	0.06(1.81)	0.04(1.42)
North America	-0.48(-3.82)	-0.41(-2.6)	0.01	-0.05(-2.33)	-0.05(-2.15)
Europe	-0.16(-2.52)	-0.13(-2.11)	0.002	-0.02(-1.96)	-0.03(-1.48)
East Asia	0.36(4.04)	0.48(3.18)	Insig	0.04(2.8)	0.05(1.42)
South Asia	0.2(7.41)	0.29(4.28)	-0.011	0.02(3.66)	0.1(2.46)
Russia, Belarus, Ukraine	-0.02(-0.87)	-0.02(-0.63)	0.001	-0.0031(-0.51)	-0.0034(-0.56)
Middle East	0.09(4.79)	0.09(2.23)	-0.01	0.01(1.52)	0.01(0.92)
Rest of the World	0.22(2.58)	0.79(1.73)	-0.008	0.05(1.26)	0.08(1.87)

Table 3: Theil-Sen estimator/slope of trends in contributions of NOx tagged metrics, wherever applicable and significant. Trend slope and significance has been estimated by an original Mann-Kendall test at 5 %-significance level using the pymannkendall python module described in Hussain and Mahmud (2019). In brackets is the trend in %/year. Note that this

trend in %/year is the trend slope relative to the first value of the absolute contribution (for year 2000), and not to be confused with the relative contribution to the total as shown in Supplementary Figures S1–S4.

Source	Emissions (TgC/yr)	Tropospheri O ₃ burden (TgO ₃)	OPE (mol-O ₃ /mol-C)	Surface-mean (ppbv)	Population weighted-mean (ppbv)
Total	1407.07	336.4	N/A	27.13	33.6
Stratosphere	N/A	74.48(22.14)	N/A	3.85(14.21)	2.74(8.16)
Aircraft	0.24(0.02)	0.03(0.01)	0.0294	0.0025(0.01)	0.0029(0.01)
Biogenic	430.93(30.63)	54.67(16.25)	0.0317	4.23(15.61)	7.54(22.45)
Biomass Burning	171.39(12.18)	14.21(4.22)	0.0208	1.21(4.45)	1.32(3.93)
Methane	492.39(34.99)	148.52(44.15)	0.0754	13.51(49.79)	13.33(39.68)
Extra Production	N/A	5.05(1.5)	N/A	0.48 (1.77)	0.68 (2.02)
Anthropogenic	301.52(21.43)	39.38(11.71)	0.0326	3.83(14.12)	7.98(23.74)
Regional contribution to Anthropogenic component					
International Shipping	0.76(0.05)	0.18(0.05)	0.0586	0.02(0.07)	0.02(0.07)
North America	36.47(2.59)	4.2(1.25)	0.0291	0.58(2.15)	0.65(1.93)
Europe	21.19(1.51)	2.5(0.74)	0.0294	0.44(1.62)	0.63(1.86)
East Asia	95.36(6.78)	11.35(3.37)	0.0297	1.06(3.9)	2.26(6.72)
South Asia	41.89(2.98)	5.69(1.69)	0.034	0.42(1.54)	2.4(7.14)
Russia; Belarus; Ukraine	7.97(0.57)	0.96(0.28)	0.03	0.17(0.64)	0.2(0.58)
Middle East	7.12(0.51)	1.27(0.38)	0.0444	0.18(0.65)	0.36(1.06)
Rest-of-the World	90.77(6.45)	13.24(3.94)	0.0365	0.96(3.54)	1.47(4.37)

Table 4: Same as Table 2 but for RC-tagged components

Source	Emissions (TgC/yr ²)	Tropospheric O ₃ burden (TgO ₃ /yr)	OPE ((mol O ₃ /mol C)/yr)	Surface mean (ppbv/yr)	Population weighted mean (ppbv/yr)
Total	3.74(0.28)	0.91(0.28)	N/A	0.1(0.37)	0.16(0.48)
Stratosphere	N/A	Insig	N/A	0.02(0.43)	Insig
Aircraft	0.004(1.74)	0.0005(1.88)	Insig	0.03(1.64)	0.03(1.59)
Biogenic	Insig	0.09(0.16)	Insig	0.01(0.29)	0.03(0.42)
Biomass Burning	Insig	Insig	Insig	Insig	Insig
Extra Production	N/A	0.03(0.64)	N/A	0.01(0.59)	0.01(0.74)
Methane	2.27(0.48)	0.63(0.43)	Insig	0.07(0.5)	0.05(0.41)
Anthropogenic	1.74(0.61)	0.32(0.84)	0.0001	Insig	0.08(1.13)
Regional contribution to Anthropogenic component					
International Shipping	0.01(1.79)	-0.003(- 1.03)	-0.0014	-0.005(- 0.76)	Insig
North America	-1.8(- 3.16)	-0.18(- 2.85)	0.0002	-0.02(- 2.68)	-0.02(- 2.55)
Europe	-0.72(- 2.53)	-0.09(- 2.54)	Insig	-0.01(- 2.43)	-0.02(- 2.22)
East Asia	1.58(2.2)	0.24(2.78)	0.0002	0.02(2.86)	0.04(2.43)
South Asia	1.19(3.76)	0.16(3.58)	Insig	0.01(3.56)	0.06(3.31)
Russia, Belarus, Ukraine	-0.06(- 0.72)	-0.01(- 0.99)	-0.0001	-0.0013(- 0.71)	-0.002(- 0.88)
Middle East	Insig	-0.02(- 1.07)	-0.0002	-0.0016(- 0.83)	-0.004(- 0.93)
Rest-of-the World	1.5(1.95)	0.22(1.84)	Insig	0.01(1.6)	0.03(2.24)

Table 5: Same as table 3 but for RC-tagged metrics