



# Satellite-observed relationships between land cover, burned area and atmospheric composition over the southern Amazon.

Emma Sands<sup>1</sup>, Richard Pope<sup>2,3</sup>, Ruth M. Doherty<sup>1</sup>, Fiona M. O'Connor<sup>4,5</sup>, Chris Wilson<sup>2,3</sup>, and Hugh Pumphrey<sup>1</sup>

<sup>1</sup>School of Geosciences, University of Edinburgh, Edinburgh, United Kingdom

<sup>2</sup>School of Earth and Environment, University of Leeds, Leeds, United Kingdom

<sup>3</sup>National Centre for Earth Observation, University of Leeds, Leeds, United Kingdom

<sup>4</sup>Met Office, Exeter, United Kingdom

<sup>5</sup>Department of Mathematics and Statistics, Global Systems Institute, University of Exeter, Exeter, United Kingdom

**Correspondence:** Emma Sands (e.g.sands@ed.ac.uk)

**Abstract.** Land surface changes can have substantial impacts on the interactions between the biosphere and atmosphere. In South America, rainforests abundantly emit biogenic volatile organic compounds (BVOCs), which coupled with pyrogenic emissions from deforestation fires, can have substantial impacts on regional air quality. We use novel and long-term satellite records of trace gases, aerosol optical depth (AOD), vegetation and burned area to characterise the impacts of biogenic and pyrogenic emissions on atmospheric composition for the period 2001 to 2019 in the southern Amazon, a region of substantial deforestation. We find that the seasonal cycle for all of the atmospheric constituents peaks in the dry season (August-October) and that year-to-year variability in carbon monoxide (CO), formaldehyde (HCHO), nitrogen dioxide (NO<sub>2</sub>), and AOD is strongly linked to burned area. We find a robust relationship between broadleaf forest cover and total column isoprene ( $R^2 = 0.59$ ), while burned area exhibits an approximate 5<sup>th</sup> root power law relationship with tropospheric column NO<sub>2</sub> ( $R^2=0.32$ ), both in the dry season. Vegetation and burned area together show a relationship with HCHO ( $R^2=0.23$ ). Overall, we provide a detailed observational quantification of biospheric process influences on southern Amazon regional atmospheric composition, which in future studies can be used to help constrain the underpinning processes in Earth System Models.

## 1 Introduction

Ten million hectares of forest were cut down globally each year over 2010-2020 (Ritchie and Roser, 2021). Land cover changes such as deforestation can substantially modify the emissions of biogenic gases and aerosols, for example biogenic volatile organic compounds (BVOCs) (Fowler et al., 2009; Pacifico et al., 2012). Additionally, one of the major causes of land cover change globally is fire (Heald and Spracklen, 2015), which also impacts atmospheric composition.

BVOCs are emitted during photosynthesis and particular plant development stages, or as a response to stresses on plants, such as droughts and insect infestations (Loreto and Fares, 2013). Estimates of the global emission of isoprene (C<sub>5</sub>H<sub>8</sub>), the globally dominant BVOC, are within the ranges of 300-600 TgC yr<sup>-1</sup> (Arneth et al., 2011; Cao et al., 2021; Szopa et al., 2021), highlighting that substantial uncertainties in its magnitude remain.



While BVOCs are associated with a wide range of vegetation, particular plants may emit different compounds more strongly or not at all, resulting in the use of emission factors to describe the sensitivity of emission strength to plant type (e.g. Guenther et al. (1995, 2012); Pacifico et al. (2011); Weber et al. (2023)). For example, isoprene is associated with tropical broadleaf trees, while monoterpenes are particularly associated with needleleaf trees (Artaxo et al., 2022), which has implications for the geographical distribution of different BVOC emissions. In the Amazon rainforest, isoprene and methanol ( $\text{CH}_3\text{OH}$ ) are the most strongly emitted biogenic compounds based on mixing ratio measurements (Yáñez-Serrano et al., 2020). Although biogenic emissions are the largest sources of isoprene, monoterpenes and methanol, they are also emitted during biomass burning (Akagi et al., 2011; Ciccioli et al., 2014; Bates et al., 2021).

Satellite measurements of formaldehyde ( $\text{HCHO}$ ), a common oxidation product of BVOCs, are often used to estimate BVOC emissions (e.g. Palmer et al. (2006); Millet et al. (2008); Marais et al. (2012); Kefauver et al. (2014); Stavrou et al. (2015); Strada et al. (2023)). However, the pyrogenic source of  $\text{HCHO}$  is more significant than the pyrogenic emission of isoprene, and it is an oxidation product of many other gases, introducing challenges to the interpretation of the data (Freitas and Fornaro, 2022; Palmer et al., 2007). Recent work has enabled the measurement of isoprene column densities from space (Fu et al., 2019; Wells et al., 2020, 2022; Palmer et al., 2022). The new isoprene measurements have created opportunities to address the uncertainties in regional isoprene emissions. This is particularly relevant for regions experiencing land cover change and in the southern hemisphere, where ground level measurements are sparse, despite the major BVOC emission sources being located in the tropical southern hemisphere (Paton-Walsh et al., 2022). We utilise these measurements alongside  $\text{HCHO}$  in this study to quantify vegetation-driven changes in composition.

The Amazon experiences significant impacts from fire activity (Reddington et al., 2015).  $\text{NO}_2$  is a trace gas measurable from space that is primarily emitted during combustion, whether that is biomass burning or anthropogenic emissions. Due to its short lifetime,  $\text{NO}_2$  can be indicative of biomass burning activity. In the context of land cover change through biomass burning, the burning of different land cover types is associated with different  $\text{NO}_2$  emission rates (Schreier et al., 2014). Wiedinmyer et al. (2023) assign the highest  $\text{NO}_2$  emission factors for burning of tropical forests, followed by savanna grasslands and lower values for crops and woody savanna. This is in contrast to earlier emission inventory estimates from Akagi et al. (2011), where the biomass burning emission factor for  $\text{NO}_x$  for tropical forests was lower than for savannas, although uncertainties were substantial.

These biomass burning  $\text{NO}_x$  emissions can play a key role in enabling ozone ( $\text{O}_3$ ) formation from VOC emissions, including BVOCs (e.g. Helas et al. (1995)). Particularly in the Amazon, a  $\text{NO}_x$ -limited region abundant in BVOCs,  $\text{O}_3$  production increases substantially over the rainforest in plumes of anthropogenic or pyrogenic emissions (Kuhn et al., 2010; Bela et al., 2015).

Fires also emit significant amounts of aerosols and carbon monoxide ( $\text{CO}$ ) (Badr and Probert, 1994; Wiedinmyer et al., 2023) and are a key factor in driving regional aerosol concentrations in the dry season (Reddington et al., 2015). In addition to the pyrogenic sources of these species, in areas such as the Amazon forest, there may be substantial biogenic contributions, for example, through the role of BVOCs in the formation and growth of secondary organic aerosol (SOA) (Artaxo et al., 2013; Shrivastava et al., 2017; Artaxo et al., 2022). SOA, as a component of particular matter air pollution, is detrimental to human



health (Kim et al., 2015). In the Amazon, the biogenic source of aerosols and CO has a greater relative contribution to regional atmospheric composition outside the wildfire season, when pyrogenic emissions decrease (Artaxo et al., 2022). Understanding the drivers of biogenic and pyrogenic emissions of O<sub>3</sub> and particulate matter precursors is important for both regional air quality and climate.

Overall, land cover change influences regional atmospheric composition through biogenic and pyrogenic sources, which can vary substantially over time and space. In this study, we aim to quantify the relevance of vegetation and fire to spatial and temporal variations in regional atmospheric composition over the last two decades, as observed using satellite remote sensing, over an area of significant land cover change: the southern Amazon.

Six measures of trace gases and aerosols in the atmosphere were chosen to represent a range of chemical species that may be impacted by biogenic and/or pyrogenic emissions. These are isoprene, methanol, HCHO, CO, NO<sub>2</sub>, and aerosol optical depth (AOD), which indicates the amount of aerosol in the atmospheric column, and will be referred to as atmospheric constituents throughout this paper. We compare this range of atmospheric constituents to vegetation and fire proxies using both new and complementary satellite datasets to build a comprehensive picture of the relative impact of both biogenic and pyrogenic sources on regional atmospheric composition. The paper will introduce the region, data and methodology in section 2, before looking at the spatial and seasonal distribution of the atmospheric constituents in the region and their links to both land cover and burned area (section 3 for results and section 4 for discussion and conclusions).

## 2 Data and Methodology


### 2.1 Study region

The southern Amazon is one of the regions that has undergone substantial deforestation in the 21<sup>st</sup> century. The region investigated in this study: 50°- 70°W, 5°- 25°S (Fig. 1); covers the majority of the "arc of deforestation", which forms the epicenter of Amazon deforestation (Santos et al., 2021; Silva Junior et al., 2021; Reddington et al., 2015). The area includes parts of the southern Amazon, as well savannas and grasslands such as the Cerrado and Sarmiento.

### 2.2 Remote sensing data sources

#### 2.2.1 Land surface data

The Moderate Resolution Imaging Spectroradiometer (MODIS) land cover data at 0.05° resolution (product MCD12C1, Friedl and Sulla-Menashe (2015), last access July 2022) was chosen as a measure of vegetation type based on its long temporal record of annual data (Table 1), and a comparison of the magnitude and timing of Amazon deforestation in three land cover datasets (see Supplement). The MODIS sensor operates on two satellites: Terra (launched in December 1999) and Aqua (launched in May 2002). For 2001-2020, the time period for which the data was used, the Terra and Aqua crossing times were, respectively, 10:30 and 13:30 local time. Both had sun synchronous, near polar orbits and an altitude of 705 km. The land cover data is available since 2001 and combines retrievals from both satellites. From the multiple land cover classifications available



**Figure 1.** Change in forest cover (% of grid cell area deforested or re/afforested) between 2001 and 2019 calculated using MODIS land cover data. The region analysed in this study is marked with the purple box.

in the MCD12C1 product, the 17-class International Geosphere-Biosphere Programme classification (IGBP, Loveland and Belward (1997)) was chosen for this project, as it has been used in relevant work, such as the FINNv2.5 fire emission inventory (Wiedinmyer et al., 2023).

The leaf area index (LAI) data is also a MODIS product (MOD15A2H, Myneni et al. (2021), last access January 2024) and can represent vegetation abundance. The LAI product is based on measurements from the Terra satellite at an 8-day temporal resolution and is available since February 2000 to present-day. We used the LAI product to calculate the annual and monthly mean values for the period 2001-2020 at 1° by 1° resolution in Google Earth Engine (Gorelick et al., 2017).

The burned area to represent fire activity was obtained from Version 4 of the Global Fire Emissions Database (GFED4, Giglio et al. (2013), last access August 2022). Monthly burned area is available at 0.25° spatial resolution from August 2000 to December 2016. The data for 2001-2016 was used in the analysis. A summary of the land surface and atmospheric composition datasets is provided in Table 1.

### 2.2.2 Atmospheric composition data

Gridded total isoprene columns for 2012-2020 at 0.5° × 0.625° spatial resolution at monthly resolution were obtained from Wells et al. (2020). These data are derived from measurements taken using the Cross-track Infrared Sounder (CrIS), a Fourier-transform spectrometer, onboard the Suomi-NPP satellite. Suomi-NPP was launched in October 2011 and has a sun-synchronous orbit and near-global twice daily coverage, with a daytime local overpass time of 13:30. Isoprene column densities were calculated using two isoprene infrared absorption features in the spectral range 890-910 cm<sup>-1</sup> and the data product has been quality assessed (Fu et al., 2019; Wells et al., 2020, 2022).



**Table 1.** Summary of satellite datasets analysed in the study. All datasets were analysed on  $1^\circ \times 1^\circ$  spatial resolution.

Variable	Source	Temporal Period	Period of Data
Land Cover	MODIS, Terra and Aqua	Annual	2001-2020
Leaf Area Index (LAI)	MODIS, Terra	Monthly/Annual	2001-2020
Burned area	GFED4	Monthly	2001-2016
Isoprene	CrIS, Suomi-NPP	Monthly	2012-2020
Methanol	IASI, Metop-B	Monthly	2008-2018
Formaldehyde (HCHO)	OMI, EOS Aura	Monthly	2005-2018
Carbon Monoxide (CO)	Mopitt, Terra	Monthly	2001-2019
Nitrogen Dioxide (NO <sub>2</sub> )	OMI, EOS Aura	Monthly	2005-2020
Aerosol Optical Depth (AOD)	MODIS, Terra	Monthly	2000-2019

The total column methanol data comes from the Infrared Atmospheric Sounding Interferometer (IASI) on board the MetOp-B satellite. MetOp-B is one of three Eumetsat MetOp satellites, which have sun-synchronous polar orbits and local overpass times of 9:30 and 21:30. The daytime (9:30) data for 2008-2018 was produced using the Infrared-Microwave-Sounding (IMS) scheme developed by the Rutherford Appleton Laboratory (RAL) (Pope et al., 2021).

110 HCHO and NO<sub>2</sub> are measured using the Ozone Monitoring Instrument (OMI) located on the Earth Observing System Aura satellite. OMI employs spectrometers in visible and ultraviolet wavelengths and provides daily global coverage since late 2004. Aura flies at 705 km and has an equator crossing time of 13:45 local time. Level 2 total column HCHO data was downloaded from the NASA Goddard Earth Sciences Data and Information Services Center (GES DISC, Chance (2007), last access: June 2022). Only pixels with the "good" main quality flag, which includes checks of fit convergence, column uncertainty and  
 115 absolute column value, and cloud cover less than 20% were used to calculate monthly mean values. We found an underlying positive trend in the HCHO data, possibly associated with instrument degradation (Wang et al., 2022), and anomalous values in 2019. Consequently, only the HCHO data for 2005-2018 was used and the monthly values were de-trended based on a remote Pacific region (see Supplement). This highlights local variations due to biogenic and pyrogenic emissions, as opposed to those driven by changes to the global background.

120 The tropospheric column NO<sub>2</sub> data is the Quality Assurance for Essential Climate Variables (QA4ECV) tropospheric NO<sub>2</sub> product (Boersma et al. (2011), last access: December 2022) available as global monthly averages at  $0.125^\circ \times 0.125^\circ$  spatial resolution. The data was downloaded for 2005-2020. The monthly mean values only include retrievals with cloud radiance fractions under 50%, which is approximately equivalent to geometric cloud fractions under 20%.

AOD is provided as part of a level 3  $1^\circ \times 1^\circ$  spatial resolution monthly product from MODIS measurements on board the  
 125 Terra satellite (MOD08\_M3, Platnick (2015), last access: May 2022). Each monthly statistically derived dataset (SDS) is based on the relevant MODIS Atmosphere Daily Global Joint Product. The quality controlled overland AOD data is available at three wavelengths: 0.47  $\mu\text{m}$ , 0.55  $\mu\text{m}$  and 0.66  $\mu\text{m}$ . The 0.47  $\mu\text{m}$  data for 2000-2019 is used throughout the analysis, as it was found to exhibit the strongest statistical relationship with the land variables (not shown).



The total column CO data is from the Measurement of Pollution in the Troposphere (MOPITT) sensor also on board the  
130 Terra satellite. Monthly mean values at  $1^\circ \times 1^\circ$  resolution were calculated for 2001-2019 from the version 7 level 2 product  
(NASA/LARC/SD/ASDC (2000), last access: December 2021). The data for 2000 was omitted due to large data gaps that year.  
Only clear-sky scenes are included in the MOPITT retrievals.

### 2.3 Analysis of remote sensing data for the southern Amazon


All observational data was re-gridded to a  $1^\circ \times 1^\circ$  horizontal resolution using linear interpolation to provide spatially consistent  
135 datasets for analysis and inter-comparison. The atmospheric composition and burned area data were all analysed on a monthly  
temporal resolution, while the land cover and LAI data were annual, with the exception of calculating the LAI seasonal cycle.  
To better understand the impacts of seasonality, we separated the monthly data into wet (February, March, April) and dry  
(August, September, October) seasons. The three month intervals were chosen based on seasonal variations in precipitation  
and compatibility with previous definitions (Barkley et al., 2009; Reddington et al., 2015). The dry season data was further  
140 separated into high burned area (monthly burned areas  $\geq 0.04\%$  grid cell area) and low burned area grid cells. The threshold  
for defining high burned area of  $\geq 0.04\%$  was chosen to ensure sufficient sample sizes in each category, while identifying areas  
with clear fire signals (see Supplement). There is a 6 km difference in elevation within the region with maximum elevations in  
the south west (SW). As some of the satellite retrievals are associated with high errors over the high altitudes of the Andes, the  
small portion of the domain  $>1000$  m a.s.l. was not included in our analysis.

145 Regression analysis was used to quantify the relationship between surface variables: land cover, LAI and burned area, and  
the atmospheric constituents. Spearman rank correlation, and ordinary least squared (OLS) and Theil-Sen regression methods  
(Fernandes and Leblanc, 2005) were utilised. Where a clear relationship was identified between an atmospheric constituent and  
land cover or burned area, the data were binned by the explanatory variable, and a weighted least squares (WLS) regression  
was used to account for the variance of the data in each bin. Throughout the text we use standard errors to represent uncertainty  
150 ranges.

## 3 Results

### 3.1 Change in atmospheric constituents, land cover and burned area through time

Over the early 21<sup>st</sup> century (2001-2019), forest cover averaged over the southern Amazon region decreases by 3.8% from  
52.0% to 48.2% of the study region (Fig. 2a). The deforestation is greatest in the north (N) of the domain at around  $10^\circ$  S  
155 (see Fig. 1). Generally across the region, forest cover reduces substantially from 2001 to 2013 with a smaller decline thereafter  
(Fig. 2b). Over the same period, savanna and grassland expand from 46.7% to 50.5% (see Supplement for separated savanna  
and grassland time series). The two land cover categories: broadleaf forest and savanna/grassland, display opposite behaviours,  
reflecting that broadleaf forest cover is being replaced by the savanna/grassland modal land cover type.



**Figure 2.** Annual mean values for (a) broadleaf forest cover percentage, (b) savanna and grassland cover percentage, (c) LAI, (d) burned area (based on monthly sums) and the atmospheric constituents: (e) isoprene, (f) methanol, (g) HCHO, (h) CO, (i) NO<sub>2</sub> and (j) AOD; averaged for the southern Amazon region for all years available for each variable. The shading represents the standard error based on all data included in calculating the annual mean: the values of each grid cell for a given year for the land cover and LAI, or the monthly values of each grid cell for burned area and the atmospheric constituents. The larger amount of data used in calculating the mean values for subplots d-j results in smaller standard errors.



Mean LAI and total annual burned area averaged over the study region do not have consistent trends over 2001-2019 and  
160 2001-2016, respectively (Fig. 2c,d). Average annual LAI values in the region fluctuate around a value of 3.1 with considerable  
inter-annual variation. As the regional estimate of LAI is dependent on all vegetation types in the domain, any decrease in  
LAI due to decreases in broadleaf forest (high LAI values) may be of a lesser magnitude than year-to-year variability in the  
vegetation overall due to, e.g., weather or disease. Burned area also exhibits substantial year-to-year variability, with maximum  
values observed in 2007 and 2010, while the least burning occurs in 2009. Since 2011, the monthly average burned area has  
165 remained below 310 km<sup>2</sup> (or 0.007% of the region) and the variability has decreased.

The year-to-year variability for annual regional-mean CO, NO<sub>2</sub> and AOD is similar to that for burned area (minima in 2009  
and maxima in 2007 and 2010; Fig. 2h,i,j). These atmospheric constituents also show reduced year-to-year variability from  
2011 onwards, with the exception of elevated values in 2015 for AOD and CO. The 2007 and 2010 maxima are also observed  
in the HCHO record, although the year-to-year variability does not reduce for this trace gas (Fig. 2g). The similarities in the  
170 temporal record suggest burned area has an influence on the domain averaged AOD, CO, NO<sub>2</sub> and HCHO.

Methanol and isoprene are also characterised by substantial inter-annual variability and neither shows a clear trend through  
time for the region-average (Fig. 2e,f). Consequently, other annually varying factors affecting these constituents, such as mete-  
orology or changes in other sources and sinks (Pacífico et al., 2009; Wohlfahrt et al., 2015), may have a greater impact than the  
decrease in broadleaf forest cover, which is limited in its spatial extent, over this time period. It is noted that the annual mean  
175 isoprene and methanol datasets do not extend as far back in time as the land cover dataset and the change in broadleaf forest  
cover is smaller during the period covered by these datasets.

### 3.2 Seasonal cycle in atmospheric composition and burned area


The regional -average seasonal cycles of the six atmospheric constituents are found to be similar with a peak in the dry  
season (August to October) for all species (Fig. 3). Uniquely, isoprene shows a secondary peak earlier in the calendar year in  
180 March/April, when the isoprene column densities reach 80% of the annual maximum (August regional mean of  $8.6 \pm 0.5 \times$   
 $10^{15}$  molecules cm<sup>-2</sup>).

Several of the constituents also have co-occurring minima. Isoprene, methanol and AOD all have the lowest values of  $4.7 \pm$   
 $0.1 \times 10^{15}$  molecules cm<sup>-2</sup>,  $0.1 \pm 0.01$  ppbv, and  $0.11 \pm 0.004$ , respectively, around June, while HCHO, CO and NO<sub>2</sub> remain  
more stable between December and June.

185 The amplitudes of the seasonal cycles of the different constituents vary. Isoprene, HCHO and CO have relatively low intra-  
annual variation, as their column amounts remain above 50% of their maxima throughout the year. In comparison, methanol,  
AOD and NO<sub>2</sub> exhibit a more extreme seasonality, with values dropping below 30% of their annual maxima. This pronounced  
dry season peak in the atmospheric constituents is consistent with the burned area seasonal cycle, which rises from  $0.02 \pm$   
 $0.002\%$  in July to  $0.06 \pm 0.009\%$  in September, before decreasing to  $0.02 \pm 0.002\%$  again in October. Burned area values  
190 remain below 10% of the September peak for the rest of the year.

The LAI seasonality is substantially different, as values remain above 70% of the maximum throughout the whole year. LAI  
is elevated at  $>3$  ( $>90\%$  of the maximum value) between April and October. The regional monthly mean drops slightly ( $<2.7$





**Figure 3.** Seasonal cycles of domain-averaged LAI, burned area, isoprene, methanol, HCHO, CO, NO<sub>2</sub> and AOD for the periods where each dataset is available between 2001-2020. All variables have been normalised (to a value of 1) against their respective regional maximum monthly mean. The shaded areas represent the standard error for each variable for a given month.

or <80%) from December to February. The lower percentage amplitude of the LAI seasonal cycle may be driven by different phenologies of the vegetation in the domain, e.g. Cerrado savanna grassland flowering is relatively consistent throughout the year, while Amazon forest vegetation tends to flower in the dry season (Morellato et al., 2013). Other research in Brazil has found that while seasonal changes in forests are associated with solar radiation, they are driven by rainfall for savannas and grasslands, resulting in opposite cycles for the different land cover types (Myneni et al., 2007). A comparison of 6 LAI datasets consistently shows that in the southern Amazon the LAI in regions of broadleaf forest cover is higher in July, compared to January, while the savanna/grassland region has the opposite LAI signal (Fang et al., 2013). Consequently, phenology-driven LAI values may peak at different times depending on the land cover types, resulting in full/partial vegetation coverage of the region throughout the year.

The magnitudes of all atmospheric constituents increase in the dry season when burned area is largest, suggestive of changes to the atmospheric chemistry due to a substantial pyrogenic source. In particular, the seasonal variability of NO<sub>2</sub> and AOD closely matches the seasonality in burned area, while methanol does so with a lag of one month. As outlined above, vegetation cover, as represented by the LAI, shows more consistent values throughout the year. Therefore, constituents with lower seasonal amplitudes are potentially more strongly linked to the seasonality of biogenic emission sources, although other factors such as atmospheric lifetime will have an important impact on the respective atmospheric constituent concentrations.

### 3.3 Spatial distribution of vegetation, fire and atmospheric composition

Across the southern Amazon, broadleaf forest, savanna and grasslands represent the dominant vegetation types. Annual average broadleaf forest cover dominates in the north west (NW), with land surface coverage typically between 80 and 100%, while




savannas and grasslands represent the main land cover (80-100% coverage) in the south east (SE) (Fig. 4). In the centre of the domain, a transition region exists where the competing phenologies typically have 40-60% coverage. The corresponding LAI spatial distribution highlights larger values (3.5 to >5.0) in the NW and lower values (2.0-3.5) in the SE. This is consistent with broadleaf forest having a larger biomass (per unit area) in comparison to savanna/grassland vegetation types.


215 For the dry season burned area data, there are sporadic hot spots peaking at >0.3%. The most coherent spatial structures are a filament of burned area values between 0.2 to 0.3% stretching along the Bolivian eastern border down into Paraguay in the S of the region, and the cluster of western Brazilian burned area values at >0.1%, though peaking at 0.3-0.5%. This latter feature is generally consistent with the reported "arc of deforestation" in the Amazon (Reddington et al., 2015; Santos et al., 2021). The hatching in Figure 4d represents at least a 2.5% decrease in broadleaf forest between 2001 and 2019, which coincides with the  
220 burned area patterns reported here. Therefore, the fire activity is related to regions undergoing deforestation and, to a certain extent, the land cover classifications. For instance, the burned area filament in Figure 4d closely follows the high (>90%)/low (<20%) savanna/grassland and broadleaf forest structure in Figure 4a & b. Thus, it is suggestive of transitional regions between biomes driven by predominately anthropogenic pyrogenic activity. Overall, the spatial distribution of burned area exhibits more localised maxima than vegetation cover.

225 The spatial distributions of multi-annual mean values of total column isoprene are similar to those of broadleaf forest cover and LAI for both the wet and dry season (c.f. Fig. 5a,b and Fig. 4b,c). Values decrease from the far NW (maximum of  $2.7 \times 10^{16}$  molecules  $\text{cm}^{-2}$  in the dry season) to  $<0.5 \times 10^{16}$  molecules  $\text{cm}^{-2}$  in the south (S), following the NW-SE transition from forest to savanna/grassland as evident in the dry season total column isoprene values (c.f. Fig. 4b and Fig. 5b). Over the broadleaf forest region, isoprene is elevated in the west (W), where the maximum forest cover occurs, compared to the east  
230 (E). The spatial similarities between broadleaf forest cover and total column isoprene amounts suggest the broadleaf forest is the dominant source of this BVOC for this region, especially in the dry season. This finding is also consistent with the short lifetime of isoprene confining the peak concentrations to the source region. The W-E gradient in mean total column isoprene over the forested region itself suggests variability in isoprene emissions within the broadleaf forest biome due to different plant species or local environmental conditions, as found by Gu et al. (2017), who identify an elevation gradient in isoprene  
235 emissions in the tropical forest N of the region studied here, probably driven by plant species distributions. In this study, the highest total column isoprene values occur at elevations of around 200 m a.s.l., with lower isoprene column densities in the N of the forest, where elevations decrease to below 100 m a.s.l.

Two other atmospheric constituents have similar distributions in the wet season to that of broadleaf forest cover: CO and AOD. CO is elevated in the NW ( $>16 \times 10^{17}$  molecules  $\text{cm}^{-2}$ , maximum of  $18.2 \times 10^{17}$  molecules  $\text{cm}^{-2}$ ), compared to the  
240 SE ( $<14 \times 10^{17}$  molecules  $\text{cm}^{-2}$ ). AOD is similarly elevated in the N (>0.2, maximum of 0.35) and reaches a minimum in the SE of the study region (<0.1) (Fig. 6a,c). The spatial patterns of CO and AOD during the wet season are consistent with the broadleaf forest acting as a source of biogenic precursors of CO and aerosols, as values increase over more densely forested areas (compare with Fig. 4b). This is consistent with biogenic sources having a greater relative impact on CO and aerosols in the wet season, when pyrogenic emissions are minimal (Artaxo et al., 2022).




**Figure 4.** The annual mean distribution for 2001-2019 of (a) savanna and grassland and (b) broadleaf forest cover, represented as the % of each grid cell covered by the respective land cover type, and (c) annual mean LAI for 2001-2019. Dry season (August-October) monthly mean burned area (% grid cell area) for 2005-2016 is shown on panel (d) along with regions where at least 2.5% of the area has been deforested marked by hatching. Areas over 1000 m a.s.l. based on the GMTED2010 digital elevation model (Danielson and Gesch, 2011) have been masked on all panels.



**Figure 5.** Regional distributions of column mean isoprene (top row (a, b), molecules  $\text{cm}^{-2}$ ), methanol (middle row (c, d), ppbv), and formaldehyde (bottom row (e, f), molecules  $\text{cm}^{-2}$ , note colorbar starts at  $0.5 \times 10^{16}$  molecules  $\text{cm}^{-2}$ ) for the wet season (left column) and dry season (right column). The wet season includes the months of February-April, while the dry season covers August-October. The time period varies with constituent. Grid cells with mean elevations over 1000 m a.s.l. are shaded grey and not included in further analysis.

245 In contrast, wet season HCHO and  $\text{NO}_2$  increase from the NW (HCHO and  $\text{NO}_2$  column values of  $\sim 0.8 \times 10^{16}$  molecules  $\text{cm}^{-2}$  and  $< 0.6 \times 10^{15}$  molecules  $\text{cm}^{-2}$ , respectively) to the SE, where HCHO reaches a maximum of  $3.2 \times 10^{16}$  molecules  $\text{cm}^{-2}$  and  $\text{NO}_2$ :  $1.36 \times 10^{15}$  molecules  $\text{cm}^{-2}$ . This spatial pattern is not consistent with biogenic emissions from the forest being the dominant controlling factor (Figures 5e and 6e, compare to Fig. 4b). Further, biomass burning is minimal in the wet season (see section 3.2), suggesting another source or sink is driving total column HCHO and tropospheric column  $\text{NO}_2$  in this season.



**Figure 6.** Regional distributions of CO (top row (a, b), molecules  $\text{cm}^{-2}$ , note colorbar starts at  $10 \times 10^{17}$  molecules  $\text{cm}^{-2}$ ), AOD (middle row (c, d)), and  $\text{NO}_2$  (bottom row (e, f), molecules  $\text{cm}^{-2}$ ) for the wet season (left column) and dry season (right column). The wet season includes the months of February-April, while the dry season covers August-October. The time period varies with constituent. Grid cells with mean elevations over 1000 m a.s.l. are shaded grey and not included in further analysis.

250 The wet season methanol concentrations are also different to that of vegetation cover. The highest methanol concentrations of  $\geq 0.4$  ppbv are recorded in the far E of the domain between  $10^\circ$  and  $16^\circ$  S (Fig. 5c).

The dry season spatial patterns of HCHO, CO and AOD differ somewhat from their wet season distributions, suggesting a change in dominant sources/sinks e.g. from vegetation to fires, transport and/or atmospheric chemistry processes. CO appears more well-mixed compared to the other trace gases as expected with its relatively longer lifetime (average tropospheric lifetime



255 of 1-3 months, Seinfeld and Pandis (2016)) compared to the other species (Fig. 6b). HCHO, CO and AOD reach maximum values (up to  $3.7 \times 10^{16}$  molecules  $\text{cm}^{-2}$ ,  $32.7 \times 10^{17}$  molecules  $\text{cm}^{-2}$  and 0.88 respectively) over the transition zone between forests and other land cover classes, although the exact locations of these peaks vary. This region is most strongly affected by deforestation and proximate to the highest average dry season burned areas (compare with Fig. 4). Total column HCHO decreases beyond this zone, consistent with a significant pyrogenic source (Freitas and Fornaro, 2022; Palmer et al., 2007),  
260 both further into the broadleaf forest ( $1.2$  to  $1.6 \times 10^{16}$  molecules  $\text{cm}^{-2}$ ) and to the S, where values drop to  $<0.6 \times 10^{16}$  molecules  $\text{cm}^{-2}$  (Fig. 5f). Minima in the AOD and CO data are also found in the S in the foot of the Andes and in the SE ( $<0.2$  and  $<18 \times 10^{17}$  molecules  $\text{cm}^{-2}$ , respectively) (Fig. 6d). Peak total column CO is found slightly further S to the HCHO maximum ( $10$ - $12^\circ\text{S}$  compared to  $8$ - $11^\circ\text{S}$ ). AOD is characterised by two maxima ( $>0.8$ ) within the same region as CO, which occur in areas of mixed land cover (forest cover  $10$ - $100\%$ ) that have experienced substantial deforestation ( $>2.5\%$  area  
265 deforested) and some burning (dry season monthly mean burned area  $\sim 0.01\%$ ) (compare Figures 6d and 4). However, this does not correspond to the locations of maximum dry season burned areas. The co-location of these maximum values with regions of deforestation and biomass burning suggests a pyrogenic source, potentially emissions from deforestation fires, is important for these constituents. These constituents were also elevated to the N of this region where the forests are dominant, suggesting the presence of a biogenic source for this region or transport of pyrogenic emissions. Particularly in the case of HCHO, the  
270 presence of  $\text{NO}_x$  associated with biomass burning could affect the HCHO yield from BVOC emissions (Langford et al., 2022), potentially increasing HCHO formation from isoprene oxidation over the broadleaf forest.

Methanol dry season concentration values are similarly elevated ( $>0.6$  ppbv) in the N over the forest to savanna/grassland transition, but they are also consistently high further into the Amazon forest, as well as in the NE. Some of the maxima are located over areas of mixed vegetation, which experienced substantial deforestation over 2001-2019, as found for HCHO, CO  
275 and AOD (compare Fig. 5d with Figures 1 and 4), but other regions do not reflect any of the studied land surface variables.

The dry season  $\text{NO}_2$  column densities had several distinct maxima  $>2.7 \times 10^{15}$  molecules  $\text{cm}^{-2}$ , the highest being  $3.5 \times 10^{15}$  molecules  $\text{cm}^{-2}$ , all located  $9$ - $16^\circ\text{S}$ . Some of these overlap with the two AOD maxima. Unlike AOD,  $\text{NO}_2$  is clearly elevated over the maximum burned areas recorded at  $15^\circ\text{S}$ ,  $65^\circ\text{W}$  and in the E of the study region (Fig. 6d,f, Fig. 4d). In regions of minimal fire activity, such as deeper into the Amazon forest and the edge of the Andes, tropospheric column  $\text{NO}_2$  is  $<0.9$   
280  $\times 10^{15}$  molecules  $\text{cm}^{-2}$ . The spatial pattern of tropospheric column  $\text{NO}_2$  closely resembles that of burned area, highlighting the close relationship between the trace gas and fire activity, owing to its much shorter lifetime (around 1 day for  $\text{NO}_x$  at the surface; Jacob (1999)) compared to CO and aerosols.

### 3.4 Influence of vegetation and fire on atmospheric composition

#### 3.4.1 Variations in atmospheric composition with broadleaf forest cover

285 In this section, the variation in atmospheric composition as a function of broadleaf forest cover (i.e. percentage cover in  $10\%$  bins) is considered for the dry and wet seasons over the whole region and the time period of co-existing data. Within the dry season, the impact of pyrogenic activity is assessed by splitting the atmospheric constituent data into "low fire" ( $\leq 0.04\%$




burned area) and "high fire" (>0.04% burned area) regimes. Our results are relatively insensitive to the burned area threshold, as well as the bin width choice (see Supplement).

290 Isoprene consistently increases with higher broadleaf forest cover in all four regimes (Fig. 7a). The spatial distributions rather than temporal variations drive this signal. Values are lower in the wet season (approximately  $0.5-0.7 \times 10^{15}$  molecules  $\text{cm}^{-2}$ ) than the dry season for broadleaf forest cover values >20%. In densely forested areas (broadleaf forest cover >90%) the dry season isoprene column density is 50% greater than in the wet season at approximately  $1.0-1.5 \times 10^{15}$  molecules  $\text{cm}^{-2}$ . The greater change between seasons in isoprene column amounts at higher forest cover suggests that isoprene emissions  
295 from broadleaf forest have a stronger seasonality than those from other vegetation types. The differences in mean isoprene column densities between areas of high and low fire activity remain within  $0.3 \times 10^{15}$  molecules  $\text{cm}^{-2}$ . Consequently, isoprene responds to the change in forest cover more than the change in burned area, highlighting the importance of its biogenic source, as suggested in section 3.3.

For  $\text{NO}_2$ , a similar pattern occurs with larger column values in the dry season (approximately  $1.5-2.0 \times 10^{15}$  molecules  
300  $\text{cm}^{-2}$ ) than in the wet season ( $\leq 0.86 \times 10^{15}$  molecules  $\text{cm}^{-2}$ ) (Fig. 7f). However, the relationship with forest cover is non-linear with peak dry season column  $\text{NO}_2$  in the 40-50% forest cover bin. When split into the two fire regimes, there is a large column  $\text{NO}_2$  difference independent of forest cover bin. When fires are low, tropospheric column  $\text{NO}_2$  ranges between  $1.2$  and  $1.3 \times 10^{15}$  molecules  $\text{cm}^{-2}$ , while in the high fire regime all column  $\text{NO}_2$  values are  $>2.5 \times 10^{15}$  molecules  $\text{cm}^{-2}$  (i.e. at least 60% larger). Overall, as noted in section 3.3, this result is suggestive of the largest  $\text{NO}_2$  emissions from pyrogenic sources in  
305 forested regions and/or transition zones between vegetation types.

Consistent with the other constituents, column methanol values are larger (0.5-0.7 ppbv) in the dry season than wet (<0.3 ppbv). While the low fire regime is very similar to the dry season average, the high fire regime column methanol values are larger ( $\geq 0.6$  ppbv) for all forest cover bins. Similarly to  $\text{NO}_2$ , though, the peak values are in the 40-50% and 70-80% forest cover bins (Fig. 6b). Therefore, unlike for isoprene, the methanol columns are not linearly linked to forest cover, but typically  
310 a larger forest cover percentage bin will have larger methanol values.

The other atmospheric constituents (Fig. 7c,d,e) also have some similarities with both isoprene and  $\text{NO}_2$ . The wet season values are much lower than those in the dry season regardless of forest cover, consistent with the seasonal cycle of the regional averages discussed in section 3.2. In the dry season, HCHO, CO and AOD increase as forest cover increases from 0 to 50% and values remain elevated at higher forest covers ( $\geq 1.5 \times 10^{16}$  molecules  $\text{cm}^{-2}$ ,  $>25.7 \times 10^{17}$  molecules  $\text{cm}^{-2}$  and  $>0.55$ ,  
315 respectively, for forest cover >50%). The high fire regime is associated with higher values of the atmospheric constituents compared to the low fire regime across all forest cover values. HCHO is elevated by at least 13%, AOD by 27%, and CO by 7-17%. The overall maxima are associated with both high burned area and high forest cover of 80-90%. In the dry season average a secondary maximum occurs for the 40-50% (40-60% for HCHO) forest cover bin, resembling that of the peak in  
320  $\text{NO}_2$  and methanol values. This result suggests both vegetation and fire are important in driving the concentration of HCHO, CO and aerosols in the region in the dry season through a combination of biogenic emissions, particularly from the forest, and pyrogenic emissions in the transition zone and in forested regions, strengthening conclusions drawn from the analysis of spatial maps in section 3.3.



**Figure 7.** Mean monthly atmospheric constituent values depending on percentage broadleaf forest cover in a given grid cell for 4 categories: High burned area dry season (H), Low burned area dry season (L), dry season (D) and wet season (W). Each subplot shows data for one atmospheric constituent: (a) isoprene ( $1 \times 10^{16}$  molecules  $\text{cm}^{-2}$ ), (b) methanol (ppbv), (c) HCHO ( $1 \times 10^{16}$  molecules  $\text{cm}^{-2}$ ), (d) CO ( $1 \times 10^{17}$  molecules  $\text{cm}^{-2}$ ), (e) AOD, and (f)  $\text{NO}_2$  ( $1 \times 10^{15}$  molecules  $\text{cm}^{-2}$ ). Each subplot includes data for years when the given atmospheric constituent, land cover and burned area datasets overlap.

### 3.4.2 Variations in atmospheric composition with burned area

In this section, the change in the atmospheric constituents with burned area is analysed, depending on season (W - wet season, 325 D - dry season) and, for the dry season, dominant land cover (F -  $\geq 50\%$  forest cover, S -  $< 50\%$  forest cover, i.e.  $\geq 50\%$  savanna/grassland).





Although isoprene total column amounts are increased during the dry season, when fire activity is at its highest in the southern Amazon (see section 3.2), there is no clear relationship between the burned area extent and the isoprene column amount (Fig. 8a), as the highest dry season total column isoprene values occur at 0-0.01% ( $0.9 \pm 0.01 \times 10^{15}$  molecules  $\text{cm}^{-2}$ ) and 0.08-0.09% ( $0.9 \pm 0.1 \times 10^{15}$  molecules  $\text{cm}^{-2}$ ) burned area. The data does further highlight the relevance of land cover type for this trace gas. Regardless of the amount of burning, for each respective burned area bin, total column isoprene is at least 140% higher in regions of dominant forest cover than in regions dominated by savannas and grasslands. Therefore, forest cover is closely connected to isoprene emissions, supporting conclusions reached in section 3.4.1, suggesting the broadleaf forest is the dominant source of this BVOC in the region in the dry season, while the pyrogenic source, limited in its spatial extent, has a more minimal influence, despite the dry season maximum (see section 3.2).

The other constituents are also elevated over the forested, compared to the savanna/grassland, region, though the relative difference in atmospheric constituent between the savanna/grassland and forest categories varies. This difference is particularly pronounced for HCHO and AOD (Fig. 8c,e). HCHO and AOD total column amounts over forests ( $1.5\text{-}2 \times 10^{16}$  molecules  $\text{cm}^{-2}$  and  $0.6\text{-}1.1$ ) are, respectively,  $\geq 24\%$  and  $\geq 35\%$  higher than those over savannas/grasslands.  $\text{NO}_2$  has the smallest difference in values between the land cover categories ( $1.4\text{-}3.1 \times 10^{15}$  molecules  $\text{cm}^{-2}$  for savanna/grassland,  $1.3\text{-}4.2 \times 10^{15}$  molecules  $\text{cm}^{-2}$  for forest) (Fig. 8f). For  $\text{NO}_2$ , AOD, CO and HCHO the differences between land cover types increase at higher burned area values.


The increase in atmospheric constituents over the forest may be driven by the emission of biogenic precursors and/or higher pyrogenic emissions when forest, as opposed to savanna/grassland, vegetation is burned. The latter is particularly relevant for  $\text{NO}_2$ , as the difference in column densities between the land cover categories is only significant at high burned area values. At  $>0.09\%$  burned area,  $\text{NO}_2$  reaches  $4.2 \pm 0.1 \times 10^{15}$  molecules  $\text{cm}^{-2}$  over forest compared to  $3.1 \pm 0.1 \times 10^{15}$  molecules  $\text{cm}^{-2}$  over savanna/grassland.

On average, most of the same atmospheric constituents ( $\text{NO}_2$ , AOD, CO and HCHO) increase with burned area in the dry season, reaching peak values at 0.08-0.09% burned area (AOD and CO) or  $>0.09\%$  burned area ( $\text{NO}_2$  and HCHO). For methanol the increase with burned area is limited to values of burned area under 0.05%. The greater column amounts of these constituents associated with both forest cover and burned area are consistent with the dry season maxima observed in (or proximate to) forested regions and over burned areas on the spatial maps in section 3.3.

Consequently, all studied atmospheric constituents are influenced by land cover type, as their abundances increase over more densely forested areas. Additionally, maximum values are reached in the presence of burning, especially at  $\geq 0.07\%$  burned area for HCHO, CO, AOD and CO in forested regions, showcasing the pyrogenic source. At low burned area values there is little difference between forests and savannas/grasslands for  $\text{NO}_2$ , highlighting that burning is the major source of the trace gas in the region during the dry season and the role of land cover is to modify the emissions where burning occurs.

### 3.5 Statistical relationships between atmospheric composition and land cover, LAI and fire

In this section, statistical relationships between land cover, LAI, and burned area, and the atmospheric constituents for the dry season are explored, to quantify the extent to which land cover variables drive atmospheric composition over the southern



**Figure 8.** Mean monthly atmospheric composition depending on percentage burned area in a given grid cell for 4 categories: Savanna/grassland dominated dry season (S), Forest dominated dry season (L), Dry season (D) and Wet season (W). Each subplot shows data for one atmospheric constituent: (a) isoprene ( $1 \times 10^{16}$  molecules  $\text{cm}^{-2}$ ), (b) methanol (ppbv), (c) HCHO ( $1 \times 10^{16}$  molecules  $\text{cm}^{-2}$ ), (d) CO ( $1 \times 10^{17}$  molecules  $\text{cm}^{-2}$ ), (e) AOD, and (f) NO<sub>2</sub> ( $1 \times 10^{15}$  molecules  $\text{cm}^{-2}$ ). Each subplot includes data for years when the given atmospheric constituent, land cover and burned area datasets overlap.

Amazon for the time period both datasets are available. Spearman rank correlation (Table 2) and OLS regression (Table 3; see section 2) are utilised. The results from both methods are very similar, with the squared Spearman rank correlation coefficients slightly higher than the equivalent OLS  $R^2$  values.

There is a strong significant positive relationship between the two vegetation variables of broadleaf forest cover and LAI, and total column isoprene over the region (Spearman's  $r = 0.77$  for both, OLS  $R^2 = 0.59$  and  $0.54$  for broadleaf forest cover and



**Table 2.** Spearman rank correlation coefficients  $r$  (with the  $r^2$  for comparison with Table 3 in brackets). For each correlation, data for the three dry season months for years when the atmospheric constituent and surface variable datasets overlap is used. In the case of annual vegetation data, all months of a given year are assumed to have the same vegetation cover. All  $r$  values are significant at the  $p = 0.05$  level.

	Isoprene	Methanol	HCHO	CO	AOD	NO <sub>2</sub>
Broadleaf forest cover	0.77 (0.59)	0.19 (0.04)	0.44 (0.19)	0.41 (0.17)	0.39 (0.15)	-0.14 (0.02)
LAI	0.77 (0.59)	0.22 (0.05)	0.49 (0.24)	0.43 (0.18)	0.38 (0.14)	-0.03 (0.00)
Burned area	-0.09 (0.01)	0.15 (0.02)	0.23 (0.05)	0.17 (0.03)	0.18 (0.03)	0.46 (0.21)

**Table 3.** OLS regression coefficients of determination ( $R^2$ ) for the dry season for single and multiple linear regression calculations. For each regression, data for the three dry season months for years when the atmospheric constituent and surface variable datasets overlap is used. In the case of annual vegetation data, all months of a given year are assumed to have the same vegetation cover. All  $R^2$  values are significant at the  $p = 0.05$  level. The equivalent scatter plots for the data used to derive the linear OLS coefficients for broadleaf forest cover vs. isoprene and burned area vs. NO<sub>2</sub> are shown in figures 9a and 10, respectively.

	Isoprene	Methanol	HCHO	CO	AOD	NO <sub>2</sub>
Broadleaf forest cover	0.59	0.02	0.14	0.12	0.08	0.00
LAI	0.54	0.04	0.18	0.14	0.09	0.00
Burned area	0.02	0.01	0.03	0.01	0.02	0.25
Forest + Burned Area	0.59	0.04	0.19	0.14	0.11	0.25
LAI + Burned Area	0.54	0.05	0.23	0.15	0.12	0.26

LAI, respectively), as expected based on sections 3.3-3.4.2. However, there is a weak negative relationship between isoprene and burned area. In contrast, tropospheric column densities of NO<sub>2</sub> exhibit a moderate positive relationship with burned area (Spearman's  $r = 0.46$ , OLS  $R^2 = 0.25$ ), but weak relationships with land cover variables over the region.

The relationships for the other atmospheric constituents are mixed and much weaker. HCHO, CO and AOD all show positive weak to moderate relationships with both broadleaf forest cover and LAI ( $r$ -values around 0.4; OLS  $R^2 = 0.08$  to 0.18), suggesting some influence of vegetation on these atmospheric constituents, particularly for HCHO. The relationships between these atmospheric constituents and burned area are considerably weaker, but also positive ( $r$  values from 0.17 for CO to 0.23 for HCHO; OLS  $R^2$  between 0.01 and 0.03). Methanol showed a similarly weak positive relationship with both land cover and fire ( $r$  values from 0.15 with burned area to 0.22 with LAI; OLS  $R^2$  between 0.01 and 0.04). These weak results, as compared to the isoprene relationship with vegetation, highlight that with the longer lifetimes of these trace gases and aerosols, especially CO and methanol (average lifetimes of 1-3 months and 5 days, respectively, Seinfeld and Pandis (2016); Bates et al. (2021)), there is more transport and mixing.

Multiple OLS regression was also performed to test whether the combination of biogenic and pyrogenic sources improved their explanatory power for the variation in atmospheric constituents. The  $R^2$  value is unchanged for isoprene, while for NO<sub>2</sub> the  $R^2$  values are minimally affected, suggesting one main emission source of vegetation and fire, respectively, for these two




atmospheric species. For the other atmospheric constituents, the combination of broadleaf forest cover or LAI and burned area increased the  $R^2$  values (as compared to the OLS linear regression results) modestly. The change is most notable for HCHO ( $R^2$  value of 0.23 compared to 0.18) and then for AOD ( $R^2$  value of 0.12 compared to 0.09). While the overall  $R^2$  values are still much lower than the  $R^2$  value for the linear relationship between isoprene and broadleaf forest cover, for HCHO the multiple regression  $R^2$  values are only slightly lower than for the  $\text{NO}_2$ -burned area relationship. This illustrates the relevance of both a biogenic and pyrogenic source to HCHO column densities, as previously observed elsewhere in Brazil (Freitas and Fornaro, 2022).

The more robust relationships between broadleaf forest cover vs. isoprene and burned area vs.  $\text{NO}_2$  were studied in more detail (Fig. 9a, Fig. 10). The dry season composition data were binned based on broadleaf forest cover (for isoprene) or burned area (for  $\text{NO}_2$ ). We used a bootstrapping approach to test the significance of the isoprene ( $\text{NO}_2$ ) increase with broadleaf forest cover (burned area) and found the results are significant at the 95% confidence level (not shown). Over the region, both binned (at 10% broadleaf forest cover intervals) and non binned data suggest that isoprene increases by  $1.1 \times 10^{15}$  molecules  $\text{cm}^{-2}$  for every 10% increase in broadleaf forest cover in the dry season (OLS  $R^2 = 0.59$ , WLS  $R^2 = 0.97$ ) (Fig. 9). Consequently, the values for total isoprene column densities over completely (100%) forested regions are on average 4 times greater than over non-forested regions (0%). In non-forested regions, isoprene concentrations reflect the local background arising from emissions from non-forest species as well as mixing and transport of forest-related emissions on short time-scales. Tree species composition, in addition to forest dynamics and environmental conditions affecting the plant's emission efficiency, could explain the variability of isoprene column amounts within each forest cover bin.

The linear burned area/ $\text{NO}_2$  relationship was found to exhibit different sensitivities depending on vegetation cover (Fig. 10). Over regions of high forest cover and high LAI values, tropospheric  $\text{NO}_2$  increases with burned area more than over locations with low forest cover and lower LAI (i.e. those dominated by savannas and grasslands, see section 3.4.2). These different sensitivities are particularly clear up to around 0.5% burned area (Fig. 10, see Fig. 8f for average conditions for burned areas of 0 to >0.1%). Extremely high burned areas tend to occur in less forested regions, though the spatial distributions (Fig. 4) suggest these are savanna/grassland fires proximate to the broadleaf forest in the central part of the region. The dominance of savanna/grassland at extremely high burned areas could decrease the variation in  $\text{NO}_2$  explainable by forest cover over the whole dataset (Tables 2 and 3).

Hence, to represent this more complex burned area/ $\text{NO}_2$  relationship we also explored the log-log relationship for  $\text{NO}_2$  ( $\log_e(\text{NO}_2)$ ) and burned area ( $\log_e(\text{BA})$ ) ( $R^2 = 0.32$  for non binned data; WLS  $R^2 = 0.90$  for binned data (at 0.25  $\log_e(\text{BA})$  intervals) ; Fig. 10). This log-log relationship, approximately a fifth root power law, indicates that the change in tropospheric column  $\text{NO}_2$  is slightly greater at low burned area percentages compared to at high burned areas. This is consistent with the findings for the different vegetation types outlined above. The relationship was found to remain relatively consistent through time by testing the relationship between the natural logarithm of burned area from GFED5 (Chen et al., 2023) and natural logarithm of the  $\text{NO}_2$  tropospheric column for the later time period 2017-2020 (not shown). For both  $\text{NO}_2$  and isoprene, the WLS linear regression results were largely unaltered when the bin sizes were halved or doubled (not shown).




**Figure 9.** The relationship between dry season monthly isoprene data and broadleaf forest cover (BFC) for 2012-2016 with the best fit ordinary least square (OLS) regression line (a). Dry season isoprene data for the same time period binned based on 10% broadleaf forest cover intervals with weighted least squares (WLS) regression results. The errorbars show the standard error for each land cover bin (b). On panel (a) and (b) the text gives the best fit linear regression equations.

#### 415 4 Discussion and conclusions

This study used new and complementary satellite observations to investigate the relationships between land cover, LAI, and burned area, and five trace gases (isoprene, methanol, HCHO, CO, NO<sub>2</sub>) and AOD in the southern Amazon, with the aim to understand the influence of vegetation and fire on atmospheric composition in a region of land cover change.

Biomass burning is identified as a potential driver of year-to-year variability in several trace gases: CO, HCHO and NO<sub>2</sub>, as well as aerosols, as suggested by the AOD record. Previous studies have identified drought years in 2005, 2007, 2010, 2012 and 2015 for this region (Pope et al., 2020; Panisset et al., 2018; Reddington et al., 2015), which may increase burning and modify the relationship between burned area and emitted trace gases and aerosols. Although 2007 and 2010 are clearly elevated for burned area, AOD, CO, NO<sub>2</sub> and HCHO, not all of the listed drought years show maxima for a given variable in this study.


Pyrogenic sources in the dry season may further drive the seasonal cycle maxima of most of the atmospheric constituents over the region. The observed mean monthly isoprene seasonal cycle differs from the other constituents with a likely vegetation-driven secondary peak in the wet season. The isoprene seasonality is consistent with greater photosynthetically active radiation



**Figure 10.** The distribution of monthly dry season tropospheric NO<sub>2</sub> columns against monthly dry season burned area % coloured by broadleaf forest cover (a) and LAI (b) for 2005-2019. The panels display the OLS regression line and associated R<sup>2</sup> value, as included in Table 3.

(PAR) in the dry season, as well as potentially greater isoprene emissions in response to water stress (Yáñez-Serrano et al., 2020). The isoprene minimum observed here in June has been previously linked to new leaf growth in the tropical forest in the transition between the wet and dry seasons, as young leaves start producing isoprene only after a few weeks (Barkley et al., 430 2009). The dry season peak for HCHO in September is also consistent with a previous satellite study; however, a secondary wet season peak was found in that work (Barkley et al., 2009). A weaker relationship between isoprene and HCHO is suggested for this region, since HCHO does not appear to respond to the variations in isoprene at the end of the wet season.

The spatial distributions across the southern Amazon further highlight the co-location of isoprene with broadleaf forest cover and high LAI, and NO<sub>2</sub> with burned area. Most of the other atmospheric constituents show characteristics of both distributions. 435 Methanol is unusual, due to a greater region of elevated column values in the N and NE of the region.



**Figure 11.** Natural logarithm of the monthly dry season NO<sub>2</sub> data ( $\log_e(\text{NO}_2)$ ) against the natural logarithm of the monthly dry season burned area ( $\log_e(\text{BA})$ ) for 2005-2016 with the results of OLS regression (a). The WLS regression analysis on the natural logarithm of monthly dry season NO<sub>2</sub> data binned based on the natural logarithm of monthly burned area percentage cover for the same time period (b). The errorbars show the standard error for each burned area bin. On panel (a) and (b) the text shows the best fit equation found by the regressions and the respective  $R^2$  value.

Broadleaf forest cover (LAI) explains 59% (54%) of the variation in total column isoprene across the study region for the dry season over 2012-2019, which increases to 97% when the isoprene data is binned based on 10% forest cover bins. In the dry season, isoprene column amounts increase linearly with tree cover. For every 10% increase in broadleaf forest cover the isoprene total column amount increases by  $11 \times 10^{15}$  molecules  $\text{cm}^{-2}$ , or around 40% of the average background (0% broadleaf forest cover) total column isoprene. This result is consistent for both the binned and original data.

In contrast, burned area explains 25% of the variability in tropospheric column NO<sub>2</sub> for the dry season. 32% of the NO<sub>2</sub> variability can be explained when the natural logarithms of each variable are used, and the relationship is well-captured when this data is binned at 0.25  $\log_e(\text{BA})$  intervals ( $R^2 = 0.90$ ). There is a stronger increase in the trace gas at lower burned area values, which is captured by the fifth root power law. Additionally, the NO<sub>2</sub> amount varies depending on burning location, with greater values of NO<sub>2</sub> for an equivalent burned area in regions with at least 50% broadleaf forest cover. These results are in agreement with the FINNv2.5 emissions inventory, which has larger biomass burning emission factors for tropical forest



compared to savanna/grassland (Wiedinmyer et al., 2023). However, the burned areas can be greater where tree cover is more sparse, highlighting the potential for substantial pyrogenic emissions from both forest and savanna/grassland regimes.

450 In the wet season low tropospheric column  $\text{NO}_2$  values over the forested region and higher values in the SE could be driven by the forest canopy acting as an  $\text{NO}_2$  sink through biological uptake, as suggested by Kang et al. (2023). Alternatively, a further  $\text{NO}_x$  emission source e.g. from soils associated with agricultural activities (Wong and Geddes, 2021) or long-range transport of anthropogenic emissions in the region of São Paulo (van der A et al., 2008), may influence  $\text{NO}_2$  concentrations.

455 The clear dry season relationships of isoprene with broadleaf forest cover and  $\text{NO}_2$  with burned area contrast with the mixed signals for the other atmospheric constituents. However, the combination of broadleaf forest cover and burned area can explain 23% of the variation in dry season total column HCHO, suggesting interactions between pyrogenic and biogenic emissions of HCHO and its precursors. The moderate correlation values of AOD and CO with the land cover variables suggest some influence of vegetation through a biogenic source that yields CO and SOA formation in August-October, although forest emissions are most relevant in the wet season (Yáñez-Serrano et al., 2020; Artaxo et al., 2022). Methanol does not exhibit a strong relationship with any land surface variable despite some observed similarities.

460 A key difference between methanol, AOD, CO, and isoprene, HCHO and  $\text{NO}_2$  is their lifetime. While isoprene, HCHO and  $\text{NO}_2$  have lifetimes of up to a day (Pacífico et al., 2009; Wells et al., 2020; Pommier, 2023; Jacob, 1999), aerosols, methanol and CO atmospheric lifetimes range from several days to months (Bates et al., 2021; Hodzic et al., 2016; Holloway et al., 2000). The extended time period the aerosols, methanol and CO are present in the atmosphere will increase the role of transport in the observed distribution, resulting in a less clear local source signal. These longer lived species, particularly CO and aerosols, can therefore be transported from more distant anthropogenic sources (e.g. Park et al. (2015); Wang et al. (2015)). However, anthropogenic emissions are thought to be minor compared to pyrogenic and biogenic sources in the study region. Anthropogenic sources are in the SE of the study area and beyond the region of interest, e.g. the large agglomerations of São Paulo and Rio de Janeiro located further to the SE (see e.g. the European Commission EDGAR v6.1 Global Air Pollution Emissions database, last access: December 2023).

470 The findings confirm the tropical broadleaf forest, as opposed to other vegetation types, as the dominant source of isoprene in the region.  $\text{NO}_2$  is predominantly driven by pyrogenic emissions in the dry season, although the land cover type modulates the emission amount. HCHO, and to a lesser extent CO and aerosols, is linked to both biogenic and pyrogenic drivers. More specific land cover categories, and/or a consideration of other factors are necessary to identify the potential biogenic sources of methanol. The study finds both land cover and fire have significant impacts on regional atmospheric composition in the southern Amazon, including modifying amounts of trace gases and aerosols that have implications for regional air quality. Therefore, having established the importance of vegetation and fire activity on South American atmospheric composition, future work could exploit these relationships for Earth System Model (ESM) evaluation, and by using ESMs to explore the underpinning processes and potential feedbacks between the biosphere and atmosphere.





*Code availability.* The code used to calculate statistics on the data and produce the plots is shared in the github repository:

480 "[https://github.com/sands-eg/SouthernAmazon\\_figures](https://github.com/sands-eg/SouthernAmazon_figures)"

*Author contributions.* RP, RMD, FOC, and ES designed the research study. ES and RP analysed the satellite data supported by advice from and constructive discussion with RMD, FOC, CW, and HP. ES prepared the manuscript with scientific, editorial and technical input from all coauthors.

*Competing interests.* At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics.

485 *Acknowledgements.* This research has been supported by the Natural Environment Research Council (NERC) through a SENSE CDT studentship (grant no. NE/T00939X/1) and through funding for the National Centre for Earth Observation (NCEO, award reference NE/R016518/1). This work used JASMIN, the UK's collaborative data analysis environment (<https://jasmin.ac.uk>). For the methanol data, RAL's NRT system processes Eumetsat Level-1 data from MetOp-B IASI, MHS, AMSU, and GOME-2 and uses ECMWF meteorological forecast data, all processed on RAL's Jasmin infrastructure. We acknowledge the free use of tropospheric NO<sub>2</sub> column data from the OMI sensor from  
490 [www.temis.nl](http://www.temis.nl). The isoprene data was provided courtesy of Kelley Wells and others at the University of Minnesota. We acknowledge the advice and support from Brian Kerridge, Richard Siddans, Lucy Ventress and Barry Latter from the Rutherford Appleton Laboratory when using the IASI methanol data in this study



## References

- 495 Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmospheric Chemistry and Physics*, 11, 4039–4072, <https://doi.org/10.5194/acp-11-4039-2011>, 2011.
- Arnth, A., Schurgers, G., Lathiere, J., Duhl, T., Beerling, D. J., Hewitt, C. N., Martin, M., and Guenther, A.: Global terrestrial isoprene emission models: sensitivity to variability in climate and vegetation, *Atmospheric Chemistry and Physics*, 11, 8037–8052, <https://doi.org/10.5194/acp-11-8037-2011>, 2011.
- 500 Artaxo, P., Rizzo, L. V., Brito, J. F., Barbosa, H. M. J., Arana, A., Sena, E. T., Cirino, G. G., Bastos, W., Martin, S. T., and Andreae, M. O.: Atmospheric aerosols in Amazonia and land use change: from natural biogenic to biomass burning conditions, *Faraday Discuss.*, 165, 203–235, <https://doi.org/10.1039/C3FD00052D>, publisher: The Royal Society of Chemistry, 2013.
- 505 Artaxo, P., Hansson, H.-C., Andreae, M. O., Bäck, J., Alves, E. G., Barbosa, H. M. J., Bender, F., Bourtsoukidis, E., Carbone, S., Chi, J., Decesari, S., Després, V. R., Ditas, F., Ezhova, E., Fuzzi, S., Hasselquist, N. J., Heintzenberg, J., Holanda, B. A., Guenther, A., Hakola, H., Heikkinen, L., Kerminen, V.-M., Kontkanen, J., Krejci, R., Kulmala, M., Lavric, J. V., de Leeuw, G., Lehtipalo, K., Machado, L. A. T., McFiggans, G., Franco, M. A. M., Meller, B. B., Morais, F. G., Mohr, C., Morgan, W., Nilsson, M. B., Peichl, M., Petäjä, T., Praß, M., Pöhlker, C., Pöhlker, M. L., Pöschl, U., Von Randow, C., Riipinen, I., Rinne, J., Rizzo, L. V., Rosenfeld, D., Silva Dias, M. A. F., Sogacheva, L., Stier, P., Swietlicki, E., Sörgel, M., Tunved, P., Virkkula, A., Wang, J., Weber, B., Yáñez-Serrano, A. M., Zieger, P., Mikhailov, E., Smith, J. N., and Kesselmeier, J.: Tropical and Boreal Forest – Atmosphere Interactions: A Review, *Tellus B: Chemical and Physical Meteorology*, <https://doi.org/10.16993/tellusb.34>, 2022.
- 510 Badr, O. and Probert, S. D.: Sources of atmospheric carbon monoxide, *Applied Energy*, 49, 145–195, [https://doi.org/10.1016/0306-2619\(94\)90036-1](https://doi.org/10.1016/0306-2619(94)90036-1), 1994.
- Barkley, M. P., Palmer, P. I., De Smedt, I., Karl, T., Guenther, A., and Van Roozendaal, M.: Regulated large-scale annual shutdown of Amazonian isoprene emissions?, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2008GL036843>, 2009.
- 515 Bates, K. H., Jacob, D. J., Wang, S., Hornbrook, R. S., Apel, E. C., Kim, M. J., Millet, D. B., Wells, K. C., Chen, X., Brewer, J. F., Ray, E. A., Commane, R., Diskin, G. S., and Wofsy, S. C.: The Global Budget of Atmospheric Methanol: New Constraints on Secondary, Oceanic, and Terrestrial Sources, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033439, <https://doi.org/10.1029/2020JD033439>, 2021.
- Bela, M. M., Longo, K. M., Freitas, S. R., Moreira, D. S., Beck, V., Wofsy, S. C., Gerbig, C., Wiedemann, K., Andreae, M. O., and Artaxo, P.: 520 Ozone production and transport over the Amazon Basin during the dry-to-wet and wet-to-dry transition seasons, *Atmospheric Chemistry and Physics*, 15, 757–782, <https://doi.org/10.5194/acp-15-757-2015>, 2015.
- Boersma, K. F., Eskes, H. J., Dirksen, R. J., van der A, R. J., Veefkind, J. P., Stammes, P., Huijnen, V., Kleipool, Q. L., Sneep, M., Claas, J., Leitão, J., Richter, A., Zhou, Y., and Brunner, D.: An improved tropospheric NO<sub>2</sub> column retrieval algorithm for the Ozone Monitoring Instrument, *Atmospheric Measurement Techniques*, 4, 1905–1928, <https://doi.org/10.5194/amt-4-1905-2011>, 2011.
- 525 Cao, Y., Yue, X., Lei, Y., Zhou, H., Liao, H., Song, Y., Bai, J., Yang, Y., Chen, L., Zhu, J., Ma, Y., and Tian, C.: Identifying the Drivers of Modeling Uncertainties in Isoprene Emissions: Schemes Versus Meteorological Forcings, *Journal of Geophysical Research: Atmospheres*, 126, <https://doi.org/10.1029/2020JD034242>, 2021.
- Chance, K.: OMI/Aura Formaldehyde (HCHO) Total Column 1-orbit L2 Swath 13x24 km V003, <https://doi.org/10.5067/Aura/OMI/DATA2015>, 2007.



- 530 Chen, Y., Hall, J., van Wees, D., Andela, N., Hantson, S., Giglio, L., van der Werf, G. R., Morton, D. C., and Randerson, J. T.: Global Fire Emissions Database (GFED5), <https://doi.org/10.5281/zenodo.7668424>, 2023.
- Ciccio, P., Centritto, M., and Loreto, F.: Biogenic volatile organic compound emissions from vegetation fires, *Plant, Cell & Environment*, 37, 1810–1825, <https://doi.org/10.1111/pce.12336>, 2014.
- Danielson, J. and Gesch, D.: Global Multi-resolution Terrain Elevation Data 2010 (GMTED2010), Tech. rep., US Geological Survey, 2011.
- 535 European Commission: EDGARv6.1, [https://edgar.jrc.ec.europa.eu/index.php/dataset\\_ap61](https://edgar.jrc.ec.europa.eu/index.php/dataset_ap61), accessed 2023-12-22.
- Fang, H., Jiang, C., Li, W., Wei, S., Baret, F., Chen, J. M., Garcia-Haro, J., Liang, S., Liu, R., Myneni, R. B., Pinty, B., Xiao, Z., and Zhu, Z.: Characterization and intercomparison of global moderate resolution leaf area index (LAI) products: Analysis of climatologies and theoretical uncertainties, *Journal of Geophysical Research: Biogeosciences*, 118, 529–548, <https://doi.org/https://doi.org/10.1002/jgrg.20051>, 2013.
- 540 Fernandes, R. and Leblanc, S. G.: Parametric (modified least squares) and non-parametric (Theil–Sen) linear regressions for predicting biophysical parameters in the presence of measurement errors, *Remote Sensing of Environment*, 95, 303–316, <https://doi.org/10.1016/j.rse.2005.01.005>, 2005.
- Fowler, D., Pilegaard, K., Sutton, M. A., Ambus, P., Raivonen, M., Duyzer, J., Simpson, D., Fagerli, H., Fuzzi, S., Schjoerring, J. K., Granier, C., Neftel, A., Isaksen, I. S. A., Laj, P., Maione, M., Monks, P. S., Burkhardt, J., Daemmgen, U., Neiryck, J., Personne, E., Wichink-Kruit, R., Butterbach-Bahl, K., Flechard, C., Tuovinen, J. P., Coyle, M., Gerosa, G., Loubet, B., Altimir, N., Gruenhage, L., Ammann, C., Cieslik, S., Paoletti, E., Mikkelsen, T. N., Ro-Poulsen, H., Cellier, P., Cape, J. N., Horváth, L., Loreto, F., Niinemets, Ü., Palmer, P. I., Rinne, J., Mitztal, P., Nemitz, E., Nilsson, D., Pryor, S., Gallagher, M. W., Vesala, T., Skiba, U., Brüggemann, N., Zechmeister-Boltenstern, S., Williams, J., O’Dowd, C., Facchini, M. C., Leeuw, G. d., Flossman, A., Chaumerliac, N., and Erisman, J. W.: Atmospheric composition change: Ecosystems–Atmosphere interactions, *Atmospheric Environment*, 43, 5193–5267, <https://doi.org/10.1016/j.atmosenv.2009.07.068>, 2009.
- 545 Freitas, A. D. and Fornaro, A.: Atmospheric Formaldehyde Monitored by TROPOMI Satellite Instrument throughout 2020 over Sao Paulo State, Brazil, *Remote Sensing*, 14, <https://doi.org/10.3390/rs14133032>, 2022.
- Friedl, M. and Sulla-Menashe, D.: MCD12C1 MODIS/Terra+Aqua Land Cover Type Yearly L3 Global 0.05Deg CMG V006, <https://doi.org/10.5067/MODIS/MCD12C1.006>, 2015.
- Fu, D., Millet, D. B., Wells, K. C., Payne, V. H., Yu, S., Guenther, A., and Eldering, A.: Direct retrieval of isoprene from satellite-based infrared measurements, *Nature Communications*, 10, <https://doi.org/10.1038/s41467-019-11835-0>, 2019.
- Giglio, L., Randerson, J. T., and van der Werf, G. R.: Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4), *Journal of Geophysical Research: Biogeosciences*, 118, 317–328, <https://doi.org/10.1002/jgrg.20042>, 2013.
- Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., and Moore, R.: Google Earth Engine: Planetary-scale geospatial analysis for everyone, *Remote Sensing of Environment*, <https://doi.org/10.1016/j.rse.2017.06.031>, 2017.
- 560 Gu, D., Guenther, A. B., Shilling, J. E., Yu, H., Huang, M., Zhao, C., Yang, Q., Martin, S. T., Artaxo, P., Kim, S., Seco, R., Stavrakou, T., Longo, K. M., Tóta, J., de Souza, R. A. F., Vega, O., Liu, Y., Shrivastava, M., Alves, E. G., Santos, F. C., Leng, G., and Hu, Z.: Airborne observations reveal elevational gradient in tropical forest isoprene emissions, *Nature Communications*, 8, 15541, <https://doi.org/10.1038/ncomms15541>, 2017.
- 565 Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W. A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: A global model of natural volatile organic compound emissions, *Journal of Geophysical Research: Atmospheres*, 100, 8873–8892, <https://doi.org/10.1029/94JD02950>, 1995.



- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geoscientific Model Development*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.
- 570 Heald, C. L. and Spracklen, D. V.: Land Use Change Impacts on Air Quality and Climate, *Chemical Reviews*, 115, 4476–4496, <https://doi.org/10.1021/cr500446g>, 2015.
- Helas, G., Lobert, J., Scharffe, D., Schäfer, L., Goldammer, J., Baudet, J., Ahoua, B., Ajavon, A.-L., Lacaux, J.-P., Delmas, R., and Andreae, M. O.: Ozone production due to emissions from vegetation burning, *Journal of Atmospheric Chemistry*, 22, 163–174, 575 <https://doi.org/10.1007/BF00708187>, 1995.
- Hodzic, A., Kasibhatla, P. S., Jo, D. S., Cappa, C. D., Jimenez, J. L., Madronich, S., and Park, R. J.: Rethinking the global secondary organic aerosol (SOA) budget: stronger production, faster removal, shorter lifetime, *Atmospheric Chemistry and Physics*, 16, 7917–7941, <https://doi.org/10.5194/acp-16-7917-2016>, 2016.
- Holloway, T., Levy II, H., and Kasibhatla, P.: Global distribution of carbon monoxide, *Journal of Geophysical Research: Atmospheres*, 105, 580 12 123–12 147, <https://doi.org/10.1029/1999JD901173>, 2000.
- Jacob, D. J.: Introduction to Atmospheric Chemistry, Princeton University Press, <http://www.jstor.org.ezproxy.is.ed.ac.uk/stable/j.ctt7t8hg>, 1999.
- Kang, R., Huang, K., Gao, T., Mulder, J., Duan, L., Wang, C., Ke, P., Yao, M., Su, C., Li, J., Zhu, W., Zhu, J., and Fang, Y.: Forest Canopy Acts as an Atmospheric NO<sub>x</sub> Sink: Results From Micrometeorological Flux Measurements, *Journal of Geophysical Research: Atmospheres*, 585 128, e2022JD037 729, <https://doi.org/10.1029/2022JD037729>, 2023.
- Kefauver, S. C., Filella, I., and Peñuelas, J.: Remote Sensing of atmospheric biogenic volatile organic compounds (BVOCs) via satellite-based formaldehyde vertical column assessments, *International Journal of Remote Sensing*, 35, 7519–7542, <https://doi.org/10.1080/01431161.2014.968690>, 2014.
- Kim, K.-H., Kabir, E., and Kabir, S.: A review on the human health impact of airborne particulate matter, *Environment International*, 74, 590 136–143, <https://doi.org/10.1016/j.envint.2014.10.005>, 2015.
- Kuhn, U., Ganzeveld, L., Thielmann, A., Dindorf, T., Schebeske, G., Welling, M., Sciare, J., Roberts, G., Meixner, F. X., Kesselmeier, J., Lelieveld, J., Kolle, O., Ciccioli, P., Lloyd, J., Trentmann, J., Artaxo, P., and Andreae, M. O.: Impact of Manaus City on the Amazon Green Ocean atmosphere: ozone production, precursor sensitivity and aerosol load, *Atmospheric Chemistry and Physics*, 10, 9251–9282, <https://doi.org/10.5194/acp-10-9251-2010>, 2010.
- 595 Langford, B., House, E., Valach, A., Hewitt, C. N., Artaxo, P., Barkley, M. P., Brito, J., Carnell, E., Davison, B., MacKenzie, A. R., Marais, E. A., Newland, M. J., Rickard, A. R., Shaw, M. D., Yáñez-Serrano, A. M., and Nemitz, E.: Seasonality of isoprene emissions and oxidation products above the remote Amazon, *Environ. Sci.: Atmos.*, 2, 230–240, <https://doi.org/10.1039/D1EA00057H>, publisher: RSC, 2022.
- Loreto, F. and Fares, S.: Biogenic Volatile Organic Compounds and Their Impacts on Biosphere–Atmosphere Interactions, in: *Developments in Environmental Science*, vol. 13, pp. 57–75, Elsevier, <https://doi.org/10.1016/B978-0-08-098349-3.00004-9>, 2013.
- 600 Loveland, T. R. and Belward, A. S.: The IGBP-DIS global 1km land cover data set, DISCover: First results, *International Journal of Remote Sensing*, 18, 3289–3295, <https://doi.org/10.1080/014311697217099>, 1997.
- Marais, E. A., Jacob, D. J., Kurosu, T. P., Chance, K., Murphy, J. G., Reeves, C., Mills, G., Casadio, S., Millet, D. B., Barkley, M. P., Paulot, F., and Mao, J.: Isoprene emissions in Africa inferred from OMI observations of formaldehyde columns, *Atmospheric Chemistry and Physics*, 12, 6219–6235, <https://doi.org/10.5194/acp-12-6219-2012>, 2012.
- 605



- Millet, D. B., Jacob, D. J., Custer, T. G., de Gouw, J. A., Goldstein, A. H., Karl, T., Singh, H. B., Sive, B. C., Talbot, R. W., Warneke, C., and Williams, J.: New constraints on terrestrial and oceanic sources of atmospheric methanol, *Atmospheric Chemistry and Physics*, 8, 6887–6905, <https://doi.org/10.5194/acp-8-6887-2008>, 2008.
- Morellato, L. P. C., Camargo, M. G. G., and Gressler, E.: A Review of Plant Phenology in South and Central America, in: *Phenology: An Integrative Environmental Science*, edited by Schwartz, M. D., pp. 91–113, Springer Netherlands, Dordrecht, [https://doi.org/10.1007/978-94-007-6925-0\\_6](https://doi.org/10.1007/978-94-007-6925-0_6), 2013.
- Myneni, R., Knyazikhin, Y., and Park, T.: MODIS/Terra Leaf Area Index/FPAR 8-Day L4 Global 500m SIN Grid V061, 2021.
- Myneni, R. B., Yang, W., Nemani, R. R., Huete, A. R., Dickinson, R. E., Knyazikhin, Y., Didan, K., Fu, R., Juárez, R. I. N., Saatchi, S. S., Hashimoto, H., Ichii, K., Shabanov, N. V., Tan, B., Ratana, P., Privette, J. L., Morisette, J. T., Vermote, E. F., Roy, D. P., Wolfe, R. E., Friedl, M. A., Running, S. W., Votava, P., El-Saleous, N., Devadiga, S., Su, Y., and Salomonson, V. V.: Large seasonal swings in leaf area of Amazon rainforests, *Proceedings of the National Academy of Sciences*, 104, 4820–4823, <https://doi.org/10.1073/pnas.0611338104>, 2007.
- NASA/LARC/SD/ASDC: MOPITT Derived CO (Near and Thermal Infrared Radiances) V007, [https://doi.org/10.5067/TERRA/MOPITT/MOP02J\\_L2.007](https://doi.org/10.5067/TERRA/MOPITT/MOP02J_L2.007), 2000.
- Pacifico, F., Harrison, S. P., Jones, C. D., and Sitch, S.: Isoprene emissions and climate, *Atmospheric Environment*, 43, 6121–6135, <https://doi.org/10.1016/j.atmosenv.2009.09.002>, 2009.
- Pacifico, F., Harrison, S. P., Jones, C. D., Arneth, A., Sitch, S., Weedon, G. P., Barkley, M. P., Palmer, P. I., Serça, D., Potosnak, M., Fu, T.-M., Goldstein, A., Bai, J., and Schurgers, G.: Evaluation of a photosynthesis-based biogenic isoprene emission scheme in JULES and simulation of isoprene emissions under present-day climate conditions, *Atmospheric Chemistry and Physics*, 11, 4371–4389, <https://doi.org/10.5194/acp-11-4371-2011>, 2011.
- Pacifico, F., Folberth, G. A., Jones, C. D., Harrison, S. P., and Collins, W. J.: Sensitivity of biogenic isoprene emissions to past, present, and future environmental conditions and implications for atmospheric chemistry, *JOURNAL OF GEOPHYSICAL RESEARCH-ATMOSPHERES*, 117, <https://doi.org/10.1029/2012JD018276>, 2012.
- Palmer, P. I., Abbot, D. S., Fu, T.-M., Jacob, D. J., Chance, K., Kurosu, T. P., Guenther, A., Wiedinmyer, C., Stanton, J. C., Pilling, M. J., Pressley, S. N., Lamb, B., and Sumner, A. L.: Quantifying the seasonal and interannual variability of North American isoprene emissions using satellite observations of the formaldehyde column, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006689>, 2006.
- Palmer, P. I., Barkley, M. P., Kurosu, T. P., Lewis, A. C., Saxton, J. E., Chance, K., and Gatti, L. V.: Interpreting satellite column observations of formaldehyde over tropical South America, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 365, 1741–1751, <https://doi.org/10.1098/rsta.2007.2042>, 2007.
- Palmer, P. I., Marvin, M. R., Siddans, R., Kerridge, B. J., and Moore, D. P.: Nocturnal survival of isoprene linked to formation of upper tropospheric organic aerosol, *Science*, 375, 562–566, <https://doi.org/10.1126/science.abg4506>, 2022.
- Panisset, J. S., Libonati, R., Gouveia, C. M. P., Machado-Silva, F., Franca, D. A., Franca, J. R. A., and Peres, L. F.: Contrasting patterns of the extreme drought episodes of 2005, 2010 and 2015 in the Amazon Basin, *International Journal of Climatology*, 38, 1096–1104, <https://doi.org/10.1002/joc.5224>, 2018.
- Park, K., Emmons, L. K., Wang, Z., and Mak, J. E.: Joint Application of Concentration and  $\delta^{18}\text{O}$  to Investigate the Global Atmospheric CO Budget, *Atmosphere*, 6, 547–578, <https://doi.org/10.3390/atmos6050547>, 2015.



- Paton-Walsh, C., Emmerson, K. M., Garland, R. M., Keywood, M., Hoelzemann, J. J., Huneus, N., Buchholz, R. R., Humphries, R. S., Altieri, K., Schmale, J., Wilson, S. R., Labuschagne, C., Kalisa, E., Fisher, J. A., Deutscher, N. M., van Zyl, P. G., Beukes, J. P., Joubert, W., Martin, L., Mkololo, T., Barbosa, C., de Fatima Andrade, M., Schofield, R., Mallet, M. D., Harvey, M. J., Formenti, P., Piketh, S. J., and Olivares, G.: Key challenges for tropospheric chemistry in the Southern Hemisphere, *Elementa: Science of the Anthropocene*, 10, 00050, <https://doi.org/10.1525/elementa.2021.00050>, 2022.
- Platnick, S., e. a.: MODIS Atmosphere L3 Monthly Product., [https://doi.org/10.5067/MODIS/MOD08\\_M3.061](https://doi.org/10.5067/MODIS/MOD08_M3.061), goddard Space Flight Center, USA, 2015.
- Pommier, M.: Estimations of NO<sub>x</sub> emissions, NO<sub>2</sub> lifetime and their temporal variation over three British urbanised regions in 2019 using TROPOMI NO<sub>2</sub> observations, *Environ. Sci.: Atmos.*, 3, 408–421, <https://doi.org/10.1039/D2EA00086E>, publisher: RSC, 2023.
- Pope, R. J., Arnold, S. R., Chipperfield, M. P., Reddington, C. L. S., Butt, E. W., Keslake, T. D., Feng, W., Latter, B. G., Kerridge, B. J., Siddans, R., Rizzo, L., Artaxo, P., Sadiq, M., and Tai, A. P. K.: Substantial Increases in Eastern Amazon and Cerrado Biomass Burning-Sourced Tropospheric Ozone, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2019GL084143>, 2020.
- Pope, R. J., Kerridge, B. J., Siddans, R., Latter, B. G., Chipperfield, M. P., Arnold, S. R., Ventress, L. J., Pimlott, M. A., Graham, A. M., Knappe, D. S., and Rigby, R.: Large Enhancements in Southern Hemisphere Satellite-Observed Trace Gases Due to the 2019/2020 Australian Wildfires, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD034892, <https://doi.org/10.1029/2021JD034892>, 2021.
- Reddington, C. L., Butt, E. W., Ridley, D. A., Artaxo, P., Morgan, W. T., Coe, H., and Spracklen, D. V.: Air quality and human health improvements from reductions in deforestation-related fire in Brazil, *Nature Geoscience*, 8, 768–771, <https://doi.org/10.1038/ngeo2535>, 2015.
- Ritchie, H. and Roser, M.: Forests and Deforestation, Our World in Data, <https://ourworldindata.org/forests-and-deforestation>, 2021.
- Santos, A. M. d., Silva, C. F. A. d., Junior, P. M. d. A., Rudke, A. P., and Melo, S. N. d.: Deforestation drivers in the Brazilian Amazon: assessing new spatial predictors, *Journal of Environmental Management*, 294, 113020, <https://doi.org/10.1016/j.jenvman.2021.113020>, 2021.
- Schreier, S. F., Richter, A., Kaiser, J. W., and Burrows, J. P.: The empirical relationship between satellite-derived tropospheric NO<sub>2</sub> and fire radiative power and possible implications for fire emission rates of NO<sub>x</sub>, *Atmospheric Chemistry and Physics*, 14, 2447–2466, <https://doi.org/10.5194/acp-14-2447-2014>, 2014.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics : From Air Pollution to Climate Change*, John Wiley & Sons, Incorporated, <https://ebookcentral.proquest.com/lib/ed/detail.action?docID=4462549>, 2016.
- Shrivastava, M., Cappa, C. D., Fan, J., Goldstein, A. H., Guenther, A. B., Jimenez, J. L., Kuang, C., Laskin, A., Martin, S. T., Ng, N. L., Petaja, T., Pierce, J. R., Rasch, P. J., Roldin, P., Seinfeld, J. H., Shilling, J., Smith, J. N., Thornton, J. A., Volkamer, R., Wang, J., Worsnop, D. R., Zaveri, R. A., Zelenyuk, A., and Zhang, Q.: Recent advances in understanding secondary organic aerosol: Implications for global climate forcing, *Reviews of Geophysics*, 55, 509–559, <https://doi.org/10.1002/2016RG000540>, 2017.
- Silva Junior, C. H. L., Pessôa, A. C. M., Carvalho, N. S., Reis, J. B. C., Anderson, L. O., and Aragão, L. E. O. C.: The Brazilian Amazon deforestation rate in 2020 is the greatest of the decade, *Nature Ecology & Evolution*, 5, 144–145, <https://doi.org/10.1038/s41559-020-01368-x>, 2021.
- Stavrou, T., Müller, J.-F., Bauwens, M., De Smedt, I., Van Roozendaal, M., De Mazière, M., Vigouroux, C., Hendrick, F., George, M., Clerbaux, C., Coheur, P.-F., and Guenther, A.: How consistent are top-down hydrocarbon emissions based on formaldehyde observations from GOME-2 and OMI?, *Atmospheric Chemistry and Physics*, 15, 11861–11884, <https://doi.org/10.5194/acp-15-11861-2015>, 2015.



- 680 Strada, S., Fernández-Martínez, M., Peñuelas, J., Bauwens, M., Stavrakou, T., Verger, A., and Giorgi, F.: Disentangling temperature and water stress contributions to trends in isoprene emissions using satellite observations of formaldehyde, 2005–2016, *Atmospheric Environment*, 295, 119530, <https://doi.org/10.1016/j.atmosenv.2022.119530>, 2023.
- Szopa, S., Naik, V., Adhikary, B., Artaxo, P., Berntsen, T., Collins, W. D., Fuzzi, S., Gallardo, L., Kiendler-Scharr, A., Klimont, Z., Liao, H., Unger, N., and Zanis, P.: Short-Lived Climate Forcers, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 817–922, Cambridge University Press, 2021.
- 685 van der A, R. J., Eskes, H. J., Boersma, K. F., van Noije, T. P. C., Van Roozendaal, M., De Smedt, I., Peters, D. H. M. U., and Meijer, E. W.: Trends, seasonal variability and dominant NO<sub>x</sub> source derived from a ten year record of NO<sub>2</sub> measured from space, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD009021>, 2008.
- 690 Wang, P., Holloway, T., Bindl, M., Harkey, M., and De Smedt, I.: Ambient Formaldehyde over the United States from Ground-Based (AQS) and Satellite (OMI) Observations, *Remote Sensing*, 14, <https://doi.org/10.3390/rs14092191>, 2022.
- Wang, Y., Jiang, J. H., and Su, H.: Atmospheric responses to the redistribution of anthropogenic aerosols, *Journal of Geophysical Research: Atmospheres*, 120, 9625–9641, <https://doi.org/10.1002/2015JD023665>, 2015.
- Weber, J., King, J. A., Sindelarova, K., and Val Martin, M.: Updated isoprene and terpene emission factors for the Interactive BVOC (iBVOC) emission scheme in the United Kingdom Earth System Model (UKESM1.0), *Geoscientific Model Development*, 16, 3083–3101, <https://doi.org/10.5194/gmd-16-3083-2023>, 2023.
- 695 Wells, K. C., Millet, D. B., Payne, V. H., Deventer, M. J., Bates, K. H., de Gouw, J. A., Graus, M., Warneke, C., Wisthaler, A., and Fuentes, J. D.: Satellite isoprene retrievals constrain emissions and atmospheric oxidation, *NATURE*, 585, 225+, <https://doi.org/10.1038/s41586-020-2664-3>, 2020.
- 700 Wells, K. C., Millet, D. B., Payne, V. H., Vigouroux, C., Aquino, C. A. B., De Mazière, M., de Gouw, J. A., Graus, M., Kurosu, T., Warneke, C., and Wisthaler, A.: Next-Generation Isoprene Measurements From Space: Detecting Daily Variability at High Resolution, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD036181, <https://doi.org/10.1029/2021JD036181>, 2022.
- Wiedinmyer, C., Kimura, Y., McDonald-Buller, E. C., Emmons, L. K., Buchholz, R. R., Tang, W., Seto, K., Joseph, M. B., Barsanti, K. C., Carlton, A. G., and Yokelson, R.: The Fire Inventory from NCAR version 2.5: an updated global fire emissions model for climate and chemistry applications, *Geoscientific Model Development*, 16, 3873–3891, <https://doi.org/10.5194/gmd-16-3873-2023>, 2023.
- 705 Wohlfahrt, G., Amelynck, C., Ammann, C., Arneth, A., Bamberger, I., Goldstein, A. H., Gu, L., Guenther, A., Hansel, A., Heinesch, B., Holst, T., Hörtnagl, L., Karl, T., Laffineur, Q., Nefel, A., McKinney, K., Munger, J. W., Pallardy, S. G., Schade, G. W., Seco, R., and Schoon, N.: An ecosystem-scale perspective of the net land methanol flux: synthesis of micrometeorological flux measurements, *Atmospheric Chemistry and Physics*, 15, 7413–7427, <https://doi.org/10.5194/acp-15-7413-2015>, 2015.
- 710 Wong, A. Y. H. and Geddes, J. A.: Examining the competing effects of contemporary land management vs. land cover changes on global air quality, *Atmospheric Chemistry and Physics*, 21, 16479–16497, <https://doi.org/10.5194/acp-21-16479-2021>, 2021.
- Yáñez-Serrano, A. M., Bourtsoukidis, E., Alves, E. G., Bauwens, M., Stavrakou, T., Llusà, J., Filella, I., Guenther, A., Williams, J., Artaxo, P., Sindelarova, K., Doubalova, J., Kesselmeier, J., and Peñuelas, J.: Amazonian biogenic volatile organic compounds under global change, *Global Change Biology*, 26, 4722–4751, <https://doi.org/10.1111/gcb.15185>, 2020.