

The potential of drone observations to improve air quality predictions by 4D-var

Hassnae Erraji¹, Philipp Franke¹, Astrid Lampert², Tobias Schuldt¹, Ralf Tillmann¹, Andreas Wahner¹, and Anne Caroline Lange¹

¹Forschungszentrum Jülich GmbH, Institute of Energy and Climate Research - Troposphere (IEK-8), Jülich, Germany

²Institute of Flight Guidance, TU Braunschweig, Braunschweig, Germany

Correspondence: A. C. Lange (ann.lange@fz-juelich.de)

Abstract. Vertical profiles of atmospheric pollutants, acquired by unmanned aerial vehicles (UAVs, known as drones), represent a new type of observation that can help to fill the existing observation gap in the planetary boundary layer (PBL). This article presents the first study of assimilating air pollutant observations from drones to evaluate the impact on local air quality analysis. The study uses the high-resolution air quality model EURAD-IM (EUROpean Air pollution Dispersion – Inverse Model), including the four-dimensional variational data assimilation system (4D-var), to perform the assimilation of ozone (O₃) and nitrogen oxide (NO) vertical profiles. 4D-var ~~takes advantage of the inverse technique and is an inverse modelling technique that~~ allows for simultaneous adjustments of initial values and emissions rates. The drone data was collected during the MesSBAR (Automated airborne measurement of air pollution levels in the near earth atmosphere in urban areas) field campaign, which was conducted in Wesseling, Germany, on 22-23 September 2021. The results show that the 4D-var assimilation of high-resolution drone measurements has a beneficial impact on the representation of regional air pollutants within the model. On both days, a significant improvement in the vertical distribution of O₃ and NO is noticed in the analysis compared to the reference simulation without data assimilation. Moreover, the validation of the analysis against independent observations shows an overall improvement in the bias, root-mean-square error, and correlation for O₃, NO, and NO₂ (nitrogen dioxide) ground concentrations at the measurement site as well as in the surrounding region. Furthermore, the assimilation allows for the deduction of emission correction factors in the area nearby the measurement site, which significantly contribute to the improvement in the analysis.

1 Introduction

In response to the increasing need for high-resolution and accurate air-quality forecasts, extended efforts to improve the performance of chemical transport models (CTM) have been made over recent decades. One of the effective means of improvement involves the use of advanced data assimilation techniques (Elbern et al., 2007; Liu et al., 2017; Klonecki et al., 2012). The aim is to combine observations and model data to obtain a better representation of the pollutants in the atmosphere as well as to optimise the input parameters, such as emissions, when considering inverse models. Although data assimilation holds significant potential for enhancing air quality modelling, its application is often still limited due to the scarcity of available observational

data. In fact, the observational data types, which are usually used for assimilation (ground-based, airborne, and satellite obser-
25 vations), are certainly valuable for enhancing forecast accuracy, but they remain insufficient due to various constraints related
to their availability, resolution, and especially their limited vertical coverage. Ground-based observations are the major source
of information for regional CTMs and are generally taken from in-situ monitoring networks. Even if they are fairly dense in
the horizontal distribution on a regional scale, no information regarding the vertical distribution of air pollutants is provided. In
contrast, lidar (light detection and ranging) remote sensing instruments and in-situ sonde measurements can provide this infor-
30 mation, but unfortunately, only a sparse and limited number of such stations exists. Similarly, [airborne-ground-based Fourier
Transform InfraRed \(FTIR\) spectrometers, which form part of the Network for the Detection of Atmospheric Composition
Change \(NDACC\), are capable of retrieving vertically resolved mixing ratios of a range of atmospheric constituents. However,
the vertical resolution of these profiles is constrained by their dependence on a priori information, and the network's spatial
coverage remains sparse \(De Mazière et al., 2018; García et al., 2021\). Multi-axis differential optical absorption spectroscopy
\(MAX-DOAS\) is also capable of retrieving trace gas and aerosol vertical profiles \(Tirpitz et al., 2021\). Airborne observations](#)
(e.g., In-service Aircraft for a Global Observing System – IAGOS, or flight campaigns) provide [high-resolution](#) vertical pro-
files during take-off and landing; however, the spatial coverage is still limited because of the high costs (Wang et al., 2022;
Petetin et al., 2018; Tillmann et al., 2022). Satellite retrievals mainly provide the total column of air pollutants, thus providing
little information on the vertical distribution of the air pollutant concentrations in the planetary boundary layer (PBL) and at
40 the Earth's surface (Martin, 2008). Consequently, a significant observational gap exists in the PBL, which is the lowest part
of the atmosphere characterized by the highest concentrations of air pollutants due to its vicinity to anthropogenic emission
sources (Scheffe et al., 2009).

Unmanned Aerial Vehicles (UAVs), also known as drones, are comparatively new measurement platforms that have begun to
be widely utilized in recent years to obtain in-situ measurements of atmospheric trace gases and aerosols within the lower
45 atmosphere (Schuyler and Guzman, 2017; Yang et al., 2023), bringing many opportunities to improve air pollution monitoring.
The increase in drone applications comes mainly from their numerous advantages, such as portability and flexibility while
being affordable. In addition, they can provide in-situ observations of various atmospheric constituents with high temporal
and vertical resolution (Lawrence and Balsley, 2013). However, drone measurements come along with some limitations as, for
instance, flights are complicated during strong wind conditions, require good visibility, and are often restricted to maximum
50 altitudes due to aviation safety reasons. Nevertheless, they can fill the existing observational gap in the PBL and provide valu-
able information on the distribution of air pollutants.

Several studies present drone campaigns that observed the atmospheric composition and meteorological parameters during the
last two decades (Villa et al., 2016; Bretschneider et al., 2022). The measured data, mostly from the PBL region, were used
for research on the atmospheric boundary layer (Wang et al., 2021), pollutants variability and distribution (Altstädter et al.,
55 2015; Illingworth et al., 2014), as well as to study the properties of aerosols (Roberts et al., 2008; Corrigan et al., 2008), and
to qualify local emissions sources (Nathan et al., 2015). Furthermore, drone campaigns have been conducted in remote areas,
such as the Arctic and Antarctic regions (Lampert et al., 2020), as well as during volcano eruptions (Diaz et al., 2012).

To our knowledge, the assimilation of drone observations has only been tested in the context of Numerical Weather Prediction

(NWP) models (Flagg et al., 2018; Leuenberger et al., 2020), and no study has yet explored their impact in the case of chemical data assimilation. Meteorological studies have shown that the assimilation of meteorological drone data has a positive impact on improving weather forecasts. This has prompted further ongoing research regarding the possibility of implementing drone observations in support of operational meteorology forecasting and for real-time data assimilation studies (O’Sullivan et al., 2021). Impact studies have revealed a large improvement in the vertical distribution of temperature, relative humidity, and wind as well as a reduction of bias and root-mean-square error (RMSE) when drone observations are assimilated using a variational data assimilation system within high-resolution NWP models (Jonassen et al., 2012; Flagg et al., 2018; Jensen et al., 2021; Sun et al., 2020; Leuenberger et al., 2020).

Given the positive impact that has been reported in the case of meteorological applications, questions arise about the potential benefits and limitations of drone observations when assimilated within a CTM. In this study, the impact of drone data assimilation on air quality analyses is investigated using the regional and high-resolution EUROpean Air pollution Dispersion – Inverse Model (EURAD-IM) with its four-dimensional variational (4D-var) data assimilation system (Elbern et al. (2007)). Vertical profiles of ozone (O_3) and nitrogen oxide (NO) collected during the MesSBAR (Automatisierte luftgestützte Messung der Schadstoffbelastung in der erdnahen Atmosphäre in urbanen Räumen / Automated airborne measurement of air pollution levels in the near earth atmosphere in urban areas) field campaign are assimilated. The potential of drone observations to improve air quality analysis and forecast is explored in a two-day case study by applying the joint optimisation of initial values and emission rates. The aim is to investigate the ability of the 4D-var system to adjust local emission rates using vertical profiles that were collected in a region characterised by diverse emission sources. This paper is structured as follows: In Sect. 2, the EURAD-IM and its 4D-var data assimilation system are presented. The MesSBAR field campaign and the experimental design are described in Sect. 3. The results of the 4D-var data assimilation experiments are discussed in Sect. 4. Finally, the summary and conclusions are given in Sect. 5.

2 The modeling system

2.1 The EURAD-IM Model

EURAD-IM (EUROpean Air pollution Dispersion – Inverse Model) is a three-dimensional high resolution Eulerian CTM simulating air pollution in the troposphere at continental to regional scale. It has been used for several scientific studies for air quality forecasting, episode scenarios, data assimilation, and inverse modelling (Deroubaix et al., 2024; Gama et al., 2019; Elbern et al., 2007; Duarte et al., 2021; Franke et al., 2022, 2024). EURAD-IM is part of the regional Copernicus Atmosphere Monitoring Service (CAMS), providing daily air quality forecasts and reanalysis over Europe, which enable continuous quality assurance using observations and inter-model evaluation (Marécal et al., 2015).

Table 1 presents a summary of the specific model settings and modules utilized in the EURAD-IM configuration employed in this study. EURAD-IM describes the transport by diffusion and advection of various trace gas components emitted both by anthropogenic and biogenic sources and considers the gas-phase chemical transformation of about 110 chemical species with 265 reactions. The MADE (Modal Aerosol Dynamics model for Europe) module is employed to investigate aerosol dynamics

Table 1. Summary of EURAD-IM configuration.

	Processes	Modules & References
Transport	Advection	Walcek scheme (Walcek, 2000)
Gas-phase Chemistry	Kinetic Chemistry mechanism	RACM-MIM (Stockwell et al., 1997)
	Dry deposition	Zhang et al. (2003) scheme
	Wet deposition	Roselle and Binkowski (1999)
	Chemistry solver	KPP (Sandu and Sander, 2006)
Aerosols	Aerosol dynamics	MADE (Ackermann et al., 1998)
	Secondary Inorganic Aerosols	HDMR (Rabitz and Aliş (1999))
	Secondary Organic Aerosols	SORGAM (Schell et al., 2001)
Emissions	Biogenic emissions	MEGAN (Guenther et al., 2012)
	Anthropogenic emissions	TNO-UBA emission inventory (Kuenen et al., 2014)
Assimilation	4D-var system	Elbern et al. (2007)
	Minimisation algorithm	L-BFGS algorithm (Liu and Nocedal, 1989)
	Background error covariance modelling	Weaver and Courtier (2001)

within EURAD-IM, providing information on aerosol size distribution and chemical composition. This module simulates the formation and transformation of both primary and secondary aerosols, considering the interactions between the gas-phase and aerosols. EURAD-IM accounts for the loss of chemical components through wet and dry deposition, as well as aerosol sedi-
95 mentation. Moreover, EURAD-IM includes a 4D-var assimilation system, as described in the subsequent section, along with the adjoint code derived from the forward code detailed in Elbern et al. (2007). The adjoint model incorporates the transport, diffusion and gas transformation processes of the chemical species as well as secondary inorganic aerosol formation.

The CTM is driven by meteorological fields from the Weather Research and Forecasting model (WRF, version 3.7, Skamarock et al. (2008)) as thermo-dynamical forcing. The ECMWF (European Centre for Medium-Range Weather Forecasts) IFS (Inte-
100 grated Forecasting System) global analysis (ERA5) is used for initialization and boundary conditions for the WRF simulations. Chemical boundary conditions are generated by the CAMS global reanalysis data set (EAC4) that is produced by the ECMWF Composition Integrated Forecasting System (C-IFS). Anthropogenic emissions used for this study are provided by the German Environment Agency (Umweltbundesamt, UBA) for Germany and by the TNO-MACC-II inventory (Kuenen et al., 2014) for the rest of Europe. The emission data set is subject to processing in the EURAD Emission Module (EEM) (Memmesheimer
105 et al., 1995) for seasonal and diurnal redistribution, as well as attributions to working days and weekends. The emission data is divided into point and area sources. The data contains emissions of gaseous air pollutants, i.e., carbon monoxide (CO), nitrogen oxides (NO_x), sulfur dioxide (SO₂), total non-methane volatile organic compounds (NMVOC), and ammonia (NH₃), and for aerosols PM₁₀ (particulate matter with a diameter <10 µm) and PM_{2.5} (particulate matter with a diameter <2.5 µm) emis-
110 sions. Biogenic emissions are calculated online using the Model of Emissions of Gases and Aerosols from Nature (MEGAN), while wild fire emissions are not considered here and did not play a role in the investigated case.

2.2 4D-Var data assimilation

The EURAD-IM data assimilation system is based on the 4D-var method as described in Elbern and Schmidt (2001) and Elbern et al. (2007). The 4D-var approach aims to determine the optimal model state by combining the prior information (e.g., provided by a forecast) with observational data over an assimilation window through the minimization of the following cost function \mathbf{J} :

$$\begin{aligned} \mathbf{J}(\mathbf{x}_0, \mathbf{e}) &= \mathbf{J}_b(\mathbf{x}_0) + \mathbf{J}_o(\mathbf{x}_0) + \mathbf{J}_e(\mathbf{e}) \\ &= \frac{1}{2}(\mathbf{x}_0 - \mathbf{x}^b)^T \mathbf{B}^{-1}(\mathbf{x}_0 - \mathbf{x}^b) + \frac{1}{2} \sum_{i=0}^n ((\mathbf{y}_i - \mathbf{H}_i \mathbf{M}_i \mathbf{x}_0)^T \mathbf{R}_i^{-1} (\mathbf{y}_i - \mathbf{H}_i \mathbf{M}_i \mathbf{x}_0)) + \frac{1}{2}(\mathbf{e} - \mathbf{e}^b)^T \mathbf{K}^{-1}(\mathbf{e} - \mathbf{e}^b) \end{aligned} \quad (1)$$

Here, the optimisation is subject to the initial conditions \mathbf{x}_0 and the emission correction factor \mathbf{e} . The cost function equation includes an additional element (in contrast to the usual 4D-var used for NWP) that accounts for emissions ($\mathbf{J}_e(\mathbf{e})$). The model state is mapped from the model space to the observation space by the observation operator \mathbf{H}_i and the model operator \mathbf{M}_i , producing the model equivalents of each observation \mathbf{y}_i . The matrices \mathbf{B} , \mathbf{R} , and \mathbf{K} represent the error covariance matrices associated with the a-priori state vector \mathbf{x}^b , the observations \mathbf{y}_i , and a-priori emissions \mathbf{e}^b , respectively. The matrix \mathbf{R} considers only diagonal elements (i.e., it ignores any error correlation between different observations) while accounting for the uncertainties in the measurements and model representation error. The matrix \mathbf{B} is estimated using error variances and the diffusion operator proposed by Weaver and Courtier (2001). Thus, \mathbf{B} can be factorized as $\mathbf{B} = \mathbf{B}^{1/2} \mathbf{B}^{T/2}$ for the use in the preconditioning of the highly underdetermined data assimilation system. The matrix \mathbf{K} is ~~specified as a defined as~~ block diagonal, ~~as presented with non-zero entries for correlations between species and near-by emissions. The variance and correlation values are provided~~ in Paschalidi (2015). The minimization of the cost function \mathbf{J} is performed through an iterative process using the Quasi-Newton limited memory L-BFGS algorithm (Liu and Nocedal, 1989), which includes the iterative integration of the forward and adjoint EURAD-IM.

3 The MesSBAR campaign analysis

3.1 Air quality measurements

The MesSBAR field campaign took place near Wesseling, Germany, on 22-23 September 2021. During these two days, a multicopter system composed of a drone and a set of low-cost air quality monitoring instruments was used to carry out vertical profile measurements of air pollutants during the morning hours. Among the instruments loaded in the multicopter, electrochemical sensors were used to monitor nitrogen oxide (NO), and a Personal Ozone Monitor (POM) was ~~employed~~ deployed for assessing ozone (O_3) concentrations. The NO drone observations have an accuracy of 35 % at 40 ppb_v with a precision of ± 2.5 ppb_v (1 σ at 30 s time resolution). POM provides an accuracy of 1.5 ppb_v and a precision of 1.5 ppb_v (1 σ at 10 s time resolution) in the observed O_3 mixing ratio range. The feasibility of using these sensors for measurements in the planetary boundary layer was discussed in (Schuldt et al., 2023; Tillmann et al., 2022). A detailed description of the development,




Figure 1. Geographic map displaying the MesSBAR measurement location, air quality ground stations, and meteorological station situated near the A555 highway. (Source: © OpenStreetMap contributors 2023. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.)

technical characteristics, and calibration of the multicopter system can be found in Bretschneider et al. (2022). The campaign's basis was located within the proximity of the A555 highway, which is a much-frequented connection between the German cities of Cologne and Bonn. The measurements were conducted above agricultural land located about 1 km south of the town of Wesseling. The city centres of Cologne and Bonn are about 15 km north and 10 km south of the measurement location, respectively (Fig.1). The Wesseling region is located within the Rhineland chemical region and is widely recognized as a leading chemical hub in Europe. Wesseling, in particular, hosts a remarkable level of industrial activity attributed to the presence of major companies operating in the chemical and petroleum sectors (source: <https://www.chemcologne.de/en/investments/the-rhineland-chemical-region>, access date: 21 February 2024).

The objective of this campaign was to capture the early morning evolution of air pollutant concentrations with the development of the PBL. Furthermore, the proximity to the highway allows for measurements of pollutants specifically originating from traffic sources.

The drone is operated by an autopilot system that uses an inertial navigation solution with an Earth related position based on GNSS data (Global Navigation Satellite System). During the measurements, the autopilot controls a constant lateral position

and a constant vertical climb rate of approximately 1 ms^{-1} . Wind affects only the attitude of the copter, but given the low
155 wind situations during this campaign, the effect on the attitude can be neglected. The drone reached a maximum altitude of
350 m. This altitude limitation was imposed by air traffic restrictions in the area due to its proximity to the Cologne /Bonn
airport. During each drone flight, two profiles were acquired: ~~the ascent and the descent~~one ascending and one descending
were done in a short period of time. For the assimilation experiments ~~conducted~~carried out with EURAD-IM, only the ~~ascent~~
ascending profiles were utilized due to their higher accuracy (Schlerf et al., 2024). The measurements during the descending
160 flights are strongly influenced by the turbulence generated by the drone's propellers, which reduces the data quality. In this
study, the vertical profiles of O_3 and NO obtained from the multicopter are utilized and assimilated within EURAD-IM. The
vertical resolution of these profiles is approximately 10 m, with 254 data points assimilated on 22 September 2021 and 257 on
23 September 2021 for both O_3 and NO . Additionally, observations from two ground-based stations situated on both sides of
highway A555 (Fig. 1) are used to validate the simulation results. Furthermore, meteorological observations from an automatic
165 weather station, located approximately 1 km south-east of the measurement site, are employed for comparing meteorological
data, especially the wind fields.

3.2 Simulations setup

The objective of this study is to investigate the impact of O_3 and NO drone profile assimilation on the air quality analysis
using high-resolution EURAD-IM simulations. The model grid has a horizontal resolution of $5 \text{ km} \times 5 \text{ km}$ and is vertically
170 divided into 30 layers defined by terrain following sigma coordinates between the surface and 100 hPa, with about 19 layers
covering the lowest 1 km of the atmosphere. The EURAD-IM domain covers central Europe, including Germany with 271
 $\times 298$ grid points. The model output is adjusted to provide forecasts with a temporal resolution of 60 s, allowing for a more
precise comparison with the high-resolution drone observations. To assess the impact of drone data assimilation on air quality
forecast, simulations are conducted both with and without data assimilation (Table 2). The joint initial value/emission rate
175 optimisation mode of EURAD-IM is activated for this purpose. Two 24-hour experiments are performed without assimilation:
one on 22 September 2021, and the other on 23 September 2021. For these experiments, the model is initialized from a
climatological chemical state with a spin-up simulation of 6 days (16-21 September 2021) prior to the campaign dates in order
to establish a chemically balanced initial state. Moreover, two additional simulations focusing on O_3 and NO data assimilation
are performed for 24 hours on 22 and 23 September 2021. The assimilation window is deliberately selected to coincide with
180 the availability of observations, aiming to minimize computational time in the simulations while also ensuring a meaningful
lead time for emission optimisation. For drone data assimilation, the observation error is considered as the sum of measurement
and representativeness errors. The measurement error for O_3 is taken as the standard deviation of the measurements. For NO ,
the error is estimated according to (Elbern et al., 2007), by defining a relative error ϵ_{rel} and a minimal absolute error ϵ_{abs} :

$$\epsilon_{meas} = \max(\epsilon_{abs}, \epsilon_{rel} \cdot y) \quad (2)$$

185 The absolute error used for NO is 2 ppb_v , and the relative error is considered to be 20 % of the observed values.
The representation error is calculated by applying the corresponding formula from (Elbern et al., 2007), which consider the

grid cell spacing (dx), the representativeness length of the measurement location (L_x), and an absolute error specific to the measured species. The formula is expressed as

$$\epsilon_{rep} = \sqrt{\frac{dx}{L_x}} \times \epsilon_{abs}. \quad (3)$$

190 The grid cell spacing (dx) corresponds to the spatial resolution of the measurement grid, while the representativeness length (L_x) indicates the effective range over which the measurement is considered representative. In this case study, L_x is set to 3 km. The absolute error (ϵ_{abs}) varies by species: it is 2 ppb_v for O₃ and 3 ppb_v for NO. For the estimation of background errors, horizontal correlation lengths of 2.5 km, 10 km, and 20 km are employed at the surface, at the top of the planetary boundary layer, and at the upper model levels, respectively.

Table 2. Model simulations presented in this paper.

Experiment name	Assimilation	Period	Assimilation Window	Assimilated Observations
REF_22SEP	no	24-hours, 22 September 2021	-	-
REF_23SEP	no	24-hours, 23 September 2021	-	-
DA_22SEP	yes	24-hours, 22 September 2021	00-11 UTC	6 drone profiles of O ₃ and NO
DA_23SEP	yes	24-hours, 23 September 2021	00-09 UTC	5 drone profiles of O ₃ and NO

195 3.3 Evaluation of the wind situation

The wind is critical parameter that governs the dispersion of air pollutants and their transport, with a direct influence on emission optimisation within the framework of inverse CTMs. The wind conditions at the observation site are evaluated for two purposes: firstly, to validate the suitability of the measurement site location for measuring local traffic emissions, and secondly, to assess the horizontal wind for applications to emission optimisation.

200 Figure 2 shows the surface wind speed and direction observed by the nearby weather station observed during the flights' operation hours. The dominant wind direction is primarily from the south-east on 22 September 2021, with a maximum speed of 1.3 ms⁻¹, while it comes from the south to south-east in the morning hours of 23 September 2021, with a maximum recorded speed of 2.0 ms⁻¹. This indicates that the observation point is strategically located downwind of the nearest traffic emission source, which enabled the multicopter to successfully capture the emissions from the highway.

205 Apart from the surface conditions during the measuring period, the two days are each characterized by a distinct wind situation, as shown in the horizontal wind profiles extracted from the WRF simulations in Fig. 2. On 22 September 2021, the wind patterns exhibit vertical wind shear throughout the day and across all levels, changing direction from the southeast/east at lower altitudes to the west/northwest at higher altitudes. However, the wind intensity remains relatively low, measuring less than 3.0 ms⁻¹. On 23 September 2021, the surface wind direction aligns with the observations during the campaign period. Nevertheless, at higher
210 levels and beyond the campaign period, westerly and south-westerly winds dominate, and their speed increases with height. The maximum speed is reached at 450 m with 12.0 ms⁻¹ between 05 UTC and 07 UTC. The difference of the wind profiles between the two days may result in variations in the assimilation results, particularly with respect to emission optimisation.




Figure 2. Observed surface wind speed and direction during the measurement period on 22 September 2021 (upper left) and 23 September 2021 (upper right). Forecast of horizontal wind profiles for different hours for the lowest 500 m at the campaign location on 22 September 2021 (bottom left) and 23 September 2021 (bottom right).

4 Results

4.1 Impact on vertical profiles

215 In order to evaluate the impact of the drone data assimilation on the air pollutants' vertical distribution and given the lack of independent vertical profiles, the simulation results are first compared to the drone observations that are assimilated. Figure 3 presents the observed O_3 and NO drone profiles as well as vertical profiles resulting from the 4D-var assimilation and the reference simulations. For both days, the 4D-var analyses agree better with the drone observations in comparison to the refer-

ence forecast for both species, which indicates the successful assimilation of the drone observations. On 22 September 2021, an underestimation by the reference simulation is observed for the O_3 levels at altitudes above 200 m, with discrepancies reaching up to 15 ppb_v, especially for the first three flights (F1, F2, and F3). The assimilation of drone profiles significantly reduces this underestimation ~~by 45%. The bias was reduced by 98% on average (-4.58 ppb_v) for F1, 36% (-0.74 ppb_v) for F2, and 41% (-1.44 ppb_v) for F3, with an average reduction of 30% (-0.73 ppb_v) across all flights (Table 3).~~ On 23 September 2021, the reference model run overestimates ~~the O_3 concentration at the concentrations at both~~ ground and near-surface levels. The ~~largest overestimation is visible for most pronounced overestimations occur during~~ the first three flights of the day (F7, F8, and F9) ~~by about~~, with discrepancies reaching up to 20 ppb_v. ~~Due to Following~~ the 4D-var assimilation, the O_3 ground bias is reduced by more than ~~85~~82% ~~. Consequently, biases in concentration are reduced by nearly 30~~(-12.49 ppb_v) for F7, ~~56% on the first day and 55~~(-2.86 ppb_v) for F8, and 25% (-0.96 ppb_v) for F9. As a result, the overall O_3 bias on the second day ~~is reduced by approximately 55%~~ (Table-3.46 ppb_v) (Table 3).

On both days, the reference simulations underestimate the NO vertical distribution at all heights, with the strongest discrepancies at ground level. Improvement due to the assimilation is accomplished mostly at surface and near-surface levels for the initial three flights of each day (F1, F2, F3, F7, F8, and F9), with more pronounced adjustments on the second day ~~, resulting in a bias decrease of up to 84 at ground level. For higher levels at the ground level, while at higher levels during these same flights,~~ the impact of the assimilation is minimal to non-existent, for instance, for the flights F7 and F8 above 150 m. ~~Contrarily, because~~ Overall, bias reductions of 24% (6.78 ppb_v), 33% (11.61 ppb_v), and 23% (8.91 ppb_v) were observed for F1, F2, and F3, respectively. On the second day, greater improvements were achieved, with reductions of 30% (4.17 ppb_v) for F7, 49% (10.1 ppb_v) for F8, and 57% (15.29 ppb_v) for F9. Because the pollutant concentrations are well mixed in the PBL, a uniformly positive impact throughout the vertical can be seen in the NO analyses of the later flights of the day (F4, F5, F6, F10, and F11). The bias is reduced by 38% (-10.81 ppb_v) for F4, 54% (-15.26 ppb_v) for F5, and 49% (-14.66 ppb_v) for F6. On the following day, the bias reduction is smaller, with a 27% (-7.48 ppb_v) reduction for F10 and 18% (-5.58 ppb_v) for F11. Overall, the 4D-var assimilation of drone observations leads to a substantial reduction ~~of more than 35~~ in NO biases, with a 36% reduction (-11.34 ppb_v) on the first day and a 35% reduction (-8.52 ppb_v) on the second day, between the reference model forecast and observations ~~on both days~~ (Table (Table 3)).

These results highlight the successful assimilation of drone observations by the EURAD-IM 4D-var system. The accuracy of these findings is further examined and discussed in Sect. 4.3 through a validation process using independent observations.




Figure 3. The vertical profiles of O_3 and NO measured by the drone system (red line), compared to the 4D-var analysis (blue line) and the reference run (black line) for all flights on 22-23 September 2021. The red shading highlights the standard deviation of the drone observations.

Table 3. O₃ and NO biases (model value minus observation) in ppb_v for each flight.

Model runs	O ₃ Vertical Profiles							NO Vertical Profiles						
	F 1	F 2	F 3	F 4	F 5	F 6	Daily bias	F 1	F 2	F 3	F 4	F 5	F 6	Daily bias
REF_22SEP	-4.65	-2.06	-3.53	-1.23	-0.91	-2.49	2.48	-27.96	-35.39	-39.34	-28.21	-28.11	-30.09	31.52
DA_22SEP	0.07	-1.32	-2.09	-0.38	-2.42	-4.20	1.75	-21.18	-23.78	-30.43	-17.40	-12.85	-15.43	20.18
	F 7	F 8	F 9	F 10	F 11	Daily bias		F 7	F 8	F 9	F 10	F 11	Daily bias	
REF_23SEP	15.20	5.12	3.81	3.64	3.86	6.33		-13.95	-20.75	-26.65	-28.03	-30.88	24.05	
DA_23SEP	2.71	-2.26	-2.85	-3.92	-2.63	2.87		-9.78	-10.65	-11.37	-20.55	-25.30	15.53	

4.2 Emission optimisation

The ~~4d-var~~4D-var data assimilation method applied here aims at finding the best representation of the pollutants combining the knowledge provided by the EURAD-IM simulations and the drone O₃ and NO profile observations. The method relies on the assumption that the ~~biggest uncertainties of~~largest uncertainties in the modelled pollutant concentrations base on uncertainties ~~of in~~ initial values and emission rates. Emission correction factors for 25 anthropogenic pollutants can be deduced from the analysis. Consequently, it is worth looking at the emission factors being analysed to gain ~~first insights for a first insight into~~ the potential to retrieve detailed information about emission assessment applying this inverse modelling technique. However, their generalization and significance should be ~~rated carefully~~carefully evaluated, mainly because of the limited number of drone profiles being available ~~-, the distributed over the whole day, the resulting~~ short assimilation windows~~selected~~, and the ~~deficiency to perform~~lack of a long-term statistical analysis.

The assimilation experiments performed with the O₃ and NO drone observations result in significant corrections of NO and NO₂ emission rates in the grids surrounding the observation site. The resulting emissions factors, which represent the ratio between the optimised emission rates and the input emission rates for each species, have variability that ranges from 1 to 4 for NO and from 1 to 6 for NO₂ in the DA_22SEP experiment. In contrast, the variability extends from 1 to 14 for both NO and NO₂ in the DA_23SEP experiment (Fig. A1). This indicates that an increase in emissions is analysed in the studied region. Figure 4 (first row) displays the original daily NO_x emissions rates and the analysed emission changes on 22 and 23 September 2021. A significant increase of NO_x emissions is obtained in the DA_22SEP results, with changes in emission rates reaching up to 16 Mgd⁻¹ in the grid cells located north and northwest of the observation site.The emission of 16 Mgd⁻¹ represents approximately 3.46% of the total daily NO_x emissions in the analyzed region, where the total daily NO_x emission is about 462 Mgd⁻¹. For DA_23SEP in contrast, the emission rates increase by up to 10 Mgd⁻¹ in the grid cells surrounding the observation site. Based on the chemical coupling with NO and O₃, carbon monoxide (CO), sulfur dioxide (SO₂), and sulfate (SO₄) emissions are optimised resulting in emission correction factors between 1 and 3 (not shown).

To interpret the results and to investigate this discrepancy between the two days, the changes in NO_x emissions are evaluated according to the emission source sectors. Figure 4 additionally shows the original NO_x emissions and the analyzed emission changes for three dominant polluter sectors in this region: power production, industry, and road transport. The original emission data set includes in total 12 GNFR (gridded Nomenclature For Reporting) sectors, while only these three sectors are substantially affected in the analysis. The DA_22SEP results indicate that ~~66-75~~66-75% of the emissions increase can be attributed to power generation and industrial activities. The remaining emission increase is mainly attributed to the road transportation sector. For the DA_23SEP results, ~~50~~almost half of the analysed emissions come from the road transport sector. In some grid cells, the additional road emissions of DA_23SEP are twice as high as those of DA_22SEP, reaching up to 6 Mgd⁻¹ compared to 1.5 Mgd⁻¹, respectively.

The area affected by the emission corrections differs for the two consecutive analysis days. This disparity lies in the different meteorological conditions, particularly in the variation of wind patterns, that occur during these days. As shown in Fig. 2 the prevailing winds in the studied region has low intensity and significant variability at the ground and high altitude on 22

280 September 2021, while on 23 September, the wind is more intense and predominantly originating from the west. This causes different dispersion situations for the pollutant during the two days.

This can be seen in Fig. 5, which shows tropospheric NO₂ columns observed by the TROPOMI (Tropospheric Monitoring Instrument) aboard the Sentinel-5 Precursor (Sentinel-5P) satellite. This data highlights that the accumulation of pollutants resulting in high NO₂ concentrations is very distinct for each individual day. On 22 September 2021, TROPOMI data show a
285 highly polluted area north and northwest of the observation site, which does not prevail on 23 September 2021. This might explain the increase in emissions rates seen in the DA_22SEP results at the north and northwest of the observation site. However, it is unfortunately not possible to directly obtain information about the NO₂ emissions from the TROPOMI data. Nevertheless, the 4D-var assimilation algorithm seems to react to the high concentrations by attributing corrections to emission increases.

These results indicate the strong effects of the wind condition on the observability of the drone measurement. Nevertheless,
290 it shows the potential that the drone observations might have for emissions optimisation, especially for emissions that are emitted at ~~high~~higher altitudes, such as power plants and industries. Drawing definitive conclusions regarding the accuracy of emissions changes is consistently challenging, primarily due to the scarcity of emissions observations. Consequently, we will validate the 4D-var analysis using independent ground-based observations, and we will analyze the contribution of emission changes to the observed improvements in order to evaluate the potential of drone observations in optimising emission rates.




Figure 4. Daily NO_x emissions within the analysed domain (first column) and the analysed NO_x emission changes on 22 September (middle column) and 23 September (last column) 2021. The rows (from top to bottom) display the total NO_x emissions, and the emissions from the public power production, industry, and road transport, respectively.




Figure 5. Maps of the TROPOMI NO₂ tropospheric columns (in molec cm^{-2}) over the studied area on the 22 September 2021 at 11:00 UTC (left) and on the 23 September 2021 at 12:18 UTC (right). Source: <https://browser.dataspace.copernicus.eu/>

295 4.3 Validation against independent observations

4.3.1 Local impact

To validate the impact of the drone data assimilation, we compare the experiment results with independent ground-based observations. Local observations from two monitoring stations located one on each side of the A555 highway but in the same grid cell as the assimilated data (Fig. 1) are used for this evaluation. Figure 6 shows the daily time series of observed O₃, NO, and NO₂ concentrations along with the modelled concentrations from both the reference and assimilation experiments. To evaluate the benefits of the drone data assimilation, the bias, RMSE (Root Mean Square Error), and Pearson correlation are examined for all experiments averaged over the assimilation window and over a 24-hour period (Table 4), using the means of the observations from the two stations as reference.

The DA_22SEP experiment performance for the O₃ concentrations is almost similar to the reference experiment (REF_22SEP). Following the analysis of Sect. 4.1, this is expected because the a priori forecast and the drone observation for near-ground O₃ concentration agree well during this day. The main improvement during the first day is seen for the NO concentrations within the assimilation window as well as during the subsequent free forecast. The assimilation of drone observations results in a strong reduction of the bias by 87 % ($-20.48 \mu\text{gm}^{-3}$) and the RMSE by 20 % ($-7.7 \mu\text{gm}^{-3}$), with an amelioration in the Pearson correlation of 0.15 over the 24-hour period. The daily NO₂ cycle is impacted by the assimilation due to its chemical coupling with O₃ and NO. Therefore, the assimilation experiment exhibits a better performance during the daytime relative to the reference experiment. However, during the late afternoon and nighttime, REF_22SEP performs better than DA_22SEP, as NO₂ is slightly overestimated. The best performance of the drone data assimilation results is obtained on 23 September 2021. A remarkable improvement in the O₃ concentration is noticed within the initial seven hours of the day. The, while a deterioration is observed between 16:00 and 24:00. The daily bias is reduced by 60 % ($-11.18 \mu\text{gm}^{-3}$) and the RMSE by

315 ~~45~~46 % (~~-11.06~~ μgm^{-3}), which also results in an improvement of the correlation by 0.22 during the assimilation window. An improvement in the assimilation results is achieved for NO concentrations. The assimilation experiment reduces the bias by ~~50~~53 % (~~-13.07~~ μgm^{-3}) and RMSE by ~~more than 27~~28 % (~~-11.59~~ μgm^{-3}), with an amelioration in the correlation by 0.5 over the 24-hour evaluation period. For NO₂, a notable improvement can be seen in the forecast from DA_23SEP compared to REF_23SEP. Within the assimilation window, the bias is reduced by 43 % (~~-7.77~~ μgm^{-3}), the RMSE by 29 % (~~-6.68~~ μgm^{-3}), and the correlation improved by 0.19.

320 These results indicate that the 4D-var assimilation of the drone observations has the potential to improve concentration of O₃, NO, and NO₂ during the early morning and daytime when optimising both the initial values and emissions rates simultaneously. The observed deterioration of the O₃ and NO₂ forecast during the late afternoon and nighttime in the DA_23SEP assimilation run is likely related to the NO_x titration process. During the night, O₃ removal is the dominant process in areas with significant NO emission sources (Sillman, 1999). Taking this into account may indicate that the drone data assimilation provides a higher estimate of NO₂ emissions during the night. Since the assimilation algorithm derives only one emission factor per day, the amplitude of the daily temporal emission profile is adjusted. ~~The assumption that the time~~ It is assumed that the temporal emission profile is more certain than the emission strength ~~constrains the optimisation to~~. Deriving e.g. hourly emission factors instead would allow for more flexible adjustments of the emissions, which would be beneficial for the nowadays strongly regulated emission sources, such as the power production (dependent on the availability of renewable energy). Previous studies demonstrated that the temporal distribution of traffic emissions significantly influences nighttime concentrations of NO₂ and O₃ (Menut et al., 2012). As the emission optimisation process maintains the same temporal variability, it is necessary to have 24-hour data assimilation to improve the nighttime O₃ and NO₂ forecasts. Moreover, an inaccurately predicted PBL height can lead to uncertainties in the O₃ and NO₂ forecasts. A full analysis of the PBL representation is however beyond the scope of this study.

Table 4. Statistical comparison of ground-ground-based observations and model outputs (REF: reference run, DA: assimilation run) for O₃, NO, and NO₂ ~~within during~~ the assimilation window ~~(and for 24h-forecast) during~~, in parentheses, the 24-hour forecast on 22-23 September 2021. The Bias and RMSE are in μgm^{-3} .

Statistics		O ₃		NO		NO ₂	
		REF	DA	REF	DA	REF	DA
22 Sep 2021	Bias	-3.91 (-6.02)	-4.37 (-8.50)	-39.93 (-23.45)	-14.52 (-2.97)	2.97 (-1.40)	27.17 (15.73)
	RMSE	10.52 (11.42)	10.93 (13.73)	53.17 (37.84)	38.44 (30.14)	13.90 (17.66)	32.08 (26.10)
	Corr	0.83 (0.92)	0.81 (0.92)	-0.14 (0.13)	-0.10 (0.28)	-0.13 (0.20)	0.16 (0.18)
23 Sep 2021	Bias	18.53 (-5.37)	-7.35 (-21.60)	-52.62 (-24.82)	-23.61 (-11.75)	-17.83 (-9.45)	10.06 (8.99)
	RMSE	24.10 (21.91)	13.04 (26.32)	66.16 (41.77)	46.93 (30.18)	22.84 (17.40)	16.16 (18.70)
	Corr	0.70 (0.71)	0.92 (0.86)	-0.28 (-0.07)	0.22 (0.56)	0.40 (0.28)	0.59 (0.49)




Figure 6. Temporal evolution of the O_3 , NO , and NO_2 concentrations as observed by the ground-stations (red line) and given by the model in the corresponding grid cell: the reference (black line) and the analysis (blue line) over the 24-hour forecast period on 22 and 23 September 2021. Green dots highlight the time of the assimilated drone profiles.

4.3.2 Regional impact

To further investigate the effect on a larger spatial scale, an additional validation is performed using independent ground-based observations from six different ground-based air quality monitoring stations situated in the vicinity of the observation site (Fig. 1, Table A1). For this validation, only stations that are impacted by the assimilation are selected. These are located at distances ranging from 12 km to 85 km from the campaign location. Given the unavailability of NO observations, this validation considers only O₃ and NO₂. Although NO₂ is not assimilated in this study, it is indirectly influenced due to chemical coupling with the observed species and via the optimised NO_x emissions. Figure 7 presents the hourly RMSE time series of O₃ concentrations for the assimilation and reference experiments, averaged over all selected stations. The individual RMSE of O₃ and NO₂ within the assimilation window, for all simulations per station, are presented in Table 5.

Figure 7 shows that the O₃ RMSE for DA_22SEP and DA_23SEP is notably lower than that REF_22SEP within the data assimilation window. Outside the assimilation window, only a small added error is noted between 11:00 and 17:00 UTC for DA_22SEP, which appears similar to the results of the local validation, while no impact is observed during the subsequent free forecast period for DA_23SEP. The largest RMSE reduction takes place at Station 59 (by 30 % (-2.26 ppb_v) on 22 September and by 40 % (-6.61 ppb_v) on 23 September) and, as well as at Station 80 (by 35 % (-2.22 ppb_v) on 22 September and by 34 % (-4.98 ppb_v) on 23 September) that. These stations are situated 12 km and 43 km north of the campaign site, respectively. The smallest reduction occurs at the stations of furthest distance, namely at Station 8 (by 5 % (-0.59 ppb_v) on 22 September and by 4 % (-0.46 ppb_v) on 23 September) and Station 179 (by 2 % (-0.73 ppb_v) on 22 September and by 7 % (-1.22 ppb_v) on 23 September), which are located approximately 85 km north-east of the campaign site. These results suggest that the positive impact of the drone data assimilation is transported to a broader area surrounding the campaign location, resulting in an improvement of O₃ concentrations across a larger area.

For NO₂, a significant RMSE reduction is found at Station 80 (with a decrease of 72 % (-7.7 ppb_v) for DA_22SEP. However, the RMSE for Station 59 and Station 53 show an increase within the assimilation window. For DA_23SEP, better results can be seen for all stations except for the rural Station 59. The best reduction is achieved at Station 80 (by 21 % (-4.16 ppb_v) and Station 114 (by 22 % (-2.80 ppb_v).

Despite the simplicity of the current assimilation approach, which only incorporates data from a single grid box, a positive effect of assimilation is apparent even for stations situated at larger distances from the drone campaign location. This is attributed to the transport of the analysis increment throughout large areas of the studied region.




Figure 7. Temporal evolution of the RMSE (model-observations) in ppb_v for O₃ calculated for the reference (black) and the data assimilation (blue) runs over the 24-hour forecast period across all ground stations on 22 September 2021 (top) and 23 September 2021 (bottom). [Green dots highlight the time of the assimilated drone profiles.](#)

Table 5. The O₃ and NO₂ RMSE between observations data and model results obtained with (DA) and without (REF) drone data assimilation. The results are shown for every ground-based station for the assimilation window. The RMSE is in ppb_v.

RMSE	DA Window		DA Window		
	REF_22SEP	DA_22SEP	REF_23SEP	DA_23SEP	
O ₃	Station 8	11.33	10.74	12.17	11.71
	Station 53	10.29	9.66	8.19	7.29
	Station 59	7.75	5.49	16.71	10.10
	Station 80	6.35	4.13	14.58	9.60
	Station 114	25.86	24.39	22.69	19.87
	Station 179	27.96	27.23	17.55	16.33
NO ₂	Station 8	18.11	17.49	24.05	22.92
	Station 53	12.85	23.81	10.26	10.77
	Station 59	24.25	44.34	16.88	24.45
	Station 80	10.63	2.93	19.59	15.43
	Station 114	24.14	25.82	12.81	10.01
	Station 179	17.78	18.04	19.85	18.08

4.4 Discussion of the potential and limitations of drone data assimilation

The analysis of the DA_22SEP and DA_23SEP experiments shows that the assimilation of drone observations has a positive impact on the vertical distribution of O₃ and NO, and on the daily cycle of O₃ and NO_x at ground level. These promising results underscore the significant potential of drone data assimilation in enhancing regional air quality analysis. Moreover, the assimilation process ~~allows to obtain~~ provides optimised emissions rates, ~~with distinct outcomes observed for~~ each day. To investigate the role of emission optimisation in the analysis improvement, Table 6 presents the cost reduction for O₃ and NO, as well as the partial costs attributed to the optimisation of the initial values (IV) $\left(\frac{J_b(\mathbf{x}_0)}{J(\mathbf{x}_0, \mathbf{e})}\right)$ and the emissions correction factors (EF) $\left(\frac{J_e(\mathbf{e})}{J(\mathbf{x}_0, \mathbf{e})}\right)$. For both assimilation experiments, the costs are reduced by more than 30 %, which confirms the successful assimilation of the drone profiles. In particular, the O₃ costs of DA_23SEP are highly reduced by 80 %, resulting in a precise alignment between the 4D-var analysis and the O₃ observations. The partial costs vary between the two days. For DA_22SEP, the costs associated with IV are more than twice that of EF, which indicates important IV adjustments and a minimal impact of the emissions changes in the cost minimisation. In contrast for DA_23SEP, the effect of optimising the emissions is higher. This indicates that a significant part of the improvement observed in the analysis is due to the optimisation of EF. Therefore, the drone observations may also have significant potential for assessing local emissions. ~~This is supported by the findings of Wu et al. (2022), affirming that observation at high altitudes can be advantageous for optimising emissions under suitable wind conditions.~~ In a recent study by Wu et al. (2022), it was demonstrated that for high-altitude observations, the efficiency of emission rate optimization is conditioned by favorable wind conditions and strong vertical diffusion.

Despite the observed improvements in the analysis, some limitations are noted. Firstly, the results reported in Sect. 4.1 show a limited impact on the NO vertical profiles on 23 September 2021. Although effective correction is achieved in the ground and near-ground levels, limited improvements are obtained for the NO concentrations at higher altitudes (above 150 m) for the first 3 profiles of the day. Figure 8 illustrates the vertically resolved analysis increment (4D-var analysis – reference run) for O₃, NO, and NO₂ on 23 September 2021. A negative O₃ increment alongside a positive NO₂ increment is noted, both exhibiting a well-developed vertical spread. The NO increment is constrained near ground level during the early hours of the day. The reason behind this is the NO_x titration process, where freshly emitted NO, including additional NO emissions resulting from emission optimisation, reacts with O₃ to produce NO₂. To achieve a better results, a larger NO increment is need. However, the NO observations from the drone exhibit high measurement errors compared to the background errors, which limits the effectiveness of assimilating this data.

Secondly, ~~Some~~ some suboptimal outcomes are observed in the free run, namely for O₃ and NO₂ ground concentration, suggesting that the advantage of the drone data assimilation is limited to the assimilation window (Fig. 6, Fig. A3, and Fig. A4). Nevertheless, this result is not surprising and is completely explainable. Initially, it is important to note that the reference model simulation already provides underestimations of O₃ peaks during the afternoon and nighttime, which may be linked to uncertainties in the boundary layer height at night, vertical diffusion, and/or emissions profiles. Through the 4D-var assimilation of drone data, adjustments are made to the NO_x emissions. However, in regions characterised by high NO_x emissions, O₃ formation exhibits reduced sensitivity to NO_x emissions but increased sensitivity to VOCs (Visser et al., 2019; Sillman, 1999).

Thus, the inability to adjust O₃ concentrations and, consequently, NO₂ in our simulations is not a limitation specific to drone data assimilation.

Table 6. The percentage of cost reduction achieved for O₃ and NO, as well as the percentage of the partial costs attributed to initial value correction (IV) and emissions correction (EF) relative to the total cost function.

	Cost reduction		Partial costs	
	O ₃	NO	EF	IV
DA_22SEP	34 %	41 %	9 %	25 %
DA_23SEP	80 %	36 %	10 %	4 %




Figure 8. Vertical cross-section of the analysis increment of O₃, NO, and NO₂ on 23 September 2021 at selected time steps. The cross-section location is the MesSBAR campaign site.

400 5 Conclusion

In this study, drone profile measurements of O₃ and NO are assimilated using the 4D-var data assimilation system of EURAD-IM. This represents the first application of drone data assimilation within a CTM. The primary objective is to assess the ability of drone observations to improve regional air quality analysis when the joint initial value and emission correction factor optimisation approach is applied. The research is conducted using data collected during the two-day MesSBAR campaign in 2021.

405 To evaluate the results, a comparison is made with ground-based observations obtained at stations very close to the drone flight base location. Moreover, regional validation is conducted using ground-based data from the the European air quality monitoring network.

The 4D-var assimilation of drone data has a positive impact on the representation of these pollutants in the PBL. First, significant improvements are noted in the O₃ and NO vertical profiles, with biases decreasing by 30 % and 55 %, respectively, on the first day and by 35 % on the second day for both species. Moreover, there is a noticeable impact on ground concentrations in the analysis. In the studied grid cell, biases are reduced by up to 60 % for O₃, 55 % for NO, and 43 % for NO₂ ground concentrations within the assimilation window. Furthermore, due to the pollution transport and the connected information propagation in the 4D-var algorithm, a positive impact is seen on the ground concentrations of O₃ and NO₂ in locations farther from the measurement site. This study also identifies the assessment of emission correction factors as one component of the analysis improvements, which underline the potential of the drone observations to be beneficial for emission optimisation.

415 There are some limitations to this study. Firstly, due to constraints in data availability, the study is restricted to assimilating drone data within a singular grid cell column. Therefore, it would be advantageous to include multiple measurement points distributed across the region, strategically positioned both upwind and downwind of emission sources. Another limitation of this study is the assimilation of data available only during a partial time window of the day. The inclusion of a more extensive observational data set covering longer periods, ideally over 24 hours to enable an extended assimilation window, would greatly enhance the optimisation of emission rates.

In conclusion, the 4D-var assimilation of drone data within the regional air quality model EURAD-IM yields promising results by improving the vertical distribution of pollutants and correcting ground concentrations. ~~As a perspective for future work, one interesting approach is~~ From a future perspective, a valuable extension of this work will be to conduct Observing System Experiments (OSE) ~~to assess the advantages and limitations of integrating drone observations into CTMs through the application of a variational dataassimilation technique~~ Simulation Experiments (OSSEs) to evaluate the added value of integrating drone-based observations into the air quality forecasting system, compared to conventional observations such as ground-based measurements and satellite data.

Data availability. The drone data from the MesSBAR campaign used in this study are publicly available by Schlerf et al. (2024) on PAN-GAEA at the following DOI: <https://doi.pangaea.de/10.1594/PANGAEA.971503>.

Author contributions. HE and ACL designed the study. HE conducted the simulations, performed the analyses under scientific supervision of ACL, PF and AW. TS and RT provided the observational profile data. The manuscript was prepared by HE with the help of all co-authors. All authors reviewed the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

435 *Acknowledgements.* The authors gratefully acknowledge all the MesSBAR project partners for their valuable efforts in conducting the campaign and processing the data used in this work. We also thank the Federal Highway Research Institute (BAST) for providing the ground-based observations and meteorological data. Financial support for the MesSBAR project was provided by the Modernity Fund mFUND of the Federal Ministry of Transport and Digital Infrastructure (BMVI) under grant agreement 19F2097. The authors also gratefully acknowledge the computing time granted through JARA on the supercomputer JURECA (Jülich Supercomputing Centre, 2021) at Forschungszentrum Jülich.

440

References

- Ackermann, I. J., Hass, H., Memmesheimer, M., Ebel, A., Binkowski, F. S., and Shankar, U.: Modal aerosol dynamics model for Europe: development and first applications, *Atmos. Environ.*, 32, 2981–2999, [https://doi.org/10.1016/S1352-2310\(98\)00006-5](https://doi.org/10.1016/S1352-2310(98)00006-5), 1998.
- Altstädter, B., Platis, A., Wehner, B., Scholtz, A., Wildmann, N., Hermann, M., Käthner, R., Baars, H., Bange, J., and Lampert, A.: ALADINA
445 - an unmanned research aircraft for observing vertical and horizontal distributions of ultrafine particles within the atmospheric boundary layer, *Atmos. Meas. Tech.*, 8, 1627–1639, <https://doi.org/10.5194/amt-8-1627-2015>, 2015.
- Bretschneider, L., Schlerf, A., Baum, A., Bohlius, H., Buchholz, M., Düsing, S., Ebert, V., Erraji, H., Frost, P., Käthner, R., Krüger, T., Lange, A. C., Langner, M., Nowak, A., Pätzold, F., Rüdiger, J., Saturno, J., Scholz, H., Schuldt, T., Seldschopf, R., Sobotta, A., Tillmann, R., Wehner, B., Wesolek, C., Wolf, K., and Lampert, A.: MesSBAR-Multicopter and Instrumentation for Air Quality Research, *Atmosphere*,
450 <https://doi.org/10.3390/atmos13040629>, 2022.
- Corrigan, C. E., Roberts, G. C., Ramana, M. V., Kim, D., and Ramanathan, V.: Capturing vertical profiles of aerosols and black carbon over the Indian Ocean using autonomous unmanned aerial vehicles, *Atmos. Chem. Phys.*, 8, 737–747, <https://doi.org/10.5194/acp-8-737-2008>, 2008.
- De Mazière, M., Thompson, A. M., Kurylo, M. J., Wild, J. D., Bernhard, G., Blumenstock, T., Braathen, G. O., Hannigan, J. W., Lambert, J.-C., Leblanc, T., McGee, T. J., Nedoluha, G., Petropavlovskikh, I., Seckmeyer, G., Simon, P. C., Steinbrecht, W., and Strahan, S. E.: The
455 Network for the Detection of Atmospheric Composition Change (NDACC): history, status and perspectives, *Atmospheric Chemistry and Physics*, 18, 4935–4964, <https://doi.org/10.5194/acp-18-4935-2018>, 2018.
- Deroubaix, A., Hoelzemann, J. J., Ynoue, R. Y., de Almeida Albuquerque, T. T., Alves, R. C., de Fatima Andrade, M., ao, W. L. A., Bouarar, I., de Souza Fernandes Duarte, E., Elbern, H., Franke, P., Lange, A. C., Lichtig, P., Lugon, L., Martins, L. D., de Arruda Moreira, G.,
460 Pedruzzi, R., Rosario, N., and Brasseur, G.: Intercomparison of Air Quality Models in a Megacity: Toward an Operational Ensemble Forecasting System for São Paulo, *J. Geophys. Res.: Atmospheres*, 129, <https://doi.org/10.1029/2022JD038179>, 2024.
- Diaz, J., Corrales, E., Madrigal, Y., Pieri, D., Bland, G., Miles, T., and Fladeland, M.: Volcano Monitoring with small Unmanned Aerial Systems, *American Institute of Aeronautics and Astronautics*, <https://doi.org/10.2514/6.2012-2522>, 2012.
- Duarte, E. D. S. F., Franke, P., Lange, A. C., Friese, E., da Silva Lopes, F. J., ao da Silva, J. J., dos Reis, J. S., Landulfo, E., e Silva, C. M. S.,
465 Elbern, H., and Hoelzemann, J. J.: Evaluation of atmospheric aerosols in the metropolitan area of São Paulo simulated by the regional EURAD-IM model on high-resolution, *Atmos. Pollut. Res.*, 12, 451–469, <https://doi.org/10.1016/j.apr.2020.12.006>, 2021.
- Elbern, H. and Schmidt, H.: Ozone episode analysis by four-dimensional variational chemistry data assimilation, *J. Geophys. Res. Atmospheres*, 106, 3569–3590, <https://doi.org/10.1029/2000JD900448>, 2001.
- Elbern, H., Strunk, A., Schmidt, H., and Talagrand, O.: Emission rate and chemical state estimation by 4-dimensional variational inversion,
470 *Atmos. Chem. Phys.*, 7, 3749–3769, <https://doi.org/10.5194/acp-7-3749-2007>, 2007.
- Flagg, D. D., Doyle, J. D., Holt, T. R., Tyndall, D. P., Amerault, C. M., Geiszler, D., Haack, T., Moskaitis, J. R., Nachamkin, J., and Eleuterio, D. P.: On the Impact of Unmanned Aerial System Observations on Numerical Weather Prediction in the Coastal Zone, *Mon. Wea. Rev.*, 146, 599–622, <https://doi.org/10.1175/MWR-D-17-0028.1>, 2018.
- Franke, P., Lange, A. C., and Elbern, H.: Particle-filter-based volcanic ash emission inversion applied to a hypothetical sub-Plinian Eyjafjallajökull eruption using the Ensemble for Stochastic Integration of Atmospheric Simulations (ESIAS-chem) version 1.0, *Geosci. Model Dev.*, 15, 1037–1060, <https://doi.org/10.5194/gmd-15-1037-2022>, 2022.

- Franke, P., Lange, A. C., Steffens, B., Pozzer, A., Wahner, A., and Kiendler-Scharr, A.: European air quality in view of the WHO 2021 guideline levels: Effect of emission reductions on air pollution exposure, *Elem. Sci. Anth.*, 12, 00127, <https://doi.org/10.1525/elementa.2023.00127>, 2024.
- 480 Gama, C., Ribeiro, I., Lange, A. C., Vogel, A., Ascenso, A., Seixas, V., Elbern, H., Borrego, C., Friese, E., and Monteiro, A.: Performance assessment of CHIMERE and EURAD-IM' dust modules, *Atmos. Pollut. Res.*, 10, 1336–1346, <https://doi.org/10.1016/j.apr.2019.03.005>, 2019.
- García, O. E., Schneider, M., Sepúlveda, E., Hase, F., Blumenstock, T., Cuevas, E., Ramos, R., Gross, J., Barthlott, S., Röhling, A. N., Sanromá, E., González, Y., Gómez-Peláez, A. J., Navarro-Comas, M., Puenteadura, O., Yela, M., Redondas, A., Carreño, V., León-Luis, S. F., Reyes, E., García, R. D., Rivas, P. P., Romero-Campos, P. M., Torres, C., Prats, N., Hernández, M., and López, C.: Twenty years of ground-based NDACC FTIR spectrometry at Izaña Observatory – overview and long-term comparison to other techniques, *Atmospheric Chemistry and Physics*, 21, 15 519–15 554, <https://doi.org/10.5194/acp-21-15519-2021>, 2021.
- 485 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.
- 490 Illingworth, S., Allen, G., Percival, C., Hollingsworth, P., Gallagher, M., Ricketts, H., Hayes, H., Åradsz, P., Crawley, D., and Roberts, G.: Measurement of boundary layer ozone concentrations on-board a Skywalker unmanned aerial vehicle, *Atmos. Sci. Lett.*, 15, 252–258, <https://doi.org/10.1002/asl2.496>, 2014.
- Jensen, A. A., Pinto, J. O., Bailey, S. C. C., Sobash, R. A., de Boer, G., Houston, A. L., Chilson, P. B., Bell, T., Romine, G., Smith, S. W., Lawrence, D. A., Dixon, C., Lundquist, J. K., Jacob, J. D., Elston, J., Waugh, S., and Steiner, M.: Assimilation of a Coordinated Fleet of Uncrewed Aircraft System Observations in Complex Terrain: EnKF System Design and Preliminary Assessment, *Mon. Wea. Rev.*, 149, 1459–1480, <https://doi.org/10.1175/MWR-D-20-0359.1>, 2021.
- 500 Jonassen, M. O., Ólafsson, H., Ágústsson, H., Ólafur Rögnvaldsson, and Reuder, J.: Improving High-Resolution Numerical Weather Simulations by Assimilating Data from an Unmanned Aerial System, *Mon. Wea. Rev.*, 140, 3734–3756, <https://doi.org/10.1175/MWR-D-11-00344.1>, 2012.
- Jülich Supercomputing Centre: JURECA: Data Centric and Booster Modules implementing the Modular Supercomputing Architecture at Jülich Supercomputing Centre, *Journal of large-scale research facilities JLSRF*, 7, A182, <https://doi.org/10.17815/jlsrf-7-182>, 2021.
- Klonecki, A., Pommier, M., Clerbaux, C., Ancellet, G., Cammas, J.-P., Coheur, P.-F., Cozic, A., Diskin, G. S., Hadji-Lazaro, J., Hauglustaine, D. A., Hurtmans, D., Khatatov, B., Lamarque, J.-F., Law, K. S., Nedelec, P., Paris, J.-D., Podolske, J. R., Prunet, P., Schlager, H., Szopa, S., and Turquety, S.: Assimilation of IASI satellite CO fields into a global chemistry transport model for validation against aircraft measurements, *Atmos. Chem. Phys.*, 12, 4493–4512, <https://doi.org/10.5194/acp-12-4493-2012>, 2012.
- 505 Kuenen, J. J. P., Visschedijk, A. J. H., Jozwicka, M., and van der Gon, H. A. C. D.: TNO-MACC-II emission inventory; a multi-year (2003-2009) consistent high-resolution European emission inventory for air quality modelling, *Atmos. Chem. Phys.*, 14, 10 963–10 976, <https://doi.org/10.5194/acp-14-10963-2014>, 2014.
- 510 Lampert, A., Altstädter, B., Bärffuss, K., Bretschneider, L., Sandgaard, J., Michaelis, J., Lobitz, L., Asmussen, M., Damm, E., Käthner, R., Krüger, T., Lüpkes, C., Nowak, S., Peuker, A., Rausch, T., Reiser, F., Scholtz, A., Zakharov, D. S., Gaus, D., Bansmer, S., Wehner, B., and Pätzold, F.: Unmanned Aerial Systems for Investigating the Polar Atmospheric Boundary Layer - Technical Challenges and Examples of Applications, *Atmosphere*, 11, 416, <https://doi.org/10.3390/atmos11040416>, 2020.

- Lawrence, D. A. and Balsley, B. B.: High-Resolution Atmospheric Sensing of Multiple Atmospheric Variables Using the DataHawk Small
515 Airborne Measurement System, *J. Atmos. Ocean. Technol.*, 30, 2352–2366, <https://doi.org/10.1175/JTECH-D-12-00089.1>, 2013.
- Leuenberger, D., Haeferle, A., Omanovic, N., Fengler, M., Martucci, G., Calpini, B., Fuhrer, O., and Rossa, A.: Improving
High-Impact Numerical Weather Prediction with Lidar and Drone Observations, *Bull. Am. Meteorol. Soc.*, 101, E1036–E1051,
<https://doi.org/10.1175/BAMS-D-19-0119.1>, 2020.
- Liu, D. C. and Nocedal, J.: On the limited memory BFGS method for large scale optimization, *Math. Program.*, 45, 503–528,
520 <https://doi.org/10.1007/BF01589116>, 1989.
- Liu, X., Mizzi, A. P., Anderson, J. L., Fung, I. Y., and Cohen, R. C.: Assimilation of satellite NO₂ observations at high spatial resolution
using OSSEs, *Atmos. Chem. Phys.*, 17, 7067–7081, <https://doi.org/10.5194/acp-17-7067-2017>, 2017.
- Marécal, V., Peuch, V.-H., Andersson, C., Andersson, S., Arteta, J., Beekmann, M., Benedictow, A., Bergström, R., Bessagnet, B., Cansado,
A., Chéroux, F., Colette, A., Coman, A., Curier, R. L., Denier van der Gon, H. A. C., Drouin, A., Elbern, H., Emili, E., Engelen, R. J.,
525 Eskes, H. J., Foret, G., Friese, E., Gauss, M., Giannaros, C., Guth, J., Joly, M., Jaumouillé, E., Josse, B., Kadygrov, N., Kaiser, J. W., Kra-
jsek, K., Kuenen, J., Kumar, U., Liora, N., Lopez, E., Malherbe, L., Martinez, I., Melas, D., Meleux, F., Menut, L., Moinat, P., Morales, T.,
Parmentier, J., Piacentini, A., Plu, M., Poupkou, A., Queguiner, S., Robertson, L., Rouil, L., Schaap, M., Segers, A., Sofiev, M., Tarasson,
L., Thomas, M., Timmermans, R., Valdebenito, A., van Velthoven, P., van Versendaal, R., Vira, J., and Ung, A.: A regional air quality fore-
casting system over Europe: the MACC-II daily ensemble production, *Geosci. Model Dev.*, 8, 2777–2813, [https://doi.org/10.5194/gmd-](https://doi.org/10.5194/gmd-8-2777-2015)
530 [8-2777-2015](https://doi.org/10.5194/gmd-8-2777-2015), 2015.
- Martin, R. V.: Satellite remote sensing of surface air quality, *Atmos. Environ.*, 42, 7823–7843,
<https://doi.org/10.1016/j.atmosenv.2008.07.018>, 2008.
- Memmesheimer, M., H. Hass, J. Tippke, and A. Ebel: Modeling of episodic emission data for Europe with the EURAD Emission Model
EEM, in: the International Speciality Conference "Regional Photochemical Measurement and modeling studies", San Diego, CA, USA,
535 1995.
- Menut, L., Goussebaile, A., Bessagnet, B., Khvorostyanov, D., and Ung, A.: Impact of realistic hourly emissions profiles on air pollutants
concentrations modelled with CHIMERE, *Atmos. Environ.*, 49, 233–244, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2011.11.057>,
2012.
- Nathan, B. J., Golston, L. M., O'Brien, A. S., Ross, K., Harrison, W. A., Tao, L., Lary, D. J., Johnson, D. R., Covington, A. N., Clark, N. N.,
540 and Zondlo, M. A.: Near-Field Characterization of Methane Emission Variability from a Compressor Station Using a Model Aircraft,
Environ. Sci. Technol., 49, 7896–7903, <https://doi.org/10.1021/acs.est.5b00705>, 2015.
- O'Sullivan, D., Taylor, S., Elston, J., Baker, C. B., Hotz, D., Marshall, C., Jacob, J., Barfuss, K., Pigué, B., Roberts, G., Omanovic, N.,
Fengler, M., Jensen, A. A., Steiner, M., and Houston, A. L.: The Status and Future of Small Uncrewed Aircraft Systems (UAS) in
Operational Meteorology, *Bull. Am. Meteorol. Soc.*, 102, E2121–E2136, <https://doi.org/10.1175/BAMS-D-20-0138.1>, 2021.
- 545 Paschalidi, Z.: Inverse Modelling for Tropospheric Chemical State Estimation by 4-Dimensional Variational Data Assimilation from Rou-
tinely and Campaign Platforms, Ph.D. thesis, University of Cologne, 2015.
- Petetin, H., Jeoffrion, M., Sauvage, B., Athier, G., Blot, R., Boulanger, D., Clark, H., Cousin, J.-M., Gheusi, F., Nedelec, P., Steinbacher,
M., and Thouret, V.: Representativeness of the IAGOS airborne measurements in the lower troposphere, *Elementa-Sci. Anthropol.*, 6,
<https://doi.org/10.1525/elementa.280>, 2018.
- 550 Rabitz, H. and Aliş, O. F.: General foundations of high-dimensional model representations, *J. Math. Chem.*, 25, 197–233,
<https://doi.org/10.1023/A:1019188517934>, 1999.

- Roberts, G. C., Ramana, M. V., Corrigan, C., Kim, D., and Ramanathan, V.: Simultaneous observations of aerosol-cloud-albedo interactions with three stacked unmanned aerial vehicles, *P. Natl. Acad. Sci. USA*, 105, 7370–7375, <https://doi.org/10.1073/pnas.0710308105>, 2008.
- Roselle, S. and Binkowski, F.: Cloud Dynamics and Chemistry, in *Science Algorithms of the EPA Models-3 Community Multiscale Air Quality (CMAQ) Modeling System*, Research Triangle Park, EPA 600/R-99-030, 1999.
- 555 Sandu, A. and Sander, R.: Technical note: Simulating chemical systems in Fortran90 and Matlab with the Kinetic PreProcessor KPP-2.1, *Atmos. Chem. Phys*, 6, 187–195, <https://doi.org/10.5194/acp-6-187-2006>, 2006.
- Scheffe, R., Philbrick, R., on Macdonald, C., Dye, T., Gilroy, M., and Carlton, A.-M.: Observational Needs for Four-dimensional Air Quality Characterization, https://cfpub.epa.gov/si/si_public_record_report.cfm?Lab=NERL&dirEntryId=213564, 2009.
- 560 Schell, B., Ackermann, I. J., Hass, H., Binkowski, F. S., and Ebel, A.: Modeling the formation of secondary organic aerosol within a comprehensive air quality model system, *J. Geophys. Res.: Atmospheres*, 106, 28 275–28 293, <https://doi.org/10.1029/2001JD000384>, 2001.
- Schlerf, A., Bretschneider, L., Schuchard, M., Düsing, S., Käthner, R., Wehner, B., Schuldt, T., Wesolek, C., Tillmann, R., Lange, A. C., Erraji, H., Krüger, T., Scholz, H., Frost, P., Sobotta, A., Baum, A., Ebert, V., Bohlius, H., Nowak, A., Langner, M., and Lampert, A.:
- 565 Validation and optimization of the drone air quality measurement system MesSBAR, <https://doi.org/10.1594/PANGAEA.971503>, 2024.
- Schuldt, T., Gkatzelis, G. I., Wesolek, C., Rohrer, F., Winter, B., Kuhlbusch, T. A. J., Kiendler-Scharr, A., and Tillmann, R.: Electrochemical sensors on board a Zeppelin NT: in-flight evaluation of low-cost trace gas measurements, *Atmos. Meas. Tech.*, 16, 373–386, <https://doi.org/10.5194/amt-16-373-2023>, 2023.
- Schuyler, T. and Guzman, M.: Unmanned Aerial Systems for Monitoring Trace Tropospheric Gases, *Atmosphere*, 8, 206, <https://doi.org/10.3390/atmos8100206>, 2017.
- 570 Sillman, S.: The relation between ozone, NO_x and hydrocarbons in urban and polluted rural environments, *Atmos. Environ.*, 33, 1821–1845, [https://doi.org/10.1016/S1352-2310\(98\)00345-8](https://doi.org/10.1016/S1352-2310(98)00345-8), 1999.
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang, X.-Y., Wang, W., and Powers, J. G.: A Description of the Advanced Research WRF Version 3, 2008.
- 575 Stockwell, W. R., Kirchner, F., Kuhn, M., and Seefeld, S.: A new mechanism for regional atmospheric chemistry modeling, *J. Geophys. Res.*, 102, 847–872, <https://doi.org/10.1029/97JD00849>, 1997.
- Sun, Q., Vihma, T., Jonassen, M. O., and Zhang, Z.: Impact of Assimilation of Radiosonde and UAV Observations from the Southern Ocean in the Polar WRF Model, *Adv. Atmos. Sci.*, 37, 441–454, <https://doi.org/10.1007/s00376-020-9213-8>, 2020.
- Tillmann, R., Gkatzelis, G. I., Rohrer, F., Winter, B., Wesolek, C., Schuldt, T., Lange, A. C., Franke, P., Friese, E., Decker, M., Wegener, R.,
- 580 Hundt, M., Aseev, O., and Kiendler-Scharr, A.: Air quality observations onboard commercial and targeted Zeppelin flights in Germany - a platform for high-resolution trace-gas and aerosol measurements within the planetary boundary layer, *Atmos. Meas. Tech.*, 15, 3827–3842, <https://doi.org/10.5194/amt-15-3827-2022>, 2022.
- Tirpitz, J.-L., Frieß, U., Hendrick, F., Alberti, C., Allaart, M., Apituley, A., Bais, A., Beirle, S., Berkhout, S., Bogner, K., Bösch, T., Bruchkouski, I., Cede, A., Chan, K. L., den Hoed, M., Donner, S., Drosoglou, T., Fayt, C., Friedrich, M. M., Frumau, A., Gast, L., Gielen, C.,
- 585 Gomez-Martín, L., Hao, N., Hensen, A., Henzing, B., Hermans, C., Jin, J., Kreher, K., Kuhn, J., Lampel, J., Li, A., Liu, C., Liu, H., Ma, J., Merlaud, A., Peters, E., Pinardi, G., Piters, A., Platt, U., Puentedura, O., Richter, A., Schmitt, S., Spinei, E., Stein Zweers, D., Strong, K., Swart, D., Tack, F., Tiefengraber, M., van der Hoff, R., van Roozendaal, M., Vlemmix, T., Vonk, J., Wagner, T., Wang, Y., Wang, Z., Wenig, M., Wiegner, M., Wittrock, F., Xie, P., Xing, C., Xu, J., Yela, M., Zhang, C., and Zhao, X.: Intercomparison of MAX-DOAS

- vertical profile retrieval algorithms: studies on field data from the CINDI-2 campaign, *Atmospheric Measurement Techniques*, 14, 1–35, <https://doi.org/10.5194/amt-14-1-2021>, 2021.
- 590 Villa, T., Gonzalez, F., Miljievic, B., Ristovski, Z., and Morawska, L.: An Overview of Small Unmanned Aerial Vehicles for Air Quality Measurements: Present Applications and Future Prospectives, *Sensors*, 16, 1072, <https://doi.org/10.3390/s16071072>, 2016.
- Visser, A. J., Boersma, K. F., Ganzeveld, L. N., and Krol, M. C.: European NO_x emissions in WRF-Chem derived from OMI: impacts on summertime surface ozone, *Atmos. Chem. Phys.*, 19, 11 821–11 841, <https://doi.org/10.5194/acp-19-11821-2019>, 2019.
- 595 Walcek, C. J.: Minor flux adjustment near mixing ratio extremes for simplified yet highly accurate monotonic calculation of tracer advection, *J. Geophys. Res.: Atmospheres*, 105, 9335–9348, <https://doi.org/10.1029/1999JD901142>, 2000.
- Wang, H., Lu, X., Jacob, D. J., Cooper, O. R., Chang, K.-L., Li, K., Gao, M., Liu, Y., Sheng, B., Wu, K., Wu, T., Zhang, J., Sauvage, B., Nédélec, P., Blot, R., and Fan, S.: Global tropospheric ozone trends, attributions, and radiative impacts in 1995–2017: an integrated analysis using aircraft (IAGOS) observations, ozonesonde, and multi-decadal chemical model simulations, *Atmos. Chem. Phys.*, 22, 13 753–13 782, <https://doi.org/10.5194/acp-22-13753-2022>, 2022.
- 600 Wang, Y.-C., Wang, S.-H., Lewis, J. R., Chang, S.-C., and Griffith, S. M.: Determining Planetary Boundary Layer Height by Micro-pulse Lidar with Validation by UAV Measurements, *Aerosol. Air. Qual. Res.*, 21, 200 336, <https://doi.org/10.4209/aaqr.200336>, 2021.
- Weaver, A. and Courtier, P.: Correlation modelling on the sphere using a generalized diffusion equation, *Q. J. Roy. Meteor. Soc.*, 127, 1815–1846, <https://doi.org/https://doi.org/10.1002/qj.49712757518>, 2001.
- 605 Wu, X., Elbern, H., and Jacob, B.: The assessment of potential observability for joint chemical states and emissions in atmospheric modelings, *Stoch. Environ. Res. Risk. Assess.*, 36, 1743–1760, <https://doi.org/10.1007/s00477-021-02113-x>, 2022.
- Yang, S., Li, X., Zeng, L., Yu, X., Liu, Y., Lu, S., Huang, X., Zhang, D., Xu, H., Lin, S., Liu, H., Feng, M., Song, D., Tan, Q., Cui, J., Wang, L., Chen, Y., Wang, W., Sun, H., Song, M., Kong, L., Liu, Y., Wei, L., Zhu, X., and Zhang, Y.: Development of multi-channel whole-air sampling equipment onboard an unmanned aerial vehicle for investigating volatile organic compounds' vertical distribution in the planetary boundary layer, *Atmos. Meas. Tech.*, 16, 501–512, <https://doi.org/10.5194/amt-16-501-2023>, 2023.
- 610 Zhang, L., Brook, J. R., and Vet, R.: A revised parameterization for gaseous dry deposition in air-quality models, *Atmos. Chem. Phys.*, 3, 2067–2082, <https://doi.org/10.5194/acp-3-2067-2003>, 2003.

Appendix A

Table A1. Information about the Ground-based monitoring stations.

Station Number	Station Code	Station Name	Distance from campaign site	Station Type	Latitude(°N)	Longitude (°E)	Altitude
8	DENW008	Dortmund-Eving	86.5 km	Suburban	51.5369	7.4575	75 m
53	DENW053	Köln-Chorweiler	28.2 km	Suburban	51.0193	6.8846	45 m
59	DENW059	Köln-Rodenkirchen	12.1 km	Rural	50.8898	6.9852	45 m
80	DENW080	Solingen-Wald	43.2 km	Rural	51.1838	7.0526	207 m
114	DENW114	Wuppertal-Langerfeld	56.8 km	Suburban	51.2776	7.2319	186 m
179	DENW179	Schwerte	82.4 km	Suburban	51.4488	7.5823	157 m




Figure A1. Emission correction factors of NO and NO₂ resulting from the conducted assimilation experiments on 22 September 2021 (a and b) and 23 September 2021 (c and d).




Figure A2. Temporal evolution of the RMSE (model-observations) in ppb_v for NO_2 calculated for the reference (black) and the analysis (blue) over the 24-hour forecast period across all ground stations on 22 September 2021 (top) and 23 September 2021 (bottom). [Green dots](#) highlight the time of the assimilated drone profiles. [The grey shade](#) illustrates the length of the assimilation window.




Figure A3. Time series of O_3 concentrations in ppb_v as measured by ground-based stations and predicted by the model. The left panel shows data from 22 September 21, 2021, while the right panel displays data from September 23 - September 2021.




Figure A4. Same as Figure A3 but for NO_2 .