



A global analysis of the fractal properties of clouds revealing anisotropy of turbulence across scales

Karlie N. Rees¹, Timothy J. Garrett¹, Thomas D. DeWitt¹, Corey Bois¹, Steven K. Krueger¹, and Jérôme C. Riedi²

¹Department of Atmospheric Sciences, University of Utah, 135 S 1460 E Rm 819, Salt Lake City, UT 84112, USA

²University of Lille, CNRS, UMR 8518 – LOA – Laboratoire d’Optique Atmosphérique, 59000 Lille, France

Correspondence: Tim Garrett (tim.garrett@utah.edu)

Abstract. The deterministic motions of clouds and turbulence, despite their chaotic nature, nonetheless follow simple statistical power-law scalings: a fractal dimension D relates individual cloud perimeters p to measurement resolution, and turbulent fluctuations scale with separation distance through the Hurst exponent \mathcal{H} . It remains uncertain whether atmospheric turbulence is best characterized by split isotropy that is three-dimensional with $\mathcal{H} = 1/3$ at small scales and two-dimensional with $\mathcal{H} = 1$ at large scales, or by wide-range anisotropic scaling with an intermediate value of \mathcal{H} . Here, we introduce an “ensemble fractal dimension” D_e – analogous to D – that relates the total cloud perimeter per domain area \mathcal{P} as seen from space to measurement resolution, and show theoretically how turbulent dimensionality and cloud edge geometry are linked through $\mathcal{H} = D_e - 1$. Observationally, by progressively coarsening the resolution of cloud mask arrays from various global satellite platforms and a numerical simulation of a tropical domain we find the scaling $D_e \sim 5/3$, or $\mathcal{H} \sim 2/3$, a value nearly consistent with a previously proposed “23/9D” anisotropic turbulent scaling. Remarkably, the same scaling links two “limiting case” estimates of \mathcal{P} evaluated at the planetary scale and the Kolmogorov microscale, as separated by 11 orders of magnitude, suggesting that cloud and turbulent behaviors are constrained by basic planetary parameters.

1 Introduction

The atmosphere is radiatively open and materially closed. Radiatively, Earth’s global mean temperature is sustained by a balance between absorption of high-intensity shortwave sunlight that is reemitted nearly isotropically at longwave frequencies to the cold of space. Materially, the total atmospheric mass is confined to the planet by gravity and can only be redistributed by turbulent circulations that mix air and moisture over a broad range of scales within the thin atmospheric layer. Clouds play important roles in determining the magnitude of both categories of flow. Geometrically speaking, cloud areas govern radiative fluxes (Zelinka et al., 2022) while the edge length or perimeter of clouds controls exchanges of air between clouds and their clear-sky environment (Zhao and Austin, 2005; Heus et al., 2008; Garrett et al., 2018).

A scientific challenge is that the seemingly objective properties of cloud area and perimeter are a function of the more subjective choice of spatial resolution ξ (defined as either the pixel side length in a satellite image or the grid spacing in a model). Clouds smaller than ξ cannot be resolved, and the square shapes in a resolved grid do not reflect more irregular cloud



structures. Even casual observations of the sky show cloud edges that are intricately complex for any plausibly resolvable scale.
25 For example, the boundary of a small cumulus cloud may appear smoothly rounded at first glance, but fine turbulent structures
become apparent when it is viewed through binoculars. The change of observational scale lengthens the cloud boundary with
clear skies, even as the total cloud area remains nearly unchanged. Because the resolution-dependent cloud perimeter is shaped
by the complex and chaotic processes of turbulent mixing and diffusion (Hentschel and Procaccia, 1984), and while air and
energy exchanges are physically independent of ξ , a resolution-based link is required to relate the two (Lovejoy et al., 2001;
30 Fielding et al., 2020).

Fractal geometry is often used as a tool for characterizing the resolution-dependent complexity of shapes. The fractal dimension D was first introduced by Richardson (1961) to characterize the complexity of political borders and was later popularized by Mandelbrot (1967) to describe how the length of a coastline changes depending on the length of the ruler used to measure it. Generally, the perimeter p around an individual fractal object can be related to the measurement resolution ξ through

$$35 \quad p \propto \xi^{1-D} \quad (1)$$

For the Euclidean case that p is independent of ξ then $D = 1$. At the other extreme, a “space-filling” curve that passes through every resolved point in a unit area has $D = 2$. Lovejoy (1982) first measured D for clouds by relating individual cloud perimeters p to cloud areas a using the expression $p \propto \sqrt{a}^D$. A measured value of $D = 1.35 \pm 0.05 \approx 4/3$ has since been adopted as the canonical value describing individual clouds (Siebesma and Jonker, 2000; Christensen and Driver, 2021).

40 A “monofractal” has a constant value of D , for the case that the scaling of p with ξ (e.g., by a power-law) is the same at all length scales, and so is “scale invariant.” For such scale invariance to apply to an atmospheric cloud field, this would require that the physics controlling cloud shapes is unchanged with measurement resolution, at least between the limits of possible cloud sizes. While clouds have been shown to be broadly scale invariant for the number distributions of cloud perimeters (DeWitt et al., 2023), cloud shapes might better be described as being multifractal, where D is a continuous function of
45 threshold used to define cloud (Lovejoy and Schertzer, 1990; Marshak et al., 1995; Lovejoy and Schertzer, 2006). Various studies have shown that D can vary considerably from cloud to cloud and even within different regions of the same cloud. For example, Batista-Tomás et al. (2016) found distinct fractal dimension values for cirrus with ragged, tenuous edges of $D = 1.37$, whereas for cumulonimbus, with smoother edges, $D = 1.18$. Other analyses of cumulus fields have found $D = 1.28$ (Zhao and Di Girolamo, 2007) and $D = 1.19$ (Mieslinger et al., 2019) using the expression $p \propto \sqrt{a}^D$. Cahalan and Joseph (1989) reported
50 $D = 1.27$ for small clouds and $D = 1.56$ for large clouds, supported by Benner and Curry (1998) who found $D = 1.23$ and $D = 1.34$ respectively. Furthermore, after reexamining the data in Lovejoy (1982), Gifford (1989) noted that D increases from 1.35 to 1.77 for the largest clouds with areas $> 2.5 \times 10^4 \text{ km}^2$. Inclusion of holes in the measured cloud areas (Peters et al., 2009) and merged clouds (Cahalan and Joseph, 1989) have been theorized to affect calculations of D .

This multifractal size and type dependence of D seems to contradict the arguments of Lovejoy (1982) that cloud fractal
55 properties are consistent across scales. However, a monofractal, scale-invariant assumption might more reasonably describe a large ensemble of clouds considered over a sufficiently long period of time and space. Indeed, the topic of whether or how scale invariance can describe atmospheric structures has been the topic of decades of debate.



The pioneering work of Richardson (1926) showed that the turbulent eddy diffusivity K , measured using the relative motion of pairs of particles separated by distance ℓ , followed a power-law with a 4/3 exponent from the millimeter scale for molecular diffusion to the length scale of atmospheric cyclones ($\ell \sim 10^3$ km), $K \propto \ell^{4/3}$, termed the Richardson “4/3 law” of atmospheric diffusion.

The scaling exponent of the diffusivity with respect to length scale can be obtained experimentally from measurements of velocity fluctuations Δv of two air parcels separated by a distance ℓ using passive scalars Θ , as a physical quantity that is affected by but does not affect the turbulent flow, such as the concentration of aerosols (Celani et al., 2002). Along one dimension x , the generalized first-order “structure function” expresses the covariance of Θ as a function of separation distance ℓ . For turbulent scalars, the function tends to be a power-law given by

$$S(\ell) = \Delta\Theta(\ell) = \langle \Theta(x + \ell) - \Theta(x) \rangle \propto \ell^{\mathcal{H}} \quad (2)$$

where brackets indicate averaging over many iterations of the experiment, and \mathcal{H} is the Hurst exponent¹ bounded by $0 < \mathcal{H} < 1$ (Hentschel and Procaccia, 1984; Lovejoy and Schertzer, 2012).

The 4/3 law was later derived using dimensional reasoning applied to the theory of 3D isotropic turbulence developed by Kolmogorov (1941). In the theory, for a fluid with kinematic viscosity ν , turbulence kinetic energy is passed along an energy cascade, from large eddies of the energy input scale L to progressively smaller eddies with a constant kinetic energy dissipation rate ε , ending at the “Kolmogorov microscale,” $\eta \sim (\nu^3/\varepsilon)^{1/4} \sim 1$ mm, a dissipation length scale where inertial and viscous forces balance. Through dimensional analysis, the covariance of air parcel velocity fluctuations was derived to be $\Delta v \approx \varepsilon^{1/3} \ell^{\mathcal{H}}$, where $\mathcal{H} = 1/3$ for the case of 3D isotropic turbulence. The dimensional approximation that $K \sim \ell v$ (Tennekes and Lumley, 1972) results in $K \sim \varepsilon^{1/3} \ell^{4/3}$, reproducing Richardson’s 4/3 power-law, and implying that the relationship between diffusivity and the Hurst exponent \mathcal{H} follows

$$K \sim \ell^{1+\mathcal{H}} \quad (3)$$

As Sect. 5 elaborates, the value of \mathcal{H} depends on the dimensionality of the turbulence. The problem that 3D turbulence cannot apply at the “flatter” planetary scales to a relatively thin troposphere has been well known. Even Kolmogorov predicted that 3D turbulence can only apply in the atmosphere at scales < 100 m. This led to the paradigm that 3D isotropic turbulence must be applicable at small scales and 2D at large scales, separated by a scale break around the depth of the troposphere (See Lovejoy (2023) for a historical review.). The contrasting case for 2D turbulence was developed for the case of an incompressible fluid (Kraichnan, 1967), where the expected value is instead $\mathcal{H} = 1$.

Figure 1 illustrates how two- and three-dimensional components in cloud structures are visible at all scales, but arguably 2D structures predominate at scale L , becoming more 3D approaching η , reflecting a scale-dependence due to large-scale stratification. Aircraft measurements of turbulent spectra of wind and temperature fluctuations have been argued to support this, where quasi-two-dimensional structures are seen at large scales and isotropic three-dimensional structures at small scales

¹The Hurst exponent has various applications in other fields, but here we employ its common usage in the field of fractal geometry to relate turbulent fluctuations with separation distance ℓ .




Figure 1. Diagram showing the similarity of rotational motions of clouds from the planetary diameter $L \sim 2a$, where structures are nearly 2D, to smaller scales with more 3D structure. At L (left), swirling features associated with synoptic-scale systems are $\sim 10^4 \text{ km}$ long and nearly 2D compared to the tropospheric depth H . At smaller resolved scales, the vertical component is more similar to the horizontal component, and thus the structure is more 3D. The images on the left are from EPIC (top) and a MODIS and VIIRS composite (bottom) for the same date. The upper right inset shows cloud features shaped by von Kármán vortices viewed near the Canary Islands obtained by VIIRS with eddy length scales $\sim 10 \text{ km}$. The image on the bottom right shows swirling clouds in a thunderstorm photographed from the ground with a length scale $\sim 0.01 \text{ km}$. The bottom inset is a cartoon depicting the smallest length scale of turbulence, the Kolmogorov microscale η , where kinetic energy is dissipated to heat through molecular diffusion K_η , with individual cloud droplets illustrated as dots with spacing to represent the cloud edge interface.

(Fiedler and Panofsky, 1970), with a scale break seen between approximately 20 km and 500 km (Nastrom et al., 1984; Gage and Nastrom, 1986). Lovejoy et al. (2007) (hereafter L07) argued that this scale break owes to vertical aircraft movements that occur when flying along isobars rather than isoheights and proposed instead that 3D isotropic turbulence is inapplicable at nearly any scale because stratification compresses the atmosphere vertically, even for scales as small as 5 m.

Specifically, L07, and more comprehensively Lovejoy and Schertzer (2018), provided evidence that the atmosphere is best characterized by a consistent intermediate turbulence regime at all scales following the framework of generalized scale in-



95 variance (GSI), which accounts for stratification in the “23/9D” elliptical model of turbulence in the atmosphere (Schertzer
and Lovejoy, 1985). Power spectra of radar reflectivity, cloud radiance, wind speed, and temperature all revealed length-
scaling exponents that lie between purely 2D and 3D turbulence cases, consistent with an intermediate turbulence regime
predicted to have a (fractal) volume dimension of $D = 2.55 = 2 + \mathcal{H}$ where $\mathcal{H} \approx 0.55$ (Schertzer and Lovejoy, 1985; Lovejoy
and Schertzer, 1985; Lovejoy et al., 1993; Lovejoy, 2021). In these cases, \mathcal{H} is calculated from the power spectrum of the
100 observed phenomenon, $E(k) \sim k^{-B}$, where $B = 2\mathcal{H} + 1$.

Simplifications of the first-order structure function have also been used to determine \mathcal{H} for properties of clouds (Pressel and
Collins, 2012; Pressel et al., 2014), and to link the dimension of turbulence to the fractal dimension through the expression
 $D = 2 - \mathcal{H}$ (Hentschel and Procaccia, 1984; Mandelbrot, 1985). Observations of scaling behaviors in clouds, whether expressed
through the fractal dimension or turbulent structure functions, point to a robust relationship between ξ , cloud geometry, and
105 turbulence. This paper explores the topic as follows. In Sect. 2, we relate the Hurst exponent to an “ensemble” fractal dimension
 D_e that defines a globally distributed cloud field and discuss in Sect. 3 a resolution coarsening procedure to measure it. Section
4 presents the values of the ensemble fractal dimension obtained using several satellite and numerical model datasets. Section
5 interprets the significance of the results by comparing them to the expected values of D_e and \mathcal{H} for 2D and 3D isotropic
turbulence, as well as for an intermediate turbulence regime that combines the two at all scales. Our findings support the
110 concept of a scale-invariant 2D-3D intermediate turbulence regime proposed by Schertzer and Lovejoy (1985), described in
detail by Lovejoy and Schertzer (2018), that we show applies to cloud perimeters over a remarkable 11 orders of magnitude
from the Kolmogorov microscale η to the planetary diameter $2a$.

2 Analytical expressions relating the perimeter of cloud ensembles to the dimension of turbulence

To explore how cloud perimeter varies with measurement resolution ξ , the total perimeter of a cloud ensemble viewed from
115 above (e.g., looking down as a satellite would view it from space) can be expressed in terms of a “perimeter density.” The
perimeter density \mathcal{P} is defined as the summed perimeters p of all clouds $P = \sum p$ normalized by the area of the horizontal
domain A_d ; that is, $\mathcal{P} = P/A_d$, a quantity analogous to the cloud fraction $\mathcal{A} = A/A_d$ where A is the total cloud area. In this
section, we show how \mathcal{P} can be related to ξ through the Hurst exponent \mathcal{H} .

2.1 Cloud perimeter and the Hurst exponent

120 In Garrett et al. (2018), the total cloud edge perimeter P of a tropical convective cloud field was estimated theoretically for
equal horizontal and vertical resolutions $\xi_H = \xi_V = \xi$ within a domain volume $V = A_d \xi_V$. To obtain P , a “mixing engine”
framework was introduced, that described cloud edge circulations consisting of coupled large-scale vertical buoyancy oscilla-
tions and horizontal turbulent exchanges as shown in Fig. 2. The derivation reflects a dimensional balance between two speeds.
In the horizontal, $v_H = K\mathcal{P}$ represents a speed of erosion or formation of cloud edge due to dissipative mixing with a character-
125 istic length scale ξ_H . The speed in the vertical direction is $v_V = \mathcal{N}\xi_V$ where \mathcal{N} is the moist adiabatic Brunt-Väisälä frequency,
and represents the speed of production of potential energy through oscillatory vertical motions. Assuming steady-state and that




Figure 2. Illustration of the theorized cloud edge mixing engine from Garrett et al. (2018). The circulations are generated from the production and dissipation of buoyant potential energy at the planetary scale. All clouds in the domain area A_d of Earth’s surface are represented as a single cylinder with total perimeter $P = \sum p$. The total available buoyant potential energy is dissipated vertically through moist adiabatic convection with vertical buoyancy speed $v_V \sim \mathcal{N}\xi_V$, and horizontally via turbulent mixing at cloud edge with speed $v_H \sim KP$. Globally, the vertical and horizontal components must balance.

the speeds of the horizontal and vertical legs of the circulation are equal, $v_V = v_H$, then it follows that

$$KP = \mathcal{N}\xi_V \quad (4)$$

Invoking mass continuity for the cloud edge circulation, $\nabla \cdot \mathbf{v} = 0$ leads to $\partial v_V / \partial \xi_V = -\partial v_H / \partial \xi_H$, and through scale analysis, $\mathcal{N}\xi_V / \xi_V \sim KP / \xi_H$. Thus, where ξ_H is the horizontal measurement resolution ξ viewed from space,

$$\mathcal{P}_\xi \sim \frac{\mathcal{N}\xi}{K_\xi} \quad (5)$$

where K_ξ is the turbulent eddy diffusivity with eddy length scale ξ .

From Eq. (3), K_ξ scales with \mathcal{H} , the value of which varies depending on the dimensionality of turbulence reflecting any anisotropy between ξ_V and ξ_H . The adjustment needed to scale K_ξ from the molecular diffusivity at η (i.e., K_η , the diffusion coefficient of air) to the resolution ξ is (Richardson, 1926; Garrett et al., 2018)

$$K_\xi = K_\eta \left(\frac{\xi}{\eta} \right)^{1+\mathcal{H}} \quad (6)$$

Substituting Eq. (6) into Eq. (5), the expected relationship relating measurement resolution to the cloud perimeter density \mathcal{P}_ξ becomes:

$$\mathcal{P}_\xi = \frac{\mathcal{N}\eta}{K_\eta} \left(\frac{\eta}{\xi} \right)^\mathcal{H} \propto \xi^{-\mathcal{H}} \quad (7)$$

An observed value of \mathcal{H} is then obtainable from measurements of \mathcal{P} as a function of ξ .

2.2 The fractal dimension of cloud ensembles

Equation (7) expresses the cloud perimeter as a function of resolution, and is thus analogous to Eq. (1) where $p \propto \xi^{1-D}$ with fractal dimension D . The canonical value for individual clouds is $D \approx 4/3$ (Lovejoy, 1982; Siebesma and Jonker, 2000; Christensen and Driver, 2021), but there are complications with this expression for D , including the aforementioned multifractal



145 nature of clouds. Additionally, D in the expression $p \propto \sqrt{a}^D$ only applies mathematically to the shape of an individual cloud or a set of identically shaped clouds (Imre, 1992).

From a climatological perspective, it is instead the ensemble of clouds with total perimeter density \mathcal{P} that governs exchanges of energy and air across cloud edges. Accordingly, we introduce an “ensemble fractal dimension” for clouds D_e analogous to Eq. (1) such that

$$150 \quad \mathcal{P}_\xi \propto \xi^{1-D_e} \quad (8)$$

implying from Eq. (5), that the scaling exponent of the diffusivity is equivalent to the ensemble fractal dimension:

$$K_\xi \propto \xi^{D_e} \quad (9)$$

The distinction between D and D_e was first raised by Mandelbrot (1977), who showed that an ensemble of fractal “islands” with power-law distributed areas a follows the Korčák Law, a survival function $S(a' > a) \propto a^{-\mathcal{K}}$ (Korčák, 1938), where the
 155 ensemble fractal dimension of the total coastline perimeter P is $D_e = 2\mathcal{K}$ confined to the bounds $1 \leq D < D_e \leq 2$.

The survival function can be related to a cumulative distribution function (CDF) through $S = 1 - CDF$ and \mathcal{K} is equivalent to the exponent of the power-law number distribution (Clauset et al., 2009). The area number distribution can be expressed as $n_a \propto a^{-(1+\alpha)}$ for clouds (Cahalan and Joseph, 1989; Benner and Curry, 1998; Wood and Field, 2011), and $\mathcal{K} \sim \alpha$. The perimeter number distribution, $n_p \propto p^{-(1+\beta)}$, is related to that for the area through $\alpha = D\beta/2$ for clouds (DeWitt et al., 2023).
 160 It follows that the ensemble fractal dimension is given by $D_e = D\beta$. The inequality $D < D_e$ requires that $\beta > 1$.

Comparing the exponents in Eqs. (7) and (8), the Hurst exponent can be related to the ensemble fractal dimension through

$$\mathcal{H} = D_e - 1 \quad (10)$$

This equation is important because it provides a means for linking satellite observations of cloud perimeter fractal properties and size distributions to the less easily seen but more physically relevant turbulent structures at cloud edge. For comparison
 165 with a LES model of a tropical cloud field resolved at 100 m scales, Garrett et al. (2018) applied a value of $\mathcal{H} = 1/3$ to Eq. (6) consistent with Richardson (1926) and the 4/3 law. Implicit in this case is an assumption of 3D isotropic turbulence at resolved scales.

However, while $D = 4/3$ is consistent with values seen for individual clouds, a larger value is required for cloud ensembles, in which case the inequality $D < D_e$ predicted by Mandelbrot (1977); DeWitt et al. (2023) applies. To allow for an adjustment
 170 to the individual fractal dimension, Hentschel and Procaccia (1984) related the perimeter fractal dimension of clouds to \mathcal{H} through the expression $D = 2 - \mathcal{H}$, but required a correction for intermittency in turbulence μ , where $D_\mu = (4 + \mu)/3 \approx 5/3$. We obtain, from Eq. (10) and $D = 2 - \mathcal{H}$, an adjustment to D for an ensemble of clouds:

$$D_e = 3 - D \quad (11)$$

The quantity $3 - D$ has been described as the fractal intermittency of turbulence within a 3D volume (Mandelbrot, 1977;
 175 Hentschel and Procaccia, 1984). Applying the canonical value of $D = 4/3$ for individual clouds leads to the expected value



of $D_e = 5/3$. This is in agreement with Hentschel and Procaccia (1984), who found that Richardson's 4/3 law only applies if the fractal dimension is $D_\mu = 5/3$, obtained by adding an intermittency correction with a value between $0.25 < \mu < 0.5$ to the value $D = 4/3$. The 5/3 value is also nearly identical to the value of $D_e = 1.68 \pm 0.06$ obtained from $D_e = D\beta$ for $\beta = 1.26$ from DeWitt et al. (2023), which applied across various satellite instruments and climate regimes. In this case, the implied
180 value of the Hurst exponent is $\mathcal{H} = 2/3$, which is between the 3D turbulence value of $\mathcal{H} = 1/3$ and the 2D turbulence value of $\mathcal{H} = 1$. Using Eq. (8) and the methods below, we observationally evaluate the applicability of the result $\mathcal{H} = 2/3$ that is associated with $D_e = 5/3$.

3 Data and methods

Equation (10) implies that the dimensionality of the turbulent structure in clouds can be inferred from observations of cloud
185 perimeters. To explore this hypothesis and the suggestion from Eq. (11) that $\mathcal{H} = 2/3$, we consider satellite imagery of clouds from platforms in polar-orbiting, geostationary, and heliocentric orbits using cloud mask algorithms. The resulting binary arrays, hereafter "cloud masks," represent cloudy pixels with a value of unity and clear skies by a value of zero. Cloudy pixels are considered connected when they are vertically or horizontally adjacent (i.e., "4-connectivity"). The edges of the domain are not included as part of the perimeter. p and a are calculated for all individual cloudy regions, which are summed and normalized
190 by domain area A_d to determine \mathcal{P} and \mathcal{A} .

The polar-orbiting data sets considered are from the instruments VIIRS and MODIS, which have native pixel resolutions ξ_N at the nadir of 0.75 km and 1 km, and capture imagery in narrow, meridional swaths. Values of ξ_N represent the pixel resolution at satellite nadir; the horizontal and vertical dimensions of each pixel are adjusted based on their distance from the nadir. The average swath size for VIIRS is 2501×12944 pixels with a domain area A_d of 2.6×10^7 km². For MODIS it is
195 1261×8120 pixels with $A_d = 2.0 \times 10^7$ km². The VIIRS and MODIS datasets include 60 and 72 cloud masks from 02 June 2021. Their respective cloud mask techniques are described by Kopp et al. (2014) and Ackerman et al. (2008). We also include 12 MODIS cloud masks with 0.25 km resolution as defined by optical reflectance values $R \geq 0.01$ described by DeWitt et al. (2023). These high-resolution cloud masks have $A_d = 5.1 \times 10^6$ km² and average image dimensions of 5048×8120 pixels obtained from 01 January 2021 through 09 January 2021.

200 Geostationary datasets are obtained from instruments denoted here by their more familiar satellite names, Himawari (instrument name: AHI), GOES-WEST (ABI), and METEOSAT-11 (SEVIRI), which provide full-disk imagery of Earth with $A_d = 1.0 \times 10^8$ km² positioned over the fixed longitudes 141°E, 137°W, and 0°, respectively. Their nadir pixel resolutions are 2 km, 2 km, and 3 km, respectively, with cloud masks as described by Derrien and Gléau (2005). For each of the geostationary datasets, 30 cloud masks were obtained from 02 June 2021 through 01 July 2021, each at approximately the local noon at the
205 satellite nadir.

To provide unique observations of global cloud coverage, we also include cloud masks from GEO-Ring and EPIC. GEO-Ring is a composite of geostationary satellite imagery (Ceamanos et al., 2021) that provides stitched satellite imagery of the surface of the Earth (excluding the poles) with $A_d = 4.4 \times 10^8$ km² at $\xi_N = 11$ km. Thirty-nine GEO-Ring cloud masks were



obtained from 02 June 2021 through 21 June 2021. EPIC obtains full-disk imagery of Earth from the DSCOVR satellite in
210 heliocentric orbit, photographing Earth as it rotates, providing coverage of all longitudes. Due to its location at the L1 Lagrange
Point in deep space, EPIC imagery has a coarser pixel resolution of 8 km. 30 EPIC cloud masks (described by described by
Yang et al. (2019)) were obtained from 01 June 2017 through 30 June 2017.

As a means to compare measurements of \mathcal{P}_ξ from satellite observations to the value derived by Garrett et al. (2018), we
include measurements of clouds from the System for Atmospheric Modeling (SAM), a high-resolution 3D LES, initialized with
215 idealized GATE Phase III campaign soundings for tropical convection. The simulation domain of $204.8 \text{ km} \times 204.8 \text{ km} \times 19 \text{ km}$
includes more than one billion grid points – often referred to as a “Giga-LES.” There are 2048 grid points in the horizontal
directions with a grid spacing of 100 m, and 256 grid points in the vertical with grid spacing ranging from 50 m to 100 m.
The simulation is integrated at two-second intervals for 24 hours. Refer to Khairoutdinov et al. (2009) for a more complete
description of the simulation. We analyze scenes hourly from hour 12 through hour 24 of the model to ensure that steady-state
220 has been reached.

In order to compare the 3D model data with 2D satellite retrievals, we define the SAM cloud masks as 3D binary arrays
for mixing ratios of non-precipitating cloud condensate $q_n > 0.01 \text{ g kg}^{-1}$. A 2D facsimile of a satellite cloud mask is created
from a vertical projection of the 3D cloud mask that represents the view from above. The 2D binary cloud mask is assigned a
cloudy pixel where the corresponding vertical columns of the 3D cloud mask have \sum_0^H (cloudy pixels) $> j$. For example, with
225 a threshold value of $j = 3$, each pixel in the 2D cloud mask is classified as cloudy if more than three of the pixels in the cor-
responding 3D vertical column are cloudy. Multiple 2D cloud masks were obtained using threshold values $j = 1, 2, 3, 5, 9, 15$.
This thresholding procedure is similar to an analysis performed by DeWitt et al. (2023) that compared differences in cloud
statistics defined by various optical depth thresholds.

To obtain values of D_e , total cloud perimeter P is calculated first at the native spatial resolution ξ_N and normalized by A_d
230 to obtain \mathcal{P}_N . The image is then artificially coarsened and the procedure is repeated. \mathcal{P}_ξ is obtained at progressively coarser
spatial resolutions $\xi > \xi_N$ such that $\xi = \xi_N k$, where k is the coarsening factor. The image is coarsened by reducing the number
of pixel elements by k^2 . Each pixel of the coarsened image is determined to be cloudy or clear by rounding the average of the
values from the corresponding pixels in the native resolution image. The values of k are chosen to be the nearest odd integers
that differ by a constant factor (e.g., $k = 2^n$). The maximum value of k for each dataset corresponds to coarsening to a single
235 pixel along the shorter dimension of the domain. Figure 3 shows an example of resolution coarsening to a single pixel for
various EPIC cloud masks.

A least squares linear regression is performed on values of $\ln \mathcal{P}_\xi$ and $\ln \xi$ to obtain the Hurst exponent \mathcal{H} and D_e from Eqs.
(7) and (8). Uncertainties in the linear regression are evaluated at the 95% confidence level.

4 Results: Cloud measurements

240 Alongside measurements of perimeter density \mathcal{P}_ξ , the more familiar quantity of cloud fraction \mathcal{A}_ξ is included as a point of
comparison. Figure 4 shows the cloud fraction and perimeter density obtained from satellite and model datasets at native




Figure 3. EPIC cloud masks shown at native resolution ξ_N and coarsened resolutions ξ to a single pixel for four cases with initial native cloud fraction between $0.48 < \mathcal{A}_N < 0.58$ (increasing from top to bottom) illustrating a bifurcation of cloud fraction with coarsening of resolution depending on the native cloud fraction to either zero or unity. Note that the single pixel case shown here has a value of $\mathcal{A} = 0.37$ rather than unity because the domain area represented by the square pixel is the disk area $A_d = \pi a^2$.

resolution ξ_N , termed \mathcal{A}_N and \mathcal{P}_N , and coarsened resolutions ξ normalized by ξ_N . Both ξ_N and the domain areas A_d span two orders of magnitude: ξ_N from 0.1 km to 11 km, and A_d from 5.1×10^6 to 4.4×10^8 km².

4.1 Measured cloud fraction \mathcal{A}

245 Global cloud fraction values \mathcal{A}_N in Fig. 4a range from 0.5 to 0.7, reflecting differences in cloud mask techniques. With progressive coarsening, \mathcal{A} changes by less than 5% before bifurcating at $\xi/\xi_N \sim 100$. As ξ approaches $\xi/\xi_N \sim 1,000$, \mathcal{A} is represented by a single pixel, with a value of either zero or unity (except for the polar-orbiting satellites, which are represented by a 1×5 line). Interestingly, geostationary cloud fraction measurements with $\mathcal{A}_N > 0.56$ approach a value of unity, whereas MODIS 0.25 km and EPIC datasets with $\mathcal{A}_N = 0.55$ instead trend toward zero.

250 This bifurcation of cloud fraction reflects that as an image of a cloud field is coarsened to a single pixel, that pixel will either have a value of zero or unity, depending on whether the native cloud fraction is greater or less than $\mathcal{A}_N = 0.55$. Figure 3 shows four examples of coarsening EPIC cloud masks to a single pixel, resulting in either a single clear or cloudy pixel. Although statistically the initial cloud fraction value is a good indicator of whether the single pixel will be cloudy or clear, it is not the only factor. For $\mathcal{A}_N = 0.55$, the single-pixel value of \mathcal{A} depends more on the initial *distribution* of clouds. Where




Figure 4. Measured cloud fraction \mathcal{A}_N (a) and perimeter density \mathcal{P}_N (b) at native measurement resolution ξ_N (solid markers), and \mathcal{A}_ξ and \mathcal{P}_ξ at coarsened resolutions ξ normalized by ξ_N (lines and hollow markers) for polar-orbiting (blue), full-disk (green), and global mosaic (yellow) satellite datasets. The SAM numerical simulations are shown as pink diamonds (with brightness scaled by threshold value j). Legend entries are sorted by increasing ξ_N with the associated negative Hurst exponent \mathcal{H} obtained from a least squares linear regression of $\ln \mathcal{P}_\xi$ and $\ln \xi$ (Eq. (7)) and uncertainties evaluated for a 95% confidence interval.

255 clouds are more evenly distributed across the globe (Fig. 3b), smaller isolated structures vanish more quickly with coarsening and approach $\mathcal{A} = 0$ for a single pixel. When clouds are more clustered (Fig. 3c), a coarsened single pixel has $\mathcal{A} = 1$.

The application of vertical pixel thresholding in SAM results in a wide range of native cloud fraction values between $0.15 < \mathcal{A}_N < 0.60$. Larger threshold values j tend to exclude small and shallow clouds, and in turn, decrease the overall cloud fraction. Bifurcation of \mathcal{A} occurs for SAM at a native value of $\mathcal{A}_N \approx 0.45$, notably smaller than the value at which bifurcation
 260 occurs for satellite datasets at $\mathcal{A}_N \approx 0.55$. This discrepancy suggests a difference between the clustering behavior of clouds viewed globally by satellite and those of modeled clouds for a region of tropical convection.

4.2 Measured cloud perimeter density \mathcal{P}

The resolution dependence of cloud perimeter density \mathcal{P} can be defined more simply than for cloud fraction \mathcal{A} . As shown in Fig. 4b, perimeter density \mathcal{P}_ξ has a power-law scaling with ξ in all datasets, independent of satellite orbit, domain size, and
 265 resolution. For $\xi > \xi_N$, \mathcal{P}_ξ is well characterized by a linear regression of $\ln \mathcal{P}_\xi$ to $\ln \xi$ (Eq. (7)). This power-law relationship $\mathcal{P}_\xi \propto \xi^{-\mathcal{H}}$ holds even past the point $\xi/\xi_N \sim 100$ where cloud fraction values \mathcal{A} tend to diverge. However, for very large $\xi/\xi_N \sim 1,000$, \mathcal{P}_ξ can deviate from the power-law regression to lower values, reflecting the fractal nature of the problem: complex cloud structures cannot be fully represented by coarse Euclidean geometries such as a single square pixel. This low bias in \mathcal{P}_ξ for values of ξ/ξ_N between ~ 100 and ~ 1000 can also be seen in the fourth and fifth columns of Fig. 3. There,
 270 for $\xi/\xi_N = 243$ the images appear pixelated, but maintain their general structure. However, for $\xi/\xi_N = 729$, the cloud mask



consists of either a single cloudy or a clear pixel. The value of ξ/ξ_N at which \mathcal{P} begins to depart from the linear regression corresponds to the coarsest resolution for which the complexity of the cloud edge can still be reliably measured.

Notably, the value of \mathcal{P}_N for the native resolution $\xi/\xi_N = 1$ does not always align with the best-fit line, especially for the case of EPIC. As discussed in DeWitt et al. (2023), EPIC employs an on-board averaging and post-processing interpolation that artificially smooths the edges of clouds to compress data for transmission. This interpolation results in the artifact that \mathcal{P}_N is lowered due to the reduced edge complexity. A similar phenomenon is observed to a lesser degree for the other satellite datasets. To avoid this issue, the \mathcal{P}_N data points are not included in the linear regression.

For satellite datasets, values of \mathcal{H} lie in the range $0.71 < \mathcal{H} < 0.84$, with a mean value of $\mathcal{H} = 0.78$ with uncertainty evaluated at the 95% confidence interval of 0.09. The ensemble fractal dimension that corresponds to the total cloud perimeter given by Eq. (10) is $D_e = \mathcal{H} + 1 = 1.78 \pm 0.09$. Calculated values of \mathcal{H} from the satellite datasets do not appear to depend on the type of satellite orbit or resolution, but they are significantly larger than those found for modeled clouds. \mathcal{P}_ξ measured from SAM follows a power-law with exponent values ranging from $0.60 < \mathcal{H} < 0.71$ depending on threshold value j . The average value of $\mathcal{H} = 0.67 \pm 0.08$ is two standard deviations smaller than the the satellite datasets.

Note that modeled values of \mathcal{H} lie closer to the value of $1/3$ expected for 3D isotropic turbulence than is the case for the satellite datasets, perhaps reflecting the smaller domain area and atmospheric regime or the subgrid-scale turbulence assumption used in LES models. In general, increasing the threshold value j (which determines the minimum vertical cloud thickness required for 2D cloud masking) leads to smaller values of \mathcal{H} , reflecting the multifractal nature of clouds. For example, for a detection threshold of $j = 0$, all cloudy pixels in the domain are considered and $\mathcal{H} = 0.71$. Meanwhile, for the highest detection threshold value of $j = 15$, and $\mathcal{H} = 0.61$. The latter case requires that only the largest overlapping cloud structures are included in the analysis, leaving most small, shallow clouds omitted. The smallest clouds are only observed with the finest resolution, resulting in a shallower linear regression slope for more highly thresholded cloud scenes.

5 Discussion

To summarize the observations, global cloud perimeter density \mathcal{P} is much more sensitive than cloud fraction \mathcal{A} to measurement resolution ξ , but the dependence is also much more simply mathematically characterized. The observed power-law scaling relating \mathcal{P} to ξ is remarkably similar for imagery from a wide range of satellite platforms. We measured an ensemble fractal dimension of $D_e = 1.78 \pm 0.09$, corresponding to a Hurst exponent of $\mathcal{H} = 0.78 \pm 0.09$. Similarly, from DeWitt et al. (2023), $D_e = D\beta \simeq 1.68 \pm 0.06$ derived from satellite observations of the perimeter distribution power-law exponent $\beta = 1.26 \pm 0.06$ and assuming $D = 4/3$ for individual clouds.

To account for how the dimensionality of turbulence may help explain the difference between the measured value of $D_e = 1.78 \pm 0.09$ for satellite observations (Fig. 4) and the theoretical value of $5/3$ implied by Eq. (11), we compare D_e with canonical values of \mathcal{H} associated with 2D, 3D, and “intermediate” turbulent regimes and explore “limiting cases” that correspond to possible upper and lower bounds of \mathcal{P}_ξ evaluated at the planetary scale and the Kolmogorov microscale.



5.1 Scaling exponents for 3D, 2D, and intermediate turbulence regimes

As introduced in Sect. 1, the theory of 3D isotropic turbulence predicts that the length dependence of turbulent diffusivity follows $K \sim \varepsilon^{1/3} \ell^{4/3}$, i.e., Richardson's 4/3 law. Within the context of resolved clouds, we express ℓ as the resolved eddy length ξ , assuming that the smallest resolved cloud features are shaped by turbulent eddies of that size. In this case, the turbulent diffusivity scaling expression for cloud edges resolved at scale ξ is

$$K_{\xi,3D} \propto \xi^{4/3} \quad (12)$$

Following from Eq. (3), $K \sim \ell^{1+\mathcal{H}}$, the implied scaling exponent for velocity fluctuations in 3D isotropic turbulence is $\mathcal{H} = 1/3$.

For 2D isotropic turbulence, where vertical motions are negligible, due to e.g., stratification, the diffusivity scaling exponent can be obtained from dimensional analysis with the conserved property being enstrophy \mathcal{E} — the integrated 2D vorticity squared — instead of ε . The dependence of \mathcal{E} on the eddy length scale ℓ is $\mathcal{E}(\ell) \sim \Phi^{2/3} \ell^3$ where Φ is the enstrophy flux density with units s^{-3} (Kraichnan, 1967; Charney, 1971). The velocity scaling exponent is $v \sim \Phi^{1/3} \ell$, and substituting v into $K \sim v\ell$, the 2D turbulent diffusivity scaling becomes

$$K_{\xi,2D} \sim \xi^2 \quad (13)$$

and from Eq. (3), the implied scaling exponent for velocity fluctuations in 2D turbulence is $\mathcal{H} = 1$.

The framework of generalized scale invariance (Schertzer and Lovejoy, 1985) allows for the derivation of an "elliptical dimension" D_{el} that applies to an "intermediate" 23/9D model of anisotropic turbulence at all scales in the atmosphere, rather than distinct regions of 2D isotropic turbulence at large scales and 3D isotropic turbulence at smaller scales. This continuous scaling accounts for the horizontal-vertical anisotropy of the atmosphere due to stratification and is determined by comparing velocity fluctuations Δv_H and Δv_V in the horizontal and vertical directions. Because stratification is only observable in vertical velocity perturbations, it has been observed that horizontal velocity fluctuations follow the 3D scaling $\Delta v_H \sim \varepsilon^{1/3} \ell^{1/3}$ up to the planetary scale (Lovejoy and Schertzer, 2018). The Bolgiano-Obukhov law (Bolgiano Jr, 1959; Obukhov, 1959) describes the corresponding vertical scaling relationship in buoyancy-forced turbulence $\Delta v_V \sim \phi^{1/5} \ell^{3/5}$, where ϕ is analogous to the potential energy dissipation rate ε in the vertical dimension with units $m^2 s^{-5}$. To account for this anisotropy in the vertical, \mathcal{H} for the combined turbulence case was derived from the ratio of the horizontal and vertical Hurst exponents $\mathcal{H}_H = 1/3$ and $\mathcal{H}_V = 3/5$, resulting in $\mathcal{H}_H/\mathcal{H}_V = 5/9 \sim 0.56$. From Eq. (10), the elliptical dimension becomes $D_{el} = 14/9 = 1.56$ (for the volume, $23/9 = 14/9 + 1$. See Lovejoy (2023) for a review.) From Eq. (3), the turbulent diffusivity for this intermediate 23/9D regime then scales as

$$K_{\xi,int} \sim \xi^{14/9} \quad (14)$$

Lovejoy et al. (2007) (L07) analyzed 5 m vertical resolution dropsonde wind datasets to determine the relationship $\Delta v_V \sim \ell_V^{\mathcal{H}_V}$ where ℓ_V is the vertical separation between measurements. The observed Hurst exponent ranged from $\mathcal{H}_V = 0.60$ — in



Theory		
Turbulence Regime	\mathcal{H} ($\mathcal{P}_\xi \sim \xi^{-\mathcal{H}}$)	D_e ($K_\xi \sim \xi^{-D_e}$)
3D Isotropic Turbulence Eq. (12)	1/3 (0.33)	4/3 (1.33)
$D_e = 5/3$ Eq. (11)	2/3 (0.67)	5/3 (1.67)
23/9D Elliptical Dimension (GSI)		
$\mathcal{H}_H/\mathcal{H}_V$ Eq. (14)	5/9 (0.56)	14/9 (1.56)
2D Turbulence Eq. (13)	1	2
Observations		
Vertical wind structure functions \mathcal{H}_V (L07)		
Tropopause: 12.6 km	0.77	1.77
Surface to 10 km	0.60 to 0.75	
Measured cloud perimeters (Figure 4b)		
Satellite	0.78 ± 0.09	1.78
SAM	0.67 ± 0.08	1.67

Table 1. Theorized values (top) of \mathcal{H} and D_e from the expressions $\mathcal{P}_\xi \sim \xi^{-\mathcal{H}}$ (Eq. 7), $K_\xi \sim \xi^{-D_e}$ (Eq. 9), and $\mathcal{H} = D_e - 1$ (Eq. 10), for the cases of 3D isotropic turbulence, $D_e = 5/3$, the 23/9D elliptical model from generalized scale invariance (GSI), and 2D turbulence. Observations (bottom) include \mathcal{H}_V from Lovejoy et al. (2007) for vertical wind profiles and the measurements obtained here shown in Fig. 4b. Values of \mathcal{H} for each case are compared in Fig. 6. Decimal values are shown alongside the derived fraction values for ease of comparison with observations.

agreement with the Bolgiano-Obukhov 3/5 scaling for $\ell_V < 1$ km — to $\mathcal{H} = 0.77$ for $\ell_V < 13$ km, the tropospheric depth. Increasing values of \mathcal{H}_V as ℓ_V approaches the tropospheric depth were argued to be consistent with more 2D turbulent structures influenced by upper-level jet shear.

Table 1 summarizes previously derived expressions for the scaling exponents \mathcal{H} and D_e for 3D, 2D and intermediate turbulence, along with their relationship to \mathcal{P} through Eq. (7), for comparison with the satellite and numerical model results obtained here. The exponent values in Eqs. (12)-(14) are labeled D_e following Eq. (9).

Observational values from Sect. 4 and from L07 are similar to the theoretically obtained values of $D_e = 5/3$ from Eq. (11) and the 23/9D model implying $D_e = 14/9 = 1.56$, and not to either of 2D or 3D isotropic turbulence. The $D_e = 5/3$ case is closest to the value of $D_e = 1.78 \pm 0.09$ obtained from satellite observations (Fig. 4b) and particularly to the range of values seen in SAM simulations ($1.61 < D_e < 1.72$).



5.2 Limiting cases: cloud perimeter density at the turbulent microscale and the planetary scale

345 Because cloud shapes and sizes are determined by objective physical processes that are independent of subjective measurement resolution, in principle it should be possible to infer information about cloud geometries from the physical properties of the planet and its atmosphere. To this end, we examine order-of-magnitude “limiting case” values for \mathcal{P} evaluated at the smallest and largest possible conceivable scales for clouds, expressed in terms of basic planetary and atmospheric parameters, and compare these with the observations shown in Fig. 4b.

350 Given the turbulent nature of fractal cloud edges, the Kolmogorov microscale η is the smallest theoretical resolution length scale ξ , for which \mathcal{P} is anticipated to be a maximum. Substituting ξ with η in Eq. (5) yields

$$\mathcal{P}_\eta = \frac{\mathcal{N}\eta}{K_\eta} \quad (15)$$

For the planetary scale (denoted with \oplus), Eq. (4) becomes $\mathcal{P}_\oplus \sim \mathcal{N}H/K_\oplus$ where H atmospheric scale height. The planetary-scale diffusivity is $K_\oplus = LU$ where $L = 2a$ and $U = \mathcal{N}H$ is the characteristic speed of the production and dissipation of moist
 355 convective potential energy in cloud edge circulations (described in Sect. 2). Thus

$$\mathcal{P}_\oplus \sim \frac{1}{2a} \quad (16)$$

This result is similar to the case where Earth is resolved as a single point source of light, or a “Pale Blue Dot,” as coined by Sagan (1994). The extremely idealized case of perimeter density resolved by a single pixel is $\mathcal{P}_\oplus = P_\oplus/A_\oplus$. Considering either a square with side length $\xi = 2a$ ($\mathcal{P} = 8a/(4a^2)$) or a circular dot with diameter $2a$ ($\mathcal{P} = 2\pi a/(\pi a^2)$), gives

$$360 \quad \mathcal{P}_{\oplus, PBD} = \frac{2}{a} \quad (17)$$

In either case, \mathcal{P}_\oplus is a function only of planetary radius a . Furthermore, each variable in Eq. (15) can be estimated from basic physical planetary properties, including the atmospheric composition, temperature, and pressure, as described in Appendix A.

5.3 Comparison between observations and theory

Figure 5 presents observations and theoretical predictions of \mathcal{P}_ξ . Theoretically derived estimates of \mathcal{P}_ξ are obtained from Eq.
 365 (7) for three cases: 3D turbulence ($\mathcal{H} = 1/3$ and $D_e = 4/3$), 2D turbulence ($\mathcal{H} = 1$ and $D_e = 2$), and the intermediate case $D_e = 5/3$. For clarity, the 23/9D case, which has a line nearly the same as the $D_e = 5/3$ case, is not included in Fig. (5). Satellite and SAM measurements are clearly aligned with the case that $D_e = 5/3$, as predicted by Eq. (11), lying distinctly between the curves corresponding to $D_e = 4/3$ for 3D isotropic turbulence and $D_e = 2$ for 2D turbulence. The limiting case for the Kolmogorov microscale \mathcal{P}_η marks the intersection of \mathcal{P}_ξ from Eq. (7) where $\xi = \eta$.

370 What is striking is how well the predicted value of $D_e = 5/3$ connects the highly idealized limiting case values of \mathcal{P}_\oplus and \mathcal{P}_η to the observed scaling for \mathcal{P}_ξ . The alignment is particularly remarkable considering that \mathcal{P}_η and \mathcal{P}_\oplus are obtained only from the physical properties of the planet and its atmosphere, and are separated by 11 orders of magnitude. This correspondence suggests that the statistical aspects of cloud geometries and atmospheric turbulence, D_e and \mathcal{H} , could in principle be inferred from knowing only a few basic physical parameters of a planet.




Figure 5. Measured perimeter density \mathcal{P}_ξ for the satellites and SAM shown as the same markers from Fig. 4, with the derived \mathcal{P}_ξ from Eq. (7) overlaid as gray ($\mathcal{H} = 1/3$ and $D_e = 4/3$ for 3D turbulence), blue ($D_e = 5/3$), and red ($\mathcal{H} = 1$ and $D_e = 2$ for 2D turbulence) dashed lines. The average scaling exponents $\overline{\mathcal{H}}$ are shown as solid green (satellite) and black (SAM) lines, with the mean and standard deviations in the legend. The limiting case value of \mathcal{P}_η from Eq. (15) is shown as a dark blue hexagram with the uncertainty indicated by shading. The limiting case \mathcal{P}_\oplus at the planetary diameter $\xi = 2a$ from Eq. (16) is a light blue hexagram and the Pale Blue Dot case from Eq. (17) is a light blue circle.




Figure 6. Visualization of theorized and observed \mathcal{H} . Theorized values of \mathcal{H} are shown as horizontal dashed lines for 2D turbulence (red), for $D_e = 5/3$ (blue), Bolgiano-Obukhov scaling (black), the 23/9D model from generalized scale invariance (GSI) (purple), and 3D turbulence (gray). Observations from Lovejoy et al. (2007) are shown (left) as horizontal gray dashed lines darkening as vertical separation distance ℓ_V increases from $\ell_V < 158$ m, to $\ell_V = 12.6$ km corresponding to the top of the atmosphere (TOA). Observations from this work (middle) are shown with symbols corresponding to Figs. 4b and 5 with the averages shown as horizontal green (satellite) and black (SAM) solid lines. On the right are markers corresponding to the slopes from \mathcal{P}_η (Eq. 15) to the values of \mathcal{P}_\oplus from Eqs. (16) and (17).

375 Figure 6 compares the observed and theoretical values of \mathcal{H} . The scaling relationship connecting microscale values (η, \mathcal{P}_η) to planetary values $(2a, \mathcal{P}_\oplus)$, as well as the scaling relationships inferred from observations lie between $1/3 < \mathcal{H} < 1$, the limits for 3D and 2D turbulence. The values are most consistent with the case $D_e = 5/3$, and to a lesser extent with the 23/9D intermediate turbulence regime obtained from generalized scale invariance (Schertzer and Lovejoy, 1985; Lovejoy et al., 2007; Lovejoy and Schertzer, 2018) with $\mathcal{H} \sim 5/9$.

380 Comparing the results here with observations of vertical wind structure functions by Lovejoy et al. (2007) (L07) in Fig. 6 and Table 1, it is worth noting that the variation in \mathcal{H} with threshold j and with ℓ_V reflects the multifractal nature of clouds. Values of \mathcal{H} for the smallest vertical separation distances in L07 ($\ell_V \sim 5$ m, $\mathcal{H} = 0.60$), and for cloud structures resolved vertically in SAM ($\xi_V \sim 100$ m, $\mathcal{H} = 0.67$), correspond most closely to the Bolgiano-Obukov scaling $\mathcal{H} \sim 0.6$. However, the value inferred from satellite observations ($\mathcal{H} = 0.78$) is most consistent with L07 ($\mathcal{H}_V = 0.77$) inferred from vertical separation distances of
 385 $\ell_V \sim H$. Despite these variations in \mathcal{H} , the observations of clouds reveal an intermediate turbulence regime that excludes both of the purely 2D or 3D isotropic turbulence cases.



6 Conclusions

The measured relationship between the ensemble cloud perimeter density \mathcal{P}_ξ seen from space and the resolution at which it is imaged ξ yields an “ensemble fractal dimension” D_e , a scaling exponent analogous to the individual cloud fractal dimension D . We conclude that D_e represents the degree to which turbulence is 2D or 3D, and corresponds simply to the Hurst exponent \mathcal{H} , the basis of a scaling law for quantifying turbulent fluctuations of atmospheric scalars, through $D_e = \mathcal{H} + 1$.

Global cloud measurements of \mathcal{P} from various satellite orbit types and a Large Eddy Simulation (LES) of tropical convection follow a consistent power-law scaling with respect to ξ across five orders of magnitude. The associated scaling exponent of $\mathcal{H} = 0.78 \pm 0.09$ that we obtained from satellite measurements lies between the theoretical values for isotropic 2D and 3D turbulence, consistent with a model of anisotropic 23/9D turbulence (Schertzer and Lovejoy, 1985).

Measured values of the ensemble fractal dimension D_e are also greater than the value of $D \sim 4/3$ that is often assumed to apply to individual clouds. The value obtained from SAM $D_e = 1.67 \pm 0.08$ is equal to the theorized value of $D_e \sim 5/3$ implied by Eq. (11)). The measured value from satellite imagery $D_e = 1.78 \pm 0.09$ is intermediate to the value of $D_e = 2$ expected for 2D turbulence and $D_e = 4/3$ for 3D turbulence. It is similar to a value of $D_e = 1.68$ suggested by DeWitt et al. (2023) for cloud ensembles, and to a theoretically derived value of $D_\mu \approx 5/3$ obtained by Hentschel and Procaccia (1984) for intermittent turbulence. The value of D_e from satellite data is significantly greater than that obtained from analysis of a detailed LES model of a tropical cloud field, suggesting natural cloud ensembles are more geometrically complex.

Values of \mathcal{P} evaluated at the Kolmogorov microscale η and the planetary diameter $2a$ purely from physical parameters lie remarkably in line with satellite observations and LES model calculations, despite being separated by 11 orders of magnitude in ξ . The value of \mathcal{P}_η was only inferred from the molecular composition, temperature, and pressure of clouds and the atmosphere, while \mathcal{P}_\oplus was inferred from the planetary radius a and the atmospheric depth H and stability \mathcal{N} .

Globally distributed, the total perimeter of clouds has a resolution dependence in satellite and numerical datasets, one that can be tethered to physically parameterized values evaluated at the Kolmogorov microscale and the planetary diameter, that points to existence of an intermediate 2D/3D turbulence regime that applies at all conceivable tropospheric scales. Observations of clouds on other planets in the solar system could help identify whether the observed scaling is specific to present-day Earth or in fact general to stratified atmospheres. Any generalization of the scaling laws could prove useful for constraining predictions of cloud behaviors in a future climate state on Earth, or for exoplanetary studies where — like Earth’s “pale blue dot” — only coarse-resolution physical parameters are available.

Code and data availability. The VIIRS and EPIC datasets were downloaded from NASA Earthdata (<https://www.earthdata.nasa.gov/>, NASA, 2023) and all others from the ICARE Data Center in Lille, France (<https://www.icare.univ-lille.fr/>, ICARE, 2023). Code used to analyze data and generate figures is available from the first author upon request.



Appendix A: Variables and parameters

Values of the parameters and variables used for calculation of \mathcal{P} in Eqs. (7) through (16) are shown in Table A1. Uncertainty in \mathcal{P}_η and η owes to the range of possible values of ε , ν , and K_η . Diffusivity and kinematic viscosity are proportional to atmospheric pressure p , so uncertainties include the range of values corresponding to the temperature T and p between the surface ($T = 300$ K, $p = 1$ bar) and the top of the troposphere ($T = 200$ K, $p = 0.1$ bar).

The Brunt-Väisälä frequency \mathcal{N} for a dry adiabat is typically expressed as a function of gravity g and vertical temperature profiles, but can also be expressed in terms of physical planetary parameters as $\mathcal{N} \sim g \left(\frac{S(1-\alpha)}{4\sigma} \right)^{-1/8} c_p^{-1/2}$ (Read et al., 2016) where S is the solar constant, α is the planetary albedo, σ is the Boltzman constant, and c_p is the specific heat at constant pressure. The value for a moist adiabat shown here is slightly less than, but of the same order of magnitude as the value for a dry adiabat of 0.01 s^{-1} (Mapes, 2001).

Parameters	Symbol	Units	Value	Notes
Planetary radius	a	km	6.37×10^3	
Scale height	H	km	8.50	
Brunt-Väisälä frequency	\mathcal{N}	s^{-1}	6.00×10^{-3}	Evaluated for a moist adiabat (Mapes, 2001)
Kolmogorov microscale	η	km	2.19×10^{-6}	$\eta \sim (\nu^3/\varepsilon)^{1/4}$
TKE dissipation rate	ε	$\text{km}^2 \text{ s}^{-3}$	3.00×10^{-9}	$10^{-10} < \varepsilon < 10^{-8}$ (Kantha and Hocking, 2011)
Kinematic viscosity of air	ν	$\text{km}^2 \text{ s}^{-1}$	1.86×10^{-11}	$1.5 \times 10^{-11} < \nu < 1.3 \times 10^{-10}$
Diffusion coefficient of air	K_η	$\text{km}^2 \text{ s}^{-1}$	2.42×10^{-11}	$2.3 \times 10^{-11} < K_\eta < 9.7 \times 10^{-11}$ (Schvertz and Brow, 1951)

Table A1. Values of variables and parameters described in the text used to determine theoretical values of \mathcal{P} shown in Fig. 5.

Author contributions. KNR: methodology, formal analysis and writing (original draft, review and editing). TJG: Conceptualization, funding acquisition, supervision, methodology, writing (review and editing). TDD: methodology and analysis, writing (review and editing). CB: writing (review and editing). SKK: funding acquisition, writing (review and editing). JCR: methodology, writing (review and editing).

430 *Competing interests.* The authors declare that they have no conflict of interest.



References

- Ackerman, S. A., Holz, R. E., Frey, R., Eloranta, E. W., Maddux, B. C., and McGill, M.: Cloud detection with MODIS. Part II: Validation, *J. Atmos. Oceanic Technol.*, 25, 1073–1086, 2008.
- Batista-Tomás, A., Díaz, O., Batista-Leyva, A., and Altshuler, E.: Classification and dynamics of tropical clouds by their fractal dimension, *Quarterly Journal of the Royal Meteorological Society*, 142, 983–988, 2016.
- 435 Benner, T. C. and Curry, J. A.: Characteristics of small tropical cumulus clouds and their impact on the environment, *Journal of Geophysical Research: Atmospheres*, 103, 28 753–28 767, 1998.
- Bolgiano Jr, R.: Turbulent spectra in a stably stratified atmosphere, *Journal of Geophysical Research*, 64, 2226–2229, 1959.
- Cahalan, R. F. and Joseph, J. H.: Fractal statistics of cloud fields, *Monthly Weather Review*, 117, 261–272, 1989.
- 440 Ceamanos, X., Six, B., and Riedi, J.: Quasi-Global Maps of Daily Aerosol Optical Depth From a Ring of Five Geostationary Meteorological Satellites Using AERUS-GEO, *Journal of Geophysical Research: Atmospheres*, 126, e2021JD034 906, 2021.
- Celani, A., Cencini, M., Mazzino, A., and Vergassola, M.: Active versus Passive Scalar Turbulence, *Phys. Rev. Lett.*, 89, 234 502, <https://doi.org/10.1103/PhysRevLett.89.234502>, 2002.
- Charney, J. G.: Geostrophic turbulence, *Journal of the Atmospheric Sciences*, 28, 1087–1095, 1971.
- 445 Christensen, H. M. and Driver, O. G.: The Fractal Nature of Clouds in Global Storm-Resolving Models, *Geophysical Research Letters*, 48, e2021GL095 746, 2021.
- Clauset, A., Shalizi, C. R., and Newman, M. E.: Power-law distributions in empirical data, *SIAM review*, 51, 661–703, 2009.
- Derrien, M. and Gléau, H. L.: MSG/SEVIRI cloud mask and type from SAFNWC, *International Journal of Remote Sensing*, 26, 4707–4732, <https://doi.org/10.1080/01431160500166128>, 2005.
- 450 DeWitt, T. D., Garrett, T. J., Rees, K. N., Bois, C., and Krueger, S. K.: Observations of climatologically invariant scale-invariance describing cloud horizontal sizes, *EGUsphere*, 2023, 1–20, <https://doi.org/10.5194/egusphere-2023-943>, 2023.
- Fiedler, F. and Panofsky, H. A.: Atmospheric scales and spectral gaps, *Bulletin of the American Meteorological Society*, 51, 1114–1120, 1970.
- Fielding, M. D., Schäfer, S. A. K., Hogan, R. J., and Forbes, R. M.: Parametrizing cloud geometry and its application in a subgrid cloud-edge erosion scheme, *Quarterly Journal of the Royal Meteorological Society*, 146, 1651–1667, <https://doi.org/https://doi.org/10.1002/qj.3758>, 2020.
- 455 Gage, K. S. and Nastrom, G. D.: Theoretical interpretation of atmospheric wavenumber spectra of wind and temperature observed by commercial aircraft during GASP, *Journal of Atmospheric Sciences*, 43, 729–740, 1986.
- Garrett, T. J., Glenn, I. B., and Krueger, S. K.: Thermodynamic constraints on the size distributions of tropical clouds, *Journal of Geophysical Research: Atmospheres*, 123, 8832–8849, 2018.
- 460 Gifford, F.: The shape of large tropospheric clouds, or “very like a whale”, *Bulletin of the American Meteorological Society*, 70, 468–475, 1989.
- Hentschel, H. and Procaccia, I.: Relative diffusion in turbulent media: the fractal dimension of clouds, *Physical Review A*, 29, 1461, 1984.
- Heus, T., Van Dijk, G., Jonker, H. J., and Van den Akker, H. E.: Mixing in shallow cumulus clouds studied by Lagrangian particle tracking, *Journal of the Atmospheric Sciences*, 65, 2581–2597, 2008.
- 465 Imre, A.: Problems of measuring the fractal dimension by the slit-island method, *Scripta metallurgica et materialia*, 27, 1713–1716, 1992.



- Kantha, L. and Hocking, W.: Dissipation rates of turbulence kinetic energy in the free atmosphere: MST radar and radiosondes, *Journal of Atmospheric and Solar-Terrestrial Physics*, 73, 1043–1051, 2011.
- 470 Khairoutdinov, M. F., Krueger, S. K., Moeng, C.-H., Bogenschutz, P. A., and Randall, D. A.: Large-eddy simulation of maritime deep tropical convection, *Journal of Advances in Modeling Earth Systems*, 1, 2009.
- Kolmogorov, A. N.: The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers, *Cr Acad. Sci. URSS*, 30, 301–305, 1941.
- Kopp, T. J., Thomas, W., Heidinger, A. K., Botambekov, D., Frey, R. A., Hutchison, K. D., Iisager, B. D., Brueske, K., and Reed, B.: The VIIRS Cloud Mask: Progress in the first year of S-NPP toward a common cloud detection scheme, *Journal of Geophysical Research: Atmospheres*, 119, 2441–2456, <https://doi.org/https://doi.org/10.1002/2013JD020458>, 2014.
- 475 Korčák, J.: Geopolitické základy Československa: jeho kmenové oblasti, 1938.
- Kraichnan, R. H.: Inertial ranges in two-dimensional turbulence, *The Physics of Fluids*, 10, 1417–1423, 1967.
- Lovejoy, S.: Area-perimeter relation for rain and cloud areas, *Science*, 216, 185–187, 1982.
- Lovejoy, S.: CloudSat Cloud Length, Thickness Distributions Again Confirm the 23/9 (2.55 D) Scaling, Stratified, Turbulence Model, in: 2021 IEEE International Geoscience and Remote Sensing Symposium IGARSS, pp. 1727–1730, IEEE, 2021.
- 480 Lovejoy, S.: Scaling, dynamical regimes and stratification: How long does weather last? How big is a cloud?, *Nonlinear Processes in Geophysics Discussions*, 2023, 1–146, 2023.
- Lovejoy, S. and Schertzer, D.: Generalized scale invariance in the atmosphere and fractal models of rain, *Water Resources Research*, 21, 1233–1250, 1985.
- 485 Lovejoy, S. and Schertzer, D.: Multifractals, universality classes and satellite and radar measurements of cloud and rain fields, *Journal of Geophysical Research: Atmospheres*, 95, 2021–2034, 1990.
- Lovejoy, S. and Schertzer, D.: Multifractals, cloud radiances and rain, *Journal of Hydrology*, 322, 59–88, 2006.
- Lovejoy, S. and Schertzer, D.: Haar wavelets, fluctuations and structure functions: convenient choices for geophysics, *Nonlinear Processes in Geophysics*, 19, 513–527, 2012.
- 490 Lovejoy, S. and Schertzer, D.: *The Weather and Climate: Emergent Laws and Multifractal Cascades*, Cambridge University Press, 2018.
- Lovejoy, S., Schertzer, D., Silas, P., Tessier, Y., and Lavallée, D.: The unified scaling model of atmospheric dynamics and systematic analysis of scale invariance in cloud radiances, in: *Annales Geophysicae*, vol. 11, pp. 119–127, 1993.
- Lovejoy, S., Schertzer, D., and Stanway, J.: Direct evidence of multifractal atmospheric cascades from planetary scales down to 1 km, *Physical Review Letters*, 86, 5200, 2001.
- 495 Lovejoy, S., Tuck, A., Hovde, S., and Schertzer, D.: Is isotropic turbulence relevant in the atmosphere?, *Geophysical Research Letters*, 34, 2007.
- Mandelbrot, B.: How long is the coast of Britain? Statistical self-similarity and fractional dimension, *Science*, 156, 636–638, 1967.
- Mandelbrot, B.: *Fractals*, Freeman San Francisco, 1977.
- Mandelbrot, B. B.: Self-affine fractals and fractal dimension, *Physica scripta*, 32, 257, 1985.
- 500 Mapes, B. E.: Water’s two height scales: The moist adiabat and the radiative troposphere, *Quarterly Journal of the Royal Meteorological Society*, 127, 2353–2366, 2001.
- Marshak, A., Davis, A., Wiscombe, W., and Cahalan, R.: Radiative smoothing in fractal clouds, *Journal of Geophysical Research: Atmospheres*, 100, 26 247–26 261, 1995.



- Mieslinger, T., Horváth, , Buehler, S. A., and Sakradzija, M.: The Dependence of Shallow Cumulus Macrophysical Properties on Large-Scale Meteorology as Observed in ASTER Imagery, *Journal of Geophysical Research: Atmospheres*, 124, 11 477–11 505, <https://doi.org/https://doi.org/10.1029/2019JD030768>, 2019.
- Nastrom, G., Gage, K., and Jasperson, W.: Kinetic energy spectrum of large-and mesoscale atmospheric processes, *Nature*, 310, 36–38, 1984.
- Obukhov, A.: Description of turbulence in terms of Lagrangian variables, *Advances in Geophysics*, 6, 113–116, 1959.
- Peters, O., Neelin, J. D., and Nesbitt, S. W.: Mesoscale convective systems and critical clusters, *Journal of the atmospheric sciences*, 66, 2913–2924, 2009.
- Pressel, K. G. and Collins, W. D.: First-order structure function analysis of statistical scale invariance in the AIRS-observed water vapor field, *Journal of climate*, 25, 5538–5555, 2012.
- Pressel, K. G., Collins, W. D., and Desai, A. R.: The spatial scale dependence of water vapor variability inferred from observations from a very tall tower, *Journal of Geophysical Research: Atmospheres*, 119, 9822–9837, 2014.
- Read, P. L., Barstow, J., Charnay, B., Chelvaniththilan, S., Irwin, P. G. J., Knight, S., Lebonnois, S., Lewis, S. R., Mendonça, J., and Montabone, L.: Global energy budgets and ‘Trenberth diagrams’ for the climates of terrestrial and gas giant planets, *Quarterly Journal of the Royal Meteorological Society*, 142, 703–720, 2016.
- Richardson, L. F.: Atmospheric diffusion shown on a distance-neighbour graph, *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, 110, 709–737, 1926.
- Richardson, L. F.: The problem of contiguity: An appendix to statistics of deadly quarrels, *General systems yearbook*, 6, 139–187, 1961.
- Sagan, C.: *Pale Blue Dot: A Vision of the Human Future in Space*, New York: Random House, Inc, 1994.
- Schertzer, D. and Lovejoy, S.: Generalised scale invariance in turbulent phenomena, *PhysicoChemical Hydrodynamics*, 6, 623–635, 1985.
- Schwartz, F. and Brow, J. E.: Diffusivity of water vapor in some common gases, *The Journal of Chemical Physics*, 19, 640–646, 1951.
- Siebesma, A. and Jonker, H.: Anomalous scaling of cumulus cloud boundaries, *Physical Review Letters*, 85, 214, 2000.
- Tennekes, H. and Lumley, J. L.: *A first course in turbulence*, MIT press, 1972.
- Wood, R. and Field, P. R.: The distribution of cloud horizontal sizes, *Journal of Climate*, 24, 4800–4816, 2011.
- Yang, Y., Meyer, K., Wind, G., Zhou, Y., Marshak, A., Platnick, S., Min, Q., Davis, A. B., Joiner, J., Vasilkov, A., Duda, D., and Su, W.: Cloud products from the Earth Polychromatic Imaging Camera (EPIC): algorithms and initial evaluation, *Atmospheric Measurement Techniques*, 12, 2019–2031, <https://doi.org/10.5194/amt-12-2019-2019>, 2019.
- Zelinka, M. D., Klein, S. A., Qin, Y., and Myers, T. A.: Evaluating Climate Models’ Cloud Feedbacks Against Expert Judgment, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD035 198, <https://doi.org/https://doi.org/10.1029/2021JD035198>, 2022.
- Zhao, G. and Di Girolamo, L.: Statistics on the macrophysical properties of trade wind cumuli over the tropical western Atlantic, *Journal of Geophysical Research: Atmospheres*, 112, 2007.
- Zhao, M. and Austin, P. H.: Life cycle of numerically simulated shallow cumulus clouds. Part II: Mixing dynamics, *Journal of the atmospheric sciences*, 62, 1291–1310, 2005.