

Source specific bias correction of US background and anthropogenic ozone modeled in CMAQ

T. Nash Skipper^{1,*}, Christian Hogrefe², Barron H. Henderson², Rohit Mathur², Kristen M. Foley², Armistead G. Russell¹

5 ¹School of Civil & Environmental Engineering, Georgia Institute of Technology, Atlanta, GA 30332, USA

²U.S. Environmental Protection Agency, Research Triangle Park, NC, 27709, USA

[*Now at U.S. Environmental Protection Agency, Research Triangle Park, NC, 27709, USA](#)

Correspondence to: Armistead G. Russell (ar70@gatech.edu)

Abstract. United States (US) background ozone (O₃) is the counterfactual O₃ that would exist with zero
10 US anthropogenic emissions. Estimates of US background O₃ typically come from chemical transport
models (CTMs), but different models vary in their estimates of both background and total O₃. Here, a
measurement-model data fusion approach is used to estimate CTM biases in US anthropogenic O₃ and
multiple US background O₃ sources, including natural emissions, long-range international emissions,
short-range international emissions from Canada and Mexico, and stratospheric O₃. Spatially and
15 temporally varying bias correction factors adjust each simulated O₃ component so that the sum of the
adjusted components evaluates better against observations compared to unadjusted estimates. The
estimated correction factors suggest a seasonally consistent positive bias in US anthropogenic O₃ in the
eastern US, with the bias becoming higher with coarser model resolution and with higher simulated total
O₃ though the bias does not increase much with higher observed O₃. Summer average US anthropogenic
20 O₃ in the eastern US was estimated to be biased high by 2, 7, and 11 ppb (11%, 32%, and 49%) for one
set of simulations at 12, 36, and 108 km resolutions and 1 and 6 ppb (10% and 37%) for another set of
simulations at 12 and 108 km resolutions. Correlation among different US background O₃ components
can increase the uncertainty in the estimation of the source-specific adjustment factors. Despite this,
results indicate a negative bias in modeled estimates of the impact of stratospheric O₃ at the surface, with
25 a western US spring average bias of -3.5 ppb (-25%) estimated based on a stratospheric O₃ tracer. This
type of data fusion approach can be extended to include data from multiple models to leverage the
strengths of different data sources while reducing uncertainty in the US background ozone estimates.

1 Introduction

United States (US) background-~~(USB)~~ ozone (O₃) is the counterfactual O₃ that would exist if US anthropogenic-~~(USA)~~ emissions were zero. The National Ambient Air Quality Standard (NAAQS) for O₃ was set at a level of 70 ppb in 2015 and may be lowered. In its recent reviews of the O₃ NAAQS, the US Environmental Protection Agency (EPA) noted the importance of USBUS background O₃ (US EPA, 2013, 2014, 2020b, a). USBUS background O₃ takes up a larger portion of the allowed ozone as the NAAQS is tightened and is a larger portion of total observed O₃ as anthropogenic precursor emissions decline (Lin et al., 2017; Guo et al., 2018; Jaffe et al., 2018). USBUS background O₃ cannot be observed (Fiore et al., 2003; Dentener et al., 2010; McDonald-Buller et al., 2011; Fiore et al., 2014; Jaffe et al., 2018; US EPA, 2013, 2014, 2020b, a). It is typically quantified using a chemical transport model (CTM), most commonly using the zero-out method in which USAUS anthropogenic emissions are set to zero. There is much uncertainty in CTM estimates of USBUS background O₃ due to model biases and differences in CTM-estimated USBUS background O₃ among different models (McDonald-Buller et al., 2011; Fiore et al., 2014; Dolwick et al., 2015; Huang et al., 2015; Guo et al., 2018; Jaffe et al., 2018). Jaffe et al. (2018) estimated that the typical uncertainty in CTM-estimated seasonal mean USBUS background O₃ is ±10 ppb.

Sources of USBUS background O₃ include naturally occurring emissions such as wildfires, biogenic VOCs, oxides of nitrogen (NO_x) from soil, lightning NO_x, stratosphere-to-troposphere exchange, and oxidation of methane (Fiore et al., 2014; Jaffe et al., 2018; US EPA, 2020a). Some portions of total O₃ contributions from soil NO_x and methane oxidation are USBUS background sources while some are anthropogenic. Soil NO_x is emitted by microbial processes in both natural and agricultural lands and is limited by availability of nitrogen in the soil. There is a pre-industrial level of methane that contributes to USBUS background O₃ formation, but any O₃ created through oxidation of methane above the pre-industrial level is anthropogenic. Soil NO_x and methane oxidation are often treated as USBUS background O₃ sources in their entirety in CTM studies due to the complexity of splitting up the natural and anthropogenic portions (US EPA, 2020a). Wildfires are treated as USBUS background O₃ sources, but the impacts of wildfires on O₃ can be affected by US anthropogenic emissions when VOCs from fires are transported over NO_x-rich urban areas, leading to enhanced O₃ production (Jaffe et al., 2013; Langford et

al., 2023; Rickly et al., 2023). USBUS background O₃ sources also include non-US anthropogenic pollution which may be from long range transport (Lin et al., 2012b) or from short range transport from neighboring countries (Wang et al., 2009).

60 In previous work (Skipper et al., 2021), we developed a bias correction method which used regression modeling to adjust CTM-simulated US Aanthropogenic and USBUS background O₃ to better align with observations and to improve agreement of differing USBUS background O₃ estimates from different model configurations. We developed spatially and temporally varying scaling factors to adjust US Aanthropogenic and USBUS background O₃. In that work, USBUS background O₃ was treated as a single quantity rather than considering different sources of USBUS background O₃ individually. A
65 consistent low bias in USBUS background O₃ in spring was identified, though the specific source of this low bias could not be identified. Here, we extend the bias correction method to estimate biases in separate components of USBUS background O₃. Separating the USBUS background O₃ components provides new insights into the inferred CTM error in USBUS background O₃ that was not possible when USBUS background O₃ was treated as a lumped quantity.

70 **2 Methods**

2.1 Chemical transport model simulations

Total O₃ (i.e., BASE-base O₃), USBUS background O₃, and individual USBUS background O₃ components are simulated at both regional and hemispheric scales using the Community Multiscale Air Quality (CMAQ) model. We use maximum daily 8-h average (MDA8) O₃ as the metric of interest since
75 this is the metric used in determining attainment of the NAAQS. References to O₃ throughout are to MDA8 O₃. CMAQ results are from two recent sets of simulations by the US EPA (Table 1). The two sets of simulations include different USBUS background O₃ components allowing us to explore how different components of USBUS background O₃ affect the bias in O₃.

80 **Table 1. Simulation names and descriptions for hemispheric-scale and regional-scale simulations. Table adapted from 2020 O₃ Policy Assessment Table 2-1 (US EPA, 2020a).**

Simulation	Description
BASE	All emission sectors are included.
ZUSA	All US anthropogenic emissions are removed including prescribed fires. ^a
ZROW	All anthropogenic emissions outside the US are removed including prescribed fires where possible (ROW = rest of world). ^b
ZCANMEX	All anthropogenic emissions from Canada and Mexico are removed including prescribed fires where possible. ^b
ZANTH	All anthropogenic emissions globally are removed including prescribed fires. ^b
STRAT	Tracer species for O ₃ injected into the upper troposphere/lower stratosphere based on CMAQ potential vorticity parameterization for stratospheric O ₃ . ^c

^a Emissions estimated to be associated with intentionally set fires (“prescribed fires”) are grouped with anthropogenic fires.

^b Only for PA simulations

^c Only for EQUATES simulations.

85

The first set of simulations was conducted for the Policy Assessment (PA) for the review of the O₃ NAAQS in 2020 (US EPA, 2020a). These simulations also support the draft PA for the reconsideration of the O₃ NAAQS. The PA simulations cover the entire year of 2016 and provide estimates of US ~~A~~anthropogenic and ~~US~~background O₃ as well as natural (~~NAT~~) and international anthropogenic (~~INTL~~) contributions to background O₃. ~~INTL~~International O₃ is also further decomposed to short-range international anthropogenic contributions from Canada and Mexico (~~CANMEX~~)(Canada+Mexico) and long-range international (~~LINTL~~) contributions from other countries. The PA simulations consist of nested simulations from hemispheric scale (Mathur et al., 2017) at 108 km horizontal resolution to continental scale at 36 km resolution to a finer continental scale at 12 km resolution.

95

background O₃ components are determined by the zero-out method in which the model is run in the same configuration as the base case but with specified emissions sources removed. The zero-out method is the most common approach for simulating background O₃, though other approaches such as sensitivity simulations and source tagging techniques have also been previously employed (Jaffe et al., 2018). The zero-out method neglects non-linear interactions between sources which can affect the simulated source contribution (Wu et al., 2009; Dolwick et al., 2015). However, the

100

zero-out method is consistent with the definition of USBUS background O₃ as the level of O₃ in the absence of US anthropogenic emissions, while sensitivity or tagging techniques would instead provide an estimate of source contributions to total simulated O₃ (including O₃ from US anthropogenic sources).

105 USBUS background O₃ is estimated by removing US anthropogenic emissions (ZUSA simulation). USAUS anthropogenic O₃ is calculated as BASE-base O₃ minus USBUS background O₃. NATNatural O₃ is estimated by removing all anthropogenic emissions (ZANTH simulation). The non-US anthropogenic O₃ contribution is estimated by removing anthropogenic emissions everywhere except the US (ZROW simulation). The INTL-international contribution is calculated as BASE-base O₃ minus O₃

110 from the ZROW simulation. CANMEXCanada+Mexico O₃ is estimated by removing Canada and Mexico anthropogenic emissions (ZCANMEX simulation). The CANMEXCanada+Mexico O₃ contribution is calculated as BASE-base O₃ minus the O₃ from the ZCANMEX simulation. LINTLLong-range international O₃ is estimated as INTL-international O₃ minus CANMEXCanada+Mexico O₃. Due to non-linear chemistry, there is some residual anthropogenic contribution to BASE-base O₃ which is not

115 attributed to US or international emissions. Descriptions of these CMAQ simulations and calculation of O₃ components are given in Tables S1 and S2. Further details of the modeling setup are available in the 2020 Policy Assessment (US EPA, 2020a).-

The second set of simulations was developed from EPA's Air QUALity TimE Series (EQUATES) project which spans 2002-2019. Additional simulations using the EQUATES modeling framework were

120 conducted for 2016–2017 to estimate USBUS background O₃ and USAUS anthropogenic O₃ using the zero-out method. The EQUATES simulations consist of hemispheric scale simulations at 108 km horizontal resolution and nested US continental scale simulations at 12 km horizontal resolution. Descriptions of these CMAQ simulations and calculation of O₃ components are given in Table S3. Further details on the model configuration for EQUATES are available from Foley et al. (2020) and Foley et al.

125 (2023). More details on both the PA and EQUATES simulations are summarized in Tables S4 and S5.

The 108 km EQUATES simulations also include an inert tracer species which serves as a proxy for simulated stratospheric O₃ contributions. Separate stratospheric O₃ contributions were not available from the PA simulations, so the EQUATES simulations provide an opportunity to assess potential biases specific to stratospheric O₃ contributions. CMAQ simulates stratospheric O₃ using a parameterization

130 based on the relationship between O₃ and potential vorticity (PV) in the upper troposphere and lower
stratosphere (UTLS) (Xing et al., 2016). The parameterization was developed using 21 years of
ozonesonde data from the World Ozone and Ultraviolet Radiation Data Centre and PV data from the
Weather Research Forecasting (WRF) model for 1990-2010. In the EQUATES 108 km simulations, the
parameterization is applied to the top model layer only. A PV tracer species tracks O₃ injected into the
135 UTLS throughout the rest of the model domain for the hemispheric simulations. The 12 km continental
simulations inherit the PV tracer species through lateral boundary conditions from the hemispheric
simulations. This tracer is subject to transport and deposition but not chemistry. We refer to the PV tracer
concentrations as ~~STRAT~~ (short for stratospheric) O₃ since it relates to the stratospheric influence, but it
only partly replicates the impact of stratospheric O₃ since it does not undergo chemical losses. The
140 sSTRATstratospheric O₃ tracer does, however, provide a measure of the spatiotemporal variability of
stratospheric O₃ impacts. We also estimate the contribution to USBUS background O₃ from sources other
than the stratosphere as USBUS background O₃ minus the STRATstratospheric O₃ tracer and refer to it
as non-stratospheric USBUS background-NOSTRAT O₃. The use of the chemically inert PV tracer to
split up stratospheric and non-stratospheric influences on USBUS background O₃ introduces uncertainty
145 as the STRATstratospheric O₃ component may be unrealistically high, especially in areas and times with
more active chemistry.

The modeling configurations of the PA and EQUATES simulations differ in some respects which
is expected to lead to some differences in simulated O₃, though they do share some of the same
configuration options. Both the PA and EQUATES simulations use a 44-layer vertical structure for
150 hemispheric scale applications (at 108 km resolution) and a 35-layer vertical structure for continental (i.e.,
36 km and 12 km resolution) applications with a vertical extent from the surface to 50 hPa and a surface
layer height of approximately 20 m for both the hemispheric and continental configurations (see Mathur
et al. (2017) for more details on these vertical layer structures). CMAQ v5.2.1 was used for the PA
simulations while CMAQ v5.3.2 was used for the EQUATES simulations. These were the latest versions
155 of CMAQ at the respective times that each set of simulations were conducted. One potential source of
differences is ~~that updates to~~ halogen chemistry ~~was updated~~ that were introduced in CMAQ v5.3 (Sarwar
et al., 2019). ~~The EQUATES hemispheric simulations therefore include losses of O₃ over seawater that~~

~~are not present in the PA hemispheric simulations which could affect O₃ transported over the Pacific in particular. These updates in the EQUATES simulations enhance halogen-mediated O₃ losses, which are strongest over the oceans. These losses are most relevant for O₃ contributions (natural and anthropogenic) that are transported long distances over oceans.~~ An intercomparison of CMAQ v5.2.1 and CMAQ v5.3.1 (which is not significantly different from CMAQ v5.3.2) showed that the newer version typically had lower O₃ compared to the older version, with mean bias ~1 ppb lower in CMAQ v5.3.1 (Appel et al., 2021). Besides the ~~addition of updates to~~ halogen chemistry, there are other differences in the chemical mechanisms used for each set of simulations. The mechanisms used for the hemispheric simulations were cb6r3_ae6_aq for the PA simulations and cb6r3m_ae7_kmtbr for the EQUATES simulations. The part of the mechanism name labeled cb6r3m indicates additional chemistry relevant in marine environments (the halogen chemistry described above); ae6 and ae7 indicate the version number for chemistry relevant to aerosols; aq and kmtbr indicate different treatments of cloud chemistry. The chemical mechanisms used for continental-scale PA and EQUATES simulations (cb6r3_ae6nvPOA_aq and cb6r3_ae7_aq) also differ in their representation of organic aerosols (Murphy et al., 2017; Pye et al., 2019; Qin et al., 2021; Appel et al., 2021) which could affect O₃ concentrations. Different versions of WRF (v3.8 for PA simulations and v4.1.1 for EQUATES simulations) employed may also contribute to differences in O₃.

Emission inputs also differ between the PA and EQUATES simulations. Different US anthropogenic emission inventories were used for the simulations. The PA simulations used an early version (sometimes called the “alpha” version) of a 2016 emissions modeling platform developed by the National Emissions Inventory Collaborative (US EPA, 2019b). The EQUATES simulations used an inventory that was developed as part of the broader EQUATES framework to model a long timeseries using consistent methods for emissions estimates (Foley et al., 2023). For emissions in Canada and Mexico, both sets of simulations use emission inventories developed by the respective national governments, though the EQUATES simulations use more recent inventories (as described by Foley et al. (2020)) than the PA simulations (as described by US EPA (2019b)). Both the PA and EQUATES simulations use the Tsinghua University inventory of emissions in China (Zhao et al., 2018). For other countries, both sets of simulations use the Hemispheric Transport of Air Pollution (HTAP) v2.2 inventory (Janssens-Maenhout et al., 2015) with scaling factors derived from the Community Emissions Data

System (CEDS) (Hoesly et al., 2018) to account for yearly changes. Differences in the anthropogenic emissions used in the two model configurations are expected to contribute to differences in simulated O₃, most notably for the different US anthropogenic emissions since we focus here on O₃ in the US.

For hemispheric-scale simulations, biogenic VOC emissions are from the Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1) (Guenther et al., 2012). The PA simulations additionally replace MEGAN emissions with emissions from the Biogenic Emission Inventory System (BEIS) (Bash et al., 2016) over North America (US EPA, 2019a). The EQUATES MEGAN emissions are obtained from a compilation by Sindelarova et al. (2014). Soil NO_x emissions for the PA hemispheric simulations are also from MEGAN with replacement by BEIS soil NO_x over North America. Soil NO_x emissions for the hemispheric EQUATES simulations are from a dataset by the Copernicus Atmosphere Monitoring Service (CAMS, 2018) based on methods by Yienger and Levy (1995). Both the PA and EQUATES simulations use BEIS for biogenic VOC and soil NO_x emissions in the continental-scale simulations. Lightning NO_x emissions for both the PA and EQUATES hemispheric simulations are from monthly climatology obtained from the Global Emissions Initiative (GEIA) and are based on Price et al. (1997). Lightning NO_x was not included in the PA continental-scale simulations, while lightning NO_x for the EQUATES continental-scale simulations is calculated using an inline module in CMAQ (Kang et al., 2019). For both PA and EQUATES, wildfire emissions outside of North America are based on the Fire Inventory from NCAR (FINN) v1.5 (Wiedinmyer et al., 2011) which provides day-specific fire emissions. Wildfires are vertically allocated with 25% of emissions distributed to the lowest two layers (~0-45 m), 35% distributed to layers 3-9 (~45-350 m), and the remaining 40% distributed to layers 10-19 (~350-2000 m) as described in the Technical Support Document for northern hemispheric emissions (US EPA, 2019a). Wildfire emissions within North America are based on the Hazard Mapping System (HMS) fire product which provides day-specific fire activity data. Emission processing for North American wildfires is further described in the Technical Support Document for North American emissions (US EPA, 2019b) (applicable to PA simulations) and Foley et al. (2023) (applicable to EQUATES simulations). Although the methods are similar, North American wildfire emissions may differ between PA and EQUATES based on the specific fire activity data that was used in each case. Fire plume injection height for North American fires is determined by an inline plume rise algorithm in CMAQ based on fire heat content (see e.g.,

215 Wilkins et al. (2022) for more details on fire plume injection height in CMAQ). Stratospheric O₃ in both the PA and EQUATES simulations is from the PV parameterization by Xing et al. (2016) (described in more detail above) in the hemispheric simulations. Stratospheric O₃ in the continental-scale simulations only comes from any stratospheric O₃ inherited from the lateral boundary conditions provided by the hemispheric simulations.

2.2 O₃ observations

220 O₃ observational data are from the Air Quality System (AQS) database, which provides data from federal, state, local, and tribal air quality monitoring networks across the US. The average precision of O₃ monitors in the AQS database was reported as 2.2% and 2.4% in 2016 and 2017, respectively, and the national average absolute bias was reported as 1.5% in both 2016 and 2017 (<https://www.epa.gov/amtic/amtic-ambient-air-monitoring-assessments>). There were ~360,000 MDA8
225 O₃ observations available per year for 2016 and 2017 from ~1250 unique monitoring sites. These numbers take into account monitoring sites where O₃ is measured by multiple instruments at the same location (as indicated in the AQS database by a parameter occurrence code). In these cases, the MDA8 O₃ observations from multiple instruments are averaged for a given site and day and treated as a single observation. The observations overrepresent the eastern US compared to the western US. About 40% of
230 daily MDA8 O₃ observations and ~36% of O₃ monitoring sites are in the western US (as defined by longitude < -97 °W). Western US sites are also overrepresented by sites in the state of California. About 40% of daily MDA8 O₃ observations and ~40% of O₃ monitoring sites in the western US are in California. The observations also overrepresent the high O₃ season of April – October (Figure 1) since many monitors are only required to be operated during the high O₃ season.

235




Figure 1. Locations of O₃ observational sites in 2016 indicated with a circle whose color shows the number of daily MDA8 O₃ observations available from each site in 2016 (top). Total number of daily MDA8 O₃ observations in each month of 2016 (bottom).

240

2.3 O₃ data fusion model

We use multivariate ordinary least squares regression to model the relationship between the individual model components and observed MDA8 O₃. Regression parameters provide estimates of the spatial and temporal model bias attributable to each individual O₃ component. The regression model for

245 ozone mixing ratio O₃ on day d and location (lon, lat, z) is formulated as follows:

$$O_3 = \sum_i \alpha_i O_{3i}^{simulated} + \varepsilon$$

Where:

$$\alpha_i = \alpha_{0,i} + \alpha_{x,i}lon + \alpha_{y,i}lat + \alpha_{z,i}z + \alpha_{sin,i} \sin(d) + \alpha_{cos,i} \cos(d)$$

d is day of year in radians

250 z is elevation above sea level

$lon, lat, z, \sin(d),$ and $\cos(d)$ are normalized to zero mean and unit standard deviation (Table S6)

$$\varepsilon \sim N(0, \sigma^2)$$

index i represents different sets of O₃ components. Specifically, we consider four sets of i :

$i \in \{US\ Anthropogenic, US\ Background\}$ (PA and EQUATES)

255 $i \in \{US\ Anthropogenic, NAT\ Natural, INTL\ International\}$ ———(PA-only)

$i \in$

$\{US\ Anthropogenic, NAT\ Natural, LINTL\ Long\ range\ international, CANMEX\ Canada + Mexico\}$ (PA-only)

$i \in$

260 $\{US\ Anthropogenic, USB_{NOSTRATNON}\ stratospheric\ US\ Background, STRAT\ Stratospheric\}$
 ———(EQUATES-only)

Each simulated O₃ component ($O_{3i}^{simulated}$) is multiplied by the alpha adjustment factor for that component (α_i), which varies as a function of space and time, to calculate an adjusted estimate of each O₃ component. The inferred model bias for a particular component is calculated as the difference between the original simulated O₃ and adjusted O₃ for that component. The individual adjusted O₃ components are summed to calculate the total adjusted O₃. The longitude and latitude terms of α_i are intended to capture the spatial variability of O₃ biases while the z term of α_i is intended to capture biases in O₃ related to elevation. The sinusoidal day of year terms of α_i are intended to capture the cyclical nature of O₃ production and to identify any seasonal dependence in O₃ biases. The modeled O₃ components do not add up to observed O₃ because of biases in the model or its inputs. The CMAQ-simulated O₃ components are adjusted by applying estimated regression coefficients to the gridded data so that the sum of the components more closely aligns with observed O₃. A more complex method (e.g., nonlinear regression or machine learning) may give a better fit to observed O₃, but the interest here is to estimate potential biases in the modeled O₃ components which is more straightforward with a linear regression. Empirical

orthogonal function (EOF) analysis was used to further explore the spatial and temporal structure of the inferred bias fields and is discussed in the SI.

A separate regression model is developed for each separate model configuration (i.e., model resolution, PA or EQUATES simulation, and USBUS background O₃ component split). There are three model resolutions and three USBUS background O₃ splits for the PA simulations, resulting in nine PA models. There are two model resolutions for the EQUATES simulations. The 12 km EQUATES data has two USBUS background O₃ splits while the 108 km EQUATES data has one USBUS background O₃ split, resulting in three EQUATES models. For the PA models, only 2016 PA simulation data are used to train the models since these simulations are for only that year. For the EQUATES models, both 2016 and 2017 EQUATES simulation data are used to train the models. The location and sampling schedule of the monitoring sites overrepresent the eastern US, low elevations, and high O₃ season which may impact how representative the results are for non-monitored locations. Overfitting of the regression model is tested using three cross-validation approaches in which the data are split in both space and time, in space only, and in time only. In the first approach (spatial and temporal withholding), 10% of all observational data are randomly selected and reserved as a test set while the remaining 90% are used as the training set. In the second approach (spatial withholding), data from 10% of randomly selected observation sites are used as a test set while data from the remaining 90% of sites is used as the training set. In the third approach (temporal withholding), data from 10% of randomly selected days of the year are used as a test set while data from the remaining 90% of days of the year are used as the training set. The root mean square error (RMSE) and mean bias for the test and training set are compared to evaluate the potential for the model to overfit the data.

3 Results and discussion

3.1 CTM results

The overall performance of MDA8 O₃ for each simulation is summarized here by the normalized mean bias (NMB) compared to O₃ monitoring sites. The 12 km PA simulations were biased high for 2016 (NMB=1.2%) while the 12 km EQUATES simulations were biased low for 2016 and 2017 (NMB=-3.7%


and -5.1%). The 36 km and 108 km PA simulations were biased high over the US for 2016 (NMB=5.2% and 10.0%). The 108 km EQUATES simulations were also biased high over the US for 2016 and 2017 (NMB=2.8% and 0.5%). The two sets of simulations are broadly consistent with one another for BASEbase, USAUS anthropogenic, and total USBUS background O₃ which are common to both. Details on the contributions from the different O₃ components in the PA and EQUATES simulations follow.

CMAQ-simulated annual average MDA8 O₃ from the PA simulations show similar results across the three different model resolutions for USBUS background O₃ sources (Figure 2; Table 2). Simulated USAUS anthropogenic O₃ tends to increase with coarser model resolution which results in corresponding increases in BASEbase O₃. NATNatural O₃ makes the largest contribution to annual average O₃ across the US with a larger contribution in the western US (~55% of BASEbase) than in the eastern US (~45% of BASEbase). USAUS anthropogenic O₃ is the second largest component of annual average O₃ with a larger contribution in the eastern US (~35% of BASEbase) than in the western US (~20% of BASEbase). There are a small number of US grid cells with negative annual averages for USAUS anthropogenic O₃. This means that USBUS background O₃ was greater than BASEbase O₃ and indicates that anthropogenic emissions suppress O₃ through NO_x titration. LINTLlong-range international sources impacts the western US (~15% of BASEbase) more strongly than the eastern US (~10% of BASEbase). Both NATnatural and LINTLlong-range international O₃ tend to be higher at higher elevations, suggesting that some of the effects from NATnatural and LINTLlong-range international are from O₃ in the free troposphere. In spring, O₃ lifetimes are longer, and trans-Pacific transport of O₃ is more likely which is consistent with the spring peak in LINTLlong-range international O₃ (Liu et al., 1987). The other components and BASEbase O₃ peak in the summer with some exceptions (Figure 3). In the southeastern US, NATnatural O₃ is lower during summer compared to surrounding areas and is lower than NATnatural O₃ in the southeastern US during spring. This is likely because O₃ loss through reaction with biogenic VOCs (which peak in the summer and are abundant in the southeastern US) reduces O₃ under the extremely low NO_x conditions with zero anthropogenic emissions. The CANMEXCanada+Mexico contribution to O₃ is small except at some locations along the border with Mexico where the contributions can be high, especially in the summer. For US grid cells within 100 km of the border with Canada, the

annual average impact is ~2 ppb while for US grid cells within 100 km of the border with Mexico, the
 330 annual average impact is ~5 ppb.

335 **Table 2. Summary of annual average of MDA8 O₃ components for the Policy Assessment set of simulations. Averages are shown for all of the US and separately for the eastern and western US with a longitude of 97 °W serving as the east-west dividing line. The mean across all grid cells within the given area is shown along with the minimum and maximum for any grid cell within the given area in parentheses. Numbers in the table are in units of ppb. Seasonal averages are provided in Table S13.**

	<u>BASE</u> Base	<u>USA</u> US anthropogenic	<u>NAT</u> Natural	<u>LINTL</u> Long- range international	<u>CANMEX</u> Canad a+Mexico
PA 12 km					
all US	39 (18, 56)	10 (-12, 23)	20 (15, 30)	6 (4, 10)	2 (-4, 9)
eastern US	39 (28, 49)	13 (2, 23)	18 (15, 21)	4 (4, 9)	1 (1, 6)
western US	40 (18, 56)	7 (-12, 23)	22 (15, 30)	7 (4, 10)	2 (-4, 9)
PA 36 km					
all US	40 (28, 62)	11 (2, 30)	20 (15, 28)	6 (4, 10)	2 (1, 16)
eastern US	40 (28, 55)	14 (4, 28)	18 (15, 21)	4 (4, 9)	1 (1, 5)
western US	40 (30, 62)	8 (2, 30)	22 (15, 28)	7 (4, 10)	2 (1, 16)
PA 108 km					
all US	42 (30, 70)	11 (3, 42)	21 (16, 28)	5 (3, 10)	2 (1, 9)
eastern US	42 (30, 70)	15 (4, 42)	19 (16, 23)	4 (3, 6)	1 (1, 4)
western US	42 (31, 54)	8 (3, 20)	23 (16, 28)	6 (3, 10)	2 (1, 9)



340






Figure 2. Annual average MDA8 O₃ from Policy Assessment CMAQ simulations. Results are shown for 12 km (top row), 36 km (middle row), and 108 km (bottom row) horizontal resolutions. O₃ concentrations include total (**BASE**base) O₃ as well as O₃ components from **USA**US anthropogenic, **NAT**natural, **LINT**long-range international, and **CANMEX**Canada+Mexico sources.

345





350 **Figure 3. Seasonal average MDA8 O₃ from Policy Assessment CMAQ simulations. Results are**
shown for 12 km horizontal resolution for winter (DJF), spring (MAM), summer (JJA), and fall
(SON). Seasonal averages for the 36 km and 108 km simulations are provided in Figures S1 and S2.
O₃ concentrations include total (BASEbase) O₃ as well as O₃ components from USAUS
anthropogenic, NATnatural, LINTLlong-range international, and CANMEXCanada+Mexico
355 **sources.**

While the annual (Figure 2, Table 2) and seasonal (Figure 3) average MDA8 O₃ contributions are
instructive about longer term contributions, compliance with the NAAQS is determined based on the
fourth highest observed MDA8 O₃, averaged over three years. We examine the fourth highest total (base)
360 MDA8 O₃ along with the contribution from each of MDA8 O₃ components on the same day (Figure 4).

The areas with the greatest fourth highest MDA8 O₃ in the base simulation mostly have large contributions from the US anthropogenic O₃ component. This includes much of California and major metropolitan areas in the rest of the US. The eastern US has a higher level of US anthropogenic O₃ outside of the metropolitan areas compared to most of the western US where US anthropogenic O₃ outside of urban areas is typically in the range of 5-20 ppb. Although the western US and eastern US have similar fourth highest MDA8 O₃ values for base O₃ (western US 60 ppb, eastern US 61 ppb for 12 km simulations), the western US has a lower average contribution from the US anthropogenic component (14 ppb) compared to the eastern US (33 ppb).

The contribution to the fourth highest MDA8 O₃ from natural O₃ is largest in parts of the western US with extreme wildfire effects. Large impacts on natural O₃ from wildfire events can be seen in Idaho, Wyoming, and California. The contribution from natural O₃ is nearly always less than the contribution from US anthropogenic O₃ in the eastern US. However, in much of the western US (excluding California and large urban areas) the contribution from natural O₃ typically exceeds that of US anthropogenic O₃. On average the natural contribution is higher in the western US than in the eastern US (western US 34 ppb, eastern US 22 ppb for 12 km simulations) which reflects the greater prevalence of wildfires in the western US, a larger background contribution from stratospheric O₃ due to the higher elevation of the western US, and a larger impact from both long-range and short-range (Canada+Mexico) international sources. The contribution from long-range international MDA8 O₃ has a maximum of 20 ppb in the western US and is typically lower in the eastern US compared to the western US on average (western US 6 ppb, eastern US 2 ppb for 12 km simulations). The seasonal average of the long-range international contribution is highest in the spring while base MDA8 O₃ is typically highest in the summer (Figure 3), so days with the highest total O₃ tend not to be the same days with the highest long-range international O₃. The contribution from Canada+Mexico MDA8 O₃ is largest in states along the southern and northern borders as expected. Contributions from Canada+Mexico tend to be small except in border areas. The average MDA8 O₃ contributions on days of the top ten highest base MDA8 O₃ levels are similar to the results for the fourth highest MDA8 O₃ shown here (Figure S3).





Figure 4. MDA8 O₃ on the day of the fourth highest base case MDA8 O₃ from Policy Assessment CMAQ simulations. Results are shown for 12 km (top row), 36 km (middle row), and 108 km (bottom row) horizontal resolutions. O₃ concentrations include total (base) O₃ as well as O₃ components from US anthropogenic, natural, long-range international, and Canada+Mexico sources.

The A second set of simulations (EQUATES) split USBUS background O₃ to different components compared to the PA simulations. The use of different USBUS background O₃ components provides additional insight into the source-specific biases in USBUS background O₃. CMAQ simulated O₃ results from the 2016 EQUATES simulations are comparable to the results from the PA simulations for the 12 km simulations, though the EQUATES simulations have slightly less O₃ from USAUS anthropogenic and more from USBUS background compared to the PA simulations (Figure 45; Table 3). USAUS anthropogenic O₃ contributed ~20% of annual average BASE-base O₃ across all US model grid cells (~25% for PA simulations). Like in the PA simulations, the contribution to USAUS anthropogenic O₃ was higher in the eastern US (~25% of BASEbase) than in the western US (~15% of BASEbase).

STRATStratospheric O₃ is higher in the western US, especially at higher elevations, which is consistent
 405 with previous studies (Jaffe et al., 2018). On average, **STRATstratospheric** O₃ is 40% of **BASE-base** O₃
 in the western US and 34% of **BASE-base** O₃ in the eastern US. **STRATStratospheric** O₃ represents an
 upper bound of stratospheric influences because the tracer species used for its calculation in this study
 does not undergo chemical losses. Non-**STRATstratospheric US background** O₃ (i.e., **USB_NOSTRAT**)
 410 contributes 47% of annual average **BASE-base** O₃ in the western US and 42% in the eastern US. **USBNon-**
stratospheric US background O₃ _NOSTRAT is likely underestimated in regions and seasons with more
 active chemistry due to the use of the chemically inert tracer species used to calculate **USBnon-**
stratospheric US background_NOSTRAT. The 108 km hemispheric CMAQ (H-CMAQ) results for the
 EQUATES and PA simulations are similar on average but do have some notable differences. The H-
 CMAQ simulations are similar in their simulation of **USBUS background** O₃. The **USAUS anthropogenic**
 415 O₃ contributions are also similar on average, though the PA simulations have higher maximum values
 compared to the EQUATES simulations which leads to higher maximum values of **BASE-base** O₃.

Table 3. Summary of annual average of MDA8 O₃ components for the EQUATES set of simulations. Averages are shown for all of the US and separately for the eastern and western US with a longitude of 97 °W serving as the east-west dividing line. The mean across all grid cells within the given area is shown along with the minimum and maximum for any grid cell within the given area in parentheses. Numbers in the table are in units of ppb. Seasonal averages are provided in Table S14.

	BASEBase	USAUS anthropogenic	USBUS background	USBNon-stratospheric US background_NO STRAT	STRATStratosp heric
EQUATES 12 km					
all US	39 (22, 51)	7 (-4, 18)	32 (24, 44)	17 (8, 23)	15 (12, 22)
eastern US	38 (30, 45)	9 (1, 15)	29 (24, 36)	16 (8, 23)	13 (12, 19)
western US	40 (22, 51)	5 (-4, 18)	35 (25, 44)	19 (12, 22)	16 (12, 22)
EQUATES 108 km					
all US	41 (31, 49)	8 (2, 18)	33 (26, 41)	---	---
eastern US	40 (31, 49)	10 (3, 18)	30 (26, 38)	---	---
western US	41 (32, 49)	6 (2, 12)	36 (29, 41)	---	---







Figure 45. Annual average MDA8 O₃ from EQUATES CMAQ simulations. Results are shown for 12 km resolution (top and middle rows) and 108 km (bottom row). O₃ concentrations include total (**BASE**base) O₃ as well as O₃ components from **USA**US anthropogenic, **USB**non-stratospheric US background-**NOSTRAT**, and **STRAT**stratospheric sources for 12 km. For both the 12 km and 108 km simulations, O₃ concentrations of **BASE**base, **USA**US anthropogenic, and total **USB**US background are also shown.


430

435

BASE-base O₃ in EQUATES is highest in the summer (Figure 56). USBUS background O₃ is the highest during spring throughout most of the US. However, in much of the Mountain West, USBUS background O₃ is highest during the summer (Figures S34 and S45). The STRATstratospheric O₃ tracer is the highest in the western US. Much of the western US has STRATstratospheric O₃ at about the same level in the spring and summer. In the southeastern US, STRATstratospheric O₃ is highest in the summer while in the northeastern US, there are similar levels of STRATstratospheric O₃ in the spring and summer. SSTRATstratospheric O₃ is elevated in the summer because of the lack of chemical sinks due to the inert tracer species used to estimate STRATstratospheric O₃. Most previous studies have indicated that stratospheric O₃ peaks in the spring (Lin et al., 2015). The stratospheric contribution to O₃ from H-CMAQ calculated using the decoupled direct method (which does account for chemical losses) also showed higher stratospheric contributions in spring than in summer (Mathur et al., 2022). The higher summer STRATstratospheric O₃ here is explained by the lack of chemical losses due to the tracer method used. Potential biases are explored further in Section 3.3. USAUS anthropogenic O₃ is highest in the summer in the eastern US and in California, consistent with the PA simulations. Non-STRATstratospheric USBUS background O₃ is relatively uniform outside of summer, though it tends to be slightly lower in the southeast and higher in the western US.

The results from both the PA and EQUATES simulations indicate that USBUS background O₃ contributes more than USAUS anthropogenic O₃ to BASE-base O₃ on an annual average basis. Simulated USBUS background O₃ is higher in the western US than in the eastern US due to greater impacts from both natural and non-domestic anthropogenic sources. Simulated USAUS anthropogenic O₃ is higher in the eastern US than in the western US due to the higher population density and consequently greater anthropogenic emissions. The contributions from USAUS anthropogenic O₃ peak in the summer which causes BASE-base O₃ to peak in the summer as well. USBUS background O₃ varies by season but is not as seasonally variable as USAUS anthropogenic O₃. These results are broadly consistent with previous efforts to quantify USBUS background and USAUS anthropogenic O₃ using CTMs (McDonald-Buller et al., 2011; Jaffe et al., 2018).





465 **Figure 56.** Seasonal average MDA8 O₃ from EQUATES CMAQ simulations. Results are shown for
 12 km horizontal resolution for winter (DJF), spring (MAM), summer (JJA), and fall (SON). O₃
 concentrations include total (**BASE**base) O₃ as well as O₃ components from **USA**US anthropogenic,
USBnon-stratospheric US background_NO**STRAT**, and **STRAT**stratospheric sources. Seasonal
 470 averages for the other **USB**US background O₃ split cases are provided in the SI (Figures S34 and
 S45).

Similarly to the PA simulations, we examine the fourth highest total (base) MDA8 O₃ along with the contribution from each of MDA8 O₃ components on the same day for the EQUATES simulations. Like in the PA simulations, the areas with the greatest fourth highest MDA8 O₃ values for base MDA8

475 O₃ tend to have larger contribution from US anthropogenic O₃ than from US background O₃. The
EQUATES fourth highest base MDA8 O₃ is slightly lower than in the PA simulations (56 ppb in the
western and 57 ppb in the eastern US compared to 60 and 61 ppb in the PA simulations at 12 km). The
US anthropogenic contribution is similarly lower in the EQUATES simulations (10 ppb in the western
US, 25 ppb in the eastern US compared to 14 and 33 ppb in the PA simulations at 12 km). The
480 contributions from US background O₃ are higher in the western US than in the eastern US on average
(eastern US 32 ppb, western US 46 ppb for 12 km simulations). The contribution from non-stratospheric
US background O₃ (western US 25 ppb, eastern US 18 ppb) is generally greater than the contribution
from stratospheric US background O₃ (western US 21 ppb, eastern US 14 ppb). The western US has larger
contributions from stratospheric O₃, long-range international O₃, and wildfires. In the EQUATES
485 simulations, the Flint Hills area of Kansas stands out as an area influenced by fires. The fires in this area
are typically prescribed burning of grasslands used for agricultural land management. While these were
included in the fire emissions for the US background O₃ simulation, prescribed burns are typically
classified as anthropogenic sources rather than background sources. The average MDA8 O₃ contributions
on days of the top ten highest base MDA8 O₃ levels are similar to the results for the fourth highest MDA8
490 O₃ shown here (Figure S6).




Figure 7. MDA8 O₃ on the day of the fourth highest base case MDA8 O₃ from EQUATES CMAQ simulations. Results are shown for 12 km resolution (top and middle rows) and 108 km (bottom row). O₃ concentrations include total (base) O₃ as well as O₃ components from US anthropogenic, non-stratospheric US background, and stratospheric sources for 12 km. For both the 12 km and 108 km simulations, O₃ concentrations of base, US anthropogenic, and total US background are also shown.

495

500 3.2 Cross-validation of regression modeling

Overfitting is tested using a cross-validation analysis as described in Section 2.2. Three different cross-validation methods are used: spatial and temporal withholding, spatial withholding, and temporal withholding. The parameters derived from the training set are then used to predict the observed O₃ in the test set. The RMSE and mean bias with respect to the true observations of both the training and test set are compared to one another (Table 4; Tables S7 and S8). For each of the three cross-validation methods, the RMSE and mean bias of the training and test sets are similar to one another. This indicates that the model is not overfitting and is generalizable to data outside of its training data, providing confidence that we can apply the regression models to the gridded CTM results to estimate the bias in O₃ and individual O₃ components across the US.

510

Table 4. Summary of performance for cross-validation of MDA8 O₃ data fusion model. Values shown are the average over all regression model cases. RMSE and mean bias statistics for individual cases are provided in Tables S7 and S8. The performance for the **BASE-base O₃ simulations prior to applying the bias adjustment is also provided for comparison.**

metric	BASE-Base simulations	spatial and temporal withholding		spatial withholding		temporal withholding	
		training	test	training	test	training	test
RMSE (ppb)	9.53	7.80	7.83	7.83	7.58	7.81	7.79
mean bias (ppb)	1.13	-0.19	-0.20	-0.19	-0.63	-0.19	0.38

515

3.3 Inferred CTM biases


The coefficients from the regression models (Tables S9 – S12) are applied to the gridded CTM data to calculate adjusted values of each O₃ component. The inferred CMAQ bias for each component is the difference between the original CMAQ-simulated value and the adjusted value. The inferred bias in **BASE-base** O₃ is the original CMAQ-simulated **BASE-base** O₃ minus the sum of adjusted O₃ components. For the PA simulations, there is a residual anthropogenic component of **BASE-base** O₃ that is not apportioned to either **USAUS anthropogenic** or **INTL-international** sources due to the effects of non-linear chemistry (Table S2). The residual anthropogenic component is equal to **BASE-base O₃ – NATnatural O₃ – INTL-international O₃ – USAUS anthropogenic O₃**. This means that the sum of biases in the individual


525 components do not add up to the bias in BASE-base O₃ as the residual anthropogenic component was not included in the adjusted O₃ results. In the PA simulations, BASE-base O₃ is inferred to be biased high in most of the Eastern US as well as in some parts of California and Arizona (Figure 68). USAUS anthropogenic O₃ is inferred to be biased high in the same areas. Reducing the amount of USAUS anthropogenic O₃ improves the fit to BASE-base O₃ which is suggestive that biases in the effects from
530 US anthropogenic emissions contribute to the high biases inferred in BASE-base O₃. The inferred high biases in BASE-base and USAUS anthropogenic O₃ increase with increasing coarseness of model resolution in the eastern US. Similarly, the high bias increases with coarser model resolution in the CANMEXCanada+Mexico component along the border with Mexico. The inferred high biases in USAUS anthropogenic O₃ in the eastern US are primarily driven by biases in the summer and fall (Table S15, 535 Figures S57-S79). Inferred eastern US USAanthropogenic O₃ biases average 2, 7, and 11 ppb in the summer and 3, 4, and 5 ppb in the fall for the 12, 36, and 108 km simulations. In the western US, where USAUS anthropogenic O₃ is mostly found to be biased low, coarser model resolution results in the summer average bias changing from slightly negative in the 12 km simulations (-0.5 ppb) to slightly positive in the 36 and 108 km simulations (+0.7 ppb and +1.0 ppb).

540 In contrast to our results showing an increase in O₃ with coarser resolution, Schwantes et al. (2022) found that O₃ tended to increase for a finer resolution simulation (~14 km vs. ~111 km over the CONUS) during the summer over urban areas using the Community Earth System Model (CESM)/Community Atmosphere Model with full chemistry (CAM-chem) model which was attributed to improvements in the spatial resolution of NO_x emissions resulting in less artificial dilution of NO_x and enhanced O₃ production.
545 Similarly, Lin et al. (2024) found that a variable resolution global model (AM4VR with horizontal resolution of 13 km over the CONUS) had increased O₃ over urban areas compared to a fixed resolution model (AM4.1 with horizontal resolution of ~100 km globally). In particular for the Los Angeles Basin and Central Valley regions of California, Lin et al. (2024) found that the increased resolution of AM4VR led to better simulation of observed O₃ levels in these areas due the finer resolution model's ability to
550 represent sharp spatial gradients in areas with NO_x-limited vs. NO_x-saturated O₃ production regimes. Our analysis of the fourth highest MDA8 O₃ levels shows similar findings over California (Figures 4 and 7). Given these previous results finding increased O₃ with finer resolution simulations, our results here

finding higher biases in USAUS anthropogenic O₃ in the eastern US with coarser resolution should be taken to apply specifically to the CMAQ model results described here rather than as a general finding on the impact of model resolution on O₃ production. Additionally, given that the finding of higher US anthropogenic O₃ with coarser model resolution does not hold for the analysis of the fourth highest MDA8 O₃ levels, this finding should be taken to apply only to longer term (e.g., annual or seasonal) averages.

There are offsetting inferred biases in the LINTLlong-range international and NATnatural O₃ components in much of the western US. The offsetting inferred biases may reflect an inability of the regression model to separate the signals from LINTLlong-range international and stratospheric O₃. LINTLLong-range international and stratospheric O₃ are expected to impact sites at similar spatial and temporal scales, with larger impacts expected at high elevations in the western US during spring. Stratospheric O₃ effects are not limited to episodic intrusion events but also come from constant entrainment of stratospheric air to the free troposphere. The impacts from LINTLlong-range international emissions are primarily from long-range transport in the free troposphere, so stratospheric O₃ and LINTLlong-range international O₃ are expected to be correlated. The regression model may be assigning bias due to stratospheric O₃ to LINTLlong-range international O₃ because the CTM-modeled LINTLlong-range international component has more correlation with the stratospheric O₃ impact than the CTM-modeled NATnatural component. This could result in the regression model adjusting LINTLlong-range international O₃ upwards (i.e., inferred negative bias) to add stratospheric O₃. The NATnatural O₃ is then adjusted downwards (i.e., inferred positive bias) in the same locations because some of the effects of stratospheric O₃ are captured in the CTM-modeled NATnatural O₃ component but need to be offset because of the O₃ that was added to the LINTLlong-range international component. This indicates a limitation of this method in that it is sensitive to correlation between modeled O₃ components. Correlation of the O₃ components is a major confounding issue in this analysis. In interpreting the results, it is necessary to consider both the inferred biases and the correlation of the components together.







580 **Figure 68.** Annual average of inferred MDA8 O₃ model bias from Policy Assessment CMAQ
simulations. Results are shown for 12 km (top row), 36 km (middle row), and 108 km (bottom row)
horizontal resolutions. O₃ concentrations include total (**BASE**base) O₃ as well as O₃ components
from **USAUS** anthropogenic, **NAT**natural, **LINTL**long-range international, and
CANMEXCanada+Mexico sources. Seasonal averages are provided in Figures S57-S79.

585

In the temporal trends of inferred **BASE**base O₃ bias, the PA simulations show a consistent low
bias in winter and spring and high bias in summer and fall which is consistent across model resolution
scales (Figure 79). There is also a consistent high bias in **USAUS** anthropogenic O₃ in summer and fall
in the eastern US which increases with coarser model resolution. Inferred bias in **USAUS** anthropogenic
590 O₃ in the western US has some small seasonal variability but is near zero on average. The seasonal patterns
of **LINTL**long-range international O₃ bias have the largest underestimate in the winter and spring and the
smallest underestimate in late summer and early fall. The temporal trend of **NAT**natural O₃ differs in the
12 km simulation compared to the 36 km and 108 km simulations. In the 12 km simulation, **NAT**natural
O₃ biases are higher in the middle of the year than in the beginning and end of the year. In the 36 km and

595 108 km simulations, the opposite is found. This change in sign is a result of changes in the spatial patterns
of NATnatural O₃ inferred bias in different seasons. In the 12 km simulation, NATnatural O₃ is inferred
to be biased low in the southern part of the US and biased high in the northern part of the US. In the 36
km and 108 km simulations NATnatural O₃ is inferred to be biased low in the eastern US and mostly
biased high in the western US, particularly in the Mountain West region. These spatial changes in the
600 seasonal average NATnatural O₃ bias are enough to change the sign of the US average temporal bias
trend. As described before, the offsetting negative LINTLlong-range international bias and positive
NATnatural O₃ bias in the high elevation areas of the western US are thought to be a result of the
regression model allocating stratospheric O₃ bias to the LINTLlong-range international O₃ signal while
removing some stratospheric O₃ from the NATnatural O₃ signal. CANMEXCanada+Mexico O₃ biases
605 are very small when averaged across the US since this source primarily affects border areas and only has
small impacts elsewhere.






610 **Figure 79.** Daily average of inferred MDA8 O₃ model bias from Policy Assessment CMAQ
simulations averaged across US model grid cells in the eastern and western US. A longitude of 97
°W is used as the dividing line between east and west. PA O₃ concentrations include total
(**BASE**base) O₃ as well as O₃ components from **USA**US anthropogenic, **NAT**natural, **LINT**long-
615 **range international**, and **CANMEX**Canada+Mexico sources. **USBUS** background indicates the sum
of biases for individual **USBUS** background components.

The spatial results for the EQUATES 12 km simulations are shown for two O₃ split cases. One
case splits **USBUS** background O₃ to **STRAT**stratospheric and non-**STRAT**stratospheric sources while
the other considers all **USBUS** background O₃ together. Results show a mostly low bias inferred in **BASE**
620 **base** O₃ throughout most of the US for the 12 km simulation (Figure 810). For the 108 km H-CMAQ
simulation there is a high bias in the eastern US and a low bias in the western US for **BASE**-**base** O₃. Like
the PA results there is a high bias in **USA**US anthropogenic O₃ in the eastern US that increases with

coarser model resolution. The inferred low bias in the STRATstratospheric O₃ component indicates that there is too little stratospheric O₃ in the western US. There is an inferred high bias in STRATstratospheric O₃ in the eastern US. The STRATstratospheric O₃ results should be interpreted with some caution because the STRATstratospheric component comes from a chemically inert tracer. The STRATstratospheric O₃ biases are partly offset by opposite biases in the non-STRATstratospheric USBUS background O₃. The low biases in STRATstratospheric O₃ and the lack of low biases in the non-STRATstratospheric USBUS background O₃ provides more evidence that the low biases in the LINTLlong-range international O₃ from the PA simulations are related to low biases in stratospheric O₃.

In the case where USBUS background O₃ is not split into STRATstratospheric and non-STRATstratospheric components, the 12 km and 108 km simulations both have low biases in USBUS background O₃, but the magnitude of bias is greater in the 12 km simulation than in the 108 km simulation. This may be a result of differences in the impacts of stratospheric O₃ at the surface level in the H-CMAQ simulation compared to the continental-scale simulation. Differences in the estimation of stratospheric O₃ impacts may arise from differences in how the vertical structure of the model in the H-CMAQ simulations is configured compared to the continental simulations. The UTLS PV O₃ scaling is turned on during the H-CMAQ simulation. For the continental simulation, PV O₃ scaling is turned off because the continental model configuration uses fewer vertical layers and a coarser vertical resolution in the UTLS compared to the H-CMAQ simulations. The stratospheric O₃ influences in the continental simulation are only those influences that are inherited from the lateral boundary conditions. Previous work indicates that O₃ in the upper layers of the continental-scale model is driven mostly by horizontal advection of the lateral boundary conditions (Hogrefe et al., 2018), meaning that if stratospheric intrusion events are captured by the hemispheric-scale simulation, the effects of these events are also expected to be captured by the continental-scale simulation. However, a sensitivity test with UTLS PV O₃ scaling turned on during the continental simulation may be an area for future study. This would require the addition of more vertical layers with finer resolution in the UTLS in the continental simulation to support the PV O₃ scaling parameterization. The differences in vertical structure of the hemispheric and continental simulations can affect the vertical mixing of stratospheric O₃ from upper layers down to the surface which may explain the differences in inferred bias of USBUS background O₃. Alternatively, the differences in USBUS

background O₃ biases could also occur due to differences in O₃ production from local ~~USB~~US background O₃ sources across model resolution scales and may not necessarily be affected by differences in stratospheric O₃.



655





Figure 810. Annual average of inferred MDA8 O₃ model bias from EQUATES CMAQ simulations. Results are shown for 12 km resolution (top and middle rows) and 108 km (bottom row). O₃ concentrations include total (**BASEbase**) O₃ as well as O₃ components from **USAUS anthropogenic**, **USBnon-stratospheric US background-NOSTRAT**, and **STRATstratospheric** sources for 12 km. For both the 12 km and 108 km simulations, O₃ concentrations of **BASEbase**, **USAUS anthropogenic**, and total **USBUS background** are also shown. Seasonal averages are provided in Figures S810-S1012.

For the EQUATES temporal results, **BASE-base** O₃ is biased low in the spring and high in the summer in the eastern US (Figure 911). In the western US, **BASE-base** O₃ is biased low throughout most of the year. Averaged across the US, bias is near zero in the summer and fall in the 12 km simulation with

high biases in the 108 km simulation during the same period (+1 ppb in summer; +2 ppb in fall). The high biases in BASE-base O₃ in the eastern US are mostly due to high biases in the USAUS anthropogenic O₃ component which peak in the summer (average +1.4 and +6.0 ppb for the 12 and 108 km simulations) and continue to be biased high into the fall (average +0.8 and +2.2 ppb for the 12 and 108 km simulations). The STRATstratospheric O₃ component is inferred to be biased low except in the summer and early fall. In the western US, STRATstratospheric O₃ bias in the summer is near zero in the summer and fall while in the eastern US, STRATstratospheric O₃ is biased high in the summer and fall. The lowest biases in STRATstratospheric O₃ occur in the winter. The STRATstratospheric O₃ biases are partially offset by opposing biases in the non-STRATstratospheric USBUS background O₃. The regression model formulation without the separate STRATstratospheric O₃ indicates that there is a low bias in USBUS background O₃ throughout most of the year in the 12 km simulation which is at its lowest in the spring. The 108 km simulations show a low bias for USBUS background O₃ in the spring and summer and high bias in the fall and winter.

In the 12 km EQUATES simulations, the STRATstratospheric O₃ tracer averages 14 ppb in the western US during spring, with a maximum spring average across all western US grid cells of 17 ppb. Using the bias correction approach developed here, we find that the spring average STRATstratospheric O₃ in the western US is biased low by 3.5 ppb, resulting in an adjusted (i.e., bias corrected) estimate of western US spring average STRATstratospheric O₃ of 17 ppb. Consistent with the low bias in stratospheric O₃ suggested here, other CTMs have estimated higher stratospheric O₃ contributions compared to those simulated here with CMAQ. The spring average of stratospheric O₃ contributions estimated with the AM3 model has been estimated at 20-25 ppb (Lin et al., 2012a; Langford et al., 2015; Lin et al., 2015). The AM3 estimates of stratospheric O₃ have sometimes been estimated to be biased high (Lin et al., 2012a) and have also been shown to lead to overestimated springtime O₃ concentrations when used as boundary conditions for regional-scale CMAQ simulations (Hogrefe et al., 2018) but at other times have been estimated to be relatively unbiased based on evaluation against observations from intensive field studies (Langford et al., 2015). The stratospheric O₃ contribution simulated by AM3 has been previously found to be higher than that of the GEOS-Chem global model (Fiore et al., 2014). Using GEOS-Chem, Zhang et al. (2014) found the spring mean stratospheric O₃ influence in the Intermountain

West to range from 8-10 ppb as estimated using the standard GEOS-Chem definition of stratospheric O₃ as described in Zhang et al. (2011) and, alternatively, found a spring mean of 12-18 ppb using a definition of stratospheric O₃ adopted from Lin et al. (2012a) (the same method used for the AM3 estimates reported here). Itahashi et al. (2020) previously found that the stratospheric O₃ representation in CMAQ was biased
700 low in the free troposphere and suggested that improvements were needed to the CMAQ representation of stratosphere to troposphere transport. Our bias adjusted estimate of western US spring mean stratospheric O₃ (17 ppb) falls in between the estimates from the default GEOS-Chem representation (8-10 ppb) and from AM3 (20-25 ppb). As these are seasonal averages, the values are more representative of the continual entrainment of stratospheric air into the troposphere rather than episodic deep
705 stratospheric intrusion events.






Figure 911. Daily average of inferred MDA8 O₃ model bias from EQUATES CMAQ simulations averaged across US model grid cells in the eastern and western US. A longitude of 97 °W is used as the dividing line between east and west. EQUATES O₃ concentrations include **BASE-base** O₃ as well as O₃ components from **USAUS anthropogenic**, **USBnon-stratospheric US background_NOSTRAT**, and **STRATstratospheric** sources for 12 km. For both the 12 km and 108 km simulations, O₃ concentrations of **BASEbase**, **USAUS anthropogenic**, and total **USBUS background** are also shown. For the case with multiple **USBUS background** O₃ components, **USBUS background** indicates the sum of biases for individual **USBUS background** components.

3.4 CTM biases by O₃ concentration


The contributions and biases of different O₃ components have been presented so far as annual or seasonal averages (Figures 2-3, 5-6, 6 and 8, and 10), as the fourth highest value which is relevant from a regulatory perspective (Figures 4 and 7), or as daily averages over US model grid cells (Figures 79 and

911). However, the relative contributions of O₃ components at different total O₃ concentrations is also of interest. For example, the relative contribution of USAUS anthropogenic and USBUS background O₃ to total O₃ may be different on days with higher total O₃ vs. days with lower total O₃. Situations where O₃ exceeds the NAAQS, which is currently set at a level of 70 ppb, are of particular interest. We analyze the different O₃ components at O₃ monitoring sites under cases when O₃ is less than 60 ppb, between 60 and 70 ppb (inclusive), and greater than 70 ppb. These concentration bins are selected because they reflect the current level of the standard (70 ppb) as well as a potential range which might be considered as the level of the standard in the future (60-70 ppb). We compare the results of the analysis when using both simulated and observed O₃ bins. Simulated O₃ has a positive bias on average when simulated O₃ is high and has a negative bias on average when observed O₃ is high, so selection bias influences these results. For this analysis, we consider the 12 km resolution simulations for the PA and EQUATES simulations. The 12 km simulations are the resolution that is typical for simulations that support regulatory analyses. Monitoring sites are split into western or eastern US using a longitude of 97 °W as the dividing line. The division to western and eastern US is done because there are differences in the contribution of US anthropogenic vs. background contributions in the two parts of the country.

The impacts of the linear regression adjustment technique at the observation sites are examined by comparing the original simulated bias to the residual bias (i.e., the sum of the adjusted individual O₃ components minus observed O₃) (Figure 1012). The change in bias from the original to residual bias is the inferred bias that has been referenced elsewhere. In all cases when O₃ is binned by simulated O₃ levels, the adjustment brings the bias closer to zero. In the eastern US, high biases at higher simulated O₃ levels were reduced for both the PA and EQUATES simulations. In the western US, low biases when simulated O₃ was below 60 ppb were brought closer to zero for both the PA and EQUATES simulations. At higher simulated O₃ levels, the PA simulations originally had high biases in the western US which were reduced in the adjusted results while the EQUATES simulations originally had low biases in the western US which were improved in the adjusted results. The effects on bias when binning by observed O₃ are mixed. In both the western and eastern US for both the PA and EQUATES simulations, the simulations were originally biased low at higher observed O₃ levels, with the EQUATES simulations more biased low than the PA simulations. The low bias is improved in the EQUATES simulations, but in the PA simulations

750 the bias is either about the same or becomes more biased low. The inability of the adjustment to improve the bias across the range of both observed and simulated O₃ levels is a limitation of this technique. The fitting of multi-axis (lat, lon, season) linear correction factors (α_i) will be strongly influenced by the larger population of lower (O₃ < 70 ppb) concentrations and will only correct the upper end if the bias structure is consistent.

755



760

Figure 1012. Bias compared to MDA8 O₃ observations of original simulations (black) and residual bias (purple) obtained as the difference between adjusted MDA8 O₃ and observations for PA (top row) and EQUATES (bottom row) simulations. The horizontal line shows the median; the box shows the 25th-75th percentiles; the whiskers show the 5th and 95th percentiles. Grey vertical lines separate the boxplots for each MDA8 O₃ concentration bin. The numbers at the bottom of each panel are the number of data points falling within each concentration bin.

For the PA simulations, the contribution from USAUS anthropogenic O₃ tends to increase with
765 higher simulated O₃ and with higher observed O₃ (Figure 4+13), indicating that domestic anthropogenic
pollution is driving the highest O₃ concentrations. The contribution from USAUS anthropogenic O₃ is
higher at eastern US sites than at western US sites due to higher anthropogenic precursor emissions in the
east. There may also be impacts on USAUS anthropogenic O₃ in the eastern US from O₃ or precursor
pollutants transported from the western to eastern US. The median USAUS anthropogenic O₃ contribution
770 is biased high (+1 ppb in the western US; +4 ppb in the eastern US) when BASE-base O₃ is between 60
and 70 ppb with higher median biases (+2 ppb in the western US; +6 ppb in the eastern US) when BASE
base O₃ exceeds 70 ppb. When observed O₃ is between 60 and 70 ppb, the median USAUS anthropogenic
O₃ contribution is biased slightly low in the western US (-0.2 ppb) and biased high in the eastern US (+2
ppb). Bias is higher in the western US when observed O₃ exceeds 70 ppb (+1 ppb) but is about the same
775 in the eastern US (+2 ppb). Inferred biases of USAUS anthropogenic O₃ are higher across the range of
simulated and observed O₃ levels in the eastern US compared to the western US.

In the western US, NATnatural O₃ tends to be higher when either simulated or observed O₃ is
greater than 60 ppb; however, the distribution of NATnatural O₃ when O₃ is above 70 ppb is similar to
the distribution of NATnatural O₃ when O₃ is between 60 and 70 ppb. In the eastern US, the distribution
780 of NATnatural O₃ is similar across the range of simulated and observed O₃ concentration bins but is
slightly higher when O₃ is greater than 60 ppb. LINTLlong-range international makes a small
contribution to O₃ across concentration bins and tends to be lower as simulated or observed O₃ increases.
CANMEXCanada+Mexico O₃ is typically very small and only makes significant contributions at a few
near-border sites (not shown). The NATnatural and LINTLlong-range international O₃ components are
785 biased slightly low at monitoring sites in the western US. For western US sites, the sum of the median
biases in USAUS anthropogenic and USBUS background (i.e., NATnatural + LINTLlong-range
international + CANMEXCanada+Mexico) O₃ at monitoring sites is negative across the simulated and
observed O₃ concentration bins but gets closer to zero at higher O₃ levels. For eastern US sites, the bias
in USAUS anthropogenic O₃ is predicted to be the main contributor to biases at high simulated O₃ when
790 simulated O₃ concentrations exceed 60 ppb. When the O₃ components are binned by observed O₃ rather
than simulated O₃, the sum of the median biases in USAUS anthropogenic and USBUS background O₃

at monitoring sites in the eastern US is negative across the range of simulated O₃ with US&US background O₃ becoming less negatively biased as observed O₃ increases and USA&US anthropogenic O₃ becoming more positively biased as observed O₃ increases.

795




Figure 1413. Contributions to MDA8 O₃ from the PA simulation (top row) and inferred biases (bottom row) of **USAUS** anthropogenic, **NAT** natural, **LINTL** long-range international, and **CANMEX** Canada+Mexico separated by both observed and simulated **BASE**-base MDA8 O₃

800

concentration at O₃ monitoring sites. The sum of NATnatural, LINTLlong-range international, and CANMEXCanada+Mexico is shown as USBUS background. The horizontal line shows the median; the box shows the 25th-75th percentiles; the whiskers show the 5th and 95th percentiles. Grey vertical lines separate the boxplots for each MDA8 O₃ concentration bin. The numbers in the bottom row of panels are the number of data points falling within each concentration bin.

For the 12 km EQUATES simulations, the USAUS anthropogenic O₃ contribution is similar to the 12 km PA results across the simulated O₃ concentration bins (Figure 1214). At higher observed O₃, the EQUATES simulations generally simulate lower USAUS anthropogenic O₃ compared to the PA simulations. Like in the PA simulations, the USAUS anthropogenic O₃ contribution increases with increasing simulated and observed O₃, meaning that domestic anthropogenic emissions are mostly driving the highest O₃ levels. There is an inferred negative bias in USAUS anthropogenic O₃ in the western US which becomes increasingly more negative as simulated or observed O₃ increases. In the eastern US, there is an inferred positive bias in USAUS anthropogenic O₃ which becomes larger at higher simulated O₃ concentrations (median bias of +0.05, +2, +4 ppb at <60, 60-70, and >70 ppb simulated O₃). There is also an inferred high bias across the range of observed O₃; however, the magnitude is smaller, and the bias does not increase much at higher levels of observed O₃ (median bias of +0.05, +0.5, and +0.6 ppb at <60, 60-70, and >70 ppb observed O₃).

The contribution from STRATstratospheric O₃ is higher in the western US than in the eastern US across simulated and observed O₃ concentrations. In the western US, STRATstratospheric O₃ tends to be higher when either observed or simulated is above 60 ppb. In the eastern US, STRATstratospheric O₃ is at similar levels across the range of simulated and observed O₃. In the western US, STRATstratospheric O₃ has a negative bias which gets closer to zero when simulated and observed O₃ is above 60 ppb. In the eastern US, STRATstratospheric O₃ has a positive bias which gets higher when simulated and observed O₃ are above 60 ppb. In both the western and eastern US, USBnon-stratospheric US background-NOSTRAT O₃ makes similar contributions across different O₃ concentrations. In the western US, USBnon-stratospheric US background-NOSTRAT O₃ has a negative bias when simulated or observed O₃ is below 60 ppb and a positive bias when O₃ is above 60 ppb. In the eastern US, USBnon-stratospheric US background-NOSTRAT O₃ has a negative bias across the range of simulated and

observed O_3 . The magnitude of the negative bias is smaller when simulated or observed O_3 is below 60 ppb than when O_3 is above 60 ppb.




Figure 1214. Contributions to MDA8 O₃ by the EQUATES simulation (top row) and inferred biases (bottom row) of USAUS anthropogenic, USBUS background–NOSTRAT no stratospheric, and STRATstratospheric separated by both observed and simulated BASE–base MDA8 O₃


840 concentration at O₃ monitoring sites. The sum of USB_{non-stratospheric} US
background–NOSTRAT and STRAT_{stratospheric} is shown as USBUS background. The line shows
the median; the box shows the 25th-75th percentiles; the whiskers show the 5th and 95th
percentiles. Grey vertical lines separate the boxplots for each MDA8 O₃ concentration bin. The
845 numbers in the bottom row of panels are the number of data points falling within each
concentration bin.

Binning the O₃ contributions and inferred biases by observed and simulated O₃ results in different
numbers of data points in each sample. In the western US, there were 4145 instances when observed O₃
exceeded 70 ppb, while there were 3302 (PA) and 627 (EQUATES) instances when simulated O₃
850 exceeded 70 ppb at a monitoring site, with a large fraction of the observed and simulated exceedances
occurring in California. In the eastern US there were 2135 instances when observed O₃ exceeded 70 ppb
with 2901 (PA) and 556 (EQUATES) instances when simulated O₃ exceeded 70 ppb. The PA simulations
more accurately simulated the number of exceedances compared to EQUATES, though this does not
consider the timing or location of exceedances. Given the different number of samples in the observed
855 vs. simulated bins and the lower number of data points for EQUATES simulated O₃ exceeding 70 ppb, it
is possible that the population of data points are when simulated O₃ exceeds 70 ppb are not spatially
representative of the population when observed O₃ exceeds 70 ppb.

For the western US, the PA simulations largely capture the spatial distribution of exceedances
seen in the observations, although the number of exceedances is underestimated (Figure ~~13~~15). The
860 exceedances from the EQUATES simulations are not very representative of the spatial distribution of
observed exceedances in the western US as there are very few sites with more than one or two exceedances
outside of California. In particular, the number of exceedances in the Denver, Colorado; Phoenix,
Arizona; Las Vegas, Nevada; and Boise, Idaho; areas are underestimated in EQUATES relative to both
the PA simulations and observations. Both the PA and EQUATES simulations underestimate the number
865 of exceedances in the state of Utah. For the eastern US, the PA simulations generally capture the spatial
distribution of observed exceedances but simulate too many exceedances. This is particularly notable in
the northeastern US and along the Gulf Coast. The EQUATES simulations underestimate the number of
exceedances, although the spatial distribution is generally similar to the observations. The degree of
spatial representativeness provides additional context for interpreting the findings for the O₃ component

870 contributions and biases binned by O_3 levels. For the western US, the findings for instances when O_3
exceeds 70 ppb are not applicable to the western US more broadly. There are a limited number of instances
when O_3 exceeds 70 ppb in the western US outside of California. These results are mostly indicative of
conditions in the Los Angeles area and in the Central Valley in California. This applies especially to the
EQUATES results, but it is also the case for the PA simulations and the observations. For the eastern US,
875 on the other hand, there is enough spatial variability in the observations as well as both sets of simulations
to interpret the findings for the eastern US more generally. These results are informative in an average
sense but are not expected to hold in all cases when applied to specific monitoring sites or to specific days
(e.g., fourth highest O_3). The biases for bins 60-70 ppb and greater than 70 ppb should be interpreted with
caution because the inferred biases apply the mean tendency to these high concentration subpopulations.

880



885

Figure 1315. Spatial distribution of the number of times MDA8 O_3 exceeded 70 ppb for observed and simulated O_3 . The circles show the locations of sites, and the color indicates the number of times MDA8 O_3 exceeds 70 ppb at each site for observations (left), PA 12 km simulation (middle), and EQUATES 12 km simulation (right). Only sites with at least one exceedance are shown. The black dotted line shows the longitude of 97° W which is used to divide west and east. Similar results for other model resolutions are shown in Figure S113.

4 Conclusions

890 In this work, we use two sets of CMAQ simulations to analyze the contributions to USBUS background O₃ from different sources. Naturally occurring sources, long-range international anthropogenic pollution, and short-range international anthropogenic pollution from Canada and Mexico are separately considered for one set of simulations. In the other set of simulations, stratospheric and non-stratospheric sources of USBUS background O₃ are also considered separately. We also consider the
895 contribution to total O₃ from US domestic anthropogenic sources. The measurement-model data fusion approach for apportioning bias to USAUS anthropogenic and USBUS background O₃ components from our previous study (Skipper et al., 2021) was extended to identify biases in separate USBUS background O₃ components. The results generally confirm previous high-level results, but provide new insights from additional components and more detailed analysis.

900 Results indicated that USAUS anthropogenic O₃ was consistently inferred to be biased high (on an annual and seasonal average basis) in the eastern US where domestic anthropogenic emissions are the dominant contributor to total O₃, with increasingly higher biases with coarser model resolution and at higher simulated O₃ concentrations. This is consistent with our previous findings. This does not necessarily imply that the trend of decreasing biases with finer resolutions would continue at resolutions
905 finer than 12 km as we have not tested this approach at those resolutions. As noted in Section 3.3, previous modeling studies examining the effects of horizontal resolution have found that O₃ increased over urban areas with finer resolution, so the findings for the effects of model resolution should be taken to apply our current results rather than as a general finding on the impacts of model resolution. Our finding that USAUS anthropogenic O₃ biases increase with higher O₃ does not hold when O₃ is binned by observed
910 rather than simulated concentrations. There is much less variation in the USAUS anthropogenic O₃ bias across the range of observed O₃ than for simulated O₃. Although the choice of binning O₃ by observed or simulated levels changes the sample of data, the results for the eastern US are generalizable to this part of the country because the samples have consistent spatial representation across the eastern US. In the western US, USAUS anthropogenic O₃ was inferred to be biased high at higher O₃ levels for the PA
915 simulations and biased low at higher O₃ levels for the EQUATES simulations. These differences are explained by the use of different emission inventories in the two sets of simulations. Regardless, the

findings for inferred O₃ biases at higher O₃ levels in the western US are not broadly applicable to the entire western US because the sample that these findings are based on is dominated by sites in California. There are relatively few sites in other states in the western US that contribute to this sample, so the results are not likely to be indicative of conditions in other parts of the western US.

The correction of USBUS background components provided consistent results with previous studies, but more detail. Like Skipper et al. (2021) and Hosseinpour et al. (2024), simulated USBUS background O₃ was inferred to be biased slightly low overall. The original simulated annual averages of USBUS background O₃ across all the PA and EQUATES modeling configurations considered here ranged from 30-33 ppb while the adjusted annual average USBUS background O₃ ranged from 31-34 ppb. The annual average of simulated US background O₃ for the hemispheric-scale (108 km resolution) and continental-scale (12 km resolution) modeling was slightly higher for the EQUATES simulations (32-33 ppb) than for the PA simulations (30-31 ppb). The differences are not explainable by the updated chemical mechanism used in EQUATES because the most relevant updates (halogen-mediated O₃ loss) tend to reduce O₃ at the northern mid-latitudes (Sarwar et al., 2019; Appel et al., 2021). The difference is also not likely due to anthropogenic emissions outside of the US, which are similar between the two sets of simulations. So, the higher US background O₃ in EQUATES likely relates to differences in the natural emissions. The EQUATES simulations used MEGAN for biogenic emissions throughout the entire Northern Hemisphere while the PA simulations used BEIS for biogenic emissions in North America and MEGAN elsewhere. The two hemispheric model configurations also used different sources for soil NO_x emissions (see Section 2.1) which could contribute to differences in US background O₃. Lightning NO_x emissions were the same in EQUATES and PA hemispheric-scale simulations, but the continental-scale PA simulations did not include lightning in the continental domain. Given that US background O₃ in both the EQUATES and PA 12 km continental-scale simulations are 1 ppb lower than their northern hemispheric counterparts, the differences in US background O₃ in the continental-scale simulations is more likely driven by the large-scale background inherited through the lateral boundary conditions than from differences in lightning NO_x configurations.

This work separated USBUS background O₃ into natural, short-range international, and long-range international components, and each had distinct seasonality to the inferred bias. Short-range

945 international (Canada+Mexico) was marginally high-biased in spring/winter and marginally low-biased
in summer. The contribution from natural and long-range international O₃ have larger seasonality, which
are slightly out of phase. Natural O₃ bias was low in winter, but high in summer peaking in July. Long-
range international O₃ was consistently low-biased with a minimum in April and a maximum (near
unbiased) in August-September. From May to October, the natural and long-range international O₃ biases
950 are largely offsetting while they are reinforcing in other parts of the year.

The seasonality of inferred long-range international bias highlights a key uncertainty in correlative
bias attribution. The biases associated with long-range international may be misattributed due to the
difficulty of the regression model formulation to isolate stratospheric influences from other natural
sources such as lightning and soil NO_x, wildfires, and biogenic VOC emissions, all of which have a high
955 degree of uncertainty. Stratospheric O₃ is expected to have similar temporal and spatial patterns to
LINTLlong-range international O₃, with contributions being higher in spring and at high elevations. It is
suspected that the regression model formulation may be assigning a negative bias in LINTLlong-range
international O₃ to make up for missing stratospheric O₃ that has a similar pattern to LINTLlong-range
international O₃ while at the same time assigning a high bias for NATnatural O₃ to reallocate some of
960 stratospheric O₃ that is present in NATnatural O₃ to LINTLlong-range international O₃ instead. Results
for the STRATstratospheric O₃ tracer in the second set of simulations support the idea that there is missing
stratospheric O₃ at the surface level in the western US as the STRATstratospheric O₃ is inferred to be
biased low. Taken together, there is an overall low bias in the simulated USBUS background O₃ that is
most pronounced in the spring. This may be a result of too little stratospheric O₃ reaching the surface.
965 Photolysis of particulate nitrate over oceans has been found to increase O₃ (Shah et al., 2023; Sarwar et
al., 2024). This process is not included in the chemical mechanism which could contribute to low biases
in O₃ during the same time of year. The potential for misattribution is not specific to the methods
employed here but is inherent to correlative bias approaches with incomplete information contained in
independent variables.

970 Analysis of the original bias and residual bias emphasize the importance of subpopulation
diversity. The correction factors are optimized for the whole population and can degrade performance at
any subpopulation (e.g., a site, a day, or a subgroup). For example, in the western US, the PA simulation

was originally high-biased for days with high predictions and low-biased for days with high observations (>70 ppb). The overall correction was downward for both populations because they are generally consistent spatially and seasonally. This means that the “corrected” model has more bias on days with high observations in the western US than the “uncorrected.” This is not unexpected but highlights that correlative adjustments should be considered as broad conclusions and should only be cautiously applied more narrowly (e.g., specific monitors or days). This is a limitation of the linear formulation as noted by Hosseinpour et al. (2024).

This work has focused only on surface O₃. We are not able to draw a conclusion as to whether the potential lack of stratospheric O₃ is a result of biases in the UTLS PV scaling in the upper layers or from errors in vertical transport from upper layers to the surface. More detailed studies that analyze the entire vertical structure, such as a recent study of CMAQ stratospheric O₃ by Itahashi et al. (2020), are needed to identify the exact causes and solutions for the surface biases identified here. Another potential area for future work is to separate stratospheric O₃ from natural sources in a set of simulations like those conducted for the O₃ Policy Assessment. This might solve the suspected issue of bias in stratospheric O₃ being allocated to long-range international emissions that may be caused by the correlation of stratospheric O₃ and long-range international impacts. While details on the spatial and temporal characteristics of biases in different O₃ components are provided here, the correlational bias attribution method employed here does not necessarily identify the specific factors that drive the biases. These results provide estimates of potential biases in USBUS background and USAUS anthropogenic O₃ that can inform more targeted future work examining the individual sources in greater detail. Additional future work could take a process-oriented approach rather than the source-oriented approach described here. A process-oriented approach would focus on how different physical and chemical processes (deposition, transport, photochemical activity, etc.) relate to biases in O₃ simulations. The role of uncertainties in O₃ deposition and in O₃ production efficiency across various chemical regimes could be examined in a more process-focused analysis. A further area for future work is to apply the data fusion bias correction method to an ensemble of USBUS background O₃ estimates from different models. This work has only used the CMAQ model. A test of the method would be to apply it to several different models to determine whether it is able to reduce the uncertainty of USBUS background O₃ estimates while also reducing bias in total O₃.

Acknowledgements

TNS and AGR received funding from the Phillips 66 Company. AGR also received funding from NASA HAQAST. Funding organizations have not dictated the topic or content of this work nor have they had any editorial role. The views expressed in this paper are those of the authors and do not necessarily represent the view or policies of the U.S. Environmental Protection Agency, the Phillips 66 Company, or NASA. We thank Benjamin Murphy and Sergey Napelenok for their comments on a draft version of the paper.

Code and data availability

The CMAQ source code is available from GitHub (<https://github.com/USEPA/CMAQ>) and Zenodo (<https://zenodo.org/doi/10.5281/zenodo.1079878>). O₃ observational data are available via the AQS website (<https://www.epa.gov/aqs>).

Author contributions

TNS: conceptualization, investigation, methodology, software, visualization, writing – original draft. CH: data curation, software, writing – review and editing. BHH: data curation, software, writing – review and editing. RM: software, writing – review and editing. KMF: data curation, software, writing – review and editing. AGR: conceptualization, methodology, resources, supervision, writing – review and editing.

Competing interests

The authors declare that they have no competing interests.

References

- 1020 Appel, K. W., Bash, J. O., Fahey, K. M., Foley, K. M., Gilliam, R. C., Hogrefe, C., Hutzell, W. T., Kang, D., Mathur, R., Murphy, B. N., Napelenok, S. L., Nolte, C. G., Pleim, J. E., Pouliot, G. A., Pye, H. O. T., Ran, L., Roselle, S. J., Sarwar, G., Schwede, D. B., Sidi, F. I., Spero, T. L., and Wong, D. C.: The Community Multiscale Air Quality (CMAQ) model versions 5.3 and 5.3.1: system updates and evaluation, *Geosci. Model Dev.*, 14, 2867-2897, 10.5194/gmd-14-2867-2021, 2021.

- 1025 Bash, J. O., Baker, K. R., and Beaver, M. R.: Evaluation of improved land use and canopy representation in BEIS v3.61 with biogenic VOC measurements in California, *Geosci. Model Dev.*, 9, 2191-2207, 10.5194/gmd-9-2191-2016, 2016.
- CAMS: Soil N emissions for 2000-present, D81.3.6.1 ([https://atmosphere.copernicus.eu/sites/default/files/2019-11/25 CAMS81 2017SC1 D81.3.6.1-201810 APPROVED Ver1.pdf](https://atmosphere.copernicus.eu/sites/default/files/2019-11/25_CAMS81_2017SC1_D81.3.6.1-201810_APPROVED_Ver1.pdf)), 2018.
- 1030 Dentener, F., Keating, T., and Akimoto, H. (Eds.): Hemispheric Transport of Air Pollution 2010, Part A: Ozone and Particulate Matter. Task Force on Hemispheric Transport of Air Pollution., *Air Pollution Studies*, No. 17 Geneva: United Nations Economic Commission for Europe., <https://doi.org/10.18356/2c908168-en>, 2010.
- Dolwick, P., Akhtar, F., Baker, K. R., Possiel, N., Simon, H., and Tonnesen, G.: Comparison of background ozone estimates over the western United States based on two separate model methodologies, *Atmospheric Environment*, 109, 282-296, <https://doi.org/10.1016/j.atmosenv.2015.01.005>, 2015.
- 1035 Fiore, A., Jacob, D. J., Liu, H., Yantosca, R. M., Fairlie, T. D., and Li, Q.: Variability in surface ozone background over the United States: Implications for air quality policy, 108, 10.1029/2003jd003855, 2003.
- Fiore, A. M., Oberman, J. T., Lin, M. Y., Zhang, L., Clifton, O. E., Jacob, D. J., Naik, V., Horowitz, L. W., Pinto, J. P., and Milly, G. P.: Estimating North American background ozone in U.S. surface air with two independent global models: Variability, uncertainties, and recommendations, *Atmospheric Environment*, 96, 284-300, <https://doi.org/10.1016/j.atmosenv.2014.07.045>, 2014.
- 1040 Foley, K., Pouliot, G., Eyth, A., Possiel, N., Aldridge, M., Allen, C., Appel, W., Bash, J., Beardsley, M., Beidler, J., Choi, D., Eder, B., Farkas, C., Gilliam, R., Godfrey, J., Henderson, B., Hogrefe, C., Koplitz, S., Mason, R., Mathur, R., Misenis, C., Pye, H., Reynolds, L., Roark, M., Roberts, S., Schwede, D., Seltzer, K., Sonntag, D., Talgo, K., Toro, C., and Vukovich, J.: EQUATES: EPA's Air QUALity TimE Series Project, 19th Annual CMAS Conference 2020.
- 1045 Foley, K. M., Pouliot, G. A., Eyth, A., Aldridge, M. F., Allen, C., Appel, K. W., Bash, J. O., Beardsley, M., Beidler, J., Choi, D., Farkas, C., Gilliam, R. C., Godfrey, J., Henderson, B. H., Hogrefe, C., Koplitz, S. N., Mason, R., Mathur, R., Misenis, C., Possiel, N., Pye, H. O. T., Reynolds, L., Roark, M., Roberts, S., Schwede, D. B., Seltzer, K. M., Sonntag, D., Talgo, K., Toro, C., Vukovich, J., Xing, J., and Adams, E.: 2002–2017 anthropogenic emissions data for air quality modeling over the United States, *Data in Brief*, 109022, <https://doi.org/10.1016/j.dib.2023.109022>, 2023.
- 1050 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471-1492, 10.5194/gmd-5-1471-2012, 2012.

- Guo, J. J., Fiore, A. M., Murray, L. T., Jaffe, D. A., Schnell, J. L., Moore, C. T., and Milly, G. P.: Average versus high surface ozone levels over the continental USA: model bias, background influences, and interannual variability, *Atmos. Chem. Phys.*, 18, 12123-12140, 10.5194/acp-18-12123-2018, 2018.
- 1055 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J. I., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geosci. Model Dev.*, 11, 369-408, 10.5194/gmd-11-369-2018, 2018.
- 1060 Hogrefe, C., Liu, P., Pouliot, G., Mathur, R., Roselle, S., Flemming, J., Lin, M., and Park, R. J.: Impacts of different characterizations of large-scale background on simulated regional-scale ozone over the continental United States, *Atmos. Chem. Phys.*, 18, 3839-3864, 10.5194/acp-18-3839-2018, 2018.
- Hosseinpour, F., Kumar, N., Tran, T., and Knipping, E.: Using machine learning to improve the estimate of U.S. background ozone, *Atmospheric Environment*, 316, 120145, <https://doi.org/10.1016/j.atmosenv.2023.120145>, 2024.
- 1065 Huang, M., Bowman, K. W., Carmichael, G. R., Lee, M., Chai, T., Spak, S. N., Henze, D. K., Darmenov, A. S., and da Silva, A. M.: Improved western U.S. background ozone estimates via constraining nonlocal and local source contributions using Aura TES and OMI observations, 120, 3572-3592, 10.1002/2014jd022993, 2015.
- Itahashi, S., Mathur, R., Hogrefe, C., and Zhang, Y.: Modeling stratospheric intrusion and trans-Pacific transport on tropospheric ozone using hemispheric CMAQ during April 2010 – Part 1: Model evaluation and air mass characterization for stratosphere–troposphere transport, *Atmos. Chem. Phys.*, 20, 3373-3396, 10.5194/acp-20-3373-2020, 2020.
- 1070 Jaffe, D. A., Wigder, N., Downey, N., Pfister, G., Boynard, A., and Reid, S. B.: Impact of Wildfires on Ozone Exceptional Events in the Western U.S, *Environmental Science & Technology*, 47, 11065-11072, 10.1021/es402164f, 2013.
- Jaffe, D. A., Cooper, O. R., Fiore, A. M., Henderson, B. H., Tonnesen, G. S., Russell, A. G., Henze, D. K., Langford, A. O., Lin, M. Y., and Moore, T.: Scientific assessment of background ozone over the US: Implications for air quality management, *Elementa-Sci. Anthropol.*, 6, 30, 10.1525/elementa.309, 2018.
- 1075 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G., Keating, T., Zhang, Q., Kurokawa, J., Wankmüller, R., Denier van der Gon, H., Kuenen, J. J. P., Klimont, Z., Frost, G., Darras, S., Koffi, B., and Li, M.: HTAP_v2.2: a mosaic of regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of air pollution, *Atmos. Chem. Phys.*, 15, 11411-11432, 10.5194/acp-15-11411-2015, 2015.

- Kang, D., Pickering, K. E., Allen, D. J., Foley, K. M., Wong, D. C., Mathur, R., and Roselle, S. J.: Simulating lightning NO production in CMAQv5.2: evolution of scientific updates, *Geosci. Model Dev.*, 12, 3071-3083, 10.5194/gmd-12-3071-2019, 2019.
- 1080 Langford, A. O., Senff, C. J., Alvarez, R. J., Brioude, J., Cooper, O. R., Holloway, J. S., Lin, M. Y., Marchbanks, R. D., Pierce, R. B., Sandberg, S. P., Weickmann, A. M., and Williams, E. J.: An overview of the 2013 Las Vegas Ozone Study (LVOS): Impact of stratospheric intrusions and long-range transport on surface air quality, *Atmospheric Environment*, 109, 305-322, <https://doi.org/10.1016/j.atmosenv.2014.08.040>, 2015.
- Langford, A. O., Senff, C. J., Alvarez II, R. J., Aikin, K. C., Ahmadov, R., Angevine, W. M., Baidar, S., Brewer, W. A., Brown, S. S., James, E. P., McCarty, B. J., Sandberg, S. P., and Zucker, M. L.: Were Wildfires Responsible for the Unusually High Surface Ozone in Colorado During 2021?, *Journal of Geophysical Research: Atmospheres*, 128, e2022JD037700, <https://doi.org/10.1029/2022JD037700>, 2023.
- 1085
- Lin, M., Horowitz, L. W., Payton, R., Fiore, A. M., and Tonnesen, G.: US surface ozone trends and extremes from 1980 to 2014: quantifying the roles of rising Asian emissions, domestic controls, wildfires, and climate, *Atmos. Chem. Phys.*, 17, 2943-2970, 10.5194/acp-17-2943-2017, 2017.
- 1090 Lin, M., Fiore, A. M., Horowitz, L. W., Langford, A. O., Oltmans, S. J., Tarasick, D., and Rieder, H. E.: Climate variability modulates western US ozone air quality in spring via deep stratospheric intrusions, *Nature Communications*, 6, 7105, 10.1038/ncomms8105, 2015.
- Lin, M., Fiore, A. M., Cooper, O. R., Horowitz, L. W., Langford, A. O., Levy II, H., Johnson, B. J., Naik, V., Oltmans, S. J., and Senff, C. J.: Springtime high surface ozone events over the western United States: Quantifying the role of stratospheric intrusions, 117, 10.1029/2012jd018151, 2012a.
- 1095 Lin, M., Fiore, A. M., Horowitz, L. W., Cooper, O. R., Naik, V., Holloway, J., Johnson, B. J., Middlebrook, A. M., Oltmans, S. J., Pollack, I. B., Ryerson, T. B., Warner, J. X., Wiedinmyer, C., Wilson, J., and Wyman, B.: Transport of Asian ozone pollution into surface air over the western United States in spring, 117, 10.1029/2011jd016961, 2012b.
- Lin, M., Horowitz, L. W., Zhao, M., Harris, L., Ginoux, P., Dunne, J., Malyshev, S., Shevliakova, E., Ahsan, H., Garner, S., Paulot, F., Pouyaei, A., Smith, S. J., Xie, Y., Zadeh, N., and Zhou, L.: The GFDL Variable-Resolution Global Chemistry-Climate Model for Research at the Nexus of US Climate and Air Quality Extremes, *Journal of Advances in Modeling Earth Systems*, 16, e2023MS003984, <https://doi.org/10.1029/2023MS003984>, 2024.
- 1100 Liu, S. C., Trainer, M., Fehsenfeld, F. C., Parrish, D. D., Williams, E. J., Fahey, D. W., Hübler, G., and Murphy, P. C.: Ozone production in the rural troposphere and the implications for regional and global ozone distributions, *Journal of Geophysical Research: Atmospheres*, 92, 4191-4207, <https://doi.org/10.1029/JD092iD04p04191>, 1987.

- 1105 Mathur, R., Kang, D., Napelenok, S. L., Xing, J., Hogrefe, C., Sarwar, G., Itahashi, S., and Henderson, B. H.: How Have Divergent Global Emission Trends Influenced Long-Range Transported Ozone to North America?, *Journal of Geophysical Research: Atmospheres*, 127, e2022JD036926, <https://doi.org/10.1029/2022JD036926>, 2022.
- Mathur, R., Xing, J., Gilliam, R., Sarwar, G., Hogrefe, C., Pleim, J., Pouliot, G., Roselle, S., Spero, T. L., Wong, D. C., and Young, J.: Extending the Community Multiscale Air Quality (CMAQ) Modeling System to Hemispheric Scales: Overview of Process Considerations and Initial Applications, *Atmos Chem Phys*, 17, 12449-12474, 10.5194/acp-17-12449-2017, 2017.
- 1110 McDonald-Buller, E. C., Allen, D. T., Brown, N., Jacob, D. J., Jaffe, D., Kolb, C. E., Lefohn, A. S., Oltmans, S., Parrish, D. D., Yarwood, G., and Zhang, L.: Establishing Policy Relevant Background (PRB) Ozone Concentrations in the United States, *Environmental Science & Technology*, 45, 9484-9497, 10.1021/es2022818, 2011.
- Murphy, B. N., Woody, M. C., Jimenez, J. L., Carlton, A. M. G., Hayes, P. L., Liu, S., Ng, N. L., Russell, L. M., Setyan, A., Xu, L., Young, J., Zaveri, R. A., Zhang, Q., and Pye, H. O. T.: Semivolatile POA and parameterized total combustion SOA in CMAQv5.2: impacts on source strength and partitioning, *Atmos. Chem. Phys.*, 17, 11107-11133, 10.5194/acp-17-11107-2017, 2017.
- 1115 Price, C., Penner, J., and Prather, M.: NO_x from lightning: 1. Global distribution based on lightning physics, *Journal of Geophysical Research: Atmospheres*, 102, 5929-5941, <https://doi.org/10.1029/96JD03504>, 1997.
- Pye, H. O. T., D'Ambro, E. L., Lee, B. H., Schobesberger, S., Takeuchi, M., Zhao, Y., Lopez-Hilfiker, F., Liu, J., Shilling, J. E., Xing, J., Mathur, R., Middlebrook, A. M., Liao, J., Welti, A., Graus, M., Warneke, C., de Gouw, J. A., Holloway, J. S., Ryerson, T. B., Pollack, I. B., and Thornton, J. A.: Anthropogenic enhancements to production of highly oxygenated molecules from autoxidation, *Proceedings of the National Academy of Sciences*, 116, 6641-6646, doi:10.1073/pnas.1810774116, 2019.
- 1120 Qin, M., Murphy, B. N., Isaacs, K. K., McDonald, B. C., Lu, Q., McKeen, S. A., Koval, L., Robinson, A. L., Efstathiou, C., Allen, C., and Pye, H. O. T.: Criteria pollutant impacts of volatile chemical products informed by near-field modelling, *Nature Sustainability*, 4, 129-137, 10.1038/s41893-020-00614-1, 2021.
- 1125 Rickly, P. S., Coggon, M. M., Aikin, K. C., Alvarez, R. J., II, Baidar, S., Gilman, J. B., Gkatzelis, G. I., Harkins, C., He, J., Lamplugh, A., Langford, A. O., McDonald, B. C., Peischl, J., Robinson, M. A., Rollins, A. W., Schwantes, R. H., Senff, C. J., Warneke, C., and Brown, S. S.: Influence of Wildfire on Urban Ozone: An Observationally Constrained Box Modeling Study at a Site in the Colorado Front Range, *Environmental Science & Technology*, 57, 1257-1267, 10.1021/acs.est.2c06157, 2023.
- 1130 Sarwar, G., Hogrefe, C., Henderson, B. H., Mathur, R., Gilliam, R., Callaghan, A. B., Lee, J., and Carpenter, L. J.: Impact of particulate nitrate photolysis on air quality over the Northern Hemisphere, *Science of The Total Environment*, 917, 170406, <https://doi.org/10.1016/j.scitotenv.2024.170406>, 2024.

- 1135 Sarwar, G., Gantt, B., Foley, K., Fahey, K., Spero, T. L., Kang, D., Mathur, R., Foroutan, H., Xing, J., Sherwen, T., and Saiz-Lopez, A.: Influence of bromine and iodine chemistry on annual, seasonal, diurnal, and background ozone: CMAQ simulations over the Northern Hemisphere, *Atmospheric Environment*, 213, 395-404, <https://doi.org/10.1016/j.atmosenv.2019.06.020>, 2019.
- 1140 Schwantes, R. H., Lacey, F. G., Tilmes, S., Emmons, L. K., Lauritzen, P. H., Walters, S., Callaghan, P., Zarzycki, C. M., Barth, M. C., Jo, D. S., Bacmeister, J. T., Neale, R. B., Vitt, F., Kluzek, E., Roozitalab, B., Hall, S. R., Ullmann, K., Warneke, C., Peischl, J., Pollack, I. B., Flocke, F., Wolfe, G. M., Hanisco, T. F., Keutsch, F. N., Kaiser, J., Bui, T. P. V., Jimenez, J. L., Campuzano-Jost, P., Apel, E. C., Hornbrook, R. S., Hills, A. J., Yuan, B., and Wisthaler, A.: Evaluating the Impact of Chemical Complexity and Horizontal Resolution on Tropospheric Ozone Over the Conterminous US With a Global Variable Resolution Chemistry Model, *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002889, <https://doi.org/10.1029/2021MS002889>, 2022.
- 1145 Shah, V., Jacob, D. J., Dang, R., Lamsal, L. N., Strode, S. A., Steenrod, S. D., Boersma, K. F., Eastham, S. D., Fritz, T. M., Thompson, C., Peischl, J., Bourgeois, I., Pollack, I. B., Nault, B. A., Cohen, R. C., Campuzano-Jost, P., Jimenez, J. L., Andersen, S. T., Carpenter, L. J., Sherwen, T., and Evans, M. J.: Nitrogen oxides in the free troposphere: implications for tropospheric oxidants and the interpretation of satellite NO₂ measurements, *Atmos. Chem. Phys.*, 23, 1227-1257, 10.5194/acp-23-1227-2023, 2023.
- Sindelarova, K., Granier, C., Bouarar, I., Guenther, A., Tilmes, S., Stavrou, T., Müller, J. F., Kuhn, U., Stefani, P., and Knorr, W.: Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years, *Atmos. Chem. Phys.*, 14, 9317-9341, 10.5194/acp-14-9317-2014, 2014.
- 1150 Skipper, T. N., Hu, Y., Odman, M. T., Henderson, B. H., Hogrefe, C., Mathur, R., and Russell, A. G.: Estimating US Background Ozone Using Data Fusion, *Environmental Science & Technology*, 55, 4504-4512, 10.1021/acs.est.0c08625, 2021.
- US EPA: Integrated Science Assessment (ISA) of Ozone and Related Photochemical Oxidants (Final Report, Feb 2013). U.S. Environmental Protection Agency, Washington, DC, EPA/600/R-10/076F, 2013.
- US EPA: Policy Assessment for the Review of the Ozone National Ambient Air Quality Standards. U.S. Environmental Protection Agency, Washington, DC, EPA-452/R-14/006., 2014.
- 1155 US EPA: Technical Support Document (TSD) Preparation of Emissions Inventories for the Version 7.1 2016 Hemispheric Emissions Modeling Platform, 2019a.
- US EPA: Technical Support Document (TSD) Preparation of Emissions Inventories for the Version 7.1 2016 North American Emissions Modeling Platform, 2019b.

- 1160 US EPA: Policy Assessment for the Review of the Ozone National Ambient Air Quality Standards. U.S. Environmental Protection Agency, Washington, DC, EPA-452/R-20-001, 2020a.
- US EPA: Integrated Science Assessment (ISA) for Ozone and Related Photochemical Oxidants (Final Report). U.S. Environmental Protection Agency. Washington, DC. EPA/600/R-20/012, 2020b.
- Wang, H., Jacob, D. J., Le Sager, P., Streets, D. G., Park, R. J., Gilliland, A. B., and van Donkelaar, A.: Surface ozone background in the United States: Canadian and Mexican pollution influences, *Atmospheric Environment*, 43, 1310-1319, <https://doi.org/10.1016/j.atmosenv.2008.11.036>, 2009.
- 1165 <https://doi.org/10.1016/j.atmosenv.2008.11.036>, 2009.
- Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model Dev.*, 4, 625-641, 10.5194/gmd-4-625-2011, 2011.
- Wilkins, J. L., Pouliot, G., Pierce, T., Soja, A., Choi, H., Gargulinski, E., Gilliam, R., Vukovich, J., and Landis, M. S.: An evaluation of empirical and statistically based smoke plume injection height parametrisations used within air quality models, *International Journal of Wildland Fire*, 31, 193-211, <https://doi.org/10.1071/WF20140>, 2022.
- 1170 <https://doi.org/10.1071/WF20140>, 2022.
- Wu, S., Duncan, B. N., Jacob, D. J., Fiore, A. M., and Wild, O.: Chemical nonlinearities in relating intercontinental ozone pollution to anthropogenic emissions, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2008GL036607>, 2009.
- Xing, J., Mathur, R., Pleim, J., Hogrefe, C., Wang, J., Gan, C. M., Sarwar, G., Wong, D. C., and McKeen, S.: Representing the effects of stratosphere–troposphere exchange on 3-D O₃ distributions in chemistry transport models using a potential vorticity-based parameterization, *Atmos. Chem. Phys.*, 16, 10865-10877, 10.5194/acp-16-10865-2016, 2016.
- 1175 <https://doi.org/10.1029/95JD00370>, 1995.
- Yienger, J. J. and Levy, H.: Empirical model of global soil-biogenic NO_x emissions, *Journal of Geophysical Research: Atmospheres*, 100, 11447-11464, <https://doi.org/10.1029/95JD00370>, 1995.
- Zhang, L., Jacob, D. J., Yue, X., Downey, N. V., Wood, D. A., and Blewitt, D.: Sources contributing to background surface ozone in the US Intermountain West, *Atmos. Chem. Phys.*, 14, 5295-5309, 10.5194/acp-14-5295-2014, 2014.
- 1180 <https://doi.org/10.1016/j.atmosenv.2011.07.054>, 2011.
- Zhang, L., Jacob, D. J., Downey, N. V., Wood, D. A., Blewitt, D., Carouge, C. C., van Donkelaar, A., Jones, D. B. A., Murray, L. T., and Wang, Y. X.: Improved estimate of the policy-relevant background ozone in the United States using the GEOS-Chem global model with 1/2 degrees x 2/3 degrees horizontal resolution over North America, *Atmospheric Environment*, 45, 6769-6776, [10.1016/j.atmosenv.2011.07.054](https://doi.org/10.1016/j.atmosenv.2011.07.054), 2011.

- 1185 Zhao, B., Zheng, H., Wang, S., Smith, K. R., Lu, X., Aunan, K., Gu, Y., Wang, Y., Ding, D., Xing, J., Fu, X., Yang, X., Liou, K.-N., and Hao, J.: Change in household fuels dominates the decrease in PM_{2.5} exposure and premature mortality in China in 2005–2015, *Proceedings of the National Academy of Sciences*, 115, 12401-12406, [10.1073/pnas.1812955115](https://doi.org/10.1073/pnas.1812955115), 2018.