

Interactions of fertilisation and crop productivity on soil nitrogen cycle microbiome and gas emissions

Laura Kuusemets^{1*}, Ülo Mander¹, Jordi Escuer-Gatius², Alar Astover², Karin Kauer², Kaido Soosaar¹, Mikk Espenberg¹

5

¹University of Tartu, Institute of Ecology and Earth Sciences, Vanemuise 46, Tartu, 51003, Estonia

²Estonian University of Life Sciences, Institute of Agricultural and Environmental Sciences, Kreutzwaldi 5, Tartu, 51014, Estonia

10 *Correspondence to:* Laura Kuusemets (laura.kuusemets@ut.ee)

Abstract. Fertilised soils are a significant source of nitrous oxide (N₂O), a highly active greenhouse gas and stratospheric ozone depleter. Nitrogen (N) fertilisers, while boosting crop yield, also lead to N₂O emissions into the atmosphere, impacting global warming. We investigated relationships between mineral N fertilisation rates and additional manure amendment with different crop types through the analysis of abundances of N cycle functional genes, soil N₂O and N₂ emissions, nitrogen use efficiency (NUE), soil physicochemical analysis and biomass production. Our study indicates that N₂O emissions are predominantly dependent on the mineral N fertilisation rate and enhance with increased mineral N fertilisation rate. Higher N₂O emissions were attained with the application of manure in comparison to mineral fertilisation. Manure amendment also increased the number of N cycle genes that are significant in the variations of N₂O. Contrary to our hypothesis, there was no significant influence of crop type on soil N₂O emissions. The study indicated that N₂O emissions were mainly related to nitrification in the soil. Quantification of nitrogen cycle functional genes also showed the potential role of denitrification, comammox and DNRA processes as a source of N₂O. Our study did not find soil moisture to be significantly linked to N₂O emissions. Results of the study provide evidence that for wheat, a fertilisation rate of 80 kg N ha⁻¹ is closest to the optimal rate for balancing biomass yield, N₂O emissions, and achieving high NUE. Sorghum showed a good potential for cultivation in temperate climate, as it showed similar biomass yield compared to the other crop types and fertilisation rates, but maintained low N₂O emissions and N losses on mineral N fertilisation rate of 80 kg N ha⁻¹.

15
20
25

1 Introduction

The rising demand for agricultural commodities and the management of agroecosystems are important factors contributing to global environmental problems. Increasing crop yield while reducing pollution from agricultural production is crucial (Abdalla *et al.*, 2019; Tilman *et al.*, 2011). Global food demand projections suggest a 50% increase in agricultural production by 2050 (compared to 2012) to feed the fast-growing human population (FAO, 2017). Enhancing agricultural production involves actions such as expanding agricultural land, applying more fertilisers, and using water resources and

30

fertilisers more effectively (Tian *et al.*, 2021). In today's agricultural practises, the applied N with fertilisation is often excessive for plant needs (Robertson and Vitousek *et al.*, 2009; Zhou *et al.*, 2016). About half of the applied N to the fields is not taken up by crops (Coskun *et al.*, 2017); which may lead to N loss in the surrounding environment. Main soil N loss mechanisms include denitrification, ammonia oxidation, N leaching, erosion of soil and ammonia (NH₃) volatilisation (Thomson *et al.*, 2012). This results in adverse ecological impacts, such as eutrophication of aquatic ecosystems and increased gaseous emissions of N into the atmosphere (Cameron *et al.*, 2013; Liu *et al.*, 2017; Whetton *et al.*, 2022). Fertilised soils are a significant source of nitrous oxide (N₂O), contributing to the greenhouse effect and ozone depletion (Ravishankara *et al.*, 2009; Shcherbak *et al.*, 2014). N₂O has 273 times higher global warming potential than carbon dioxide (CO₂) over a 100-year timescale (IPCC, 2021). Even without adding N fertiliser in the current season or year, background N₂O emissions (BNEs) may still occur. BNEs are caused by different N sources, including residual N in the soil from previous years' N application, deposition from the atmosphere, biological N₂ fixation and mineralised N from plant residues (Gu *et al.*, 2007; Kim *et al.*, 2013, Abdalla *et al.*, 2022).

The key microbial processes leading to soil N loss are nitrification and denitrification (Thomson *et al.*, 2012). In agriculture, N fertilisers added to the soil can be lost due to these processes (Saud *et al.*, 2022). Nitrification was traditionally viewed as a two-step process carried out by separate functional groups of microorganisms, oxidising ammonium (NH₄⁺) sequentially to nitrite (NO₂⁻) and nitrate (NO₃⁻) under aerobic conditions (Kuypers *et al.*, 2018; Koch *et al.*, 2019; Nardi *et al.*, 2020). However, in 2015, a significant advancement in our understanding of nitrification occurred with the discovery that a single microorganism, through the comammox (complete ammonia oxidation) process, can perform both nitrification steps (Daims *et al.*, 2015; Van Kessel *et al.*, 2015). Nitrification can reduce N availability for plant uptake by up to 50%, primarily due to NO₃⁻ leaching and N₂O emissions (Beeckman *et al.*, 2018). Synthetic fertilizers containing NH₃ offer an immediate substrate for ammonia oxidizers, thus accelerating the nitrification process (Ayiti & Babalola, 2022). Also, fertilizers that raise soil pH can significantly enhance the nitrification rate, as increasing soil pH from 4.8 to 6.7 can boost nitrification rates by 30 times (DeForest & Otuya, 2020).

Denitrification is a microbially-catalysed process under oxygen-limited condition responsible for transforming NO₃⁻ sequentially to gaseous forms of N: nitric oxide, N₂O and atmospheric N (Philippot *et al.*, 2007; Zaman *et al.*, 2012). The input of N fertilisers affects the soil's mineral N pool by providing larger amounts of available N for nitrification and denitrification processes, contributing to N₂O emissions (Engel *et al.*, 2010). Dissimilatory nitrate reduction to ammonium (DNRA) supplies NH₄⁺ to the soil, conserves bioavailable N and prevents the leaching of NO₃⁻ (Bai *et al.*, 2020; Pandey *et al.*, 2020). DNRA competes with denitrification in NO₃⁻-reducing processes (Putz *et al.*, 2018). Similarly to denitrification and nitrification processes, DNRA can also be a source of N₂O, although the quantities are modest (Rütting *et al.*, 2011; Stremińska *et al.*, 2012; Zaman *et al.*, 2012). Carbon to nitrogen ratio (C/N) and C/NO₃⁻ are recognised as the main environmental factors controlling, which nitrate-reducing process is favoured as DNRA and denitrifying microbes compete for NO₃⁻ and carbon sources (Bai *et al.*, 2020). DNRA is dominant in the presence of a high C/N ratio and low NO₃⁻ availability, while the denitrification process favours a low ratio of C/N and C/NO₃⁻ (Bai *et al.*, 2020; Pandey *et al.*, 2020).

These processes are mediated by different functional marker genes, including archaeal, bacterial and comammox *amoA* genes for nitrification, *nrfA* genes for DNRA and *nosZ* clade I and II, *nirK*, *nirS* genes for denitrification (Zaman *et al.*, 2012; Hu *et al.*, 2015; Zhang *et al.*, 2021).

70 C3 photosynthesis, a dominant pathway among plants and found in wheat and barley, uses the Calvin-Benson pathway, while an alternative, the Hatch-Slack pathway, is used by C4 plants like sorghum and maize (Hibberd and Quick, 2002; Ehleringer and Cerling, 2002; Ehleringer, 1979; Ledvinka, 2022). In C3 plants, water loss through transpiration during CO₂ uptake is a risk in hot and water-limited conditions (Joshi *et al.*, 2022; Stevens *et al.*, 2022). However, C4 plants, with higher water use efficiency and greater tolerance to hot and dry environments, make the cultivation of sorghum and other drought-
75 tolerant plants likely to expand in regions affected by droughts (Anderson *et al.*, 2020). Due to climate change, sorghum, as a resilient plant, is considered a novel crop for temperate Europe (Schaffasz *et al.*, 2019). Only a limited number of studies have compared N₂O emissions between different crop species. Abdalla *et al.* (2022) found that crop type has significant effect ($p < 0.05$) on the BNE values from soil. Furthermore, Bouwman *et al.* (2002) also found that crop type has a significant influence on N₂O emissions. However, study including 372 sites showed that cover crops did not have significant ($p > 0.05$)
80 effect on direct N₂O emissions (Abdalla *et al.*, 2019).

Previous studies on long-term fertilisation experiments have mostly focused on fertilisation's yield effects and changes in soil organic matter (Cvetkov and Tajnšek, *et al.*, 2009; Hijbeek *et al.*, 2017; Káš *et al.*, 2010; Spiegel *et al.*, 2010; Tajnšek *et al.*, 2013). Improved management of arable soils holds significant potential for mitigating greenhouse gas emissions, as agroecosystems contribute ca 66% of total anthropogenic N₂O emissions (Davidson and Kanter, 2014; Paustian *et al.*, 2016; Shen *et al.*, 2021). Efficient mitigation of N loss requires a comprehensive understanding of microbial processes related to N₂O emissions in agricultural soils (Davidson and Kanter, 2014; Shen *et al.*, 2021).

The general objectives of the study were to evaluate temporal patterns of gaseous N loss, link N-cycle processes with abundances of functional N cycle genes in arable soil, and evaluate the performance of different crops (including novel crop in Northern Europe) in terms of biomass production and N₂O emissions under mineral and organic fertilisation. The
90 following hypotheses were tested: (1) crop type significantly affects N₂O emissions; (2) nitrification is the primary pathway of soil N₂O production due to aerobic conditions; (3) in arable soil, low soil moisture results in reduced N₂O losses; (4) amendment of manure fertiliser increases soil N₂O emissions and affects the abundances of functional N cycle genes.

2 Material and methods

2.1 Field experiment description

95 The field study was conducted on the International Organic Nitrogen Long-term Fertilisation Experiment (IOSDV; Internationaler Organischer Stickstoff Dauerdüngungs Versuch) experimental field. The experimental site is located near Tartu, southern Estonia, Northern Europe (58°22'30" N, 26°39'48" E). The experiment was set up as a three-field crop

rotation experiment in 1989 to investigate the long-term effects of mineral and organic fertilisation on crop responses of various crops and soil properties.

100 In 2022, the average temperature in the area was -2.0 °C in winter, 4.6 °C in spring, 18.1 °C in summer and 7.2 °C in autumn. The mean annual precipitation was 531 mm (Republic of Estonia Environment Agency, 2023) in 2022. A climate diagram for the area during the study period is in Figure S1 in the Supplementary materials.

The soil type is *Stagnic Luvisol* combined with *Fragic Glossic Retisol* (IUSS WG WRB 2015). The thickness of the humus layer is $27-32$ cm. Soil texture by FAO classification is sandy loam: 57.86% sand (>0.063 mm), 33.58% silt ($0.063-0.002$ mm) and 8.55% clay (<0.002 mm). Soil bulk density was in range of 1.5 to 1.6 g cm⁻³ with slightly lower values for manure treatment plots. The average pH levels in spring 2022 were 5.4 for barley plots, 5.3 for wheat plots, 5.6 for sorghum plots without manure amendment, and 6.2 for sorghum plots with manure amendment.

The experiment was organised into 12 plots in a systematic block design (Figure 1) with three sampling spots per plot. Every plot was 50 m² in size. The crop species studied were spring barley (cultivar “Elmeri”), sorghum (*Sorghum bicolor* x *Sorghum sudanense*, cultivar “SUSU”), and spring wheat (cultivar “Mistral”). Initially, the crop rotation was potato–spring wheat–spring barley (Astover *et al.*, 2016). In 2019, potato was replaced with sorghum-sudangras hybrid.

The fertiliser treatment consisted of mineral N fertilisation and mineral fertilisation with farmyard manure amendment. All fertilisation treatments are applied continuously from year 1989, when the experimental site was established. Three mineral N fertiliser treatment rates were studied: 0 , 80 and 160 kg N ha⁻¹. The farmyard manure rate added to the sorghum plots was 40 t ha⁻¹ of manure (231.2 kg N ha⁻¹). The mineral fertiliser applied was ammonium nitrate (NH₄NO₃) and organic fertiliser was farmyard manure. The farmyard manure was cattle dung with straw bedding, freely fermented before use 6-8 months in heap. The chemical properties (C, N, P, K) of manure added in 2022 and during the last ten years are presented in Table S1 in the Supplementary materials. The farmyard manure treatment with mineral fertilisation was applied only to sorghum. Manure treatment is amended with solid farmyard manure (40 t ha⁻¹) in every third year before sorghum/potato. The main management activities and timing in the field are displayed in Table S2 in Supplementary materials.




Figure 1: Satellite view of the study site with study plots (from Maa-amet). Each plot constituted of three sampling spots indicated as white dots. N0 – 0 kg N ha⁻¹, N80 – 80 kg N ha⁻¹, N160 – 160 kg N ha⁻¹ as mineral fertiliser.

125

2.2 Gas sampling for N₂O flux analyses

The field study was conducted during the growing season from April 2022 to October 2022. Sampling took place on 15 different dates, starting on April 27th and ending on October 12th (every week until the end of June and then twice a month until the end of September). Gas samples for N₂O flux analysis were collected on all fifteen fieldwork days. N₂O gas sampling was carried out using the static chamber method (Hutchinson and Livingston, 1993). Polyvinyl chloride chambers (Ø 50 cm, volume 65 L) were placed on top of the collars during the gas sampling. Chamber extensions were used for some treatments of sorghum on four occasions as the chambers alone were too small to accommodate the growing crops. Pre-vacuumed 50 ml glass vials were used for gas sampling. Gas samples were collected at 20 minutes intervals for one hour (0, 20, 40, 60 min). The concentration of N₂O in the collected air was measured in the Biogeochemical Cycling Research Laboratory in the Department of Geography, University of Tartu, with the gas chromatograph Shimadzu GC-2014 (Kyoto, Japan), equipped with electron capture and flame ionisation detectors (Poole, 2015).

130

135

2.3. Soil sampling and physicochemical analyses

Soils were sampled for chemical and microbiological analyses six times (April 27th, May 9th, June 2nd, July 7th, September 2nd, October 12th). Soil sampling was conducted after gas sampling. Soil samples were collected close to collars with a soil probe from the top 10 cm of the soil. Three auger samples from each point (both bulk and rhizosphere soil were sampled)

140

were collected for one composite sample for chemical and microbiological analyses. All in all, 216 samples were collected for chemical analyses and 144 samples for microbial analyses. Until chemical and microbiological analyses, samples were stored at +4 °C and -20 °C, respectively. In addition to soil sampling, soil temperature (°C) at a 10 cm depth was measured with a temperature logger (Comet Systems Ltd., Rožnov pod Radhoštěm, Czech Republic) and soil moisture (m³/m³) was recorded using water content reflectometers (model CS615, Campbell Scientific Inc., Logan, UT, USA). The soil samples were analysed for total carbon (C_{tot}), total nitrogen (N_{tot}), nitrate-nitrogen (NO₃⁻-N), and ammonium-nitrogen (NH₄⁺-N) concentrations in the Soil Science and Agrochemistry Laboratory of Estonian University of Life Sciences. N_{tot} and C_{tot} analyses were done by Dumas method (International Organization for Standardization, 1998) with dry combustion on a VarioMAX CNS elemental analyser (ELEMENTAR, Elementar Analysensysteme GmbH, Langenselbold, Germany). NO₃⁻-N analyses were done according to EPA (United States Environmental Protection Agency) method 9056: determination of inorganic anions by ion chromatography. NH₄⁺-N analyses were done according to Thermo Fisher Application Note 141 (AU204: Determination of Inorganic Cations and Ammonium in Environmental Waters Using a Compact Ion Chromatography System) using ion chromatography. Soil pH was measured using a glass-electrode pH meter in a water solution of 1:2.5. Total phosphorus (P) and potassium (K) concentration in manure were determined through acid digestion using a sulfuric acid solution (van Reeuwijk, 2002).

The hot-water extractable C (HWEOC) represents the readily mineralising C fraction and was determined on dry soil samples by a modified method of Haynes and Francis (1993) in two steps. In the first step the soil was shaken with deionized water at room temperature for 1 h. After that the soil suspension put into the thermostat at 80 °C for 16 h. The mixture was centrifuged for 10 min at 8000 rpm and filtered through a 0.45-µm membrane filter (25-mm diameter, nylon, Agilent®). The HWEOC concentration was determined from the extracts by the VarioMaX CNS analyzer (ELEMENTAR, Elementar Analysensysteme GmbH, Langenselbold, Germany).

2.4 Total biomass

The total biomass (above- and below-ground) was measured at the maturity phase on the harvest day of each crop (Supplementary Table S2). The above-ground biomass was cut from the ground level in a 0.2 m² area near each collar. The belowground biomass samples were taken with a soil auger (Ø 34 cm). Frasier *et al.* (2016) provides a more detailed description of the method used for below-ground biomass measurement. The sampling depth extended to the plowing depth, where most of the roots are found, up to a depth of 18 cm. Samples were stored at +4 °C until the roots were washed on a sieve (mesh size 0.5 mm).

Dry matter yield was determined after drying the biomass (including roots) at 70 °C to constant weight. The straw and grains were separated before weighing as air dry. The biomass (straw, grain, roots) were milled and the N_{tot} content was determined by the Dumas method with dry combustion on a VarioMAX CNS elemental analyser (ELEMENTAR, Elementar Analysensysteme GmbH, Langenselbold, Germany).

175 2.5 Soil microbial analyses

2.5.1 DNA extraction

DNA was extracted from 0.25 g of soil samples using the DNeasy® PowerSoil® Pro Kit (Qiagen, Hilden, Germany) following the manufacturer's instructions. The difference from the instruction was the homogenisation of samples with homogeniser, Precellys 24 (Bertin Technologies, Montaigne-le-Bretonneux, France), for 20 s at the rate of 5000 rpm. The concentration and quality of the extracted DNA were evaluated with an Infinite 200 M spectrophotometer (Tecan AG, Männedorf, Switzerland). The extracted DNA was stored in a freezer at -20°C .

2.5.2 Quantification of gene copies using qPCR

Quantification of the 16S rRNA genes of bacteria and archaea, along with the quantification of nitrification (bacterial, archaeal, and comammox *amoA*), denitrification (*nirS*, *nirK*, *nosZI*, and *nosZII*) and DNRA (*nrfA*) genes was done using quantitative polymerase chain reaction (qPCR). qPCR reactions were performed by The Rotor-Gene Q thermocycler (Qiagen). The reaction mixture of 10 μL consisted of extracted DNA (1 μL), gene-specific forward and reverse primers, Maxima SYBR Green Master mix reagent (5 μL ; Thermo Fisher Scientific, Waltham, MA, USA) and distilled water. Each sample was amplified two times. All of the qPCR assays included two DNA-free negative control samples. Details on thermal cycling conditions and used primers are added in Table S3 in Supplementary Materials. The Rotor-Gene® Q software v. 2.0.2 (Qiagen) and LinRegPCR v. 2020.2. were used to assess the qPCR results. The amount of gene copies was calculated using standard curve ranges, and results were presented in gene copies per gram of dry matter (copies/g dw). Espenberg *et al.* (2018) provides a more detailed description of the used qPCR methodology.

195 2.6 Statistical analyses and modelling

Statistical software programs Statistica (v. 7.1) and R (v. 4.0.4) were used for statistical analyses and visualising the data. Principal component analysis (PCA) were conducted on soil physicochemical parameters and microbiological data (abundance of functional marker genes) with the “FactoMineR” (Lê *et al.*, 2008) and “factoextra” (Kassambara *et al.*, 2020) packages in the software R. Analysis of variance (ANOVA) with post-hoc Tukey HSD test was used (cumulative N_2O emission values are meeting the assumptions of parametric test) to find statistically significant differences between different fertilisation rates, use of manure and crop types.

Spearman's rank correlation coefficient measured the association between N_2O and N_2 emissions, gene abundances and environmental factors. Random Forest classification analysis was conducted using Boruta v. 8.0 (Kursa and Rudnicki, 2010) to identify the gene parameters that best predicted N_2O fluxes.

205 Nitrogen use efficiency (NUE, kg DM kg⁻¹ N⁻¹) was calculated as the biomass production per unit of N applied (Pandey *et al.*, 2001; Supplementary Methodology S1). The N₂ emissions were estimated from the measured N₂O emissions using the N₂:N₂O ratio, which was calculated as proposed in the DAYCENT model (Parton *et al.*, 2001), with the equations described in Del Grosso *et al.* (2000) (Supplementary Methodology S2), where the ratio N₂:N₂O is a function of the content of NO₃⁻ in the soil, CO₂ emissions, and water-filled pore space (WFPS). The change of soil N content (kg N ha⁻¹) was calculated according to Sainju (2017), as the difference between the initial and final soil total N contents (Supplementary Methodology S3). N losses are calculated by subtracting N outputs and change of soil N content from N inputs (Sainju, 2017; Escuer-Gatius *et al.*, 2022; Supplementary Methodology S4).


Linear mixed-effects model (LMM) was used to investigate differences in N₂O emissions, cumulative N₂ emissions and gene parameters between different crop types and fertilisation rates using the R package “nlme.” For N₂O emissions and gene parameters, spatial (different fertilisation rate) and temporal (sampling dates) effects were used as random effects. For N₂ emissions, spatial effect (different fertilisation rate) was used as a random effect. The Kruskal–Wallis test and the post-hoc Tukey HSD test were used to compare the N₂O, cumulative N₂ emission and gene parameter (not meeting the assumptions of parametric test) values between different crop types and fertilisation rates. Due to limited number of observations in the case of total dry weight biomass, N content in biomass, LMM was not possible to apply for statistical differences between different fertilisation rates and crop types.

3 Results

3.1 Soil physicochemical characteristics and biomass production

The NH₄⁺-N content in soil decreased on most of the plots at the beginning of the study period, while NO₃⁻-N content was increasing (Supplementary Figure S2). Fertilised plots had higher soil N_{tot}, C_{tot}, NO₃⁻-N and NH₄⁺-N content compared to non-fertilised plots according to the principal component analysis (PCA) (Figure 2; Supplementary Figure S2 and Figure S3). For sorghum without manure amendment plots (Figure 2C), NO₃⁻-N and NH₄⁺-N contents were more different from each other compared to sorghum with manure amendment plots, where NO₃⁻-N and NH₄⁺-N contents were relatively similar (Figure 2D). HWEOC concentrations were higher in sorghum plots with farmyard manure amendment compared to sorghum plots without manure amendment.

230 Soil moisture ranged from 0.02 m³/m³ to 0.32 m³/m³ with an average of 0.23 m³/m³ over the study period (Supplementary Figure S4). There were no significant correlations between soil moisture and N₂O emissions (Supplementary Table S4). Soil moisture was not significantly linked to gene copy numbers across all crop types, except *nirS*. A climate diagram for the area during the study period is presented in Figure S1 in the Supplementary materials.



235

Figure 2: Principal components analysis (PCA) ordination plots demonstrate the grouping of fertilisation rates according to physicochemical parameters for different crop type. N₂O is added as a supplementary variable. The month indicates the sampling time. Abbreviations: Ctot – total carbon content of soil; Ntot – total nitrogen content of soil; HWEOC – hot-water extractable organic carbon.

240

The total dry biomass of barley ranged between 2.6 to 6.4 t ha⁻¹, and wheat between 4.6 to 8.5 t ha⁻¹ depending on the mineral N fertilisation rate (Figure 3). For sorghum without manure amendment, the total dry biomass varied between 2.3 and 7.1 t ha⁻¹, and for sorghum with manure amendment, the total dry biomass varied between 8.2 and 11.7 t ha⁻¹.

245

The biomass production was higher per unit area of crop growth with higher fertiliser input (Figure 3A). Total biomass was significantly positively correlated with N_{tot} (p<0.01), C_{tot} (p<0.05) and NO₃⁻-N (p<0.001) levels in soil (Supplementary Table S5). Also, higher N fertilisation rate caused an increase in N content in the crop biomass (Figure 3B).

250




Figure 3: Total dry weight biomass (aboveground + belowground) produced per unit area according to crop types and fertilisation rates. Error bars show standard errors.

255

The highest values of nitrogen use efficiency (NUE) were obtained from wheat plots, and the lowest from sorghum plots. The average NUE of wheat plots at fertilisation rate 80 was 0.84, and at rate 160, it was 0.64. For sorghum plots with manure amendment, NUE at mineral N fertilisation rate 0 was 0.15, at rate 80 was 0.16, and at rate 160 was 0.25. For sorghum plots without manure amendment, the average NUE at fertilisation rate 80 was 0.12, and at rate 160, it was 0.25. The NUE for barley plots at fertilisation rate 80 was 0.35, and at fertilisation rate 160, it was 0.45. The highest estimated N losses occurred on sorghum plots with manure amendment (Supplementary Table S6). In general, wheat plots at different fertilisation rates lost more N compared to sorghum plots without manure amendment. The lowest estimated N losses occurred on barley plots.

260

3.2 Nitrogen cycle genes

265

The abundances of N cycling genes on plots with different fertilisation rates and crop species show different patterns throughout the study period (Supplementary Figures S5, S6, S7, and S8). The principal component analysis (PCA) of the N cycle genes abundances showed differences between sites with different fertilisation rates (Figure 4). There were greater differences in gene abundances between three different mineral N fertilisation rates in sorghum plots compared to barley and wheat plots (Figure 4). For sorghum without manure amendment (Figure 4C), archaeal 16S rRNA and *nosZII* gene abundances were highest for fertilisation rate 80 compared to rate 0 ja 160 ($p < 0.001$), but for sorghum with manure

270

amendment (Figure 4D), the highest archaeal 16S rRNA and *nosZII* gene abundances were for fertilisation rate 160 compared to rate 0 ($p < 0.001$) and 80 ($p < 0.05$). For all sorghum plots, comammox *amoA* gene abundance was highest on non-fertilised plots. However, fertilised wheat and barley plots had higher comammox *amoA* gene abundance compared to non-fertilised plots.

275





Figure 4: Principal components analysis (PCA) ordination plots demonstrate the grouping of fertilisation rates according to functional marker genes abundances for different crop type. N_2O is added as a supplementary variable. The month shows the sampling time. Abbreviations: *bact16S* – bacterial 16S rRNA gene; *arch16S* – archaeal 16S rRNA gene; *amoAbact* – bacterial *amoA* gene; *amoAarch* – archaeal *amoA* gene; *amoAcomammox* – comammox *amoA* gene.

280

3.3 N_2O emissions

The N_2O emissions over the course of the study period show that different fertilisation rates influence N_2O emissions, and the highest N_2O emissions tend to be emitted from the highest N fertiliser treatment (160 kg N ha⁻¹) (Figure 5). N_2O emissions among all crop species tended to be higher during the first part of the study period (spring and early summer). Taken together, the highest average N_2O emissions for barley plots were measured in the middle of May, for sorghum plots without and with manure in the middle of June, and for wheat plots at the beginning of June.

285



290 **Figure 5:** N₂O emissions ($\mu\text{g N m}^{-2} \text{h}^{-1}$) according to crop types and fertilisation rates during the study period.

Throughout the study period, cumulative N₂O and N₂ emissions were highest in plots with the highest fertilization rate, except sorghum plots with manure amendment (Figure 6A, B). For wheat and barley plots, there is a clear pattern of increasing N₂O emissions with increasing fertilisation rates.

295 For barley plots, cumulative N₂O emissions did not significantly differ between fertilisation rates 0 and 80 (Figure 6A). However, N₂O emissions on barley plots were significantly higher at fertilisation rate 160 than at rate 0 and 80 ($p < 0.05$). Similarly, for wheat plots, cumulative N₂O emissions were also significantly higher at fertilisation rate 160 compared to rates 0 ($p < 0.05$) and 80 ($p < 0.05$); however, fertilisation rates 0 and 80 did not significantly differ from each other. For plots with sorghum without manure, cumulative N₂O emissions at fertilisation rate 160 were significantly higher compared to fertilisation rates 0 ($p < 0.05$) and 80 ($p < 0.05$). For sorghum with manure plots, cumulative N₂O emissions at fertilisation rate 160 was significantly different compared to fertilisation rate 0 ($p < 0.05$) and 80 ($p < 0.05$).

For barley plots, the cumulative N₂ emissions were significantly higher at fertilisation rate 160 compared to rates 0 ($p < 0.05$) and 80 ($p < 0.05$) (Figure 6B). For wheat, sorghum with and without manure plots, cumulative N₂ emissions emitted from all three fertilisation rates did not significantly differ from each other.

305




Figure 6: Cumulative N₂O and N₂ emissions according to crop types and fertilisation rates. Error bars show standard errors. Letters above the boxes indicate statistically significant differences at significance level $p < 0.05$ according to a post-hoc Tukey HSD test. Lowercase letters indicate comparisons within crop types. Uppercase letters indicate comparisons of the same fertilisation rate over all crop types.

310

3.4 Relationships between environmental and genetic parameters and N emissions

Mineral N fertilisation rate ($p < 0.001$) and manure amendment ($p < 0.01$) significantly influenced N₂O emissions (Table 1). Crop type did not significantly influence N₂O emissions. The mineral N fertilisation effect on N₂O emissions ($\omega^2 = 0.528$) was larger compared to the effects of crop type ($\omega^2 = 0.021$) and manure amendment ($\omega^2 = 0.121$).

315


320

325 **Table 1:** Results of ANOVA testing the effects of crop type, mineral N fertilisation rate and manure amendment on cumulative N₂O fluxes. Significance is indicated as *** – 0.001; ** – 0.01; * – 0.05; ns – not significant.

	Df	F value	Pr (>F)
Crop type	2	0.957	0.39544
Mineral N fertilisation rate	2	23.995	5.97×10 ⁻⁷ ***
Manure amendment	1	11.020	0.00237 **
Residuals	30		

330 **Random Forest classification analysis** for the N₂O emissions from wheat plots considered bacterial *amoA*, archaeal *amoA*, *nosZI* and *nosZII* genes relevant (Figure 7). For barley plots, bacterial *amoA*, comammox *amoA*, bacterial 16S rRNA, *nirK*, *nirS* and *nosZII* were deemed as important genes in the **variations** of N₂O emissions. For sorghum without manure amendment plots, bacterial *amoA*, comammox *amoA*, archaeal 16S rRNA and *nirK* genes were considered important for the N₂O emissions. For sorghum with manure amendment plots, archaeal *amoA*, bacterial *amoA*, comammox *amoA*, *nirK*, *nirS*, *nosZII*, *nosZI* and *nrfA* genes were considered important for the N₂O emissions.

335



340

Figure 7: Results of feature selection in predicting the genes that are important in the variations of N₂O emissions. Important factors are indicated in green, unimportant factors in red, and shadow variables (the random shadow copies of features (noise) will be created to test the feature against those copies to determine if it is better than the noise, and therefore significant) in blue. Abbreviations: bact16S – bacterial 16S rRNA gene; arch16S – archaeal 16S rRNA gene; amoAabact – bacterial *amoA* gene; amoAaarch – archaeal *amoA* gene; amoAacomammox – comammox *amoA* gene.

345

The relationships between gene abundances and N₂O emissions showed that the ratio of *amoA/nir* was in a significant positive correlation with N₂O emissions ($\rho=0.20$; $p<0.001$). Furthermore, the ratio of *nosZ/nir* was also significantly positively correlated with N₂O emissions ($\rho=0.21$; $p<0.001$). *nirS* genes were positively correlated with N₂O emissions over all crops species plots ($\rho=0.19$; $p<0.05$). N₂O emissions from barley plots had also a strong positive correlation with *nirS* gene abundance ($\rho=0.58$; $p<0.001$). On wheat plots, *nosZII* genes were negatively correlated with N₂O emissions ($\rho=-0.46$; $p<0.01$). The correlation matrix is provided as Table S7 in Supplementary materials.

350

The relationship between N₂ emissions and *nrfA* genes showed that N₂ emissions were negatively correlated to *nrfA* genes over all crop types ($\rho=-0.39$; $p<0.05$). *nosZII* genes were positively correlated with N₂ emissions on plots with wheat ($\rho=0.85$; $p<0.01$).

355

4 Discussion

Mineral N fertilisation positively influenced biomass increase in all three crop types (Figure 3A), with similar findings observed in other IOSDV experiments by Csitári *et al.* (2021) and Tajnšek *et al.* (2013). The results also showed a significant positive correlation between biomass production and soil NO_3^- -N, C_{tot} and N_{tot} content, explaining higher biomass production in fertilised soil, as N limitation is the most influential factor constraining crop growth (Mengel and Kirkby, 2001). Furthermore, increasing mineral N fertilisation led to higher N accumulation in the biomass (Figure 3). The higher N content in biomass can be explained by applying N at rates that exceed crop needs for optimal yield, leading to an increase in crop protein content (Serret *et al.*, 2008; Mengel and Kirkby, 2001).

The sorghum plots without fertilisation yielded 2.3 t ha^{-1} , while those with only manure amendment produced an additional 5.9 t ha^{-1} of total dry biomass (Figure 3A), consistent with the results from Spiegel *et al.* (2010). N_{tot} and C_{tot} were also higher on sorghum plots with manure amendment compared to plots with only mineral fertilisation (Figure S3), which can explain higher biomass production. The positive effect of manure amendment could be attributed to increased availability of nutrients. Meta-analysis by Hijbeek *et al.* (2017), covering 20 long-term experiments (including the IOSDV experimental site used in our study) in Europe, reported that organic input does not necessarily guarantee increased crop yields. Although Hijbeek *et al.* (2017) also found that in specific cases, like spring sown cereals and sandy soils, the use of organic inputs led to an increase in crop yield.

In various ecosystems, N cycle genes have been linked to N_2O emissions (Butterbach-Bahl *et al.*, 2013; Espenberg *et al.*, 2018; Harter *et al.*, 2014). The significant positive correlation between the ratio of *amoA/nir* and N_2O emissions ($\rho=0.20$, $p<0.001$) in our study indicates that nitrification potential was higher than denitrification potential and thereby N_2O emissions were mainly related to nitrification in the soil. Previous studies have also used the ratio of *amoA* to *nir* genes to study N-cycle processes (Kazmi *et al.*, 2023; Tang *et al.*, 2018; Zhu *et al.*, 2018). Additionally, an initial decrease in NH_4^+ -N content in soil was observed, suggesting NH_4^+ consumption (nitrification) and mineral N uptake by plants (Supplementary Figure S2). A simultaneous increase in NO_3^- -N accompanied by a decrease in NH_4^+ -N was recorded, likely resulting from the nitrification production process.

nirS genes exhibited a positive correlation with N_2O emissions across all crops species plots ($\rho=0.19$; $p<0.05$), suggesting that while nitrification is predominant, denitrification is also evident. This finding aligns with results from several other agricultural studies, which also reported a significant positive correlation between *nirS* genes and N_2O emissions (Castellano-Hinojosa *et al.*, 2020; Cui *et al.*, 2016). Additionally, the ratio of *nosZ* to *nir* genes (*nosZ/nir*) was positively correlated with N_2O emissions ($\rho=0.21$, $p<0.001$). It highlights the importance of complete denitrifiers that have a capacity to convert N_2O to N_2 . Since N_2O emissions are increasing with a high abundance of the *nosZ* gene, this positive correlation may be also related to N_2O emissions being emitted from nitrification.

For all plots, one or more functional marker genes related to nitrification and denitrification were identified as important in the variations of N_2O emissions (Figure 7), emphasizing the significance of both processes in N_2O emissions. Comammox

was also recognized as an important process in N₂O emissions, except in wheat plots, indicating its potential important role. Additionally, Li *et al.* (2019) demonstrated an order of magnitude higher abundance of comammox *Nitrospira* clade A compared to ammonia-oxidizing archaea and ammonia-oxidizing bacteria in fertilised agricultural soil. More functional marker genes show significance in the variations of N₂O with manure compared to other treatments (Figure 7), indicating that a greater number of N cycle processes are relevant in plots with manure. Additionally, *nosZI*, *nosZII* and *nirS* genes were identified as important in the variations of N₂O emissions for sorghum with manure amendment, but not for only mineral fertiliser sorghum plots, which indicates significance of denitrification in these plots. Previous studies also suggest a higher denitrification potential from manure treatment, highlighting the importance of denitrifying microorganisms in manure-fertilised plots (Clark *et al.*, 2012; Wan *et al.*, 2023). The increased denitrification rate in manure-amended plots may be due to improved soil water retention promoting denitrification and increased availability of labile C content, which is the energy source for denitrifiers (Lazcano *et al.*, 2021; Rayne and Aula, 2020). Our results also support higher labile C content in plots with manure amendment (Figure 2). Furthermore, sorghum with manure plots were the only plots where the *nrfA* gene was identified as an important gene in N₂O emissions, suggesting that manure amendment is likely enhancing the rate of DNRA process.

The negative relationship between *nrfA* gene and N₂ emissions is suggesting that the DNRA process is not contributing to N₂ emissions. The DNRA process, which is mediated by *nrfA* gene, is beneficial as it supplies NH₄⁺ to the soil and conserves bioavailable N (Bai *et al.*, 2020; Pandey *et al.*, 2020). In addition, significant positive correlation between *nosZII* genes and N₂ emissions on plots with wheat indicates that there is likely a potential production of N₂ due to high abundance of *nosZII* genes that reduce N₂O into inert N₂. It is also supported by negative correlation between *nosZII* genes and N₂O emissions ($\rho=-0.46$; $p<0.01$) on wheat plots. It indicates *nosZII* genes role in reducing N₂O emissions (Graf *et al.*, 2014). Jones *et al.* (2014) demonstrated that the abundance and phylogenetic diversity of *nosZII* community is an important factor driving the soil's N₂O sink capacity.

Agricultural soils typically act as a source of N₂O (Davidson and Kanter, 2014), as shown in this study. The three mineral N fertilisation rates investigated influenced N₂O emissions, with N₂O emissions increasing with higher mineral N application rate for all three crop species (Figure 5, 6A). This can be attributed to higher available N levels with increased fertilisation rates for processes contributing to N₂O emissions (Engel *et al.*, 2010), as N₂O emissions showed a strong positive correlation with both NO₃⁻-N and NH₄⁺-N levels in soil. Prior studies have also highlighted a positive relationship between soil N₂O emissions and mineral N content (Sosulski *et al.*, 2014; Yao *et al.*, 2009; Yuan *et al.*, 2022). Furthermore, among the investigated factors, the mineral N fertilisation rate was the primary determinant of cumulative N₂O emissions (Table 1), indicating that soil N₂O emissions are mainly linked to the excess N added with mineral fertiliser in the cropping system (Supplementary Table S6).

Our study found that crop type did not significantly affect cumulative N₂O emissions, while the effects of mineral N fertilisation rate and manure amendment on cumulative N₂O emissions were significant (Table 1). It suggests that N₂O emissions from soil are more closely related to the excess N in the cropping system than the crop type. Study including 372

425 sites also showed that cover crops did not have significant ($p>0.05$) effect on N_2O emissions (Abdalla *et al.*, 2019). However, some studies have shown a significant effect of crop type on N_2O emissions (Bouwman *et al.*, 2002; Kaiser and Ruser, 2000). Manure amendment significantly impacted N_2O emissions (Table 1). Additionally, mineral fertiliser plus manure amendment showed higher soil N_2O emissions compared to mineral fertiliser alone for sorghum. This can be attributed to the overall higher mineral input of N into the cropping system in mineral fertiliser plus manure plots compared to mineral fertiliser-only plots ($231.2 \text{ kg N ha}^{-1}$ was added extra), enhancing N_2O production. In addition to providing nitrifiable N compounds, manure incorporation improves soil conditions for nitrification and denitrification by increasing moisture and adding C to the soil (Chadwick *et al.*, 2000). While the increase in moisture with manure was not detectable from our study, it may be explained by the slow evolution of soil properties over previous years in the 33-year-long fertilisation experiment. Moreover, manure can enhance the activity of soil microbes, oxygen consumption, and the development of anaerobic zones in the soil, favouring denitrification (Akiyama and Tsuruta, 2003).

430 Soil microbial processes leading to N_2O production are influenced by soil water content, as it directly affects oxygen availability for nitrification and denitrification processes. The recorded lowest soil moisture contents for barley and wheat plots on 7th of July (Supplementary Figure S4) likely explain the lowest N_2O emissions on that date (Figure 5). Previous studies on N_2O emissions and soil moisture dynamics have reported similar trend (Yamulki *et al.*, 1995; Yuan *et al.*, 2022). Additionally, Thapa *et al.* (2017) reported a reduction in N_2O emissions from wheat fields, which could be due to soil salinity interfering with nitrification and denitrification processes (Dang *et al.*, 2016). Dry soils may lead to microorganisms experiencing cell dehydration and increased soil salinity, hindering soil microbial activity and, therefore, the production of gaseous N emissions (Haj-Amor *et al.*, 2022; Schimel *et al.*, 2018). Although our study did not find significant correlations between soil moisture and N_2O emissions or most of the functional marker genes.

440 Considering climate changes and population growth, N_2O management should be aligned with the future need to increase crop yield and sustain rapidly increasing human population. Biomass production increased with fertilisation rate (Figure 3), except for wheat plots. In our study, the biomass production on wheat plots between fertilisation rates 80 kg N ha^{-1} and 160 kg N ha^{-1} had very similar biomass values. However, long-term fertilisation experiments (IOSDV) by Káš *et al.* (2010) achieved highest wheat yields from N fertilisation rate of 160 kg N ha^{-1} . Our study shows increasing N_2O emissions at higher fertilisation rate on wheat plots (Figure 6A), indicating potential overfertilisation and suggesting fertilisation rate at 80 kg N ha^{-1} as the optimal fertilisation rate. In addition, the highest nitrogen use efficiency (NUE) was observed at fertilisation rate 80 kg N ha^{-1} for wheat (NUE = 0.84), indicating a balance between low N_2O emissions and high yield. In India, Chaturvedi *et al.* (2006) conducted similar fertilisation experiments with N fertilisation rates of 0, 25, 50, 75, 100, and 125 kg N ha^{-1} , and identified the highest N input rate as optimal.

450 On fertilisation rate of 160 kg N ha^{-1} , N_2O emissions significantly increased compared to lower rates, but this rate also results in higher total dry biomass (Figure 3; Figure 6). The fertilisation rate of 80 kg N ha^{-1} for sorghum plots with only mineral N fertiliser amendment appears optimal with low N_2O emissions and N losses (Supplementary Table S6). However, in sorghum plots without manure amendment, NUE values are low (160 kg N ha^{-1} NUE = 0.25; 80 kg N ha^{-1} NUE = 0.12).

5 Conclusions

The results of our study (part of the 33 year old IOSDV experiment) showed that the mineral N fertilisation rate was the dominant factor determining cumulative N₂O emissions. The study observed an increase in N₂O emissions with an elevated mineral N fertilisation rate, attributed to higher NO₃⁻-N and NH₄⁺-N levels in fertilised soil. Higher N₂O emissions were measured during spring and early summer when mineral N fertilisers and farmyard manure was applied. These findings supported our hypothesis of higher N₂O emissions on sorghum plots under mineral fertiliser plus manure treatment compared to only mineral fertiliser treatment. Additionally, the number of N-cycle genes that are significant in the variations of N₂O emissions also increased with manure amendment. Contrary to our hypothesis, crop type did not have significant effect on N₂O emissions in this study.

N₂O emissions were mostly caused by nitrification with potential contribution from denitrification, comammox and DNRA processes. Plots with manure amendment exhibited a greater impact of N-cycle microbial processes on N₂O emissions, compared to plots with other crop types. Soil moisture showed no correlation with N₂O emissions and most of the functional marker gene abundances. Nonetheless, the lowest N₂O emissions and functional marker gene abundances were recorded during periods of low soil moisture, suggesting a decrease in N₂O under such conditions.

For wheat, a high NUE value and low N₂O emissions, coupled with relatively high crop yield, suggest that a fertilisation rate of 80 kg N ha⁻¹ is optimal. Similarly, in sorghum plots with only mineral N fertiliser amendment, a fertilisation rate of 80 kg N ha⁻¹ resulted in low N₂O emissions and N losses considering comparable biomass production with other crop types.

Author contributions. ME, AA, and ÜM designed the experiment and developed the methodology. LK and JEG carried out the fieldwork. LK analysed the results, performed data visualization, and wrote the original manuscript. JEG and ME participated in data analyses and assisted with paper editing. All authors were involved in revising the paper for submission and contributed to its improvement.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Acknowledgements. We thank Triin Teesalu for assisting with soil sampling for chemical analyses. Thanks to Tõnu Tõnutare and Kristi Kõva for conducting soil chemical analysis. We would also like to acknowledge Avo Toomsoo for operating the fertilisation experiment and maintenance of the study field.

Financial support. The study was supported by the Estonian Research Council (grants number PRG352 and PRG2032), European Research Council (ERC) under the grant agreement No 101096403 (MLTOM23415R), European Commission through the HORIZON-WIDERA ‘Living Labs for Wetland Forest Research’ Twinning project No 101079192 and the European Regional Development Fund (Centres of Excellence EcolChange, TK131, and AgroCropFuture, TK200).

490 **References**

- Abdalla, M., Hastings, A., Cheng, K., Yue, Q., Chadwick, D., Espenberg, M., Truu, J., Rees, R. M., & Smith, P.: A critical review of the impacts of cover crops on nitrogen leaching, net greenhouse gas balance and crop productivity. *Global Change Biology*, 25(8), 2530–2543. <https://doi.org/10.1111/gcb.14644>, 2019.
- 495 Abdalla, M., Shang, Z., Espenberg, M., Cui, X., Mander, Ü., & Smith, P.: Impacts of crop type, management and soil quality indicators on background nitrous oxide emissions (BNE) from Chinese croplands: A quantitative review and analysis. *Environmental Science: Atmospheres*, 2(4), 563–573. <https://doi.org/10.1039/D2EA00033D>, 2022.
- Akiyama, H., & Tsuruta, H.: Effect of organic matter application on N₂O, NO, and NO₂ fluxes from an Andisol field. *Global Biogeochemical Cycles - GLOBAL BIOGEOCHEM CYCLE*, 17, 11–1. <https://doi.org/10.1029/2002GB002016>, 2003.
- 500 Anderson, R., Bayer, P. E., & Edwards, D.: (2020). Climate change and the need for agricultural adaptation. *Current Opinion in Plant Biology*, 56, 197–202. <https://doi.org/10.1016/j.pbi.2019.12.006>, 2020.
- 505 Andrews, M., & Lea, P. J.: Our nitrogen ‘footprint’: The need for increased crop nitrogen use efficiency. *Annals of Applied Biology*, 163(2), 165–169. Scopus. <https://doi.org/10.1111/aab.12052>, 2013.
- Asseng, S., Ewert, F., Martre, P., Rötter, R. P., Lobell, D. B., Cammarano, D., Kimball, B. A., Ottman, M. J., Wall, G. W., White, J. W., Reynolds, M. P., Alderman, P. D., Prasad, P. V. V., Aggarwal, P. K., Anothai, J., Basso, B., Biernath, C., 510 Challinor, A. J., De Sanctis, G., ... Zhu, Y.: Rising temperatures reduce global wheat production. *Nature Climate Change*, 5(2), 2. <https://doi.org/10.1038/nclimate2470>, 2015.
- Astover, A., Szajdak, L. W., & Kölli, R.: Impact of Long-Term Agricultural Management and Native Forest Ecosystem on the Chemical and Biochemical Properties of Retisols’ Organic Matter. In L. W. Szajdak (Ed.), *Bioactive Compounds in* 515 *Agricultural Soils* (pp. 149–171). Springer International Publishing. https://doi.org/10.1007/978-3-319-43107-9_7, 2016.
- Ayiti, O. E., & Babalola, O. O.: Factors Influencing Soil Nitrification Process and the Effect on Environment and Health. *Frontiers in Sustainable Food Systems*, 6. [10.3389/fsufs.2022.821994](https://doi.org/10.3389/fsufs.2022.821994), 2022.
- 520 Bai, R., Fang, Y.-T., Mo, L.-Y., Shen, J.-P., Song, L.-L., Wang, Y.-Q., Zhang, L.-M., & He, J.-Z. Greater promotion of DNRA rates and nrfA gene transcriptional activity by straw incorporation in alkaline than in acidic paddy soils. *Soil Ecology Letters*, 2(4), 255–267. Scopus. <https://doi.org/10.1007/s42832-020-0050-6>, 2020.

- 525 Beeckman, F., Motte, H., & Beeckman, T.: Nitrification in agricultural soils: Impact, actors and mitigation. *Current Opinion in Biotechnology*, 50, 166–173. Scopus. <https://doi.org/10.1016/j.copbio.2018.01.014>, 2018.
- Bouwman, A. F., Boumans, L. J. M., & Batjes, N. H.: Modeling global annual N₂O and NO emissions from fertilized fields. *Global Biogeochemical Cycles*, 16(4), 28-1-28–29. <https://doi.org/10.1029/2001GB001812>, 2002.
- 530 Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R., & Zechmeister-Boltenstern, S.: Nitrous oxide emissions from soils: How well do we understand the processes and their controls? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1621), 20130122. <https://doi.org/10.1098/rstb.2013.0122>, 2013.
- 535 Cameron, K. C., Di, H. J., & Moir, J. L.: Nitrogen losses from the soil/plant system: A review. *Annals of Applied Biology*, 162(2), 145–173. Scopus. <https://doi.org/10.1111/aab.12014>, 2013.
- Castellano-Hinojosa, A., Correa-Galeote, D., González-López, J., & Bedmar, E. J.: Effect of nitrogen fertilisers on nitrous oxide emission, nitrifier and denitrifier abundance and bacterial diversity in closed ecological systems. *Applied Soil Ecology*, 145, 103380. <https://doi.org/10.1016/j.apsoil.2019.103380>, 2020.
- 540 Chadwick, D. R., Pain, B. F., & Brookman, S. K. E.: Nitrous Oxide and Methane Emissions following Application of Animal Manures to Grassland. *Journal of Environmental Quality*, 29(1), 277–287. <https://doi.org/10.2134/jeq2000.00472425002900010035x>, 2000.
- 545 Chaturvedi, I.: Effects of different nitrogen levels on growth, yield and nutrient uptake of wheat (*Triticum aestivum* L.). *International Journal of Agricultural Science* 2, 372–374, 2006.
- 550 Clark, I. M., Buchkina, N., Jhurreea, D., Goulding, K. W. T., & Hirsch, P. R.: Impacts of nitrogen application rates on the activity and diversity of denitrifying bacteria in the Broadbalk Wheat Experiment. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1593), 1235–1244. <https://doi.org/10.1098/rstb.2011.0314>, 2012.
- Coskun, D., Britto, D. T., Shi, W., & Kronzucker, H. J.: Nitrogen transformations in modern agriculture and the role of biological nitrification inhibition. *Nature Plants*, 3. <https://doi.org/10.1038/nplants.2017.74>, 2017.
- 555 Csitári, G., Tóth, Z., & Kökény, M.: Effects of Organic Amendments on Soil Aggregate Stability and Microbial Biomass in a Long-Term Fertilization Experiment (IOSDV). *Sustainability*, 13(17), 17. <https://doi.org/10.3390/su13179769>, 2021.

Cui, P., Fan, F., Yin, C., Song, A., Huang, P., Tang, Y., Zhu, P., Peng, C., Li, T., Wakelin, S. A., & Liang, Y.: Long-term organic and inorganic fertilization alters temperature sensitivity of potential N₂O emissions and associated microbes. *Soil Biology and Biochemistry*, 93, 131–141. <https://doi.org/10.1016/j.soilbio.2015.11.005>, 2016.

Cvetkov, M., & Tajnšek, A.: Soil organic matter changes according to the application of organic and mineral fertilizers within long-term experiments. *Acta Agriculturae Slovenica*, 93, 311–320, 2009.

565 Daims, H., Lebedeva, E. V., Pjevac, P., Han, P., Herbold, C., Albertsen, M., Jehmlich, N., Palatinszky, M., Vierheilg, J., Bulaev, A., Kirkegaard, R. H., Von Bergen, M., Rattei, T., Bendinger, B., Nielsen, P. H., & Wagner, M.: Complete nitrification by Nitrospira bacteria. *Nature*, 528(7583), 504–509. <https://doi.org/10.1038/nature16461>, 2015.

570 Dang, D. M., Macdonald, B., Warneke, S., White, I.: Available carbon and nitrate increase greenhouse gas emissions from soils affected by salinity. *Soil Research*, 55(1), 47-57. <https://doi.org/10.1071/SR16010>, 2016.

Davidson, E. A., & Kanter, D.: Inventories and scenarios of nitrous oxide emissions. *Environmental Research Letters*, 9(10). Scopus. <https://doi.org/10.1088/1748-9326/9/10/105012>, 2014.

575 Deforest, J. L., & Otuya, R. K.: Soil nitrification increases with elevated phosphorus or soil pH in an acidic mixed mesophytic deciduous forest. *Soil Biology and Biochemistry*, 142. <https://doi.org/10.1016/j.soilbio.2020.107716>, 2020.

Del Grosso, S. J., Parton, W. J., Mosier, A. R., Ojima, D. S., Kulmala, A. E., & Phongpan, S.: General model for N₂O and N₂ gas emissions from soils due to denitrification. *Global Biogeochemical Cycles*, 14(4), 1045–1060. <https://doi.org/10.1029/1999GB001225>, 2000.

Ehleringer, J. R.: Photosynthesis and Photorespiration: Biochemistry, Physiology, and Ecological Implications. *HortScience*, 14(3), 217–222. <https://doi.org/10.21273/HORTSCI.14.3.217>, 1979.

585 Ehleringer, J.R., Cerling, T. E.: C₃ and C₄ photosynthesis. T. Munn (Ed.), *Encyclopedia of Global Environmental Change*, John Wiley & Sons, Ltd, Chichester, 186-190, 2002.

Engel, R., Liang, D. L., Wallander, R., & Bembenek, A.: Influence of urea fertiliser placement on nitrous oxide production from a silt loam soil. *Journal of Environmental Quality*, 39(1), 115–125. Scopus. <https://doi.org/10.2134/jeq2009.0130>, 2010.

Escuer-Gatius, J., Löhmus, K., Shanskiy, M., Kauer, K., Vahter, H., Mander, Ü., Astover, A., Soosaar, K. Critical points for closing the carbon and nitrogen budgets in a winter rapeseed field. *Nutrient Cycling in Agroecosystems*, 122, 289–311. <https://doi.org/10.1007/s10705-022-10202-8>, 2022.

595

Espenberg, M., Truu, M., Mander, Ü., Kasak, K., Nõlvak, H., Ligi, T., Oopkaup, K., Maddison, M., & Truu, J.: Differences in microbial community structure and nitrogen cycling in natural and drained tropical peatland soils. *Scientific Reports*, 8(1), 1. <https://doi.org/10.1038/s41598-018-23032-y>, 2018.

600 FAO. *The future of food and agriculture: Trends and challenges*. Food and Agriculture Organization of the United Nations, 2017.

Frasier, I., Noellemeyer, E., Fernández, R., & Quiroga, A.: Direct field method for root biomass quantification in agroecosystems. *MethodsX*, 3, 513-159. <https://doi.org/10.1016/j.mex.2016.08.002>, 2016.

605

Graf, D. R. H., Jones, C. M., Hallin, S.: Intergenomic Comparisons Highlight Modularity of the Denitrification Pathway and Underpin the Importance of Community Structure for N₂O Emissions. *PLoS One*, 9(12). <https://doi.org/10.1371/journal.pone.0114118>, 2014.

610 Gu, J., Zheng, X., Wang, Y., Ding, W., Zhu, B., Chen, X., Wang, Y., Zhao, Z., Shi, Y., & Zhu, J.: Regulatory effects of soil properties on background N₂O emissions from agricultural soils in China. *Plant and Soil*, 295(1), 53–65. <https://doi.org/10.1007/s11104-007-9260-2>, 2007.

615 Haj-Amor, Z., Araya, T., Kim, D-G, Bouri, S., Lee, J., Ghiloufi, W., Yang, Y., Kang, H., Jhariya, M. K., Banerjee, A., Lal, R.: Soil salinity and its associated effects on soil microorganisms, greenhouse gas emissions, crop yield, biodiversity and desertification: A review. *Science of The Total Environment*, 843, 156946. <https://doi.org/10.1016/j.scitotenv.2022.156946>, 2022.

620 Harter, J., Krause, H.-M., Schuettler, S., Ruser, R., Fromme, M., Scholten, T., Kappler, A., & Behrens, S.: Linking N₂O emissions from biochar-amended soil to the structure and function of the N-cycling microbial community. *The ISME Journal*, 8(3), 3. <https://doi.org/10.1038/ismej.2013.160>, 2014.

Haynes, R. J., & Francis, G. S.: Changes in microbial biomass C, soil carbohydrate composition and aggregate stability induced by growth of selected crop and forage species under field conditions. *Journal of Soil Science*, 44(4), 665–675. Scopus. <https://doi.org/10.1111/j.1365-2389.1993.tb02331.x>, 1993.

Hibberd, J. M., & Quick, W. P.: Characteristics of C4 photosynthesis in stems and petioles of C3 flowering plants. *Nature*, 415(6870), 451–454. Scopus. <https://doi.org/10.1038/415451a>, 2002.

630 Hijbeek, R., van Ittersum, M. K., ten Berge, H. F. M., Gort, G., Spiegel, H., & Whitmore, A. P.: Do organic inputs matter – a meta-analysis of additional yield effects for arable crops in Europe. *Plant and Soil*, 411(1), 293–303. <https://doi.org/10.1007/s11104-016-3031-x>, 2017.

Hu, H.-W., Chen, D., & He, J.-Z.: Microbial regulation of terrestrial nitrous oxide formation: Understanding the biological pathways for prediction of emission rates. *FEMS Microbiology Reviews*, 39(5), 729–749. Scopus. <https://doi.org/10.1093/femsre/fuv021>, 2015.

International Organization for Standardization. Determination of total nitrogen content by dry combustion ("elemental analysis"). ISO Standard No. 13878:1998. <https://www.iso.org/standard/23117.html>, 1998.

640

IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2391 pp. <https://doi.org/10.1017/9781009157896>

IUSS Working Group WRB. World Reference Base for Soil Resources 2014 (update 2015). International Soil Classification System for Naming Soils and Creating Legends for Soil Maps. World Soil Resources Report No. 106. Rome, Food and Agriculture Organisation of the United Nations, 2015.

650

Jones, C. M., Spor, A., Brennan, F. P., Breuil, M.-C., Bru, D., Lemanceau, P., Griffiths, B., Hallin, S., Philippot, L.: Recently identified microbial guild mediates soil N₂O sink capacity. *Nature Climate Change*, 4, 801–805. <https://doi.org/10.1038/nclimate2301>, 2014.

- 655 Joshi, J., Stocker, B. D., Hofhansl, F., Zhou, S., Dieckmann, U., & Prentice, I. C.: Towards a unified theory of plant photosynthesis and hydraulics. *Nature Plants*, 8(11), 1304–1316. Scopus. <https://doi.org/10.1038/s41477-022-01244-5>, 2022.
- Kaiser, E.-A., & Ruser, R.: Nitrous oxide emissions from arable soils in Germany—An evaluation of six long-term field
660 experiments. *Journal of Plant Nutrition and Soil Science*, 163(3), 249–259. [https://doi.org/10.1002/1522-2624\(200006\)163:3<249::AID-JPLN249>3.0.CO;2-Z](https://doi.org/10.1002/1522-2624(200006)163:3<249::AID-JPLN249>3.0.CO;2-Z), 2000.
- Káš, M., Haberle, J., & Matějková, S.: Crop productivity under increasing nitrogen rates and different organic fertilization systems in a long-term IOSDV experiment in the Czech Republic. *Archives of Agronomy and Soil Science*, 56(4), 451–461.
665 <https://doi.org/10.1080/03650340903369392>, 2010.
- Kassambara, A.; Mundt, F.: Factoextra: Extract and Visualise the Results of Multivariate Data Analyses. R Package Version 1.0.7. Available online: <https://CRAN.R-project.org/package=factoextra> (accessed on 18 August 2023), 2020.
- 670 Kaur, B., Kaur, G., & Asthir, B.: Biochemical aspects of nitrogen use efficiency: An overview. *Journal of Plant Nutrition*, 40(4), 506–523. <https://doi.org/10.1080/01904167.2016.1240196>, 2017.
- Kazmi, F.A., Espenberg, M., Pärn, J., Masta, M., Ranniku, R., Thayamkottu, S., Mander, Ü.: Meltwater of freeze-thaw cycles drives N₂O-governing microbial communities in a drained peatland forest soil. *Biology and Fertility of Soils*.
675 <https://doi.org/10.1007/s00374-023-01790-w>, 2023.
- Kim, D.-G., Giltrap, D., & Hernandez-Ramirez, G.: Background nitrous oxide emissions in agricultural and natural lands: A meta-analysis. *Plant and Soil*, 373(1), 17–30. <https://doi.org/10.1007/s11104-013-1762-5>, 2013.
- 680 Koch, H., van Kessel, M. A. H. J., & Lückner, S.: Complete nitrification: Insights into the ecophysiology of comammox Nitrospira. *Applied Microbiology and Biotechnology*, 103(1), 177–189. <https://doi.org/10.1007/s00253-018-9486-3>, 2019.
- Kursa, M. B., & Rudnicki, W. R.: Feature Selection with the Boruta Package. *Journal of Statistical Software*, 36, 1–13. <https://doi.org/10.18637/jss.v036.i11>, 2010.
- 685 Kuypers, M. M. M., Marchant, H. K., & Kartal, B.: The microbial nitrogen-cycling network. *Nature Reviews Microbiology*, 16(5), 263–276. <https://doi.org/10.1038/nrmicro.2018.9>, 2018.

- Lazcano, C., Zhu-Barker, X., Decock, C.: Effects of Organic Fertilizers on the Soil Microorganisms Responsible for N₂O Emissions: A Review. *Microorganisms*, 9(5), 983. <https://doi.org/10.3390/microorganisms9050983>, 2021.
- 690
- Lê, S., Josse, J., & Husson, F.: FactoMineR: An R Package for Multivariate Analysis. *Journal of Statistical Software*, 25, 1–18. <https://doi.org/10.18637/jss.v025.i01>, 2008.
- Ledvinka, H. D., Toghyani, M., Tan, D. K. Y., Khoddami, A., Godwin, I. D., & Liu, S. Y.: The Impact of Drought, Heat and Elevated Carbon Dioxide Levels on Feed Grain Quality for Poultry Production. *Agriculture*, 12(11), 11. <https://doi.org/10.3390/agriculture12111913>, 2022.
- 695
- Li, C., Hu, H.-W., Chen, Q.-L., Chen, D., & He, J.-Z.: Comammox Nitrospira play an active role in nitrification of agricultural soils amended with nitrogen fertilizers. *Soil Biology and Biochemistry*, 138, 107609. <https://doi.org/10.1016/j.soilbio.2019.107609>, 2019.
- 700
- Liu, B., Asseng, S., Müller, C., Ewert, F., Elliott, J., Lobell, D. B., Martre, P., Ruane, A. C., Wallach, D., Jones, J. W., Rosenzweig, C., Aggarwal, P. K., Alderman, P. D., Anothai, J., Basso, B., Biernath, C., Cammarano, D., Challinor, A., Deryng, D., ... Zhu, Y.: Similar estimates of temperature impacts on global wheat yield by three independent methods. *Nature Climate Change*, 6(12), 12. <https://doi.org/10.1038/nclimate3115>, 2016.
- 705
- Liu, S., Wang, J. J., Tian, Z., Wang, X., & Harrison, S.: Ammonia and greenhouse gas emissions from a subtropical wheat field under different nitrogen fertilisation strategies. *Journal of Environmental Sciences*, 57, 196–210. <https://doi.org/10.1016/j.jes.2017.02.014>, 2017.
- 710
- Maa-amet. (2023). <https://xgis.maaamet.ee/xgis2/page/app/maainfo>
- Mahato, A.: Climate Change and its Impact on Agriculture. *International Journal of Scientific and Research Publications*, 4(4), 2014.
- 715
- Mengel, K., Kirkby, E. A.: Principles of plant nutrition. 5th ed., Kluwer Academic Publishers, Netherlands, 2001.
- Nardi, P., Laanbroek, H. J., Nicol, G. W., Renella, G., Cardinale, M., Pietramellara, G., Weckwerth, W., Trinchera, A., Ghatak, A., & Nannipieri, P.: Biological nitrification inhibition in the rhizosphere: Determining interactions and impact on microbially mediated processes and potential applications. *FEMS Microbiology Reviews*, 44(6), 874–908. Scopus. <https://doi.org/10.1093/femsre/fuaa037>, 2020.
- 720

- 725 Ostberg, S., Schewe, J., Childers, K., & Frieler, K.: Changes in crop yields and their variability at different levels of global warming. *Earth System Dynamics*, 9(2), 479–496. <https://doi.org/10.5194/esd-9-479-2018>, 2018.
- 730 Pandey, C. B., Kumar, U., Kaviraj, M., Minick, K. J., Mishra, A. K., & Singh, J. S. DNRA: A short-circuit in biological N-cycling to conserve nitrogen in terrestrial ecosystems. *Science of the Total Environment*, 738. Scopus. <https://doi.org/10.1016/j.scitotenv.2020.139710>, 2020.
- 735 Pandey, R. K., Maranville, J. W., & Bako, Y.: Nitrogen Fertilizer Response and Use Efficiency for Three Cereal Crops in Niger. *Communications in Soil Science and Plant Analysis*, 32(9–10), 1465–1482. <https://doi.org/10.1081/CSS-100104206>, 2001.
- 740 Parton, W. J., Holland, E. A., Del Grosso, S. J., Hartman, M. D., Martin, R. E., Mosier, A. R., Ojima, D. S., & Schimel, D. S.: Generalized model for NO_x and N₂O emissions from soils. *Journal of Geophysical Research: Atmospheres*, 106(D15), 17403–17419. <https://doi.org/10.1029/2001JD900101>, 2001.
- 745 Pathak, T. B., Maskey, M. L., Dahlberg, J. A., Kearns, F., Bali, K. M., & Zaccaria, D.: Climate change trends and impacts on California Agriculture: A detailed review. *Agronomy*, 8(3). Scopus. <https://doi.org/10.3390/agronomy8030025>, 2018.
- 750 Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G. P., Smith, P.: Climatesmart soils. *Nature*, 532, 49–57. <https://doi.org/10.1038/nature17174>, 2016.
- 755 Philippot, L., Hallin, S., & Schloter, M.: Ecology of Denitrifying Prokaryotes in Agricultural Soil. *Advances in Agronomy* (Vol. 96, pp. 249–305). Academic Press. [https://doi.org/10.1016/S0065-2113\(07\)96003-4](https://doi.org/10.1016/S0065-2113(07)96003-4), 2007.
- 760 Poole, C. F.: Ionization-based detectors for gas chromatography. *Journal of Chromatography A*, 1421, 137–153. <https://doi.org/10.1016/j.chroma.2015.02.061>, 2015.
- 765 Putz, M., Schleusner, P., Rütting, T., & Hallin, S.: Relative abundance of denitrifying and DNRA bacteria and their activity determine nitrogen retention or loss in agricultural soil. *Soil Biology and Biochemistry*, 123, 97–104. <https://doi.org/10.1016/j.soilbio.2018.05.006>, 2018.
- 770 Ravishankara, A.R., Daniel, J.S., Portmann, R.W.: Nitrous Oxide (N₂O): The Dominant Ozone-Depleting Substance Emitted in the 21st Century. *Science*, 326 (5949), 123–125. <https://doi.org/10.1126/science.1176985>, 2009.

Rayne, N., Aula, L.: Livestock Manure and the Impacts on Soil Health: A Review. *Soil Systems*, 4(4), 64. <https://doi.org/10.3390/soilsystems4040064>, 2020.

760 Republic of Estonia Environment Agency. Meteorological Yearbook of Estonia 2022. Available online: <https://kaur.maps.arcgis.com/sharing/rest/content/items/1c8b8f59e7184619b4455b2beceacd6a/data> (accessed on 18 August 2023), 2023.

Robertson, G. P., & Vitousek, P. M. Nitrogen in Agriculture: Balancing the Cost of an Essential Resource. *Annual Review of Environment and Resources*, 34(1), 97–125. <https://doi.org/10.1146/annurev.environ.032108.105046>, 2009.

Ruser, R., Schulz, R.: The effect of nitrification inhibitors on the nitrous oxide (N₂O) release from agricultural soils—a review. *Journal of Plant Nutrition and Soil Science*, 178(2), 171-188. <https://doi.org/10.1002/jpln.201400251>, 2015.

770 Rütting, T., Boeckx, P., Müller, C., & Klemmedtsson, L.: Assessment of the importance of dissimilatory nitrate reduction to ammonium for the terrestrial nitrogen cycle. *Biogeosciences*, 8(7), 1779–1791. <https://doi.org/10.5194/bg-8-1779-2011>, 2011.

Sainju, U. M.: Determination of nitrogen balance in agroecosystems. *MethodsX*, 4, 199–208. <https://doi.org/10.1016/j.mex.2017.06.001>, 2017.

Saud, S., Wang, D., Fahad, S.: Improved Nitrogen Use Efficiency and Greenhouse Gas Emissions in Agricultural Soils as Producers of Biological Nitrification Inhibitors. *Frontiers in Plant Science*. <https://doi.org/10.3389/fpls.2022.854195>, 2022.

780 Schaffasz, A., Windpassinger, S., Friedt, W., Snowdon, R., Wittkop, B.: Sorghum as a Novel Crop for Central Europe: Using a Broad Diversity Set to Dissect Temperate-Adaptation. *Agronomy*, 9(9), 535. <https://doi.org/10.3390/agronomy9090535>, 2019.

Schimel, J.P.: Life in Dry Soils: Effects of Drought on Soil Microbial Communities and Processes. *Annual Review of Ecology, Evolution, and Systematics*, 49, 409-432. <https://doi.org/10.1146/annurev-ecolsys-110617-062614>, 2018.

Serret, M. d, Ortiz-Monasterio, I., Pardo, A., & Araus, J. L.: The effects of urea fertilisation and genotype on yield, nitrogen use efficiency, $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ in wheat. *Annals of Applied Biology*, 153(2), 243–257. <https://doi.org/10.1111/j.1744-7348.2008.00259.x>, 2008.

Shcherbak, I., Millar, N., & Robertson, G. P.: Global metaanalysis of the nonlinear response of soil nitrous oxide (N₂O) emissions to fertilizer nitrogen. *Proceedings of the National Academy of Sciences*, 111(25), 9199–9204. <https://doi.org/10.1073/pnas.1322434111>, 2014.

795 Shen, H., Shiratori, Y., Ohta, S., Masuda, Y., Isobe, K., & Senoo, K.: Mitigating N₂O emissions from agricultural soils with fungivorous mites. *The ISME Journal*, 15(8), 8. <https://doi.org/10.1038/s41396-021-00948-4>, 2021.

Sosulski, T., Szara, E., Stepien, W., & Szymańska, M.: Nitrous oxide emissions from the soil under different fertilization systems on a long-term experiment. *Plant, Soil and Environment*, 60, 481–488. <https://doi.org/10.17221/943/2013-PSE>,
800 2014.

Spiegel, H., Dersch, G., Baumgarten, A., & Hösch, J.: The International Organic Nitrogen Long-term Fertilisation Experiment (IOSDV) at Vienna after 21 years. *Archives of Agronomy and Soil Science*, 56(4), 405–420. <https://doi.org/10.1080/03650341003645624>, 2010.

805

Stevens, N., Bond, W., Feurdean, A., & Lehmann, C. E. R.: Grassy Ecosystems in the Anthropocene. *Annual Review of Environment and Resources*, 47(1), 261–289. <https://doi.org/10.1146/annurev-environ-112420-015211>, 2022.

Stremińska, M. A., Felgate, H., Rowley, G., Richardson, D. J., & Baggs, E. M.: Nitrous oxide production in soil isolates of
810 nitrate-ammonifying bacteria. *Environmental Microbiology Reports*, 4(1), 66–71. <https://doi.org/10.1111/j.1758-2229.2011.00302.x>, 2012.

Tajnsšek, A., Čergan, Z., & Čeh, B.: Results of the long-term field experiment IOSDV Jable at the beginning of the 21st century. *Archives of Agronomy and Soil Science*, 59(8), 1099–1108. <https://doi.org/10.1080/03650340.2012.697996>, 2013.

815

Tang, Y., Yu, G., Zhang, X., Wang, Q., Ge, J., Liu, S.: Changes in nitrogen-cycling microbial communities with depth in temperate and subtropical forest soils. *Applied Soil Ecology*, 124, 218–228. <https://doi.org/10.1016/j.apsoil.2017.10.029>,
2018.

820 Thapa, R., Chatterjee, A., Wick, A., Butcher, K.: Carbon Dioxide and Nitrous Oxide Emissions from Naturally Occurring Sulfate-Based Saline Soils at Different Moisture Contents. *Pedosphere*, 27(5), 868–876. [https://doi.org/10.1016/S1002-0160\(17\)60453-3](https://doi.org/10.1016/S1002-0160(17)60453-3), 2017.

- Thomson, A. J., Giannopoulos, G., Pretty, J., Baggs, E. M., & Richardson, D. J.: Biological sources and sinks of nitrous oxide and strategies to mitigate emissions. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1593), 1157–1168. <https://doi.org/10.1098/rstb.2011.0415>, 2012.
- 825
- Tian, X., Engel, B. A., Qian, H., Hua, E., Sun, S., & Wang, Y.: Will reaching the maximum achievable yield potential meet future global food demand? *Journal of Cleaner Production*, 294, 126285. <https://doi.org/10.1016/j.jclepro.2021.126285>,
830 2021.
- Tilman, D., Balzer, C., Hill, J., & Befort, B. L.: Global food demand and the sustainable intensification of agriculture. *Proceedings of the National Academy of Sciences of the United States of America*, 108(50), 20260–20264. Scopus. <https://doi.org/10.1073/pnas.1116437108>, 2011.
- 835
- Van Kessel, M. A. H. J., Speth, D. R., Albertsen, M., Nielsen, P. H., Op Den Camp, H. J. M., Kartal, B., Jetten, M. S. M., & Lüscher, S.: Complete nitrification by a single microorganism. *Nature*, 528(7583), 555–559. Scopus. <https://doi.org/10.1038/nature16459>, 2015.
- 840 **Van Reeuwijk, L. P.: Procedures for soil analysis. Technical Paper No. 9, 6th Edition, FAO/ISRIC, Wageningen, the Netherlands. 120 pp, 2002.**
- Wan, Z., Wang, L., Huang, G., Rasul, F., Awan, M. I., Cui, H., Liu, K., Yu, X., Tang, H., Wang, S., & Xu, H.: NirS and nosZII bacterial denitrifiers as well as fungal denitrifiers are coupled with N₂O emissions in long-term fertilized soils.
845 *Science of The Total Environment*, 897, 165426. <https://doi.org/10.1016/j.scitotenv.2023.165426>, 2023.
- Whetton, R. L., Harty, M. A., Holden, N. M.: Communicating Nitrogen Loss Mechanisms for Improving Nitrogen Use Efficiency Management, Focused on Global Wheat. *Nitrogen*, 3(2), 213-246. <https://doi.org/10.3390/nitrogen3020016>, 2022.
- 850 Yamulki, S., Goulding, K. W. T., Webster, C. P., & Harrison, R. M.: Studies on no and N₂O fluxes from a wheat field. *Atmospheric Environment*, 29(14), 1627–1635. [https://doi.org/10.1016/1352-2310\(95\)00059-8](https://doi.org/10.1016/1352-2310(95)00059-8), 1995.
- Yao, Z., Zhou, Z., Zheng, X., Xie, B., Mei, B., Wang, R., Butterbach-Bahl, K., & Zhu, J.: Effects of organic matter incorporation on nitrous oxide emissions from rice-wheat rotation ecosystems in China. *Plant and Soil*, 327, 315–330.
855 <https://doi.org/10.1007/s11104-009-0056-4>, 2009.

Yuan, J., Yan, L., Li, G., Sadiq, M., Rahim, N., Wu, J., Ma, W., Xu, G., & Du, M.: Effects of conservation tillage strategies on soil physicochemical indicators and N₂O emission under spring wheat monocropping system conditions. *Scientific Reports*, 12. <https://doi.org/10.1038/s41598-022-11391-6>, 2022.

860

Zaman, M., Nguyen, M. L., Šimek, M., Nawaz, S., Jamil Khan, Babar, M. N., & Zaman, S.: Emissions of Nitrous Oxide (N₂O) and Di-Nitrogen (N₂) from the Agricultural Landscapes, Sources, Sinks, and Factors Affecting N₂O and N₂ Ratios. In *Greenhouse Gases-Emission, Measurement and Management*; Liu, G., Ed.; InTech Europe: Rijeka, Croatia, 2012. <https://doi.org/10.5772/32781>, 2012.

865

Zhang, Q., Wu, Z., Zhang, X., Duan, P., Shen, H., Gunina, A., Yan, X., & Xiong, Z.: Biochar amendment mitigated N₂O emissions from paddy field during the wheat growing season. *Environmental Pollution*, 281. Scopus. <https://doi.org/10.1016/j.envpol.2021.117026>, 2021.

870 Zhou, M., Zhu, B., Brüggemann, N., Dannenmann, M., Wang, Y., & Butterbach-Bahl, K.: Sustaining crop productivity while reducing environmental nitrogen losses in the subtropical wheat-maize cropping systems: A comprehensive case study of nitrogen cycling and balance. *Agriculture, Ecosystems & Environment*, 231, 1–14. <https://doi.org/10.1016/j.agee.2016.06.022>, 2016.

875 Zhu, W., Wang, C., Hill, J., He, Y., Tao, B., Mao, Z., Wu, Q.: A missing link in the estuarine nitrogen cycle?: Coupled nitrification-denitrification mediated by suspended particulate matter. *Scientific Reports*, 8, 2282. <https://doi.org/10.1038/s41598-018-20688-4>, 2018.

880