

The Impact of Cloud Microphysics and Ice Nucleation on Southern Ocean Clouds Assessed with Single Column Modeling and Instrument Simulators

Andrew Gettelman^{1,2}, Richard Forbes³, Roger Marchand⁴, Chih-Chieh Chen¹, and Mark Fielding³

¹National Center for Atmospheric Research, Boulder, CO, USA

²Pacific Northwest National Laboratory, Richland, WA, USA

³European Centre for Medium Range Weather Forecasts, Reading, UK

⁴University of Washington, Seattle, WA, USA

Correspondence: A. Gettelman (andrew.gettelman@pnnl.gov)

Abstract. Supercooled liquid clouds are common at higher latitudes (especially over the Southern Ocean) and are critical for constraining climate projections. We take advantage of the Macquarie Island Cloud and Radiation Experiment (MICRE) to perform an analysis of observed and simulated cloud processes over the Southern Ocean in a region and season dominated by supercooled liquid clouds. Using a single-column version of the European Centre for Medium-range Weather Forecast's (ECMWF) Integrated Forecast System (IFS), we compare two different cloud microphysical schemes to ground based observations of cloud, precipitation and radiation over a 2.5 month period (January 1 - March 17, 2017). Both schemes are able to reproduce aspects of the cloud and radiation observations during MICRE to within the uncertainty of the data, when the thermodynamic profile is prescribed with relaxation. There are differences in water mass and representation of reflectivity between the schemes. A sensitivity study of the cloud microphysics schemes, one a bulk one-moment scheme, and the other a two-moment scheme with prediction of mass and number, indicates that several key processes create differences between the schemes. Surface radiative fluxes and total water path are highly sensitive to the formation and fall speed of precipitation. The prediction of hydrometeor number with the two-moment scheme yields a better comparison with observed reflectivity and radiative fluxes, despite predicting higher liquid water contents than observed. With the two-moment scheme, we are also able to test the sensitivity of the results to the input of liquid Cloud Condensation Nuclei (CCN) and Ice Nuclei (IN). The cloud properties and resulting radiative effects are found to be sensitive to the CCN and IN concentrations. More CCN and IN increase liquid and ice water paths respectively. Thus both the dynamic environment and aerosols, integrated through the cloud microphysics, are important for properly representing Southern Ocean cloud radiative effects.

1 Introduction

Supercooled liquid water, defined as water below the freezing point of 0°C (273K), is common in colder regions of the planet (Hu et al., 2010). Freezing at temperatures between 0 and -40°C (273 and 233K) typically require very high ice supersaturations and/or Ice Nucleating Particles, also called Ice Nuclei (IN). IN are relatively rare (McCluskey et al., 2018), leading to the presence of supercooled liquid and an important role for secondary ice (number) production processes (Järvinen et al.,

2022). Recent work has made clear that supercooled liquid is important for both weather and climate. For example, improving supercooled liquid has been shown to improve forecasts of 2m temperatures at high-latitudes over land (Forbes and Ahlgrim, 25 2014) and cold air outbreaks at high latitudes (Forbes et al., 2015), while Bodas-Salcedo et al. (2012) showed a deficiency of supercooled liquid in several climate models and demonstrated how this deficiency resulted in substantial errors in the radiative budget of the Southern Ocean. More generally, Bodas-Salcedo et al. (2019) and Gettelman et al. (2019b) illustrated in two different models how supercooled liquid clouds can impact cloud feedbacks and the climate sensitivity of the planet to an imposed forcing.

30 There is a longstanding bias in climate (Trenberth and Fasullo, 2010) and weather forecast (Forbes et al., 2015) models of too few Southern Ocean clouds and too much surface absorption of solar radiation. To reduce cloud model biases at mid- to high-latitudes, it is necessary to focus on the representation of cloud phase and precipitation, which is handled by the cloud microphysical scheme in a large scale model. In this manuscript we present observational and model comparisons over a season for supercooled liquid clouds over the Southern Ocean. We attempt to reproduce the observations in a single column version 35 of the ECMWF IFS (Integrated Forecast System) with cloud microphysics schemes from both a weather forecast model (the ECMWF IFS) and a climate model (Community Earth System Model version 2, CESM2).

The unique aspects to this study are several. First, we compare two ‘operational’ cloud microphysics codes in a region with complex cloud regimes within the same modelling framework. Forbes and Tompkins (2011) describe a bulk, one-moment (predicted mass only) scheme designed for weather forecasting. The other scheme (Gettelman et al., 2019a) is a two-moment 40 (predicted mass and number) scheme designed for climate simulation. We seek to understand what is similar between the schemes, what is different and how the differences affect model biases relative to the observations. In particular, is a two-moment scheme more representative of observations than the 1-moment IFS scheme. When comparing to observations, we also consider the limits of the observations in detecting clouds, as well as the assumptions used in the GCM for cloud cover and sub-grid representation of clouds which add uncertainty to the comparisons. We use a constrained model set up, with a 45 single column model forced by observed winds and temperatures over a 75 day austral summer period. This allows a statistical representation of a series of events and regimes, while also being able to look at detailed process rates and a close comparison to observations. Finally, we focus on clouds with extensive supercooled liquid, a long-standing issue for weather and climate models.

Section 2 discusses the simulation set up, the microphysical parametrization schemes, and the observations. Section 3 50 presents the results, including comparisons to the observations, and comparisons between the two schemes. We also present sensitivity studies of the model configurations to understand the role of different microphysical processes and why the schemes differ. A summary is in Section 4 and conclusions are in Section 5.

2 Methodology

Here we first describe the observations (Section 2.1), followed by the IFS Single Column Model, the IFS microphysical and 55 the MG3 microphysical schemes, the model radar simulator in (Section 2.2), and details of the simulations (Section 2.3).

2.1 MICRE Observations

During the Macquarie Island Cloud and Radiation Experiment (MICRE), observations were collected from March 2016 to March 2018 at Macquarie Island (54.5° S, 158.9°E), located about half-way between New Zealand and Antarctica. The project is described in detail in McFarquhar et al. (2021), and included deployment of a suite of ground-based instruments including a
60 ceilometer, surface rain disdrometer, microwave radiometer, and broadband shortwave (SW) and longwave (LW) radiometers, as well as an upward looking W-band cloud radar and depolarization lidar. Most of the instruments suffered periods with significant down time, and we focus on the 75-day period from January 1 to March 17, 2017, when upward looking cloud radar and microwave radiometer observations are available, in addition to twice-daily radiosondes and broadband radiometer fluxes. We focus on observations of surface downward radiation, radar reflectivity, and microwave-radiometer-derived liquid
65 water path, as well as the ancillary radiosonde temperature data. The MICRE broadband radiometer SW and LW fluxes have been evaluated and compared with Clouds and the Earth's Radiant Energy System (CERES) satellite synoptic (SYN) product (Doelling et al., 2013; Rutan et al., 2015) and the surface Energy Balanced and Filled (surface EBAF, Kato et al., 2018) fluxes by Hinkelman and Marchand (2020). The MICRE instruments location was on a narrow point of land and is generally representative of maritime conditions. The location is representative of the Southern Ocean, with frequent frontal passages and
70 different cloud types.

2.2 Model Description

In this study we run a version of the Integrated Forecast System (IFS) (ECMWF, 2019) in single column mode with two different cloud microphysical schemes. Both are 'bulk' schemes representing hydrometeors with a mean mass, and one is a two-moment scheme that also represents the mean number of hydrometeors. The schemes however often use similar bulk
75 formulations for process rates, either solely mass based, or with an assumed size in the one-moment (mass only) scheme. This makes for an interesting and direct set of process rate comparisons, and highlights where two-moment schemes may be necessary to capture different atmospheric process sensitivities and regimes.

2.2.1 IFS Single Column Model

The Single Column Model (SCM) is a standalone vertical column version of the IFS with the same suite of physical parameterizations as the global model. Time-varying surface and advective forcings can be specified as well as a relaxation towards a specified time-varying state with a given timescale. The particular version used here is IFS Cycle 46r1, used operationally at
80 ECMWF from June 2019 to June 2020. There are 137 vertical levels, the same as the operational global model, with a layer depth increasing from 20m to 170m in the lowest 2km and around 300m in the rest of the troposphere. Only an overview of the parameterizations is given here and further details can be found in the IFS 46r1 documentation (ECMWF, 2019).

85 The IFS has prognostic variables for cloud fraction, specific humidity, specific cloud liquid, cloud ice, rain and snow water contents. The sub-grid cloud parameterization is based on Tiedtke (1993) with sources and sinks from the vertical advection, radiation and convection parameterizations. Cloud fraction and saturation adjustment are determined by of the sub-grid cloud

parameterization scheme and is the same for both microphysics schemes. Supersaturation with respect to ice is allowed and the assumptions are described in Tompkins et al. (2007). The cloud scheme is tightly coupled with the convection parameterization with detrainment of cloud fraction, condensate and precipitation from sub-grid convective updraughts. Vertical advection due to convectively generated sub-grid subsidence within a convecting grid cell is represented, as well as turbulent erosion of cloud fraction and condensate at sub-grid cloud edges. The cloud and precipitation microphysics is described in the next subsection.

The parameterization of shallow, mid-level and deep convection is based on the mass-flux approach (Tiedtke, 1989; Bechtold et al., 2008, 2014). The turbulent mixing scheme follows the Eddy-Diffusivity Mass-Flux (EDMF) framework, with a K-diffusion turbulence closure and a mass-flux component to represent the non-local eddy fluxes in unstable boundary layers (Siebesma et al., 2007; Köhler et al., 2011). The radiation scheme (ecRad) is described in Hogan and Bozzo (2018) with the gas optics from the Rapid Radiation Transfer Model (RRTMG, Mlawer et al., 1997; Iacono et al., 2008). Cloud-radiation interactions are taken into account using the McICA (Monte Carlo Independent Column Approximation) method (Morcrette et al., 2008). Surface exchange and gravity wave drag are also represented (Balsamo et al., 2009; Lott and Miller, 1997; Beljaars et al., 2004; Orr et al., 2010).

2.2.2 IFS Microphysics

The IFS microphysics is a one-moment bulk scheme with prognostic variables for the mass of 4 classes of hydrometeor (liquid, ice, rain, snow) as described in Forbes and Tompkins (2011) and Forbes et al. (2011) with various modifications, particularly to improve the representation of mixed-phase boundary layer clouds (Forbes and Ahlgrimm, 2014) and warm-rain processes (Ahlgrimm and Forbes, 2014). The numerical formulation of the hydrometeor sedimentation follows an implicit upstream approach with parameterized fall speed for rain (Sachidananda and Zrnica, 1986; Abel and Boutle, 2012) and fixed fallspeeds for snow (1 m/s) and ice (0.13 m/s).

2.2.3 MG3 Microphysics

The Morrison-Gottelman microphysics scheme version 3 (MG3) is a two-moment bulk microphysics scheme with prognostic variables for mass and number concentration of 5 classes of hydrometeors (liquid, ice, rain, snow, graupel) as described by Gottelman et al. (2019a). MG3 is based on the original Morrison et al. (2005) scheme, adapted and modified extensively for climate models (Morrison and Gottelman, 2008), with extensions for ice nucleation (Gottelman et al., 2010) and prognostic precipitation (Gottelman and Morrison, 2015). The scheme without graupel is used in the Community Earth System Model version 2 (CESM2, Danabasoglu et al., 2020). The scheme compares well to mesoscale schemes by Morrison et al. (2005) and Thompson and Eidhammer (2014) for idealized shallow and convective cloud cases, as shown by Gottelman et al. (2019a).

For this study MG3 has been adapted for use in the IFS SCM as an alternative to the IFS parameterization of microphysical processes and sedimentation. The rest of the model is kept the same including the saturation adjustment, sub-grid cloud parameterization and convection interactions.

The additional MG3 prognostic variables for mass and number concentration of all 5 classes of hydrometeors have been added to the IFS SCM. The two-moment MG3 scheme can run with fixed hydrometeor number concentration, or use activated

number concentration rates ($\# \text{ s}^{-1}$) for liquid drops (cloud condensation nuclei, CCN) and ice crystals (ice nuclei, IN). Since the IFS does not have an explicit representation of aerosols or an activation scheme, for this work we assume constant CCN and IN activation rates of $1 \text{ cm}^{-3} \text{ s}^{-1}$ and $5 \text{ L}^{-1} \text{ s}^{-1}$ respectively for all altitudes at every timestep. These rates are just used to initialize new particles in the microphysics when the model physics condenses water or ice and we test the sensitivity of the results to the activation/nucleation rate.

2.2.4 Radar Simulator

The radar simulator, described in full in Fielding and Janisková (2020), provides the model-equivalent radar reflectivity factor to compare with observations. It has been adapted to run in-line within the SCM and, wherever possible, uses assumptions consistent with the two microphysics schemes used in this study. The simulator runs in ‘one-moment’ mode when the IFS microphysics scheme is used and ‘two-moment’ mode when the MG3 scheme is used. When in two-moment mode, the simulator computes bulk scattering properties by combining the MG3 particle size distributions and densities (Morrison and Gettelman, 2008) with the prognostic number concentration, mass and the same single scattering properties as the one-moment simulator described in Fielding and Janisková (2020).

The size distribution of hydrometeors is chosen such that it is consistent with the predicted masses (and prognostic number concentrations when in ‘two-moment’ mode). The radar reflectivity and attenuation is calculated by the radar simulator (primarily using Mie theory) using the double-column approach as described in Fielding and Janisková (2020), partitioning the signal into a uniform cloudy layer and a uniform clear layer. The hydrometeors are assumed to be uniformly distributed within each species’ cloud or precipitation fraction. The attenuation by hydrometeors depends on the overlap of clouds and precipitation; the simulations here assume maximum-random overlap. This has a large effect on the simulated reflectivity, as will be discussed later. Attenuation due to gaseous attenuation is also included based on the models of Liebe (1985) and Liebe et al. (1992). Details can be found in Fielding and Janisková (2020). Attenuation does not include a cut-off for any height dependent Minimum Detectable Signal (MDS), which will also affect the comparisons with observed reflectivity.

2.3 Simulation Setup

The simulations are set up by generating forcing files for the location of Macquarie Island station (54.5°S , 158.9°E) from January to March 2017. The forcing comes from CESM2 simulations relaxed to the MERRA2 reanalysis (Molod et al., 2015), similar to that used by Gettelman et al. (2020) for the Southern Ocean Clouds, Radiation, Aerosol, Transport Experimental Study (SOCRATES) project. The IFS SCM is relaxed back to the forcing profile with an 8 hour relaxation time for winds, temperature and humidity to keep the boundary layer structure from drifting too far from the radiosonde observations. Longer relaxation timescales, or no relaxation at all, caused the simulations to drift by several $^{\circ}\text{C}$ in the upper part of the boundary layer, resulting in significant biases in clouds relative to the observations. This is a key feature: getting boundary layer structure correct is vital for constraining cloud radiative effects. There is no relaxation of cloud variables and these are allowed to freely evolve during the simulation. The base time step for the simulations is 225s, but we also investigate the time step sensitivity of

both schemes by running at 60s and 900s (the time step used in the global forecast model at ECMWF depends on resolution but typically ranges from 225s to 900s).

155 Simulations are repeated with the IFS and MG3 microphysics for comparison, and a series of sensitivity tests are performed, modifying different aspects of the MG3 code (see section 3.2.3). The CESM2 version of the MG3 microphysics was initially implemented in the SCM, but first comparisons highlighted significant differences in the profiles of rain that were due to differences in a few basic assumptions in the IFS. To allow a more meaningful comparison of the process rates, it was decided to implement these IFS assumptions in the MG3 scheme. Rain evaporation is modified to more closely match the IFS, with
160 a scaling factor of 0.3 and a relative humidity threshold between 80% and 90% before evaporation can occur, both currently required in the IFS to reduce excessive evaporation in the operational IFS. The threshold for rain freezing is changed from -40°C in CESM2 to the value of -5°C in the IFS, so all precipitation below -5°C is treated as snow. In addition, MG3 did not originally calculate a fall speed for layers with no hydrometeors, so sedimenting condensate can enter a layer of zero velocity. A fall speed correction was added to ensure that if there are hydrometeors above, the fall speed is not zero (this particularly affects
165 rain). The sedimentation of precipitation is changed to an implicit monotonic scheme (Harris et al., 2020; Zhou et al., 2019) to reduce timestep sensitivity. Finally, there is one change for the ice microphysics: mixed-phase ice nuclei are determined by the Meyers et al. (1992) empirical function of temperature as in Morrison et al. (2005) and the IFS, rather than the Hoose et al. (2010) classical nucleation theory scheme used in CESM2. These changes together define the ‘base’ version of the MG3 scheme used for the comparison with the IFS in section 3, although the original CESM2 version of the MG3 scheme is also
170 included for comparison in the sensitivity study (section 3.2.3).

3 Results

First we compare the IFS and MG3 microphysics (modified from the CESM2 version as described in section 2.3) to observations from MICRE. We then examine the differences between the IFS and MG3 microphysics, including their timestep sensitivity, differences in the balance of process rates between the schemes, and finally sensitivity tests with the MG3 scheme.

175 3.1 Comparison to Observations

In this subsection, we compare model simulations to surface and satellite based observations. We begin by focusing on short-wave and longwave radiative fluxes, followed by in-cloud liquid water path (LWP) and radar reflectivity (obtained from the model via a radar simulator as described in section 2).

3.1.1 Top-of-Atmosphere and Surface Radiation

180 Figure 1 shows the downward surface shortwave (SW, Figure 1a) and longwave (LW, Figure 1b) radiative fluxes for the 75-day time series, for the surface radiometer (SFC, black line), the CERES SYN product retrieved from combined geostationary and MODIS satellite data (for the SW flux, SYN, blue), and the simulation results from the MG3 (orange) and IFS (green) microphysical schemes. Although there are short periods during which the model simulated fluxes depart noticeably from

the surface and satellite data, there is generally good agreement (on average). Differences are not easy to distinguish because
185 they are small and it is the average difference that is really important (hence the importance of the 90 day record). Hourly
correlations for both simulations with the observations are ~ 0.75 for the SW flux and ~ 0.6 for the LW, and the mean SW flux
for both simulations lies between the mean values from the surface and satellite data sets. The one-sigma sampling uncertainty
in the surface and satellite daily mean SW flux is about 17 Wm^{-2} and 5 Wm^{-2} for the LW. The sampling uncertainty is
190 estimated as the standard deviation divided by the square root of the number of days, effectively treating each day as equivalent
to a single independent sample. The difference between the surface and satellite mean SW flux might be due to a bias in one
or both data sets, or due to differences in the field of view (for example, it is possible that it is slightly cloudier or clouds
are less optically transparent over the island site than the nearby ocean) but we do not know for certain the source of the
mean difference. In short, the small differences between all of the mean SW fluxes shown in Figure 1a are not significant with
respect to the sampling uncertainty. Sampling and systematic uncertainties in the surface and satellite fluxes are discussed in
195 more detail by Hinkelman and Marchand (2020). Figure 1b also shows the strong correspondence between the simulated and
observed surface downward longwave fluxes. Here the SYN retrievals are not shown because there is a significant bias in the
CERES SYN retrievals at night (Hinkelman and Marchand, 2020). The small differences ($< 5 \text{ Wm}^{-2}$) between the mean model
LW fluxes and the observed mean value are likewise not significant with respect to the sampling uncertainty.

We examine the diurnal cycle of surface and top of atmosphere (TOA) SW and LW fluxes in Figure 2. For both the models
200 and the observational datasets, hours containing only low clouds occurred about 35% of the time and completely clear hours
only about 2%, with the remainder containing some hydrometeors above 2 km (most often with hydrometer both above and
below 2km). In Figure 2 each dataset is processed independently, and we have not restricted the low cloud times to those hours
in which only low cloud occur at the same time in both surface observations and model simulations. Doing so significantly
decreases the total number of hours and increases the estimated sampling uncertainties, but does not otherwise qualitatively
205 change the results. We examine the observed and simulated radar reflectivity and associated vertical profiles of cloud occurrence
in more detail later in this section.

The simulated diurnal cycle of surface downward SW (Figure 2e,f) and LW (Figure 2g,h) fluxes compare well with the
surface and satellite data; and the good agreement holds for low cloud periods (Figure 2f,h). There is no discernible diurnal
cycle in the observed or modeled surface LW flux (Figure 2g,h). The mean surface SW downward flux is about 30 Wm^{-2}
210 larger during low cloud periods (Figure 2f) as compared with all times (Figure 2e), while the LW downward flux is about 8
 Wm^{-2} smaller (Figure 2g,h). This demonstrates the significant role played by both shallow and deeper clouds in the radiative
energy budget, and the models capture this well. The modeled outgoing TOA SW flux (Figure 2a,b) also agrees reasonably
well with the CERES SYN product. The model (daily) means are within about 15 Wm^{-2} of the SYN mean, with the CERES
SYN product being a bit smaller (consistent with the downward CERES SYN surface SW flux being a bit larger than the model
215 simulated surface SW flux). For clarity, the sampling uncertainty is not shown on the model simulated curves. If displayed, the
one-sigma uncertainty in the model mean diurnal cycle would just overlap with the uncertainty shown for the observations. A
paired data test (not shown) indicates the 15 Wm^{-2} difference between the models and SYN data is significant at the one-sigma
level. Nonetheless, it remains a reasonable possibility that the 15 Wm^{-2} difference is just a result of sampling uncertainty. The




Figure 1. Time series of hourly averaged values: (a) downward surface shortwave (SW) flux, (b) downward longwave (LW) flux, and (c) liquid water path (LWP). For each dataset the legend gives the mean value, and for the SW and LW flux, the estimated one-sigma uncertainty in the mean is also given. For LWP, the legend gives the mean value for all times followed by the mean restricted to periods where only low clouds are present (see text). Surface observations (Black Solid and tick), MG3 microphysics (Orange Solid), IFS microphysics (Green Dashed). The CERES SYN data (Purple Solid and cross) and average values are also restricted to data collected between 9 and 16 hours local time to avoid biases associated with low solar zenith angles (see text).

220 same is not true for the outgoing TOA LW flux (Figure 2c). In the LW, while there is good agreement between the model
simulated outgoing TOA LW and the CERES SYN product during periods with low clouds for both simulations (Figure 2d),
under all sky conditions (Figure 2c) the IFS microphysical scheme has too little outgoing LW flux (and this is significant at
the two-sigma level in a paired data test). The underestimate in the IFS flux becomes even clearer when restricted to hours
where only high-level hydrometeors are present (not shown). As will be shown later, an examination of radar reflectivity fields
likewise reveals the simulations with the IFS microphysics have a higher occurrence of cloud above 6 km and we will return
225 to this topic after an examination of the liquid water path.

3.1.2 Liquid Water Path

The timeseries of LWP is shown in Figure 1c, and the mean diurnal cycle for all and low cloud conditions is shown in Figure 2i
and j. Both figures depict the mean in-cloud LWP, with the model in-cloud value taken as the grid mean value divided by
the model prognostic cloud fraction. Retrievals of the LWP based on the surface microwave radiometer (MWR) are only
230 accurate when the microwave radiometer is dry (when it has not recently rained or even drizzled heavily), and so the surface
(SFC) MWR data include only periods when low-and-non-precipitating cloud are present. The SFC LWP was derived using
a physical-iterative approach (Marchand et al., 2003), with a structural uncertainty of about 15 to 30 gm^{-2} (owing primarily
to uncertainties in the underlying microwave spectroscopy and the treatment of supercooled liquid water absorption). Here the
structural uncertainty is larger than the sampling uncertainty. On a minor note, the very small differences between the mean
235 LWP values given in the legends in Figure 1c and Figure 2i arise because in Figure 2i the data are first averaged in each hour
before the overall (daily) mean is taken and the number of samples is not constant throughout the day, but depends on when
low cloud conditions occur.

The CERES SYN mean (and median) values given in the legend of Figure 2i,j are restricted to data collected between 9
and 16 hours local time. The diurnal cycle plots show there are large nonphysical "spikes" in the CERES SYN LWP before
240 and after this time, approaching sunrise and sunset. During daylight, the CERES SYN cloud property retrieval is based on
visible and near infrared imagery and assumes that clouds (whether they are liquid or ice phase) are sufficiently spatially
homogeneous that scattering at these wavelengths can be treated using a one-dimensional (plane-parallel) approximation.
Details on the CERES retrieval methodology are given in Minnis et al. (2011, 2008) and largely follow the approach used
in CERES-MODIS products. The plane-parallel approximation is widely known to work poorly at large solar zenith angles
245 or large view angles, and we speculate that these spikes are a result of this approximation not working well. The CERES
SYN product does use an infrared-only algorithm at night, and we also speculate that the lower SYN LWP values observed
during nighttime are also a retrieval artifact (and may play a role in the nighttime downward LW surface bias in the SYN
product noted by Hinkelman and Marchand (2020)). As a point of comparison, we have also plotted (red bars in Figure 2i,j)
the mean plus/minus one-standard-error in the operational MODIS MOD06 and MYD06 mean LWP (Platnick et al., 2003,
250 2017), different than the CERES-MODIS algorithm. While MODIS, on board the Terra and Aqua satellites, orbits in a sun-
synchronous orbit, Macquarie Island is far enough south that it is often observed on the edges of the MODIS swath resulting
in local overpasses that can be somewhat earlier or later than the nominal 10:30 and 13:30 equator crossing times. The mean




Figure 2. Mean diurnal cycles. Outgoing TOA (a,b) SW and (c,d) LW flux. Downward surface (e,f) SW and (g,h) LW flux, and LWP. Left column (a,c,e,g,i) shows diurnal averages for all times and the right column (b,d,f,h,j) for periods when only low cloud (hydrometeors < 2 km) are present base don hourly average (simulated or observed) radar reflectivity. In each panel, shading that surrounds the surface or satellite values depicts the estimated one-sigma sampling uncertainty for a particular hour over all days. Sampling uncertainty for all datasets is of similar magnitude. The diurnal cycle of LWP is also shown for all times (i) and just low cloud periods (j). CERES SYN retrieved LWP in purple (which combines geostationary and MODIS data). MODIS only (red symbols) as described in the text. The legend gives the daily mean for each dataset, and for LWP both the daily mean and median are given. CERES SYN means and medians are restricted to data collected between 9 and 16 hours local time (see text).

MOD06 and MYD06 LWP, which are processed independently of the CERES project, are in reasonable agreement with the CERES SYN data.

255 As is apparent in Figure 2i,j, the MG3 microphysics results in a LWP that is too large on average, even when restricted to periods of low clouds. The time series of LWP in Figure 1c shows that the MG3 scheme results in much larger LWP during periods when the IFS scheme is relatively large (more than 300 g m^{-2} , e.g. near 350 and 1200 hours since January 1), while at other times it is noticeably lower than IFS and the observational data (e.g. near 250, 800 and 1000 hours since January 1). How is it that the MG3 shortwave radiative fluxes show little bias with respect to observations while there is a large bias
260 in LWP? Part of the answer to this apparent contradiction is that the distribution of LWP is highly skewed, where a small occurrence of large LWP events substantially increases the mean. The median values of MG3 LWP are comparable to that for the observational dataset and higher than the IFS (the mean / median values are given in the legend of Figure 2i,j). The nonlinear relationship between albedo and LWP is also a factor. Figure 3 plots the TOA SW albedo as a function of the LWP for the low cloud periods. The MG3 simulation clearly has more high LWP events, but the high LWP does not substantially
265 increase the SW albedo because the albedo begins to saturate when the LWP is much above about 300 g m^{-2} with little change in albedo (from 0.55-0.65) for large changes in LWP. The partially-offsetting greater occurrence of low LWP events in MG3 is also evident in Figure 3. We note that ice condensate also has a major effect on the SW albedo in deeper clouds and likely reduces the impact of the large LWP events on albedo, but the cloud condensate below 2 km is predominantly liquid.

3.1.3 Radar Reflectivity

270 Figure 4 illustrates the observed (Figure 4a) and simulated reflectivity from the simulations with MG3 (Figure 4b) and IFS (Figure 4c) microphysics. Reflectivity is estimated using a radar simulator as described in Section 2. Liquid, ice, rain and snow and their attenuation are included. In this figure, the observed and simulated reflectivity are averaged over an hour (in power or Z-space), with the hourly values given here as equivalent radar reflectivity dBZe (where $\text{dBZe} = 10 \log_{10}(Ze)$). As with the SW and LW fluxes, there is good correspondence between the simulations and observations with both depicting synoptic events
275 with deep cloud layers that have high reflectivity, and periods with persistent low cloud. However, compared to the observations the models show a greater occurrence of high reflectivity near the surface related to rain (red colors), especially beneath the deep clouds, and a greater occurrence of low reflectivity (light blue colors) in the upper troposphere related to ice cloud. Here, the radar simulator does NOT represent the effect of the ground-based radar's Minimum Detectable Signal (MDS) such that the simulated radar reflectivity is sometime below the reflectivity the actual radar can measure. As we will discuss in more
280 detail below, this depends strongly on the altitude (distance from the radar). Also note that rain on the radar radome can result in attenuation of the radar signal, so with periods of significant rain, more attenuation can be expected than has been included in the radar simulator. Thus we do not expect the observed radar reflectivity to definitively characterize high altitude clouds. The TOA radiative fluxes (Figure 2a,c) are a better measure of the climate effect of the high clouds.

To better understand these differences, we look at reflectivity-height histograms (often called a Contoured Frequency Altitude Diagram, or CFAD). Figure 5a shows the reflectivity-height histogram for the surface (vertically pointing W-band) radar,
285 followed below by simulated histograms using the MG3 (Figure 5c) and IFS (Figure 5e) microphysics. The reflectivity data




Figure 3. TOA SW albedo from CERES SYN as a function of in-cloud liquid water path (LWP) for low-cloud periods during daylight (local time from 9 to 16 hours) from CERES Observations (purple cross), MG3 simulation (orange circles) and IFS simulation (green diamond). Legend shows the mean/median LWP.




Figure 4. Hourly mean reflectivity from (a) observations and SCM simulations with (b) MG3 and (c) IFS microphysics.

shown here are not hourly-averages. Rather the observational surface radar data has a temporal resolution of about 12 seconds (and so represents a horizontal grid scale less than one km), while the radar simulator reflectivities depend on the assumed model sub-grid distribution (more on this below). The surface radar histogram has very few detections below (to the left of) the minimum detectable signal (MDS) of the radar (violet line). For reference, the surface radar MDS is also plotted on the model simulated histograms (Figure 5c,e). The model simulations contain a large occurrence of hydrometeors above 5 km that the surface radar would be unable to detect due to the reduction of the MDS with height (violet line in Figure 5a,c,e). To a degree, this detection limit explains the larger occurrence of low-reflectivity hydrometeors in the model simulations (noted in connection with the time series in Figure 4). The IFS microphysics produces more low and high reflectivity, high-altitude cloud than the MG3 microphysics (Figure 5c). While the surface radar does not help to evaluate the models in this regard, we note that earlier it was found that the IFS microphysics produces too little outgoing LW flux (during periods with deeper clouds) while the MG3 microphysics compared well to the CERES SYN product (Figure 2c). This suggests that the MG3 microphysics is likely better representing this high-altitude cloud, likely because the MG3 two-moment scheme allows for an evolution of the size distribution of ice crystals, which better represents variable sedimentation and precipitation processes for ice.

Above (to the right of) the surface radar MDS (the violet line), both simulations (Figure 5c,e) produce a higher occurrence of hydrometeors than observed (Figure 5a): note the many orange and red colors in the simulated reflectivity-height histograms that are absent in the observations. This is also demonstrated in Figure 5b, which displays the vertical profile of hydrometeor occurrence counting only detections above the radar MDS. In addition to attenuation, the occurrence of hydrometeors above the MDS is strongly affected by the model sub-grid representation. Specifically, in the model radar simulator, clouds and precipitation are assumed to be uniformly distributed in the horizontal (e.g., liquid and ice water content do not vary horizontally) and to have the same fractional coverage when both exists (meaning the precipitation fraction equals cloud fraction). The cloud fraction is given by the largest of (i) the model prognostic cloud fraction, (ii) the stratiform precipitation fraction, or (iii) a fixed value of 10% representing convective precipitation. So for example, if the model prognostic cloud fraction is 78% and convective precipitation is present (at a given vertical level) then both the cloud and precipitation are assumed to cover 78% of the region (at the given vertical level) and the amount of "in-cloud" condensate will be equal to the grid-mean amount divided by 0.78. This uniform horizontal distribution has a significant affect on the simulated radar reflectivities. An analysis by Hillman et al. (2018) found these approximations cause the simulated radar reflectivity histograms to narrow around a "characteristic curve" where the reflectivity associated with precipitation most frequently occurs. In Figure 5a, the characteristic curve for the observations is depicted by the solid black line. The effect of the model assumption is especially evident near the surface, below about 2 km. Here the surface radar shows two distinct modes, a low reflectivity mode with a peak near -25 dBZe and a high reflectivity mode with a peak near 0 dBZe. This is also clear in Figure 5f (black line), which plots the mean observed reflectivity distribution below 2 km for both the observations and the simulations. The presence of these two modes is typical of most regions (Marchand et al., 2009) and is associated with non-precipitating (low reflectivity) cloud (left) and precipitation (high reflectivity, right). On a minor note, these modes are much less distinct over the Southern Ocean in the spring through fall, when there is more frequently drizzle, as well as frozen (snow) and mixed-phase precipitation with intermediate reflectivity factors (not shown). The simulations (especially with the IFS microphysics) show very little low reflectivity cloud near the

surface (Figure 5f). This is because some precipitation is almost always present in the simulations and even a small quantity of precipitation can generate a significant radar reflectivity. Thus while the model simulated total hydrometeor occurrence near the surface is only slightly larger than observed (Figure 5b), the assumption that precipitation (when present) is coincident with cloud results in a large overestimate in the occurrence of reflectivity factors larger than about -20 dBZe and an underestimate below -20 dBZe.

Of course, one could make a completely different assumption about the horizontal distribution of condensate and about the co-occurrence of cloud and precipitation (i.e. not maximize the horizontal cloud and precipitation overlap) in the radar simulator, and thereby obtain a simulated histogram that more closely matches the observations. But doing so only in the context of the simulator and without accounting for this sub-grid variability in a consistent way throughout the microphysics scheme is problematic. In our view, the point of using a radar simulator in the present analysis is to help evaluate the model physics, rather than trying to predict what a radar might observe. While in some sense the conclusion that cloud-and-precipitation shouldn't be maximally overlapped is obvious; on the positive side, the analysis shows the radiative effect of clouds in the simulations is reasonable (Figure 2) and also points to the potential value of these radar data in helping develop more sophisticated microphysics schemes that can represent sub-grid variability (and such schemes would nominally be better able to account for cloud-aerosol-precipitation interactions).

Returning to the topic of the characteristic curve, Hillman et al. (2018) suggest that while the distribution of reflectivities around the curve is sensitive to the sub-grid distribution, the position of the characteristic curve is not very sensitive to this representation. For comparison purposes the observed characteristic curve (shown by the solid black line) is plotted on top of the simulated histograms in Figure 5c,e, with the simulated characteristic curves given by the solid white lines. With the IFS microphysics (Figure 5e), there is good agreement between the simulated and observed characteristic curves between 1 and 5 km, suggesting that typical modeled precipitating ice water content (and the underlying parameterized/fixed particle size in the one-moment scheme) are quite reasonable. With the MG3 microphysics (Figure 5c), on the other hand, the characteristic curve is shifted to smaller reflectivity between 1 and perhaps 4 km. As we will see in the next section, the mean ice water content in the MG3 and IFS experiments are similar, and the difference in the characteristic curve is likely due in some combination to the additional attenuation caused by having too much liquid water in MG3 or the particle size (which is not fixed) being too small. In both experiments, below 1 km the characteristic reflectivity is below (left of) the observations and there is a large occurrence of reflectivity values between -15 and -5 dBZe relative to the observations. This is consistent with the early discussion of precipitation being spread over too wide an area (too frequent and consequently too light). Above 5 km, the MDS is likely affecting the observed characteristic curve in the observations and shouldn't be compared directly with the model, but as noted above, IFS clearly has larger volume of ice and a higher occurrence of smaller reflectivity factors.

3.2 Model Simulation Differences

In this section we explore the differences between the MG3 and IFS microphysics schemes in more detail. This SCM framework provides a good platform for a detailed analysis of two very different microphysical schemes in a large scale model. We start with examining the difference in hydrometeor profiles and timestep sensitivity of the two schemes (Section 3.2.1). We then look




Figure 5. Radar reflectivity-height histograms from (a) surface (vertically pointing W-band) radar, and from simulations using a radar-simulator with the (c) MG3 and (e) IFS microphysics. Colors indicate the fraction of time that hydrometeors are detected at the specified altitude and reflectivity. Histogram bins are 0.5 km x 5 dBZe. Violet line shows the Minimum Detectable Signal (MDS) of the surface radar. Black and white lines show the characteristic curve (see text) for observations (black, same in all panels) and simulations (white). (b) profile of total hydrometeor occurrence, including only detections above the MDS (violet line). (d) Mean distribution of radar reflectivity for hydrometeors above 2 km altitudes (note occurrence in OBS is strongly affected by MDS below about -20 dBZe) . (f) Same as (d) but includes only hydrometeors below 2 km (occurrence in OBS is unaffected by MDS above about -35 dBZe)

at the differences between process rates in MG3 and IFS microphysics (Section 3.2.2). Being able to compare the processes in detail is a unique feature of the SCM forced thermodynamics framework. Finally we look at configuration differences between the MG3 and IFS schemes that can be adjusted (Section 3.2.3), including some additional MG3 process sensitivity tests.

3.2.1 Mean Hydrometeor Profile and Timestep Sensitivity

360 Figure 6 shows the mean profiles of key hydrometeors for the IFS and MG3 microphysical schemes. There is overall more supercooled liquid water (SLW, Figure 6A) in the MG3 scheme than the IFS, consistent with the LWP discussion in Section 3.1.2, with almost a factor of two more at altitudes below 750hPa, but less SLW above. Sea surface temperature is about 5°C and the average freezing level is about 900hPa or 0.8km. Total ice (ice, snow and graupel, Figure 6B) is quite similar between the two schemes, as is the total precipitation (rain, snow and graupel, Figure 6C). A common problem is a sensitivity of microphysical
365 processes to the length of the timestep (Gettelman et al., 2015; Santos et al., 2021), particularly for sedimentation, with various different numerical solutions proposed with different accuracy and stability. Figure 6 also shows the sensitivity of the mean profile of supercooled liquid water, total ice and precipitation, as well as liquid and ice number concentration (for MG3) to the timestep used in the MG3 and IFS microphysics schemes. A range of timesteps, specifically 60, 225 and 900s are shown for each microphysics scheme. The default used in the baseline simulations described above is 225s. The IFS microphysics is
370 insensitive to timestep, due to the fact that it has all the microphysical process calculations inside an implicit sedimentation loop for each model level. As the CESM2 version of the MG3 scheme has a significant timestep sensitivity (not shown), a modification was made to change the sedimentation numerics from an explicit to an implicit calculation (as discussed in section 3.2.3). It is the modified MG3 scheme that is used here and Fig. 6 shows it is quite stable with respect to timestep, with some differences in peak SLW (Figure 6A) and in ice number (Figure 6E), particularly for the long (900s) timestep.

375 3.2.2 Microphysical Process Rates

The two schemes have many similarities but also significant differences in the formulation of the microphysical process rates, and in this section we explore the balance of individual process rates for each hydrometeor (cloud liquid, ice, rain and snow) in both schemes. We do not break this down by regime (e.g. low clouds), but look at process rates for all conditions. A longer descriptive name for each process rate (and which scheme uses it) is contained in Table A1 (Liquid and Ice) and Table A2
380 (Rain and Snow).

Figure 7A and B illustrate different process rates that sum to give the total sink (loss) tendency for cloud liquid water. These are the microphysical process rates after the cloud scheme has produced large scale condensation by removing supersaturation (i.e. the source term for cloud condensate which therefore has the same formulation for both simulations). MG3 represents more microphysical processes than the IFS due primarily to the more detailed representation of ice nucleation and the ad-
385 dition of graupel. The mixed phase vapor deposition (or Wegener-Bergeron-Findeisen or ‘Bergeron’) process for the growth of ice particles is the most important loss process for supercooled liquid water droplets at upper levels in both schemes. At lower altitudes the collision (accretion) of snow with liquid droplets (‘Accre Snow’) dominates and rain collision-coalescence (‘Accretion’) below that at warmer temperatures. The major difference for liquid water is the vapor deposition process onto




Figure 6. Vertical time averages of A) Supercooled Liquid Water, B) Total Ice (ice+snow+graupel), C) Total Precipitation, D) Liquid Number Concentration and E) Ice Number concentration, for Reference IFS simulations (dotted) and MG3 microphysics simulations (solid) at different time steps.

390 snow (‘Berg Snow’) in MG3; a process that is not active in the IFS. The IFS accretion onto snow compensates for the missing process to some extent, but still ends up less than the sum of the accretion and the Bergeron vapor deposition onto snow in MG3 at altitudes above 750hPa. There are several other terms that are different with relatively small magnitudes, such as the autoconversion of liquid to rain, rime splintering and graupel collection of liquid.

Ice process rates have similar dominant processes in both microphysics schemes (Figure 7C,D), with a source from vapor deposition (‘Bergeron’) offset by a loss of ice through autoconversion to snow (‘Autoconv $Q_i > Q_s$ ’). However, the Bergeron process is much more active at lower altitudes/warmer temperatures for MG3 compared to the IFS (as also seen in Figure 7A,B), with a similar difference in magnitude for ice autoconversion to snow which compensates. The more active Bergeron process in MG3 is likely due to the larger supercooled liquid water present at these lower levels (Figure 6A). In MG3 (Figure 7D), there is also loss from accretion of ice onto snow (‘Accret $Q_i > Q_s$ ’) throughout the depth of the cloud from 300 to 800 hPa, and a source due to deposition onto ice (‘Ice Sub/Dep’) at high altitudes around 400 hPa. The ice sedimentation in the IFS




Figure 7. Profiles of (A,B) cloud liquid water and (C,D) cloud ice process rates, A,C) for the IFS microphysics and B,D) for MG3. Process rates are described in Table A1.

400 shows a small loss in the upper part of the cloud and gain lower down, whereas the net redistribution is much smaller in MG3; a difference likely due to the different sedimentation numerics in the two schemes.

Rain process rates are illustrated in Figure 8A,B. Rain sedimentation ('QR SED') is the major loss for precipitation, larger in the IFS, and snow melting ('Melt Snow') and accretion of liquid cloud droplets ('Accre') are the main rain source terms. Snow melting is more active nearer the surface in the MG3 scheme. There is not much rain evaporation in the PBL which is
405 consistent with large reflectivity in the lowest kilometre (Fig 5e). This is due to the specific rain evaporation assumptions in the IFS and modified MG3 (see section 2.3). There is even less rain evaporation in MG3 due to the deeper melting layer, and the two-moment scheme allowing rapidly falling rain drops rather than slower falling drizzle.

There are some differences in snow process rates (Figure 8C,D). The general balance at upper levels is a sedimentation loss ('QS SED') and an ice autoconversion ('Autoconv $Q_i > Q_s$ ') source, representing aggregation of ice crystals. Snow fall speeds
410 are fixed in the IFS and set to be the same in MG3, making the sedimentation rate similar. MG3 also has a source from the vapor deposition onto snow ('Berg > Snow'), and loss from snow sublimation/evaporation ('Evap') (larger in MG3 than IFS). At lower levels, the sedimentation source term for snow falling from above ('QS SED') is balanced by a melting loss term to rain ('Melt Q_s '), which is just the inverse of the melting source term for rain (Figure 8A,B). The shallower melting layer in IFS is also evident.

415 In summary, both microphysics schemes produce a largely similar balance of process rates. Given that some of the microphysical process formulations are similar between the two schemes and that there is some convergence that has been imposed by adjusting the MG3 scheme towards the IFS (see Section 2.3), this is perhaps not surprising. However, there are also some differences in the formulations that result in different hydrometeor profiles, particularly for the radiatively important supercooled liquid water. As shown in Figure 6, the mean profiles of ice and snow are very similar between the two schemes (panels
420 B,C). There is clearly compensation between some of the processes, such as the higher Bergeron process rate for ice growth and snow accretion rates in the IFS which compensates for the lack of a Bergeron process for snow in the IFS compared to MG3 (Figure 7A,B). This compensation results in a similar supercooled liquid water profile at altitudes above 750hPa, apart from for the MG3 900s timestep simulation, (Figure 6A). However, at altitudes below 750hPa, the profile of supercooled liquid water is more than a factor of two different, with much higher values in MG3 compared to the IFS (Figure 6A). The higher
425 cloud liquid accretion rates for both snow and rain below 750hPa in the IFS (Figure 7A,B) are likely leading to the lower SLW in the IFS, which agrees more closely with the observations (Figure 2i,j). At lower levels below 850hPa, there are also differences in rain in the precipitation profile (Figure 6C). For sedimenting precipitation, differences can come from higher in the column, but given that the ice and snow hydrometeor profiles are very similar, differences in the low level precipitation must therefore be due to the local impact of processes, particularly the melting rate of snow to rain, evaporation of rain, and
430 differences in the rain fall speed.

3.2.3 Modifications and Sensitivity

In order to further understand the impacts of the changes to the MG3 scheme and differences with the IFS, a series of sensitivity experiments are performed changing individual aspects of the MG3 microphysics. The first set of changes (Figure 9)




Figure 8. Profiles of (A,B) rain and (C,D) snow process rates, A,B) for the IFS microphysics and C,D) for MG3. Process rates are described in Table A2.

investigates the impact of the initial modifications from the original CAM6 version to the base MG3 scheme used for all the
435 simulations described so far, as described in Section 2.3. These changes were made for this study to remove some of the most
obvious differences due to numerical implementation and basic threshold and other assumptions, to allow a more meaningful
comparison of the process rates between the two microphysics schemes. The second set of changes investigates the sensitivity
to assumptions that affect several of the important microphysical process rates (Figure 10).

The sensitivity experiments are summarized in Table 1 and are described in more detail below. Figures 9 & 10 show the
440 impacts on time-averaged vertical profiles for various quantities and the impacts on time-averaged single level quantities are
shown in Figure 11. For reference, the figures also show the ‘IFS’, the modified MG3 used for this study (‘MG3 Base’), on
which all of the sensitivity experiments are based, and the original ‘MG3 CAM6’ settings used with the MG3 microphysics
in the CAM6 atmospheric component of the coupled climate model CESM2. In general, the MG3 CAM6 settings are outliers
for various quantities and far away from the IFS and ‘MG3 Base’ results. It is therefore notable that significant differences can
445 occur from the numerical implementation of the sedimentation and microphysical threshold assumptions.

(i) Sensitivity to MG3 modifications from CAM6

We first assess changes to the fall speed of hydrometeors (‘Fallspeed’) and numerical formulation of the sedimentation of
precipitation (‘Explicit Fall’) in MG3. The ‘Fallspeed’ sensitivity test removes a numerical fix for zero fall speed in the column
450 when no precipitation is initialized there in MG3, and ‘Explicit Fall’ reverts the change from an implicit sedimentation of
precipitation calculation, which is more stable across time-steps, back to the explicit scheme in CAM6. In addition, the IFS
and MG3 Base have constant fall speeds for ice and snow, which are reverted to the MG3 CAM6 fall speed formulation (fall
speed varies as a function of hydrometeor size) in the ‘Sediment’ test. The ‘Fallspeed’ and ‘Explicit Fall’ slightly decrease the
total amount of ice (Figure 9B) and the ‘Sediment’ change increases the (supercooled) cloud liquid water at altitudes above
455 700 hPa (Figure 9A). A reversion to allow the immediate evaporation of sedimenting condensate (‘Evapsed’) has negligible
impact.

A series of changes to harmonize evaporation was also conducted (‘Old Evap’) as the IFS uses a relative humidity threshold
of 90% before evaporation starts, and the IFS has a similar evaporation formula as MG3 but is multiplied by a factor of 0.3.
The rain freezing threshold (‘Rain Freeze’) has small effects on total precipitation (Figure 9C) near 850hPa in the temperature
460 range just below -5°C . Finally the IFS and MG3 Base do not have a threshold RH for ice nucleation (RH for IN) whereas
MG3 CAM6 has an RH threshold of 105% with respect to ice for ice nucleation to occur. Reverting the RH threshold for
ice nucleation (RH for IN) increases liquid mass (Figure 9A) and number (Figure 10D) near 600hPa and reduces ice number
(Figure 9E) significantly between 300–600hPa by making it harder to form ice. Each of these experiments is a single reversion
experiment that removes these elements from MG3 base in IFS to get back to MG3 CAM6. However, the combination of these
465 effects is important. The test ‘Old Rain’ combines the Rain Freeze, Sediment and Fallspeed tests, and it is this combination
that reduces the unrealistic increase in precipitation (rain) mass near the surface in MG3 CAM6 (Figure 9C).

The implicit formulation for sedimentation as in MG3 Base has lower LWP (Figure 11A) than MG3 CAM6, and shortwave
(Figure 11E) and longwave (Figure 11F) downward radiation more similar to the IFS which also has an implicit formulation.

Table 1. Sensitivity tests applied to MG3 simulations

Name	Description
MG3 Base	Base MG3 code
MG3 CAM6	MG3 code with CAM6 settings
Changes to revert MG3 Base to MG3 CAM6	
Fallspeed	Remove fallspeed correction that ensures non-zero sedimentation
Explicit Fall	Revert from implicit to explicit fallspeed
Sediment	Variable snow and ice fall speed
Evapsed	Allow evaporation of sedimenting condensate.
Old Evap	Original RH threshold for evaporation and un-scaled evaporation
Rain Freeze	Revert rain freezing temp to -40°C
RH for IN	Use 105% RH over ice threshold for ice nucleation (rather than no threshold)
Old Rain	Combines Rain Freeze, Fallspeed and Sediment
Other Sensitivity Tests	
Ac3Au1.5	Sub-grid heterogeneity scaling for Accretion (x3) and Autoconversion (x1.5) following IFS
IFS KK	Use original constants for Autoconversion following IFS
Au in Ac	Allow accretion to see recently autoconverted rain
No Graupel	Graupel off (riming forms snow not graupel)
No Meyers	Removal of Meyers et al. (1992) ice nucleation
IN/5	Ice nucleation of 1 L ⁻¹ per timestep
IN*5	Ice nucleation of 25 L ⁻¹ per timestep
CCN*4	Liquid activation rate of 4x10 ⁴ m ⁻³ s ⁻¹
CCN/4	Liquid activation rate of 0.25x10 ⁴ m ⁻³ s ⁻¹
N=cnst	Constant number concentrations: N _c = 200 cm ⁻³ , N _i =50 L ⁻¹ , N _r =5 L ⁻¹ , N _s =1 L ⁻¹ , N _g =0.5 L ⁻¹

The largest difference between MG3 CAM6 and the MG3 Base is due generally to the different fall speed estimates (mostly affecting total ice) and rain freezing (affecting total ice and precipitation). Note that the SW downward radiation (Figure 11E) is generally inversely proportional to the LWP (Figure 11A) and to a lesser extent total cloud cover (Figure 11C). The spread in the LW downward flux is relatively small (6 W m⁻², Figure 11F), relative to the spread in SW downward flux (35 W m⁻², Figure 11E).

Finally, it is worth noting that supercooled liquid water above 700hPa is much higher in the MG3 CAM6 version (Figure 9A). This is due to the fall speed changes ('Sediment', which is also in the 'Old Rain' test) combining in a non-linear way with the change to the relative humidity used for ice nucleation ('RH for IN').

(ii) Additional MG3 process sensitivity tests




Figure 9. Vertical time averages of A) Supercooled liquid water, B) Total ice (ice+snow+graupel), C) Total precipitation, D) Liquid number concentration, E) Ice number concentration, for the MG3 base version (black solid) and various MG3 microphysics sensitivity tests as noted in the legend to differentiate MG3 and IFS microphysics. The reference IFS microphysics is included in A,B,C (red solid).

The second set of tests is performed with further modifications to the ‘MG3 Base’ code to understand its sensitivity relative to the IFS.

One of the most important aspects of the microphysics is the generation of precipitation. For liquid clouds, this is defined by the collision-coalescence process of interacting hydrometeors of different sizes. In bulk microphysics schemes like the IFS and MG3, this process has to be highly parameterized since the different sized drops are not explicitly represented. The parameterizations are for autoconversion (formation of rain from cloud water) and accretion (the removal of cloud water by rain). In MG3 and IFS, these parameterizations come from Khairoutdinov and Kogan (2000), a regression fit to a bin microphysical model. In MG3, the parameters for autoconversion were adjusted to better match recent observations, with resulting reduced sensitivity to droplet number, whereas in the IFS the original Khairoutdinov and Kogan (2000) parameters are used (‘IFS KK’). In the IFS there is also an enhancement factor of 1.5 for autoconversion and 3 for accretion (‘Ac3Au1.5’) representing the effects of subgrid heterogeneity of cloud and precipitation. We test the MG3 scheme with these factors. The scaling of autoconversion and accretion (‘Ac3Au1.5’) significantly reduces cloud water, as expected, (Figure 11A) and slightly




Figure 10. Vertical time averages of A) Supercooled liquid water, B) Total ice (ice+snow+graupel), C) Total precipitation, D) Liquid number concentration, E) Ice number concentration, for the MG3 base version (black solid) and various MG3 microphysics sensitivity tests as noted in the legend to understand MG3 sensitivity.

increases ice water path (Figure 11B), with corresponding increases in the downward SW radiation (Figure 11E). Changing MG3 to the IFS parameters for Khairoutdinov and Kogan (2000) ('IFS KK') has more of an impact higher up, reducing SLW at altitudes above 800hPa, but with the major impact seen as a large increase in ice number (Figure 10E). This seems to result from a conversion process from liquid (possibly freezing rain) to ice at supercooled temperatures. These settings produce a simulation with LWP and surface SW downward radiation closer to observations (and IFS microphysics), largely due to lower LWP (Figure 11A). We also test the MG3 scheme with a change that allows the autoconverted rain to be immediately used by the accretion process ('Au in Ac'), which slightly enhances accretion but has little overall effect on the simulations. The CAM6 settings result in the highest LWP (Figure 11A), which does enhance the surface LW down (Figure 11F) to be closer to the observations (but note the difference is only a few Wm^{-2}), but produces much too small surface SW down (Figure 11E).

Two changes to the mixed phase have substantial impacts. Turning off the prognostic graupel (rimed ice particles) in the MG3 simulation results in a significant increase in ice mass (Figure 10B) and precipitation (Figure 10C). This is due to the removal of a loss process for rain onto graupel, which now occurs only onto snow, a less efficient process, and leads to higher




Figure 11. Time averages of A) Liquid water path, B) Ice water path, C) Cloud fraction, D) Surface stratiform rain rate, E) Surface SW down and F) Surface LW down for MG3 microphysics sensitivity tests as noted on the plot. MG3 base case in black. Reference IFS plotted in Red. MICRE observations for LWP, and SW and LW radiation in Blue.

snow mass. Removing the Meyers et al. (1992) mixed phase ice nucleation significantly reduces ice number (Figure 10E) and IWP (Figure 11B), as well as supercooled liquid at higher altitudes (Figure 10A).

505 Next, we examine the role of number concentrations for liquid and ice in the simulations. Because MG3 typically runs with an aerosol model, it requires the activation of liquid drops and ice crystals to set the hydrometeor number concentrations, unless they are set to a constant. In the IFS there is no explicit representation of activation of aerosol, so for the MG3 simulations a constant rate of cloud condensation nuclei (CCN) and ice nuclei (IN) production is defined. The ice and liquid mass and number concentrations are sensitive to these rates and this will naturally create differences between MG3 and IFS microphysics. 510 Indeed, increasing the CCN production rate ('CCN*4') results in many more (Figure 10D) and hence smaller cloud water drops, increasing SLW (Figure 10A) and averaged LWP (Figure 11A), and reducing downward SW (Figure 11E). The opposite effects are seen for reductions in the CCN production rate ('CCN/4'). CCN and IN do vary by an order of magnitude over the S. Ocean (McFarquhar et al., 2021). There are also tests performed adjusting the constant for the rate of production of ice nuclei (IN/5 and IN*5). More IN slightly increases IWP (Figure 11B,C) with opposite effects for reducing IN.

515 We also perform a simulation with fixed number concentrations for all species in the MG3 scheme (rather than fixed production rates), as specified in Table 1. Fixed number concentration ignores any drop or ice nucleation, resets the prognostic number to a set value every timestep, and uses this throughout the microphysics. Fixed number concentrations ('N=cnst') have a substantial effect on the simulations if they are not set to reasonable values. Here the number concentrations have been adjusted to produce similar number concentrations as the prognostic MG3 Base simulation. Nonetheless, even with similar LWP, average 520 liquid number concentrations are decreased (Figure 10D) similar to the CCN/4 experiment. Note that the average number is lower than the fixed drop number due to averaging and number depletion.

In summary, in MG3 there is sensitivity of LWP and radiation to the fall speed. The autoconversion and accretion processes that dominate rain formation and remaining LWP are also important for surface radiation. Changes to the activation of cloud drops (CCN) also affects cloud water and rain partitioning, and hence LWP and radiation. Finally, the ice phase is also im- 525 portant, with the inclusion of graupel changing the loss process of rain to make it less efficient and increases the snow mass (included in IWP) significantly. Ice nucleation and mixed phase ice nucleation ('No Meyers') is important for ice water path and surface radiation. Note that none of the experiments reduces the MG3 LWP to the level of the Obs or IFS (Figure 11A), but with further combination of some of these tests, like the conversion and accretion scaling, fixed number concentration at low number or large size and/or reduced CCN, the MG3 LWP could be further reduced to match the IFS more closely.

530 4 Summary

Simulations are performed with the IFS single column model with two "operational" microphysics parametrizations (the one-moment IFS scheme used at ECMWF for NWP, and the two-moment MG3 scheme used in the CAM6 climate model). Both are able to reproduce many of the characteristics, evolution and time-mean variability of Southern Ocean clouds observed during the MICRE campaign at Macquarie Island during the period January–March 2017, in a regime with significant amounts of 535 supercooled liquid cloud, and in a maritime location representative of the broader Southern Ocean.

Assessing the fidelity of GCM clouds with respect to observations requires dealing with uncertainty in the observations and the model. Observations have their own biases, such as the minimum detectable signal and loss of signal in precipitation for the radar (in addition to attenuation). The IFS simulation results (for both microphysical schemes) when compared to observations are sensitive to the model assumptions, such as the definition of cloud cover, and the sub-grid treatment of clouds and hydrometeor size distributions. We have carefully considered these uncertainties in performing these evaluations.

Surface radiation is well represented compared to the ground-based radiometer and satellite data sets with radiative flux biases within or close to the uncertainty of the observations in both the LW and SW. In particular, in the shortwave solar spectrum, there is no “too few/too bright” problem that has previously been seen in some global models (Wall et al., 2017; Kuma et al., 2020). Hydrometeor occurrence (cloud fraction) is slightly higher than observed when radar simulator output is compared to observations.

There are large uncertainties in the observed surface (passive microwave) LWP retrievals, largely due to the presence of precipitation, but for low level non-precipitating clouds, the surface radiometer values at Macquarie Island are consistent with satellite estimates based on geostationary (Himawari-8) and MODIS polar orbiting satellites. Mean LWP for the IFS microphysics compares well to both surface and satellite LWP retrievals for low clouds, while mean LWP from the MG3 microphysics is too large. An analysis of the distribution of LWP shows that the MG3 microphysics produces more high LWP events than the observations and IFS simulations. Median LWP values for MG3 are comparable to the IFS and the observations. Because the radiative effects of condensed water saturate when the LWP is larger than $\sim 400 \text{ g m}^{-2}$ (cloud albedo does not change much), the large LWP events do not translate into a large SW radiation bias in the MG3 scheme.

A comparison of radar reflectivity with the vertically pointing radar at Macquarie Island shows the simulations capture the overall characteristic shape of the reflectivity-height distribution, although the mean frequency of occurrence of hydrometeors (with reflectivity higher than the surface radar minimum detectable signal) is too high, at least above the boundary layer. In general, the simulations contain a higher occurrence of reflectivity factors larger than -15 dBZ compared to the observations. While some of this difference is expected due to attenuation of the observed radar reflectivity that results from rain on the radar radome, the difference also occurs for periods when there is no surface rain. Rather this difference in the distribution of reflectivity factors is largely driven by the model assumption that when precipitation is present, it occupies the same area (fraction) as that predicted for cloud. A better model sub-grid representation of cloud and precipitation could reduce biases in the radiation and addresses this “too frequent, too light” precipitation problem. While this problem is certainly not unique to the Southern Ocean (Stephens et al., 2010), it is perhaps especially pertinent because of the high occurrence of light precipitation in the region. This also highlights that this too light precipitation issue may be related to sub-grid assumptions in models, not a fundamental flaw in the cloud physical process rates.

The single column model is a partially constrained testing environment and for long simulations requires forcing to keep the temperature, humidity and wind profile from drifting too far from the observations. However, the cloud and microphysical response to the forcing is not constrained and thus it is a useful platform for assessing different microphysical schemes. In a test without any relaxation of the temperature, the PBL deviates from observations, and significant SW and LW radiation biases emerge on the order of tens of W m^{-2} with both the IFS or MG3 microphysics. While advective forcing of wind and

temperature is included in the SCM, advection of hydrometeors is not included and could create errors. However, the timescale for microphysical processes is shorter than an advection time scale (for a 10m/s wind and a 50km GCM gridbox, this implies a 2500s advection timescale, but the microphysical timesteps are 225s), and means the issue should be a small one. The good reproduction of observed cloudiness (Figure 4) and LWP (Figure 1) indicates that this is the case.

575 A series of sensitivity tests modifying the numerical and physical representation of various cloud and precipitation processes in the MG3 scheme are performed to understand the differences from the IFS microphysics scheme. When MG3 is modified to use formulations similar to the IFS (the 'MG3 Base' microphysics scheme), the average ice, snow and rain hydrometeor profiles are remarkably similar, with the main difference being double the amount of supercooled liquid water on average in MG3. The detailed process rate analysis illustrates that the schemes have a different discretization of process rates (with MG3 being
580 more complex), and that the phase transition between liquid and ice, as well as precipitation formation and sedimentation are the major cause of differences between the schemes, despite significant compensation between processes (e.g., the IFS has a higher vapor deposition for ice growth which compensates for a lack of vapor deposition onto snow). The sensitivity simulations confirm that differences in the autoconversion to rain and accretion onto rain loss processes for liquid water at sub-freezing temperatures are the primary cause of the differences in LWP (which is mostly SLW), and the MG3 scheme could
585 be adjusted to enhance the liquid loss process and reduce the LWP in deeper clouds. Finally, for cloud liquid and cloud ice formation, there are clearly large sensitivities to the specification of the CCN and IN production rates. Although tuning these in the SCM can bring closer agreement, a full comparison in the 3D model with interactive aerosol activation is required for the impact to be fully assessed.

5 Conclusions

590 The SCM is a good framework for testing cloud physics. Part of getting clouds and cloud radiative effects correct is getting the PBL structure right and the correct coupling of the thermodynamics to the cloud physics, as well as the microphysical processes themselves. With constrained dynamics, the cloud radiative effects stay close to observations and we do not see the large scale biases seen in many global models in the Southern Ocean, indicating that coupled dynamics and microphysical processes are important.

595 The required complexity of microphysical parametrization is a key question for future development of atmospheric models, and one of the motivations for the comparison here of the one-moment (mass) IFS with the higher complexity two-moment (mass and number concentration) MG3 scheme including graupel. Representing a graupel (rimed ice) hydrometeor in MG3 has a small impact with more precipitation and less SLW than without graupel, but there was not a significant amount of graupel present in the type of clouds at this location and a larger difference would be expected in regimes with more deep convection.
600 Regarding the two-moment versus one-moment representation, it is difficult to disentangle the impacts on the mean fields of the two-moment representation in MG3 from the differences in microphysical process formulation between the schemes as there can be compensation between processes, and numerical formulation and different tunings of microphysical process rates can dominate. However, the representation of supercooled liquid is better in the two-moment scheme. Despite similar ice and

precipitation mass profiles, there are differences in the radar reflectivity profiles which are at least partly due to the additional
605 degrees of freedom from the representation of particle number concentration in MG3. The two moment scheme as a result
likely produces a slightly better reproduction of observed radiative fluxes.

These results have several important messages for 3D modeling, which should be tested in future work. The correct PBL
structure (the temperature and stability profile) is necessary for getting clouds and radiation right. There is little bias in the
constrained SCM. The major differences between microphysics schemes lie in precipitation formation, sedimentation of hy-
610 drometeors and freezing processes. The two moment scheme is sensitive to the number of condensation and ice nuclei. The
representation of supercooled liquid is better in the two-moment scheme. The additional degrees of freedom from the two
moment scheme representation of variable particle number concentration improves the simulation of reflectivity and likely
produces a slightly better reproduction of observed radiative fluxes. Finally, the sub-grid assumptions for cloud and precipi-
tation impact both the ability to compare to observations, as well as the model evolution itself for both one and two moment
615 schemes. One promise of higher resolution 'storm-resolving' models without fractional cloud and precipitation is to reduce the
need to make these assumptions and enable more direct evaluation against observations.

Code and data availability. Simulation output used in this manuscript, and specific code for MG3 microphysics in the IFS is available in a
zenodo archive. DOI:10.5281/zenodo.10037746 (<https://zenodo.org/records/10037746>). Data used for comparison is available through the
noted references in the text. All MICRE data are available from the Atmospheric Radiation Measurement (ARM) data archive (www.arm.gov),
620 a U.S. Department of Energy (DOE) Office of Science user facility managed by the Biological and Environmental Research Program. Specif-
ically, data from MICRE can be found using the data discovery tool for the "MCQ" site.

The IFS source code is available subject to a licence agreement with ECMWF. ECMWF member-state weather services and approved
partners will be granted access. The OpenIFS single column model code is also available for educational and academic purposes via an
OpenIFS licence (see <http://www.ecmwf.int/en/research/projects/openifs>).

625 **Appendix A: Appendix**

Author contributions. AG, CCC, RF and MF developed and evolved the MG3 microphysics code for the open IFS. AG performed the
simulations. Analysis and figure development were conducted by AG, RM, RF and MF. AG wrote the manuscript with editorial input and
sections written by all.

Competing interests. The authors state there are no competing interests

Table A1. Microphysical Process Rates: Liquid and Ice

Name	Scheme Used	Description
Liquid		
Liq Sed	IFS, MG3	Sedimentation of Liquid
Cond Resid	MG3	Condensation Residual
Imm Frz	MG3	Immersion Freezing of Liquid
Cnt Frz	MG3	Contact Freezing of Liquid
Bergeron	IFS, MG3	Liquid Vapor Deposition onto Ice
Melt I	IFS, MG3	Melting of Ice
Homo Frz	IFS, MG3	Homogenous Freezing of Liquid
Rime Splint	MG3	Rime Splintering
Berg Snow	IFS, MG3	Liquid Deposition onto Snow
Accret	IFS, MG3	Accretion of Liquid onto Rain
Accre Snow	IFS, MG3	Accretion of Liquid onto Snow
Autoconv	IFS, MG3	Autoconversion of Liquid to Rain
Coll Graupel	MG3	Collection of Liquid by Graupel
Coll Snow	MG3	Collection of Liquid by Snow
Mult Qc>Grau	MG3	Ice multiplication of liquid to Graupel
Ice		
Ice Sed	IFS, MG3	Sedimentation of Ice
Sed Resid	MG3	Ice sedimentation residual
Ice Sub/Dep	MG3	Ice Sublimation and Deposition
Imm Frz	MG3	Immersion Freezing to Ice
Cnt Frz	MG3	Immersion Freezing to Ice
Bergeron	IFS, MG3	Vapor Deposition onto Ice
Melt I	IFS, MG3	Melting of ice
Homo Frz	IFS, MG3	Homogenous Freezing to Ice
Rime Splint	MG3	Rime Splintering of Ice
Autoconv Qi>Qs	IFS, MG3	Autoconversion of Ice to Snow
Accret Qi>Qs	MG3	Accretion of Ice to Snow
Mult Qc>Grau	MG3	Ice Multiplication Liquid to Graupel
Mult Qr>Grau	MG3	Ice multiplication Rain to Graupel
Dep Nuc	MG3	Deposition Nucleation of Ice
Accre Rain>Ice	IFS, MG3	Accretion of Rain to Ice

630 *Acknowledgements.* The Pacific Northwest National Laboratory is operated for the U.S. Department of Energy by the Battelle Memorial Institute under contract DE-AC05-76RL01830. This work was supported in part by the U.S. Department of Energy Atmospheric System

Table A2. Microphysical Process Rates: Rain and Snow

Name	Scheme Used	Description
Rain		
QR SED	IFS, MG3	Rain Sedimentation
Accre	IFS, MG3	Accretion to Rain
Auto	IFS, MG3	Autoconversion to Rain
Accre Rain	IFS, MG3	Accretion of Rain to Snow
Frz Rain > Snow	IFS, MG3	Rain Freezing
Mult Qr>Grau	MG3	Ice Multiplication Rain to Graupel
Coll Qr> Grau	MG3	Collection of Rain by Graupel
Coll Qr by Qs>Grau	MG3	Collection of Rain by Snow
Accre Rain>Ice	IFS, MG3	Accretion of Rain by Ice
Evap Rain	IFS, MG3	Evaporation of Rain
Melt Snow	IFS, MG3	Melting of Snow to Rain
Snow		
QS SED	IFS, MG3	Snow Sedimentation
Accrete Rain	IFS, MG3	Accretion of Rain to Snow
Freeze Rain > Snow	IFS, MG3	Freezing of Rain to Snow
Berg > Snow	IFS, MG3	Liquid Deposition to Snow
Accret Liq	IFS, MG3	Accretion of Liquid to Snow
Evap	IFS, MG3	Evaporation (Sublimation) of Snow
Autoconv Qi>Qs	IFS, MG3	Autoconversion of Ice to Snow
Accre Qi>Qs	MG3	Accretion of Ice onto Snow
Accre Qr>Qs	IFS, MG3	Accretion of Rain onto Snow
Qg coll Qs	MG3	Graupel Collection of Snow
Melt Qs	IFS, MG3	Melting of Snow to Rain
Vap Dep Snow	MG3	Vapor Deposition onto Snow

Research program through Grant DE-SC0016225. The National Center for Atmospheric Research is supported by the U.S. National Science Foundation.

References

- 635 Abel, S.J. and Boutle, I.A.: An Improved Representation of the Raindrop Size Distribution for Single-Moment Microphysics Schemes, *Quarterly Journal of the Royal Meteorological Society*, 138, 2151–2162, 2012.
- Ahlgrimm, M. and Forbes, R.: Improving the Representation of Low Clouds and Drizzle in the ECMWF Model Based on ARM Observations from the Azores, *Monthly Weather Review*, 142, 668–685, 2014.
- Balsamo, G., Beljaars, A., Scipal, K., Viterbo, P., van den Hurk, B., Hirschi, M., and Betts, A. K.: A Revised Hydrology for the ECMWF
640 Model: Verification from Field Site to Terrestrial Water Storage and Impact in the Integrated Forecast System, *Journal of hydrometeorology*, 10, 623–643, 2009.
- Bechtold, P., Kohler, M., Jung, T., Doblas-Reyes, F., Leutbecher, M., Rodwell, M. J., Vitart, F., and Balsamo, G.: Advances in Simulating Atmospheric Variability with the ECMWF Model: From Synoptic to Decadal Time-Scales, *Quarterly Journal of the Royal Meteorological Society*, 134, 1337–1351, 2008.
- 645 Bechtold, P., Semane, N., Lopez, P., Chaboureaud, J.-P., Beljaars, A., and Bormann, N.: Representing Equilibrium and Nonequilibrium Convection in Large-Scale Models, *Journal of the Atmospheric Sciences*, 71, 734–753, 2014.
- Beljaars, A. C. M., Brown, A. R., and Wood, N.: A New Parametrization of Turbulent Orographic Form Drag, *Quarterly Journal of the Royal Meteorological Society*, 130, 1327–1347, <https://doi.org/10.1256/qj.03.73>, 2004.
- Bodas-Salcedo, A., Williams, K.D., Field, P.R., and Lock, A.P.: The Surface Downwelling Solar Radiation Surplus over the Southern Ocean
650 in the Met Office Model: The Role of Midlatitude Cyclone Clouds, *J. Climate*, 25, 7467–7486, 2012.
- Bodas-Salcedo, A., Mulcahy, J. P., Andrews, T., Williams, K. D., Ringer, M. A., Field, P. R., and Elsaesser, G. S.: Strong Dependence of Atmospheric Feedbacks on Mixed-Phase Microphysics and Aerosol-Cloud Interactions in HadGEM3, *Journal of Advances in Modeling Earth Systems*, 11, 1735–1758, <https://doi.org/10.1029/2019MS001688>, 2019.
- Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R.,
655 Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Versteijn, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001916, <https://doi.org/10.1029/2019MS001916>, 2020.
- 660 Doelling, D. R., Loeb, N. G., Keyes, D. F., Nordeen, M. L., Morstad, D., Nguyen, C., Wielicki, B. A., Young, D. F., and Sun, M.: Geostationary Enhanced Temporal Interpolation for CERES Flux Products, *Journal of Atmospheric and Oceanic Technology*, 30, 1072–1090, <https://doi.org/10.1175/JTECH-D-12-00136.1>, 2013.
- ECMWF: IFS Documentation CY46R1 - Part IV: Physical Processes, in: IFS Documentation CY46R1, no. 4 in IFS Documentation, ECMWF, <https://doi.org/10.21957/xphfxep8c>, 2019.
- 665 Fielding, M. D. and Janisková, M.: Direct 4D-Var Assimilation of Space-Borne Cloud Radar Reflectivity and Lidar Backscatter. Part I: Observation Operator and Implementation, *Quarterly Journal of the Royal Meteorological Society*, 146, 3877–3899, <https://doi.org/10.1002/qj.3878>, 2020.
- Forbes, R., Greer, A., Loniz, K., and Ahlgrimm, M.: Reducing Systematic Errors in Cold-Air Outbreaks, *ECMWF Newsletter*, pp. 17–22, 2015.

- 670 Forbes, R. M. and Ahlgrimm, M.: On the Representation of High-Latitude Boundary Layer Mixed-Phase Cloud in the ECMWF Global Model, *Monthly Weather Review*, 142, 3425–3445, <https://doi.org/10.1175/MWR-D-13-00325.1>, 2014.
- Forbes, R. M., Tompkins, A. M., and Untch, A.: A New Prognostic Bulk Microphysics Scheme for the IFS, European Centre for Medium-Range Weather Forecasts, 2011.
- Forbes, R.M. and Tompkins, A.: An Improved Representation of Cloud and Precipitation, *ECMWF Newsl*, 129, 2011.
- 675 Gettelman, A. and Morrison, H.: Advanced Two-Moment Bulk Microphysics for Global Models. Part I: Off-Line Tests and Comparison with Other Schemes, *Journal of Climate*, 28, 1268–1287, <https://doi.org/10.1175/JCLI-D-14-00102.1>, 2015.
- Gettelman, A., Liu, X., Ghan, S. J., Morrison, H., Park, S., Conley, A. J., Klein, S. A., Boyle, J., Mitchell, D. L., and Li, J.-L. F.: Global Simulations of Ice Nucleation and Ice Supersaturation with an Improved Cloud Scheme in the Community Atmosphere Model, *J. Geophys. Res.*, 115, D18 216, <https://doi.org/10.1029/2009JD013797>, 2010.
- 680 Gettelman, A., Morrison, H., Santos, S., Bogenschutz, P., and Caldwell, P. M.: Advanced Two-Moment Bulk Microphysics for Global Models. Part II: Global Model Solutions and Aerosol–Cloud Interactions, *J. Climate*, 28, 1288–1307, <https://doi.org/10.1175/JCLI-D-14-00103.1>, 2015.
- Gettelman, A., Hannay, C., Bacmeister, J. T., Neale, R. B., Pendergrass, A. G., Danabasoglu, G., Lamarque, J.-F., Fasullo, J. T., Bailey, D. A., Lawrence, D. M., and Mills, M. J.: High Climate Sensitivity in the Community Earth System Model Version 2 (CESM2), *Geophysical Research Letters*, 46, 8329–8337, <https://doi.org/10.1029/2019GL083978>, 2019a.
- 685 Gettelman, A., Mills, M. J., Kinnison, D. E., Garcia, R. R., Smith, A. K., Marsh, D. R., Tilmes, S., Vitt, F., Bardeen, C. G., McInerney, J., Liu, H.-L., Solomon, S. C., Polvani, L. M., Emmons, L. K., Lamarque, J.-F., Richter, J. H., Glanville, A. S., Bacmeister, J. T., Phillips, A. S., Neale, R. B., Simpson, I. R., DuVivier, A. K., Hodzic, A., and Randel, W. J.: The Whole Atmosphere Community Climate Model Version 6 (WACCM6), *Journal of Geophysical Research: Atmospheres*, 124, 12 380–12 403, <https://doi.org/10.1029/2019JD030943>, 2019b.
- 690 Gettelman, A., Bardeen, C. G., McCluskey, C. S., Järvinen, E., Stith, J., Bretherton, C., McFarquhar, G., Twohy, C., D’Alessandro, J., and Wu, W.: Simulating Observations of Southern Ocean Clouds and Implications for Climate, *Journal of Geophysical Research: Atmospheres*, 125, e2020JD032 619, <https://doi.org/10.1029/2020JD032619>, 2020.
- Harris, L., Zhou, L., Lin, S.-J., Chen, J.-H., Chen, X., Gao, K., Morin, M., Rees, S., Sun, Y., Tong, M., Xiang, B., Bender, M., Benson, R., Cheng, K.-Y., Clark, S., Elbert, O. D., Hazelton, A., Huff, J. J., Kaltenbaugh, A., Liang, Z., Marchok, T., Shin, H. H., and Stern, W.: GFDL
- 695 SHIELD: A Unified System for Weather-to-Seasonal Prediction, *Journal of Advances in Modeling Earth Systems*, 12, e2020MS002 223, <https://doi.org/10.1029/2020MS002223>, 2020.
- Hillman, B. R., Marchand, R. T., and Ackerman, T. P.: Sensitivities of Simulated Satellite Views of Clouds to Subgrid-Scale Overlap and Condensate Heterogeneity, *Journal of Geophysical Research: Atmospheres*, 123, 7506–7529, <https://doi.org/10.1029/2017JD027680>, 2018.
- Hinkelman, L. M. and Marchand, R.: Evaluation of CERES and CloudSat Surface Radiative Fluxes Over Macquarie Island, the Southern
- 700 Ocean, *Earth and Space Science*, 7, e2020EA001 224, <https://doi.org/10.1029/2020EA001224>, 2020.
- Hogan, R. J. and Bozzo, A.: A Flexible and Efficient Radiation Scheme for the ECMWF Model, *Journal of Advances in Modeling Earth Systems*, 10, 1990–2008, 2018.
- Hoose, C., Kristjánsson, J. E., Chen, J.-P., and Hazra, A.: A Classical-Theory-Based Parameterization of Heterogeneous Ice Nucleation by Mineral Dust, Soot, and Biological Particles in a Global Climate Model, *J. Atmos. Sci.*, 67, 2483–2503, <https://doi.org/10.1175/2010JAS3425.1>, 2010.
- 705

- Hu, Y., Rodier, S., Xu, K.-m., Sun, W., Huang, J., Lin, B., Zhai, P., and Josset, D.: Occurrence, Liquid Water Content, and Fraction of Supercooled Water Clouds from Combined CALIOP/IIR/MODIS Measurements, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD012384>, 2010.
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative Forcing by Long-Lived Greenhouse Gases: Calculations with the AER Radiative Transfer Models, *JGR*, 113, <https://doi.org/10.1029/2008JD009944>, 2008.
- Järvinen, E., McCluskey, C. S., Waitz, F., Schnaiter, M., Bansemer, A., Bardeen, C. G., Gettelman, A., Heymsfield, A., Stith, J. L., Wu, W., D'Alessandro, J. J., McFarquhar, G. M., Diao, M., Finlon, J. A., Hill, T. C. J., Levin, E. J. T., Moore, K. A., and DeMott, P. J.: Evidence for Secondary Ice Production in Southern Ocean Maritime Boundary Layer Clouds, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD036411, <https://doi.org/10.1029/2021JD036411>, 2022.
- Kato, S., Rose, F. G., Rutan, D. A., Thorsen, T. J., Loeb, N. G., Doelling, D. R., Huang, X., Smith, W. L., Su, W., and Ham, S.-H.: Surface Irradiances of Edition 4.0 Clouds and the Earth's Radiant Energy System (CERES) Energy Balanced and Filled (EBAF) Data Product, *J. Climate*, 31, 4501–4527, <https://doi.org/10.1175/JCLI-D-17-0523.1>, 2018.
- Khairoutdinov, M. F. and Kogan, Y.: A New Cloud Physics Parameterization in a Large-Eddy Simulation Model of Marine Stratocumulus, *Monthly Weather Review*, 128, 229–243, 2000.
- Köhler, M., Ahlgrimm, M., and Beljaars, A.: Unified Treatment of Dry Convective and Stratocumulus-Topped Boundary Layers in the ECMWF Model, *Quarterly Journal of the Royal Meteorological Society*, 137, 43–57, 2011.
- Kuma, P., McDonald, A. J., Morgenstern, O., Alexander, S. P., Cassano, J. J., Garrett, S., Halla, J., Hartery, S., Harvey, M. J., Parsons, S., Plank, G., Varma, V., and Williams, J.: Evaluation of Southern Ocean Cloud in the HadGEM3 General Circulation Model and MERRA-2 Reanalysis Using Ship-Based Observations, *Atmospheric Chemistry and Physics*, 20, 6607–6630, <https://doi.org/10.5194/acp-20-6607-2020>, 2020.
- Liebe, H. J.: An Updated Model for Millimeter Wave Propagation in Moist Air, *Radio Science*, 20, 1069–1089, <https://doi.org/10.1029/RS020i005p01069>, 1985.
- Liebe, H. J., Rosenkranz, P. W., and Hufford, G. A.: Atmospheric 60-GHz Oxygen Spectrum: New Laboratory Measurements and Line Parameters, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 48, 629–643, [https://doi.org/10.1016/0022-4073\(92\)90127-P](https://doi.org/10.1016/0022-4073(92)90127-P), 1992.
- Lott, F. and Miller, M. J.: A New Subgrid-Scale Orographic Drag Parametrization: Its Formulation and Testing, *Quarterly Journal of the Royal Meteorological Society*, 123, 101–127, 1997.
- Marchand, R., Ackerman, T., Westwater, E. R., Clough, S. A., Cady-Pereira, K., and Liljegren, J. C.: An Assessment of Microwave Absorption Models and Retrievals of Cloud Liquid Water Using Clear-Sky Data, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2003JD003843>, 2003.
- Marchand, R., Haynes, J., Mace, G. G., Ackerman, T., and Stephens, G.: A Comparison of Simulated Cloud Radar Output from the Multiscale Modeling Framework Global Climate Model with CloudSat Cloud Radar Observations, *Journal of Geophysical Research: Atmospheres*, 114, <https://doi.org/10.1029/2008JD009790>, 2009.
- McCluskey, C. S., Hill, T. C. J., Humphries, R. S., Rauker, A. M., Moreau, S., Stratton, P. G., Chambers, S. D., Williams, A. G., McRobert, I., Ward, J., Keywood, M. D., Harnwell, J., Ponsonby, W., Loh, Z. M., Krummel, P. B., Protat, A., Kreidenweis, S. M., and DeMott, P. J.: Observations of Ice Nucleating Particles Over Southern Ocean Waters, *Geophysical Research Letters*, 45, 11,989–11,997, <https://doi.org/10.1029/2018GL079981>, 2018.

- McFarquhar, G. M., Bretherton, C., Marchand, R., Protat, A., DeMott, P. J., Alexander, S. P., Roberts, G. C., Twohy, C. H., Toohey, D., Siems, S., Huang, Y., Wood, R., Rauber, R. M., Lasher-Trapp, S., Jensen, J., Stith, J., Mace, J., Um, J., Järvinen, E., Schnaiter, M.,
745 Gettelman, A., Sanchez, K. J., McCluskey, C. S., Russell, L. M., McCoy, I. L., Atlas, R., Bardeen, C. G., Moore, K. A., Hill, T. C. J.,
Humphries, R. S., Keywood, M. D., Ristovski, Z., Cravigan, L., Schofield, R., Fairall, C., Mallet, M. D., Kreidenweis, S. M., Rainwater,
B., D'Alessandro, J., Wang, Y., Wu, W., Saliba, G., Levin, E. J. T., Ding, S., Lang, F., Truong, S. C. H., Wolff, C., Haggerty, J., Harvey,
M. J., Klekociuk, A., and McDonald, A.: Observations of Clouds, Aerosols, Precipitation, and Surface Radiation over the Southern
Ocean: An Overview of CAPRICORN, MARCUS, MICRE and SOCRATES, *Bulletin of the American Meteorological Society*, -1, 1–92,
750 <https://doi.org/10.1175/BAMS-D-20-0132.1>, 2021.
- Meyers, M. P., DeMott, P. J., and Cotton, W. R.: New Primary Ice-Nucleation Parameterizations in an Explicit Cloud Model, *J. Applied Met.*,
31, 708–721, 1992.
- Minnis, P., Trepte, Q. Z., Sun-Mack, S., Chen, Y., Doelling, D. R., Young, D. F., Spangenberg, D. A., Miller, W. F., Wielicki, B. A., Brown,
R. R., Gibson, S. C., and Geier, E. B.: Cloud Detection in Nonpolar Regions for CERES Using TRMM VIRS and Terra and Aqua MODIS
755 Data, *IEEE Transactions on Geoscience and Remote Sensing*, 46, 3857–3884, <https://doi.org/10.1109/TGRS.2008.2001351>, 2008.
- Minnis, P., Sun-Mack, S., Young, D. F., Heck, P. W., Garber, D. P., Chen, Y., Spangenberg, D. A., Arduini, R. F., Trepte, Q. Z., Smith, W. L.,
Ayers, J. K., Gibson, S. C., Miller, W. F., Hong, G., Chakrapani, V., Takano, Y., Liou, K.-N., Xie, Y., and Yang, P.: CERES Edition-2
Cloud Property Retrievals Using TRMM VIRS and Terra and Aqua MODIS Data—Part I: Algorithms, *IEEE Transactions on Geoscience
and Remote Sensing*, 49, 4374–4400, <https://doi.org/10.1109/TGRS.2011.2144601>, 2011.
- 760 Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative Transfer for Inhomogeneous Atmospheres:
RRTM, a Validated Correlated-k Model for the Longwave, *Journal of Geophysical Research: Atmospheres*, 102, 16 663–16 682,
<https://doi.org/10.1029/97JD00237>, 1997.
- Molod, A., Takacs, L., Suarez, M., and Bacmeister, J.: Development of the GEOS-5 Atmospheric General Circulation Model: Evolution
from MERRA to MERRA2, *Geosci. Model Dev.*, 8, 1339–1356, <https://doi.org/10.5194/gmd-8-1339-2015>, 2015.
- 765 Morcrette, J. J., Barker, H. W., Cole, J. N. S., Iacono, M. J., and Pincus, R.: Impact of a New Radiation Package, McRad, in the ECMWF Integrated
Forecasting System, *Monthly weather review*, 136, 4773–4798, 2008.
- Morrison, H. and Gettelman, A.: A New Two-Moment Bulk Stratiform Cloud Microphysics Scheme in the NCAR Community Atmosphere
Model (CAM3), Part I: Description and Numerical Tests, *J. Climate*, 21, 3642–3659, 2008.
- Morrison, H., Curry, J. A., and Khvorostyanov, V. I.: A New Double-Moment Microphysics Parameterization for Application in Cloud and
770 Climate Models. Part I: Description, *jas*, 62, 1665–1677, 2005.
- Orr, A., Bechtold, P., Scinocca, J., Ern, M., and Janiskova, M.: Improved Middle Atmosphere Climate and Forecasts in the ECMWF Model
through a Nonorographic Gravity Wave Drag Parameterization, *Journal of Climate*, 23, 5905–5926, 2010.
- Platnick, S., King, M. D., Ackerman, S. A., Menzel, W. P., Baum, B. A., Riédi, J. C., and Frey, R. A.: The MODIS Cloud Products: Algorithms
and Examples from Terra, *Geoscience and Remote Sensing, IEEE Transactions on*, 41, 459–473, 2003.
- 775 Platnick, S., Meyer, K. G., King, M. D., Wind, G., Amarasinghe, N., Marchant, B., Arnold, G. T., Zhang, Z., Hubanks, P. A., Holz, R. E.,
Yang, P., Ridgway, W. L., and Riedi, J.: The MODIS Cloud Optical and Microphysical Products: Collection 6 Updates and Examples From
Terra and Aqua, *IEEE Transactions on Geoscience and Remote Sensing*, 55, 502–525, <https://doi.org/10.1109/TGRS.2016.2610522>, 2017.
- Rutan, D. A., Kato, S., Doelling, D. R., Rose, F. G., Nguyen, L. T., Caldwell, T. E., and Loeb, N. G.: CERES Synoptic Product: Methodology
and Validation of Surface Radiant Flux, *Journal of Atmospheric and Oceanic Technology*, 32, 1121–1143, [https://doi.org/10.1175/JTECH-](https://doi.org/10.1175/JTECH-
780 D-14-00165.1)
D-14-00165.1, 2015.

- Sachidananda, M. and Zrnica, D.S.: Differential Propagation Phase Shift and Rainfall Rate Estimation, *Radio science*, 21, 235–247, 1986.
- Santos, S. P., Caldwell, P. M., and Bretherton, C. S.: Cloud Process Coupling and Time Integration in the E3SM Atmosphere Model, *Journal of Advances in Modeling Earth Systems*, 13, e2020MS002359, <https://doi.org/10.1029/2020MS002359>, 2021.
- 785 Siebesma, A. P., Soares, P. M., and Teixeira, J.: A Combined Eddy-Diffusivity Mass-Flux Approach for the Convective Boundary Layer, *Journal of the atmospheric sciences*, 64, 1230–1248, 2007.
- Stephens, G. L., L'Ecuyer, T., Forbes, R., Gettelman, A., Golaz, J.-C., Bodas-Salcedo, A., Suzuki, K., Gabriel, P., and Haynes, J.: Dreary State of Precipitation in Global Models, *J. Geophys. Res.*, 115, D24 211, <https://doi.org/10.1029/2010JD014532>, 2010.
- Thompson, G. and Eidhammer, T.: A Study of Aerosol Impacts on Clouds and Precipitation Development in a Large Winter Cyclone, *J. Atmos. Sci.*, 71, 3636–3658, <https://doi.org/10.1175/JAS-D-13-0305.1>, 2014.
- 790 Tiedtke, M.: Representation of Clouds in Large-Scale Models, *Monthly Weather Review*, 121, 3040–3061, [https://doi.org/10.1175/1520-0493\(1993\)121<3040:ROCILS>2.0.CO;2](https://doi.org/10.1175/1520-0493(1993)121<3040:ROCILS>2.0.CO;2), 1993.
- Tiedtke, MICHAEL.: A Comprehensive Mass Flux Scheme for Cumulus Parameterization in Large-Scale Models, *Monthly weather review*, 117, 1779–1800, 1989.
- Tompkins, A. M., Gierens, K., and Rädcl, G.: Ice Supersaturation in the ECMWF Forecast System, *qjrm*, 133, 53–63, <https://doi.org/10.1002/qj.14>, 2007.
- 795 Trenberth, K. E. and Fasullo, J. T.: Simulation of Present-Day and Twenty-First-Century Energy Budgets of the Southern Oceans, *J. Climate*, 23, 440–454, <https://doi.org/10.1175/2009JCLI3152.1>, 2010.
- Wall, C. J., Hartmann, D. L., and Ma, P.-L.: Instantaneous Linkages between Clouds and Large-Scale Meteorology over the Southern Ocean in Observations and a Climate Model, *Journal of Climate*, 30, 9455–9474, <https://doi.org/10.1175/JCLI-D-17-0156.1>, 2017.
- 800 Zhou, L., Lin, S.-J., Chen, J.-H., Harris, L. M., Chen, X., and Rees, S. L.: Toward Convective-Scale Prediction within the Next Generation Global Prediction System, *Bulletin of the American Meteorological Society*, 100, 1225–1243, <https://doi.org/10.1175/BAMS-D-17-0246.1>, 2019.