

The impact of dehydration and extremely low HCl values in the Antarctic stratospheric vortex in mid-winter on ozone loss in spring

Yiran Zhang-Liu¹, Rolf Müller^{1,5}, Jens-Uwe Grooß^{1,5}, Sabine Robrecht^{1,2}, Bärbel Vogel^{1,5}, Abdul Mannan Zafar^{1,3}, and Ralph Lehmann⁴

¹Institute of Climate and Energy Systems (ICE-4), Forschungszentrum Jülich, Jülich, Germany

²Deutscher Wetterdienst, Offenbach, Germany

³Biotechnology Research Center, Technology Innovation Institute, Masdar City, Abu Dhabi, United Arab Emirates

⁴Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Potsdam, Germany

⁵Center for Advanced Simulation and Analytics (CASA), Forschungszentrum Jülich, Jülich, Germany

Correspondence: Rolf Müller (ro.mueller@fz-juelich.de) and Yiran Zhang-Liu (yiranz.luna@gmail.com)

Abstract. Simulations of Antarctic chlorine and ozone chemistry [in previous work](#) show that in the core of the Antarctic vortex (16–18 km, 85–55 hPa, 390–430 K) HCl null cycles (initiated by reactions ~~and of Cl with~~ CH₄ [and](#) CH₂O) are effective. These HCl null cycles ~~allow HCl cause~~ [HCl molar](#) mixing ratios to remain very low throughout Antarctic winter/[spring](#) and ozone destroying chlorine (ClO_x) to remain enhanced, so that rapid ozone depletion proceeds. ~~Sensitivity studies show that the reaction is important for the efficacy of the HCl null cycle initiated by the reaction and that using the current kinetic recommendations instead of earlier ones has little impact on the simulations. Dehydration in Antarctica~~ [Here we investigate the impact of the observed dehydration in Antarctica, which](#) strongly reduces ice formation and the uptake of HNO₃ from the gas phase; however the efficacy of HCl null cycles is not affected. ~~Further, the effect of~~ [Moreover, also when using](#) the observed very low HCl [molar](#) mixing ratios in Antarctic winter ~~are considered as~~ [initial value](#); HCl null cycles are efficient in maintaining low HCl (and high ClO_x) throughout ~~Antarctic winter~~ [winter/spring](#). ~~Further, the reaction~~ CH₃O₂ + ClO [is important for the efficacy of the HCl null cycle initiated by the reaction](#) CH₄ + Cl. ~~Using the current kinetic recommendations instead of earlier ones has very little impact on the simulations.~~ All simulations presented here for the core of the Antarctic vortex show extremely low minimum ozone values (below 50 ppb) in late September/early October in agreement with observations.

15 *Copyright statement.* TEXT

1 Introduction

The Antarctic ozone hole is a phenomenon of substantially reduced polar ozone that has reoccurred every winter and spring over Antarctica ~~since for~~ about four decades (~~Jones and Shanklin, 1995; Müller et al., 2008; Bodeker and Kremser, 2021; WMO, 2022; Klekociuk~~ [\(Jones and Shanklin, 1995; Müller et al., 2008; Bodeker and Kremser, 2021; WMO, 2022; Klekociuk et al., 2022; Johnson et al., 2023; R](#)
20 . In many years, the Antarctic ozone hole shows ~~a broad altitude region of~~ very low ozone in the altitude region of 14-

21 km ($\sim 380\text{-}550\text{ K}$ [potential temperature](#)) (Solomon et al., 2005; Jurkat et al., 2017; Johnson et al., 2023). In exceptional years, sudden stratospheric warmings occur in the Antarctic (2002 and 2019) causing unusually low ozone depletion (e.g. Müller et al., 2008; Grooß et al., 2005; Smale et al., 2021)(e.g., Müller et al., 2008; Grooß et al., 2005; Smale et al., 2021). Substantial polar ozone loss [may can](#) also occur in the Arctic; albeit the ozone loss shows a much stronger year-to-year variability (e.g., Müller et al., 2008; Johansson et al., 2019; Wohltmann et al., 2020; Dameris et al., 2021; Grooß and Müller, 2021; von der Gathen et al., 2021; Ardra et al., 2022). Polar ozone depletion is ultimately driven by chlorine and bromine substances released to the atmosphere as a result of human activities. (~~Although there are also bromine and chlorine substances with natural sources, W~~
~~Notwithstanding there are also bromine and chlorine substances with natural sources (WMO, 2022).~~ The release of the human made chlorine and bromine substances to the atmosphere has led to a substantial increase in the atmospheric halogen loading in the latter half of the last century; human made halogen compounds started increasing substantially in the atmosphere since the 1960s. Consequently, with the stratospheric halogen loading declining since about 2000, the first signs of recovery of both the Antarctic ozone hole and global ozone levels are observed (e.g., Várai et al., 2015; Kuttippurath and Nair, 2017; Strahan and Douglass, 2018; WMO, 2022; Bodeker and Kremser, 2021; Stone et al., 2021; Weber et al., 2022; Johnson et al., 2023).

~~Chemistry-climate~~ [For a successful simulation of the total column ozone field in the Antarctic vortex in spring, both the stratospheric chlorine and bromine chemistry need to be represented correctly in a model as well as the dynamical isolation of the Antarctic vortex \(e.g., Sonnabend et al., 2024\).](#) Current ~~chemistry-climate~~ models allow many characteristics of the global total column ozone field to be reproduced, but there is a considerable spread among models in the predictions of the absolute ozone column and the simulation of the Antarctic ozone hole is often not satisfactory. Possible reasons could be deficiencies in the model dynamics or in the stratospheric chemistry scheme of chemistry-climate models (Struthers et al., 2009; Dhomse et al., 2018). Such issues also impact the reliability and the accuracy of projections of the recovery of the Antarctic ozone hole under different climate scenarios for the future including climate intervention (Jöckel et al., 2016; Dhomse et al., 2018; Tilmes et al., 2021).

For polar stratospheric ozone depletion to occur, chlorine (which mostly prevails in the stratosphere in the form of the reservoir species HCl and ClONO₂) needs to be converted to an ozone destroying form. That is, HCl and ClONO₂ need to be “activated” by heterogeneous reactions on polar stratospheric clouds (PSCs) ~~and or~~ cold sulphate aerosol particles (Portmann et al., 1996; Solomon, 1999; Shi et al., 2001; Drdla and Müller, 2012; WMO, 2022; Tritscher et al., 2021). Ozone depletion occurs with the return of sunlight to the polar region; this time period is characterised by maintenance of high levels of active chlorine (e.g., Santee et al., 2005; Santee et al., 2008; Solomon et al., 2015; Nedoluha et al., 2016; Jurkat et al., 2017; Wohltmann et al., 2017; Müller et al., 2018; Santee et al., 2005; Santee et al., 2008; Solomon et al., 2015; Nedoluha et al., 2016; Jurkat et al., 2017; Wohltmann et al., 2017; Müller et al., 2018). [When PSCs occur, HNO₃ is sequestered in PSC particles and thus removed from the gas-phase. If the HNO₃ containing PSC particles sediment, permanent denitrification in the polar stratosphere occurs \(e.g., de Laat et al., 2024\).](#) PSCs are present in the Antarctic lower stratosphere throughout winter until early October, whereas in the Arctic PSCs occur with much greater year-to-year variability (Pitts et al., 2009; Spang et al., 2018).

In the initial step of chlorine activation, in the heterogeneous reaction



the available ClONO₂ is titrated against HCl (e.g., Solomon et al., 1986; Wohltmann et al., 2017). In the Antarctic lower stratosphere, the initial concentrations of HCl are greater than those of ClONO₂ (Jaeglé et al., 1997; Santee et al., 2008; Nakajima et al., 2020), so that there is no full activation in this step. Then a period of relatively little chemical change in polar night follows (“sleeping chemistry”). ~~Later, in-~~

60 In austral spring, the strongest ozone depletion occurs; during this period in the core of the Antarctic vortex at altitudes of 16–18 km (85–55 hPa, 390–430 K), high values of active chlorine (ClO_x = Cl + ClO + 2 × Cl₂O₂) are maintained in spite of increasingly rapid formation of HCl in the gas phase (Müller et al., 2018). The through reactions of Cl with CH₄ and CH₂O (Müller et al., 2018). During this period the most rapid ozone depletion occurs. For such conditions, the maintenance of high ClO_x values ~~during this period~~ is accomplished by effective reaction cycles (“HCl null cycles”) in which deactivation (i.e. 65 production of HCl) is immediately balanced by the heterogeneous reaction of HCl with HOCl



(Crutzen et al., 1992; Prather, 1992), which occurs on the surfaces of nitric acid trihydrate (NAT) and ice particles or within supercooled (liquid) ternary solutions and cold liquid aerosol particles. This period of high Further, the reaction



70 is essential for the HCl null cycle initiated by the reaction of Cl with CH₄ (Crutzen et al., 1992; Zafar et al., 2018, see also AR1-AR8 in app

However, at altitudes somewhat greater than 18 km (55 hPa, 430 K) and for conditions in the lower stratosphere closer to the edge of the polar vortex, HNO₃ will not continuously be sequestered in PSCs, so that periods with enhanced gas-phase concentrations of HNO₃ (compared to the vortex core) will occur. Under such conditions, more NO₂ will be available in the 75 gas-phase (e.g., de Laat et al., 2024), enhancing the production of ClONO₂, so that reaction R1 will have a stronger impact on chlorine chemistry. As a result, the chemistry of HCl null cycles will be more complex.

The period of strongly enhanced ClO_x and strong ozone loss in the Antarctic ends with a very rapid formation of HCl leading to a practically complete conversion of ClO_x to HCl through the reactions of Cl with CH₄ and CH₂O (i.e., to deactivation) (e.g., Crutzen et al., 1992; Douglass et al., 1995; Groöb et al., 1997; Groöb et al., 2011; Nakajima et al., 2020). However, very rarely, 80 when the Antarctic vortex is perturbed by a sudden stratospheric warming ~~;(2002 and 2019), there is less ozone depletion and~~ significant deactivation into ClONO₂ may also occur ~~in Antarctica~~ (Groöb et al., 2005; Smale et al., 2021).

Heterogeneous chlorine activation, enhanced concentrations of active chlorine and subsequent ozone loss occur frequently in the polar regions ~~and—under exceptional circumstances—~~. Under exceptional circumstances chlorine activation also occurs in the mid-latitudes for conditions of low temperatures and enhanced water vapour. The surfaces for heterogeneous reactions 85 might be provided for example by stratospheric ice particles, stratospheric sulphate aerosol particles (potentially enhanced by volcanic eruptions or climate intervention) or by wildfire smoke injected into the stratosphere (e.g. Solomon et al., 1997; Tilmes et al., 2008; von Hobe et al., 2011; Klobas et al., 2017; Robrecht et al., 2019, 2021; Tilmes et al., 2021; Ohneiser et al., 2022; Santee et al., 2022).

In the present study, we perform sensitivity analyses, exploring the influence of different parameters on the rate of these HCl null cycles and the resulting ozone loss. We extend earlier work (Grooß et al., 2011; Müller et al., 2018; Zafar et al., 2018) investigating the chemical processes in the core of the Antarctic vortex, in the lower stratosphere (16–18 km, 85–55 hPa, 390–430 K), where extremely low ozone ~~mixing ratios~~ molar mixing ratios in spring are reached regularly (Solomon et al., 2005; Johnson et al., 2023). ~~We use now~~ (Molar mixing ratios are identical to volume mixing ratios in the case of an ideal gas). As in the earlier work (Grooß et al., 2011; Müller et al., 2018), we rely on a detailed examination of a single trajectory and an analysis of multi-trajectory simulations. Here we do *not* employ a three-dimensional model version (see also section 2.1 below), which is based on global or hemispheric meteorological fields and includes atmospheric mixing (e.g., Poshyvailo et al., 2018; Grooß and Müller, 2021; Sonnabend et al., 2024). We now use the most recent recommendation (Burkholder et al., 2020) of chemical kinetics and photochemical data. ~~And we focus on~~

In particular, we take into account the impact of the observed Antarctic dehydration (e.g., Kelly et al., 1989; Vömel et al., 1995; Nedoluha et al., 2002; Jiménez et al., 2006; Ivanova et al., 2008; Rolf et al., 2015), which was neglected in earlier work (Müller et al., 2018; Zafar et al., 2018). Further, the impact of very low HCl molar mixing ratios in Antarctic winter (Wohltmann et al., 2017; Grooß et al., 2018) is now considered. Both dehydration and very low HCl molar mixing ratios are clearly observed in the atmosphere.

Taking into account the observed dehydration in the Antarctic vortex (see also section 2.2.1 below for details) reduces substantially the occurrence of ice clouds in the model. ~~A Ice clouds are very efficient in sequestering HNO₃ from the gas-phase (e.g. Hynes et al., 2002), thus a~~ lower occurrence of ice clouds in the model reduces substantially the uptake of gas-phase HNO₃ on ice particles. ~~–~~

; however we find that the efficacy of HCl null cycles is not affected. Assuming an HCl mixing ratio of zero after polar night while keeping total inorganic chlorine (Cl_y) constant (Wohltmann et al., 2017; Grooß et al., 2018, see also section 2.2.2 below for details) approximately takes into account the observed very low HCl molar mixing ratios in the Antarctic vortex in mid-winter. This assumption leads to very low HCl molar mixing ratios throughout Antarctic ~~spring~~ winter and spring, but the efficacy of HCl null cycles is again not affected.

The simulations presented here show that neither of these two assumptions (dehydration and very low HCl molar mixing ratios) has a very strong effect on the simulated chemical ozone depletion compared to earlier work (Müller et al., 2018; Zafar et al., 2018); similarly, using the most recent recommendation (Burkholder et al., 2020) has little impact. Severe ozone depletion to values below 50 ppb is simulated (consistent with observations) for the South Pole in late September and early October.

In summary, our box-model calculations of Antarctic chlorine and ozone chemistry corroborate earlier findings that HCl null cycles, in the core of the vortex in the lower stratosphere in spring are effective in allowing high levels of active chlorine to be maintained and rapid ozone loss to proceed. ~~The reaction of is important for the HCl null cycle initiated by the reaction~~ ~~–~~ We show here that these conclusions are not changed when current kinetic recommendations (Burkholder et al., 2020) are employed or when dehydration and very low HCl ~~mixing ratios~~ molar mixing ratios, both observed in polar winter, are taken into account.

2 Methods

125 2.1 Chemical model

The simulations reported here were performed with the Chemical Lagrangian model of the Stratosphere (CLaMS, McKenna et al., 2002; Grooß et al., 2005, 2018) with a set-up following closely the one used ~~in earlier studies (see Grooß et al., 2011; Müller et al., 20~~
earlier (Grooß et al., 2011; Müller et al., 2018; Zafar et al., 2018). Briefly, here, the stratospheric chemistry is calculated for
particular air parcels along three-dimensional trajectories. To integrate the differential equations representing the set of chem-
130 ical reactions considered here, we use the solver SVODE (Brown et al., 1989). Chemical rate constants and photolysis cross
sections generally are taken from the most current recommendation (Burkholder et al., 2020), but the earlier recommendations
by Sander et al. (2011) were used for comparison. Photolysis rates are calculated in spherical geometry (Becker et al., 2000).

Heterogeneous chemistry in the model (~~see Grooß et al., 2011; Müller et al., 2018, for details~~) is assumed to occur on the
surface of ice and ~~nitric acid trihydrate (NAT, NAT~~ (with a particle density of $3 \cdot 10^{-3} \text{ cm}^{-3}$), as well as in supercooled
135 liquid ternary particles (HNO₃/H₂SO₄/H₂O) and cold liquid binary (H₂SO₄/H₂O) particles. The occurrence of particles in
the model is determined by the temperature of the air mass. NAT particles are assumed to form at a supersaturation of 10 from
liquid ternary solutions or from ice evaporation. Ice is formed in the model at the equilibrium temperature (no supersaturation).
The initial density of liquid (binary) aerosol particles is assumed to be 10 cm^{-3} . The condensable material for liquid ternary
particles, NAT and ice is determined from the equilibrium with the gas-phase. The temperature dependent reaction probabilities
140 in liquid ternary particles (HNO₃/H₂SO₄/H₂O) and cold liquid binary (H₂SO₄/H₂O) particles (~~Burkholder et al., 2020~~) are
determined from recent recommendations (Burkholder et al., 2020). See also Grooß et al. (2011) and Müller et al. (2018) for
further details.

2.2 Trajectory and chemical set-up

Here trajectories for the austral winter 2003 are used, which are defined by the location and time of minimum ozone measured
145 by ozone sondes at the South Pole. Forward and backward trajectories are calculated from the South Pole ~~measurements at~~
different days (Grooß et al., 2011; Müller et al., 2018). We focus on one particular trajectory passing through the location of an
ozone sonde measurement at South Pole; 14 ppb O₃ at 74 hPa (391 K) on 24 September 2003 (Grooß et al., 2011). The same
trajectory was investigated in earlier work (Müller et al., 2018; Zafar et al., 2018) and is referred to below as the reference
trajectory ~~(see also section 3.3.1)~~.

150 Meteorological data were taken from operational analyses of the European Centre for Medium-range Weather Forecasts
(ECMWF) and diabatic descent rates of the air-parcels were calculated using a radiation code (Zhong and Haigh, 1995) and
climatological ozone and water vapour profiles (Grooß and Russell, 2005).

The initial molar mixing ratios for the reference trajectory for the main trace gases on 1 June are listed in Table 1; with the
exception of H₂O they are the same as in earlier work (Müller et al., 2018; Zafar et al., 2018). The initial value for ClONO₂ is
155 extremely low. These initial conditions imply the assumption that for the air parcels in question, the initial step of heterogeneous
chlorine activation (reaction R1) has already occurred, so that ClO_x values are enhanced and the HCl molar mixing ratio is

O ₃	2.2 ppm
H ₂ O	2.05 ppm
CH ₄	1.2 ppm
HNO ₃	4.5 ppb
HCl	1.05 ppb
ClO _x	1.01 ppb
ClONO ₂	12 ppt
HOCl	4.65 ppt
Br _y	17 ppt
CO	16 ppb

Table 1. Initial molar mixing ratios (for 1 June) of atmospheric trace gases used for the CLaMS simulation along the reference trajectory.

lower than at the beginning of the winter. Further, ~~denitrification~~ Antarctic denitrification (e.g., de Laat et al., 2024) is also represented in the initial conditions by assuming $\text{HNO}_3 = 4.5$ ppb. The sensitivity of the results of the simulations ~~on~~ to the initial ozone and initial HNO₃ molar mixing ratios, as well as the impact of assumptions on the chemistry of methylhypochlorite (CH₃OCl) and the methyl peroxy radical (CH₃O₂) has been discussed in previous work (Müller et al., 2018; Zafar et al., 2018).

2.2.1 Initial water vapour

There is one important exception to the initial values used previously, namely the initial value of water vapour. Assuming $\text{H}_2\text{O} = 4.1$ ppm (Müller et al., 2018; Zafar et al., 2018) is an appropriate estimate for 1 June; however such a value means essentially that ~~dehydration of the Antarctic stratosphere is neglected. However, every year irreversible dehydration irreversible~~ dehydration which occurs thereafter through ice particle sedimentation ~~occurs is neglected. Dehydration occurs every year in the Antarctic,~~ with the removal of water vapour from the air at sufficiently low (ice-formation) temperatures (e.g., Jiménez et al., 2006; Tritscher et al., 2019, 2021). Particles of different sizes will sediment at different rates (Müller and Peter, 1992). Further, there is a year-to-year variability in the extent and timing of the severity of Antarctic dehydration (e.g., Nedoluha et al., 2002) and the dehydration is not uniform throughout the Antarctic vortex (Kelly et al., 1989; Ivanova et al., 2008). Nonetheless, strong dehydration in the Antarctic winter vortex ~~(in June, July, August)~~ has been reported consistently both for in-situ and remote sensing measurements as well as for model simulations (Kelly et al., 1989; Vömel et al., 1995; Nedoluha et al., 2002; Jiménez et al., 2006; Ivanova et al., 2008; Schoeberl and Dessler, 2011; Rolf et al., 2015; Poshyvailo et al., 2018; Tritscher et al., 2019, 2021).

Kelly et al. (1989) report minimum values of H₂O based on aircraft measurements down to 1.5 ppm (at about ≈ 350 K) and H₂O molar mixing ratios of 2.0-2.4 ppm for an isentropic flight on 430 K on 2 September 1987. Kelly et al. (1989) also report temperatures corresponding to ice saturation molar mixing ratios of ~ 2 ppm over an altitude range 350-450 K in late August at the South Pole. Vömel et al. (1995) conducted a series of balloon measurements with a frost point hygrometer from

McMurdo station (in 1990) and South Pole (1990-1994) and find an average [molar](#) mixing ratio of H₂O of 2.3 ppm between 16 and 18 km. Ivanova et al. (2008) report airborne H₂O measurements in the core of the Antarctic vortex on 21 and 23 September and 2 and 8 October 1999 during the APE-GAIA campaign – measurements at 410-430 K cover the range between 1.8-2.5 ppm [molar](#) mixing ratio of H₂O. Jiménez et al. (2006), based on observations by the Microwave Limb Sounder (MLS) report reductions of H₂O up to ~ 3 ppm – they find H₂O [molar](#) mixing ratios at ~ 17 km (440 K) at about 80°S equivalent latitude on 15 September of 2-2.2 ppm water vapour.

Here (for about 430 K) we assume an initial [molar](#) mixing ratio of H₂O = 2.05 ppm, which accounts for the observed dehydration in Antarctica [during winter](#); 2.05 ppm of water vapour will remain in the gas-phase if a temperature of ~ 185 K is reached in the atmosphere. A value for initial H₂O = 2.05 is also close to the value of initial H₂O = 2.2 ppm employed in earlier work (Crutzen et al., 1992) [and is appropriate for conditions in late winter and early spring in the Antarctic](#).

2.2.2 Initial HCl

For Antarctic HCl, a discrepancy between simulations and observations between May and July was reported (Wohlmann et al., 2017; Grooß et al., 2018); during this period model simulations significantly overestimate the observed HCl [molar](#) mixing ratios. The process causing this discrepancy between observations and simulations is not known at this point in time ([but it must be a process missing in the models](#)). Different options include an increased uptake of HCl into PSC particles (Wohlmann et al., 2017), a temperature bias of the underlying meteorological analyses, and unknown (possibly heterogeneous) chemical reactions (Grooß et al., 2018).

Here we do not investigate a process missing in models (Wohlmann et al., 2017; Grooß et al., 2018); rather our choice of the initial conditions is aimed at estimating the maximum possible effect of the very low HCl [molar](#) mixing ratios on HCl null cycles, ClO_x, and ozone depletion. This is the case when assuming initial HCl to be zero, with a corresponding increase in ClO_x, i.e., initial Cl_y is unchanged in the simulations.

2.3 Kinetic and photochemical parameters

Simulation	Initial HCl	Initial H ₂ O	Kinetics (JPL recomm.)	Colour
S1	HCl = 1.05 ppb	H ₂ O = 4.1 ppm	(Sander et al., 2011)	magenta
S2	HCl = 1.05 ppb	H ₂ O = 4.1 ppm	(Burkholder et al., 2020)	ochre
S3	HCl = 1.05 ppb	H ₂ O = 2.05 ppm	(Burkholder et al., 2020)	blue
S4	HCl = 0.0 ppb	H ₂ O = 2.05 ppm	(Burkholder et al., 2020)	red

Table 2. Employed assumptions for four different box-model simulations (S1-S4) a long the reference trajectory. The colours refer to those used in Figs. 2, 4, and 5 below.

In the results presented below, the sensitivity of Antarctic chlorine chemistry and ozone loss in spring to dehydration, to low early winter HCl [molar](#) mixing ratios and to different recommendations for kinetic parameters (Sander et al., 2011; Burkholder

et al., 2020) is explored. The different model simulations (S1-S4) and the employed assumptions (as well as the colours used in the figures below) are summarised in Table 2.

2.3.1 Recent recommendations of chemical kinetic and photochemical data

205 Müller et al. (2018) and Zafar et al. (2018) presented results of box-model simulations for the lowermost stratosphere in the core of the Antarctic vortex based on earlier recommendations of chemical kinetic and photochemical data by the Jet Propulsion Laboratory (JPL, Sander et al., 2011, simulation S1 in Table 2). Simulation S1 is shown here to establish a link to earlier work. Otherwise (simulations S2-S4) we use here chemical kinetics and photochemical data from the most recent recommendation (Burkholder et al., 2020).

210 The observed partitioning of ClO and Cl₂O₂ in the Antarctic stratosphere is well represented when the recommendations by Burkholder et al. (2015) are used (Canty et al., 2016). As the rate constants affecting chlorine chemistry were not changed substantially between the recommendations by Burkholder et al. (2015) and ~~(Burkholder et al., 2020)~~ [the Burkholder et al. \(2020\)](#), polar chlorine chemistry is also well represented by the most recent recommendation [\(see also appendix B\)](#). The largest remaining uncertainties in the ozone depletion rate are caused by uncertainties in the kinetic parameters influencing the Cl₂O₂ photolysis rate (Kawa et al., 2009; Canty et al., 2016; Wohltmann et al., 2017).

2.3.2 The rate constant of the reaction ClO + CH₃O₂

Figure 1. The temperature dependent rate constant ([in cm³ molecule⁻¹ s⁻¹](#)) for reaction R3 (ClO + CH₃O₂ → products) from a variety of sources. The recommendation by Sander et al. (2011) is shown as a black solid line (JPL 11) and the red line (JPL 15/20) is for Burkholder et al. (2015) (and Burkholder et al., 2020) with the corrected value $A = 1.8 \times 10^{-11}$. Also shown are recent measurements (dashed line, Leather et al., 2012); (blue line, Ward and Rowley, 2016).

~~Compared to the listing in recommendations (Burkholder et al., 2015, 2020)~~ [The temperature dependence of the rates of bimolecular reactions are given by the Arrhenius equation](#)

$$k = A \cdot \exp\left(-\frac{E_a}{R \cdot T}\right) \quad (1)$$

220 [here \$k\$ is the rate constant \(in cm³ molecule⁻¹ s⁻¹\), \$A\$ is a pre-exponential factor \(with the same unit as \$k\$ \), \$T\$ is temperature \(in K\), \$R\$ is the universal gas-constant, and \$E_a\$ may be interpreted as the molar activation energy of the reaction. In recommendations \(Burkholder et al., 2015, 2020\), commonly values for \$A\$ \(in cm³ molecule⁻¹ s⁻¹\) and \$E_a/R\$ \(in K\) are listed. In the simulations reported below, we corrected the rate constant of ~~the reaction → prod.*0;~~ \[reactionR3 \\(ClO + CH₃O₂ → prod.\\)\]\(#\) compared to the listing in re](#)

225 the latter value is listed incorrectly (~~an obvious a typo~~; J. Burkholder, pers. comm.) both in Burkholder et al. (2015) and Burkholder et al. (2020). The temperature dependent rate constant is listed in Sander et al. (2011) as:

$$k_{\text{jp11}}(T) = 3.3 \times 10^{-12} \cdot \exp\left(-\frac{1}{115} \cdot \frac{1}{T} \cdot \frac{115}{T}\right) \quad (3)$$

~~where T is temperature.~~ This recommendation was updated ~~by Burkholder et al. (2015) and Burkholder et al. (2020), but~~ (Burkholder et al., 2015, 2020) and the correct equation is:

230 $k_{\text{jp15/20}}(T) = 1.8 \times 10^{-11} \cdot \exp\left(-\frac{1}{600} \cdot \frac{1}{T} \cdot \frac{600}{T}\right) \quad (4)$

~~where T is temperature in Kelvin.~~ We show (Fig. 1) the reaction rate constant in ~~equation Eq. 4~~ against the recommendation reported by Sander et al. (2011) and other recent measurements (Leather et al., 2012; Ward and Rowley, 2016) (Fig. 1).

$k(298\text{K})$	Source
$2.243 \cdot 10^{-12}$	Sander et al. (2011)
$2.404 \cdot 10^{-12}$	Burkholder et al. (2015, 2020)
$2.399 \cdot 10^{-12}$	Leather et al. (2012)
$2.552 \cdot 10^{-12}$	Ward and Rowley (2016)

Table 3. The rate constant k (in $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) of reaction R3 ($\text{ClO} + \text{CH}_3\text{O}_2 \rightarrow \text{products}$) at 298 K. Note that the correct value for the A-factor is: $A = 1.8 \times 10^{-11}$ (which is not listed correctly in recommendations, see eq. ??).

Using the incorrect A-factor would result in a rate constant $k_{\text{jp15/20}}$ which is inconsistent with laboratory measurements. ~~Further~~ However, when calculating the reaction rate at room temperature $k(298\text{K})$, the formulation of ~~equation Eq. 4~~ yields 235 consistent results with earlier work and also reproduces the $k(298\text{K})$ value reported ~~in by~~ Burkholder et al. (2015, 2020), see Table 3.

~~A simulation employing The impact of using the incorrect A-factor in the rate constant of reaction R3 leads to underestimating the rate constant for reaction R3 by a factor of ten. Assuming this incorrect value (for R3) in simulation S4 yields a slower HCl increase in early October and the minimum ozone mixing ratio reached is 30.6 ppb instead of 5.7 ppb in simulation S4.~~

240 ~~Assuming a too low (by a factor of ten) value for reaction R3 is not equivalent to neglecting reaction R3 completely. Removing reaction R3 completely from the set of reactions considered here leads to a weaker rate of ozone depletion and to a much larger value for minimum ozone (more than 100 ppb), whereas the simulated minimum ozone is ≈ 6 , when reaction R3 is taken into account correctly.~~

245 ~~Further, the temporal development of HCl is very different; when neglecting reaction R3, the increase in HCl is much less steep. Using the value for reaction R3 listed in the recommendations (Burkholder et al., 2015, 2020) shows very low HCl values~~

on 10 September (below 0.1 ppb), whereas HCl on this day is about 0.6 ppb when reaction R3 is neglected. Using the correct value for reaction R3 (eq. 4) results in even lower HCl values (close to zero) model simulations is discussed in section 3.3.2 below.

3 Results

250 3.1 Comparing the chemical kinetic and photochemical recommendations by Sander et al., 2011 and Burkholder et al., 2020

A comparison is shown (Fig. 2) for the box model simulation (S1) based on the kinetic and photochemical recommendations by Sander et al. (2011) ~~, (Müller et al., 2018; Zafar et al., 2018, which were used in earlier work)~~ (which were used in earlier work, Müller et al. and Burkholder et al. (2020), the most recent recommendation (S2). The results for HCl, ClO_x, HOCl, ClONO₂, and O₃ reported earlier (magenta lines in Fig. 2; simulation S1 in Table 2) showed that at 16–18 km (85–55 hPa) in the core of the vortex, high levels of active chlorine are maintained by HCl null cycles, where the formation of HCl is balanced by immediate reactivation (Müller et al., 2018; Zafar et al., 2018). The strongest ozone loss rates occur in September. The results when using the most recent recommendation (Burkholder et al., 2020) are very similar to those reported earlier (Fig. 2). However, levels of HCl in September are somewhat lower (and thus ClO_x somewhat higher, resulting in somewhat stronger ozone loss) when using the recommendations by Sander et al. (2011). The maximum difference between the two runs using different kinetic recommendations (Fig. 2) is less than 0.025 ppm (or 25 ppb) ~~although the relative difference is substantial. Box-model simulations along a trajectory passing through the location of the ozone-sonde observation at South Pole of 14 ppb on 74 hPa (391 K) on 24 September 2003 (Groß et al., 2011; Müller et al., 2018; Zafar et al., 2018, reference trajectory similar as in earlier studies) using the recommendations from Sander et al. (2011) (simulation S1, magenta lines) and Burkholder et al. (2020) (simulation S2, ochre lines).~~ (See also Table 2). Top panel (a) shows HCl (b) , (c) HOCl, (d) , (e) , and (f) ozone (log-scale). The results for simulation S1 using Sander et al. (2011) are identical to those presented earlier (Müller et al., 2018; Zafar et al., 2018). Overall, the differences between simulations S1 and S2 are minor (Fig. 2); see also section 2.3.1.

3.2 The impact of initial water vapour

~~In Fig. 4~~

270 3.2.1 Formation of ice particles

Two simulations are compared for a different initialisation of H₂O, namely H₂O = 4.11 ppm (simulation S2) and H₂O = 2.05 ppm (a more realistic initialisation for H₂O, simulation S3); both simulations are employing ~~Burkholder et al. (2020)~~ the most recent kinetic recommendations (Burkholder et al., 2020). For an initial water vapour molar mixing ratio of 2.05 ppm, both the area of ice PSCs and the molar mixing ratio of gas-phase water vapour are substantially smaller than for an initial water vapour molar mixing ratio of 4.11 ppm (Fig. 3). In particular, the simulated ice surface for H₂O_{ini} = 2.05 ppm is substantially




Figure 2. Box-model simulations along a trajectory passing through the location of the ozone sonde observation at South Pole of 14 ppb on 74 hPa (391 K) on 24 September 2003 (Grooß et al., 2011; Müller et al., 2018; Zafar et al., 2018, reference trajectory similar as in earlier studies) using the recommendations from Sander et al. (2011) (simulation S1, magenta lines) and Burkholder et al. (2020) (simulation S2, ochre lines). (See also Table 2). Top panel (a) shows HCl (b) ClO_x, (c) HOCl, (d) ClONO₂, (e) Cl₂O₂, and (f) ozone (log-scale). The results for simulation S1 using Sander et al. (2011) are identical to those presented earlier (Müller et al., 2018; Zafar et al., 2018).



Figure 3. The simulated temporal development of ice surface (top panel, a), gas-phase water vapour concentrations-mixing ratios (middle panel, b) and gas-phase HNO₃ concentrations-mixing ratios (bottom panel, c). The results for both initial water vapour molar mixing ratio of 4.11 ppm (simulation S2, ochre lines) and 2.05 ppm (simulation S3, blue lines) are shown. (See also Table 2 for further details).

280 ~~lower-smaller~~ than for $\text{H}_2\text{O}_{\text{ini}} = 4.11$ ppm. ~~There should be observational consequences of the very different ice surfaces in simulations S2 and S3 (Fig. 3), i.e. observations should allow discriminating between the hypotheses about initial water vapour in simulations S2 and S3. A significant difference between the runs for a different initial water vapour molar mixing ratio is a different concentration of gas-phase HNO_3 – caused by the uptake of HNO_3 onto ice particles (Fig. 3, bottom). The enhanced gas-phase HNO_3 (for initial water vapour molar mixing ratio of 2.05 ppm) allows more NO_x to be released to the gas phase and (in sun-light) leads to somewhat more formation of ClONO_2 (Fig. 4).~~

285 However, the very different ice ~~surface-between-the-two-simulations-has~~ surfaces between simulations S2 and S3 have remarkably little impact on the temporal development of HCl and active chlorine ~~with-(although there are a few periods with more ClONO_2 for an initial water vapour molar mixing ratio of 4.11 ppm, Fig. 4). This has consequences for chemical ozone depletion (Fig. 4). This-There is a slightly lower minimum value of ozone (≈ 10 ppb lower) for an initial water vapour molar mixing ratio of 4.11 ppm. The simulated ozone values for an initial water vapour molar mixing ratio of 2.05 ppm are very similar to those for an initial water vapour molar mixing ratio of 4.11 ppm in June and July and are ≈ 50 ppb higher between August and mid-September; the largest difference (≈ 100 ppb) in ozone molar mixing ratios is reached in mid to late September.~~

290 ~~This finding~~ is consistent with the notion that the ~~actual-rate constant~~ of the heterogeneous reactions (~~but-not-the-occurrence-heterogeneous-reactions~~) within HCl null cycles is of little relevance for the efficacy of the HCl null cycles ~~-(Müller et al., 2018)~~. ~~Although it is important for the efficacy of the HCl null cycles that temperatures are sufficiently low so that particles are present and heterogeneous reactions occur.~~ The rate constant of heterogeneous reactions is influenced strongly by the type of the available PSC particles. The efficacy of the HCl null cycles, however, ~~are-is~~ limited by the ~~rate-of-the-reaction-rates-of-the-reactions~~ of Cl with CH_4 and CH_2O (see appendix A). ~~The HCl null cycles are relevant for the maintenance of high levels of active chlorine (Müller et al., 2018)-~~

295 ~~Further~~In consequence, a substantial difference in initial water vapour ~~mixing-ratios~~ molar mixing ratios in simulations S2 and S3 does not result in a substantial difference of polar chlorine chemistry and ozone loss (Fig. 4). ~~There is a slightly lower minimum value of ozone-~~

300 3.2.2 The eruption of the Hunga volcano

The initial water vapour in the Antarctic vortex assumed here and for the related model simulations (Table 1 and section 3.2.1) is discussed below regarding the interpretation of water vapour injections into the stratosphere by volcanic eruptions. In January 2022, the eruption of the Hunga underwater volcano injected a huge, unprecedented in the observational record, amount of water vapour into the mid-stratosphere (Wohltmann et al., 2023; Fleming et al., 2024; Zhou et al., 2024).

305 The impact of this water vapour enhancement on Antarctic ozone has been assessed through model studies. Fleming et al. (2024) find that the excess H_2O is projected to increase polar stratospheric clouds and springtime halogen-ozone loss, enhancing the Antarctic ozone hole by 25–30 DU. Wohltmann et al. (2023) find that the direct chemical effect of the increased water vapour on vortex average Antarctic ozone depletion in June through October was minor (less than 4 DU). Zhou et al. (2024) confirm this conclusion, but find somewhat more ozone loss caused by the injected water vapour (≈ 10 ppb lower) for an initial water




Figure 4. Similar as Fig. 2 but using the recommendations from Burkholder et al. (2020). Results are compared for a different initialisation of H₂O; H₂O = 4.11 ppm (simulation S2, ochre lines) and H₂O = 2.05 ppm (simulation S3, blue lines). (See also Table 2).

310 ~~vapour mixing ratio of 4.11 ppm. The simulated ozone values for an initial water vapour mixing ratio of 2.05 ppm are very similar to those for an initial water vapour mixing ratio of 4.11 ppm in June and July and are ≈ 50 ppb higher between August and mid-September; the largest *difference* (≈ 100 ppb) in ozone mixing ratios is reached in mid-to-late September~~ DU) at the vortex edge.

315 The impact of the stratospheric water vapour enhancement through the Hunga eruption on Antarctic ozone has further been assessed in the analysis of satellite observations (Santee et al., 2024). It was observed that the Hunga eruption increased the vertical extent of PSC formation and chlorine activation in early Austral winter in the Antarctic vortex in 2023 (the Antarctic season influenced most strongly by the Hunga eruption). Nonetheless, ozone depletion in the Antarctic in 2023 was unremarkable throughout the lower stratosphere (Santee et al., 2024). The observation of a small impact of water vapour injected into the stratosphere on polar ozone loss is consistent with the notion put forward in this paper (Fig. 3) that low

320 temperatures in the vortex, which occur regularly in the Antarctic, limit the atmospheric water vapour to the water vapour saturation pressure over ice and thus remove any anomalies through dehydration before they can affect ozone loss.

~~A significant difference between the runs for a different initial water vapour mixing ratio is a different concentration of gas-phase — caused by the uptake of onto ice particles (The minor impact of the huge water vapour injections into the stratosphere by the Hunga volcano on Antarctic ozone in the 2023 season (Wohlmann et al., 2023; Fleming et al., 2024; Zhou et al., 2024;~~
325 is consistent with the small impact of initial water vapour in mid-winter and the subsequent formation of ice PSC particles in the model simulations presented here (section 3.2.1 and Fig. 3, bottom). The enhanced gas-phase (for initial water vapour mixing ratio of 2.05 ppm) allows more to be released to the gas phase and (in sun-light) leads to somewhat more formation of
). First, the low temperatures in the lower stratosphere in the core of the Antarctic vortex determine mid-winter water vapour (independent of the amount of water vapour present at the time of the formation of the vortex). Second, even if higher water
330 vapour mixing ratios prevailed in mid-winter, chlorine activation and chemical ozone loss remain practically unaltered (Fig. 4).

3.3 ~~Impact~~ The impact of initial HCl

3.3.1 Reference simulation

We conducted a simulation (for $\text{H}_2\text{O}_{\text{initial}} = 2.05$ ppm) assuming $\text{HCl}_{\text{initial}} = 0$ (see Sec. 2.2), which corresponds to an increase in initial active chlorine ($\text{ClO}_{\text{x,initial}} = 2.26$ ppb); that is we assume $\text{Cl}_y = \text{const.}$ (Fig. 5). Assuming an initial value of $\text{HCl} = 0$
335 for early June (red lines in Fig. 5) resembles the conditions in the atmosphere (Wohlmann et al., 2017; Grooß et al., 2018, see also section 2.2 above), while an initial value of $\text{HCl} = 1.05$ ppb (blue lines in Fig. 5) is closer to HCl values in current model simulations. Therefore, we refer to simulation S4 here as the reference simulation.

As expected, for more initial ClO_{x} , there is a somewhat stronger ozone depletion (Fig. 5, bottom panel, red line). The difference in ozone between the two simulations is below ≈ 100 ppb in June and July but increases to ≈ 400 ppb in September.
340 However, the ozone minimum values reached, differ only by ≈ 10 ppb. Overall, the difference in absolute ozone depletion between the two simulations (S3 and S4) is moderate (albeit not in relative terms) in accordance with the conclusions by Grooß et al. (2018).

~~However, HOCl~~ However, there is clearly an earlier onset of ozone depletion when $\text{HCl}_{\text{initial}} = 0$ is employed, with the difference between simulation S3 and S4 notable in late August/early September. Although these different temporal developments
345 of ozone (and ClO_{x}) are obvious in our Lagrangian simulation, it will not be simple to detect such a behaviour in satellite observations, where spatial averages over large horizontal and vertical scales are measured for a given point in time.

Further, HOCl molar mixing ratios are substantially higher for $\text{HCl}_{\text{initial}} = 0$ (in particular from mid-June to mid-August).
~~Further~~ The reduction of HOCl via the heterogeneous reaction R2 is suppressed for low HCl (Fig. 5). Indeed, the HCl molar
mixing ratios remain low from June to mid September indicating that the HCl null cycles are effective in maintaining low
350 HCl molar mixing ratios (Müller et al., 2018). When ozone molar mixing ratios reach extremely low values in late September, HCl molar mixing ratios increase rapidly, which occurs a few days earlier in the case of $\text{HCl}_{\text{initial}} = 0$ (Fig. 5). Differences in




Figure 5. Similar as Fig. 4 (for initial $\text{H}_2\text{O} = 2.05$ ppm) but comparing the results for simulation S3 (blue lines) with the results for simulation S4 (red lines), where a different initialisation of HCl and active chlorine (ClO_x) was used. The kinetic recommendations of Burkholder et al. (2020) are used. (See also Table 2).

the temporal development of HOCl, ClO_x , and HCl between simulations S3 and S4 are greater than for ozone and thus might possibly be simpler to detect in satellite observation than a different temporal development of ozone.

3.3.2 The impact of the reaction $\text{CH}_3\text{O}_2 + \text{ClO}$

355 Reaction R3 is essential for the HCl null cycle initiated by the reaction of Cl with CH_4 (see AR1-AR8 in appendix A) and thus has a certain importance for “ozone hole” chemistry (e.g. Crutzen et al., 1992; Zafar et al., 2018). Here we report further results of simulation S4 exploring the impact of assuming the incorrect A-factor in the rate constant of reaction R3 (see section 2.3.2 above). Clearly, the incorrect A-factor leads to underestimating the rate constant for reaction R3 by a factor of ten. Assuming this incorrect value (for R3) in simulation S4 yields a slower HCl increase in late September and the minimum ozone molar
360 mixing ratio reached is 30.6 ppb instead of 5.7 ppb in simulation S4 (Fig. 6).




Figure 6. The impact of the formulation of reaction $\text{CH}_3\text{O}_2 + \text{ClO}$ (see section 2.3.2) in simulation S4. Red line shows the results for simulation S4 (as in Fig. 5), dotted black line the results assuming the incorrect A-factor (see section 2.3.2) and the solid black line neglecting reaction $\text{CH}_3\text{O}_2 + \text{ClO}$ (R3) entirely. Top panel shows molar mixing ratios for HCl, bottom panel for ozone.

Of course, assuming a too low (by a factor of ten) value for the rate constant of reaction R3 is not equivalent to neglecting reaction R3 completely. Removing reaction R3 completely from the set of reactions considered here leads to a weaker rate of ozone depletion. Under these conditions, the minimum ozone is more than 100 ppb, whereas the simulated minimum ozone is 5.7 ppb, when reaction R3 is taken into account correctly (Fig. 6). Further, the temporal development of HCl is very different; when neglecting reaction R3, the increase in HCl starts earlier and is much less steep (Fig. 6). Using the value for the rate constant of reaction R3 as listed in the recommendations (Burkholder et al., 2015, 2020) shows very low HCl values on 10 September (below 0.1 ppb), whereas HCl on this day is about 0.6 ppb when reaction R3 is neglected. Using the correct value for reaction R3 (Eq. 4) results in even lower HCl values (close to zero).

3.4 Multi-trajectory simulations

Similar as in earlier work (Grooß et al., 2011; Müller et al., 2018) In the discussions above, one particular (reference) trajectory was considered. However, this trajectory is representative for the conditions in the core of the Antarctic vortex at 16–18 km (85–55 hPa, 390–430 K). To demonstrate this, we selected here twenty one trajectories passing the South Pole (in late September/early October) at the 400 K potential temperature level; these trajectories include diabatic descent and latitude variations. We This selection is the same as in earlier work (Grooß et al., 2011; Müller et al., 2018). For the simulations we use the most recent kinetic recommendations (Burkholder et al., 2020) (Burkholder et al., 2020, with the corrected rate constant of R3). From early August to early October these trajectories show roughly the same diabatic descent of ≈ 10 K, similarly as the reference trajectory discussed above. However, over this period, the different trajectories show strong variations in latitude (and thus exposure to sunlight); the latitude varies between the South Pole and $\approx 65^\circ\text{S}$ with some equatorward excursions to $\approx 60^\circ\text{S}$ and, sometimes to $\approx 55^\circ\text{S}$.

The initial values (for 1 August) were taken from Grooß et al. (2011); briefly denitrification and dehydration are taken into account, total chlorine (Cl_y) is deduced from observations (correlation with N_2O) and the HCl initialisation is taken from a climatology based on ACE-FTS (Atmospheric Chemistry Experiment – Fourier Transform Spectrometer) measurements (Jones et al., 2012). Overall, these initial conditions correspond well with the assumptions made for simulation S4. Since the initial value of Cl_y is taken from (Grooß et al., 2011) Grooß et al. (2011), also the initial ClO_x is different for the individual trajectories.

The results of the multi-trajectory simulations (Fig. 7) show a certain variability in the ozone loss rate, which depends strongly on solar insolation and on the initial value of ClO_x ; also note that initial ozone is different for the individual trajectories. The minimum ozone is reached for the individual trajectories between late September and early October. There is also a certain variability of the temporal development of HCl, with some trajectories showing intermittent increases in HCl for certain periods (but no complete deactivation).

However Nonetheless, all trajectories show strongly enhanced values of ClO_x over the period of strong ozone loss in August and September, consistent with suppressed values of HCl (Fig. 7). Minimum ozone values for all trajectories are very low; below ≈ 100 ppb for most and below 10 ppb for several trajectories (Fig. 7). The As in the reference simulation, the period of rapid ozone loss (driven by high levels of ClO_x) ends abruptly with chlorine deactivation through very rapid formation of HCl



Figure 7. Results from multi-trajectory simulations (21 trajectories) of the CLaMS box-model. The trajectories are consistently initialised (for 1 August) as in Grooß et al. (2011), which corresponds well with the assumptions made for simulation S4 (see text for details). Simulations were performed for a set of trajectories passing the South Pole at 400 K (Grooß et al., 2011). The results of the box-model simulations are shown for the time period from 1 August to 1 November 2003. Panel (a) shows HCl, panel (b) ClO_x, and panel (c) ozone. Individual trajectories are shown in different colour to allow them to be distinguished more easily.

395 (e.g., Crutzen et al., 1992; Douglass et al., 1995; Grooß et al., 1997; Müller et al., 2018). After deactivation, HCl values remain high and practically unchanged in the box model simulation.

4 Discussion

Simulations of Antarctic chlorine and ozone chemistry for the core of the Antarctic vortex (16–18 km, 85–55 hPa, 390–430 K) indicate that HCl null cycles (C1 and C2, see appendix A) are effective throughout winter and spring (Grooß et al., 2011; Müller et al., 2018). The HCl null cycles require sufficiently low temperatures so that heterogeneous reactions (in particular reaction R2) have significant reaction rates. Further, a significant rate of the reaction $\text{CH}_3\text{O}_2 + \text{ClO}$ (reaction R3) is important for the efficacy of the HCl null cycle C1 (~~as discussed earlier, Crutzen et al., 1992; Zafar et al., 2018~~)(as discussed earlier, Crutzen et al., 1992; Zafar et al., 2018). The HCl null cycles allow HCl molar mixing ratios to be maintained at very low values so that rapid ozone depletion proceeds until the deactivation of ClO_x into HCl.

405 A low value of water vapour in mid winter is suggested here ($\text{H}_2\text{O}_{\text{ini}} = 2.05$ ppm, see also section 2.2.1) to account for the observed dehydration in the Antarctic vortex (e.g., Kelly et al., 1989; Vömel et al., 1995; Nedoluha et al., 2002). This leads to substantially less formation of ice particles (Fig. 3) and thus to a substantially lower rate of heterogeneous reactions on ice as well as less uptake of HNO_3 from the gas phase compared to earlier work (Müller et al., 2018; Zafar et al., 2018). However, ozone depletion is not strongly affected, consistent with the results of Kirner et al. (2015), who used the chemistry-climate model ECHAM5/MESSy for Atmospheric Chemistry (EMAC) to investigate the impact of different types of PSCs on Antarctic chlorine activation and ozone loss. Kirner et al. (2015) find that heterogeneous chemistry on liquid particles is responsible for more than 90% of the ozone depletion in Antarctic spring and that heterogeneous chemistry on ice particles causes less than 5 DU of additional column ozone depletion. The conclusion that heterogeneous chlorine chemistry is dominated by reactions on liquid particles is supported also by other work (e.g., Wegner et al., 2012; Grooß and Müller, 2021; Tritscher et al., 2021).

415 Further, Antarctic winter MLS observations during the period of the onset of chlorine activation (between May and July in austral winter) and ground-based measurements at Syowa station in early July (Nakajima et al., 2020) indicate that very low molar mixing ratios of HCl prevail in the vortex, which are not well reproduced by model simulations (Wohltmann et al., 2017; Grooß et al., 2018). Possible reasons for these very low molar mixing ratios of HCl are discussed in detail above (section 2.2.2).

420 Accounting for very low HCl values in mid-winter in the initial values of our simulations (see also section 2.2.2), we find very low HCl molar mixing ratios throughout winter and spring. This result is consistent with HCl values being maintained at very low values through the efficacy of the HCl null cycles (C1 and C2, see appendix A). Further, ~~while ozone depletion is somewhat enhanced under these conditions,~~ ozone depletion is not strongly affected by the initial values of HCl (and also the minimum values of Antarctic ozone reached are similar) consistent with Grooß et al. (2018).

The results of our simulations corroborate earlier findings that effective HCl null cycles (C1 and C2; see Appendix A) allow high levels of active chlorine to be maintained in the Antarctic lower stratosphere during the period of strong ozone depletion. During this period, HCl production rates in the gas-phase are high (and increase with decreasing ozone, Grooß et al., 2011; Müller et al., 2018).

430 The sensitivity investigations in the present study show the following. First, using the most recent recommendation for chemical kinetic and photochemical data (Burkholder et al., 2020), does not change the results of the simulations substantially compared to earlier work (where Sander et al., 2011, was used).

Second, the HCl null cycles require the heterogeneous reaction R2 ($\text{HCl} + \text{HOCl} \rightarrow \text{Cl}_2 + \text{H}_2\text{O}$) to proceed at a substantial rate. Further, the gas-phase reaction R3 ($\text{ClO} + \text{CH}_3\text{O}_2 \rightarrow \text{prod.}$) is essential (for the null cycle initiated by the reaction
435 $\text{CH}_4 + \text{Cl}$, see also appendix A and section 2.3.2). If reaction R3 ~~were~~ was neglected, HCl molar mixing ratios in early September of ≈ 0.6 ppb are simulated, instead of HCl values close to zero. Further simulated minimum ozone is more than 100 ppb instead of ≈ 6 ppb with reaction R3.

Third, taking into account the observed dehydration in the Antarctic lower stratosphere in winter (see section 2.2), which was neglected in earlier work (Müller et al., 2018; Zafar et al., 2018), substantially reduces the occurrence of ice clouds in
440 the model, but does not affect strongly the results of chlorine chemistry and ozone loss. The most important impact of the simulated difference in the occurrence of ice clouds in the model is caused by the uptake of HNO_3 from the gas-phase into ice particles. The HNO_3 uptake is smaller, when less ice particle surface is available. If the observed dehydration is taken into account in the simulations, a slightly higher minimum value of ozone (≈ 10 ppb higher) is simulated.


Finally, the maximum impact of the observed (but unexplained) observation of extremely low values of HCl during polar
445 night (Wohlmann et al., 2017; Grooß et al., 2018) was investigated based on the current simulations. This is done by assuming an HCl molar mixing ratio of zero after polar night on 1 June (while keeping Cl_y constant). These assumptions lead to a temporal development of the chlorine chemistry that is different ~~than when~~ from that assuming a higher initial HCl; in particular HOCl molar mixing ratios are enhanced from about mid-June to mid-August. Further, ClO_x is enhanced throughout winter and spring, and HCl molar mixing ratios remain very low until rapid chlorine deactivation occurs into HCl. However, the overall
450 ozone depletion in the Antarctic lower stratosphere is not strongly affected (consistent with Grooß et al., 2018); the simulated ozone minimum values differ by ≈ 10 ppb. Overall, our simulations indicate extremely low minimum ozone values at the South Pole (below 50 ppb) in late September/early October in agreement with observations (Solomon et al., 2005; Grooß et al., 2011; Johnson et al., 2023).

Code availability. The CLaMS model is accessible via <https://jugit.fz-juelich.de/clams/CLaMS> (CLaMS, 2022).


455 *Data availability.* The model results presented here are attached to the paper as an electronic supplement (in netcdf format).

Appendix A: HCl null cycles



The HCl null cycles listed below are responsible for the maintenance of high levels of active chlorine throughout Antarctic spring and were reported earlier (Müller et al., 2018; Zafar et al., 2018); they are repeated here for reference. Cycle C1 (Crutzen et al., 1992; Müller et al., 2018) starts with HCl production in the reaction $\text{CH}_4 + \text{Cl}$



Further, the formation of HCl in the reaction



leads to the following cycle (C2, Müller et al., 2018):



Appendix B: Changes in JPL2015 and JPL2020

460 In Burkholder et al. (2020) a new algorithm for the formulation of termolecular reactions (association dissociation) was introduced, which is used here. Note the correction (compared to the values listed in the recommendations) to the reaction $\text{ClO} + \text{CH}_3\text{O}_2 \rightarrow \text{prod.}$ discussed in detail in section 2.3.




Table A1. Changes between recent recommendations; Sander et al. (2011) versus Burkholder et al. (2015).




Table A2. Changes between recent recommendations; Burkholder et al. (2015) versus Burkholder et al. (2020).

Author contributions. Y.Z.-L., J.-U.G., and A.M.Z. conducted the simulations with the CLaMS box model that are reported here. R.M. and J.-U.G. conceived and designed the research project. The issues regarding the rate constant of the reaction $\text{ClO} + \text{CH}_3\text{O}_2$ (as discussed in section 2.3) were first raised by R.L.. All co-authors discussed the results and contributed to formulating the manuscript.

Competing interests. J.-U.G. and R.M. are editors of ACP; otherwise the authors declare that they have no competing interests

Acknowledgements. This paper originates from the Master’s thesis of Y. Z-L.; we thank Prof. E. Rühl from the Free University of Berlin for supervising this thesis work. We also thank J. Burkholder for very helpful comments on the rate constant of reaction R3. We [are grateful to M. Chipperfield and D. Kinnison for comments on the manuscript.](#) We thank the European Centre for Medium-range Weather Forecasts
470 (ECMWF) for providing meteorological data sets. This research was partly funded by the “Pilot Lab Exascale Earth System Modelling” project of the Helmholtz association. [We thank the three anonymous reviewers of the paper very much for helpful comments that led to an improved version of the paper.](#)

References

- Ardra, D., Kuttippurath, J., Roy, R., Kumar, P., Raj, S., Müller, R., and Feng, W.: The unprecedented ozone loss in the Arctic winter and spring of 2010/2011 and 2019/2020, *ACS Earth and Space Chemistry*, sp-2021-003338, <https://doi.org/10.1021/acsearthspacechem.1c00333>, 2022.
- 475
- Becker, G., Groöß, J.-U., McKenna, D. S., and Müller, R.: Stratospheric photolysis frequencies: Impact of an improved numerical solution of the radiative transfer equation, *J. Atmos. Chem.*, *37*, 217–229, <https://doi.org/10.1023/A:1006468926530>, 2000.
- Bodeker, G. E. and Kremser, S.: Indicators of Antarctic ozone depletion: 1979 to 2019, *Atmos. Chem. Phys.*, *21*, 5289–5300, <https://doi.org/10.5194/acp-21-5289-2021>, 2021.
- 480
- Brown, P. N., Byrne, G. D., and Hindmarsh, A. C.: VODE: A variable coefficient ODE solver, *SIAM J. Sci. Stat. Comput.*, *10*, 1038–1051, 1989.
- Burkholder, J. B., Sander, S. P., Abbatt, J. P. D., Barker, J. R., Huie, R. E., Kolb, C. E., Kurylo, M. J., Orkin, V. L., Wilmouth, D. M., and Wine, P. H.: Chemical kinetics and photochemical data for use in atmospheric studies, Evaluation Number 18, JPL Publication 15-10, 2015.
- 485
- Burkholder, J. B., Sander, S. P., Abbatt, J. P. D., Barker, J. R., Cappa, C., Crouse, J. D., Dibble, T. S., Huie, R. E., Kolb, C. E., Kurylo, M. J., Orkin, V. L., Percival, C. J., Wilmouth, D. M., and Wine, P. H.: Chemical kinetics and photochemical data for use in atmospheric studies, Evaluation Number 19, JPL Publication 19-5, <http://jpldataeval.jpl.nasa.gov>, 2020.
- Canty, T. P., Salawitch, R. J., and Wilmouth, D. M.: The kinetics of the ClOOCl catalytic cycle, *J. Geophys. Res.*, *121*, 13 768–13 783, <https://doi.org/10.1002/2016JD025710>, 2016.
- 490
- Crutzen, P. J., Müller, R., Brühl, C., and Peter, T.: On the potential importance of the gas phase reaction $\text{CH}_3\text{O}_2 + \text{ClO} \rightarrow \text{ClOO} + \text{CH}_3\text{O}$ and the heterogeneous reaction $\text{HOCl} + \text{HCl} \rightarrow \text{H}_2\text{O} + \text{Cl}_2$ in “ozone hole” chemistry, *Geophys. Res. Lett.*, *19*, 1113–1116, <https://doi.org/10.1029/92GL01172>, 1992.
- Dameris, M., Loyola, D. G., Nützel, M., Coldewey-Egbers, M., Lerot, C., Romahn, F., and van Roozendaal, M.: Record low ozone values over the Arctic in boreal spring 2020, *Atmos. Chem. Phys.*, *21*, 617–633, <https://doi.org/10.5194/acp-21-617-2021>, 2021.
- 495
- de Laat, A., van Geffen, J., Stammes, P., van der A, R., Eskes, H., and Veeckind, J. P.: The Antarctic stratospheric nitrogen hole: Southern Hemisphere and Antarctic springtime total nitrogen dioxide and total ozone variability as observed by Sentinel-5p TROPOMI, *Atmos. Chem. Phys.*, *24*, 4511–4535, <https://doi.org/10.5194/acp-24-4511-2024>, 2024.
- Dhomse, S. S., Kinnison, D., Chipperfield, M. P., Salawitch, R. J., Cionni, I., Hegglin, M. I., Abraham, N. L., Akiyoshi, H., Archibald, A. T., Bednarz, E. M., Bekki, S., Braesicke, P., Butchart, N., Dameris, M., Deushi, M., Frith, S., Hardiman, S. C., Hassler, B., Horowitz, L. W., Hu, R.-M., Jöckel, P., Josse, B., Kirner, O., Kremser, S., Langematz, U., Lewis, J., Marchand, M., Lin, M., Mancini, E., Marécal, V., Michou, M., Morgenstern, O., O’Connor, F. M., Oman, L., Pitari, G., Plummer, D. A., Pyle, J. A., Revell, L. E., Rozanov, E., Schofield, R., Stenke, A., Stone, K., Sudo, K., Tilmes, S., Visionsi, D., Yamashita, Y., and Zeng, G.: Estimates of ozone return dates from Chemistry-Climate Model Initiative simulations, *Atmos. Chem. Phys.*, *18*, 8409–8438, <https://doi.org/10.5194/acp-18-8409-2018>, 2018.
- 500
- Douglass, A. R., Schoeberl, M. R., Stolarski, R. S., Waters, J. W., Russell III, J. M., Roche, A. E., and Massie, S. T.: Interhemispheric differences in springtime production of HCl and ClONO₂ in the polar vortices, *J. Geophys. Res.*, *100*, 13 967–13 978, <https://doi.org/10.1029/95JD00698>, 1995.
- 505
- Drdla, K. and Müller, R.: Temperature thresholds for chlorine activation and ozone loss in the polar stratosphere, *Ann. Geophys.*, *30*, 1055–1073, <https://doi.org/10.5194/angeo-30-1055-2012>, 2012.

- 510 Fleming, E. L., Newman, P. A., Liang, Q., and Oman, L. D.: Stratospheric temperature and ozone impacts of the Hunga Tonga-Hunga Ha'apai water vapor injection, *J. Geophys. Res.*, 129, e2023JD039298, <https://doi.org/10.1029/2023JD039298>, 2024.
- Groß, J.-U. and Müller, R.: Simulation of record Arctic stratospheric ozone depletion in 2020, *J. Geophys. Res.*, 126, e2020JD033339, <https://doi.org/10.1029/2020JD033339>, 2021.
- Groß, J.-U. and Russell, J. M.: Technical note: A stratospheric climatology for O₃, H₂O, CH₄, NO_x, HCl, and HF derived from HALOE
515 measurements, *Atmos. Chem. Phys.*, 5, 2797–2807, <https://doi.org/10.5194/acp-5-2797-2005>, 2005.
- Groß, J.-U., Pierce, R. B., Crutzen, P. J., Grose, W. L., and Russell III, J. M.: Re-formation of chlorine reservoirs in southern hemisphere polar spring, *J. Geophys. Res.*, 102, 13 141–13 152, <https://doi.org/10.1029/96JD03505>, 1997.
- Groß, J.-U., Günther, G., Müller, R., Konopka, P., Bausch, S., Schlager, H., Voigt, C., Volk, C. M., and Toon, G. C.: Simulation of denitrification and ozone loss for the Arctic winter 2002/2003, *Atmos. Chem. Phys.*, 5, 1437–1448, 2005.
- 520 Groß, J.-U., Konopka, P., and Müller, R.: Ozone chemistry during the 2002 Antarctic vortex split, *J. Atmos. Sci.*, 62, 860–870, 2005.
- Groß, J.-U., Brautzsch, K., Pommrich, R., Solomon, S., and Müller, R.: Stratospheric ozone chemistry in the Antarctic: What controls the lowest values that can be reached and their recovery?, *Atmos. Chem. Phys.*, 11, 12 217–12 226, <https://doi.org/10.5194/acp-11-12217-2011>, 2011.
- Groß, J.-U., Müller, R., Spang, R., Tritscher, I., Wegner, T., Chipperfield, M. P., Feng, W., Kinnison, D. E., and Madronich, S.: On the discrepancy of HCl processing in the core of the wintertime polar vortices, *Atmos. Chem. Phys.*, pp. 8647–8666, <https://doi.org/10.5194/acp-18-8647-2018>, 2018.
- 525 Hynes, R. G., Fernandez, M. A., and Cox, R. A.: Uptake of HNO₃ on water-ice and coadsorption of HNO₃ and HCl in the temperature range 210–235 K, *J. Geophys. Res.*, 107, 4797, <https://doi.org/https://doi.org/10.1029/2001JD001557>, 2002.
- Ivanova, E. V., Volk, C. M., Riediger, O., Klein, H., Sitnikov, N. M., Ulanovskii, A. E., Yushkov, V. A., Ravegnani, F., Möbius, T., and
530 Schmidt, U.: A quasi-Lagrangian coordinate system based on high resolution tracer observations: implementation for the Antarctic polar vortex, *Atmos. Chem. Phys. Discuss.*, 8, 16 123–16 173, <https://doi.org/10.5194/acpd-8-16123-2008>, 2008.
- Jaeglé, L., Webster, C. R., May, R. D., Scott, D. C., Stimpfle, R. M., Kohn, D. W., Wennberg, P. O., Hanisco, T. F., Cohen, R. C., Proffitt, M. H., Kelly, K. K., Elkins, J., Baumgardner, D., Dye, J. E., Wilson, J. C., Pueschel, R. F., Chan, K. R., Salawitch, R. J., Tuck, A. F., Hovde, S. J., and Yung, Y. L.: Evolution and stoichiometry of heterogeneous processing in the Antarctic stratosphere., *J. Geophys. Res.*,
535 102, 13 235–13 253, <https://doi.org/10.1029/97JD00935>, 1997.
- Jiménez, C., Pumphrey, H. C., MacKenzie, I. A., Manney, G. L., Santee, M. L., Schwartz, M. J., Harwood, R. S., and Waters, J. W.: EOS MLS observations of dehydration in the 2004–2005 polar winters, *Geophys. Res. Lett.*, 33, L16806, <https://doi.org/10.1029/2006GL025926>, 2006.
- Jöckel, P., Tost, H., Pozzer, A., Kunze, M., Kirner, O., Brenninkmeijer, C., Brinkop, S., Duy, S. C., Dyroff, C., Eckstein, J., Frank, F.,
540 Garny, H., Gottschaldt, K.-D., Graf, P., Grewe, V., Kerkweg, A., Kern, B., Matthes, S., Mertens, A., Meul, S., Neumaier, M., Nützel, M., Oberländer-Hayn, S., Ruhnke, R., Runde, T., Sander, R., Scharffe, D., and Zahn, A.: Earth System integrated Modelling (ESCiMo) with the Modular Earth Submodel System (MESSy) version 2.51, *Geosci. Model Dev.*, 9, 1153–1200, 2016.
- Johansson, S., Santee, M. L., Groß, J.-U., Höpfner, M., Braun, M., Friedl-Vallon, F., Khosrawi, F., Kirner, O., Kretschmer, E., Oelhaf, H., Orphal, J., Sinnhuber, B.-M., Tritscher, I., Ungermann, J., Walker, K. A., and Woiwode, W.: Unusual chlorine partitioning in the 2015/16
545 Arctic winter lowermost stratosphere: observations and simulations, *Atmos. Chem. Phys.*, 19, 8311–8338, <https://doi.org/10.5194/acp-19-8311-2019>, 2019.

- Johnson, B. J., Cullis, P., Booth, J., Petropavlovskikh, I., McConville, G., Hassler, B., Morris, G. A., Sterling, C., and Oltmans, S.: South Pole Station ozonesondes: variability and trends in the springtime Antarctic ozone hole 1986–2021, *Atmos. Chem. Phys.*, 23, 3133–3146, <https://doi.org/10.5194/acp-23-3133-2023>, 2023.
- 550 Jones, A., Walker, K. A., Jin, J. J., Taylor, J. R., Boone, C. D., Bernath, P. F., Brohede, S., Manney, G. L., McLeod, S., Hughes, R., and Daffer, W. H.: Technical Note: A trace gas climatology derived from the Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS) data set, *Atmos. Chem. Phys.*, 12, 5207–5220, <https://doi.org/10.5194/acp-12-5207-2012>, 2012.
- Jones, A. E. and Shanklin, J. D.: Continued decline of total ozone over Halley, Antarctica, since 1985, *Nature*, 376, 409–411, <https://doi.org/10.1038/376409a0>, 1995.
- 555 Jurkat, T., Voigt, C., Kaufmann, S., Grooss, J. U., Ziereis, H., Doernbrack, A., Hoor, P., Bozem, H., Engel, A., Boenisch, H., Keber, T., Hueneke, T., Pfeilsticker, K., Zahn, A., Walker, K. A., Boone, C. D., Bernath, P. F., and Schlager, H.: Depletion of ozone and reservoir species of chlorine and nitrogen oxide in the lower Antarctic polar vortex measured from aircraft, *Geophys. Res. Lett.*, 44, 6440–6449, <https://doi.org/10.1002/2017GL073270>, 2017.
- Kawa, S. R., Stolarski, R. S., Newman, P. A., Douglass, A. R., Rex, M., Hofmann, D. J., Santee, M. L., and Frieler, K.: Sensitivity of polar stratospheric ozone loss to uncertainties in chemical reaction kinetics, *Atmos. Chem. Phys.*, 9, 8651–8660, <https://doi.org/10.5194/acp-9-8651-2009>, 2009.
- 560 Kelly, K. K., Tuck, A. F., Murphy, D. M., Proffitt, M. H., Fahey, D. W., Jones, R. L., McKenna, D. S., Loewenstein, M., Podolske, J. R., Strahan, S. E., Ferry and K. R. Chan and J. F. Vedder, G. V., Gregory, G. L., Hypes, W. D., McCormick, M. P., Browell, E. V., and Heidt, L. E.: Dehydration in the lower Antarctic stratosphere during late winter and early spring, 1987, *J. Geophys. Res.*, 94, 11 317–11 357, 1989.
- 565 Kirner, O., Müller, R., Ruhnke, R., and Fischer, H.: Contribution of liquid, NAT and ice particles to chlorine activation and ozone depletion in Antarctic winter and spring, *Atmos. Chem. Phys.*, 15, 2019–2030, <https://doi.org/10.5194/acp-15-2019-2015>, 2015.
- Klekociuk, A., Tully, M., Krummel, P., Henderson, S., Smale, D., Querel, R., Nichol, S., Alexander, S., Fraser, P., and Nedoluha, G.: The Antarctic Ozone Hole during 2020, *Journal of Southern Hemisphere Earth System Science*, 72, 19–37, <https://doi.org/doi:10.1071/ES21015>, 2022.
- 570 Klobas, E. J., Wilmouth, D. M., Weisenstein, D. K., Anderson, J. G., and Salawitch, R. J.: Ozone depletion following future volcanic eruptions, *Geophys. Res. Lett.*, 44, 7490–7499, <https://doi.org/https://doi.org/10.1002/2017GL073972>, 2017.
- Kuttippurath, J. and Nair, P. J.: The signs of Antarctic ozone hole recovery, *Sci. Rep.*, 7, 585, <https://doi.org/10.1038/s41598-017-00722-7>, 2017.
- 575 Leather, K. E., Bacak, A., Wamsley, R., Archibald, A. T., Husk, A., Shallcross, D. E., and Percival, C. J.: Temperature and pressure dependence of the rate coefficient for the reaction between ClO and CH₃O₂ in the gas-phase, *Phys. Chem. Chem. Phys.*, 14, 3425–3434, <https://doi.org/10.1039/C2CP22834C>, 2012.
- McKenna, D. S., Groß, J.-U., Günther, G., Konopka, P., Müller, R., Carver, G., and Sasano, Y.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS): 2. Formulation of chemistry scheme and initialization, *J. Geophys. Res.*, 107, 4256, <https://doi.org/10.1029/2000JD000113>, 2002.
- 580 Müller, R. and Peter, T.: The numerical modelling of the sedimentation of polar stratospheric cloud particles, *Ber. Bunsenges. Phys. Chem.*, 96, 353–361, 1992.
- Müller, R., Groß, J.-U., Lemmen, C., Heinze, D., Dameris, M., and Bodeker, G.: Simple measures of ozone depletion in the polar stratosphere, *Atmos. Chem. Phys.*, 8, 251–264, <https://doi.org/10.5194/acp-8-251-200>, 2008.

- 585 Müller, R., Grooß, J.-U., Zafar, A. M., Robrecht, S., and Lehmann, R.: The maintenance of elevated active chlorine levels in the Antarctic lower stratosphere through HCl null cycles, *Atmos. Chem. Phys.*, 18, 2985–2997, <https://doi.org/10.5194/acp-18-2985-2018>, 2018.
- Nakajima, H., Murata, I., Nagahama, Y., Akiyoshi, H., Saeki, K., Kinase, T., Takeda, M., Tomikawa, Y., Dupuy, E., and Jones, N. B.: Chlorine partitioning near the polar vortex edge observed with ground-based FTIR and satellites at Syowa Station, Antarctica, in 2007 and 2011, *Atmos. Chem. Phys.*, 20, 1043–1074, <https://doi.org/10.5194/acp-20-1043-2020>, 2020.
- 590 Nedoluha, G. E., Bevilacqua, R. M., and Hoppel, K. W.: POAM III measurements of dehydration in the Antarctic and comparison with the Arctic, *J. Geophys. Res.*, 107, 2002.
- Nedoluha, G. E., Connor, B. J., Mooney, T., Barrett, J. W., Parrish, A., Gomez, R. M., Boyd, I., Allen, D. R., Kotkamp, M., Kremser, S., Deshler, T., Newman, P., and Santee, M. L.: 20 years of ClO measurements in the Antarctic lower stratosphere, *Atmos. Chem. Phys.*, 16, 10 725–10 734, <https://doi.org/10.5194/acp-16-10725-2016>, 2016.
- 595 Ohneiser, K., Ansmann, A., Kaifler, B., Chudnovsky, A., Barja, B., Knopf, D. A., Kaifler, N., Baars, H., Seifert, P., Villanueva, D., Jimenez, C., Radenz, M., Engelmann, R., Veselovskii, I., and Zamorano, F.: Australian wildfire smoke in the stratosphere: the decay phase in 2020/2021 and impact on ozone depletion, *Atmos. Chem. Phys.*, 22, 7417–7442, <https://doi.org/10.5194/acp-22-7417-2022>, 2022.
- Pitts, M. C., Poole, L. R., and Thomason, L. W.: CALIPSO polar stratospheric cloud observations: second-generation detection algorithm and composition discrimination, *Atmos. Chem. Phys.*, 9, 7577–7589, 2009.
- 600 Portmann, R. W., Solomon, S., Garcia, R. R., Thomason, L. W., Poole, L. R., and McCormick, M. P.: Role of aerosol variations in anthropogenic ozone depletion in the polar regions, *J. Geophys. Res.*, 101, 22 991–23 006, 1996.
- Poshyvailo, L., Müller, R., Konopka, P., Günther, G., Riese, M., Podglajen, A., and Ploeger, F.: Sensitivities of modelled water vapour in the lower stratosphere: temperature uncertainty, effects of horizontal transport and small-scale mixing, *Atmos. Chem. Phys.*, 18, 8505–8527, <https://doi.org/10.5194/acp-18-8505-2018>, 2018.
- 605 Prather, M. J.: More rapid ozone depletion through the reaction of HOCl with HCl on polar stratospheric clouds, *Nature*, 355, 534–537, 1992.
- Robrecht, S., Vogel, B., Grooß, J.-U., Rosenlof, K., Thornberry, T., Rollins, A., Krämer, M., Christensen, L., and Müller, R.: Mechanism of ozone loss under enhanced water vapour conditions in the mid-latitude lower stratosphere in summer, *Atmos. Chem. Phys.*, 19, 5805–5833, <https://doi.org/10.5194/acp-19-5805-2019>, 2019.
- 610 Robrecht, S., Vogel, B., Tilmes, S., and Müller, R.: Potential of future stratospheric ozone loss in the midlatitudes under global warming and sulfate geoengineering, *Atmos. Chem. Phys.*, 21, 2427–2455, <https://doi.org/10.5194/acp-21-2427-2021>, 2021.
- Rolf, C., Afchine, A., Bozem, H., Buchholz, B., Ebert, V., Guggenmoser, T., Hoor, P., Konopka, P., Kretschmer, E., Müller, S., Schlager, H., Spelten, N., Sumińska-Ebersoldt, O., Ungermann, J., Zahn, A., and Krämer, M.: Transport of Antarctic stratospheric strongly dehydrated air into the troposphere observed during the HALO-ESMVal campaign 2012, *Atmos. Chem. Phys.*, 15, 9143–9158, <https://doi.org/10.5194/acp-15-9143-2015>, 2015.
- 615 Roy, R., Kumar, P., Kuttippurath, J., and Lefevre, F.: Chemical ozone loss and chlorine activation in the Antarctic winters of 2013–2020, *Atmos. Chem. Phys.*, 24, 2377–2386, <https://doi.org/10.5194/acp-24-2377-2024>, 2024.
- Sander, S. P., Friedl, R. R., Barker, J. R., Golden, D. M., Kurylo, M. J., Wine, P. H., Abbatt, J. P. D., Burkholder, J. B., Kolb, C. E., Moortgat, G. K., Huie, R. E., and Orkin, V. L.: Chemical kinetics and photochemical data for use in atmospheric studies, JPL Publication 10-6, 2011.
- 620 Santee, M., Manney, G., Lambert, A., Millan, L., Livesey, N., Pitts, M., Froidevaux, L., Read, W., and Fuller, R.: The Influence of Stratospheric Hydration from the Hunga Eruption on Chemical Processing in the 2023 Antarctic Vortex, ESS Open Archive, <https://doi.org/10.22541/essoar.170542085.55151307/v1>, 2024.

- Santee, M. L., Manney, G. L., Livesey, N. J., Foidevaux, L., MacKenzie, I. A., Pumphrey, H. C., Read, W. G., Schwartz, M. J., Waters, J. W., and Harwood, R. S.: Polar processing and development of the 2004 Antarctic ozone hole: First results from MLS on Aura, *Geophys. Res. Lett.*, 32, L12817, <https://doi.org/10.1029/2005GL022582>, 2005.
- 625 Santee, M. L., MacKenzie, I. A., Manney, G. L., Chipperfield, M. P., Bernath, P. F., Walker, K. A., Boone, C. D., Froidevaux, L., Livesey, N. J., and Waters, J. W.: A study of stratospheric chlorine partitioning based on new satellite measurements and modeling, *J. Geophys. Res.*, 113, D12307, <https://doi.org/10.1029/2007JD009057>, 2008.
- Santee, M. L., Lambert, A., Manney, G. L., Livesey, N. J., Froidevaux, L., Neu, J. L., Schwartz, M. J., Millán, L. F., Werner, F., Read, W. G., Park, M., Fuller, R. A., and Ward, B. M.: Prolonged and Pervasive Perturbations in the Composition of the Southern Hemisphere Midlatitude Lower Stratosphere From the Australian New Year's Fires, *Geophys. Res. Lett.*, 49, e2021GL096270, <https://doi.org/https://doi.org/10.1029/2021GL096270>, e2021GL096270 2021GL096270, 2022.
- 630 Schoeberl, M. R. and Dessler, A. E.: Dehydration of the stratosphere, *Atmos. Chem. Phys.*, 11, 8433–8446, <https://doi.org/10.5194/acp-11-8433-2011>, 2011.
- 635 Shi, Q., Jayne, J. T., Kolb, C. E., Worsnop, D. R., and Davidovits, P.: Kinetic model for reaction of ClONO₂ with H₂O and HCl and HOCl with HCl in sulfuric acid solutions, *J. Geophys. Res.*, 106, 24 259–24 274, <https://doi.org/10.1029/2000JD000181>, 2001.
- Smale, D., Strahan, S. E., Querel, R., Frieß, U., Nedoluha, G. E., Nichol, S. E., Robinson, J., Boyd, I., Kotkamp, M., Gomez, R. M., Murphy, M., Tran, H., and McGaw, J.: Evolution of observed ozone, trace gases, and meteorological variables over Arrival Heights, Antarctica (77.8°S, 166.7°E) during the 2019 Antarctic stratospheric sudden warming, *Tellus B: Chemical and Physical Meteorology*, 73, 1–18, <https://doi.org/10.1080/16000889.2021.1933783>, 2021.
- 640 Solomon, S.: Stratospheric ozone depletion: A review of concepts and history, *Rev. Geophys.*, 37, 275–316, <https://doi.org/10.1029/1999RG900008>, 1999.
- Solomon, S., Garcia, R. R., Rowland, F. S., and Wuebbles, D. J.: On the depletion of Antarctic ozone, *Nature*, 321, 755–758, 1986.
- Solomon, S., Borrmann, S., Garcia, R. R., Portmann, R., Thomason, L., Poole, L. R., Winker, D., and McCormick, M. P.: Heterogeneous chlorine chemistry in the tropopause region, *J. Geophys. Res.*, 102, 21 411–21 429, 1997.
- 645 Solomon, S., Portmann, R. W., Sasaki, T., Hofmann, D. J., and Thompson, D. W. J.: Four decades of ozonesonde measurements over Antarctica, *J. Geophys. Res.*, 110, D21311, <https://doi.org/10.1029/2005JD005917>, 2005.
- Solomon, S., Kinnison, D., Bandoro, J., and Garcia, R.: Simulation of polar ozone depletion: An update, *J. Geophys. Res.*, 120, 7958–7974, <https://doi.org/10.1002/2015JD023365>, 2015.
- 650 Sonnabend, J., Grooß, J.-U., Ploeger, F., Hoffmann, L., Jöckel, P., Kern, B., and Müller, R.: Lagrangian transport based on the winds of the icosahedral nonhydrostatic model (ICON), *Meteorol. Z.*, accepted, 2024.
- Spang, R., Hoffmann, L., Müller, R., Grooß, J.-U., Tritscher, I., Höpfner, M., Pitts, M., Orr, A., and Riese, M.: A climatology of polar stratospheric cloud composition between 2002 and 2012 based on MIPAS/Envisat observations, *Atmos. Chem. Phys.*, 18, 5089–5113, <https://doi.org/10.5194/acp-18-5089-2018>, 2018.
- 655 Stone, K. A., Solomon, S., Kinnison, D. E., and Mills, M. J.: On Recent Large Antarctic Ozone Holes and Ozone Recovery Metrics, *Geophys. Res. Lett.*, 48, <https://doi.org/10.1029/2021GL095232>, 2021.
- Strahan, S. E. and Douglass, A. R.: Decline in Antarctic Ozone Depletion and Lower Stratospheric Chlorine Determined From Aura Microwave Limb Sounder Observations, *Geophys. Res. Lett.*, 45, 382–390, <https://doi.org/https://doi.org/10.1002/2017GL074830>, 2018.

- Struthers, H., Bodeker, G. E., Austin, J., Bekki, S., Cionni, I., Dameris, M., Giorgetta, M. A., Grewe, V., Lefèvre, F., Lott, F., Manzini, E.,
660 Peter, T., Rozanov, E., and Schraner, M.: The simulation of the Antarctic ozone hole by chemistry-climate models, *Atmos. Chem. Phys.*,
9, 6363–6376, <https://doi.org/10.5194/acp-9-6363-2009>, 2009.
- Tilmes, S., Müller, R., Salawitch, R. J., Schmidt, U., Webster, C. R., Oelhaf, H., Russell III, J. M., and Camy-Peyret, C. C.: Chemical ozone
loss in the Arctic winter 1991–1992, *Atmos. Chem. Phys.*, 8, 1897–1910, 2008.
- Tilmes, S., Richter, J. H., Kravitz, B., MacMartin, D. G., Glanville, A. S., Visionsi, D., Kinnison, D. E., and Müller, R.: Sensitivity of total col-
665 umn ozone to stratospheric sulfur injection strategies, *Geophys. Res. Lett.*, 48, e2021GL094058, <https://doi.org/10.1029/2021GL094058>,
2021.
- Tritscher, I., Grooß, J.-U., Spang, R., Pitts, M. P., Poole, L. R., Müller, R., and Riese, M.: Lagrangian simulation of ice particles and resulting
dehydration in the polar winter stratosphere, *Atmos. Chem. Phys.*, 19, 543–563, <https://doi.org/10.5194/acp-19-543-2019>, 2019.
- Tritscher, I., Pitts, M. C., Poole, L. R., Alexander, S. P., Cairo, F., Chipperfield, M. P., Grooß, J.-U., Höpfner, M., Lambert, A., Luo, B. P.,
670 Molleker, S., Orr, A., Salawitch, R., Snels, M., Spang, R., Woivode, W., and Peter, T.: Polar Stratospheric Clouds: Satellite Observations,
Processes, and Role in Ozone Depletion, *Rev. Geophys.*, 59, <https://doi.org/10.1029/2020RG000702>, 2021.
- Várai, A., Homonnai, V., Jánosi, I. M., and Müller, R.: Early signatures of ozone trend reversal over the Antarctic, *Earth’s Future*, 3, 95–109,
<https://doi.org/10.1002/2014EF000270>, 2015.
- Vömel, H., Oltmans, S. J., Hofmann, D. J., Deshler, T., and Rosen, J. M.: The evolution of the dehydration in the Antarctic stratospheric
675 vortex, *Geophys. Res. Lett.*, 100, 13 919 – 13 926, 1995.
- von der Gathen, P., Kivi, R., Wohltmann, I., Salawitch, R. J., and Rex, M.: Climate change favours large seasonal loss of Arctic ozone, *Nat.*
Commun., 12, 3886, <https://doi.org/10.1038/s41467-021-24089-6>, 2021.
- von Hobe, M., Grooß, J.-U., Günther, G., Konopka, P., Gensch, I., Krämer, M., Spelten, N., Afchine, A., Schiller, C., Ulanovsky, A., Sitnikov,
N., Shur, G., Yushkov, V., Ravegnani, F., Cairo, F., Roiger, A., Voigt, C., Schlager, H., Weigel, R., Frey, W., Borrmann, S., Müller, R., and
680 Stroh, F.: Evidence for heterogeneous chlorine activation in the tropical UTLS, *Atmos. Chem. Phys.*, 11, 241–256, 2011.
- Ward, M. K. M. and Rowley, D. M.: Kinetics of the $\text{ClO} + \text{CH}_3\text{O}_2$ reaction over the temperature range $T = 250\text{--}298$ K, *Phys. Chem. Chem.*
Phys., 18, 13 646–13 656, <https://doi.org/10.1039/C6CP00724D>, 2016.
- Weber, M., Arosio, C., Coldewey-Egbers, M., Fioletov, V. E., Frith, S. M., Wild, J. D., Tourpali, K., Burrows, J. P., and Loyola, D.: Global
total ozone recovery trends attributed to ozone-depleting substance (ODS) changes derived from five merged ozone datasets, *Atmos.*
685 *Chem. Phys.*, 22, 6843–6859, <https://doi.org/10.5194/acp-22-6843-2022>, 2022.
- Wegner, T., Grooß, J.-U., von Hobe, M., Stroh, F., Sumińska-Ebersoldt, O., Volk, C. M., Hösen, E., Mitev, V., Shur, G., and Müller, R.:
Heterogeneous chlorine activation on stratospheric aerosols and clouds in the Arctic polar vortex, *Atmos. Chem. Phys.*, 12, 11 095–11 106,
<https://doi.org/10.5194/acp-12-11095-2012>, 2012.
- WMO: Scientific assessment of ozone depletion: 2022, GAW Report No. 278, Geneva, Switzerland, 2022.
- 690 Wohltmann, I., Lehmann, R., and Rex, M.: A quantitative analysis of the reactions involved in stratospheric ozone depletion in the polar
vortex core, *Atmos. Chem. Phys.*, 17, 10 535–10 563, <https://doi.org/10.5194/acp-17-10535-2017>, 2017.
- Wohltmann, I., von der Gathen, P., Lehmann, R., Maturilli, M., Deckelmann, H., Manney, G. L., Davies, J., Tarasick, D., Jepsen, N., Kivi,
R., Lyall, N., and Rex, M.: Near complete local reduction of Arctic stratospheric ozone by severe chemical loss in spring 2020, *Geophys.*
Res. Lett., 47, <https://doi.org/10.1029/2020GL089547>, 2020.
- 695 Wohltmann, I., Santee, M. L., Manney, G. L., and Millán, L. F.: The chemical effect of increased water vapor from the Hunga Tonga-Hunga
Ha’apai eruption on the Antarctic ozone hole, *Geophys. Res. Lett.*, 51, e2023GL106980, <https://doi.org/10.1029/2023GL106980>, 2023.

- Zafar, A. M., Müller, R., Groß, J.-U., Robrecht, S., Vogel, B., and Lehmann, R.: The relevance of reactions of the methyl peroxy radical (CH_3O_2) and methylhypochlorite (CH_3OCl) for Antarctic chlorine activation and ozone loss, *Tellus B: Chemical and Physical Meteorology*, 70, 1–18, <https://doi.org/10.1080/16000889.2018.1507391>, 2018.
- 700 Zhong, W. and Haigh, J. D.: Improved Broadband Emissivity Parameterization for Water Vapor Cooling Rate Calculations, *J. Atmos. Sci.*, 52, 124–138, [https://doi.org/10.1175/1520-0469\(1995\)052<0124:IBEPFW>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<0124:IBEPFW>2.0.CO;2), 1995.
- Zhou, X., Dhomse, S. S., Feng, W., Mann, G., Heddell, S., Pumphrey, H., Kerridge, B. J., Latter, B., Siddans, R., Ventress, L., Querel, R., Smale, P., Asher, E., Hall, E. G., Bekki, S., and Chipperfield, M. P.: Antarctic Vortex Dehydration in 2023 as a Substantial Removal Pathway for Hunga Tonga-Hunga Ha’apai Water Vapor, *Geophys. Res. Lett.*, 51, e2023GL107630 2023GL107630, 705 <https://doi.org/https://doi.org/10.1029/2023GL107630>, 2024.