

Systematic evaluation of the predictability of different Mediterranean cyclone categories

Benjamin Doiteau^{1,2}, Florian Pantillon¹, Matthieu Plu², Laurent Descamps², and Thomas Rieutord^{2,3}

¹Laboratoire d'Aérodynamique, Université de Toulouse, CNRS, UPS, IRD, Toulouse, France

²CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France

³Met Éireann, Dublin, Ireland

Correspondence: Benjamin Doiteau (benjamin.doiteau@aero.obs-mip.fr)

Abstract.

Cyclones are essential components of the weather patterns in the densely populated Mediterranean region, providing necessary rainfall for both the environment and human activities. The most intense of them also lead to natural disasters because of their strong winds and heavy precipitation. Identifying sources of errors in the predictability of Mediterranean cyclones is therefore essential to better anticipate and prevent their impact. The aim of this work is to characterise the medium-range cyclone predictability in the Mediterranean. Here, it is investigated in a systematic framework using the European Centre for Medium-Range Weather Forecasts fifth generation reanalysis (ERA5), and ensemble reforecasts in a homogeneous configuration over the 2001-2021 period. First, a reference dataset of 1960 cyclones is obtained for the period by applying a tracking algorithm to the ERA5 reanalysis. Then the predictability is systematically evaluated in the ensemble reforecasts. It is quantified using a new probabilistic score based on the error distribution of cyclone location and intensity (mean sea level pressure). The score is firstly computed for the complete dataset and then for different categories of cyclones based on their intensity, deepening rate, motion speed and on the geographic area and the season in which they occur. When crossing the location and intensity errors with the different categories, the conditions leading to a poorer or better predictability are discriminated. The motion speed of cyclones appears to be determinant in the predictability of the location, the slower the cyclone, the better the forecast location. Particularly, the position of stationary lows located in the Gulf of Genoa is remarkably well predicted. The intensity of deep and rapid-intensification cyclones, occurring mostly during winter, is for its part particularly poorly predicted. This study provides the first systematic evaluation of the cyclone predictability in the Mediterranean and opens the way to identify the key processes leading to forecast errors in the region.

1 Introduction

Extratropical cyclones are fundamental components of weather patterns in the mid-latitudes. The associated frontal systems provide the majority of the necessary rainfall (Hawcroft et al., 2012) but can also be at the origin of damaging storms (*e.g.* Roberts et al., 2014). A good representation of extratropical cyclones in numerical weather prediction systems is therefore essential to prevent their negative impacts, and identifying sources of forecast error is an important step to understand the processes leading to a poor predictability and improve forecasts.

25 In the Mediterranean, extratropical cyclones are generally smaller and with shorter lifetime than in other larger basins (Campins et al., 2011). However, they are at the origin of most of the high-impact weather events in the area, including intense rainfall (e.g. Flaounas et al., 2018), windstorms (e.g. Lfarh et al., 2023) and compound events (e.g. Raveh-Rubin and Wernli, 2016). The location of the Mediterranean between the tropics and the mid-latitudes, as well as the high mountain chains enclosing the basin, make it the site of complex interactions. The influence of Alpine lee-cyclogenesis (Trigo et al., 2002) and
30 Rossby wave breaking coming from the Atlantic (Raveh-Rubin and Flaounas, 2017) is clearly established in the formation of cyclones in the western part of the basin. Mediterranean cyclogenesis can also be influenced by other mountain ranges, the presence of both polar and subtropical jets, the entrance of Atlantic cyclones into the basin, or heat lows over land (see Flaounas et al., 2022, for a review).

Using a piecewise inversion of the potential vorticity equation, Flaounas et al. (2021) showed that intense Mediterranean
35 cyclones are influenced by two kinds of processes. On the one hand, the intrusion of a potential vorticity streamer in the upper-troposphere, related to deviation of the polar jet and to Rossby wave breaking, is identified as a principal dynamical contribution to cyclogenesis. On the other hand, diabatic processes, and in particular latent heat release, are important in the lower troposphere, where they act as a source of potential vorticity, reinforcing the cyclonic circulation. The relatively warm Mediterranean Sea can also lead to the formation of tropical-like cyclones, called medicanes, which received interest of the
40 scientific community in recent years (e.g. Miglietta et al., 2021). These phenomena can produce severe winds and rainfall, as in the cases of Ianos in September 2020 (Lagouvardos et al., 2022) or Daniel in September 2023. However, they are very rare with 1 to 2 events per year (Cavicchia et al., 2014). Thus, the statistical signal of medicanes can be considered as negligible. Therefore, the present study mainly focuses on the predictability of extratropical cyclones in the Mediterranean.

Limitations in the representation of cyclogenesis processes in numerical weather prediction systems can lead to forecast
45 errors propagating through lead times. Additionally, and beyond errors associated with the quality of the numerical model, the chaotic nature of the atmosphere leads to an intrinsic limit of predictability (Lorenz, 1969). More precisely, slight differences in the initial conditions can lead to radically different states of the atmosphere at increasing lead times. The forecast error is therefore due to both limitations in the representation of physical processes in the numerical model and to the chaotic nature of the atmosphere (practical and intrinsic predictability, respectively; see Melhauser and Zhang, 2012). In the following study, the
50 'practical predictability' will be denoted by 'predictability' for simplicity. Earlier work by Zhang et al. (2007) in an idealized baroclinic wave simulation, and by Baumgart et al. (2019) in hemispheric-wide simulations of PV structures, identified three phases in forecast error growth. In a first phase, errors in the representation of diabatic processes dominate in the first 12 h lead time. In a second phase, they are projected to the upper troposphere between 12 h and 2 days by tropospheric divergence. In a third phase, after 2 days lead time, the error growth is dominated by the upper-troposphere dynamics.

55 Ensemble prediction systems have been developed to provide an estimation of the forecast error growth. They offer a measure of forecast uncertainty and different possible scenarios from perturbed initial conditions and model parametrisations (Leutbecher and Palmer, 2008). This is crucial for extreme weather events, which are hardly sampled especially at longer lead times, ensemble prediction providing more robust results than a single deterministic forecast. For these reasons, ensemble prediction systems have long been proved useful for the early detection of extratropical cyclones and their associated hazards

60 (Buizza and Hollingsworth, 2002) or to assess the sensitivity of tropical cyclone genesis to the initial conditions (Torn and Cook, 2013). In the Mediterranean, studies based on ensemble forecasts revealed large uncertainty in the formation of medicane case studies and pointed its origin in error growth along the Rossby wave guide over the North Atlantic a few days ahead (Pantillon et al., 2013; Portmann et al., 2020).

To the best of the authors' knowledge, there is currently no systematic identification of error sources in the predictability of
65 Mediterranean cyclones. For instance, earlier work highlighted the crucial representation of upper-level dynamical precursors in the western Mediterranean (Argence et al., 2008; Vich et al., 2011) or cloud processes and air-sea interactions for medicanes (Miglietta et al., 2015; Tous et al., 2013) but these results relied on case studies. Using ensemble forecasts, Di Muzio et al. (2019) suggested the existence of a predictability barrier for the formation of several medicanes, but these rare events may not be representative of the broad spectrum of Mediterranean cyclones. Noteworthy, Picornell et al. (2011) assessed the determin-
70 istic forecast quality for more than 1000 extratropical cyclones during a whole year and found that the mean error in location increased from 50 km at 12 h to 118 km at 48 h lead time. However, the results were limited to relatively short forecast ranges and were not linked with the cyclone characteristics.

On a broader scale, Froude et al. (2007a, b) were among the first to investigate the predictability of extratropical cyclones in a systematic framework. They tracked the cyclones in global forecast data for two winter and two summer periods and defined
75 errors in both location and intensity based on maximum relative vorticity compared to analysis data. For the location, they found out that the error increases almost linearly at a rate of 1.25 geodesic degrees per day. For the intensity, they highlighted differences between summer and winter cyclones. In particular, intense storms occurring during the winter period were less accurately predicted, which was attributed to an incorrect representation of their vertical structure. More recent studies followed a similar approach and showed a systematic slow bias in the position of North Atlantic cyclones and a weak bias in intensity
80 of the deepest ones (Pirret et al., 2017; Pantillon et al., 2017). They explored links between the predictability and the dynamics of cyclogenesis but faced a robustness issue due to limited samples.

In this paper, ensemble reforecasts are used to systematically identify errors in the location and intensity of Mediterranean cyclones. The forecast model covers a 20-year period with the same configuration, which allows extracting statistically robust signals. The aim of the paper is to characterise the cyclone predictability in the Mediterranean region. Their representation
85 in an ensemble prediction system is discussed and the cyclone characteristics leading to a poorer or better predictability are identified. In particular, errors in the prediction of the cyclone location and intensity are evaluated for several categories of cyclones, based on their geographical location and seasonality, their intensity, deepening rate and motion speed.

The article is structured as follows. In Section 2, the data, cyclone tracking methods and tools to evaluate the predictability are described. The catalogue of Mediterranean cyclones and the associated climatology is presented in Section 3. The predictability
90 is then firstly evaluated for the whole dataset in Section 4, and secondly for specific categories of cyclones in Section 5. Section 6 contains a summary of the main results and the conclusion of the study.

2 Data and methods

2.1 Data for the reference tracks: ERA5 reanalysis

Reanalyses assimilate historical observation data spanning decades with a fixed assimilation scheme and forecast model. ERA5
95 (Hersbach et al., 2020) is the fifth reanalysis produced by the European Center for Medium-Range Forecasts (ECMWF). It is
based on the Integrated Forecast System (IFS, cycle 41r2), and includes models for atmosphere, land surface and ocean waves.
The horizontal resolution of the atmospheric model is about 31 km at mid-latitude, and it has 137 vertical levels from the
surface to 0.01 hPa. The reanalysis products are available globally with hourly resolution, from 1940 to present. In this study,
ERA5 is used from 2001 to 2021 with $0.25^\circ \times 0.25^\circ$ horizontal grid to produce a reference set of cyclone tracks on a domain
100 covering the Mediterranean (25° N - 50° N, 15° W - 45° E; see Fig. 1).

2.2 Tracking method for the reference tracks: the AYRAULT algorithm

Before investigating the predictability of Mediterranean cyclones, the first step is to produce a reference set of tracks. The
tracking method is based on the Ayrault (1998) algorithm (later AYRAULT), which has been implemented in the open-source
TRAJECT software (Plu and Joly, 2023). Originally designed for Atlantic cyclones in coarse model data, AYRAULT had to be
105 adapted for this study. As stated before, Mediterranean cyclones are generally smaller and have shorter lifetimes than those in
the Atlantic (Campins et al., 2011), and ERA5 has a higher spatio-temporal resolution than any previous reanalysis used with
the algorithm. Therefore, the parameters have been retuned specifically for both ERA5 and the Mediterranean region, starting
from the values used in Sanchez-Gomez and Somot (2018).

The main idea of AYRAULT is to track cyclones firstly in the relative vorticity field at 850 hPa. The horizontal wind is
110 then used at both 700 hPa and 850 hPa to choose the best following tracking point in the direction of cyclone propagation.
Finally, the track points are paired with the mean sea level pressure (MSLP) field. In the following, a time step is denoted by
 t , the relative vorticity field at 850 hPa by ζ , the zonal and meridional wind fields by u and v , respectively. AYRAULT can be
separated into five steps:

- 115 (1) Data preparation: a moving average with Gaussian weights is applied to ζ at 850 hPa and to u, v at 850 hPa and 700 hPa
to remove noisy features into these fields. The characteristic length in the weight decay is 225 km for ζ (to keep a
sufficient number of relevant vorticity cores), and 280 km for the wind fields (to keep the environmental wind and avoid
the vortex wind anomaly).
- (2) Detection of ζ -maxima: local maxima are detected in the ζ smoothed field. A single maximum (the strongest one) is
retained within a radius of 300 km.
- 120 (3) Loop over successive time steps: for every ζ -maximum at time t , a corresponding maximum at time $t + 1$ is searched for
using a three-steps method. First, the ζ -maximum at time t is advected by the wind at both 850 hPa and 700 hPa, giving
two guess positions for time $t + 1$. In a second step, a new ζ -maximum at time $t + 1$ is searched in the neighbourhood

of the two guessed points, within a radius of 300 km. Third and last, two quality criteria, based on the distance between the guessed point and the new ζ location and on the ζ value variation, must be fulfilled in order to keep a vortex core at $t + 1$. A cyclone track is finally defined by the successive positions of ζ maxima at every time step.

- (4) Pairing with MSLP: for every point belonging to the track, the local minimum of MSLP located within a 3° -square centred on the ζ -maximum becomes the new track point.
- (5) Validation criteria: the tracking process is stopped if the value of the ζ -maximum is less than $10^{-4} s^{-1}$ or if the MSLP minimum is greater than 1015 hPa. Among all tracks, only those which last for longer than 24 h and reach at least 1005 hPa along their lifetime are retained. This last criterion avoids most of the artefact cyclones. Indeed, some of the cases with their deepest MSLP over 1005 hPa appear to be local secondary lows caused by stronger storms crossing Northern Europe. Finally, an additional criterion is applied to only retain tracks entering into either the Mediterranean Sea or the Black Sea.

The Mediterranean-adapted version of AYRAULT previously described has been successfully tested with a slightly different configuration in an intercomparison of 10 tracking methods applied on ERA5 (Flaounas et al., 2023). The produced dataset remained close to the consensus between all algorithms in the spatial and seasonal distributions of cyclones. In the present study, our dataset is used as a reference instead of the consensus produced by Flaounas et al. (2023) for two principal reasons. First, the latter contains only 206 tracks in the highest confidence level (*i.e.* consensus of the 10 algorithms), which is not enough for a systematic study. At the mean confidence level (*i.e.* consensus of 5 over 10 algorithms), on the 2001-2021 period and with the same thresholds used here on the pressure and on the location of cyclones, 1231 tracks are detected in Flaounas et al. (2023), against 2853 with AYRAULT. Second, AYRAULT is conceptually similar to the tracking algorithm applied to the reforecasts (see Section 2.4), which reduces the influence of the tracking method on the results to focus on the predictability.

2.3 Data for the reforecast tracks: the IFS ensemble

Reforecasts are forecasts made retrospectively starting from historical initial conditions with a fixed model version. They are a key tool to investigate the predictability of the Mediterranean cyclones previously tracked in ERA5. The ECMWF ensemble reforecasts used here are constituted of 10 perturbed + 1 control members based on the IFS model (cycle 47r3) and initialized from ERA5 (Vitart et al., 2019). Initial perturbations on the reanalysis are constructed from the ERA5 ensemble data assimilation and singular vectors. Additionally, the model uncertainties are represented using a stochastically perturbed parametrisation tendency scheme (Buizza et al., 1999). The reforecasts used here cover an historical period of 20 years from October 2001 to October 2021, during which they are initialised every Monday and Thursday at 0000 UTC, leading to a total of about 2000 base times. The output spatial resolution of 0.25° is identical to the one in the ERA5 reanalysis. For each base time, a forecast output is available every 6 h (temporal resolution coarser than ERA5). Despite the maximum lead time of 14 days available with constant resolution in the reforecasts, the maximum lead time is restricted in this study to 144 h (6 days) because of the short lifetime of Mediterranean cyclones, considering that only less than 1 % of the cyclones of our reference dataset last longer than 6 days. The small number of ensemble members able to produce cyclone tracks at longer lead times (see Section 4.1) is

also pleading in favour of a limitation on the maximum lead time. Note that the same cyclone can be tracked in two successive forecast initialisations. When this happens, the two forecasts are treated independently.

2.4 Tracking method for the reforecast tracks: the VDG algorithm

In the reforecasts, the tracking of the cyclones is made with another algorithm (van der Grijn (2002); hereafter VDG), developed at the ECMWF and originally designed for the operational tracking of tropical cyclones. The VDG algorithm, also implemented in the open-source TRAJECT software (Plu and Joly, 2023), is similar to the previously applied (AYRAULT), as it also uses MSLP, the ζ smoothed field at 850 hPa and the horizontal wind at 850 hPa and 700 hPa. The main difference between the two algorithms is that VDG starts the tracking from a given geographical point or from an existing track. Cyclones detected in ERA5 are consequently directly linked with the reforecast by construction of VDG, as the position of the cyclone in the reforecast at the initial time $r(0)$, is directly dependent on the presence of a reference track at the same time. Applying AYRAULT to the reforecasts would have required an additional step for matching forecasted and observed cyclones, bringing more complexity.

At initialisation time, a ζ -maximum is searched for in the reforecast field, in the neighbourhood of the reference track calculated in ERA5. The tracking in VDG is then independent of the reference track, and is based on a combination of past movement and steering flow vector V_{av} defined as the layer average of the local wind fields at 850 hPa and 700 hPa. In the following, r and r_{fg} are respectively the positions of the cyclone and of the first guess. The initial step apart, the VDG algorithm can be divided as follows:

- (1) First guest: the steering flow V_{av} and the past movement $r(t) - r(t - 1)$ vectors are combined to obtain the first guess position of the next tracking point r_{fg} using the equation $r_{fg}(t + 1) = r(t) + w[r(t) - r(t - 1)] + (1 - w)V_{av} \delta t$, where w is a weight parameter ranging from 0 to 1 depending on the temporal resolution of the forecast δt , and here set to 0.4. *N.B.* at the first time step, only the steering flow vector is used (there is no past movement).
- (2) Detection of the ζ -maximum: a maximum is searched for in the ζ field within a square of 5° centred around the first guess.
- (3) Pairing with MSLP: another search is performed for the MSLP minimum within a same square of 5° , centred this time on the ζ -maximum. The location of this MSLP point finally becomes the next track point $r(t + 1)$.
- (4) Stopping criteria: the tracking of the cyclone is stopped when the value of the vorticity maximum ζ is less than the corresponding threshold of $10^{-4} s^{-1}$ or when the value of the MSLP minimum is greater than 1015 hPa, as in AYRAULT. This last criterion also implies that the tracking begins only if a MSLP minimum is found below the pressure threshold. The validation criteria assuring that cyclones last longer than 24 h and reach at least 1005 hPa along their lifetime, which were applied with AYRAULT to construct the reference dataset, are not applied here in the reforecasts.

185 2.5 Tracking algorithms comparison and final reference dataset

As demonstrated by Flaounas et al. (2023), using different cyclone tracking methods often leads to different results in the Mediterranean. In this study, 2853 cyclones are detected with AYRAULT in ERA5 on the 2001-2021 period, while cyclones are detected in the reforecasts using VDG starting from the reference tracks previously built. Using non-identical tracking methods for the reference and the reforecast tracks can introduce biases into the analysis. To assess the robustness of the results, VDG is also applied to the ERA5 data, using the tracks detected by AYRAULT as a reference. Note that VDG is applied to 6 h ERA5 data for consistency with the temporal resolution of the reforecasts for which it is tuned for. For each track detected by both algorithms, the difference in terms of location and intensity are calculated for all simultaneous track points. For 85 % of the tracks, no difference is found between the two algorithms. However, for 10 % of the dataset, the distance between AYRAULT and VDG tracks reaches almost 200 km at the time of minimum MSLP. To avoid this discrepancy, tracks are removed from the reference dataset if they are detected in ERA5 by AYRAULT but not by VDG (206 tracks), or if the maximal distance between them reaches more than 40 km (687 tracks). With these two criteria, the two algorithms provide identical tracks in 99 % of the dataset for both location and intensity. The following results are based on the remaining 1960 cyclones tracks that satisfy these two criteria.

2.6 Predictability metrics

200 The predictability is investigated using errors and spread in both location and intensity. The relationship between mean error and spread is used to verify the ensemble reliability before proceeding with further quantification of the predictability. For a reliable ensemble, one should expect mean error and spread to be comparable in magnitude.

In the following, errors are calculated by comparing the location and the intensity of each ensemble member with the corresponding reference track in ERA5, at each time t of the cyclone lifetime. The spread is for its part calculated from the pairwise difference between the members of the ensemble.

To assess the predictability of the cyclone location, we use the total track error (TTE) as defined in Froude et al. (2007b); Leonardo and Colle (2017). The TTE is also decomposed into an along-track error (ATE) and a cross-track error (CTE). A positive (negative) ATE stands for a forecast track ahead (behind) of the reference track, while a positive (negative) CTE stands for a forecast track on the left-hand side (on the right-hand side) of the reference track. Track errors (TTE, ATE and CTE) are calculated for each member individually and are presented in Section 4. Additionally, \overline{TTE} is here defined for each forecast cyclone as the mean of the TTEs of the members at each time t of the cyclone lifetime. The spread in location (hereafter σ_{loc}) is determined by averaging the distance between each pair of members as follows:

$$\sigma_{loc}(t) = \frac{1}{N(N-1)/2} \sum_{1 \leq i < j \leq N} d(r^i(t), r^j(t)) \quad (1)$$

where N is the number of members in which the cyclone is detected by the tracking algorithm at time t , r^i (r^j) is the position of the cyclone in the i -th member (in the j -th member) and d is the geodesic distance between the two positions.

Regarding the cyclone intensity, the MSLP error (hereafter MSLPE) is defined for each member as the difference between the MSLP of the member and the MSLP of the reference track at the same time. Unlike errors on the location, MSLPEs can also be negative. Consequently, $\langle \text{MSLPE} \rangle$ is defined as the root-mean-square of the MSLPEs over the members, for a specific track and at a specific time t of the cyclone lifetime. The spread in MSLP (hereafter σ_{int}) is for its part determined from the

220 root-mean-square of the differences between each pair of members as follows:

$$\sigma_{int}(t) = \sqrt{\frac{1}{N(N-1)/2} \sum_{1 \leq i < j \leq N} (p^i(t) - p^j(t))^2} \quad (2)$$

where p^i (p^j) is the MSLPE of the i -th member (j -th member).

An additional metric is defined to compare distributions of TTE or MSLPE between different categories of cyclones (see Section 5). In a preliminary step, for each category of cyclone, a cumulative density function (CDF) of errors is constructed

225 by taking into account every member of every cyclone track found at each lead time τ . CDFs of errors are then compared in a framework close to the continuous ranked probability score (CRPS) described in Candille et al. (2007). The metric denoted here by Cumulative Density Function Error (later CDFE) measures the distance between a CDF of errors and a virtual null-error distribution (100 % of the errors equal to 0):

$$CDFE(F_\tau) = \int [F_\tau(x) - 1_{x \geq 0}]^2 dx \quad (3)$$

230 where $F_\tau(x)$ is the CDF of the errors (either TTEs or MSLPEs) at a specific lead time τ and $1_{x \geq 0}$ stands for the Heaviside step function. Note that the CDFE metric has the same dimension as the variable on which it is applied. A higher (smaller) CDFE indicates a poorer (better) predictability. At each lead time, the statistical significance is evaluated using the Kolmogorov-Smirnov test, which in our case determines if two CDFs of errors are similar or not at a confidence level of 95%. This will ensure the robustness of the difference in the predictability of several categories of cyclones presented with the CDFE

235 metric.

3 Climatology of the reference dataset

This section provides the climatology of our reference dataset, based on the Mediterranean cyclones tracked with AYRAULT in ERA5 data and satisfying the two criteria of Section 2.5. In particular, the spatial distribution, the seasonal cycle, the intensity and the motion speed of cyclones are presented. Figure 1a shows the ground elevation over the Mediterranean and toponyms

240 that will be used in this manuscript.

3.1 Spatial distribution

For the whole 2001-2021 period, a total of 1960 cyclones are detected in the Mediterranean region, *i.e.* about 100 cyclones per year on average. The colour shading in Figure 1b accounts for the number of tracks having at least one track point within

a radius of 100 km divided by the total number of tracks. The figure can thus be seen as the relative frequency of cyclone
245 occurrence in our reference dataset, regardless of their stage of development. This spatial distribution is not homogeneous, as
the majority of cyclones are concentrated in preferred regions. In particular, six regions of interests, designed to cover equal
areas, are here identified by visual examination of the spatial distribution.

The six preferred regions concentrate 63 % of the cyclones of the dataset. The most active of them is the West Mediterranean
(22 %). It includes the Gulf of Genoa, in the lee of the Alps, which is recognized as the most cyclogenetic area (Trigo et al.,
250 2002). Then come the regions of the Adriatic (11 %), the East Mediterranean (10 %), the Black Sea (7 %) and the Sahara (7 %),
and finally the Middle East (6 %). The importance of the Alps in the formation of the West Mediterranean cyclones is clearly
established (Trigo et al., 2002). Horvath et al. (2008) show that lee cyclogenesis is also the dominant formation process for
Adriatic cyclones, whether they form in the Gulf of Genoa or in the Adriatic itself. The same orographic processes are known
to play a role in the formation of Saharan cyclones in the lee of Atlas mountains (Winstanley, 1972; Alpert and Ziv, 1989),
255 while Thorncroft and Flocas (1997) and Prezerakos et al. (2006) mostly highlighted the importance of interactions between
the polar and the subtropical jets in cyclogenesis in this particular region. For the Black Sea, and generally in the eastern parts
of the Mediterranean, Trigo et al. (2002) argued that cyclones are formed by different processes. In particular, they stated that
surface cyclones in the Black Sea seem to be associated with an upper trough in the west of the region, advecting vorticity
toward a relatively warm sea. Similar processes are found in the Aegean. The same authors argued that cyclones in the Middle
260 East are the manifestation of the extensions of the Asian trough in late spring.

The overall spatial distribution of our dataset is in agreement with previous studies (Alpert et al., 1990; Trigo et al., 1999;
Maheras et al., 2001; Campins et al., 2011; Lionello et al., 2016; Aragão and Porcù, 2022; Flaounas et al., 2023). However, two
minor differences remain. First, the hotspot in the western Atlas mountains and the high density of cyclones over the Iberian
Peninsula described in the literature do not appear here. This is mainly due to the criteria used to construct our dataset by
265 removing weak thermal lows with a pressure threshold of 1005 hPa on the one hand, and removing cyclones that do not enter
into either the Mediterranean Sea or the Black Sea on the other hand. Second, the high density of cyclones found here in the
Adriatic is not highlighted in the majority of previous studies.




Figure 1. (a) Elevation map over the Mediterranean domain with the toponyms mentioned in the text. (b) Relative frequency of Mediterranean cyclones, based on ERA5 over the 2001-2021 period, defined as the percentage of cyclones having a track point within a radius of 100 km. Regions of interest are framed by black boxes. Note that the shading scale is not linear.

3.2 Seasonal cycle

Figure 2 shows the number of cyclones striking any of the six regions of interest during each month of the year, averaged over the 20 years of our dataset. One can see that the number of cyclones in the Mediterranean is highly dependent on the season. The peak activity spans from November to May, while the period from June to October experiences fewer occurrences. However, this general trend is also dependent on the region considered.




Figure 2. Monthly number of cyclones in the six regions defined in Fig. 1b. Cyclones are counted at their minimum MSLP point and averaged over the 20-year period.

In the West Mediterranean and in the Adriatic, the cold season generally experiences more cyclones. Horvath et al. (2008) came to the same conclusion for the majority of Adriatic cyclones, while highlighting the importance of a subcategory of summer cyclones for their association with high-impact weather. In the East Mediterranean, more cyclones are also found during the cold part of the year. Saharan cyclones clearly exhibit a peak of occurrence in April and May, in agreement with previous studies (Winstanley, 1972; Alpert et al., 1990; Trigo et al., 2002). The Black Sea has a unique seasonal cycle, with higher activity during a long period spanning from December to July, with a peak in March. The presence of those cyclones during a large part of the year was already observed in Trigo et al. (1999). For the case of Middle East cyclones, higher occurrences are found here from March to May, while Trigo et al. (2002) found the peak of activity in August.

3.3 Intensity and deepening rate

Figure 3a shows the spatial distribution of the 10 % deepest cyclones of the reference dataset. They are mainly concentrated in the West Mediterranean and in the Adriatic, while some deep cyclones are found in the north-western parts of the Black Sea. The West Mediterranean and the Adriatic are also two hotspots for rapid intensification when looking at the deepening

285 rates (not shown). While cyclones in these two areas are influenced by the Atlantic (Raveh-Rubin and Flaounas, 2017), the origin of deep cyclones in the north-western Black Sea remains unclear. In this last region, cyclones do not experience rapid intensification, suggesting other processes of cyclogenesis. The shallowest cyclones are for their part concentrated in the Gulf of Genoa, highlighting the wide spectrum of intensities in this particular region. The other shallow cyclones are found mainly in the East Mediterranean and in the eastern parts of the Black Sea (not shown).

290 Figure 3b presents the typical seasonal cycle for three intensity-based categories of Mediterranean cyclones. Shallow and medium-intensity cases are more present during spring and exhibit a flat minimum from July to November. The 10 % deepest cyclones (green curve) show a more pronounced seasonal cycle, with very few cyclones during the warm part of the year, and a peak of activity from November to March. The similar pattern is observed for rapid-intensification cyclones, which are found almost exclusively during the cold part of the year (not shown).




Figure 3. (a) Relative frequency of Mediterranean cyclones, defined as the percentage of the 10 % deepest cyclones having a track point within a radius of 100 km. Note that the shading scale is not linear. (b) Monthly mean number of cyclones in the 3 categories of intensity. Each category contains 10 % of the dataset, *i.e.*, the 10 % deepest (green curve), the 10 % around median intensity (orange curve) and the 10 % shallowest cyclones (blue curve).

The motion speed of a cyclone is defined here by the median speed along its whole lifetime. According to our calculations on the reference dataset, Mediterranean cyclones move on average eastward at a median motion speed of 25 km h^{-1} . However, the variability is large and the fastest 5 % are moving at speeds greater than twice the median. Figure 4 shows the spatial distribution of the cyclones in each of the three motion speed categories: the 10 % fastest (Fig. 4a), the 10 % around the median
300 speed (Fig. 4b) and the 10 % slowest (Fig. 4c). The strong changes in spatial patterns between the different motion speed-based categories highlight the close relationship between the region in which the Mediterranean cyclone evolves and its motion speed.

The fastest cyclones (Fig. 4a) can be found in several particular areas. First, cyclones originating from the Sahara are clearly marked along an axis from the south of the Atlas mountains to the Ionian Sea. Cyclones in this region are also the fastest, with a median speed of 30 km h^{-1} . This result is in agreement with previous studies, which often highlight the high velocities
305 of Saharan cyclones compared to other Mediterranean lows (Alpert and Ziv, 1989; Kouroutzoglou et al., 2011). Second, fast Atlantic cyclones enter into the West Mediterranean, mainly from the Bay of Biscay and through the straight of Gibraltar. Third, another group of fast cyclones crosses the western Black Sea. Fourth and last, two other favourable regions for fast cyclones are found in the northern Adriatic and in western Greece. Medium speed cyclones (Fig. 4b) are for their part mainly located over sea, in the West Mediterranean, in the Adriatic and in the Ionian Sea. Finally, the slowest cyclones (Fig. 4c) are
310 clearly concentrated in the Gulf of Genoa, with median motion speed around 17 km h^{-1} . Some quasi-stationary lows can also be found in the eastern parts of the Black Sea. The location of these last quasi-stationary lows is contrasting with the fast cyclones observed over the western Black Sea (Fig. 4a), suggesting two different processes of cyclogenesis in the Black Sea region.




Figure 4. Relative frequency of occurrences (as defined in Fig. 3a) for the three motion speed-based categories. Each category contains 10 % of the dataset, *i.e.*, the 10 % fastest (a), the 10 % around the median speed (b) and the 10 % slowest cyclones (c). The black boxes are the regions of interests defined in Fig. 1b. Note that the shading scale is not linear.

4 Evaluation of the ensemble reforecasts

315 This section is dedicated to the evaluation of the Mediterranean cyclones representation in the reforecasts. Errors in the location and intensity, as defined in Section 2, are firstly evaluated by taking the tracks detected in ERA5 as reference, while the reliability of the ensemble reforecasts is assessed in a second step.

4.1 Location and intensity errors

To evaluate the reforecasts, both errors in location and intensity are considered. In Figure 5, distributions of errors are computed
320 at each lead time by taking into account the individual error of each member of the ensemble, for the entire dataset. The large number of 1960 cyclone tracks ensures the results to be robust. The mean number of members in which a cyclone is found by VDG decreases approximately linearly with lead time (orange curve in Fig. 5). While more than 9 members out of 11 detect a cyclone at the initial time, less than 4 members are remaining on average after 144 h lead time.

The TTEs distribution is presented for each lead time up to 144 h (Fig. 5a). Both median error and interquartile range are
325 increasing with lead time, with 50 % of the TTEs spanning from 80 km to 220 km after 72 h lead time. It is noticeable that the error growth is slower than linear at high lead times and seems to exhibit two phases: in the first 78 h, the median TTE increases of about 40 km per day, while from 84 h lead time and beyond it increases at a smaller rate of about 20 km per day. This behaviour can be explained by two different reasons. Firstly, the construction of VDG constrains the tracking to begin in the neighbourhood of the reference track. Consequently, when the lead time increases, the proportion of cyclones
330 followed since early lead times (*i.e.*, which forecast track may have diverged from the reference track) decreases in comparison to the ones followed since long lead times (*i.e.*, which forecast track is still in the neighbourhood of the reference track, by construction). It results that the error growth tends to decline with increasing lead time. The second process has to do with the error saturation. For long enough lead times, an ensemble forecast should ultimately converge to the climatological distribution. Consequently, the mean and median errors are expected to increase at a smaller rate at long lead times and saturate ultimately
335 at constant values.

Overall, the growth rate of 40 km per day in the first 78 h lead time is remarkably close to the 43 km per day found by Picornell et al. (2011) in the Mediterranean. The authors used for their part the ECMWF operational deterministic model during the 2006-2007 period and evaluated errors only during the first 48 h, which may explain the comparable error growth despite the older model version used in their study. In the extratropical Northern Hemisphere, and using the operational ensemble predic-
340 tion system of the ECMWF from January to July 2005, Froude et al. (2007b) found a much higher mean error growth rate of 1.25° (about 137 km) per day, almost constant until 7 days lead time. The coarser resolution of the ensemble prediction system used in their study (about 80 km) and the particular characteristics of Mediterranean cyclones could explain this difference in the mean error growth rate.

As presented in Section 2.6, the TTE can be decomposed into ATE and CTE. The ATE exhibits a weak and constant bias
345 of -15 km at 72 h lead time and beyond, indicating that forecast tracks have slow propagation speed bias compared to the reference (not shown). It is in agreement with Froude et al. (2007a), who highlighted that forecast cyclones in the IFS model

are on average getting too slow by about 1 km per hour compared to the analysis. Pirret et al. (2017) and Pantillon et al. (2017) found a systematic slow bias in the prediction of 60 and 25 severe European storms, respectively. The little bias found here in the ATE, however, is much smaller than in the previously mentioned studies. Regarding the CTE, a weak positive systematic shift is observed, growing at a constant rate of 4 km per day, indicating a weak shift to the left of the track (not shown). When looking into absolute values of ATE and CTE, it appears that the TTE is the result of an equivalent contribution of both components.

Errors in intensity (MSLPE) are presented in Fig. 5b. The bias reaches quickly -0.5 hPa in the first 12 h lead time, and forecasts continue to deviate at a very slow rate of -0.1 hPa per day until 144 h. This is pleading in favour of a well-centred error distribution of the ensemble reforecasts with a slight overestimation of the cyclone intensity, as in Froude et al. (2007b). After 72 h of forecast, 50 % of the MSLPEs are between -2.5 hPa and 1.5 hPa, and the interquartile range grows linearly of 0.9 hPa until the last lead time. When looking at absolute MSLPE (not shown), a little linear bias of 0.6 hPa per day is observed. Froude et al. (2007b) highlighted an even smaller bias around 0.2 hPa per day for the extratropical Northern Hemisphere. It could indicate a better prediction of the intensity of cyclones in other basins compared to the Mediterranean, however, the small magnitude of these biases should be considered.




Figure 5. Distributions of (a) total track errors (TTEs) and (b) MSLP errors (MSLPEs) relative to ERA5 as a function of lead time. Means are depicted by the blue circles, medians by the red lines, the first to third quartiles by grey boxes, and the minima and maxima by black whiskers. The orange curve is the mean number of members in which a cyclone is detected.

4.2 Reliability of the ensemble reforecasts

The reliability of the ensemble reforecasts is evaluated for both intensity and location by comparing for each cyclone and at a specific lead time the spread and the mean error of the ensemble, as defined in Section 2.6. One expects the mean error to be

close to the spread for a reliable ensemble, while a mean error greater (smaller) than the spread indicates an under-dispersive (over-dispersive) ensemble prediction system.




Figure 6. Spread-skill relationship at 72h lead time. The blue shading represents the number of cyclones populating each bin. (a) Mean of the TTEs of the members, denoted \overline{TTE} , compared to the spread in location, denoted σ_{loc} . The bin length is equal to 8 km. (b) Root mean square of the MSLPEs of the members, denoted by $\langle MSLPE \rangle$, compared to the spread in intensity, denoted σ_{int} . The bin length is equal to 0.2 hPa. The red curve represents an idealised, perfectly reliable set of ensemble reforecasts with a mean error equivalent to the spread. Percentages indicate the proportion of cyclones above and below the diagonal, respectively.

Figure 6 presents a comparison between the spread and the mean error of the ensemble at 72 h lead time, for the location (Fig. 6a) and for the intensity (Fig. 6b). Similar observations can be made for both aspects: firstly, the ensemble is reasonably reliable, with an identifiable linear relationship between spread and mean error (correlation coefficient equal to 0.65). Secondly, there is a slight but noticeable over-dispersion, with about 60 % of cyclone forecasts presenting a spread greater than the mean error. Finally, the mean error-over-spread ratio is equal to 1.12 for the location and 1.21 for the intensity, while the median ratio is equal to 0.90 in both cases. This indicates that while the ensemble tends to be over-dispersive in most forecasts, some of them are totally off, with a mean error much greater than the spread. It is noticeable that the opposite case with a spread much greater than the mean error is not really observed. Note that these three conclusions remain valid for all lead times (not shown).

375 5 Predictability of different categories of Mediterranean cyclones

In the previous section, the predictability was evaluated considering the complete dataset. In this section, cyclones are categorised following different features in order to determine the factors leading to a systematically better or poorer predictability. In particular, differences in the predictability are identified depending on the region, the season, the seasonality, the intensity and the motion speed of the cyclones.

The mean number of ensemble members in which a cyclone is detected (later denoted by number of members) is a key measure to investigate, as a high (low) number of members indicates a high (low) predictability. In Figure 7, the results are presented for different categories and are compared to the general behaviour of Mediterranean cyclones (shown by the black circles).




Figure 7. Difference in the mean number of ensemble members in which a cyclone is detected in the reforecasts: (a) for the six regions defined in Fig. 1b, (b) for seasonal categories (*e.g.* DJF stands for December January February), (c) for the intensity-based categories defined in Fig. 3b and (d) for the motion speed categories defined in Fig. 4. The mean number of members computed from the complete dataset is recalled by the black circles.

In Figure 7a., the number of members is presented as functions of the lead time for the regional categories of Fig. 1b. Most of the categories follow the general behaviour, except for the Sahara and the Middle East. In these two regions, the number of members quickly falls in the first 12 h lead time (particularly in the Middle East), and then decreases at a smaller rate until 144 h. An apparent diurnal cycle is visible with a lower number of members every 24 h, around 1200 UTC, corresponding to the warmest part of the day in these regions. The season in which the cyclone occurs has also an impact on the number of

members. In summer in particular (green curve on Fig. 7b.), the number of members decreases quickly in the first 18 h and then
 390 at a smaller rate until the maximum lead time. This is not the case for the other seasons, during which the decrease in number
 of members follows the general behaviour. Differences are also visible for different categories of intensity in Fig. 7c. The
 number of members detecting a cyclone is greater for deep cyclones, lower for shallow ones, and follows the general behaviour
 for medium-intensity cyclones. Finally, in Fig. 7d, the number of members is presented for three motion speed categories.
 Although differences are small, the slow and fast categories almost always lie below the general behaviour of the complete
 395 dataset, which is more closely followed by the medium-speed category.

Overall, the number of members falls down relatively strongly in the first 12 h for Saharan and Middle East cyclones (in
 warm regions) and in the first 18 h lead time for summer cyclones. The number of members in which a cyclone is detected is
 also lower for the shallowest cyclones between 18 h and 108 h lead time. The predictability in terms of number of members
 therefore seems to be linked with the intensity of the cyclones, which are often shallow during summer. Deep winter cyclones
 400 are for their part better predicted using this metric.

5.2 From CDFs of errors to CDFE scores

For each cyclone categorisation, CDFs of errors in both location (TTE) and intensity (MSLPE) are used to compute the CDFE
 metric presented in Section 2 at a specific lead time. It should be noted that the CDFE has the same unit as the variable
 considered. The greater (smaller) the CDFE, the poorer (better) the predictability of the cyclone category.




Figure 8. CDFs of the errors in location (a) and in the intensity (b) at 72 h lead time for the six regions defined in Fig. 1b.

405 To illustrate the approach, Fig. 8 presents CDFs of errors for the six regional categories presented in Section 3. In this
 representation, a category of cyclone is better predicted than another when the shape of its CDF of errors better resembles the
 Heaviside step function. At 72 h lead time and for the TTE (Fig. 8a), the East Mediterranean is the region in which cyclones
 are the least accurately predicted (orange curve), while the West Mediterranean cyclones have the smallest errors (blue curve).
 This is highlighted by the CDFE metric, with scores ranging from 51.8 km for the West Mediterranean to 94.2 km for the East

410 Mediterranean. In terms of MSLPE (Fig. 8b), Middle East cyclones are the best predicted with a CDFE of 0.42 hPa, while the Black Sea is the region in which the intensity of cyclones is the least accurately predicted at this particular lead time, with a CDFE equal to 0.61 hPa. In the next subsections, CDFE scores are computed at each lead time in order to compare the predictability between several categories of cyclones along the complete forecast duration considered.

5.3 Differences between regional categories

415 As shown in Fig. 1b, the spatial distribution of Mediterranean cyclones is not homogeneous, and six regions have been defined according to their cyclone density. Figure 9a presents the differences in the predictability of the cyclone location, using the CDFE metric applied at each lead time on the TTEs distributions of the six regions (colour curves). It immediately appears that the location of cyclones is the best predicted in the West Mediterranean at lead times beyond 42 h. The statistical significance of the difference between this region and any other is verified between 42 h and 120 h, except with the Middle East at 78–90 h lead time. The poorest predicted categories, namely the Adriatic and East Mediterranean cyclones, are in fact following the mean behaviour of the complete dataset (black circles) in the first 78 h. The difference between the best and the worst category is also noticeable and reaches more than 50 km at 144 h.




Figure 9. Differences in the predictability between the regions defined in Fig. 1b, for (a) and (c) the total track error (TTE), for (b) and (d) the MSLP error (MSLPE). The statistical significance is tested between each pair of categories, and results appear in thick lines when the considered category is significantly different from every other. CDFE scores computed from the complete dataset are represented by the black circles.

In Fig. 9b, the differences in the predictability considering the MSLPE are presented for the complete set of six regions. Regional differences are observed, in particular between the best and the poorest predicted categories. The Middle East is the region in which the intensity of cyclones is the best predicted at each lead time, probably linked with the absence of deep cyclones in this region (see Fig. 3a). A clear diurnal cycle is also observed, with local CDFE maxima at 66 h, 90 h, 108 h and 132 h lead time, corresponding to local times of 3 p.m. to 9 p.m. While the coarse temporal resolution of 6 h does not allow a precise timing of this behaviour, it seems that cyclones in this region are experiencing greater errors during the warm

part of the day. The cyclones in the Black Sea are the poorest predicted in the first 72 h, and a diurnal cycle is observed with two pronounced maxima at 36 h and 60 h, corresponding to the afternoon in this region. Trigo et al. (2002) already identified diurnal cycles in summer cyclones developing over northern Africa, the Iberian Peninsula, the Black Sea, and over the Middle East. The maximum intensity was reached during the afternoon, while cyclolysis generally occurred in the early morning. The reason for the diurnal cycle of errors shown here could be linked with the representation of the convective processes, often occurring during the afternoons of summer days.

435 5.4 Differences between seasonal categories

Another possible categorisation of Mediterranean cyclones is based on the seasonality. As previously visualised, Fig. 10a presents the CDFE score for the TTE and Fig. 10b for the MSLPE. In terms of errors of location, winter cyclones (December-January-February) are generally less well predicted than summer ones (June-July-August), except at 24–42 h. The results are statistically significant for these two extreme seasons in the first 84 h (not shown), but differences remain under 25 km before 120 h lead time. Consequently, the season in which the cyclone occurs does not appear to be determinant in the predictability of its location.




Figure 10. Differences in the predictability between the seasons (*e.g.* DJF stands for December January February), for (a) the total track error (TTE) and (b) the MSLP error (MSLPE). The statistical significance is tested between each pair of categories, and results appear in thick lines when the considered category is significantly different from every other. CDFE scores computed from the complete dataset are represented by the black circles.

Differences are more pronounced for the intensity, and they are statistically significant between winter and summer cyclones from 42 h until maximum lead time (not shown). CDFE scores in the autumn and spring follow the general behaviour of all Mediterranean cyclones (black circles), while errors are greater than average in winter and smaller than average in summer.

Differences in the predictability for different intensity-based categories are shown in Fig. 11. Considering the location, the predictability is the poorest for deep cyclones between 66 h and beyond (green curve in Fig. 11a). Meanwhile, location errors are independent of the deepening rate (Fig. 11c).




Figure 11. (a) and (b): Differences in the predictability between 3 intensity-based categories of Mediterranean cyclones following their minimum MSLP, namely the 10 % shallowest, the 10 % around median intensity and the 10 % deepest. (c) and (d): Same as (a) and (b) but based on the deepening rate, defined as the difference between the MSLP at the time of maximum intensity and 12 hours before. (a) and (c): Results for the total track error (TTE). (b) and (d): Results for the MSLP error (MSLPE). The statistical significance is tested between each pair of categories, and results appear in thick lines when the considered category is significantly different from every other. CDFE scores computed from the complete dataset are represented by the black circles.

In terms of MSLPE, deep cyclones are clearly poorer predicted than average after 66 h lead time (green curve in Fig. 11b). It is in agreement with Pantillon et al. (2017) and Pirret et al. (2017), who both showed a poor prediction of the intensity of the severe European storms they investigated. However, it should be noted that on average, the intensity of deep storms in our dataset is slightly over-predicted from 108 h onward (not shown), while it is slightly under-predicted in these two previous

studies. This difference could find an explanation in the region considered but also in the samples of studied cases, as Pantillon et al. (2017) and Pirret et al. (2017) find a slight under-prediction in a dataset of 25 and 60 extreme North Atlantic storms, respectively, while 280 less extreme Mediterranean cyclones are represented here in the deep cyclones' category. Regarding the two other categories, shallow cyclones are not necessarily better predicted than the medium category, and the difference is not always significant. The same conclusions can be drawn from the deepening rate (Fig. 11d), where rapid intensification cyclones strike out with intensity errors greater than in the other categories after 66 h lead time.

To summarise, the predictability is significantly poorer in terms of MSLPE for deep cyclones, at 66 h lead time and beyond. The same conclusions can be drawn from the deepening rate, but differences are not statistically significant. As seen in Section 3.3, these poorly predicted cyclones tend to form during the cold part of the year (Fig. 3b), in agreement with the poorest predictability of winter cyclones shown in section 5.4. They are also mainly located in the West Mediterranean and in the Adriatic, with a direct influence of the Atlantic (see Fig. 3a).

5.6 Differences between motion speed categories

It has been demonstrated in Fig. 4 that different motion speed-based categories of Mediterranean cyclones have different spatial distributions. It is consequently expected that differences will also appear in the predictability, which varies between regions (Fig. 9). Figure 12a presents the CDFE metric for a motion speed-based categorisation of cyclones.




Figure 12. Differences in the predictability between 3 motion speed-based categories, namely the 10 % slowest, the 10 % around median motion speed and the 10 % fastest moving cyclones, respectively. (a) for the total track error (TTE) and (b) for the MSLP error (MSLPE). The statistical significance is tested between each pair of categories, and results appear in thick lines when the considered category is significantly different from every other. CDFE scores computed from the complete dataset are represented by the black circles.

The link between the motion speed of the cyclone and the predictability of its location is remarkable, and differences are statistically significant from 12 h to 54 h lead time: the faster the cyclone, the poorer the predictability. The slow cyclones (blue curve) are clearly better predicted than any others. The difference with the two other categories is statistically significant

and increases with lead time, reaching almost 100 km after 120 h of forecast. The particularly good predictability of these slow cyclones has to be linked with the spatial distribution highlighted in Fig. 4c. Indeed, these quasi-stationary lows are in a vast majority concentrated in the Gulf of Genoa in the West Mediterranean, which is where the cyclone location is the best predicted (see Fig. 9). This result has to be compared with the predictability of the West Mediterranean cyclones' intensity, which is not particularly well predicted. It suggests the existence of at least two different types of cyclones in this particular region. The first is made of slow cyclones (Fig.4c) with a good predictability in terms of location and a fair predictability in terms of intensity. The second one is made of fast moving cyclones (Fig.4a) with a poor predictability in terms of intensity and a fair predictability in terms of location.

Unlike for the location, the motion speed of the cyclones does not play an important role in the predictability of the intensity (Fig. 12b) in the first 78 h. At lead times longer than 78 h, the fastest cyclones are the worst predicted, but the difference with the other categories is not statistically significant beyond 96 h, and does not allow building any robust conclusion.

6 Summary and conclusions

The predictability of extratropical cyclones can be highly variable from a case to another. Here, an approach based on the use of both reanalysis and ensemble reforecasts with a fixed model configuration over 20 years makes it possible to investigate the predictability of Mediterranean cyclones in a systematic framework.

Cyclones are first tracked in the ERA5 reanalysis, providing a large reference dataset of 1960 cyclones over the 2001-2021 period. Their spatial distribution is in agreement with most of the previous climatological studies, confirming the inhomogeneity in the distribution of Mediterranean cyclones. Six preferred regions concentrating 63 % of the dataset are identified, the Gulf of Genoa being the main hotspot in the region. In comparison to previous studies, a higher density of cyclones is found in the Adriatic. A clear seasonal cycle is highlighted, with a higher occurrence during the cold part of the year. The cold season is also more favourable to the development of intense cyclones, which mainly occur in the West Mediterranean, in the Adriatic, and in the north-western parts of the Black Sea.

Reference cyclones are then tracked in the homogeneous set of ensemble reforecasts for the same period. The predictability is evaluated in terms of errors in both cyclone location and intensity. Comparable magnitudes between mean error and spread indicate a reasonably good reliability of the IFS ensemble reforecasts for Mediterranean cyclones. A slight over-dispersion of the ensemble can however be observed at every lead time, whether in the location or in the intensity. It should also be noted that while the ensemble spread is slightly greater than the mean error in most forecasts, some cyclones remain very poorly predicted with median and mean errors that can be more than 4 times greater than the ensemble spread.

Considering the entire set of cyclones, it is shown that the median location error seems to grow at two different rates with increasing lead time. In the first about 3 days, the error grows at a nearly constant rate of 40 km per day, comparable to the one found in Picornell et al. (2011) for Mediterranean cyclones. The growth rate is however two times smaller for longer lead times. This behaviour is attributed to the progressive saturation of errors with lead time and to the limitation inherent to the verification of tracks against the reference. In terms of intensity error, the bias reaches quickly -0.5 hPa at 12 h lead

time, and forecasts continue to deviate at a slow rate of -0.1 hPa per day until the maximum lead time. This indicates a slight
505 overestimation of the intensity of forecast cyclones, in agreement with Froude et al. (2007b) for North Atlantic cyclones. This
result should be regarded with some caution, as reforecasts are not compared with observational data but with reanalysis data,
which may underestimate the actual cyclone intensity.

Looking at different categories of Mediterranean cyclones allows determining several factors contributing to a better or
poorer predictability. It is shown that the mean number of members in which the cyclone is detected is dependent on the
510 cyclone intensity. In particular, deep winter cyclones are detected in more members than shallower summer cyclones. In a
further step, the errors are summarized for the large number of cyclone forecasts by introducing a newly-defined CDFE score,
which is the CRPS applied to the error distributions of location (TTEs) and intensity (MSLPEs).

In terms of cyclone location, the motion speed appears to be a key factor. In particular, the slowest Mediterranean cyclones,
which are mainly located in the Gulf of Genoa are much better predicted than any other category, at every lead time. The
515 impact of such quasi-stationary cyclones can be considerable, as they can cause large amounts of accumulated precipitation in
the same area. The predictive skill in their location is therefore important. For their part, the location of the fastest cyclones is
relatively poorly predicted in the first 54 h lead time. To the authors' best knowledge, it is the first time that a link between the
cyclone motion speed and predictability is highlighted. The intensity of the cyclone also plays a role, and the location of deep
cyclones is less accurately predicted than in shallower categories, for lead times greater than 66 h.

Two factors leading to differences in the predictability of the cyclone intensity are clearly established. First, errors in the
520 intensity of deep cyclones are significantly greater than in any other category between 66 h and 108 h lead time. It is in
agreement with Froude et al. (2007a), who have shown a relatively poorer predictability for intense cyclones in the extratropical
Northern Hemisphere. This result is also observed here for the deepening rate, where the prediction of rapid-intensification
cyclones is the poorest, however, this result is not always statistically robust. A second important factor in the prediction of the
525 intensity is the season in which the cyclone occurs. Winter cyclones are indeed less accurately predicted than summer ones.
The difference between these two seasonal categories increases with lead time and is significant from 42 h until 144 h lead
time. In fact, the two factors are strongly related, as the deepest Mediterranean cyclones occur almost exclusively during the
cold part of the year. The forecast skill for the intensity of those strong winter cyclones is important, as some of them account
for the most destructive windstorms in the Mediterranean (*e.g.*, storm Klaus: Liberato et al., 2011). Froude et al. (2007a, b)
530 suggested that errors in the intensity of deep cyclones could originate from an incorrect representation of their vertical structure,
as the vertical tilt is known to play a major role in storm development. This hypothesis has to be verified systematically for the
Mediterranean.

In this study, the predictability has been quantified in a systematic framework for several categories of Mediterranean cy-
clones. The motion speed of the cyclone, its intensity, the season and the region in which it occurs are all playing a role. Further
535 investigations could focus on the physical processes responsible for the loss of predictability. In particular, the quantitative im-
portance of baroclinic and diabatic processes in the poor predictability of deep Mediterranean cyclones should be addressed.
Indeed, both the representation of latent heat release in the first forecast hours, and the location of Rossby wave breaking at
high lead times (several days), may be responsible for part of the loss of predictability of Mediterranean cyclones. It could

also be interesting to find a physical explanation to the remarkable good predictability of the shallow cyclones in the West
540 Mediterranean.

Code and data availability. The tracking algorithms are available in the open-source TRAJECT software <https://github.com/UMR-CNRM/Traject>. The ERA5 reanalysis is available through the Climate Data Store <https://cds.climate.copernicus.eu/>. The IFS reforecasts are available on the MARS Catalogue <https://apps.ecmwf.int/mars-catalogue/> (restricted access). Cyclones tracks are available on request.

Author contributions. BD performed the analysis and wrote the initial draft. FP prepared the reanalysis and reforecast data. MP developed
545 the tracking algorithms. TR provided expertise on statistical parts of this manuscript and LD on the evaluation of the ensemble prediction system. All authors participated in designing the study and preparing the final draft of the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This work was funded by Région Occitanie and Météo-France through project PREVIMED. It also received support
550 from the French National Research Agency under Grant ANR-21CE01-0002 and from COST Action CA19109 "MedCyclones: European Network for Mediterranean Cyclones in weather and climate". The authors thank the ECMWF and Copernicus for making the ERA5 reanalysis and the reforecasts available, as well as two anonymous reviewers for their constructive comments that helped improve the manuscript.

References

- Alpert, P. and Ziv, B.: The Sharav Cyclone: Observations and some theoretical considerations, *J. Geophys. Res.*, 94, 18 495–18 514, <https://doi.org/10.1029/JD094iD15p18495>, 1989.
- 555 Alpert, P., Neeman, B. U., and Shay-El, Y.: Climatological analysis of Mediterranean cyclones using ECMWF data, *Tellus*, 42, 65–77, <https://doi.org/10.1034/j.1600-0870.1990.00007.x>, 1990.
- Aragão, L. and Porcù, F.: Cyclonic activity in the Mediterranean region from a high-resolution perspective using ECMWF ERA5 dataset, *Clim. Dynam.*, 58, 1293–1310, <https://doi.org/10.1007/s00382-021-05963-x>, 2022.
- Argence, S., Lambert, D., Richard, E., Chaboureaud, J.-P., and Söhne, N.: Impact of initial condition uncertainties on the predictability of heavy rainfall in the Mediterranean: a case study, *Quart. J. Roy. Meteorol. Soc.*, 134, 1775–1788, <https://doi.org/10.1002/qj.314>, 2008.
- 560 Ayrault, F.: Environnement, structure et évolution des dépressions météorologiques : réalité climatologique et modèles types, Ph.D. thesis, <http://www.theses.fr/1998TOU30033>, 1998.
- Baumgart, M., Ghinassi, P., Wirth, V., Selz, T., Craig, G. C., and Riemer, M.: Quantitative View on the Processes Governing the Upscale Error Growth up to the Planetary Scale Using a Stochastic Convection Scheme, *Mon. Weather Rev.*, 147, 1713–1731, <https://doi.org/10.1175/MWR-D-18-0292.1>, 2019.
- 565 Buizza, R. and Hollingsworth, A.: Storm prediction over Europe using the ECMWF Ensemble Prediction System, *Meteorol. Appl.*, 9, 289–305, <https://doi.org/10.1017/S1350482702003031>, 2002.
- Buizza, R., Milleer, M., and Palmer, T. N.: Stochastic representation of model uncertainties in the ECMWF ensemble prediction system, *Quart. J. Roy. Meteorol. Soc.*, 125, 2887–2908, <https://doi.org/10.1002/qj.49712556006>, 1999.
- 570 Campins, J., Genovés, A., Picornell, M. A., and Jansà, A.: Climatology of Mediterranean cyclones using the ERA-40 dataset, *Int. J. Climatol.*, 31, 1596–1614, <https://doi.org/10.1002/joc.2183>, 2011.
- Candille, G., Côté, C., Houtekamer, P. L., and Pellerin, G.: Verification of an Ensemble Prediction System against Observations, *Mon. Weather Rev.*, 135, 2688–2699, <https://doi.org/10.1175/MWR3414.1>, 2007.
- Cavicchia, L., von Storch, H., and Gualdi, S.: A long-term climatology of medicanes, *Climate Dynamics*, 43, 1183–1195, <https://doi.org/10.1007/s00382-013-1893-7>, 2014.
- 575 Di Muzio, E., Riemer, M., Fink, A. H., and Maier-Gerber, M.: Assessing the predictability of Medicanes in ECMWF ensemble forecasts using an object-based approach, *Quart. J. Roy. Meteorol. Soc.*, 145, 1202–1217, <https://doi.org/10.1002/qj.3489>, 2019.
- Flaounas, E., Kotroni, V., Lagouvardos, K., Gray, S. L., Rysman, J.-F., and Claud, C.: Heavy rainfall in Mediterranean cyclones. Part I: contribution of deep convection and warm conveyor belt, *Clim. Dynam.*, 50, 2935–2949, <https://doi.org/10.1007/s00382-017-3783-x>,
- 580 2018.
- Flaounas, E., Gray, S. L., and Teubler, F.: A process-based anatomy of Mediterranean cyclones: from baroclinic lows to tropical-like systems, *Weather Clim. Dynam.*, 2, 255–279, <https://doi.org/10.5194/wcd-2-255-2021>, 2021.
- Flaounas, E., Davolio, S., Raveh-Rubin, S., Pantillon, F., Miglietta, M. M., Gaertner, M. A., Hatzaki, M., Homar, V., Khodayar, S., Korres, G., Kotroni, V., Kushta, J., Reale, M., and Ricard, D.: Mediterranean cyclones: current knowledge and open questions on dynamics, prediction, climatology and impacts, *Weather Clim. Dynam.*, 3, 173–208, <https://doi.org/10.5194/wcd-3-173-2022>, 2022.
- 585 Flaounas, E., Aragón, L., Bernini, L., Dafis, S., Doiteau, B., Flocas, H., Gray, S. L., Karwat, A., Kouroutzoglou, J., Lionello, P., Miglietta, M. M., Pantillon, F., Pasquero, C., Patlakas, P., Picornell, M. A., Porcù, F., Priestley, M. D. K., Reale, M., Roberts, M. J., Saaroni, H.,

- Sandler, D., Scoccimarro, E., Sprenger, M., and Ziv, B.: A composite approach to produce reference datasets for extratropical cyclone tracks: application to Mediterranean cyclones, *Weather Clim. Dynam.*, 4, 639–661, <https://doi.org/10.5194/wcd-4-639-2023>, 2023.
- 590 Froude, L. S. R., Bengtsson, L., and Hodges, K. I.: The Predictability of Extratropical Storm Tracks and the Sensitivity of Their Prediction to the Observing System, *Mon. Weather Rev.*, 135, 315–333, <https://doi.org/10.1175/MWR3274.1>, 2007a.
- Froude, L. S. R., Bengtsson, L., and Hodges, K. I.: The Prediction of Extratropical Storm Tracks by the ECMWF and NCEP Ensemble Prediction Systems, *Mon. Weather Rev.*, 135, 2545–2567, <https://doi.org/10.1175/MWR3422.1>, 2007b.
- Hawcroft, M. K., Shaffrey, L. C., Hodges, K. I., and Dacre, H. F.: How much Northern Hemisphere precipitation is associated with extratropical cyclones?, *Geophys. Res. Lett.*, 39, <https://doi.org/doi.org/10.1029/2012GL053866>, 2012.
- 595 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quart. J. Roy. Meteorol. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 600 Horvath, K., Lin, Y.-L., and Ivančan-Picek, B.: Classification of Cyclone Tracks over the Apennines and the Adriatic Sea, *Mon. Weather Rev.*, 136, 2210–2227, <https://doi.org/10.1175/2007MWR2231.1>, 2008.
- Kouroutzoglou, J., Flocas, H. A., Keay, K., Simmonds, I., and Hatzaki, M.: Climatological aspects of explosive cyclones in the Mediterranean, *Int. J. Climatol.*, 31, 1785–1802, <https://doi.org/10.1002/joc.2203>, 2011.
- 605 Lagouvardos, K., Karagiannidis, A., Dafis, S., Kalimeris, A., and Kotroni, V.: Ianos—A Hurricane in the Mediterranean, *Bull. Amer. Meteorol. Soc.*, 103, E1621 – E1636, <https://doi.org/10.1175/BAMS-D-20-0274.1>, 2022.
- Leonardo, N. M. and Colle, B. A.: Verification of Multimodel Ensemble Forecasts of North Atlantic Tropical Cyclones, *Weather Forecast.*, 32, 2083–2101, <https://doi.org/10.1175/WAF-D-17-0058.1>, 2017.
- Leutbecher, M. and Palmer, T.: Ensemble forecasting, *Journal of Computational Physics*, 227, 3515–3539, <https://doi.org/doi.org/10.1016/j.jcp.2007.02.014>, 2008.
- 610 Lfarh, W., Pantillon, F., and Chaboureau, J.-P.: The Downward Transport of Strong Wind by Convective Rolls in a Mediterranean Windstorm, *Mon. Weather Rev.*, 151, 2801–2817, <https://doi.org/10.1175/MWR-D-23-0099.1>, 2023.
- Liberato, M. L. R., Pinto, J. G., Trigo, I. F., and Trigo, R. M.: Klaus – an exceptional winter storm over northern Iberia and southern France, *Weather*, 66, 330–334, <https://doi.org/doi.org/10.1002/wea.755>, 2011.
- 615 Lionello, P., Trigo, I. F., Gil, V., Liberato, M. L. R., Nissen, K. M., Pinto, J. G., Raible, C. C., Reale, M., Tanzarella, A., Trigo, R. M., Ulbrich, S., and Ulbrich, U.: Objective climatology of cyclones in the Mediterranean region: a consensus view among methods with different system identification and tracking criteria, *Tellus*, 68, 29391, <https://doi.org/10.3402/tellusa.v68.29391>, 2016.
- Lorenz, E. N.: The predictability of a flow which possesses many scales of motion, *Tellus*, 21, 289–307, <https://doi.org/doi.org/10.1111/j.2153-3490.1969.tb00444.x>, 1969.
- 620 Maheras, P., Flocas, H., Patrikas, I., and Anagnostopoulou, C.: A 40 year objective climatology of surface cyclones in the Mediterranean region: spatial and temporal distribution, *Int. J. Climatol.*, 21, 109–130, <https://doi.org/10.1002/joc.599>, 2001.
- Melhauser, C. and Zhang, F.: Practical and Intrinsic Predictability of Severe and Convective Weather at the Mesoscales, *J. Atmos. Sci.*, 69, 3350–3371, <https://doi.org/10.1175/JAS-D-11-0315.1>, 2012.
- Miglietta, M. M., Mastrangelo, D., and Conte, D.: Influence of physics parameterization schemes on the simulation of a tropical-like cyclone in the Mediterranean Sea, *Atmos. Res.*, 153, 360–375, <https://doi.org/10.1016/j.atmosres.2014.09.008>, 2015.
- 625

- Miglietta, M. M., Carnevale, D., Levizzani, V., and Rotunno, R.: Role of moist and dry air advection in the development of Mediterranean tropical-like cyclones (medicanes), *Quart. J. Roy. Meteorol. Soc.*, 147, 876–899, <https://doi.org/10.1002/qj.3951>, 2021.
- Pantillon, F., Knippertz, P., and Corsmeier, U.: Revisiting the synoptic-scale predictability of severe European winter storms using ECMWF ensemble reforecasts, *Nat. Hazards Earth Syst. Sci.*, 17, 1795–1810, <https://doi.org/doi.org/10.5194/nhess-17-1795-2017>, 2017.
- 630 Pantillon, F. P., Chaboureaud, J.-P., Mascart, P. J., and Lac, C.: Predictability of a Mediterranean Tropical-Like Storm Downstream of the Extratropical Transition of Hurricane Helene (2006), *Mon. Weather Rev.*, 141, 1943–1962, <https://doi.org/10.1175/MWR-D-12-00164.1>, 2013.
- Picornell, M. A., Jansà, A., and Genovés, A.: A tool for assessing the quality of the Mediterranean cyclone forecast: a numerical index, *Nat. Hazards Earth Syst. Sci.*, 11, 1787–1794, <https://doi.org/doi.org/10.5194/nhess-11-1787-2011>, 2011.
- 635 Pirret, J. S. R., Knippertz, P., and Trzeciak, T. M.: Drivers for the deepening of severe European windstorms and their impacts on forecast quality, *Quart. J. Roy. Meteorol. Soc.*, 143, 309–320, <https://doi.org/10.1002/qj.2923>, 2017.
- Plu, M. and Joly, B.: Traject, <https://github.com/UMR-CNRM/Traject>, 2023.
- Portmann, R., González-Alemán, J. J., Sprenger, M., and Wernli, H.: How an uncertain short-wave perturbation on the North Atlantic wave guide affects the forecast of an intense Mediterranean cyclone (Medicane Zorbas), *Weather Clim. Dynam.*, 1, 597–615, <https://doi.org/doi.org/10.5194/wcd-1-597-2020>, 2020.
- 640 Prezerakos, N. G., Flocas, H. A., and Brikas, D.: The role of the interaction between polar and subtropical jet in a case of depression rejuvenation over the Eastern Mediterranean, *Meteorol. Atmos. Phys.*, 92, 1436–5065, <https://doi.org/10.1007/s00703-005-0142-y>, 2006.
- Raveh-Rubin, S. and Flaounas, E.: A dynamical link between deep Atlantic extratropical cyclones and intense Mediterranean cyclones, *Atmospheric Science Letters*, 18, 215–221, <https://doi.org/doi.org/10.1002/asl.745>, 2017.
- 645 Raveh-Rubin, S. and Wernli, H.: Large-scale wind and precipitation extremes in the Mediterranean: dynamical aspects of five selected cyclone events, *Quart. J. Roy. Meteorol. Soc.*, 142, 3097–3114, <https://doi.org/10.1002/qj.2891>, 2016.
- Roberts, J. F., Champion, A. J., Dawkins, L. C., Hodges, K. I., Shaffrey, L. C., Stephenson, D. B., Stringer, M. A., Thornton, H. E., and Youngman, B. D.: The XWS open access catalogue of extreme European windstorms from 1979 to 2012, *Nat. Hazards Earth Syst. Sci.*, 14, 2487–2501, <https://doi.org/10.5194/nhess-14-2487-2014>, 2014.
- 650 Sanchez-Gomez, E. and Somot, S.: Impact of the internal variability on the cyclone tracks simulated by a regional climate model over the Med-CORDEX domain, *Climate Dynamics*, 51, 1005–1021, <https://doi.org/10.1007/s00382-016-3394-y>, 2018.
- Thorncroft, C. D. and Flocas, H. A.: A Case Study of Saharan Cyclogenesis, *Mon. Weather Rev.*, 125, 1147–1165, [https://doi.org/10.1175/1520-0493\(1997\)125<1147:ACSOSC>2.0.CO;2](https://doi.org/10.1175/1520-0493(1997)125<1147:ACSOSC>2.0.CO;2), 1997.
- Torn, R. D. and Cook, D.: The Role of Vortex and Environment Errors in Genesis Forecasts of Hurricanes Danielle and Karl (2010), *Mon. Weather Rev.*, 141, 232–251, <https://doi.org/10.1175/MWR-D-12-00086.1>, 2013.
- 655 Tous, M., Romero, R., and Ramis, C.: Surface heat fluxes influence on medicane trajectories and intensification, *Atmos. Res.*, 123, 400–411, <https://doi.org/10.1016/j.atmosres.2012.05.022>, 2013.
- Trigo, I. F., Davies, T. D., and Bigg, G. R.: Objective Climatology of Cyclones in the Mediterranean Region, *J. Climate*, 12, 1685–1696, [https://doi.org/10.1175/1520-0442\(1999\)012<1685:OCOCIT>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<1685:OCOCIT>2.0.CO;2), 1999.
- 660 Trigo, I. F., Bigg, G. R., and Davies, T. D.: Climatology of Cyclogenesis Mechanisms in the Mediterranean, *Mon. Weather Rev.*, 130, 549–569, [https://doi.org/10.1175/1520-0493\(2002\)130<0549:COCMIT>2.0.CO;2](https://doi.org/10.1175/1520-0493(2002)130<0549:COCMIT>2.0.CO;2), 2002.
- van der Grijn, G.: Tropical cyclone forecasting at ECMWF: new products and validation, <https://doi.org/10.21957/e8525o38f>, 2002.

- Vich, M., Romero, R., and Brooks, H. E.: Ensemble prediction of Mediterranean high-impact events using potential vorticity perturbations. Part I: Comparison against the multiphysics approach, *Atmos. Res.*, 102, 227–241, <https://doi.org/10.1016/j.atmosres.2011.07.017>, 2011.
- 665 Vitart, F., Balsamo, G., Bidlot, J.-R., Lang, S., Tsonevsky, I., Richardson, D., and Alonso-Balmaseda, M.: Use of ERA5 reanalysis to initialise re-forecasts proves beneficial, <https://doi.org/10.21957/g71fv083lm>, 2019.
- Winstanley, D.: SHARAV, *Weather*, 27, 146–160, <https://doi.org/10.1002/j.1477-8696.1972.tb04279.x>, 1972.
- Zhang, F., Bei, N., Rotunno, R., Snyder, C., and Epifanio, C. C.: Mesoscale Predictability of Moist Baroclinic Waves: Convection-Permitting Experiments and Multistage Error Growth Dynamics, *J. Atmos. Sci.*, 64, 3579–3594, <https://doi.org/10.1175/JAS4028.1>, 2007.