

Surface processes and drivers of the snow water stable isotopic composition at Dome C, East Antarctica – a multi-datasets and modelling analysis

Inès Ollivier^{1,2}, Hans Christian Steen-Larsen¹, Barbara Stenni³, Laurent Arnaud⁴, Mathieu Casado²,
5 Alexandre Cauquoin⁵, Giuliano Dreossi³, Christophe Genthon⁶, Bénédicte Minster², Ghislain Picard⁴,
Martin Werner⁷ and Amaëlle Landais²

¹Geophysical Institute, University of Bergen, and Bjerknes Centre for Climate Research, Bergen, Norway

²Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, Gif sur Yvette, France

10 ³Department of Environmental Sciences, Informatics and Statistics, Ca' Foscari University of Venice, Mestre Venice, Italy

⁴Université Grenoble Alpes, CNRS, INRAE, IRD, Grenoble INP, IGE, Grenoble, France

⁵Institute of Industrial Science, The University of Tokyo, Kashiwa, Japan

⁶Laboratoire de Météorologie Dynamique, LMD/IPSL, Sorbonne Université-CNRS, Paris, France

⁷Alfred Wegener Institute (AWI), Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany

15 *Correspondence to:* Inès Ollivier (ines.ollivier@uib.no)

Abstract. Water stable isotope records in polar ice cores have been largely used to reconstruct past local temperatures and other climatic information such as evaporative source region conditions of the precipitation reaching the ice core sites. However, recent studies have identified post-depositional processes taking place at the ice sheet's surface modifying the original precipitation signal and challenging the traditional interpretation of ice core isotopic records. In this study, we use a
20 combination of existing and new datasets of the precipitation, snow surface and subsurface isotopic compositions ($\delta^{18}\text{O}$ and d-excess), meteorological parameters, ERA5 reanalyses, outputs from the isotope-enabled climate model ECHAM6-wiso, and a simple modelling approach to investigate the transfer function of water stable isotopes from precipitation to the snow surface and subsurface at Dome C, in East Antarctica. We first show that water vapor fluxes at the surface of the ice sheet result in a net annual sublimation of snow, from 3.1 to 3.7 mm water equivalent per year between 2018 and 2020,
25 corresponding to 12 to 15% of the annual surface mass balance. We find that the precipitation isotopic signal cannot fully explain the mean, nor the variability of the isotopic composition observed in the snow, from annual to intra-monthly timescales. We observe that the mean effect of post-depositional processes over the study period enriches the snow surface in $\delta^{18}\text{O}$ by 3.0‰ to 3.3‰ and lowers the snow surface d-excess by 3.4‰ to 3.5‰ compared to the incoming precipitation isotopic signal. We also show that the mean isotopic composition of the snow subsurface is not statistically different from
30 that of the snow surface, indicating the preservation of the mean isotopic composition of the snow surface in the top centimetres of the snowpack. This study confirms previous findings about the complex interpretation of the water stable

isotopic signal in the snow and provides the first quantitative estimation of the impact of post-depositional processes on the snow isotopic composition at Dome C, a crucial step for the accurate interpretation of isotopic records from ice cores.

1 Introduction

35 Polar ice cores have been widely used in paleoclimate studies to reconstruct past atmospheric conditions, up to 800 000 years back in time (EPICA community members 2004). Within the ice matrix of the core, $\delta^{18}\text{O}$ and δD (Craig, 1961) measurements have been commonly used as a proxy for past atmospheric temperatures based on the observed relationships between the local atmospheric temperature and both the isotopic composition of precipitation samples (Dansgaard, 1964) and the snow across spatial transects in Antarctica (Lorius et al., 1969, Masson-Delmotte et al., 2008).

40 The second order parameter deuterium excess (d-excess), defined as the deviation from the existing linear relationship between $\delta^{18}\text{O}$ and δD ($\text{d-excess} = \delta\text{D} - 8 \times \delta^{18}\text{O}$, Dansgaard, 1964), is driven by physical processes involving non-equilibrium, or kinetic fractionation of the different isotopes. The d-excess measured in ice cores has been interpreted as a proxy for moisture origin (Masson-Delmotte et al., 2005) and conditions at the moisture source region, such as sea-surface temperatures and relative humidity above the ocean's surface (Merlivat and Jouzel, 1979; Jouzel et al., 1982; Vimeux et al.,
45 1999; Stenni et al., 2001; Uemura et al., 2008, 2012; Landais et al., 2021; Steen-Larsen et al., 2014a). Further kinetic processes along the distillation path of an air mass have been identified to contribute to the d-excess signal in precipitation, such as condensation in supersaturated conditions (Jouzel and Merlivat, 1984) or mixing of air masses from different origins (Risi et al., 2013).

50 The reconstruction of the climatic parameters from the water isotopic records in polar ice cores relies on the assumption that the isotopic composition of precipitation is preserved from snowfall to burial and transformation into ice. However, this has been challenged by recent field studies highlighting the significant role of post-depositional processes at the surface of both the Greenland and Antarctic ice sheets modifying the isotopic composition of precipitation after snowfall (Touzeau et al., 2016; Münch et al., 2017; Casado et al., 2018, 2021; Hughes et al., 2021; Wahl et al., 2021, 2022; Zuhr et al., 2023). The
55 post-depositional processes commonly proposed to affect the water isotopes at the ice sheet's surface include (i) water vapor exchanges between the snow and the lower atmosphere through sublimation and condensation cycles, (ii) wind redistribution and (iii) diffusion of water vapor within the snowpack.

On the Greenland Ice Sheet, Steen-Larsen et al. (2014b) provided the first evidence of a co-variation of the snow surface and the lower atmosphere water vapor isotopic compositions during precipitation-free periods in the summertime, suggesting
60 seasonal vapor exchanges between the snow and the atmosphere. Wahl et al. (2021) later measured a depleted sublimation humidity flux compared to the snow surface, showing that fractionation of water isotopes was taking place during sublimation. Including fractionation during sublimation in a simple model also improved the prediction of the day-to-day variability in the snow isotopic composition during summertime (Wahl et al., 2022). Additional laboratory and modelling

studies showed that sublimation leads to an enrichment in $\delta^{18}\text{O}$ together with a lowering of d-excess in the snow surface and
65 in the firn (Hughes et al., 2021; Dietrich et al., 2023).

In addition, diffusion of water vapor within the snowpack is driven by temperature and isotopic gradients and affects the
isotopic composition of the snow and firn continuously (Johnsen et al., 2000; Gkinis et al., 2014). Field studies on both the
Greenland and Antarctic Ice Sheets identified snow metamorphism associated with water vapor diffusion within the top
layers of the snow to affect the snow isotopic composition ($\delta^{18}\text{O}$ and d-excess) after snowfall (Casado et al., 2021; Harris
70 Stuart et al., 2023).

Lastly, the wind blowing at the surface of the ice sheet leads to a heterogeneous accumulation by redistributing the snow on
the surface (Libois et al., 2014; Picard et al., 2019; Zuhr et al., 2021) which impacts the build-up of the isotopic signal in the
snow (Zuhr et al., 2023). Wind is also hypothesized to impact the snowpack isotopic composition through forced pumping
and ventilation of the snowpack (Town et al., 2008).

75 At Dome C, on the East Antarctic Plateau, previous studies have focused on qualitative description of the impact of post-
depositional processes on the snow surface at Dome C (Casado et al., 2018), monitoring the atmospheric water vapor, snow
surface and precipitation isotopic compositions (Casado et al., 2016; Touzeau et al., 2016; Stenni et al., 2016; Dreossi et al.,
2024a) or exploring the isotopic signature of snow metamorphism (Casado et al., 2021). However, a comprehensive
80 understanding of the formation of the isotopic signal in the snow, is still missing.

In this study we address the transfer function of stable water isotopes from precipitation to the snow surface and subsurface
at Dome C, from intra-monthly to multi-annual timescales. We use a combination of existing and new datasets of the
isotopic composition of precipitation, snow surface and snow subsurface over five consecutive years (2017-2021), ERA5
reanalysis products, outputs from the isotope-enabled climate model ECHAM6-wiso and a simple modelling approach to
85 investigate the contribution of precipitation to the variability observed in the snow surface and subsurface isotopic
composition. In addition, we use the meteorological parameters measured continuously on-site to estimate the magnitude of
sublimation and condensation fluxes between the surface and the lower atmosphere over three consecutive years (2018-
2020) and qualitatively evaluate their impact on the snow isotopic composition at Dome C.

2 Data and methods

90 2.1 Geographical settings

Dome C is located on the East Antarctic Plateau (75.1°S, 123.3°E), at 3233 m a.s.l. and 1000 km from the coast and is the
site where the permanent research station Concordia is installed (see location on Fig. 1a). The site is characterized by a mean
annual temperature of about -52°C (Genthon et al., 2021a) and a low accumulation rate of about 2.5 cm w.e. year⁻¹ (Genthon
et al., 2015). Due to the very small local slope and the location on a dome, the site is not subjected to strong katabatic winds,
95 the mean annual wind speed close to the surface is about 4 m s⁻¹ (Genthon et al., 2021a).




Figure 1. (a) Aerial view of Concordia station. The coloured circles indicate the location of the meteorological measurements used in this study (Sect. 2.4) and in red (rectangle and snowflake) the location of the samples presented in this study (Sect. 2.2 and 2.3). The wind rose for the five-year period 2017-2021 of the wind at 3 m is shown in the upper right corner. Background image from CNES (Pléiade satellite image of Concordia Station, Antarctica, CNES 2016, Distribution Airbus Defence and Space). (b) Snow sampling scheme taking place in the red rectangle in panel (a) (described in Sect. 2.2). SD stands for sampling day.

2.2 Snow surface and subsurface sampling

The regular sampling of the top few centimetres of the snow started in November 2013 and the sampling of a subsurface layer was added in 2017. Since then, the sampling strategy remained the same. In this study we focus on the 5-year period 2017-2021, where both surface and subsurface samples are available for analysis (see also Table 1).

The sampling takes place in the clean area about 800 m upwind of the main buildings (see location on Fig. 1a), twice a week and all-year round. The samples are taken at two different locations 50 m apart along a straight line, 3 to 10 m next to the line of the previous sampling day. At each location along the line, a small vertical snowpit is dug (about 20 cm deep). The snow surface and subsurface are collected with two 50 mL Corning tubes, placed horizontally from the snowpit wall, at the surface and just below. The two snow samples correspond to depths of 0 to 1 cm deep for the surface sample and from 1 to 4 cm for the subsurface sample, although the exact sampling depths was not recorded and may have slightly varied with the change of operator throughout the years or within one year due for instance to hard snow. The sampling scheme is illustrated in Fig. 1b.

Once the snow samples are collected, the tubes are sealed to prevent air exchange with the surrounding atmosphere and stored at temperatures well below freezing. The samples are shipped back once a year to LSCE (CNRS Paris-Saclay) to measure their isotopic composition with a laser spectrometer PICARRO L2130-i in liquid mode. We report the snow isotopic composition with delta-notation in ‰ (Craig, 1961) with respect to the Vienna Standard Mean Ocean Water (VSMOW) (Gonfiantini, 1978). The associated uncertainty (one standard deviation including quality control samples,

standards and sample replicates) on these measurements is $\pm 0.2\text{‰}$ for $\delta^{18}\text{O}$ and $\pm 0.7\text{‰}$ for δD , which yields an uncertainty
120 of $\pm 0.9\text{‰}$ for d-excess (one standard deviation).

2.3 Precipitation sampling

Since 2008, precipitation samples have been collected in the vicinity of Concordia station, as part of different projects
operated by the Italian Antarctic Research Programme (PNRA). Part of this long time series have been published in Stenni et
al. (2016) (2008-2010) and Dreossi et al. (2024a) (up to 2017). Here we extend the record of the precipitation isotopic
125 composition to 2021 and use the time series from 2017 to 2021 for our analysis (see also Table 1).

Precipitation samples are collected every day on a wooden platform (bench) 1 m above the ground, covered by a PTFE
surface and shielded by an 8 cm rail. The bench is situated about 800 m upwind of the main buildings (see location in Fig.
1a). The samples are collected at 10 AM local time (UTC+8), although it has varied throughout the years depending on the
operator. All the snow laying on the bench is collected, whether it is precipitation (including diamond dust), blown snow or
130 air hoar from atmospheric condensation. It cannot be ruled out that some of the samples might have undergone sublimation
especially during the summertime, because of exposure to 24-hour solar radiation before sample collection (Stenni et al.,
2016; Dreossi et al., 2024a). Each precipitation sample collected is weighted, and we use these weights as approximate
estimates of the precipitation amounts (details in Sect. 2.5).

After collection, the plastic bag containing the sample is sealed and stored at temperatures well below freezing before annual
135 shipment to the Ca'Foscari University of Venice, Italy, where the isotopic composition of the samples is measured with a
PICARRO laser spectrometers (L2130-i and L2140-i). We report the snow isotopic composition with delta-notation in ‰
(Craig, 1961) with respect to the Vienna Standard Mean Ocean Water (VSMOW) (Gonfiantini, 1978). The associated
uncertainty (one standard deviation of quality standard replicates) on these measurements is $\pm 0.1\text{‰}$ for $\delta^{18}\text{O}$ and $\pm 0.7\text{‰}$ for
 δD , which yields an uncertainty of $\pm 0.8\text{‰}$ for d-excess (one standard deviation).

140 2.4 Meteorological parameters

2.4.1 Atmospheric monitoring

Atmospheric parameters are measured continuously at Dome C by different weather stations and instruments installed
nearby Concordia station. For the 2017-2021 period of interest of this study, we used meteorological observations both from
an Automatic Weather Station operated by the PNRA (referred as AWSIT hereinafter) and from a 42 m meteorological
145 tower (referred as USt hereinafter). A summary of the meteorological parameters used in this study is available in Table 1.

The AWSIT is located about 800 m upwind of Concordia station (see location in Fig. 1a) and has been operating since 2005.
Hourly data from the AWSIT is available in Grigioni et al. (2022). In this study, we use the atmospheric pressure measured
at 1 m above the surface with a Vaisala PTB100 and a 3-month period of the atmospheric temperature measured at 1.5 m by

a Vaisala HMP45D. To match the same time step of the observations from the USt described below, we linearly interpolated
150 the data to 30 min.

The USt is part of the CALVA project and located about 1 km of Concordia station (see location in Fig. 1a). Meteorological instruments are installed at six different levels in the atmosphere measuring continuously for more than ten years (Genthon et al., 2021a). Due to snow accumulation, the sensors installed on the lowest level of the tower, at 3 m above the surface, were 40 cm closer to the surface in 2021 than in 2017. This height change was not considered here, and the lowest level of the
155 tower is referred to as the 3 m level. All atmospheric parameters are sampled at 30 s intervals, however in this study we use the 30 min averages.

The atmospheric temperature is measured by a PT100 in a Vaisala HMP155 combined sensor (thermo hygrometer) placed in a mechanically aspirated shield (Young 43502) and the wind speed and direction are measured by Young 05103 aerovanes. The quality-controlled (QC) data for temperature and wind speed from 2010 to 2019 is available in Genthon et al. (2021b, c).
160 Here we use this dataset from 2017 to 2019 and extend the record up to 2021 using the data from the same instruments available on the CALVA project website (see Table 1). Since the quality-control of this additional period (2019-2021, referred as non-QC) is not guaranteed, we compared the QC and non-QC datasets during an overlapping period (not shown). We found a linear correlation (Pearson correlation coefficient) of 1.0 between the datasets for both temperature and wind speed. Therefore, we use the temperature and wind records from the USt between 2017 and 2021 in our analysis. Note that to
165 fill a 3-month period of missing temperature data from the USt in 2021 (August to October), we use the temperature measured by the AWSIT (described above). The linear correlation (Pearson correlation coefficient) between the two temperature records during overlapping periods is 0.99 (not shown). The mean temperature and wind speed over the period 2017-2021 are summarized in Table 1. The dominant wind direction over the same period is shown in the windrose in Fig. 1a.

170 Atmospheric water vapor content (or humidity) is also measured continuously by sensors installed on the meteorological tower. At Dome C, the surface atmosphere is very cold and frequently above saturation (Genthon et al., 2017), and in these conditions the standard humidity sensors fail to accurately measure the true atmospheric humidity content. To cope with this issue, a modified HMP155 was designed and installed on the USt at 3 m above the ground, and proved its utility to measure atmospheric moisture accurately, capture supersaturation conditions and expand the temperature operating range of the
175 humidity sensor (Genthon et al., 2017, 2022). The sensor reports atmospheric humidity with respect to liquid (RH_{wrtl}) water even at temperature below 0°C, we therefore convert RH_{wrtl} to the relative humidity with respect to ice (RH_{wrti}), as in Genthon et al. (2017) and Vignon et al. (2022) (see also Supplement S1). The relative humidity at 3 m in the atmosphere and at 30 min resolution over the 2018-2021 period is available in Genthon et al. (2021d) and an analysis of the dataset is available in Genthon et al. (2022). Vignon et al. (2022) further provides estimations of the uncertainties for RH_{wrti} associated
180 with the temperature and humidity measurements. In this study, we use this 3-year atmospheric humidity record to estimate water vapor fluxes at the snow surface. The method is described in the following section. The mean RH_{wrti} between 2018 and 2020 is indicated in Table 1.

185 **Table 1.** Summary of samples and meteorological parameters used in this study and the reference to the data. The average values for temperature and wind speed are calculated over the 2017-2021 period; the relative humidity with respect to ice, the atmospheric pressure and the surface temperature are calculated over 2018-2020.

Project	Type/depth or height	Sampling rate/averaging time step	Average	Reference to dataset
NIVO	Snow samples: surface (0-1 cm) and subsurface (1-4 cm)	2x / week	–	Landais et al. (2024)
PRE-REC/ WHETSTONE	Precipitation samples	daily	–	Dreossi et al. (2024b)
CALVA (USt)	3 m temperature	30 min	-52.1°C	Genthon et al. (2021b) & CALVA website (see data availability section)
CALVA (USt)	3 m wind speed	30 min	3.9 m s ⁻¹	Genthon et al. (2021c) & CALVA website (see data availability section)
CALVA (USt)	3 m RH _{wrti}	30 min	104.5%	Genthon et al. (2021d)
AMCO (AWSIT)	Pressure at 1 m	1 h	642.6 hPa	Grigioni et al. (2022)
NIVO (CNR4)	Surface temperature	10 min	-55.4°C	* see data availability section

2.4.1 Estimation of water vapor flux

During the period of interest of this study, no direct Eddy-Covariance (EC) fluxes measurements were available at Dome C. We instead make use of the standard atmospheric parameters measured on site (described in the previous section) to apply the bulk method as described in Genthon et al. (2017) and estimate water vapor fluxes between the surface and the 3 m atmospheric level. We report the 30 min averages vapor fluxes during the 3-year period 2018-2020 in mm water equivalent per timestep.

The bulk method is based on the Monin-Obukhov (MO) Similarity Theory (Monin and Obukhov, 1954) and relies on several assumptions, which may not hold over the Antarctic Plateau (Vignon et al., 2016). Nevertheless, this method is still commonly used as the parametrization of surface turbulent fluxes in global and regional climate models (e.g. MAR model, Gallée and Schayes 1994) and have been compared against Eddy-Covariance measurements both at Dome C on the East Antarctic Ice Sheet (sensible heat fluxes, Vignon et al., 2016) and at EastGRIP on the Greenland Ice Sheet (water vapor fluxes, Dietrich et al., 2024). It requires the following parametrizations: (1) the choice of roughness length for momentum (z_0), (2) the choice of functions representing the atmospheric stability and (3) the calculation of the roughness lengths for water vapor (z_{0q}) and heat (z_{0t}). In their sensitivity study on the parametrization for sensible heat flux estimations at Dome C, Vignon et al. (2016) recommend the use of the stability functions from Holtslag and De Bruin (1988, referred hereinafter as H88) for stable cases and the functions from Högström (1996) for unstable cases. They also recommend the use of a constant z_0 of 0.56 mm, which corresponds to the average value observed with an EC system over one year at Dome C (Vignon et al., 2016). For the parametrization of z_{0q} and z_{0t} , we use the same approach as in Genthon et al. (2017) and King et al. (2001) where $z_0 = z_{0q} = z_{0t}$. We used this parametrization (H88, $z_0 = z_{0q} = z_{0t} = 0.56 \times 10^{-3}$ m) as the reference parametrization. In

addition, and similarly as in Vignon et al. (2016) and Genthon et al. (2017), we computed the water vapor fluxes using three other stability functions for stable conditions and a range of z_0 to estimate the sensitivity of the final flux calculations on the parametrization (fluxes computed 16 times, see Table 2 for stability functions and range of z_0). The range of z_0 tested corresponds to the observed range over one year at Dome C (Vignon et al., 2016).

210

Table 2. Set of roughness lengths for momentum z_0 and stability functions for stable conditions used to compute water vapor fluxes with the bulk method. The reference parametrization is highlighted in bold.

Roughness length for momentum z_0	0.01×10^{-3} m 0.56×10^{-3} m 1×10^{-3} m 6.3×10^{-3} m
Stability function for stable conditions	Holtslag and De Bruin (1988) – H88 Lettau (1979) Grachev et al. (2007) King and Anderson (1994)

215 To compute the water vapor fluxes, the bulk method requires temperature, wind speed, specific humidity, and pressure at the chosen atmospheric level (3 m here), as well as the snow surface temperature and the specific humidity at the surface.

For the atmospheric level, we use the temperature, wind and humidity sensors installed at 3 m above the surface on the USt together with the atmospheric pressure measured by the AWSIT (measurements described in Sect. 2.4.1). The formulas from Murphy and Koop (2005) are used to convert RH_{wrti} into specific humidity. To guarantee that the stationary conditions
220 required to apply the bulk method are met, we removed all 30 min temperature and wind speed data for which the differences in temperature and wind speed with the previous half-hour were above 2°C and 1.1 m s^{-1} , respectively (Vignon et al., 2016). This represents 4% of the whole dataset.

For the surface level, the snow surface temperature is computed from upward and downward longwave radiative fluxes with the same method as in Vignon et al. (2016) (their Eq. 1), using the same snow emissivity of 0.99 (value given by Brun et al.,
225 2011, used in Vignon et al., 2016 and Genthon et al., 2017). We use the longwave radiative measurements from a CNR4 radiometer installed approximately 500 m away from the USt (see location in Fig. 1a). We average the data over 30 min to match the time resolution of the atmospheric measurements (originally 10-min resolution). Note that to fill a 3-month period of missing data in the CNR4 record at the end 2020, we use the data provided by the BSRN network (Lupi et al., 2021). The BSRN data was corrected on the CNR4 data during overlapping periods beforehand, due to a shift identified in the upward
230 longwave flux measured by the BSRN pyrgeometer from December 2019 onwards. The mean surface temperature over the 2018-2021 time-period is reported in Table 1. The specific humidity at the snow surface is converted from the surface temperature using the formulas from Murphy and Koop (2005) and assuming saturation.

In total, due to gaps in the different input datasets and the removal of non-stationary data, the missing data in the bulk estimations represents 9% of the whole dataset.

235 2.5 Snow Isotopic Signal Generator (SISG)

To evaluate the contribution of precipitation to the isotopic variability observed in the snow surface and subsurface samples collected at Dome C (described in Sect. 2.2), we use a simple modelling approach to create synthetic snow layers solely based on the incoming precipitation. This approach was used in Casado et al. (2018, 2021) for a similar purpose, but focused on the top layer of the snowpack. Here we re-implemented the same simple model and added the snow subsurface.

240 The model (referred to as the Snow Isotopic Signal Generator – SISG) simulates snow layers by stacking precipitation events until the thickness of the stacked precipitation reaches the depths of the snow surface and subsurface layers given as input of the model. The isotopic composition of each snow layer is then calculated as the weighted average (by precipitation amounts) isotopic composition of all precipitation events necessary to build the snow layers. We choose the input snow layers depths to be 0 to 1 cm for the surface layer and 1 to 4 cm for the subsurface layer to match the snow samples collected
 245 at Dome C. We run the model at daily resolution over the 5-year period 2017-2021 and retrieve the model results for the same days as the observations. Table 5 summarizes the five model experiments performed with different inputs for the precipitation isotopic composition and precipitation amounts (described in Sect. 2.5.1 and 2.5.2).

250 **Table 3.** Experiments performed with the SISG model, with respective inputs for the daily precipitation amounts (described in Sect. 2.5.1) and the daily isotopic composition of precipitation (described in Sect. 2.5.2). Note that the precipitation amounts from the observations, ERA5 and ECHAM6-wiso are scaled to match the mean annual accumulation observed at Dome C.

Experiment	Daily precipitation amount	Daily precipitation isotopic composition
“Iso from T & cst accu”	Constant	Assuming constant iso vs temperature relationship
“Iso from T & ERA5 accu”	ERA5	Assuming constant iso vs temperature relationship
“Iso from wg. mm & obs accu”	Observations	Precipitation-weighted mean annual isotope cycle
“Iso from ar. mm & obs accu”	Observations	Arithmetic mean annual isotope cycle
“Iso from ECHAM6 & ECHAM6 accu”	ECHAM6-wiso	Modelled by ECHAM6-wiso

2.5.1 Daily precipitation amounts

255 The first SISG experiment uses a time series with a constant daily precipitation amount, calculated by dividing the mean annual accumulation at Dome C by 365 days. We use a mean annual accumulation of 2.5 cm w.e. year⁻¹ estimated from stake measurements (Genthon et al., 2015). This corresponds to 8 cm year⁻¹ using a snow density of 320 kg m⁻³, a typical value for the snow surface at Dome C (Picard et al., 2014; Genthon et al., 2015; Leduc-Leballeur et al., 2017). Hereinafter, we use this same snow density to convert precipitation amounts from snow water equivalent (SWE in mm w.e.) to snow depths (mm of snow) and inversely.

260 The second experiment uses the precipitation amount time series from ERA5 reanalysis (Hersbach et al., 2020, dataset available in Hersbach et al., 2023). We use the 24h-average of the hourly snowfall rate data for the grid point nearest Concordia station as an input to the model.

The third and fourth experiments use the observed precipitation amounts. As in Kopec et al. (2019), we assume that the weight of the daily precipitation samples collected on site for isotopic analysis (Sect. 2.3) is proportional to the amount of precipitation that has fallen over the day.

The fifth experiment uses the precipitation amounts given by the isotope-enabled global circulation model (GCM) ECHAM6-wiso (described in Cauquoin et al., 2019). The simulation was performed at a spatial resolution of 0.9° and nudged to ERA5 reanalyses (Cauquoin and Werner, 2021). The daily precipitation amounts were extracted for the grid point closest to Dome C. As the days with very low precipitation rates are not considered as “precipitation days”, any precipitation rate below $0.0016 \text{ mm w.e. day}^{-1}$ is set to zero. Because East Antarctica is a very dry region, this threshold was chosen to be ten times lower than the one commonly used for rain gauges ($0.5 \text{ mm w.e. month}^{-1}$, e.g. used for the Global Network of Isotopes in Precipitation data). Note that such a threshold was not applied to the precipitation amounts given by ERA5. The ECHAM6-wiso outputs are available in Cauquoin and Werner (2024).

Because the cumulative sum of the precipitation amounts according to the observations, ERA5 and ECHAM6-wiso was too low (18, 18, and 32 cm of snow after five years, respectively) compared to the mean accumulation over five years at Dome C (40 cm of snow), we scaled up all daily precipitation amounts in the three time series to match the mean annual accumulation (8 cm year^{-1}), acknowledging that this value is still underestimated because of net annual sublimation (Sect. 3.1). These scaled time series were used as inputs for the SISG model. The comparison of the three time series is shown in Fig. 5 (Sect. 3.3.1).

2.5.2 Daily precipitation isotopic composition

Due to some gaps in the daily precipitation samples collected at Dome C (described in Sect. 2.3), the time series of isotopic composition cannot be used as a direct input to the SISG model. Instead, we generate three artificial time series based on (1) the atmospheric temperature, (2) the precipitation-weighted mean annual isotope cycle in precipitation and (3) the arithmetic mean annual isotope cycle in precipitation. A comparison of these three artificial time series and the daily observations of the precipitation isotopic composition is shown in Fig. S4 (Supplement S4.1).

The first and second SISG experiment uses the artificial time series based on the atmospheric temperature. The precipitation isotopic composition is calculated from the atmospheric temperature using the linear relationships between $\delta^{18}\text{O}$ and δD of the precipitation samples and the 3 m daily mean temperature (Eq. 1 and 2 in Sect. 3.3.3). Deuterium excess is then calculated from the theoretical values of $\delta^{18}\text{O}$ and δD .

The third SISG experiment uses an artificial time series where all days in each month have the same isotopic composition than the corresponding monthly precipitation-weighted mean isotopic composition calculated over five years (results presented in Sect. 3.3.2, Fig. 6c and f).

The fourth experiment uses an artificial time series where all days in each month have the same isotopic composition than the corresponding monthly arithmetic mean isotopic composition calculated over five years (results presented in Sect. 3.3.2, Fig. 6b and e).

The fifth experiment uses the daily precipitation isotopic composition modelled by ECHAM6-wiso. To prevent any unrealistic values because of a numerical effect when the precipitation amounts are very low, days with precipitation amounts below 0.0016 mm w.e. (Section 2.5.1) were associated with missing values of the precipitation isotopic composition. In addition, all data points outside of the 5-year average \pm three standard deviation range were discarded (six data points for $\delta^{18}\text{O}$ and 20 data points for d-excess over five years). A comparison of the observed daily precipitation isotopic composition and ECHAM6-wiso simulations is shown in Fig. 6 (Sect. 3.3.2).

3 Results

3.1 Surface water vapor flux

The water vapor flux from the surface to the lower atmosphere for the period 2018-2020 estimated using the meteorological parameters measured at Dome C and the bulk method (described in Sect. 2.4.2) are shown in Fig. 2.




Figure 2. Water vapor flux at Dome C during the period 2018-2020 (positive for sublimation, negative for condensation). Panel (a) displays the 3-year time series of daily mean water vapor flux in mm water equivalent. The red line corresponds to the reference parametrization and the grey shading corresponds to the 10-90th percentile range of all parametrizations (described in Sect. 2.4.2). Panel (b) shows the net annual water vapor flux (sum of sublimation and condensation over one year). The red stars indicate the reference parametrization, and the three other black markers indicate the results using the stability function H88 with different z_0 . The whiskers of the grey boxplots indicate the 10-90th percentile range of all parametrizations, and the grey circles indicate the outliers outside of this range. The secondary axis to the right in blue gives water vapor fluxes in cm of snow, using a snow density of 320 kg m^{-3} to convert SWE to snow height.

During the 3-year period 2018-2020, the daily mean water vapor flux calculated with the reference parametrization varied from -0.05 (condensation) to 0.35 (sublimation) mm water equivalent per day (red line in Fig. 2a). The seasonality of water vapor fluxes over this period is characterised by sublimation during the summer months while little condensation is observed during the rest of the year. This seasonal pattern is observed independently of the parametrization used in the bulk method, which only affects the magnitude of the fluxes (grey shading in Fig. 2a).

The net annual water vapor flux between 2018 and 2020 is positive, meaning a net annual sublimation of snow, regardless of which parametrization is used (grey boxplots in Fig. 2b). In 2018, 2019, and 2020, water vapor fluxes calculated with the

reference parametrization led to a net mass loss of 3.7, 3.6, and 3.1 mm w.e., respectively (red stars in Fig. 2b), which is slightly higher than the net water vapor flux of 2.8 mm w.e. in 2015 (Genthon et al., 2017). These values correspond to 1.2, 1.1, and 1.0 cm of snow, respectively, using a snow density of 320 kg m^{-3} to convert from SWE to snow height. They are doubled when using a roughness length for momentum of $6.3 \times 10^{-3} \text{ m}$ instead of $0.56 \times 10^{-3} \text{ m}$ (black pentagons in Fig. 2b) and divided by approximately two when using a roughness length for momentum of $0.01 \times 10^{-3} \text{ m}$ (black triangles in Fig. 2b). The net annual water vapor fluxes are increased by 0.1 mm w.e. using a sensor height of 2 m above the surface instead of 3 m to consider height changes of the sensors (Sect. 2.4.1). During the summer periods only (from November to February, both months included), water vapor fluxes led to a net sublimation of 4.4, 4.1, and 3.8 mm w.e. in 2018, 2019, and 2020, respectively (not shown).

3.2 Snow isotopic composition

3.2.1 Temporal variations over five years

The five-year time series of the snow surface and subsurface isotopic composition ($\delta^{18}\text{O}$ and d-excess) is displayed in Fig. 3, together with the respective mean annual cycle (monthly means) for each layer over the whole period.




Figure 3. Observations of the snow surface and subsurface isotopic compositions at Dome C. Panels (a) and (c) show the 5-year time series (2017-2021) for $\delta^{18}\text{O}$ and d-excess (dxs) respectively. The circles represent the horizontal average between the two samples taken at the two locations 50 m apart on the sampling lines (described in Sect. 2.2) and the solid lines show a 30-day running mean of these horizontal averages. The shaded area represents the 30-day running average of the spatial spread between the two samples taken at the two locations 50 m apart on the sampling lines (one standard deviation). Panels (b) and (d) show the mean annual cycles (monthly means) of

the snow surface and subsurface layers calculated over the period 2017-2021 for $\delta^{18}\text{O}$ and d-excess respectively. The shaded area represents the inter-annual variability around the mean annual cycle (one standard deviation). The symbols on the right vertical axis indicate the mean isotopic composition of the snow surface and subsurface layers across all years. In all four panels, the darker colours correspond to the surface samples and the lighter colours to the subsurface samples.

345 Considering all samples collected during the five-year period, the $\delta^{18}\text{O}$ in the snow has a large amplitude, with values ranging from -60.9‰ to -45.1‰ in the surface layer and from -59.8‰ to -44.7‰ in the subsurface layer (circles in Fig. 3a). Both snow layers have a higher $\delta^{18}\text{O}$ during the summertime and lower values in the wintertime (lines in Fig. 3a).

350 This seasonality is further visible in their respective mean annual cycles (Fig. 3b), with the snow surface $\delta^{18}\text{O}$ highest in February (-47.8‰) and lowest in October (-53.6‰, dark blue in Fig. 3b). Compared to the snow surface, the mean annual cycle in the snow subsurface has a smaller amplitude and is shifted in time, with a maximum in March (-49.5‰) and a minimum in November (-53.1‰, light blue in Fig. 3b).

355 The temporal variation of the snow surface $\delta^{18}\text{O}$ is characterized by sharp increases during the summertime followed by slow decreases through the winter, which is particularly clear for the summers 2017-2018, 2018-2019 and 2019-2020 (dark blue line in Fig. 3a). This asymmetric seasonal pattern has been previously identified by Casado et al. (2018) for earlier years and is reflected in the mean annual cycle of the snow surface $\delta^{18}\text{O}$ (dark blue in Fig. 3b). A similar pattern is visible in the snow subsurface, although with a reduced amplitude (light blue in Fig. 3a and b).

360 The variations of $\delta^{18}\text{O}$ in the snow surface and the subsurface generally follow each other, except for specific periods when the surface and subsurface differ by several per mill, for example at the beginning of 2020 (solid lines and shaded areas in Fig. 3a). This difference between the two snow layers is reflected in their respective mean annual cycles and is the largest during the summer (Fig. 3b).

365 The overall mean $\delta^{18}\text{O}$ of the snow surface and subsurface is $-51.0 \pm 0.2\text{‰}$ and $-51.4 \pm 0.1\text{‰}$, respectively (dark and light blue dots in Fig. 3b). The uncertainty around these mean values corresponds to the standard error of the mean (SEM), calculated using the effective number of independent samples in the time series (Bretherton et al., 1999, Supplement S2).

370 As for $\delta^{18}\text{O}$, the snow surface and subsurface show large variations in d-excess over the five-year period (Fig. 3c). Considering all samples collected over the period, d-excess ranges from -0.9‰ to 21.0‰ in the snow surface and from 2.8‰ to 21.1‰ in the snow subsurface (dots in Fig. 3c). In opposition to $\delta^{18}\text{O}$, high d-excess values are encountered in the wintertime and lower d-excess in the summertime (lines in Fig. 3c).

375 This seasonality in d-excess is further reflected in the mean annual cycles of both snow layers (Fig. 3d). The snow surface d-excess is the highest in July (12.7‰) and the lowest in January (6.0‰, dark purple in Fig. 3d). As for $\delta^{18}\text{O}$, the mean annual cycle in the snow subsurface has also a smaller amplitude compared to the surface layer, with a maximum in July (12.2‰) and a minimum in February (8.9‰, violet in Fig. 3d). However, contrary to $\delta^{18}\text{O}$, the mean annual cycle in the snow subsurface does not show a clear time lag compared to the surface. Instead, the variations in the subsurface follows the ones of the surface, apart from the summer months of January and December where the subsurface has larger d-excess than the

surface layer (Fig. 3d). This summertime difference in d-excess between the surface and subsurface layers is also visible in the time series, in particular during the summer 2019-2020 (Fig. 3c).


380 Lastly, the variations in d-excess in the snow surface do not have the same pattern as for $\delta^{18}\text{O}$ (sharp increase in summertime and slow decrease in the wintertime). Instead, the d-excess in the snow surface shows a more symmetric seasonal evolution than $\delta^{18}\text{O}$ (dark purple line in Fig. 3c). This symmetry is reflected in the mean annual cycle in d-excess of the snow surface (Fig. 3d).

385 The overall mean d-excess of the snow surface and subsurface is $10.4 \pm 0.2\%$ and $10.8 \pm 0.1\%$, respectively (dark purple and violet dots in Fig. 3d). The uncertainty around these mean values corresponds to the SEM, calculated the same way as for $\delta^{18}\text{O}$.

3.2.2 Vertical difference between the snow surface and subsurface

In the previous section, we identified a seasonal pattern in the vertical difference between the snow surface and subsurface isotopic compositions ($\delta^{18}\text{O}$ and d-excess). The vertical difference is defined here as the isotopic composition of the surface layer minus the isotopic composition of the subsurface layer.

390 Considering all samples collected during the five-year period (displayed as dots in Fig. 3a and c), the minimum difference in $\delta^{18}\text{O}$ between the snow surface and the subsurface is -7.5% (surface depleted in $\delta^{18}\text{O}$ compared to subsurface) and the maximum difference is 9.1% (surface enriched in $\delta^{18}\text{O}$ compared to subsurface). The corresponding minimum and maximum values in the vertical difference in d-excess are -10.6% and 12.3% , respectively.



395 **Figure 4.** Mean annual cycle (monthly means) of the observed vertical difference between the snow surface and subsurface isotopic composition at Dome C. Panel (a) shows the mean annual cycle of the vertical difference in $\delta^{18}\text{O}$, panel (b) shows the mean annual cycle of the vertical difference in d-excess. In both panels, the black diamonds correspond to the overall monthly means calculated over the 2017-2021 period. The shaded area shows the inter-annual variability of individual years around the overall monthly means (one standard deviation) and the coloured markers show the monthly means for each individual year.

400

The mean annual cycles of the vertical difference in $\delta^{18}\text{O}$ and d-excess between the two depths show a clear seasonal pattern (Fig. 4a and b). The snow surface is relatively enriched in $\delta^{18}\text{O}$ from November to April and relatively depleted from May to October compared to the subsurface (black diamonds in Fig. 4a). The maximum vertical difference is 2.7‰ and occurs in January, the minimum difference is -1.0‰ and occurs in September. However, compared to the inter-annual variability within the averaging period (coloured markers and shaded area in Fig. 4a), only the months of January, February, September, and December show a substantial vertical difference in $\delta^{18}\text{O}$ between the snow layers.

In opposition to $\delta^{18}\text{O}$, the snow surface has a lower d-excess in the summer months of January and December, with a maximum monthly mean difference of -3.2‰ in January (black diamonds in Fig. 4b). The minimum difference of 0.9‰ is found in June and is, however, negligible compared to the inter-annual variability within the averaging period (coloured markers and shaded area in Fig. 4b).

3.3 Precipitation amounts and isotopic composition in observations, ECHAM6-wiso and ERA5 reanalyses

3.3.1 Precipitation amounts

To evaluate whether ERA5 reanalysis data and ECHAM6-wiso results correctly capture the precipitation amounts at Dome C, we compare them with the observations in Fig. 5 (all three time series scaled to the observed mean annual accumulation, Sect. 2.5).




Figure 5. Precipitation amounts at Dome C from observations, ERA5 reanalyses and ECHAM6-wiso simulation outputs. Panel (a) shows the mean annual cycles (monthly means) of the daily precipitation amounts (in mm w.e. per day) calculated over the period 2017-2021. Panel (b) shows the cumulative sum of precipitation over 5 years, reported as percentage of the total amount of precipitation (black and coloured thick lines), and the cumulative sum of days (black and coloured dotted lines) against the daily snowfall rate in mm w.e. per day. The red dashed lines guide the reading of the plot to get the snowfall rate and percentage of days responsible for 50% of the total snowfalls. Note the linear x-axis between 0 and 0.05 mm w.e. day⁻¹ and logarithmic above.

Over the whole 5-year period, the precipitation amounts in the observations, ERA5 and ECHAM6-wiso have a comparable
425 seasonal amplitude: from 0.03 to 0.1 mm w.e. day⁻¹ for the observations (black in Fig. 5a), from 0.04 to 0.11 mm w.e. day⁻¹
for ERA5 (blue in Fig. 5a) and from 0.04 to 0.1 mm w.e. day⁻¹ for ECHAM6-wiso (light green in Fig. 5a). The three
precipitation time series show a very similar seasonality, with an increase in snowfall rate at the end of the summertime
(from January to February) and in the middle of the winter (from June to July), as well as a decrease from July to December
(Fig. 5a).

430 All three precipitation cumulative sums given by the observations, ERA5 and ECHAM6-wiso show a similar shape with a
faster increase with increasing snowfall rates (plain lines in Fig. 5b). The cumulative sums from ERA5 and ECHAM6-wiso
are superposed in the whole range of snowfall rates, whereas the precipitation cumulative sum is lower for the observations
up to 0.35 mm w.e. day⁻¹, where the three curves meet.

In the observations, 50% of the total snowfalls over five years is due to precipitation events with a snowfall rate above 0.32
435 mm w.e. day⁻¹ (red dashed lines and black plain line in Fig. 5b), which represent only 6% of all days within the 5-year period
(red dashed lines and black dotted line in Fig. 5b). It should be noted that these results are dependent on the precipitation
samples collected on site, which have biases (i.e. too little precipitation to collect or blown snow instead of true precipitation,
Sect. 2.3). Similarly, in ERA5 precipitation, 50% of the total snowfalls is due to precipitation events with snowfall rates
above 0.27 mm w.e. day⁻¹ which corresponds to 7% of all days (Fig. 5b). In ECHAM6-wiso simulations, 50% of the total
440 snowfalls is due to precipitation events with snowfall rates above 0.28 mm w.e. day⁻¹ which corresponds to 6% of all days
(Fig. 5b).

For all three time series (observations, ERA5 and ECHAM6-wiso), we find that the largest precipitation events described
above (contributing to 50% of the total accumulation) occur alongside higher temperatures than average. In the observations,
the mean temperature during all precipitation days within the 5-year period and with snowfall rates above 0.32 mm w.e. per
445 day is 2.8°C warmer than the mean temperature over the whole period (-52.1°C). For ERA5, the mean temperature (given by
ERA5) during the largest precipitation events with snowfall rates above 0.27 mm w.e. per day is 8.1°C above the mean
temperature of -49.9°C. Lastly, for ECHAM6-wiso, the mean temperature (given by the model) during the largest
precipitation events with snowfall rates above 0.28 mm w.e. per day is 5.7°C above the mean temperature of -51.1°C. These
results are in agreement with previous studies (Kino et al., 2021; Servettaz et al., 2023).

450 3.3.2 Precipitation isotopic composition


The daily temporal variability of the precipitation isotopic composition ($\delta^{18}\text{O}$ and d-excess) from both observations and
ECHAM6-wiso simulations is presented in Fig. 6, together with the corresponding mean annual cycles over the same period.
In this section, all mean values across the whole period are given with an uncertainty corresponding to the SEM (see Sect.
3.2.1).

455 The observed precipitation $\delta^{18}\text{O}$ shows a large seasonal cycle, ranging from -82.6‰ to -21.8‰, with higher values in the
summertime and lower values in the wintertime (dark blue dots in Fig. 6a), following the atmospheric temperature (grey line

in Fig. 6a). The mean value over the whole period is $-56.2 \pm 0.5\text{‰}$ (dark blue dot in Fig. 6b). In comparison, the daily precipitation $\delta^{18}\text{O}$ modelled by ECHAM6-wiso show a similar seasonality as the observations, with higher and lower $\delta^{18}\text{O}$ in the summertime and wintertime respectively, and a comparable amplitude with values ranging from -82.9‰ to -22.8‰ (blue triangles in Fig. 6a). However, the mean modelled $\delta^{18}\text{O}$ in precipitation over the whole period is higher than the observed one ($-52.7 \pm 0.5\text{‰}$, blue triangle in Fig. 6b). The precipitation-weighted overall means are higher than the arithmetic means, for both observations and ECHAM6-wiso simulations ($-53.4 \pm 0.5\text{‰}$ and $-45.7 \pm 0.5\text{‰}$, respectively, dark blue dot and blue triangle in Fig. 6c).

The observed mean annual cycle (arithmetic monthly means) in the precipitation $\delta^{18}\text{O}$ is characterised by the highest $\delta^{18}\text{O}$ in December (-46.5‰) and the lowest $\delta^{18}\text{O}$ in June (-62.0‰ , dark blue dots in Fig. 6b). In comparison, the modelled mean annual cycle shows a similar seasonality but systematically higher than the observed one, ranging from -59.9‰ in June to -38.4‰ in December (blue triangles in Fig. 6b). The difference between the observations and ECHAM6-wiso results is especially large in the summertime (November to January), with a maximum difference of 8.1‰ in December.

The observed and modelled precipitation-weighted $\delta^{18}\text{O}$ mean annual cycles also show a similar seasonality (Fig. 6c). However, the bias between ECHAM6-wiso and observations is visible throughout the whole year, with a maximum difference of 10.2‰ in October (Fig. 6c). The large difference between the observations and ECHAM6-wiso during the summertime (arithmetic mean, Fig. 6b) decreases when the precipitation $\delta^{18}\text{O}$ is weighted by the precipitation amounts (Fig. 6c).




475 **Figure 6.** Observed and modelled precipitation isotopic composition at Dome C. Panels (a) and (d) show the 5-year time series (2017-2021) for $\delta^{18}\text{O}$ and d-excess (dxs) respectively. The dots represent the daily samples collected in the field (sample description in Sect. 2.3) and the triangles represent the daily precipitation isotopic composition modelled by ECHAM6-wiso (description of simulations in Sect.

2.5). In grey in panel (a) is displayed the atmospheric temperature measured at 3 m (described in Sect. 2.4.1). Panels (b) and (e) show the observed (dots) and modelled by ECHAM6-wiso (triangles) mean annual cycle, calculated over the period 2017-2021 (arithmetic means) for $\delta^{18}\text{O}$ and d-excess, respectively. The shaded area represents the inter-annual variability around the mean annual cycle (one standard deviation). The markers on the right vertical axis indicate the mean observed and simulated precipitation isotopic composition over the whole period. Panels (c) and (f) are the same as (b) and (e) but showing the weighted monthly means (by precipitation amounts).

The d-excess in the precipitation samples also shows a large seasonal amplitude, with values ranging from -38.6‰ to 65.9‰ (dark purple dots in Fig. 6d). d-excess is in anti-phase to $\delta^{18}\text{O}$, with maximum values found in the wintertime and minimum values in the summertime, which have been previously identified by Stenni et al. (2016) and Dreossi et al. (2024a). The mean observed precipitation d-excess over the whole period is equal to $15.2 \pm 0.5\text{‰}$ (dark purple dot in Fig. 6e). In comparison, the daily precipitation d-excess modelled by ECHAM6-wiso has a lower amplitude with values ranging from -26.0‰ to 37.2‰ (violet triangles in Fig. 6d). The overall mean modelled d-excess is also lower than the observations with a value of $5.8 \pm 0.3\text{‰}$ (violet triangle in Fig. 6e). The precipitation-weighted overall means are lower than the arithmetic means, for both observations and ECHAM6-wiso results ($12.2 \pm 0.5\text{‰}$ and $5.1 \pm 0.3\text{‰}$, respectively, dark purple dot and violet triangle in Fig. 6f).

The observed mean annual cycle (arithmetic monthly means) in the precipitation d-excess is characterised by the lowest value in December (0.1‰) and the highest value in June (22.3‰ , dark purple dots in Fig. 6e). In comparison, the mean annual cycle in the modelled precipitation d-excess has a lower amplitude than the observed one and a different timing in the minimum and maximum (from 1.0‰ in October to 10.1‰ in March, violet triangles in Fig. 6e). As opposed to $\delta^{18}\text{O}$, the difference between the observations and ECHAM6-wiso simulations is large in the wintertime (April to October), with a maximum difference of 15.9‰ in June.

Compared to the arithmetic mean annual cycles for d-excess, the bias between the precipitation-weighted monthly means in ECHAM6-wiso and the observations is reduced in the winter but increased in the summer months January and December (Fig. 6f). The maximum difference is also lowered to 12.6‰ and found later in the year (August, Fig. 6f).



505 **Figure 7.** Daily observed precipitation $\delta^{18}\text{O}$ (a) and d-excess (b) versus daily modelled precipitation isotopic composition at Dome C. All precipitation samples collected between 2017 and 2021 are shown (described in Sect. 2.3), together with the corresponding daily values from ECHAM6-wiso simulation results (described in Sect. 2.5). Note that due to missing observations and post-processing of ECHAM6-wiso outputs, 886/1826 days are shown in panel (a) and 882/1826 days in panel (b). The coloured lines correspond to the linear fits between the observations and the model results (slope coefficients given in legends, both linear regression coefficients are significant with a p-value < 0.001).

510

We evaluate the performance of ECHAM6-wiso to model the observed daily precipitation isotopic composition in Fig. 7. The daily precipitation $\delta^{18}\text{O}$ modelled by ECHAM6-wiso shows a good agreement with the observations, with a linear regression slope of 0.84 ± 0.03 (1.0 being the perfect fit), a Pearson correlation coefficient of 0.65 and a root mean square error (RMSE) of 8.8‰ (Fig. 7a). The model mostly overestimates the observations with an increase in bias towards more depleted values (Fig. 7a). For d-excess, the ECHAM6-wiso model results only poorly represent the observations, with a linear regression slope of 0.1 ± 0.02 , a Pearson correlation coefficient of 0.17 and a RMSE of 16.4‰ (Fig. 7b). These results are similar to the comparison of the precipitation samples and ECHAM6-wiso simulation results between 2008 and 2017 (RMSE=6.1‰ for $\delta^{18}\text{O}$ and 13.6‰ for d-excess, Dreossi et al., 2024a).

515

3.3.3 Isotope versus temperature relationships in precipitation

520 From the isotopic composition of the daily precipitation samples collected on site and the corresponding daily average temperature measured at 3 m above the surface (described in Sect. 2.4.1), we determine the following linear relationships over the 2017-2021 time-period:

$$\delta^{18}\text{O}_p = 0.47 \pm 0.01 \times T_{3m} - 31.2 \pm 0.6 \text{‰} \quad (1)$$

$$\delta\text{D}_p = 3.3 \pm 0.1 \times T_{3m} - 262 \pm 4 \text{‰} \quad (2)$$

525 Equation (1) has a coefficient of determination (R^2) of 0.62 and Eq. (2) has a R^2 of 0.63. Both linear regression slopes are significant (p-values < 0.001, Fig. A1). There are no significant changes in the linear relationships if using the daily mean atmospheric temperature of the day before the sample collection day instead of the daily mean temperature of the sampling day itself (Fig. A1). The slopes in Eq. (1) and (2) are also comparable to the ones found for the 2008-2010 period (0.49 ± 0.02 for $\delta^{18}\text{O}$, Stenni et al., 2016) and for the 2008-2017 period (0.52 ± 0.01 for $\delta^{18}\text{O}$, 3.52 ± 0.07 for δD , Dreossi et al., 530 2024a).

In ECHAM6-wiso simulations, the linear relationships between the modelled precipitation isotopic composition and the modelled temperature differ from the observed ones:

$$\delta^{18}\text{O}_{p, \text{ECHAM6wiso}} = 0.68 \pm 0.02 \times T_{2m, \text{ECHAM6wiso}} - 17.0 \pm 0.9 \text{‰} \quad (3)$$

$$\delta\text{D}_{p, \text{ECHAM6wiso}} = 5.3 \pm 0.1 \times T_{2m, \text{ECHAM6wiso}} - 134 \pm 6 \text{‰} \quad (4)$$

535 These relationships are also established over the period from 2017 to 2021. Equation (3) has a R^2 of 0.54 and Eq. (4) has a R^2 of 0.57. Both linear regression slopes are significant (p-values < 0.001, Fig. A2).

3.4 Contribution of precipitation to the snow isotopic composition

In this section we investigate the contribution of the precipitation isotopic composition to the intra-monthly and seasonal variability in the snow $\delta^{18}\text{O}$ and d-excess at Dome C, by comparing the observations and the results from the SISG experiments (described in Sect. 2.5).

3.4.1 Snow surface

Some of the features observed in the snow surface $\delta^{18}\text{O}$ variability are reproduced by the different experiments performed with the SISG model (Fig. 8a). For example, the observed summer $\delta^{18}\text{O}$ values in the snow surface are correctly reproduced in four out of five experiments (“iso from T & cst accu”, “iso from T & ERA5 accu”, “iso from wg. mm & obs accu”, “iso from ar. mm & obs accu”). All five experiments correctly reproduce the seasonality in $\delta^{18}\text{O}$ observed in the snow surface, with higher values in the summer and lower values in the winter (Fig. 8a and b). In addition, the two model experiments “iso from T & ERA5 accu” and “iso from ECHAM6 & ECHAM6 accu” reproduce some of the short-term increases in the snow surface $\delta^{18}\text{O}$ during the wintertime, such as the events in August 2018, July 2020, or August 2021 (orange shadings in Fig. 8a). The other three experiments fail to reproduce these events because of the inputs given to the SISG model: either a constant daily precipitation that weighs equally the days with high or low $\delta^{18}\text{O}$ (“iso from T & cst accu”) or a daily isotopic composition that does not vary within one month and fails to represent the high $\delta^{18}\text{O}$ events (“iso from wg. mm & obs accu” and “iso from ar. mm & obs accu”, Fig S4 in Supplement S4.1).

However, regardless of the model experiment, the amplitude of the seasonal cycle in the snow surface $\delta^{18}\text{O}$ is systematically overestimated compared to the observations (Fig. 8a and b). All five experiments fail to reproduce the slow decrease in $\delta^{18}\text{O}$ during the wintertime observed in the snow surface. This is particularly visible for the winters 2019 and 2020 where all simulated snow surface has too low $\delta^{18}\text{O}$ values (Fig. 8a). Most of the short-term variations in the snow $\delta^{18}\text{O}$ occurring within the month are also not reproduced by any of the experiments (Fig. 8a). In addition, the experiment “iso from ECHAM6 & ECHAM6 accu” gives too high $\delta^{18}\text{O}$ compared to the observations throughout the whole time-period (blue-green lines in Fig. 8a and b), which is consistent with the positive bias between the precipitation-weighted $\delta^{18}\text{O}$ modelled by ECHAM6-wiso and the observations, identified in Sect. 3.3.2 (Fig. 6c).

As for $\delta^{18}\text{O}$, all model experiments fail to reproduce the variability in d-excess observed in the snow surface. Although in some of the experiments (“iso from T & cst accu”, “iso from T & ERA5 accu”, “iso from wg. mm & obs accu”, “iso from ar. mm & obs accu”), the low summer d-excess in the snow surface is well captured (Fig. 8c), all the simulated mean annual cycles have a too large amplitude (Fig. 8d). The experiment “iso from ECHAM6 & ECHAM6 accu” provides worse results than the other experiments (pink line in Fig. 8c and d), which can be explained by the poor representation of the observed precipitation-weighted annual cycle in d-excess by ECHAM6-wiso simulations, identified in Sect. 3.3.2 (Fig. 6f).




Figure 8. Comparison between observed and simulated snow surface (0 to 1 cm depth) isotopic compositions at Dome C. Panels (a) and (c) show the 5-year time series (2017-2021) of $\delta^{18}\text{O}$ and d-excess (dxs) respectively. The black dots represent the horizontal average between the two samples taken at the two locations 50 m apart on the sampling lines (described in Sect. 2.2, also presented in Fig. 3). The coloured lines (dotted and plain) represent the different SISG model experiments (described in Table 3). In panel (a), the orange vertical shadings highlight three specific events (August 2018, July 2020, August 2021). Panels (b) and (d) show the mean annual cycles (monthly means) of $\delta^{18}\text{O}$ and d-excess, respectively, calculated over the period 2017-2021. The black dots and line represent the observations and the shaded area represents the inter-annual variability around the mean annual cycle (one standard deviation). The coloured lines (dotted and plain) represent the different SISG experiments (described in Table 3). The markers on the right vertical axis indicate the mean isotopic composition of the observed and simulated snow surface over 5 years (see Table S2 in Supplement S4.2).

To evaluate which SISG experiment best represent the observed snow surface isotopic composition, we perform a linear regression between all modelled and observed monthly means $\delta^{18}\text{O}$ and d-excess in the five-year period. The linear regression slopes (a) and the RMSE between the model results and the observations are summarized in Table 4.

The best representation of the observed $\delta^{18}\text{O}$ in the snow surface is given by the experiment “iso from wg. mm & obs accu” (a=1.1, RMSE=3.4‰, Table 4). The experiment “iso from T & cst accu” results in the highest RMSE of all configurations

585 tested (RMSE=6.9‰, Table 4), which is expected due to the intermittent nature of precipitation at Dome C. In addition, the experiment “iso from ECHAM6 & ECHAM6 accu” gives a high RMSE (6.3‰, Table 4). For d-excess, out of the five model experiments, “iso from wg. mm & obs accu” gives the lowest RMSE (3.9‰, Table 4). As for $\delta^{18}\text{O}$, the experiment “iso from T & cst accu” results in the highest RMSE (8.8‰, Table 4). Lastly, the experiment “iso from ECHAM6 & ECHAM6 accu” has a RMSE of 7.1‰ and a non-significant slope compared to the observations (p-value > 0.05).

590

Table 4. Linear regression slope (a) and RMSE between the observed and modelled monthly mean isotopic composition of the surface layer ($\delta^{18}\text{O}$ in normal font, d-excess in parenthesis and italic). The sample size is 60 for all experiments except “iso from wg. mm & obs accu” and “iso from ar. mm & obs accu” (n=59). All linear slopes are significant (p-value < 0.05), except the ones marked with an asterisk (*, p-value > 0.05).

595


Experiment	“Iso from T & cst accu”	“Iso from T & ERA5 accu”	“Iso from wg. mm & obs accu”	“Iso from ar. mm & obs accu”	“Iso from ECHAM6 & ECHAM6 accu”
a	1.4 (<i>1.4</i>)	1.2 (<i>1.0</i>)	1.1 (<i>0.9</i>)	1.3 (<i>1.1</i>)	1.1 (<i>0*</i>)
RMSE (‰)	6.9 (<i>8.8</i>)	3.7 (<i>5.3</i>)	3.4 (<i>3.9</i>)	5.8 (<i>5.5</i>)	6.3 (<i>7.1</i>)

3.4.2 Snow subsurface

600 For $\delta^{18}\text{O}$ in the snow subsurface, we find comparable results as for the surface layer. We find that all five model experiments fail to capture the short-term variations in the snow subsurface, and all experiments result in a too-large amplitude in the mean annual cycle over five years (Fig. 9a and b). The systematic positive bias given by the model experiment “iso from ECHAM6 & ECHAM6 accu” is also visible in the simulated subsurface layer (Fig. 9a and b). In addition, all five model experiments provide a mean annual cycle shifted compared to the observed one, with a maximum $\delta^{18}\text{O}$ occurring one to two months later than the observed maximum (Fig. 9b).

605 For d-excess, we find that four out of five of the model experiments (“iso from T & cst accu”, “iso from T & ERA5 accu”, “iso from wg. mm & obs accu”, “iso from ar. mean & obs accu”) give too high d-excess variations over time, which is reflected in the too large amplitude of the modelled mean annual cycle compared to the observed one (Fig. 9c and d). As for $\delta^{18}\text{O}$, the annual cycle in d-excess modelled in these three experiments is shifted, with the minimum in d-excess found two months later than the observed one (Fig. 9d).

610 On the other hand, although giving too low d-excess compared to the observations, the experiment “iso from ECHAM6 & ECHAM6 accu” results in a similar mean annual cycle as in the observations, with minimum d-excess in the summertime and maximum d-excess during the wintertime (pink line in Fig. 9d). However, because the precipitation d-excess modelled by ECHAM6-wiso poorly represent the observations (Fig. 6e and f), we argue that the good resemblance (besides the negative bias) between the modelled and observed d-excess in the snow subsurface is due to compensating effects between the precipitation d-excess and precipitation amounts.



615

620

Figure 9. Comparison between observed and simulated snow subsurface (1 to 4 cm depth) isotopic compositions at Dome C. Panels (a) and (c) show the 5-year time series (2017-2021) of $\delta^{18}\text{O}$ and d-excess (dxs) respectively. The grey dots represent the horizontal average between the two samples taken at the two locations 50 m apart on the sampling lines (described in Sect. 2.2, also presented in Fig. 3). The coloured lines (dotted and plain) represent the different SISG model experiments (described in Table 3). Panels (b) and (d) show the mean annual cycles (monthly means) of $\delta^{18}\text{O}$ and d-excess, respectively, calculated over the period 2017-2021. The grey dots and line represent the observations and the shaded area represents the inter-annual variability around the mean annual cycle (one standard deviation). The coloured lines (dotted and plain) represent the different SISG experiments (described in Table 3). The markers on the right vertical axis indicate the mean isotopic composition of the observed and simulated snow subsurface over 5 years (see Table S2 in Supplement S4.2).

625

As for the snow surface, we perform a linear regression between all modelled and observed monthly means $\delta^{18}\text{O}$ and d-excess. The linear slopes (a) and the RMSE between the model results and the observations are summarized in Table 5.

Contrary to the surface layer, the best representation of $\delta^{18}\text{O}$ in the subsurface layer is given by the experiment “iso from T & ERA5 accu” ($a=0.8$, $\text{RMSE}=3.3\text{‰}$, Table 5). The model experiment “iso from ECHAM6 & ECHAM6 accu” gives the best correlation with observations, although with the highest RMSE ($a=1.1$, $\text{RMSE}=6.7\text{‰}$, Table 5).

630

For d-excess in the snow subsurface, only the experiment “iso from ECHAM6 & ECHAM6 accu” gives a significant slope with the observations ($a=0.8$, $\text{RMSE}=6.0\text{‰}$, Table 5) while all other experiments have a non-significant linear slope with the

observations (p-value > 0.05, Table 5). However, as stated above, we argue that this good agreement between the simulated snow subsurface in the experiment “iso from ECHAM6 & ECHAM6 accu” is due to compensating effects between the precipitation d-excess and precipitation amounts.

635

Table 5. Linear regression slope (a) and RMSE between the observed and modelled monthly mean isotopic composition of the subsurface layer ($\delta^{18}\text{O}$ in normal font, d-excess in parenthesis and italic). The sample size is 60 for all experiments except “iso from wg. mm & obs accu” and “iso from ar. mm & obs accu” (n=54). All linear slopes are significant (p-value < 0.05), except the ones marked with an asterisk (*, p-value > 0.05).

640

Experiment	“Iso from T & cst accu”	“Iso from T & ERA5 accu”	“Iso from wg. mm & obs accu”	“Iso from ar. mm & obs accu”	“Iso from ECHAM6 & ECHAM6 accu”
a	1.0 (<i>0.5*</i>)	0.8 (<i>0.2*</i>)	0.5 (<i>0.4*</i>)	0.7 (<i>0.3*</i>)	1.1 (<i>0.8</i>)
RMSE (%)	6.2 (<i>8.4</i>)	3.3 (<i>5.1</i>)	3.5 (<i>3.5</i>)	6.1 (<i>6.0</i>)	6.7 (<i>6.0</i>)

4 Discussion

4.1 Limits of the methodological approach

4.1.1 Reliability of the datasets

Our study of the snow water stable isotopic composition is based on snow samples collected regularly all-year round at Dome C. In Fig. 3, 8 and 9, we use the average of the two samples collected 50 m apart (Sect. 2.2) to describe the variability in $\delta^{18}\text{O}$ and d-excess in the snow surface and subsurface. To ensure that the observed variations reflect a signal of local weather and climate rather than random noise, we analysed the temporal variations of the two samples independently (locations 1 & 2, Fig. S2 in Supplement S3). We see that the variations in $\delta^{18}\text{O}$ and d-excess at the sampling locations 1 and 2 generally follow each other. Additionally, a wavelet coherence analysis (Grinsted et al., 2004) revealed significant in-phase coherence between both locations for $\delta^{18}\text{O}$ and d-excess in the snow surface and subsurface samples, over approximately 120 days and above throughout the 5-year period (Fig. S3 in Supplement S3). Note that this is a bit less evident for the d-excess in the subsurface samples. We also see some episodic coherence between the two locations at 30 to 60 days. We argue that this shared signal between both locations indicates that the intra-annual to multi-annual variations in $\delta^{18}\text{O}$ and d-excess observed in the snow samples reflect a true temporal climate or weather driven signal. This brings confidence in using this dataset to investigate the formation of the isotopic signal in the snow. At shorter time scales (intra-monthly), the $\delta^{18}\text{O}$ and d-excess variations that are not shared by both locations most likely reflect the spatial heterogeneity of the surface, arising from snow erosion and redistribution leading to an accumulation by patches (Picard et al., 2019). This is one aspect of post-depositional processes that is further discussed in Sect. 4.2.1.

Certain limitations should also be mentioned concerning the precipitation isotopic composition time series presented in Fig. 6. As detailed in Sect. 3.2, all the snow laying on the bench is collected, whether it is precipitation (including diamond dust), blown snow or hoar formed from atmospheric condensation. In addition, some of the samples might have been affected by ad hoc sublimation, especially during the summertime, due to their exposure to the atmosphere before sample collection. As

660

665 already pointed out by Dreossi et al. (2024a), the precipitation dataset might also be biased towards higher precipitation events, because it was not possible to collect or measure the isotopic composition of the sample when very little snow was present on the table, due to too little precipitation during the day or because the sample was blown away. Lastly, some days with precipitation are missing in the dataset, because the sample was simply not collected, due to harsh conditions for example. This also affects the observed precipitation amounts presented in Fig. 5, as they are inferred from the weights of the samples collected on site (Sect. 2.3 and 2.5.1).

670 Considering the points discussed and to avoid overinterpreting the differences in the isotopic compositions of the individual precipitation and snow samples, we instead use the full datasets presented here to quantify the mean effect of post-depositional processes over five years (Sect. 4.2.2).

4.1.2 SISG setup and inputs

675 In Sect. 3.4.1 and 3.4.2, we showed that none of the five SISG experiments fully represent the variability observed in the snow surface and subsurface isotopic composition ($\delta^{18}\text{O}$ and d-excess). However, it should be mentioned that the results of the simple model used in our study come with some limitations.

680 The model was designed with a fixed depth for both snow samples to best represent the observations, although the depths of the samples collected in the field might have varied over time (Sect. 2.2). Changing the sample depths in the SISG model does not significantly improve the agreement between the model results and the observations (Table S3 in Supplement S4.3), however implementing a varying depth for each time step of the model could improve the concordance between the synthetic and observed snow layers.

The inputs of the model might also explain some of the discrepancies between the observed and synthetic snow layers. For example, there are fewer small precipitation events in the observations than given by ERA5 and ECHAM6-wiso (Fig. 5b), which reflects the possible bias of the observations towards higher precipitation events (Sect. 4.1.1). This could lead to an underestimation of the contribution of smaller precipitation events (e.g. diamond dust) to the total accumulation and explain the differences between the experiments “iso from wg. mm and obs accu” and “iso from ar. mm and obs accu” and the observations (Fig. 8 and 9). On the other hand, it might be ERA5 and ECHAM6-wiso that overestimate the contribution of smaller precipitation events to the total accumulation, due for example to too frequent precipitation days with low precipitation rates. Some snowfall events in ERA5 and ECHAM6-wiso could also be occurring too early or too late and explain why the short-term increases and decreases in the snow isotopic composition are not reproduced in the experiments using ERA5 and ECHAM6-wiso precipitation amounts (Fig. 8a and c, Fig. 9a and c). In addition, we scaled up all three precipitation time series to match the mean annual accumulation of 8 cm year⁻¹ at Dome C (Sect. 2.5), although this value can vary and estimations over other time periods have given higher mean accumulation rates (e.g. Frezzotti et al., 2005). One way to improve the model results would be to scale the precipitation time series with a different mean annual accumulation rate for each individual year using recent data from the GLACIOCLIM stake network (Genthon et al., 2015; <https://glacioclim.osug.fr/>). Lastly, due to missing data in the observations of the daily precipitation isotopic composition, we

695

used an artificial time series in three out of five experiments (Sect. 2.5), but none of these represent accurately the observations (Fig. S4 in Supplement S4.1). This could also partly explain the differences between the observed and synthetic snow layers.

700 For the different reasons stated above, we don't quantify the effect of post-depositional processes by comparing directly the model results and the observations, but rather use it as a tool to discuss the role of the different surface processes on the snow isotopic composition (Sect. 4.2.1).

4.2 Transfer of the isotopic signal between precipitation, snow surface and subsurface

In this section we discuss the potential processes at the surface of the ice sheet responsible for the discrepancies between the precipitation signal and the observed snow surface and subsurface layers and assess the overall impact of these processes on
705 the transfer of the isotopic signal between precipitation to the snow.

4.2.1 Short-term, inter-annual and seasonal variability in the snow isotopic composition

Several studies have shown that water vapor fluxes between the lower atmosphere and the snow can modify the isotopic composition of the snow surface (Casado et al., 2021; Hughes et al., 2021; Wahl et al., 2021, 2022) and deeper firn cores (Dietrich et al., 2023). Although the impact depends on the isotopic composition of the atmospheric water vapor above,
710 sublimation generally leads to an enrichment in $\delta^{18}\text{O}$ of the snow surface and a lowering of its d-excess. In addition, Casado et al. (2018) hypothesized that the inter-annual variability of the summer snow surface isotopic composition is related to the strength of metamorphism and surface vapor fluxes.

We showed that at Dome C, sublimation occurs during the summertime (Fig. 2a) and that there is a net annual sublimation of snow (Fig. 2b), which means that the surface snow isotopic composition must have been affected by these fluxes. We find
715 that in the $\delta^{18}\text{O}$ vs δD domain, most of the snow surface samples collected in December and January are below the precipitation samples (i.e. with a lower d-excess) collected in the same months (not shown), indicating an effect of sublimation on the snow surface isotopic composition in the summertime. In addition, we observe some inter-annual variability in the vertical difference between the snow surface and subsurface isotopic compositions (Fig. 4). This difference between the layer exposed to the atmosphere and the subsurface could be explained by the inter-annual variability in vapor
720 fluxes (Fig. 2a). For example, the snow surface is most enriched in $\delta^{18}\text{O}$ compared to the subsurface in December 2019 and January-February 2020 (Fig. 4). In parallel, we observe strong sublimation occurring in November and December 2019 (Fig. 2a), probably enriching the snow surface in $\delta^{18}\text{O}$, and leading to a much more enriched snow surface compared to the subsurface in the samples collected in the following months. Yet, the highest sublimation rate occurring in December 2018 (Fig. 2a) is not necessarily reflected in the vertical difference between the snow surface and subsurface layers (Fig. 4). This
725 could be explained by the length of the sublimation period, critical for the overall impact on the snow surface (Hughes et al., 2021; Wahl et al., 2022).

The mean seasonal cycle in the difference between the surface and subsurface isotopic compositions can, on the other hand, most likely be explained by the depths of the samples that integrate the precipitation events fallen during the last couple of months for the surface sample and up to the previous winter for the subsurface sample (assuming no removal or redistribution and no compaction of snow, Fig. S5 in Supplement S4.4). The higher $\delta^{18}\text{O}$ and lower d-excess in the summer precipitation compared to the winter precipitation (Fig. 6) explains why the mean summer snow surface is enriched in $\delta^{18}\text{O}$ and has a lower d-excess than the snow subsurface (Fig. 4). This integration process can also explain, at first order, why the maximum and minimum monthly mean $\delta^{18}\text{O}$ in the precipitation, the snow surface and the snow subsurface do not occur at the same time (January and May for the precipitation-weighted, Fig. 6c; February and October for the snow surface; and March and November for the snow subsurface, Fig. 3b).

If we would include vapor fluxes in the simple SISG model, the amplitude of the modelled seasonal cycle in the snow surface would be increased (summer months enriched in $\delta^{18}\text{O}$ and reduced in d-excess), increasing the discrepancy with the observations (Fig. 8b and d). This shows that an additional process reducing the amplitude of the mean annual cycle in the snow surface takes place. Diffusion is a good candidate.

Indeed, the isotopic composition of the snowpack is affected by diffusion of water molecules along isotopic and temperature gradients (Johnsen et al., 2000; Gkinis et al., 2014). Implementing this process in the model would possibly reduce the too large amplitude of the seasonal cycles in the synthetic snow layers to match the observations (Fig. 8b and d, 9b and d). Diffusion might also partly explain the difference observed at the seasonal scale between the precipitation and the snow surface isotopic compositions (Fig. 3b and d, Fig. 6c and f), smoothing the incoming high-amplitude seasonal signal of precipitation and leading to a snow surface enriched in $\delta^{18}\text{O}$ (and with lower d-excess) in the wintertime compared to precipitation. In addition, some of the short-term variations observed in the snow subsurface that are not explained by the incoming precipitation (Fig. 9a and c) could instead be explained by changes occurring in the snow surface and diffused downwards, as suggested by Casado et al. (2018).

Lastly, although Dome C is not affected by strong katabatic winds, surface winds can still be strong enough to erode and redistribute the snow (Libois et al., 2014). In a study of the process of snow accumulation at Dome C, Picard et al. (2019) showed that the snow surface changes very frequently due to snow erosion and accumulation by small patches (10% of the surface only is affected by each precipitation events), and that the snow in a patch can be as old as one year at the surface, while another patch was deposited by the last precipitation event. In the snow samples collected at Dome C, we observe higher differences in $\delta^{18}\text{O}$ and d-excess between two consecutive sampling days of the snow surface and subsurface when the wind speed increases from one sampling day to another (Fig. S6 in Supplement S5), as well as a decrease in $\delta^{18}\text{O}$ and d-excess vertical difference between the snow surface and the subsurface (Fig. S7 in Supplement S5). This could indicate that some of the short-term variations observed in the snow surface isotopic composition that cannot be explained by new precipitation falling on the surface (Fig. 3, 8a and c) be caused instead by wind erosion and redistribution.

The wind blowing at the surface of the ice sheet during or beside snowfall could lead to a mixing of new precipitation with already deposited snow. This process was proposed by Casado et al. (2018) to explain the slow decrease in the surface snow

$\delta^{18}\text{O}$ during the winter, a pattern observed in our study (Fig. 3a and b) and that cannot be explained by precipitation only (Fig. 8a and b). A recent laboratory study also suggests that snow metamorphism during wind transport (“airborne snow metamorphism”) has the potential to impact the snow isotopic composition in both $\delta^{18}\text{O}$ and d-excess (Wahl et al., 2024). This could be an additional process occurring at Dome C explaining some of the differences between the precipitation and snow isotopic signals, including in the wintertime.

Overall, our results show evidence that the snow isotopic composition is affected by post-depositional processes at different timescales. Disentangling the contribution of the different processes described above on the final isotopic signal found in the snow is beyond the scope of this study and requires the use of an isotope-equipped snowpack model that includes post-depositional processes at the surface, such as the one developed recently by Wahl et al. (2022) and Dietrich et al. (2023).

4.2.2 Mean effect of post-depositional processes

The datasets presented in this study permit to quantify the overall impact of post-depositional processes on the snow isotopic composition. Table 6 summarizes the mean isotopic composition over five years of precipitation (weighted average by the observed precipitation amounts, see Sect. 2.5.1), the snow surface, and the snow subsurface. The mean isotopic composition of precipitation excluding all samples with negative d-excess and modelled by ECHAM6-wiso are also provided. All mean values are given with their respective standard error (see Sect. 3.2.1 and Supplement S2) and 95% confidence interval (Student t-test, Supplement S2).

Table 6. Five-year mean isotopic composition of precipitation (weighted by the observed precipitation amounts) and snow samples ($\delta^{18}\text{O}$ in normal font, d-excess in parenthesis and italic). The mean values are given with their respective standard error (see Sect. 3.2.1) and in brackets their 95% confidence interval (Student t-test, Supplement S2).

	Observations	Observations (excl. samples with dxs < 0)	ECHAM6-wiso
Precipitation-weighted (‰)	-53.4 ± 0.5 [1.0] <i>(12.2 ± 0.5 [1.0])</i>	-54.3 ± 0.4 [0.9] <i>(13.9 ± 0.4 [0.8])</i>	-45.7 ± 0.5 [0.9] <i>(5.1 ± 0.3 [0.5])</i>
Snow surface (‰)	-51.0 ± 0.2 [0.4] <i>(10.4 ± 0.2 [0.5])</i>	–	–
Snow subsurface (‰)	-51.4 ± 0.1 [0.3] <i>(10.8 ± 0.1 [0.3])</i>	–	–

As shown in Section 3.3.2, the observed $\delta^{18}\text{O}$ precipitation-weighted mean over five years is higher than the overall mean (-53.4 and -56.2% , respectively). This is explained by the lowest $\delta^{18}\text{O}$ values in precipitation being associated with smaller precipitation amounts (Fig. S8 in Supplement S6), and therefore weigh less when computing the precipitation-weighted average. The opposite applies for d-excess: the weighted overall mean is lower than the arithmetic overall mean (12.2 and 15.2% , respectively) because high d-excess values are associated with lower precipitation amounts (Fig. S8 in Supplement S6).

Now because the snow surface layer represents the amount of snow accumulated over a certain period, its mean isotopic composition should reflect the weighted mean (by precipitation amounts) isotopic composition of precipitation. Therefore,

we express the mean effect of post-depositional processes at Dome C as the difference between the observed 5-year weighted mean isotopic composition of precipitation and the observed 5-year mean isotopic composition of the snow surface.

795 Considering all precipitation and snow samples, the snow surface $\delta^{18}\text{O}$ is 2.4‰ higher than in precipitation and the snow surface d-excess is 1.8‰ lower than in precipitation (Table 6). However, to exclude any imprint of sublimation on the precipitation isotopic composition, we re-compute the mean isotopic composition of precipitation discarding all the samples with a d-excess below zero, as in Steen-Larsen et al. (2011). Although the zero threshold might be arbitrary, it is supported by laboratory and field studies that showed a decrease in d-excess during snow sublimation and metamorphism (Hughes et al., 2021; Casado et al., 2021; Harris Stuart et al., 2023). In addition, Stenni et al. (2016) and Dreossi et al. (2024a) already
800 stated that some of the samples collected in the summertime at Dome C might have been affected by sublimation and we observe in the precipitation dataset presented here that the samples with very negative d-excess were collected during the summertime (e.g. December 2019, Fig. 6d) when sublimation occurred (Fig. 2a). We therefore argue that removing all precipitation samples with a d-excess below zero improves the representation of the true precipitation falling at Dome C. Now considering this new precipitation time series, the $\delta^{18}\text{O}$ in the snow surface is 3.3‰ higher than in precipitation and the
805 d-excess in the snow surface is 3.5‰ lower than in precipitation (Table 6). Both differences are significant on a 95% confidence level (Table 6).

The weighted mean isotopic composition of precipitation is computed using the observed precipitation amounts, estimated from the weights of the samples collected on site (Sect. 2.3 and 2.5.1). However, as discussed in Sect. 4.1.3, these amounts might not accurately represent the precipitation falling at Dome C. Using the precipitation amounts given by ERA5 instead,
810 the weighted mean $\delta^{18}\text{O}$ of precipitation is -54.0‰ and 13.8‰ for d-excess, which leads to similar differences between the precipitation and the snow surface: $\delta^{18}\text{O}$ in the snow surface is 3.0‰ higher than in precipitation and d-excess in the snow surface is 3.4‰ lower than in precipitation.

The discrepancies between the precipitation and the snow surface show that the mean isotopic composition of precipitation is not preserved from snowfall to the snow surface. On average over a 5-year period at Dome C, post-depositional processes
815 lead to an enrichment in $\delta^{18}\text{O}$ of the snow surface by 3.0‰ to 3.3‰ (depending on which precipitation amounts are considered) and a lowering of d-excess in the snow surface by 3.4‰ to 3.5‰ compared to the precipitation signal. It is still to be determined the individual contribution of the different post-depositional processes (discussed in Sect. 4.2.1) on the isotopic difference observed between the precipitation and the snow surface.

In contrast to the isotopic difference between the precipitation and snow surface, there is no significant difference between
820 the mean snow surface and subsurface isotopic compositions, for both $\delta^{18}\text{O}$ and d-excess (Table 6). This shows that despite a seasonality in the vertical difference between the two snow layers (discussed in Sect. 4.2.1), the mean isotopic composition of the snow surface layer is preserved in the top few centimetres of the snowpack.

From the sampling of a 2 m snow pit at Dome C, Touzeau et al. (2016) reported mean values (plus-minus standard error of the mean) of $-51.1 \pm 0.2\text{‰}$ and $9.1 \pm 0.2\text{‰}$ for the $\delta^{18}\text{O}$ and d-excess profiles, respectively. As found in our study for the

825 upper layers of the snow, the average isotopic composition of the snow pit is enriched in $\delta^{18}\text{O}$ and has a lower d-excess than the mean incoming precipitation isotope signal.

The weighted-mean isotopic composition of precipitation modelled by ECHAM6-wiso over the whole period is 7.7‰ higher in $\delta^{18}\text{O}$ and 7.1‰ lower in d-excess than the observed precipitation (Table 6, Fig. 6c and f). These large differences result from the biases identified in the model (Sect. 3.3.2, Fig. 6 and 7), combined with the distribution of the daily isotopic values against precipitation amounts (Fig. S8 in Supplement S6), which influences both the overall mean and the seasonal difference between the observations and the model (Fig. 6 and Supplement S6). This shows the limitations of using ECHAM6-wiso simulations to interpret the isotopic composition of the snow at Dome C. A thorough investigation of the biases in ECHAM6-wiso is beyond the scope of this study, but they might arise from different processes in the model, such as the environmental conditions at the moisture source region, the moisture transport and pathway, the supersaturation parametrization or the condensation height and temperature at Dome C. The fourteen-year record of the precipitation isotopic composition (Dreossi et al., 2024a and this study) gives the opportunity to evaluate isotope-enabled GCMs and can be used to improve the tuning of the empirical parameterization of supersaturation in polar clouds (e.g. Risi et al., 2013).

4.2.3 Seasonal $\delta^{18}\text{O}$ versus temperature relationships in precipitation and snow

840 In precipitation, the $\delta^{18}\text{O}$ -temperature relationship is determined between the $\delta^{18}\text{O}$ in the daily precipitation samples and the corresponding daily atmospheric temperature (Sect. 3.3.3, Eq. (1)). For the snow surface and subsurface, we determine the relationships between the $\delta^{18}\text{O}$ composition of each snow sample and the weighted-average temperature (by precipitation amounts) over the entire period during which precipitation accumulated and formed the snow sample (Fig. A3). We use the observed precipitation amounts to determine this averaging period (Fig. S5 in Supplement S4.4). Table 7 summarizes the linear regression slopes (a) with their standard errors and the coefficient of determination (R^2).

Table 7. Summary of mean seasonal $\delta^{18}\text{O}$ versus temperature linear relationships in precipitation and snow samples between 2017 and 2021. The slopes (a) are given with their standard error and all three slopes are significant (p-value < 0.001).

	Precipitation	Snow surface	Snow subsurface
a (‰ °C ⁻¹)	0.47 ± 0.01	0.17 ± 0.01	0.12 ± 0.02
R ²	0.62	0.25	0.1

850

We find a mean seasonal $\delta^{18}\text{O}$ -temperature slope in precipitation of $0.47 \pm 0.01\text{‰ }^\circ\text{C}^{-1}$, in agreement with previous studies for the same location for earlier years (Stenni et al., 2016; Dreossi et al., 2024a). We find a mean seasonal slope of $0.17 \pm 0.01\text{‰ }^\circ\text{C}^{-1}$ in the snow surface (top 1 cm) and a mean seasonal slope of $0.12 \pm 0.02\text{‰ }^\circ\text{C}^{-1}$ in the snow subsurface (1 to 4 cm, Table 7). The snow surface slope coincides with the one found for the snow top 1-2 mm at Dome C ($0.14 \pm 0.03\text{‰ }^\circ\text{C}^{-1}$, Touzeau et al., 2016) but is lower than the ones found by Casado et al. (2018) for similar samples in earlier years (slopes ranging from 0.22 to $0.49\text{‰ }^\circ\text{C}^{-1}$). This difference can be due to uncertainties on the precipitation amounts used to determine

the averaging period of temperature (i.e. some precipitation days with colder temperatures are not captured), the method used to determine the averaging period of temperature (i.e. assumes that snowfall is not redistributed or removed and does not include snow compaction, Supplement S4.4), or due to the different method used in Casado et al. (2018) where they compute the slope as the ratio of the maximum amplitude in $\delta^{18}\text{O}$ and the maximum amplitude in temperature.

Our results indicate that deriving a temporal slope from seasonal variations of $\delta^{18}\text{O}$ in the snow surface to be used for ice core studies is not consistent with other in-situ and proxy temperature reconstructions. First, the snow surface $\delta^{18}\text{O}$ is not showing a symmetric seasonal cycle. Instead, the $\delta^{18}\text{O}$ minimum is shifted towards spring (Fig. 3, Sect. 3.2.1) while the minimum temperature occurs in winter. Secondly, our determination of the seasonal relationship between the snow $\delta^{18}\text{O}$ and the atmospheric temperature results in a weaker correlation and a lower slope ($0.12 - 0.17\text{‰ } ^\circ\text{C}^{-1}$, Table 7) than what is found for precipitation ($0.47\text{‰ } ^\circ\text{C}^{-1}$, Table 7). Using the $\delta^{18}\text{O}$ -temperature slope found in the upper layers of the snow for temperature reconstructions from ice cores would lead to unrealistic glacial-interglacial temperature change exceeding 30°C . The change in the seasonal $\delta^{18}\text{O}$ -temperature relationship between the precipitation and the upper layers of the snow shows the difficulty to interpret quantitatively the $\delta^{18}\text{O}$ variations in the snow at Dome C in terms of temperature. Instead, it provides the opportunity to document the effects of post-depositional processes at this site and improve the future quantitative interpretation of water isotopic records in ice cores from Dome C for longer timescales (decadal and longer timescales).

4.3 Long-term perspectives for the interpretation of ice core isotope records

The classical paleothermometer approach (Lorius and Merlivat, 1975) to determine past temperature variations from the water isotopic profiles in ice cores uses the present-day spatial slope between the water isotopic composition of the snow surface and the local temperature. However, as illustrated in Casado et al. (2017), this spatial slope differs from the various estimates of the temporal slope between temperature and $\delta^{18}\text{O}$ found in the literature and in this study, determined for different regions and timescales. These discrepancies show the need of calibration of the isotopic paleothermometer. Such calibration has been done using alternative methods at the glacial-interglacial scale (e.g. borehole thermometry and firn properties, Buizert et al., 2021), or using isotope-enabled GCMs to determine the relationship between temperature and water isotopes during past periods (e.g. Werner et al., 2018). However, in these models, the absence of explicit modelling of how the water isotopic signal is archived in the snow and firn limits their use for paleo-reconstructions.

To progress towards an accurate quantitative interpretation of isotopes in ice cores, we recommend developing a *proxy system model* (Evans et al., 2013) that can be coupled to isotope-enabled GCMs. A first step toward such a model has been developed recently by Wahl et al. (2022) and Dietrich et al. (2023) for Greenland and includes mechanical processes leading to the recording of isotopes at the ice sheet's surface and in firn cores. The datasets presented in our study can therefore be useful to calibrate and validate this kind of model at intra-annual, seasonal, and inter-annual timescales at the deep drilling site of the EPICA Dome C ice core.

5 Conclusions

890 We have presented the compilation of new and existing datasets of the isotopic composition of precipitation, snow surface and subsurface together with meteorological parameters, reanalysis products, model outputs and a simple modelling approach to investigate the origin of the stable water isotopic signal in the upper layers of the snowpack at Dome C, East Antarctica.

From in-situ meteorological observations, we have quantified the amount of water vapor fluxes between the snow and the
895 lower atmosphere. Our results show that vapor fluxes contribute to the surface mass balance at Dome C with a net annual mass loss of snow from 3.1 to 3.7 mm w.e. year⁻¹ between 2018 and 2020, which corresponds to 12 to 15% of the annual surface mass balance. Sublimation is relatively strong in the summertime and there is small condensation in the wintertime.

From the comparison between the snow surface and subsurface isotopic compositions and a simple modelling approach, we show that the precipitation deposited onto the ice sheet cannot explain the variability observed in the snow isotopic
900 composition ($\delta^{18}\text{O}$ and d-excess) at seasonal to intra-monthly timescales, highlighting the existence of post-depositional processes affecting the snow isotopic composition.


We quantified the cumulative effect of post-depositional processes on the snow surface over five years by comparing the mean isotopic compositions of the precipitation and the snow layers. Our results show that post-depositional processes lead to an enrichment of the snow surface (top 1 cm) in $\delta^{18}\text{O}$ by 3.0‰ to 3.3‰ (depending on the precipitation amounts
905 considered) and a lowering of the snow surface d-excess by 3.4‰ to 3.5‰. In addition, our results show that the mean isotopic composition of the snow subsurface (1 to 4 cm depth) is not significantly different than the snow surface, which indicates that the mean isotopic composition of the snow surface layer is preserved in the top centimetres of the snowpack and that the processes altering the precipitation isotopic signal mainly take place in the top one centimetre of the snow.

In both the observations, ERA5 reanalyses and ECHAM6-wiso simulation, we showed that 50% of the accumulation at
910 Dome C over five years occurs during large but rare precipitation events associated with warmer temperatures than average, leading to a warm bias in the $\delta^{18}\text{O}$ record of precipitation compared to the mean annual temperature. We further find different seasonal relationships between the atmospheric temperature and $\delta^{18}\text{O}$ in the precipitation and the snow, showing the difficulty to interpret the variations of $\delta^{18}\text{O}$ in the snow at Dome C.


Overall, our results show that post-depositional processes at the ice sheet's surface have an impact on the isotopic signal
915 (both $\delta^{18}\text{O}$ and d-excess) found in the upper layers of the snowpack at Dome C, East Antarctica. The datasets presented in our study open the possibility to calibrate and validate *proxy system models* including post-depositional processes that are needed to quantitatively attribute the different mechanisms building up the isotopic signal in the snow surface and to accurately reconstruct the climatic information from the water stable isotope records in ice cores.

920


Appendix A



925 **Figure A1.** Daily observed precipitation $\delta^{18}\text{O}$ (a) and δD (b) versus observed daily mean temperature at Dome C. All precipitation samples collected between 2017 and 2021 are shown against the corresponding daily mean temperature (T3m, dark colours) and the daily mean temperature of the day before (T3m shifted, light colours). The linear regressions with the associated coefficients of determination (R^2) are shown. Both linear slopes are significant (p -values < 0.001).



930 **Figure A2.** Daily precipitation $\delta^{18}\text{O}$ (a) and δD (b) modelled by ECHAM6-wiso versus daily mean temperature modelled by ECHAM6-wiso. The linear regressions with the associated coefficients of determination (R^2) are shown. Both linear slopes are significant (p -values < 0.001).



935 **Figure A3.** Bi-weekly observed snow surface (a) and subsurface (b) $\delta^{18}\text{O}$ versus temperature at Dome C. All snow samples collected between 2017 and 2021 are shown against the weighted-average temperature (by precipitation amounts) over the period corresponding to the accumulation of the snow samples (see Sect. 4.2.3). The linear regressions with the associated coefficients of determination (R^2) are shown. Both linear slopes are significant (p-values < 0.001).

940 *Code availability.* The code for the estimation of water vapor fluxes with the bulk method is available at <https://doi.org/10.5281/zenodo.13833912> and the SISG model code is available at <https://doi.org/10.5281/zenodo.13833981>.

Data availability. All the observational and meteorological datasets used in this study are available in public repositories. The isotopic composition of the snow surface and subsurface samples between 2017 and 2021 is available in Landais et al. (2024) (<https://doi.pangaea.de/10.1594/PANGAEA.971486>). The 2017-2021 dataset of the isotopic composition and weights of the daily precipitation samples is available in Dreossi et al. (2024b) (<https://doi.pangaea.de/10.1594/PANGAEA.972031>). The atmospheric temperature dataset is available in Genthon et al. (2021b) (<https://doi.pangaea.de/10.1594/PANGAEA.932512>) and at <https://web.lmd.jussieu.fr/~cgenthon/SiteCALVA/CalvaData.html>. The wind speed dataset is available in Genthon et al. (2021c) (<https://doi.pangaea.de/10.1594/PANGAEA.932513>) and at <https://web.lmd.jussieu.fr/~cgenthon/SiteCALVA/CalvaData.html>. The atmospheric relative humidity dataset is available in Genthon et al. (2021d) (<https://doi.pangaea.de/10.1594/PANGAEA.939421>). The atmospheric pressure dataset is available in Grigioni et al. (2022) (<https://doi.org/10.12910/DATASET2022-002>). The radiative fluxes dataset from the BSRN network is available in Lupi et al. (2021) (<https://doi.pangaea.de/10.1594/PANGAEA.935421>). The CNR4 radiative fluxes dataset is available in Arnaud and Picard (2024) (<https://doi.org/10.18709/perscido.2024.11.ds415>).

955

Author contribution. IO, HCSL, BS and AL designed the study and contributed to the analysis. IO performed the formal analysis and visualization. HCSL, BS and AL supervised the study. MC designed the first version of the simple model (SISG) and provided inputs to the study. GP and LA acquired and shared radiation data (CNR4) and coordinated the snow sample collection executed by the winter-over staff. CG acquired meteorological data (USt) and provided inputs to the study. 960 GD and IO performed the isotopic analysis of the precipitation samples. BM performed the isotopic analysis of the snow samples. AC and MW performed the ECHAM6-wiso simulation. IO wrote the first draft of the manuscript and all authors contributed to reviews and edits.

Competing interests. The authors declare that they have no conflict of interest.

965

Acknowledgements. This publication was generated in the frame of the DEEPICE project. The project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 955750. Snow samples and radiation data (CNR4) have been collected within the frame of the French Polar Institute (IPEV) project NIVO 1110. Precipitation samples and its isotopic analysis have been conducted in the framework of 970 the projects PNRA 2013/AC3.05 (PRE-REC) and PNRA18_00031 (WHETSTONE) of the Italian National Antarctic Research Program "Programma Nazionale di Ricerche in Antartide" (PNRA) funded by MIUR (now MUR). We acknowledge using data from the project CALVA 1013 and GLACIOCLIM observatories supported by the French Polar Institute (IPEV) and the Observatoire des Sciences de l'Univers de Grenoble (OSUG) (<https://web.lmd.jussieu.fr/~cgenthon/SiteCALVA/CalvaData.html>) and thank Etienne Vignon from Laboratoire de 975 Météorologie Dynamique (Paris, France) for his valuable input on the bulk method and vapor fluxes estimations. We also acknowledge using data from the Italian Antarctic Meteo-Climatological Observatory (IAMCO, <https://www.climantartide.it/>) collected in the framework of the PNRA/IPEV 'Routine Meteorological Observation at Station Concordia' project and from the Baseline Surface Radiation Network (BSRN, <https://bsrn.awi.de/>). AL has received funding from the European Research Council under the European Union Horizon 2020 Excellent Science program (H2020/20192024)/ERC grant agreement no. 817493 (ERC ICORDA). HCSL has received funding from the 980 European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme: Starting Grant SNOWISO (grant agreement no. 759526). We also thank the logistics staff and winter-over crews at Concordia station for all the field sampling and instrumental maintenance during investigated periods and L. Dietrich for proofreading this article.

985

Financial support. This research has been supported by the European Union's Horizon 2020 research and innovation programme (grant no. 955750).

References

- Arnaud, L. and Picard, G.: Timeseries of shortwave and longwave radiation measurements at Dome C in Antarctica (2018-2020), PerSciDO [dataset], <https://doi.org/10.18709/perscido.2024.11.ds415>, 2024.
- 990 Bretherton, C. S., Widmann, M., Dymnikov, V. P., Wallace, J. M., and Bladé, I.: The Effective Number of Spatial Degrees of Freedom of a Time-Varying Field, *J. Climate*, 12, 1990–2009, [https://doi.org/10.1175/1520-0442\(1999\)012<1990:TENOSD>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<1990:TENOSD>2.0.CO;2), 1999.
- Brun, E., Six, D., Picard, G., Vionnet, V., Arnaud, L., Bazile, E., Boone, A., Bouchard, A., Genthon, C., Guidard, V., Moigne, P. L., Rabier, F., and Seity, Y.: Snow/atmosphere coupled simulation at Dome C, Antarctica, *J. Glaciol.*, 57, 721–736, <https://doi.org/10.3189/002214311797409794>, 2011.
- 995 Buizert, C., Fudge, T. J., Roberts, W. H. G., Steig, E. J., Sherriff-Tadano, S., Ritz, C., Lefebvre, E., Edwards, J., Kawamura, K., Oyabu, I., Motoyama, H., Kahle, E. C., Jones, T. R., Abe-Ouchi, A., Obase, T., Martin, C., Corr, H., Severinghaus, J. P., Beaudette, R., Epifanio, J. A., Brook, E. J., Martin, K., Chappellaz, J., Aoki, S., Nakazawa, T., Sowers, T. A., Alley, R. B., Ahn, J., Sigl, M., Severi, M., Dunbar, N. W., Svensson, A., Fegyveresi, J. M., He, C., Liu, Z., Zhu, J., Otto-Bliesner, B. L., Lipenkov, V. Y., Kageyama, M., and Schwander, J.: Antarctic surface temperature and elevation during the Last Glacial Maximum, *Science*, 372, 1097–1101, <https://doi.org/10.1126/science.abd2897>, 2021.
- 1000 Casado, M., Landais, A., Masson-Delmotte, V., Genthon, C., Kerstel, E., Kassi, S., Arnaud, L., Picard, G., Prie, F., Cattani, O., Steen-Larsen, H.-C., Vignon, E., and Cermak, P.: Continuous measurements of isotopic composition of water vapour on the East Antarctic Plateau, *Atmos. Chem. Phys.*, 16, 8521–8538, <https://doi.org/10.5194/acp-16-8521-2016>, 2016.
- 1005 Casado, M., Orsi, A., and Landais, A.: On the limits of climate reconstruction from water stable isotopes in polar ice cores, *PAGES Mag.*, 25, 146–147, <https://doi.org/10.22498/pages.25.3.146>, 2017.
- Casado, M., Landais, A., Picard, G., Münch, T., Laepple, T., Stenni, B., Dreossi, G., Ekaykin, A., Arnaud, L., Genthon, C., Touzeau, A., Masson-Delmotte, V., and Jouzel, J.: Archival processes of the water stable isotope signal in East Antarctic ice cores, *The Cryosphere*, 12, 1745–1766, <https://doi.org/10.5194/tc-12-1745-2018>, 2018.
- 1010 Casado, M., Landais, A., Picard, G., Arnaud, L., Dreossi, G., Stenni, B., and Prié, F.: Water Isotopic Signature of Surface Snow Metamorphism in Antarctica, *Geophys Res Lett*, 48, <https://doi.org/10.1029/2021GL093382>, 2021.
- Cauquoin, A. and Werner, M.: High-Resolution Nudged Isotope Modeling With ECHAM6-Wiso: Impacts of Updated Model Physics and ERA5 Reanalysis Data, *J Adv Model Earth Syst*, 13, <https://doi.org/10.1029/2021MS002532>, 2021.
- Cauquoin, A. and Werner, M.: ECHAM6-wiso and ECHAM5-wiso isotope daily data at Dome Concordia station, East Antarctica, Zenodo [dataset], <https://doi.org/10.5281/zenodo.11468043>, 2024.
- 1015 Cauquoin, A., Werner, M., and Lohmann, G.: Water isotopes – climate relationships for the mid-Holocene and preindustrial period simulated with an isotope-enabled version of MPI-ESM, *Clim. Past*, 15, 1913–1937, <https://doi.org/10.5194/cp-15-1913-2019>, 2019.
- Craig, H.: Standard for Reporting Concentrations of Deuterium and Oxygen-18 in Natural Waters, *Science*, 133, 1833–1834, <https://doi.org/10.1126/science.133.3467.1833>, 1961.
- 1020 Dansgaard, W.: Stable isotopes in precipitation, *Tellus*, 16, 436–468, <https://doi.org/10.3402/tellusa.v16i4.8993>, 1964.
- Dietrich, L. J., Steen-Larsen, H. C., Wahl, S., Jones, T. R., Town, M. S., and Werner, M.: Snow-Atmosphere Humidity Exchange at the Ice Sheet Surface Alters Annual Mean Climate Signals in Ice Core Records, *Geophysical Research Letters*, 50, e2023GL104249, <https://doi.org/10.1029/2023GL104249>, 2023.

- 1025 Dietrich, L. J., Steen-Larsen, H. C., Wahl, S., Faber, A.-K., and Fettweis, X.: On the importance of the humidity flux for the surface mass balance in the accumulation zone of the Greenland Ice Sheet, *The Cryosphere*, 18, 289–305, <https://doi.org/10.5194/tc-18-289-2024>, 2024.
- Dreossi, G., Masiol, M., Stenni, B., Zannoni, D., Scarchilli, C., Ciardini, V., Casado, M., Landais, A., Werner, M., Cauquoin, A., Casasanta, G., Del Guasta, M., Posocco, V., and Barbante, C.: A decade (2008–2017) of water stable isotope composition of precipitation at Concordia Station, East Antarctica, *The Cryosphere*, 18, 3911–3931, <https://doi.org/10.5194/tc-18-3911-2024>, 1030 2024a.
- Dreossi, G., Stenni, B., Masiol, M., Zannoni, D., and Ollivier, I.: Water stable isotopic composition of precipitation at Concordia Station (Dome C), East Antarctica (2017–2021), PANGAEA [dataset], <https://doi.pangaea.de/10.1594/PANGAEA.972031>, 2024b.
- Evans, M. N., Tolwinski-Ward, S. E., Thompson, D. M., and Anchukaitis, K. J.: Applications of proxy system modeling in high resolution paleoclimatology, *Quaternary Science Reviews*, 76, 16–28, <https://doi.org/10.1016/j.quascirev.2013.05.024>, 2013.
- 1035 Frezzotti, M., Pourchet, M., Flora, O., Gandolfi, S., Gay, M., Urbini, S., Vincent, C., Becagli, S., Gragnani, R., Proposito, M., Severi, M., Traversi, R., Udisti, R., and Fily, M.: Spatial and temporal variability of snow accumulation in East Antarctica from traverse data, *J. Glaciol.*, 51, 113–124, <https://doi.org/10.3189/172756505781829502>, 2005.
- Gallée, H. and Schayes, G.: Development of a Three-Dimensional Meso- γ Primitive Equation Model: Katabatic Winds Simulation in the Area of Terra Nova Bay, Antarctica, *Monthly Weather Review*, 122, 671–685, [https://doi.org/10.1175/1520-0493\(1994\)122<0671:DOATDM>2.0.CO;2](https://doi.org/10.1175/1520-0493(1994)122<0671:DOATDM>2.0.CO;2), 1040 1994.
- Genthon, C., Six, D., Scarchilli, C., Ciardini, V., and Frezzotti, M.: Meteorological and snow accumulation gradients across Dome C, East Antarctic plateau: Meteorological and snow accumulation gradients at Dome C, *Int. J. Climatol.*, 36, 455–466, <https://doi.org/10.1002/joc.4362>, 2015.
- Genthon, C., Piard, L., Vignon, E., Madeleine, J.-B., Casado, M., and Gallée, H.: Atmospheric moisture supersaturation in the near-surface atmosphere at Dome C, Antarctic Plateau, *Atmos. Chem. Phys.*, 17, 691–704, <https://doi.org/10.5194/acp-17-691-2017>, 2017.
- Genthon, C., Veron, D., Vignon, E., Six, D., Dufresne, J.-L., Madeleine, J.-B., Sultan, E., and Forget, F.: 10 years of temperature and wind observation on a 45 m tower at Dome C, East Antarctic plateau, *Earth Syst. Sci. Data*, 13, 5731–5746, <https://doi.org/10.5194/essd-13-5731-2021>, 2021a.
- Genthon, C., Veron, D., Vignon, E., Six, D., Dufresne, J. L., Madeleine, J.-B., Sultan, E., and Forget, F.: Ten years of shielded ventilated atmospheric temperature observation on a 45-m tower at Dome C, East Antarctic plateau, PANGAEA [dataset], <https://doi.org/10.1594/PANGAEA.932512>, 2021b.
- 1050 Genthon, C., Veron, D., Vignon, E., Six, D., Dufresne, J. L., Madeleine, J.-B., Sultan, E., and Forget, F.: Ten years of wind speed observation on a 45-m tower at Dome C, East Antarctic plateau, PANGAEA [dataset], <https://doi.org/10.1594/PANGAEA.932513>, 2021c.
- 1055 Genthon, C., Veron, D., Vignon, E., Madeleine, J.-B., and Piard, L.: Water vapor observation at 3 m height above ground at Dome C, East Antarctic plateau (2018–2020), PANGAEA [dataset], <https://doi.org/10.1594/PANGAEA.939421>, 2021d.
- Genthon, C., Veron, D. E., Vignon, E., Madeleine, J.-B., and Piard, L.: Water vapor in cold and clean atmosphere: a 3-year data set in the boundary layer of Dome C, East Antarctic Plateau, *Earth Syst. Sci. Data*, 14, 1571–1580, <https://doi.org/10.5194/essd-14-1571-2022>, 2022.

- 1060 Gkinis, V., Simonsen, S. B., Buchardt, S. L., White, J. W. C., and Vinther, B. M.: Water isotope diffusion rates from the NorthGRIP ice core for the last 16,000 years – Glaciological and paleoclimatic implications, *Earth and Planetary Science Letters*, 405, 132–141, <https://doi.org/10.1016/j.epsl.2014.08.022>, 2014.
- Gonfiantini, R.: Standards for stable isotope measurements in natural compounds, *Nature*, 271, 534–536, <https://doi.org/10.1038/271534a0>, 1978.
- 1065 Grachev, A. A., Andreas, E. L., Fairall, C. W., Guest, P. S., and Persson, P. O. G.: SHEBA flux–profile relationships in the stable atmospheric boundary layer, *Boundary-Layer Meteorol*, 124, 315–333, <https://doi.org/10.1007/s10546-007-9177-6>, 2007.
- Grigioni, P., Camporeale, G., Ciardini, V., De Silvestri, L., Iaccarino, A., Proposito, M., and Sarchilli, C.: Dati meteorologici della Stazione meteorologica CONCORDIA presso la Base CONCORDIA STATION (Dome C), ENEA [dataset], <https://doi.org/10.12910/DATASET2022-002>, 2022.
- 1070 Grinsted, A., Moore, J. C., and Jevrejeva, S.: Application of the cross wavelet transform and wavelet coherence to geophysical time series, *Nonlin. Processes Geophys.*, 11, 561–566, <https://doi.org/10.5194/npg-11-561-2004>, 2004.
- Harris Stuart, R., Faber, A.-K., Wahl, S., Hörhold, M., Kipfstuhl, S., Vasskog, K., Behrens, M., Zuhr, A. M., and Steen-Larsen, H. C.: Exploring the role of snow metamorphism on the isotopic composition of the surface snow at EastGRIP, *The Cryosphere*, 17, 1185–1204, <https://doi.org/10.5194/tc-17-1185-2023>, 2023.
- 1075 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., De Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.: The ERA5 global reanalysis, *Quart J Royal Meteorol Soc*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 1080 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J.-N.: ERA5 hourly data on single levels from 1940 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [dataset], <https://doi.org/10.24381/cds.adbb2d47>, 2023.
- Högström, U.: Review of some basic characteristics of the atmospheric surface layer, *Boundary-Layer Meteorology*, 78, 215–246, 1996.
- 1085 Holtslag, A. A. M. and De Bruin, H. A. R.: Applied Modeling of the Nighttime Surface Energy Balance over Land, *J. Appl. Meteor.*, 27, 689–704, [https://doi.org/10.1175/1520-0450\(1988\)027<0689:AMOTNS>2.0.CO;2](https://doi.org/10.1175/1520-0450(1988)027<0689:AMOTNS>2.0.CO;2), 1988.
- Hughes, A. G., Wahl, S., Jones, T. R., Zuhr, A., Hörhold, M., White, J. W. C., and Steen-Larsen, H. C.: The role of sublimation as a driver of climate signals in the water isotope content of surface snow: laboratory and field experimental results, *The Cryosphere*, 15, 4949–4974, <https://doi.org/10.5194/tc-15-4949-2021>, 2021.
- 1090 Johnsen, S. J., Clausen, H. B., Cuffey, K. M., Hoffmann, G., Schwander, J., and Creyts, T.: Diffusion of stable isotopes in polar firn and ice: the isotope effect in firn diffusion, *Physics of Ice Core Records*, 121–140, <https://doi.org/10.7916/D8KW5D4X>, 2000.
- Jouzel, J., Merlivat, L., and Lorius, C.: Deuterium excess in an East Antarctic ice core suggests higher relative humidity at the oceanic surface during the last glacial maximum, *Nature*, 299, 688–691, <https://doi.org/10.1038/299688a0>, 1982.
- Jouzel, J. and Merlivat, L.: Deuterium and Oxygen 18 in Precipitation: Modeling of the Isotopic Effects During Snow Formation, *J. Geophys. Res.*, 89, 11749–11757, <https://doi.org/10.1029/JD089iD07p11749>, 1984.
- 1095 King, J. C. and Anderson, P. S.: Heat and water vapour fluxes and scalar roughness lengths over an Antarctic ice shelf, *Boundary-Layer Meteorol*, 69, 101–121, <https://doi.org/10.1007/BF00713297>, 1994.

- King, J. C., Anderson, P. S., and Mann, G. W.: The seasonal cycle of sublimation at Halley, Antarctica, *J. Glaciol.*, 47, 1–8, <https://doi.org/10.3189/172756501781832548>, 2001.
- 1100 Kino, K., Okazaki, A., Cauquoin, A., and Yoshimura, K.: Contribution of the Southern Annular Mode to Variations in Water Isotopes of Daily Precipitation at Dome Fuji, East Antarctica, *JGR Atmospheres*, 126, e2021JD035397, <https://doi.org/10.1029/2021JD035397>, 2021.
- Kopec, B. G., Feng, X., Posmentier, E. S., and Sonder, L. J.: Seasonal Deuterium Excess Variations of Precipitation at Summit, Greenland, and their Climatological Significance, *JGR Atmospheres*, 124, 72–91, <https://doi.org/10.1029/2018JD028750>, 2019.
- 1105 Landais, A., Stenni, B., Masson-Delmotte, V., Jouzel, J., Cauquoin, A., Fourré, E., Minster, B., Selmo, E., Extier, T., Werner, M., Vimeux, F., Uemura, R., Crotti, I., and Grisart, A.: Interglacial Antarctic–Southern Ocean climate decoupling due to moisture source area shifts, *Nat. Geosci.*, 14, 918–923, <https://doi.org/10.1038/s41561-021-00856-4>, 2021.
- Landais, A., Casado, M., Ollivier, I., and Minster, B.: Isotopic composition of surface and subsurface snow at Dome C, East Antarctica (2017–2021), PANGAEA [dataset], <https://doi.pangaea.de/10.1594/PANGAEA.971486>, 2024.
- 1110 Leduc-Leballeur, M., Picard, G., Macelloni, G., Arnaud, L., Brogioni, M., Mialon, A., and Kerr, Y. H.: Influence of snow surface properties on L-band brightness temperature at Dome C, Antarctica, *Remote Sensing of Environment*, 199, 427–436, <https://doi.org/10.1016/j.rse.2017.07.035>, 2017.
- Lettau, H. H.: Wind and temperature profile prediction for diabatic surface layers including strong inversion cases, *Boundary-Layer Meteorol.*, 17, 443–464, <https://doi.org/10.1007/BF00118610>, 1979.
- 1115 Libois, Q., Picard, G., Arnaud, L., Morin, S., and Brun, E.: Modeling the impact of snow drift on the decameter-scale variability of snow properties on the Antarctic Plateau, *J. Geophys. Res. Atmos.*, 119, 11,662–11,681, <https://doi.org/10.1002/2014JD022361>, 2014.
- Lorius, C. and Merlivat, L.: Distribution of mean surface stable isotopes values in East Antarctica; Observed changes with depth in coastal area, *General Assembly of the International Union of Geodesy and Geophysics*, 1975.
- Lorius, C., Merlivat, L., and Hagemann, R.: Variation in the mean deuterium content of precipitations in Antarctica, *J. Geophys. Res.*, 74, 7027–7031, <https://doi.org/10.1029/JC074i028p07027>, 1969.
- 1120 Lupi, A., Lanconelli, C., and Vitale, V.: Basic and other measurements of radiation at Concordia station (2006–01 et seq), PANGAEA [dataset], <https://doi.org/10.1594/PANGAEA.935421>, 2021.
- Masson-Delmotte, V., Landais, A., Stievenard, M., Cattani, O., Falourd, S., Jouzel, J., Johnsen, S. J., Dahl-Jensen, D., Sveinbjornsdottir, A., White, J. W. C., Popp, T., and Fischer, H.: Holocene climatic changes in Greenland: Different deuterium excess signals at Greenland Ice Core Project (GRIP) and NorthGRIP, *J. Geophys. Res.*, 110, 2004JD005575, <https://doi.org/10.1594/PANGAEA.935421>, 2005.
- 1125 Masson-Delmotte, V., Hou, S., Ekaykin, A., Jouzel, J., Aristarain, A., Bernardo, R. T., Bromwich, D., Cattani, O., Delmotte, M., Falourd, S., Frezzotti, M., Gallée, H., Genoni, L., Isaksson, E., Landais, A., Helsen, M. M., Hoffmann, G., Lopez, J., Morgan, V., Motoyama, H., Noone, D., Oerter, H., Petit, J. R., Royer, A., Uemura, R., Schmidt, G. A., Schlosser, E., Simões, J. C., Steig, E. J., Stenni, B., Stievenard, M., van den Broeke, M. R., van de Wal, R. S. W., van de Berg, W. J., Vimeux, F., and White, J. W. C.: A Review of Antarctic Surface Snow Isotopic Composition: Observations, Atmospheric Circulation, and Isotopic Modeling*, *Journal of Climate*, 21, 3359–3387, <https://doi.org/10.1175/2007JCLI2139.1>, 2008.
- 1130 Merlivat, L. and Jouzel, J.: Global climatic interpretation of the deuterium-oxygen 18 relationship for precipitation, *J. Geophys. Res.*, 84, 5029–5033, <https://doi.org/10.1029/JC084iC08p05029>, 1979.

- 1135 Monin, A. S. and Obukhov, A. M.: Basic laws of turbulent mixing in the surface layer of the atmosphere, *Trudy Geofiz Inst*, 24, 163–187, 1954.
- Münch, T., Kipfstuhl, S., Freitag, J., Meyer, H., and Laepple, T.: Constraints on post-depositional isotope modifications in East Antarctic firn from analysing temporal changes of isotope profiles, *The Cryosphere*, 11, 2175–2188, <https://doi.org/10.5194/tc-11-2175-2017>, 2017.
- 1140 Murphy, D. M. and Koop, T.: Review of the vapour pressures of ice and supercooled water for atmospheric applications, *Q. J. R. Meteorol. Soc.*, 131, 1539–1565, <https://doi.org/10.1256/qj.04.94>, 2005.
- Picard, G., Royer, A., Arnaud, L., and Fily, M.: Influence of meter-scale wind-formed features on the variability of the microwave brightness temperature around Dome C in Antarctica, *The Cryosphere*, 8, 1105–1119, <https://doi.org/10.5194/tc-8-1105-2014>, 2014.
- Picard, G., Arnaud, L., Caneill, R., Lefebvre, E., and Lamare, M.: Observation of the process of snow accumulation on the Antarctic Plateau by time lapse laser scanning, *The Cryosphere*, 13, 1983–1999, <https://doi.org/10.5194/tc-13-1983-2019>, 2019.
- 1145 Risi, C., Landais, A., Winkler, R., and Vimeux, F.: Can we determine what controls the spatio-temporal distribution of d-excess and 17O-excess in precipitation using the LMDZ general circulation model?, *Clim. Past*, 9, 2173–2193, <https://doi.org/10.5194/cp-9-2173-2013>, 2013.
- Servettaz, A. P. M., Agosta, C., Kittel, C., and Orsi, A. J.: Control of the temperature signal in Antarctic proxies by snowfall dynamics, *The Cryosphere*, 17, 5373–5389, <https://doi.org/10.5194/tc-17-5373-2023>, 2023.
- 1150 Steen-Larsen, H. C., Masson-Delmotte, V., Sjolte, J., Johnsen, S. J., Vinther, B. M., Bréon, F.-M., Clausen, H. B., Dahl-Jensen, D., Falourd, S., Fettweis, X., Gallée, H., Jouzel, J., Kageyama, M., Lerche, H., Minster, B., Picard, G., Punge, H. J., Risi, C., Salas, D., Schwander, J., Steffen, K., Sveinbjörnsdóttir, A. E., Svensson, A., and White, J.: Understanding the climatic signal in the water stable isotope records from the NEEM shallow firn/ice cores in northwest Greenland, *J. Geophys. Res.*, 116, D06108, <https://doi.org/10.1029/2010JD014311>, 2011.
- 1155 Steen-Larsen, H. C., Sveinbjörnsdóttir, A. E., Peters, A. J., Masson-Delmotte, V., Guishard, M. P., Hsiao, G., Jouzel, J., Noone, D., Warren, J. K., and White, J. W. C.: Climatic controls on water vapor deuterium excess in the marine boundary layer of the North Atlantic based on 500 days of in situ, continuous measurements, *Atmos. Chem. Phys.*, 14, 7741–7756, <https://doi.org/10.5194/acp-14-7741-2014>, 2014a.
- 1160 Steen-Larsen, H. C., Masson-Delmotte, V., Hirabayashi, M., Winkler, R., Satow, K., Prié, F., Bayou, N., Brun, E., Cuffey, K. M., Dahl-Jensen, D., Dumont, M., Guillevic, M., Kipfstuhl, S., Landais, A., Popp, T., Risi, C., Steffen, K., Stenni, B., and Sveinbjörnsdóttir, A. E.: What controls the isotopic composition of Greenland surface snow?, *Clim. Past*, 10, 377–392, <https://doi.org/10.5194/cp-10-377-2014>, 2014b.
- Stenni, B., Masson-Delmotte, V., Johnsen, S., Jouzel, J., Longinelli, A., Monnin, E., Röthlisberger, R., and Selmo, E.: An Oceanic Cold Reversal During the Last Deglaciation, *Science*, 293, 2074–2077, <https://doi.org/10.1126/science.1059702>, 2001.
- 1165 Stenni, B., Sarchilli, C., Masson-Delmotte, V., Schlosser, E., Ciardini, V., Dreossi, G., Grigioni, P., Bonazza, M., Cagnati, A., Karlicek, D., Risi, C., Udisti, R., and Valt, M.: Three-year monitoring of stable isotopes of precipitation at Concordia Station, East Antarctica, *The Cryosphere*, 10, 2415–2428, <https://doi.org/10.5194/tc-10-2415-2016>, 2016.
- Touzeau, A., Landais, A., Stenni, B., Uemura, R., Fukui, K., Fujita, S., Guilbaud, S., Ekaykin, A., Casado, M., Barkan, E., Luz, B., 1170 Magand, O., Teste, G., Le Meur, E., Baroni, M., Savarino, J., Bourgeois, I., and Risi, C.: Acquisition of isotopic composition for surface snow in East Antarctica and the links to climatic parameters, *The Cryosphere*, 10, 837–852, <https://doi.org/10.5194/tc-10-837-2016>, 2016.

- Town, M. S., Warren, S. G., Walden, V. P., and Waddington, E. D.: Effect of atmospheric water vapor on modification of stable isotopes in near-surface snow on ice sheets, *J. Geophys. Res.*, 113, 2008JD009852, <https://doi.org/10.1029/2008JD009852>, 2008.
- 1175 Uemura, R., Matsui, Y., Yoshimura, K., Motoyama, H., and Yoshida, N.: Evidence of deuterium excess in water vapor as an indicator of ocean surface conditions, *J. Geophys. Res.*, 113, 2008JD010209, <https://doi.org/10.1029/2008JD010209>, 2008.
- Uemura, R., Masson-Delmotte, V., Jouzel, J., Landais, A., Motoyama, H., and Stenni, B.: Ranges of moisture-source temperature estimated from Antarctic ice cores stable isotope records over glacial–interglacial cycles, *Clim. Past*, 8, 1109–1125, <https://doi.org/10.5194/cp-8-1109-2012>, 2012.
- 1180 Vignon, E., Genthon, C., Barral, H., Amory, C., Picard, G., Gallée, H., Casasanta, G., and Argentini, S.: Momentum- and Heat-Flux Parametrization at Dome C, Antarctica: A Sensitivity Study, *Boundary-Layer Meteorol.*, 162, 341–367, <https://doi.org/10.1007/s10546-016-0192-3>, 2016.
- Vignon, E., Raillard, L., Genthon, C., Del Guasta, M., Heymsfield, A. J., Madeleine, J.-B., and Berne, A.: Ice fog observed at cirrus temperatures at Dome C, Antarctic Plateau, *Atmos. Chem. Phys.*, 22, 12857–12872, <https://doi.org/10.5194/acp-22-12857-2022>,
1185 2022.
- Vimeux, F., Masson, V., Jouzel, J., Stievenard, M., and Petit, J. R.: Glacial–interglacial changes in ocean surface conditions in the Southern Hemisphere, *Nature*, 398, 410–413, <https://doi.org/10.1038/18860>, 1999.
- Wahl, S., Steen-Larsen, H. C., Reuder, J., and Hörhold, M.: Quantifying the Stable Water Isotopologue Exchange Between the Snow Surface and Lower Atmosphere by Direct Flux Measurements, *JGR Atmospheres*, 126, e2020JD034400, <https://doi.org/10.1029/2020JD034400>, 2021.
1190
- Wahl, S., Steen-Larsen, H. C., Hughes, A. G., Dietrich, L. J., Zuhr, A., Behrens, M., Faber, A. -K., and Hörhold, M.: Atmosphere-Snow Exchange Explains Surface Snow Isotope Variability, *Geophysical Research Letters*, 49, <https://doi.org/10.1029/2022GL099529>, 2022.
- Wahl, S., Walter, B., Aemisegger, F., Bianchi, L., and Lehning, M.: Identifying airborne snow metamorphism with stable water isotopes, *EGU sphere* [preprint], <https://doi.org/10.5194/egusphere-2024-745>, 2024.
1195
- Werner, M., Jouzel, J., Masson-Delmotte, V., and Lohmann, G.: Reconciling glacial Antarctic water stable isotopes with ice sheet topography and the isotopic paleothermometer, *Nat Commun*, 9, 3537, <https://doi.org/10.1038/s41467-018-05430-y>, 2018.
- Zuhr, A. M., Münch, T., Steen-Larsen, H. C., Hörhold, M., and Laepple, T.: Local-scale deposition of surface snow on the Greenland ice sheet, *The Cryosphere*, 15, 4873–4900, <https://doi.org/10.5194/tc-15-4873-2021>, 2021.
- 1200 Zuhr, A. M., Wahl, S., Steen-Larsen, H. C., Hörhold, M., Meyer, H., and Laepple, T.: A Snapshot on the Buildup of the Stable Water Isotopic Signal in the Upper Snowpack at EastGRIP on the Greenland Ice Sheet, *JGR Earth Surface*, 128, e2022JF006767, <https://doi.org/10.1029/2022JF006767>, 2023.