

## Supplementary materials

### S1 Conversion of $RH_{wrti}$ to $RH_{wrti}$

As in Genthon et al. (2017) and Vignon et al. (2022), we use the combination of the three parameters measured simultaneously by the modified HMP155 ( $T_{heat}$ ,  $T_{amb}$  and  $RH_{wrti}$ ) to recalculate  $RH_{wrti}$ . This method assumes that the partial vapor pressure of water vapor is the same whether the air sampled by the sensor is heated or not.  $RH_{wrti}$  is computed as follows:

$$RH_{wrti} = \frac{RH_{wrti} * p_{sat,i}(T_{heated})}{p_{sat,i}(T_{ambient})} \quad (S1)$$

with  $p_{sat,i}$  and  $p_{sat,i}$  the equations for saturation vapor pressure as a function of temperature from Murphy and Koop (2005) (their Eq. 7 and 10) and  $RH_{wrti}$ ,  $T_{heated}$  and  $T_{ambient}$  given by sensor (see also Fig. 1 in Genthon et al., 2022).

### S2 Calculation of standard error of the mean (SEM) and 95% confidence interval (Student t-test)

The standard error of the mean is computed as follows:

$$SEM = \frac{std}{\sqrt{N^*-1}} \quad (S2)$$


The mean values are given with the 95% confidence interval (Student t-test) computed as follows:

$$\mu = \bar{x} \pm t_c \cdot \frac{SEM}{\sqrt{N^*-1}} \quad (S3)$$

In both Eq. (S2) and (S3),  $N^*$  is computed as in Bretherton et al. 1999:

$$N^* = N \cdot \frac{(1-r(\Delta t)^2)}{(1+r(\Delta t)^2)} \quad (S4)$$

with  $r(\Delta t)^2$  from the autocorrelation function (ACF) plots for each variable. As an example, for the  $\delta^{18}O$  in the surface snow,  $r(\Delta t)^2 = 0.66^2$  (correlation coefficient at lag 1, red dots in Fig. S1).



**Figure S1.** Autocorrelation function for  $\delta^{18}O$  in the surface snow.

Table S1 summarizes the values for  $N^*$  and  $t_c$  for each variable used to calculate the SEM and the 95% confidence interval.


**Table S1.** Summary of  $N^*$  and Student t-test  $t_c$  values for each variable.

	$N^*$	$t_c$
Precipitation $\delta^{18}\text{O}$ (observations)	244	1.97
Precipitation d-excess (observations)	572	1.964
Precipitation (excl. samples with $d_{xs} < 0$ ) $\delta^{18}\text{O}$ (observations)	246	1.97
Precipitation (excl. samples with $d_{xs} < 0$ ) d-excess (observations)	550	1.964
Snow surface $\delta^{18}\text{O}$ (observations)	193	1.972
Snow surface d-excess (observations)	256	1.969
Snow subsurface $\delta^{18}\text{O}$ (observations)	354	1.967
Snow subsurface d-excess (observations)	393	1.966
Precipitation $\delta^{18}\text{O}$ (ECHAM6-wiso)	479	1.965
Precipitation d-excess (ECHAM6-wiso)	1069	1.96

### S3 Inputs to SISG model and simulations results

#### S3.1 Isotopic composition of precipitation

Figure S2 shows the comparison between the observed isotopic composition of precipitation (daily samples collected at Dome C) and the three artificial timeseries used as inputs to the SISG model.



**Figure S2.** Comparison between the precipitation  $\delta^{18}\text{O}$  (a) and d-excess (b) in the daily samples collected at Dome C (circles) and the three artificial timeseries of the precipitation isotopic composition based on the atmospheric temperature (red), the precipitation-weighted mean annual isotope cycle (blue-green) and the arithmetic mean annual isotope cycle (blue).

### S3.2 Mean isotopic composition of observed and modelled snow layers

Table S2 summarizes the mean isotopic composition (over the five-year period 2017-2021) of the observed and modelled (five experiments with the SISG model) surface and subsurface snow layers.

**Table S2.** Mean isotopic composition of the observed and simulated snow surface over 5 years ( $\delta^{18}\text{O}$  in normal font, d-excess in parenthesis and italic).

	Observations	“Iso from T & cst accu”	“Iso from T & ERA5 accu”	“Iso from wg. mm & obs accu”	“Iso from ar. mm & obs accu”	“Iso from ECHAM6 & ECHAM6 accu”
Surface layer (%)	-51.0 <i>(10.4)</i>	-56.1 <i>(17.4)</i>	-53.2 <i>(13.9)</i>	-53.0 <i>(11.1)</i>	-55.7 <i>(14.0)</i>	-45.8 <i>(5.0)</i>
Subsurface layer (%)	-51.4 <i>(10.8)</i>	-56.3 <i>(17.5)</i>	-53.5 <i>(14.4)</i>	-53.8 <i>(12.1)</i>	-56.8 <i>(15.5)</i>	-45.5 <i>(5.1)</i>

### S3.3 Sensitivity tests on sample depths in the SISG model


We performed the experiment “iso from T and ERA5 accu” (isotopic composition of precipitation calculated from the atmospheric temperature and precipitation amounts from ERA5) with varying the surface and subsurface samples depths: 0-0.5 cm and 0.5-3.5 cm, 0-1.5 cm and 0.5-4.5 cm, 0-2 cm and 2-5 cm, 0-3 cm and 3-6 cm. The reference depths are 0-1 cm and 1-4 cm. In the current version of the model, the samples depths cannot overlap. The linear regression slopes (a) and RMSE between all modelled and observed monthly means  $\delta^{18}\text{O}$  and d-excess in the five-year period are summarized in Table S3.

**Table S3.** Linear regression slope (a) and RMSE between the observed and modelled monthly mean isotopic composition of the subsurface layer ( $\delta^{18}\text{O}$  in normal font, d-excess in parenthesis and italic) for different sample depths in the model. All linear slopes are significant (p-value < 0.05), except the ones marked with an asterisk (\*, p-value > 0.05).

	Depths	0-0.5-3.5 cm	<b>0-1-4 cm (ref)</b>	0-1.5-4.5 cm	0-2-5 cm	0-3-6 cm
Surface layer	a	1.1 <i>(1.1)</i>	<b>1.2 <i>(0.9)</i></b>	1.1 <i>(0.7)</i>	1.0 <i>(0.5)</i>	0.7 <i>(0.3*)</i>
	RMSE (‰)	4.4 <i>(5.7)</i>	<b>3.7 <i>(5.3)</i></b>	3.3 <i>(5.2)</i>	3.2 <i>(5.1)</i>	3.0 <i>(5.2)</i>
Subsurface layer	a	0.9 <i>(0.6)</i>	<b>0.8 <i>(0.2*)</i></b>	0.5 <i>(-0.2*)</i>	0.4* <i>(-0.5)</i>	-0.1* <i>(-1.0)</i>
	RMSE (‰)	2.9 <i>(4.6)</i>	<b>3.2 <i>(5.1)</i></b>	3.7 <i>(5.5)</i>	4.0 <i>(5.9)</i>	4.4 <i>(6.3)</i>


### S3.4 Time span corresponding to samples depths

We retrieve from the SISG model the period of precipitation corresponding to the snow samples depths (0 to 1 cm depth for the surface layer and 1 to 4 cm depth for the subsurface layer) for different daily precipitation amounts (constant, observed, from ERA5 and from ECHAM6-wiso simulations). The period corresponds to the number of days necessary to build the snow layers. It assumes that the snowfall events are deposited on top of each other without any removal or redistribution and doesn't include the effect of snow compaction.




**Figure S3.** Average time span necessary to build the snow samples (0-1 cm for the surface and 1-4 cm for the subsurface), depending on the daily precipitation amounts. Panel (a) shows the mean (over the five-year period 2017-2021) number of days necessary to build a snow layer 1 cm thick (i.e. number of days “included” in the surface sample). Panel (c) shows the mean number of days necessary to build the subsurface layer (1-4 cm depth, excluding the surface sample). Panels (b) and (d) show which months are included in each snow layer. For example, a snow surface sample taken at the beginning of January integrates, on average over five years, precipitation events fallen in November and December prior to sample collection (using ERA5 precipitation amounts). A subsurface sample taken at the beginning of January integrates precipitation events fallen between June and October prior to sample collection.

## S4 Snow isotopic composition and wind speed



**Figure S4.** Average difference in  $\delta^{18}O$  (a, b) and d-excess (c, d) between two consecutive sampling of the snow surface (a, c) and subsurface (b, d) during periods where the mean (and maximum) wind speed at 3 m is above a certain percentile. Wind percentiles are calculated over the whole 2017-2021 period.



**Figure S5.** Average difference in  $\delta^{18}O$  (a) and d-excess (b) between snow surface and subsurface during periods where the mean (and maximum) wind speed at 3 m is above a certain percentile. Wind percentiles are calculated over the whole 2017-2021 period.

## References for supplementary materials

- Bretherton, C. S., Widmann, M., Dymnikov, V. P., Wallace, J. M., and Bladé, I.: The Effective Number of Spatial Degrees of Freedom of a Time-Varying Field, *J. Climate*, 12, 1990–2009, [https://doi.org/10.1175/1520-0442\(1999\)012<1990:TENOSD>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<1990:TENOSD>2.0.CO;2), 1999.
- Genthon, C., Piard, L., Vignon, E., Madeleine, J.-B., Casado, M., and Gallée, H.: Atmospheric moisture supersaturation in the near-surface atmosphere at Dome C, Antarctic Plateau, *Atmos. Chem. Phys.*, 17, 691–704, <https://doi.org/10.5194/acp-17-691-2017>, 2017.
- Genthon, C., Veron, D. E., Vignon, E., Madeleine, J.-B., and Piard, L.: Water vapor in cold and clean atmosphere: a 3-year data set in the boundary layer of Dome C, East Antarctic Plateau, *Earth Syst. Sci. Data*, 14, 1571–1580, <https://doi.org/10.5194/essd-14-1571-2022>, 2022.
- Murphy, D. M. and Koop, T.: Review of the vapour pressures of ice and supercooled water for atmospheric applications, *Q. J. R. Meteorol. Soc.*, 131, 1539–1565, <https://doi.org/10.1256/qj.04.94>, 2005.
- Vignon, E., Raillard, L., Genthon, C., Del Guasta, M., Heymsfield, A. J., Madeleine, J.-B., and Berne, A.: Ice fog observed at cirrus temperatures at Dome C, Antarctic Plateau, *Atmos. Chem. Phys.*, 22, 12857–12872, <https://doi.org/10.5194/acp-22-12857-2022>, 2022.