

Comparison of shortwave radiation dynamics between boreal forest and open peatland pairs in southern and northern Finland

Otso Peräkylä¹, Erkkä Rinne², Ekaterina Ezhova¹, Anna Lintunen^{1,3}, Annalea Lohila^{1,2}, Juho Aalto^{3,4}, Mika Aurela², Pasi Kolari¹, and Markku Kulmala¹

¹Institute for Atmospheric and Earth System Research / Physics, Faculty of Science, University of Helsinki, Finland

²Climate System Research, Finnish Meteorological Institute, PL 503, FI-00101 Helsinki, Finland

³Institute for Atmospheric and Earth System Research / Agricultural and Forest Sciences, Faculty of Agriculture and Forestry, University of Helsinki, Finland

⁴Department of Forest Sciences, Faculty of Agriculture and Forestry, University of Helsinki, P.O. Box 27, FI-00014 Helsinki, Finland

Correspondence: Otso Peräkylä (otso.perakyla@helsinki.fi) and Markku Kulmala (markku.kulmala@helsinki.fi)

Abstract.

Snow cover plays a key role in determining the albedo, and thus the shortwave radiation balance, of a surface. The effect of snow on albedo is modulated by land use: tree canopies break the uniform snow layer, and lower the albedo, as compared to an open ground. This results in a higher fraction of shortwave radiation being absorbed in forests. At seasonally snow-covered high latitudes, this lowering of the albedo has been suggested to offset some or all of the climate cooling effect of the carbon stored by forests. We used long-term in situ measurements to study the albedo and shortwave radiation balance of two pairs of sites, each consisting of an open peatland and a forest. One pair is located in northern and one in southern Finland in the boreal zone. We found that both forest sites had a low, constant albedo during the snow-free period. In contrast, both peatland sites had a higher snow-free albedo, with a clear seasonal cycle. ~~This seasonal cycle was presumably caused by changing near-infrared albedo, as the albedo for photosynthetically active radiation was considerably more constant over the season~~ The albedo was found to depend on the diffuse fraction of the incoming radiation, with contrasting dependencies observed in summer and winter. The thinning of the southern forest site, resulting in a significant reduction of the leaf area index, increased the albedo especially in the snow covered period. During the snow-covered period, the peatland sites again had higher albedo than the forest sites. The transition between the high and low albedo upon snow accumulation and especially snowmelt was more abrupt at the peatland sites. ~~The annual difference in absorbed shortwave radiation~~ In the northern pair, the forest site absorbed on average 0.47 GJ m⁻² more (around 23% more) energy from the incoming shortwave radiation than the peatland site annually, whereas in the southern pair, the forest site absorbed on average 0.37 GJ m⁻² more (around 14% more) than the peatland site. The difference in the annual absorbed energy between the peatland and the forest site was greater in the northern site-pair, pair due to longer snow cover duration. This was partially offset by the greater difference in snow-free albedos and higher solar radiation at the southern site pair. ~~Annual variation in the differences in~~ The annual difference in the absorbed shortwave radiation between ~~forest and peatland sites~~ the forest and the peatland site varied considerably between the years (from 0.37 to 0.61 GJ m⁻² for the northern pair, and from 0.20 to 0.53 GJ m⁻² for the southern pair). The annual variation was mainly

controlled by the snow ~~melt date~~ cover duration in the spring at the peatland sites. These findings have implications for the future climate, as snow cover continues to evolve under global warming.

25 *Copyright statement.* TEXT

1 Introduction

Surface albedo (hereafter albedo) is the fraction of incident global broadband shortwave radiation that a surface reflects (Liang et al., 2010). The albedo, together with the amount of incident shortwave radiation, determines the shortwave energy input to the surface, typically the main mechanism by which the surface gains energy (Eugster et al., 2000; Trenberth et al., 2009; Liang et al., 2010). ~~The surface albedo differs from the top-of-atmosphere albedo in that the latter also includes the contribution of atmospheric scattering and absorption, which can dampen the effect of changes in the surface (Stephens et al., 2015).~~

Changes in albedo play a key role in major earth system processes and feedbacks (Charney et al., 1975; Sagan et al., 1979; Henderson-Sellers and Wilson, 1983; Courel et al., 1984; Curry et al., 1995; Déry and Brown, 2007; Zeng and Yoon, 2009; Loew, 2014; Pithan and Mauritsen, 2014). Snow typically has a very high reflectance in near-UV and visible range, and as a result snow cover is especially important for determining the albedo (Wiscombe and Warren, 1980). In the boreal zone, the presence of trees has a major impact on the wintertime albedo, as the tree canopy breaks up the uniform snow surface, substantially lowering the albedo (Betts and Ball, 1997; Baldocchi et al., 2000; Eugster et al., 2000; Kuusinen et al., 2012; Essery, 2013; Thackeray et al., 2014; Manninen et al., 2022). As a result, the difference in the between winter- and summertime albedo is much smaller in forests than in open areas in the boreal zone (Betts and Ball, 1997; Baldocchi et al., 2000; Eugster et al., 2000).

In addition to the properties of the surface, the albedo also depends on the properties of the incoming shortwave radiation. For example, the fraction of diffuse radiation and the solar zenith angle affect the albedo (Wiscombe and Warren, 1980; Eugster et al., 2000; Yang et al., 2008; Wang et al., 2015; Qu et al., 2015). ~~The diffuse fraction can also be different for different wavelengths of the incoming radiation (Ezhova et al., 2018).~~ The albedo under fully diffuse radiation is known as the white-sky albedo, while that under only direct beam radiation is the black-sky albedo (Qu et al., 2015). The albedo under ambient lighting conditions, the blue-sky albedo, can be approximated as a combination of these (Qu et al., 2015).

Afforestation is often considered an effective climate change mitigation option due to the potential of forests to store carbon (Goymer, 2018). However, many global studies have argued that in the boreal zone, the climate benefit of the sequestered carbon is either partially or fully negated by decreases in albedo, and thus increased absorption of sunlight solar radiation (Betts, 2000; Bala et al., 2007; Scott et al., 2018). Similar results have been found in Finland, where extensive peatland areas have been drained for forestry use (Lohila et al., 2010; Gao et al., 2014). In the case of drained peatlands, the decomposition of peat and the associated greenhouse gas emissions play a key role in determining the total climate effect of the land use change (Lohila et al., 2010). In contrast to greenhouse gases, which have a global effect, changes in albedo have a strong local impact

on the climate (Betts, 2000). Additionally, the effect of changing albedo on local temperatures is most prominently seen in the
55 springtime, when the solar radiation levels are rapidly increasing but snow cover is still present (Lohila et al., 2010; Gao et al.,
2014).

Snow affects the climate in many ways (Cohen and Rind, 1991). Climate change is already affecting the cryosphere, includ-
ing snow cover, and the impacts are projected to increase with future warming (Intergovernmental Panel On Climate Change
(IPCC), 2022). The extent and duration of snow cover are changing, but these changes are not spatially uniform (Anttila et al.,
60 2018; Bormann et al., 2018; Brown and Mote, 2009; Derksen and Brown, 2012; Manninen et al., 2019; Pulliainen et al., 2020)
Snow cover is affected by various factors, such as total precipitation, phase of precipitation (snow/rain), and occurrence of
melting events during winter, which are impacted by climate change differently (Räsänen, 2021). As a result, the changes in
snow cover in colder and warmer regions, e.g. in different parts of North Europe, sometimes ~~have opposite trends due~~ respond
differently to warming (Räsänen, 2021).

65 Satellite observations are often used to study both temporal and spatial variation in albedo (e.g. Qu et al., 2015; Anttila
et al., 2018). However, satellite products are limited in spatial resolution, with one pixel often consisting of multiple land cover
types (Kuusinen et al., 2013; Hovi et al., 2019). Disentangling the information into separate land cover types is possible, but
typically requires prior information on land cover types (Kuusinen et al., 2013). In addition, satellite estimates of albedo are
typically of lower quality in the wintertime (~~Kuusinen et al., 2013; Hovi et al., 2019~~). due to e.g. cloud cover and low solar
70 elevation angles (Kuusinen et al., 2013; Hovi et al., 2019). The latter, through low solar radiation, also decreases the quality of
in-situ measurements. However, in-situ measurements of albedo are available even under cloud cover.

To study the effect of snow and land use on albedo and shortwave radiation balance in detail, we analysed long-term observa-
tional in-situ data from two site pairs in Finland, each consisting of a forest and an open peatland, located closely together. One
pair is located in southern Finland, and the other in the northern part of the country, ~~above~~ north of the Arctic Circle (Fig. 1).
75 Each site had continuous measurements of both up- and downwelling shortwave (SW) and photosynthetically active radiation
(PAR). The proximity of the sites within each pair to each other allowed us to isolate the effect of land cover on the albedo and
shortwave radiation balance from other effects, such as changes in snowfall and cloudiness due to atmospheric circulation.

Specifically, we aimed to answer the following research questions:

1. How does albedo differ between adjacent forest and peatland sites, and between southern and northern boreal sites?
- 80 2. What determines the temporal ~~behaviour of albedo~~ variation of albedo on seasonal time scales?
3. How do these differences ~~affect yearly~~ in albedo affect the magnitude and variability of annual energy inputs from
shortwave radiation?




Figure 1. Map of a) the location of the sites in Finland, b) the northern Halssiaapa-Halssikangas site pair and c) the southern Siikaneva-Hyytiälä site pair. Peatland sites in red circles, forest sites in green triangles. Note differing scales. Data from the National Land Survey of Finland aerial photographs, accessed 1/2024.

2 Methods

2.1 Measurement sites

85 We used data from two pairs of measurement sites in the boreal zone, each pair consisting of a forested site and an open peatland site located closely together. The pairs were the Hyytiälä forest (Hari and Kulmala, 2005; Kolari et al., 2022; Aalto et al., 2023b) and Siikaneva-1 open peatland (Aurela et al., 2007; Alekseychik et al., 2017; Rinne et al., 2018) sites in southern Finland, located 5.6 km from each other, and the Halssikangas forest and Halssiaapa open peatland sites (Linkosalmi et al., 2016) at the Sodankylä Arctic Space Centre, located 0.8 km apart, north of the Arctic Circle (Fig. 1). Both forest sites are covered by evergreen conifer forest dominated by *Pinus sylvestris* (*L.*), while both peatland sites are mainly covered by *Sphagnum* mosses, sedges and shrubs (Table 1). The southern Hyytiälä forest has a substantially higher leaf area index (LAI) than the northern Halssikangas ([2.1 m² m⁻² before, 1.6 m² m⁻² after thinning in Hyytiälä compared to 1.37 m² m⁻² in Halssikangas](#), Table 1). The Hyytiälä forest was thinned in 2020, removing some 40% of foliar mass (Aalto et al., 2023b), with a similar reduction in LAI, on which forest albedo has been found to depend (Lukeš et al., 2013).

90

Table 1. Description of the measurement sites. More details on the [measurements-instrumentation and measurement heights](#) in Table A1.

	Siikaneva	Hyytiälä	Halsiaapa	Halssikangas
Latitude (WGS84)	N61.832707	N61.847411	N67.367070	N67.361866
Longitude (WGS84)	E24.192758	E24.294731	E26.651170	E26.637728
Height (m asl)	162	181	180	179
Ecosystem type	Oligotrophic fen	Forest	Mesotrophic fen	Forest
Dominant tree species		<i>Pinus sylvestris</i>		<i>Pinus sylvestris</i>
Tree height (m)		19.0 ^a , 20.9 ^b		14.8 (2022)
Tree density (ha ⁻¹)		934 ^a , 439 ^b		2100 (2022)
Basal area (m ² ha ⁻¹)		31 ^a , 18 ^b	22 (2011)	
Leaf area index (LAI) (m ² m ⁻²), projected		2.1 (2019), 1.6 (2022) ^{cde}		1.37 (2022) ^d
Soil type	Peat	Podzol	Peat	Podzol
Reference	Aurela et al. (2007); Aleksyechik et al. (2005); (2017); Rinne et al. (2022); (2018)	Hari and Kulmala (2005); Kolari et al. (2022); Aalto et al. (2023b)	Linkosalmi et al. (2016)	Linkosalmi et al. (2016)

^aBefore thinning, Aalto et al. (2023b)

^bAfter thinning, Aalto et al. (2023b)

^cHyytiälä LAI is evolving rapidly, see Kolari et al. (2022). The stand was thinned in 2020, reducing LAI, with partial recovery during subsequent years.

^dCalculated from hemispherical images, includes the overstory. Small trees, such as spruces that were removed in Hyytiälä during the thinning, are underrepresented in the values, and the [understory-forest floor vegetation](#) is absent.

^eMammarella et al. (2023)

95 2.2 [Instrumentation](#)

[On each of the sites, up-and downwelling shortwave radiation was measured with either four-component net radiometers, or separate pyranometers for up- and downwelling radiation. The measurements were conducted from meteorological masts, with the instruments located higher at the masts for the forest sites and lower for the open peatland sites \(Appendix Table A1\). In addition, both up- and downwelling PAR was measured at each of the sites, and diffuse downwelling radiation at the forest sites.](#)

100 [For the diffuse radiation, only diffuse PAR was available for Hyytiälä, and only diffuse shortwave radiation for Halssikangas. These measurements were supplemented by automatic measurements of snow depth at each of the sites \(Table A1\).](#)

2.3 Data processing

For each of the site pairs, we chose the investigated time period to maximise the number of full years with as continuous data cover from both sites as possible. For the southern Hyytiälä forest site, radiation measurements were available from multiple

105 heights: the heights were chosen to maximise data coverage (Table A1). The resulting time periods were from year 2016 to

2023 for the southern site pair, and 2013 to 2023 for the northern site pair. In addition to the radiation data, we also used snow depth data from each of the sites. The data were downloaded as either 30-minute averages, or 10-minute averages which were averaged to a 30-minute time step. If one or two values within the 30-minute time step were missing, the average was calculated from the remaining values.

110 ~~In addition to~~ Using the 30-minute values, we ~~also~~ calculated daily means of the parameters ~~based on the 30-minute data~~. For the calculation of daily values, if any 30-minute value was missing, that day was counted as missing as well. There were three exceptions:

1. For the years 2013 and 2014 in the northern Halssiaapa peatland, any negative values had been coded as missing. These mainly occurred during nights, when the actual radiation is near zero. For these years, all nighttime values (solar zenith angle over 90°) were changed to zero prior to the calculation of daily averages.
115
2. The Halssiaapa radiation data often had a single 30-minute value missing from an otherwise complete day. These gaps were linearly interpolated.
3. The gaps during morning and evening hours were often longer. ~~With zenith angles over~~ When the solar zenith angle was larger than 87° , the interpolation was changed to fill a gap of a maximum of two 30-minute values.

120 For the calculation of daily values, any non-missing data points during the night (solar zenith angle over 90°) were changed to zero to reduce random variation. The daily albedo values were calculated as the ratio of the daily mean reflected radiation to daily mean global radiation, not as an average of 30-minute albedos.

During the winter, the sensors for incoming radiation may get covered with snow. This problem is less pronounced for the down-facing sensors for reflected radiation. This cover will lead to lower-than-expected measurements of the incoming radiation. As a result, the albedo will be overestimated. Some differences in the incoming radiation between the sites are
125 expected, due to, for example, local clouds. However, any consistent differences may be indicative of problems with the sensor. We used the ratio of the daily average global ~~radiations~~ radiation values within a site pair to diagnose these snow-covered cases, excluding any days where the incoming radiation at a site was over 20% lower than at its pair. Some cases with a snow-covered sensor may still remain, leading to artificially high albedo values. In addition, any days where the average reflected radiation
130 was higher than the average incoming radiation, were excluded from analysis.

The primary focus in the snow depth analysis was on the data from the peatland sites. The snow depth data was missing for some days (Fig. A1). To obtain a continuous time series for the peatland sites, the snow depth data was interpolated linearly for the missing parts.

2.4 Hyytiälä global radiation correction

135 There were systematic, time-dependent differences between the daily mean global radiation values measured in Hyytiälä and in Siikaneva. In the first two years of measurements, Hyytiälä values were consistently lower, and in the last two years, Hyytiälä values were consistently higher than those in Siikaneva. These differences indicate a drift in the calibration of the Hyytiälä

140 global radiation sensor. We corrected Hyytiälä's global radiation by applying a year-by-year correction factor. These factors were derived as the slopes of the total least squares fits to the values of radiation in Hyytiälä and Siikaneva for each year separately. The global radiation values in Hyytiälä were then multiplied by these correction factors.

2.5 Net shortwave radiation

To assess the effect of the different albedo values on the energy budgets, we calculated the net shortwave radiation (i.e. the difference between the global and reflected solar radiation) for each site. We then took the difference of these net values between the sites: the net shortwave radiation of the peatland site was subtracted from that of the forest site. With this convention, positive differences mean that the forest site absorbs more energy from solar radiation. The net shortwave radiation is proportional to the incoming radiation, multiplied by the absorbed fraction (one minus albedo):

$$R_n^{\text{SW}} = R_d^{\text{SW}}(1 - \alpha), \quad (1)$$

where R_n^{SW} is the net shortwave radiation, R_d^{SW} is the incoming radiation, and α is the albedo (Liang et al., 2010). As a result, the difference in the net shortwave radiation across two sites, 1 and 2, is

$$150 \quad R_{n,1}^{\text{SW}} - R_{n,2}^{\text{SW}} = R_{d,1}^{\text{SW}}(1 - \alpha_1) - R_{d,2}^{\text{SW}}(1 - \alpha_2) = R_{d,1,2}^{\text{SW}}(\alpha_2 - \alpha_1). \quad (2)$$

The last step applies if the incoming radiation is the same across the sites. As a result, the difference in net shortwave radiation depends on the incoming radiation, and the difference in the albedo.

To test how the snow-cover duration affects the net SW radiation on an annual scale, we calculated annual averages of the difference in net SW radiation from August to the next year's July. We chose this time frame to include one full winter in each averaging period. The radiation data included some gaps, so the difference in the net SW radiation between the sites was not continuous (Fig. 2). This might bias the annual energy balances if there are data missing over different periods between years. We therefore gapfilled the difference values of the net SW radiation using a data model constructed in Section A1 (Fig. A2). This resulted in 86% of the data being measured for Hyytiälä-Siikaneva, and 78% for Halssikangas-Halssiaapa, with the rest being gapfilled. These gapfilled data were then averaged to annual averages. In addition, we defined a day as snow-covered if the snow depth in the peatland site was greater than 1 cm. Based on this classification, we calculated the number of days with snow cover in the spring. This number almost always corresponded directly to the snow melt day of the year.

2.6 Linear models

We used linear regression models to interpret the data. These were used for assessing the effect of the thinning of the Hyytiälä forest on albedo, for gapfilling the difference in net SW radiation, and for assessing the effect of different factors on the annual differences in the net SW radiation between the forest and the open peatland sites. In the models, we used interaction terms to allow for different classes of observations (e.g. snow-covered vs. non-snow-covered) to have different slopes.

3 Results and discussion

3.1 General features of radiation at the sites

The global radiation measured at each of the sites at the southern site pair was broadly similar (Fig. A3 a). The same holds for the northern pair (Fig. A3 b). Differences in the 30-minute values are most probably caused mainly by local clouds: one site may be shaded, while the other one receives sunlight. This effect is more pronounced in the southern Hyytiälä-Siikaneva pair, possibly due to their greater geographical separation (5.6 km vs. 0.8 km). The regression line very close to a 1:1 line shows that these effects tend to cancel each other out: neither site is on average substantially sunnier than the other. This is supported by taking daily averages of the global radiation: these fall still closer to the 1:1 line (Fig. A3). All in all, the incoming shortwave radiation within the site pairs was very similar, and a direct comparison between the pairs is reasonable.

The similarity of the global radiation between the southern Hyytiälä forest and Siikaneva peatland is evident also in time series (Fig. 2 a and b). The reflected radiation, on the other hand, showed different patterns for the two sites. For Hyytiälä, it typically had a smooth annual cycle, with a minor peak in spring and a summer maximum. For the southern Siikaneva peatland, the annual cycle had a distinct two-peak shape. A first, higher maximum was observed in the springtime, and a second, lower maximum, occurred slightly later than the summertime maximum in global radiation. This indicates a ~~radieally~~ much higher albedo in the springtime, when the ground is covered by snow (Fig. 2 c). The strength of the springtime peak varied substantially between years. As an example, in 2018, with a consistent and long-lasting snow cover, the peak was high and continued long. In contrast, in early 2020, the snow cover was thinner and more sporadic, and the springtime peak in reflected radiation was correspondingly lower. Even during the snow-free period in the summer, the reflected radiation at Siikaneva was higher than in Hyytiälä, indicating a higher snow-free albedo. Similar features were seen in the northern Halssiaapa-Halssikangas site pair (Fig. 2 c and d). Here, even the forest site consistently saw its highest reflected radiation values towards the end of the spring. During this time, the ground was still snow-covered, and the global radiation was increasing rapidly. Again, this effect of the reflective snow cover was seen much more prominently in the reflected radiation at the peatland than at the forest site. In contrast to the southern peatland, the well-defined spring peak at the northern peatland site was present during each of the measured years, due to more persistent snow cover at the northern sites.

3.2 Effect of snow cover on reflected radiation and albedo

The reflected solar radiation in the southern Hyytiälä forest increased as a function of both global radiation and snow depth in the forest (Fig. 3 a). Especially high reflected radiation was seen at high diffuse ~~light fractions~~ fractions of radiation and/or high solar zenith angles (Fig. A4). When there was no snow on the ground, the reflected radiation was most often very close to 10% of the global radiation, indicating an albedo of 0.1 (Fig. 3 a). With snow cover, the reflected radiation increased substantially, but also became more variable, with a less linear dependence on the global radiation. At the Siikaneva peatland, the reflected fraction in the snow-free period was higher than in Hyytiälä, close to 16%. ~~It also had more variation than at the Hyytiälä forest (Fig. 3 b).~~ In contrast to the moderate increase at the forest site, the reflected solar radiation at Siikaneva was dramatically increased by snow cover, even at high global radiation levels (Fig. 3 b). For snow depths over ~~some~~ approximately 20 cm, the




Figure 2. Time series of the global and reflected shortwave radiation in (a) Hyytiälä (southern forest), (b) Siikaneva (southern peatland), (d) Halssikangas (northern forest) and (e) Halssiaapa (northern peatland), and snow depth in (c) Siikaneva and (f) Halssiaapa. The sign convention is chosen so that reflected radiation is negative. Within each site pair the x-axis is shared, and all the sites have the same y-axis for the radiation values. The radiation values are 30-minute-daily averages, while the snow depth data is daily average with interpolation of missing values. The snow depth data for Halssiaapa was only available until September 2023.

200 reflected radiation was typically over 70% of the global radiation, and albedo thus over 0.7. ~~The distinct bimodal distribution of the reflected solar radiation, now as a function of the global radiation, was clearly seen in the scatter plot as well.~~ Very similar features were seen for the northern Halssiaapa-Halssikangas site pair (Figs. 3 c and d). The forest albedo was typically around 0.11 for the snow-free period, with a substantial but variable increase in the snow-covered period (Fig. 3 c). Again, ~~the peatland albedo was during the snow-free period, the peatland had a~~ higher and more variable ~~in the snow-free period~~ albedo than the
 205 ~~forest site~~, now around 0.13 (Fig. 3 d). The peatland albedo increased dramatically with snow cover, up to values around 0.8, and there was a clear bimodal distribution of the reflected radiation. In the forest site, the snow cover increased the reflected radiation, but in a more continuous way. These albedo values for ~~the snow-free period for~~ both the southern and northern forest and peatland sites agree well with prior results, ~~such as both snow-free and snow-covered. For example, during snow-free and snow-covered periods, respectively, Bright et al. (2018) report the~~ values of 0.15 – 0.16 ~~and 0.68 – 0.77~~ for shrubs and mosaic
 210 herbaceous areas, 0.13 ~~and 0.57 – 0.59~~ for wetlands, and 0.10 – 0.12 ~~and 0.16 – 0.42~~ for pine forests (Bright et al., 2018).




Figure 3. The reflected global radiation as a function of the global radiation for a) Hyytiälä, b) Siikaneva, c) Halssikangas and d) Halssiaapa, 30-minute-daily values. The points are coloured by the snow depth measured at each site: note differing y-axis scales for forest and open peatland sites. Two example lines are drawn in each figure, corresponding to illustrative albedo values that are different for each site. The peatland sites both have a well-defined upper edge of the plot, corresponding to albedos around 0.8, while the forested sites have much less clear edges, and the values are chosen as illustrative examples.

3.3 Seasonal dynamics of albedo and influencing factors

Due to snow cover, all of the sites saw their highest albedo in the winter (Fig. 4). For the forest sites Hyytiälä and Halssikangas, the transition between the high, snow-covered albedo and the low, snow-free albedo was relatively gradual. In contrast, the peatland sites had an abrupt springtime transition from high to low albedo, on a time scale of about a week. The snow melt in the northern sites happened much later in the year, leading to high albedo being observed longer towards the summer. The transition in the autumn was still steep, but less so than in the spring. Both northern sites had a higher maximum albedo as compared to the corresponding southern sites. This difference is especially pronounced for the forest sites. Part of this difference may be due to the higher solar zenith angles at the northern sites. On all of the sites, there was more absolute variation in the wintertime albedo as compared to the summertime albedo (Fig. 4). This is partly due to actual changes in the albedo, but partly due to the low levels of radiation, and therefore higher uncertainties in the albedo determination. In addition, the less persistent snow cover at the southern sites was visible especially at Siikaneva, where the 5th percentile albedo in the wintertime was often very low as compared to typical values. For the vast majority of time, within a site pair, the albedo at the peatland site was higher than that at the forest site (Figs. 4 c and f, Fig.-??). Each of the sites also showed a decreasing trend in albedo when moving from winter to spring, the trend being stronger at the northern sites (Fig. 4). As a result, the difference in




Figure 4. Typical annual cycles of the albedo in a) Hyytiälä, b) Siikaneva, d) Halssikangas and e) Halssiaapa. Difference between the peatland and the forest site in panels c and f. The statistics are calculated for all of the daily values belonging to each week number, over all of the years. The Hyytiälä-Siikaneva data set includes eight years, the Halssikangas-Halssiaapa data set includes 12 years.

225 albedo between the peatland and forest site within each pair was remarkably constant during the snow-covered period (Figs. 4
c and f).

Albedo depends not only on snow cover, but also on e.g. snow grain size, melting state and impurities (Wiscombe and
Warren, 1980; Warren and Wiscombe, 1980; Gardner and Sharp, 2010; Dang et al., 2015; Wang et al., 2020), for which we
had no direct observations. Typically, these effects act so that the albedo of fresh snow is highest, and decreases with age
230 (Baker et al., 1990). The dependence of albedo on snow depth showed a hysteresis behaviour with respect to the day of year
(DOY): early in the winter increasing snow depth rapidly increases the albedo to values close to one, while in the springtime,
decreasing snow depth was associated with a lowering of albedo (Fig. A5). This may be due to the effect of snow age: during
the winter, snowfall events lead to the increasing snow depth seen in Fig. 2. Accumulation of fresh snow keeps the albedo high.
In the spring, even if the ground is snow-covered, the melting of snow lowers the albedo. Similar features were seen in Fig. 4.
235 The highest albedos seen at the northern Halssiaapa site, in the deep winter, were higher than those at the southern Siikaneva
(Figs. 4 and A5). This may be partly due to the high solar zenith angle. The springtime albedos are comparable to each other.

~~Dependence of daily albedo on snow depth at the peatland sites, (a) southern Siikaneva and (b) northern Halssiaapa. The
points are coloured by the day of year (DOY). The springtime values in Halssiaapa around DOY 100, with a high albedo but
a low snow depth, are associated with times when the peatland snow depth measurement has reached zero, but snow is still
240 detected at the forest site (Fig. ??)~~

Each of the sites had a clear minimum in the albedo in the snow-free period (Fig. 4). For both of the forest sites, there was little ~~interannual or within-year inter- or intra-annual~~ variation in this snow-free albedo, with an average albedo around 0.11 throughout the summer for both sites (Fig. 4 a and d). The ~~albedo for both sites showed a slight increase over the summer~~Hyytiälä forest site had slightly lower albedo during the summer, as compared to Halssikangas. Similar temporal behaviour of the forest albedo has been observed by, for example, ~~Aurela et al. (2015)~~Betts and Ball (1997); Aurela et al. (2015) and Kuusinen et al. (2012). The ~~very similar values for the snow-free albedo for the forest sites are somewhat unexpected, given their substantially different LAI (Table 1), and that albedo has been found to vary with changing LAI (Lukeš et al., 2013). However, other studies have found little change in albedo for large variations in LAI (Bright et al., 2018).~~small difference in the summertime albedo between the Hyytiälä and Halssikangas sites is consistent with their differing LAI values (Lukeš et al., 2013; Bright et al.

In contrast, the peatland sites showed a distinct seasonal cycle in the snow-free albedo, with a minimum of around 0.12 for Siikaneva and 0.08 for Halssiaapa, just after snow melt (Fig. 4 b and e). This value increased to around 0.19 for Siikaneva and 0.16 for Halssiaapa by late August, with a subsequent gradual decrease until snowfall. Very similar annual cycles have been observed for boreal peatland sites by Aurela et al. (2002, 2015); Nousu et al. (2024). Linkosalmi et al. (2016) reported seasonal development of the peatland vegetation phenology at Halssiaapa. This has a similar temporal pattern to the development of the snow-free peatland albedo in Fig. 4, and possibly explains the variation in the peatland summer albedo. Hovi et al. (2019) observed similar seasonal ~~eyes~~cycle for snow-free albedo, but for boreal forest sites. However, their study was based on satellite data, and it is possible that there was contribution from open peatlands or deciduous trees as well. Similarly, Yan et al. (2021) found deciduous forest albedo to increase with greening of the forest over the growing season. Nousu et al. (2024) found snow-free seasonal cycles for albedo at both boreal forests and peatland sites that corresponded to those found here, while the modelled albedo in their study, being a prescribed parameter, was constant. Betts and Ball (1997) observed similar annual cycle for the albedo of two boreal grasslands, but with a constant albedo over the summer.

Part of the variation in the albedo was caused by differences in the fraction of incoming diffuse radiation (Fig. 5). In the summertime, increasing fraction of diffuse radiation decreased the albedo at all the sites except for Hyytiälä. This is consistent with observations that the white-sky albedo is typically lower than the black-sky albedo (Yang et al., 2008). For Siikaneva and Halssiaapa, the diffuse fraction also affected the seasonal cycle of the albedo: the albedo with a low fraction of diffuse radiation, corresponding to a clear sky albedo, increased towards the end of summer, and then stayed at the same level, while with high diffuse radiation, corresponding to a white-sky albedo, the albedo started decreasing after the summer peak (Fig. A6). Similar patterns have been observed for satellite-derived albedo above forests by Hovi et al. (2019). In the wintertime, the effect of increasing diffuse radiation seems to be the opposite: increasing diffuse fraction increased albedo (Fig. 5). Fresh snow typically has a higher albedo than old, and snowfall is typically associated with overcast conditions, when the diffuse fraction is high. This effect may explain some of the albedo increases in high diffuse fraction conditions.

In contrast to the shortwave albedo, which had a clear snow-free seasonal cycle at the peatland sites, the PAR albedo was much more constant over the course of the snow-free period (Fig. 6). Similar results have been found for another northern peatland by Aurela et al. (2001). This means that the changes in the shortwave albedo are due to changing reflectance outside




Figure 5. The dependence of albedo on the diffuse fraction of incoming radiation. The points are coloured by the snow depth. For the southern pair, the ratio is of diffuse and total photosynthetically active radiation (PAR) in Hyytiälä, and for the northern pair, of diffuse shortwave radiation at the Sodankylä Tähtelä observatory to global shortwave radiation at the forest site.

the PAR wavelength range, likely in the near infrared. Hovi et al. (2019) observed similar patterns for multiple forest sites. ~~Linkosalmi et al. (2016) reported seasonal development of the peatland vegetation phenology at Halssiaapa. This has a similar temporal pattern to the development of the snow-free peatland albedo in Fig. 4, possibly explaining the variation in the peatland summer albedo.~~

280 3.4 Effect of the Hyytiälä forest thinning on albedo

The Hyytiälä forest stand was thinned in the beginning of 2020, removing ~~some~~ approximately 40% of the leaf mass (Aalto et al., 2023b). As the snow conditions before and after thinning were not identical, comparison of wintertime albedo without considering snow is potentially misleading. ~~Accounting for snow, we observed an increase in springtime reflected radiation after the thinning. We modelled the reflected radiation, explaining it via global radiation and albedo using a linear model.~~

285 Reflected radiation is directly proportional to global radiation with the slope of the fit representing albedo (Fig. 7). We included two binary interaction terms: whether the snow cover was over 10 cm, and whether the forest was already thinned. This allowed us to estimate the slope separately for snow-covered and snow-free, thinned and non-thinned data points. We observed an increased albedo in both the snow-free and snow-covered times, with a more pronounced effect in the snow-covered period (Fig. 7).

290 Higher maximum snow depths were observed after the thinning, but for similar snow depth-global radiation combinations, especially at high global radiation values, the albedoes after the thinning were higher (Figs. A7 and ??). This can probably




Figure 6. The dependence of PAR albedo seasonal cycle at the different sites, with focus on the diffuse fraction of incoming radiation snow-free period (y-axis clipped at 0.15). The points are coloured by the ratio Halssikangas reflected PAR sensor appears to be mounted at a slight angle, registering also part of diffuse and global incoming radiation. For the southern pair, the ratio This is evident from the increase of diffuse and total photosynthetically active radiation (the measured PAR) in Hyytiälä albedo both with increasing solar zenith angles towards the autumn, and for with increasing diffuse fraction. Both of these increase the northern pair, amount of diffuse shortwave incoming radiation at the Sodankylä Tähtelä observatory coming from close to global shortwave radiation at the forest site horizon. During In the winter, the radiation values are very low, and Hyytiälä PAR albedo and diffuse fraction, as a ratio there are clear interannual differences. Part of two small numbers these are due to sensor replacement in late 2019, uncertain see Fig. A8.

be attributed to lesser canopy coverage, with more light radiation reflected from the snow-covered ground. In contrast, there was no readily discernible change in the snow-free albedo after the thinning, even though the reduction of leaf area would be expected to increase the albedo somewhat (Lukeš et al., 2013; Launiainen et al., 2016). This is in line with the finding that the northern forest site, with a lower LAI, also had a very similar albedo. A slight increase was observed in the PAR-albedo after the thinning, with a recovery in the subsequent years (Fig. A8).

3.5 Effect of land use and snow cover on shortwave radiation balances

The On a daily time scale, the difference in the net SW radiation between the forest and peatland site depended strongly on both the global radiation and the snow cover at the peatland site, indicative of the albedo difference (Eq. (2), Fig. 8). The presence of snow, combined with the global radiation, explains the difference in net SW radiation at well, without accounting for e.g. the melting state and grain size mentioned earlier (Fig. A11). In contrast, the snow-free points show a much greater




Figure 7. The reflected shortwave radiation in Hyttiälä as a function of global shortwave radiation, coloured by the snow depth. Unfilled markers have snow depth data missing. a) Time up to end of January 2020, before the thinning, b) after the thinning, starting from April 2020. Open circles have no snow depth value available, and are typically snow-free points. The albedo values are from a linear model in which the p-values of all predictors are below 1×10^{-6} . Albedo values are presented in Figs Fig. A7 and ??.

scatter due to the summertime variation in the peatland albedo (Fig. A12). We constructed a model that included the effect of global radiation, snow cover, and DOY as a proxy for the peatland phenology development in the summertime to explain the difference in net SW radiation between the sites (Section A1 for details). This model explained the variation in the daily net SW radiation differences satisfactorily, especially at the northern Halssikangas-Halssiaapa site pair (Fig. A13, R^2 of the model 0.871 for the Hyttiälä-Siikaneva pair and 0.972 for the Halssikangas-Halssiaapa pair). The model was used to gapfill the SW balance difference for calculation of cumulative sums and annual values (Fig. A2).

The net SW radiation at forest sites typically followed a bell-shaped annual cycle (Fig. A9). In contrast, the peatland sites had a clearly lower net shortwave radiation in spring, when the incoming radiation was rapidly increasing and the peatland albedo was substantially higher due to the snow cover. Typically, the forest site started absorbing more shortwave radiation than the peatland earlier in the spring in the southern pair than in the northern pair (Fig. 9 a), due to the earlier increase of global radiation. As the spring progressed, this north-south difference was reversed as snow in Siikaneva started melting (Fig. 9 a). This results in the difference in the albedos between the peatland and forest site persisting longer into spring (Fig. 4 c and f) in the north, at a time when the global radiation is already high and rapidly increasing (Fig. A10). Finally, in the summer, the difference between the forest and the peatland was again larger in the southern pair, mainly due to the higher albedo of




Figure 8. The difference in the net SW radiation between (a) Hyytiälä and Siikaneva and (b) Halssikangas site and Halssiaapa as a function of the mean global radiation over the sites. Daily averages, colour of the markers by the snow depth at the peatland site.

315 Siikaneva as compared to Halssiaapa (Fig. 9 a). Over the whole year, the difference in the absorbed radiation between the forest and the peatland was typically greater in the northern site pair (Fig. 9 b). This difference mainly arises in the springtime, due to the longer snow-cover duration in the north. On the annual scale, the clear majority of the forest-peatland net shortwave radiation difference at the northern pair was caused by snow in springtime overlapping with relatively high incoming radiation level. In addition to this, there was a minor contribution from the peatland having a higher summertime albedo than the forest.

320 In contrast, in the southern pair, roughly half of the average annual difference comes from the summertime. This is caused by the shorter snow-cover duration, and longer snow-free period. In addition, the difference in snow-free albedos at the southern pair is greater than at the northern pair. The average annual absorbed shortwave radiation, calculated as a sum of median daily values, is $3.15 GJ m^{-2}$ for Hyytiälä, $2.68 GJ m^{-2}$ for Siikaneva, $2.53 GJ m^{-2}$ for Halssikangas and $2.03 GJ m^{-2}$ for Halssiaapa (Fig. A9). Note that these values are not based on gapfilled data, and can be skewed by uneven missing data. They are thus not

325 directly comparable to the difference values reported next.

The difference in the net SW radiation between (a) Hyytiälä and Siikaneva and (b) Halssikangas site and Halssiaapa as a function of the mean global radiation between the sites. Daily averages, colour of the markers by the snow depth at the peatland site.

3.6 Annual variation in net shortwave radiation

(a) The typical annual cycle in the net SW radiation differences between Hyytiälä and Siikaneva and Halssikangas site and Halssiaapa. Positive values mean that the forest absorbs more shortwave radiation than the peatland. For each week, the average net SW difference was calculated as the median of all daily values belonging to that week over the years. Interannual variation shown in Fig. A9. (b) same as a, but cumulative from the start of the year, and calculated for each day of year.




Figure 9. (a) The typical annual cycle in the net SW radiation differences between Hyytiälä and Siikaneva in dashed blue, and Halssikangas and Halssiaapa in red. Positive values mean that the forest absorbs more shortwave radiation than the peatland. For each week, the average net SW difference was calculated as the median of all daily values belonging to that week over the years. Interannual variation shown in Fig. A9. (b) same as a, but cumulative from the start of the year, and calculated for each day of year. Thin lines are individual years. The bold lines are cumulative values calculated of the median years. Note that the median annual difference presented in this way differs from the median calculated from the cumulative individual years. All data shown are gapfilled.

330 On a daily time scale, the difference in the net SW radiation between the forest and peatland site depended strongly on both the global radiation and the snow cover at the peatland site, indicative of the albedo difference (Eq. (2), Fig. 8). The presence of snow, combined with the global radiation, explains the difference in net SW radiation well, without accounting for e.g. the melting state and grain size mentioned earlier (Fig. A11). In contrast, the snow-free points show a much greater scatter due to the summertime variation in the peatland albedo (Fig. A12). We constructed a model that included the effect of global radiation, snow cover, and DOY as a proxy for the peatland phenology development in the summertime to explain the difference in the annual cumulative difference in the net SW radiation between the sites (Section A1 for details). This model explained the variation in the net SW radiation differences satisfactorily, especially forest and the peatland site was on average higher at the northern Halssikangas-Halssiaapa site pair (Fig. A13). The model was used to gapfill the SW balance difference for calculation of annual values (Fig. A2).




Figure 10. (a) The annual average difference in the net SW radiation between Hyytiälä and Siikaneva and Halssikangas site and Halssiaapa as a function of the number of snow-covered days at the peatland site. (b) same as a, but net SW differences only calculated over the springtime (up to DOY 130 in Hyytiälä-Siikaneva and 150 in Halssikangas-Halssiaapa). Colour of the markers by the average snow depth in the peatland site over the snow-covered period.

340 The annual average difference in the net SW radiation between the forest and the peatland site mean 0.47 GJ m^{-2} , ranging from 0.37 GJ m^{-2} to 0.61 GJ m^{-2}) than at the southern site pair (mean 0.37 GJ m^{-2} , ranging from 0.2 GJ m^{-2} to 0.54 GJ m^{-2}). The annual difference depended on the snow melt day, with later snow melt corresponding to larger difference (Fig. 10 a). The dependence was even clearer when only considering the springtime net SW difference (Fig. 10 b). In addition, when considering only the springtime, both sites fell closer to similar linear behaviour (Fig. 10 b). This indicates that a) there are interannual differences in the net SW differences outside the spring period and b) that outside the spring period, Siikaneva, on average, absorbs more SW radiation than Halssiaapa, as compared to their respective forest sites. This is also supported by Fig. 9. The earliest snow melt day in Fig. 10 in the Halssikangas-Halssiaapa pair is from the spring 2021, when the snow melted from the peatland measurement location around two weeks earlier than from the forest. This indicates a spatially inhomogeneous snow cover, which is not seen in the Halssiaapa snow-depth measurements. This can explain why that spring differs from the pattern observed during other years. Both sites show a clear correlation between the springtime net SW difference and the snow melt day, and the dependence seems rather linear in the years studied (Fig. 10 b).

345
350

We constructed a linear model of the annual difference in the net shortwave radiation as a function of the peatland snow cover duration in the spring time, the mean peatland albedo in summer, and mean global radiation in the summer. The first two variables explain the variation in the difference of the net shortwave radiation within the site pairs ($R^2 = 0.803$, Fig.

355 10). Additionally, the southern pair receives more solar radiation in general, explaining the difference between the sites. For
the springtime difference, the snowmelt date as a sole predictor gives an R^2 of 0.793. We also attempted to separate
the effect of other variables on the annual difference in the net shortwave radiation. However, many of the potential other
variables, such as global radiation and fraction of diffuse radiation, are intercorrelated. In addition, the total number of years,
360 especially for a separate site pair, is relatively low compared to the total number of potential variables. This creates problems
of multicollinearity and under-determination in a linear model. As a result, our choice of the snow cover duration, summertime
albedo, and summertime global radiation, is ultimately based on physical intuition.

Finally, Northern Europe is unusually warm for its latitude, and correspondingly receives little ~~sunlight~~ solar radiation as compared to other parts of the boreal biome with a similar temperature regime. This is also seen in snow-cover duration: similar snow-cover durations are typically seen in more southern locations (Brown and Mote, 2009; Bormann et al., 2018). It is conceivable that the difference between open peatlands and forests in these more southern locations could be larger than the ones observed here due to increased solar radiation.

4 Conclusions

The albedo at each of the sites was most affected by snow cover, with clearly higher albedo observed during the winter as compared to summer. The difference between the snow-free and snow-covered values was higher for the peatland sites than for the forest sites, as expected. At both forest sites, the snow-free albedo was rather constant at 0.11 over the snow-free period. In contrast, the peatland sites showed a clear seasonal cycle in the snow-free albedo, increasing from around 0.12 for the southern site and 0.08 for the northern site, right after snow melt, up to 0.19 and 0.16, by late summer, respectively. This seasonal cycle was absent from PAR albedo, indicating that the seasonal cycle is mainly due to changes in reflectance in the near infrared region. In addition, the albedo at the open peatland sites was more sensitive to changes in the diffuse fraction of incoming radiation than that at the forest sites.

When comparing the annual shortwave radiation budgets, the forest site absorbed more shortwave radiation than the peatland site at both the northern and the southern site pair. For the northern site pair, this difference was predominantly caused by higher springtime albedo at the peatland, while at the southern site, the higher summertime albedo also made a substantial contribution. In total, the difference between the forest and the peatland site was larger at the northern pair. This was caused by the snow cover persisting longer into the spring, when incoming radiation is rapidly increasing. The total surface energy balance depends on other properties of the ecosystem, such as its ability to dissipate heat back to the atmosphere. However, understanding of the SW radiation balance is critical when assessing energy balance changes due to evolving snow cover.

The interannual variation in the differences of absorbed shortwave radiation between the forest and peatland sites are well explained by the date of snow melt, with earlier snow melt corresponding to smaller differences. The relationship is clearer still when only considering the SW radiation balance over the spring period. Any changes in snow-cover duration will therefore affect the surface energy balances of forest and peatland sites. As the study sites are located in Finland, with an unusually mild




Figure A1. Measured and gapfilled daily snow depth averages from Siikaneva and Halssiaapa.

climate for its latitude, it is possible that the effect of land use on the shortwave radiation budget is larger than observed here at comparable, more southern sites.

Data availability. The data for Siikaneva and Hyytiälä was downloaded from SmartSMEAR (<https://smear.avaa.csc.fi/>, Aalto et al., 2023a; 390 Alekseychik et al., 2023), and for Halssiaapa and Halssikangas from <https://litdb.fmi.fi> and <https://en.ilmatieteenlaitos.fi/download-observations>. The data are available under Creative Commons 4.0 Attribution (CC BY 4.0) and Creative Commons Attribution-NonCommercial 4.0 (CC BY-NC 4.0) licenses, respectively.

Appendix A: Appendix

~~The daily albedo at Halssiaapa as a function of snow depth at Halssiaapa. Colouring by snow depth at the forest site: the values 395 with zero snow depth but high albedo are explained by spatially inhomogeneous snow cover, when snow has already melted from the Halssiaapa measurement location, but is still present in the vicinity.~~

~~Albedo in Hyytiälä as a function of global radiation, coloured by snow depth. a) Time up to end of January 2020, before the thinning, b) after the thinning, starting from April 2020.~~

Table A1. Description of the measurements.

	Siikaneva	Hyttiälä	Halssiaapa	Halssikangas
Global radiation	Kipp & Zonen CNR4	Middleton SK08 ^a , EQ08 ^b	Kipp & Zonen CMA11	Kipp & Zonen CM11
Height (m)	3	18 ^a , 35 ^b	2	45
Site code	SII1	HYY	SUO0006	MET0002
Reflected shortwave	Kipp & Zonen CNR4	Middleton SK08	Kipp & Zonen CMA11	Kipp & Zonen CM11
Height (m)	3	125	2	45
Site code	SII1	HYY	SUO0006	MET0002
PAR	Li-Cor Li-190R	Li-Cor Li-190SZ	Kipp & Zonen PAR Lite	Li-Cor LI190SZ
Height (m)	3	18 ^c , 35 ^d	1.8	45
Site code	SII1	HYY	SUO0009	MET0002
Reflected PAR	Li-Cor Li-190R	Li-Cor Li-190R ^a , Li-190SZ ^b	Kipp & Zonen PAR Lite	Li-Cor LI190SZ
Height (m)	3	67	1.8	45
Site code	SII1	HYY	SUO0010	MET0002
Diffuse		Delta-T BF3/BF5 (PAR)		Kipp & Zonen CM11 (shortwave)
Height (m)		18 ^c , 35 ^d		20
Site code		HYY		Sodankylä Tähtelä
Snow depth	Campbell ultrasonic	Jenoptik SHM30	Campbell SR50	Campbell SR50
Site code	SII1	HYY	SUO0003	MET0002
<u>Water table depth (cm)</u>	<u>Druck PDCR1830^e</u> <u>Campbell CS451^f</u>		<u>Trafag</u>	
<u>Site code</u>	<u>SII1</u>		<u>SUO0010</u>	
<u>Soil water content m³ m⁻³</u>	<u>Delta PR2</u>			
<u>Site code</u>	<u>SII1</u>			

^auntil 9/2017^bfrom 9/2017^cuntil 2/2017^dfrom 2/2017^euntil 2016^ffrom 2017




Figure A2. Measured and gapfilled daily net SW radiation differences. The majority of the data used for calculating annual averages is measured (86% for Hyytiälä-Siikaneva, 78% for Halssikangas-Halssiaapa).

A1 Modelling the net SW radiation difference between the peatland and the forest sites

400 We found that the difference in the net SW radiation between the forest and the peatland site mainly depends on the average global radiation between the sites and the snow cover at the peatland site, in line with Eq. (2) and snow cover being the main determinant for the albedo difference between the sites (Fig. 8). To model the net SW radiation difference between the sites, we split the data to snow cover below and above 30 cm. We fitted a robust regression model with daily mean global radiation and whether the snow depth exceeded 30 cm as predictors, with a different slope for the above 30 cm snow points (Fig. A14).

405 Points with an albedo over 0.3 at the peatland site, but no snow recorded at the location, were excluded from the model (19 days for the northern pair, and 1 day for the southern pair). With this convention, the slope essentially describes the albedo difference between the forest and the peatland site (Eq. (2)). ~~Wintertime points are generally better described by this model~~ The model produces generally more accurate predictions in winter than in summertime. There remains a dependence of the net SW radiation difference on the day of year, caused especially by the seasonal change in the snow-free albedo at the peatland

410 sites (Figs. A12 and A15). This is caused by the increase of the peatland albedo towards late summer shown in Fig. 4 d. This increase may be due to the peatland vegetation phenology observed by Linkosalmi et al. (2016). Another possibility is that the peatland becomes drier over the summer, affecting albedo. The dryness of the site, measured by the water table depth (WTD), also shows a connection to albedo (Fig. A16). However, over the summer (snow depth below 5 cm and DOY up to 250), DOY shows a higher correlation with the residuals as compared to the WTD for both site pairs (Fig. A17).




Figure A3. The correspondence of the global radiation between each site pair: a) Siikaneva and Hyytiälä, and b) Halssiaapa and the Halssikangas. Peatland site on the y-axis, with the forest site on the x-axis. For each pair both 30-minute average and daily median radiation data are plotted. In addition, a total least squares (TLS) linear regression to the 30 min values is plotted, and the squared Pearson correlation coefficient (r^2 , coefficient of determination) is given.

415 We still added to the model the day of year to describe the summertime change in the peatland phenology. The snow depth used [for the gapfilling as an explanatory variable in the model](#) was also interpolated linearly to fill some missing values (Fig. A1). When the global radiation measurement was available at both sites, the mean of the two was used: if it was only available at one site, that value was used. After gapfilling, the majority of the data was still measured (Fig. A2).

420 *Author contributions.* OP and MK conceived and designed the study. OP conducted the data analysis and visualisation, and wrote the manuscript. MA and PK took part in the data collection. All authors contributed to the interpretation of the results and read and commented on the manuscript.

Competing interests. The authors declare that they have no conflict of interest.




Figure A4. The effect of diffuse fraction and [solar](#) zenith angle on the reflected radiation at the forest sites. [The ratio of the reflected to incoming shortwave radiation defines the albedo.](#)

Acknowledgements. We acknowledge the following projects: ACCC Flagship funded by the Academy of Finland grant number 337549 (UH) and 337552 (FMI), Academy professorship funded by the Academy of Finland (grant no. 302958), Academy of Finland projects no. 425 1325656, 311932, 334792, 316114, 325647, 325681, 347782, the Strategic Research Council (SRC) at the Academy of Finland (#352431), “Quantifying carbon sink, CarbonSink+ and their interaction with air quality” INAR project funded by Jane and Aatos Erkkö Foundation, “Gigacity” project funded by Wihuri foundation, European Research Council (ERC) project ATM-GTP Contract No. 742206, and European Union via Non-CO₂ Forcers and their Climate, Weather, Air Quality and Health Impacts (FOCI). University of Helsinki support via ACTRIS-HY is acknowledged. Support of the technical and scientific staff in Hyytiälä are acknowledged. We thank Anna Kontu for [assistance with the Sodankylä data.](#)




Figure A5. The peatland albedo as a function of forest daily albedo and peatland snow depth in at the peatland sites, (a) the southern Siikaneva and (b) the northern Halssiaapa. The points are coloured by the day of year (DOY). The springtime values in Halssiaapa around DOY 100, with a high albedo but a low snow depth, are associated with the times when the peatland snow depth measurement has reached zero, but snow is still detected at the forest site pair.




Figure A6. Albedo divided into white-sky and clear sky conditions. For a) Hyytiälä and b) Siikaneva, clear sky is defined as the fraction of diffuse PAR in Hyytiälä under 0.4, and white-sky as the ratio over 0.85. In c) Halssikangas and d) Halssiaapa the limits are 0.35 and 0.95, but for the diffuse shortwave radiation.

PAR albedo at the different sites, with focus on the snow-free period. The Halssikangas reflected PAR sensor appears to be mounted at a slight angle, registering also part of incoming radiation. In the Hyytiälä PAR albedo there are clear interannual differences, part of these are due to sensor replacement in late 2019, see Fig. A8




Figure A7. Albedo in Hyytiälä as a function of snow depth, coloured by global radiation. a) Time up to end of January 2020, before the thinning, b) after the thinning, starting from April 2020.




Figure A8. Clear sky PAR albedo in Hyytiälä, coloured by year. The jump in the PAR albedo from 2017 to 2018 is caused by replacement of the sensor for reflected PAR in 9/2017.




Figure A9. Typical annual cycles of the net SW radiation in the a) Hyytiälä, b) Siikaneva, c) difference of Hyytiälä and Siikaneva, d) Halssikangas, e) Halssiaapa and f) difference of Halssikangas and Halssiaapa. The statistics are calculated for all of the daily values belonging to each week number, over all of the years. The data are not gap-filled, and therefore the data coverage for each panel differs. The Hyytiälä-Siikaneva data set includes eight years, the Halssikangas-Halssiaapa data set includes 12 years. The average annual absorbed shortwave radiation, calculated as a sum of daily medians, is 3.15 GJ m^{-2} for Hyytiälä, 2.68 GJ m^{-2} for Siikaneva, 2.53 GJ m^{-2} for Halssikangas and 2.03 GJ m^{-2} for Halssiaapa. The average annual difference between Hyytiälä and Siikaneva, calculated in the same way, is 0.35 GJ m^{-2} , and between Halssikangas and Halssiaapa 0.45 GJ m^{-2} . Due to the differing data coverages, these numbers are not directly comparable.




Figure A10. The albedo difference (peatland - forest) as a function of the incoming radiation. The product of ~~these two parameters~~ the global radiation and the difference in albedo determines the difference in the net SW radiation between the sites (Eq. (2)).




Figure A11. The difference in the net SW radiation between (a) Hyytiälä and Siikaneva and (b) Halssikangas site and Halssiaapa as a function of the mean global radiation between-over the sites, when the snow depth at the peatland site is at least 30 cm. Daily averages, colour of the markers by the snow depth at the peatland site.




Figure A12. The difference in the net SW radiation between (a) Hyytiälä and Siikaneva and (b) Halssikangas site and Halssiaapa as a function of the mean global radiation between-over the sites. Daily averages, colour of the markers by DOY. Only days with snow depth at the peatland site of less than 30 cm are included.




Figure A13. The ~~fitted-predicted~~ vs. observed net SW difference values difference in (a) Hyytiälä and Siikaneva and (b) Halssikangas site and Halssiaapa. ~~Fitted-Predicted~~ points based on linear model with global radiation, snow cover and DOY as predictors, 1:1 line drawn as a reference




Figure A14. The difference in the net SW radiation between the Halssikangas and Halssiaapa sites as a function of the mean global radiation ~~between~~over the sites. Daily averages, colour of the markers by the snow depth in Halssiaapa. Fitted points based on linear model with different slopes for snow cover categories less and over 30 cm.




Figure A15. The residuals from the model in Fig. A14 plotted against day of year (DOY). High residuals are seen around snow melt, when the snow at the peatland measurement point has melted, but scattered snow cover still remains. In addition, there is a clear pattern with respect to DOY, both before, but especially after snowmelt. This is possibly due to the development of vegetation phenology, also visible in the summertime albedo in Fig. 4 d




Figure A16. The dependence of peatland summertime albedo on DOY, water table depth, and soil moisture.




Figure A17. The residuals from the model in Fig. A14 plotted against day of year and water table depth, for daily snow depth below 5 cm and DOY up to 250.




Figure A18. The difference in the net SW radiation between (a) Hyytiälä and Siikaneva and (b) Halssikangas site and Halssiaapa as a function of the mean global radiation ~~between~~^{over} the sites. Daily averages, colour of the markers by the snow depth in at the peatland site. Fitted points based on linear model with global radiation, snow cover and DOY as predictors.

References

- Aalto, J., Aalto, P., Keronen, P., Kolari, P., Rantala, P., Taipale, R., Kajos, M., Patokoski, J., Rinne, J., Ruuskanen, T., Leskinen, M., Laakso, H., Levula, J., Pohja, T., Siivola, E., Kulmala, M., and Ylivinkka, I.: SMEAR II Hyytiälä forest meteorology, greenhouse gases, air quality and soil, <https://doi.org/10.23729/23DD00B2-B9D7-467A-9CEE-B4A122486039>, artwork Size: 1806590642 Pages: 1806590642, 2023a.
- 435 Aalto, J., Anttila, V., Kolari, P., Korpela, I., Isotalo, A., Levula, J., Schiestl-Aalto, P., and Bäck, J.: Hyytiälä SMEAR II forest year 2020 thinning tree and carbon inventory data, <https://doi.org/10.5281/zenodo.8283057>, 2023b.
- Alekseychik, P., Kolari, P., Rinne, J., Haapanala, S., Laakso, H., Taipale, R., Matilainen, T., Salminen, T., Levula, J., and Eeva-Stiina Tuittila: SMEAR II Siikaneva 1 wetland meteorology and soil, <https://doi.org/10.23729/08D89ADA-D152-4C8B-8DB4-AE8A8F17F825>, 2023.
- Alekseychik, P. K., Korrensalo, A., Mammarella, I., Vesala, T., and Tuittila, E.-S.: Relationship between aerodynamic roughness
440 length and bulk sedge leaf area index in a mixed-species boreal mire complex, *Geophysical Research Letters*, 44, 5836–5843, <https://doi.org/10.1002/2017GL073884>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2017GL073884>, 2017.
- Anttila, K., Manninen, T., Jääskeläinen, E., Riihelä, A., and Lahtinen, P.: The Role of Climate and Land Use in the Changes in Surface Albedo Prior to Snow Melt and the Timing of Melt Season of Seasonal Snow in Northern Land Areas of 40°N–80°N during 1982–2015, *Remote Sensing*, 10, 1619, <https://doi.org/10.3390/rs10101619>, 2018.
- 445 Aurela, M., Laurila, T., and Tuovinen, J.-P.: Seasonal CO₂ balances of a subarctic mire, *Journal of Geophysical Research: Atmospheres*, 106, 1623–1637, <https://doi.org/10.1029/2000JD900481>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2000JD900481>, 2001.
- Aurela, M., Laurila, T., and Tuovinen, J.-P.: Annual CO₂ balance of a subarctic fen in northern Europe: Importance of the wintertime efflux, *Journal of Geophysical Research: Atmospheres*, 107, <https://doi.org/10.1029/2002JD002055>, 2002.
- Aurela, M., Riutta, T., Laurila, T., Tuovinen, J.-P., Vesala, T., Tuittila, E.-S., Rinne, J., Haapanala, S., and Laine, J.: CO₂ exchange of a sedge
450 fen in southern Finland—the impact of a drought period, *Tellus B*, 59, 826, <https://doi.org/10.1111/j.1600-0889.2007.00309.x>, 2007.
- Aurela, M., Lohila, A., Tuovinen, J.-P., Hatakka, J., Penttilä, T., and Laurila, T.: Carbon dioxide and energy flux measurements in four northern-boreal ecosystems at Pallas, *Boreal Environment Research*, 20, 455–473, 2015.
- Baker, D. G., Ruschy, D. L., and Wall, D. B.: The Albedo Decay of Prairie Snows, *Journal of Applied Meteorology and Climatology*, 29, 179–187, [https://doi.org/10.1175/1520-0450\(1990\)029<0179:TADOPS>2.0.CO;2](https://doi.org/10.1175/1520-0450(1990)029<0179:TADOPS>2.0.CO;2), publisher: American Meteorological Society Section:
455 *Journal of Applied Meteorology and Climatology*, 1990.
- Bala, G., Caldeira, K., Wickett, M., Phillips, T. J., Lobell, D. B., Delire, C., and Mirin, A.: Combined climate and carbon-cycle effects of large-scale deforestation, *Proceedings of the National Academy of Sciences*, 104, 6550–6555, <https://doi.org/10.1073/pnas.0608998104>, publisher: Proceedings of the National Academy of Sciences, 2007.
- Baldocchi, D., Kelliher, F. M., Black, T. A., and Jarvis, P.: Climate and vegetation controls on boreal zone energy exchange, *Global Change
460 Biology*, 6, 69–83, <https://doi.org/10.1046/j.1365-2486.2000.06014.x>, 2000.
- Betts, A. K. and Ball, J. H.: Albedo over the boreal forest, *Journal of Geophysical Research: Atmospheres*, 102, 28 901–28 909, <https://doi.org/10.1029/96JD03876>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/96JD03876>, 1997.
- Betts, R. A.: Offset of the potential carbon sink from boreal forestation by decreases in surface albedo, *Nature*, 408, 187–190, <https://doi.org/10.1038/35041545>, number: 6809 Publisher: Nature Publishing Group, 2000.
- 465 Bormann, K. J., Brown, R. D., Derksen, C., and Painter, T. H.: Estimating snow-cover trends from space, *Nature Climate Change*, 8, 924–928, <https://doi.org/10.1038/s41558-018-0318-3>, number: 11 Publisher: Nature Publishing Group, 2018.

- Bright, R. M., Eisner, S., Lund, M. T., Majasalmi, T., Myhre, G., and Astrup, R.: Inferring Surface Albedo Prediction Error Linked to Forest Structure at High Latitudes, *Journal of Geophysical Research: Atmospheres*, 123, 4910–4925, <https://doi.org/10.1029/2018JD028293>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2018JD028293>, 2018.
- 470 Brown, R. D. and Mote, P. W.: The Response of Northern Hemisphere Snow Cover to a Changing Climate, *Journal of Climate*, 22, 2124–2145, <https://doi.org/10.1175/2008JCLI2665.1>, publisher: American Meteorological Society Section: *Journal of Climate*, 2009.
- Charney, J., Stone, P. H., and Quirk, W. J.: Drought in the Sahara: A Biogeophysical Feedback Mechanism, *Science*, 187, 434–435, <https://doi.org/10.1126/science.187.4175.434>, publisher: American Association for the Advancement of Science, 1975.
- Cohen, J. and Rind, D.: The Effect of Snow Cover on the Climate, *Journal of Climate*, 4, 689–706, <https://www.jstor.org/stable/26196419>,
 475 publisher: American Meteorological Society, 1991.
- Courel, M. F., Kandel, R. S., and Rasool, S. I.: Surface albedo and the Sahel drought, *Nature*, 307, 528–531, <https://doi.org/10.1038/307528a0>, number: 5951 Publisher: Nature Publishing Group, 1984.
- Curry, J. A., Schramm, J. L., and Ebert, E. E.: Sea Ice-Albedo Climate Feedback Mechanism, *Journal of Climate*, 8, 240–247, [https://doi.org/10.1175/1520-0442\(1995\)008<0240:SIACFM>2.0.CO;2](https://doi.org/10.1175/1520-0442(1995)008<0240:SIACFM>2.0.CO;2), publisher: American Meteorological Society Section: *Journal of*
 480 *Climate*, 1995.
- Dang, C., Brandt, R. E., and Warren, S. G.: Parameterizations for narrowband and broadband albedo of pure snow and snow containing mineral dust and black carbon, *Journal of Geophysical Research: Atmospheres*, 120, 5446–5468, <https://doi.org/10.1002/2014JD022646>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2014JD022646>, 2015.
- Derksen, C. and Brown, R.: Spring snow cover extent reductions in the 2008–2012 period exceeding climate model projections, *Geophysical*
 485 *Research Letters*, 39, <https://doi.org/10.1029/2012GL053387>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2012GL053387>, 2012.
- Déry, S. J. and Brown, R. D.: Recent Northern Hemisphere snow cover extent trends and implications for the snow-albedo feedback, *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2007GL031474>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2007GL031474>, 2007.
- 490 Essery, R.: Large-scale simulations of snow albedo masking by forests, *Geophysical Research Letters*, 40, 5521–5525, <https://doi.org/10.1002/grl.51008>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/grl.51008>, 2013.
- Eugster, W., Rouse, W. R., Pielke Sr, R. A., Mcfadden, J. P., Baldocchi, D. D., Kittel, T. G. F., Chapin, F. S., Liston, G. E., Vidale, P. L., Vaganov, E., and Chambers, S.: Land-atmosphere energy exchange in Arctic tundra and boreal forest: available data and feedbacks to climate, *Global Change Biology*, 6, 84–115, <https://doi.org/10.1046/j.1365-2486.2000.06015.x>, 2000.
- 495 Ezhova, E., Ylivinkka, I., Kuusk, J., Komsaare, K., Vana, M., Krasnova, A., Noe, S., Arshinov, M., Belan, B., Park, S.-B., Lavrič, J. V., Heimann, M., Petäjä, T., Vesala, T., Mammarella, I., Kolari, P., Bäck, J., Rannik, Ü., Kerminen, V.-M., and Kulmala, M.: Direct effect of aerosols on solar radiation and gross primary production in boreal and hemiboreal forests, *Atmospheric Chemistry and Physics*, 18, 17 863–17 881, <https://doi.org/10.5194/acp-18-17863-2018>, 2018.
- Gao, Y., Markkanen, T., Backman, L., Henttonen, H. M., Pietikäinen, J.-P., Mäkelä, H. M., and Laaksonen, A.: Biogeophysical impacts of peatland forestation on regional climate changes in Finland, *Biogeosciences*, 11, 7251–7267, <https://doi.org/10.5194/bg-11-7251-2014>,
 500 publisher: Copernicus GmbH, 2014.
- Gardner, A. S. and Sharp, M. J.: A review of snow and ice albedo and the development of a new physically based broadband albedo parameterization, *Journal of Geophysical Research: Earth Surface*, 115, <https://doi.org/10.1029/2009JF001444>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2009JF001444>, 2010.

- 505 Goymer, P.: A trillion trees, *Nature Ecology & Evolution*, 2, 208–209, <https://doi.org/10.1038/s41559-018-0464-z>, number: 2 Publisher: Nature Publishing Group, 2018.
- Hari, P. and Kulmala, M.: Station for Measuring Ecosystem–Atmosphere Relations (SMEAR II), *Boreal Environment Research*, 10, 315–322, 2005.
- Henderson-Sellers, A. and Wilson, M. F.: Surface albedo data for climatic modeling, *Reviews of Geophysics*, 21, 1743–1778, <https://doi.org/10.1029/RG021i008p01743>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/RG021i008p01743>, 1983.
- 510 Hovi, A., Lindberg, E., Lang, M., Arumäe, T., Peuhkurinen, J., Sirparanta, S., Pyankov, S., and Rautiainen, M.: Seasonal dynamics of albedo across European boreal forests: Analysis of MODIS albedo and structural metrics from airborne LiDAR, *Remote Sensing of Environment*, 224, 365–381, <https://doi.org/10.1016/j.rse.2019.02.001>, 2019.
- Intergovernmental Panel On Climate Change (IPCC): *The Ocean and Cryosphere in a Changing Climate: Special Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 1 edn., ISBN 978-1-00-915796-4 978-1-00-915797-1, <https://doi.org/10.1017/9781009157964>, 2022.
- 515 Kolari, P., Aalto, J., Levula, J., Kulmala, L., Ilvesniemi, H., and Pumpanen, J.: Hyytiälä SMEAR II site characteristics, <https://doi.org/10.5281/zenodo.5909681>, 2022.
- Kuusinen, N., Kolari, P., Levula, J., Porcar-Castell, A., Stenberg, P., and Berninger, F.: Seasonal variation in boreal pine forest albedo and effects of canopy snow on forest reflectance, *Agricultural and Forest Meteorology*, 164, 53–60, <https://doi.org/10.1016/j.agrformet.2012.05.009>, 2012.
- 520 Kuusinen, N., Tomppo, E., and Berninger, F.: Linear unmixing of MODIS albedo composites to infer subpixel land cover type albedos, *International Journal of Applied Earth Observation and Geoinformation*, 23, 324–333, <https://doi.org/10.1016/j.jag.2012.10.005>, 2013.
- Launiainen, S., Katul, G. G., Kolari, P., Lindroth, A., Lohila, A., Aurela, M., Varlagin, A., Grelle, A., and Vesala, T.: Do the energy fluxes and surface conductance of boreal coniferous forests in Europe scale with leaf area?, *Global Change Biology*, 22, 4096–4113, <https://doi.org/10.1111/gcb.13497>, 2016.
- 525 Liang, S., Kaicun Wang, Xiaotong Zhang, and Martin Wild: Review on Estimation of Land Surface Radiation and Energy Budgets From Ground Measurement, Remote Sensing and Model Simulations, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 3, <https://doi.org/10.1109/JSTARS.2010.2048556>, 2010.
- 530 Linkosalmi, M., Aurela, M., Tuovinen, J.-P., Peltoniemi, M., Tanis, C. M., Arslan, A. N., Kolari, P., Böttcher, K., Aalto, T., Rainne, J., Hatakka, J., and Laurila, T.: Digital photography for assessing the link between vegetation phenology and CO₂ exchange in two contrasting northern ecosystems, *Geoscientific Instrumentation, Methods and Data Systems*, 5, 417–426, <https://doi.org/10.5194/gi-5-417-2016>, publisher: Copernicus GmbH, 2016.
- Loew, A.: Terrestrial satellite records for climate studies: how long is long enough? A test case for the Sahel, *Theoretical and Applied Climatology*, 115, 427–440, <https://doi.org/10.1007/s00704-013-0880-6>, 2014.
- 535 Lohila, A., Minkinen, K., Laine, J., Savolainen, I., Tuovinen, J.-P., Korhonen, L., Laurila, T., Tietäväinen, H., and Laaksonen, A.: Forestation of boreal peatlands: Impacts of changing albedo and greenhouse gas fluxes on radiative forcing, *Journal of Geophysical Research: Biogeosciences*, 115, <https://doi.org/10.1029/2010JG001327>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2010JG001327>, 2010.
- Lukeš, P., Stenberg, P., and Rautiainen, M.: Relationship between forest density and albedo in the boreal zone, *Ecological Modelling*, 261–262, 74–79, <https://doi.org/10.1016/j.ecolmodel.2013.04.009>, 2013.
- 540 Mammarella, I., Aalto, J., Back, J., Kolari, P., Laakso, H., Levula, J., Matilainen, T., Pihlatie, M., Pumpanen, J., Taipale, R., and Vesala, T.: ETC L2 ARCHIVE, Hyytiälä, 2017-12-31–2023-10-31, <https://hdl.handle.net/11676/cZZrPJD8ZKO-nqyp4QBVWYtC>, 2023.

- Manninen, T., Aalto, T., Markkanen, T., Peltoniemi, M., Böttcher, K., Metsämäki, S., Anttila, K., Pirinen, P., Leppänen, A., and Arslan, A. N.: Monitoring changes in forestry and seasonal snow using surface albedo during 1982–2016 as an indicator, *Biogeosciences*, 16, 223–240, <https://doi.org/10.5194/bg-16-223-2019>, publisher: Copernicus GmbH, 2019.
- Manninen, T., Roujean, J.-L., Hautecoeur, O., Riihelä, A., Lahtinen, P., Jääskeläinen, E., Siljamo, N., Anttila, K., Sukuvaara, T., and Korhonen, L.: Airborne Measurements of Surface Albedo and Leaf Area Index of Snow-Covered Boreal Forest, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD035376, <https://doi.org/10.1029/2021JD035376>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2021JD035376>, 2022.
- Nousu, J.-P., Lafaysse, M., Mazzotti, G., Ala-aho, P., Marttila, H., Cluzet, B., Aurela, M., Lohila, A., Kolari, P., Boone, A., Fructus, M., and Launiainen, S.: Modeling snowpack dynamics and surface energy budget in boreal and subarctic peatlands and forests, *The Cryosphere*, 18, 231–263, <https://doi.org/10.5194/tc-18-231-2024>, publisher: Copernicus GmbH, 2024.
- Pithan, F. and Mauritsen, T.: Arctic amplification dominated by temperature feedbacks in contemporary climate models, *Nature Geoscience*, 7, 181–184, <https://doi.org/10.1038/ngeo2071>, number: 3 Publisher: Nature Publishing Group, 2014.
- Pulliaainen, J., Luoju, K., Derksen, C., Mudryk, L., Lemmetyinen, J., Salminen, M., Ikonen, J., Takala, M., Cohen, J., Smolander, T., and Norberg, J.: Patterns and trends of Northern Hemisphere snow mass from 1980 to 2018, *Nature*, 581, 294–298, <https://doi.org/10.1038/s41586-020-2258-0>, number: 7808 Publisher: Nature Publishing Group, 2020.
- Qu, Y., Liang, S., Liu, Q., He, T., Liu, S., and Li, X.: Mapping Surface Broadband Albedo from Satellite Observations: A Review of Literatures on Algorithms and Products, *Remote Sensing*, 7, 990–1020, <https://doi.org/10.3390/rs70100990>, number: 1 Publisher: Multidisciplinary Digital Publishing Institute, 2015.
- Räisänen, J.: Snow conditions in northern Europe: the dynamics of interannual variability versus projected long-term change, *The Cryosphere*, 15, 1677–1696, <https://doi.org/10.5194/tc-15-1677-2021>, publisher: Copernicus GmbH, 2021.
- Rinne, J., Tuittila, E.-S., Peltola, O., Li, X., Raivonen, M., Alekseychik, P., Haapanala, S., Pihlatie, M., Aurela, M., Mammarella, I., and Vesala, T.: Temporal Variation of Ecosystem Scale Methane Emission From a Boreal Fen in Relation to Temperature, Water Table Position, and Carbon Dioxide Fluxes, *Global Biogeochemical Cycles*, 32, 1087–1106, <https://doi.org/10.1029/2017GB005747>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2017GB005747>, 2018.
- Sagan, C., Toon, O. B., and Pollack, J. B.: Anthropogenic Albedo Changes and the Earth's Climate, *Science*, 206, 1363–1368, <https://doi.org/10.1126/science.206.4425.1363>, publisher: American Association for the Advancement of Science, 1979.
- Scott, C. E., Monks, S. A., Spracklen, D. V., Arnold, S. R., Forster, P. M., Rap, A., Äijälä, M., Artaxo, P., Carslaw, K. S., Chipperfield, M. P., Ehn, M., Gilardoni, S., Heikkinen, L., Kulmala, M., Petäjä, T., Reddington, C. L. S., Rizzo, L. V., Swietlicki, E., Vignati, E., and Wilson, C.: Impact on short-lived climate forcers increases projected warming due to deforestation, *Nature Communications*, 9, 157, <https://doi.org/10.1038/s41467-017-02412-4>, number: 1 Publisher: Nature Publishing Group, 2018.
- Stephens, G. L., O'Brien, D., Webster, P. J., Pilewski, P., Kato, S., and Li, J.-l.: The albedo of Earth, *Reviews of Geophysics*, 53, 141–163, <https://doi.org/10.1002/2014RG000449>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2014RG000449>, 2015.
- Thackeray, C. W., Fletcher, C. G., and Derksen, C.: The influence of canopy snow parameterizations on snow albedo feedback in boreal forest regions, *Journal of Geophysical Research: Atmospheres*, 119, 9810–9821, <https://doi.org/10.1002/2014JD021858>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2014JD021858>, 2014.
- Trenberth, K. E., Fasullo, J. T., and Kiehl, J.: Earth's Global Energy Budget, *Bulletin of the American Meteorological Society*, 90, 311–324, <https://doi.org/10.1175/2008BAMS2634.1>, publisher: American Meteorological Society Section: Bulletin of the American Meteorological Society, 2009.

- Wang, D., Liang, S., He, T., Yu, Y., Schaaf, C., and Wang, Z.: Estimating daily mean land surface albedo from MODIS data, *Journal of Geophysical Research: Atmospheres*, 120, 4825–4841, <https://doi.org/10.1002/2015JD023178>, 2015.
- Wang, X., Shi, T., Zhang, X., and Chen, Y.: An Overview of Snow Albedo Sensitivity to Black Carbon Contamination and Snow Grain Properties Based on Experimental Datasets Across the Northern Hemisphere, *Current Pollution Reports*, 6, 368–379, <https://doi.org/10.1007/s40726-020-00157-1>, 2020.
- 585 Warren, S. G. and Wiscombe, W. J.: A Model for the Spectral Albedo of Snow. II: Snow Containing Atmospheric Aerosols, *Journal of the Atmospheric Sciences*, 37, 2734–2745, [https://doi.org/10.1175/1520-0469\(1980\)037<2734:AMFTSA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<2734:AMFTSA>2.0.CO;2), publisher: American Meteorological Society Section: *Journal of the Atmospheric Sciences*, 1980.
- Wiscombe, W. J. and Warren, S. G.: A Model for the Spectral Albedo of Snow. I: Pure Snow, *Journal of the Atmospheric Sciences*, 37, [2712–2733](https://doi.org/10.1175/1520-0469(1980)037<2712:AMFTSA>2.0.CO;2), [https://doi.org/10.1175/1520-0469\(1980\)037<2712:AMFTSA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<2712:AMFTSA>2.0.CO;2), publisher: American Meteorological Society Section: *Journal of the Atmospheric Sciences*, 1980.
- 590 Yan, H., Wang, S., Dai, J., Wang, J., Chen, J., and Shugart, H. H.: Forest Greening Increases Land Surface Albedo During the Main Growing Period Between 2002 and 2019 in China, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033582, <https://doi.org/10.1029/2020JD033582>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2020JD033582>, 2021.
- 595 Yang, F., Mitchell, K., Hou, Y.-T., Dai, Y., Zeng, X., Wang, Z., and Liang, X.-Z.: Dependence of Land Surface Albedo on Solar Zenith Angle: Observations and Model Parameterization, *Journal of Applied Meteorology and Climatology*, 47, 2963–2982, <https://doi.org/10.1175/2008JAMC1843.1>, publisher: American Meteorological Society Section: *Journal of Applied Meteorology and Climatology*, 2008.
- Zeng, N. and Yoon, J.: Expansion of the world’s deserts due to vegetation-albedo feedback under global warming, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2009GL039699>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2009GL039699>, 2009.
- 600