

Always on My Mind: Indications of Post-Traumatic Stress Disorder Among Those Affected by the 2021 Flood Event in the Ahr Valley, Germany

Marie-Luise Zenker¹, Philip Bubeck¹, Annegret H. Thielen¹

5 ¹Institute of Environmental Science and Geography, University of Potsdam, Karl-Liebknecht-Straße 24-25, 14476 Potsdam, Golm, Germany

Correspondence to: Marie-Luise Zenker (marie-luise.zenker@uni-potsdam.com)

Abstract.

The devastating floods that swept through the Ahr Valley in July 2021 left indelible marks on the region's landscape and communities. Beyond the visible damage, experience from other events suggests an increase in mental health issues among those affected. However, there is a lack of data and understanding regarding the impact of flooding on mental health in Germany. Therefore, this study aims to determine how much the flooding in 2021 affected the population's mental well-being. For this purpose, a household-level survey (n = 516) was conducted in the district of Ahrweiler, Rhineland-Palatinate, Germany's most affected region, one year after the flood event, specifically in June/July 2022. The survey employed a short epidemiological screening scale to assess the prevalence of individuals showing indications of post-traumatic stress disorder (PTSD). Using binary logistic regression analyses, we identify risk and protective factors that may have played a role in the development of PTSD to find intervention points for supporting those affected. Our findings indicate significant mental health issues one year after the flood event, with 28.2% of respondents showing indications of PTSD. Furthermore, this study has uncovered essential risk factors for developing indications of PTSD after flooding: female gender, getting seriously injured or sick during the event and feeling left alone to cope with flood impacts. The study emphasizes that severe flooding, such as the 2021 flood, results in new health-related needs that demand attention. As a result, care methods should be adapted to tackle the prevalence and risk factors connected with PTSD in the affected population, e.g., by providing targeted aftercare for individuals who were injured or became sick during the flood event.

25 **Keywords.** mental health, flood, recovery, resilience, trauma, public health.

1 Introduction

Floods have been the most prevalent type of natural hazards globally. They are accountable for 1.65 billion people affected, 104,614 fatalities, and economic losses of US\$651 billion from 2001 to 2019 (CRED and UNDRR, 2021). As the number of people living in flood-prone areas and the value of their assets continue to increase, these numbers are expected to rise in the future (Kron et al., 2019; Jongman et al., 2012). Further, due to climate change, extreme precipitation is expected to increase

(S.I. Seneviratne et al., 2021), leading to an even higher risk of flooding (Tradowsky et al., 2023; Alfieri et al., 2015; Jongman et al., 2014).

Floods can cause significant damage to properties, belongings, and critical infrastructure and disrupt essential services. These events can also result in injuries, infections, and loss of life. Moreover, they can have a lasting impact on the mental health and well-being of those affected (e.g., Thieken et al., 2023, 2016; Tapsell, 2010). Tapsell (2010, p. 407) has stated that “[...] the majority of their problems are just beginning, and people have to cope with the significant aftermath of the flood, not only at a practical level but also socially and psychologically.”

The extent of flood impacts became evident again in July 2021 when severe floods hit Western European countries (Tradowsky et al., 2023). In Germany, the federal states of Rhineland-Palatinate, North Rhine-Westphalia, Bavaria, and Saxony were impacted, with an overall damage of 33 billion Euros and around 190 flood fatalities. The Ahr Valley in Rhineland-Palatinate was the most affected, with 135 flood-related fatalities and one missing person (Thieken et al., 2023a). Many people lost their homes and loved ones, suffered injuries, or were still preoccupied with rebuilding and financial recovery one year after the event. Experiencing such traumatic events can have a long-lasting impact on a person's mental health.

Various studies have explored the effects of flooding on individuals' well-being and quality of life. Others have shown a considerable rise in the number of individuals suffering from mental disorders in the aftermath of flood disasters. These disorders may include posttraumatic stress disorder (PTSD), depression, or anxiety. Among these, PTSD has been the most extensively researched (Galea et al., 2005; Stanke et al., 2012; Fernández et al., 2015; Golitaleb et al., 2022; Keya et al., 2023). PTSD is the most common mental disorder that can develop after a traumatic event (Dreßing and Foerster, 2021). It can occur due to a person's exposure to a potentially traumatic event, which is characterized by the following two conditions: First, the person experienced or witnessed an event that posed a severe threat of death or serious injury for him or herself or another person. Second, the reaction to that event included intense fear, helplessness, or horror (American Psychiatric Association, 1994).

Systematic reviews and meta-analyses have revealed a broad range of PTSD prevalences in the aftermath of disasters. Recent studies conducted by Keya et al. (2023) and Golitaleb et al. (2022) have found that the prevalence of PTSD ranges from 2.6% to 52% and from 18.6% to 40.3% (mean 29.4%), respectively, with most of the studies being conducted in Asia. However, there is also literature available on PTSD after flooding in Europe, with the majority of publications originating from the UK, showing prevalences from 6.6% to 27.9% (Graham et al., 2019; Jermacane et al., 2018; Paranjothy et al., 2011; Mason et al., 2010; Tunstall et al., 2006). Further, a study from France focused on how personality traits and cognitive emotion regulation strategies interact with PTSD after experiencing a flood (Puechlong et al., 2020). Limited research has been conducted in Germany. After the flooding event in 2002, there was a study about mental health in patients who had to be evacuated from a heart disease center in Dresden (Nitschke et al., 2006). Another study was conducted after the flooding event of 2013, showing a PTSD prevalence of 20.4% one-year post-event. They named difficulties or the absence of support (financial as well as psycho-social interventions) in the aftermath of disasters as predictors for negative mental health (Apel and Coenen, 2021). However, these particular flood events in 2002 and 2013 were mainly slow-rising fluvial floods, with return periods of less

65 than 300 and 500 years at the Mulde River (Kreibich et al., 2005; Merz et al., 2014). These floods were less severe compared to the fast-onset floods of 2021, where the discharge of the Ahr River at the Altenahr gauge far exceeded the 1000-year return period (Vorogushyn et al., 2022). However, some regions in the Osterzgebirge experienced similar dynamics during the 2002 floods.

In addition to examining the number of people showing indications of PTSD after a flood event, many studies have assessed factors that might affect the onset of mental health problems. Socio-demographic factors have been studied extensively, indicating that females are at a higher risk of developing such issues than men (Paranjothy et al., 2011; Mason et al., 2010; Tunstall et al., 2006; Norris et al., 2002). Additionally, the economic situation of a person or household was found to be important, as households with low income seem to be more susceptible to suffering from mental health issues (Graham et al., 2019; Lamond et al., 2015; Paranjothy et al., 2011; Mason et al., 2010; Tapsell, 2010; Galea et al., 2007; Tunstall et al., 2006). Other personal factors can also play a role, such as personal traits (Puechlong et al., 2020) or the health status before the event (Paranjothy et al., 2011; Mason et al., 2010; Tunstall et al., 2006). By contrast, there are inconsistent findings for age (Norris et al., 2002; Galea et al., 2007; Stanke et al., 2012; Fernández et al., 2015; Graham et al., 2019). Further, it is important to take into account the characteristics of a flood event, such as the depth of water or the flow velocity (Tunstall et al., 2006; Paranjothy et al., 2011; Lamond et al., 2015; Bubeck and Thielen, 2018). In addition, the circumstances and stress experienced during a flood event, as a result of these flood characteristics, can affect whether a person exhibits signs of mental health issues, for example, if they were injured or required rescue (Galea et al., 2007; Wiseman et al., 2013).

Moreover, the period following a traumatic event can be strongly linked to negative mental health. According to Tunstall et al. (2006), this impairment to health can result from the reconstruction process, caused by several factors such as a disruption of the daily routine, lack of or insufficient insurance coverage, and the unavailability of skilled labor. The most commonly reported cause related to the rebuilding process is significant financial loss (Galea et al., 2007; Paranjothy et al., 2011; Jermacane et al., 2018; Bubeck and Thielen, 2018). Other factors that may contribute to the development of PTSD include displacement (Paranjothy et al., 2011; Galea et al., 2007; Tunstall et al., 2006; Nitschke et al., 2006) or disruption to essential services (Bei et al., 2013; Paranjothy et al., 2011; Tunstall et al., 2006). Also, different coping mechanisms influence mental health (Bubeck and Thielen, 2018; Lamond et al., 2015; Mason et al., 2010). Although there is already a body of relevant literature, relationships between these factors are complex. Studies do not always agree on the significance of influencing factors or the direction of influence (Lamond et al., 2015).

Altogether, there is already a body of relevant literature on the impacts of flooding on mental health. However, relationships between these factors are complex and have not been fully investigated or understood yet; for instance, there is still limited knowledge about the reconstruction time and lasting psychosocial effects and recovery processes following flooding. In particular, there is little research on the mental health impacts of flooding in Germany, which makes it difficult to understand how much the flooding of 2021 affected the population.

Although flooding represents a significant natural hazard in Germany (Munich Re, 2018), the region lacks a comprehensive framework to address disasters' social and psychological impacts, which other countries have already established. For example,

the US recently launched the “Post-Disaster Mental Health Response Act” (U.S. Government, 2021-2022), Australia has the
100 “Australian National Disaster Mental Health and Wellbeing Framework” (National Mental Health Commission, 2023), and
England established its guidelines on dealing with psychosocial and mental health after disasters already in 2009 (DH
Emergency Preparedness Division, 2009). Such a framework could help address the mental health needs of those affected by
flooding. This lack of attention from policymakers and a noticeable gap in (fundamental) research can lead to insufficient
support for individuals struggling with mental health issues. Additionally, such situations can create new demands that require
105 immediate attention. However, we cannot identify the additional support needs without further research.

To address this gap, we conducted a household-level survey one year after the flood event in the Ahrweiler district, Germany's
most affected region. The survey used a short screening scale to determine the prevalence of PTSD and binary regression
analyses to identify factors that influenced the indications of suffering from PTSD. Our aim was twofold: to determine the
share of people in the heavily affected Ahrweiler district who exhibited signs of PTSD following the flood and to identify any
110 risk or protective factors that may have played a role in the development of their PTSD. The study is crucial because it reflects
the outstanding severity of the flood event in Europe, which lets us expect that a significant portion of the population
experienced conditions that match those of PTSD. The study provides information on how much support needs to be provided
in the aftermath of severe floods to help make societies more flood resilient. Further, with our study, we aim to contribute to
developing a guideline for Germany on handling flood impacts on mental health.

115 **2 Case study, sampling, measurement, and methods**

2.1 Case study

Persistent and regionally very pronounced rainfall over Western Europe from the 12th to 19th of July, 2021, led to catastrophic
flood events, particularly in the steep Ahr Valley in Rhineland-Palatinate (Tradowsky et al., 2023; Junghänel et al., 2021).
According to the German Weather Service (Junghänel et al., 2021), local rainfall maxima exceeded 150 mm in 24 hours in
120 some locations, resulting in quickly rising flash floods that caused unparalleled human and economic losses in Germany
(Thieken et al., 2023a, b; Dietze et al., 2022). In the Ahr Valley alone, 135 lives were lost and one person is still missing
(Thieken et al., 2023a). For the whole of Germany, around 190 people were reported dead (Thieken et al., 2023a). The only
flooding in Germany that exceeded this number was the storm surge from 1962 along the North Sea coast (Paprotny et al 2018;
Thieken et al. 2023). One possible reason for the high death toll was that the flood forecasting, warning, and response system
125 (FFWRS) failed in areas affected by quickly rising flash floods, particularly in the Ahr Valley, catching many people by
surprise during the night (Thieken et al., 2023b; Fekete and Sandholz, 2021). At least 770 people were injured, and several
had to be dramatically rescued or were reported missing for several days (Kron et al., 2022). Also, economic losses were
unprecedented, amounting to as much as €33 billion for Germany (Munich Re, 2022). This number by far exceeds direct
economic losses reported for previous large-scale river floods along the Elbe and Danube in 2002 and 2013 (Kron et al., 2022).
130 In the Ahr Valley, critical infrastructure, such as railway tracks and bridges, was heavily impacted. A total of 103 bridges

along the Ahr River were destroyed and need to be reconstructed in the coming years. The Deutsche Bahn (2022) reported that restoring train infrastructures and services will take years.




Figure 1. Overview map of the study area. Own illustration based on material of © BfG 2023, © EuroGeographics and TurkStat 2021, © GeoBasis-DE / BKG 2024.

135

2.2 Sampling

To understand the impact of such an extreme flood event on mental health, we surveyed residents in the heavily affected district of Ahrweiler in Rhineland-Palatinate. Affected households were sampled from the residents who had applied for emergency financial relief (“Soforthilfe”). The federal state of Rhineland-Palatinate initiated this financial relief program to cover residents' urgent necessities in the flood's direct aftermath. Affected households from the district of Ahrweiler could apply for emergency relief until the 10th of September, 2021. In total, more than 17,000 applications were submitted. Around 12,300 applications for immediate relief were granted and more than € 25 million were distributed (Statistical Office RP, 2021a). The main reasons for denying applications were duplicate applications from the same household and incorrectly filed applications from companies and applicants who did not have their principal residence in the district of Ahrweiler. The financial relief was distributed by Ahrweiler's district authorities (Statistical Office RP, 2021).

140

145

With the support of the district administration, every third household that had applied for financial emergency relief was randomly selected and invited to participate in an online survey between June and August 2022, specifically 5250 households in total. They were contacted with a letter signed by the district administrator. To ensure that we also included respondents without internet access and those who did not want to complete the questionnaire online, residents could also request a printed version. Twenty-one people completed the paper version of the questionnaire. About 40 letters did not reach the addressees because they could not be delivered. In total, 516 residents completed the questionnaire, equaling a response rate of 9.9%. For further details on the sampling procedure and the total sample, see (Truedinger et al., 2023).

Altogether, the questionnaire contained 76 questions addressing the following topics: mental health status – especially screening for PTSD – recovery and reconstruction, social vulnerability, and opinions about flood risk management. Out of 516 completed questionnaires, only 411 could be used for screening PTSD due to missing values for the symptom questions. Since these questions are essential for calculating the indications of PTSD, the remaining 105 questionnaires had to be excluded from further analyses. Of the 411 respondents, 54.5% were male, 44.8% female and 0.7% did not answer. This indicates a slight gender bias in the sample population, as Ahrweiler's overall population was evenly split between women (50.5%;) and men (49.5%) in 2022 (Statistical Office RP, 2022). The sample is mainly representative with respect to age (when removing people under 18 from official statistics). It includes participants ranging from their twenties to over 80 years old. The majority of people belonged to the age group of 50-59 years, which also applied to Ahrweiler (Statistical Office RP, 2022). The median net monthly household income in the sample was € 2,600-3,599. Official statistics reported a yearly disposable household net income *per resident* in the district of Ahrweiler of € 24,342 in 2019 (Statistical Office RLP, 2021b). While the numbers are not directly comparable, the income range appears plausible. This survey was approved by the Ethical Committee of the University of Stuttgart in July 2022 (see Truedinger et al., 2023).

2.3 Measurement and Methods

2.3.1 Screening PTSD after the flooding event

To detect indications of post-traumatic stress disorder (PTSD) among affected residents, we used the German short epidemiological screening scale by Siegrist and Maercker (2010). It is based on the scale developed by Breslau et al. (1999), which is closely in line with the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) (Siegrist and Maercker, 2010; Breslau et al., 1999; American Psychiatric Association, 1994).

Symptomatology of PTSD includes trauma re-experience, emotional numbing or avoidance, and hyperarousal (American Psychiatric Association, 1994; Frans et al., 2005). The screening tool employed in this study comprises seven items, five of which relate to symptoms of avoidance and numbing and two to the hyperarousal group (Siegrist and Maercker, 2010; Breslau et al., 1999). The short version of the screening tool was empirically derived from 17 items (symptoms) for PTSD according to the DSM-IV, which best identified respondents with PTSD in a telephone survey in the US (Breslau et al., 1999). Good sensitivity, specificity, test-retest reliability, and internal consistency of the short screening scale have been demonstrated by

various studies (Siegrist and Maercker, 2010; Breslau et al., 1999; Kimerling et al., 2006). A validation of the English version of the short screening tool revealed a high agreement with the extended US version (Breslau et al., 1999).

180 Kimerling et al. (2006) evaluated Breslau's 7-item scale by comparing the self-administered results of 134 participants with the results from clinical interviews of the same patients carried out by psychologists using the Clinician-Administered PTSD Scale (CAPS). They concluded that the short screening scale was suitable for detecting probable PTSD prevalence without a lengthy or specific trauma assessment (Kimerling et al., 2006), which is the goal of this study.

The screening tool assesses how often the seven symptoms have occurred in the last four weeks, with a four-point answering
185 scale ranging from "not at all" to "5 times per week/almost always". A complete version of the 7-item scale is provided in Table A1. A symptom is considered to exist if the respondent indicates the highest or second-highest answering option (i.e., "2-4 times a week/half of the time" or "5 times per week/almost always"). The PTSD score ranging from 0 to 7 is calculated by summing up the outcomes of all seven items (symptoms). An indication of PTSD then exists if the respondent reports four or more symptoms. The cut-off point of four is derived from empirical studies and is commonly applied, as it has been proven
190 to best discriminate between patients with and without PTSD (Siegrist and Maercker, 2010; Kimerling et al., 2006; Breslau et al., 1999). Breslau et al. (1999, p. 908) reported that a score of 4 or greater "defined positive cases of PTSD with a sensitivity of 80%, specificity of 97%, positive predictive value of 71%, and negative predictive value of 98%."

As stated in Section 1, there is a variety of research on factors influencing PTSD. For this study, we used an exploratory approach, which allowed us to identify stressors specific to the Ahr Valley region to gain a more nuanced understanding of the
195 region's vulnerability and mental health status. Potential stressors were related to the conditions for probable PTSD according to the DSM-IV (American Psychiatric Association, 1994) and empirical findings from other studies exploring the impact of PTSD following flood events (e.g. Galea et al., 2007; Lamond et al., 2015).

We grouped the potential influencing factors into four categories: (a) pre-existing conditions, (b) flood characteristics, (c) stressors during the event, and (d) post-event conditions and stress. The first category refers to conditions that describe a person
200 or community before the flooding event, such as socio-demographic factors. In our study, gender, age, household income, size, and ownership were used. Further, personality traits and experiential knowledge describe a person or household before this event happened. The second category includes water level and flow velocity to describe the magnitude of the flood event. The third category is still considered to describe the conditions during the flooding event and were induced by the flood characteristics. These encompass whether respondents had to be rescued, if they were uncertain about the safety of loved ones
205 or lost someone due to the flooding event, or if they or their relatives became injured or sick due to the flood. The fourth category includes variables defining stress and coping mechanisms after the event. This involves the incurred asset damage and its repair or replacement, e.g. replacing the damaged building or contents or obtaining damage compensation within a year after the event. The amount of financial loss is also taken into account. Due to the event, many individuals had to leave their homes and have not yet been able to return. Some plan to relocate. Their personal feelings and perceptions are also considered
210 part of this category. For a comprehensive list of all variables used, please refer to Table 1.

2.3.2 Statistical analyses

IBM SPSS Version 29.0.0.0 statistics software was used for data preparation and analysis. Descriptive analyses were performed to gain an overview of the data, and correlation analyses were performed to test for multicollinearity and coherence of different variables.

215 As stated in Section 2.3.1, the questionnaire contained various questions. Overall, 26 questions were taken into account that could influence the development of PTSD. To simplify this vast number of variables, a principal component analysis (PCA) was performed to reduce the enormous number of potentially important variables from the dataset and reduce dimensions (Norris and Lecavalier, 2010), and avoid multicollinearity. Therefore, oblique (specifically oblimin) rotation was used, since this is recommended as suitable for only partially independent variables in social and behavioral research (Norris and
220 Lecavalier, 2010). This approach condensed two variables representing personality traits into a singular factor. Additionally, three variables were consolidated into “experiential knowledge,” which encapsulates participants' understanding and experience of floods and warnings (Table 1). Following this approach, a total of 23 predictors were available for analysis. Further, the existence of a PTSD indication (yes/no) was subsequently used in a logistic regression analysis to understand the influence of various potential stressors and protective factors. These stressors were explored empirically through exploratory
225 research. We developed separate binary logistic regression models for each of the four categories, considering the binary indication for PTSD or non-PTSD cases as the dependent variable. Subsequently, an overall model was conducted incorporating variables from each of the four models that proved significant at a 10% significance level. Additionally, two predicting variables underwent a sensitivity analysis, namely whether a respondent was injured or got sick and the respondent's perceived severity of negative effects on the future. One reason for testing the explanatory power of
230 getting injured or sick was the uncertainty in causality. Some respondents may have been diagnosed with a mental health disorder due to the flood event before the survey, but that was not what the particular question was trying to assess. This question aimed to determine whether the respondents had experienced infections, injuries, distress, etc. The second sensitivity analysis was undertaken because the perceived severity of negative effects on the future matched the definition of PTSD and could also be classified as a symptom (American Psychiatric Association, 1994). The two predictors were removed from their
235 category and the overall model. Nagelkerke's R^2 and Akaike Information Criterion (AIC) determined whether the predictors strongly affected the models or could be neglected.

Table 1. Comprehensive list of all variables used for the analyses. Note: Interquartile Range (IQR).


variables included	variable definition	values
pre-event conditions		
gender	The respondent's gender [0: women - reference; 1: men].	female 45%; male 55% (n = 408)
age	The respondents' ages ["18-19 years" 1-8 "> 80 years"].	median 50-59 years; (n = 409)
household income	The total income received by all members of the household within a month ["< € 900" 1-8 "> € 5,000"].	median "€ 2,600-3,599" (n = 356)
household size	The number of individuals in the respondent's household.	median 2; IQR 1 (n = 408)
ownership	The respondent is the tenant [0] or owner [1] of his/her home.	tenant 33%; owner 67% (n = 411)
personality trait	A combined factor of self-reported stress resistance " <i>I am someone who can deal with stress in a relaxed manner</i> " ["not true at all" 1-6 "very true"] and a sense of security " <i>I am someone for whom it is important to live in a safe environment. I avoid anything that could jeopardize my safety</i> " ["very true" 1-6 "not true at all"].	median 0.16, IQR 0.12
experience-based knowledge	A combined factor of the resident's flood experience, knowledge of living in a flood-prone area, and reception of a warning before the event.	median -0.66, IQR 1.33
conditions & stressors during the event: flood characteristics		
water depth	Self-reported approximate maximum water height outside the respondent's building ["there was no water inside/outside the building" 0-6 "> 4 m"].	median "> 0.5 to 1 m" (n = 398)
flow velocity	Self-reported velocity of water outside the house ["calmly flowing" 1-6 "flowing torrentially"].	median 3; IQR 1 (n = 381)
conditions & stressors during the event: stressors		
person had to be rescued	The respondent was rescued during the event [0: no; yes: 1].	yes 12% (n = 406)
person suffered serious injury/illness	The respondent him-/herself suffered from serious injury or sickness [0: no; yes: 1].	yes 11% (n = 407)
family/friends suffered serious injury/illness	A respondent's family member or close friend suffered from serious injury or sickness [0: no; yes: 1].	yes 23% (n = 404)
uncertainty about safety of family/friends	The respondent was unsure about the safety of certain family members or close friends [0:no; yes: 1].	yes 69% (n = 410)
loss of a family member/ close friend	The respondent lost a family member or close friend due to flooding [0:no; yes: 1].	yes 15% (n = 403)
post-event conditions & stress		
financial damage	Incurred damage to inventory/household contents and building/property ["no financial damage" 0-9 "> € 100,000"].	median "> € 100,000" (n = 396)
building repaired/contents replaced	The respondents' damaged building or content is already repaired/replaced ["still significant deficiencies" 1-6 "completely"].	median 3; IQR 3 (n = 392)
damage compensation completed	The damage compensation is ongoing [0] or completed [1].	completed 34% (n = 369)
return home possible	The respondent could return to his/her home ["after 1-2 days" 1-6 "not yet"]	median "after 1-2 days" (n = 411)
plans to move	The respondent intends to move elsewhere [0: no; yes: 1].	yes 7% (n = 340)
displacement	The respondent had to leave his/her home due to a summons, an official order, or damage [0: no; yes: 1].	yes 41% (n = 406)
feeling of being left alone to cope with flood impacts	The respondent feels left alone to cope with the flood impacts ["not at all" 1-6 "totally agree"].	median 2; IQR 2 (n = 401)
perceived probability of being flooded	The respondent believes he/she is likely to be affected by a flood in the future ["not likely" 1-6 "very likely"].	median 3; IQR 2 (n = 392)
perceived severity of being flooded	The respondent believes he/she is likely to be negatively affected by a flood in the future ["not bad" 1-6 "very bad"].	median 5; IQR 2 (n = 400)

3. Results & Discussion

240 Experiencing such a dramatic flood event represents an enormous mental burden for those affected. At the time of the survey, i.e. around one year after the event, about 42% of the respondents reported that they still (very) strongly felt burdened by the event.

3.1 Distribution of PTSD symptoms

245 Fig. 2 illustrates the frequency of symptoms based on the short screening scale. Our findings indicate that 67.9% of respondents experienced at least one (out of seven) PTSD symptom one-year post-event. Indeed, 5.1% experienced all seven symptoms. Fig. 2 displays these symptoms recognized as indicative of developing PTSD (see Section 2.3.1). Nearly half of the respondents experienced symptoms at least once a week, with even higher percentages reporting sleep disturbances and loss of interest in activities. About 70% of the participants reported problems sleeping at least once a week, and 20% experienced them almost daily. Additionally, many participants reported no longer enjoying activities that they used to enjoy before the flooding. The least commonly reported symptom was avoidance of thoughts, feelings, and places that reminded them of the event (Fig. 2).
250 Our study found significantly higher indications of PTSD in six out of seven symptoms than were found in a 2005 epidemiological study representative of Germany's total population using the same PTSD screening tool (Table 2). While sleep problems and loss of interest were also the most common symptoms reported one year after the flooding event, the avoidance of thoughts, feelings, and people associated with flooding was less frequent in our study.



255 **Figure 2. Distribution of PTSD symptoms reported using the PTSD screening scale by Siegrist and Maercker (2010) about one year after the occurrence of the flooding event. n = 411 for each bar.**

When comparing symptoms between genders in our study (Table 2), six out of seven symptoms were significantly higher for women. The only symptom that did not show statistically significant differences was the avoidance of thoughts, feelings, and people associated with the flooding. We assume that the ongoing reconstruction work, dealing with finance issues and insurance, and visible damages to buildings and infrastructure make it impossible to ignore the enormous impacts of the event, which were still visible at the time of the survey. Additionally, the flooding event still receives attention in national and local media, and ongoing research continuously strives to understand the circumstances better to help those affected.

Table 2. Mean values of the PTSD symptoms with possible answers ranging from 1 (“not at all”) to 4 (“5 times a week/almost always”) compared to a representative study of the total German population from 2010 measuring PTSD with the same screening tool (Siegrist and Maercker, 2010). Pearson-chi-squared test indicates gender differences with p-values of <0.05 *; <0.01 **; <0.001 *.**

PTSD symptom	District of Ahrweiler June/July 2022			A Representative Study of Germany (Siegrist and Maercker, 2010)
	Women	Men	Total	
avoidance of thoughts and feelings	1.85	1.64	1.73	1.99
loss of interest	2.38	2.08	2.21*	1.77
feeling of alienation	1.91	1.61	1.74**	1.48
dullness/numbness	2.07	1.76	1.89**	1.53
unfulfilled future plans	2.09	1.84	1.96*	1.62
sleep disturbances	2.59	2.17	2.36**	1.91
nervousness	2.14	1.72	1.91***	1.77

The conducted correlation analysis between all seven PTSD symptoms shows a significant correlation between each symptom at a 10% level (Table 3). Strong co-morbidity can be seen for loss of interest and feelings of alienation ($r = 0.58$; $p < 0.01$), loss of interest and avoidance ($r = 0.54$; $p < 0.01$), and sleep disturbances and nervousness ($r = 0.58$). These co-morbidity levels suggest that selecting a single outcome variable in regression modeling is practical, given the likelihood that similar factors will influence all outcomes, thereby reducing co-linearity. Moreover, the co-morbidity suggests that individuals encountering one symptom are more likely to experience multiple ones. This is supported by Lamond et al. (2015), who tested the co-morbidity of different psycho-social symptoms after a flood in 2007 in the UK. Moreover, several studies suggest a substantial co-morbidity of psychiatric disorders (Galatzer-Levy et al., 2013; Brady et al., 2000). Especially depression, anxiety disorders, and substance use are known to share symptoms with PTSD (Galatzer-Levy et al., 2013; Brady et al., 2000). In our study, we found an overlap with depression symptoms regarding sleep disturbances, avoidance, and loss of interest in activities, whereas anxiety, avoidance, and increased arousal overlap with PTSD symptoms. This highlights the complexity of mental health disorders and their co-existence, implying further that individuals who exhibit intense symptoms of PTSD may

experience additional mental health impacts in addition to those related to PTSD. Altogether, the high co-morbidity levels indicate that the flooding interrupted the affected individuals' daily routines.

285

Table 3. Correlation analysis of symptoms of PTSD with one another and the potentially influencing factors for PTSD used in this study. Pearson correlation coefficient; two-sided significance test indicating p-values of <0.05 *; <0.01 **; <0.001 *.**

variable	N	1	2	3	4	5	6	7
<i>symptoms</i>								
1. avoidance of thoughts and feelings	411	-						
2. loss of interest	411	.554***	-					
3. feeling of alienation	411	.488***	.580***	-				
4. dullness/numbness	411	.425***	.505***	.530***	-			
5. unfulfilled future plans	411	.324***	.410***	.479***	.506***	-		
6. sleep disturbances	411	.432***	.472***	.450***	.533***	.485***	-	
7. nervousness	411	.479***	.416***	.508***	.495***	.382***	.579***	-
<i>possible influencing factors</i>								
gender	408	-.107*	-.138**	-.150**	-.147**	-.113*	-.185**	-.199**
age	409	.043	-.047	-.058	-.106*	-.078	-.013	.009
household income	356	-.116*	-.056	-.116*	-.049	-.106*	-.177**	-.156**
household size	408	.022	.000	.017	.091	.052	-.002	-.015
ownership	411	-.024	.033	-.079	-.049	-.075	-.043	-.113*
personality trait	411	-.051	-.010	-.023	-.024	-.116*	-.042	.005
experience-based knowledge	369	-.090	-.107*	-.087	-.151**	-.100	-.107*	-.108*
water depth	406	.099*	.051	.114*	.094	.086	.049	.095
flow velocity	407	.146**	.093	.118*	.189**	.222**	.225**	.201**
person had to be rescued	404	.173**	.133**	.157**	.134**	.167**	.160**	.175**
person suffered serious injury/illness	410	.105*	.089	.173**	.151**	.110*	.165**	.167**
family/friends suffered serious injury/illness	403	.218**	.127*	.145**	.142**	.182**	.160**	.146**
uncertainty about the safety of family/friends	398	.066	.082	.078	.085	.148**	.185**	.084
loss of a family member/close friend	381	.135**	.098	.122*	.107*	.111*	.252**	.215**
financial damage	396	.058	.052	.031	.046	.057	.076	-.016
building repaired/contents replaced	392	-.182**	-.134**	-.188**	-.140**	-.262**	-.269**	-.160**
damage compensation completed	369	-.014	-.090	-.003	-.041	-.037	-.068	.028
return home possible	411	.046	.111*	.084	.078	.142**	.096	.119*
plans to move	340	.112*	.109*	.186**	.186**	.287**	.147**	.155**
displacement	406	.041	.101*	.072	.051	.096	.044	.108*
feeling of being left alone to cope with flood impacts	401	.234**	.253**	.300**	.264**	.244**	.232**	.201**
perceived probability of being flooded	392	.141**	.139**	.225**	.121*	.145**	.188**	.168**
perceived severity of being flooded	400	.243**	.257**	.253**	.284**	.338**	.338**	.289**

Table 3 shows a further correlation analysis between the seven PTSD symptoms and all the predictor variables used in this study, showing significant relationships between feeling alienated and feeling left alone to cope with the flood impacts ($r =$

0.30; $p < 0.01$), as well as for sleep disturbance and perceived severity of adverse effects of the flood in the future ($r = 0.34$; $p < 0.01$). Experiencing such a traumatic event is linked to the fear of recurrence and of being strongly affected in the future. This fear is strongly associated with adverse mental health, as was also discovered by Puechlong et al. (2020) as well as Mason, Andrews, and Upton (2010). Catastrophizing, a non-adaptive cognitive emotion regulation strategy, is also a risk factor for PTSD that is connected to the above-mentioned factors (Puechlong et al., 2020).

3.2 Prevalence of flood-related PTSD

A total of 71.8% of respondents did not exhibit indications of PTSD, according to the epidemiological screening tool. However, with a prevalence of 28.2%, it is evident that a significant portion of respondents displayed indications of PTSD approximately one year after the flooding event. The prevalence was significantly higher for women (36%, 184 female participants) than for men (22%, 224 male participants, $p < 0.01$).

The estimated prevalence of PTSD was substantially higher than typically found in the total German population. An epidemiological study conducted in 2016 (Maercker et al., 2018) found a one-month prevalence of 1.5%. In contrast, the Robert Koch Institute reported a 12-month prevalence of 2.3% for PTSD in Germany, with women being more affected (3.6%) than men (0.9%) (Jacobi et al., 2014). Another study of six European countries showed even smaller prevalence rates in the population of 0.9% (Alonso et al., 2004).

However, when comparing the estimated prevalence to the literature in which PTSD was examined after flooding events, findings are very heterogeneous, ranging from 2.6 to 52% after flooding (Golitaleb et al., 2022; Keya et al., 2023). It is important to note that most of these studies were conducted in Asia, where circumstances, susceptibilities, and methods of handling floods may differ from those in Europe. This could affect the outcomes of the studies. Nevertheless, our results fit the average of 29% given by Golitaleb et al. (2022) and seem comparable to the other findings.

It is also essential to account for the time elapsed after a traumatic event when considering the prevalence of PTSD. The percentage of people experiencing mental health symptoms can vary considerably after six months compared to those who have experienced a flood one, two, or more years ago (Mason et al., 2010; Jermacane et al., 2018; Graham et al., 2019; Zhong et al., 2018). Moreover, the overall severity of the event can contribute to differences in prevalence (Laudan et al., 2020). For Germany, the study of (Apel and Coenen, 2021) discovered PTSD prevalences of 20.4% a full 12 months after the flood of June 2013. However, as stated in Section 1, this flooding differed in type and severity compared to this case study.

In addition to possible differences in regions and cultures, it could be that the method of assessing PTSD can contribute to the variation in prevalence, for example, when diagnostic criteria are updated (e.g., DSM-III to DSM-IV to DSM-V) (Schwarz and Kowalski, 1991; Wang et al., 2000; Galea et al., 2005).

It is important to note that we were unable to filter out individuals who may have already been experiencing symptoms of PTSD before the flooding since we did not gather information on their pre-flood mental health status. However, information on respondents' pre-existing health status before the disaster can be crucial in determining the exact impacts of the flooding event, because physical impairments and chronic health conditions are known to influence mental health problems (Galea et

al., 2005; Tunstall et al., 2006). Additionally, the still ongoing COVID-19 pandemic and the war in Ukraine may have also had an impact on the prevalence of PTSD. Further, self-selection bias is possible, since those affected and randomly selected could voluntarily participate in the survey. Since individuals with severe mental health issues may have chosen not to complete the survey, this could result in an underestimation of the true prevalence of PTSD one year after the flooding event. However, we took steps in the survey introduction to inform participants about the possibility of re-traumatization. We also recommended that only those who felt comfortable participating in the survey.

3.3 Factors influencing the development of PTSD

The flood caused significant damage to many of the households that were surveyed. Half of the households suffered financial losses of more than € 100,000. The water had entered the cellar of every household, while 47.2% had water on the outside of their homes up to two meters, or even higher (40.1%). Over 40% of respondents had to vacate their homes due to the damage, and 14.4% were still unable to return home at the time of the survey, i.e., one year after the event. Additionally, 90% of respondents reported that the damaged buildings or contents had not been fully repaired or replaced. All of these and more conditions and stressors could influence the mental health status of those affected. The regression analyses presented in what follows reveal the relationship between potential factors and PTSD.

3.3.1 Pre-existing conditions

In this model, seven variables were tested, explaining 8% of the variance (Nagelkerke's $R^2 = 0.08$). As can be seen in Table 4, gender is a significant pre-existing demographic predictor of PTSD with an odds ratio (OR) of 0.4 (95% CI 0.2-0.8, $p < 0.01$), showing that men are less likely to show indications of PTSD, while women are more susceptible to developing PTSD. This is consistent with previous research, which has often shown gender to be a factor in the development of mental health issues (e.g., Paranjothy et al., 2011; Mason et al., 2010; Tunstall et al., 2006; Norris et al., 2002). However, this can have different explanations, mainly depending on social norms. On the one hand, women tend to be more sensitive and can express and show feelings more easily (Tunstall et al., 2006). They typically take on the primary caregiver role and are generally more concerned about their community (Pulcino et al., 2003; Tunstall et al., 2006). On the other hand, men are often associated with the stereotype of being strong and not showing or talking about their vulnerabilities (Turner et al., 2013).

The household income and household size have a significant correlation with disaster-related PTSD, as per the model (OR 0.8, 95% CI 0.7-1.0; OR 1.3, 95% CI 1.0-1.7). This indicates that households with lower incomes are more likely to experience adverse mental health outcomes in the aftermath of disasters. The literature also suggests that income is a crucial factor associated with negative mental health effects after disasters (Graham et al., 2019; Lamond et al., 2015; Paranjothy et al., 2011; Mason et al., 2010; Tunstall et al., 2006; Galea et al., 2005). On the other hand, high income can be seen as a protective factor. However, age, ownership, personality, and experience-based knowledge had no significant effects (Table 4) in this model.

355 **Table 4. Results of the binary regression analysis for pre-existing conditions considering the binary indication for PTSD or non-PTSD cases. Note.** n = 320, Nagelkerkes R² = 0.08, AIC = 379.40; CI = confidence interval; LL = lower limit; UL = upper limit.

explanatory variable	OR	95% CI		P
		LL	UL	
gender	.50	.30	.84	.009
age	1.04	.85	1.26	.73
household income	.85	.72	1.00	.048
household size	1.29	.99	1.68	.06
ownership	1.00	.57	1.75	.99
personality trait	.74	.50	1.08	.12
experience-based knowledge	.84	.64	1.10	.20

3.2 Conditions and stressors during the event

The flood characteristics define the initial shock and the event's magnitude and play a crucial role in determining the impact. There are several known parameters (Thieken et al., 2005); however, only water depth and flow velocity were recorded in this study, and 5% of the variance in the data was explained (Table 5).

360 The results for flow velocity in this model show a significant effect on the development of indications for PTSD (OR 1.3, 95% CI 1.1-1.6; p = 0.001). This is in line with Bubeck and Thieken (2018), where flood velocity significantly predicted a negative recovery for the (fluvial) flood of 2013 in Germany. However, water depth does not impact PTSD in our study, which is contradictory to other findings (Lamond et al., 2015; Lowe et al., 2013; Paranjothy et al., 2011).

365 **Table 5. Results of the binary regression analysis for flood characteristics during the event considering the binary indication for PTSD or non-PTSD cases. Note.** n = 381, Nagelkerkes R² = 0.05, AIC = 455.33; CI = confidence interval; LL = lower limit; UL = upper limit.

explanatory variable	OR	95% CI		P
		LL	UL	
water depth	1.01	.84	1.20	.95
flow velocity	1.32	1.11	1.56	.001

370 The event's severity is also reflected in respondents' personal experiences during the flood. Five stressors proposed by Galea et al. (2007) were tested within this model (see also Table 1). Altogether, these predictors explain 8% of the variance in the data.

At 75.9%, the vast majority of respondents reported experiencing at least one of the five stressors (see conditions & stressors during the event: stressor items in Table 1, mean 1.3, SD 1.1, n = 390), with uncertainty about the safety of a family member or close friend being the most prevalent at 69%. Nevertheless, this particular variable does not serve as a predictor for PTSD,

375 as its prevalence is comparable among residents who both do and do not exhibit indications of PTSD (see Table 6). In contrast, respondents who suffered from a severe injury or got sick due to the flooding show a high OR of 3.0 (95% CI 1.5-6.0, $p < 0.01$), meaning that if those affected are physically traumatized during the event, the likelihood of experiencing PTSD increases by 200%. Further, the sensitivity analysis reveals that this predictor contributes significantly to the model's performance: with injury/sickness predictor $R^2 = 0.08$, AIC 447.39 (see Table 6); without injury $R^2 = 0.06$, AIC = 458.29 (see Table B1).

380 Assuming the respondents related the question to the event seems reasonable since the questions before and after also referred to it. However, it cannot be completely ruled out that the question was misunderstood, and the illness referred to in the answer was a diagnosed mental health disorder due to the flood event prior to the survey. Therefore, we have decided to keep it in the model and investigate this question more thoroughly in the future. Nevertheless, becoming seriously injured during a flooding event is solid, tangible evidence a person can feel and see, and matches the definition of PTSD according to the DSM-IV

385 (American Psychiatric Association, 1994). According to Wiseman et al. (2013), PTSD is the most common disorder that people experience after a traumatic event, especially if they have been physically injured, regardless of the severity of the injury. Similar results were found by Galea et al. (2007). Although Bromet et al. (2017) found a significant effect in their univariate model, it was not present in their multivariate models.

Further, the loss of a family member or close friend emerges as a significant PTSD predictor for residents (OR 2.0, 95% CI

390 1.0-3.8, $p < 0.05$) in this model. This finding is consistent with the literature (Keyes et al., 2014; Atwoli et al., 2017; Bromet et al., 2017). No influence was detected on the stressors requiring rescue or a family member's serious illness or injury.

395 **Table 6. Results of the binary regression analysis for stressors based on Galea et al. (2007) during the event considering the binary indication for PTSD or non-PTSD cases. Note. n = 390, Nagelkerkes $R^2 = 0.08$, AIC = 447.39; CI = confidence interval; LL = lower limit; UL = upper limit.**

explanatory variable	OR	95% CI		p
		LL	UL	
person had to be rescued	.93	.45	1.93	.85
person suffered severe injury/illness	3.00	1.51	5.96	.002
family/friends suffered severe injury/illness	1.18	.65	2.13	.59
uncertainty about the safety of family/friends	1.26	.74	2.14	.40
loss of a family member/close friend	1.98	1.02	3.84	.04

3.3 Post-event conditions & stress

In summary, the fourth model deals with variables that occur after the initial shock of a flood event has subsided and the reconstruction and recovery process begins. The model is represented by eight variables and explains 26% of the variation in the data, making it the most meaningful so far. This underlines the importance of recovery processes for the well-being and

400 health of disaster-affected regions.

One year after the event, 38.3% of respondents displaying indications of PTSD expressed the feeling of being left alone to cope with flood impacts, compared to 18.2% of residents without indications of PTSD. Table 7 shows an OR of 1.5 (95% CI 1.2-1.8, $p < 0.000$), implying that respondents who feel left alone to cope with flood impacts are 50% more likely to show indications of PTSD. There is limited evidence in the literature regarding this factor. Nonetheless, most studies suggest that
405 having a support system of friends and neighbors can be helpful in protecting against mental health issues during the coping process (Norris et al., 2002; Tunstall et al., 2006; Stanke et al., 2012).

As shown in Table 7, the perceived probability of being negatively affected by future flooding was also found to have an impact on developing indications for PTSD. To further investigate this variable, a sensitivity analysis was conducted to test whether the predictor of the perceived severity of the negative future flooding aligns with the symptoms of PTSD according
410 to the DSM-IV. The regression analysis demonstrated that this predictor contributes to the accuracy of the model (with predictor $R^2 = 0.26$ AIC 290.48, see Table 7; without predictor $R^2 = 0.27$, AIC = 298.12; see B2), which is why we left the variable in the model.

Planning to move after the event significantly correlates with PTSD, showing the highest OR in this model at 4.7 (95% CI 1.4-15.7, $p < 0.01$). This suggests that PTSD could be a contributing factor leading individuals to move to a new home, making it
415 a consequence of the disorder.

The perceived probability of experiencing the adverse effects of a possible future flood has no significant influence (OR 1.5, 95% CI 1.1-2.0, $p < 0.01$).

Contrary to other findings, in our study, damage does not have a significant influence on PTSD. Jermacane et al. (2018) and Galea et al. (2007) discovered that severe financial loss resulting from flooding was linked to a higher incidence of PTSD.
420 According to Van Der Velden et al. (2023), both pre- and post-trauma financial difficulties can increase the risk of mental health problems such as PTSD. However, due to the damage scale used in this survey, the extent of financial losses was not determined precisely, leading to a small variance in the data sample.

Furthermore, there is no evidence to suggest that the lack of fully repaired or replaced damaged buildings or contents, incomplete damage compensation, no possibility to return home, displacement, or the perceived probability of future flooding
425 has any impact on PTSD (see Table 7).

Table 7. Results of the binary regression analysis for post-condition and stress considering the binary indication for PTSD or non-PTSD cases. Note. n = 278, Nagelkerkes R² = 0.26, AIC = 290.48; CI = confidence interval; LL = lower limit; UL = upper limit.

explanatory variable	OR	95% CI		p
		LL	UL	
financial damage	.89	.74	1.08	.25
building repaired/contents replaced	.87	.71	1.05	.15
damage compensation completed	.63	.28	1.44	.28
return home possible	1.09	.89	1.35	.40
plans to move	4.66	1.39	15.66	.013
displacement	.55	.13	2.43	.43
feeling of being left alone to cope with flood impacts	1.46	1.21	1.77	.000
perceived probability of being flooded	1.11	.90	1.38	.31
perceived severity of being flooded	1.48	1.12	1.95	.006

3.4 Overall model

430 Finally, the overall model integrates predictors from all four categories presented so far, significantly above the 10% significance level. It explains 28% (Nagelkerke's R² = 0.28) of the variation in the data and accurately classifies 79.1% of the categories. The model comprises nine variables, spanning all four models in Tables 4 to 7. In the overall model, three influencing factors demonstrate significant explanatory power with a p-value of less than 0.05: seriously suffering from severe injury or illness, feeling left alone to cope with flood impacts, and gender (Table 8).

435 During the flood event, residents who experienced an injury or got sick show a high odds ratio of 2.68 (95% CI 1.02-7.07, p = 0.046), meaning that if those affected were physically traumatized during the event, the likelihood of developing PTSD increased by 168%. Further, the sensitivity analysis in B1 reveals that this stressor needs to be considered and better investigated. As discussed in Section 3.2, this variable has a strong association with PTSD, according to the literature (Wiseman et al., 2013; Galea et al., 2007). This suggests that it is essential to monitor the mental health of individuals who
 440 have been injured during a flooding event and provide necessary treatment to help stabilize not only their physical but also their mental well-being. To exclude misinterpretations, the PTSD assessment should be accompanied by questions on illnesses and injuries induced by the flood, as well as information about the pre-health status.

The odds ratio of 1.5 (95% CI 1.22-1.86, p < 0.000) suggests that individuals who feel left alone to cope with flood impacts are 50% more likely to suffer from PTSD. Additionally, the significant positive correlation (r = 0.30; p < 0.01) between feelings
 445 of alienation and the feeling to being left alone during the coping process (see Table 3) indicates a lack of support during the recovery and reconstruction phase, which is an important entry point for better post-event disaster management. Half of the survey respondents (54.4%, n = 357) reported difficulties finding available building surveyors, craftsmen or materials. Additionally, managing finances seems to be a source of stress for many, with 18% (n = 357) reporting that the disbursement was insufficient or that the insurance did not acknowledge the damage and did not disburse (10.1%, n = 357). However, a

450 similar share stated they had enough reserves themselves. Moreover, 14.8% (n = 357) had to wait for their landlord or community association to act before repairs were viable.

One attempt to support those affected by the disaster was establishing a trauma help center (“Trauma Hilfe Zentrum im Ahrtal”) a few months after the event happened. The center is located in one of the most affected areas and offers psychosocial counseling services. According to Dr. Scharping (personal communication, March 4, 2024), head of the center, 1,475
455 individual counseling sessions had been held and 70 group offers were utilized by September 2022. More than two years after the event, more people still seek help. We believe this center is important to support the reconstruction process and should be maintained long-term.

According to our study, gender is the most significant demographic factor, with an OR of 0.41 (CI 95% 0.21-0.80, $p < 0.01$). This aligns with previous research, which indicates that women are at a higher risk, often due to societal norms (refer to Section
460 3.1).

Table 8. Results of the binary regression analysis for the final model considering the binary indication for PTSD or non-PTSD cases.
Note. n = 253, Nagelkerkes $R^2 = 0.28$, AIC = 252.85; CI = confidence interval; LL = lower limit; UL = upper limit.

explanatory variable	OR	95% CI		p
		LL	UL	
gender	.41	.21	.80	.009
household income	.85	.69	1.05	.13
household size	1.08	.77	1.51	.65
flow velocity	1.28	.78	2.10	.33
person suffered severe injury/illness	2.68	1.02	7.07	.046
loss of a family member / close friend	1.19	.49	2.89	.70
plans to move	2.87	.78	10.57	.11
feeling of being left alone to cope with flood impacts	1.50	1.22	1.86	.000
perceived severity of being flooded	1.32	.99	1.74	.056

Although the plan to move after the event shows the greatest effect size (OR 2.87, see Table 8) on PTSD among the residents,
465 with a p-value of 0.11, it does not seem to be a significantly strong predictor. This is probably due to the small sample size of respondents who plan to move (6.8%, 28 of n = 340). The high confidence interval (CI) indicates a disagreement and ongoing discussion among the stakeholders (politicians, authorities, residents, etc.) about relocation and resettlement after the flooding event. However, the willingness to move to another location permanently is somewhat influenced by other variables, such as ownership and the housing situation at the time of the survey, and does not influence the indications of the development of
470 PTSD. Respondents who could stay in their homes were less willing to move than respondents who had to move temporarily, and homeowners were less inclined to move than tenants (Truedinger et al., 2023).

Several participants in the survey raised concerns about the potential adverse effects of floods in the future. However, this was not strongly significant in the model (OR 1.32, 95% CI 0.99-1.74, $p = 0.06$). The correlation analysis showed that this factor

is significantly linked to sleeping disturbances ($r = 0.34$; $p < 0.01$, see Table 3), the most frequently stated symptom among
475 respondents, as well as unfulfilled future plans ($r = 0.34$; $p < 0.01$). The sensitivity analysis shows $R^2 = 0.28$ and $AIC = 252.85$
with the predictor and $R^2 = 0.27$ and $AIC = 261.27$ without the predictor, indicating further investigation of this variable.
Mason et al. (2010) suggest that experiencing a threatening situation like a flood is linked to the fear of recurrence, which
results in being strongly affected in the future. So, negatively perceiving future flood risk can harm a person's mental health,
which is also supported by Puechlong et al. (2020). This feeling and associated symptoms require proper treatment, as ignoring
480 them may worsen the symptoms and lead to even worse mental health impacts in the future. Further, Babcicky et al. (2021)
propose that such psychosocial indicators, e.g. the perceived probability and fear of being flooded again in the future, need
consideration when regulating losses and supporting the recovery process rather than only looking at damages.

Although the model included the factors listed as significant in the previous sections (3.1 to 3.3), the variables of household
income and size, flow velocity, and the loss of a family member or close friend did not produce any significant results in the
485 overall model.

The relationship between predictors and PTSD is often complex, and it is not always clear to what extent predictors are
dependent on each other or even on other circumstances. This lack of clarity is also why different studies only sometimes agree
on the significance of the predictors or the direction of influence. Other researchers have also addressed this issue (Galea et
al., 2005; Lamond et al., 2015). Further, assessing cause and effect for potential influencing factors is difficult.

490 **4. Conclusion**

Our findings highlight the major psychosocial challenges those affected during and after the flooding event faced. Given the
severity of the flooding, we assumed that a large number of residents would experience negative mental health impacts, such
as PTSD. Our study provides evidence of such effects on the mental health of affected residents one year after the flood event
in the heavily affected district of Ahrweiler. Around 70% of those affected by the flood reported suffering from sleep
495 disturbances and a lack of enjoyment in their activities at least once a week, which underlines the event's impact on their daily
lives. Additionally, we have estimated that 28% of respondents from the heavily affected district of Ahrweiler are displaying
indications of PTSD one year after the event based on the short epidemiological screening scale.

Further, by analyzing possible influencing factors that could contribute to PTSD, this study has uncovered several essential
factors. Firstly, women seem to be particularly at risk of experiencing PTSD symptoms. Secondly, during the flood event itself,
500 the most critical factor contributing to PTSD indications was whether someone became injured or sick due to the event. Thirdly,
in the aftermath of the flood, the feeling of being left alone to cope with flood impacts was the most significant factor
contributing to PTSD indications. Altogether, the findings show that such a catastrophic flood can lead to new health-related
needs that require attention. Psychosocial offers should be adjusted to address the high prevalence and risk factors associated
with PTSD in the affected population. On the one hand, our results show that long-term support is needed. On the other hand,
505 targeted interventions are required, especially during and after the flood event. The first implication is a targeted aftercare

program for those injured during the flood event. Furthermore, the findings suggest a significant need for practical assistance during reconstruction. A helpful step was to establish a trauma support center. To ensure that those affected who, for whatever reason, are unable to seek help on their own receive the help they need, it would be beneficial to introduce a visiting assistance program where qualified professionals go door-to-door and inquire whether they can be of any assistance (known as “aufsuchende Hilfe” in German). Further, results showed that overcoming bureaucratic obstacles, finding available building surveyors and craftsmen, and managing finances can all hinder a speedy recovery process. Nevertheless, there is a need for further research into coping strategies. Additional research is also needed to show the causalities of different influencing factors. Additionally, more longitudinal studies are necessary, ideally with the same sample, to determine the timeframe of negative mental health impacts and to get more insights into recovery processes. The insights can be utilized to assess the effectiveness of various measures. Finally, we propose integrating the long-term mental health-related impacts of floods into overall flood risk management, as it is an essential part of the recovery process following a flood, yet often overlooked in many regions of the world (see section 1 for selected countries with a framework). Developing a framework to support the mental health of those impacted by floods can help in building more resilient communities. Furthermore, while German Flood Risk Management includes the term "avoidance and reduction of adverse effects on human health," it lacks further explanation and recommendations. This study provides information to contribute to such a framework by highlighting the prevalence and risk factors associated with PTSD following the 2021 flood in the Ahr Valley, Germany.

Author contribution. MLZ, PB, and AT conceptualized the study and participated in data collection. MLZ conducted the analyses. PB contributed to parts of the methods sections, while MLZ authored and visualized the initial draft, which underwent review and editing by co-authors. MLZ finalized the manuscript for submission, under the supervision of PB and AT.

Competing interests. The authors declare that they have no conflict of interest.

Funding. This research was funded by the Federal Ministry of Education and Research (Bundesministerium für Bildung und Forschung – BMBF) under the Project “KAHR - Climate Adaptation, Floods and Resilience”, grant number (FKZ) 01LR2102I.

Acknowledgements. We thank all respondents for participating in the survey.

Appendix

Appendix A. PTSD-screening scale.

535 **Table A1. Short PTSD-screening scale used in the questionnaire. Original by Breslau et al. (1999), German translation and adaption by Siegrist and Maercker (2010).**

	<i>not at all</i>	<i>max. once a week</i>	<i>2 to 4 times a week/ half the time</i>	<i>5 times a week/ almost always</i>
<i>Did you avoid being reminded of this experience by avoiding certain places, people or activities?</i>	•	•	•	•
<i>Did you lose interest in activities that were once important or enjoyable?</i>	•	•	•	•
<i>Did you begin to feel more isolated or distant from other people?</i>	•	•	•	•
<i>Did you find it hard to have love or affection for other people?</i>	•	•	•	•
<i>Did you begin to feel there was no point in planning for the future?</i>	•	•	•	•
<i>After this experience were you having more trouble than usual falling asleep or staying asleep?</i>	•	•	•	•
<i>Did you become jumpy or get easily startled by ordinary noises or movements?</i>	•	•	•	•

Appendix B. Sensitivity analyses.

540

Table B1. Sensitivity analysis showing the results of the binary regression analysis for stressors based on Galea u. a. (2007), without the predictor “suffering severe injury/illness” during the event considering the binary indication for PTSD or non-PTSD cases.

explanatory variable	OR	95% CI		p
		LL	UL	
<i>conditions & stressors during the event: stressors</i>				
person had to be rescued	1.10	.55	2.21	.78
family/friends suffered severe injury/illness	1.38	.78	2.45	.27
uncertainty about the safety of family/friends	1.19	.70	2.01	.52
loss of a family member/close friend	2.41	1.28	4.54	.007
n = 393, Nagelkerkes R ² = 0.06, AIC = 458.29.				
<i>overall model</i>				
gender	.41	.21	.79	.008
household income	.84	.68	1.03	.09
household size	1.08	.78	1.49	.65
flow velocity	1.35	.82	2.21	.23
loss of a family member/close friend	1.48	.64	3.41	.36
plans to move	3.69	1.07	12.71	.04
feeling of being alone to cope with flood impacts	1.51	1.23	1.87	.000
perceived severity of being flooded	1.33	1.01	1.76	.04
n = 254, Nagelkerkes R ² = 0.26, AIC = 255.33.				

545 **Table B2. Sensitivity analysis showing the results of the binary regression analysis for post-event conditions and stress, without the predictor “perceived severity of being flooded” considering the binary indication for PTSD or non-PTSD cases.**

explanatory variable	OR	95% CI		p
		LL	UL	
<i>post-event conditions & stress</i>				
financial damage	.96	.80	1.15	.67
building/contents replaced	.88	.73	1.06	.18
damage compensation completed	.63	.29	1.41	.26
return home possible	1.10	.90	1.35	.34
plans to move	5.94	1.76	2.11	.004
displacement	.55	.13	2.30	.41
feeling of being left alone to cope with flood impacts	1.52	1.26	1.83	.000
perceived probability of being flooded	1.21	.99	1.48	.06
n = 282, Nagelkerkes R ² = 0.22, AIC = 298.12.				
<i>overall model</i>				
gender	.37	.20	.71	.003
household income	.82	.67	1.01	.07
household size	1.16	.84	1.60	.36
flow velocity	1.29	.80	2.08	.30
suffering severe injury/illness	2.63	1.01	6.86	.049
loss of a family member/close friend	1.39	.59	3.25	.45
plans to move	4.33	1.24	15.15	.02
feeling of being left alone to cope with flood impacts	1.54	1.25	1.89	.000
n = 261, Nagelkerkes R ² = 0.27, AIC = 261.27.				

References

- Alfieri, L., Burek, P., Feyen, L., and Forzieri, G.: Global warming increases the frequency of river floods in Europe, *Hydrol. Earth Syst. Sci.*, 19, 2247–2260, <https://doi.org/10.5194/hess-19-2247-2015>, 2015.
- Alonso, J., Angermeyer, M. C., Bernert, S., Bruffaerts, R., Brugha, T. S., Bryson, H., de Girolamo, G., Graaf, R., Demyttenaere, K., Gasquet, I., Haro, J. M., Katz, S. J., Kessler, R. C., Kovess, V., Lépine, J. P., Ormel, J., Polidori, G., Russo, L. J., Vilagut, G., Almansa, J., Arbabzadeh-Bouchez, S., Autonell, J., Bernal, M., Buist-Bouwman, M. A., Codony, M., Domingo-Salvany, A., Ferrer, M., Joo, S. S., Martínez-Alonso, M., Matschinger, H., Mazzi, F., Morgan, Z., Morosini, P., Palacín, C., Romera, B., Taub, N., and Vollebergh, W. A. M.: Prevalence of mental disorders in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project, *Acta Psychiatr. Scand. Suppl.*, 21–27, <https://doi.org/10.1111/j.1600-0047.2004.00327.x>, 2004.
- American Psychiatric Association (Ed.): Diagnostic and statistical manual of mental disorders. DSM-IV., American Psychiatric Publishing, Inc., 1994.
- Apel, D. and Coenen, M.: Mental health and health-related quality of life in victims of the 2013 flood disaster in Germany – A longitudinal study of health-related flood consequences and evaluation of institutionalized low-threshold psycho-social support, *Int. J. Disaster Risk Reduct.*, 58, 6, <https://doi.org/10.1016/j.ijdrr.2021.102179>, 2021.
- Atwoli, L., Stein, D. J., King, A., Petukhova, M., Aguilar-Gaxiola, S., Alonso, J., Bromet, E. J., De Girolamo, G., Demyttenaere, K., Florescu, S., Maria Haro, J., Karam, E. G., Kawakami, N., Lee, S., Lepine, J.-P., Navarro-Mateu, F., O’Neill, S., Pennell, B.-E., Piazza, M., Posada-Villa, J., Sampson, N. A., Ten Have, M., Zaslavsky, A. M., Kessler, R. C., and on behalf of the WHO World Mental Health Survey Collaborators: Posttraumatic stress disorder associated with unexpected death of a loved one: Cross-national findings from the world mental health surveys: ATWOLI ET AL., *Depress. Anxiety*, 34, 315–326, <https://doi.org/10.1002/da.22579>, 2017.
- Babcicky, P., Seebauer, S., and Thaler, T.: Make it personal: Introducing intangible outcomes and psychological sources to flood vulnerability and policy, *Int. J. Disaster Risk Reduct.*, 58, 102169, <https://doi.org/10.1016/j.ijdrr.2021.102169>, 2021.
- Bei, B., Bryant, C., Gilson, K.-M., Koh, J., Gibson, P., Komiti, A., Jackson, H., and Judd, F.: A prospective study of the impact of floods on the mental and physical health of older adults, *Aging Ment. Health*, 17, 992–1002, <https://doi.org/10.1080/13607863.2013.799119>, 2013.
- Brady, K. T., Killeen, T. K., Brewerton, T., and Lucerini, S.: Comorbidity of psychiatric disorders and posttraumatic stress disorder, *J. Clin. Psychiatry*, 61 Suppl 7, 22–32, 2000.
- Breslau, N., Peterson, E. L., Kessler, R. C., and Schultz, L. R.: Short screening scale for DSM-IV posttraumatic stress disorder, *Am. J. Psychiatry*, 156, 908–911, <https://doi.org/10.1176/ajp.156.6.908>, 1999.
- Bromet, E. J., Atwoli, L., Kawakami, N., Navarro-Mateu, F., Piotrowski, P., King, A. J., Aguilar-Gaxiola, S., Alonso, J., Bunting, B., Demyttenaere, K., Florescu, S., de Girolamo, G., Gluzman, S., Haro, J. M., de Jonge, P., Karam, E. G., Lee, S., Kovess-Masfety, V., Medina-Mora, M. E., Mneimneh, Z., Pennell, B.-E., Posada-Villa, J., Salmerón, D., Takeshima, T., and Kessler, R. C.: Post-traumatic stress disorder associated with natural and human-made disasters in the World Mental Health Surveys, *Psychol. Med.*, 47, 227–241, <https://doi.org/10.1017/S0033291716002026>, 2017.
- Bubeck, P. and Thieken, A. H.: What helps people recover from floods? Insights from a survey among flood-affected residents in Germany, *Reg. Environ. Change*, 18, 287–296, <https://doi.org/10.1007/s10113-017-1200-y>, 2018.

- 585 Bundesamt für Kartographie und Geodäsie (BKG): GeoBasis-DE: Verwaltungsgebiete 1:5 000 000, Stand 01.01. (VG5000 01.01.), dl-de/by-2-0, GeoBasis-DE/BKG [data set], <https://gdz.bkg.bund.de/index.php/default/digitale-geodaten/verwaltungsgebiete/verwaltungsgebiete-1-5-000-000-stand-01-01-vg5000-01-01.html> , 2024.
- Bundesanstalt für Gewässerkunde (BfG): Waterbody-DE. Waterbody as ESRI file geodatabase, ETRS89 (non INSPIRE conformant), WasserBLICK/BfG und Zuständige Behörden der Länder [data set],
 590 <https://geoportal.bafg.de/inspire/download/HY/waterbody/datasetfeed.xml>, 2023.
- CRED and UNDRR: Human costs of disasters. An overview of the last 20 years, 2021.
- Deutsche Bahn: Zerstörungen in historischem Ausmaß: DB zieht nach Flutkatastrophe Zwischenbilanz: https://www.deutschebahn.com/de/presse/pressestart_zentrales_uebersicht/Zerstoerungen-in-historischem-Ausmass-DB-zieht-nach-Flutkatastrophe-Zwischenbilanz-6868360, last access: 17 May 2022.
- 595 DH Emergency Preparedness Division: Emergency Planning Guidance: Planning for the Psychosocial and Mental Health Care of People Affected by Major Incidents and Disasters, DH Emergency Preparedness Division, London, 2009.
- Dietze, M., Bell, R., Ozturk, U., Cook, K. L., Andermann, C., Beer, A. R., Damm, B., Lucia, A., Fauer, F. S., Nissen, K. M., Sieg, T., and Thieken, A. H.: More than heavy rain turning into fast-flowing water – a landscape perspective on the 2021 Eifel floods, <https://doi.org/10.5194/egusphere-2022-7>, 2022.
- 600 Dreßing, H. and Foerster, K.: Traumafolgestörungen in ICD-10, ICD-11 und DSM-5, *Forensische Psychiatr. Psychol. Kriminol.*, 15, 47–53, <https://doi.org/10.1007/s11757-020-00645-6>, 2021.
- Eurostat/GISCO: NUTS, European Commission - Eurostat/GISCO, EuroGeographics and TurkStat [data set], <https://ec.europa.eu/eurostat/de/web/gisco/geodata/reference-data/administrative-units-statistical-units/nuts>, 2021.
- 605 Fekete, A. and Sandholz, S.: Here Comes the Flood, but Not Failure? Lessons to Learn after the Heavy Rain and Pluvial Floods in Germany 2021, *Water*, 13, 3016, <https://doi.org/10.3390/w13213016>, 2021.
- Fernández, A., J. Black, Mairwen K. Jones, L. Wilson, L. Salvador-Carulla, T. Astell-Burt, and D. Black: Flooding and Mental Health: A Systematic Mapping Review, undefined, 2015.
- Frans, Ö., Rimmö, P. -A., Åberg, L., and Fredrikson, M.: Trauma exposure and post-traumatic stress disorder in the general population, *Acta Psychiatr. Scand.*, 111, 291–290, <https://doi.org/10.1111/j.1600-0447.2004.00463.x>, 2005.
- Galatzer-Levy, I. R., Nickerson, A., Litz, B. T., and Marmar, C. R.: Patterns Of Lifetime PTSD Comorbidity: A Latent Class Analysis: Research Article: Patterns of Lifetime PTSD Comorbidity, *Depress. Anxiety*, 30, 489–496, <https://doi.org/10.1002/da.22048>, 2013.
- Galea, S., Nandi, A., and Vlahov, D.: The epidemiology of post-traumatic stress disorder after disasters, *Epidemiol. Rev.*, 27, 78–91, <https://doi.org/10.1093/epirev/mxi003>, 2005.
 615
- Galea, S., Brewin, C. R., Gruber, M., Jones, R. T., King, D. W., King, L. A., McNally, R. J., Ursano, R. J., Petukhova, M., and Kessler, R. C.: Exposure to hurricane-related stressors and mental illness after Hurricane Katrina, *Arch. Gen. Psychiatry*, 64, 1427–1434, <https://doi.org/10.1001/archpsyc.64.12.1427>, 2007.
- Golitaleb, M., Mazaheri, E., Bonyadi, M., and Sahebi, A.: Prevalence of Post-traumatic Stress Disorder After Flood: A Systematic Review and Meta-Analysis, *Front. Psychiatry*, 13, 890671, <https://doi.org/10.3389/fpsy.2022.890671>, 2022.
 620

- Graham, H., White, P., Cotton, J., and McManus, S.: Flood- and Weather-Damaged Homes and Mental Health: An Analysis Using England's Mental Health Survey, *Int. J. Environ. Res. Public Health*, 16, 3256, <https://doi.org/10.3390/ijerph16183256>, 2019.
- 625 Jacobi, F., Höfler, M., Strehle, J., Mack, S., Gerschler, A., Scholl, L., Busch, M. A., Maske, U., Hapke, U., Gaebel, W., Maier, W., Wagner, M., Zielasek, J., and Wittchen, H.-U.: Psychische Störungen in der Allgemeinbevölkerung: Studie zur Gesundheit Erwachsener in Deutschland und ihr Zusatzmodul Psychische Gesundheit (DEGS1-MH), *Nervenarzt*, 85, 77–87, <https://doi.org/10.1007/s00115-013-3961-y>, 2014.
- 630 Jermacane, D., Waite, T. D., Beck, C. R., Bone, A., Amlôt, R., Reacher, M., Kovats, S., Armstrong, B., Leonardi, G., James Rubin, G., and Oliver, I.: The English National Cohort Study of Flooding and Health: the change in the prevalence of psychological morbidity at year two, *BMC Public Health*, 18, 330, <https://doi.org/10.1186/s12889-018-5236-9>, 2018.
- Jongman, B., Ward, P. J., and Aerts, J. C. J. H.: Global exposure to river and coastal flooding: Long term trends and changes, *Glob. Environ. Change*, 22, 823–835, <https://doi.org/10.1016/j.gloenvcha.2012.07.004>, 2012.
- 635 Jongman, B., Hochrainer-Stigler, S., Feyen, L., Aerts, J. C. J. H., Mechler, R., Botzen, W. J. W., Bouwer, L. M., Pflug, G., Rojas, R., and Ward, P. J.: Increasing stress on disaster-risk finance due to large floods, *Nat. Clim. Change*, 4, 264–268, <https://doi.org/10.1038/nclimate2124>, 2014.
- Junghänel, T., Bissolli, P., Daßler, J., Fleckenstein, R., Imbery, F., Janssen, W., Kaspar, F., Lengfeld, K., Leppelt, T., Rauthe, M., Rauthe-Schöch, A., Rocek, M., Walawender, E., and Weigl, E.: Hydro-klimatologische Einordnung der Stark- und Dauerniederschläge in Teilen Deutschlands im Zusammenhang mit dem Tiefdruckgebiet „Bernd“ vom 12. bis 19. Juli 2021., 2021.
- 640 Keya, T. A., Leela, A., Habib, N., Rashid, M., and Bakthavatchalam, P.: Mental Health Disorders Due to Disaster Exposure: A Systematic Review and Meta-Analysis, *Cureus*, <https://doi.org/10.7759/cureus.37031>, 2023.
- Keyes, K. M., Pratt, C., Galea, S., McLaughlin, K. A., Koenen, K. C., and Shear, M. K.: The Burden of Loss: Unexpected Death of a Loved One and Psychiatric Disorders Across the Life Course in a National Study, *Am. J. Psychiatry*, 171, 864–871, <https://doi.org/10.1176/appi.ajp.2014.13081132>, 2014.
- 645 Kimerling, R., Ouimette, P., Prins, A., Nisco, P., Lawler, C., Cronkite, R., and Moos, R. H.: Brief report: Utility of a short screening scale for DSM-IV PTSD in primary care, *J. Gen. Intern. Med.*, 21, 65–67, <https://doi.org/10.1111/j.1525-1497.2005.00292.x>, 2006.
- 650 Kreibich, H., Thieken, A. H., Petrow, Th., Müller, M., and Merz, B.: Flood loss reduction of private households due to building precautionary measures – lessons learned from the Elbe flood in August 2002, *Nat. Hazards Earth Syst. Sci.*, 5, 117–126, <https://doi.org/10.5194/nhess-5-117-2005>, 2005.
- Kron, W., Eichner, J., and Kundzewicz, Z. W.: Reduction of flood risk in Europe – Reflections from a reinsurance perspective, *J. Hydrol.*, 576, 197–209, <https://doi.org/10.1016/j.jhydrol.2019.06.050>, 2019.
- Kron, W., Bell, R., Thiebes, B., and Thieken, A.: The July 2021 flood disaster in Germany, Tokyo, 2022.
- 655 Lamond, J. E., Joseph, R. D., and Proverbs, D. G.: An exploration of factors affecting the long term psychological impact and deterioration of mental health in flooded households, *Environ. Res.*, 140, 325–334, <https://doi.org/10.1016/j.envres.2015.04.008>, 2015.

- Laudan, J., Zöller, G., and Thieken, A. H.: Flash floods versus river floods – a comparison of psychological impacts and implications for precautionary behaviour, *Nat. Hazards Earth Syst. Sci.*, 20, 999–1023, <https://doi.org/10.5194/nhess-20-999-2020>, 2020.
- 660 Lowe, D., Ebi, K. L., and Forsberg, B.: Factors increasing vulnerability to health effects before, during and after floods, *Int. J. Environ. Res. Public Health*, 10, 7015–7067, <https://doi.org/10.3390/ijerph10127015>, 2013.
- Maercker, A., Hecker, T., Augsburger, M., and Kliem, S.: ICD-11 Prevalence Rates of Posttraumatic Stress Disorder and Complex Posttraumatic Stress Disorder in a German Nationwide Sample, *J. Nerv. Ment. Dis.*, 206, 270–276, <https://doi.org/10.1097/NMD.0000000000000790>, 2018.
- 665 Mason, V., Andrews, H., and Upton, D.: The psychological impact of exposure to floods, *Psychol. Health Med.*, 15, 61–73, <https://doi.org/10.1080/13548500903483478>, 2010.
- Merz, B., Elmer, F., Kunz, M., Mühr, B., Schröter, K., and Uhlemann-Elmer, S.: The extreme flood in June 2013 in Germany, *Houille Blanche*, 100, 5–10, <https://doi.org/10.1051/lhb/2014001>, 2014.
- Munich Re: Schadenereignisse in Deutschland 1980 – 2017. *NatCatSERVICE.*, 2018.
- 670 Munich Re: Hurricanes, cold waves, tornadoes: Weather disasters in USA dominate natural disaster losses in 2021, <https://www.munichre.com/en/company/media-relations/media-information-and-corporate-news/media-information/2022/natural-disaster-losses-2021.html>, last access: 22 December 2023, 2022.
- National Mental Health Commission: NATIONAL DISASTER MENTAL HEALTH AND WELLBEING FRAMEWORK. Supporting Australians’ mental health through disaster, National Mental Health Commission, 2023.
- 675 Nitschke, M., Einsle, F., Lippmann, C., Simonis, G., Köllner, V., and Strasser, R. H.: Emergency evacuation of the Dresden Heart Centre in the flood disaster in Germany 2002: perceptions of patients and psychosocial burdens, *Int. J. Disaster Med.*, 4, 118–124, <https://doi.org/10.1080/15031430601138290>, 2006.
- Norris, F. H., Friedman, M. J., and Watson, P. J.: 60,000 disaster victims speak: Part II. Summary and implications of the disaster mental health research, *Psychiatry*, 65, 240–260, <https://doi.org/10.1521/psyc.65.3.240.20169>, 2002.
- 680 Norris, M. and Lecavalier, L.: Evaluating the Use of Exploratory Factor Analysis in Developmental Disability Psychological Research, *J. Autism Dev. Disord.*, 40, 8–20, <https://doi.org/10.1007/s10803-009-0816-2>, 2010.
- Paranjothy, S., Gallacher, J., Amlôt, R., Rubin, G. J., Page, L., Baxter, T., Wight, J., Kirrage, D., McNaught, R., and Sr, P.: Psychosocial impact of the summer 2007 floods in England, *BMC Public Health*, 11, 145, <https://doi.org/10.1186/1471-2458-11-145>, 2011.
- 685 Puechlong, C., Weiss, K., Le Vigouroux, S., and Charbonnier, E.: Role of personality traits and cognitive emotion regulation strategies in symptoms of post-traumatic stress disorder among flood victims, *Int. J. Disaster Risk Reduct.*, 50, 101688, <https://doi.org/10.1016/j.ijdrr.2020.101688>, 2020.
- Pulcino, T., Galea, S., Ahern, J., Resnick, H., Foley, M., and Vlahov, D.: Posttraumatic Stress in Women after the September 11 Terrorist Attacks in New York City, *J. Womens Health*, 12, 809–820, <https://doi.org/10.1089/154099903322447774>, 2003.
- 690 Schwarz, E. D. and Kowalski, J. M.: Posttraumatic stress disorder after a school shooting: effects of symptom threshold selection and diagnosis by DSM-III, DSM-III-R, or proposed DSM-IV, *Am. J. Psychiatry*, 148, 592–597, <https://doi.org/10.1176/ajp.148.5.592>, 1991.

- 695 S.I. Seneviratne, X. Zhang, M. Adnan, W. Badi, C. Dereczynski, A. Di Luca, S. Ghosh, I. Iskandar, J. Kossin, S. Lewis, F. Otto, I. Pint, M. Sato, S.M. Vicente-Serranno, M. Wehn, and B. Zho: Weather and Climate Extreme Events in a Changing Climate. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, in: *Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Intergovernmental Panel On Climate Change, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1513–1766, <https://doi.org/10.1017/9781009157896>, 2021.
- 700 Siegrist, P. and Maercker, A.: Deutsche Fassung der Short Screening Scale for DSM-IV Posttraumatic Stress Disorder. *Aktueller Stand der Validierung, Trauma Gewalt*, 4, 208–213–213, 2010.
- Stanke, C., Murray, V., Amlôt, R., Nurse, J., and Williams, R.: The effects of flooding on mental health: Outcomes and recommendations from a review of the literature, *PLoS Curr.*, 4, e4f9f1fa9c3cae, <https://doi.org/10.1371/4f9f1fa9c3cae>, 2012.
- 705 Statistical Office Rhineland-Palatinate (RP): Soforthilfe RLP 2021. <https://www.statistik.rlp.de/de/link/>, last access: 2 February 2024. 2021a.
- Statistical Office Rhineland-Palatinate (RP): Verfügbares Einkommen 2019 in allen kreisfreien Städten und Landkreisen gestiegen: https://www.statistik.rlp.de/no_cache/de/einzelansicht/news/detail/News/3341/, last access: 2 February 2024. 2021b.
- 710 Statistical Office Rhineland-Palatinate (RP): Mein Kreis, meine kreisfreie Stadt. Landkreis Ahrweiler: <https://infothek.statistik.rlp.de/MeineHeimat/content.aspx?id=101&l=1&g=07131&tp=1027>, last access: 2 February 2024. 2022.
- Tapsell, S.: Socio-Psychological Dimensions of Flood Risk Management, in: *Flood Risk Science and Management*, edited by: Pender, G. and Faulkner, H., Wiley, 407–428, <https://doi.org/10.1002/9781444324846.ch20>, 2010.
- 715 Thieken, A., Bubeck, P., and Zenker, M.-L.: Fatal incidents during the flood of July 2021 in North Rhine-Westphalia, Germany: what can be learnt for future flood risk management?, *J. Coast. Riverine Flood Risk*, 2, <https://doi.org/10.59490/jcrfr.2023.0005>, 2023a.
- Thieken, A. H., Müller, M., Kreibich, H., and Merz, B.: Flood damage and influencing factors: New insights from the August 2002 flood in Germany, *Water Resour. Res.*, 41, 2005WR004177, <https://doi.org/10.1029/2005WR004177>, 2005.
- 720 Thieken, A. H., Bessel, T., Kienzler, S., Kreibich, H., Müller, M., Pisi, S., and Schröter, K.: The flood of June 2013 in Germany: how much do we know about its impacts?, *Nat. Hazards Earth Syst. Sci.*, 16, 1519–1540, <https://doi.org/10.5194/nhess-16-1519-2016>, 2016.
- Thieken, A. H., Bubeck, P., Heidenreich, A., Von Keyserlingk, J., Dillenaar, L., and Otto, A.: Performance of the flood warning system in Germany in July 2021 – insights from affected residents, *Nat. Hazards Earth Syst. Sci.*, 23, 973–990, <https://doi.org/10.5194/nhess-23-973-2023>, 2023b.
- 725 Tradowsky, J. S., Philip, S. Y., Kreienkamp, F., Kew, S. F., Lorenz, P., Arrighi, J., Bettmann, T., Caluwaerts, S., Chan, S. C., De Cruz, L., De Vries, H., Demuth, N., Ferrone, A., Fischer, E. M., Fowler, H. J., Goergen, K., Heinrich, D., Henrichs, Y., Kaspar, F., Lenderink, G., Nilson, E., Otto, F. E. L., Ragone, F., Seneviratne, S. I., Singh, R. K., Skålevåg, A., Termonia, P., Thalheimer, L., Van Aalst, M., Van Den Bergh, J., Van De Vyver, H., Vannitsem, S., Van Oldenborgh, G. J., Van Schaeybroeck, B., Vautard, R., Vonk, D., and Wanders, N.: Attribution of the heavy rainfall events leading to severe flooding in Western Europe during July 2021, *Clim. Change*, 176, 90, <https://doi.org/10.1007/s10584-023-03502-7>, 2023.
- 730

- Truedinger, A. J., Jamshed, A., Sauter, H., and Birkmann, J.: Adaptation after Extreme Flooding Events: Moving or Staying? The Case of the Ahr Valley in Germany, *Sustainability*, 15, 1407, <https://doi.org/10.3390/su15021407>, 2023.
- Tunstall, S., Tapsell, S., Green, C., Floyd, P., and George, C.: The health effects of flooding: social research results from England and Wales, *J. Water Health*, 4, 365–380, <https://doi.org/10.2166/wh.2006.031>, 2006.
- 735 Turner, L. R., Alderman, K., Huang, C., and Tong, S.: Impact of the 2011 Queensland floods on the use of tobacco, alcohol and medication, *Aust. N. Z. J. Public Health*, 37, 396, <https://doi.org/10.1111/1753-6405.12093>, 2013.
- U.S. Government: Post-Disaster Mental Health Response Act, HR 5703S 37677 117th Congr., 2021-2022.
- 740 Van Der Velden, P. G., Contino, C., Muffels, R., Verheijen, M. S., and Das, M.: The impact of pre- and post-trauma financial problems on posttraumatic stress symptoms, anxiety and depression symptoms, and emotional support: A prospective population-based comparative study, *J. Anxiety Disord.*, 96, 102714, <https://doi.org/10.1016/j.janxdis.2023.102714>, 2023.
- Vorogushyn, S., Apel, H., Kemter, M., and Thielen, A.: Statistical and hydraulic analysis of flood hazard in the Ahr valley, Germany considering historical floods, *display*, <https://doi.org/10.5194/iahs2022-660>, 2022.
- 745 Wang, X., Gao, L., Shinfuku, N., Zhang, H., Zhao, C., and Shen, Y.: Longitudinal Study of Earthquake-Related PTSD in a Randomly Selected Community Sample in North China, *Am. J. Psychiatry*, 157, 1260–1266, <https://doi.org/10.1176/appi.ajp.157.8.1260>, 2000.
- Wiseman, T., Foster, K., and Curtis, K.: Mental health following traumatic physical injury: An integrative literature review, *Injury*, 44, 1383–1390, <https://doi.org/10.1016/j.injury.2012.02.015>, 2013.
- 750 Zhong, S., Yang, L., Toloo, S., Wang, Z., Tong, S., Sun, X., Crompton, D., FitzGerald, G., and Huang, C.: The long-term physical and psychological health impacts of flooding: A systematic mapping, *Sci. Total Environ.*, 626, 165–194, <https://doi.org/10.1016/j.scitotenv.2018.01.041>, 2018.