

ClimeApp: Data processing tool for monthly, global climate data from the ModE-RA paleo-reanalysis, 1422 to 2008 CE

Richard Warren^{1,2}, Niklaus Bartlome^{1,2}, Noémie Wellinger^{1,3}, Jörg Franke^{2,3}, Ralf Hand^{2,3}, Stefan Brönnimann^{2,3}, and Heli Huhtamaa^{1,2}

5 ¹Institute of History, University of Bern, Bern, CH-3012, Switzerland

²Oeschger Centre for Climate Change Research, University of Bern, Bern, CH-3012, Switzerland

³Institute of Geography, University of Bern, CH-3012, Switzerland




Correspondence to: Niklaus Bartlome (niklaus.bartlome@unibe.ch)
or Richard Warren (richard.warren@unibe.ch)

10 **Abstract:** ClimeApp is a newly developed web-based processing tool for the state-of-the-art ModE-RA paleo-climate reanalysis. It presents temperature, precipitation and pressure reconstructions with global coverage and monthly resolution for the period 1422 to 2008 C.E. These can be visualized as maps or timeseries and compared with historical or other climate-related information through composite, correlation and regression functions. ClimeApp allows access to three data sets: 1. ModE-RA, a reanalysis that is created by assimilating early
15 instrumental documentary and proxy data into an ensemble of climate model simulations. 2. ModE-Sim, the native version of the underlying ensemble simulations, i.e. prior to data assimilation. 3. ModE-RAclim, an alternative version of the reanalysis product. Together, these allow researchers to separate the effects of model simulations and observations on the reanalysis. The app is designed to allow quick data processing for climatologists and easy use for non-climatologists. Specifically, it aims to help bring climate into the humanities,
20 where climatological data still has huge potential to advance research. This paper outlines the development, processing and applications of ClimeApp, and presents an updated analysis of the calamitous Tambora volcanic eruption and the 1816 ‘year without a summer’ in Europe, using the new ModE datasets. ClimeApp is available at <https://mode-ra.unibe.ch/climeapp/>.

1 Introduction

25 Interdisciplinary research is a great facilitator of scientific progress. It allows researchers to address all aspects of a problem and take a holistic view not limited to one specialised field. To assess, for instance, the impact of a volcanic eruption such as Mount Tambora in 1815, volcanologists may study the eruption itself (Kandlbauer and Sparks, 2014), climatologists use numerical models to assess the climate impact (Raible et al., 2016), historians look into administrative records for social impacts (Krämer, 2015), and economists and epidemiologists look for
30 effects on trade and the spread of disease (Wood, 2014). By combining their results, researchers can obtain unique insights relevant to all fields, as demonstrated by Brönnimann and Krämer (2016). To bridge disciplines as different as history and climate science, it is crucial to make data accessible to non-specialists. ClimeApp is a web-based application that achieves this by giving researchers without a climatology background the possibility to add climate reconstruction data to their sources and analysis. It facilitates quick and easy processing of the
35 ModE-RA climate reanalysis and associated datasets (Valler et al., 2024), allowing both climatologists and other researchers to evaluate the data without time-consuming coding. Functions for both casual investigation, as well as detailed statistical and source analysis, are built in. It is also possible to upload your own research – be it

prices, harvest yields or mortality – as a timeseries to compare with the ModE-RA dataset. ClimeApp, developed using the Shiny R package (Chang et al., 2024), stands out for its simplicity and accessibility in presenting and processing complex data. This paper summarises the ModE-RA, ModE-Sim and ModE-RAclim datasets used in ClimeApp and outlines the main features of ClimeApp’s internal structure and external interface. We then examine the 1815 Tambora eruption and the following ‘year without a summer’ in Europe to showcase the functions and applications of both ClimeApp and the ModE data. This demonstrates how reanalysis data can be used to distinguish between internally and externally forced variability of the climate system. We conclude by expanding on the app’s potential for both the humanities and climate sciences. Detailed reference material for ClimeApp data processing can be found in the appendices.



50 **Figure 1.** Layout, functions and data flow of ClimeApp. (a) ClimeApp, displaying a global temperature anomaly map. (b) ClimeApp showing the same global temperatures as a timeseries. (c) Summary of the data sources used by ClimeApp, along with its five primary functions and current options for further data analysis.

2 Data Sets

2.1 ModE-Sim

ModE-Sim is an ensemble of global climate simulations, driven by external forcings. The ensemble is produced
55 by inputting reconstructed volcanic aerosols, total solar irradiation and greenhouse gases into the ECHAM6
atmosphere general circulation model (Stevens et al., 2013), with observed/reconstructed sea surface temperature
(SST) and sea ice acting as boundary conditions. The different ensemble members are produced by using
different versions of the SST and volcanic forcings, which accounts for the uncertainty in the reconstructions.
Each member in the ModE-Sim ensemble therefore represents a possible climate state that is physically
60 compatible (from the model's perspective) with the prescribed external forcings and boundary conditions (Hand
et al., 2023). The ModE-Sim ensemble, which is the basis/prior for ModE-RA (see below), is therefore a
combination of externally forced signals and internal variability predicted by the model physics. The ModE-Sim
ensemble mean, used by ClimeApp, is the average over all ensemble members. Averaging tends to “smooth out”
most of the internal variability, but does retain signals caused by external forcing, e.g. the climate's reaction to a
65 volcanic eruption. On its own, ModE-Sim can be used as a tool to study climate variability, but in combination
with ModE-RA it can also help climatologists identify how observations affect the final reanalysis (see our case
study, section 4.1).

2.2 ModE-RA

At the time of writing, the Modern Era Reanalysis (ModE-RA) is the most comprehensive reconstruction of the
70 monthly global climate of the past 600 years (Valler et al., 2024). By assimilating a huge database of climate
observations, each ModE-Sim ensemble member is corrected/updated towards the observations, weighted by
individual uncertainties in both observations and simulations. The number of observations increases rapidly
through time, rising from a few thousand natural proxies and historical documents in the 15th century to over
100,000 (mostly instrumental) measurements by the early 20th century. These climate observations are
75 assimilated biannually for each year. Natural proxies include tree rings, ice cores, corals, speleothems and lake
sediments, while historical proxies are gathered from weather diaries, phenological data and early ship records.
Where and when records are available, this allows observation-based reconstruction of all four seasons,
improving on previous studies which mostly focussed on the summer season. If no observations are available,
ModE-RA is identical to ModE-Sim. Note that the ensemble mean of ModE-Sim mainly represents the model's
80 response to external forcings, but the assimilation of observations into ModE-RA brings back internal variability.
This also means that there is increasing variability in the ModE-RA ensemble mean over time (as the number of
observations increases). The spatial horizontal resolution for ModE-RA is 1.875° (longitude) by 1.865°
(latitude), while the temporal resolution is one month.

2.3 ModE-RAclim


85 ModE-RAclim (Valler et al., 2024) is an alternative version of ModE-RA, designed to focus on the observations,
by minimising its input from the ModE-Sim climate models. In ModE-RA, observations are assimilated into
time-aligned ensemble members from ModE-Sim (i.e. for the year 1800, the basis for ModE-RA is the externally
forced ModE-Sim states for 1800). By contrast, ModE-RAclim uses the same assimilation process (see Valler et
al., 2024 for more detail), but using non-time-aligned ModE-Sim members. Instead, for each year, a random

90 sample of 100 years from any of the ModE-Sim ensemble members is used to form the ModE-RAclim ensemble,
 prior to assimilation. This means that in ModE-RAclim, the externally forced signal in the model simulations is
 removed from the ensemble and only added back if it appears in the observations. Hence, following a volcanic
 eruption, ModE-RA and ModE-Sim will both include the model response to stratospheric aerosol forcing, but
 any volcanic signal in ModE-RAclim will exclusively result from the observations. ModE-RAclim is therefore
 95 useful for distinguishing whether observed climate anomalies are a consequence of the external forcing that was
 used for the model simulations or a result of data assimilation (see our case study, section 4.1).

3 Interface

3.1 Building a Shiny app – User Interface (UI), Server and Helpers

100 R Shiny Apps have become popular not only for research in general (Gebauer et al., 2023) but particularly in
 climate science. They provide a powerful set of tools for quickly creating and deploying simple applications for
 presenting or processing data. Other scientific applications designed using R Shiny are described in: (Sousa,
 2019; Fajardo et al., 2020; Möller et al., 2020). There are three main components to a Shiny app: The local
 server where all the necessary data is stored, the R environment and the web interface.



105

Figure 2. Internal Structure of ClimeApp, showing how the data storage, R environment and web interface interact.

In ClimeApp, ModE data from an external server is stored locally on the ClimeApp server, including some pre-processed data to reduce loading times. All the server preparation, such as loading the R-libraries, unpacking the data, loading in parameters and predefined functions is done in a supporting R-file called *helpers.R*. The main


110 *app.R* file consists of a *UI* section, where all the elements are defined, named and positioned, and a *Server*
section where processing of inputs and plot generation is performed. When in use, the *Server* and web interface
interact with each other reactively, with inputs from the web elements processed in real time and then output as
plots or data tables. All web elements have preset inputs to generate an initial plot, which can then be altered by
the user.

115 **3.2 Usability**

The ClimeApp web interface is designed to streamline data selection, providing all necessary information and
options without being overly complex. The initial *Welcome* page displays general information about the ModE
database and ClimeApp and gives access to five main tabs, one for each of the primary functions shown in
Figure 1. We deliberately decided to supply only functions that are commonly used in historical climatology, but
120 allow users to easily download selected ModE data for further processing. Input and customization options are
selected within a tab, appearing on the same screen as the plots. For new users, it is vital not to overload the
interface with information - be it with text or advanced options. To avoid this, we work with help texts that
explain how plots and data are calculated, but only when selected. The interface is also streamlined by hiding
many UI elements until they are required. Non-standard tools such as custom months, single year maps and
125 additional features to customize plots are only visible when the relevant customization option or tick box is
selected.

3.3 Customization

Having generated a plot, users have several options to customize their maps and timeseries. ClimeApp is
designed to create graphics suitable for publication without the need for further editing in other programmes. The
130 app supplies three sets of tools to customize plots: *Customize your map/timeseries*; *Custom features*; and *Custom
statistics*. Under *Customize your plot*, the user can change titles, subtitles and the plot's axis. Once selected, axis
values will stay fixed even after a plot has been changed, making it easy to quickly compare different regions
and time periods. *Custom features* adds and removes points, lines and highlights to/from the plot and includes a
location search function (Tennekes, 2021) that can add labelled, geocoded points. *Custom statistics* allows users
135 to add the *SD ratio* or *% sign match* statistics (see Appendix 2 for more information) as overlays to their maps or
add percentiles and moving averages to their timeseries. All customization stays, even when plots are altered or
different variables selected, saving time when creating multiple plots. Customization and plot inputs can also be
downloaded, saved and re-uploaded to quickly recreate previous plots.



(b)

Customize your map

Map customization

Axis customization:
 Default Custom

Set your axis values:
 to

Hide axis completely

Title customization:
 Default Custom

Custom map title:

Custom map subtitle (e.g. Ref-Period)

Custom features

Enable custom features

Select a feature type:
 Point Highlight

Add custom points
 Enter location/coordinates or double click on map
 Enter a location:

Point label:

Point longitude:

Point latitude:

Point shape:

Point colour:

Point size:

Custom statistics

Enable custom statistics

Choose custom statistic:
 SD ratio % sign match

SD ratio <

140

Figure 3. Example of a customized map from ClimeApp, showing: (a) An anomaly map using several custom features (customized title, custom axis, labelled point, highlight box and statistical overlay showing regions that match a chosen criteria); (b). ClimeApp’s map customization options.

3.4 Uploading/downloading data and plots

145

Nearly all the plots in ClimeApp can be downloaded in multiple file formats. Plot data can also be exported for use in other programmes. In the *Anomalies* tab, advanced users can download multiple Mode variables as a

single NetCDF file, while in the *Regression* tab, a statistical summary of the linear regression is also available for download. Users can upload their own data in three of the main tabs: *Composites* can be created from an imported set of years, while tables of annual data can be uploaded for *Correlation* or *Regression* analysis.

150 Uploaded files can be compared with either ModE variables or further user data, making ClimeApp a useful tool for performing simple correlation and regression analysis on your own material. Since files must be correctly formatted for ClimeApp to understand them, when uploading, users are shown an example image demonstrating how to correctly structure their file.

3.5 Reactive functions and pre-processing

155 One major benefit of Shiny apps is instant updating – as soon as a user selects new values, ClimeApp will automatically update its plots to match the new selection. This is timesaving and allows researchers to explore the climate data quickly and efficiently. Exceptions have been made for changing the geographical area, adding points, lines and highlights and using *Annual Cycles* function, to allow all information to be entered before updating the plot. To enable plots to instantly update, ClimeApp uses R Shiny’s *reactive functions*. These
160 functions track all the variables they depend on, re-executing if one of these variables changes. This means that only sections of code where a change has been made are re-run, reducing processing power and increasing response time. Reactive functions chain - each passing an output to a further reactive function until the final reactive output is passed to the UI, updating the onscreen plot or table.

165 To further reduce processing time, ClimeApp employs pre-processed data for common user selections. These include five seasonal averages: December-January-February (DJF); March-April-May (MAM); June-July-August (JJA); September-October-November (SON); and Annual; and four variables: Temperature; precipitation; sea level pressure; and 500 hPa geopotential height. Annual means for each are already calculated and stored on the ClimeApp server, allowing them to be plotted without further computation. This considerably
170 reduces the memory and time required for data processing and is especially valuable when users are working on large geographical areas or over long time periods.

4 Case Study – Tambora reanalysed

In 1815, on the Indonesian island of Sumbawa, Mount Tambora erupted. Large volumes of sulphur were injected into the stratosphere, forming a layer of aerosols around the globe. This layer considerably reduced incoming
175 solar radiation, significantly affecting the global and particularly European climate (Brönnimann and Krämer, 2016). Using the latest and most comprehensive climate data, ModE-RA, ModE-Sim and ModE-RAclim, and the tools in ClimeApp, we can take a new look at the eruption's impact.

4.1 Tambora in ModE-RA, ModE-Sim and ModE-RAclim

The year of 1816, following Tambora, has become known as the ‘year without a summer’, due to the extreme
180 global temperature anomalies that followed the eruption. Using ClimeApp’s *Anomalies* function, we can see the unusually cold summer temperatures over Europe in all three ModE datasets:





Figure 4. (a), (b) & (c) European temperature anomalies for June, July and August, 1816, as compared to reference period 1799 to 1821. Showing results from ModE-RA, ModE-Sim and ModE-RAclim, respectively. (d) The standard deviation ratio (or SD ratio) between the ensembles of ModE-RA and ModE-Sim (see Source Analysis and Further Statistical Functions in Appendix 2 for more information).

A comparison between the datasets shows that the 1816 anomalies in ModE-RA and ModE-RAclim are essentially the same, while ModE-Sim reports considerably smaller temperature changes. This suggests that either certain mechanics of the Tambora eruption are not captured by the ModE-Sim ensemble, or that only part of the observed cooling can be explained by aerosol forcing. The latter might indicate that lower temperatures were partly due to internal climate variability. This would support the suggestion by Auchmann et al. (2013) and Brönnimann and Krämer (2016) that additional cooling in the 1810s may have been attributable to the long-term effects of the 1809 ‘unknown eruption’, low solar activity and internal decadal variability in the ocean-atmosphere system.

The similarity between ModE-RA and ModE-RAclim results from the profusion of data sources for this period and region - 4778 European observations were assimilated for April to September in 1816 alone (see Figure 6). These constrain ModE-RA and ModE-RAclim, but not ModE-Sim. We can see this from the SD ratios in Figure 4d. The SD ratio measures the difference in spread between the ensembles of ModE-RA and ModE-Sim. An SD ratio of 1 means that both ensembles have the same standard deviation, i.e. nothing was assimilated or observations had no effect, while an SD ratio of 0 indicates an overconfident reanalysis since no uncertainty would remain. In general however, a lower SD ratio indicates where more information has been assimilated. In 4d, we can see this over central and northern Europe, where the lower SD ratio suggests a more reliable reanalysis, or at least one that is closer to the observations.

However, the SD ratio should not be the only measure of the dataset’s reliability. If we use ClimeApp to view global precipitation rather than temperature in 1816, we can see the disparity between the ModE datasets reverse:




210 **Figure 5.** (a), (b) & (c) Global precipitation anomalies for June, July and August 1816, as compared to reference period 1799 to 1821. Showing results from ModE-RA, ModE-Sim and ModE-RAclim respectively. (d) Standard deviation (SD) ratio of ModE-RA ensemble to ModE-Sim ensemble, showing significant constraint by observations only over Europe, coastal North America and SE India.

Here, ModE-RA very closely matches ModE-Sim, but not ModE-RAclim. This is because, while the tropical monsoon should have been affected by Tambora (Marti and Ernst, 2009), there are few equatorial observations to capture this. ModE-RAclim is therefore unable to reconstruct significant changes in tropical precipitation. ModE-Sim meanwhile, includes physical mechanisms predicting extreme monsoon changes from volcanic aerosol forcing, and passes these on to ModE-RA. Interestingly, in the one tropical location that is more constrained - southern India (see Fig. 5a) - ModE-RAclim does more closely match the predictions from ModE-Sim. Results from both precipitation and temperature broadly agree with previous reconstructions (Brönnimann, 2015) and modelling (Wegmann et al., 2014), though there are some regional differences in precipitation which might reflect improved modelling or more comprehensive observations.

215
220

Assimilated Observations - Apr. to Sept. 1816


Total Sources = 7470



225 **Figure 6.** ClimeApp *ModE-RA sources* plot, showing all assimilated observations for April to September 1816. By comparing the sources plot to the SD ratios (Figs. 4d and 5d), we can see which observations significantly constrain the ModE-RA ensemble for a particular variable. For example, instrumental data (red) noticeably constrains JJA temperatures in Europe (Fig. 4d), but the documentary proxies (yellow) in Africa have only a minor effect on the ensemble (possibly because the proxies correlate only weakly with boreal summer temperatures).

4.2 Compositing volcanic years

230 To compare Tambora with other volcanic eruptions, we can use the *Composites* function in ClimeApp to view the temperature anomalies in 1816 alongside a composite of the anomalies following other major eruptions:




235 **Figure 7.** (a) Composite of JJA temperature anomalies following the 12 largest volcanic eruptions, excluding Tambora, between 1422 and 2008, as measured by their impact on global stratospheric aerosol optical depth (Toohey and Sigl, 2017). Composited years are 1454, 1459, 1596, 1601, 1642, 1696, 1784, 1810, 1832, 1836, 1884 and 1992, corresponding to the maximum volcanic forcing following each eruption. Anomalies are calculated with respect to the three years preceding each composite year. (b) Temperature anomalies in 1816, following Tambora, as compared to the three years prior.

In the ModE-RA dataset, we see considerably larger anomalies following Tambora than in the more general composite of other eruptions. While this could suggest other, non-volcanic effects on the European climate, it is likely that the more individual patterns of each eruption are smoothed out in the composite, generating the more homogeneous weaker cooling. In this case, compositing highlights the forced part of the volcanic signal, similar to running an ensemble of simulations. This would explain the similarity between Figure 7a and Figure 4b (the ensemble mean from ModE-Sim). The disparity may also partly result from the volcanic forcing following Tambora being approximately twice that of the composite sample or the fact that the composite volcanic years were also generally less constrained than 1816 and therefore closer to ModE-Sim.

4.3 Removing the volcanic signal?

Using ClimeApp, we can also attempt to remove the volcanic signal altogether from ModE-RA. The volcanic forcing data used in ModE-Sim (Toohey and Sigl, 2017), measures global volcanic forcing through reconstructed stratospheric aerosol optical depth (SAOD). ClimeApp's linear *Regression* function can build a simple statistical model linking SAOD to European JJA temperatures. In principle this can 'remove' associated effects (at least according to the linear regression model), leaving the 'residual' temperature variation unrelated to volcanic forcing (Fig. 8c):



255 **Figure 8.** (a) Timeseries showing the ‘original’ European JJA temperatures (1422-2008) used to train the model and the predicted ‘trend’ in temperatures from the regression model. Note that both timeseries show the spatial average for each year, while the actual regression model consists of an original and trend for each point on the map. (b) Map showing the linear regression coefficients linking JJA temperature and global SAOD. (c) Residual European JJA temperatures in 1816.

The residuals (Fig. 8c) show significant non-volcanic anomalies in 1816, that generally match the results from the anomaly and composite analysis. The regression coefficients (Fig. 8b) agree with the results from ModE-Sim, predicting a mostly homogenous cooling over the European landmass in response to aerosol forcing. However, any further conclusions are limited by the assumption of a linear relation between SAOD and JJA temperatures. This is unlikely, given the complex atmospheric dynamics governing the climate response to the radiative forcing (Brönnimann, 2015). Furthermore, the limited number of major eruptions reduces the reliability of any relationship drawn from a simple comparison of the data.

4.4 Comparing ModE-RA to historical data

Tambora didn't only affect the climate, human society was also significantly impacted. At the turn of the 19th century, grain prices were the most important indicator to measure the state of an economy (Ljungqvist et al., 2022). Even if other factors such as demand, military conflicts and the quality of the harvest played a role, quantitative supply was an important determinant of price (Krämer, 2015). We would therefore expect a link between price data and temperatures during the growing season. This can be tested using a sample of bread price data from Lucerne, Switzerland and ClimeApp's *Correlation* function:




Figure 9. Timeseries correlation showing JJA temperatures from 1750-1850 over central Europe and the average price of bread (“Weissbrot”) in Lucerne, Switzerland for the following year (Haas-Zumbühl, 1903). Timeseries correlation of $r = -0.328$, with a p value < 0.01 . Prices are in Rappen per kg. Note that the following year bread price was used due to the lagged effect of temperature during the growing season on market prices.

The timeseries correlation gives us an inverse correlation of $r = -0.328$. This implies a weak but general correlation between the growing season temperatures and grain prices, where low temperatures presumably led to a poor season and a spike in bread prices the following year. The connection is particularly visible after Tambora, where we see both a sharp rise in prices (in 1817) and a sudden temperature drop during the summer season of 1816. This coincided with unusually high precipitation, which may explain why we do not see a similar price spike in 1813 or 1821 when precipitation was moderate. Our results agree with previous work on Tambora, which demonstrated how adverse weather in Switzerland in 1816 played a crucial role in massively reduced crop yields. In the following year, this led to an increase in prices for grain and bread and widespread famine, particularly in eastern Switzerland (Brönnimann and Krämer, 2016).

5 Outlook

5.1 Potential for humanities

It is well known how intertwined climate and society have been over the last 1000 years (Pfister and Wanner, 2021). Naturally, this has important implications for all sciences that concern themselves with humanity, but especially for history. An understanding of the climate can complement traditional historical approaches, adding a crucial dimension to our understanding of the past. As demonstrated in our case study, anomaly plots for specific years can illustrate how different regions were affected by extreme climate events. Correlations can measure statistical connections between climate and historical variables, ranging from crop yields and mortality rates to price data. Knowledge of climate history can also give us perspective on the current climate crisis, with historical examples employed as case studies for the impacts of a changing climate on human societies (Lamb, 2002; Degroot et al., 2021). However, there are still challenges to interweaving the two disciplines. These include the heterogeneity of approaches, the diversity of disciplinary perspectives and often mutually unintelligible terminology (White et al., 2023). We can innovate to overcome these difficulties though however. There exists a growing collection of databases and tools for accessing documentary sources and climate data for use in historical climatology. Some primary examples include tambora.org (Riemann et al., 2015), Euro-Climhist (Pfister et al., 2017) and Climate Explorer (Trouet and van Oldenborgh, 2013). Each tool fulfils a different niche. Tambora.org and Euro-Climhist both provide a vast number of original sources and indexed data from (predominantly) human archives. This qualitative data - usually documentary data, but in the case of Euro-Climhist also visual sources such as paintings – is extremely helpful for investigating and interpreting singular events or climatological trends. However, it is often sporadic and temporally and geographically inhomogeneous, hence the advantage of modelling and reanalysis products like ModE-RA and ModE-Sim. Climate Explorer meanwhile, is an excellent tool for accessing and analysing some older climate reconstructions, as well as modern instrumental data. However, its vast array of options and data can often be confusing for new users, and it does not give access to the very latest reanalysis data.

This is where ClimeApp is able to provide a complimentary tool, utilizing a smaller range of data and functions, but via a far more user-friendly interface. ClimeApp provides highly customizable plots, along with helpful explanations where necessary. It also, uniquely, allows users to view all the sources behind the reanalysis data. Combined with the standard deviation ratios (see case study, section 4.1), these allow detailed assessment of the reliability and applicability of the reanalysis. For historians, this possibility to make source-attributed, publication suitable plots without coding skills is invaluable. Furthermore, for researchers using quantitative data, ClimeApp can be a convenient tool for correlation or regression analysis without needing external programmes. However, the most important feature of ClimeApp is that it uses the most extensive and modern global reanalysis for the climate of the last 600 years, which can be directly compared with historical socio-economic time series. In general, the broad appeal of ClimeApp for the humanities lies in its simplification of accessing, visualising and analysing the latest reanalysis data.

5.2 Potential for climate sciences

We have already demonstrated some of ClimeApp's potential for climate science in our case study, but a few further points are worth making here. First and foremost, ClimeApp can save researchers considerable time

325 creating field and timeseries plots that would otherwise have to be manually coded. As it incorporates many
analysis methods used in historical climatology and paleoclimatology, climatologists studying the age before
instrumental measurements will find the ModE data invaluable. Our case study showed how contrasting ModE-
RA, ModE-Sim and ModE-RAclim can give constructive insights into the causes of certain climate anomalies,
while the source plots and SD ratio data can be used to assess the reliability and the limitations of ModE data.
330 For the developers of ModE-RA, ClimeApp is particularly useful for testing their data, allowing them to quickly
visualise and compare the datasets. Other researchers can use the app to assess new paleoclimate data, using the
Correlation function to map correlations between their data and ModE-RA. For students and lecturers, we hope
that ClimeApp makes advanced reanalysis data easily accessible for exercises and project work. Finally,
ClimeApp can be a useful template for developing other Shiny projects, particularly those for processing and
335 plotting complex data. The source code for the application is openly available (see Code availability) and can be
freely adapted for other applications.

6 Conclusions

This paper summarized the functionality and potential of the new ClimeApp web application. It demonstrates
how simple applications can be powerful tools to make specialized data available to all. The application provides
340 historians, climatologists and other researchers quick access to state-of-the-art ModE climate reanalysis.
Through a user-friendly interface, students and scientists can view and analyse the historical global climate with
just a few clicks. Furthermore, and uniquely, ClimeApp allows detailed investigation of the sources used in a
climate reconstruction. The *ModE-RA sources* tool displays the type and location of all proxies and documentary
sources used to constrain the ModE-RA climate models, while the *SD ratio* statistical analysis quantifies their
345 effect on the final data. By re-examining the Tambora eruption and the ‘year without a summer’ in 1816, we
demonstrated how combining results from ModE-RA, ModE-Sim and ModE-RAclim can help us interpret the
reanalysis data and begin to separate different factors affecting the paleoclimate. We also showed how the
correlation and regression functions can be used to combine ModE-RA with independent data to identify
statistical relationships between climatic and non-climatic variables. ClimeApp, created and developed by
350 historians, geographers and climate scientists, is as good an example as any of the value of bridging science and
humanities to advance interdisciplinary research. In this paper we have focussed on the app’s potential in our
respective fields, but possible applications could be imagined in many of the social and physical sciences. Our
hope is that researchers from various disciplines will benefit from using ClimeApp and ModE-RA, finding
innovative and enlightening ways to integrate climate data into their own research.

355 7 Appendices

Appendix 1: R libraries used by ClimeApp

```
library(shiny)  
library(ncdf4)  
library(maps)  
360 library(shinyWidgets)  
library(RColorBrewer)
```

```
library(shinyjs)
library(bslib)
library(readxl)
365 library(xlsx)
library(DT)
library(zoo)
library(colourpicker)
library(tmaptools)
370 library(ggplot2)
library(sf)
library(shinylogs)
library(shinycssloaders)

375 ClimeApp v1.0 uses R-version 4.3.2.
```

Appendix 2: ClimeApp functions and data processing

Behind the Shiny interface, the processing and analysis done by ClimeApp is relatively straightforward.

ClimeApp utilises the set of R libraries in Appendix 1 to extract and process the raw ModE data into a format selected by the user.

380 **Anomalies**

The anomaly map function shows the spatial distribution of climate anomalies averaged over a user-selected year range and month range. For example, June, July, August (JJA), 1501 to 1600 if your focus is boreal summer in the 16th century. The anomalies are created from 3 data products:

- 385 1. *Annual Means* – a timeseries of annual means for each point on the map, created by averaging absolute ModE values across the selected month range.
2. *Reference Means* – a single reference mean for each point on the map, created by averaging *annual means* across a chosen reference year range.
3. *Annual Anomalies* – a timeseries of annual anomalies for each point on the map, created by subtracting the *reference means* from the *annual means*.

390 The final anomalies shown are the time-averaged annual anomalies. These are plotted using the base R plotting functions along with the coastlines and borders from the *maps* package. The anomaly timeseries is generated by averaging the annual anomalies for each year.

For reference, the calculations behind each data product are as follows:

395 The *annual mean* for a single year and single point on the map is given by the equation

$$AnnualMean = \overline{AbsoluteValues(M)}$$

where M is the selected month range.

400 The *reference mean* for a single year and point is given by

$$ReferenceMean = \overline{AnnualMeans(Y_{ref})}$$

where Y_{ref} is the selected reference year range.

The *annual anomaly* for a single year and point is given by:

405
$$AnnualAnomaly = AnnualMean - ReferenceMean$$

Note that in the case of ModE-RAclim, the base data is already in anomaly format, so anomalies are merely calculated by subtracting time-averaged anomalies from each other.

410 The anomalies presented on the anomaly map and in the anomaly map data are given by

$$Anomaly(map) = \overline{AnnualMeans(Y)} - ReferenceMean = \overline{AnnualAnomalies(Y)}$$

where Y is the selected year range.

Anomalies presented on the timeseries map and timeseries data are given by

415
$$Anomaly(timeseries) = \overline{AnnualMeans(Lon, Lat)} - \overline{ReferenceMeans(Lon, Lat)}$$

$$= \overline{AnnualAnomalies(Lon, Lat)}$$

where Lon and Lat are the selected longitude and latitude range.

Composites

420 ClimeApp's composite maps show the time-averaged anomalies for a set of non-consecutive years, which can be entered or uploaded by the user. The anomaly reference period can be a fixed set of consecutive years, a custom set of non-consecutive years or an individual reference period generated for each year based on the X (a number of years chosen by the user) years prior. Calculations and plotting are performed in the same way as for anomalies, except for anomalies compared to X years prior (XYP):

- 425
1. *XYP Reference Means* – a set of reference means for each point on the map, one for each user-selected year. Calculated by averaging the X preceding *annual means*.
 2. *XYP Annual Anomalies* – a set of annual anomalies for each point on the map. Created by subtracting the corresponding *reference mean* from each *annual mean*.

430 To give an indication of the consistency of anomalies over the set of years in the composite, ClimeApp contains a '% sign match' statistical tool. This marks regions where the *annual anomalies* that form the composite agree in their sign more often than a user-defined threshold, given in percent. For example, for a composite of five years, with anomalies of -1°C, -5°C, 1°C, 15°C and -3°C, the displayed mean would be a positive 1.4°C, but only 40% of the years would match this, since 3 are in fact negative.

Correlation

435 The correlation function allows users to generate a map of correlation coefficients, comparing either ModE variables or user-uploaded timeseries. Using the *cor()* function from the *stats* R package (R Core Team, 2022), it can employ either the Pearson or Spearman's Ranks correlation method. If both variables are in 'field' format, i.e. gridded map data, it performs a timeseries correlation of the *annual*

440 *means* for each point on the map with the corresponding *annual means* for the second variable. If one
variable is a timeseries however, it correlates each set of *annual means* with the same timeseries. In
addition to the map, ClimeApp also produces a correlation timeseries, showing an annual timeseries of
both variables (spatially averaged in the case of ModE variables) and a single correlation coefficient and
p-value, calculated from those timeseries. The p-value shows the probability that the correlation was
445 produced by random chance rather than an actual relationship between the variables. $p < 0.05$ is
generally recommended for drawing legitimate conclusions.

Regression

In ClimeApp, regression operates in a similar way to correlation, performing a multiple linear
regression analysis on a set of *annual means*. Using *lm()* from the *stats* R package, one or more
independent variable timeseries are fitted to the dependent variable timeseries for each point on the map
450 according to the model

$$V_{Dependent} = \beta_1 V_{Independent1} + \beta_2 V_{Independent2} + \dots + \alpha + Residual$$

where β is the coefficient and α is the intercept. ClimeApp then plots the spatial average of the
dependent variable, trend ($\beta_1 V_{Independent1} + \beta_2 V_{Independent2} + \dots$) and residual as a timeseries. Provided the
dependent variable is a field, maps of the coefficients for each independent variable can be produced, as
455 can maps of the p-values and residuals for each year.

Annual Cycles

This function shows the spatially averaged monthly ModE values over a given year or set of years. In
the case of a set of years, these can be presented individually or as an average.

Source Analysis and Further Statistical Functions

460 The accuracy of ModE-RA is dependent on the availability and reliability of observations to constrain
the model ensemble of ModE-Sim. To capture this, ClimeApp includes tools for visualizing the sources
used to create ModE-RA and ModE-RAclim and the standard deviation (SD) ratio of the ModE-RA and
ModE-Sim ensembles. The ModE-RA sources are presented as a semi-annual map showing the data
points assimilated for each half-year, grouped by type and variable (see Figure 6). This allows the user
465 to see where proxy, documentary or instrumental observations were integrated into the reconstruction
and any gaps in the data. The SD ratio meanwhile, is the standard deviation of the ModE-Sim ensemble
divided by the standard deviation of ModE-RA after the assimilation of observations:

$$SDratio = \frac{\sigma_{ModE-RAEnsemble}}{\sigma_{ModE-Ensemble}}$$

This gives a value between 0 and 1 for each month and grid point, with 1 showing no constraint (i.e. the
470 ModE-RA output is the same as that of ModE-Sim and entirely generated from the models) and lower
values showing increasing constraint by observations, meaning there are either more observations or
that they are more ‘trusted’ by the reconstruction. The temporal mean of the SD ratio can be presented
in ClimeApp as a contour map or grid-point overlay on the anomaly maps.

475 On timeseries plots, users have the option to add percentiles and moving averages. The moving
 averages are calculated using a rolling mean of timeseries values over a number of years selected by the
 user (default 11). To generate the percentiles, a Shapiro-Wilk test (Shapiro and Wilk, 1965) is first
 conducted on the timeseries data. If the data is normally distributed, which is rare for ModE timeseries,
 then percentiles are calculated from the mean and standard deviation of the timeseries using the *qnorm()*
 480 function from the *stats* package. If the distribution is non-normal, ClimeApp instead finds the value
 corresponding to the quantile matching the users selection (i.e. for the 0.95 percentile, it returns values
 that 5% of all values are above/below), using the *quantile()* function from the *stats* package.

Appendix 3: Current limitations and future development

The data available in ClimeApp is currently limited to ModE-RA, ModE-Sim and ModE-RAclim, and only four
 485 variables within those datasets. There are limited options for users to upload and process other data, but the
 format of this data is currently restricted to annual time series. The web app’s ability to host a large number of
 users at the same time is also constrained by current processing power. To address some of these issues and to
 add further functionality to ClimeApp, a four stage development plan has been devised:

1. Facilitation	<ul style="list-style-type: none"> • Interactive map to look at ModE-RA sources as an access point for the ModE-RA feedback archive, a database detailing each used source • Several video tutorials to facilitate the use of ClimeApp • Obtain funding and increase available processing power to reduce loading times and facilitate multiple simultaneous users
2. Optimization	<ul style="list-style-type: none"> • Optimized and stream-lined plotting (e.g. various projections for global maps, pacific centred plotting) • More options for customization (e.g. for regression and correlation) • Ability to export georeferenced raster and vector files for usage in GIS
3. Implementation	<ul style="list-style-type: none"> • Access to individual ModE-Sim ensemble members for more detailed study • Access to more variables from ModE-RA, such as wind speed and direction • Possibility for users to upload their own georeferenced data for purposes of plotting, averaging and correlation/regression against the ModE-RA data
4. Cooperation	<ul style="list-style-type: none"> • API for other web-based research environments such as nodegoat (van Bree and Kessels, 2013)

490 Appendix 4: Experience documentation, outreach and feedback

ClimeApp underwent extensive internal and external testing before being deployed for general use. The beta
 version was introduced during a summer school in summer 2023 (Huhtamaa and Hibberts, 2024) and later
 trialled at several workshops and conferences, with both historians and climatologists. The initial user
 interactions and resulting valuable feedback included (but was not limited to): The addition of help texts to
 495 explain the different functions and options; the implementation of an interactive map of all ModE-RA sources to
 allow detailed source evaluation; faster data processing; more options for composite analysis; the addition of
 reference maps to see the reference and absolute values for anomalies; customization options for plots; and

facility for multiple simultaneous users. This feedback was documented and either implemented or road mapped for future development (see Appendix 3).

500

The current version of ClimeApp has already been included in the curriculum of two courses at the University of Bern: Brönnimann, 2023, Climatology III (Climate variability and change); and Huhtamaa, 2023, Climate and Society in History. It will also be included in further courses at the University of Bern from 2024 onwards including a three session workshop introducing ClimeApp and its applications.

505

The full application was launched in early 2024 and presented at the EGU conference in Vienna and the Climate of the Past and Societal Responses to Environmental Changes conference in Bern.

ClimeApp has its own feedback and suggestion email address (climeapp.hist@unibe.ch), presented on the *Welcome* page, where users can report any issues and suggest improvements. These are then considered by the developers before being added to the application's Trello page (<https://trello.com/b/3hKu3RIL/climeapp-development>), where users to track their suggestions and see what we are currently working on. These contributions are vital for making ClimeApp as useful and user-friendly as possible.

510

Code availability

The essential code of ClimeApp – *app.R* and *helpers.R* – is available on the projects GitHub page.

515 https://github.com/ClimeApp/ClimeApp_development

Data availability

The Mode-RA database can be downloaded at: https://www.palaeo-ra.unibe.ch/data_access/

Author contributions

NB and RW conceptualized, created and developed ClimeApp as part of their respective PhD projects. They conducted research and analysed data for the case study, produced all graphs and figures and contributed equally to the manuscript. NW co-developed ClimeApp, is responsible for the mail-feedback and manages the Trello page. JF and RH wrote the SD ratio function used in ClimeApp, set up the university server and gave technical advice on the ModE data and contributed to the writing of the manuscript. HH and SB encouraged and promoted the creation of the application and helped with the scientific outreach for the project. All authors discussed the project and its results, helped with the quality control of the final product and provided comments on the manuscript.

525

Competing Interests

The authors declare that they have no conflict of interest.

Disclaimer

530 Acknowledgements

The ECHAM6 simulations (ModE-Sim) were performed at the Swiss Supercomputer Centre (CSCS) and the University of Bern Linux Cluster (UBeLix).

Luc Billaud: Graphics designer and creator of the ClimeApp logo.

Financial Support

535 NB, RW, NW and HH were supported by the Swiss State Secretariat for Education, Research and Innovation (contract number MB22.00030) and Swiss National Science Foundation (grant number PZ00P1_201953). JF, RH and SB received funding from the European Union's Horizon 2020 (grant number 787574).

References

- Auchmann, R., Arfeuille, F., Wegmann, M., Franke, J., Barriendos, M., Prohom, M., Sanchez-Lorenzo, A.,
540 Bhend, J., Wild, M., Folini, D., Štěpánek, P., and Brönnimann, S.: Impact of volcanic stratospheric aerosols on diurnal temperature range in Europe over the past 200 years: Observations versus model simulations, *JGR Atmos.*, 118, 9064–9077, <https://doi.org/10.1002/jgrd.50759>, 2013.
- Brönnimann, S.: *Climatic Changes Since 1700*, 55, Springer, Cham, 2015.
- Brönnimann, S.: Infopage for Course - Climatology III (Climate variability and change), University of Bern,
545 https://www.climatestudies.unibe.ch/courses/courseinfo/index_eng.html?id=54, last access: 26 January 2024.
- Brönnimann, S. and Krämer, D.: Tambora and the "Year Without a Summer" of 1816. A perspective on earth and human systems science, *Geographica Bernensia*, 90, Geographica Bernensia, 2016.
- Chang, W., Cheng, J., Allaire, J. J., Sievert, C., Schloerke, B., Xie, Y., Allen, J., McPherson, J., Dipert, A., and
550 Borges, B.: shiny: Web Application Framework for R [code], <https://doi.org/10.32614/CRAN.package.shiny>, 2024.
- Degroot, D., Anchukaitis, K., Bauch, M., Burnham, J., Carnegie, F., Cui, J., Luna, K. de, Guzowski, P., Hambrecht, G., Huhtamaa, H., and others: Towards a rigorous understanding of societal responses to climate change, *Nature*, 591, 539–550, <https://doi.org/10.1038/s41586-021-03190-2>, 2021 .
- 555 Fajardo, J., Corcoran, D., Roehrdanz, P. R., Hannah, L., and Marquet, P. A.: GCM compareR: A web application to assess differences and assist in the selection of general circulation models for climate change research, *Methods Ecol. Evol.*, 11, 656–663, <https://doi.org/10.1111/2041-210X.13360>, 2020.
- Gebauer, J. E., Adler, J., and DGKL working group "Digital Competence": Using Shiny apps for statistical analyses and laboratory workflows, *J. Lab. Med.*, 47, 149–153, <https://doi.org/10.1515/labmed-2023-0020>,
560 2023.
- Haas-Zumbühl, F.: Die Kernpreise und Brotpreise in Luzern von 1601 bis 1900: Beilage Nr. 16, *Zeitschrift für schweizerische Statistik*, 39, 369–374, 1903.

- Hand, R., Samakinwa, E., Lipfert, L., and Brönnimann, S.: ModE-Sim – a medium-sized atmospheric general circulation model (AGCM) ensemble to study climate variability during the modern era (1420 to 2009), *Geosci. Model Dev.*, 16, 4853–4866, <https://doi.org/10.5194/gmd-16-4853-2023>, 2023.
- 565 Huhtamaa, H.: Infopage for Course - Climate and Society in History, University of Bern, https://www.climatestudies.unibe.ch/courses/courseinfo/index_eng.html?id=166, last access: 26 January 2024.
- Huhtamaa, H. and Hibberts, A.: Young researchers explore climate histories, *PAGES Mag.*, 32, 54, <https://doi.org/10.22498/pages.32.1.54>, 2024.
- 570 Kandlbauer, J. and Sparks, R. S. J.: New estimates of the 1815 Tambora eruption volume, *J. Volcanol. Geotherm. Res.*, 286, 93–100, <https://doi.org/10.1016/j.jvolgeores.2014.08.020>, 2014.
- Krämer, D.: *Menschen grasten nun mit dem Vieh": Die letzte grosse Hungerkrise der Schweiz 1816/17*, Wirtschafts-, Sozial- und Umweltgeschichte, 4, Schwabe Verlag, 2015.
- 575 Lamb, H. H.: *Climate, history and the modern world*, Second Edition, Routledge, 2002.
- Ljungqvist, F. C., Thejll, P., Christiansen, B., Seim, A., Hartl, C., and Esper, J.: The significance of climate variability on early modern European grain prices, *Climetrica*, 16, 29–77, <https://doi.org/10.1007/s11698-021-00224-7>, 2022.
- Marti, J. and Ernst, G. G. J.: *Volcanoes and the Environment*, Cambridge University Press, 2009.
- 580 Möller, M., Boutarfa, L., and Strassemeyer, J.: PhenoWin-An R Shiny application for visualization and extraction of phenological windows in Germany, *Comput. Electron. Agric.*, 175, 1–6, <https://doi.org/10.1016/j.compag.2020.105534>, 2020.
- Nominatim: Open-source geocoding with OpenStreetMap data, <https://nominatim.org/>, last access: 26 January 2024.
- 585 Pfister, C. and Wanner, H.: *Climate and Society in Europe: The Last Thousand Years*, Haupt Verlag, 2021.
- Pfister, C., Rohr, C., and Jover, A.: Euro-Climhist: eine Datenplattform der Universität Bern zur Witterungs-, Klimaund Katastrophengeschicht, *Wasser Energie Luft*, 109, 45–48, <https://doi.org/10.5169/seals-941608>, 2017.
- Posit Team: *RStudio: Integrated Development Environment for R* [code], PBC, Boston, MA, 2024.
- 590 R Core Team: *R: A Language and Environment for Statistical Computing* [code], Vienna, Austria, 2022.
- Raible, C. C., Brönnimann, S., Auchmann, R., Brohan, P., Frölicher, T. L., Graf, H.-F., Jones, P., Luterbacher, J., Muthers, S., Neukom, R., and others: Tambora 1815 as a test case for high impact volcanic eruptions: Earth system effects, *WIREs Clim. Change*, 7, 569–589, <https://doi.org/10.1002/wcc.407>, 2016.
- Riemann, D., Glaser, R., Kahle, M., and Vogt, S.: The CRE tambora. org-new data and tools for collaborative research in climate and environmental history, *Geosci. Data J.*, 2, 63–77, <https://doi.org/10.1002/gdj3.30>, 2015.
- 595 Shapiro, S. S. and Wilk, M. B.: An analysis of variance test for normality (complete samples), *Biometrika*, 52, 591–611, <https://doi.org/10.2307/2333709>, 1965.
- Sousa, J. A. N. P. de: *Exploratory Analysis of Meteorological Data*, Mestrado Integrado, Department of Computer Science, University of Porto, 2019.
- 600 Stevens, B., Giorgetta, M., Esch, M., Mauritsen, T., Crueger, T., Rast, S., Salzmann, M., Schmidt, H., Bader, J., Block, K., and others: Atmospheric component of the MPI-M earth system model: ECHAM6, *J. Adv. Model. Earth Syst.*, 5, 146–172, <https://doi.org/10.1002/jame.20015>, 2013.

- Tennekes, M.: tmaptools: Thematic Map Tools [code], <https://doi.org/10.32614/CRAN.package.tmaptools>, 2021.
- 605 Toohey, M. and Sigl, M.: Volcanic stratospheric sulfur injections and aerosol optical depth from 500 BCE to 1900 CE, *Earth Syst. Sci. Data*, 9, 809–831, <https://doi.org/10.5194/essd-9-809-2017>, 2017.
- Trouet, V. and van Oldenborgh, G. J.: KNMI Climate Explorer: A Web-Based Research Tool for High-Resolution Paleoclimatology, *Tree-Ring Res.*, 69, 3–13, <https://doi.org/10.3959/1536-1098-69.1.3>, 2013.
- Valler, V., Franke, J., Brugnara, Y., Samakinwa, E., Hand, R., Lundstad, E., Burgdorf, A.-M., Lipfert, L.,
610 Friedman, A. R., and Brönnimann, S.: ModE-RA: a global monthly paleo-reanalysis of the modern era 1421 to 2008, *Sci. Data*, 11, 1–19, <https://doi.org/10.1038/s41597-023-02733-8>, 2024.
- van Bree, P. and Kessels, G.: nodegoat: a web-based data management, network analysis & visualisation environment, LAB1100, <https://nodegoat.net/>, last access: 26 January 2024, 2013.
- Wegmann, M., Brönnimann, S., Bhend, J., Franke, J., Folini, D., Wild, M., and Luterbacher, J.: Volcanic
615 Influence on European Summer Precipitation through Monsoons: Possible Cause for “Years without Summer”, *J. Climate*, 27, 3683–3691, <https://doi.org/10.1175/JCLI-D-13-00524.1>, 2014.
- White, S., Pei, Q., Kleemann, K., Dolák, L., Huhtamaa, H., and Camenisch, C.: New perspectives on historical climatology, *WIREs Clim. Change*, 14, 1–17, <https://doi.org/10.1002/wcc.808>, 2023.
- Wood, G. D.: *Tambora: The Eruption that Changed the World*, Princeton University Press, 2014.

620