

The Co-benefits of a Low-Carbon Future on PM_{2.5} and O₃ Air Pollution in Europe

Connor J. Clayton¹, Daniel R. Marsh², Steven T. Turnock^{3,4}, Ailish M. Graham¹, Kirsty J. Pringle⁵, Carly L. Reddington¹, Rajesh Kumar⁶, and James B. McQuaid¹

¹. University of Leeds, Institute of Climate and Atmospheric Science, Woodhouse, Leeds, LS2 9JT

². University of Leeds, School of Physics and Astronomy, Woodhouse, Leeds, LS2 9JT

³. Met Office Hadley Centre, Fitzroy Rd, Exeter EX1 3PB

⁴. University of Leeds, Met Office Strategic (LUMOS) Research Group, Woodhouse, Leeds, LS2 9JT

⁵. University of Edinburgh, Sustainable Software Institute, Old College, South Bridge, Edinburgh EH8 9YL

⁶. National Center for Atmospheric Research, 1850 Table Mesa Dr, Boulder, CO 80305, USA

Correspondence: Connor J. Clayton (eecjc@leeds.ac.uk)

Abstract. There is considerable academic interest in the potential for air quality improvement as a co-benefit of climate change mitigation. Few studies use regional air quality models for simulating future co-benefits, but many use global chemistry-climate model output. Using regional atmospheric chemistry could provide a better representation of air quality changes than global chemistry-climate models, especially by improving the representation of elevated urban concentrations. We use a detailed regional atmospheric chemistry model (WRF-Chem v 4.2) to model European air quality in 2050 compared to 2014 following three climate change mitigation scenarios. We represent different climate futures by using air pollutant emissions and chemical boundary conditions (from CESM2-WACCM output) for three Shared Socioeconomic Pathways (SSP1-2.6, SSP2-4.5, SSP3-7.0; a high, medium and low mitigation pathway).

10

We find that in 2050, following SSP1-2.6, mean population-weighted PM_{2.5} concentrations across European countries reduces by 52% compared to 2014. Whilst under SSP2-4.5, this average reduction is 34%. The smallest average reduction was 18% by following SSP3-7.0. Maximum 6-monthly-mean daily-maximum 8 h (6mDM8h) ozone (O₃) is reduced across Europe by 15% following SSP1-2.6, and 3% following SSP2-4.5, but increases by 13% following SSP3-7.0. This demonstrates clear co-benefits of climate mitigation. The additional resolution allows us to analyse regional differences and identify key sectors. We find that mitigation of agricultural emissions will be key for attaining meaningful co-benefits of mitigation policies, evidenced by the importance of changes in NO₃ aerosol mass to determining future PM_{2.5} air quality and changes in CH₄ emissions to future O₃ air quality.

15

20 1 Introduction

Air pollution is a major public health issue worldwide. The health impacts are usually attributed to two air pollutants - PM_{2.5} (any airborne non-gaseous particle under 2.5 microns in diameter) which can be both a primary or secondary air pollutant and ozone (O₃) which is a secondary pollutant. Primary sources of PM_{2.5} include a range of natural and anthropogenic sources. As a secondary pollutant, it can be formed from emissions of species such as ammonia (NH₃), sulfur dioxide (SO₂) and nitrogen oxides (NO_x). Tropospheric O₃ is a secondary pollutant formed by photochemical reactions involving volatile organic compounds (VOCs) including methane (CH₄), nitrogen oxides (NO_x) and carbon monoxide (CO) in the presence of sunlight. Air pollution has consequences wider than just mortality, such as economic cost (Vandyck et al. 2020) and reduced crop yields (Lobell et al. 2022). Both PM_{2.5} and O₃ are linked with climate change; many sources of primary PM_{2.5} and O₃ precursors are also sources of long-lived greenhouse gases. Additionally, air pollutants themselves have an impact on climate forcing through several pathways including directly affecting the radiative balance of the atmosphere, modifying the albedo of clouds and glaciers and increasing cloud lifetime. (Von Schneidmesser et al. 2020; Peace et al. 2020).

Exposure to air pollution contributes to about 6.7 million deaths per year (World Health Organisation, 2023), 4.2 million of which are from ambient outdoor air pollution. In Europe, PM_{2.5} is responsible for an estimated 368,000 deaths per year (Juginovic et al. 2021) and the annual mean mortality rate from air pollution in Europe of 133 deaths per 100,000 people exceeds the global mean of 120 deaths per 100,000 (Lelieveld et al. 2020). Additionally, the European Environment Agency (2023) reported that in 2022, 96% of Europe's urban population was exposed to PM_{2.5} concentrations above the World Health Organisation's guideline value of 5 µg/m³ (World Health Organisation, 2021).

Improving air quality in Europe is feasible: primary air pollution responds quickly to air pollutant emissions reductions, potentially resulting in lower population exposure. Notably, some secondary air pollutants such as O₃ can worsen depending on emissions reductions in some circumstances, but reducing emissions largely leads to an overall air quality benefit. Due to reductions in anthropogenic emissions of air pollutants, PM_{2.5} air quality in most of Europe has improved over the past half-century; between 1960 and 2009, population-weighted PM_{2.5} concentrations in the European Union decreased by 55.3% (Butt et al. 2017). Similarly, the responsiveness of air pollution to emissions changes was demonstrated by the changes in PM_{2.5} air quality from national to global scales during the COVID-19 pandemic (Jephcote et al. 2021; Venter et al. 2021; Saha et al. 2022; Putaud et al. 2023). The speed of this response to changes in emissions indicates that considerable improvements in air quality can be achieved when air pollutant emissions are reduced. Conversely, O₃ concentrations in Europe have increased in the latter half of the 20th century and early 21st century (Turnock et al. 2020) despite considerable reductions in local, anthropogenic O₃ precursor emissions. This is potentially due to increased intercontinental transport of O₃ precursors (Guerreiro et al. 2014). Despite improving trends in PM_{2.5}, O₃ concentrations may increase due to reduced NO_x emissions causing reduced titration of O₃ (Miyazaki et al.2021). A different approach may therefore be required to reduce exposure to O₃.

Greenhouse gas mitigation policies may also result in lower air pollution emissions and subsequent improved air quality (Turnock et al. 2020; Vandyck et al. 2020). This co-benefit is often suggested as a motivator to encourage faster and stronger climate mitigation from policymakers, including at the regional level, because it turns the concept of climate change mitigation from a diffuse, global-scale requirement to something that can provide measurable, near-term local benefits (von Schneidemesser et al. 2020). Existing research suggests that air quality co-benefits of climate mitigation could occur in Europe. For example, Turnock et al. (2020) find that across Europe, PM_{2.5} concentrations would decrease by the middle of this century for a range of future scenarios and be stronger in scenarios with greater mitigation. Reddington et al. (2023) find that reductions in PM_{2.5} across Europe following a sustainable scenario (SSP1-1.9) could improve health across the continent. Fenech et al. (2021) find similar focusing on the UK. This may not be true for all pollutants and all scenarios. Findings differ for O₃, for which Turnock et al. (2020) project an increase in Europe following scenarios with limited climate change mitigation and Fenech et al. (2021) project an increase compared to the present in all scenarios.

The interactions between air quality and climate change mitigation policies are complicated and non-linear (Von Schneidemesser et al. 2015). Surface level O₃, for example, may worsen following NO_x reductions, as discussed previously (Miyazaki et al. 2021). Complexity is also added by climatic impacts of surface level O₃ (Archibald et al. 2020) and secondary organic aerosol (Scott et al. 2018; Raes et al. 2010). Climate change will also affect the prevailing meteorological conditions, and impact the dispersion of air pollutants, thereby affecting human exposure (Graham et al. 2020). It is also not a given that all climate mitigation strategies will reduce emissions of primary air pollutants as it depends on the mitigation strategy used. Modelling is therefore needed to understand how climate change mitigation and air quality might interact, considering differing strategies for climate mitigation.

Previous research into the linkages between climate change and air quality largely uses the air pollutant emissions associated with CMIP5 as model input (e.g. Silva et al. 2016; Kumar et al. 2018; Fenech et al. 2021), which are linked with the Representative Concentration Pathways (RCPs) (van Vuuren et al. 2011). RCPs are pathways of greenhouse gas concentrations over the 21st Century that result in different radiative forcing endpoints in 2100. Some more recent research (e.g. Rao et al. 2017; Turnock et al. 2020; Reddington et al. 2023) uses the CMIP6 (the successor to CMIP5) emissions, which work with the Shared Socioeconomic Pathways (SSPs) (O'Neill et al. 2017). SSPs expand the range of pathways and also provide different narratives of socioeconomic development, meaning they factor in the role of socioeconomic development in more detail than the RCPs. They also further expand on the link between pollution and climate; describing how air pollution control progresses following the narratives, which is then fed into the air pollutant emissions used in CMIP6 (Rao et al. 2017) and so the SSPs used in CMIP6 will provide a better assessment (Coelho et al. 2023). Using scenarios for research this way does have disadvantages: the RCPs and SSPs are optimised for climate modelling, not air quality modelling and over-reliance on them in the literature may reduce the use of more specific scenarios.

The difficulty of modelling air quality and climate simultaneously makes modelling air quality/ climate mitigation co-benefits challenging. Many studies using the SSPs use the output from global climate models and/or Earth system models (e.g. Turnock et al. 2020, Allen et al. 2020), or "reduced form" models that generalise over large regions (Rao et al. 2017).
90 These types of models may have less detailed chemistry schemes than regional air quality models. They also tend to have a coarser horizontal resolution than regional air quality models. This is important for air quality research to simulate chemical processes that impact on air pollutants at local and urban scales (Adedeji et al. 2020, Fenech et al. 2018, Goto et al. 2016). Despite this, global chemistry-climate models have tended to be used for future projections of air quality due to prohibitive computational requirements for running multi-decadal simulations with regional air quality models. Some studies using CMIP6
95 output (Turnock et al. 2023; Reddington et al. 2023) are making progress in improving the representation and resolution of present-day air quality by combining CMIP6 output with observational and reanalysis data, however, the approaches taken by these studies still use a coarser grid for future simulations.

Some regions are over-represented in regional air quality and climate mitigation co-benefits studies, notably China and India
100 (Von Schneidemesser et al. 2020). Examples include Kumar et al. (2018) and Chowdhury et al. (2020) for India and Cheng et al. (2021) and Conibear et al. (2022) for China. We chose to focus our domain on Europe as it is an under-represented region in the literature. Although studies that focus on Europe or subregions of European countries exist, they largely use CMIP5 emissions instead of CMIP6 (e.g. Fenech et al. 2021 and Sa et al. 2016) or have the primary focus of quantifying the impacts of climate change itself on air quality as opposed to emissions change (Tainio et al. 2013; Tarin-Carrasco et al. 2019).

105

2 Method

Here, we explore the potential mid-century air quality impacts in Europe following the emissions changes from three up-to-date SSPs (SSP1-2.6, SSP2-4.5, and SSP3-7.0) using a state-of-the-art regional atmospheric chemistry model. This aims to help us understand the implications of these updated emissions changes on a sub-regional scale in the European domain.

110 2.1 Model Description

We use the Weather Research and Forecasting coupled with Chemistry model version 4.2. (WRF-Chem). This is an Eulerian, grid-based atmospheric chemistry model. Grell et al. (2005) provide a general model description. We use WRF-Chem at 30 km horizontal resolution with 38 vertical levels up to 50 hPa and a domain of 100*100 grid boxes ranging from latitudes 32° North to 60° North and longitudes from 22° West to 30° East in the North of the domain, narrowing to 13° West to 19° East in
115 the Mediterranean (Supplementary Figure A1). Note that the model domain does not cover all of Europe and for the purpose of this study we define "Europe" as 13 countries: Germany, the UK, France, Spain, Italy, the Netherlands, the Czech Republic, Hungary, Poland, Slovakia, Ireland, Slovenia and Portugal. These countries have a combined population of approximately 380 million and represent a range of sources of primary air pollutants and environmental conditions that will affect air quality. We

chose the model, resolution and domain in order to capture the changes at the regional and country level; while the resolution
120 is not fine enough to fully represent the chemistry at city scale, it is sufficient to demonstrate urban peaks or elevated concentrations of air pollutants near some power stations. We chose Europe as a domain in part due to the fewer studies in this area, but also due to the concentration of national and regional administrative areas and thereby improve the policy-relevance of our work. A 30km horizontal resolution allows a compromise between global and local scale models by allowing us to increase our domain size to cover most of Europe while also representing air quality in smaller regions more realistically than global models.

125

We simulate a present-day air quality control with gridded 2014 emissions used in CMIP6 (Hoesly et al. 2018) and simulations with anthropogenic emissions representing 2050 for each of SSP1-2.6, SSP2-4.5 and SSP3-7.0 (Feng et al. 2020). These emissions are created based upon the SSPs, meaning they include similar assumptions, for example the projected land use changes factor into the emissions. The model parameters are shown in Table 1. For all scenarios the meteorology was
130 fixed at 2014 conditions using meteorological initial and boundary conditions from ECMWF ERA5 (Hersbach et al. 2020). The aerosol-radiation feedback is switched on in the model, however as the simulations are frequently nudged to meteorology, there is no meaningful meteorological difference between the scenarios. It is well established that in Europe, the impact of emissions changes on future $PM_{2.5}$ air quality is likely to far eclipse the impact of climate change (Colette et al. 2013; Chemel et al. 2014; Doherty et al. 2017). Additionally, in Europe, even O_3 pollution may not be sensitive to changes in climate (Zanis
135 et al. 2022), thus we are confident that not factoring in meteorological changes is a worthwhile trade-off to allow us to use a detailed model in WRF-Chem at a relatively fine resolution.

2014 was chosen as this is the most recent year of historical emissions data from the emissions inventory used in CMIP6. We also use CMIP6 output from CESM2-WACCM (Danabasoglu, 2019) simulations to provide initial and chemical boundary
140 conditions. To simulate chemistry, a scheme described by Hodzic & Knote (2014) is used that combines MOZART-4 gas phase chemistry, which includes 85 gas-phase species, 157 gas phase reactions and 39 photolysis reactions (this scheme and the included reactions are provided by Emmons et al. 2010) with the MOSAIC aerosol chemistry scheme described initially by Zaveri et al. (2008). This provides detailed chemistry for a range of aerosol species including nitrate from ammonium nitrate (NO_3), sulfate (SO_4), organic carbon (OC), black carbon (BC), ammonium from other sources (NH_4), sodium and chloride, all
145 in four size bins up to 10 microns in diameter. The combined scheme described by Hodzic & Knote (2014) enhances these by including aqueous chemistry, improved treatment of monoterpenes and hydrocarbons, and updating the mechanism calculating secondary organic aerosols. $PM_{2.5}$ in the model is the sum of the total dry aerosol mass in 3 smallest size bins (up to 2.5 microns diameter) of the above aerosol components and "other inorganics" (OIN) which largely consists of dust. The full range of model inputs are shown in Table 1.

150

Parameter	Scheme Used	Source
Chemical Scheme	MOZART-MOSAIC 4-bin Aerosol w/Aqueous Chemistry	Hodzic and Knote (2014)
Biogenic Emissions	MEGAN v2.0.4	Guenther et al.(2006)
Fire Emissions	FINNv1.5	Wiedinmyer et al. (2010)
Natural Dust Emissions	GOCART	Chin et al. (2000), Ginoux et al. (2001)
Meteorological Boundary Conditions	ECMWF ERA5	Hersbach et al. (2020)
Chemical Boundary Conditions	CESM2-WACCM CMIP6 Simulations	Danabasoglu (2019)

Table 1. Model parameters used and sources.

2.2 Emissions associated with CMIP6

We simulated 3 different emissions scenarios: SSP1-2.6 represents a scenario with accelerated mitigation of greenhouse gases and sustainable societal development. SSP2-4.5 is a "middle of the road" scenario in which the trajectory of greenhouse gas mitigation does not accelerate or decelerate strongly and there are no great changes in uptake of sustainable behaviours. SSP3-7.0 is a scenario in which regional rivalry hampers greenhouse gas mitigation and sustainable development. The assumptions in air pollutant controls mirror the trajectories of greenhouse gas emissions in each scenario, with some non-linearity or deviation in particular species to match the scenario narrative. These are explained by Rao et al. (2017); In summary, SSP1 assumes an acceleration in pollution control progress, SSP3 a deceleration, and SSP2 neither a notable acceleration or deceleration from present-day controls. 1 shows how the emissions of key species change in future scenarios compared to the present day, demonstrating how the narrative scenarios translate to emissions data. Here the non-methane VOCs are grouped. We see that SSP1-2.6 has considerably lower emissions of all pollutant species compared to the present day. SSP2-4.5 and SSP3-7.0 have lower reductions in emissions overall, but notably differing trajectories for NH₃ emissions compared to SSP1-2.6, which increase compared to the present following both scenarios. The NH₃ emissions increases are largest in rural regions, including most of France and Spain and Northern Poland. Both scenarios have NH₃ emissions decreases in Paris, The Rhine-Ruhr region and coastal parts of Spain and Italy. There are differences between SSP3-7.0 and SSP2-4.5 for CH₄ emissions which are mitigated following SSP2-4.5 but worsen following SSP3-7.0 and CO, which is heavily mitigated following SSP1-2.6 and SS2-4.5 but reduces only minimally following SSP3-7.0.

Table 2 shows the European total emissions of air pollutants assumed for the present-day scenario in 2014 and in the scenarios we perform simulations for 2050, taken from the input emissions files from Hoesly et al. (2018) for the present-day and Feng et al. (2020) for the future scenarios. All the emissions files were at 50 km horizontal resolution. The emissions were then regridded to 30km using a standard bilinear regridding method.




Figure 1. Relative change of model domain average annual emissions from 2014 to each of the future scenarios in 2050.

Pollutant	2014 emissions	SSP1-2.6 (2050)	SSP2-4.5 (2050)	SSP3-7.0 (2050)
Black Carbon	0.37	0.07	0.11	0.24
CO	24	9	13	23.7
NH ₃	4.9	4.1	5.8	5.9
NO _x	12.9	3.4	8.9	9
SO ₂	4.56	1.2	2	2.9
Organic Carbon	1.24	0.28	0.55	0.8
NMVOC	8.1	3.3	6.3	6.7
CH ₄	22.5	9.7	13	24.5

Table 2. European (domain defined above) total emissions of air pollutants in 2014 from CMIP6 and 2050 from ScenarioMIP SSP1-2.6, SSP2-4.5 and SSP3-7.0, all expressed in Mt/yr

As the CMIP6 emissions do not include a component of inorganic PM_{2.5} or PM₁₀ directly emitted as anthropogenic dust (as
 175 required for WRF-Chem) we created files to simulate this fraction using linear regression based on the anthropogenic carbon

monoxide emissions using the relationship between EDGAR-HTAPv2 carbon monoxide and PM_{2.5}/ PM₁₀ (which represent anthropogenic dust) emissions (Janssens-Maenhout et al. 2015). This methodology has been used previously by Kumar et al. (2018) and Wu et al. (2019). These input files are referred to in the rest of the text as anthropogenic dust emissions. To generate emissions of individual non-methane VOC (NMVOC) chemical species we use scaling factors derived from ratio of individual
180 NMVOCs to total NMVOCs in the EDGAR-HTAPv2 emissions inventory (Huang et al. 2017). This provided a greater spectrum of speciated VOCs than the scaling factors used by Hoesly et al. (2018) and Feng et al. (2020).

2.3 Model Output

We use hourly output from each the year-long WRF-Chem simulations for each species (O₃, CO, CH₄, SO₂, NO₂, nitrogen
185 oxide (NO), NH₃ and PM_{2.5} dry aerosol mass, and separate files for the individual PM_{2.5} components, NO₃, NH₄, SO₄, OC, BC, sodium and chloride). All air pollutant output was analysed only at surface level. For some analysis, we weighted PM_{2.5} and O₃ by population using the formula outlined in Abdul Shakor et al. (2020). We used time-varying gridded population projections for each SSP from Jones & O'Neill (2016). To represent the present-day population, the SSP2 population projection for 2020 was used. This was to allow for a consistent source for all population data as there was no data for 2014.

190

2.4 Model Validation

The present-day simulation for 2014 was validated against PM_{2.5}, O₃ and other aerosol component observations (as detailed in Table 3) from the European Modelling and Evaluation Programme (EMEP) as this features sites for a range of species across Europe. Sites with an altitude above 1 km were excluded as measurement and modelling of air quality in complex terrain is
195 challenging and frequently less accurate (Giovannini et al. 2020). We used spatial linear interpolation to extract data from our gridded model output to compare to the locations of the observation sites. The sites used are shown on Supplementary figure A1.

Comparisons of modelled and observed O₃ and PM_{2.5} are shown in Figures 2 and 3. Figure 2 shows observed and simulated monthly mean PM_{2.5} (a) and O₃ (b) colour-coded by the observation station. Simulated monthly O₃ data shows a slight under-
200 estimation (mean absolute bias compared to observations was -3.40 µg/m³) overall. This underestimation is generally larger in the observation sites in Germany (Schmucke, Neuglobsow and Waldhof), however, an overestimation is seen in sites closer to the Mediterranean (Saint-Nazaire, Barcarrota). There were no O₃ observations available for Vredepeel, Cabauw Wielsekade, Guipry and Melpitz. PM_{2.5} showed an overestimation compared to observations, with a mean absolute bias of 7.98 µg/m³. The sites with the largest overestimation were Cabauw Wielsekade, Harwell, and Vredepeel. The overestimation was smaller
205 in sites such as San Pablo de los Montes, Barcarrota and Penausende. The Eastern European observation sites all showed a bias lower than the average. As all observation sites use different monitoring technology and all overestimate PM_{2.5}, we expect this to be an artifact of the model.




Figure 2. Comparison of modelled $\text{PM}_{2.5}$ (a) and O_3 (b) to ground-based Observations. Monthly means are used for both species and the units are $\mu\text{g}/\text{m}^3$. Model data is interpolated from the latitude and longitude coordinates of the observation site. The middle line on each plot represents what the data would look like if observations and model were equal.

Figure 3 is a comparison of monthly mean modelled data compared to observations (averaged over all sites). The model represented the seasonal cycle, with higher $\text{PM}_{2.5}$ in spring and autumn, matching when the emissions peaked. The model $\text{PM}_{2.5}$ overestimation was larger in spring and autumn and smaller in summer. Simulated O_3 showed seasonal biases, The model underestimated in winter and spring but overestimated during summer and autumn.

Turnock et al. (2020) reported an underestimation of $\text{PM}_{2.5}$ compared to observations in Europe in a similar period (2005-2014). This is likely because of the additional emission source of $\text{PM}_{2.5}$ in our simulations and the coarse resolution of Turnock et al. (2020)'s simulations. Conversely, $\text{PM}_{2.5}$ overestimations have been seen in other studies using CMIP6 emissions to drive regional models, such as Cheng et al. (2021) simulating over China, who found that nitrate, sulfate and ammonium $\text{PM}_{2.5}$ were




Figure 3. Seasonal cycle in 2014 of a) $PM_{2.5}$ and b) O_3 comparison of monthly mean modelled data to observations (average of all sites). Number of sites = 26 for $PM_{2.5}$ and 21 for O_3 . Both pollutants are measured in $\mu g/m^3$. The shaded areas represent variation from the mean across sites as $2 \times$ standard deviation.

overestimated compared to observations by 30-60%.

220 Further validation of $PM_{2.5}$ components (comparison of modelled values with ground-based observations) was conducted to
 diagnose the difference between model and observations. These are shown in Table 3. The total bias in $PM_{2.5}$ is greater than the
 combined bias of the individual aerosol species. Although not every observation site measured each species and therefore the
 proportions cannot be assumed to be the same at each site, this implies that a different source may account for much of the bias.
 A large proportion of the bias likely comes from the OIN (dust) component. This agrees with previous research; Im et al. (2015)
 225 find that WRF-Chem setups using GOCART over-produce dust and that the partitioning of dust particulates between $PM_{2.5}$
 and PM_{10} when MOSAIC aerosols does not reflect real conditions over Europe. Similarly, Georgiou et al. (2018) validating
 WRF-Chem over Cyprus show that WRF-Chem simulations with MOSAIC aerosols can result in a significant overestimation
 of $PM_{2.5}$, largely driven by the dust scheme. Additionally, some of the overestimation in the OIN component is likely the result
 of the derived anthropogenic dust emissions as this is calculated from the CO emissions and it would explain the larger overall
 230 $PM_{2.5}$ overestimation in polluted, urban regions.

Pollutant	N (sites)	Model Mean ($\mu\text{g}/\text{m}^3$)	Observations Mean ($\mu\text{g}/\text{m}^3$)	Bias (model-obs) ($\mu\text{g}/\text{m}^3$)	Root Mean Square error
PM _{2.5}	26	17.9	10.1	7.8	9.1
Nitrate (NO ₃)	14	5.8	3.1	2.7	4
Ammonium (NH ₄)	10	1.5	1.2	0.3	0.6
Organic Carbon (OC)	24	3	2.9	0.1	1.5
sulfate (SO ₄)	10	1.3	3.4	-2.1	2.5
Black Carbon	13	0.8	0.4	0.4	0.6
Chloride	13	1.4	0.2	1.2	1.5
Sodium	14	1.9	0.2	1.7	1.9

Table 3. Mean difference between annual mean model output and annual mean ground-based observations for different PM_{2.5} components across monitoring sites in Europe

Table 3 shows that the model overestimates NO₃ aerosol by 2.8 $\mu\text{g}/\text{m}^3$ and underestimates SO₄ aerosol by -2.1 $\mu\text{g}/\text{m}^3$ when compared to the observation sites. Overestimation of NO₃ aerosol matches the findings of other WRF-Chem studies including Cheng et al. (2021) and Balzarini et al. (2015), however, both of these studies found SO₄ overestimation as opposed to underestimation. The NO₃ overestimation may be the result of high NH₃ emissions over much of the year for the emissions used in CMIP6 in comparison to other emissions inventories. When compared to EDGAR-HTAPv3 (Crippa et al. 2023) CMIP6 NH₃ emissions were lower during February, March, and April, but higher the rest of the year. Similarly, the CMIP6 emissions of nitrogen oxides (NO_x) are generally higher than EDGAR-HTAPv3 in urban regions, which may also contribute to the overestimation in NO₃ aerosols.

240

3 Results

3.1 Changes in PM_{2.5}

Table 4 shows the European annual mean PM_{2.5} in the present day and the change from this in the future scenarios. In general, annual mean PM_{2.5} reduced in all future scenarios compared to the present day. The future reduction in European annual mean PM_{2.5} of 38% in SSP1-2.6 was far greater than the 11% following SSP3-7.0. There are differences in the pattern when population-weighting is applied; overall, population exposure to PM_{2.5} increases slightly following SSP3-7.0 despite the domain-wide decrease. This suggests that the majority of the increases in PM_{2.5} are in highly populated areas

Spatially (Figure 4 - shows percentage changes between the present and future scenarios. Supplementary Figure A2 shows the absolute changes), we see greater reductions in PM_{2.5} in urban and industrial regions than the domain average. Both indus-

250

		Change from present-day in $\mu\text{g}/\text{m}^3$ (% in brackets)		
Domain statistic	Present day concentration ($\mu\text{g}/\text{m}^3$)	SSP1-2.6	SSP2-4.5	SSP3-7.0
Minimum	4.9	-1.5 (-31)	-0.61 (-13)	-0.1 (-2)
Mean	19.4	-7.3 (-38)	-4.2 (-21.6)	-2.1 (-10.8)
Maximum	44.4	-21.3 (-48)	-17.6 (-40)	-5.6 (-12.6)
Population-weighted mean	17.5	-8.6 (49)	-5.2 (30)	+0.9 (5)

Table 4. Annual mean $\text{PM}_{2.5}$ whole domain change statistics for each future scenario in 2050 compared to the present day baseline (the left-hand column). For future scenarios, the raw change for each of these is shown in $\mu\text{g}/\text{m}^3$ followed by the percentage change in brackets.

trial (for example the Po Valley, Northern Italy and the Rhine-Ruhr, North-West Germany) and urban regions see strong $\text{PM}_{2.5}$ reductions under SSP1-2.6 and SSP2-4.5 (although these are far larger following SSP1-2.6). Conversely, under SSP3-7.0, only urban regions see considerable $\text{PM}_{2.5}$ reductions. This could be explained by a larger reduction in industrial $\text{PM}_{2.5}$ emissions following SSP2-4.5 than SSP3-7.0 (-56% compared to -47%) and a larger reduction in OC emissions (-72% compared to -65%). Both SSP2-4.5 and SSP3-7.0 have small percentage increases in industrial NO_x and SO_2 emissions and relatively consistent NH_3 emissions when compared to the present day, suggesting that it is not differences in the NO_x and NH_3 ratio that cause the differences between SSP2-4.5 and SSP3-70 in industrial regions.

Additionally, under SSP3-7.0, localised areas of worsening air quality are seen, including around East Yorkshire, UK (worsening up to $4 \mu\text{g}/\text{m}^3$) and Zeeland and South Holland, the Netherlands (worsening up to $2 \mu\text{g}/\text{m}^3$). Some of these localised increases correspond with the locations of major combustion power plants, including Drax (North Yorkshire, UK) and Belchatow (Central Poland). This is because the emissions scenarios assume that power generation emissions increase up to mid-century compared to the present day following SSP3-7.0, they drop following SSP2-4.5 but this is approximately half the reduction that is predicted following SSP1-2.6. All scenarios show slightly worsening $\text{PM}_{2.5}$ air quality of up to $2 \mu\text{g}/\text{m}^3$ near Gijon, on the North coast of Spain. This reflects worsening NO_x emissions in all scenarios at this location.




Figure 4. (a) Modelled annual mean PM_{2.5} ($\mu\text{g}/\text{m}^3$) using 2014 emissions. (b),(c) and (d) are the percentage change from (a) the annual mean PM_{2.5} ($\mu\text{g}/\text{m}^3$) simulated using 2050 emissions for SSP1-2.6, SSP2-4.5 and SSP3-7.0 respectively.

The PM_{2.5} reductions in SSP1-2.6 are larger across most of the domain compared to the other scenarios. This is most notable across Central/Eastern Europe (e.g. Germany, Poland, the Czech Republic, Austria). This is potentially because these regions have a larger proportion of anthropogenic PM_{2.5} sources than natural sources. Smaller improvements are projected in countries such as Portugal and Ireland where natural sources of PM_{2.5} dominate. In the regions where the reductions in SSP1-2.6 compared to SSP2-4.5 and SSP3-7.0 are the largest (such as Central/Eastern Germany), the size of the reduction is likely the result of the combined reduction in NH₃ and NO_x emissions in SSP1-2.6 (domain-wide, -25% and -70%, respectively).

Across the domain, SSP2-4.5 and SSP3-7.0 have increases in NH₃ emissions (both approximately 20%) and reductions in NO_x (both approx -30%) (Figure 1). Where only NO_x is reduced, in some regions (usually NO_x-abundant regions) the increased oxidising capacity of the atmosphere can result in increased formation of secondary organic aerosol, thus limiting the efficacy of emissions reductions. This impact can be mitigated by joined reductions of both NO_x and NH₃ (Clappier et al. 2021). This is supported by the near-universal decreases of over 60% of NO₃ PM_{2.5} following SSP1-2.6, which is the result of a domain-wide reduction of 18% of agricultural NO_x emissions and 7% of agricultural NH₃ emissions. While following SSP2-4.5 and SSP3-7.0, NO₃ PM_{2.5} does decrease universally, however, the reductions are much larger in urban regions than rural. Notably, both of these scenarios have approximately 20% increases in agricultural NH₃ emissions and increases in agricultural NO_x emissions (41% following SSP2-4.5, 10% following SSP3-7.0). This suggests that the difference in agricultural emissions will be a large driver of the extra reductions in PM_{2.5} following SSP1-2.6 and mitigation of emissions in this sector will be key to achieving improved air quality in Europe. It should however be noted, that some rural regions (e.g Western France, Scotland, Wales and most of Spain) do still have overall NH₃ emissions increases following SSP1-2.6. This may explain the smaller reductions in PM_{2.5} in these regions (Supplementary Figures A3 & A4).

Reductions in SO₂ emissions also contribute to the reduced PM_{2.5}. The countries that tend to see the largest decreases in PM_{2.5} concentrations following the future scenarios are Central/European countries. These countries, particularly in urban areas usually have a higher contribution from SO_x aerosols (including SO₄, which will cover much of these aerosols in the chemistry scheme) to PM_{2.5} (Zauli-Sajani et al. 2024). SO_x aerosol is formed by reactions between SO₂, NO_x and NH₃. For example, atmospheric sulfuric acid is formed when SO₂ reacts with OH radicals. The sulfuric acid can then react with NH₃ to form SO₄ particulates (Clappier et al. 2021). Unlike NH₃ and NO_x, there is limited non-linearity of SO₄ PM_{2.5} reductions resulting from mitigating SO₂ emissions (Clappier et al. 2021). This means that they retain efficacy in reducing PM_{2.5} despite trends in NH₃ and NO_x reductions that may work against each other when only one is present. SO₂ emissions reduce in all scenarios by approximately 30% for both SSP2-4.5 and SSP3-7.0 and by approximately 75% following SSP1-2.6 (Figure 1). This may explain why PM_{2.5} reductions are most consistently seen in Central/Eastern Europe across all the future scenarios - here, reductions in SO₂ emissions (see Supplementary Figure A5) lower PM_{2.5} in the sulfate fraction to a greater degree than in Western Europe. In Western Europe, following SSP2-4.5 and SSP3-7.0, the reductions in NO_x in the absence of NH₃ hamper PM_{2.5} reductions. This impact is reduced in Eastern Europe, where NH₃ and NO_x provide a lesser contribution to

PM_{2.5} compared to SO₂, meaning Eastern Europe has larger PM_{2.5} reductions following SSP2-4.5 and SSP3-7.0.

All countries show overall decreases in population-weighted PM_{2.5} (Figure 5). The magnitude of which varies greatly based on the scenario. Following SSP1-2.6, the percentage decrease ranges from 22.7% in Ireland to 68.6% in Hungary. Other countries with decreases in population-weighted mean PM_{2.5} following SSP1-2.6 greater than 50% include Slovenia, Slovakia and Germany. These countries also see the greatest reductions in PM_{2.5} following the other scenarios, for example, the largest reduction following SSP3-7.0 is in Slovenia, at nearly 25%. This may be due to reductions in residential emissions, which see the greatest reduction in the proportion of total emissions in Slovenia and are reduced in all scenarios. This is in keeping with literature that identifies the importance of reducing residential sector emissions to improve PM_{2.5} concentrations in Europe, especially Eastern Europe (Zauli-Sajani et al. 2024) For most countries in the domain, the agricultural sector contributes most to total air pollutant emissions.

In addition to benefiting the least following SSP1-2.6, Ireland benefits the least following SSP2-4.5 with a reduction of 5%. Similar to Ireland, Portugal and Spain do not benefit as much from the emissions changes compared to others. Ireland even shows an increase in population-weighted PM_{2.5} following SSP3-7.0 of nearly 4%, which is also seen in the UK. What this suggests is that the benefits are concentrated in countries where anthropogenic sources dominate PM_{2.5} concentrations in the present day. As coastal island countries, Ireland and the UK likely have a greater proportional quantity of natural sea salt aerosol making up PM_{2.5} - previous literature suggests that sea salt PM_{2.5} can reach up to 300km inland and produce up to 5 $\mu\text{g}/\text{m}^3$ of PM_{2.5} (Manders et al. 2010). Spain and Portugal are likely to have large proportions of natural dust PM_{2.5} due to proximity to North Africa.

Figure 5 also compares the population-weighted mean to the World Health Organisation annual mean PM_{2.5} guideline value of 5 $\mu\text{g}/\text{m}^3$. It suggests that following SSP1-2.6, many countries could see PM_{2.5} exposure reduce below interim target values (guidelines the WHO suggest as targets to aim for before reaching the guideline value), representing a significant potential benefit for human health. Even the emissions reductions from SSP1-2.6 do not result in annual mean population-weighted PM_{2.5} concentrations under this guideline, but when factoring in the average overestimation of PM_{2.5} from our model, it is likely that this is achievable in some locations. Notably, while countries where PM_{2.5} is dominated by natural sources see less improvement, these have among the lowest PM_{2.5} population exposure in the present day. This means that the benefits of the emissions changes are primarily seen in the countries that most need them. The whole domain average also moves below the WHO interim target 1 of 10 $\mu\text{g}/\text{m}^3$ following SSP1-2.6, after a reduction of almost 50%.

Figure 6 shows the seasonal cycle of PM_{2.5} components averaged across the entire model domain. SSP1-2.6 has a much lower contribution of anthropogenic PM_{2.5} than SSP2-4.5 and SSP3-7.0, driven by the emissions reductions shown in Figure 1. Figure 6 also shows that it is the change in these anthropogenic species, particularly in NO₃ and OC, that drive the differences between the future scenarios, with NO₃ alone reducing total PM_{2.5} following SSP1-2.6 by over 5 $\mu\text{g}/\text{m}^3$ throughout much of the




Figure 5. Population-weighted PM_{2.5} in a selection of European countries following different scenarios compared to WHO guideline values. The lines labelled with "IG" are WHO interim guidelines. All Europe refers to a combined population-weighted mean of the 13 countries on the Figure

year. The importance of NO₃ aerosol in the future scenarios to determining total PM_{2.5} implies that NH₃ and NO_x emissions reductions will be key to improving future air quality. All future scenarios show overall reductions in NO_x emissions, which can limit the formation of NO₃ and NH₄ particulates (Pusede et al. 2016), however, it is only SSP1-2.6 that shows a significant reduction in NO₃ aerosol, likely because it is the only scenario where NH₃ emissions reduce compared to the present day.

340 This suggests that agriculture will be a key sector for attaining air quality co-benefits as agriculture is a major source of NH₃ emissions. The reduction in OC concentrations is proportionally far larger under SSP1-2.6 than the other scenarios, potentially

due to the trajectories in power sector OC emissions.




Figure 6. Seasonal cycle of domain average PM_{2.5} over each simulation by component, (a) is the 2014 simulation, (b),(c) and (d) are 2050 from each of SSP1-2.6, SSP2-4.5 and SSP3-7.0 respectively.

Our findings are in agreement with other work in the area, in that we find that air quality co-benefits of climate mitigation are likely for PM_{2.5}. When compared to Fenech et al. (2021) for example (who used CMIP5 emissions and focused on the UK),

we see that both studies project $PM_{2.5}$ that strong mitigation will result in air quality co-benefits for the UK. The reductions we project for the most comparable scenarios (SSP1-2.6/ RCP 2.6) are larger (-7.3 as opposed to $-2.2 \mu g/m^3$ or approximately 38% vs 25%). We see a diverging trend for the most pessimistic scenarios - while they see reductions in $PM_{2.5}$ concentrations following RCP8.5, we see worsening $PM_{2.5}$ following SSP3-7.0.

350

Notably, our model simulated high present-day $PM_{2.5}$ in urban regions (e.g.Paris, Madrid and London) compared to surrounding areas. It also produced elevated $PM_{2.5}$ in heavily industrial regions of Europe such as the Po Valley and the Rhine-Ruhr (Figure 4a). It is the changes in these regions that stand out in the other panels of Figure 4. What this suggests, is that our methodology allows us to better represent changes on a local level than work using climate model output. This can be shown when our work is compared to Turnock et al. (2020) (who used CMIP6 output on a global domain) in Figure 7, which shows the difference in change between the present and 2050 following SSP3-7.0 for both $PM_{2.5}$ and O_3 . Figure 7 shows that we see similar spatial changes excluding different trends in $PM_{2.5}$ across the Iberian Peninsula and most of the British Isles. This comparison shows how the finer spatial resolution allows us to see localised elevated concentrations of pollution, whereas pollutants are distributed more evenly over the coarser resolution of global models. We also see greater improvements in $PM_{2.5}$ overall than Turnock et al. (2020), for example, for SSP1-2.6 our domain improvement of $7 \mu g/m^3$ exceeds by more than double theirs of approximately $3 \mu g/m^3$. This highlights that using air pollutant concentrations from global model simulations may underestimate the extent of future changes in air quality.

365

Geographically the reductions in European $PM_{2.5}$ are lower than other studies in more polluted regions. Cheng et al. (2021) find a reduction in population-weighted mean $PM_{2.5}$ in China between 2020 and 2050 following SSP1-2.6 of between 20 and $25 \mu g/m^3$ (from approximately $42 \mu g/m^3$ in the present-day scenario). However, this reduction is similar to the average relative reduction of 52% across European countries that we find. Studies on future air quality in India also find that scenarios with a greater focus on sustainability result in reductions in surface air pollution (Chowdhury et al. (2020); Kumar et al.2018) although methodological differences make direct comparisons with these studies challenging. What these comparisons suggest is that Europe could see similar relative air quality co-benefits to other regions following future sustainability scenarios.

370

3.2 Changes in O_3

Maximum 6-monthly-mean daily-maximum 8h (6mDM8h) O_3 is marginally higher concentrations in Mediterranean regions, including Italy, Southern Spain and the French Riviera, which get more sunlight (Figure 8 shows percentage changes between the present and future simulations - Supplementary Figure A6 shows the absolute changes). In the simulations using future scenarios, O_3 largely reduces following SSP1-2.6 (mean reduction of approximately 15%), SSP2-4.5 shows variation across the domain; while the mean change is a reduction of approximately 3%, increases are seen in most of England, the Benelux region and North-West Germany. 6mDM8h O_3 increases across most of the domain following SSP3-7.0 (mean increase of approximately 13%). This is not universal; small decreases of up to 5% are seen in most of the Mediterranean regions with

375




Figure 7. a) Percentage change annual mean PM_{2.5} between present day and 2050 following SSP3-7.0 from our simulations with WRF-Chem b) The same from Turnock et al. (2020) using CMIP6 multi-model output. c) and d) Same as the above, but O₃ compared. Note that this is annual mean O₃ and thus, c) differs from panel d) of Figure 10

380 high present-day O₃. Despite the lack of peaks in the present-day simulation, in the future simulations, O₃ pollution does not reduce as much in urban regions as much of the rest of the domain following SSP1-2.6. The same regions show increases following SSP2-4.5 (and some, such as around Barcelona and Naples show increases following SSP1-2.6) where much of the rest of the domain has reductions in surface level O₃ and the increases are higher than surrounding areas following SSP3-7.0.

The differing trends in CH₄, CO and NO_x (Figure 1) between the scenarios may explain the difference in O₃. It is well established in the literature that in urban areas reductions in NO_x emissions can cause increases in surface level O₃, including within Europe (Lee et al.2020; Finch & Palmer 2020). This is due to increased titration of O₃, in which reduced NO concentrations due to the reductions in NO_x emissions result in less destruction of O₃ molecules (Monks et al 2015). This is




Figure 8. 6mDM8h O₃ calculated as the highest 6-month mean of the highest rolling 8-hour O₃ in 24-hour periods in the O₃ output for each scenario. (a) shows this metric for the CMIP6 2014 simulation. (b),(c) and (d) show the percentage change from this for SSP1-2.6, SSP2-4.5 and SSP3-7.0 respectively.

supported by our results as the largest O₃ increases are seen in the urban areas with the largest reductions in NO₂, such as Naples, Italy and Barcelona, Spain (Supplementary Figure A7. This effect is likely to be amplified in autumn and winter, when emissions reduction policies are likely to have the most impact. Supplementary Figures A9-A12 support this, showing that the

390

		Change from present-day in $\mu\text{g}/\text{m}^3$ (% in brackets)		
Domain Statistic	Present day concentration ($\mu\text{g}/\text{m}^3$)	SSP1-2.6	SSP2-4.5	SSP3-7.0
Minimum	23.9	+6.4 (+26.7)	+8.3 (+34.7)	+12.3 (+51.5)
Mean	66.2	-9.8 (-15)	+1.7 (+2.6)	+8.7 (+13.1)
Maximum	97.4	-13.3 (-24)	-0.6 (0.6)	+7.6 (+7.8)
Population-weighted mean	52.7	-3 (-7)	+5.3 (+10)	+8.8 (+16.7)

Table 5. Annual mean O_3 whole domain change statistics for each future scenario compared to the present day baseline (the left-hand column). For future scenarios, the absolute change for each of these is shown in $\mu\text{g}/\text{m}^3$ followed by the percentage change in brackets.

O_3 increases are largest during the winter months. This is why the annual mean O_3 concentration changes (Supplementary Figure A8) follow a different pattern to 6mDM8h O_3 .

Reductions in NMVOCs could be having similar effects to the NO_x reductions in some regions. In all scenarios, reductions in NO_x emissions are stronger than NMVOC emissions, however following SSP2-4.5 and SSP3-7.0, the proportional gap is much larger. This suggests that in VOC-limited regimes (which are generally urban areas), where reductions in NO_x are more likely to exacerbate O_3 , we may see a larger O_3 response. Notably, the emissions changes may cause different patterns in NO_x and VOC-limited regimes that may impact on the O_3 response, for example, Liu et al. (2022) find that in Europe following SSP3-7.0, the percentage of VOC-limited areas drops from nearly 80% to 27% in winter and from 37% to under 3% in summer. If our simulations have a similar change in sensitivity this could suggest that a different precursor, such as CH_4 , primarily drives the increases in O_3 following SSP3-7.0, which suggests that agriculture will be a key sector for determining future O_3 pollution.

In SSP1-2.6, this impact of reduced NO_x emissions causing O_3 increases is masked by O_3 increases from another source; likely much stronger decreases in CO and CH_4 emissions than SSP2-4.5 and SSP3-7.0. What this means is that while all scenarios assume increased pollution control, the additional focus given to climate mitigation (e.g. reducing CH_4 emissions) and the more sustainable socioeconomic development in SSP1-2.6 and SSP2-4.5 has potential air quality co-benefits by outweighing any impacts on O_3 from reduced NO_x . O_3 is considerably more impactful on health during peak season due to the high thresholds needed to affect population health on a large scale, thus most parts of Europe will see reduced impact of O_3 pollution following both SSP1-2.6 and SSP2-4.5.

410

We find a considerably higher percentage increase in annual mean O_3 in 2050 following SSP3-7.0 compared to that found for Europe by Turnock et al. (2020) (Figure 7). Once again, the difference in resolution is clear here as we see much larger increases in O_3 around certain urban regions, whereas Turnock et al. (2020) show a smaller trend distributed over larger areas. As with $\text{PM}_{2.5}$, the finer resolution we use could prove valuable, especially for understanding health impacts and trends in urban areas. As shown by the contrasting trends in urban and rural areas for O_3 pollution following some scenarios, being able

415

to represent these changes is valuable.

Compared to studies focusing on regions outside of Europe, our findings are similar to those reported by Zhang et al. (2017) who use WRF-Chem to simulate the impact of RCPs on air pollutants over the USA. They find overall small decreases in
420 6mDM8h O₃ (up to 2 ppb) except in some large urban areas following RCP 4.5, where they report increases of up to 10 ppb. Following SSP2-4.5, we see a similar trend (average reduction of -1.5 ppb, but localised increases around some cities and industrialised areas).

4 Limitations

425 There are limitations with this study to be considered - primary among these is the overestimation of PM_{2.5} concentrations in the present-day. While these are mostly systemic, they should be taken into account when considering the absolute concentrations reported, especially when compared to air quality guideline values. It is likely that the percentage changes we see would be less drastic had the model not overestimated PM_{2.5}. As the model overestimates NO₃, and underestimates SO₄ PM_{2.5} we may overestimate the impacts on changes in NH₃ and NO_x emissions, particularly in the agriculture sector on future air quality
430 compared to other sectors, such as industry.

The resolution that we use is designed for the region/country scale and not the urban scale. 30km horizontal resolution is unlikely to faithfully capture atmospheric chemistry at city scale. Although we can represent the locations urban and industrial peaks, be aware that the model may not simulate chemistry at this scale as effectively as a finer-scale model.

435

We also do not control for the impacts of population change in the future scenarios for Table 4, Figure 6 or Table 5.

5 Summary and Conclusions

We use emissions data for three SSPs (SSP1-2.6, SSP2-4.5 and SSP3-7.0), representing very different climate futures, to simulate air quality in Europe in 2050 compared to the present day. This work uses WRF-Chem v4.2 with a much more detailed
440 chemistry scheme and finer grid resolution than much of the previous work using SSPs to provide a more detailed assessment of potential air quality co-benefits on a regional scale.

We show that PM_{2.5}, while expected to reduce compared to the present day across most of Europe in all future scenarios,
445 shows by far the biggest reductions in scenarios with a greater focus on sustainability, and therefore more stringent emissions reductions. We find that in 2050, following SSP1-2.6, mean population-weighted PM_{2.5} concentrations across European coun-

tries reduces by 52% compared to 2014. Whilst under SSP2-4.5, this average reduction is 34%. The smallest average reduction was 18% by following SSP3-7.0. The additional benefits we see from following SSP1-2.6 are likely due to emissions reductions in the agricultural and industrial sectors.

450

We also show a different sign of change O_3 across the scenarios, demonstrating that in the more sustainable scenario, SSP1-2.6 (and to a lesser extent, SSP2-4.5), much of Europe will see reduced 6mDM8h O_3 concentrations, whereas 6mDM8h O_3 will worsen following SSP3-7.0. This is likely driven by a combination of reduced NO_x and increased CH_4 emissions. This demonstrates the importance of reducing CH_4 alongside other O_3 precursor species to avoid reducing the efficacy of overall
455 air pollutant controls, caused by focusing entirely on $PM_{2.5}$ and NO_x without also considering the impacts on O_3 , as evident from the increases in O_3 concentrations during the COVID-19 lockdowns where large reductions in NO_x emissions occurred with smaller or no effect on CH_4 (Jephcote et al. 2021; Miyazaki et al. 2021).

We find that using a regional atmospheric chemistry model provides us with the ability to analyse in more detail where air
460 quality in Europe could change in response to the scenarios and that the patterns in air quality changes using this methodology differ from what you get using climate model output. From that, we can make a more informed hypothesis as to why air pollutants respond the way they do based on sector-specific emissions changes. We demonstrate the value that can be added using this methodology, for example by providing country-specific population-weighted mean changes, which may be more useful to regional and national policymakers. This demonstrates the importance of a combined approach to modelling air quality
465 co-benefits using both global and regional models.

To conclude, our results suggest that air quality co-benefits will be seen if society follows a pathway in which environmental sustainability is a priority, particularly in terms of mitigating climate change. This implies there are potential public health benefits, although the results of this may differ from those of other studies, so further studies to calculate the health benefits
470 are important.

6 Data Availability

Data is available on an online repository at Zenodo (DOI: 10.5281/zenodo.10781398). Data can alternatively be accessed by contacting Connor Clayton. The emissions data is publicly available on the Input4MIPs website (<https://aims2.llnl.gov/search/input4mips/>)

475

Appendix A: Supplementary Figures




Figure A1. The domain input into WRF-Chem v4.2 for our simulations at 30 km resolution. The observation sites used for model validation are also shown.




Figure A2. Annual mean PM_{2.5} for each scenario. (a) shows this metric for the CMIP6 2014 simulation. (b),(c) and (d) show the absolute change from this for SSP1-2.6, SSP2-4.5 and SSP3-7.0 respectively.




Figure A3. Annual mean NH_3 emissions for each scenario. (a) shows this metric for 2014. (b),(c) and (d) show the percentage change from this for SSP1-2.6, SSP2-4.5 and SSP3-7.0 in 2050 respectively.




Figure A4. Annual mean NO_x emissions for each scenario. (a) shows this metric for 2014. (b),(c) and (d) show the percentage change from this for SSP1-2.6, SSP2-4.5 and SSP3-7.0 in 2050 respectively.




Figure A5. Annual mean SO₂ emissions for each scenario. (a) shows this metric for 2014. (b),(c) and (d) show the percentage change from this for SSP1-2.6, SSP2-4.5 and SSP3-7.0 in 2050 respectively.




Figure A6. 6mDM8h O₃ for each scenario. (a) shows this metric for the CMIP6 2014 simulation. (b),(c) and (d) show the absolute change from this for SSP1-2.6, SSP2-4.5 and SSP3-7.0 respectively.




Figure A7. Annual mean NO₂ for each scenario. (a) shows this metric for the CMIP6 2014 simulation. (b),(c) and (d) show the percentage change from this for SSP1-2.6, SSP2-4.5 and SSP3-7.0 respectively.




Figure A8. Annual mean O_3 for each scenario. (a) shows this metric for the CMIP6 2014 simulation. (b),(c) and (d) show the percentage change from this for SSP1-2.6, SSP2-4.5 and SSP3-7.0 respectively.




Figure A9. Mean O₃ in January, February and December for each scenario. (a) shows this metric for the CMIP6 2014 simulation. (b),(c) and (d) show the percentage change from this for SSP1-2.6, SSP2-4.5 and SSP3-7.0 respectively.




Figure A10. Mean O₃ in March, April and May for each scenario. (a) shows this metric for the CMIP6 2014 simulation. (b),(c) and (d) show the percentage change from this for SSP1-2.6, SSP2-4.5 and SSP3-7.0 respectively.




Figure A11. Mean O₃ in June, July and August for each scenario. (a) shows this metric for the CMIP6 2014 simulation. (b),(c) and (d) show the percentage change from this for SSP1-2.6, SSP2-4.5 and SSP3-7.0 respectively.




Figure A12. Mean O_3 in September, October and November for each scenario. (a) shows this metric for the CMIP6 2014 simulation. (b),(c) and (d) show the percentage change from this for SSP1-2.6, SSP2-4.5 and SSP3-7.0 respectively.

Author contributions. Connor J. Clayton: Performing model simulations, transformed emissions files to work with WRF-Chem, creating the figures and writing the paper.

480 James B. McQuaid, Daniel R. Marsh, Steven T. Turnock and Kirsty J. Pringle: Devising the main conceptual ideas, supervising project and advising on developing methodology, writing, and interpreting results. CESM2-WACCM boundary conditions were developed and provided by Daniel R. Marsh. Steven T. Turnock provided model output from previous work to compare with these results.

Ailish M. Graham: Assisting with model setup and code to produce some figures. developed code to make chemical boundary conditions
485 work for WRF-Chem.

Carly J. Reddington: Advising on optimising WRF-Chem setup, population-weighting, use of the reanalysis product and interpreting results.

490 Rajesh Kumar: Advising on transforming SSP emissions to work with WRF-Chem and method of using linear regression to estimate primary PM_{2.5} emissions.

Competing interests. The authors declare that they have no conflict of interest

Acknowledgements. We would like to acknowledge Lansinoh Laboratories Inc. and The Priestley Centre for Climate Futures at the University of Leeds for funding this work and the Met Office for additional funding, supervision and advice. We would also like to thank the Centre
495 for Environmental Modelling and Computation at the University of Leeds for maintaining WRF-Chem at Leeds and providing assistance with implementing the changes to the model inputs needed for this project. We would also like to acknowledge the use of University of Leeds Advanced Research Computing (ARC4) facilities for all model simulations that were used in this project. Rajesh Kumar's contribution to this study is based upon work supported by the NSF National Center for Atmospheric Research, which is a major facility sponsored by the U.S. National Science Foundation under Cooperative Agreement No. 1852977. This work was supported by the UK Natural Environment
500 Research Council (NERC), Grant No. NE/T010401/1. The contributions of Steven Turnock were funded by the Met Office Climate Science for Service Partnership (CSSP) China project under the International Science Partnerships Fund (ISPF).

References

- Abdul Shakor, A. S. A., Pahrol, M. A., and Mazeli, M. I.: Effects of Population Weighting on PM10 Concentration Estimation, *Journal of Environmental and Public Health*, 2020, 1-11, 10.1155/2020/1561823, 2020.
- Adedeji, A. R., Dagar, L., Petra, M. I., De Silva, L. C., and Tao, Z.: Sensitivity of WRF-Chem model resolution in simulating particulate matter in South-East Asia, 10.5194/acp-2019-692, 2020.
- Air Quality in Europe 2022: <https://www.eea.europa.eu/publications/air-quality-in-europe-2022>, last access: 15/02/.
- Amann, M., Kiesewetter, G., Schöpp, W., Klimont, Z., Winiwarter, W., Cofala, J., Rafaj, P., Höglund-Isaksson, L., Gomez-Sabriana, A., Heyes, C., Purohit, P., Borken-Kleefeld, J., Wagner, F., Sander, R., Fagerli, H., Nyiri, A., Cozzi, L., and Pavarini, C.: Reducing global air pollution: the scope for further policy interventions, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 378, 20190331, 10.1098/rsta.2019.0331, 2020.
- Archibald, A. T., Turnock, S. T., Griffiths, P. T., Cox, T., Derwent, R. G., Knote, C., and Shin, M.: On the changes in surface ozone over the twenty-first century: sensitivity to changes in surface temperature and chemical mechanisms, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 378, 20190329, 10.1098/rsta.2019.0329, 2020.
- Balzarini, A. P., G; Honzak, L; Zabkar, R; Curci, G; Forkel, R; Hirtl, M; San Jose, R; Tuccella, P; Grell, G.A.: WRF-Chem model sensitivity to chemical mechanisms choice in reconstructing aerosol optical properties, *Atmospheric Environment*, 115, 2015.
- Ban, J. L., K; Wang, Q; Li, T: Climate change will amplify the inequitable exposure to compound heatwave and ozone pollution, *One Earth*, 5, 2022.
- Butt, E. W., Turnock, S. T., Rigby, R., Reddington, C. L., Yoshioka, M., Johnson, J. S., Regayre, L. A., Pringle, K. J., Mann, G. W., and Spracklen, D. V.: Global and regional trends in particulate air pollution and attributable health burden over the past 50 years, *Environmental Research Letters*, 12, 10.1088/1748-9326/aa87, 2017.
- Chemel, C., Fisher, B. E. A., Kong, X., Francis, X. V., Sokhi, R. S., Good, N., Collins, W. J., and Folberth, G. A.: Application of chemical transport model CMAQ to policy decisions regarding PM2.5 in the UK, *Atmospheric Environment*, 82, 410-417, 10.1016/j.atmosenv.2013.10.001, 2014.
- Cheng, J., Tong, D., Liu, Y., Yu, S., Yan, L., Zheng, B., Geng, G., He, K., and Zhang, Q.: Comparison of Current and Future PM _{2.5} Air Quality in China Under CMIP6 and DPEC Emission Scenarios, *Geophysical Research Letters*, 48, 10.1029/2021gl093197, 2021.
- Chin, M., Savoie, D. L., Huebert, B. J., Bandy, A. R., Thornton, D. C., Bates, T. S., Quinn, P. K., Saltzman, E. S., and De Bruyn, W. J.: Atmospheric sulfur cycle simulated in the global model GOCART: Comparison with field observations and regional budgets, *Journal of Geophysical Research: Atmospheres*, 105, 24689-24712, 10.1029/2000jd900385, 2000.
- Clappier, A. T., P; Beekman, M; Putaud, J.P; de Meij, A.: Impact of SO_x, NO_x and NH₃ emission reductions on PM_{2.5} concentrations across Europe: Hints for future measure development, *Environment International*, 156, 2021.
- Coelho, S. R., S; Fernades, A.P; Lopes, M; Carvalho, D.: How the New Climate Scenarios Will Affect Air Quality Trends: An Exploratory Research, *Urban Climate*, 49, 2023.
- Colette, A., Bessagnet, B., Vautard, R., Szopa, S., Rao, S., Schucht, S., Klimont, Z., Menut, L., Clain, G., Meleux, F., Curci, G., and Rouil, L.: European atmosphere in 2050, a regional air quality and climate perspective under CMIP5 scenarios, *Atmospheric Chemistry and Physics*, 13, 7451-7471, 10.5194/acp-13-7451-2013, 2013.
- Conibear, L., Reddington, C. L., Silver, B. J., Arnold, S. R., Turnock, S. T., Klimont, Z., and Spracklen, D. V.: The contribution of emission

- 540 sources to the future air pollution disease burden in China, *Environmental Research Letters*, 17, 064027, 10.1088/1748-9326/ac6f6f, 2022.
- Crippa, M., Janssens-Maenhout, G., Dentener, F., Guizzardi, D., Sindelarova, K., Muntean, M., Rita, and Granier, C.: Forty years of improvements in European air quality: regional policy-industry interactions with global impacts, *Atmospheric Chemistry and Physics*, 16, 3825-3841, 10.5194/acp-16-3825-2016, 2016.
- 545 Crippa, M., Guizzardi, D., Butler, T., Keating, T., Wu, R., Kaminski, J., Kuenen, J., Kurokawa, J., Chatani, S., Morikawa, T., Pouliot, G., Racine, J., Moran, M. D., Klimont, Z., Manseau, P. M., Mashayekhi, R., Henderson, B. H., Smith, S. J., Suchyta, H., Muntean, M., Solazzo, E., Banja, M., Schaaf, E., Pagani, F., Woo, J.-H., Kim, J., Monforti-Ferrario, F., Pisoni, E., Zhang, J., Niemi, D., Sassi, M., Ansari, T., and Foley, K.: The HTAP_v3 emission mosaic: merging regional and global monthly emissions (2000–2018) to support air quality modelling and policies, *Earth System Science Data*, 15, 2667-2694, 10.5194/essd-15-2667-2023, 2023.
- 550 Danabasoglu, G. WCRP CMIP6: National Center for Atmospheric Research (NCAR) CESM2-WACCM model output collection: <https://catalogue.ceda.ac.uk/uuid/108b11504ef44f4fb0bb66273e261709>, 2019
- Doherty, R. M., Heal, M. R., and O'Connor, F. M.: Climate change impacts on human health over Europe through its effect on air quality, *Environmental Health*, 16, 10.1186/s12940-017-0325-2, 2017.
- Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J. F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., 555 Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geoscientific Model Development*, 3, 43-67, 10.5194/gmd-3-43-2010, 2010.
- Fenech, S., Doherty, R. M., Heaviside, C., Vardoulakis, S., Macintyre, H. L., and O'Connor, F. M.: The influence of model spatial resolution on simulated ozone and fine particulate matter for Europe: implications for health impact assessments, *Atmospheric Chemistry and Physics*, 18, 5765-5784, 10.5194/acp-18-5765-2018, 2018.
- 560 Fenech, S., Doherty, R. M., O'Connor, F. M., Heaviside, C., Macintyre, H. L., Vardoulakis, S., Agnew, P., and Neal, L. S.: Future air pollution related health burdens associated with RCP emission changes in the UK, *Science of The Total Environment*, 145635, 10.1016/j.scitotenv.2021.145635, 2021.
- Feng, L., Smith, S. J., Braun, C., Crippa, M., Gidden, M. J., Hoesly, R., Klimont, Z., Van Marle, M., Van Den Berg, M., and Van Der Werf, G. R.: The generation of gridded emissions data for CMIP6, *Geoscientific Model Development*, 13, 461-482, 10.5194/gmd-13-461-2020, 2020.
- 565 Finch, D. P. P., P.I: Increasing ambient surface ozone levels over the UK accompanied by fewer extreme events, *Atmospheric Environment*, 237, 2020.
- Finch, D. P. P., P.I: Increasing ambient surface ozone levels over the UK accompanied by fewer extreme events, *Atmospheric Environment*, 237, 2020.
- Fuller, R., Landrigan, P. J., Balakrishnan, K., Bathan, G., Bose-O'Reilly, S., Brauer, M., Caravanos, J., Chiles, T., Cohen, A., Corra, L., 570 Cropper, M., Ferraro, G., Hanna, J., Hanrahan, D., Hu, H., Hunter, D., Janata, G., Kupka, R., Lanphear, B., Lichtveld, M., Martin, K., Mustapha, A., Sanchez-Triana, E., Sandilya, K., Schaeffli, L., Shaw, J., Seddon, J., Suk, W., Téllez-Rojo, M. M., and Yan, C.: Pollution and health: a progress update, *The Lancet Planetary Health*, 6, e535-e547, 10.1016/s2542-5196(22)00090-0, 2022.
- Georgiou, G. K., Christoudias, T., Proestos, Y., Kushta, J., Hadjinicolaou, P., and Lelieveld, J.: Air quality modelling in the summer over the eastern Mediterranean using WRF-Chem: chemistry and aerosol mechanism intercomparison, *Atmospheric Chemistry and Physics*, 575 18, 1555-1571, 10.5194/acp-18-1555-2018, 2018.
- Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S. J.: Sources and distributions of dust aerosols simulated with the GOCART model, *Journal of Geophysical Research: Atmospheres*, 106, 20255-20273, 10.1029/2000jd000053, 2001.

Giovannini, L. F., E; Karl, T; Rotach, M.W; Staquet, C; Trini Castelli, S; Zardi, D: Atmospheric Pollutant Dispersion over Complex Terrain: Challenges and Needs for Improving Air Quality Measurements and Modeling, *Atmosphere*, 11, 2020.

580 Goto, D., Ueda, K., Ng, C. F. S., Takami, A., Ariga, T., Matsuhashi, K., and Nakajima, T.: Estimation of excess mortality due to long-term exposure to PM_{2.5} in Japan using a high-resolution model for present and future scenarios, *Atmospheric Environment*, 140, 320-332, 10.1016/j.atmosenv.2016.06.015, 2016.

Graham, A. M., Pringle, K. J., Arnold, S. R., Pope, R. J., Vieno, M., Butt, E. W., Conibear, L., Stirling, E. L., and McQuaid, J. B.: Impact of weather types on UK ambient particulate matter concentrations, *Atmospheric Environment: X*, 5, 100061, 10.1016/j.aeaoa.2019.100061, 585 2020.

Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled “online” chemistry within the WRF model, *Atmospheric Environment*, 39, 6957-6975, 10.1016/j.atmosenv.2005.04.027, 2005.

Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chemistry and Physics*, 6, 3181-3210, 10.5194/acp-6-590 3181-2006, 2006.

Guerreiro, C. B. B., Foltescu, V., and De Leeuw, F.: Air quality status and trends in Europe, *Atmospheric Environment*, 98, 376-384, 10.1016/j.atmosenv.2014.09.017, 2014.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., 595 Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999-2049, 10.1002/qj.3803, 2020.

Hodzic, A; Knote., C; : MOZART gas-phase chemistry with MOSAIC aerosols, 2014.

600 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J.-I., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geoscientific Model Development*, 11, 369-408, 10.5194/gmd-11-369-2018, 2018.

Hystad, P. Y., S; Brauer, M: Air pollution health impacts: the knowns and unknowns for reliable global burden calculations, *Cardiovascular Research*, 116, 2020. 605

Im, U. B., R; Solazzo, E; Kioutsioukis, I; Badia, A; Balzarini, A; Baro, R; Bellasio, R; Brunner, D; Chemel, C; Curci, G; Denier van der Gon, H; Flemming, J; Forkel, R; Giordano, L; Jimenez-Guerrero, P; Hirtl, M; Hodzic, A; Honzak, L; Jorba, O; Galmarini, S: Evaluation of operational online-coupled regional air quality models over Europe and North America in the context of AQMEII phase 2. Part II: Particulate Matter, *Atmospheric Environment*, 2014.

610 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G., Keating, T., Zhang, Q., Kurokawa, J., Wankmüller, R., Denier Van Der Gon, H., Kuenen, J. J. P., Klimont, Z., Frost, G., Darras, S., Koffi, B., and Li, M.: HTAP_v2.2: a mosaic of regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of air pollution, *Atmospheric Chemistry and Physics*, 15, 11411-11432, 10.5194/acp-15-11411-2015, 2015.

Jephcote, C. H., A.L; Adams, K; Gulliver, J: Changes in air quality during COVID-19 ‘lockdown’ in the United Kingdom, *Environmental Pollution*, 272, 2021. 615

- Jiang, J; Aksoyoglu, S; Ciarelli, G; Baltensperger, U; Prevot, A.S.H.: Changes in ozone and PM_{2.5} in Europe during the period of 1990–2030: Role of reductions in land and ship emissions, *Science of the Total Environment*, 741, 2020.
- Jones, B. and O'Neill, B. C.: Spatially explicit global population scenarios consistent with the Shared Socioeconomic Pathways, *Environmental Research Letters*, 11, 2016.
- 620 Juginović, A., Vuković, M., Aranza, I., and Biloš, V.: Health impacts of air pollution exposure from 1990 to 2019 in 43 European countries, *Scientific Reports*, 11, 10.1038/s41598-021-01802-5, 2021.
- Kim, S. E., Xie, Y., Dai, H., Fujimori, S., Hijioka, Y., Honda, Y., Hashizume, M., Masui, T., Hasegawa, T., Xu, X., Yi, K., and Kim, H.: Air quality co-benefits from climate mitigation for human health in South Korea, *Environment International*, 136, 105507, 10.1016/j.envint.2020.105507, 2020.
- 625 Kumar, R., Barth, M. C., Pfister, G. G., Delle Monache, L., Lamarque, J. F., Archer-Nicholls, S., Tilmes, S., Ghude, S. D., Wiedinmyer, C., Naja, M., and Walters, S.: How Will Air Quality Change in South Asia by 2050?, *Journal of Geophysical Research: Atmospheres*, 123, 1840-1864, 10.1002/2017jd027357, 2018.
- Lee, J. D. D., WS; Finch, D. P; Wilde, S. E; Palmer, P.I: UK surface NO₂ levels dropped by 42% during the COVID-19 lockdown: impact on surface O₃, *Atmos Chem Phys*, 20, 15743–15759, 2020.
- 630 Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., and Pozzer, A.: The contribution of outdoor air pollution sources to premature mortality on a global scale, *Nature*, 525, 367-371, 10.1038/nature15371, 2015.
- Lelieveld, J., Pozzer, A., Pöschl, U., Fnais, M., Haines, A., and Münzel, T.: Loss of life expectancy from air pollution compared to other risk factors: a worldwide perspective, *Cardiovascular Research*, 116, 1910-1917, 10.1093/cvr/cvaa025, 2020.
- Liu, Z., Doherty, R. M., Wild, O., O'Connor, F. M., and Turnock, S. T.: Tropospheric ozone changes and ozone sensitivity from the present day to the future under shared socio-economic pathways, *Atmospheric Chemistry and Physics*, 22, 1209-1227, 10.5194/acp-22-1209-2022, 2022.
- 635 Lobell, D. B., Di Tommaso, S., and Burney, J. A.: Globally ubiquitous negative effects of nitrogen dioxide on crop growth, *Science Advances*, 8, 10.1126/sciadv.abm9909, 2022.
- Manders, A. M. M. S., M; Querol, X; Albert, M.F.M.A; Vercauteren, J; Kuhlbusch, T.A.J; Hoogerbrugge, R.: Sea salt concentrations across the European continent, *Atmospheric Environment*, 44, 2010.
- 640 Miyazaki, K., Bowman, K., Sekiya, T., Takigawa, M., Neu, J. L., Sudo, K., Osterman, G., and Eskes, H.: Global tropospheric ozone responses to reduced NO_x emissions linked to the COVID-19 worldwide lockdowns, *Science Advances*, 7, eabf7460, 10.1126/sciadv.abf7460, 2021.
- Monks, P. S., Archibald, A. T., Colette, A., Cooper, O., Coyle, M., Derwent, R., Fowler, D., Granier, C., Law, K. S., Mills, G. E., Stevenson, D. S., Tarasova, O., Thouret, V., Von Schneidmesser, E., Sommariva, R., Wild, O., and Williams, M. L.: Tropospheric ozone and its precursors from the urban to the global scale from air quality to short-lived climate forcer, *Atmospheric Chemistry and Physics*, 15, 8889-8973, 10.5194/acp-15-8889-2015, 2015.
- 645 O'Neill, B. C., Krieglner, E., Ebi, K. L., Kemp-Benedict, E., Riahi, K., Rothman, D. S., Van Ruijven, B. J., Van Vuuren, D. P., Birkmann, J., Kok, K., Levy, M., and Solecki, W.: The roads ahead: Narratives for shared socioeconomic pathways describing world futures in the 21st century, *Global Environmental Change*, 42, 169-180, 10.1016/j.gloenvcha.2015.01.004, 2017.
- 650 Peace, A. H., Carslaw, K. S., Lee, L. A., Regayre, L. A., Booth, B. B. B., Johnson, J. S., and Bernie, D.: Effect of aerosol radiative forcing uncertainty on projected exceedance year of a 1.5 °C global temperature rise, *Environmental Research Letters*, 15, 0940a0946, 10.1088/1748-9326/aba20c, 2020.

655 Pusede, S. E., Duffey, K. C., Shusterman, A. A., Saleh, A., Laughner, J. L., Wooldridge, P. J., Zhang, Q., Parworth, C. L., Kim, H., Capps, S. L., Valin, L. C., Cappa, C. D., Fried, A., Walega, J., Nowak, J. B., Weinheimer, A. J., Hoff, R. M., Berkoff, T. A., Beyersdorf, A. J., Olson, J., Crawford, J. H., and Cohen, R. C.: On the effectiveness of nitrogen oxide reductions as a control over ammonium nitrate aerosol, *Atmospheric Chemistry and Physics*, 16, 2575-2596, 10.5194/acp-16-2575-2016, 2016.

660 Putaud, J.-P., Pisoni, E., Mangold, A., Hueglin, C., Sciare, J., Pikridas, M., Savvides, C., Ondracek, J., Mbengue, S., Wiedensohler, A., Weinhold, K., Merkel, M., Poulain, L., Van Pinxteren, D., Herrmann, H., Massling, A., Nordstroem, C., Alastuey, A., Reche, C., Pérez, N., Castillo, S., Sorribas, M., Adame, J. A., Petaja, T., Lehtipalo, K., Niemi, J., Riffault, V., De Brito, J. F., Colette, A., Favez, O., Petit, J.-E., Gros, V., Gini, M. I., Vratolis, S., Eleftheriadis, K., Diapouli, E., Denier Van Der Gon, H., Yttri, K. E., and Aas, W.: Impact of 2020 COVID-19 lockdowns on particulate air pollution across Europe, *Atmospheric Chemistry and Physics*, 23, 10145-10161, 10.5194/acp-23-10145-2023, 2023.

665 Raes, F., Liao, H., Chen, W.-T., and Seinfeld, J. H.: Atmospheric chemistry-climate feedbacks, *Journal of Geophysical Research*, 115, 10.1029/2009jd013300, 2010.

Rao, S., Klimont, Z., Smith, S. J., Van Dingenen, R., Dentener, F., Bouwman, L., Riahi, K., Amann, M., Bodirsky, B. L., Van Vuuren, D. P., Aleluia Reis, L., Calvin, K., Drouet, L., Fricko, O., Fujimori, S., Gernaat, D., Havlik, P., Harmsen, M., Hasegawa, T., Heyes, C., Hilaire, J., Luderer, G., Masui, T., Stehfest, E., Strefler, J., Van Der Sluis, S., and Tavoni, M.: Future air pollution in the Shared Socio-economic Pathways, *Global Environmental Change*, 42, 346-358, 10.1016/j.gloenvcha.2016.05.012, 2017.

670 Reddington, C. L., Turnock, S. T., Conibear, L., Forster, P. M., Lowe, J. A., Ford, L. B., Weaver, C., Van Bavel, B., Dong, H., Alizadeh, M. R., and Arnold, S. R.: Inequalities in Air Pollution Exposure and Attributable Mortality in a Low Carbon Future, *Earth's Future*, 11, 10.1029/2023ef003697, 2023.

Sa, E; Martins, H; Ferreira, J; Marta-Almeida, M; Rocha, A; Carvalho, A; Freitas, S; Borrego, C;: Climate change and pollutant emissions impacts on air quality in 2050 over Portugal, *Atmospheric Environment*, 131, 2016.

675 Scott, C. E., Arnold, S. R., Monks, S. A., Asmi, A., Paasonen, P., and Spracklen, D. V.: Substantial large-scale feedbacks between natural aerosols and climate, *Nature Geoscience*, 11, 44-48, 10.1038/s41561-017-0020-5, 2018.

Silva, R. A., West, J. J., Lamarque, J.-F., Shindell, D. T., Collins, W. J., Dalsoren, S., Faluvegi, G., Folberth, G., Horowitz, L. W., Nagashima, T., Naik, V., Rumbold, S. T., Sudo, K., Takemura, T., Bergmann, D., Cameron-Smith, P., Cionni, I., Doherty, R. M., Eyring, V., Josse, B., Mackenzie, I. A., Plummer, D., Righi, M., Stevenson, D. S., Strode, S., Szopa, S., and Zengast, G.: The effect of future ambient air pollution on human premature mortality to 2100 using output from the ACCMIP model ensemble, *Atmospheric Chemistry and Physics*, 16, 9847-9862, 10.5194/acp-16-9847-2016, 2016.

680 Tainio, M., Juda-Rezler, K., Reizer, M., Warchałowski, A., Trapp, W., and Skotak, K.: Future climate and adverse health effects caused by fine particulate matter air pollution: case study for Poland, *Regional Environmental Change*, 13, 705-715, 10.1007/s10113-012-0366-6, 2013.

685 Tarín-Carrasco, P., Morales-Suárez-Varela, M., Im, U., Brandt, J., Palacios-Peña, L., and Jiménez-Guerrero, P.: Isolating the climate change impacts on air-pollution-related-pathologies over central and southern Europe – a modelling approach on cases and costs, *Atmospheric Chemistry and Physics*, 19, 9385-9398, 10.5194/acp-19-9385-2019, 2019.

Turnock, S. T., Reddington, C. L., West, J. J., and O'Connor, F. M.: The Air Pollution Human Health Burden in Different Future Scenarios That Involve the Mitigation of Near-Term Climate Forcers, *Climate and Land-Use, GeoHealth*, 7, 10.1029/2023gh000812, 2023.

690 Turnock, S. T., Allen, R. J., Andrews, M., Bauer, S. E., Deushi, M., Emmons, L., Good, P., Horowitz, L., John, J. G., Michou, M., Nabat, P., Naik, V., Neubauer, D., O'Connor, F. M., Olivié, D., Oshima, N., Schulz, M., Sellar, A., Shim, S., Takemura, T., Tilmes, S., Tsigaridis,

- K., Wu, T., and Zhang, J.: Historical and future changes in air pollutants from CMIP6 models, *Atmospheric Chemistry and Physics*, 20, 14547-14579, 10.5194/acp-20-14547-2020, 2020.
- 695 Van Donkelaar, A. H., Melanie S; Bindle, Liam; Brauer, Michael; Brook, Jeffrey R; Garay, Michael J; Hsu, N. Christina; Kalashnikova, Olga V; Kahn, Ralph A; Lee, Colin; Levy, Robert C; Lyapustin, Alexei; Sayer, Andrew M; Martin, Randall V: Monthly Global Estimates of Fine Particulate Matter and Their Uncertainty, *Environmental Science & Technology*, 55, 2021.
- Van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J., and Rose, S. K.: The representative concentration pathways: an overview, *Climatic Change*, 109, 5-31, 10.1007/s10584-011-0148-z, 2011.
- 700 Vandyck, T., Keramidas, K., Tchung-Ming, S., Weitzel, M., and Van Dingenen, R.: Quantifying air quality co-benefits of climate policy across sectors and regions, *Climatic Change*, 163, 1501-1517, 10.1007/s10584-020-02685-7, 2020.
- Venter, Z. S., Aunan, K., Chowdhury, S., and Lelieveld, J.: Air pollution declines during COVID-19 lockdowns mitigate the global health burden, *Environmental Research*, 192, 110403, 10.1016/j.envres.2020.110403, 2021.
- 705 Vohra, K., Vodonos, A., Schwartz, J., Marais, E. A., Sulprizio, M. P., and Mickley, L. J.: Global mortality from outdoor fine particle pollution generated by fossil fuel combustion: Results from GEOS-Chem, *Environmental Research*, 110754, 10.1016/j.envres.2021.110754, 2021.
- Von Schneidmesser, E., Monks, P. S., Allan, J. D., Bruhwiler, L., Forster, P., Fowler, D., Lauer, A., Morgan, W. T., Paasonen, P., Righi, M., Sindelarova, K., and Sutton, M. A.: Chemistry and the Linkages between Air Quality and Climate Change, *Chemical Reviews*, 115, 3856-3897, 10.1021/acs.chemrev.5b00089, 2015.
- 710 Von Schneidmesser, E., Driscoll, C., Rieder, H. E., and Schiferl, L. D.: How will air quality effects on human health, crops and ecosystems change in the future?, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 378, 20190330, 10.1098/rsta.2019.0330, 2020.
- Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geoscientific Model Development*, 4, 625-641, 10.5194/gmd-4-625-2011, 2011.
- 715 WHO global air quality guidelines. Particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. Geneva: World Health Organization; 2021.
- World Health Organisation (2022) WHO Ambient Air Quality Factsheet [online], available at: [https://www.who.int/news-room/factsheets/detail/ambient-\(outdoor\)-air-quality-and-health](https://www.who.int/news-room/factsheets/detail/ambient-(outdoor)-air-quality-and-health)
- 720 Wu, X., Xu, Y., Kumar, R., and Barth, M.: Separating Emission and Meteorological Drivers of Mid-21st-Century Air Quality Changes in India Based on Multiyear Global-Regional Chemistry-Climate Simulations, *Journal of Geophysical Research: Atmospheres*, 124, 13420-13438, 10.1029/2019jd030988, 2019.
- Zanis, P. A., D; Turnock, S. T; Naik, V; Szopa, S; Georgoulias, A.K; Bauer, S. E; Deushi, M; Horowitz, L. W; Keeble, J; Le Sager, P; O'Connor, F M; Oshima, N; Tsigaridis, K; Van Noije, T: Climate change penalty and benefit on surface ozone: a global perspective based on CMIP6 earth system models, *Environmental Research Letters*, 17, 2022.
- 725 Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), *Journal of Geophysical Research: Atmospheres*, 113, 10.1029/2007jd008782, 2008.
- Zhang, Y., Smith, S. J., Bowden, J. H., Adelman, Z., and West, J. J.: Co-benefits of global, domestic, and sectoral greenhouse gas mitigation for US air quality and human health in 2050, *Environmental Research Letters*, 12, 114033, 10.1088/1748-9326/aa8f76, 2017.

