

# Assessing the impact of future altimeter constellations in the Met Office global ocean forecasting system

Robert R. King<sup>1</sup>, Matthew J. Martin<sup>1</sup>, Lucile Gaultier<sup>2</sup>, Jennifer Waters<sup>1</sup>, Clément Ubelmann<sup>3</sup>, and Craig Donlon<sup>4</sup>

<sup>1</sup>Met Office, Exeter, UK

<sup>2</sup>OceanDataLab, Plouzané, France

<sup>3</sup>DATLAS, France

<sup>4</sup>ESTEC, ESA, Noordwijk, Netherlands

**Correspondence:** Robert R. King (robert.r.king@metoffice.gov.uk)

## Abstract.

Satellite altimeter measurements of Sea [Level Anomaly \(SLA\)](#)[Surface Height \(SSH\)](#) are a crucial component of current operational ocean forecasting systems. The launch of the [SWOT](#)[Surface Water and Ocean Topography \(SWOT\)](#) wide-swath altimeter mission is bringing a step change in our observing capacity with 2-dimensional mesoscale structures now able to be observed over the global ocean. Proposals are now being considered for the make-up of the future altimeter constellation. In this study we use Observing System Simulation Experiments (OSSEs) to compare the impact of additional altimeter observations from two proposed future satellite constellations. We focus on the expected impact on the Met Office operational ocean analysis and forecasting system of assimilating an observation network including either 12 nadir altimeters or 2 wide-swath altimeters.

Here we show that an altimeter constellation of 12 nadir altimeters produces greater reductions in the errors for SSH, surface currents, temperature and salinity fields compared to a constellation of 2 wide-swath altimeters. The impact is greatest in the dynamic Western Boundary Current regions where the nadir altimeters can reduce the SSH RMS ([root mean square](#)) error by half, while the wide-swath altimeter only reduces this by one-quarter. A comparison of the spatial scales resolved in daily SSH fields also highlights the superiority of the nadir constellation in our forecasting system. We also highlight the detrimental impact spatially-correlated errors could have on the immediate use of wide-swath altimeter observations. However, we still achieve promising impacts from the assimilation of wide-swath altimetry and work is ongoing to develop improved methods to account for spatially-correlated observation errors within our data assimilation scheme.

*Copyright statement.* The works published in this journal are distributed under the Creative Commons Attribution 4.0 License. This licence does not affect the Crown copyright work, which is re-usable under the Open Government Licence (OGL). The Creative Commons Attribution 4.0 License and the OGL are interoperable and do not conflict with, reduce or limit each other.

## 1 Introduction

Measurements of sea ~~level anomaly (SLA)~~ surface height (SSH) from satellite altimeters have been available for more than 30 years and are routinely assimilated into ocean reanalysis and forecasting systems (Le Traon et al., 2017; Davidson et al., 2019). They play a crucial role in constraining ~~models of~~ the mesoscale ocean dynamics in these systems, allowing estimates of the past, current and future state of the ocean to be made. Until recently, ~~the SLA data from altimeters were altimeters~~ only measured along a ~~single~~ track directly under the path of the satellites (the nadir track). The nature of these nadir altimeters means that the smallest scales which can be resolved along-track are between about 35 km and 55 km depending on the altimeter (Pujol et al., 2023). While multiple satellites are available, there is still not enough data to fully constrain the mesoscale ~~ocean dynamics~~ due to the large gaps between tracks on any particular day. Ballarotta et al. (2019) estimated the spatial and temporal scales of maps of ~~SLA~~ Sea Level Anomaly (SLA) created using three nadir altimeters, with the mean effective spatial resolution at mid-latitudes estimated to be about 200 km.

The Surface Water and Ocean Topography (SWOT) satellite mission (Morrow et al., 2019) was launched in December 2022 with ~~near-real time~~ some data products already available and near-realtime swath data products expected to be available sometime in 2024. It is the first satellite mission to measure ~~SLA~~ SSH across a 120 km wide swath. There is a 20 km gap in the middle of the swath, in the centre of which a nadir instrument measures the ~~SLA~~ SSH. SWOT is expected to observe the 2-D structure of mesoscale ocean processes down to between 15–30 km in wavelength, depending on the sea-state, but will have a 21-day repeat time so will only revisit the same ocean features on relatively long time-scales. SWOT is an experimental mission to demonstrate the concept of swath altimetry and is expected to provide very useful information which can be assimilated into ocean forecasting systems to improve the accuracy of their analyses and forecasts.

There are plans to include more than one wide-swath altimeter (WiSA) ~~on board as part of~~ the Sentinel-3 Next Generation (S3-NG) operational mission which is likely to be flying in around 2030. Different combinations of nadir and swath altimeters are being considered by ESA for S3-NG and the aim of the work described here is to contribute information about the impact of different altimeter constellations on operational ocean forecasts. The ~~main two~~ two main options which are studied here include a constellation of 12 nadir altimeters, and a constellation of two ~~WiSA~~ wide-swath altimeter satellites.

The impact of potential future observations on data assimilation systems is traditionally assessed by running Observing System Simulation Experiments (OSSEs, Fujii et al., 2019) and that is the approach taken in this study. In an OSSE a "nature run", usually a high-resolution model run without data assimilation, is used as a representation of the true ocean. Observations are simulated using the nature run fields for both the existing observing systems as well as the new data type. This simulation of observations includes a representation of the errors expected from each observation type. The simulated observations from the existing observing systems are assimilated into a model which is usually different to the one used to generate the nature run (with different initial conditions, different surface atmospheric forcing, and sometimes a different resolution), to generate a "control" experiment. Another experiment is run assimilating the simulated observations for the existing observing systems and the new observing system. Both the control run and the ~~OSSE run~~ experiments assimilating additional observations can be

compared to the nature run fields in order to assess the reduction in error expected from assimilating the new data in addition  
55 to the existing observing systems. See for example Halliwell et al. (2017) for a complete description of the approach.

Recent examples of the application of OSSEs to study the impact of SWOT data include the work of King and Martin  
60 (2021) in a regional high-resolution (1.5 km) ocean forecasting system, and the work of Benkiran et al. (2021) and Tchonang  
et al. (2021) in a global  $1/12^\circ$  resolution ocean forecasting system. King and Martin (2021) included experiments showing  
spatially correlated errors.

One of the limitations of the OSSE approach is that the results are dependent on the model, data assimilation scheme and  
other aspects of the experimental set-up (such as the realism of the simulated observation errors). Also, operational ocean  
forecasting systems are continually developed, so the impact in today's systems will likely be different to the impact in the  
systems when data from future missions are actually available. To address the forecasting-system dependence of the results,  
65 coordinated OSSEs using more than one system can be carried out (see for example Martin et al. (2020)). The project to which  
this work contributes developed a coordinated framework and ~~the~~ OSSEs were run using the Met Office ocean forecasting  
system (described here) and the Mercator Ocean International ocean forecasting system ~~(described by Benkiran et al., in~~  
~~prep.)~~ (described by Benkiran et al., 2024).

The Forecasting Ocean Assimilation Model (FOAM) is the Met Office's ocean forecasting system and consists of global  
70 and regional configurations. The global ocean/sea-ice system (Barbosa Aguiar et al., 2024) is run at  $1/12^\circ$  resolution while  
a regional system around the UK is run at 1.5 km resolution (Tonani et al., 2019). A lower resolution ( $1/4^\circ$ ) version of the  
global configuration is used as part of the Met Office's coupled numerical weather prediction (NWP) system (Guiavarc'h et al.,  
2019). This is used to produce ocean reanalyses and as part of the Met Office's coupled seasonal prediction system (GloSea,  
MacLachlan et al., 2015). These forecasting systems assimilate ~~SLA-SSH~~ data from nadir altimeters, sea surface temperature  
75 (SST) data from satellites and in situ platforms, in situ profiles of temperature and salinity from various sources including  
Argo and gliders, and sea-ice concentration data from satellites (in the global systems only). These data are assimilated using  
a 3D-Var version of the NEMOVAR data assimilation software (Waters et al., 2015; Mirouze et al., 2016) together with the  
NEMO ocean (Madec et al., 2022) and CICE sea-ice models (Hunke and Lipscombe, 2010). In this study we consider the  
impact of different ~~SLA-altimeter~~ constellations on the global configurations.

80 Details of the OSSE framework used in this study are described in section 2 including the ~~nature~~ Nature run, the simulation of  
observations, the ocean model and data assimilation system used in the assimilation experiments, and details of the experiments.  
Section 3 presents the results of the OSSEs in terms of the impact of the different altimeter constellations on the sea surface  
height (SSH), surface currents, temperature and salinity. It also describes an assessment of the impact on the constrained time  
and space scales. While the results largely focus on the  $1/12^\circ$  global system, a comparison of the results from the  $1/4^\circ$  and  
85  $1/12^\circ$  resolution versions of the system is also presented. Most of the results are from the assimilation of ~~WiSA-wide-swath~~  
altimeter data without the expected correlated observation errors, but we show some results from experiments which include  
these correlated errors in the  $1/4^\circ$  system to highlight some of the issues expected from more realistic data. Finally a discussion  
of the results is given in section 4 and our conclusions ~~are given~~ in section 5.

## 2 Experiment design

90 Observing System Simulation Experiments (OSSEs) require several components: a ~~nature~~ Nature run, observations simulated from the ~~nature~~ Nature run, and additional OSSE runs using a different model setup into which the simulated observations are assimilated (Hoffman and Atlas, 2016; Halliwell et al., 2017). The ~~nature~~ Nature run provides the truth against which other experiments are assessed and from which we sample observations. In this section we describe the ~~nature~~ Nature run used here, the simulation of the observations, the model and data assimilation system used in the OSSEs, and the experimental set-up.

### 95 2.1 Nature run

To draw useful conclusions about changes to the observing system, the ~~nature~~ Nature run needs to be a realistic representation of the ocean. Additionally, the OSSEs must use an ocean model which differs in enough respects so that there is sufficient error growth between the ~~nature~~ Nature run and OSSEs to emulate the differences between an operational system and the real ocean. Ideally, an OSSE would use a separate lower-resolution ocean model with different parameterisations, different surface 100 boundary inputs, and be initialised from a different (though realistic) state.

In this work the ~~nature~~ Nature run was chosen to be a 1/12° global free-running NEMO model (with no data assimilation), the same resolution as the OSSEs, taken from Mercator Ocean and described by Gasparin et al. (2019) and Lellouche et al. (2018). The ~~nature run model~~–Nature run differs from the OSSE ~~model~~ in the version of ~~NEMO~~–the NEMO ocean model (version 3.1 for the ~~nature~~ Nature run), ~~the encompassing changes to the bottom friction, vertical mixing schemes, and the~~ 105 ~~introduction of a non-linear free surface; the~~ parameter settings used, ~~the~~; ~~the sea-ice model (LIM2 for the~~ Nature run, ~~CICE~~ for the OSSEs); ~~the~~ surface forcing and the initial conditions. The ~~nature~~ Nature run was initialised on 11th October 2006 from the EN4 dataset (Good et al., 2013) and was forced at the surface by atmospheric fields from the real-time atmospheric analysis produced at ECMWF<sup>1</sup>. Here we consider the period starting in January 2009. The realism of the ~~nature~~ Nature run in terms of large-scale variability was assessed by Gasparin et al. (2018).

### 110 2.2 Simulation of observations

#### 2.2.1 Standard observation types

The standard set of observations currently assimilated operationally include in situ and satellite SST data, satellite nadir-altimeter along-track ~~SLA~~–SSH data, in situ profiles of temperature and salinity, and satellite sea-ice concentration (SIC) data. Simulated observations were generated for each of these data types using model data from the ~~nature~~ Nature run described above. The positions of the observations were taken from real-world observation coverage representative of a recent period. Realistic observation errors were generated for each type using methods described by Mao et al. (2020) for SST and SIC, and by Gasparin et al. (2019) for in situ profiles. ~~Briefly, this involved adding representation errors by randomly selecting the date either three days before or after the observation date, then using these time-shifted Nature run values in~~

---

<sup>1</sup><https://www.ecmwf.int/en/forecasts/datasets/set-i>

120 the interpolation process (instead of the correct date). This produces larger errors in regions with higher variability, which is  
desirable for generating realistic representation errors and is the same method used by Gasparin et al. (2019) for the AtlantOS  
inter-comparison. Uncorrelated instrumental errors were also added to each observation. These were created by randomly  
sampling from a Gaussian distribution with zero mean and an appropriate standard deviation for each observation type.  
The standard deviations used for the synthetic SST (0.1–0.5 K) and in situ profiles (0.01–0.05 K, 0.01–0.05 psu) varied  
with platform type and are detailed fully in Mao et al. (2020) and Table 4 of Gasparin et al. (2019), respectively. For SIC  
125 observations only representation error was added, but this accounts for uncertainties in the marginal ice zone, so the errors are  
much larger in this area than elsewhere.

### 2.2.2 Altimeter observations

As mentioned in the introduction, the ~~main two~~ two main options for S3-NG altimeter constellations considered here are  
130 i) a constellation of 12 nadir altimeters flying in equidistant Sentinel-3A/B-like orbits and ii) a constellation of two ~~WiSAs~~  
wide-swath altimeters (each with its own nadir altimeter). We also include a baseline altimeter constellation consisting of 3  
altimeters meant to be representative of the current Sentinel-3 satellites (A and B) and Sentinel-6. The two S3-NG constellations  
are assimilated in addition to the nadir altimeter data from Sentinel-6. The data assimilated in the different experiments is  
summarised in Table 1. Other satellite altimeters are also likely to be producing data at the same time as S3-NG and Sentinel-6,  
but since we do not know their likely characteristics we focus ~~only on the Sentinel-1 on Sentinel-6 in conjunction with either 2~~  
135 wide-swath or 12 additional nadir altimeters.

Simulated data of ~~SLA~~ SSH were generated for the existing nadir altimeter constellation as well as for the ~~WiSAs-wide-swath~~  
altimeters using the SWOT simulator described by Gaultier et al. (2016). The ~~nature~~ Nature run data described above were fed  
into the simulator and it generated observations with realistic errors. The error budget for the ~~baseline nadir altimeters was based~~  
~~on the errors in the real datasets for Sentinel-3A, Sentinel-3B and for Jason-3 (which is representative of Sentinel-6)~~. The nadir  
140 altimeter altimeters used the error spectrum computed for current altimeter missions (defined in Esteban-Fernandez, 2014)  
which includes components resulting from instrumental errors, the residual path delay error from the wet-troposphere correction,  
and the sea-state bias. The error spectrum was computed from level-3 products which already had the typical corrections applied  
to real altimeter observations (removal of tides, dynamic atmosphere correction, and long wavelength errors) allowing us to  
directly simulate level-3-like altimeter data. The resulting RMS error in our simulated nadir altimeter observations is  $\sim$ 1.4 cm.  
145 The nadir altimeter data were generated with a sampling of 6 km, similar to the ~~near-real time~~ near-realtime product assimilated  
in our operational systems. For the constellation of 12 nadir altimeters, data were simulated assuming the same error budget as  
for the Sentinel-3 satellites and also with a sampling of 6 km.


For ~~WiSA~~ the wide-swath altimeters, the error budget was generated using a simulation for each component of the error  
expected from SWOT, including phase error, roll error, timing error, KaRIn noise, baseline dilation error and residual path  
150 delay error from the wet troposphere correction (see Gaultier et al., 2016). Each of these components of the error are available  
separately ~~in the files~~, so experiments could be run with different error components. In the main OSSE experiments described  
below, only the spatially uncorrelated components of the error budget (the KaRIn noise and the residual path delay error)

were included in the assimilated data. This is a major simplification and we discuss the impact of this later in the paper. The simulated ~~WiSA-wide-swath altimeter~~ data were generated for the 120 km wide swath (with a gap in the middle of 20 km) 155 at 2 km resolution and were averaged to generate "super-observations" at 10 km resolution, similar to the resolution of the model grid. ~~Since the WiSA~~ The KaRIn noise was generated using a Gaussian distribution with the resulting 2x2 km gridded observations having an RMS of 1.7 cm, reducing to 0.3 cm for the 10 km resolution super-observations. The residual path delay error was simulated by generating a random realisation of a wet troposphere following a spectrum derived from the AMSR instrument and then correcting it using the estimation of the wet troposphere on two beams. The RMS of the residual path delay 160 error was 0.4 cm which was negligibly affected by the observation averaging. The sum of these two components gives a total RMS error in the 10 km resolution observations assimilated in our main experiments of  $\sim$ 0.5 cm, compared 1.4 cm for the nadir altimeter observations. Since the wide-swath altimeter data are expected to suffer from ~~problems-larger errors~~ in high wave conditions, observations were removed when the significant wave height (SWH) was greater than 8 m (a wave model was also generated in conjunction with the main ~~nature~~ ~~Nature~~ run which provided additional inputs to the SWOT simulator). For real 165 data it is likely that the accuracy of the data will suffer at lower SWH than this threshold (see Peral et al., 2015). We note that we have chosen to retain the 6 km resolution of the nadir altimeter observations despite generating "super-observations" at 10 km resolution for the wide-swath altimeter observations. This was a deliberate choice to reflect the pragmatic decisions we intend to make with real wide-swath altimeter observations, that is we are likely to continue to assimilate the nadir observations at the along-track resolution provided, but will average the wide-swath data to approximately match the grid-scale of our model.

170 Our Control experiment assimilated on average 188k altimeter observations per day. With the super-obbing applied to the wide-swath altimeter observations, our 2WiSA experiment assimilated on average 970k altimeter observations per day (including Sentinel-6, the two wide-swath altimeters and the nadir altimeter component of each wide-swath altimeter). On the other hand, the NADIR experiment with Sentinel-6 and an additional 12 nadir altimeters assimilated on average 831k altimeter 175 observations per day. An example of the observational coverage of ~~SLA~~ ~~SSH~~ from the two scenarios studied is shown in Fig. 1. This figure shows that the ~~WiSA-wide-swath altimeter~~ data increase the coverage of the ocean on each day, and that there are regions within the swath that are measured with high spatial resolution, but that ~~significant~~ ~~large~~ gaps still exist. While the ~~WiSA-wide-swath altimeter~~ data measure regions of the ocean with high spatial resolution, there is a trade-off between the two scenarios in terms of spatial and temporal coverage, and particularly in the regularity of the coverage in time and space.

### 2.3 Ocean model and data assimilation scheme

180 The ocean model used in the OSSEs is NEMO at version 3.6 coupled to the ~~CICE~~ sea-ice model CICE version 5.2.1. The configuration of the ocean model is called GO6 and is described in detail by Storkey et al. (2018) while the sea-ice model configuration is called GSI8.1 and is described by Ridley et al. (2018). This ocean/sea-ice model configuration is the version currently used operationally in the global FOAM system. It is available at two different resolutions, one at  $1/4^\circ$  resolution (called ORCA025) and one at  $1/12^\circ$  resolution (called ORCA12). The main results presented in this work are from runs of 185 the ORCA12 system, but some additional experiments have been run using ORCA025 to investigate the impact of correlated errors in the ~~WiSA-wide-swath altimeter~~ observations.



**Figure 1.** Example observation coverage from the two altimeter constellation scenarios (left: [12xnadir12\\_x\\_nadir](#), right: [2xWiSA2\\_x\\_wide-swath](#)). Global 1-day coverage is shown on the top row, along with 1-day (middle) and 7-day (bottom row) coverage over the Gulf Stream region.

The data assimilation system used in FOAM is called NEMOVAR (Waters et al., 2015) and is an incremental multivariate 3DVar-FGAT (First Guess at Appropriate Time) scheme. A key feature of the scheme is that it uses physical balance relationships to transfer information between the different model variables, as described by Weaver et al. (2005). This means that 190 SSH observations are used to estimate corrections to the model's SSH, but also affect the subsurface density field and the horizontal velocities. Similarly, observations of the subsurface temperature and salinity are used to adjust the model's density structure and affect the SSH and horizontal velocities. These different observation types are ~~all assimilated together, and the best analysis produced to make use of them all at the same time~~ assimilated simultaneously to produce a single analysis for each 1-day assimilation window.

195 Another aspect of NEMOVAR is that the information at observation locations is spread spatially (horizontally and vertically) to produce corrections to the model fields at surrounding locations. This is done through the so-called background error correlations, and these are modelled efficiently in NEMOVAR using an implicit diffusion operator (Weaver et al., 2016). In the operational FOAM system, the horizontal correlations are specified using a combination of two functions with distinct length-scales as described in Mirouze et al. (2016). The small-scale errors have a length-scale which depends on the first baroclinic 200 Rossby radius and varies from 25 km to 300 km while the large-scale errors have a 400 km length-scale. Recent work has shown that the altimeter assimilation produces improved results without the longer of these two length-scales, and an update to the FOAM system is being prepared which removes the longer length-scale for temperature (Carneiro et al., 2024). This change is used in the experiments described here.

205 Satellite observations often suffer from biases and NEMOVAR includes the facility to correct these observation biases for SST data (as described by While and Martin, 2019) and for SSH (as described by Lea et al., 2008). For SSH data the bias correction in FOAM consists of two aspects. The first is that the mean dynamic topography (MDT) needed to compare the 210 observations of ~~SLA-SSH (relative to the geoid)~~ with the model's SSH contains errors which result in a bias in the observations. In the idealised OSSE set-up used here however, we produce observations of SSH directly without the need of an external MDT. The estimate of this MDT bias term was therefore initialised to zero and will account for differences in the mean SSH in the two 215 versions of ORCA12 (for the ~~nature~~ Nature run and OSSEs) used in the experiments. At the end of the experiments described later, this MDT bias was small in both magnitude and scale, as expected. The second component of the SSH bias correction is ~~related to~~ designed to account for differences between the modelled and observed SSH due to errors in the ~~pre-processing of operational SLA observations which remove some of the high-frequency signals, representation and removal of high-frequency atmospheric effects~~ particularly at mid- to high-latitudes (poleward of 40°). ~~The~~ As the simulated SSH data ~~produced using the SWOT simulator removes these signals in a similar way to the real observations, so we retain~~ were produced from a Nature run which used a different source of surface forcing compared to the OSSEs, we retained the second bias correction term in these experiments.

220 The FOAM system uses a ~~one-day time window for the data assimilation. All observations valid on a particular day are collected and read into a one-day forecast of NEMO, during which the model counterparts of the observations are calculated~~ 1-day assimilation window, meaning that an analysis is produced daily using observations over a 24-hour period. The observation operator in NEMO is used to calculate a model counterpart to every observation at the nearest model ~~time-step~~. The

225 ~~observations and their model counterparts are then read into NEMOVAR, timestep and interpolated to the observation location. The innovations (the difference between the observation and the model counterparts) are used by NEMOVAR together with gridded information about the model state for use in estimating the multivariate balance relationships, and information about the background and observation error covariances. Note that we use the same observation and background errors here as in our operational system.~~ The analysis increments generated by NEMOVAR (the corrections to the model state) are then read into another run of NEMO over the same day, during which a fraction of the increments are added in on each time-step using Incremental Analysis Updates (IAU; Bloom et al., 1996).

## 2.4 Experimental set-up

230 The OSSE period was chosen to be from January through to July 2009. Three main runs were carried out with the FOAM-ORCA12 model and assimilation system described in the previous sub-section. The first is the Control experiment which assimilated all the standard observations, including a representative nadir altimeter constellation consisting of Sentinel-3A, 3B and Sentinel-6. The second is an experiment which assimilated data from an additional 12 nadir altimeters, in addition to the Sentinel-6 altimeter and the standard observations assimilated in the Control, which we call NADIR. The third is the 2WISA 235 experiment which assimilated the simulated data from two ~~WiSAs~~ wide-swath altimeters described in section 2.2.2 in addition to the Sentinel-6 altimeter and the standard observations assimilated in the Control.

240 The Control, NADIR and 2WISA experiments were started from the same initial condition, which was different to that used in the ~~nature~~ Nature run. This came from a previous reanalysis of the FOAM system valid on 1st January 2009. A three-week spin-up of the system with assimilation of the simulated standard observations was carried out and the Control, NADIR and 2WISA experiments then started from the same initial conditions on 21st January 2009. These experiments were forced at the surface by atmospheric fields coming from the ERA-5 reanalysis produced by ECMWF (Hersbach et al., 2020) which are different to those used in the ~~nature~~ Nature run (which used the real-time ECMWF atmospheric analysis). A summary of the different experiments is given in Table 1.

**Table 1.** Experimental set-up. The standard observations include satellite and in situ SST, in situ profiles of temperature and salinity, and sea-ice concentration.

Experiments	Model configuration	Atmospheric forcing	Assimilated observations						
			Standard obs	S6	S3A&B	12xS3	2xWiSA		
Nature run	NEMOv3.1/LIM, ORCA12	real-time ECMWF	-	-	-	-	-	-	-
Control	NEMOv3.6/CICE, ORCA12	ERA-5	Y	Y	Y	-	-	-	-
NADIR	NEMOv3.6/CICE, ORCA12	ERA-5	Y	Y	-	Y	-	-	-
2WISA	NEMOv3.6/CICE, ORCA12	ERA-5	Y	Y	-	-	-	-	Y

245 ~~To demonstrate that the OSSE framework used here is representative of our operational system assimilating real observations, we have compared the innovation statistics between our Control run and our operational system. While the observational~~

coverage in the two systems is not a perfect match, it was designed to be similar and this allows a fairer comparison than using innovation statistics from the operational system and the full field RMSE from the OSSE Control experiment.


The Control experiment ~~has a globally averaged SSH~~ SSH innovations have a globally-averaged root-mean-square error (RMSE) of just over 6 cm (as shown in Fig. 2). This is consistent with the level of error in the operational FOAM system, estimated by comparing the one day forecasts with observations. Globally averaged RMSE (RMS) of  $\sim 5.8$  cm broadly consistent with an RMS of 6.4 cm in our operational system. Globally-averaged innovation RMS for temperature in the Control peaks at about  $0.80.75^{\circ}\text{C}$  at 100 m depth and decays to around  $0.2^{\circ}\text{C}$  at 1500 m depth, which is also similar to the operational FOAM system. ~~The where the innovation RMS is around  $0.7$ – $1.0^{\circ}\text{C}$  at 100 m depth and  $\sim 0.2^{\circ}\text{C}$  at 1500 m depth. The full-field~~ salinity RMSE in the Control experiment peaks at the surface ~~with a much larger value (at over 1.5 psu )~~ compared to the errors in the operational FOAM system which are estimated to be of the order of 0.25 dominated by regions of high salinity variability and which are sparsely sampled by in situ profiles. However, when comparing innovation statistics globally-averaged innovation RMS for salinity in the Control peaks at about 0.18 psu at the surface (compared to Argo observations). This discrepancy is likely due to the sampling differences of comparisons with the Nature run (which measures the error everywhere), as opposed to the Argo salinity data which do not sample well the regions of highest salinity variability, e.g. in river outflows, at very high latitudes, and in regional seas. Nevertheless, the overall levels of RMSE and decays to around 0.025 psu at 1500 m depth, similar to the operational FOAM system where the innovation RMS is around 0.2 psu at the surface and  $\sim 0.025$  psu at 1500 m depth. The overall innovation errors in the Control experiment are consistent with the errors seen in our operational system, so the idealised experimental framework used here can be viewed as representative of the real system.

### 3 Results

In this section, we detail the impact of assimilating the simulated observations by comparing each experiment with the truth provided by the full 4-D fields from the Nature run. This grid-point-by-grid-point comparison allows us to examine the impact of the assimilation over the full domain ~~meaning we are not hindered by incomplete observation sampling as would be the case with a system assimilating real observations, unlike in reality where our knowledge of the true ocean state is limited~~. We first describe the impact of each experiment on the SSH, the 3D temperature and salinity, and the surface currents. We then describe the impact on the temporal and spatial resolution of SSH features in each experiment and the differences seen when running the experiments with a lower resolution model. Finally, we also show the impact of the ~~expected potential~~ large spatially-correlated errors in the ~~WiSA-wide-swath altimetry~~ data.

#### 3.1 Impact on SSH

The globally-averaged impact on SSH of assimilating additional altimeter observations is ~~significantly positive~~ positive overall in both the NADIR and 2WISA experiments (see Fig. 2). A greater reduction in the SSH RMSE is apparent in the NADIR experiment where the percentage reduction reaches  $\sim 16\%$  compared to  $\sim 10\%$  in the 2WISA experiment. The Western Boundary Current (WBC) regions dominate the SSH variability and, as shown in Fig. 3, are the regions where the additional altimeter




**Figure 2.** Globally-averaged SSH RMSE for the  $1/12^\circ$  experiments (left) along with the percentage reduction in the SSH RMSE compared to the Control experiment (right).

observations have the greatest impact. Broadly, the impact is again greater in the NADIR experiment compared to 2WISA. **While both show some small**


280 **Both experiments show** regions of degradation under the Antarctic sea-ice, **the**. Even though we have no SSH observations in sea-ice covered areas, the long background error correlation length-scale produces changes to the (highly variable) SSH under the sea-ice. While this emulates what happens in our operational system, these experiments have highlighted that we should restrict the spreading of this information under the ice. **The** 2WISA experiment additionally shows a confined region of degradation in the **north-eastern** **north-east** Pacific in the SSH RMSE compared to the Control. **This** occurs in a region of 285 **very low intrinsic SSH variability** (as shown in the map of the monthly RMS of SSH from the Nature run in Fig 4) which is not captured in the background errors in our data assimilation system resulting in noise being added in this region. Although there are a few other areas with a similarly low SSH variability (north of the Antarctic Circumpolar Current and in the mid-Atlantic), the **north-east** Pacific feature aligns with the boundary where we stop applying the full SSH balance (where the temperature stratification is less than 5K). An inspection of daily SSH increments shows large length-scale increments between altimeter 290 swaths which align with this boundary in the 2WISA experiment only. It appears that the interaction of very low SSH variability with the transition from applying balanced SSH increments to only barotropic increments leads to spurious SSH changes in this region.

295 **Monthly SSH RMSE from July 2009 for the Control run (top) and the difference in RMSE compared to the Control for the NADIR (bottom left) and 2WISA (bottom right) runs for the  $1/12^\circ$  experiments. Negative values imply a reduction in RMSE for the experiment (NADIR or 2WISA) compared to the Control.**


Focussing on the Gulf Stream region (Fig. 5), we see that the NADIR experiment shows a **significant** **large** reduction in RMSE **from the 12 nadir altimeters** (49% over the region shown) from assimilating observations from the 13 nadir altimeters



**Figure 3.** Monthly SSH RMSE from July 2009 for the Control run (top) and the difference in RMSE compared to the Control for the NADIR (bottom left) and 2WISA (bottom right) runs for the  $1/12^\circ$  experiments. Negative values imply a reduction in RMSE for the experiment (NADIR or 2WISA) compared to the Control.




**Figure 4.** Globally-averaged RMS of SSH RMSE for from the  $1/12^\circ$  experiments (left) along Nature run over July 2009. The same field is shown in both panels, but with a smaller colour range on the percentage reduction in the SSH RMSE compared right to emphasise the Control experiment (right) low variability regions.



**Figure 5.** Monthly SSH RMSE difference from July 2009 compared to the Control for the NADIR (left) and 2WISA (right) runs for the Gulf Stream region. Negative values imply a reduction in RMSE for the experiment (NADIR or 2WISA) compared to the Control.

(Sentinel-6 plus the additional 12 altimeters) compared to the Control's three nadir altimeters with almost no areas of degradation. In contrast, while the 2WISA experiment reduces the SSH RMSE overall, ~~and significantly in some places,~~ (24% over 300 the region shown), there are also areas along the main Gulf Stream path and its extension where the SSH RMSE is degraded. This experiment assimilated one nadir altimeter in conjunction with the two ~~WiSAs~~ wide-swath altimeters which results in an uneven spatial sampling on any given day (as shown in Fig. 1). ~~While~~


The daily SSH increments shown in Fig. 6 illustrate the effect of the different spatial sampling in the NADIR and 2WISA experiments. The relatively wide spacing of the altimeter swaths in the 2WISA experiment over our 1-day assimilation window 305 produces short length-scale increments near the observation locations and longer length-scale unbalanced SSH increments in the regions between altimeter swaths. In contrast, the relatively close spacing of the altimeter tracks from the 13 nadir altimeters (in the NADIR experiment) over our 1-day assimilation window produces predominantly small-scale SSH increments. The cumulative effect of this is shown by the RMS of the SSH increments over a 21-day period (the repeat cycle of the wide-swath altimeters) in Fig. 6 which indicate that the data assimilation is introducing more variability in the 2WISA experiment than 310 in the NADIR experiment. While the wide-swath altimeter data will constrain the ~~WiSA data will allow a good analysis of the~~ SSH in the vicinity of the data on a particular day, there will be a number of days at any given location which are not sampled by the data, during which time the ~~errors in the model will grow~~ only constraint we have on the SSH comes from the correlations with distant locations. This makes it much harder for the data assimilation to constrain the mesoscale eddy field at 315 all locations with the 2WISA constellation, compared to the constellation of ~~12-13~~ nadir altimeters assimilated in the NADIR experiment, which while having a less detailed picture of the SSH in particular locations, has a more even sampling on each daily assimilation cycle.



**Figure 6.** SSH increments for 20090701 (top) and RMS of SSH increments for the 21-day period from 20090701–20090721 from the Control (left), NADIR (middle) and 2WISA experiments (right).

### 3.2 Impact on subsurface T/S

The assimilation of 12 nadir altimeters significantly 13 nadir altimeters (Sentinel-6 plus the additional 12 altimeters) reduces the globally averaged RMSE for temperature profiles to a large extent compared to the Control with only three nadir altimeters, 320 as shown in Fig. 7. This reduction happens at most depths, but is particularly strong between about 100 m to 1000 m with a peak percentage reduction of  $\sim$ 8% at 400-600 m. In contrast, the reduction in global RMSE for temperature profiles when assimilating the data from two WiSA instruments wide-swath altimeters plus one nadir altimeter is only  $\sim$ 1-2% in the upper 600 m. This is probably due to the regions where we saw degradations in the SSH RMSE (e.g., in the north-east Pacific and parts of the ACC) also having degradations in the impact on temperature profiles, which offsets any improvements seen 325 elsewhere. In the Gulf Stream region where there is an overall improvement in SSH RMSE in the 2WISA experiment there is also an improvement in the temperature profile RMSE at all depths above about 1000 m, peaking at  $\sim$ 8% at about 600 m depth. The NADIR experiment has an even larger reduction in temperature RMSE in this region with improvements of over 20% at the same depth. The balances in our data assimilation scheme allow altimeter observations of the SSH to introduce subsurface 330 changes to the temperature and salinity. However, previous experiments have shown that the assimilation of in situ profiles and altimeter observations can sometimes work against one another (King et al., 2018). With such a large increase in the altimeter observations, the balanced changes applied to subsurface temperature and salinity may dominate over the changes due to the in situ observations leading to degradations in some regions over some depths, such as those seen for temperature below 1000 m in the Gulf Stream region.



**Figure 7.** Global (top) and Gulf Stream region (bottom) profiles of the percentage improvement in RMSE compared to the Control experiment for temperature (left) and salinity (right) for the NADIR and 2WISA runs in the 1/12° system. [Note that the scales on the x-axis vary in each panel.](#)

The global salinity RMSE is [not significantly marginally](#) affected by the assimilation of either of the additional altimeter 335 constellations. There is a slight improvement of <0.25% in the NADIR experiment relative to the Control at most depths, but a slight degradation of <0.25% in the 2WISA experiment. However, in the Gulf Stream region, the impact of the additional observations is positive at depths of ~300-900 m with up to a 25% reduction in salinity RMSE in the NADIR experiment and ~8% in the 2WISA experiment.

### 3.3 Impact on surface currents


340 The assimilation of altimeter data produces corrections to the geostrophic component of the velocities and so, given the SSH improvements we saw in section 3.1, we would expect to see improvements to the geostrophic currents but ~~no significant~~ little impact on the ageostrophic component of the velocities. Therefore, here we follow the procedure we have used for the other variables and compare the daily mean surface velocities from the OSSEs to the Nature run.

345 The time-series of ~~globally-averaged~~ globally-averaged RMSE for the u- and v-components of the surface current velocity in Fig. 8 shows that the Control run velocities take about 4 months to reduce to a stable level whilst the velocity RMSE in the NADIR and 2WISA experiments reduces to a stable level within about a month. The NADIR experiment has a consistently lower RMSE than the 2WISA experiment for both the u- and v-velocity components throughout the experiments. Overall, in the final 2 months of the experiments, the NADIR experiment has 9% and 11% reductions in u- and v-velocity RMSE respectively, while the 2WISA experiment has 4% and 7% reductions in u- and v-velocity RMSE respectively.


350 Focussing on the last month of the experiments (July 2009) in Fig. 9 we see a similar pattern in the changes in RMSE over the globe for surface currents to the one we saw for SSH, though there are more regions of degradation in the surface currents than for SSH. In the Amazon outflow and Somali Current regions for instance there are some degradations in the surface current RMSE. The assimilation scheme used in FOAM does not generate balanced velocity increments near the equator so assimilation of extra ~~SLA-SSH~~ data in those regions could generate a model response which degrades the velocities. This 355 appears to be the case in both the NADIR and 2WISA experiments, though the regions of degradation are quite small overall.

360 Looking in more detail at the Gulf Stream region, the position of the Gulf Stream is noticeably different between the Nature ~~Run-run~~ and the Control, particularly along the coasts of North and South Carolina (see top panel of Fig. 10). This is improved in the 2WISA run, but ~~is more significantly improved to a larger extent~~ in the NADIR experiment. The misplacement of the main Gulf Stream path in the Control and the differing improvements in the NADIR and 2WISA runs is highlighted by the dipole in the surface current mean error seen in each experiment (see second row of Fig. 10). The magnitude of the dipole reduces in the 2WISA run (compared to that in the Control), but is smaller again in the NADIR experiment.


365 In addition to the change in the position of the main Gulf Stream path, there is the expected misplacement of individual eddies in the Control compared to the Nature ~~Run-run~~. For instance, in the Nature run there are two eddies present in the ~~southwest~~ south-west part of this area to the east of the main Gulf Stream path. However, only one of these eddies is resolved in the Control run. Both eddies appear in the 2WISA run, but with more diffuse structure than in the Nature run, while both eddies are resolved in the NADIR run and are qualitatively similar to the Nature run. Similarly, in the ~~northeast~~ north-east corner of this area where there is more variability than closer to the coast, it appears that the Control run cannot sufficiently initialise the position of mesoscale features. This is somewhat improved in the 2WISA run, but the NADIR run shows a much improved qualitative match to the Nature run. It is also clear from the maps of the surface current RMSE and the reduction in 370 the surface current RMSE (see bottom two rows of Fig. 10) that there is an improvement across almost this entire area in the NADIR experiment, while the 2WISA experiment shows ~~significant~~ regions of slight degradation compared to the Control.




**Figure 8.** Globally-averaged time-series of the u- and v-components of surface current velocity RMSE for the Control, NADIR, and 2WISA runs (top), and the corresponding percentage reduction in RMSE compared to the Control (bottom) in the  $1/12^\circ$  system. Note that the y-axis in the upper plots does not start from zero to help highlight the differences between the experiments.



**Figure 9.** Monthly surface current speed RMSE difference from July 2009 compared to the Control for the NADIR (left) and 2WISA (right) runs.



**Figure 10.** Comparison of the monthly (July 2009) mean surface current speed (top row) in the Gulf Stream region near the North American coast for the Nature Run (top left), Control (2nd column), NADIR (3rd column) and 2WISA experiments (4th column). Also shown is the surface current speed mean error (second row) and RMSE (third row) for the Control, NADIR and 2WISA experiments along with the difference in the surface current speed RMSE compared to the Control for the NADIR and 2WISA experiments.



**Figure 11.** Frequency-wavelength powerspectral density score for the Gulf Stream region SSH for the Control (left), NADIR (middle), and 2WISA (right) runs in the 1/12° system. The 0.5 contour defines a boundary between the resolved and unresolved scales.


### 3.4 Impact on constrained scales

As mentioned earlier, the **WiSA-wide-swath altimeter** satellites will measure the ocean with high resolution within their swath, but the repeat cycle is long (21 days) so that their ability to constrain the ocean dynamics on different time and space scales


375 is not obvious a priori. To investigate how the assimilation of the different altimeter constellations constrains the model's SSH at different time and space scales **the power spectra** we have used two power spectra-based metrics which use the ratio of the spectral content of the error (the SSH difference between each experiment and the Nature run) and the spectral content of the true signal (from the Nature run) to determine a signal-to-noise ratio. First, the frequency-wavenumber 2D power spectrum of the errors in the OSSEs were calculated, with a focus on the Gulf Stream region (defined here as 40–80W, 30–50N). This was 380 done using the method of [Le Guillou et al. \(2021\)](#) and software of [Ballarotta et al. \(2019\)](#)<sup>2</sup> and used the daily SSH fields from the full 6-month duration of the OSSEs.

Fig. 11 shows the 2D power spectral density (PSD) score which compares the power spectrum of the SSH error (i.e., the difference between a specific OSSE and the Nature Run) with the power spectrum of the Nature Run SSH for the Control, NADIR and 2WISA experiments. A PSD score of 0.5 corresponds to the space-time scales where the signal 385 is twice the magnitude of the error and is used here to define the boundary between resolved and unresolved space-time scales. All the experiments have large errors at small time-scales and short length-scales, while the large time and space scales are well constrained (have low errors). However, there is a clear improvement in the constrained scales in both NADIR and 2WISA experiments compared to the Control. For the larger spatial wavelengths of 4° and above there is a significant 390 large improvement in the time-scales constrained by assimilating observations from the 12 nadir constellation compared to assimilating the two WiSA observations from the two wide-swath altimeter satellites. However, there is an indication that at spatial wavelengths of 2–4° the 2WISA experiment constrains the errors at smaller time-scales than the NADIR experiment, though the results are rather noisy in this part of the spectrum.

<sup>2</sup>[https://github.com/ocean-data-challenges/2020a\\_SSH\\_mapping\\_NATL60/blob/master/README.md](https://github.com/ocean-data-challenges/2020a_SSH_mapping_NATL60/blob/master/README.md)



**Figure 12.** Effective spatial resolution of the daily SSH fields from each of the experiments (Control run top, NADIR middle left, 2WISA middle right) along with the gain in the effective spatial resolution compared to the Control for the NADIR (bottom left) and 2WISA runs (bottom right).



**Figure 13.** Zonal average over the Atlantic basin of the effective spatial resolution of the daily SSH fields from each of the experiments.

To further investigate the ability of the two different altimeter constellations to constrain the ocean dynamics, we used the technique of Ballarotta et al. (2019) which uses the wavenumber power spectrum to calculate an ‘effective spatial resolution’ using the daily mean SSH fields from each day of the experiments. The somewhat arbitrary term of ‘effective resolution’ is useful here to compare similar experiments with the same metric. However, as discussed by Ballarotta et al. (2019), the scale of resolved ocean features is generally around one-quarter of this ‘effective resolution’, suggesting that a spatial resolution of 100 km corresponds to resolving features of around 25 km diameter. The method used emulates what can be done with real observations and analysis/forecast fields of SSH. This involves estimating the spatial resolution by calculating the ratio of the power spectral density of the SSH error (the difference between the OSSE and the simulated observation) to the power spectral density of the observations. For the observations, we used the simulated AltiKa observations before any errors were added, i.e., they were samples of the truth from the Nature Run. The gridded SSH data from each experiment were interpolated to the positions of the AltiKa observations. The along-track and interpolated OSSE data were then split into overlapping 1500 km

405 segments every 300 km. Finally, the globe was segmented into  $10^\circ \times 10^\circ$  boxes every  $1^\circ$  and all segments within each box were used to compute the power spectral density. The effective resolution at ~~each point the centre of each box~~ was then given by the wavelength where the ratio described ~~above was was equal to~~ 0.5.

The daily effective spatial resolution is shown in Figure 12 for the Control, NADIR and 2WISA experiments. Also shown is the gain in effective resolution in the NADIR and 2WISA experiments compared to the Control. The ~~somewhat arbitrary term of ‘effective resolution’ is useful hereto compare similar experiments with the same metric. However, as discussed by Ballarotta et al. (2019), the scale of resolved ocean features is generally around one-quarter of this ‘effective resolution’, suggesting that a spatial resolution of 100 localised degradation in the north-east Pacific noted in Section km corresponds to resolving features of around 25 km diameter.~~

410 ~~The 3.1 is again evident here. In fact, the effective resolution highlights the localised nature of this degradation. Despite this there are still large improvements generally in the 2WISA experiment compared to the Control. To assess these without being affected by this region, we computed the~~ zonal average of the effective resolution over the Atlantic basin (shown in Figure 13). ~~This~~ highlights the difficulty in constraining the ocean dynamics near the equator in all experiments. The additional observations in the NADIR and 2WISA experiments cannot change this very close to the equator. However, in both the NADIR and 2WISA experiments, there is a clear improvement in the spatial resolution at mid- to high-latitudes with a gain in resolution of up to around 50%. As with the earlier results, the NADIR experiment appears superior overall with a larger gain in the 415 resolution than in the 2WISA experiment across all ocean basins. This superior improvement in the NADIR experiment is particularly evident in all of the Western Boundary Currents.

420 ~~Frequency-wavelength powerspectral-density score for the Gulf Stream region SSH for the Control (left), NADIR (middle), and 2WISA (right) runs in the  $1/12^\circ$  system. The 0.5 contour defines a boundary between the resolved and unresolved scales.~~

425 ~~Effective spatial resolution of the daily SSH fields from each of the experiments (Control Run top, NADIR middle left, 2WISA middle right) along with the gain in the effective spatial resolution compared to the Control for the NADIR (bottom left) and 2WISA runs (bottom right).~~

~~Zonal average over the Atlantic basin of the effective spatial resolution of the daily SSH fields from each of the experiments.~~

### 3.5 Impact of model resolution and correlated errors

430 To understand the impact in the different operational configurations we run at the Met Office, we repeated the experiments described above (Control, NADIR, and 2WISA runs) using our  $1/4^\circ$  resolution ORCA025 system. Similarly to the ORCA12 experiments, these were initialised using a reanalysis of the ORCA025 system valid on 1st January 2009 with a 3-week spin-up where there was assimilation of the standard set of observations. Each experiment was then started from 21st January from the same initial condition.

435 The absolute SSH RMSE was higher in each of the ORCA12 experiments compared to the lower resolution (ORCA025) counterpart, as expected due to the double penalty effect (Rossa et al., 2008). However, the percentage reduction in SSH RMSE shows the NADIR experiment to be superior to the 2WISA experiment in both the high and low resolution systems. Although the absolute improvement in RMSE is largest in the low resolution system, in both systems the percentage reduction is 60%


greater in the NADIR experiment than in the 2WISA experiment. Specifically, while the NADIR and 2WISA experiments reduced the SSH RMSE compared to the Control by 16% and 10% in the experiments with the  $1/12^\circ$  system detailed above, 440 the reductions were 28% and 18% respectively in the experiments with the  $1/4^\circ$  system (see Fig. 14).

Unlike for SSH, the velocity RMSE is of similar magnitude in the Control run of both the high and low resolution systems. Initially the ORCA025 NADIR experiment is slightly superior to the 2WISA experiment but by the final 2 months of the runs there is a similar impact of 2% and 7% reduction in the  $u$  ~~and~~ RMSE and  $v$  RMSE, respectively, showing that the higher resolution system is better able to constrain the surface current RMSE with additional observations and the NADIR experiment 445 observations allow a greater reduction than the 2WISA observations.


The impacts of assimilating wide-swath observations presented so far have used simulated observations which included only the uncorrelated KaRIn errors and ~~residual the residual path delay errors from the~~ wet troposphere correction errors, but none of the expected correlated errors. ~~To investigate the potential impact of these correlated errors we have performed three experiments assimilating wide-swath altimeter observations with the (computationally cheaper)  $1/4^\circ$  resolution ORCA025 system. 2WISA assimilates the uncorrelated wide-swath observations, 2WISA\_CORR assimilates wide-swath observations which include the full correlated errors, and 2WISA\_CORR\_TRIM assimilates the observations with correlated errors but discards observations in the outer half of the swath where the correlated errors are largest. No changes were made to our data assimilation scheme for these experiments.~~ For these additional experiments observations were again simulated using the SWOT simulator tool of Gaultier et al. (2016) and included estimates of the phase, roll, timing, and baseline dilation errors in 450 addition to the KaRIn errors and residual wet troposphere correction errors described in Section 2.2.2 and in King and Martin (2021). ~~The level-2 crossover calibration (as described in Dibarboure et al. (2022)) was applied to correct for the simulated roll errors resulting in a small ( $\sim 2$  cm) residual roll error. The total RMS errors in the 10 km resolution observations with correlated errors was  $\sim 3.9$  cm (but with large extrema), compared to 0.5 cm with only the uncorrelated KaRIn errors and the residual path delay errors.~~ Although techniques are being developed to minimise the magnitude of these correlated errors in real 455 data, such methods may be more limited when applied to near-realtime data which is required for operational ocean prediction. Consequently, these experiments aim to highlight the potential limitations due to correlated errors.

~~To investigate the potential impact of these correlated errors we have performed three experiments assimilating wide-swath altimeter observations with the  $1/4^\circ$  resolution ORCA025 system. 2WISA assimilates the uncorrelated wide-swath observations, 2WISA\_CORR assimilates wide-swath observations which include the full correlated errors, and 2WISA\_CORR\_TRIM assimilates the observations with correlated errors but discards observations in the outer half of the swath where the correlated errors are largest.~~

While the assimilation of simulated observations from 2 wide-swath altimeters without correlated errors (2WISA) reduced the SSH RMSE by 28%, when the full correlated errors were applied this reduction was only 5%. Furthermore, when only observations from the inner half of the swath were used, to discard the largest magnitude correlated errors, the reduction was 470 still only 6%. For surface currents, the addition of the correlated errors in the wide-swath altimeter observations results in worse performance and in fact degrades the RMSE compared to the Control even when the observations are restricted to the inner half of the swath. The  $u$  ~~and~~  $v$  RMSE was reduced by 2% and 7%, respectively, when assimilating observations without correlated



**Figure 14.** Globally-averaged percentage reduction in SSH RMSE for the  $0.25^\circ$  experiments compared to the Control.



**Figure 15.** Globally-averaged time-series of the percentage reduction in RMSE compared to the Control for the u- (left) and v-components of surface current velocity for the  $0.25^\circ$  experiments.

errors, but increased by ~~2%~~ and 1% when assimilating the restricted observations with correlated errors, and increased by ~~6%~~ and 8% ~~for u and v~~ when assimilating the full swath with correlated errors. This highlights the need to develop improved

475 methods to represent correlated observation errors in data assimilation systems such as those of Guillet et al. (2019).

#### 4 Discussion

The experiments described here show that a superior impact is found when assimilating the 12-nadir altimeter constellation. However, similar experiments by the Mercator Ocean International group find the opposite result (Benkiran et al., 2024), that is in their system a greater impact is found when assimilating the 2-wide-swath altimeter constellation. The additional observations assimilated in the two experiments differ in a number of ways which could contribute to the different impacts we have found: the gridding of the observations differs (as explained in Sect. 2.2.2), although both are at slightly higher resolution than the model grid-scale the wide-swath altimeter provides the ability to resolve across-track features; and the spatial and temporal gaps between the tracks/swaths differ markedly. However, an important difference between our system and the Mercator system is the time window used in the data assimilation schemes. The Met Office system uses a 1-day assimilation window (compared to 7-days in the Mercator system) which leads to a less regular spatial sampling of altimeter observations within each analysis window. It appears that a longer assimilation window may be more conducive to initialising mesoscale structures which are relatively static over the window. Whilst such a change could improve results for SSH, this would have to be balanced with the impact on other variables as in an earlier study we found than a shorter assimilation window improved results for SST with little impact on other variables (Lea et al., 2015). However, given the difference in the daily SSH increments between the NADIR and 2WISA experiments it may be possible to make better use of the wide-swath altimeter observations with a 1-day assimilation window by altering the balance between the short and long length-scale increments.

There was also a specific region of degradation found in the north-east Pacific affecting the SSH and surface current RMSE fields. This region has very low intrinsic SSH variability which is not accounted for in the background errors in our data assimilation system resulting in adding noise in this region. This area of degradation also aligns with the boundary where we stop applying the full SSH balance and with large length-scale SSH increments in the 2WISA experiment. It appears that the interaction of very low SSH variability with the transition from applying balanced SSH increments to only barotropic increments leads to a degradation which is localised to the north-east Pacific. The comparison of the effective resolution in Fig. 12 further confirms that the effect is localised and did not affect the impact of assimilation in other regions, nor the overall order of the impact or our conclusions.

The main experiments reported here were performed without spatially-correlated errors included in the simulated wide-swath altimeter observations. However, such correlated errors are expected in real wide-swath data and so additional experiments were run including these error components. The calibration of observations before public release may be able to address major components of these error correlations, but it is not yet clear how effective this will be, nor whether it will be possible for the near-realtime data products required for assimilation into operational systems. Our experiments showed that while the assimilation of simulated observations from 2 wide-swath altimeters without correlated errors (2WISA) reduced the SSH RMSE by more than 25%, when the full correlated errors were applied this reduction was only 5%. We further explored the impact of assimilating only the inner portion of the swath where correlated errors are lower, but the effect was minimal with an SSH RMSE reduction of 6% compared to the Control. The impact of the inclusion of correlated errors was greater for surface currents where we found the RMSE was reduced when assimilating observations without correlated errors by 2% and 7% for the u- and v-velocity components, but was increased by 6% and 8% when correlated errors were included. In this case, restricting the observations to the inner swath limited the increase in RMSE to 2% and 1% for the u- and v-velocities. We therefore

stress the importance of developing improved methods to represent correlated observation errors in data assimilation systems (e.g., Guillet et al., 2019) and testing these with real wide-swath altimeter data. Such data assimilation developments will be a crucial aspect in extracting the full potential of SWOT and other upcoming wide-swath altimeter missions. Additionally, 515 we suggest that the impact of assimilating wide-swath altimetry on the surface currents should be closely monitored as this is an important physical parameter for many users which is difficult to verify due to the sparseness of available observations (Aijaz et al., 2023).

## 5 Conclusions

With the recent launch of the SWOT wide-swath altimetry mission and the on-going planning of the future altimeter constellation, 520 we have used an Observing System Simulation Experiment (OSSE) to investigate the potential impact of two proposed next generation altimeter scenarios in an operational ocean analysis and forecasting system. ~~While OSSEs have some important limitations, they allow us to assess and prepare for the assimilation of new and additional observations within operational systems.~~

In this study we found that the assimilation of additional altimeter observations has a clear positive impact in both experiments 525 with additional nadir altimeters (NADIR) and additional wide-swath altimeters (2WISA). The SSH RMSE is reduced by 10% in our 2WISA experiment compared to the Control, while the NADIR experiment has a superior reduction in SSH RMSE of 16%. The greatest impact from the additional observations is seen in dynamic regions such as the Western Boundary Currents, for example in the Gulf Stream region the SSH RMSE is reduced by 24% in the 2WISA experiment and by 49% in the NADIR experiment. A similar impact is seen on the temperature and salinity fields where the NADIR experiment is 530 generally superior to 2WISA. While the NADIR experiment shows a reduction in the global temperature RMSE of ~8% at around 500m, the reduction is only 1–2% in the 2WISA experiment. The global salinity RMSE was ~~not significantly improved or degraded relative to the Control~~ only marginally affected in either the NADIR or 2WISA experiments. However, the impacts were again more pronounced in the Western Boundary Currents with reductions in the Gulf Stream region of up to 8 and 20% for temperature RMSE and up to 8 and 25% for salinity RMSE in the 2WISA and NADIR experiments respectively.

535 The global surface current fields again show a similar impact with the NADIR experiment having a consistently lower RMSE for both the u- and v-components of surface velocity throughout the experiments. By the final 2 months, the NADIR experiment has 9% and 11% reductions in the u- and v-velocity RMSE, while the 2WISA experiment has 4% and 7% reductions in u- and v-velocity RMSE respectively. A comparison of monthly mean surface current fields in the Control, NADIR and 2WISA experiments to those from the Nature ~~Run run~~ showed that the NADIR experiment better corrected the Gulf Stream path and 540 better initialised the positions and strengths of individual eddies both closer to the the North American coast and in the Gulf Stream extension region further from the coast. The improvement in the initialisation of mesoscale features is an important aspect expected from the increased observation coverage from either altimeter constellation. Knowledge of the surface currents is important for a number of users of operational ocean forecasts, but there is a significant gap in our ability to observe and

constrain this field. Until there are satellite observations of the total surface current vectors, altimetry will remain our most  
545 important tool for constraining surface currents in our forecasting systems.

A comparison of the spatial and temporal scales resolved in each of the experiments highlighted a clear improvement in both the NADIR and 2WISA experiments compared to the Control. The temporal resolution of features appears superior in the NADIR experiment for length-scales greater than  $4^{\circ}$  while at smaller spatial scales the 2WISA experiment appears to constrain the errors at shorter time-scales than the NADIR experiment. Focussing on the spatial resolution of daily SSH fields showed a  
550 clear improvement over the Control at mid- to high-latitudes with a gain in resolution of up to around 50%, with the NADIR experiment again superior overall with a larger gain in the resolution compared to 2WISA, which was particularly evident in the Western Boundary ~~Currents~~ Current regions.

~~While the 2WISA experiment showed an overall lower impact than the NADIR experiment, a specific region of degradation was found in the northern Pacific affecting the SSH and surface current RMSE fields. We have found that the component of our SLA bias correction scheme which is intended to correct for errors in the surface forcing (particularly at high latitudes) inhibited the assimilation of the wide-swath altimeter observations in this region. Although we were unable to find why this affected only the 2WISA experiment and not the NADIR, we found that the effect was localised to the region of degradation in the northern Pacific and did not affect the impact of assimilation in other regions, nor the overall order of the impact or our conclusions. The additional observations assimilated in the two experiments differ in a number of ways which could contribute to the different impacts we have found: the gridding of the observations differs (as explained in Sect. 2.2.2), although both are at slightly higher resolution than the model grid scale; the wide-swath altimeter provides the ability to resolve across-track features; and the spatial and temporal gaps between the tracks/swaths differs markedly.~~

~~Although the experiments described here show that a superior impact is found when assimilating the 12-nadir altimeter constellation, similar experiments by the Mercator Ocean International group find the opposite result (Benkiran et al., 2024), that is in their system a greater impact is found when assimilating the 2-wide-swath altimeter constellation. An important difference between our system and the Mercator system is the time window used in the data assimilation schemes. The Met Office system uses a 1-day assimilation window, compared to 7-days in the Mercator system. Therefore, the most important difference between the two sets of additional observations may be the regularity of the sampling. While the total number of observations in a single data assimilation window is similar between the two altimeter constellations, the NADIR experiment with 12 nadir altimeters sacrifices high spatial resolution for higher temporal resolution and more even sampling, while the 2WISA experiment has the high spatial resolution, but with much larger gaps between individual swaths.~~

~~The main experiments reported here were performed without spatially correlated errors included in the simulated wide-swath altimeter observations. However, such correlated errors are expected in real wide-swath data and so additional experiments were run including these error components. The calibration of observations before public release may be able to address major components of these error correlations, but it is not yet clear how effective this will be, nor whether it will be possible for the near real-time data products required for assimilation into operational systems. Our experiments showed that while the assimilation of simulated observations from 2 wide-swath altimeters without correlated errors (2WISA) reduced the SSH RMSE by more than 25%, when the full correlated errors were applied this reduction was only 5%. We further explored~~

580 the impact of assimilating only the inner portion of the swath where correlated errors are lower, but the effect was minimal with an SSH RMSE reduction of 6% compared to the Control. The impact of the inclusion of correlated errors was greater for surface currents where we found the u/v RMSE was reduced when assimilating observations without correlated errors by 27% for the u- and v-velocity components, but was increased by 68% when correlated errors were included. In this case, restricting the observations to the inner swath limited the increase in RMSE to 21% for the u- and v-velocities. We therefore stress the importance of developing improved methods to represent correlated observation errors in data assimilation systems 585 (e.g., Guillet et al., 2019) and testing these with real wide-swath altimeter data. Such data assimilation developments will be a crucial aspect in extracting the full potential of SWOT and other upcoming wide-swath altimeter missions. Additionally, we suggest that the impact of assimilating wide-swath altimetry on the surface currents should be closely monitored as this is an important physical parameter for many users which is difficult to verify due to the sparseness of available observations (Aijaz et al., 2023).

590 The European Space Agency (ESA) is planning its future altimeter constellation through the Sentinel-3 Next-Generation Topography mission and ~~have~~has chosen to develop a solution involving both nadir and wide-swath altimeters. While the experiments detailed here show a superior impact from an additional 12 nadir altimeters compared to 2 wide-swath altimeters, we still show promising impacts from the new wide-swath altimeters. It must also be noted that the results from OSSEs provide 595 an estimate of the impact in the specific system tested. These results do not reflect the best achievable impact, but rather an estimate of the impact in our current system. Future model and assimilation changes tailored to the observation network is expected to lead to a greater impact. Work is now underway to prepare our operational systems to assimilate real data from the SWOT wide-swath altimeter mission, including accounting for correlated errors in the assimilation scheme, which will test our ability to make good use of ~~WiSA~~wide-swath altimeter data to deliver improved ocean forecasts to users.

600 *Data availability.* The nature of the 4-D data generated in running the model experiments requires a large tape storage facility. These data are of the order of 300 TB (terabytes). The model data can be made available upon request from the authors.

*Author contributions.* RRK, MJM, LG, JW, CU and CD contributed to the design of the project. RRK, MJM and JW worked on the experimental design, RRK ran the experiments, LG simulated the observations, RRK, MJM, JW and CU contributed to the analysis, and the manuscript writing was led by RRK and MJM. All authors have read and agreed to the published version of the manuscript.

mandatory

605 *Competing interests.* The authors declare that they have no conflict of interest.

mandatory even if you declare that no competing interests are present

*Acknowledgements.* Funding support from the European Space Agency is gratefully acknowledged. We also acknowledge our collaborators in developing NEMOVAR at CERFACS, INRIA and ECMWF.

## References

610 Ajaz, S., Brassington, G. B., Divakaran, P., Regnier, C., Drevillon, M., Maksymczuk, J., and Peterson, K. A.: Verification and intercomparison of global ocean Eulerian near-surface currents, *Ocean Modelling*, 186, 102241, <https://doi.org/https://doi.org/10.1016/j.ocemod.2023.102241>, 2023.

615 Ballarotta, M., Ubelmann, C., Pujol, M.-I., Taburet, G., Fournier, F., Legeais, J.-F., Faugère, Y., Delepoule, A., Chelton, D., Dibarbour, G., and Picot, N.: On the resolutions of ocean altimetry maps, *Ocean Science*, 15, 1091–1109, <https://doi.org/10.5194/os-15-1091-2019>, 2019.

620 Barbosa Aguiar, A., Bell, M. J., Blockley, E., Calvert, D., Crocker, R., Inverarity, G., King, R. R., Lea, D. J., Maksymczuk, J., Martin, M. J., Price, M., Waters, J., While, J., Siddorn, J., and Smout-Day, K.: The Met Office Forecast Ocean Assimilation Model (FOAM) using a 1/12 degree grid for global forecasts, *Quarterly Journal of the Royal Meteorological Society*, submitted, 2024.

625 Benkiran, M., Ruggiero, G., Greiner, E., Le Traon, P.-Y., Rémy, E., Lellouche, J. M., Bourdalle-Badie, R., Drillet, Y., and Tchonang, B.: Assessing the Impact of the Assimilation of SWOT Observations in a Global High-Resolution Analysis and Forecasting System Part 1: Methods, *Frontiers in Marine Science*, 8, 947, <https://doi.org/10.3389/fmars.2021.691955>, 2021.

630 Benkiran, M., Le Traon, P.-Y., Rémy, E., and Drillet, Y.: Impact of two high resolution altimetry mission concepts for ocean forecasting, *EGUphere*, 2024, 1–19, <https://doi.org/10.5194/eguphere-2024-420>, 2024.

635 Bloom, S. C., Takacs, L. L., da Silva, A. M., and Ledvina, D.: Data Assimilation Using Incremental Analysis Updates, *Mon. Wea. Rev.*, 124, 1256–1271, [https://doi.org/10.1175/1520-0493\(1996\)124%3C1256:dauiau%3E2.0.co;2](https://doi.org/10.1175/1520-0493(1996)124%3C1256:dauiau%3E2.0.co;2), 1996.

640 Carneiro, D. M., Waters, J., Lea, D., Martin, M., While, J., Weaver, A., Vidard, A., Guiavarc'h, C., Storkey, D., Ford, D., Blockley, E., Baker, J., Haines, K., Price, M., Bell, M., and Renshaw, R.: Updates to the global FOAM system including model and data assimilation changes, Met Office Technical Report, 659, <https://www.metoffice.gov.uk/research/library-and-archive/publications/science/weather-science-technical-reports>, 2024.

645 Davidson, F., Alvera-Azcárate, A., Barth, A., Brassington, G. B., Chassignet, E. P., Clementi, E., De Mey-Frémaux, P., Divakaran, P., Harris, C., Hernandez, F., Hogan, P., Hole, L. R., Holt, J., Liu, G., Lu, Y., Lorente, P., Maksymczuk, J., Martin, M., Mehra, A., Melsom, A., Mo, H., Moore, A., Oddo, P., Pascual, A., Pequignet, A.-C., Kourafalou, V., Ryan, A., Siddorn, J., Smith, G., Spindler, D., Spindler, T., Stanev, E. V., Staneva, J., Storto, A., Tanajura, C., Vinayachandran, P. N., Wan, L., Wang, H., Zhang, Y., Zhu, X., and Zu, Z.: Synergies in Operational Oceanography: The Intrinsic Need for Sustained Ocean Observations, *Frontiers in Marine Science*, 6, <https://doi.org/10.3389/fmars.2019.00450>, 2019.

650 Dibarbour, G., Ubelmann, C., Flamant, B., Briol, F., Peral, E., Bracher, G., Vergara, O., Faugère, Y., Soulat, F., and Picot, N.: Data-driven calibration algorithm and pre-launch performance simulations for the swot mission, *Remote Sensing*, 14, 6070, 2022.

655 Esteban-Fernandez, D.: SWOT project mission performance and error budget document, JPL Doc. JPL D-79084, 83, 2014.

660 Fujii, Y., Rémy, E., Zuo, H., Oke, P., Halliwell, G., Gasparin, F., Benkiran, M., Loose, N., Cummings, J., Xie, J., et al.: Observing system evaluation based on ocean data assimilation and prediction systems: On-going challenges and a future vision for designing and supporting ocean observational networks, *Frontiers in Marine Science*, 6, 417, 2019.

665 Gasparin, F., Greiner, E., Lellouche, J.-M., Legalloudec, O., Garric, G., Drillet, Y., Bourdallé-Badie, R., Traon, P.-Y. L., Rémy, E., and Drévillon, M.: A large-scale view of oceanic variability from 2007 to 2015 in the global high resolution monitoring and forecasting system at Mercator Océan, *Journal of Marine Systems*, 187, 260–276, <https://doi.org/https://doi.org/10.1016/j.jmarsys.2018.06.015>, 2018.

645 Gasparin, F., Guinehut, S., Mao, C., Mirouze, I., Rémy, E., King, R. R., Hamon, M., Reid, R., Storto, A., Le Traon, P.-Y., et al.: Requirements for an integrated in situ Atlantic Ocean observing system from coordinated observing system simulation experiments, *Frontiers in Marine Science*, 6, 83, 2019.

Gaultier, L., Ubelmann, C., and Fu, L.-L.: The challenge of using future SWOT data for oceanic field reconstruction, *Journal of Atmospheric and Oceanic Technology*, 33, 119–126, 2016.

650 Good, S. A., Martin, M. J., and Rayner, N. A.: EN4: Quality controlled ocean temperature and salinity profiles and monthly objective analyses with uncertainty estimates, *Journal of Geophysical Research: Oceans*, 118, 6704–6716, 2013.

Guiavarc'h, C., Harris, C., Lea, D. J., Roberts-Jones, J., Ryan, A., and Ascione, I.: Assessment of ocean analysis and forecast from an atmosphere-ocean coupled data assimilation operational system, *Ocean Science Discussions*, 2019, 1–26, <https://doi.org/10.5194/os-2018-170>, 2019.

655 Guillet, O., Weaver, A. T., Vasseur, X., Michel, Y., Gratton, S., and Gürol, S.: Modelling spatially correlated observation errors in variational data assimilation using a diffusion operator on an unstructured mesh, *Quarterly Journal of the Royal Meteorological Society*, 145, 1947–1967, 2019.

Hallowell, G. R., Mehari, M. F., Le Hénaff, M., Kourafalou, V. H., Androulidakis, I. S., Kang, H. S., and Atlas, R.: North Atlantic Ocean OSSE system: Evaluation of operational ocean observing system components and supplemental seasonal observations for potentially 660 improving tropical cyclone prediction in coupled systems, *Journal of Operational Oceanography*, 10, 154–175, 2017.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmonds, A., Soci, C., Abdalla, S., Abellán, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Vil-laume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/https://doi.org/10.1002/qj.3803>, 2020.

665 Hoffman, R. N. and Atlas, R.: Future observing system simulation experiments, *Bulletin of the American Meteorological Society*, 97, 1601–1616, 2016.

Hunke, E. C. and Lipscombe, W. H.: CICE: The Los Alamos sea ice model, Documentation and software users manual, Version 4.1 (LA-CC-06012), T-3 Fluid Dynamics Group, Los Alamos National Laboratory, Los Alamos, US, 2010.

670 King, R. R. and Martin, M. J.: Assimilating realistically simulated wide-swath altimeter observations in a high-resolution shelf-seas forecasting system, *Ocean Science*, 17, 1791–1813, 2021.

King, R. R., While, J., Martin, M. J., Lea, D. J., Lemieux-Dudon, B., Waters, J., and O'Dea, E.: Improving the initialisation of the Met Office operational shelf-seas model, *Ocean Modelling*, 130, 1 – 14, <https://doi.org/https://doi.org/10.1016/j.ocemod.2018.07.004>, 2018.

675 Le Guillou, F., Metref, S., Cosme, E., Ubelmann, C., Ballarotta, M., Le Sommer, J., and Verron, J.: Mapping altimetry in the forthcoming swot era by back-and-forth nudging a one-layer quasigeostrophic model, *Journal of Atmospheric and Oceanic Technology*, 38, 697–710, 2021.

Le Traon, P.-Y., Dibarboure, G., Jacobs, G., Martin, M., Rémy, E., and Schiller, A.: Use of satellite altimetry for operational oceanography, in: *Satellite altimetry over oceans and land surfaces*, pp. 581–608, CRC Press, 2017.

680 Lea, D., Drecourt, J.-P., Haines, K., and Martin, M.: Ocean altimeter assimilation with observational-and model-bias correction, *Quarterly Journal of the Royal Meteorological Society*, 134, 1761–1774, 2008.

Lea, D. J., Mirouze, I., Martin, M. J., King, R. R., Hines, A., Walters, D., and Thurlow, M.: Assessing a New Coupled Data Assimilation System Based on the Met Office Coupled Atmosphere-Land-Ocean-Sea Ice Model, *Monthly Weather Review*, 143, 4678–4694, <https://doi.org/10.1175/MWR-D-15-0174.1>, 2015.

685 Lellouche, J.-M., Greiner, E., Le Galloudec, O., Garric, G., Regnier, C., Drevillon, M., Benkiran, M., Testut, C.-E., Bourdalle-Badie, R., Gasparin, F., et al.: Recent updates to the Copernicus Marine Service global ocean monitoring and forecasting real-time 1/12 degree high-resolution system, *Ocean Science*, 14, 1093–1126, 2018.

MacLachlan, C., Arribas, A., Peterson, K. A., Maidens, A., Fereday, D., Scaife, A. A., Gordon, M., Vellinga, M., Williams, A., Comer, R. E., Camp, J., Xavier, P., and Madec, G.: Global Seasonal forecast system version 5 (GloSea5): a high-resolution seasonal forecast system, *Quarterly Journal of the Royal Meteorological Society*, 141, 1072–1084, [https://doi.org/https://doi.org/10.1002/qj.2396](https://doi.org/10.1002/qj.2396), 2015.

690 Madec, G., Bourdallé-Badie, R., Chanut, J., Clementi, E., Coward, A., Ethé, C., Iovino, D., Lea, D., Lévy, C., Lovato, T., Martin, N., Masson, S., Mocavero, S., Rousset, C., Storkey, D., Müller, S., Nurser, G., Bell, M., Samson, G., Mathiot, P., Mele, F., and Moulin, A.: NEMO ocean engine, <https://doi.org/10.5281/zenodo.6334656>, 2022.

Mao, C., King, R. R., Reid, R., Martin, M. J., and Good, S. A.: Assessing the Potential Impact of Changes to the Argo and Moored Buoy 695 Arrays in an Operational Ocean Analysis System, *Frontiers in Marine Science*, 7, 588 267, 2020.

Martin, M. J., Remy, E., Tranchant, B., King, R. R., Greiner, E., and Donlon, C.: Observation impact statement on satellite sea surface salinity data from two operational global ocean forecasting systems, *Journal of Operational Oceanography*, 15, 87–103, <https://doi.org/10.1080/1755876X.2020.1771815>, 2020.

Mirouze, I., Blockley, E. W., Lea, D. J., Martin, M. J., and Bell, M. J.: A multiple length scale correlation operator for ocean data assimilation, 700 *Tellus A: Dynamic Meteorology and Oceanography*, 68, 29 744, 2016.

Morrow, R., Fu, L.-L., Arduhin, F., Benkiran, M., Chapron, B., Cosme, E., d'Ovidio, F., Farrar, J. T., Gille, S. T., Lapeyre, G., Le Traon, P.-Y., Pascual, A., Ponte, A., Qiu, B., Raschke, N., Ubelmann, C., Wang, J., and Zaron, E. D.: Global Observations of Fine-Scale Ocean Surface Topography With the Surface Water and Ocean Topography (SWOT) Mission, *Frontiers in Marine Science*, 6, <https://doi.org/10.3389/fmars.2019.00232>, 2019.

705 Peral, E., Rodríguez, E., and Esteban-Fernández, D.: Impact of surface waves on SWOT's projected ocean accuracy, *Remote Sensing*, 7, 14 509–14 529, 2015.

Pujol, M.-I., Dupuy, S., Vergara, O., Sánchez Román, A., Faugère, Y., Prandi, P., Dabat, M.-L., Dagneaux, Q., Lievin, M., Cadier, E., Dibarboure, G., and Picot, N.: Refining the Resolution of DUACS Along-Track Level-3 Sea Level Altimetry Products, *Remote Sensing*, 15, <https://doi.org/10.3390/rs15030793>, 2023.

710 Ridley, J. K., Blockley, E. W., Keen, A. B., Rae, J. G., West, A. E., and Schroeder, D.: The sea ice model component of HadGEM3-GC3. 1, *Geoscientific Model Development*, 11, 713–723, 2018.

Rossa, A., Nurmi, P., and Ebert, E.: Overview of methods for the verification of quantitative precipitation forecasts, in: *Precipitation: Advances in measurement, estimation and prediction*, pp. 419–452, Springer, 2008.

715 Storkey, D., Blaker, A. T., Mathiot, P., Megann, A., Aksenov, Y., Blockley, E. W., Calvert, D., Graham, T., Hewitt, H. T., Hyder, P., et al.: UK Global Ocean GO6 and GO7: A traceable hierarchy of model resolutions, *Geoscientific Model Development*, 11, 3187–3213, 2018.

Tchonang, B. C., Benkiran, M., Le Traon, P.-Y., Jan van Gennip, S., Lellouche, J. M., and Ruggiero, G.: Assessing the Impact of the Assimilation of SWOT Observations in a Global High-Resolution Analysis and Forecasting System 2013 Part 2: Results, *Frontiers in Marine Science*, 8, 1208, <https://doi.org/10.3389/fmars.2021.687414>, 2021.

Tonani, M., Sykes, P., King, R. R., McConnell, N., Péquignet, A.-C., O'Dea, E., Graham, J. A., Polton, J., and Siddorn, J.: The impact of  
720 a new high-resolution ocean model on the Met Office North-West European Shelf forecasting system, *Ocean Science*, 15, 1133–1158,  
<https://doi.org/10.5194/os-15-1133-2019>, 2019.

Waters, J., Lea, D. J., Martin, M. J., Mirouze, I., Weaver, A., and While, J.: Implementing a variational data assimilation system  
725 in an operational 1/4 degree global ocean model, *Quarterly Journal of the Royal Meteorological Society*, 141, 333–349,  
<https://doi.org/10.1002/qj.2388>, 2015.

Weaver, A., Tshimanga, J., and Piacentini, A.: Correlation operators based on an implicitly formulated diffusion equation solved with the  
725 Chebyshev iteration, *Quarterly Journal of the Royal Meteorological Society*, 142, 455–471, 2016.

Weaver, A. T., Deltel, C., Machu, É., Ricci, S., and Daget, N.: A multivariate balance operator for variational ocean data assimilation, *Quarterly  
730 Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography*, 131, 3605–3625, 2005.

While, J. and Martin, M. J.: Variational bias correction of satellite sea-surface temperature data incorporating observations of the bias,  
Quarterly Journal of the Royal Meteorological Society, 145, 2733–2754, 2019.