

# Drivers of decadal trends of the ocean carbon sink in the past, present, and future in Earth system models

Jens Terhaar<sup>1,2,3</sup>

<sup>1</sup>Woods Hole Oceanographic Institution, Woods Hole, MA, USA

5 <sup>2</sup>Climate and Environmental Physics, Physics Institute, University of Bern, Bern, Switzerland

<sup>3</sup>Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

*Correspondence to:* Jens Terhaar (jens.terhaar@unibe.ch)

**Abstract.** The ocean and the land biosphere ~~and the ocean~~ are the two major sinks of anthropogenic carbon at present. When anthropogenic carbon emissions become zero and temperatures stabilizes, the ocean is projected to become the dominant and only global natural sink of carbon. Despite the ocean's importance for the carbon cycle and hence the climate, ~~observing the ocean carbon sink and detecting anthropogenic changes over time remain challenging because~~ uncertainties of the decadal variability of this carbon sink and the underlying drivers of this decadal variability remain large ~~because~~ observing the ocean carbon sink and detecting anthropogenic changes over time remain challenging. The main tools that are used to provide annually resolved estimates of the ocean carbon sink over the last decades are global observation-based  $p\text{CO}_2$  products that extrapolate sparse  $p\text{CO}_2$  observations in space and time and global ocean biogeochemical models forced with atmospheric reanalysis data. However, these tools (i) are limited in time over the last 3 to 7 decades, which hinders statistical analyses of the drivers of decadal trends, (ii) are all based on the same internal climate state, which makes it impossible to separate externally and internally forced contributions to decadal trends, and (iii) cannot assess the robustness of the drivers in the future, especially when carbon emissions decline or cease entirely. Here, I use an ensemble of 12 Earth System Models (ESMs) from phase 6 of the Coupled Model Intercomparison Project (CMIP6) to understand drivers of decadal trends of the past, present and future ocean carbon sink. The simulations by these ESMs span the period from 1850 to 2100 and include 4 different future Shared Socioeconomic Pathways (SSPs), from low emissions and high mitigation to high emissions and low mitigation. Using this ensemble, I show that 80% of decadal trends in the multi-model mean ocean carbon sink can be explained by changes in decadal trends of atmospheric  $\text{CO}_2$  as long as the ocean carbon sink remains smaller than  $4.5 \text{ Pg C yr}^{-1}$ . The remaining 20% are due to internal climate variability and ocean heat uptake, which results in a loss of carbon from the ocean. When the carbon sink exceeds  $4.5 \text{ Pg C yr}^{-1}$ , which only occurs in the high emission SSP3-7.0 and SSP5-8.5, atmospheric  $\text{CO}_2$  rises faster, climate change accelerates, and the ocean overturning and the chemical capacity to take up carbon from the atmosphere reduce, so that decadal trends in the ocean carbon sink become substantially smaller than estimated based on changes in atmospheric  $\text{CO}_2$  trends. The breakdown of this relationship in both high emission pathways also implies that the decadal increase in the ocean carbon sink is effectively limited to be  $\sim 1 \text{ Pg C yr}^{-1} \text{ dec}^{-1}$  in these pathways, even if the trend in atmospheric  $\text{CO}_2$  continues to increase. Previously proposed drivers, such as the atmospheric  $\text{CO}_2$  or the growth rate of

atmospheric CO<sub>2</sub> can explain trends in the ocean carbon sink for specific time periods, for example during exponential atmospheric CO<sub>2</sub> growth, but fail when emissions start to decrease again. The robust relationship over a large ESM ensemble of 12 different ESMs also suggests that very large positive and negative decadal trends of the ocean carbon sink by some pCO<sub>2</sub> products are highly unlikely, and that the change in the decadal trends of the ocean carbon sink around 2000 is likely substantially smaller than estimated by these pCO<sub>2</sub> products.

## 1 Introduction

The ocean has taken up around one quarter of all anthropogenic CO<sub>2</sub> emissions from land use change and fossil fuels since the beginning of the industrial revolution (Friedlingstein et al., 2023; Gruber et al., 2023; Terhaar et al., 2022b). As such, it is, in addition to the land biosphere, one of the two major natural sinks of carbon in the earth system. If land-use change emissions are considered part of the land carbon sink, the land becomes almost neutral and the ocean carbon sink becomes the only major natural carbon sink (Friedlingstein et al., 2023). Once temperatures stabilize, the ocean will become the dominant global natural sink of carbon (Silvy et al., 2024) and will store more than half of the anthropogenically emitted CO<sub>2</sub> in around 1000 years (Joos et al., 2013). By taking up carbon from the atmosphere, the ocean effectively slows down global warming (IPCC, 2021) and will contribute to stabilizing global temperatures over the next centuries if emissions reach near-zero (Terhaar et al., 2023; MacDougall et al., 2020). Here we, I define the ocean carbon sink as in the Global Carbon Budget as the change in air-sea CO<sub>2</sub> flux due to anthropogenic carbon emissions and anthropogenic climate change in comparison to a relatively stable pre-industrial state (Friedlingstein et al., 2023). Consequently, ‘anthropogenic’ refers to direct effects from anthropogenic emissions and the indirect effect to the anthropogenically caused climate change.

The overall magnitude of the ocean carbon sink is mainly determined by the ocean overturning circulation, i.e., the rate at which surface waters with increased anthropogenic carbon content can be transported to the deep ocean and be replaced by waters with low anthropogenic carbon content (Sarmiento et al., 1992; Caldeira and Duffy, 2000; Orr et al., 2001). Furthermore, the magnitude of the ocean carbon sink is influenced by surface ocean capacity to take up more anthropogenic carbon, which itself is determined by the surface ocean carbonate chemistry and especially the alkalinity (Broecker et al., 1979; Terhaar et al., 2022b). Over the historical period, the change in atmospheric CO<sub>2</sub> has been the main driver of changes in the ocean carbon sink and is assumed to be approximately proportional to the strength of the ocean carbon sink (Mikaloff Fletcher et al., 2006; Gruber et al., 2009; Terhaar et al., 2021b). However, this linear relationship between the strength of the ocean carbon sink and atmospheric CO<sub>2</sub> is only assumed to work under exponential atmospheric CO<sub>2</sub> growth (Raupach, 2013; Raupach et al., 2014); Ridge and McKinley, 2021).

Over the last decades, the relatively steady growth of the ocean carbon sink has been weakened by outgassing of natural carbon due to warming and climate change (Joos et al., 1999; McNeil and Matear, 2013; Frölicher et al., 2015) and is superimposed by decadal variability and trends of the ocean carbon sink, i.e., a reduction in the 1990s and an increase since 2000 (Lovenduski et al., 2008, 2007; Le Quéré et al., 2007; Landschützer et al., 2015, 2016). A consensus of the drivers of these trends is still not reached and possible explanations for these different trends are changes in wind and pressure systems (Le Quéré et al., 2007; Keppler and Landschützer, 2019), variability in the ocean circulation and ventilation (DeVries et al., 2017), or changes in the atmospheric CO<sub>2</sub> growth rate and ocean surface temperature due to the eruption of Mount Pinatubo (McKinley et al., 2020; Frölicher et al., 2011). Moreover, recent studies suggest that the observational-based decadal variability over the last decades might be overestimated (Gloege et al., 2021; Hauck et al., 2023).

Despite the importance of the ocean carbon sink for the global climate and carbon cycle, observing or simulating the ocean carbon sink is still challenging. The two main tools to estimate the annually-resolved ocean carbon sink over the past ~~four~~three to seven decades, to provide an annual update every year within the Global Carbon Budget, and to understand the drivers of the magnitude and trends of the ocean carbon sink are observation-based *p*CO<sub>2</sub> products and global ocean biogeochemical models (GOBMs) (Friedlingstein et al., 2023; Hauck et al., 2023). Observation-based *p*CO<sub>2</sub> products extrapolate ~~relatively~~ sparse observations of surface ocean *p*CO<sub>2</sub> using statistical methods and/or machine learning to create global monthly maps of surface ocean *p*CO<sub>2</sub> (Fay et al., 2021; Gregor and Gruber, 2021; Chau et al., 2022; Rödenbeck et al., 2015; Watson et al., 2020; Landschützer et al., 2015); ~~;~~ Gloege et al., 2022; Bennington et al., 2022a, 2022b). These monthly *p*CO<sub>2</sub> maps are then used to estimate the global ocean carbon uptake. In addition, an estimate of pre-industrial natural outgassing of CO<sub>2</sub> due to the difference of riverine carbon input and carbon sequestration in sediments (Lacroix et al., 2020; Regnier et al., 2022) has to be added to estimate the change in the air-sea CO<sub>2</sub> flux with respect to pre-industrial conditions. Global ocean biogeochemical models (GOBMs) (Orr et al., 2001; Hauck et al., 2020; Terhaar et al., ~~in press~~2024) simulate the ocean carbon sink while being forced with past observed atmospheric CO<sub>2</sub> and observation-based reanalysis data, such as wind, humidity, precipitation, temperatures (Hersbach et al., 2020; Tsujino et al., 2018).

The estimates of both product classes vary in magnitude and decadal trends, with *p*CO<sub>2</sub> products estimating a larger magnitude of the ocean carbon sink and also generally larger decadal trends over the last two decades (DeVries et al., 2023; Friedlingstein et al., 2023). One reason for the low carbon sink in ~~*p*CO<sub>2</sub> products~~GOBMs might be the starting year of these simulations that is often later than the beginning of the simulation and the thus slightly too high different pre-industrial reference period and *p*CO<sub>2</sub> in the ocean (Terhaar et al., ~~in press~~2024; Bronselaer et al., 2017). Another reason ~~is~~for the low carbon sink are existing biases in ~~the~~GOBMs in their simulated ocean circulation, especially the Southern Ocean and Atlantic Ocean overturning, and biases in the surface ocean carbonate chemistry ~~in GOBMs that~~(Terhaar et al., 2024). Similar biases were previously identified as drivers of ~~uncertainties and biases~~ in ESM ensembles (Terhaar et al., 2022b, 2021b; Goris et al., 2018). As opposed to the magnitude, the differences in the decadal trends between ~~both~~*p*CO<sub>2</sub> products and GOBMs might be due to ~~sparse~~

amountuneven sampling of observations in space and in time, especiallye.g., few observations in the 1980s and 1990s and few observations in the southern hemisphere, as demonstrated with a subset of  $p\text{CO}_2$  products evaluated with output from a GOBM (Hauck et al., 2023;) and ESMs (Gloege et al., 2021). In addition to differences in trends between from  $p\text{CO}_2$  products and GOBMs, no consensus has yet been made with respect to the underlying drivers of the decadal trends in the ocean carbon sink (Friedlingstein et al., 2023; DeVries et al., 2023; Gruber et al., 2023). The detection of these drivers with  $p\text{CO}_2$  products, GOBMs, and other methods such as data assimilation models (DeVries et al., 2017), is difficult due to the relatively short time period over which enough  $p\text{CO}_2$  observations and atmospheric reanalysis data exist, due to the relative homogeneity of drivers over this period, e.g., constantly increasing atmospheric  $\text{CO}_2$ , and the absence of an alternative climate state against which these ocean carbon sink estimates can be compared.

Here, I use an ensemble of 12 ESMs from phase 6 of the Coupled Model Intercomparison Project (CMIP6) (Table 1) to provide a new perspective on potential drivers of the decadal trends of the ocean carbon sink ~~from phase 6 of the Coupled Model Intercomparison Project (CMIP6) (Table 1). For the analyses of decadal drivers of the ocean carbon sink, i.e., the underlying causes of its multi-decadal variability.~~ Fully coupled ESMs are another tool to quantify and understand the ocean carbon sinks (e.g., Joos et al., 1999; McNeil and Matear, 2013; Frölicher et al., 2015; Goris et al., 2018; Terhaar et al., 2022b, 2021b). As ESMs are fully coupled and not forced with atmospheric reanalysis data, they do not simulate the same inter-annual internal climate variability as  $p\text{CO}_2$  products and GOBMs do and their biases of the surface ocean physics and biogeochemistry are thus larger than surface ocean biases of GOBMs (Terhaar et al., 2022b; Terhaar et al., 2024). However, ESMs have distinctive advantages compared to  $p\text{CO}_2$  products and GOBMs for the analyses of decadal drivers of the ocean carbon sink because (1) they cover a period of 251 years from 1850 to 2100, (2) cover at least four different future scenarios, and (3) they all have a different internal climate state. The long time-period with different climate states in each model gives ample material to perform statistical analyses and the different future scenarios allow to test the robustness of drivers of the decadal variability of the ocean carbon sink under continuously rising and under strongly decreasing carbon emission trajectories. Furthermore, using different Earth System Models (e.g., Goris et al., 2018; Terhaar et al., 2022b, 2021b; McKinley et al., 2023) in comparison to large ensembles of one ESM (Fay et al., 2023; McKinley et al., 2016) avoids the risk of having a common bias in that one ESM, which might wrongly influence the analysis. Using the ESM ensemble from CMIP6, I will present how potential drivers of the ocean carbon sink, i.e., the atmospheric  $\text{CO}_2$  and its growth rate, ocean heat uptake, and climate variability drive trends in the ocean carbon sink from 1850 to 2100 in these models.

125

## 2 Methods and Datasets

### 2.1 Earth system model ensemble

In this study, I use an ensemble of 12 ESMs from CMIP6 (Table 1). All ESMs from CMIP6 that provide the necessary model output for the following analysis were chosen. For each ESM, only the first ensemble member is used as averaging over multiple ensemble members would have removed variability and using different numbers of ensemble members per ESM would have biased results towards the ESMs with more ensemble members.

**Table 1:** CMIP6 models used in this study and the corresponding model groups.

Model name	Modelling centre	References
ACCESS-ESM1-5	Commonwealth Scientific and Industrial Research Organisation (CSIRO)	(Ziehn et al., 2020)
CanESM5 CanESM5-CanOE	Canadian Centre for Climate Modelling and Analysis	(Christian et al., 2022)
CESM2 CESM2-WACCM	Community Earth System Model contributors	(Danabasoglu et al., 2020)
CMCC-ESM2	Centro euro-Mediterraneo sui Cambiamenti Climatici	(Lovato et al., 2022)
IPSL-CM6A-LR	Institut Pierre Simon Laplace (IPSL)	(Boucher et al., 2020)
MPI-ESM1-2-HR MPI-ESM1-2-LR	Max-Planck-Institute for Meteorology	(Mauritsen et al., 2019; Gutzjahr et al., 2019)
NorESM2-LM NorESM2-MM	Norwegian Climate Centre	(Tjiputra et al., 2020)
UKESM1-0-LL	Met Office Hadley Centre	(Sellar et al., 2020)

### 2.2 Calculating the ocean carbon sink

The annually averaged ocean carbon sink was calculated from concentration-driven historical simulations from CMIP (1850-2014) and four different concentration-driven Shared Socioeconomic Pathways (SSPs) (2015-2100): the low-emission high-mitigation SSP1-2.6, the high-emission low-mitigation SSP5-8.5, and the two intermediate pathways SSP2-4.5 and SSP3-7.0 (Riahi et al., 2017). To account for drifts in the historical and SSP simulations, a linear fit was calculated over the annual carbon sink over the years of the pre-industrial control run that correspond to the years 1850 to 2100 in the historical and SSP simulations. The linear change in the carbon sink in the pre-industrial simulations since 1850 was then subtracted from the historical and SSP simulations.

Furthermore, ESMs have biases in the magnitude of the ocean carbon sink due to biases their respective circulation and surface ocean carbonate chemistry that also affect the size of the decadal trends, i.e., a negative bias in the magnitude of the carbon sink also introduces a negative bias in the decadal trends. To statistically compare the decadal trends of the carbon sink over

the here-used ESM ensemble, the global estimate of the ocean carbon sink was adjusted for each ESM with respect to biases in its circulation and surface ocean carbonate chemistry following Terhaar et al. (2022). First, the Revelle factor, the inter-frontal Southern Ocean sea surface salinity, and the AMOC strength were calculated for each model. Afterwards, a multi-linear fit was performed with the three observation-based quantities as predictors and the average ocean carbon sink from 1850 to 2100 as target variables (the period from 2015 to 2100 was used four times for each of the four SSPs). Finally, the biases in each predictor with respect to observation-based estimates of these predictors are calculated and used to adjust the simulated ocean carbon sink based on the determined constants from the multi-linear fit. Overall, this result in an adjustment of  $10 \pm 7\%$  (i.e., increased ocean carbon uptake) for the here used model ensemble. The adjustment corrects for known biases in the models' circulations and surface ocean carbonate chemistry and hence reduced differences in the overall magnitude of the simulated carbon sink between ESMs (Terhaar et al., 2022b). This reduction in the difference in the magnitude of the carbon sink also reduces differences between the magnitude of trends and slightly improves the relationships found here as it ( $r^2$  in Figure 3 would have been 0.83 without adjustment instead of 0.91 with adjustment). Nevertheless, the results would quantitatively and qualitatively almost identical with and without that adjustment.

160

The ocean carbon sink was also calculated for each of the five major ocean basins (Atlantic Ocean, Pacific Ocean, Indian Ocean, Arctic Ocean, and Southern Ocean) using the RECCAP2 biome mask (DeVries et al., 2023; Terhaar et al., ~~in~~ [press2024](#)), which is a slightly adapted version of a previously developed biome mask (Fay and McKinley, 2014). Regionally, no bias adjustments were performed as it still remains largely unclear how biases in circulation and carbonate chemistry affect the regional ocean carbon sink estimates.

165

### 2.3 Atmospheric CO<sub>2</sub> and growth rate

The annually averaged atmospheric CO<sub>2</sub> over the historical period and for each SSP was taken from the CMIP6 forcing files (Meinshausen et al., 2020, 2017). The atmospheric CO<sub>2</sub> growth rate in each year was calculated as the difference in atmospheric CO<sub>2</sub> in that year and the year before.

### 170 2.4 Estimating the effect of climate change and ocean heat uptake on the ocean carbon sink

The effect of climate change and ocean heat uptake on the ocean carbon sink in ESMs was calculated based on additional idealized simulations provided by five of the twelve ESMs in the ensemble (ACCESS-ESM1-5, CanESM5, MRI-ESM2-0, NorESM2-LM, UKESM1-0-LL) within the CMIP6 framework. These five ESMs made historical simulations, called 'bgc', where the change in atmospheric CO<sub>2</sub> had no effect on climate change but the carbon cycle still 'sees' the increase in atmospheric CO<sub>2</sub>. However, other non-CO<sub>2</sub> radiative agents (aerosols, CH<sub>4</sub>, N<sub>2</sub>O, etc.) still ~~effect~~ [affect](#) the climate in these simulations. These 'bgc' simulations were only made for SSP5-8.5 ('ssp585-bgc') and not for the other SSPs. The difference of the normal historical and SSP5-8.5 simulations (including effects from CO<sub>2</sub> and non-CO<sub>2</sub> radiative agents) and the additional 'bgc' simulations quantifies the heat and carbon fluxes that are solely due to the CO<sub>2</sub>-induced climate change and warming.

175

## 180 2.5 Climate modes

To assess the climate variability across the ensemble of the ESM, annual averages of three climate modes were calculated for each ESM over the 251 years of the pre-industrial control simulation: (1) The Atlantic Multi-decadal Oscillation (AMO), (2) the Niño 3.4 index, and (3) the Marshall Southern Annular Mode (SAM) index. The AMO was calculated based on SST anomalies in the North Atlantic between 0 and 80°N. The Niño 3.4 index was calculated based on SST anomalies in the tropical  
185 Pacific region from 5°S to 5°N and from 170°W to 120°W. The Marshall SAM index was calculated as anomalies of the zonal pressure difference between the latitudes of 40S and 65S. Anomalies for each index in ESMs were calculated by removing a linear fit over the 251 years of the pre-industrial control simulation.

In addition, observation-based estimates of each climate mode were used. The observation-based AMO index  
190 ([https://climatedataguide.ucar.edu/sites/default/files/2022-03/amo\\_monthly.txt](https://climatedataguide.ucar.edu/sites/default/files/2022-03/amo_monthly.txt)) and the Niño 3.4 index ([https://psl.noaa.gov/gcos\\_wgsp/Timeseries/Data/nino34.long.data](https://psl.noaa.gov/gcos_wgsp/Timeseries/Data/nino34.long.data)) are based on HadISST1 (Rayner et al., 2003). The Marshall SAM index has been calculated based on twelve stations, six stations at ~40°S and six stations at ~65°S (Marshall, 2003). To compare each observation-based index to the simulated index in the pre-industrial control simulations, the timeseries of the observation-based indexes have been detrended by subtracting a linear trend over the respective observation-based index  
195 estimate.

## 2.6 Decadal trends

Decadal trends of different variables are here defined as the slope of linear fits over ten years.

## 2.7 Coefficient of determination, p-values, and Bayes factor

To determine the strength of correlations, the coefficient of correlation was calculated throughout this study ( $r^2$ ). In addition,  
200 the p-value was calculated to test the hypothesis that the trend change in atmospheric CO<sub>2</sub> from one decade to another decade is a significant driver of trends of the global and regional ocean carbon sink. A p-value larger than 0.1 indicates little or no evidence for that hypothesis exists, a p-value from 0.1 to 0.05 indicates weak evidence or a suggestion of evidence, a p-value from 0.05 to 0.01 indicates evidence or modest evidence and a p-value from 0.01 to 0.001 indicates strong evidence (Held and Ott, 2018). In addition, an upper bound for the Bayes factor can be calculated following Halsey (2019). Throughout the  
205 manuscript, the p-values are never larger than 1e-89 resulting in Bayes factors that are at least 1e86. Based on the Bayes factor the hypothesis that the trend change in atmospheric CO<sub>2</sub> from one decade to another decade is a significant driver of trends of the global and regional ocean carbon sink is 1e86 more likely than the hypothesis that the trend change in atmospheric CO<sub>2</sub> from one decade to another decade is not a significant driver of trends of the global and regional ocean carbon sink. As p-values are that small and Bayes factors are that high, I simply refrain to report that the p-values are smaller than 0.001.

### 3 The influence of atmospheric CO<sub>2</sub> on the ocean carbon sink


#### 3.1 Atmospheric CO<sub>2</sub>

Over the historical period of CMIP6 simulations from 1850 to 2014, the annually averaged global ocean carbon sink has increased approximately proportional to the rise in atmospheric CO<sub>2</sub> (Figure 1a, b, c). Due to the exponential rise in atmospheric CO<sub>2</sub>, the cumulative ocean carbon sink is also approximately proportional to the rise in atmospheric CO<sub>2</sub>. However, these quasi-linear relationships did not hold from 1920 (atmospheric CO<sub>2</sub> of 304 ppm) to 1960 (317 ppm) and from 1990 (354 ppm) to ~~2000 (369 ppm)~~ 1995 (360 ppm) when the ocean carbon sink did not increase while the atmospheric CO<sub>2</sub> continued to increase. These periods manifest themselves as ‘jumps’ in the linear relationship between the atmospheric CO<sub>2</sub> and the cumulative ocean carbon sink (indicated by light grey shadings in Figure 1d).


After 2014, when the historical period in CMIP6 ends and SSPs start, the link between atmospheric CO<sub>2</sub> and the ocean carbon sink depends strongly on the future scenario of atmospheric CO<sub>2</sub>. The linear relationship between the annually averaged carbon sink and atmospheric CO<sub>2</sub> breaks down under all scenarios. Under SSP5-8.5, a pathway with continuous increase in emissions (Riahi et al., 2017) and exponentially growing atmospheric CO<sub>2</sub> (Fig. 1a), the increase in the ocean carbon sink per increase in atmospheric CO<sub>2</sub> reduces until the ocean carbon sink reaches a maximum just above 6 Pg C yr<sup>-1</sup> (Fig 1b), which is not exceeded even if atmospheric CO<sub>2</sub> rises (Fig 1e). Under SSP3-7.0, a pathway with slightly smaller emissions and atmospheric CO<sub>2</sub> than SSP5-8.5, the ocean carbon sink also converges to a maximum but at around 5 Pg C yr<sup>-1</sup>. Under SSP2-4.5, CO<sub>2</sub> emissions start to decline around 2050 (Riahi et al., 2017) and atmospheric CO<sub>2</sub> stabilizes around 600 ppm by 2100 (Fig 1a). Although atmospheric CO<sub>2</sub> stabilizes, the ocean carbon sink reduces strongly (Fig 1b). Under SSP1-2.6, atmospheric CO<sub>2</sub> does not only stabilize but starts to reduce by 2080, leading to a strong reduction of the ocean carbon sink (Fig. 1a). In comparison, the relationship between the cumulative ocean carbon sink and atmospheric CO<sub>2</sub> remains almost linear in the two high-emission pathways (SSP3-7.0 and SSP5-8.5) although the slope reduces with warming (Fig. 1f). For the two low-emission pathways (SSP1-2.6 and SSP2-4.5), the relationship breaks down as the ocean continuously takes up carbon, even when atmospheric CO<sub>2</sub> stabilizes and decreases (Fig. 1e, f).




(a)




(b)




(c)




(d)




(e)



(f)





**Figure 1: The relationship between atmospheric CO<sub>2</sub> and the global ocean carbon sink.** (a) The annually averaged atmospheric CO<sub>2</sub> that was used to force the ESMs from CMIP6 based on observation-based estimates from 1850 to 2014 (black) and based on four different SSPs (SSP1-2.6 in blue, SSP2-4.5 in orange, SSP3-7.0 in red, and SSP5-8.5 in brown) from 2015 to 2100. (b) The resulting ocean carbon sink as simulated by 12 ESMs (Table 1) after being adjusted for biases in circulation and surface ocean carbonate chemistry following Terhaar et al. (2022). The thick lines indicate multi-model means and the shading the 1- $\sigma$  standard deviation across the model ensemble. Relationships between atmospheric CO<sub>2</sub> and the annually averaged ocean carbon sink (c) for the historical period until 2014 and (e) for the 21<sup>st</sup> century from 2015 onwards, as well as between atmospheric CO<sub>2</sub> and the cumulative ocean carbon sink (d) for the historical period until 2014 and (f) for the 21<sup>st</sup> century from 2015 onwards. The light grey shadings in (a) – (d) indicate the time periods from 1920 to 1960 and from 1990 to 1995.


240

245

### 3.2 Atmospheric CO<sub>2</sub> growth rate

As an alternative to the atmospheric CO<sub>2</sub>, the growth rate of atmospheric CO<sub>2</sub> was proposed as a key driver for the strength of the ocean carbon sink (McKinley et al., 2017, 2020). Over the historical period, the atmospheric CO<sub>2</sub> growth rate appears to be weakly linearly related to the square root of the strength of the ocean carbon sink (Fig 2c). This relationship weakens after ~1960 when the prescribed atmospheric CO<sub>2</sub> growth rate is based on direct atmospheric CO<sub>2</sub> observations and not, as before, on relatively smooth observation-based estimates from proxies (light grey shading in Fig. 2a and grey dots in Fig. 2c, d). The direct observations capture the strong inter-annual variability of the atmospheric CO<sub>2</sub> growth rate that cannot be reconstructed by observation-based estimates from proxies. However, even this relationship between the atmospheric CO<sub>2</sub> growth rate and the square root of the strength of the ocean carbon sink breaks down in the 1920s and 1940s (pale green dots in Fig. 2c, d) when the growth rate is around zero over around a decade each time (pale green shading in Fig. 2a), but the ocean carbon sink does not go back close to zero but remains almost stable (pale green shading in Fig. 2b).


Over the 21<sup>st</sup> century, the relationship between the ocean carbon sink and the atmospheric CO<sub>2</sub> growth rate breaks down (Fig 2c). As long as CO<sub>2</sub> emissions and atmospheric CO<sub>2</sub> growth rate rise, as they do under SSP3-7.0 and SSP5-8.5 (Fig 2a), the relationship flattens (Fig. 2a). However, the strength of the relationship varies between SSP3-7.0 and SSP5-8.5. Under SSP5-8.5, the relationship also breaks down in the last two decades when the atmospheric CO<sub>2</sub> growth stabilizes but the ocean



carbon sink weakens.

Under SSP1-2.6 and SSP2-4.5 with declining emissions or declining and even negative atmospheric CO<sub>2</sub> growth rates, the ocean carbon sink reduces but not along the same path as it increased over the historical period (Fig. 2c).

265



270

275

**Figure 2: The relationship between the atmospheric CO<sub>2</sub> growth rate and the global ocean carbon sink.** (a) The annually averaged atmospheric CO<sub>2</sub> growth rate based on atmospheric CO<sub>2</sub> forcing files from CMIP6, which are based on observation-based estimates from 1850 to 2014 (black) and based on four different SSPs (SSP1-2.6 in blue, SSP2-4.5 in orange, SSP3-7.0 in red, and SSP5-8.5 in brown) from 2015 to 2100. (b) The ocean carbon sink as simulated by 12 ESMs (Table 1) after being adjusted for biases in circulation and surface ocean carbonate chemistry following Terhaar et al. (2022). The thick lines indicate multi-model means and the shading the 1- $\sigma$  standard deviation across the model ensemble. Relationships between atmospheric CO<sub>2</sub> growth rate and the annually averaged ocean carbon sink (c) for the entire period from 1850 to 2100 and (d) only for historical period until 2014. -The light grey shading in (a) indicates the period where direction atmospheric CO<sub>2</sub> observations are available and the pale green shading in (a) and (b) and the pale green dots in (c) and (d) indicate the 1990s and 1940s. The zero growth rate and ocean carbon sink in (a) and (b) are shown as black dashed lines.

280

Over the 21<sup>st</sup>-century, the relationship between the ocean carbon sink and the square root of the atmospheric CO<sub>2</sub> growth rate depends on the scenario and breaks down under the two low emission scenarios (SSP1-2.6 and SSP2-4.5) (Fig 2e). As long as CO<sub>2</sub> emissions and atmospheric CO<sub>2</sub> growth rate rise, as they do under SSP3-7.0 and SSP5-8.5 (Fig 2a), the relationship holds (Fig. Under SSP1-2.6 and SSP2-4.5 and declining emissions and declining and even negative atmospheric CO<sub>2</sub> growth rates, the ocean carbon sink reduces but not along the same path as it increased over the historical period (Fig. 2e).

### 3.3 Changes in atmospheric CO<sub>2</sub> growth rate determine changes in decadal trends of the ocean carbon sink

#### 3.3.1 Global relationship

285 Although neither the atmospheric CO<sub>2</sub> nor its growth rate can quantify the strength of the ocean carbon sink over various time period and different trajectories of atmospheric CO<sub>2</sub>, the atmospheric CO<sub>2</sub> growth rate can nevertheless be used to understand changes in the ocean carbon sink on decadal timescales, i.e., decadal trends of the ocean carbon sink. For the period from 1980 to 2018, it has been shown that a ~~smaller increase~~ slowing of the growth rate in comparison to a linear trend has led to a stagnation of the increase of the ocean carbon sink and that an accelerated increase of the growth rate has led to a strongly increasing carbon sink (McKinley et al., 2020).

290


Over longer time periods and different future SSPs, ESMs provide more such examples where changes in the growth rate of atmospheric CO<sub>2</sub> led to changes in the decadal trends of the simulated ocean carbon sink (Figure 2). Around 1915, the atmospheric CO<sub>2</sub> growth rate changes from an increase to a decrease, at the same time the ocean carbon sink stops increasing and starts to decrease. Then in 1930, the atmospheric CO<sub>2</sub> growth rate increases, and the ocean carbon sink also starts to increase simultaneously. Then, in 1940 the atmospheric CO<sub>2</sub> growth rate decreases again, and the ocean carbon sink also decreases at the same time. Similarly, the atmospheric CO<sub>2</sub> growth rate changes from a positive trend to a negative trend in 1990, exactly when the ocean carbon sink also starts to slow down. When the atmospheric CO<sub>2</sub> growth rate increases again, the ocean carbon sink also increases. Over the 21<sup>st</sup> century, the same relationship continues. Under SSP2-4.5, the atmospheric CO<sub>2</sub> growth rate slows ~~down~~ until 2050 and the positive trend in the ocean carbon sink weakens. Once the atmospheric CO<sub>2</sub> growth declines, the trend in the ocean carbon sink becomes negative. ~~However, over longer time periods and under decreasing atmospheric CO<sub>2</sub>,~~

295  
300

~~Although a~~ As a slowing or acceleration of the growth rate in comparison to a theoretical linear trend as in McKinley et al. (2020) is not anymore possible over ~~longer time periods and under decreasing~~ exponential growth or when atmospheric CO<sub>2</sub> peaks, I here generalize the idea of McKinley et al. (2020) that a slowing or acceleration of the atmospheric CO<sub>2</sub> growth rate drives the trends of the ocean carbon sink by defining such slowing or acceleration as the difference in the growth rate in a given decade with respect to the preceding decade. When defining slowing or acceleration of the atmospheric CO<sub>2</sub> growth rate that way, a clear relationship ( $r^2=0.91$ ) emerges indeed over the entire historical period and all four future scenarios over the 21<sup>st</sup> century (excluding years where the ocean carbon sink exceeds 4.5 Pg C yr<sup>-1</sup>) between changes in the atmospheric CO<sub>2</sub> growth rate ~~from one decade to another~~ and the decadal trend of the multi-model average of the ocean carbon sink (Fig. 3). Changes in the atmospheric CO<sub>2</sub> growth rate are here defined as the change in trends (linear fit over one decade) of atmospheric CO<sub>2</sub> from one decade to the next. Trends in the ocean carbon sink are a linear fit over annual values of the global ocean carbon sink in the 2<sup>nd</sup> decade. It thus appears that it is the change in the growth rate in comparison to the previous decade that appears to drive the decadal trends of the ocean carbon sink and not the difference to an expected linear trend. If, for example, the

305  
310

315 growth rate strongly reduced from one decade to another, the ocean carbon sink would show a negative trend. If the growth rate than stays at that lower level, the carbon sink would not decline further but stabilize at its new level. This relationship even holds when CO<sub>2</sub> emissions decline strongly as under SSP1-2.6.



320

**Figure 3: The relationship between changes in the atmospheric CO<sub>2</sub> growth rate and decadal trends of the global ocean carbon sink for the multi-model mean.**

325

(a) Decadal trends of the multi-model mean ocean carbon sink compared to changes in decadal trends in atmospheric CO<sub>2</sub>, which represent the decadal averaged growth rate of atmospheric CO<sub>2</sub>. The dark blue to yellow circles without a surrounding black line show multi-model averages for all years of the historical period from 1850 to 2014 and for all years from 2015 to 2100 for all four SSPs. All decades over from 1850 to 2100 are shown, i.e., 2000-2009, 2001-2010, 2002-2011, etc. The brown line shows a linear fit for all years when the global ocean carbon sink is smaller than 4.5 Pg C yr<sup>-1</sup> and the brown shading is the 1-σ projection uncertainty. The dots with black lines around them show values from the respective ensemble means of the pCO<sub>2</sub> products (pink) and GOBMs (orange) from the Global Carbon Budget 2023 (Friedlingstein et al., 2023) for the three decades between 1990 and 2020. Small deviations from the relationship in SSP1-2.6 are marked by 'SSP1-2.6'.

330

(b) The simulated ocean carbon sink in comparison to the expected ocean carbon sink based on the relationship in (a) and the prescribed trend change in atmospheric CO<sub>2</sub> in the simulations.

However, this relationship between changes in the atmospheric CO<sub>2</sub> growth rate and the decadal trend of the multi-model average of the ocean carbon sink breaks down if the ocean carbon sink is larger than 4.5 Pg C yr<sup>-1</sup> (Figure 3, r<sup>2</sup> starts to reduce if years with an ocean carbon sink larger than 4.5 Pg C yr<sup>-1</sup> are included). The breakdown likely occurs because climate change and associated ocean heat uptake and circulation changes become so large that effects on the natural carbon sink reduce the trend in the ocean carbon sink substantially enough. Thus, it is not the carbon uptake of 4.5 Pg C yr<sup>-1</sup> itself that causes the breakdown of the relationship but the combined impact of an increasing Revelle factor (Revelle and Suess, 1957) and climate change (Joos et al., 1999; McNeil and Matear, 2013; Frölicher et al., 2015). In SSPs from CMIP6, the combined impact is large enough to affect the here identified relationship when the ocean carbon sink is around 4.5 Pg C yr<sup>-1</sup>. The breakdown of the relationship also implies that the decadal trends in the ocean carbon sink cannot exceed 0.78±0.10 Pg C yr<sup>-1</sup> dec<sup>-1</sup> (the uncertainty is the 1-σ standard deviation across the ESM ensemble in the decade when the multi-model mean decadal trend is largest). Thus, if the ocean carbon sink is below 4.5 Pg C yr<sup>-1</sup> and if its magnitude 10 years ago and the change in the decadal trends of atmospheric CO<sub>2</sub> between the last two decades (20 to 10 years ago and 10 years to now) is known, the absolute ocean carbon sink this year can be determined (Fig 3b).

345

### 3.3.2 Regional relationships

The relationship between changes in the atmospheric CO<sub>2</sub> growth rate and the global ocean carbon sink holds in all large five ocean basins (Fig. 4a-j) as it has also done from 1980 to 2018 (McKinley et al., 2020). The ~~correlation~~ coefficient of determination (r<sup>2</sup>) is larger than 0.84 in the Atlantic, Pacific, Indian, and Southern Ocean. Only the Arctic Ocean has a smaller correlation coefficient of 0.66.

350

In the Arctic Ocean, the carbon sink has been shown to be already substantially more affected by climate change than in any other ocean basin (Yasunaka et al., 2023). In the future, when sea ice will disappear and the Arctic will continue to warm faster than any other region, the importance of climate change for the Arctic Ocean carbon sink will likely remain relatively large, for example through freshening (Terhaar et al., 2021a) and a change in the seasonal cycle of pCO<sub>2</sub> (Orr et al., 2022), and hence reduce the importance of changes in the atmospheric CO<sub>2</sub> for trends in the ocean carbon sink.

355


In the Southern Ocean, the simulated trends in the ocean carbon also slightly differ from the expected trends based on changes in trends of atmospheric CO<sub>2</sub> in three brief periods (Fig. 4i). From 1995 to 2005 over the historical period and from 2030 to 2050 under SSP1-2.6, the decadal trend in the ocean carbon sink is larger than expected, whereas it is smaller than expected from 2080 to 2100 under SSP1-2.6. The differences under SSP-1.2.6 are even visible for the global carbon sink (Fig 3a). As the difference is occurring in the multi-model mean, it appears to be a forced response and not a response that is linked to the particular state of the climate in one of the models. The time periods where the differences are visible globally (2030-2050 and 2080 to 2100 under SSP1-2.6) are the times when the growth in atmospheric CO<sub>2</sub> stops and when it starts to decrease in that

360

365 scenario (Fig 1c). As the atmospheric CO<sub>2</sub> growth rate changes quickly in these periods (Fig. 2a), first by changing into a  
decreasing phase and then transitioning into a stabilizing phase. ~~It~~, it appears that a fast transition of the trend change in  
atmospheric CO<sub>2</sub> temporarily leads to differences in the expected relationship. If the trend change in atmospheric CO<sub>2</sub>  
decreases fast, the trend in ocean carbon sink remains larger than expected and if the trend change in atmospheric CO<sub>2</sub> increases  
fast, the trend in ocean carbon sink remains smaller than expected. However, the drivers behind the divergence from the  
370 expected decadal trend of the multi-model mean in from 1995 to 2005 in the Southern Ocean remain unclear and should be  
analysed in future research.

Despite these small differences, the overall relationship between changes in decadal trends in the atmospheric CO<sub>2</sub> and decadal  
trends in the local and global ocean carbon sink is very strong ( $r^2 > 0.84$ , apart from the Arctic Ocean) and demonstrates how  
375 atmospheric CO<sub>2</sub> is the main driver of the externally forced decadal trends of the ocean carbon sink.





380

**Figure 4: The relationship between changes in the atmospheric CO<sub>2</sub> growth rate and decadal trends of the ocean carbon sink in the five major ocean basins.** Decadal trends in the ocean carbon sink in the (a) Atlantic Ocean, (c) Pacific Ocean, (e) Indian Ocean, (g) Arctic Ocean, and (i) Southern Ocean compared to changes in decadal trends in atmospheric CO<sub>2</sub>, which represent the decadal averaged growth rate of atmospheric CO<sub>2</sub>. The dark blue to yellow circles without a surrounding black line show multi-model averages for all years of the historical period from 1850 to 2014 and for all years from 2015 to 2100 for all four SSPs. The brown line shows a linear fit for all years when the global ocean carbon sink is smaller than 4.5 Pg C yr<sup>-1</sup> and the brown shading is the 1-σ projection uncertainty. The dots with black lines around them show values from *p*CO<sub>2</sub> products (pink) and GOBMs (orange) from the Global Carbon Budget 2023 (Friedlingstein et al., 2023) for the three decades between 1990 and 2020. The simulated ocean carbon sink in the (b) Atlantic Ocean, (d) Pacific Ocean, (f) Indian Ocean, (h) Arctic Ocean, and (j) Southern Ocean in comparison to the expected ocean carbon sink based on the respective relationships in (a, c, e, g, i) and the prescribed trend change in atmospheric CO<sub>2</sub> in the simulations. The *p*-value for each regional relationship is smaller than 0.001.

385


390

|

#### 4 The importance of climate variability on decadal trends on the ocean carbon sink

395 Internal climate and ocean variability in ESMs can strongly affect the decadal trends in ESMs (Li and Ilyina, 2018) and hence reduces the strength of the relationship between changes in decadal trends in the atmospheric CO<sub>2</sub> and decadal trends in the ocean carbon sink. To quantify the importance of climate variability, I calculated the relationship between changes in decadal trends in the atmospheric CO<sub>2</sub> and decadal trends in the ocean carbon sink not for the multi-model mean but for the individual ESMs. When calculating the linear fit over the results from the individual ESMs, the correlation factor only slightly reduces from  $r^2=0.91$  to  $r^2=0.80$  (Fig 5). Although Li and Ilyina (2018) showed that 50-70 ensemble members are needed to remove internal variability for one given decade, the relationship here remains strong because it is based on at least 34 decades for each ESM (16 historical plus 18 future decades (SSP1-2.6 and SSP2-4.5)). Thus, two ESMs alone already provide the necessary 50-70 decades that were shown to be necessary (Li and Ilyina, 2018). The 1- $\sigma$  prediction interval around the linear fit is 0.16 Pg C yr<sup>-1</sup> dec<sup>-1</sup>, meaning that 68% of all trends will be within  $\pm 0.16$  Pg C yr<sup>-1</sup> dec<sup>-1</sup> of the predicted trend based on decadal trends in the atmospheric CO<sub>2</sub>, 95 % will be within  $\pm 0.31$  Pg C yr<sup>-1</sup> dec<sup>-1</sup> of the predicted trend, and virtually all trends (99.7%) will be within  $\pm 0.47$  Pg C yr<sup>-1</sup> dec<sup>-1</sup> of the predicted trend. The largest simulated trend in the ocean carbon sink in one of the ESMs is 0.96 Pg C yr<sup>-1</sup> dec<sup>-1</sup>. This is within the 2- $\sigma$  range of the largest trend as diagnosed by the multi-model mean  $0.78 \pm 0.10$  Pg C yr<sup>-1</sup> dec<sup>-1</sup>.

410



415


420

**Figure 5: The relationship between changes in the atmospheric CO<sub>2</sub> growth rate and decadal trends of the global ocean carbon sink for individual ESMs.** (a) Decadal trends in the ocean carbon sink for all ESMs individually compared to changes in decadal trends in atmospheric CO<sub>2</sub>, which represent the decadal averaged growth rate of atmospheric CO<sub>2</sub>. The dark blue to yellow circles without a surrounding black line show multi-model averages for all years of the historical period from 1850 to 2014 and for all years from 2015 to 2100 for all four SSPs. The brown line shows a linear fit for all years when the global ocean carbon sink is smaller than 4.5 Pg C yr<sup>-1</sup> and the brown shading is the 1- $\sigma$  projection uncertainty. The dots with black lines around them show values from individual pCO<sub>2</sub> products (pink) and GOBMs (orange) from the Global Carbon Budget 2023 (Friedlingstein et al., 2023) for the three decades between 1990 and 2020. (b) The simulated ocean carbon sink in comparison to the expected ocean carbon sink based on the relationship in (a) and the prescribed trend change in atmospheric CO<sub>2</sub> in the simulations.

The range of simulated trends in ocean carbon sink with different internal climate variability encompasses the ocean carbon sink trend estimates of GOBMs from the Global Carbon Budget 2023 (Friedlingstein et al., 2023) but the trend estimates of the  $p\text{CO}_2$  products exceed the range that is simulated by ESMS. For the decade from 1990 to 1999, 7 out of 10 GOBMs fall within the  $\pm 1\text{-}\sigma$  range and the remaining 3 GOBMs fall within the  $\pm 2\text{-}\sigma$  range. In comparison, only 3 out of 8  $p\text{CO}_2$  products fall within the  $\pm 1\text{-}\sigma$  range, 2  $p\text{CO}_2$  products fall within the  $\pm 3\text{-}\sigma$  range, 1  $p\text{CO}_2$  product falls within the  $\pm 4\text{-}\sigma$  range, 2  $p\text{CO}_2$  products fall within the  $\pm 5\text{-}\sigma$  range. For the decade from 2000 to 2009, 5 GOBMs fall within the  $\pm 1\text{-}\sigma$  range, 4 GOBMs fall within the  $\pm 2\text{-}\sigma$  range, and the final GOBM falls within the  $\pm 3\text{-}\sigma$  range. In comparison, only 2  $p\text{CO}_2$  products fall within the  $\pm 1\text{-}\sigma$  range, 1  $p\text{CO}_2$  product falls within the  $\pm 2\text{-}\sigma$  range, 2  $p\text{CO}_2$  products fall within the  $\pm 3\text{-}\sigma$  range, 1  $p\text{CO}_2$  products fall within the  $\pm 5\text{-}\sigma$  range, 1  $p\text{CO}_2$  products fall within the  $\pm 6\text{-}\sigma$  range, and 1  $p\text{CO}_2$  product falls within the  $\pm 7\text{-}\sigma$  range. For the decade from 2010 to 2019, 9 GOBMs fall within the  $\pm 1\text{-}\sigma$  range and the remaining one falls within the  $\pm 2\text{-}\sigma$  range. In comparison, only 1  $p\text{CO}_2$  products fall within the  $\pm 1\text{-}\sigma$  range, 2  $p\text{CO}_2$  products fall within the  $\pm 2\text{-}\sigma$  range, 3  $p\text{CO}_2$  products fall within the  $\pm 3\text{-}\sigma$  range, 1  $p\text{CO}_2$  product falls within the  $\pm 4\text{-}\sigma$  range, and 1  $p\text{CO}_2$  product falls within the  $\pm 5\text{-}\sigma$  range. In general, the  $p\text{CO}_2$  product estimates of the decadal trends are not randomly distributed across the possible range that the ESMS suggest. Instead,  $p\text{CO}_2$  products systematically overestimate the magnitude of the respective trends that is suggested by ESMS, i.e., a too small negative trend in the 1990s and a too high positive trend in the 2000s and 2010s.

## 5 Imprint of climate change and ocean heat uptake on the ocean carbon sink

In addition to atmospheric CO<sub>2</sub> and internal climate variability, climate change and ocean heat uptake also affects the ocean carbon sink and potentially its decadal trends. The ocean heat uptake, for example, causes ~~to~~ changes in the ocean circulation such as stratification and outgassing of natural carbon from the ocean due to increasing temperatures and reduced solubility (Fig. 6). Across the five ESMs that performed the simulations to quantify the effect of ocean heat uptake on the natural carbon in the ocean (see Methods), the loss of natural carbon from the ocean to the atmosphere is related to the ocean heat uptake via a 2<sup>nd</sup> degree polynomial function under strong radiative forcing (SSP5-8.5) (Fig. 6a). Although annual variability hides part of this relationship, the relationship emerges strongly for decadal averages (Fig. 6b).



**Figure 6: The relationship between natural carbon loss and ocean heat uptake and its link to the Atlantic Meridional Overturning Circulation.** (a) Annually and (b) decadal averaged air-sea CO<sub>2</sub> flux solely caused by ocean heat uptake (for details see Methods) vs the annually averaged ocean heat uptake in five different ESMs (ACCESS-ESM1-5 in blue, CanESM5 in orange, MRI-ESM2-0 in green, NorESM2-LM in purple, and UKESM1-0-LL in brown). A 2<sup>nd</sup> degree polynomial function (coloured lines) was fitted over the decadal averaged values of each ESM. (c) The CO<sub>2</sub> flux per ocean heat uptake, calculated for each model from the fitted 2<sup>nd</sup> degree polynomial function at an ocean heat uptake at 35 ZJ yr<sup>-1</sup>, against the Atlantic Meridional Overturning Circulation (AMOC) at 26.5°N calculated in each ESM from 2004 to 2018 (historical plus SSP5-8.5 simulations). The vertical dashed black line and the grey shading indicate the averaged observed AMOC at the RAPID array from 2004 to 2018 (McCarthy et al., 2020).

While each of the five ESMs suggests that the loss of natural carbon from the ocean to the atmosphere is related to the ocean heat uptake via a 2<sup>nd</sup> degree polynomic function, the amount of carbon loss due per ocean heat uptake varies across ESMs (Fig 460 5b). The main reason for differences likely the different changes in ocean circulation and stratification due to ocean heat uptake in each ESM. One of the parts of the ocean overturning circulation that is expected to change strongly with climate change and ocean warming in ESMs is the Atlantic Meridional Overturning Circulation (Weijer et al., 2020). Across a large ensembles of ESMs from CMIP6, it has been shown that ESMs with an already stronger Atlantic Meridional Overturning Circulation also show a stronger reduction in the Atlantic Meridional Overturning Circulation (Weijer et al., 2020). The larger overturning 465 reduction thus causes the models with a higher Atlantic Meridional Overturning Circulation to lose more natural carbon loss for the same heat uptake (Fig. 5c). Based on this linear relationship, it would be possible to constrain the loss of carbon per heat uptake with observations of the present-day Atlantic Meridional Overturning Circulation. However, using only five ESMs to quantify a linear relationship is likely not yielding a robust relationship so that I abstain from constraining the loss of carbon per heat uptake. Nevertheless, the observed Atlantic Meridional Overturning Circulation at 26.5°N is close to the average of 470 the simulated Atlantic Meridional Overturning Circulation at 26.5°N in ESMs suggesting that the multi-model average sensitivity of air-sea CO<sub>2</sub> fluxes to heat uptake are a good approximation of the real-world sensitivity.


Unfortunately, CMIP6 only provides simulations that allow to quantify the ocean natural carbon sink response to ocean warming for SSP5-8.5 and not for other scenarios where the ocean warming slows down or even stabilizes. Thus, it remains 475 impossible for now to quantify the effect of ocean heat uptake for other scenarios and to test if the here identified relationship is robust. However, as differences in the decline of the Atlantic Meridional Overturning Circulation are similar across all these scenarios although the ocean heat uptake is much smaller in the low emission scenarios (Weijer et al., 2020), the sensitivity of carbon loss to heat uptake might be larger in low-emission scenarios. As such, changes in the ocean heat uptake and its trend might well cause changes in the anthropogenic ocean carbon sink via the outgassing of marine natural carbon pool. Although 480 these changes are likely small as decadal averaged ocean heat uptake does not change quickly, these changes might still be partly responsible for differences between the decadal trend of the ocean carbon sink that were expected based on changes in trend of atmospheric CO<sub>2</sub> and the simulated ocean carbon sink, especially those in SSP1-2.6 globally (Fig. 3) and in the Southern Ocean (Fig. 4i). To verify this hypothesis, CMIP simulations that allow to quantify the ocean natural carbon sink response to ocean warming would have to be made for other scenarios than SSP5-8.5.

485

## 6 Potential caveats and limitations

The strong dependence of decadal trends in ocean carbon sinks on the change of the atmospheric CO<sub>2</sub> growth rate from one decade to the other was here identified across an ensemble of state-of-the-art ESMs from CMIP6. The robustness of this relationship depends on the model's ability to represent the internal climate variability and might also be biased if the entire  
490 model ensemble is biased, for example due to relatively coarse resolution or a common unrealistic representation of the physics or biogeochemistry.

If, for example, the internal climate variability on decadal timescales was underestimated by the here-used ESMs, the predictability of the decadal trends in the ocean carbon sink by changes in the growth rate of CO<sub>2</sub> would be overestimated. A  
495 prerequisite for ESMs to simulate the contribution of the natural variability to decadal trends of the ocean carbon sink is that they also simulate the size of the decadal trends of internal climate modes that are known to affect the variability of the ocean carbon sink most. The major climate modes that are known to influence the decadal variability of the ocean carbon sink are the Niño 3.4 index (Feely et al., 1999; Ishii et al., 2014; McKinley et al., 2004), Atlantic Multi-decadal Oscillation (Breedon and McKinley, 2016; Keppler et al., 2023), and Marshall Southern Annular Mode (Le Quéré et al., 2007; Gruber et al., 2019b;  
500 Landschützer et al., 2015; Lovenduski et al., 2008; Thompson and Solomon, 2002; Lenton and Matear, 2007; Hauck et al., 2013). The decadal trends of the Niño 3.4 index in ESMs is 17 (-19 to 53) % larger than the decadal trends of the observation-based estimates of the Niño 3.4 index (the numbers in parenthesis indicates the standard deviation across ESMs) (Fig 7b), the decadal trends of the Atlantic Multi-decadal Oscillation in ESMs are 13 (-7 to 33) % larger than the decadal trends in the observation-based estimate (Fig 7c), and the decadal trends of the Marshall Southern Annular Mode in ESMs are 52 (30 to  
505 75) % larger than the decadal trends in the observation-based estimate (Marshall, 2003) (Fig 7d). The relatively large decadal trends of climate modes in ESMs suggest that the ESMs are indeed capable of simulating the internal climate variability on decadal timescales. ~~Thus~~ **As the decadal trends in climate mode are larger or equal to the observed ones**, there is no indication that the decadal variability of the ocean carbon sink in ESMs (Fig 7a) might be too small because of a too small internal climate variability in ESMs as previously hypothesized ~~(by Gruber et al., (2023),)~~ **based on small carbon-climate feedbacks in**  
510 **idealized scenarios without variability in the atmospheric CO<sub>2</sub> growth (Arora et al., 2020).**



515 **Figure 7: Timeseries and decadal trends of the ocean carbon sink and climate modes in earth system models compared to observations.** (a) The globally integrated annual air-sea CO<sub>2</sub> flux in the pre-industrial control simulations for 12 ESMs (thin blue lines) and the multi-model average (thick blue line). The same is shown for (b) the Niño 3.4 index, (c) the Atlantic Multi-decadal Oscillation, and (d) the Marshall Southern Annular Mode. For the three climate modes, observation-based estimates are shown based on HadISST1 for (b, c) and based on Marshall (2003) for (d). The decadal trends of these observation-based estimates (orange numbers) are compared to the decadal trends of ESM estimates (blue numbers indicating average and standard deviation across the ESM ensemble).


520



However, in addition to physical climate-driven variability, which is the dominant driver of variability of the ocean carbon sink (Doney et al., 2009), there is also biology- and biogeochemical-driven climate-related variability in the air-sea CO<sub>2</sub> fluxes (Ostle et al., 2022; Doney et al., 2009; Keller et al., 2012) for example due to changes in net primary production or  
525 remineralization caused by changes in nutrient supply, temperature, and oxygen. Over the North Atlantic, it has been shown that biogeochemical variability is also strongly influenced by climate modes, such as the Atlantic Multi-decadal Oscillation (Ostle et al., 2022). Nevertheless, GOBMs suggest that the influence of physical variability exceeds the influence of biogeochemical variability (Doney et al., 2009; DeVries et al., 2023). Despite different representations of the biogeochemistry and biology across the models from RECCAP2 (DeVries et al., 2023; Rodgers et al., 2023), they all simulate a similar inter-  
530 annual and decadal variability and trends in the ocean carbon sink (DeVries et al., 2023; Terhaar et al., ~~in press~~2024) as they are forced with historical atmospheric reanalyses products and share the same internal climate modes. Although the similarity of all GOBMs when forced with historical reanalysis strongly suggest that the physical impact on decadal variability exceeds the biogeochemical impact, detailed regional analyses of the biogeochemical-driven climate-related variability in the air-sea CO<sub>2</sub> fluxes (Ostle et al., 2022; Keller et al., 2015), which exceed the scope of this manuscript, are necessary. Overall, the  
535 dominance of physical variability over biogeochemical variability and the larger decadal trends of climate modes in ESMs than in the real world suggest that the ESMs do not underestimate the natural variability of the ocean carbon sink [although it always remains possible that the pCO<sub>2</sub> products are capturing real signals that are not yet simulated.](#)

Although the ~~here-used~~-ESMs from CMIP6 [used here](#) simulate even larger decadal trends of important climate modes, they  
540 might still underestimate decadal trends of the ocean carbon sink driven by climate variability because of their resolution that has increased over the past decades but is still too coarse to explicitly resolve mesoscale ocean eddies. Higher resolved ESMs are still computationally too expensive to be run within the CMIP framework with sufficiently long spin-ups that are necessary for these models to be in equilibrium (Séférian et al., 2016; Gupta et al., 2013). In a few studies with less simulations than required for CMIP, higher-resolved ocean models have been shown to affect the ocean carbon sink and physics and their  
545 variability (Lachkar et al., 2007, 2009; Dufour et al., 2015; Griffies et al., 2015). While it remains impossible to evaluate the effect of higher resolution over a large ensemble of ESMs, as such an ensemble does not exist yet, I tested the here-identified relationship with the highest-resolved earth system model within CMIP6, GFDL-CM4 (Held et al., 2019), which has a horizontal resolution of 0.25° that allows to resolve eddies in tropical and subtropical oceans but still has to parametrize some eddy activity in subpolar and polar oceans. GFDL-CM4 had not been included in the overall analyses as it did not provide  
550 simulations under SSP1-2.6 and SSP3-7.0, presumably because of its large computational costs. The trends in the ocean carbon sink in GFDL-CM4 lie mostly within  $\pm 1\sigma$  of the relationship between changes in the atmospheric CO<sub>2</sub> growth rate and trends in the ocean carbon sink, with only a few decades being in the  $\pm 2\sigma$  range (Figure 8). As for the other ESMs, the relationship in GFDL-CM4 only holds if the ocean carbon sink remains below 4.5 Pg C yr<sup>-1</sup>. Although a potential change in this relationship at an even higher resolution cannot be excluded with certainty until simulations with higher resolution are performed, the

555 robustness of the relationship even for higher-resolved ESMs such as GFDL-CM4 gives no indication that the relationship will  
not hold at even higher resolution.



560 **Figure 8: The relationship between changes in the atmospheric CO<sub>2</sub> growth rate and decadal trends of the global ocean carbon**  
**in a high-resolution model. (a)** Decadal trends of the multi-model mean ocean carbon sink compared to changes in decadal trends in  
atmospheric CO<sub>2</sub>, which represent the decadal averaged growth rate of atmospheric CO<sub>2</sub>. The dark blue to yellow circles without a  
surrounding black line show multi-model dec averages for all years of the historical period from 1850 to 2014 and for all years from 2015 to  
2100 for all four SSPs. The brown line shows a linear fit for all years when the global ocean carbon sink is smaller than 4.5 Pg C yr<sup>-1</sup>  
565 and the brown shading is the 1-σ projection uncertainty. The red dots with black lines around them show values for the high-resolution ESM  
GFDL-CM4 (Held et al., 2019) under the historical simulation and SSP2-4.5 and SSP5-8.5.


## 7 Discussion and Conclusion

570 The analysis with ESM suggests that changes in the atmospheric growth rate of CO<sub>2</sub> can indeed explain most of the decadal trends of the ocean carbon sink, as previously proposed by McKinley et al. (2020). ESMs support the hypothesis by McKinley et al. (2020) that the weak decadal trend in the 1990s and the stronger trend in the 2000s is mainly driven by changes in the atmospheric CO<sub>2</sub> growth rate (Figures 3 and 5). The importance of atmospheric CO<sub>2</sub> extends over all ocean basins, as also previously shown by McKinley et al. (2020). While McKinley et al. (2020) have focused on the last decades and suggested  
575 that the trends in the ocean carbon sink depends on differences on the atmospheric growth rate of CO<sub>2</sub> compared to the long-term trend of the growth rate, I could [generalize this idea here and](#) show that it is the change in the growth rate compared to the previous decade that drives the trends of the ocean carbon sink [over a wide range of timescales and SSPs](#). Moreover, this analysis here extends the timeline of previous analysis and shows how atmospheric CO<sub>2</sub> drives trends in the ocean carbon sink on a range of different future scenarios, from high-mitigation low emission scenarios to high-emission scenarios. In addition,  
580 the use of ESMs allowed to quantify the link between changes in the atmospheric CO<sub>2</sub> growth rate and decadal trends of the ocean carbon, which allows to better separate the effect of internal and external forcing of past decadal trends of the ocean carbon sink. However, if atmospheric CO<sub>2</sub> rises too high and the impact of climate change on the ocean carbon sink increases, atmospheric CO<sub>2</sub> is not anymore the dominant driver of trends of the ocean carbon sink due to changes in the buffer factor and ocean ventilation (Revelle and Suess, 1957; Heinze et al., 2015; Joos et al., 1999; McNeil and Matear, 2013; Frölicher et al.,  
585 2015).

Although atmospheric CO<sub>2</sub> is here shown to be the main driver of the decadal trends in the ocean carbon sink, climate variability also plays an important role for the decadal trends of the ocean carbon sink. With a standard-deviation of  $\pm 0.16$  Pg C yr<sup>-1</sup> across all ESMs, climate variability drives 17% of the trend in the ocean carbon sink when the growth rate of atmospheric  
590 CO<sub>2</sub> is largest and drives all changes in carbon trends when the change in the growth rate of atmospheric CO<sub>2</sub> is zero. Known drivers of this internal climate variability are for example El Niño (Feely et al., 1999; Ishii et al., 2014; McKinley et al., 2004), the Atlantic Multi-decadal Oscillation (Breedon and McKinley, 2016; Keppler et al., 2023), changes in the overturning circulation (DeVries et al., 2017), and changes in the Southern Annual Mode linked to changes in Southern Ocean winds (Le Quéré et al., 2007; Keppler and Landschützer, 2019; Landschützer et al., 2015) and stronger consequent upwelling of older  
595 waters (Lovenduski et al., 2008, 2007), as well as changes in the Southern Ocean stratification (Gruber et al., 2019b). Across the here-used ESMs, the variability of decadal trends is highest in the Southern Ocean, followed by the tropical regions, and again followed by the Northern subpolar gyres (Figure 9); confirming that the decadal trends of the ocean carbon sink are indeed most variable due to internal climate variability in the regions where they are expected based on the previous studies mentioned above. ESMs simulate decadal trends of important climate modes that are even larger than their observation-based  
600 counterparts, which suggests that ESMs also capture the climate-driven decadal trends of the ocean carbon sink. As ESMs slightly overestimate decadal trends of important climate modes and still suggest that changes in the atmospheric growth rate

are the dominant drivers of decadal trends of the ocean carbon sink, climate variability and associated changes in ocean circulation appear to not be the first order driver of decadal trends of the ocean carbon sink over the last decades as previously suggested (Landschützer et al., 2015; DeVries et al., 2017, 2019).

605



610 **Figure 9: Variability of the decadal trends of the zonally integrated ocean carbon sink in earth system models.** The multi-model mean (thick blue line) and the 1- $\sigma$  standard deviation of the variability of the zonally integrated ocean carbon sink across the 251 years of the pre-industrial control simulation in each of the across all 12 ESMs (Table 1) (thin blue lines). In addition, the multi-model mean (thick blue line) maximum and minimum variability in the 1- $\sigma$  standard deviation across all 12 ESMs (blue shading) is shown at each latitude (thin blue lines).

615 The here presented results have implications for previous estimates of the ocean carbon sink, especially those from  $p\text{CO}_2$  products that suggested very strong decadal trends of the ocean carbon sink (Landschützer et al., 2015; Gruber et al., 2019b,

2023). The trend estimates of the ocean carbon sink by  $p\text{CO}_2$  products is larger than the likely trends based on the here-identified relationship between changes in the atmospheric  $\text{CO}_2$  growth rate and the decadal trends of the ocean carbon sink. Thus,  $p\text{CO}_2$  products either overestimate decadal trends of the ocean carbon sink or ESMs underestimate these trends. In each decade from 1990 to 2019, there are five out of eight  $p\text{CO}_2$  products from the Global Carbon Budget (Friedlingstein et al., 2023) that estimate decadal trends that are outside of the  $2\text{-}\sigma$  range that is estimated based on ESMs here (Figure 5), giving these results a likelihood of less than 5% to occur if the ESM results are indeed robust. Some  $p\text{CO}_2$  products estimate trends that are within the  $5\text{-}\sigma$ ,  $6\text{-}\sigma$ , and  $7\text{-}\sigma$  ranges that corresponds to events that occur once every 4776 years ( $5\text{-}\sigma$ ), once every 1.38 million years ( $6\text{-}\sigma$ ), and once every 1.07 billion years ( $7\text{-}\sigma$ ). While it is already extremely unlikely that decadal trends in all three decades from 1990 to 2020 lie outside the  $2\text{-}\sigma$  range, the estimates within the  $5\text{-}\sigma$  to  $7\text{-}\sigma$  range are virtually impossible based on the ESM-derived range. Only two  $p\text{CO}_2$  products (NIES-ML3 from Zeng et al. (2022) and OS-ETHZ-Gracer from Gregor and Gruber (2021)) lie within the  $1\text{-}\sigma$  and  $2\text{-}\sigma$  ranges in the 1990s and 2000s, and only very slightly above the  $2\text{-}\sigma$  range in the 2010s. The slightly higher trend in the 2010s in these products may very well be a consequence of the uneven sampling in space and time (Hauck et al., 2023). While the trends in these two  $p\text{CO}_2$  products are closer to what is expected based on ESMs, only an in-depth analysis will eventually allow with to judge the performance of each  $p\text{CO}_2$  product with certainty.

Here, I have demonstrated that ESMs are capable of simulating the size of decadal trends of important climate modes that have strong impact on the variability of the ocean carbon sink (Figure 7) and that higher resolution does not alter the identified relationship (Figure 8). While this analysis does not guarantee that ESMs do not underestimate the decadal trends in the ocean carbon sink, it suggests that ESMs can simulate the size of the variability of the ocean carbon sink. This conclusion also challenges earlier findings that GOBMs might underestimate decadal trends of the ocean carbon sink (DeVries et al., 2019). Other studies (Gloege et al., 2021; Hauck et al., 2023) support the hypothesis of an overestimation of decadal trends in the ocean carbon sink by  $p\text{CO}_2$  products. Hauck et al. (2023) have recently demonstrated with one GOBM and two  $p\text{CO}_2$  products that sampling biases of  $p\text{CO}_2$  have caused trends in the ocean carbon sink to be overestimated. A similar finding has been made previously, when data from one GOBM, which was sampled in the same way as the real-world ocean was sampled, was extrapolated by one  $p\text{CO}_2$  products to reconstruct the ocean carbon sink. This so-reconstructed ocean carbon sink by the  $p\text{CO}_2$  products also had a larger variability than the directly simulated ocean carbon sink by the GOBM (Gloege et al., 2021). Thus, it appears that most of the  $p\text{CO}_2$  products have and still overestimate decadal trends of the global ocean carbon sink. Therefore, estimates of the variability and size of the flux of natural carbon based on the difference of the total air-sea  $\text{CO}_2$  flux from  $p\text{CO}_2$  products and the change of interior ocean anthropogenic carbon (Müller et al., 2023; Gruber et al., 2019a), defined in this special case only as the additional carbon from increasing atmospheric  $\text{CO}_2$  and not from climate change, are also likely too large.

650 The here found dependence of the ocean carbon sink on atmospheric CO<sub>2</sub> also has implications for studies that extrapolate present-day observation-based estimates of the ocean carbon sink back in time over the entire historical period to estimate a cumulative ocean carbon sink since the beginning of the pre-industrial revolution using the difference of atmospheric CO<sub>2</sub> since pre-industrial times as a scaling factor (Gruber et al., 2009; Mikaloff Fletcher et al., 2006). While this scaling works approximately for most of the historical period, it breaks down from 1920 to 1960 and in the 1990s (Figure 1). In addition, 655 such estimates might be highly sensitive to the year for which the ocean carbon sink was estimated based on observations. If that year falls in one of these anomalous periods, as the year 1995 in Mikaloff Fletcher et al. (2006), the scaling might be biased low or high. Therefore, these extrapolations of present-day fluxes over the historical period should be used with caution or with a slightly more complex extrapolation method that takes the change in the atmospheric CO<sub>2</sub> growth rate into account.

660 The importance for changes of the atmospheric CO<sub>2</sub> growth rate for the trends of the global ocean carbon sink also affects our understanding of the uncertainty of the ocean carbon sink and the role of internal variability in the future. Previous studies have used CMIP simulations with prescribed atmospheric CO<sub>2</sub> to quantify the importance of internal variability for the uncertainty of the projections of the ocean carbon sink in comparison to the importance of model and scenario uncertainty (Gooya et al., 2023; Lovenduski et al., 2016; Schlunegger et al., 2020). As these prescribed atmospheric CO<sub>2</sub> timeseries in 665 CMIP simulations are much smoother than observed atmospheric CO<sub>2</sub> timeseries (Fig. 1), changes in the atmospheric CO<sub>2</sub> growth rate are also much smaller. Thus, these concentration driven CMIP SSPs suppress the internal variability of the atmospheric CO<sub>2</sub> growth rate caused by variabilities in atmospheric temperature, precipitation, El Niño, and volcanic eruptions (Keeling et al., 1995; Kuo et al., 1990; Raupach et al., 2008; Zeng et al., 2005; Bacastow, 1976; Yang and Wang, 2000). The suppressed variability of the atmospheric CO<sub>2</sub> growth rate in concentration driven SSPs also suppresses the variability of the 670 ocean carbon sink in the future, leading to an underestimation of the importance of the internal variability in ESMs for the overall uncertainty of ocean carbon sink projections over the 21<sup>st</sup> century (Gooya et al., 2023; Lovenduski et al., 2016; Schlunegger et al., 2020). This underestimation of the variability of the ocean carbon sink due to prescribed atmospheric CO<sub>2</sub> can be avoided if ESMs were run in an emission-driven mode that automatically introduces a strong variability of the of the atmospheric CO<sub>2</sub> growth rate, as in the model intercomparison project using the Adaptive Emission Reduction Approach 675 (Terhaar et al., 2022a; Silvy et al., 2024).

While changes in the atmospheric CO<sub>2</sub> growth rate and ocean heat uptake might allow to estimate changes in the decadal variability of the ocean carbon sink, it remains still unknown how climate variability and individual modes can be used to predict inter-annual variability of the near-term ocean carbon sink (Lovenduski et al., 2019). In addition, other external 680 forcings, such as volcanic eruptions, are an important factor to the inter-annual variability of the ocean carbon sink but also contribute to decadal trends (McKinley et al., 2017; Fay et al., 2023; Frölicher et al., 2011, 2013).

The influence of changes in the atmospheric CO<sub>2</sub> growth rate on the ocean carbon sink also has profound implications on the near-term future of the ocean carbon sink. With less strongly increasing or even peaking carbon emissions the atmospheric CO<sub>2</sub> growth rate will also peak and potentially decline. The growth rate of atmospheric CO<sub>2</sub> in Mauna Loa has shown a robust negative trend since 2016 and the last time that the growth rate of atmospheric CO<sub>2</sub> has been as small as in 2022 was the year 2008 (<https://gml.noaa.gov/ccgg/trends/gr.html>). If this change from a rise of the atmospheric CO<sub>2</sub> growth rate towards a decline of the atmospheric CO<sub>2</sub> growth rate continues, the forced trend in the ocean carbon sink will also be negative. In addition, GOBMs and *p*CO<sub>2</sub> products suggest that the internal climate variability has led to particularly positive trends of the ocean carbon sink in the 2000s and 2010s (Figs 3 & 5, orange and pink dots lie above the derived relationship). This climate variability will eventually reverse at some point and lead to a larger decline of the ocean carbon sink. Furthermore, ocean heat uptake is projected to increase over the next decade or two independent of the chosen future pathway, also leading to stronger outgassing of carbon from the ocean in the near future and negative trends in the ocean carbon sink (Fig. 6). Although CMIP6 ESMs tend to simulate a smaller ocean heat content over the last two decades (Lyu et al., 2021), this might not be an overestimation by heat uptake of the ESMs but an especially low uptake due to climate variability. Indeed, the recent strong increase in ocean heat content and sea surface temperatures in 2023 (Cheng et al., 2024) might be the beginning of a shift from a period of low heat uptake due to climate variability to a period of high ocean heat uptake. Together, the decreasing atmospheric CO<sub>2</sub> growth rate, the potential change in internal climate variability, and the increased ocean heat uptake will likely cause a substantial negative trend of the ocean carbon sink over the next decade. If, however, emissions and atmospheric CO<sub>2</sub> will rise, the continuous increase in the atmospheric CO<sub>2</sub> growth rate will cause the ocean carbon sink to increase as well.

Overall, this study demonstrate how ESMs can be used to better understand the past and future of the ocean carbon sink and drivers of its variability. They hence provide not only a valuable addition to *p*CO<sub>2</sub> products and GOBMs, but also a unique tool to statistically assess uncertainties and drivers of variability, also potentially in the interior ocean (Müller et al., 2023). The robustness in these results is further corroborated by their capability to simulate the size of decadal trends of important climate modes.

### **Data availability**

710 The Earth system model output used in this study is available via the Earth System Grid Federation (<https://esgf-node.ipsl.upmc.fr/projects/esgf-ipsl/>, last access: 1 June 2022).

### **Competing interests**

The author has declared that he has no competing interests.

### **Acknowledgments**

715 First, I want to thank Thomas L. Frölicher, Fortunat Joos, Roland Séférian, and Jens D. Müller for their helpful comments on the manuscript and during discussions. Furthermore, I acknowledge funding from the Woods Hole Oceanographic Institution Postdoctoral Scholar Program, and the Swiss National Science Foundation under grant # PZ00P2\_209044 (ArcticECO).

720



## References

- 725 Arora, V. K., Katavouta, A., Williams, R. G., Jones, C. D., Brovkin, V., Friedlingstein, P., Schwinger, J., Bopp, L., Boucher, O., Cadule, P., Chamberlain, M. A., Christian, J. R., Delire, C., Fisher, R. A., Hajima, T., Ilyina, T., Joetzjer, E., Kawamiya, M., Koven, C. D., Krasting, J. P., Law, R. M., Lawrence, D. M., Lenton, A., Lindsay, K., Pongratz, J., Raddatz, T., Séférian, R., Tachiiri, K., Tjiputra, J. F., Wiltshire, A., Wu, T., and Ziehn, T.: Carbon–concentration and carbon–climate feedbacks in CMIP6 models and their comparison to CMIP5 models, *Biogeosciences*, 17, 4173–4222, [https://doi.org/10.5194/bg-17-4173-](https://doi.org/10.5194/bg-17-4173-2020)  
730 2020, 2020.
- Bacastow, R. B.: Modulation of atmospheric carbon dioxide by the Southern Oscillation, *Nature*, 261, 116–118, <https://doi.org/10.1038/261116a0>, 1976.
- 735 Bennington, V., Galjanic, T., and McKinley, G. A.: Explicit Physical Knowledge in Machine Learning for Ocean Carbon Flux Reconstruction: The pCO<sub>2</sub>-Residual Method, *J. Adv. Model. Earth Sy.*, 14, 3345, <https://doi.org/10.1029/2021ms002960>, 2022a.
- Bennington, V., Gloege, L., and McKinley, G. A.: Variability in the global ocean carbon sink from 1959 to 2020 by correcting  
740 models with observations, *Geophys. Res. Lett.*, 49, e2022GL098632, <https://doi.org/10.1029/2022GL098632>, 2022b.
- Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Balkanski, Y., Bastrikov, V., Bekki, S., Bonnet, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Caubel, A., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., D’Andrea, F., Davini, P., de Lavergne, C., Denvil, S., Deshayes, J., Devilliers, M., Ducharne, A., Dufresne, J.-L., Dupont, E., Éthé, C., Fairhead, L.,  
745 Falletti, L., Flavoni, S., Foujols, M.-A., Gardoll, S., Gastineau, G., Ghattas, J., Grandpeix, J.-Y., Guenet, B., Guez E., L., Guilyardi, E., Guimberteau, M., Hauglustaine, D., Hourdin, F., Idelkadi, A., Jousaume, S., Kageyama, M., Khodri, M., Krinner, G., Lebas, N., Levavasseur, G., Lévy, C., Li, L., Lott, F., Lurton, T., Luyssaert, S., Madec, G., Madeleine, J.-B., Maignan, F., Marchand, M., Marti, O., Mellul, L., Meurdesoif, Y., Mignot, J., Musat, I., Otlé, C., Peylin, P., Planton, Y., Polcher, J., Rio, C., Rochetin, N., Rousset, C., Sepulchre, P., Sima, A., Swingedouw, D., Thiéblemont, R., Traore, A. K.,  
750 Vancoppenolle, M., Vial, J., Vialard, J., Viovy, N., and Vuichard, N.: Presentation and Evaluation of the IPSL-CM6A-LR Climate Model, *J. Adv. Model. Earth Syst.*, 12, e2019MS002010, <https://doi.org/https://doi.org/10.1029/2019MS002010>, 2020.
- Breeden, M. L. and McKinley, G. A.: Climate impacts on multidecadal pCO<sub>2</sub> variability in the North Atlantic: 1948-2009,  
755 *Biogeosciences*, 13, 3387–3396, <https://doi.org/10.5194/bg-13-3387-2016>, 2016.

Broecker, W. S., Takahashi, T., Simpson, H. J., and Peng T.-H.: Fate of Fossil Fuel Carbon Dioxide and the Global Carbon Budget, *Science*, 206, 409–418, <https://doi.org/10.1126/science.206.4417.409>, 1979.

760 Bronselaer, B., Winton, M., Russell, J., Sabine, C. L., and Khatiwala, S.: Agreement of CMIP5 Simulated and Observed Ocean Anthropogenic CO<sub>2</sub> Uptake, *Geophys. Res. Lett.*, 44, 12, 212–298, 305, <https://doi.org/https://doi.org/10.1002/2017GL074435>, 2017.

Caldeira, K. and Duffy, P. B.: The Role of the Southern Ocean in Uptake and Storage of Anthropogenic Carbon Dioxide, *Science.*, 287, 620–622, <https://doi.org/10.1126/science.287.5453.620>, 2000.

Chau, T. T. T., Gehlen, M., and Chevallier, F.: A seamless ensemble-based reconstruction of surface ocean *p*CO<sub>2</sub> and air–sea CO<sub>2</sub> fluxes over the global coastal and open oceans, *Biogeosciences*, 19, 1087–1109, <https://doi.org/10.5194/bg-19-1087-2022>, 2022.

770

Cheng, L., Abraham, J., Trenberth, K. E., Boyer, T., Mann, M. E., Zhu, J., Wang, F., Yu, F., Locarnini, R., Fasullo, J., Zheng, F., Li, Y., Zhang, B., Wan, L., Chen, X., Wang, D., Feng, L., Song, X., Liu, Y., Reseghetti, F., Simoncelli, S., Gouretski, V., Chen, G., Mishonov, A., Reagan, J., Von Schuckmann, K., Pan, Y., Tan, Z., Zhu, Y., Wei, W., Li, G., Ren, Q., Cao, L., and Lu, Y.: New Record Ocean Temperatures and Related Climate Indicators in 2023, *Adv. Atmos. Sci.*, <https://doi.org/10.1007/s00376-024-3378-5>, 2024.

775

Christian, J. R., Denman, K. L., Hayashida, H., Holdsworth, A. M., Lee, W. G., Riche, O. G. J., Shao, A. E., Steiner, N., and Swart, N. C.: Ocean biogeochemistry in the Canadian Earth System Model version 5.0.3: CanESM5 and CanESM5-CanOE, *Geosci. Model Dev.*, 15, 4393–4424, <https://doi.org/10.5194/gmd-15-4393-2022>, 2022.

780

Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Vertenstein, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2 (CESM2), *J. Adv. Model. Earth Syst.*, 12, <https://doi.org/https://doi.org/10.1029/2019MS001916>, 2020.

785

DeVries, T., Holzer, M., and Primeau, F.: Recent increase in oceanic carbon uptake driven by weaker upper-ocean overturning, *Nature*, 542, 215–218, <https://doi.org/10.1038/nature21068>, 2017.

790

DeVries, T., Le Quéré, C., Andrews, O., Berthet, S., Hauck, J., Ilyina, T., Landschützer, P., Lenton, A., Lima, I. D., Nowicki, M., Schwinger, J., and Séférian, R.: Decadal trends in the ocean carbon sink, *Proc. Natl. Acad. Sci.*, 116, 11646–11651, <https://doi.org/10.1073/pnas.1900371116>, 2019.

795

DeVries, T., Yamamoto, K., Wanninkhof, R., Gruber, N., Hauck, J., Müller, J. D., Bopp, L., Carroll, D., Carter, B. R., Chau, T. T. T., Doney, S. C., Gehlen, M., Gloege, L., Gregor, L., Henson, S., Kim, J. H., Iida, Y., Ilyina, T., Landschützer, P., Le Quéré, C., Munro, D. R., Nissen, C., Patara, L., Perez, F. F., Resplandy, L., Rodgers, K. B., Schwinger, J., Séférian, R., Sicardi, V., Terhaar, J., Triñanes, J., Tsujino, H., Watson, A. J., Yasunaka, S., and Zeng, J.: Magnitude, trends, and variability of the global ocean carbon sink from 1985-2018, *Global Biogeochem. Cycles*, 2023.

800

Doney, S. C., Lima, I., Feely, R. A., Glover, D. M., Lindsay, K., Mahowald, N., Moore, J. K., and Wanninkhof, R.: Mechanisms governing interannual variability in upper-ocean inorganic carbon system and air–sea CO<sub>2</sub> fluxes: Physical climate and atmospheric dust, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 56, 640–655, <https://doi.org/https://doi.org/10.1016/j.dsr2.2008.12.006>, 2009.

805

Dufour, C. O., Griffies, S. M., de Souza, G. F., Frenger, I., Morrison, A. K., Palter, J. B., Sarmiento, J. L., Galbraith, E. D., Dunne, J. P., Anderson, W. G., and Slater, R. D.: Role of Mesoscale Eddies in Cross-Frontal Transport of Heat and Biogeochemical Tracers in the Southern Ocean, *J. Phys. Oceanogr.*, 45, 3057–3081, <https://doi.org/10.1175/JPO-D-14-0240.1>, 2015.

810

Fay, A. R. and McKinley, G. A.: Global open-ocean biomes: mean and temporal variability, *Earth Syst. Sci. Data*, 6, 273–284, <https://doi.org/10.5194/essd-6-273-2014>, 2014.

815

Fay, A. R., Gregor, L., Landschützer, P., McKinley, G. A., Gruber, N., Gehlen, M., Iida, Y., Laruelle, G. G., Rödenbeck, C., Roobaert, A., and Zeng, J.: SeaFlux: harmonization of air–sea CO<sub>2</sub> fluxes from surface *p*CO<sub>2</sub> data products using a standardized approach, *Earth Syst. Sci. Data*, 13, 4693–4710, <https://doi.org/10.5194/essd-13-4693-2021>, 2021.

820

Fay, A. R., McKinley, G. A., Lovenduski, N. S., Eddebbbar, Y., Levy, M. N., Long, M. C., Olivarez, H. C., and Rustagi, R. R.: Immediate and long-lasting impacts of the Mt. Pinatubo eruption on ocean oxygen and carbon inventories, *Global Biogeochem. Cycles*, n/a, e2022GB007513, <https://doi.org/https://doi.org/10.1029/2022GB007513>, 2023.

Feely, R. A., Wanninkhof, R., Takahashi, T., and Tans, P.: Influence of El Niño on the equatorial Pacific contribution to atmospheric CO<sub>2</sub> accumulation, *Nature*, 398, 597–601, <https://doi.org/10.1038/19273>, 1999.

Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J., Landschützer, P., Le Quéré, C., Lujckx, I. T., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Barbero, L., Bates, N. R., Becker, M., Bellouin, N., Decharme, B., Bopp, L., Brasika, I. B. M., Cadule, P., Chamberlain, M. A., Chandra, N., Chau, T.-T.-T., Chevallier, F., Chini, L. P., Cronin, M., Dou, X., Enyo, K., Evans, W., Falk, S., Feely, R. A., Feng, L., Ford, D. J., Gasser, T., Ghattas, J., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Hefner, M., Heinke, J., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Joos, F., Kato, E., Keeling, R. F., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Lan, X., Lefèvre, N., Li, H., Liu, J., Liu, Z., Ma, L., Marland, G., Mayot, N., McGuire, P. C., McKinley, G. A., Meyer, G., Morgan, E. J., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O'Brien, K. M., Olsen, A., Omar, A. M., Ono, T., Paulsen, M., Pierrot, D., Pockock, K., Poulter, B., Powis, C. M., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Séférian, R., et al.: Global Carbon Budget 2023, *Earth Syst. Sci. Data*, 15, 5301–5369, <https://doi.org/10.5194/essd-15-5301-2023>, 2023.

Frölicher, T. L., Joos, F., and Raible, C. C.: Sensitivity of atmospheric CO<sub>2</sub> and climate to explosive volcanic eruptions, *Biogeosciences*, 8, 2317–2339, <https://doi.org/10.5194/bg-8-2317-2011>, 2011.

Frölicher, T. L., Joos, F., Raible, C. C., and Sarmiento, J. L.: Atmospheric CO<sub>2</sub> response to volcanic eruptions: The role of ENSO, season, and variability, *Global Biogeochem. Cycles*, 27, 239–251, <https://doi.org/10.1002/gbc.20028>, 2013.

Frölicher, T. L., Sarmiento, J. L., Paynter, D. J., Dunne, J. P., Krasting, J. P., and Winton, M.: Dominance of the Southern Ocean in Anthropogenic Carbon and Heat Uptake in CMIP5 Models, *J. Clim.*, 28, 862–886, <https://doi.org/10.1175/JCLI-D-14-00117.1>, 2015.

Gloege, L., McKinley, G. A., Landschützer, P., Fay, A. R., Frölicher, T. L., Fyfe, J. C., Ilyina, T., Jones, S., Lovenduski, N. S., Rodgers, K. B., Schlunegger, S., and Takano, Y.: Quantifying Errors in Observationally Based Estimates of Ocean Carbon Sink Variability, *Global Biogeochem. Cycles*, 35, e2020GB006788, <https://doi.org/10.1029/2020GB006788>, 2021.

Gloege, L., Yan, M., Zheng, T., and McKinley, G. A.: Improved Quantification of Ocean Carbon Uptake by Using Machine Learning to Merge Global Models and pCO<sub>2</sub> Data, *J. Adv. Model. Earth Sy.*, 14, e2021MS002620, <https://doi.org/10.1029/2021MS002620>, 2022.

- 860 Gooya, P., Swart, N. C., and Hamme, R. C.: Time-varying changes and uncertainties in the CMIP6-ocean carbon sink from global to local scale, *Earth Syst. Dyn.*, 14, 383–398, <https://doi.org/10.5194/esd-14-383-2023>, 2023.
- Goris, N., Tjiputra, J. F., Olsen, A., Schwinger, J., Lauvset, S. K., and Jeansson, E.: Constraining Projection-Based Estimates of the Future North Atlantic Carbon Uptake, *J. Clim.*, 31, 3959–3978, <https://doi.org/10.1175/JCLI-D-17-0564.1>, 2018.
- 865 Gregor, L. and Gruber, N.: OceanSODA-ETHZ: a global gridded data set of the surface ocean carbonate system for seasonal to decadal studies of ocean acidification, *Earth Syst. Sci. Data*, 13, 777–808, <https://doi.org/10.5194/essd-13-777-2021>, 2021.
- Griffies, S. M., Winton, M., Anderson, W. G., Benson, R., Delworth, T. L., Dufour, C. O., Dunne, J. P., Goddard, P., Morrison, A. K., Rosati, A., Wittenberg, A. T., Yin, J., and Zhang, R.: Impacts on Ocean Heat from Transient Mesoscale Eddies in a Hierarchy of Climate Models, *J. Clim.*, 28, 952–977, <https://doi.org/10.1175/JCLI-D-14-00353.1>, 2015.
- 870 Gruber, N., Gloor, M., Mikaloff Fletcher, S. E., Doney, S. C., Dutkiewicz, S., Follows, M. J., Gerber, M., Jacobson, A. R., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S. A., Sarmiento, J. L., and Takahashi, T.: Oceanic sources, sinks, and transport of atmospheric CO<sub>2</sub>, *Global Biogeochem. Cycles*, 23, <https://doi.org/https://doi.org/10.1029/2008GB003349>, 2009.
- 875 Gruber, N., Clement, D., Carter, B. R., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Key, R. M., Kozyr, A., Lauvset, S. K., Lo Monaco, C., Mathis, J. T., Murata, A., Olsen, A., Perez, F. F., Sabine, C. L., Tanhua, T., and Rik, W.: The oceanic sink for anthropogenic CO<sub>2</sub> from 1994 to 2007, *Science*, 363, 1193–1199, <https://doi.org/10.1126/science.aau5153>, 2019a.
- 880 Gruber, N., Landschützer, P., and Lovenduski, N. S.: The variable southern ocean carbon sink, <https://doi.org/10.1146/annurev-marine-121916-063407>, 3 January 2019b.
- 885 Gruber, N., Bakker, D. C. E., DeVries, T., Gregor, L., Hauck, J., Landschützer, P., McKinley, G. A., and Müller, J. D.: Trends and variability in the ocean carbon sink, *Nat. Rev. Earth Environ.*, <https://doi.org/10.1038/s43017-022-00381-x>, 2023.
- Gupta, A. Sen, Jourdain, N. C., Brown, J. N., and Monselesan, D.: Climate Drift in the CMIP5 Models, *J. Clim.*, 26, 8597–8615, <https://doi.org/https://doi.org/10.1175/JCLI-D-12-00521.1>, 2013.
- 890 Gutjahr, O., Putrasahan, D., Lohmann, K., Jungclauss, J. H., von Storch, J.-S., Brüggemann, N., Haak, H., and Stössel, A.: Max Planck Institute Earth System Model (MPI-ESM1.2) for the High-Resolution Model Intercomparison Project (HighResMIP), *Geosci. Model Dev.*, 12, 3241–3281, <https://doi.org/10.5194/gmd-12-3241-2019>, 2019.

Halsey, L. G.: The reign of the p-value is over: what alternative analyses could we employ to fill the power vacuum?, *Biol. Lett.*, 15, 20190174, <https://doi.org/10.1098/rsbl.2019.0174>, 2019.

895

Hauck, J., Völker, C., Wang, T., Hoppema, M., Losch, M., and Wolf-Gladrow, D. A.: Seasonally different carbon flux changes in the Southern Ocean in response to the southern annular mode, *Global Biogeochem. Cycles*, 27, 1236–1245, <https://doi.org/https://doi.org/10.1002/2013GB004600>, 2013.

900 Hauck, J., Zeising, M., Le Quéré, C., Gruber, N., Bakker, D. C. E., Bopp, L., Chau, T. T. T., Gürses, Ö., Ilyina, T., Landschützer, P., Lenton, A., Resplandy, L., Rödenbeck, C., Schwinger, J., and Séférian, R.: Consistency and Challenges in the Ocean Carbon Sink Estimate for the Global Carbon Budget, *Front. Mar. Sci.*, 7, <https://doi.org/10.3389/fmars.2020.571720>, 2020.

905 Hauck, J., Nissen, C., Landschützer, P., Rödenbeck, C., Bushinsky, S., and Olsen, A.: Sparse observations induce large biases in estimates of the global ocean CO<sub>2</sub> sink: an ocean model subsampling experiment, *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.*, 381, 20220063, <https://doi.org/10.1098/rsta.2022.0063>, 2023.

Heinze, C., Meyer, S., Goris, N., Anderson, L., Steinfeldt, R., Chang, N., Le Quéré, C., and Bakker, D. C. E.: The ocean carbon  
910 sink – impacts, vulnerabilities and challenges, *Earth Syst. Dyn.*, 6, 327–358, <https://doi.org/10.5194/esd-6-327-2015>, 2015.

Held, I. M., Guo, H., Adcroft, A., Dunne, J. P., Horowitz, L. W., Krasting, J., Shevliakova, E., Winton, M., Zhao, M., Bushuk, M., Wittenberg, A. T., Wyman, B., Xiang, B., Zhang, R., Anderson, W., Balaji, V., Donner, L., Dunne, K., Durachta, J., Gauthier, P. P. G., Ginoux, P., Golaz, J.-C., Griffies, S. M., Hallberg, R., Harris, L., Harrison, M., Hurlin, W., John, J., Lin, P., Lin, S.-J., Malyshev, S., Menzel, R., Milly, P. C. D., Ming, Y., Naik, V., Paynter, D., Paulot, F., Ramaswamy, V., Reichl, B., Robinson, T., Rosati, A., Seman, C., Silvers, L. G., Underwood, S., and Zadeh, N.: Structure and Performance of GFDL's  
915 CM4.0 Climate Model, *J. Adv. Model. Earth Syst.*, 11, 3691–3727, <https://doi.org/https://doi.org/10.1029/2019MS001829>, 2019.

920 Held, L. and Ott, M.: On p-values and Bayes factors, *Annu. Rev. Stat. Appl.*, 5, 393–419, <https://doi.org/10.1146/annurev-statistics-031017-100307>, 2018.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L.,  
925 Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P.,

Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999–2049, <https://doi.org/https://doi.org/10.1002/qj.3803>, 2020.

930 IPCC: Summary for Policymakers, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press, 2021.

935

Ishii, M., Feely, R. A., Rodgers, K. B., Park, G.-H., Wanninkhof, R., Sasano, D., Sugimoto, H., Cosca, C. E., Nakaoka, S., Telszewski, M., Nojiri, Y., Mikaloff Fletcher, S. E., Niwa, Y., Patra, P. K., Valsala, V., Nakano, H., Lima, I., Doney, S. C., Buitenhuis, E. T., Aumont, O., Dunne, J. P., Lenton, A., and Takahashi, T.: Air–sea CO<sub>2</sub> flux in the Pacific Ocean for the period 1990–2009, *Biogeosciences*, 11, 709–734, <https://doi.org/10.5194/bg-11-709-2014>, 2014.

940

Joos, F., Plattner, G.-K., Stocker, T. F., Marchal, O., and Schmittner, A.: Global Warming and Marine Carbon Cycle Feedbacks on Future Atmospheric CO<sub>2</sub>, *Science*, 284, 464–467, <https://doi.org/10.1126/science.284.5413.464>, 1999.

Joos, F., Roth, R., Fuglestedt, J. S., Peters, G. P., Enting, I. G., von Bloh, W., Brovkin, V., Burke, E. J., Eby, M., Edwards, 945 N. R., Friedrich, T., Frölicher, T. L., Halloran, P. R., Holden, P. B., Jones, C., Kleinen, T., Mackenzie, F. T., Matsumoto, K., Meinshausen, M., Plattner, G.-K., Reisinger, A., Segschneider, J., Shaffer, G., Steinacher, M., Strassmann, K., Tanaka, K., Timmermann, A., and Weaver, A. J.: Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis, *Atmos. Chem. Phys.*, 13, 2793–2825, <https://doi.org/10.5194/acp-13-2793-2013>, 2013.

950

Keeling, C. D., Whorf, T. P., Wahlen, M., and van der Plichtt, J.: Interannual extremes in the rate of rise of atmospheric carbon dioxide since 1980, *Nature*, 375, 666–670, <https://doi.org/10.1038/375666a0>, 1995.

Keller, K. M., Joos, F., Raible, C. C., Cocco, V., Frölicher, T. L., Dunne, J. P., Gehlen, M., Bopp, L., Orr, J. C., Tjiputra, J., 955 Heinze, C., Segschneider, J., Roy, T., and Metzl, N.: Variability of the ocean carbon cycle in response to the North Atlantic Oscillation, *Tellus B Chem. Phys. Meteorol.*, 64, 18738, <https://doi.org/10.3402/tellusb.v64i0.18738>, 2012.

Keller, K. M., Joos, F., Lehner, F., and Raible, C. C.: Detecting changes in marine responses to ENSO from 850 to 2100 C.E.: Insights from the ocean carbon cycle, *Geophys. Res. Lett.*, 42, 518–525, 960 <https://doi.org/https://doi.org/10.1002/2014GL062398>, 2015.

Keppler, L. and Landschützer, P.: Regional Wind Variability Modulates the Southern Ocean Carbon Sink, *Sci. Rep.*, 9, 7384, <https://doi.org/10.1038/s41598-019-43826-y>, 2019.

965 Keppler, L., Landschützer, P., Lauvset, S. K., and Gruber, N.: Recent Trends and Variability in the Oceanic Storage of Dissolved Inorganic Carbon, *Global Biogeochem. Cycles*, 37, e2022GB007677, <https://doi.org/https://doi.org/10.1029/2022GB007677>, 2023.

Kuo, C., Lindberg, C., and Thomson, D. J.: Coherence established between atmospheric carbon dioxide and global  
970 temperature, *Nature*, 343, 709–714, <https://doi.org/10.1038/343709a0>, 1990.

Lachkar, Z., Orr, J. C., Dutay, J.-C., and Delecluse, P.: Effects of mesoscale eddies on global ocean distributions of CFC-11, CO<sub>2</sub>, and  $\Delta^{14}\text{C}$ , *Ocean Sci.*, 3, 461–482, <https://doi.org/10.5194/os-3-461-2007>, 2007.

975 Lachkar, Z., Orr, J. C., and Dutay, J.-C.: Seasonal and mesoscale variability of oceanic transport of anthropogenic CO<sub>2</sub>, 6, 2509–2523, <https://doi.org/10.5194/bg-6-2509-2009>, 2009.

Lacroix, F., Ilyina, T., and Hartmann, J.: Oceanic CO<sub>2</sub> outgassing and biological production hotspots induced by pre-industrial river loads of nutrients and carbon in a global modeling approach, *Biogeosciences*, 17, 55–88, <https://doi.org/10.5194/bg-17-55-2020>, 2020.  
980

Landschützer, P., Gruber, N., Haumann, F. A., Rödenbeck, C., Bakker, D. C. E., van Heuven, S., Hoppema, M., Metzl, N., Sweeney, C., Takahashi, T., Tilbrook, B., and Wanninkhof, R.: The reinvigoration of the Southern Ocean carbon sink, *Science*, 349, 1221–1224, <https://doi.org/10.1126/science.aab2620>, 2015.  
985

Landschützer, P., Gruber, N., and Bakker, D. C. E.: Decadal variations and trends of the global ocean carbon sink, *Global Biogeochem. Cycles*, 30, 1396–1417, <https://doi.org/https://doi.org/10.1002/2015GB005359>, 2016.

Lenton, A. and Matear, R. J.: Role of the Southern Annular Mode (SAM) in Southern Ocean CO<sub>2</sub> uptake, *Global Biogeochem. Cycles*, 21, <https://doi.org/https://doi.org/10.1029/2006GB002714>, 2007.  
990

Li, H., and Ilyina, T.: Current and future decadal trends in the oceanic carbon uptake are dominated by internal variability. *Geophys. Res. Lett.*, 45, 916–925, <https://doi.org/10.1002/2017GL075370>, 2018.



- 995 Lovato, T., Peano, D., Butenschön, M., Materia, S., Iovino, D., Scoccimarro, E., Fogli, P. G., Cherchi, A., Bellucci, A., Gualdi, S., Masina, S., and Navarra, A.: CMIP6 Simulations With the CMCC Earth System Model (CMCC-ESM2), *J. Adv. Model. Earth Syst.*, 14, e2021MS002814, <https://doi.org/https://doi.org/10.1029/2021MS002814>, 2022.
- Lovenduski, N. S., Gruber, N., Doney, S. C., and Lima, I. D.: Enhanced CO<sub>2</sub> outgassing in the Southern Ocean from a positive  
1000 phase of the Southern Annular Mode, *Global Biogeochem. Cycles*, 21, <https://doi.org/https://doi.org/10.1029/2006GB002900>, 2007.
- Lovenduski, N. S., Gruber, N., and Doney, S. C.: Toward a mechanistic understanding of the decadal trends in the Southern Ocean carbon sink, *Global Biogeochem. Cycles*, 22, <https://doi.org/https://doi.org/10.1029/2007GB003139>, 2008.  
1005
- Lovenduski, N. S., McKinley, G. A., Fay, A. R., Lindsay, K., and Long, M. C.: Partitioning uncertainty in ocean carbon uptake projections: Internal variability, emission scenario, and model structure, *Global Biogeochem. Cycles*, 30, 1276–1287, <https://doi.org/https://doi.org/10.1002/2016GB005426>, 2016.
- 1010 Lovenduski, N. S., Yeager, S. G., Lindsay, K., and Long, M. C.: Predicting near-term variability in ocean carbon uptake, *Earth Syst. Dyn.*, 10, 45–57, <https://doi.org/10.5194/esd-10-45-2019>, 2019.
- Lyu, K., Zhang, X., and Church, J. A.: Projected ocean warming constrained by the ocean observational record, *Nat. Clim. Chang.*, <https://doi.org/10.1038/s41558-021-01151-1>, 2021.  
1015
- McCarthy, G. D., Brown, P. J., Flagg, C. N., Goni, G., Houpert, L., Hughes, C. W., Hummels, R., Inall, M., Jochumsen, K., Larsen, K. M. H., Lherminier, P., Meinen, C. S., Moat, B. I., Rayner, D., Rhein, M., Roessler, A., Schmid, C., and Smeed, D. A.: Sustainable Observations of the AMOC: Methodology and Technology, *Rev. Geophys.*, 58, e2019RG000654, <https://doi.org/10.1029/2019RG000654>, 2020.  
1020
- MacDougall, A. H., Frölicher, T. L., Jones, C. D., Rogelj, J., Matthews, H. D., Zickfeld, K., Arora, V. K., Barrett, N. J., Brovkin, V., Burger, F. A., Eby, M., Eliseev, A. V, Hajima, T., Holden, P. B., Jeltsch-Thömmes, A., Koven, C., Mengis, N., Menviel, L., Michou, M., Mokhov, I. I., Oka, A., Schwinger, J., Séférian, R., Shaffer, G., Sokolov, A., Tachiiri, K., Tjiputra, J., Wiltshire, A., and Ziehn, T.: Is there warming in the pipeline? A multi-model analysis of the Zero Emissions Commitment  
1025 from CO<sub>2</sub>, 17, 2987–3016, <https://doi.org/10.5194/bg-17-2987-2020>, 2020.
- Marshall, G. J.: Trends in the Southern Annular Mode from Observations and Reanalyses, *J. Clim.*, 16, 4134–4143, [https://doi.org/https://doi.org/10.1175/1520-0442\(2003\)016<4134:TITSAM>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0442(2003)016<4134:TITSAM>2.0.CO;2), 2003.

1030 Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C., Hohenegger, C., Ilyina, T., Jahns, T., Jimenéz-de-la-Cuesta, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W. A., Nabel, J. E. M. S., Nam, C. C. W., Notz, D., Nyawira, S.-S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J.,  
1035 Popp, M., Raddatz, T. J., Rast, S., Redler, R., Reick, C. H., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von Storch, J.-S., Tian, F., Voigt, A., Vrese, P., Wieners, K.-H., Wilkenskjeld, S., Winkler, A., and Roeckner, E.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and Its Response to Increasing CO<sub>2</sub>, *J. Adv. Model. Earth Syst.*, 11, 998–1038, <https://doi.org/https://doi.org/10.1029/2018MS001400>, 2019.

1040

McKinley, G. A., Follows, M. J., and Marshall, J.: Mechanisms of air-sea CO<sub>2</sub> flux variability in the equatorial Pacific and the North Atlantic, *Global Biogeochem. Cycles*, 18, <https://doi.org/https://doi.org/10.1029/2003GB002179>, 2004.

McKinley, G. A., Pilcher, D. J., Fay, A. R., Lindsay, K., Long, M. C., and Lovenduski, N. S.: Timescales for detection of  
1045 trends in the ocean carbon sink, *Nature*, 530, 469–472, <https://doi.org/10.1038/nature16958>, 2016.

McKinley, G. A., Fay, A. R., Lovenduski, N. S., and Pilcher, D. J.: Natural Variability and Anthropogenic Trends in the Ocean Carbon Sink, *Ann. Rev. Mar. Sci.*, 9, 125–150, <https://doi.org/10.1146/annurev-marine-010816-060529>, 2017.

1050 McKinley, G. A., Fay, A. R., Eddebbar, Y. A., Gloege, L., and Lovenduski, N. S.: External Forcing Explains Recent Decadal Variability of the Ocean Carbon Sink, *AGU Adv.*, 1, e2019AV000149, <https://doi.org/https://doi.org/10.1029/2019AV000149>, 2020.

1055 [McKinley, G. A., Bennington, V., Meinshausen, M., & Nicholls, Z.: Modern air-sea flux distributions reduce uncertainty in the future ocean carbon sink, \*Environmental Research Letters\*, 18\(4\), 044011, <https://doi.org/https://doi.org/10.1088/1748-9326/acc195>, 2023.](https://doi.org/https://doi.org/10.1088/1748-9326/acc195)

McNeil, B. I. and Matear, R. J.: The non-steady state oceanic CO<sub>2</sub> signal: its importance, magnitude and a novel way to detect it, 10, 2219–2228, <https://doi.org/10.5194/bg-10-2219-2013>, 2013.

1060

Meinshausen, M., Vogel, E., Nauels, A., Lorbacher, K., Meinshausen, N., Etheridge, D. M., Fraser, P. J., Montzka, S. A., Rayner, P. J., Trudinger, C. M., Krummel, P. B., Beyerle, U., Canadell, J. G., Daniel, J. S., Enting, I. G., Law, R. M., Lunder,

- 1065 C. R., O'Doherty, S., Prinn, R. G., Reimann, S., Rubino, M., Velders, G. J. M., Vollmer, M. K., Wang, R. H. J., and Weiss, R.: Historical greenhouse gas concentrations for climate modelling (CMIP6), *Geosci. Model Dev.*, 10, 2057–2116, <https://doi.org/10.5194/gmd-10-2057-2017>, 2017.
- 1070 Meinshausen, M., Nicholls, Z. R. J., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., Beyerle, U., Gessner, C., Nauels, A., Bauer, N., Canadell, J. G., Daniel, J. S., John, A., Krummel, P. B., Luderer, G., Meinshausen, N., Montzka, S. A., Rayner, P. J., Reimann, S., Smith, S. J., van den Berg, M., Velders, G. J. M., Vollmer, M. K., and Wang, R. H. J.: The shared socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to 2500, *Geosci. Model Dev.*, 13, 3571–3605, <https://doi.org/10.5194/gmd-13-3571-2020>, 2020.
- 1075 Mikaloff Fletcher, S. E., Gruber, N., Jacobson, A. R., Doney, S. C., Dutkiewicz, S., Gerber, M., Follows, M., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S. A., and Sarmiento, J. L.: Inverse estimates of anthropogenic CO<sub>2</sub> uptake, transport, and storage by the ocean, *Global Biogeochem. Cycles*, 20, <https://doi.org/https://doi.org/10.1029/2005GB002530>, 2006.
- 1080 Müller, J. D., Gruber, N., Carter, B., Feely, R., Ishii, M., Lange, N., Lauvset, S. K., Murata, A., Olsen, A., Pérez, F. F., Sabine, C., Tanhua, T., Wanninkhof, R., and Zhu, D.: Decadal Trends in the Oceanic Storage of Anthropogenic Carbon From 1994 to 2014, *AGU Adv.*, 4, e2023AV000875, <https://doi.org/https://doi.org/10.1029/2023AV000875>, 2023.
- 1085 Orr, J. C., Maier-Reimer, E., Mikolajewicz, U., Monfray, P., Sarmiento, J. L., Toggweiler, J. R., Taylor, N. K., Palmer, J., Gruber, N., Sabine, C. L., Le Quéré, C., Key, R. M., and Boutin, J.: Estimates of anthropogenic carbon uptake from four three-dimensional global ocean models, *Global Biogeochem. Cycles*, 15, 43–60, <https://doi.org/https://doi.org/10.1029/2000GB001273>, 2001.
- Orr, J. C., Kwiatkowski, L., and Pörtner, H.-O.: Arctic Ocean annual high in *p*CO<sub>2</sub> could shift from winter to summer, *Nature*, 610, 94–100, <https://doi.org/10.1038/s41586-022-05205-y>, 2022.
- 1090 Ostle, C., Landschützer, P., Edwards, M., Johnson, M., Schmidtke, S., Schuster, U., Watson, A. J., and Robinson, C.: Multidecadal changes in biology influence the variability of the North Atlantic carbon sink, *Environ. Res. Lett.*, 17, 114056, <https://doi.org/10.1088/1748-9326/ac9ecf>, 2022.
- 1095 Le Quéré, C., Rödenbeck, C., Buitenhuis, E. T., Conway, T. J., Langenfelds, R., Gomez, A., Labuschagne, C., Ramonet, M., Nakazawa, T., Metz, N., Gillett, N., and Heimann, M.: Saturation of the Southern Ocean CO<sub>2</sub> Sink Due to Recent Climate Change, *Science*, 316, 1735–1738, <https://doi.org/10.1126/science.1136188>, 2007.

- Raupach, M. R.: The exponential eigenmodes of the carbon-climate system, and their implications for ratios of responses to forcings, *Earth Syst. Dyn.*, 4, 31–49, <https://doi.org/10.5194/esd-4-31-2013>, 2013.
- 1100 Raupach, M. R., Canadell, J. G., and Le Quéré, C.: Anthropogenic and biophysical contributions to increasing atmospheric CO<sub>2</sub> growth rate and airborne fraction, *Biogeosciences*, 5, 1601–1613, <https://doi.org/10.5194/bg-5-1601-2008>, 2008.
- Raupach, M. R., Gloor, M., Sarmiento, J. L., Canadell, J. G., Frölicher, T. L., Gasser, T., Houghton, R. A., Le Quéré, C., and Trudinger, C. M.: The declining uptake rate of atmospheric CO<sub>2</sub> by land and ocean sinks, *Biogeosciences*, 11, 3453–3475, 1105 <https://doi.org/10.5194/bg-11-3453-2014>, 2014.
- Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, *J. Geophys. Res. Atmos.*, 108, <https://doi.org/https://doi.org/10.1029/2002JD002670>, 2003.
- 1110
- Regnier, P., Resplandy, L., Najjar, R. G., and Ciais, P.: The land-to-ocean loops of the global carbon cycle, *Nature*, 603, 401–410, <https://doi.org/10.1038/s41586-021-04339-9>, 2022.
- Revelle, R. and Suess, H. E.: Carbon Dioxide Exchange Between Atmosphere and Ocean and the Question of an Increase of 1115 Atmospheric CO<sub>2</sub> during the Past Decades, 9, 18–27, <https://doi.org/https://doi.org/10.1111/j.2153-3490.1957.tb01849.x>, 1957.
- Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O’Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., KC, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, 1120 T., Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview, *Glob. Environ. Chang.*, 42, 153–168, <https://doi.org/https://doi.org/10.1016/j.gloenvcha.2016.05.009>, 2017.
- 1125
- Ridge, S. M. and McKinley, G. A.: Ocean carbon uptake under aggressive emission mitigation, *Biogeosciences*, 18, 2711–2725, <https://doi.org/10.5194/bg-18-2711-2021>, 2021.
- Rödenbeck, C., Bakker, D. C. E., Gruber, N., Iida, Y., Jacobson, A. R., Jones, S., Landschützer, P., Metz, N., Nakaoka, S., 1130 Olsen, A., Park, G.-H., Peylin, P., Rodgers, K. B., Sasse, T. P., Schuster, U., Shutler, J. D., Valsala, V., Wanninkhof, R., and

- Zeng, J.: Data-based estimates of the ocean carbon sink variability – first results of the Surface Ocean  $p\text{CO}_2$  Mapping intercomparison (SOCOM), *Biogeosciences*, 12, 7251–7278, <https://doi.org/10.5194/bg-12-7251-2015>, 2015.
- Rodgers, K. B., Schwinger, J., Fassbender, A. J., Landschützer, P., Yamaguchi, R., Frenzel, H., Stein, K., Müller, J. D., Goris, N., Bushinsky, S. M., Chau, T. T. T., Gehlen, M., Gallego, M. A., Gloege, L., Gregor, L., Gruber, N., Hauck, J., Iida, Y., Ishii, M., Keppler, L., Kim, J.-E., Schlunegger, S., Sharma, S., Tjiputra, J., Toyama, K., and Vaittinada Ayar, P.: Seasonal variability of the surface ocean carbon cycle: a synthesis, *Global Biogeochem. Cycles*, 2023.
- Sarmiento, J. L., Orr, J. C., and Siegenthaler, U.: A perturbation simulation of  $\text{CO}_2$  uptake in an ocean general circulation model, *J. Geophys. Res. Ocean.*, 97, 3621–3645, <https://doi.org/https://doi.org/10.1029/91JC02849>, 1992.
- Schlunegger, S., Rodgers, K. B., Sarmiento, J. L., Ilyina, T., Dunne, J. P., Takano, Y., Christian, J. R., Long, M. C., Frölicher, T. L., Slater, R., and Lehner, F.: Time of Emergence and Large Ensemble Intercomparison for Ocean Biogeochemical Trends, *Global Biogeochem. Cycles*, 34, e2019GB006453, <https://doi.org/https://doi.org/10.1029/2019GB006453>, 2020.
- Séférian, R., Gehlen, M., Bopp, L., Resplandy, L., Orr, J. C., Marti, O., Dunne, J. P., Christian, J. R., Doney, S. C., Ilyina, T., Lindsay, K., Halloran, P. R., Heinze, C., Segschneider, J., Tjiputra, J., Aumont, O., and Romanou, A.: Inconsistent strategies to spin up models in CMIP5: implications for ocean biogeochemical model performance assessment, *Geosci. Model Dev.*, 9, 1827–1851, <https://doi.org/10.5194/gmd-9-1827-2016>, 2016.
- Sellar, A. A., Walton, J., Jones, C. G., Wood, R., Abraham, N. L., Andrejczuk, M., Andrews, M. B., Andrews, T., Archibald, A. T., de Mora, L., Dyson, H., Elkington, M., Ellis, R., Florek, P., Good, P., Gohar, L., Haddad, S., Hardiman, S. C., Hogan, E., Iwi, A., Jones, C. D., Johnson, B., Kelley, D. I., Kettleborough, J., Knight, J. R., Köhler, M. O., Kuhlbrodt, T., Liddicoat, S., Linova-Pavlova, I., Mizielinski, M. S., Morgenstern, O., Mulcahy, J., Neining, E., O'Connor, F. M., Petrie, R., Ridley, J., Rioual, J.-C., Roberts, M., Robertson, E., Rumbold, S., Seddon, J., Shepherd, H., Shim, S., Stephens, A., Teixeira, J. C., Tang, Y., Williams, J., Wiltshire, A., and Griffiths, P. T.: Implementation of U.K. Earth System Models for CMIP6, *J. Adv. Model. Earth Syst.*, 12, e2019MS001946, <https://doi.org/https://doi.org/10.1029/2019MS001946>, 2020.
- Silvy, Y., Frölicher, T. L., Terhaar, J., Joos, F., Burger, F. A., Lacroix, F., Allen, M., Bernadello, R., Bopp, L., Brovkin, V., Buzan, J. R., Cadule, P., Dix, M., Dunne, J., Friedlingstein, P., Georgievski, G., Hajima, T., Jenkins, S., Kawamiya, M., Kiang, N. Y., Lapin, V., Lee, D., Lerner, P., Mengis, N., Monteiro, E. A., Paynter, D., Peters, G. P., Romanou, A., Schwinger, J., Sparrow, S., Stofferahn, E., Tjiputra, J., Tourigny, E., and Ziehn, T.: AERA-MIP: Emission pathways, remaining budgets and carbon cycle dynamics compatible with 1.5°C and 2°C global warming stabilization, 2024, 1–47, <https://doi.org/10.5194/egusphere-2024-488>, 2024.

Terhaar, J., Torres, O., Bourgeois, T., and Kwiatkowski, L.: Arctic Ocean acidification over the 21st century co-driven by anthropogenic carbon increases and freshening in the CMIP6 model ensemble, *Biogeosciences*, 18, 2221–2240, <https://doi.org/10.5194/bg-18-2221-2021>, 2021a.

1170 Terhaar, J., Frölicher, T., and Joos, F.: Southern Ocean anthropogenic carbon sink constrained by sea surface salinity, *Sci. Adv.*, 7, 5964–5992, <https://doi.org/10.1126/sciadv.abd5964>, 2021b.

Terhaar, J., Frölicher, T. L., Aschwanden, M. T., Friedlingstein, P., and Joos, F.: Adaptive emission reduction approach to reach any global warming target, *Nat. Clim. Chang.*, 12, 1136–1142, <https://doi.org/10.1038/s41558-022-01537-9>, 2022a.

Terhaar, J., Frölicher, T. L., and Joos, F.: Observation-constrained estimates of the global ocean carbon sink from Earth System Models, *Biogeosciences*, 19, 4431–4457, <https://doi.org/10.5194/bg-19-4431-2022>, 2022b.

1180 Terhaar, J., Frölicher, T. L., and Joos, F.: Ocean acidification in emission-driven temperature stabilization scenarios: the role of TCRE and non-CO<sub>2</sub> greenhouse gases, *Environ. Res. Lett.*, 18, <https://doi.org/10.1088/1748-9326/acaf91>, 2023.

Terhaar, J., Goris, N., Müller, J. D., DeVries, T., Gruber, N., Hauck, J., Perez, F. F., and Séférian, R.: Assessment of global ocean biogeochemical models for ocean carbon sink estimates in RECCAP2 and recommendations for future studies, *Global Biogeochem. Cycles*, *in press* [Cycles](https://doi.org/10.1029/2023MS003840), 16, e2023MS003840, <https://doi.org/10.1029/2023MS003840>, 2024.

Thompson, D. W. J. and Solomon, S.: Interpretation of Recent Southern Hemisphere Climate Change, *Science*, 296, 895–899, <https://doi.org/10.1126/science.1069270>, 2002.

1190 Tjiputra, J. F., Schwinger, J., Bentsen, M., Morée, A. L., Gao, S., Bethke, I., Heinze, C., Goris, N., Gupta, A., He, Y.-C., Olivié, D., Seland, Ø., and Schulz, M.: Ocean biogeochemistry in the Norwegian Earth System Model version 2 (NorESM2), *Geosci. Model Dev.*, 13, 2393–2431, <https://doi.org/10.5194/gmd-13-2393-2020>, 2020.

1195 Tsujino, H., Urakawa, S., Nakano, H., Small, R. J., Kim, W. M., Yeager, S. G., Danabasoglu, G., Suzuki, T., Bamber, J. L., Bentsen, M., Böning, C. W., Bozec, A., Chassignet, E. P., Curchitser, E., Boeira Dias, F., Durack, P. J., Griffies, S. M., Harada, Y., Ilicak, M., Josey, S. A., Kobayashi, C., Kobayashi, S., Komuro, Y., Large, W. G., Le Sommer, J., Marsland, S. J., Masina, S., Scheinert, M., Tomita, H., Valdivieso, M., and Yamazaki, D.: JRA-55 based surface dataset for driving ocean–sea-ice models (JRA55-do), *Ocean Model.*, 130, 79–139, <https://doi.org/10.1016/j.ocemod.2018.07.002>, 2018.

Watson, A. J., Schuster, U., Shutler, J. D., Holding, T., Ashton, I. G. C., Landschützer, P., Woolf, D. K., and Goddijn-Murphy, L.: Revised estimates of ocean-atmosphere CO<sub>2</sub> flux are consistent with ocean carbon inventory, *Nat. Commun.*, 11, 4422, <https://doi.org/10.1038/s41467-020-18203-3>, 2020.

Weijer, W., Cheng, W., Garuba, O. A., Hu, A., and Nadiga, B. T.: CMIP6 Models Predict Significant 21st Century Decline of the Atlantic Meridional Overturning Circulation, *Geophys. Res. Lett.*, 47, e2019GL086075, <https://doi.org/https://doi.org/10.1029/2019GL086075>, 2020.

Yang, X. and Wang, M.: Monsoon ecosystems control on atmospheric CO<sub>2</sub> interannual variability: Inferred from a significant positive correlation between year-to-year changes in land precipitation and atmospheric CO<sub>2</sub> growth rate, *Geophys. Res. Lett.*, 27, 1671–1674, <https://doi.org/https://doi.org/10.1029/1999GL006073>, 2000.

Yasunaka, S., Manizza, M., Terhaar, J., Olsen, A., Yamaguchi, R., Landschützer, P., Watanabe, E., Carroll, D., Adiwara, H., Müller, J. D., and Hauck, J.: An assessment of CO<sub>2</sub> uptake in the Arctic Ocean from 1985 to 2018, *Global Biogeochem. Cycles*, 2023.

Zeng, N., Mariotti, A., and Wetzel, P.: Terrestrial mechanisms of interannual CO<sub>2</sub> variability, *Global Biogeochem. Cycles*, 19, <https://doi.org/https://doi.org/10.1029/2004GB002273>, 2005.

Zeng, J., Iida, Y., Matsunaga, T., and Shirai, T.: Surface ocean CO<sub>2</sub> concentration and air-sea flux estimate by machine learning with modelled variable trends, *Front. Mar. Sci.*, 9, 989233, <https://doi.org/10.3389/fmars.2022.989233>, 2022.

Ziehn, T., Chamberlain, M. A., Law, R. M., Lenton, A., Bodman, R. W., Dix, M., Stevens, L., Wang, Y.-P., and Srbinovsky, J.: The Australian Earth System Model: ACCESS-ESM1.5, *J. South. Hemisph. Earth Syst. Sci.*, 70, 193–214, 2020.