

Overview: Quasi-Lagrangian observations of Arctic air mass transformations – Introduction and initial results of the HALO-($\mathcal{A}\mathcal{C}$)³ aircraft campaign

Manfred Wendisch¹, Susanne Crewell², André Ehrlich¹, Andreas Herber³, Benjamin Kirbus¹, Christof Lüpkes³, Mario Mech², Steven J. Abel⁴, Elisa F. Akansu⁵, Felix Ament⁶, Clémantyne Aubry^{7,8}, Sebastian Becker¹, Stephan Borrmann^{9,10}, Heiko Bozem¹⁰, Marlen Brückner¹, Hans-Christian Clemen⁹, Sandro Dahlke³, Georgios Dekoutsidis⁷, Julien Delanoë⁸, Elena De La Torre Castro^{7,10,11}, Henning Dorff⁶, Regis Dupuy¹², Oliver Eppers⁹, Florian Ewald⁷, Geet George^{13,14}, Irina V. Gorodetskaya¹⁵, Sarah Grawe⁵, Silke Groß⁷, Jörg Hartmann³, Silvia Henning⁵, Lutz Hirsch¹³, Evelyn Jäkel¹, Philipp Joppe^{9,10}, Olivier Jourdan¹², Zsafia Jurányi³, Michail Karalis¹⁶, Mona Kellermann⁵, Marcus Klingebiel¹, Michael Lonardi^{1,17}, Johannes Lucke^{7,11}, Anna Luebke¹, Maximilian Maahn¹, Nina Maherndl¹, Marion Maturilli³, Bernhard Mayer¹⁸, Johanna Mayer⁷, Stephan Mertes⁵, Janosch Michaelis^{3,19}, Michel Michalkov⁵, Guillaume Mioche¹², Manuel Moser^{7,10}, Hanno Müller¹, Roel Neggers², Davide Ori², Daria Paul², Fiona Paulus², Christian Pils⁵, Felix Pithan³, Mira Pöhlker^{5,1}, Veronika Pörtge¹⁸, Maximilian Ringel⁶, Nils Risse², Gregory C. Roberts²⁰, Sophie Rosenberg¹, Johannes Röttenbacher¹, Janna Rückert²¹, Michael Schäfer¹, Jonas Schaefer⁵, Vera Schemann², Imke Schirmacher², Jörg Schmidt¹, Sebastian Schmidt²², Johannes Schneider⁹, Sabrina Schnitt², Anja Schwarz¹, Holger Siebert⁵, Harald Sodemann^{23,24}, Tim Sperzel^{1,25}, Gunnar Spreen²¹, Bjorn Stevens¹³, Frank Stratmann⁵, Gunilla Svensson¹⁶, Christian Tatzelt⁵, Thomas Tuch⁵, Timo Vihma²⁶, Christiane Voigt^{7,10}, Lea Volkmer¹⁸, Andreas Walbröl², Anna Weber¹⁸, Birgit Wehner⁵, Bruno Wetzel⁵, Martin Wirth⁷, and Tobias Zinner¹⁸

¹Leipziger Institut für Meteorologie (LIM), Universität Leipzig, Leipzig, Deutschland

²Institut für Geophysik und Meteorologie (IGM), Universität zu Köln, Köln, Deutschland

³Alfred-Wegener-Institut, Helmholtz-Zentrum für Polar- und Meeresforschung (AWI), Bremerhaven & Potsdam, Deutschland

⁴Met Office, Exeter, United Kingdom

⁵Leibniz-Institut für Troposphärenforschung (TROPOS), Leipzig, Deutschland

⁶Meteorologisches Institut, Universität Hamburg, Hamburg, Germany

⁷Institut für Physik der Atmosphäre, Deutsches Zentrum für Luft- und Raumfahrt (DLR), Oberpfaffenhofen, Deutschland

⁸Laboratoire Atmosphères, Milieux et Observations Spatiales (LATMOS), Centre National de la Recherche Scientifique (CNRS), Guyancourt, France

⁹Abteilung für Partikelchemie, Max-Planck-Institut für Chemie (MPIC), Mainz, Deutschland

¹⁰Institut für Physik der Atmosphäre (IPA), Johannes Gutenberg-Universität, Mainz, Deutschland

¹¹Faculteit Luchtvaart- en Ruimtevaarttechniek, Technische Universiteit Delft, Delft, Nederland

¹²Laboratoire de Météorologie Physique (LaMP), Université Clermont Auvergne, Centre National de la Recherche Scientifique (CNRS), Clermont-Ferrand, France

¹³Max-Planck-Institut für Meteorologie (MPIM), Hamburg, Deutschland

¹⁴Now at: Delft University of Technology (TU Delft), Delft, Netherlands

¹⁵Centro de Estudos do Ambiente e do Mar (CESAM), Universidade de Aveiro, Aveiro, Portugal

¹⁶Department of Meteorology and Bolin Centre for Climate research, Stockholm University, Stockholm, Sweden

¹⁷Now at: Extreme Environments Research Laboratory (EERL), Ecole Polytechnique Fédérale de Lausanne (EPFL), Sion, Switzerland

¹⁸Meteorologisches Institut, Ludwig-Maximilians-Universität München, München, Deutschland

¹⁹Now at: Maritime Klimatologie, Maritim-klimatologische Analysen und Produkte, Deutscher Wetterdienst (DWD), Hamburg, Deutschland

²⁰Scripps Institution of Oceanography, University of California San Diego, La Jolla, USA

²¹Institut für Umweltphysik (IUP), Universität Bremen, Bremen, Deutschland

²²Department of Atmospheric and Oceanic Sciences, Laboratory for Atmospheric and Space Physics (LASP), University of Colorado, Boulder, USA

²³Geophysical Institute, University of Bergen, Bergen, Norway

²⁴Bjerknes Centre for Climate Research, Bergen, Norway

²⁵Now at: Deutscher Wetterdienst, Offenbach am Main, Deutschland

²⁶Finnish Meteorological Institute (FMI), Helsinki, Finland

Correspondence: Manfred Wendisch (m.wendisch@uni-leipzig.de)

Abstract. ~~The global~~ Global warming is amplified in the Arctic. However, numerical models struggle to represent key processes that determine Arctic weather and climate. To collect data that help to constrain ~~weather and climate models, which often do not realistically represent the enhanced Arctic warming, the~~ the models, the HALO-($\mathcal{A}\mathcal{C}$)³ aircraft campaign was conducted ~~in March and April 2022~~ over the Norwegian and Greenland Seas, the Fram Strait, and the central Arctic Ocean .

5 ~~Observations were made over areas of open ocean, the marginal sea ice zone, and the central Arctic sea ice. Two low-flying and one long-range, high-altitude research aircraft have been employed. Whenever possible, the three aircraft were flown in collocated formation.~~ in March and April 2022. The campaign focused on one specific challenge posed by the models: The reasonable representation of transformations of air masses during their meridional transport into ~~(northward by and out of the Arctic via northward~~ moist and warm air intrusions ~~,~~ (WAIs) and ~~out of (southward via southward~~ marine cold air outbreaks

10 ~~,~~ (CAOs) (CAOs). ~~Observations were made over areas of open ocean, the marginal sea ice zone, and the central Arctic sea ice. Two low-flying and one long-range, high-altitude research aircraft were flown in collocated formation whenever possible. To follow the Arctic. To observe the~~ air mass transformations, a quasi-Lagrangian flight strategy using trajectory calculations was realized enabling to sample the ~~moving air mass same, moving air~~ parcels twice along their trajectories. ~~Eight~~ Seven distinct WAI and 12 CAO cases were probed ~~extensively.~~ From the quasi-Lagrangian measurements, we have ~~derived~~ quantified the

15 ~~diabatic heating and moistening of the moving air masses during CAOs and WAIs, the development of cloud macrophysical and microphysical properties along the southward pathways of the air masses during CAOs, and the moisture budget of WAIs. As an example result, we have obtained typical values of the surface-driven diabatic heating between 1–3/cooling and moistening/drying of the transported air masses. During CAOs, maximum values of 3 K h⁻¹ and of the near-surface moistening between 0.05–0.3 warming and 0.3 g kg⁻¹ h⁻¹ within the lowest about 0.5 moistening were obtained below 1 km altitude.~~ From

20 the observations of WAIs, ~~a weak diabatic cooling~~ diabatic cooling rates of up to 0.4 K h⁻¹ and a moisture loss of up to 0.1 g kg⁻¹ h⁻¹ from the ground to about ~~5.5~~ 5 km altitude were derived. Furthermore, the development of cloud macrophysical (cloud top height and horizontal cloud cover) and microphysical (liquid water path, precipitation, ice index) properties along the southward pathways of the air masses were documented during CAOs, and the moisture budget during a specific WAI event was estimated. In addition, we discuss the statistical frequency of occurrence of the different thermodynamic phases of Arctic

25 low-level clouds, the interaction of Arctic cirrus with sea ice ~~and~~ water vapor, and ~~aerosol particles, and the characteristic~~
the characteristics of microphysical and chemical properties of Arctic aerosol particles. Finally, we provide proof of concept
to measure mesoscale divergence and subsidence in the Arctic using data from dropsondes released during ~~circular flight~~
~~patterns~~the flights.

1 Introduction

30 In 2017, anthropogenic ~~global warming~~ warming quantified by the globally and annually averaged near-surface air temperature
reached around 1 K above the pre-industrial levels (Masson-Delmotte et al., 2021). ~~To date, human-caused climate change~~
~~has resulted in 1.2~~level (Masson-Delmotte et al., 2021). ~~In 2022, the human-induced warming averaged 1.26 K of warming~~
~~(Forster et al., 2023)~~. ~~Data over the decade 2013-2022 (Forster et al., 2023)~~. For 2023, the data published by the Copernicus
Climate Change Service show that ~~in 2023, on some days, the daily global average temperature was more than~~on almost 50 %
35 of days in that year the anthropogenic warming exceeded the values of the pre-industrial period (1850–1900) by at least 1.5 K
~~warmer than during pre-industrial times~~(<https://climate.copernicus.eu/global-climate-highlights-2023>). The advancing global
warming triggers numerous feedback mechanisms ~~in~~within the Earth's climate system, most of which are not fully accounted
for in ~~climate~~corresponding numerical models (Ripple et al., 2023). Of the 41 important feedback loops identified by Ripple
et al. (2023), at least a quarter cause ~~relevant~~distinct, mainly amplifying effects in the Arctic. Prominent examples of these
40 Arctic-relevant feedback mechanisms are the Planck, water vapor, surface albedo, and cloud effects. This makes the Arctic one
of the "hot spots" of global climate change (Overland et al., 2011).

~~As a result of global warming and these interlinked feedback mechanisms, a new Arctic has emerged over the last three~~
~~decades (Jeffries et al., 2013). The rapidly progressing changes in many Arctic climate parameters are obviously imprinted in~~
~~properties such as Arctic sea ice (extent, thickness, age), sea surface temperature, near-surface air temperature, Greenland ice~~
45 ~~sheet and glaciers, terrestrial snow cover, and permafrost, precipitation, primary biological productivity of the Arctic Ocean~~
~~and others (Druckenmiller et al., 2022). This enhanced response of the Arctic climate system to global warming is commonly~~
~~referred to as Arctic amplification (Serreze and Francis, 2006; Graverson et al., 2008; Serreze et al., 2009; Serreze and Barry, 2011; Previd~~
~~et al., 2014).~~

~~One important~~One obvious indication of Arctic amplification is the up to four times faster increase in Arctic near-surface
50 air temperature compared to global warming over the last three to four decades, which fits only poorly into the scatter of the
multi-model ensemble results of the Coupled Model Intercomparison Project (CMIP), Phase 5 (CMIP5) and Phase 6 (CMIP6)
~~(Holland and Landrum, 2021; Rantanen et al., 2022). Chylek et al. (2022) concludes that after 1999, Arctic amplification appears~~
~~to be caused by internal variability, which is not appropriately represented in CMIP6 models.~~(Holland and Landrum, 2021; Rantanen et al.,
2022). Further obvious signs of Arctic amplification are the faster-than-expected remarkable decline of sea ice cover of the Arctic
55 Ocean since around 1970, especially in late summer (Stroeve et al., 2007; Olonscheck et al., 2019; Serreze and Meier, 2019;
Screen, 2021), and the ~~gradual~~ thawing of the permafrost soils (Beer et al., 2020). These changes have important consequences
for the living conditions of the local Arctic human population, as well as for the flora and fauna of the Arctic. They also imply

potentially far-reaching economic impacts for fishing in Arctic waters, transoceanic shipping routes, tourism, and the extraction of natural resources ([Alvarez et al., 2020](#); [Melia et al., 2016](#)).

60 Arctic amplification has long been understood to be a feature of global climate change (Manabe and Wetherald, 1975). More recently, knowledge and understanding of the processes and feedback mechanisms governing Arctic amplification have improved considerably (Previdi et al., 2021; Smith et al., 2021; Taylor et al., 2022; Wendisch et al., 2023a). Nevertheless, the current ability to model them is still limited and, therefore, future model-based projections of Arctic climate changes are highly uncertain (Smith et al., 2019; Cohen et al., 2020; Block et al., 2020; Linke et al., 2023). In particular, the model representations
65 of the effects and development of clouds (Pithan et al., 2014; Wendisch et al., 2019; Kretzschmar et al., 2020; Stevens and Kluft, 2023), and of the interactions of the atmosphere with sea ice, snow on sea ice, and ocean physics as well as biogeochemical feedback processes are challenging (Rinke et al., 2019; Huang et al., 2019; Pefanis et al., 2020). In addition, the role of aerosol particles in Arctic amplification has not been sufficiently investigated (Schmale et al., 2021; Dada et al., 2022; Gong et al., 2023).

70 For more than a decade, there is some debate as to whether climate changes in the Arctic will impact the weather and climate in the mid-latitudes (Cohen et al., 2014). Several dynamic processes in the lower and upper troposphere and the stratosphere may lead to a weaker or stronger jet stream (Francis and Vavrus, 2015; Blackport and Screen, 2020; Yuval and Kaspi, 2020), with consequences on the meandering (amplitude) and persistence of the Rossby waves. These dynamical effects influence the meridional transport of heat, moisture, and momentum through northward warm air intrusions (WAIs¹), including
75 so-called atmospheric rivers (ARs), and southward cold air outbreaks (CAOs²). More frequent WAIs could further enhance Arctic warming (Pithan et al., 2018; Nash et al., 2018), whereas CAOs may contribute to cold events in mid-latitudes. Recent studies suggest that despite of the Arctic warming, cold winter events in mid-latitudes have remained nearly as extreme and as common as decades ago (Cohen et al., 2023; Nygård et al., 2023). ~~However, the~~ [The](#) results of the Polar Amplification Model Intercomparison Project (PAMIP) show very little evidence of linkages (Smith et al., 2022).

80 ~~In any case,~~ Rossby waves realize the meridional transport of air masses into and out of the Arctic. It is estimated that WAIs increase total column water vapor and cloud prevalence in the Arctic winter by about 70 % and 30 %, respectively (~~Johansson et al., 2017a~~)([Johansson et al., 2017b](#)). As a result, stronger thermal-infrared downward radiation reduces the net surface radiative cooling and increase the near-surface air temperature in winter by about 5 K (Johansson et al., 2017a). This warming could trigger an earlier onset of melting and more melt ponds, which would reduce the surface albedo. In addition,
85 particles and pollution are transported into the Arctic during WAIs, which may influence cloud properties (Bossioli et al., 2021).

In spite of the high impact they have on the Arctic climate, there are problems in modeling air mass transformations during meridional transport (Sato et al., 2016; Pithan et al., 2016; Dimitrelos et al., 2020). In particular, large-scale models have difficulties representing important thermodynamic processes driving Arctic air mass transformations, including the evolution
90 of microphysical properties of mixed-phase clouds (McCoy et al., 2015; Pithan et al., 2014; Tan and Storelvmo, 2019), the

¹In our perspective, WAIs include the northward transport of both warm and humid air.

²In this paper we restrict ourselves to marine CAOs; [our project does not investigate land surfaces such as the Greenland ice sheets](#).

development of turbulent fluxes under stable stratification (Tjernström et al., 2005; Holtslag et al., 2013; Gryanik and Lüpkes, 2023), and the response of snow-covered sea ice to atmospheric forcing (Pithan et al., 2023). These processes control the response of the Arctic to climatic forcing, and their realistic representation in models is crucial for understanding the behavior and feedback mechanisms of the Arctic climate system (Block et al., 2020; Taylor et al., 2022).

95 While the large-scale conditions that favor the development of CAOs can be well predicted on sub-seasonal time scales, a better understanding of when and how CAOs lead to the development of polar lows is necessary to predict these events, which often have large impacts (Polkova et al., 2021). Furthermore, improvements to the observing system and the understanding and model representation of small-scale synoptic features and processes are needed to advance polar predictions on daily to seasonal time scales (Jung et al., 2016). Moreover, given the lack of in-situ observations, models are often evaluated using
100 reanalysis data. However, the fidelity of such atmospheric reanalyses can suffer from a low number of assimilated observations, as well as biases inherited from its driving model (Tjernström and Graversen, 2009).

As a consequence, dedicated ~~quasi-Lagrangian~~ observations of WAIs and CAOs would be helpful to improve the model capabilities in order to realistically represent processes that determine air mass transformations during meridional ~~transport transports~~ into and out of the Arctic (Wendisch et al., 2021). Lagrangian measurements are well suited for this purpose. The Lagrange approach assumes that the observations are made in relation to a coordinate system that moves together with the air mass. In this way, the changes in the properties of the same air parcel can be observed along its pathway. In contrast, the observations from an Eulerian perspective refer to a locally fixed coordinate system, so that the properties of successive, different air parcels are measured from a fixed position as a time series. Previous observations of air mass transformations in the Arctic have mostly been conducted in the ~~framework-of-case-studies-Eulerian framework~~ at locally fixed, ground-based
105 positions (~~Eulerian point of view~~) partly combined with ship, aircraft, or satellite data. Examples include case studies based on data from the Surface Heat Budget of the Arctic Ocean (SHEBA) (Uttal et al., 2002) and the Multidisciplinary drifting Observatory for the Study of Arctic Climate (MOSAiC) (Shupe et al., 2022; Kirbus et al., 2023; Svensson et al., 2023) ship expeditions. Due to the lack of geostationary satellite data, the development of Arctic cloud properties can only be analyzed by polar orbiting satellite observations from an Eulerian perspective assuming stationary conditions (Murray-Watson et al., 2023).
115 However, the Eulerian approach does not permit the required observations of temporal air mass-transforming processes.

Only very few ~~quasi-Lagrangian-Lagrangian~~ aircraft-based studies have been carried out ~~so far in the past~~ (Boettcher et al., 2021), and none in the Arctic. ~~Furthermore, several studies use atmospheric reanalysis to identify and discuss WAIs or CAOs (You et al., 2021b; Kirbus et al., 2024).~~ Some authors have ~~put combined~~ local icebreaker and airborne or satellite observations ~~into~~ within a Lagrangian framework by means of trajectories based on reanalysis (Tjernström et al., 2019; Ali and Pithan,
120 2020; You et al., 2021a; Kirbus et al., 2023; Mateling et al., 2023). Another approach was performed in earlier aircraft-based campaigns with measurements along the mean wind direction in CAO conditions and WAIs (Hartmann et al., 1997; Brümmer and Thiemann, 2002; Vihma et al., 2003; Lüpkes et al., 2012; Chechin et al., 2013). These authors have focused on the local development around the Fram Strait and the close marginal sea ice zone (MIZ) with measurements above sea ice and open ocean. A disadvantage was that the quasi-Lagrangian analysis had to assume stationary conditions. This strategy turned
125 out to be helpful, e.g., for the analysis and development of turbulence parametrizations. They could be validated by using

them in mesoscale models and comparing their results with the observations. But then it was always difficult to identify the impact of changing inflow conditions during the necessary 3-6 hours model runs, a drawback, which can be avoided by purely Lagrangian measurements. Furthermore, several studies use atmospheric reanalysis to identify and discuss WAIs or CAOs (You et al., 2021b; Kirbus et al., 2024).

130 Therefore, we have conducted the HALO–(\mathcal{AC})³ aircraft campaign (HALO: High Altitude and Long Range Research Aircraft, (\mathcal{AC})³: Project on Arctic Amplification: Climate Relevant Atmospheric and Surface Processes, and Feedback Mechanisms, see <https://halo-ac3.de/>). Based on the open issues in modeling air mass transformations during meridional transport into and out of the Arctic, the HALO–(\mathcal{AC})³ mission pursued two general objectives (Wendisch et al., 2021). The first was to jointly use HALO and the Polar 5 (P5) and Polar 6 (P6) research aircraft to perform "quasi-Lagrangian"³ observations of
135 air mass transformations during WAIs and CAOs – an approach that has not been tried before in the Arctic. The second was to test the ability of numerical atmospheric models to reproduce the measurements taken from the aircraft. The benchmarked models can then for example be applied to investigate linkages between Arctic amplification and mid-latitude weather. This paper describes efforts carried out in support of the first objective.

The article is structured in five sections. After the introduction ~~-, the mainly utilized (Sect. 1), the~~ three research aircraft
140 mainly utilized in our campaign and their instrumentation, as well as ~~the collocated flights their partly collocated flight patterns~~ are described in Sect. 2. The unique quasi-Lagrangian measurement strategy successfully applied during the campaign is presented in Sect. 3. Some initial results from HALO–(\mathcal{AC})³ and ongoing analysis are discussed in Sect. 4. The following themes are elaborated: Air mass transformations during WAIs and CAOs (Sect. 4.1), Arctic clouds (Sect. 4.2) and aerosol particles (Sect. 4.3), and a proof of concept to measure mesoscale divergence and subsidence in the Arctic (Sect. 4.4). A ~~summary~~
145 ~~and some conclusions~~ concise summary and a brief outlook are given in Sect. 5. ~~Three appendices cover some interesting supplementary aspects of the HALO–(\mathcal{AC})³ aircraft campaign.~~

2 Aircraft, instrumentation, and flight pattern

Three research aircraft were mainly involved in the HALO–(\mathcal{AC})³ ~~aircraft~~-campaign: HALO (High Altitude and Long Range Research Aircraft), Polar 5 (P5), and Polar 6 (P6). HALO is operated by the German Aerospace Center (Deutsches Zentrum
150 für Luft- und Raumfahrt, DLR). It was based in Kiruna (northern Sweden, geographical coordinates: 67.85° N, 20.22° E). The P5 and P6 were stationed at Longyearbyen (Svalbard, Norway, 78.24° N, 15.49° E). These two aircraft belong to the Alfred Wegener Institute, Helmholtz Center for Polar and Marine Research (AWI). In addition, the British Facility for Airborne Atmospheric Measurements (FAAM) and the French Avions de Transport Régional (ATR) aircraft were concurrently based in Kiruna measuring partly in coordination with HALO, P5, and P6. The ATR was operating from 22 March to 3 April 2022, and had
155 some similar objectives as HALO, but with the addition of using in-situ water isotope measurements to characterize processing of water vapor in these weather systems. The FAAM aircraft operated from 7 March to 1 April 2022, and focused on making in-situ measurements of the development of CAOs from the sea ice around Svalbard to the Scandinavian coastline. FAAM

³The term "quasi-Lagrangian" is introduced in Sect. 3.2 in detail.

was fitted with a range of instrumentation to measure the thermodynamic, aerosol, cloud, and precipitation properties within CAOs. ~~An overview of the instrumentation and all data collected during the campaign period is given by Ehrlich et al. (2024).~~
160 Furthermore, intensive ground-based measurements were carried out at the [permanent German-French AWIPEV research base](#) ~~in~~ [operated by German Alfred Wegener Institute \(AWI\) and the French Polar Institute Paul Émile Victor \(IPEV\) at Ny-Ålesund](#) ~~-(Svalbard, 78.92° N, 11.92° E),~~ including additional observations with a tethered balloon (Lonardi et al., 2024). [An overview of the instrumentation and all data collected during the campaign period is given by Ehrlich et al. \(2024\) \(to be submitted to Earth Syst. Sci. Data\).](#)

165 ~~In the following, the three main aircraft used during HALO (AC)³ are described.~~ HALO – a Gulfstream G550 – has sufficient range/endurance (up to 9000 km and 10 hours) for quasi-Lagrangian air mass observations. HALO is capable to lift up to three tons of state-of-the-art meteorological and remote sensing instruments up to 15 km altitude to observe the complete vertical tropospheric air mass column, including water vapor, aerosol particles, clouds, precipitation and surface properties. HALO was equipped with a unique remote sensing payload that has matured in several campaigns in the past (Stevens et al.,
170 2019; Konow et al., 2021). The instrumentation includes a 26-channel microwave radiometer, a 35 GHz Doppler radar, aerosol and water vapor lidar, spectral and broadband solar and thermal-infrared radiation sensors, which are upward and downward looking, imaging and polarization camera spectrometers in the solar and thermal-infrared spectral ranges, and dropsondes. HALO was operated from Kiruna between 7 March and 12 April 2022. 17 research flights (RFs) were conducted (RF02–RF18) with a total flight time of 147 hours (Fig. 1a). In total, 330 dropsondes were released ~~during the RFs~~ from HALO
175 (Tab. 1).

The low-flying P5 and P6 (Wesche et al., 2016) aircraft are Basler BT-67 (DC-3) types with a ceiling of up to 6 km, and a range of about 2300 km. Each of the two planes can carry a scientific payload ~~that can weigh up to~~ [with a maximum weight of 1 ton](#). P5 provided active and passive remote sensing measurements to characterize clouds, precipitation, aerosol particles, trace gases, and surface properties from atop, similar to HALO. The instrumentation of P5 included a 94 GHz radar, an aerosol
180 lidar, passive microwave, and radiation sensors. In addition, P5 carried in-situ instrumentation to derive turbulence parameters and energy fluxes (Mech et al., 2019; Schirmacher et al., 2023). ~~Dropsondes were also released from P5.~~

P6 focused on in-situ measurements in the lower troposphere ~~of~~ below 2–4 km altitude in cloudy and cloud-free conditions. The measurements of P6 aimed to determine radiative and turbulent energy fluxes as well as to investigate smaller-scale processes. For this purpose, P6 was equipped with in-situ probes to measure cloud and precipitation particles (droplets, ice
185 crystals) and cloud residuals, aerosol particles, radiation, chemistry, and trace gas properties. For cloud observations, the P6 aircraft was equipped with a Cloud Droplet Probe (CDP), Cloud Imaging Probe (CIP), 2D-Stereo Probe (2D-S), and Precipitation Imaging Probe (PIP) for cloud particle counting and sizing, along with a Polar Nephelometer (PN) for measuring scattering properties and phase discrimination (Wendisch and Brenguier, 2013; Kirschler et al., 2023; De La Torre Castro et al., 2023). The total data set of in-situ cloud measurements with P6 below 1.5 km sums up to about 21 hours, where about 15.5
190 hours were spent above the open ocean and 3.5 hours over sea ice.

The P5 and P6 aircraft followed a [proven carefully designed](#) measurement strategy with vertically stacked, collocated remote sensing measurements above clouds (P5) and in-situ sampling inside clouds at lower altitudes (P6). Experiences with this




Figure 1. Flight paths (a) of HALO operating from Kiruna (KRN), and (b) of Polar 5 (P5) and Polar 6 (P6) aircraft based in Longyearbyen (LYR).

measurement strategy were collected during the Arctic Cloud Observations Using airborne measurements during polar Day (ACLOUD) campaign performed in 2017 (Wendisch et al., 2019). During HALO-(AC)³ both aircraft performed 13 RFs with 195 53 (P5) and 63 (P6) flight hours, respectively (Fig. 1b). 141 drosondes were released during the flights of P5 (Tab. 1). Some coordinated flights between HALO and the P5 and P6 aircraft including closely vertically collocated flight segments were conducted. In addition, several joint flights of HALO, the ATR, and FAAM planes were realized.

3 Quasi-Lagrangian-Lagrangian air mass sampling

3.1 General approach Ideal Lagrangian observations using balloons

200 True Lagrangian observations would require flying embedded in a moving air mass parcel and recording data characterizing the changes of the air mass properties during its movement. The closest approach to achieving this would be balloons continuously accompanying measurements would require instruments that are (i) either embedded in the moving air parcel and taking in-situ data along its transport path way, or (ii) flying in higher altitudes and accompanying the moving air parcel by remote sensing observations from above. The most suited instrument platform to achieving approach (i) would be a balloon drifting with the moving air parcel and carrying devices continuously recording the air mass and simultaneously drifting along with

205

it transformation along its transport pathway. However, ~~such balloon measurements also suffer some balloon measurements suffer from several inevitable~~ drawbacks (Businger et al., 1996; Johnson et al., 2000; Businger et al., 2006; Roberts et al., 2016). ~~Because of their~~ For example, even balloons have some non-zero inertia, ~~drifting balloon observations do not pose a fully Lagrangian approach. In addition;~~ thus they do not serve as perfect instrument carrier to realize approach (i). Furthermore, due to vertical wind shear ~~of wind speed and direction~~, the air ~~mass trajectories follow different pathways in different altitudes.~~ ~~It quickly~~ parcel trajectories differ with altitude and change height levels along their pathways. Therefore, it becomes a non-trivial task to select an altitude level to follow. ~~Furthermore, the payload of such balloons is very limited. with the balloon.~~ Also, the mass of the instrumental payload that can be lifted by balloons appears quite limited. These and further shortcomings remain when using a balloon as instrumental platform carrying out remote sensing observations from above in approach (ii).

215 3.2 Quasi-Lagrangian approach applying aircraft and trajectories

Here, we ~~avoid these drawbacks~~ try to avoid most of the drawbacks of balloon-borne air mass observations by following a ~~new strategy termed~~ dedicated measurement strategy termed "quasi-Lagrangian ~~and that accounts, e. g., for the height dependence of trajectories.~~" and applying approach (ii). Instead of relying on the limited payload capacity of balloons, we employ three ~~different~~ research aircraft, equipped with ~~instrumentation that has shown its~~ comprehensive instrumentation that have proven ~~their~~ potential in state-of-the-art atmospheric and Earth's surface observations. Naturally, ~~such aircraft move~~ aircraft fly much faster than the relatively slow air masses moving with the respective wind speeds. To mitigate this ~~problem~~ mismatch, we design ~~appropriate~~ sophisticated flight patterns aiming to sample ~~air mass the very same air~~ parcels along their ~~pathway as often as possible~~ pathways at least twice during the same or consecutive flights. To realize this idea we utilize air ~~mass~~ parcel trajectory calculations to ~~determine the pathway~~ project (during the flight planning) or reanalyze (after the campaign) the pathways of the ~~air mass~~ parcels. In case the air ~~mass trajectory initially sampled~~ parcel trajectory is crossing the flight path ~~at any level below a second time~~ two times, we define this as a quasi-Lagrangian match.

This ~~approach~~ quasi-Lagrangian technique is exemplified by the sketch presented in Fig. 2 for the case of a WAI. At a time t_1 , we observe a vertical air Column 1 consisting of stacked air ~~mass~~ parcels. One such air ~~mass~~ parcel is indicated as a blue cube ~~in this sketch~~. A number of height-resolving remote sensing instruments aboard the high-flying HALO and P5, ~~such as~~ ~~radar, lidar and microwave radiometer, and dropsondes~~ and dropsondes released during flight characterize the properties of the air ~~mass~~ parcels at different altitudes ~~within Column 1~~ at t_1 . In addition, we ~~directly probe the air mass parcels with the P6 aircraft collecting~~ probe selected air parcels with in-situ ~~data~~ instruments installed on P6. Besides many other quantities, dry potential air temperature (θ), specific humidity (q), air temperature (T), and radar reflectivity (Z) are measured by remote sensing and in-situ instruments.

235 In the next step, we use forward trajectories (dashed arrows in Fig. 2) to follow the ~~pathway~~ pathways of the individual air ~~mass~~ parcels of the vertical air column. To calculate the trajectories, we define a horizontal circular area with a radius of 30 km in the center of the air ~~mass~~ parcel. As an example, we refer to the dashed ellipse within the blue cube of Column 1. Starting points are evenly spaced horizontally every 10 km, which results in about 30 regularly distributed points per starting altitude. From the starting points, simulations of the 30 forward-trajectories per starting altitude are performed to project the

240 average movement of the corresponding air **mass**-parcel (for example the blue cube in Fig. 2). Subsequently, we follow the same approach for each air **mass**-parcel within Column 1 to project the average movement of the air parcels as a function of altitude.

The vertical geometric thickness of the individual air **mass**-parcels is assumed as 5hPa, which also corresponds to the vertical resolution of the horizontal circular areas. The top of the vertical column is defined as 250 hPa (approximate flight altitude of HALO), which results in a total of 150 (750hPa divided by 5hPa) air **mass**-parcels and horizontal circular start areas for the trajectories. For each of the 150 air **mass**-parcels in Column 1, 4500 trajectories are started (150 parcels \times 30 regularly spaced initial points). These trajectories describe the movements of the air **mass**-parcels in Column 1, which were sampled by the three aircraft. Now, the whole procedure is repeated along the entire flight track with a temporal resolution of one minute. In the case of HALO, the approximate flight time during the HALO-(AC)³ campaign was about 8 hours per RF, which means that $4500 \times 480 = 2.2 \times 10^6$ air **mass**-parcel trajectories have been calculated for each HALO flight during the campaign.

~~Flight planning to realize quasi-Lagrangian observations required substantial efforts for the mission coordination. To combine flight tracks with the projected trajectories, the Mission Support Tool was used (Bauer et al., 2022).~~

3.3 Trajectories to find matches

255 ~~The actual~~ The computation of the forward-trajectories of the air **mass**-parcels was performed using the Lagrangian analysis tool (LAGRANTO) (Sprenger and Wernli, 2015). During the campaign, the trajectory calculations were based on the Integrated Forecast System (IFS) of the European Centre for Medium-Range Weather Forecasts (ECMWF) wind product. For processing the data after the campaign (for this paper) we have applied the Fifth Generation ECMWF Atmospheric Reanalysis (ERA5) (Hersbach et al., 2020). Both methods (IFS- and ERA5-based) have assimilated the dropsonde profile observations of thermodynamic and wind data taken during the flight (Table 1). Trajectories were calculated 60 hours forward in time. IFS and ERA5 were retrieved on 137 model levels, which are vertically spaced between the surface and top-of-atmosphere on a regular $0.25^\circ \times 0.25^\circ$ latitude/longitude grid with a one-hourly temporal resolution. The improved performance of ERA5 compared to alternative atmospheric reanalyses data has been shown by Graham et al. (2019a, b).

265 ~~Figure 2 illustrates the Flight planning to realize quasi-Lagrangian approach used for observations required substantial efforts for the mission coordination. To combine flight tracks with the projected trajectories, both were included in our Mission Support Tool (Bauer et al., 2022). Then the flight planning.~~ The flight plans were prepared such that there are enhanced chances to meet the air **mass**-parcel indicated as a blue cube a second time along its trajectory during the flight. This case is illustrated in Column 2 of Fig. 2 where actually the same blue cube ~~was encountered again within Column 1 was encountered a second~~ by the aircraft at time t_2 . Such a quasi-Lagrangian match is counted only (i) after at least 60 min of air **mass**-parcel drift, and (ii) if the air **mass**-parcel trajectory crosses the aircraft track within a 30 km radius. Each quasi-Lagrangian match enables to compare the air mass properties measured in Column 1 (subscript 1) with those measured during the quasi-Lagrangian match in Column 2 (subscript 2). Thus, the temporal tendency of a thermodynamic or cloud property ψ (e.g., dry potential air temperature θ , specific humidity q , and air temperature T) that characterizes the rate of change of a particular air mass ~~parcel~~ property along

its trajectory can be quantified using the following equation:

$$275 \quad \frac{\Delta\psi}{\Delta t} = \frac{\psi_2 - \psi_1}{t_2 - t_1}. \quad (1)$$




Figure 2. Illustration of the quasi-Lagrangian approach adopted during the HALO-(AC)³ airborne aircraft campaign. At an initial time t_1 and within the atmospheric Column 1, an air mass parcels-parcel (blue eubescube) are-is observed using remote sensing and in-situ instrumentation installed on HALO and P5, and P6. Trajectories are initiated-simulated that describe the movement of the air mass-parcels (dashed arrows). If they cross the flight path of the aircraft at a later time t_2 , the air mass-parcel can be sampled a second time within the atmospheric Column 2. This approach enables to-observe-observing the changes of the properties of the air mass-parcel (for example, θ , q , T , and Z in the time increment $(t_2 - t_1)$ along its trajectory).

3.3 Example and statistics of matches

In-the-following, we We illustrate the quasi-Lagrangian procedure for two consecutive HALO research flights (RF03 and RF04) performed within two days after each other consecutive days (13 and 14 March 2022). In Fig. 3, the notable corridor of increased northward water vapor transport (IVT/IVT) indicates a substantial WAI event. During RF03 on 13 March, the HALO flight path (blue line) was designed to cover the WAI by a horizontal zig-zag flight pattern (FigFigs. 3a-ba-c). HALO took off shortly after 8 UTC in Kiruna. The forward-trajectory-simulations-were-started forward trajectories were simulated along the HALO flight path with a one-minute resolution (black lines), whereby only a subset of simulated-trajectories-is-depicted-in-the

figure with start times corresponding to the HALO flight time series. Only those trajectories are depicted in Figs. 3a-c that later on during RF04 matched the HALO flight path a second time. These trajectories evolved until HALO landed in Kiruna around 17 UTC. During the night, the trajectories advanced indicating a continued transport of the humid air masses pole-ward (Fig. 3e). HALO took off on the second day (RF04) on 14 March around 10 UTC and crossed the trajectories that had been initiated the day before (Fig. 3d-f). Altogether, about 80,000 matches of air mass parcels (5 hPa vertical and 13-15 km horizontal size) at different altitudes have been obtained during these two consecutive HALO flights.

Warm Air Intrusion (WAI): 13 and 14 March 2022




Figure 3. Time series (based on ERA5) of geographic maps including the integrated water vapor transport (IVT , in red colors, with the length of arrows being proportional to the magnitude of IVT), and the HALO flight tracks on 13 March 2022 (RF03, blue line) and 14 March 2022 (RF04, red line). Black lines indicate the horizontal projections of the evolving matching trajectories. The blue-filled circles represent the trajectory start points, and the filled orange circles show where trajectories cross the second flight track.

The procedure described above was applied to all cases when two HALO flights took place on consecutive days. This included not only WAIs or atmospheric rivers (ARs), but also CAOs, Arctic cirrus (AC) and polar low (PL) cases (Walbröl

et al., 2023). The respective statistics of the number of quasi-Lagrangian matches are given in Fig. 4a. Numerous cases have been identified where individual air mass parcels, sampled on day one have been encountered a second time on day two. In a similar way, individual flights performed during a single day were analyzed (Fig. 4b). In these cases, matches between different flight sections along the individual flight were identified. Overall, the number of matches in both scenarios (two flights on consecutive days, one flight during one day) is more than sufficient for a statistical evaluation, scientific analysis, and discussion. This data set of quasi-Lagrangian matches provides a new quality-an unprecedented quantity of possibilities to observe and study atmospheric air mass transformations.




Figure 4. Number of matches for the two scenarios: (a) Two flights on consecutive days, and (b) one flight during one day. The analysis includes moist and warm air intrusions (WAI), atmospheric rivers (ARs), marine cold air outbreaks (CAOs), as well as Arctic cirrus (AC), and polar low (PL) cases.

4 Initial results and ongoing analysis

In this section, we provide the first results obtained from the measurements conducted during the HALO-(AC)³ aircraft campaign with respect to air mass transformations (thermodynamic tendencies) quantified during CAOs and WAIs, Arctic cloud and aerosol characteristics, and atmospheric turbulence measurements evolution during CAOs, the moisture budget during an example WAI, as well as some characteristic properties of Arctic clouds and aerosol particles. Furthermore, we discuss a proof of concept to measure mesoscale divergence and subsidence in the Arctic. ~~In the three appendices, we add partly preliminary, but nonetheless very interesting supplementary discussion and results from the HALO-(AC)³ campaign: Appendix A: A hypothesis on humidity fields in WAIs, Appendix B: The particular mode structure of the Arctic radiant energy budget, and Appendix C: An extraordinary CAO case covered during HALO-(AC)³.~~

4.1 Air mass transformations during CAOs and WAIs

4.1.1 Thermodynamic tendencies

To quantify-derive the tendencies of thermodynamic properties of air-mass-the moving air parcels, such as the dry potential air temperature (θ), specific humidity (q), and air temperature (T), we apply ~~the quasi-Lagrangian match analysis using~~ Eq. 1

with $\psi = \theta, q, z, T$. Corresponding results are depicted in Fig. 5. ~~Here, the vertical profiles of the numbers of, which illustrates the number of occurrence (counts) of certain values of the $\theta, q,$ and T tendencies are illustrated (Eq. 1), color bar) of the temporal tendencies $\Delta\theta/\Delta t, \Delta q/\Delta t,$ and $\Delta T/\Delta t$ as a function of altitude. In addition, the vertical displacement of the air mass-parcels along its trajectory between t_1 and t_2 is plotted. ~~The tendencies for the CAOs as $\Delta z/\Delta t$. The results for the CAO cases~~ (Figs. 5a–d) are derived for trajectories from the north to south, ~~i.e., therefore in general general~~ warming and moistening tendencies at low levels ~~occurs-were obtained~~ due to heat and moisture surface fluxes increasing from sea ice to open ocean. The WAI tendencies (Figs. 5e–h) concern the opposite sense, i.e., from south to north.~~

It is important to note that the diabatic heating and moistening presented in ~~FigFigs.~~ 5a-d merge the quasi-Lagrangian matches over open ocean exclusively (CAOs), whereby all available matches (over open ocean and sea ice) are considered in the tendencies for WAIs. In CAOs, major air mass transformations occur over the open ocean due to intense surface turbulent heat fluxes driven by temperature and humidity gradients, whereas the preconditioning over Arctic sea ice typically involves rates one order of magnitude smaller (Papritz and Spengler, 2017; Kirbus et al., 2024). Thus, in this article we focus only on the CAO processes setting in over open ocean. On the contrary, during WAIs intense air mass transformations through turbulent, radiative, and cloud processes can set in over open ocean, the marginal sea ice zone, as well as the sea ice (Woods and Caballero, 2016; Johansson et al., 2017a; You et al., 2022). Therefore, we do not restrain the analysis of air temperature and moisture changes during WAIs to any surface type.

Instead of the air temperature T , we first investigate the change-rate-temporal tendency of the dry potential air temperature θ , which is insensitive to dry adiabatic vertical movements of the air mass-parcel during transport. Furthermore, θ is characteristic of a moving air mass and changes only in response to diabatic processes. These include cloud evolution (release or consumption of latent heat), surface influences (such as turbulent and energy fluxes), as well as radiative processes. Using θ instead of T thus quantifies the influence of processes we are most interested in, namely the cloud and surface effects.

For the CAO cases (~~panels a and b~~), within the layer between the surface and about 0.51 km altitude, a surface-driven diabatic heating between $1\text{--}3 \text{ K h}^{-1}$ (Fig. 5a) and a moistening between $0.05\text{--}0.3 \text{ g kg}^{-1} \text{ h}^{-1}$ (Fig. 5b) are observed. Air mass-parcel trajectories are descending throughout most of the vertical column with a wider spread of upward and downward motion in the atmospheric boundary layer (ABL) (Fig. 5c). For the WAI observations, a weak diabatic cooling of up to 0.4 K h^{-1} (Fig. 5e), and a moisture loss of up to $0.1 \text{ g kg}^{-1} \text{ h}^{-1}$ (Fig. 5f) are observed, both reaching from the surface to heights to about 55.5 km. For a specific intense CAO (RF12; 1 April 2022, not shown here), Kirbus et al. (2024) derived a maximum diabatic heating larger than 6 K h^{-1} close to the ocean surface just downwind of the MIZ. Values of moisture uptake of more than $0.3 \text{ g kg}^{-1} \text{ h}^{-1}$ were observed in this study from Kirbus et al. (2024).

For the CAOs, slight subsidence of the air mass-parcels during transport is derived (Fig. 5c). If air temperature tendencies are evaluated instead of dry potential air temperature θ , ~~the~~ adiabatic warming effects due to subsidence for the CAO cases become apparent throughout the entire vertical column up to 8 km altitude (Fig. 5d), whereas no ~~such distinct~~ adiabatic warming is obvious ~~in from~~ Fig. 5a. No clear ascent/descent trends of the air parcels are obvious from Fig. 5g.




Figure 5. Vertical profiles of the number of occurrences (counts) of temporal tendencies of (a) dry potential air temperature (diabatic heating/cooling, $\Delta\theta/\Delta t$), (b) specific humidity (moistening/drying, $\Delta q/\Delta t$), (c) air mass-parcel ascent/descent ($\Delta z/\Delta t$), and (d) air temperature ($\Delta T/\Delta t$) for marine cold air outbreaks (CAOs) sampled with HALO on 20, 21, 28, 29, 30 March and 1 April 2022. Panels (e) to (h) illustrate the same temporal tendencies for moist and warm air intrusions (WAIs) observed with HALO between 12-16 March 2022. Vertical solid red lines indicate abscissa values of zero, tilted dashed red lines in panels a, b, e, d indicate the height range of main air mass modification.

Similar to Fig. 5, the cloud reflectivity Z measured by radar on HALO was used to follow the cloud evolution during CAOs and WAIs (not shown). In case of CAOs, clouds evolve mainly in lower altitudes (below 3 km). In case of WAI, cloud dissipation dominates mostly below 6 km altitude.

4.1.2 Development of cloud properties during CAOs

Firstly, we investigated how the preconditions over sea ice influence the crucial initial cloud formation using targeted research flights by ~~the P5~~, which statistically sampled the developing roll convection just behind the MIZ (80–100 % sea ice concentration). Contrary to previous ~~airborne aircraft~~ studies that followed the air mass downstream along developing cloud streets, the P5 flew multiple legs orthogonal to them. In this way, the roll convection forming the cloud streets could be identified from radar profiles and characterized statistically with respect to their macrophysical and microphysical cloud properties using multiple instruments (Schirmacher et al., 2024). Because air mass transformation is mainly triggered by the exposure of air to open water surfaces, we used backward trajectories to assign each measurement to its fetch, i.e., the horizontal distance the air mass traveled over open water until reaching P5. Two cases of CAOs ~~with many similarities but different strengths of different strength~~ observed on the 1st and 4th of April 2022 were analyzed ~~in detail by Schirmacher et al. (2024) and by Schirmacher et al. (2024). The results~~ are summarized briefly here.

The evolution of crucial parameters characterizing the cloud and precipitation development along the two CAOs ~~within the first 170 km is illustrated~~ as a function of fetch ~~is illustrated~~ in Fig. 6 ~~within the first 170 km~~. Cloud streets form approximately 15 km downwind of the sea ice edge reaching almost 100 % cloud cover for a fetch ~~exceeding of~~ 20 km in both cases. Due to the strong surface fluxes, the ABL grows quickly ~~along the cloud pathway~~, and cloud top height (~~CTH~~~~CTH~~) increases by about 4 m per km in the strong CAO case (1 April 2022), and only half as much for the weaker CAO (4 April 2022). Both CAO events feature mixed-phase clouds with the stronger case showing ~~about~~ twice the liquid water path (~~LWPLWP~~). Precipitation sets in after a fetch of about 30 km with a slightly later onset for the weaker event. The combination of remote sensing ~~from data measured with instruments installed on P5~~ with in-situ measurements ~~collected~~ by P6 ~~also allows us devices allows~~ to investigate the ice growth process (~~Maherndl et al., 2024~~) ~~as reported by Maherndl et al. (2024)~~. For the stronger CAO case, we ~~also~~ detect stronger riming, which occurs on a horizontal scale similar to the roll circulation.

To understand how CAOs develop from their initial phase along their way further south, HALO with its long range can provide valuable insights. ~~For the stronger CAO, images of spatial CTH and cloud thermodynamic phase were gathered the spectrometer of the Munich Aerosol Cloud Scanner (specMACS) instrument (Ewald et al., 2016; Weber et al., 2024) west of Svalbard during HALO RF12.~~ The spectral slope phase index defined by Ehrlich et al. (2008) was used to derive information about cloud thermodynamic phase from the measurements ~~-(Fig. 7a)~~. Values of the spectral slope phase index smaller than about 20 indicate pure liquid water, and larger values mixed-phase or ice clouds. ~~A transition from pure liquid water to mixed-phase clouds occurs within the first two hours after passing the MIZ.~~ In addition to the thermodynamic phase, we have retrieved ~~CTH~~~~CTH~~ from the specMACS observations with a stereographic method (Kölling et al., 2019; Volkmer et al., 2024) ~~-(Fig. 7b)~~. This data was ~~again~~ combined with back trajectories to calculate the time (~~instead of fetch~~) the measured air mass traveled above the open ocean after passing the MIZ (~~here 80~~ ~~abscissa in Fig. %~~). ~~Figure 7a-b~~. ~~For the stronger CAO case, images of spatial CTH and cloud thermodynamic phase were gathered the spectrometer of the Munich Aerosol Cloud Scanner (specMACS) instrument (Ewald et al., 2016; Weber et al., 2024) west of Svalbard during HALO RF12. Figures 7 shows c-e~~ illustrate three example images of the polarization cameras showing how the clouds in their initial phase organize in cloud

Two Cold Air Outbreaks (CAOs): Strong—1 April 2022, ~~weak—4~~ Weak—4 April 2022




Figure 6. Development of ~~geometrical-macrophysical~~ and microphysical cloud properties as a function of fetch on 1 April (red line, strong CAO) and 4 April (blue line, weak CAO) 2022 as measured with instruments installed on P5: (a) Cloud top height (~~CTH~~CTH), (b) horizontal cloud cover per minute measured by the radar MiRAC and lidar AMALi, (c) liquid water path (~~LWP~~LWP), and (d) precipitation rate (~~P~~P) at 150 m height (~~S~~). The shaded areas indicate the 5 % and 95 % quantiles of the distributions ~~with~~ as a function of fetch. Figure adapted from Schirmacher et al. (2024).

streets and then develop into closed cells with increasing distance to the MIZ. While the initial phase with cloud tops up to 1 km was already captured by P5, the HALO observations show that ~~CTH~~CTH continues to increase up to 2 km. ~~A transition from pure liquid-water to mixed-phase clouds occurs within the first two hours after passing the MIZ.~~

Cold Air Outbreak (CAO): 1 April 2022




Figure 7. (a) Spectral slope phase index (ice index) derived from measurements of the spectrometer of the Munich Aerosol Cloud Scanner (specMACS) on 01 April 2022 as a function of the time the air mass traveled above open ocean. (b) Same as (a) but with cloud top height (*CTH*) from stereographic reconstruction. (c) to (e) Example RGB (red-green-blue) images at points indicated by colored crosses in panels (a) and (b).

385 ~~But how much liquid and ice water is contained in CAOs? von Lerber et al. (2022) noted the difficulty of quantifying snowfall in CAOs both from models, which heavily rely on parameterizations, and also in spaceborne radar, as CAO clouds are often within the blind zone of Cloudsat. LWP is~~ *LWP* is a key parameter within the energy and water cycle. Therefore, it is important to understand how *LWP* and its spatial distribution develop during CAOs. However, observing *LWP* is prone to high uncertainties, especially in the Arctic, leading to about a factor of two difference in satellite retrievals between passive microwave and solar radiation retrievals (Lohmann and Neubauer, 2018). ~~Our~~ *LWP* measurements from P5 and HALO offer opportunities to better constrain satellite observations. Exemplary, Fig. 8b illustrates the flight path of HALO a shows the time series of radar reflectivity measured along the flight track of RF08 (Fig. 8b) on 21 March 2022 (RF08) when, with high reaching clouds belonging to the Shapiro-Keyser cyclone over Svalbard (Shapiro and Keyser, 1990). During this flight, HALO probed a CAO in its initial state close to the MIZ flying parallel to the sea ice edge, and perpendicular to developing cloud rolls, repeating the pattern with increased distance from the MIZ and finally straight towards Kiruna. To illustrate the

395 quality of different *LWP* data sets from space and aircraft, we focus on a flight leg where HALO sampled the transition

to cellular convection (Fig. 8c). ~~Cloud~~The cloud radar measurements show a clear rise in ~~CTH~~CTH for this leg and growing cells southwards(~~at later times~~). ~~LWP~~LWP values retrieved from the HAMP instrument (Mech et al., 2014) have maximum values around 300 g m^{-2} within cells, while close to the sea ice edge, maximum values hardly reach 100 g m^{-2} . ~~(Fig. 8d)~~. They clearly resolve the individual cells, which is not possible from spaceborne microwave radiometry due to their coarse resolution.

400 Note, that the HAMP retrieval only includes the cloud contribution, and thus no enhanced values occur in precipitation. This leg was coordinated with the British FAAM aircraft, which focused on in-situ measurements of the cloud and sub-cloud layer along this coordinated track enabling future joint analysis. Due to the time shift between the MODIS (Moderate Resolution Imaging Spectroradiometer) measurement and the HALO flight track, convective cells are shifted between MODIS and HALO measurements. Nevertheless, it becomes clear that MODIS generally shows higher ~~LWP~~LWP values for these warm cloud

405 conditions. Our comprehensive measurements from different platforms will be used to further investigate the reasons for this long-standing problem.

The combination of various remotely sensed measurements with back trajectory calculation for the targeted CAO ~~flight~~flights by P5, P6, and HALO bears strong potential to further investigate CAO cloud development and transitions. Future analysis will include further retrieval development of liquid and ice clouds, e.g., from specMACS, HAMP, and detailed evaluation

410 of satellite data. Most importantly, the comprehensive measurements provide solid reference data to test high-resolution models that are able to resolve the complex circulation involved in CAOs.

4.1.3 Moisture budget during WAIs

An ultimate test of our understanding of the atmospheric water cycle is ~~to be able to~~provided by checking the ability to close the water budget. ~~On 15 March 2022~~For this test, a research flight (RF05 ~~) was on 15 March 2022~~ was specifically dedicated

415 to optimally ~~determining~~determine the moisture budget components (Eq. 2), including their accuracy for a strong WAI event. These data should serve as a critical test for the respective simulations using the ICON (Icosahedral Nonhydrostatic) model. The governing equation for the moisture budget is given by the local change of the integrated water vapor (IWV):

$$\frac{\partial IWV}{\partial t} - \frac{dIWV}{dt} = \underline{E} - \underline{P} - \vec{\nabla} \cdot IVT + \underline{E - P + \epsilon} = -(\underline{ADV + DIV_{mass}}) + \underline{E - P + \epsilon}, \quad (2)$$

with t the time, E the evaporation, P the precipitation rate, IVT the integrated water vapor transport, and ϵ the residual.

420 $\vec{\nabla} \cdot IVT$ describes the divergence of the IVT ; this quantity ~~is~~was derived from the sum of the ~~dynamic mass divergence~~(DIV_{mass}) and the integral of the horizontal moisture advection (ADV) and the dynamic mass divergence (DIV_{mass}).

The flight pattern (Fig. 9a) chosen to assess the moisture budget ~~includes~~included two legs perpendicular to the flow (thick light green lines, cross-flow), and one internal leg (thick blue line). The moisture flux across the two cross-flow flight legs ~~is~~was estimated from the wind and humidity observations provided by dropsondes (large white triangles). From the difference

425 between exported and imported moisture fluxes determined by the dropsonde observations along the cross-flow flight paths, we estimate the internal divergence of moisture flux ($\vec{\nabla} \cdot IVT$) was evaluated as one key component of the atmospheric moisture budget. Along the internal leg, measurements of radar, microwave radiometer, and dropsondes were used to derive precipitation

Cold Air Outbreak (CAO): 21 March 2022




Figure 8. (a) Time series of radar reflectivity along the flight track of RF08 on 21 March 2022 ~~with high-reaching clouds belonging to the Shapiro-Keyser cyclone over Svalbard (Shapiro and Keyser, 1990),~~ times of dropsonde launches are indicated by black-dotted vertical lines, (b) Flight pattern of HALO (white line) and FAAM (green line) during RF08 ~~on-top-of~~ along the MODIS (Moderate Resolution Imaging Spectroradiometer) overpath (Terra image from 10:45-10:50 UTC, MOD02HKM - Level 1B Calibrated Radiances). (c) Zoom into radar measurements from last leg, which was ~~coheated~~ collocated with FAAM (green line). (d) Time series of liquid water path (LWP) retrieved from HALO and MODIS (MOD06 - Cloud Optical Properties, two-channel retrieval using band 7 ($2.1 \mu\text{m}$) and band 6 ($1.6 \mu\text{m}$)) ~~along the zoom of flight track of RF08.~~ Dotted-Dashed lines show the temporal mean.

rate (P), evaporation (E), integrated water vapor (IWV), and its temporal tendency ($\frac{dIWV}{dt}$). The simulations with ICON ~~are~~ were performed in the domain enclosed by the red dashed line in Fig. 9a with a horizontal resolution of 2.4 km.

430 Fig. 9b compares the moisture budget components derived from the HALO observations (full triangles) and ICON simulations (full dots) along the HALO track. The ICON-based and observational estimates ~~derived~~ agree reasonably well in the quantification of moisture tendency due to mass convergence and surface evaporation, while there are discrepancies regarding the temporal tendency of water vapor. A potential explanation might be the substantial dissipation of the WAI during the flight. Future work will focus on understanding the causes of these discrepancies, closing the moisture budget in the observations, and

435 identifying the major processes for the correct representation of WAIs in models. Here we ~~can also will~~ exploit RF02, RF03, RF04, and RF06 where meteorological conditions and the flight pattern of HALO are well suited to estimate the local moisture tendency and directly compare it with the ICON simulations.

Warm Air Intrusion (WAI): 15 March 2022




Figure 9. (a) HALO flight pattern (gray dashed lines and colored thick lines) and ICON (Icosahedral Nonhydrostatic) model domain (framed by the orange dashed line) for the WAI sampled during RF05 on 15 March 2022. Along the two cross-flow flight sections (thick light green lines), dropsonde observations are indicated by large triangles, whereas small triangles represent the dropsonde releases during the remaining flight sections. The internal leg is indicated by the blue solid line. For the ICON domain, simulated integrated water vapor transport (IVT) values are illustrated in red color. (b) For the eastern part of the flight pattern, the moisture budget components from the HALO observations (triangles) are compared to those simulated by ICON (circles), with respect to their contribution to the moisture budget in mm h^{-1} . Moisture advection (ADV), mass divergence (DIV_{mass}), precipitation rate (P), evaporation (E), integrated water vapor (IWV), and its temporal tendency ($d(IWV)/dt$) are compared. Vertical lines indicate the uncertainties for each component.

4.2 Arctic clouds

4.2.1 Arctic low-level Low-level clouds: Thermodynamic phase distribution

440 ~~The In addition to the spectral slope phase index derived from remote sensing measurements (Fig. 7a), we have determined the thermodynamic phase of the clouds is determined from the microphysical cloud properties derived from the from in-situ measured~~ particle size distribution data in the size range from $2.8 \mu\text{m}$ to $6.4 \mu\text{m}$ (Moser et al., 2023). The ~~measured resulting~~

fractions of ice, mixed-phase, and liquid water clouds are shown as a function of altitude in Fig. 10, classified into cloud measurements over open ocean and ~~over~~ sea ice. Due to the decrease of the temperature with altitude, the fraction of ice clouds over ~~the open~~ ocean increases with altitude for altitudes larger than about 500 m. Over sea ice, the ABL shows a high fraction of ~~pure~~ ice and liquid water. In contrast, the cloud characteristics over the ~~open~~ ocean are more variable in height, as mixed-phase ~~clouds~~ and pure liquid ~~water~~ clouds are detected over the whole altitude range.

These results are obtained from cloud measurements conducted in different meteorological conditions, including CAOs, convergence lines, and polar lows. ~~For a more precise statistical analysis of the cloud properties collected over the ocean, a normalized cloud structure is beneficial. Near future~~ Future studies will evaluate microphysical properties, including the total number concentration ~~;~~ ~~cloud and~~ particle effective diameter ~~;~~ ~~and of the cloud droplets, as well as~~ the cloud water content, ~~and relate the data to environmental~~. ~~These data will be investigated and discussed in relation to the prevailing synoptic conditions. The results of further statistical thermodynamic and microphysical analyses on of low-level Arctic clouds measured cloud measurements obtained during HALO-(AC)³ will be compared with a previous data set collected in a similar season and checked for representativeness.~~ previous data acquired in the past similar seasons and synoptic situations. Furthermore, ~~the method to detect thermodynamic phase in Arctic mixed-phase clouds with in-situ particle measurements as describes in Moser et al. (2023) will be used to validate existing remote sensing algorithms, such as that of Shupe et al. (2008).~~




Figure 10. Fraction of detected cloud particle types resolved by altitude. Cloud types (ice: ice clouds; mpc: mixed-phase clouds; liquid: liquid clouds), shown for (a) clouds over the open ocean, and (c) clouds over the sea ice. (b) and (d) show the distribution of measurements in altitude. Thermodynamic phase classification was performed according to the algorithm presented by Moser et al. (2023).

To understand the conditions and feedback mechanisms, that maintain the persistence of the inherently unstable mixture of super-cooled liquid water cloud droplets and ice crystals ~~of mixed-phase clouds~~, a three-dimensional characterization of the thermodynamic phase partitioning ~~in the clouds~~ is required. For this purpose, the radar-lidar retrieval framework VarCloud was used to ~~retrieve derive~~ ice cloud microphysical properties of mixed-phase clouds (Aubry et al., 2024). An example for the simultaneous retrieval of cloud ice and liquid water microphysical properties is given in Fig 11, which shows the cloud particle (liquid water droplets and ice crystals) effective radius ~~retrieved obtained~~ from combined radar-lidar measurements collected

during the HALO RF06 on 16 March 2022. The [figure-plot](#) shows the cloud cross-section within a decaying WAI. On top of the leading marine stratocumulus deck ([around 10:30 UTC](#)) and embedded in the trailing ice cloud layer ([between 10:15 and 10:20 UTC](#)), the combination of strong lidar with unremarkable cloud radar returns indicates the presence of layers of super-cooled water. These two regions with embedded super-cooled liquid water layers and liquid water-topped ice clouds represent two distinct types of super-cooled liquid water. While the long-lived nature of the latter is well understood, the presence of super-cooled layers embedded within deep ice clouds requires further investigations that are planned for future work.




Figure 11. Simultaneous [retrieval-retrievals](#) of effective radius of cloud ice crystals ($r_{e,ice}$) and liquid water droplets ($r_{e,liq}$) using radar-lidar measurements from the HALO research flight RF06 performed on 16 March 2022.

470 4.2.2 [Arctic-cirrusCirrus: The impact-Impact of surface ,properties and water vapor, and aerosol](#)

The highly reflecting sea ice surface [strongly](#)-modifies the radiative effects of clouds in [general in](#) both the solar and thermal-infrared spectral ranges (Stapf et al., 2020; Becker et al., 2023). [Here we focus on the surface impact on cirrus transmissivity and emissivity.](#) Compared to the open ocean, the high surface albedo and low surface skin temperature of sea ice increase the relevance of surface properties for the cloud radiative effects. Figure 12a shows the brightness temperature field measured by the VELOX (Video airborne Longwave Observations within siX channels) instrument (Schäfer et al., 2022) in a broadband wavelength channel ranging from $7.7 \mu\text{m}$ to $12.0 \mu\text{m}$. For comparison, the time series of the broadband thermal-infrared net irradiance measured by the broadband (solar and thermal-infrared) irradiance sensor called the Broadband AirCrAft RaDiometer Instrumentation (BACARDI) (Ehrlich et al., 2024) ([to be submitted to Earth Syst. Sci. Data](#)) is shown in Fig. 12b. The brightness temperature field shows a tendency to lower values during the first half of this flight section (up to 30 km distance), which is caused by an increased ice water path and [the a](#) reduced emission by the cold cirrus. However, the structure of the sea ice is still imprinted in the measurements (e.g., at 10 km and 20 km distance) indicating the high transmissivity of the cirrus. [The](#)

Beyond 40 km distance, the cirrus is thinning towards the end of the flight section. The warmer areas of open leads and the thin ice of nilas increase and the emitted upward radiation is governed by the surface, which is characterized by a mixture of pack ice and leads with relatively warm open water and young sea ice of a few centimeters thickness (nilas), whose surface is also warmer than the surfaces of pack ice and cirrus. Thus, we see an increase of the emitted upward radiance beyond 40 km in this region. The thermal-infrared net irradiance shows higher values over the cirrus due to a reduced emission compared to the warmer surface. This difference quantifies the top of the atmosphere warming effect of the cirrus, which reaches in this specific case up to about 30 W m^{-2} . However, due to the hemispheric integrating view of BACARDI, the surface variability is not obvious in the broadband irradiance but still might impact the total cirrus radiative effect. Hence, to estimate the total cirrus radiative effect, cirrus and surface inhomogeneities need to be considered.




Figure 12. a) Two-dimensional brightness temperature field measured by VELOX (Video airborne Longwave Observations within six channels) along the flight track in the spectral range of $7.7 \mu\text{m}$ to $12 \mu\text{m}$. b) Time series of the thermal-infrared (TIR) net irradiance, $F_{\text{net,TIR}} = F_{\text{TIR}}^{\downarrow} - F_{\text{TIR}}^{\uparrow}$, measured by the Broadband AirCrAft RaDiometer Instrumentation (BACARDI) for the same flight section.

WAIs lead to an increase of relative humidity and enhance aerosol particle concentrations. Both components can impact the evolution of the cirrus radiative effect. WAIs also often contain enhanced number concentrations of aerosol particles. Both components can impact the evolution of cirrus radiative effects. Therefore, it is important to characterize cirrus clouds in the Arctic and their changes due to increased impact from mid-latitude air masses. Studies on the distribution of relative humidity with respect to ice (RHi) inside and around Arctic and WAIs cirrus cirrus that formed in WAIs have been performed during HALO-(AC)³ using combined aerosol, cloud, and water vapor measurements from the water vapor differential absorption lidar (WALEs) (Wirth et al., 2009), together with temperature information from the model analysis. Particular attention was paid to the vertical distribution of RHi within the cirrus as well as on differences with respect to ice supersaturation giving an estimate of the dominant ice formation processes. From the vertical profiles of cloud properties and RHi we found that cirrus

formed in air masses transported into the Arctic by WAIs have a larger vertical extent compared to cirrus formed in Arctic air masses (Dekoutsidis et al., 2023).

WAI cirrus clouds are characterized by high ice supersaturation throughout their whole-vertical profile. Figure 13 shows an example of the vertical extent and the RHi within and around a WAI cirrus measured during RF03 on 13 March 2022. From the backscatter ratio (upper-left) one can derive Fig. 13a) the vertical extent of the cirrus can be derived. A typical value to distinguish cloud backscatter ratio to distinguish the cirrus from cloud-free pixels is around three (Groß et al., 2014; Urbanek et al., 2017). Applying these values one can see that the WAI cirrus can have this threshold, it becomes obvious from the example shown in Fig. 13a that for this particular WAI cirrus case a vertical extent from about 3 km altitude has been obtained. The cirrus is associated with large enhanced values of RHi (lower-left Fig. 13b). Values of 140 % and larger are reached, and the majority of data points within the cloud shows supersaturation with respect to ice. This becomes even more clearly visible when looking at the combined distribution of backscatter ratio and RHi (Fig. 13, rightc). High values of RHi within the cloud (backscatter ratio > larger than 3) are were found with a peak of the RHi distribution at about 110 %. Even values exceeding the threshold of homogeneous homogenous freezing have been found-identified inside and around the WAI cirrus. This is in accordance with former findings of Gierens et al. (2020) who used radiosonde measurements to study cirrus clouds in the Arctic.




Figure 13. Cross-section of (a) backscatter coefficient ratio at 532 nm, and (b) relative humidity with respect to ice (RHi) for RF03 on 13 March 2022 during a WAI. (c) shows a histogram of joint occurrence of RHi and backscatter coefficient ratio at 532 nm. The relative humidity was calculated from WALES water vapor measurements and model temperature field.

4.3 Arctic aerosol particles

Sources, abundance, and properties of Arctic aerosol particles in general, and cloud condensation nuclei (CCN) in particular, are not comprehensively monitored. Therefore, aerosol measurements were performed during the HALO-($\mathcal{A}\mathcal{C}$)³ aircraft campaign using data obtained from in-situ aerosol instrumentation installed on board of P6. Particles were sampled behind a well-
 520 characterized aerosol inlet (Leaith et al., 2016) and a Counterflow Virtual Impactor (CVI) (Ogren et al., 1985), showing comparable sampling characteristics when both were operated as an aerosol inlet (Ehrlich et al., 2019). Among others, in-situ black carbon (BC) measurements were performed by a single particle soot photometer (SP2) installed behind the CVI. Further details on the applied instrumentation are given by Ehrlich et al. (2024) ~~-(to be submitted to *Earth Syst. Sci. Data*).~~

We have derived typical values of microphysical aerosol properties measured during HALO-($\mathcal{A}\mathcal{C}$)³. ~~We-Here we~~ present
 525 averaged data from the research flights RF08–RF13 of P6 (1–10 April 2022), including periods over open ocean and sea ice. The analysis ~~shows has shown~~ a median total aerosol particle number concentration N_{total} of 303 cm^{-3} (inter-quartile range(IQR)-: ~~IQR =~~ $207\text{--}419 \text{ cm}^{-3}$; Figs. 14a and 14d), and a median CCN number concentration N_{CCN} (measured at 0.1 % supersaturation) of 155 cm^{-3} (IQR ~~:=~~ $81\text{--}204 \text{ cm}^{-3}$; Figs. 14b and 14e). No obvious change of ~~partiele-number-concentration with-altitude-is-~~ N_{total} ~~with altitude became~~ evident up to roughly 1000 m ~~above-sea-levelaltitude~~, and only a slight decrease
 530 with height can be observed above. Average CCN hygroscopicity (κ measured at 0.1 % supersaturation, Figs. 14c and 14f) is in the range typical for mostly inorganic aerosol ~~particles~~ mixed with organic material (median ~~values of~~ κ : 0.50; IQR: 0.40–0.68), featuring slightly higher values above 1000 m.




Figure 14. Relative occurrence of (a) total aerosol particle number concentration N_{total} , (b) Cloud condensation nuclei (CCN) number concentration N_{CCN} , and (c) CCN hygroscopicity κ . Vertical-binned averages of (d) N_{total} , (e) N_{CCN} , and (f) κ . P6 research flights RF08 to RF13 are considered.

Furthermore, we have ~~measured-collected and analyzed~~ cloud droplet residuals (CDR) ~~from in-cloud-sampling~~ using the CVI installed on board of P6 (Ehrlich et al., 2019). We have compared the CDR ~~aerosol-properties-properties (measured inside~~
 535 ~~the clouds) to those of ambient aerosol particles~~ collected in the ABL (below cloud) and in the free troposphere (above cloud). ~~A size-distribution-example-CDR and ambient aerosol particle number size distributions~~ representative for the HALO-(\mathcal{A})³ conditions ~~for low-clouds-in case of low-level clouds probed~~ over open ocean ~~is-are~~ shown in Fig. 15. Due to the identical shape of the CDR and particle distributions below cloud, it is concluded that the cloud is formed and sustained by the activation of ABL particles at the cloud base. A major influence of cloud-forming particles entrained from the free troposphere above the
 540 cloud can be excluded since the respective size distribution appears different. ~~This finding is-~~ ~~which appears~~ similar to the ACLOUD results over open ocean (Wendisch et al., 2019). However, the absence of clouds over sea ice during HALO-(\mathcal{A})³ did not allow a comparison with ACLOUD results, where entrainment of cloud-forming particles from the free troposphere was suggested. This analysis will be continued to look for dependencies on the distance to the sea ice edge, the chemical composition of CDR and out of cloud particles, and changing meteorological conditions during the HALO-(\mathcal{A})³ campaign.




Figure 15. Normalized ~~particle~~ number size ~~distribution-distributions~~ of cloud droplet residuals (CDR, ~~measured inside cloud, blue line~~) and ~~of~~ ambient aerosol particles ~~measured inside- (blue)above clouds-green line,~~ below ~~(red)clouds-red line~~ and ~~above (green)-clouds-~~, as ~~measured~~ over open ocean during research flight RF01 with P6 (20 March 2022). ~~The distributions have been normalized with respect to the integrated (total) CDR or ambient particle number concentration.~~

545 We also look at the chemical composition of ~~aerosol-particles and cloud residuals-and~~ ~~the CDR and ambient particles and~~ their variation with season. For this purpose, we compare observations collected during ACLOUD (late spring/early summer) (Wendisch et al., 2019) with ~~our-the~~ data available from HALO-(\mathcal{A})³ (late winter/early spring). The single particle mass spectrometer ALABAMA (Brands et al., 2011; Clemen et al., 2020) was used to ~~measure-quantify~~ the chemical composition and size of single particles in the size range of 0.23–3 μm (50 % cutoff diameter with respect to detection efficiency). The
 550 ALABAMA was connected to the CVI inlet allowing for sampling of CDR and ambient aerosol particles, dependent on the counter-flow settings. Simultaneous measurements of trace gases such as CO, CO₂, and O₃ were used to identify different air mass origins, e.g., ~~to distinguish polluted from non-polluted air masses.~~ The results ~~(Fig-16)~~ show a large abundance of

particulate amines in ambient air and a dominance of amines in CDR during late spring/early summer (ACLOUD), emphasizing the importance of marine biogenic sources for summertime Arctic cloud processes ~~-(Fig. 16)~~. In contrast, amine-containing particles were rarely observed during late winter/early spring (HALO-(AC)³).

Furthermore, the abundance of elemental carbon (EC) containing particles was proven to be higher during HALO-(AC)³, accompanied by higher CO mixing ratios than during ACLOUD, indicating the presence of anthropogenic pollution typical for the ‘Arctic Haze’ season in spring. These differences between the data obtained during ACLOUD and HALO-(AC)³ reflect the general ~~observation that ACLOUD was~~ conclusion that ACLOUD measurements were dominated by WAIs, while CAOs predominated during the HALO-(AC)³ measurement flights of P6. The composition of CDR is similar to that of the particles in ambient air during HALO-(AC)³, but with a higher contribution of fresh and aged sea salt particles to the CDR, while in summer the amine-containing particles dominate the CDR. In future work, a detailed analysis of different air mass situations (WAIs versus CAOs) combined with air mass history analysis (e.g., air mass trajectories) will be ~~included~~ performed to investigate the sources of the identified particle types.




Figure 16. Number fraction of different particle types for ambient particles and cloud droplet residuals (CDR) during the aircraft missions ACLOUD and HALO-(AC)³ analyzed by the ALABAMA. amm.sulf.: ammonium sulfate, DCA: dicarboxylic acids, van.: Vanadium, EC: elemental carbon. Particle types are assigned by the dominating peaks in the mass spectra.

565 4.4 Proof of concept to measure mesoscale divergence and subsidence in the Arctic

ABL cloud transformations at high latitudes play a key role in the Arctic and are partially controlled by large-scale dynamics such as divergence and subsidence. During several research flights (RFs) of HALO-(AC)³, we have successfully applied a measurement technique using the data from multiple dropsonde releases in circular or quadratic flight patterns to estimate

mesoscale properties including divergence, associated subsidence, pressure gradients, and advective tendencies (Paulus et al., 2024). As illustrated in Fig. 17a, HALO RF10 and RF11 probed a weak CAO ~~in-applying~~ a quasi-Lagrangian ~~way, sampling it at multiple points~~ approach. The air mass was sampled at multiple locations during two days along its southbound trajectory from the central Arctic into the Fram Strait. The four selected ~~locations~~ places were determined before take-off, using trajectory estimates based on forecast data. Mesoscale flight patterns surrounding these locations were then incorporated in the HALO flight plan, in principle allowing the calculation of mesoscale gradients across the area (Bony and Stevens, 2019).

The scientific objective of this effort is twofold. Firstly, we aim to test the hypothesis that this novel sampling technique (the sondes) can reliably yield mesoscale profiles of (thermo-)dynamic circulation properties also at high latitudes, in particular in cold, transforming low level air masses. Various studies have reported encouraging results with this method in marine subtropical areas (George et al., 2021; Bony and Stevens, 2019) and the mid-latitudes (Li et al., 2022). We show that the method works equally well at high latitudes, given the absence of large-scale weak temperature gradients (Charney, 1963; Sobel et al., 2001) and the (partially) associated high transience in synoptic weather. The second objective is to achieve a dataset of circulation properties sampled by HALO at mesoscales that is suitable for driving scientific process models of clouds in CAOs along the trajectory. While CAO cases for Large-Eddy Simulations (LES) and Single-Column Models (SCM) have been generated before (de Roode et al., 2019), typically the observational data to realistically constrain the simulation in the upstream areas was lacking. Previous LES studies in the Arctic have shown a strong dependence of Arctic mixed layers on larger-scale forcings, in particular subsidence (Neggers et al., 2019; Dimitrelos et al., 2023), prioritizing the need for such observational data. The HALO-(AC)³ campaign supplies this crucial data for the first time, a unique opportunity for exclusively forcing scientific process model experiments for this weather regime with mesoscale data sampled along the trajectory. This has not been achieved before and would represent a step forward in anchoring Arctic process model studies in reality.

Figures 17b-e show the vertical profiles of subsidence as derived from dropsonde data from four “mesoscale circle” flight segments that were flown at selected locations along the two-day trajectory. The northernmost circle was sampled by HALO RF10, while the southern three circles were flown a day later by RF11. The third HALO circle on the trajectory (RF11 C02) located just south of the sea ice edge was also probed in collocation by the P5 and P6 aircraft, providing additional in-situ measurements of clouds, turbulence, radiation, and aerosol properties. We obtain robust profiles of mesoscale subsidence at all four circle sites. For reference the measured profiles are cross-compared with ERA5 reanalysis data, reporting less-than-optimal agreement in general. At circle RF11 C02 the subsidence profile sampled by HALO is reproduced by independent dropsonde data from the P5 aircraft, conducted an hour later in the same area and covering only the lowest few kilometers below the P5 aircraft. This agreement, in particular, suggests that the HALO profiles of subsidence are realistic, and provides proof of principle for the applicability of the mesoscale dropsonde technique also at high latitudes.

The encouraging results obtained for this CAO case are fully described by Paulus et al. (2024). They motivate taking the next step and configure LES and SCM experiments for this case that are purely based on HALO measurements. Apart from divergence and associated subsidence, the data also yields profiles of horizontal pressure gradients (or geostrophic wind) and advective tendencies of temperature, humidity, and momentum. The in-situ and independent P5 and P6 data can well be used

to evaluate the process model simulations, yielding a complete package for realistic model studies of this CAO. This research
 605 is currently in progress.




Figure 17. (a) Overview of mesoscale flight patterns flown by HALO (dark-blue-green circles) and P5 (green-red square pattern) sampling a low-level-low-level air mass on 29 and 30 March 2022 at four locations along its southbound trajectory (light-blue-yellow lines). Dropsonde launch locations are marked by crosses (green if dropped by HALO, red crosses for sondes dropped by P5). (b)–(e) Profiles of pressure velocity Ω calculated from dropsonde data (colored time of first and last dropsonde launch indicated) released from HALO and P5 (solid lines) and ERA5 reanalysis data (grey-dashed).

5 Summary and outlook

In this paper, we illustrate the application of a quasi-Lagrangian aircraft measurement approach to observe air mass transformations in the Arctic. The data were collected during the HALO–(AC)³ aircraft campaign that took place between Scandinavia and the North Pole in March and April 2022. We have mainly employed three research aircraft with distinct objectives: The
 610 low-flying (mainly within and below clouds) Polar 6 (P6) was equipped with in-situ instrumentation to measure aerosol, cloud, and precipitation properties, whereas the higher-flying (mostly above clouds) Polar 5 (P5) aircraft conducted remote sensing observations of water vapor, aerosol particle, cloud, and precipitation characteristics. The third research aircraft, the High Altitude and Long Range Research Aircraft (HALO) was flying in about 10 km altitude. Whenever possible, HALO was flying closely collocated with P5 and P6. The in-situ and remote sensing payloads of the three aircraft were complemented by
 615 numerous dropsonde launches from P5 and HALO.

The focus of the observations was on air mass transformations during moist and warm air intrusions (WAIs) and marine cold air outbreaks (CAOs). The processes during these transformations were observed in a novel quasi-Lagrangian manner, probing the same air mass-parcel twice on its way into or out of the Arctic, either on the same flight (e.g., at its beginning and end) or on two flights during consecutive days. To plan such flights, dedicated air mass-parcel trajectory simulations along the

620 flight paths were employed. A substantial number of matches of the same air ~~mass~~ parcels was obtained enabling statistical analysis of the temporal tendencies of potential temperature and moisture during WAIs and CAOs. For the CAO cases, a strong surface-driven diabatic heating between $1\text{--}3\text{ K h}^{-1}$ and a near-surface moistening between $0.05\text{--}0.3\text{ g kg}^{-1}\text{ h}^{-1}$ were observed ~~within the lowest about 0.5~~close to the ground below about 1 km. For the WAI observations, a weak diabatic cooling of up to 0.4 K h^{-1} and a moisture loss of up to $0.1\text{ g kg}^{-1}\text{ h}^{-1}$ were obtained from the ground to about ~~5.5~~5.5 km altitude.

625 ~~Furthermore, we~~We followed the evolution of cloud properties (cloud top height and cover, liquid water path, precipitation rate, thermodynamic phase, and ~~radar-cloud~~cloud reflectivity) during CAOs. We have shown that for a strong CAO case, cloud tops are higher, and more liquid water-topped clouds exist. The liquid water path and mean ~~radar-reflectivity~~cloud reflectivity measured by radar increase compared to a case of a weaker CAO. In addition, we see how the cloud parameters evolved with distance over the open sea, with the ~~ABL-deepening and CTH-~~atmospheric boundary layer deepening and cloud top height
630 rising. We observed that for the stronger CAO case, the characteristic features such as the formation of cloud streets and the onset of precipitation occur closer to the sea ice edge.

~~Furthermore, the~~The moisture budget of a WAIs was quantified for the case of a strong WAI. ICON-based and observational estimates of the moisture budget components agree reasonably well for the ~~moisture-tendency~~temporal tendency of moisture due to mass convergence and surface evaporation, while there are remarkable discrepancies regarding the local tendency of
635 water vapor in ICON compared to the HALO data.

The vertical distribution of the thermodynamic phase in low-level Arctic clouds was quantified in a statistical manner using in-situ cloud measurements carried out over the open ocean and sea ice. The clouds over sea ice are dominated by the ice phase. In the upper part of the ~~ABL~~atmospheric boundary layer, liquid clouds are frequently detected. Over sea ice, only a small fraction of the observed clouds is attributed to mixed-phase clouds. In contrast, over the open ocean, the cloud phase
640 distribution is more variable in height, as mixed-phase clouds and pure liquid clouds are detected over the entire altitude range. Some reasons for the longevity of mixed-phase clouds were discussed by using a three-dimensional characterization of the thermodynamic phase partitioning.

Typical Arctic aerosol characteristics were quantified, and the chemical composition of the aerosol particles was studied. Values of the median total aerosol particle number concentration of 303 cm^{-3} , and a median ~~CCN~~cloud condensation nuclei
645 number concentration at 0.1 % supersaturation of 155 cm^{-3} were derived. No altitude trend in particle numbers is evident up to roughly 1000 m above sea level, but a decreasing ~~trend~~tendency was sometimes observed higher up. With regard to the chemical composition of the aerosol particles, a large abundance of particulate amines in ambient air and a dominance of amines in cloud droplet residuals (~~CDR~~) during late spring/early summer is shown, emphasizing the importance of marine biogenic sources for summertime Arctic cloud processes. In contrast, amine-containing particles were rarely observed during
650 late winter/early spring.

It was shown that circular or quadratic flight patterns with sufficiently frequent dropsonde releases provide appropriate data to estimate mesoscale gradients, which can be used to derive subsidence and advective tendencies in the Arctic. These data are highly valuable, for example, to make available data to constrain the initial conditions for large eddy simulations avoiding the use of numerical models with a coarser resolution.

655 ~~Delivering data for evaluating and testing numerical models and reanalysis is a major objective of the HALO-(AC)³ campaign. This objective will further be pursued using the wealth of observations from the campaign.~~

Analysis of the results of the HALO-(AC)³ aircraft campaign is ongoing, and resulting papers will be published in a Special Issue of Atmospheric Chemistry and Physics (ACP) on "HALO-(AC)³ – an airborne campaign to study air mass transformations during warm-air intrusions and cold-air outbreaks", see: https://acp.copernicus.org/articles/special_issue1272.html and other peer-reviewed scientific journals. Three specific examples of topics that are currently being worked on are given below:

- 665 – We will investigate the hypothesis that a secondary circulation acts to spread out moisture from the initial river-like intrusion in the cross-flow direction. This hypothesized circulation consists of uplift in the core of the intrusions, divergence in the upper troposphere above the core, and subsidence of drier air on the flanks of the intrusion. We will investigate the typical structure of the atmospheric moisture field, how it evolves during a WAIs, and what are the governing drivers of these processes.
- 670 – The particular mode structure of the Arctic radiant energy budget (REB) will be analyzed as a function of altitude, surface type (sea ice or open ocean), cloudy or cloud-free conditions, and thermodynamic properties (temperature lapse rate, horizontal temperature gradient between sea ice and open ocean). The observations will be confronted with model results that often struggle to correctly represent the mode structure of the REB, e.g., due to limitations in the treatment of sub-grid processes including clouds and the sea ice albedo. Therefore, detailed observation-model comparisons are envisioned to identify potential misrepresentations of properties affecting the REB.
- 675 – We will integrate measurements obtained during two CAO events observed during HALO-(AC)³ (29 March and 9 April 2022) with literature data. The 9 April case was extraordinary for two reasons. Firstly, it was characterized by cloud-free conditions along the entire 180 km North-South flight track across the MIZ to the open ocean, which is very rare and unusual. Only at the southernmost position, convective clouds appeared. Secondly, the southernmost position was influenced by the front of a polar low, which was a remnant of a polar low over Fram Strait on the preceding day. We will discuss how the results of our observations on these two days fit to earlier measurements.

680 Evaluating and testing numerical weather and climate models as well as reanalysis products using the comprehensive measurements conducted during the HALO-(AC)³ campaign is the second major objective of this project. This goal will further be pursued in future studies using the wealth of available observations from the campaign.

Data availability. All data collected during the HALO-(AC)³ aircraft campaign are being published by Ehrlich et al. (2024) (to be submitted to *Earth Syst. Sci. Data*).

685 *Author contributions.* All authors provided text and figures and contributed to the editing of the article, and to the analysis and discussion of the results.

Competing interests. One of the co-authors (Birgit Wehner) is a member of the editorial board of ACP.

Acknowledgements. We gratefully acknowledge the funding by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project Number 268020496 – TRR 172, within the framework of the Transregional Collaborative Research Center “Arctic Amplification: Climate Relevant Atmospheric and Surface Processes, and Feedback Mechanisms (AC)³”. The authors are grateful to AWI for providing and operating the two Polar 5 and Polar 6 aircraft. We thank the crews and the technicians of the three research aircraft for excellent technical and logistical support. The generous funding of the flight hours for the Polar 5 and Polar 6 aircraft by AWI, and for HALO by DFG, Max-Planck-Institut für Meteorologie (MPIM), and Deutsches Zentrum für Luft- und Raumfahrt (DLR) is greatly appreciated. We are further grateful for funding of project grant number 316646266 by DFG within the framework of Priority Program SPP 1294 to promote research with HALO. Oliver Eppers, Philipp Joppe, and Johanna Mayer acknowledge funding by the DFG – TRR 301 – Project-ID 428312742. Hans-Christian Clemen acknowledges funding by the DFG, project No. 442647984 (SCHN 1138/8-1). Timo Vihma acknowledges funding from the European Commission Horizon 2020 project Polar Regions in the Earth System project (PolarRES, 101003590). Gunilla Svensson and Michail Karalis acknowledge funding from the Swedish Research Council (VR, project 2020-04064). Manuel Moser is grateful for funding by DFG SPP 2115 PROM (VO 1504/5-1), Johannes Lucke appreciates funding within the EU project SENS4ICE (grant agreement number 824253), and Elena De La Torre Castro acknowledges support by DFG SPP HALO 1294 (VO 1504/7-1). Harald Sodemann acknowledges funding from the European Research Council (Consolidator Grant Nr. 773245). This publication was supported by the Open Access Publishing Fund of Leipzig University.

Appendix A: A hypothesis on humidity fields in WAIs

~~We investigate the typical structure of the atmospheric moisture field, how it typically evolves during a WAI, and what processes drive this development. Relative humidity varies strongly and determines the structure of the moisture field in the cross-flow perspective going from 100% in the core of the WAI to around 50% within no more than 100 km distance from the core. Temperature determines the structure of the moisture field in an along-flow direction, as relative humidity is at saturation throughout the core of the intrusion. In the ABL and close to the sea ice edge, temperatures change by 3–4 K over a distance of about 150 km. Clearly an atmospheric river is channeled in the Fram Strait between sea ice (20°W) and Svalbard (0°). In future work we will investigate the hypothesis that a secondary circulation acts to spread out moisture from the initial river-like intrusion in the cross-flow direction (Fig. ??). This hypothesized circulation consists of uplift in the core of the intrusions, divergence in the upper troposphere above the core, and convergence and subsidence of drier air on the flanks of the intrusion.~~

~~Zonal-vertical cross-section of relative humidity with respect to liquid water during a moist and warm air intrusion (WAI) on the basis of ERA5 data. Arrows indicate hypothesized circulation.~~

Appendix B: The particular mode structure of the Arctic radiant energy budget

715 The radiant energy budget (REB) in the Arctic depends on different surface, cloud, and thermodynamic properties (Sedlar et al., 2011; Stapf
720 At the surface, Wendisch et al. (2023b) identified four modes (regimes) of the REB, depending on the surface type (sea ice or
open ocean) and the atmospheric state (cloudy or cloud-free conditions). Models often struggle to correctly represent this mode
structure due to limitations in the treatment of sub-grid processes including clouds and the sea ice albedo (Kretzschmar et al., 2020; Solomon
725 Consequently, detailed observation-model comparisons are helpful to identify potential misrepresentations of properties
affecting the REB.

During HALO-(AC)³, the coordinated combination of measurements from three aircraft operating in different altitude
regimes allowed to extend the statistical analysis of the Arctic REB to higher altitudes. Beside REB measurements performed
by P6 close to the surface, the REB above the ABL clouds and close to the tropopause was observed by P5 and HALO,
respectively (Fig. ??). The common mode structure (cloudy and cloud-free, over sea ice and open ocean) is most prominently
730 illustrated in the near-surface REB (modes indicated by 3, 4, and 5, Fig. ??c). Modes 4 and 5 result from the observations in
cloudy and cloud-free conditions above P6 when flying over the open ocean. Mode 3 corresponds to cloud-free situations over
sea ice (only few clouds were observed over sea ice). For P5 (mid-level regime, Fig. ??b), clouds were not present above the
aircraft, leaving a combined mode for weaker emitting cases above sea ice or low clouds (mode 1) and a mode for cloud-free
observations above the open ocean (mode 2). The REB in high altitudes shows a continuous transition between modes 1 and
735 2. In general, the thermal-infrared net irradiance becomes more negative with increasing altitude. Furthermore, the surface and
top-of-atmosphere (TOA) REBs obtained from ERA5 along the flight tracks are compared to the P6 and HALO measurements
(not shown). The near-surface thermal-infrared REB is less negative than in the P6 observations. The TOA REB in ERA5 is
slightly shifted towards lower solar net irradiances and is more narrow than the distribution of the HALO observations due to
the lower horizontal ERA5 resolution. Such comparisons will help to constrain the REB in numerical models which aim to
740 reproduce HALO-(AC)³ observations.

Two-dimensional probability density function of the solar (normalized by the solar downward irradiance) and thermal-infrared
net irradiances observed in different altitudes: (a) HALO, (b) P5, and (c) P6 flight altitudes. Mode numbers: 1—above clouds
or sea ice, 2—above open ocean, 3—cloud-free over sea ice, 4—cloud-free over open ocean, 5—cloudy over open ocean.

Appendix C: An extraordinary CAO case covered during HALO-(AC)³

740 HALO-(AC)³ has provided an opportunity to expand the limited data base of airborne CAO studies that are known from the
literature (Brümmer, 1997, 1999; Chechin et al., 2013; Dahlke et al., 2022; Geerts et al., 2022; Schirmacher et al., 2024). Here
we integrate our data obtained during two CAO events with measurements following approximately a North-South trajectory
(29 March and 9 April 2022) with literature data. The 9 April case was extraordinary for two reasons. Firstly, it was characterized
by cloud-free conditions along the entire 180 km North-South flight track across the MIZ to the open ocean, which is very
745 rare and unusual. Only at the southernmost position, convective clouds appeared. There and over sea ice, vertical profile
measurements were performed with P6 (RF12). Secondly, the southernmost position was influenced by the front of a polar

low, which was a remnant of a polar low over Fram Strait on the preceding day. Figure ?? shows profiles of mean and turbulent thermodynamic quantities measured over open sea during the two HALO-(AC)³ flights in comparison to previous measurements. For 29 March 2022, the results fit well to the earlier measurements except some peaks in the momentum fluxes, which could be traced back to orographic influences of Svalbard mountains at the northern coast. However, on 9 April (the cloudless day), unusually high temperatures and untypically low wind speeds were observed leading to extraordinarily weak turbulent convection over open sea when compared to previous CAO events.

Comparison of atmospheric boundary layer (ABL) profiles, with z representing altitude, of mean and turbulent quantities (normalized by the ABL height z_1) obtained by airborne measurements during CAOs on 29 March (orange) and 9 April (red) 2022 during HALO-(AC)³. Data from previous aircraft campaigns (ARTIST, Hartmann et al. (1999), and AFLUX, Mech et al. (2022)) are indicated by black lines with different symbols. TKE is the turbulent kinetic energy, and $\overline{w'^2}$ represents the vertical velocity variance.

References

- Ali, S. M. and Pithan, F.: Following moist intrusions into the Arctic using SHEBA observations in a Lagrangian perspective, *Q. J. R. Meteorol. Soc.*, 146, 3522–3533, <https://doi.org/10.1002/qj.3859>, 2020.
- Alvarez, J., Yumashev, D., and Whiteman, G.: A framework for assessing the economic impacts of Arctic change, *Ambio*, 49, 407–418, <https://doi.org/10.1007/s13280-019-01211-z>, 2020.
- Aubry, C., Delanoë, J., Groß, S., Ewald, F., Tridon, F., Jourdan, O., and Mioche, G.: Lidar-radar synergistic method to retrieve ice, supercooled and mixed-phase clouds properties, *Atmos. Meas. Technol. Discuss.*, <https://doi.org/10.5194/amt-2023-252>, 2024.
- 765 Bauer, R., Grooß, J.-U., Ungermann, J., Bär, M., Geldenhuys, M., and Hoffmann, L.: The Mission Support System (MSS v7.0.4) and its use in planning for the SouthTRAC aircraft campaign, *Geosci. Model Dev.*, 15, 8983–8997, <https://doi.org/10.5194/gmd-15-8983-2022>, 2022.
- Becker, S., Ehrlich, A., Schäfer, M., and Wendisch, M.: Airborne observations of the surface cloud radiative effect during different seasons over sea ice and open ocean in the Fram Strait, *Atmos. Chem. Phys.*, 23, 7015–7031, <https://doi.org/10.5194/acp-23-7015-2023>, 2023.
- 770 Beer, C., Zimov, N., Olofsson, J., Porada, P., and Zimov, S.: Protection of Permafrost Soils from Thawing by Increasing Herbivore Density, *Sci. Rep.*, 10, 4170, <https://doi.org/10.1038/s41598-020-60938-y>, 2020.
- Blackport, R. and Screen, J. A.: Insignificant effect of Arctic amplification on the amplitude of midlatitude atmospheric waves, *Sci. Adv.*, 6, 2880, <https://doi.org/10.1126/sciadv.aay2880>, 2020.
- Block, K., Schneider, F., Mülmenstädt, J., Salzmann, M., and Quaas, J.: Climate models disagree on the sign of total radiative feedback in the Arctic, *Tellus A: Dyn. Meteorol. Oceanogr.*, 72:1, 1–14, <https://doi.org/10.1080/16000870.2019.1696139>, 2020.
- 775 Boettcher, M., Schäfler, A., Sprenger, M., Sodemann, H., Kaufmann, S., Voigt, C., Schlager, H., Summa, D., Di Girolamo, P., Nerini, D., Germann, U., and Wernli, H.: Lagrangian matches between observations from aircraft, lidar and radar in a warm conveyor belt crossing orography, *Atmos. Chem. Phys.*, 21, 5477–5498, <https://doi.org/10.5194/acp-21-5477-2021>, 2021.
- Bony, S. and Stevens, B.: Measuring Area-Averaged Vertical Motions with Dropsondes, *J. Atmos. Sci.*, 76, 767–783, <https://doi.org/10.1175/JAS-D-18-0141.1>, 2019.
- 780 Bossioli, E., Sotiropoulou, G., Methymaki, G., and Tombrou, M.: Modeling extreme warm-air advection in the Arctic during summer: The effect of mid-latitude pollution inflow on cloud properties, *J. Geophys. Res.*, 126, <https://doi.org/10.1029/2020JD033291>, 2021.
- Brands, M., Kamphus, M., Böttger, T., Schneider, J., Drewnick, F., Roth, A., Curtius, J., Voigt, C., Borbon, A., Beekmann, M., Bourdon, A., Perrin, T., and Borrmann, S.: Characterization of a Newly Developed Aircraft-Based Laser Ablation Aerosol Mass Spectrometer (ALABAMA) and First Field Deployment in Urban Pollution Plumes over Paris During MEGAPOLI 2009, *Aerosol Sci. Technol.*, 45, 46–64, <https://doi.org/10.1080/02786826.2010.517813>, 2011.
- Brümmer, B.: Boundary Layer Mass, Water, and Heat Budgets in Wintertime Cold-Air Outbreaks from the Arctic Sea Ice, *Mon. Wea. Rev.*, 125, 1824–1837, [https://doi.org/10.1175/1520-0493\(1997\)125<1824:BLMWAH>2.0.CO;2](https://doi.org/10.1175/1520-0493(1997)125<1824:BLMWAH>2.0.CO;2), 1997.
- Brümmer, B.: Roll and Cell Convection in Wintertime Arctic Cold-Air Outbreaks, *J. Atmos. Sci.*, 56, 2613–2636, [https://doi.org/10.1175/1520-0469\(1999\)056<2613:RACCIW>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<2613:RACCIW>2.0.CO;2), 1999.
- 790 Brümmer, B. and Thiemann, S.: The Atmospheric Boundary Layer In An Arctic Wintertime On-Ice Air Flow, *Boundary-Layer Meteorol.*, 104, 53–72, <https://doi.org/10.1023/A:1015547922104>, 2002.
- Businger, S., Chiswell, S. R., Ulmer, W. C., and Johnson, R.: Balloons as a Lagrangian measurement platform for atmospheric research, *J. Geophys. Res.*, 101, 4363–4376, <https://doi.org/10.1029/95JD00559>, 1996.

- 795 Businger, S., Johnson, R., and Talbot, R.: Scientific Insights from Four Generations of Lagrangian Smart Balloons in Atmospheric Research, *Bull. Amer. Meteorol. Soc.*, 87, 1539–1554, <http://www.jstor.org/stable/26217226>, 2006.
- Charney, J. G.: A note on large-scale motions in the tropics, *J. Atmos. Sci.*, 20, 607–609, [https://doi.org/10.1175/1520-0469\(1963\)020<0607:ANOLSM>2.0.CO;2](https://doi.org/10.1175/1520-0469(1963)020<0607:ANOLSM>2.0.CO;2), 1963.
- Chechin, D. G., Lüpkes, C., Repina, I. A., and Gryanik, V. M.: Idealized dry quasi 2-D mesoscale simulations of cold-air outbreaks over
800 the marginal sea ice zone with fine and coarse resolution, *J. Geophys. Res. Atmos.*, 118, 8787–8813, <https://doi.org/10.1002/jgrd.50679>, 2013.
- Chylek, P., Folland, C., Klett, J. D., Wang, M., Hengartner, N., Lesins, G., and Dubey, M. K.: Annual mean Arctic amplification 1970–2020: Observed and simulated by CMIP6 climate models, *Geophys. Res. Lett.*, 49, e2022GL099371, <https://doi.org/10.1029/2022GL099371>, 2022.
- 805 Clemen, H.-C., Schneider, J., Klimach, T., Helleis, F., Köllner, F., Hünig, A., Rubach, F., Mertes, S., Wex, H., Stratmann, F., Welti, A., Kohl, R., Frank, F., and Borrmann, S.: Optimizing the detection, ablation, and ion extraction efficiency of a single-particle laser ablation mass spectrometer for application in environments with low aerosol particle concentrations, *Atmos. Meas. Technol.*, 13, 5923–5953, <https://doi.org/10.5194/amt-13-5923-2020>, 2020.
- Cohen, J., Screen, J. A., Furtado, J. C., Barlow, M., Whittleston, D., Coumou, D., Francis, J. A., Dethloff, K., Entekhabi, D., Overland, J. E.,
810 and Jones, J.: Recent Arctic amplification and extreme mid-latitude weather, *Nat. Geosci.*, 7, 627–637, <https://doi.org/10.1038/ngeo2234>, 2014.
- Cohen, J., Zhang, X., Francis, J., Jung, T., Kwok, R., Overland, J., Ballinger, T., Bhatt, U. S., Chen, H. W., Coumou, D., Feldstein, S., Handorf, D., Henderson, G., Ionita, M., Kretschmer, M., Laliberte, F., Lee, S., Linderholm, H. W., Maslowski, W., Peings, Y., Pfeiffer, K., Rigor, I., Semmler, T., Stroeve, J., Taylor, P. C., Vavrus, S., Vihma, T., Wang, S., Wendisch, M., Wu, Y., and Yoon, J.: Divergent consensuses on
815 Arctic amplification influence on midlatitude severe winter weather, *Nat. Clim. Change*, 10, 20–29, <https://doi.org/10.1038/s41558-019-0662-y>, 2020.
- Cohen, J., Agel, L., Barlow, M., and Entekhabi, D.: No detectable trend in mid-latitude cold extremes during the recent period of Arctic amplification, *Commun. Earth Environ.*, 4, <https://doi.org/10.1038/s43247-023-01008-9>, 2023.
- Dada, L., Angot, H., Beck, I., Baccarini, A., Quéléver, L. L. J., Boyer, M., Laurila, T., Brasseur, Z., Jozef, G., de Boer, G., Shupe, M. D.,
820 Henning, S., Bucci, S., Dütsch, M., Stohl, A., Petäjä, T., Daellenbach, K. R., Jokinen, T., and Schmale, J.: A central Arctic extreme aerosol event triggered by a warm air-mass intrusion, *Nat. Commun.*, 13, <https://doi.org/10.1038/s41467-022-32872-2>, 2022.
- Dahlke, S., Solbès, A., and Maturilli, M.: Cold Air Outbreaks in Fram Strait: Climatology, Trends, and Observations During an Extreme Season in 2020, *J. Geophys. Res. Atmos.*, 127, e2021JD035741, <https://doi.org/10.1029/2021JD035741>, 2022.
- De La Torre Castro, E., Jurkat-Witschas, T., Afchine, A., Grewe, V., Hahn, V., Kirschler, S., Krämer, M., Lucke, J., Spelten, N., Wernli, H.,
825 Zöger, M., and Voigt, C.: Differences in microphysical properties of cirrus at high and mid-latitudes, *Atmos. Chem. Phys.*, 23, 13167–13189, <https://doi.org/10.5194/acp-23-13167-2023>, 2023.
- de Roode, S. R., Frederikse, T., Siebesma, A. P., Ackerman, A. S., Chylik, J., Field, P. R., Fricke, J., Gryschka, M., Hill, A., Honnert, R., Krueger, S. K., Lac, C., Lesage, A. T., and Tomassini, L.: Turbulent Transport in the Gray Zone: A Large Eddy Model Intercomparison Study of the CONSTRAIN Cold Air Outbreak Case, *J. Adv. Model. Earth Syst.*, <https://doi.org/10.1029/2018MS001443>, 2019.
- 830 Dekoutsidis, G., Groß, S., Wirth, M., Krämer, M., and Rolf, C.: Characteristics of supersaturation in midlatitude cirrus clouds and their adjacent cloud-free air, *Atmos. Chem. Phys.*, 23, 3103–3117, <https://doi.org/10.5194/acp-23-3103-2023>, 2023.

- Dimitrelos, A., Ekman, A. M., Caballero, R., and Savre, J.: A sensitivity study of Arctic air-mass transformation using large eddy simulation, *J. Geophys. Res.*, 125, e2019JD031738, <https://doi.org/10.1029/2019JD031738>, 2020.
- 835 Dimitrelos, A., Caballero, R., and Ekman, A. M. L.: Controls on Surface Warming by Winter Arctic Moist Intrusions in Idealized Large-Eddy Simulations, *J. Climate*, 36, 1287–1300, <https://doi.org/10.1175/JCLI-D-22-0174.1>, 2023.
- Druckenmiller, M. L., Thoman, R. L., and Moon, T. A.: Arctic Report Card 2022: Executive Summary, NOAA Arctic Report Card 2021, 22–01, <https://doi.org/10.25923/yjx6-r184>, 2022.
- Ehrlich, A., Bierwirth, E., Wendisch, M., Gayet, J.-F., Mioche, G., Lampert, A., and Heintzenberg, J.: Cloud phase identification of Arctic boundary-layer clouds from airborne spectral reflection measurements: Test of three approaches, *Atmos. Chem. Phys.*, 8, 7493–7505, 840 <https://doi.org/10.5194/acp-8-7493-2008>, 2008.
- Ehrlich, A., Wendisch, M., Lüpkes, C., Buschmann, M., Bozem, H., Chechin, D., Clemen, H.-C., Dupuy, R., Eppers, O., Hartmann, J., Herber, A., Jäkel, E., Järvinen, E., Jourdan, O., Kästner, U., Kliesch, L.-L., Köllner, F., Mech, M., Mertes, S., Neuber, R., Ruiz-Donoso, E., Schnaiter, M., Schneider, J., Stapf, J., and Zanatta, M.: A comprehensive in situ and remote sensing data set from the Arctic CLOUD Observations Using airborne measurements during polar Day (ACLOUD) campaign, *Earth Syst. Sci. Data*, 11, 1853–1881, 845 <https://doi.org/10.5194/essd-11-1853-2019>, 2019.
- Ehrlich, A., Luebke, A. E., Klingebiel, M., Dorff, H., Walbröl, A., Becker, S., Schäfer, M., Rosenburg, S., Müller, H., Kirbus, B., Maahn, M., Mahernndl, N., Ewald, F., Schäfler, A., Pörtge, V., Weber, A., and Wendisch, M.: A comprehensive in situ and remote sensing data set collected during the HALO-(AC)³ airborne campaign, for Submission to *Earth Syst. Sci. Data*, 2024.
- Ewald, F., Kölling, T., Baumgartner, A., Zinner, T., and Mayer, B.: Design and characterization of specMACS, a multipurpose hyperspectral cloud and sky imager, *Atmos. Meas. Technol.*, 9, 2015–2042, <https://doi.org/10.5194/amt-9-2015-2016>, 2016.
- 850 Forster, P. M., Smith, C. J., Walsh, T., Lamb, W. F., Lamboll, R., Hauser, M., Ribes, A., Rosen, D., Gillett, N., Palmer, M. D., Rogelj, J., von Schuckmann, K., Seneviratne, S. I., Trewin, B., Zhang, X., Allen, M., Andrew, R., Birt, A., Borger, A., Boyer, T., Broersma, J. A., Cheng, L., Dentener, F., Friedlingstein, P., Gutiérrez, J. M., Gütschow, J., Hall, B., Ishii, M., Jenkins, S., Lan, X., Lee, J.-Y., Morice, C., Kadow, C., Kennedy, J., Killick, R., Minx, J. C., Naik, V., Peters, G. P., Pirani, A., Pongratz, J., Schleussner, C.-F., Szopa, S., Thorne, P., 855 Rohde, R., Rojas Corradi, M., Schumacher, D., Vose, R., Zickfeld, K., Masson-Delmotte, V., and Zhai, P.: Indicators of Global Climate Change 2022: annual update of large-scale indicators of the state of the climate system and human influence, *Earth Syst. Sci. Data*, 15, 2295–2327, <https://doi.org/10.5194/essd-15-2295-2023>, 2023.
- Francis, J. A. and Vavrus, S. J.: Evidence for a wavier jet stream in response to rapid Arctic warming, *Environ. Res. Lett.*, 10, 014005, <https://doi.org/10.1088/1748-9326/10/1/014005>, 2015.
- 860 Geerts, B., Giangrande, S. E., McFarquhar, G. M., Xue, L., Abel, S. J., Comstock, J. M., Crewell, S., DeMott, P. J., Ebell, K., Field, P., Hill, T. C. J., Hunzinger, A., Jensen, M. P., Johnson, K. L., Juliano, T. W., Kollias, P., Kosovic, B., Lackner, C., Luke, E., Lüpkes, C., Matthews, A. A., Neggers, R., Ovchinnikov, M., Powers, H., Shupe, M. D., Spengler, T., Swanson, B. E., Tjernström, M., Theisen, A. K., Wales, N. A., Wang, Y., Wendisch, M., and Wu, P.: The COMBLE Campaign: A Study of Marine Boundary Layer Clouds in Arctic Cold-Air Outbreaks, *Bull. Amer. Meteorol. Soc.*, 103, E1371 – E1389, <https://doi.org/10.1175/BAMS-D-21-0044.1>, 2022.
- 865 George, G., Stevens, B., Bony, S., Pincus, R., Fairall, C., Schulz, H., Kölling, T., Kalen, Q. T., Klingebiel, M., Konow, H., Lundry, A., Prange, M., and Radtke, J.: JOANNE: Joint dropsonde Observations of the Atmosphere in tropical North Atlantic meso-scale Environments, *Earth Syst. Sci. Data*, 13, 5253–5272, <https://doi.org/10.5194/essd-13-5253-2021>, 2021.
- Gierens, R., Kneifel, S., Shupe, M. D., Ebell, K., Maturilli, M., and Löhnert, U.: Low-level mixed-phase clouds in a complex Arctic environment, *Atmos. Chem. Phys.*, 20, 3459–3481, <https://doi.org/10.5194/acp-20-3459-2020>, 2020.

- 870 Gong, X., Zhang, J., Croft, B., Yang, X., Frey, M. M., Bergner, N., Chang, R. Y.-W., Creamean, J. M., Kuang, C., Martin, R. V., Ranjithkumar, A., Sedlacek, A. J., Uin, J., Willmes, S., Zawadowicz, M. A., Pierce, J. R., Shupe, M. D., Schmale, J., and Wang, J.: Arctic warming by abundant fine sea salt aerosols from blowing snow, *Nat. Geosci.*, 16, 768–774, <https://doi.org/10.1038/s41561-023-01254-8>, 2023.
- Graham, R., Cohen, L., Ritzhaupt, N., Segger, B., Graverson, R., Rinke, A., Walden, V. P., Granskog, M. A., and Hudson, S. R.: Evaluation of six atmospheric reanalyses over Arctic sea ice from winter to early summer, *J. Clim.*, 32, 4121–4143, <https://doi.org/10.1175/JCLI-D-18-0643.1>, 2019a.
- 875 Graham, R. M., Hudson, S. R., and Maturilli, M.: Improved performance of ERA5 in Arctic gateway relative to four global atmospheric reanalyses, *Geophys. Res. Lett.*, 46, 6138–6147, <https://doi.org/10.1029/2019GL082781>, 2019b.
- Graverson, R. G., Mauritsen, T., Tjernström, M., Källén, E., and Svensson, G.: Vertical structure of recent Arctic warming, *Nature*, 451, 53–56, <https://doi.org/10.1038/nature06502>, 2008.
- 880 Groß, S., Wirth, M., Schäfler, A., Fix, A., Kaufmann, S., and Voigt, C.: Potential of airborne lidar measurements for cirrus cloud studies, *Atmos. Meas. Technol.*, 7, 2745–2755, <https://doi.org/10.5194/amt-7-2745-2014>, 2014.
- Gryanik, V. M. and Lüpkes, C.: A Package Momentum and Heat Transfer Coefficients for the Stable Surface Layer Extended by New Coefficients over Sea Ice, *Boundary-Layer Meteorol.*, 187, 41–72, <https://doi.org/10.1007/s10546-022-00730-9>, 2023.
- Hartmann, J., Kottmeier, C., and Raasch, S.: Roll vortices and boundary-layer development during a cold air outbreak, *Bound.-Lay. Meteorol.*, 84, 45–65, <https://doi.org/10.1023/A:1000392931768>, 1997.
- 885 Hartmann, J., Albers, F., Argentini, S., Bochert, A., Bonafé, U., Cohrs, W., Conidi, A., Freese, D., Georgiadis, T., Ippoliti, A., Kaleschke, L., Lüpkes, C., Maixner, U., Mastrantonio, G., Ravegnani, F., Reuter, A., Trivellone, G., and Viola, A.: Arctic Radiation and Turbulence Interaction Study (ARTIST), https://doi.org/10.2312/BzP_0305_1999, 1999.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 890 Holland, M. M. and Landrum, L.: The emergence and transient nature of Arctic amplification in coupled climate models, *Front. Earth Sci.*, 9, <https://doi.org/10.3389/feart.2021.719024>, 2021.
- 895 Holtslag, B., Svensson, G., Baas, P., Basu, S., Beare, R., Beljaars, A., Bosveld, F., Cuxart, J., Lindvall, J., Steeneveld, G.-J., Tjernström, M., and Van de Wiel, B.: Stable Atmospheric Boundary Layers and Diurnal Cycles: Challenges for Weather and Climate Models, *Bull. Amer. Meteorol. Soc.*, 94, 1691–1706, <https://doi.org/10.1175/BAMS-D-11-00187.1>, 2013.
- Huang, Y., Dong, X., Bailey, D. A., Holland, M. M., Xi, B., DuVivier, A. K., Kay, J. E., Landrum, L. L., and Deng, Y.: Thicker clouds and accelerated Arctic sea ice decline: The atmosphere-sea ice interactions in spring, *Geophys. Res. Lett.*, 46, 6980–6989, <https://doi.org/10.1029/2019GL082791>, 2019.
- 900 Jeffries, M. O., Overland, J. E. E., and Perovich, D. K.: The Arctic shifts to a new normal, *Phys. Today*, 66, 35–40, <https://doi.org/10.1063/PT.3.2147>, 2013.
- Johansson, A. M., King, J. A., Doulgeris, A. P., Gerland, S., Singha, S., Spreen, G., and Busche, T.: Combined observations of Arctic sea ice with near-coincident colocated X-band, C-band, and L-band SAR satellite remote sensing and helicopter-borne measurements, *J. Geophys. Res.*, 122, 669–691, <https://doi.org/10.1002/2016jc012273>, 2017a.
- 905

- Johansson, E., Devasthale, A., Tjernström, M., Ekman, A. M. L., and L'Ecuyer, T.: Response of the lower troposphere to moisture intrusions into the Arctic, *Geophys. Res. Lett.*, 44, 2527–2536, <https://doi.org/10.1002/2017GL072687>, 2017b.
- Johnson, R., Businger, S., and Baerman, A.: Lagrangian air mass tracking with smart balloons during ACE-2, *Tellus B: Chem. Phys. Meteorol.*, 52, 321–334, <https://doi.org/10.3402/tellusb.v52i2.16114>, 2000.
- Jung, T., Gordon, N. D., Bauer, P., Bromwich, D. H., Chevallier, M., Day, J. J., Dawson, J., Doblas-Reyes, F., Fairall, C., Goessling, H. F., Holland, M., Inoue, J., Iversen, T., Klebe, S., Lemke, P., Losch, M., Makshtas, A., Mills, B., Nurmi, P., Perovich, D., Reid, P., Renfrew, I. A., Smith, G., Svensson, G., Tolstykh, M., and Yang, Q.: Advancing Polar Prediction Capabilities on Daily to Seasonal Time Scales, *Bull. Am. Meteorol. Soc.*, 97, 1631–1647, <https://doi.org/10.1175/BAMS-D-14-00246.1>, 2016.
- 915 Kirbus, B., Chylik, J., Ehrlich, A., Becker, S., Schäfer, M., Neggers, R., and Wendisch, M.: Analysis of an Arctic cold air outbreak during autumn and related air mass transformations forced by surface changes and advection in higher altitudes, *Elem. Sci. Anth.*, 11, <https://doi.org/10.1525/elementa.2023.00079>, 2023.
- Kirbus, B., Schirmacher, I., Klingebiel, M., Schäfer, M., Ehrlich, A., Slättberg, N., Lucke, J., Moser, M., Müller, H., and Wendisch, M.: Thermodynamic and cloud evolution in a cold-air outbreak during HALO-(AC)³: quasi-Lagrangian observations compared to the ERA5 and CARRA reanalyses, *Atmos. Chem. Phys.*, 24, 3883–3904, <https://doi.org/10.5194/acp-24-3883-2024>, 2024.
- 920 Kirschler, S., Voigt, C., Anderson, B. E., Chen, G., Crosbie, E. C., Ferrare, R. A., Hahn, V., Hair, J. W., Kaufmann, S., Moore, R. H., Painemal, D., Robinson, C. E., Sanchez, K. J., Scarino, A. J., Shingler, T. J., Shook, M. A., Thornhill, K. L., Winstead, E. L., Ziemba, L. D., and Sorooshian, A.: Overview and statistical analysis of boundary layer clouds and precipitation over the western North Atlantic Ocean, *Atmos. Chem. Phys.*, 23, 10731–10750, <https://doi.org/10.5194/acp-23-10731-2023>, 2023.
- 925 Kölling, T., Zinner, T., and Mayer, B.: Aircraft-based stereographic reconstruction of 3-D cloud geometry, *Atmos. Meas. Technol.*, 12, 1155–1166, <https://doi.org/10.5194/amt-12-1155-2019>, 2019.
- Konow, H., Ewald, F., George, G., Jacob, M., Klingebiel, M., Kölling, T., Luebke, A. E., Mieslinger, T., Pörtge, V., Radtke, J., Schäfer, M., Schulz, H., Vogel, R., Wirth, M., Bony, S., Crewell, S., Ehrlich, A., Forster, L., Giez, A., Göttsche, F., Groß, S., Gutleben, M., Hagen, M., Hirsch, L., Jansen, F., Lang, T., Mayer, B., Mech, M., Prange, M., Schnitt, S., Vial, J., Walbröl, A., Wendisch, M., Wolf, K., Zinner, T.,
- 930 Zöger, M., Ament, F., and Stevens, B.: EUREC⁴A's HALO, *Earth Syst. Sci. Data*, 13, 5545–5563, <https://doi.org/10.5194/essd-13-5545-2021>, 2021.
- Kretzschmar, J., Stapf, J., Klocke, D., Wendisch, M., and Quaas, J.: Employing airborne radiation and cloud microphysics observations to improve cloud representation in ICON at kilometer-scale resolution in the Arctic, *Atmos. Chem. Phys.*, 20, 13145–13165, <https://doi.org/10.5194/acp-20-13145-2020>, 2020.
- 935 Leaitch, W. R., Aliabadi, A., Korolev, A. A., Burkart, J., Willis, M. D., Abbatt, J. P. D., Bozem, H., Hoor, P., Köllner, F., Schneider, J., Herber, A., Konrad, C., and Brauner, R.: Effects of 20–100nm particles on liquid clouds in the clean summertime Arctic, *Atmos. Chem. Phys.*, 16, 11107–11124, <https://doi.org/10.5194/acp-16-11107-2016>, 2016.
- Li, X.-Y., Wang, H., Chen, J., Endo, S., George, G., Cairns, B., Chellappan, S., Zeng, X., Kirschler, S., Voigt, C., Sorooshian, A., Crosbie, E., Chen, G., Ferrare, R. A., Gustafson, W. I., Hair, J. W., Kleb, M. M., Liu, H., Moore, R., Painemal, D., Robinson, C., Scarino, A. J., Shook, M., Shingler, T. J., Thornhill, K. L., Tornow, F., Xiao, H., Ziemba, L. D., and Zuidema, P.: Large-Eddy Simulations of Marine Boundary Layer Clouds Associated with Cold-Air Outbreaks during the ACTIVATE Campaign. Part I: Case Setup and Sensitivities to Large-Scale Forcings, *J. Atmos. Sci.*, 79, 73–100, <https://doi.org/10.1175/JAS-D-21-0123.1>, 2022.
- Linke, O., Quaas, J., Baumer, F., Becker, S., Chylik, J., Dahlke, S., Ehrlich, A., Handorf, D., Jacobi, C., Kalesse-Los, H., Lelli, L., Mehrdad, S., Neggers, R. A., Riebold, J., Garfias, P. S., Schnierstein, N., Shupe, M. D., Smith, C., Spreen, G., Verneuil, B., Vinjamuri, K. S.,

- 945 Vountas, M., and Wendisch, M.: Constraints on simulated past Arctic amplification and lapse-rate feedback from observations, *Atmos. Chem. Phys.*, 23, 9963–9992, <https://doi.org/10.5194/acp-23-9963-2023>, 2023.
- Lohmann, U. and Neubauer, D.: The importance of mixed-phase and ice clouds for climate sensitivity in the global aerosol–climate model ECHAM6-HAM2, *Atmos. Chem. Phys.*, 18, 8807–8828, <https://doi.org/10.5194/acp-18-8807-2018>, 2018.
- Lonardi, M., Akansu, E. F., Ehrlich, A., Mazzola, M., Pilz, C., Shupe, M. D., Siebert, H., and Wendisch, M.: Tethered balloon-borne
950 observations of thermal-infrared irradiance and cooling rate profiles in the Arctic atmospheric boundary layer, *Atmos. Chem. Phys.*, 24, 1961–1978, <https://doi.org/10.5194/acp-24-1961-2024>, 2024.
- Lüpkes, C., Vihma, T., Birnbaum, G., Dierer, S., Garbrecht, T., Gryanik, V. M., Gryschka, M., Hartmann, J., Heinemann, G., Kaleschke, L., Raasch, S., Savijürvi, H., Schlünzen, K. H., and Wacker, U.: Mesoscale modelling of the Arctic atmospheric boundary layer and its interaction with sea ice, vol. 43, chap. ARCTIC Climate Change - The ACSYS Decade and Beyond (edited by Peter Lemke and
955 Hans-Werner Jacobi), Springer, Atmospheric and Oceanographic Sciences Library, <https://doi.org/10.1007/978-94-007-2027-5>, 2012.
- Maherndl, N., Moser, M., Lucke, J., Mech, M., Risse, N., Schirmacher, I., and Maahn, M.: Quantifying riming from airborne data during the HALO-(AC)³ campaign, *Atmos. Meas. Technol.*, 17, 1475–1495, <https://doi.org/10.5194/amt-17-1475-2024>, 2024.
- Manabe, S. and Wetherald, R. T.: The effects of doubling the CO₂ concentration on the climate of a general circulation model, *J. Atmos. Sci.*, 32, 3–15, [https://doi.org/10.1175/1520-0469\(1975\)032<0003:TEODTC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1975)032<0003:TEODTC>2.0.CO;2), 1975.
- 960 Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B.: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, UK, <https://doi.org/10.1017/9781009157896>, 2021.
- Mateling, M. E., Pettersen, C., Kulie, M. S., Ecuyer, L., and Tristan, S.: Marine Cold-Air Outbreak Snowfall in the North Atlantic: A
965 CloudSat Perspective, *J. Geophys. Res.*, 128, e2022JD038 053, <https://doi.org/10.1029/2022JD038053>, 2023.
- McCoy, D. T., Hartmann, D. L., Zelinka, M. D., Ceppi, P., and Grosvenor, D. P.: Mixed-phase cloud physics and Southern Ocean cloud feedback in climate models, *J. Geophys. Res.*, 120, 9539–9554, <https://doi.org/10.1002/2015JD023603>, 2015.
- Mech, M., Orlandi, E., Crewell, S., Ament, F., Hirsch, L., Hagen, M., Peters, G., and Stevens, B.: HAMP – the microwave package on the High Altitude and LOng range research aircraft (HALO), *Atmos. Meas. Technol.*, 7, 4539–4553, <https://doi.org/10.5194/amt-7-4539-2014>, 2014.
- 970 Mech, M., Kliesch, L.-L., Anhäuser, A., Rose, T., Kollias, P., and Crewell, S.: Microwave Radar/radiometer for Arctic Clouds (MiRAC): first insights from the ALOUD campaign, *Atmos. Meas. Technol.*, 12, 5019–5037, <https://doi.org/10.5194/amt-12-5019-2019>, 2019.
- Mech, M., Ehrlich, A., Herber, A., Lüpkes, C., Wendisch, M., Crewell, S., Becker, S., Boose, Y., Chechin, D., Dupuy, R., Gourbeyre, C., Hartmann, J., Jäkel, E., Jourdan, O., Kliesch, L.-L., Kulla, B. S., Mioche, G., Moser, M., Risse, N., Schäfer, M., Klingebiel, M.,
975 Stapf, J., and Voigt, C.: MOSAiC-ACA and AFLUX: Arctic airborne campaigns characterizing the exit area of MOSAiC, *Sci. Data*, 9, <https://doi.org/10.1038/s41597-022-01900-7>, 2022.
- Melia, N., Haines, K., and Hawkins, E.: Sea ice decline and 21st century trans-Arctic shipping routes, *J. Geophys. Res.*, 43, 9720–9728, <https://doi.org/10.1002/2016GL069315>, 2016.
- Moser, M., Voigt, C., Jurkat-Witschas, T., Hahn, V., Mioche, G., Jourdan, O., Dupuy, R., Gourbeyre, C., Schwarzenboeck, A., Lucke, J.,
980 Boose, Y., Mech, M., Borrmann, S., Ehrlich, A., Herber, A., Lüpkes, C., and Wendisch, M.: Microphysical and thermodynamic phase analyses of Arctic low-level clouds measured above the sea ice and the open ocean in spring and summer, *Atmos. Chem. Phys.*, 23, 7257–7280, <https://doi.org/10.5194/acp-23-7257-2023>, 2023.

- Murray-Watson, R. J., Gryspeerdt, E., and Goren, T.: Investigating the development of clouds within marine cold-air outbreaks, *Atmos. Chem. Phys.*, 23, 9365–9383, <https://doi.org/10.5194/acp-23-9365-2023>, 2023.
- 985 Nash, D., Waliser, D., Guan, B., Ye, H., and Ralph, F. M.: The role of atmospheric rivers in extratropical and polar hydroclimate, *J. Geophys. Res.: Atmos.*, 123, 6804–6821, <https://doi.org/10.1029/2017JD028130>, 2018.
- Neggers, R. A. J., Chylík, J., Egerer, U., Griesche, H., Schemann, V., Seifert, P., Siebert, H., and Macke, A.: Local and remote controls on Arctic mixed-layer evolution, *J. Adv. Model. Earth Syst.*, 11, 2214–2237, <https://doi.org/10.1029/2019MS001671>, 2019.
- Nygård, T., Papritz, L., Naakka, T., and Vihma, T.: Cold wintertime air masses over Europe: where do they come from and how do they
990 form?, *Weath. Clim. Dyn.*, 4, 943–961, <https://doi.org/10.5194/wcd-4-943-2023>, 2023.
- Ogren, J. A., Heintzenberg, J., and Charlson, R. J.: In-situ sampling of clouds with a droplet to aerosol converter, *Geophys. Res. Lett.*, 12, 121–124, <https://doi.org/10.1029/GL012i003p00121>, 1985.
- Olonscheck, D., Mauritsen, T., and Notz, D.: Arctic sea-ice variability is primarily driven by atmospheric temperature fluctuations, *Nat. Geosci.*, 12, 430–434, <https://doi.org/10.1038/s41561-019-0363-1>, 2019.
- 995 Overland, J. E., Wood, K. R., and Wang, M.: Warm Arctic–cold conditions: Impacts of the newly open Arctic Sea, *Polar Res.*, 30, 15,787, <https://doi.org/10.3402/polar.v30i0.15787>, 2011.
- Papritz, L. and Spengler, T.: A Lagrangian Climatology of Wintertime Cold Air Outbreaks in the Irminger and Nordic Seas and Their Role in Shaping Air-Sea Heat Fluxes, *J. Clim.*, 30, 2717 – 2737, <https://doi.org/10.1175/JCLI-D-16-0605.1>, 2017.
- Paulus, F. M., Karalis, M., George, G., Svensson, G., Wendisch, M., and Neggers, R. A. J.: Airborne measurements of mesoscale divergence
1000 at high latitudes during HALO-(AC)³, Submitted to *J. Atmos. Sci.*, 2024.
- Pefanis, V., Losa, S. N., Losch, M., Janout, M. A., and Bracher, A.: Amplified Arctic Surface Warming and Sea Ice Loss Due to Phytoplankton and Colored Dissolved Material, *Geophys. Res. Lett.*, 47, e2020GL088 795, <https://doi.org/10.1029/2020GL088795>, 2020.
- Pithan, F., Medeiros, B., and Mauritsen, T.: Mixed-phase clouds cause climate model biases in Arctic wintertime temperature inversions, *Clim. Dyn.*, 43, 289–303, <https://doi.org/10.1007/s00382-013-1964-9>, 2014.
- 1005 Pithan, F., Ackerman, A., Angevine, W. M., Hartung, K., Ickes, L., Kelley, M., Medeiros, B., Sandu, I., Steeneveld, G.-J., Sterk, H. A. M., Svensson, G., Vaillancourt, P. A., and Zadra, A.: Select strengths and biases of models in representing the Arctic winter boundary layer over sea ice: the Larcform 1 single column model intercomparison, *J. Adv. Model. Earth Syst.*, 8, 1345–1357, <https://doi.org/10.1002/2016MS000630>, 2016.
- Pithan, F., Svensson, G., Caballero, R., Chechin, D., Cronin, T., Ekman, A., Neggers, R. A. J., Shupe, M., Solomon, A., Tjernström, M., and Wendisch, M.: Role of air-mass transformations in exchange between the Arctic and mid-latitudes, *Nat. Geosci.*, 11, 805–812, <https://doi.org/10.1038/s41561-018-0234-1>, 2018.
- 1010 Pithan, F., Athanase, M., Dahlke, S., Sánchez-Benítez, A., Shupe, M. D., Sledd, A., Streffing, J., Svensson, G., and Jung, T.: Nudging allows direct evaluation of coupled climate models with in situ observations: a case study from the MOSAiC expedition, *Geosci. Model Dev.*, 16, 1857–1873, <https://doi.org/10.5194/gmd-16-1857-2023>, 2023.
- 1015 Polkova, I., Afargan-Gerstman, H., Domeisen, D. I. V., King, M. P., Ruggieri, P., Athanasiadis, P., Dobrynin, M., Aarnes, O., Kretschmer, M., and Baehr, J.: Predictors and prediction skill for marine cold-air outbreaks over the Barents Sea, *Q. J. R. Meteorol. Soc.*, 147, 2638–2656, <https://doi.org/10.1002/qj.4038>, 2021.
- Previdi, M., Smith, K. L., and Polvani, L. M.: Arctic amplification of climate change: a review of underlying mechanisms, *Environ. Res. Lett.*, 16, 093 003, <https://doi.org/10.1088/1748-9326/ac1c29>, 2021.

- 1020 Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Commun. Earth Environ.*, 3, 168, <https://doi.org/10.1038/s43247-022-00498-3>, 2022.
- Rinke, A., Knudsen, E., Mewes, D., Dorn, W., Handorf, D., Dethloff, K., and Moore, J. C.: Arctic summer sea-ice melt and related atmospheric conditions in coupled regional climate model simulations and observations, *J. Geophys. Res.*, 124, 6027–6039, <https://doi.org/10.1029/2018JD030207>, 2019.
- 1025 Ripple, W. J., Wolf, C., Lenton, T. M., Gregg, J. W., Natali, S. M., Duffy, P. B., Rockström, J., and Schellnhuber, H. J.: Many risky feedback loops amplify the need for climate action, *One Earth*, 6, 86–91, <https://doi.org/10.1016/j.oneear.2023.01.004>, 2023.
- Roberts, T. J., Dütsch, M., Hole, L. R., and Voss, P. B.: Controlled meteorological (CMET) free balloon profiling of the Arctic atmospheric boundary layer around Spitsbergen compared to ERA-Interim and Arctic System Reanalyses, *Atmos. Chem. Phys.*, 16, 12 383–12 396, <https://doi.org/10.5194/acp-16-12383-2016>, 2016.
- 1030 Sato, Y., Miura, H., Yashiro, H., Goto, D., Takemura, T., Tomita, H., and Nakajima, T.: Unrealistically pristine air in the Arctic produced by current global scale models, *Sci. Rep.*, 6, 26 561, <https://doi.org/10.1038/srep26561>, 2016.
- Schäfer, M., Wolf, K., Ehrlich, A., Hallbauer, C., Jäkel, E., Jansen, F., Luebke, A. E., Müller, J., Thoböll, J., Rösenthaller, T., Stevens, B., and Wendisch, M.: VELOX – a new thermal infrared imager for airborne remote sensing of cloud and surface properties, *Atmos. Meas. Technol.*, 15, 1491–1509, <https://doi.org/10.5194/amt-15-1491-2022>, 2022.
- 1035 Schirmacher, I., Kollias, P., Lamer, K., Mech, M., Pfitzenmaier, L., Wendisch, M., and Crewell, S.: Assessing Arctic low-level clouds and precipitation from above – a radar perspective, *EGUsphere*, pp. 1–30, <https://doi.org/10.5194/egusphere-2023-636>, [preprint], 2023.
- Schirmacher, I., Schnitt, S., Klingebiel, M., Mahernndl, N., Kirbus, B., and Crewell, S.: Clouds and precipitation in the initial phase of marine cold air outbreaks as observed by airborne remote sensing, To be submitted to *Special Issue of Atmos. Chem. Phys.*, 2024.
- 1040 Schmale, J., Zieger, P., and Ekman, A. M. L.: Aerosols in current and future Arctic climate, *Nat. Clim. Change*, 11, 95–105, <https://doi.org/10.1038/s41558-020-00969-5>, 2021.
- Screen, J. A.: An ice-free Arctic: What could it mean for European weather?, *Weather*, 76, 327–328, <https://doi.org/10.1002/wea.4069>, 2021.
- Sedlar, J., Tjernström, M., Mauritsen, T., Shupe, M. D., Brooks, I. M., Persson, P. O. G., Birch, C. E., Leck, C., Sirevaag, A., and Nicolaus, M.: A transitioning Arctic surface energy budget: the impacts of solar zenith angle, surface albedo and cloud radiative forcing, *Clim. Dyn.*, 37, 1643–1660, <https://doi.org/10.1007/s00382-010-0937-5>, 2011.
- 1045 Serreze, M. C. and Barry, R. C.: Processes and impacts of Arctic amplification: A research synthesis, *Glob. Planet. Change*, 77, 85–96, <https://doi.org/10.1016/j.gloplacha.2011.03.004>, 2011.
- Serreze, M. C. and Francis, J. A.: The Arctic Amplification debate, *Clim. Change*, 76, 241–264, <https://doi.org/10.1007/s10584-005-9017-y>, 2006.
- 1050 Serreze, M. C. and Meier, W. N.: The Arctic’s sea ice cover: trends, variability, predictability, and comparisons to the Antarctic, *Annals of the New York Academy of Sciences*, 1436, 36–53, <https://doi.org/10.1111/nyas.13856>, 2019.
- Serreze, M. C., Barrett, A. P., Stroeve, J. C., Kindig, D. N., and Holland, M. M.: The emergence of surface-based Arctic amplification, *The Cryosphere*, 3, 11–19, <https://doi.org/10.5194/tc-3-11-2009>, 2009.
- Shapiro, M. A. and Keyser, D.: Fronts, Jet Streams and the Tropopause, pp. 167–191, American Meteorological Society, Boston, MA, https://doi.org/10.1007/978-1-944970-33-8_10, 1990.
- 1055

- Shupe, M. D., Daniel, J. S., de Boer, G., Eloranta, E. W., Kollias, P., Long, C. N., Luke, E. P., Turner, D. D., and Verlinde, J.: A focus on mixed-phase clouds: The status of ground-based observational methods, *Bull. Am. Meteorol. Soc.*, 89, 1549–1562, <https://doi.org/10.1175/2008BAMS2378.1>, 2008.
- 1060 Shupe, M. D., Rex, M., Blomquist, B., Persson, P. O. G., Schmale, J., Uttal, T., Althausen, D., Angot, H., Archer, S., Bariteau, L., Beck, I., Bilberry, J., Bussi, S., Buck, C., Boyer, M., Brasseur, Z., Brooks, I. M., Calmer, R., Cassano, J., Castro, V., Chu, D., Costa, D., Cox, C. J., Creamean, J., Crewell, S., Dahlke, S., Damm, E., de Boer, G., Deckelmann, H., Dethloff, K., Dütsch, M., Ebell, K., Ehrlich, A., Ellis, J., Engelmann, R., Fong, A. A., Frey, M. M., Gallagher, M. R., Ganzeveld, L., Gradinger, R., Graeser, J., Greenamyre, V., Griesche, H., Griffiths, S., Hamilton, J., Heinemann, G., Helmig, D., Herber, A., Heuzé, C., Hofer, J., Houchens, T., Howard, D., Inoue, J., Jacobi, H.-W., Jaiser, R., Jokinen, T., Jourdan, O., Jozef, G., King, W., Kirchgaessner, A., Klingebiel, M., Krassovski, M., Krumpfen, T., Lampert, A.,
- 1065 Landing, W., Laurila, T., Lawrence, D., Lonardi, M., Loose, B., Lüpkens, C., Maahn, M., Macke, A., Maslowski, W., Marsay, C., Maturilli, M., Mech, M., Morris, S., Moser, M., Nicolaus, M., Ortega, P., Osborn, J., Pätzold, F., Perovich, D. K., Petäjä, T., Pilz, C., Pirazzini, R., Posman, K., Powers, H., Pratt, K. A., Preußner, A., Quéféver, L., Radenz, M., Rabe, B., Rinke, A., Sachs, T., Schulz, A., Siebert, H., Silva, T., Solomon, A., Sommerfeld, A., Spreen, G., Stephens, M., Stohl, A., Svensson, G., Uin, J., Viegas, J., Voigt, C., von der Gathen, P., Wehner, B., Welker, J. M., Wendisch, M., Werner, M., Xie, Z., and Yue, F.: Overview of the MOSAiC expedition: Atmosphere, *Elem. Sci.*
- 1070 *Anth.*, 10, <https://doi.org/10.1525/elementa.2021.00060>, 2022.
- Smith, A. J. P., Smith, D. M., Cohen, J., and Jones, M. W.: Arctic warming amplifies climate change and its impacts, *Zenodo*, <https://doi.org/10.5281/zenodo.5596791>, 2021.
- Smith, D. M., Screen, J. A., Deser, C., Cohen, J., Fyfe, J. C., García-Serrano, J., Jung, T., Kattsov, V., Matei, D., Msadek, R., Peings, Y., Sigmond, M., Ukita, J., Yoon, J.-H., and Zhang, X.: The Polar Amplification Model Intercomparison Project (PAMIP) contribution to CMIP6: investigating the causes and consequences of polar amplification, *Geosci. Model Dev.*, 12, 1139–1164, <https://doi.org/10.5194/gmd-12-1139-2019>, 2019.
- 1075 Smith, D. M., Eade, R., Andrews, M. B., Ayres, H., Clark, A., Chripko, S., Deser, C., Dunstone, N. J., García-Serrano, J., Gastineau, G., Graff, L. S., Hardiman, S. C., He, B., Hermanson, L., Jung, T., Knight, J., Levine, X., Magnusdottir, G., Manzini, E., Matei, D., Mori, M., Msadek, R., Ortega, P., Peings, Y., Scaife, A. A., Screen, J. A., Seabrook, M., Semmler, T., Sigmond, M., Streffing, J., Sun, L., and Walsh, A.: Robust
- 1080 but weak winter atmospheric circulation response to future Arctic sea ice loss, *Nat. Commun.*, 13, 727, <https://doi.org/10.1038/s41467-022-28283-y>, 2022.
- Sobel, A. H., Nilsson, J., and Polvani, L. M.: The Weak Temperature Gradient Approximation and Balanced Tropical Moisture Waves, *J. Atmos. Sci.*, 58, 3650–3665, [https://doi.org/10.1175/1520-0469\(2001\)058<3650:TWTGAA>2.0.CO;2](https://doi.org/10.1175/1520-0469(2001)058<3650:TWTGAA>2.0.CO;2), 2001.
- Solomon, A., Shupe, M. D., Svensson, G., Barton, N. P., Batrak, Y., Bazile, E., Day, J. J., Doyle, J. D., Frank, H. P., Keeley, S., Remes, T.,
- 1085 and Tolstykh, M.: The winter central Arctic surface energy budget: A model evaluation using observations from the MOSAiC campaign, *Elem. Sci. Anth.*, 11, 00 104, <https://doi.org/10.1525/elementa.2022.00104>, 2023.
- Sprenger, M. and Wernli, H.: The LAGRANTO Lagrangian analysis tool - version 2.0, *Geosci. Model Dev.*, 8, 2569–2586, <https://doi.org/10.5194/gmd-8-2569-2015>, 2015.
- Stapf, J., Ehrlich, A., Jäkel, E., Lüpkens, C., and Wendisch, M.: Reassessment of shortwave surface cloud radiative forcing in the Arctic: consideration of surface-albedo–cloud interactions, *Atmos. Chem. Phys.*, 20, 9895–9914, <https://doi.org/10.5194/acp-20-9895-2020>, 2020.
- 1090 Stapf, J., Ehrlich, A., and Wendisch, M.: Influence of Thermodynamic State Changes on Surface Cloud Radiative Forcing in the Arctic: A Comparison of Two Approaches Using Data From AFLUX and SHEBA, *J. Geophys. Res. Atmos.*, 126, e2020JD033 589, <https://doi.org/10.1029/2020JD033589>, 2021.

- 1095 Stevens, B. and Kluff, L.: A colorful look at climate sensitivity, *Atmos. Chem. Phys.*, 23, 14 673–14 689, <https://doi.org/10.5194/acp-23-14673-2023>, 2023.
- Stevens, B., Ament, F., Bony, S., Crewell, S., Ewald, F., Gross, S., Hansen, A., Hirsch, L., Jacob, M., Kölling, T., Konow, H., Mayer, B., Wendisch, M., Wirth, M., Wolf, K., Bakan, S., Bauer-Pfundstein, B., Brueck, M., Delanoë, J., Ehrlich, A., Farrell, D., Forde, M., Göttele, F., Grob, H., Hagen, M., Jäkel, E., Jansen, F., Klepp, C., Klingebiel, M., Mech, M., Peters, G., Rapp, M., Wing, A. A., and Zinner, T.: A high-altitude long-range aircraft configured as a cloud observatory—the NARVAL expeditions, *Bull. Am. Meteorol. Soc.*, 100, 1061–1077, <https://doi.org/10.1175/BAMS-D-18-0198.1>, 2019.
- 1100 Stroeve, J., Holland, M. M., Meier, W., Scambos, T., and Serreze, M.: Arctic sea ice decline: Faster than forecast, *Geophys. Res. Lett.*, 34, <https://doi.org/10.1029/2007GL029703>, 2007.
- Svensson, G., Murto, S., Shupe, M. D., Pithan, F., Magnusson, L., Day, J. J., Doyle, J. D., Renfrew, I. A., Spengler, T., and Vihma, T.: Warm air intrusions reaching the MOSAiC expedition in April 2020—The YOPP targeted observing period (TOP), *Elem. Sci. Anth.*, 11, 00016, <https://doi.org/10.1525/elementa.2023.00016>, 2023.
- 1105 Tan, I. and Storelvmo, T.: Evidence of Strong Contributions From Mixed-Phase Clouds to Arctic Climate Change, *Geophys. Res. Lett.*, 46, 2894–2902, <https://doi.org/10.1029/2018GL081871>, 2019.
- Taylor, P. C., Boeke, R. C., Boisvert, L. N., Feldl, N., Henry, M., Huang, Y., Langen, P. L., Liu, W., Pithan, F., Sejas, S. A., and Tan, I.: Process drivers, inter-model spread, and the path forward: A review of amplified Arctic warming, *Front. Earth Sci.*, 9, 758361, <https://doi.org/10.3389/feart.2021.758361>, 2022.
- 1110 Tjernström, M. and Graversen, R. G.: The vertical structure of the lower Arctic troposphere analysed from observations and the ERA-40 reanalysis, *Q. J. R. Meteorol. Soc.*, 135, 431–443, <https://doi.org/10.1002/qj.380>, 2009.
- Tjernström, M., Zagar, M., Svensson, G., Cassano, J. J., Pfeifer, S., Rinke, A., Wyser, K., Dethloff, K., Jones, C., Semmler, T., and Shaw, M.: Modelling the Arctic Boundary Layer: An Evaluation of Six Arctic Regional-Scale Models using Data from the Sheba Project, *Bound.-Layer Meteorol.*, 117, 337–381, <https://doi.org/10.1007/s10546-004-7954-z>, 2005.
- 1115 Tjernström, M., Shupe, M. D., Brooks, I. M., Achtert, P., Prytherch, J., and Sedlar, J.: Arctic Summer Air Mass Transformation, Surface Inversions, and the Surface Energy Budget, *J. Clim.*, 32, 769–789, <https://doi.org/10.1175/JCLI-D-18-0216.1>, 2019.
- Urbanek, B., Groß, S., Schäfler, A., and Wirth, M.: Determining stages of cirrus evolution: a cloud classification scheme, *Atmos. Meas. Technol.*, 10, 1653–1664, <https://doi.org/10.5194/amt-10-1653-2017>, 2017.
- 1120 Uttal, T., Curry, J. A., McPhee, M. G., Perovich, D. K., Moritz, R. E., Maslanik, J. A., Guest, P. S., Stern, H., Moore, J. A., Turenne, R., Heiberg, A., Serreze, M. C., Wylie, D. P., Persson, O. G., Paulson, C. A., Halle, C., Morison, J. H., Wheeler, P. A., Makshtas, A., Welch, H., Shupe, M. D., Intrieri, J. M., Stamnes, K., Lindsey, R. W., Pinkel, R., Pegau, W. S., Stanton, T. P., and Grenfeld, T. C.: Surface heat budget of the arctic ocean, *Bull. Am. Meteorol. Soc.*, 83, 255–276, [https://doi.org/10.1175/1520-0477\(2002\)083<0255:SHBOTA>2.3.CO;2](https://doi.org/10.1175/1520-0477(2002)083<0255:SHBOTA>2.3.CO;2), 2002.
- 1125 Vihma, T., Hartmann, J., and Lüpkes, C.: A case study of an on-ice air flow over the Arctic marginal sea ice zone, *Bound.-Layer Meteorol.*, 107, 189–217, <https://doi.org/10.1023/A:1021599601948>, 2003.
- Volkmer, L., Kölling, T., Zinner, T., and Mayer, B.: Consideration of the cloud motion for aircraft-based stereographically derived cloud geometry and cloud top heights, *Atmos. Meas. Technol. Discuss.*, 2024, 1–13, <https://doi.org/10.5194/amt-2024-19>, 2024.
- 1130 von Lerber, A., Mech, M., Rinke, A., Zhang, D., Lauer, M., Radovan, A., Gorodetskaya, I., and Crewell, S.: Evaluating seasonal and regional distribution of snowfall in regional climate model simulations in the Arctic, *Atmos. Chem. Phys.*, 22, 7287–7317, <https://doi.org/10.5194/acp-22-7287-2022>, 2022.

- Walbröl, A., Michaelis, J., Becker, S., Dorff, H., Gorodetskaya, I., Kirbus, B., Lauer, M., Mahernndl, N., Maturilli, M., Mayer, J., Müller, H., Rückert, J., Schirmacher, I., Slättberg, N., Röttenbacher, J., Ehrlich, A., Wendisch, M., and Crewell, S.: Environmental conditions during the HALO-(AC)³ campaign in the North Atlantic sector of the Arctic, EGU sphere, <https://doi.org/10.5194/egusphere-2023-668>, [preprint], 2023.
- 1135
- Weber, A., Kölling, T., Pörtge, V., Baumgartner, A., Rammeloo, C., Zinner, T., and Mayer, B.: Polarization upgrade of specMACS: calibration and characterization of the 2D RGB polarization-resolving cameras, *Atmos. Meas. Technol.*, 17, 1419–1439, <https://doi.org/10.5194/amt-17-1419-2024>, 2024.
- Wendisch, M. and Brenguier, J.-L.: *Airborne Measurements for Environmental Research – Methods and Instruments*, Wiley–VCH Verlag GmbH & Co. KGaA, Weinheim, Germany, Weinheim, Germany, ISBN: 978-3-527-40996-9, 2013.
- 1140
- Wendisch, M., Macke, A., Ehrlich, A., Lüpkes, C., Mech, M., Chechin, D., Barrientos, C., Bozem, H., Brückner, M., Clemen, H.-C., Crewell, S., Donth, T., Dupuy, R., Ebell, K., Egerer, U., Engelmann, R., Engler, C., Eppers, O., Gehrman, M., Gong, X., Gottschalk, M., Gourbeyre, C., Griesche, H., Hartmann, J., Hartmann, M., Herber, A., Herrmann, H., Heygster, G., Hoor, P., Jafariserajehlou, S., Jäkel, E., Järvinen, E., Jourdan, O., Kästner, U., Kecorius, S., Knudsen, E. M., Köllner, F., Kretzschmar, J., Lelli, L., Leroy, D., Maturilli, M., Mei, L., Mertes, S., Mioche, G., Neuber, R., Nicolaus, M., Nomokonova, T., Notholt, J., Palm, M., van Pinxteren, M., Quaas, J., Richter, P., Ruiz-Donoso, E., Schäfer, M., Schmieder, K., Schnaiter, M., Schneider, J., Schwarzenböck, A., Seifert, P., Shupe, M. D., Siebert, H., Spreen, G., Stapf, J., Stratmann, F., Vogl, T., Welti, A., Wex, H., Wiedensohler, A., Zanatta, M., Zeppenfeld, S., Dethloff, K., and Heinold, B.: The Arctic Cloud Puzzle: Using ACLOUD/PASCAL Multi-Platform Observations to Unravel the Role of Clouds and Aerosol Particles in Arctic Amplification, *Bull. Am. Meteorol. Soc.*, 100 (5), 841–871, <https://doi.org/10.1175/BAMS-D-18-0072.1>, 2019.
- 1145
- Wendisch, M., Handorf, D., Tegen, I., Neggers, R. A. J., and Spreen, G.: Glimpsing the ins and outs of the Arctic atmospheric cauldron, *EOS*, 102, <https://doi.org/10.1029/2021EO155959>, 2021.
- 1150
- Wendisch, M., Brückner, M., Crewell, S., Ehrlich, A., Notholt, J., Lüpkes, C., Macke, A., Burrows, J. P., Rinke, A., Quaas, J., Maturilli, M., Schemann, V., Shupe, M. D., Akansu, E. F., Barrientos-Velasco, C., Bärfuss, K., Blechschmidt, A.-M., Block, K., Bougoudis, I., Bozem, H., Böckmann, C., Bracher, A., Bresson, H., Bretschneider, L., Buschmann, M., Chechin, D. G., Chylik, J., Dahlke, S., Deneke, H., Dethloff, K., Donth, T., Dorn, W., Dupuy, R., Ebell, K., Egerer, U., Engelmann, R., Eppers, O., Gerdes, R., Gierens, R., Gorodetskaya, I. V., Gottschalk, M., Griesche, H., Gryanik, V. M., Handorf, D., Harm-Altstädter, B., Hartmann, J., Hartmann, M., Heinold, B., Herber, A., Herrmann, H., Heygster, G., Höschel, I., Hofmann, Z., Hälemann, J., Hünerbein, A., Jafariserajehlou, S., Jäkel, E., Jacobi, C., Janout, M., Jansen, F., Jourdan, O., Jurányi, Z., Kalesse-Los, H., Kanzow, T., Käthner, R., Kliesch, L. L., Klingebiel, M., Knudsen, E. M., Kovács, T., Körtke, W., Krampe, D., Kretzschmar, J., Kreyling, D., Kulla, B., Kunkel, D., Lampert, A., Lauer, M., Lelli, L., von Lerber, A., Linke, O., Löhnert, U., Lonardi, M., Losa, S. N., Losch, M., Maahn, M., Mech, M., Mei, L., Mertes, S., Metzner, E., Mewes, D., Michaelis, J., Mioche, G., Moser, M., Nakoudi, K., Neggers, R., Neuber, R., Nomokonova, T., Oelker, J., Papakonstantinou-Presvelou, I., Pätzold, F., Pefanis, V., Pohl, C., van Pinxteren, M., Radovan, A., Rhein, M., Rex, M., Richter, A., Risse, N., Ritter, C., Rostosky, P., Rozanov, V. V., Ruiz-Donoso, E., Saavedra-Garfias, P., Salzmänn, M., Schacht, J., Schäfer, M., Schneider, J., Schnierstein, N., Seifert, P., Seo, S., Siebert, H., Soppa, M. A., Spreen, G., Stachlewska, I. S., Stapf, J., Stratmann, F., Tegen, I., Viceto, C., Voigt, C., Vountas, M., Walbröl, A., Walter, M., Wehner, B., Wex, H., Willmes, S., Zanatta, M., and Zeppenfeld, S.: Atmospheric and Surface Processes, and Feedback Mechanisms Determining Arctic Amplification: A Review of First Results and Prospects of the (AC)³ Project, *Bull. Am. Meteorol. Soc.*, 104, E208–E242, <https://doi.org/10.1175/bams-d-21-0218.1>, 2023a.
- 1160
- 1165

- Wendisch, M., Stapf, J., Becker, S., Ehrlich, A., Jäkel, E., Klingebiel, M., Lüpkes, C., Schäfer, M., and Shupe, M. D.: Effects of variable ice-ocean surface properties and air mass transformation on the Arctic radiative energy budget, *Atmos. Chem. Phys.*, 23, 9647–9667, <https://doi.org/10.5194/acp-23-9647-2023>, 2023b.
- 1170 Wesche, C., Steinhage, D., and Nixdorf, U.: Polar aircraft Polar5 and Polar6 operated by the Alfred Wegener Institute, *J. Large-Scale Res. Facilities*, 2, <https://doi.org/10.17815/jlsrf-2-153>, 2016.
- Wirth, M., Fix, A., Mahnke, P., Schwarzer, H., Schrandt, F., and Ehret, G.: The airborne multi-wavelength water vapor differential absorption lidar WALES: system design and performance, *Appl. Phys. B*, 96, 201–213, <https://doi.org/10.1007/s00340-009-3365-7>, 2009.
- 1175 Woods, C. and Caballero, R.: The Role of Moist Intrusions in Winter Arctic Warming and Sea Ice Decline, *J. Clim.*, 29, 4473–4485, <https://doi.org/10.1175/JCLI-D-15-0773.1>, 2016.
- You, C., Tjernström, M., and Devasthale, A.: Warm-air advection over melting sea-ice: A Lagrangian case study, *Bound.-Layer Meteorol.*, 179, 99–116, <https://doi.org/10.1007/s10546-020-00590-1>, 2021a.
- You, C., Tjernström, M., and Devasthale, A.: Eulerian and Lagrangian views of warm and moist air intrusions into summer Arctic, *Atmos. Res.*, 256, 105 586, <https://doi.org/10.1016/j.atmosres.2021.105586>, 2021b.
- 1180 You, C., Tjernström, M., and Devasthale, A.: Warm and moist air intrusions into the winter Arctic: a Lagrangian view on the near-surface energy budgets, *Atmos. Chem. Phys.*, 22, 8037–8057, <https://doi.org/10.5194/acp-22-8037-2022>, 2022.
- Yuval, J. and Kaspi, Y.: Eddy Activity Response to Global Warming–Like Temperature Changes, *J. Climate*, 33, 1381 – 1404, <https://doi.org/10.1175/JCLI-D-19-0190.1>, 2020.

Table 1. Overview of HALO-(AC)³ research flights (RFs) including the RF number and information on the coordination of HALO (High Altitude and Long Range Research Aircraft) with the P5 (Polar 5), P6 (Polar 6), FAAM (Facility for Airborne Atmospheric Measurements), and ATR (Avions de Transport Régional) aircraft. Furthermore, the synoptic situation (warm air intrusion, WAI, cold air outbreak, CAO, Arctic cirrus, AC, and polar low, PL) and overpasses of P5 over AWIPEV research base in Ny-Ålesund are indicated. The number of successfully launched dropsondes is given together with the number of dropsondes used in the Global Telecommunication System (GTS) data assimilation. [The direct transmission to the GTS was setup for HALO only. Therefore, all dropsondes listed here submitted to GTS are from HALO.](#)

	Research Flight (RF) Number			Coordination of HALO with ...	Synoptic Situation	P5 over AWIPEV	Number of Dropsondes		
	HALO	P5	P6				P5	HALO	GTS
WARM & HUMID PERIOD									
DAY IN 2022									
12 March	02	–	–		WAI		–	20	–
13 March	03	–	–		WAI		–	21	–
14 March	04	–	–		WAI		–	9	–
15 March	05	–	–		WAI		–	25	3
16 March	06	–	–	FAAM	WAI		–	23	19
20 March	07	01	01	P5, P6	WAI	YES	12	17	13
COLD & DRY PERIOD									
DAY IN 2022									
21 March	08	–	–	FAAM	CAO		–	13	13
22 March	–	02,03	02		CAO	YES	12	–	–
24 March	–	–	03		CAO		–	–	–
25 March	–	04	–		CAO	YES	5	–	–
26 March	–	–	04		CAO		–	–	–
28 March	09	05	05	P5, P6	CAO	YES	15	16	16
29 March	10	06,07	06	P5, P6, ATR, FAAM	CAO	YES	5	18	10
30 March	11	08	07	P5, P6, ATR, FAAM	CAO	YES	15	32	32
01 April	12	09	08	P5, P6	CAO	YES	18	41	41
04 April	13	10	09	P5, P6	CAO		14	13	11
05 April	–	11	10		CAO		10	–	–
07 April	14	12	–	P5	AC	YES	17	15	10
08 April	15	–	11	P6	PL		–	21	5
09 April	–	–	12		CAO/PL		–	–	–
10 April	16	13	13	P5, P6	AC/WAI		18	22	21
11 April	17	–	–		AC		–	7	6
12 April	18	–	–		AC		–	17	16