

The effect of carbonate mineral additions on biogeochemical conditions in surface sediments and benthic-pelagic exchange fluxes

Kadir Biçe^{1,*}, Tristen Myers Stewart^{2,#}, George Waldbusser², Christof Meile^{1,*}

5 ¹ Department of Marine Sciences, University of Georgia, Athens GA 30602

² College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, Corvallis, OR 97331

now at: Pacific Northwest National Laboratory, Sequim, WA 98382

~ now at: Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA 02139

Correspondence to: Kadir Biçe (bicekadir@gmail.com), Christof Meile (cmeile@uga.edu)

10 **Abstract.** Coastal sediments are hotspots of biogeochemical processes that are impacting subsurface and overlying water conditions. Fluid composition in sediments is altered through the mineralization of organic matter which under oxic conditions, further lowers both pH and the carbonate saturation state. As a potential mitigation strategy for this sediment acidification, we explored the effects of mineral additions to coastal sediments. We experimentally quantified carbonate mineral dissolution kinetics of carbonate shells suitable for field application, and then integrated this data into a reactive transport model that
15 represents early diagenetic cycling of C, O, N, S and Fe, and traces total alkalinity, pH and saturation state of CaCO₃. Model simulations were carried out to delineate the impact of mineral type and amount added, porewater mixing and organic matter mineralization rates on sediment alkalinity and its flux to the overlying water. Model results showed that the added minerals undergo initial rapid dissolution and generate saturated conditions demonstrating the potential of alkalinity enhancement in mitigating surface sediment acidification. Aragonite dissolution led to higher total alkalinity concentrations than calcite.
20 Simulations of carbonate mineral additions to sediment environments with low rates of organic matter mineralization exhibited a substantial increase in mineral saturation state compared to sediments with high CO₂ production rates, highlighting the environment-specific extent of the effect of mineral addition. Our work indicates that carbonate additions have the potential to effectively buffer surficial sediments over multiple years, yielding biogeochemical conditions that counteract the detrimental effect of low pH sediment conditions on larval recruitment, and potentially increase benthic alkalinity fluxes to support marine
25 carbon dioxide removal (mCDR) in the overlying water.

1. Introduction

In coastal systems, ocean-freshwater mixing, terrestrial inputs, and active biogeochemical transformations (Cai et al., 2017; Soetaert et al., 2007) lead to natural fluctuations in pH that are large and decoupled from the decrease in pH observed in the
30 open ocean due to ocean acidification (Hofmann et al., 2011). Especially in shallow water environments, sediments play a significant role as benthic fluxes can constitute a substantial alkalinity source to the overlying water, increasing its buffering

capacity (Rassmann et al., 2016; Krumins et al., 2013; Cyronak et al., 2013b). Sediments receive organic matter (OM) from the overlying water which fuels the dissolution of carbonate minerals and early diagenetic processes. Aerobic respiration and other processes consuming dissolved O₂ can lead to steep pH decreases within the oxygenated sediment and the interaction of
35 OM mineralization and mineral dissolution determines the balance of the porewater carbonate system (Burdige et al., 2010; Jourabchi et al., 2005). Mineral dissolution and early diagenesis are connected through the changes in carbonate system which impacts mineral saturation states and hence the rate of dissolution (e.g., Morse et al., 2007). Apart from its impact on mineral dissolution, OM mineralization is also a significant contributor to alkalinity generation under suboxic and anoxic conditions (Hu & Cai, 2011a). These early diagenetic processes determine the composition of the porewaters, which affect benthic
40 calcifying organisms; furthermore, they can affect the biogeochemical conditions in the overlying water through benthic fluxes releasing nutrients and alkalinity.

In surface sediments, porewater pH is generally lower than in the overlying water, in particular when sediment O₂ uptake is elevated due to burrowing and irrigation activities by benthic macrofauna (e.g. Aller, 1982). This exposes benthic organisms that live in the upper layers of the sediment to more corrosive conditions (Green et al., 2013; Waldbusser & Salisbury, 2014).
45 Combined with abundant OM, enhanced supply of O₂ into the sediment also leads to elevated fluxes of alkalinity and DIC to the water column (Mucci et al., 2000; Burdige, 2007).

In this study, we investigate the effects of the addition of carbonate minerals in the coastal sediments focusing on benthic alkalinity generation and release in a shallow water environment. Such ocean alkalinity enhancement (OAE) aims to buffer increased dissolved CO₂ in the ocean by dissolving carbonate or silicate minerals, producing alkalinity. OAE is mainly studied
50 for its effect on ocean-atmosphere CO₂ exchange, the impact on organisms (Renforth & Henderson, 2017) and its potential to mitigate ocean acidification and lower atmospheric CO₂ concentrations (marine carbon dioxide removal; mCDR). Previous studies have shown that increasing ocean alkalinity through enhanced weathering has the potential to reverse global ocean acidification in an intermediate CO₂ emission scenario (RCP 4.5) and reduce its impacts if emissions were to continue to rise throughout the 21st century (RCP 8.5) (Taylor et al., 2016). To date, OAE implementations mainly target the open ocean and
55 are aimed at identifying the potential of scaling up the mCDR efforts to a global scale (Fennel et al., 2023). Recent studies also focused on the impact of OAE on marine ecosystems such as fish and plankton communities (Goldenberg et al., 2024; Paul et al., 2024; Ferderer et al., 2022). Only a few studies focused on coastal OAE and even fewer on sedimentary applications (Green et al., 2009, 2013; Hangx & Spiers, 2009; Fuhr et al., 2024), where the effectiveness of OAE mainly depends on the dissolution rate of the mineral which is controlled by the mineral and the chemistry of the water and physical properties of the environment
60 (Hartmann et al., 2013; Montserrat et al., 2017). One benefit of implementing OAE in coastal sediments is the generally higher rate of organic matter remineralization. This elevated CO₂ generation aids mineral dissolution, given most overlying marine waters are typically super-saturated with respect to carbonate minerals. Here, we simulate this effect of adding carbonate minerals to the sediment using a reactive transport model. We use measurements of carbonate mineral dissolution rates to assess the extent and duration of buffering in surficial sediments, and the impact on benthic alkalinity fluxes. Building on the

65 previous work by Krumins et al. (2013) that explored the interaction of carbonate chemistry with early diagenesis in coastal
sediments, our work identifies factors that determine the impact of sediment alkalization on the spatial and temporal
distribution of buffering by exploring the effect of mineral type, amount, organic matter remineralization rate and bioturbation.

2. Methods

2.1. Dissolution experiments

70 Dissolution rates for the minerals used in this model were derived from lab-based experiments detailed in Myers (2022). We
provide pertinent details below. Two biomineral calcium carbonates, calcitic oyster shells *Crassostrea gigas* (below referred
to as “bio-calcite”) and aragonitic clam soft-shell clam shells, *Mya arenaria*, (“bio-aragonite”) were collected from Oregon
Oyster Farm in Newport, Oregon, USA and from Broad Cove in Yarmouth, Maine, USA, respectively. The biomineral samples
were briefly cleaned, dried, crushed and sieved through a 2000 μm sieve and retained on 710 μm , for an approximately median
75 grain size of 1 mm. We measured the mineral dissolution rate across four saturation states experimentally; $\Omega_{\text{calcite}} = 0.25$ (low),
0.45 (mid), 0.85 (hi) and 1.03 (saturated); carbonate chemistry conditions are detailed in Table S1. A flow-through, feedback-
controlled with pH monitoring and CO_2 injection seawater system similar to Waldbusser et al. (2011) was used for
manipulations at the Hatfield Marine Science Center (HMSC), Newport, OR, USA (additional details in Myers, 2022). pH
measurements from each manipulation tank were compared against a benchtop meter daily during the experiment. pH was
80 additionally compared against a reference sample in which pCO_2 and TCO_2 , in situ temperature and salinity were measured to
calculate pH on the NBS scale. pH (all tanks) and temperature (low and pre-treatment tanks) data were recorded every ten
minutes. The coefficient of determination (R^2) between measured benchtop pH data and the pH data calculated from
 $\text{pCO}_2/\text{TCO}_2$ measurements for header tanks in all experiments was consistently greater than 0.9 across all experiments (Figure
S2). Plug-flow style dissolution chambers were 2.5 cm ID, 12.7 cm long, schedule 40 PVC pipe, with 150 μm Nitex mesh on
85 either end of the upright chambers. Approximately 50 (± 0.1) grams of biomineral were used in each chamber, water was
initially added slowly and agitated to remove air bubbles prior to initiating flow. Flowrates were controlled to 17 (± 1.2) mL
 min^{-1} and experiments were run for three to four days to verify consistent alkalinity concentrations, as observed in trials
conducted to optimize the experiment (Myers, 2022). Using the inlet and outlet measured alkalinity concentrations from an
individual chamber ($\text{mol alkalinity L}^{-1}$), the measured flow rate (L d^{-1}), and the known amount of sample added ($\text{g}_{\text{mineral}}$), the
90 change in alkalinity in $\text{mol alkalinity g}_{\text{mineral}}^{-1} \text{d}^{-1}$ was recorded as (Eq. 1):

$$\Delta = \frac{([\text{alkalinity}]_{\text{outlet}} - [\text{alkalinity}]_{\text{inlet}})}{(\text{mineral amount})} * \text{flow rate}$$

The early diagenetic model (see below) uses a rate law in which the volumetric dissolution rate R_d (in $\text{mol}_{\text{mineral}} \text{m}^{-3} \text{sediment} \text{d}^{-1}$)
depends on the concentration of the mineral ($\text{mol}_{\text{mineral}} \text{m}^{-3} \text{sediment}$), the saturation state Ω (= ion concentration product/solubility
95 K_{sp}) and the dissolution rate constant k_d (d^{-1}) (Jourabchi et al., 2008):

$$R_d = k_d[\text{Mineral}](1 - \Omega)^n \quad (2)$$

In the experiment, the dissolving mineral is the entire solid phase, so that $g_{\text{mineral}} = g_{\text{solid phase}}$. Hence, the mineral concentration in $\text{mol}_{\text{mineral}}/g_{\text{solid phase}}$ is the molecular weight of the mineral (MW_{mineral}) and k_d can be calculated from the measured Δ as:

$$k_d = \frac{\Delta * MW_{\text{mineral}}}{\frac{dT_A}{dR} * (1 - \Omega)^n} \quad (3)$$

100 where dTA/dR is the mineral to alkalinity stoichiometric ratio in the dissolution reaction (+2 $\text{mol}_{\text{alkalinity}}/\text{mol}_{\text{mineral}}$; Table 3) and n is the rate order.

2.2. Carbonate Chemistry Measurements

Alkalinity

Open-cell alkalinity titrations were conducted over the course of the experiment, utilizing a two-point titration after Edmond
105 & Gieskes (1970), as modified by Waldbusser et al. (2011). A micrometer burette was used with a micro pH probe with
precisions of 0.002 mL and ± 0.02 , respectively. The probe was calibrated with NBS calibration standards (pH = 4, 7, and 10)
daily before being used to take measurements. As outlined in Waldbusser et al. (2011), Baker Analyzed Reagent grade 0.0995-
0.1005 normal HCl acid was used as the titrant with reagent grade NaCl to a final molar concentration of 0.7 to minimize gap
junction potentials during titrations. Analytical precision and accuracy were compared against triplicate titrations and Dickson
110 CRM batch 142, respectively, and found to have an average coefficient of variance of 0.07% and typically $\pm 16 \mu\text{mol}$ or less,
for precision and accuracy.

Dissolved Inorganic Carbon, $p\text{CO}_2$, and Carbonate Calculations

$p\text{CO}_2/\text{TCO}_2$ samples were collected in clean amber glass ~350 ml bottles preserved with 30 microliters of saturated HgCl_2 and
sealed with polyurethane-lined crimp-sealed metal caps. Analysis via the Burkator was carried out to measure $p\text{CO}_2$ and
115 TCO_2 via near-infrared detection following the procedure of Bandstra et al. (2006), modified for discrete samples as in Hales
et al. (2005), Barton et al. (2012), and Hales et al. (2017). Liquid and gas standards were employed to ensure the accuracy of
 $p\text{CO}_2/\text{TCO}_2$ measurements. The system can resolve the accuracy and precision of TCO_2 concentrations within $\pm 0.02\%$ and
2% for $p\text{CO}_2$ (Hales et al., 2017). Carbonate chemistry parameters were calculated using dissociation constants from Millero
(2010). Salinity was measured on the sample bottles following $p\text{CO}_2/\text{TCO}_2$ measurements via a salinometer (8400B “Autosal”
120 Laboratory Salinometer; Guildline Instruments, Sorrento, FL, US). Calculated carbonate chemistry variables presented in
association with the mineral dissolution experiments were computed using CO2calc and the K_1 and K_2 carbonic acid
dissociation constants from Lueker et al. (2000), K_{HSO_4} constants from Dickson (1990), sulfate constants from Morris & Riley
(1966), fluorine constants from Riley (1965), and boron constants from Uppström (1974).

2.3. Reactive transport modeling

125 A one-dimensional diagenetic model was developed to simulate the distribution of chemical species with depth in the sediment
over time. The model domain extended 20 cm deep into the sediment discretized into 50 intervals with a linearly increasing
grid size of 1 mm at the top and 1 cm at the bottom. The model included 13 state variables, representing chemicals involved
in the main early diagenetic reactions, and mineral dissolution/precipitation (Table 1). At the upper boundary, the

130 concentrations of solutes and fluxes of solids were imposed, while at the bottom boundary, no gradient conditions were imposed for all chemicals.

Table 1: State variables and their upper boundary conditions. Seawater compositions are computed with *aquaenv* (Hofmann et al. 2010) for the pressure, temperature and salinity given in Table 2.

Variable	Notation	Boundary condition
Oxygen	O ₂	0.2 mM
Sulfate	SO ₄ ²⁻	Seawater composition
Total sulfide	TS	0 mM
Dissolved inorganic carbon	DIC	equilibrium with a given alkalinity and pH 7.96
Total alkalinity	TA	1.9 mEq/L (measured at Yaquina Bay)
Calcium	Ca ²⁺	Seawater composition
Magnesium	Mg ²⁺	Seawater composition
Iron	Fe ²⁺	0 mM
Ammonium	NH ₄ ⁺	0 mM
Nitrate	NO ₃ ⁻	0.02 mM
Carbonate mineral (s)	CaCO ₃	F = 0 μmol cm ⁻² yr ⁻¹
Iron oxides (s)	Fe(OH) ₃	F = 1 μmol cm ⁻² yr ⁻¹
Iron sulfide (s)	FeS	F = 0 μmol cm ⁻² yr ⁻¹

135 The spatio-temporal evolution of the concentrations was described by an advection-diffusion equation with the addition of bioturbation and bioirrigation and the effect of reactions. Bioturbation was treated as a diffusive process (Boudreau, 1997) and bioirrigation was described as a non-local exchange between porewater at depth and the overlying water (Boudreau, 1984). The governing equations were:

$$\phi \frac{\partial C_i}{\partial t} = -\frac{\partial \phi u C_i}{\partial x} + \frac{\partial}{\partial x} \left(\phi (D_i + D_b) \frac{\partial C_i}{\partial x} \right) + \sum_r s_{ir} R_r + B_i \quad (4)$$

$$(1 - \phi) \frac{\partial C_j}{\partial t} = -\frac{\partial (1 - \phi) v C_j}{\partial x} + \frac{\partial}{\partial x} \left((1 - \phi) D_b \frac{\partial C_j}{\partial x} \right) + \sum_r s_{jr} R_r \quad (5)$$

140 where t is time, x is depth in sediment, ϕ is porosity, C_i (C_j) is the concentration of the solute i (in mol/volume pore fluid; solid j in mol/volume solid phase), u is the burial velocity for solutes (v for solids), D_i is the diffusion coefficient of the solute i , D_b is the depth bioturbation coefficient, R_r is the rate of production/consumption through reaction r (in mass/total volume and time), s is the stoichiometric coefficient for solute i or solid j in reaction r and B_i represents bioirrigation for solute i .

145 Porosity is constant with time but depth-dependent and exponentially decreases with depth reflecting steady state sediment compaction (Eqn. 6):

$$\phi_x = \phi_\infty + (\phi_0 - \phi_\infty)e^{-\frac{x}{\gamma}} \quad (6)$$

where ϕ_∞ and ϕ_0 are the porosities at infinite depth and at the top of the sediment, respectively, and γ is the e-folding distance for the porosity. Burial velocities were corrected for compaction (Berner, 1980):

$$u_x = v_\infty \frac{\phi_\infty}{\phi} \quad (7)$$

150

$$v_x = v_\infty \frac{(1 - \phi_\infty)}{(1 - \phi)} \quad (8)$$

Diffusion coefficients were calculated using the R package *marelac* after Boudreau (1997) at a given salinity, temperature and pressure (Soetaert et al., 2010) and corrected for tortuosity (Boudreau, 1996) so that:

$$D_i = \frac{D_{aq,i}}{1 - 2\ln(\phi_x)} \quad (9)$$

155 where D_{aq} is the molecular diffusion coefficient of solute i in solution. The bioturbation coefficient decreases exponentially with depth below the mixed layer,

$$D_b = D_{b0}e^{-\max(0, x-x_m)} \quad (10)$$

where x_m is the depth of mixed layer and D_{b0} is the bioturbation coefficient at the SWI, calculated as a function of water depth (Middelburg et al., 1997).

160 Bioirrigation is defined as an exchange between the surface layer and deeper layers of the sediment which was constrained by a bioirrigation coefficient exponentially decreasing with depth:

$$B_x = \alpha_0 e^{-\frac{x}{\varepsilon}} \phi_x (C_{i,0} - C_{i,x}) \quad (11)$$

where α_0 is the bioirrigation coefficient at the SWI, ε is the e-folding distance for bioirrigation and $C_{i,x}$ is the concentration of solute i at depth x .

Table 2: Environmental parameters

Parameter	Value
Salinity	35
Temperature	12 deg C
Overlying water pH _(total)	7.96
Pressure	1.013 bar
Seawater density	1.027 kg L ⁻¹
Sediment density	2.65 g cm ⁻³ (Boudreau, 1997)
Porosity decay with depth (γ)	5 cm (Rooze et al., 2016)
Porosity at the surface (ϕ_0)	0.8
Porosity at infinite depth (ϕ_∞)	0.6
Sedimentation at infinite depth (v_∞)	0.5 cm yr ⁻¹ (Middelburg et al., 1997)

Bioturbation mixing coefficient at the top (D_{b0})	30 cm ² yr ⁻¹ (Middelburg et al., 1997, water depth <5m)
Bioturbation e-folding distance	1 cm (Soetaert et al., 1996)
Depth of mixed layer (x_m)	4 cm (Rooze et al., 2016)
Bioirrigation rate constant at the top (α_0)	200 yr ⁻¹ (Wang & Van Cappellen, 1996; Meile & Van Cappellen, 2003)
Bioirrigation decay with depth (ϵ)	3.5 cm (Thullner et al., 2009)

165

Reactions considered in the model include organic matter mineralization with aerobic and anaerobic pathways, reoxidation of reduced species and mineral formation/dissolution reactions (Table 3). Each reaction has an impact on the total alkalinity (TA), which were taken from previous studies (Wolf-Gladrow et al., 2007; Dickson, 1981).

170 Organic matter is mineralized using O₂, NO₃⁻, iron (oxyhydr)oxides (FeOx) and SO₄²⁻ as terminal electron acceptors and modeled with Monod kinetics. The OM mineralization rate (R_C^0) was adjusted to yield a TA flux of ~1200 μmol cm⁻² yr⁻¹. This value was chosen to match open core top incubations measured in Yaquina Bay (Myers, 2022), with the resulting R_C^0 value being comparable to earlier measurements by D'Andrea & DeWitt (2009) (~200-460 μmol C cm⁻³ yr⁻¹), and higher than values reported for deeper coastal environments (Wang & Van Capellen, 1996; Krumins et al., 2013). Reactions considered further include nitrification, iron oxidation, sulfide oxidation with O₂ and FeOx, and iron sulfide oxidation, as well as the
175 formation/dissolution of CaCO₃ minerals and FeS (Table 3). Following Rooze et al. (2020), at each time step, DIC is speciated to calculate CO₃²⁻ concentration and pH from DIC and alkalinity, assuming seawater concentrations of borate and negligible contributions of phosphate, silicic acid, nitrite and hydrogen fluoride and other acids or bases.

Table 3: Reactions included in the model. Benthic primary production is assumed to be negligible, and organic matter has
180 Redfield stoichiometry (a = 1, b = 16/106, c = 1/106). Alkalinity production or consumption in each reaction is determined following Wolf-Gladrow et al. (2007). The total mineralization rate is defined as $R_C = R_C^0 \cdot \exp(-x/\gamma_{om})$ after Wang & Van Cappellen (1996) where γ_{om} is e-folding distance for OM mineralization.

Reaction	dTA/dR	dDIC/dR	Reaction rate
<i>Organic Matter Mineralization</i>			
(CH ₂ O) _a (NH ₃) _b (H ₃ PO ₄) _c + a O ₂ => a CO ₂ + b NH ₄ ⁺ + c HPO ₄ ²⁻ + (-b+2c) H ⁺ + a H ₂ O	b - c	a	$R_{O_2}^m = RC [O_2] / (K_{O_2}^m + [O_2])$
(CH ₂ O) _a (NH ₃) _b (H ₃ PO ₄) _c + 4a/5 NO ₃ ⁻ + (4a/5 + b - 2c) H ⁺ => a CO ₂ + b NH ₄ ⁺ + c	4a/5 + b - c	a	$R_{NO_3}^m = (RC - R_{O_2}^m) [NO_3] / (K_{NO_3}^m + [NO_3])$

$\text{HPO}_4^{2-} + 2a/5 \text{N}_2 + 7a/5 \text{H}_2\text{O}$			
$(\text{CH}_2\text{O})_a(\text{NH}_3)_b(\text{H}_3\text{PO}_4)_c + 4a \text{Fe}(\text{OH})_3 + (8a+b-2c) \text{H}^+ \Rightarrow a \text{CO}_2 + b \text{NH}_4^+ + c \text{HPO}_4^{2-} + 4a \text{Fe}^{2+} + 11a \text{H}_2\text{O}$	$b - c + 2*4a$	a	$R_{\text{FeOx}}^m = (RC - R_{\text{O}_2}^m - R_{\text{NO}_3}^m) [\text{FeOx}] / (K_{\text{FeOx}}^m + [\text{FeOx}])$
$(\text{CH}_2\text{O})_a(\text{NH}_3)_b(\text{H}_3\text{PO}_4)_c + a/2 \text{SO}_4^{2-} + (a/2 + b - 2c) \text{H}^+ \Rightarrow a \text{CO}_2 + b \text{NH}_4^+ + c \text{HPO}_4^{2-} + a/2 \text{HS}^- + a \text{H}_2\text{O}$	$b - c + a$	a	$R_{\text{SO}_4}^m = (RC - RC_{\text{O}_2}^m - R_{\text{NO}_3}^m - R_{\text{FeOx}}^m) [\text{SO}_4] / (K_{\text{SO}_4}^m + [\text{SO}_4])$
<i>Secondary reactions</i>			
$\text{NH}_4^+ + 2\text{O}_2 \Rightarrow \text{NO}_3^- + 2\text{H}^+ + \text{H}_2\text{O}$	-2	0	$R_{\text{NH}_4}^o = k_{\text{NH}_4}^o [\text{NH}_4][\text{O}_2]$
$\text{Fe}^{2+} + 5/2 \text{H}_2\text{O} + 1/4 \text{O}_2 \Rightarrow \text{Fe}(\text{OH})_3 + 2\text{H}^+$	-2	0	$R_{\text{Fe}}^o = k_{\text{Fe}}^o [\text{Fe}^{2+}][\text{O}_2]$
$\text{HS}^- + 2\text{O}_2 \Rightarrow \text{SO}_4^{2-} + \text{H}^+$	-2	0	$R_{\text{TS}}^o = k_{\text{TS}}^o [\text{TS}][\text{O}_2]$
$\text{HS}^- + 8\text{Fe}(\text{OH})_3 + 15\text{H}^+ \Rightarrow \text{SO}_4^{2-} + 8\text{Fe}^{2+} + 20\text{H}_2\text{O}$	+14	0	$R_{\text{TS}}^{of} = k_{\text{TS}}^{of} [\text{TS}][\text{FeOx}]$
$\text{FeS} + 2\text{O}_2 \Rightarrow \text{Fe}^{2+} + \text{SO}_4^{2-}$	0	0	$R_{\text{FeS}}^o = k_{\text{FeS}}^o [\text{FeS}][\text{O}_2]$
<i>Mineral precipitation/dissolution</i>			
$\text{FeS} + 2\text{H}^+ \rightleftharpoons \text{Fe}^{2+} + 2\text{HS}^-$	+2 (dis)/-2 (prec)	0	$\Omega_{\text{FeS}} = [\text{Fe}][\text{TS}] / ([\text{H}^+] K_{\text{FeS}}^{sp} \rho_{\text{sw}}^2)$ $R_{\text{FeS}} = \begin{cases} k_{\text{FeS}}^f (\Omega_{\text{FeS}} - 1), & \text{if } \Omega_{\text{FeS}} \geq 1 \\ k_{\text{FeS}}^d [\text{FeS}] (1 - \Omega_{\text{FeS}}), & \text{if } \Omega_{\text{FeS}} < 1 \end{cases}$
$\text{CaCO}_3 \rightleftharpoons \text{Ca}^{2+} + \text{CO}_3^{2-}$	+2 (dis)/-2 (prec)	+1 (dis)/-1 (prec)	$\Omega_{\text{M}} = [\text{Ca}^{2+}][\text{CO}_3] / (K_{\text{M}}^{sp} \rho_{\text{sw}}^2)$ $R_{\text{M}} = \begin{cases} k_{\text{M}}^f (\Omega_{\text{M}} - 1), & \text{if } \Omega_{\text{M}} \geq 1 \\ k_{\text{M}}^d [\text{Mineral}] (1 - \Omega_{\text{M}})^n, & \text{if } \Omega_{\text{M}} < 1 \end{cases}$

185 **Table 4: Reaction constants.** Carbonate dissolution rate constants determined experimentally in this study are described in the results.

Reaction parameters	Value	Source
R_C^0	400 $\mu\text{mol C cm}^{-3} \text{ yr}^{-1}$	Calibrated to match experimental TA data
γ_{om}	5 cm	(Wang & Van Capellen, 1996)
$K_{O_2}^m$	0.02 mM	(Rooze et al., 2016)
$K_{NO_3}^m$	0.004 mM	(Rooze et al., 2016)
K_{FeOx}^m	130 mM	(Rooze et al., 2016)
$K_{SO_4}^m$	1.6 mM	(Rooze et al., 2016)
$k_{NH_4}^o$	5000 $\text{mM}^{-1} \text{ yr}^{-1}$	(Rooze et al., 2016)
k_{Fe}^o	140 000 $\text{mM}^{-1} \text{ yr}^{-1}$	(Rooze et al., 2016)
k_{TS}^o	160 $\text{mM}^{-1} \text{ yr}^{-1}$	(Rooze et al., 2016)
k_{TS}^{of}	6 $\text{mM}^{-1} \text{ yr}^{-1}$	(Rooze et al., 2016)
k_{FeS}^o	300 $\text{mM}^{-1} \text{ yr}^{-1}$	(Rooze et al., 2016)
K_{FeS}^{sp}	$10^{-2.2} (\text{mol/kg})^2$	Wang & Van Capellen (1996)
k_{FeS}^f	$6 \cdot 10^{-6} \text{ mol g}^{-1} \text{ yr}^{-1}$	Wang & Van Capellen (1996)
k_{FeS}^d	10^{-3} yr^{-1}	Wang & Van Capellen (1996)
$K_{calcite}^{sp}$	$4.31 \cdot 10^{-7} (\text{mol/kg})^2$	f(salinity) (Mucci, 1983)
$K_{aragonite}^{sp}$	$6.82 \cdot 10^{-7} (\text{mol/kg})^2$	f(salinity) (Mucci, 1983)
$k_M^f(\text{calcite})$	10^4 mM yr^{-1}	(Luff & Wallman, 2003)
$k_M^f(\text{aragonite})$	10^5 mM yr^{-1}	(Luff & Wallman, 2003)
$k_M^d(\text{calcite})$	5.05 yr^{-1}	(this study)
$k_M^d(\text{aragonite})$	5.05 yr^{-1}	(this study)

Application

To establish initial conditions, each simulation was first run to steady state with a zero-deposition flux of calcium carbonate. After that, 8 or 16 weight % (weight of mineral/ total weight of the sediment) CaCO_3 were added to the top 2 cm of sediment, representing the sudden, one-time addition of the buffering carbonate minerals. To assess the magnitude and temporal evolution of the impact of the alkalization experiment on sediment biogeochemistry and benthic fluxes, transient simulations using these new initial conditions were then carried out, covering periods of up to 50 years.

190

We tested the impact of several environmental factors on the buffering effect of carbonate mineral additions to surficial sediments in comparison to the baseline addition scenario (Table 5). Specifically, we lowered the overlying water pH from 7.96 to 7.8, representing the impact of ocean acidification. The depth of the bioturbated layer affects the distribution of minerals added, which may impact the effectiveness of the extent of buffering; it was varied from the relatively shallow mixing depth of 4cm in the baseline implementation to the global average depth of bioturbation of 10 cm (Boudreau, 1997). We also varied the deposition flux of iron (oxy)hydroxides from 1 to 10 $\mu\text{mol}/\text{cm}^2/\text{yr}$ to assess differences between sites that vary in the extent of coupling between Fe, S and C cycles. And as the rate of organic matter mineralization has a major impact on sediment redox conditions and porewater composition, R_c^0 was varied from the baseline value of 400 to 200 $\mu\text{mol cm}^{-3} \text{ yr}^{-1}$ to represent environments with different O_2 penetration depths. Finally, simulations were carried out to assess the impact of the type of mineral in the system (calcite vs. aragonite). The spin-up simulations were specific to each model run, and as such represent the environmental conditions (e.g., high vs. low rate of mineralization, or a setting in which either aragonite or calcite are the only carbonate mineral considered). Simulations were performed in R using ReacTran package (Soetaert & Meysman, 2012).

205

Table 5. Sensitivity analyses. Rows represent the scenario tested. Columns show the parameters changed.

Scenario/Parameter	R_c^0 ($\mu\text{mol cm}^{-3} \text{ yr}^{-1}$)	$\text{Fe}(\text{OH})_3$ flux ($\mu\text{mol cm}^{-2} \text{ yr}^{-1}$)	Depth of mixed layer (cm)	Mineral type	pH in overlying water
Baseline	400	1	4	Calcite	7.96
Low OW pH	400	1	4	Calcite	7.8
High ML	400	1	10	Calcite	7.96
High Fe Flux	400	10	4	Calcite	7.96
Low R_c^0	200	1	4	Calcite	7.96
Aragonite	400	1	4	Aragonite	7.96

CO₂ drawdown

We estimated the potential impact of enhanced sediment TA fluxes on atmospheric CO_2 uptake in an enclosed bay using a simple box model. Water exchange is due to the tides, and riverine impact was assumed negligible, reflecting for example, the strong coupling of Yaquina Bay with the coastal ocean (Brown & Ozretich, 2009).

To estimate the order of magnitude of the impact of mineral additions on CO_2 uptake, we used a simple mass balance approach similar to Brenner et al. (2016), which focuses on the ratio of DIC over TA as the controlling factor of CO_2 uptake (Egleston et al., 2010). We calculated the mass balance for TA and DIC prior to and 2 years after the mineral addition and attributed the difference in atmospheric CO_2 exchange to the mineral addition. As described in detail in the Supplementary Information, we first estimated tidal water exchange from water level measurements. The characteristics of the inflowing marine (alkalinity =

215

2.2 mmol L⁻¹, pH = 8.0) and outflowing bay water (alkalinity = 1.9 mmol L⁻¹, pH = 7.96) were chosen to be representative of a coastal setting such as Yaquina Bay and were informed by unpublished temperature, salinity, and pH data (pers. comm. G. Waldbusser). Combined with our modeled benthic exchange fluxes, this allowed us to estimate the alkalinity production in the water column (R_{TA} ; Eq. 12). At steady state, the alkalinity balance is given by:

$$F_i^{TA} - F_o^{TA} + F_b^{TA} + R^{TA} = 0 \quad (12)$$

where F_i^{TA} is the tidal influx of TA as a function of water exchange rate and oceanic TA ($F_i^{TA} = v[TA_{ocean}]$), F_o^{TA} is the tidal outflux of TA from the Bay to the ocean as a function of tidal velocity and TA of the study site ($F_o^{TA} = v[TA_{site}]$), F_b^{TA} is the benthic flux computed by the reactive transport model and R^{TA} is the areal net rate of water column TA production/consumption resulting from primary productivity and respiration.

Next, we used the alkalinity and pH to calculate the DIC and converted the net alkalinity production rate into net DIC production in the water column, assuming Redfield stoichiometry. This finally allowed us to estimate the CO₂ exchange flux (F_a^{DIC}) necessary to close the DIC mass balance for benthic fluxes with and without mineral additions to quantify its effect on ocean CO₂ uptake.

The mass balance for DIC is given by:


$$F_i^{DIC} - F_o^{DIC} + F_b^{DIC} + F_a^{DIC} + R^{DIC} = 0 \quad (13)$$

where F_i^{DIC} is the tide-driven influx of DIC ($F_i^{DIC} = v[DIC_{ocean}]$), F_o^{DIC} is the outflux of DIC ($F_o^{DIC} = v[DIC_{bay}]$), F_b^{DIC} is the benthic flux of DIC, F_a^{DIC} is the atmospheric CO₂ uptake (negative values indicate outgassing) and R^{DIC} is the net areal reaction rate producing/consuming DIC.

3. Results & Discussion

3.1. Mineral Dissolution Rates

Dissolution rates of minerals were highest under highly undersaturated conditions (Figure 1). Dissolution rates are approximately proportional to the degree of saturation. Thus, the dissolution rate constant k_d for each mineral was estimated using Eqns. (1) and (2) with a dissolution order (n) of 1. However, we acknowledge the large variability in mineral dissolution dynamics (especially n) in previous studies (see e.g., Adkins et al., 2021; Subhas et al., 2018), which is a critical factor for the evolution of the dissolution rate over time. Our results are in line with previous data synthesis studies focused on mineral dissolution (e.g., Morse et al., 2007; Jourabchi et al., 2005; 2008). Kinetic constant for aragonite dissolution was previously reported as 5 yr⁻¹ (Luff & Wallmann, 2003) which was comparable to the 5.2 yr⁻¹ derived from our measurements. For $n=1$, Jourabchi et al. (2005), reported a dissolution rate constant of 3.65 yr⁻¹ while our estimation for bio-calcite was 5 yr⁻¹. Measured dissolution rates did not show significant differences between two mineral types (t-test with p-value > 0.05) therefore, a kinetic constant for both minerals was calculated as 5.05 yr⁻¹.



250 **Figure 1:** Saturation state for aragonite and calcite vs. their respective dissolution rates (mmol alkalinity g_{solid}⁻¹ d⁻¹). Line shows the fit with n=1. R²: coefficient of determination. p-val: statistical significance of the fit. Shaded area represents 95% confidence interval.

255 3.2. Early Diagenetic Modeling

To assess the impact of sediment alkalization through the addition of carbonate minerals to the sediment, early diagenetic model simulations were carried out that describe the spatio-temporal distribution of porewater and solid phase constituents. Prior to the addition of minerals, the simulated porewater profiles show sequential depletion of electron acceptors with depth, with O₂, NO₃⁻ and reactive FeOx only present in the top 5 cm (solid lines in Figure 2). SO₄²⁻ is present throughout the depth range and only decreases by about 2.5 mM. As a result of mineralization reactions, the concentrations of reduced porewater

260

constituents (TS, NH_4^+ , Fe^{2+}) increase with depth (Figure 2). Reduced dissolved iron (Fe^{2+}) peaked around 1 cm then decreased with depth due to the precipitation of FeS, which in our baseline simulation limited the build-up of hydrogen sulfide (TS) and dissolved reduced iron (Fe^{2+}). These porewater profiles reflect the sequential use of electron acceptors in the mineralization of organic matter, through aerobic respiration and anaerobic pathways such as denitrification, dissimilatory iron and sulfate reduction. The dominant OM mineralization pathway was SO_4^{2-} reduction (78.9%) followed by aerobic mineralization (17.2%), denitrification (3.9%) and iron reduction (0.06%).

DIC and TA show pronounced concentration increases with depth, reflecting OM mineralization (Figure 2 i,j). The pH profile exhibits a rapid decrease in the top centimeter to about 7, showing the impact of CO_2 production without the concurrent consumption of protons through aerobic mineralization (Table 3; Jahnke et al., 1994; Green et al., 2013). Below the oxic layer, active anaerobic OM mineralization pathways such as iron and sulfate reduction can generate considerable amounts of TA if the reaction products are not oxidized (Thomas et al., 2009). This was observed in the buildup of reduced species such as TS (discussed below) which limited the drop in pH and increased saturation state (from 0.5 to > 0.8) in the deeper layers of the sediment (Figure 2 k,l). pH values were in the lower part of the range reported in Krumins et al. (2013), reflecting the lack of carbonates buffering the pH prior to the mineral additions. Sulfate reduction produced the majority of the TA and DIC (Figure 3), consistent with the findings of Krumins et al. (2013), Brenner et al. (2016) and Gimenez (2018). Aerobic mineralization produced considerable amounts of DIC in line with previous work (Thamdrup & Canfield, 2000) while generating less TA than anaerobic processes and mineral dissolution (Figure 3), leading to a low porewater pH (Soetaert et al., 2007; Hofmann et al., 2011). DIC and TA production by denitrification and in particular dissimilatory iron reduction was substantially less. With a high $d\text{TA}/dR$ value (Table 3), TS oxidation with FeOx also contributed to alkalinity production. Consumption of TA was dominated by secondary redox reactions that oxidize reduced species with O_2 . Due to fast kinetics, Fe^{2+} oxidation with O_2 , nitrification and precipitation of FeS were the major TA consuming reactions. Brenner et al. (2016) identified sulfide oxidation and nitrification as the dominant alkalinity consuming processes in the shallow water sediments in Southern North Sea. This discrepancy may be due to the potentially substantial impact of sulfide oxidation on alkalinity dynamics in organically rich near shore sediments (Krumins et al. 2013). The oxidation of reduced products of organic matter mineralization such as ammonium, hydrogen sulfide and dissolved Fe^{2+} with O_2 all lower alkalinity, albeit using different amounts of molecular oxygen. However, the oxidation of hydrogen sulfide with iron oxide as electron donor produces a large amount of alkalinity, that either accumulates if the reduced iron produced is captured in the sediment or is removed if the Fe^{2+} subsequently reacts with O_2 in the oxic zone (see Table 3). Coupling of SO_4^{2-} reduction with FeS precipitation has been shown to be an important mechanism for TA burial on continental shelves (Gustafsson et al., 2019). However, in our simulations, FeS formation was limited by Fe availability, which led to the buildup of hydrogen sulfide in the porewater.

Effect of carbonate mineral additions

After adding carbonate minerals, the concentration profiles of primary redox species largely remained unchanged (Figure 2). This is because the organic matter mineralization rate expressions (Table 3) do not contain dependencies on the carbonate system or pH, as is common in early diagenetic models (e.g., Jourabchi et al., 2008; Morse & Mackenzie, 1990). Mineral

295 addition slightly decreased Fe^{2+} while increasing FeS concentrations. This is due to TA production due to carbonate dissolution, which elevates the pH and increases the saturation state of FeS (Table 3), promoting FeS formation which eventually can lead to the long-term removal of reduced sulfur through pyrite burial (Hu & Cai, 2011a)

The added carbonate minerals started to dissolve in our model, producing 1 DIC and 2 TA per carbon (Burdige et al., 2010; Zeebe & Wolf-Gladrow, 2001), so that both the pH (Figure 2k) and porewater saturation state (Figures 2l and 4B) initially
 300 increased (Cai et al., 2011). After 2 years, CO_2 production still lowered pH in the oxic layer and prevented full saturation. However, Ω was always above 0.9 and reached 1 at and below a depth of approximately 2.5 cm and subsequent decrease was observed below 12 cm. Following the addition of carbonates, the extent of the undersaturated conditions was constrained to the top 2 cm only, while in the natural buffering scenario reported in Krumins et al. (2013) porewater remained undersaturated in the top 10 cm. This demonstrates the potential of OAE in mitigating surface sediment acidification and providing potential
 305 benefits for calcifiers living in the upper sediment layers. As mineral addition counteracts porewater acidification, it generates biogeochemical conditions conducive for the survival of juvenile bivalves.





Figure 2. Simulated baseline porewater profiles of (a) O_2 , (b) NO_3^- , (c) FeOx, (d) SO_4^{2-} , (e) NH_4^+ , (f) Fe^{2+} , (g) TS, (h) FeS, (i) DIC, (j) TA, (k) pH and (l) Ω_{calcite} with 8% calcite addition. The solid lines denote the steady state profiles prior to the

310 addition of calcite to the top 2 cm, while the dashed line indicate the simulated profiles 2 years after the mineral addition
(some are coinciding with the solid lines and hence not visible). Dissolved constituents are in mmol L^{-1} porewater, while
solids are given in $\mu\text{mol cm}^{-3}$ solid phase.

Our results are in line with literature showing that the main alkalinity producing processes were calcite mineral dissolution,
315 sulfate reduction, denitrification and sulfide oxidation with iron oxides (Krumins et al., 2013; Berelson et al., 2007). Although
organic matter mineralization reactions generate alkalinity, they have lower dTA/dDIC ratios than mineral dissolution on a per
carbon basis (Table 3), and the TA producing mineralization reactions are mostly coupled to TA consuming reoxidation
reactions. Therefore, the net effect of (some) mineralization pathways on TA production can be limited (Krumins et al., 2013).
For example, the amount of TA generated by benthic denitrification was almost balanced out by NH_4^+ oxidation (Figure 3)
320 which results in the small effect of N cycling on alkalinity. This is in line with previous studies showing strong coupling
between oxidation/reduction of N in the marine environment (Middelburg et al., 1996) and its weak impact on buffering
capacity of the overlying water (Hu & Cai, 2011b). Since reoxidation reactions are dependent on the availability of O_2 ,
enhanced biological mixing can also increase TA consumption as previously demonstrated by Rao et al. (2014) for Fe^{2+}
oxidation by O_2 . Although total production of TA was around $1300 \mu\text{mol cm}^{-2} \text{yr}^{-1}$ (Figure 3, baseline with bio-calcite), total
325 TA fluxes from bio-calcite were slightly lower, which points to the importance of consumption of TA in the oxic layer as
previously reported in Krumins et al. (2013). Finally, before mineral addition, we observed a small production of calcite at the
top of the sediment due to oversaturated conditions in overlying water. After the addition however, added mineral dissolved
in deeper layers which led to net dissolution and generation of alkalinity in the sediment column.




330 **Figure 3.** DIC and TA production (positive) and consumption (negative) by different early diagenetic processes integrated over the entire model domain for steady state ($t = 0$; before addition) and 2 years after the addition.

Temporal evolution of buffering with different minerals and amounts

335 Longer-term simulations (50 yrs) were performed to investigate the evolution of the buffering with time, considering different amounts and types of minerals added. We focused on the biogeochemical conditions in the top 10 cm of sediment, because the surficial sediment is most relevant for juvenile bivalves impacted by acidification. Figure 4 shows the temporal evolution of TA and saturation state following the addition of minerals to the steady state conditions at $t=0$. Depth-averaged TA concentrations in top 10 cm peaked around the 2nd year after addition, with a significant drop starting around year 10 for the 8% mineral addition and around year 15 for the 16% addition and then return to steady state levels after more than 30 years (Figure 4A). Doubling the amount of mineral added generated slightly higher TA and extended the length of the peak buffering period. The simulation of aragonite addition produced higher alkalinity concentrations than that of calcite by approximately

340

0.3 mM at the peak time reflecting its higher solubility. However, TA levels in aragonite application dropped faster after approximately 10 and 15 yrs and approached the initial pre-mineral application levels after about 30 and 40 years, for 8% and 16% additions, respectively. In contrast, following the addition of calcite, TA concentrations remained elevated over longer time periods. Averaging over 4 cm yielded the same patterns (with slightly lower TA concentrations) as shown in Figure 4 (not shown).



350 **Figure 4.** Temporal evolution of depth-averaged TA after addition of bio-aragonite and bio-calcite (A), and saturation states
355 (B) in baseline simulation. Concentrations and saturation states are averaged over the top 10 cm of sediment. Colors show
the type of mineral added. Solid and dashed lines show the addition of 16% and 8% respectively.

Being less stable than calcite, aragonite had a slightly faster dissolution rate constant and a larger solubility product (Burdige et al., 2010). Therefore, it generated higher TA concentrations and benthic fluxes early on. Because of its higher solubility, it initially took aragonite slightly longer to reach saturation than calcite (i.e., orange curve reaches maximum slightly after the

blue one). As the rate of dissolution is set to be proportional not only to the degree of undersaturation but also the carbonate mineral concentration (Eq. 2), doubling the mineral addition had a more pronounced effect for aragonite which was approaching saturation more slowly than calcite (Figure 4B). Burial of carbonate minerals below 10 cm led to a faster decrease in TA in aragonite due to its higher dissolution. Initial conditions in 50 years simulations with calcite had higher saturation state than aragonite implementation therefore, after the impact of addition got buried, calcite levelled at higher saturation state and TA than aragonite.

Sensitivity analyses

The impact of mineral additions on the generation of alkalinity depends not only on the type of mineral added, but also on which early diagenetic processes dominate (Figure 3). The relative magnitude of these processes depends on a number of factors, including the rate of organic matter mineralization, the extent of bioturbation, the deposition fluxes of potential electron acceptors such as iron oxyhydroxides, and the composition of the overlying water. Varying these factors in our model, we explore their impact on both the environmental conditions in the surficial sediment as well as the impact on benthic alkalinity fluxes.

In our simulations, under all scenarios, mineral saturation states reached similar values at the end of the 2 years with a low R_c^0 scenario generating the highest values (blue dots in Figure 5A). However, saturation was reached fastest in the deeper mixed layer scenario which then decreased by years 1 and 2. Baseline, high FeOx flux and low R_c^0 scenarios showed a similar time course, while the addition of aragonite instead of calcite exhibited the slowest increase in saturation state (Figure 5A). Depth-averaged TA concentrations were highest following the addition of aragonite (Figure 5B). Deepening the bioturbation layer did not have much of an impact, the high FeOx flux scenario produced slightly lower TA than the baseline and the lowest amounts of alkalinity were produced by the low R_c^0 scenario. pH increased the most and fastest in the aragonite implementation, followed by low R_c^0 conditions (Figure 5C). Although the deep mixed layer scenario increased pH faster than the baseline and high FeOx flux scenarios, these three scenarios converged to a similar pH as the impact of mineral dissolution developed. The highest benthic TA fluxes were generated during the dissolution of aragonite while the lowest values were produced in the low R_c^0 scenario (Figure 5D). Baseline and deeper bioturbation layer scenarios produced very similar results while high FeOx flux scenario produced slightly lower TA fluxes. Lowering overlying water pH led to decreases in saturation state and pH while increasing concentration and flux of alkalinity.




Figure 5. Buffering response after carbonate mineral additions for a range of environmental conditions. Concentrations are averaged over the top 10cm. Scenarios listed on the x axis (from left to right) represent: baseline simulation; baseline with lower overlying water pH representing ocean acidification scenario (7.8); baseline with deeper mixed layer (10 cm); baseline with higher Fe flux ($10 \mu\text{mol cm}^{-2} \text{yr}^{-1}$); baseline with lower R_c^0 ($200 \mu\text{mol cm}^{-3} \text{yr}^{-1}$); baseline with aragonite instead of calcite.

A lower pH in the overlying water that represents ocean acidification led to a lower pH in the porewater and less saturated conditions in surface sediments (Figure 5). This promoted mineral dissolution and therefore led to increased TA production and flux to the overlying water. This demonstrates the sensitivity of buffering towards the overlying water conditions. Deeper bioturbation led to faster downward mixing and subsequent dissolution of the added mineral which then led to rapid increase in TA, pH and saturation state (also seen in dpH/dTA , Figure S4). Higher mixing led to faster transport of minerals inducing enhanced dissolution in year 0.5 (i.e., Figure 5, higher buffering 0.5 than 2 years after mineral addition). However, faster transport extended the depth of buffering and diluted dissolution products, which caused a slight decrease in pH and saturation

state after 2 years at the top 10 cm (Figure 5). Increased FeOx input to the sediment led to enhanced FeOx reduction which produced TA. However, increased reoxidation of Fe²⁺ led to decreased availability of O₂ and consumption of TA. This balanced out the additional TA produced through FeOx reduction and led to an additional net consumption of ~46 μmol cm⁻² yr⁻¹ of TA. Overall, increased iron oxide supply led to slight decreases in TA concentrations and fluxes, and small increases in pH showing an effect similar to decreasing the OM mineralization rate (Figure 5). TA production and flux was significantly lower in low OM reaction rate scenario (Figure 5D) which showed the critical impact of OM mineralization on TA dynamics. The lower rate of OM mineralization decreases TA production, both through the reduction of overall mineralization rates, and the reduction of the relative contribution of anaerobic mineralization pathways like dissimilatory sulfate reduction. The low rate of aerobic mineralization also led to reduced H⁺ production, which in turn slows carbonate dissolution (Morse & Mackenzie, 1990). However, even with lower dissolution, this situation led to higher pH than the baseline suggesting that the artificial buffering might not be needed to maintain saturated conditions in environments where OM reactivity is low. In deeper layers where the surficial mineral addition is not as effective, TA and mineral saturation states will be lower due to lower anaerobic TA production. Notably, the effect of changing the OM mineralization rate is not linear, and halving the rate reduced the benthic alkalinity flux by a factor of about 2.5. In our simulations, the lower OM mineralization rate led to depth-integrated reduction by 29% for aerobic mineralization, 19% for denitrification, 55% for sulfate reduction and 380% for mineral dissolution. Aside from the impact on the partitioning of OM mineralization between metabolic pathways, this mainly reflects that changes in mineralization rates lead to differences in the porewater conditions, which in turn alter the rate of mineral dissolution/precipitation. Due to the higher solubility and hence larger degree of undersaturation, initial aragonite dissolution was faster than calcite. Hence, aragonite produced the largest TA in 2 years, generated highest benthic TA fluxes, and increased the pH the most. The addition of aragonite produced more buffering than the scenario with lower OM mineralization which shows the potential of artificial buffering overcoming the acidification impact due to OM mineralization.

3.3 Limitations and challenges

Our model demonstrates the potential for coastal sediment alkalization to improve the biogeochemical conditions for juvenile calcifying organisms living in surficial sediments. However, alkalinity fluxes are sensitive to overlying water conditions, temporal changes in overlying water should be considered. These changes include the impact of short-term diel patterns and small-scale spatial heterogeneity on benthic fluxes (Gadeken et al., 2023), seasonal changes such as temperature (Rao et al., 2014) and long-term changes such as anthropogenic impacts (Pacella et al., 2018; e.g. eutrophication, Cai et al., 2011). In addition, in particular in near-shore shallow water environments, one may have to take into consideration the input of groundwater (Cyronak et al., 2013a), advective flow through sand ripples (Huettel et al., 1998) and other factors driving advective flow in marine sediments (Santos et al., 2012), which can impact sediment biogeochemistry and alkalinity fluxes. Furthermore, the prediction of the time course of buffering by mineral dissolution may need to consider dissolution rate expressions that account for changes in mineral surface areas, rather than bulk concentrations (Morse et al., 2007), and the

potential for washing away or enhancing dissolution (Meysman & Montserrat, 2017) of minerals deposited to the sediment in
430 high-energy environments.

Our results indicate a lasting buffering effect of mineral additions. This finding depends on the kinetics of the mineral
dissolution rate, in particular the reaction order. Given the range of observations (Adkins et al., 2021; Subhas et al., 2018), the
temporal evolution of the dissolution rates in the field needs to be explored further. In addition, our model assumes that the
minerals are mixed into the upper 2 cm of sediment, matching recent field manipulation experiments in Yaquina Bay. However,
435 an important consideration is how the minerals are applied to the surface sediment. For example, a simple ‘top-dressing’
approach would involve less disturbance to the sediment and less labor than surface raking, but may be washed away more
easily, and react less readily with the metabolic acids produced in the underlying sediment. Thus, optimal mineral application
and subsequent effects clearly requires field trials across a range of hydrodynamic settings

3.4 Potential impact of benthic fluxes on marine CO₂ uptake

440 Benthic alkalinity fluxes are critical for water column C dynamics especially in shallow water environments where water
column and atmospheric interactions are more sensitive to processes occurring in the sediment (Brenner et al., 2016). The
significant decrease in shells due to overharvesting has likely altered alkalinity cycling in estuarine waters, such as Chesapeake
Bay (Waldbusser et al., 2013). One outcome of this benthic-pelagic connection is the impact of benthic fluxes on oceanic CO₂
uptake which was estimated to have significant influence in the North Sea (Thomas et al., 2009), and the potential for shallow
445 water sediment alkalization to alter overlying water.

We estimated the potential impact of enhanced sediment TA fluxes on atmospheric CO₂ uptake in Yaquina Bay, OR (USA)
using the diagenetic model output (baseline scenario) in a simple box model at the peak buffering period (~2 years after
addition). The bay was found to be net heterotrophic, with tidal exchange and net respiration in the water column impacting
air-sea CO₂ flux the most. The impact of sediment buffering on the air-sea CO₂ was estimated as the difference in atmospheric
450 CO₂ fluxes between before and after the mineral addition. Mineral addition resulted in a TA flux enhancement of ~126 μmol
cm⁻² yr⁻¹ and a decrease in CO₂ flux by approximately 58 μmol cm⁻² yr⁻¹, which was comparable to previous studies (Brenner
et al., 2016). Scaling to a 10 m water depth (or per liter of water), this stoichiometrically estimated marine carbon dioxide
removal (mCDR) due to the application and subsequent dissolution of carbonate minerals by the sediments results in a similar
amount of mCDR as the current estimates of the anthropogenic C (C_{anth}) concentration in the US Pacific coast waters, 30-60
455 μmol L⁻¹ (Feely et al. 2016, Pacella et al. 2024). It is important to note three key points in this estimate: (1) the uptake of C_{anth}
by marine waters from anthropogenic activities results in an increase of DIC without any concurrent change in alkalinity, thus
acidification. (2) Alkalinity enhancement shifts the carbonate speciation and allows increased uptake of atmospheric CO₂ by
marine waters. Thus, ocean alkalinity enhancement increases the total amount of C_{anth} in seawater. However, while C_{anth} and
DIC both increase with OAE, (3) the mineral dissolution due to sediment respiration results in an increase in the TA:DIC ratio,
460 thus providing the mCDR benefit of atmospheric removal as well as mitigating existing acidification impacts. The annual
mCDR effect modelled here, while notable compared to the total anthropogenic impact, is however miniscule compared to the

total air-sea exchange flux, which is driven by tidal exchange of DIC and alkalinity between the bay and the coastal ocean and biogeochemical processes in the water column. The signal to noise of this potentially significant effect also highlights the existing measurement challenges in accurately discerning OAE effects within coastal systems.

465 **4. Conclusions**

Our study successfully combines lab experiments and modeling to investigate the impact of coastal alkalinity enhancement, points to critical aspects to be considered and presents the potential of carbonate additions as a mitigation strategy for the effects of climate change in coastal settings. We demonstrated that addition of minerals increased pH and saturation state of the sediment over the timespan of a few months and had the potential to persist over 30-40 years after which subsequent additions might be needed to sustain buffering. During the peak buffering period, porewater in the top 10 cm of the sediment was almost fully saturated with respect to the mineral added, which implies favorable conditions for benthic calcifying organisms. We showed that two main pathways of producing TA in the sediment were sulfate reduction and mineral dissolution with sulfate reduction having a larger impact. We also demonstrated that the type of mineral was an important factor affecting benthic fluxes and that the application of an increased amount of minerals was effective in extending the duration rather than the strength of the buffering.

Our analysis suggests that buffering through mineral additions to coastal sediments can establish biogeochemical conditions that are conducive for the growth and development of calcifying benthic organisms. However, the effect of mineral additions can vary substantially between environments. Sensitivity analyses demonstrated that the OM mineralization had a significant impact on mineral dissolution, pH and mineral saturation. The type of mineral added also showed significant effect with aragonite producing more TA and higher pH. Increased mixing depth led to a faster increase in saturation state while increased iron flux caused reduction in TA concentration and benthic fluxes. Overall, the addition of minerals with higher dissolution rates would be most effective especially in environments with high OM degradation and undersaturated water conditions. Similarly, the type of mineral and the environmental conditions affect mCDR potential through benthic fluxes. Importantly, the implementation of OAE can lead to unforeseen effects, including ecological feedback such as changes in species composition (Bach et al., 2019; Köhler et al., 2013), and the effectiveness of carbonate additions may depend strongly on site characteristics. Thus, studies should combine field, lab and modeling components (Meysman & Montserrat, 2017) to improve predictive capabilities and decrease uncertainty.

Code Availability

490 The reaction transport model can be found at: https://bitbucket.org/MeileLab/kadir_oa/src/master/

Data Availability

Data for this manuscript is available in Myers (2022).

Author contribution

Lab experiments were designed by GW and TM, carried out by TM with the guidance of GW. KB developed the numerical
495 model with guidance of CM and carried out the simulations. KB prepared the initial draft of the manuscript with significant
input from CM, TM and GW.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

500 The authors would like to thank the ClimateWorks Foundation for funding and Jurjen Rooze for input on the model
implementation.

References

- Adkins, J. F., Naviaux, J. D., Subhas, A. V., Dong, S., & Berelson, W. M.: The dissolution rate of CaCO₃ in the ocean. *Annual Review of Marine Science*, 13(1), 57-80, 2021.
- 505 Aller, R. C.: The effects of macrobenthos on chemical properties of marine sediment and overlying water. In *Animal-sediment relations: the biogenic alteration of sediments* (pp. 53-102). Boston, MA: Springer US, 1982.
- Bach, L. T., Gill, S. J., Rickaby, R. E., Gore, S., & Renforth, P.: CO₂ removal with enhanced weathering and ocean alkalinity enhancement: potential risks and co-benefits for marine pelagic ecosystems. *Frontiers in Climate*, 1, 7, 2019
- Bandstra, L., Hales, B., & Takahashi, T.: High-frequency measurements of total CO₂: Method development and first
510 oceanographic observations. *Marine Chemistry*, 100(1-2), 24-38, 2006.
- Barton, A., Hales, B., Waldbusser, G. G., Langdon, C., & Feely, R. A.: The Pacific oyster, *Crassostrea gigas*, shows negative correlation to naturally elevated carbon dioxide levels: Implications for near-term ocean acidification effects. *Limnology and oceanography*, 57(3), 698-710, 2012.
- Berelson, W. M., Balch, W. M., Najjar, R., Feely, R. A., Sabine, C., & Lee, K.: Relating estimates of CaCO₃ production,
515 export, and dissolution in the water column to measurements of CaCO₃ rain into sediment traps and dissolution on the sea floor: A revised global carbonate budget. *Global Biogeochemical Cycles*, 21(1), 2007.
- Berner, R. A.: *Early diagenesis: a theoretical approach*. 241 p. Princeton University Press, 1980.
- Boudreau, B. P.: On the equivalence of nonlocal and radial-diffusion models for porewater irrigation. *Journal of Marine Research*, 42(3), 731-735, 1984
- 520 Boudreau, B. P.: The diffusive tortuosity of fine-grained unlithified sediments. *Geochimica et Cosmochimica Acta*, 60(16), 3139-3142, 1996
- Boudreau, B. P.: *Diagenetic models and their implementation*. 414 p. (Vol. 505). Berlin: Springer, 1997.
- Brenner, H., Braeckman, U., Le Guitton, M., & Meysman, F. J.: The impact of sedimentary alkalinity release on the water column CO₂ system in the North Sea. *Biogeosciences*, 13(3), 841-863, 2016.
- 525 Brown, C. A., & Ozretich, R. J.: Coupling between the coastal ocean and Yaquina Bay, Oregon: Importance of oceanic inputs relative to other nitrogen sources. *Estuaries and Coasts*, 32, 219-237, 2009.

- Burdige, D. J.: Preservation of organic matter in marine sediments: controls, mechanisms, and an imbalance in sediment organic carbon budgets? *Chemical Reviews*, 107(2), 467-485, 2007.
- Burdige, D. J., Hu, X., & Zimmerman, R. C.: The widespread occurrence of coupled carbonate dissolution/precipitation in surface sediments on the Bahamas Bank. *American Journal of Science*, 310(6), 492-521, 2010.
- 530 Cai, W. J., Hu, X., Huang, W. J., Murrell, M. C., Lehrter, J. C., Lohrenz, S. E., ... & Gong, G. C.: Acidification of subsurface coastal waters enhanced by eutrophication. *Nature geoscience*, 4(11), 766-770, 2011.
- Cai, W. J., Huang, W. J., Luther III, G. W., Pierrot, D., Li, M., Testa, J., ... & Kemp, W. M.: Redox reactions and weak buffering capacity lead to acidification in the Chesapeake Bay. *Nature Communications*, 8(1), 369, 2017.
- 535 Cyronak, T., Santos, I. R., Erler, D. V., & Eyre, B. D.: Groundwater and porewater as major sources of alkalinity to a fringing coral reef lagoon (Muri Lagoon, Cook Islands). *Biogeosciences*, 10(4), 2467-2480, 2013a.
- Cyronak, T., Santos, I. R., McMahon, A., & Eyre, B. D.: Carbon cycling hysteresis in permeable carbonate sands over a diel cycle: Implications for ocean acidification. *Limnology and Oceanography*, 58(1), 131-143, 2013b.
- D'Andrea, A. F., & DeWitt, T. H.: Geochemical ecosystem engineering by the mud shrimp *Upogebia pugettensis* (Crustacea: Thalassinidae) in Yaquina Bay, Oregon: Density-dependent effects on organic matter remineralization and nutrient cycling. *Limnology and Oceanography*, 54(6), 1911-1932, 2009.
- 540 Dickson, A. G.: An exact definition of total alkalinity and a procedure for the estimation of alkalinity and total inorganic carbon from titration data. *Deep Sea Research Part A. Oceanographic Research Papers*, 28(6), 609-623, 1981.
- Dickson, A. G.: Standard potential of the reaction: $\text{AgCl (s)} + 12\text{H}_2 \text{(g)} = \text{Ag (s)} + \text{HCl (aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *The Journal of Chemical Thermodynamics*, 22(2), 113-127, 1990.
- 545 Edmond, J. M., & Gieskes, J. M. T. M.: On the calculation of the degree of saturation of sea water with respect to calcium carbonate under in situ conditions. *Geochimica et cosmochimica acta*, 34(12), 1261-1291, 1970.
- Egleston, E. S., Sabine, C. L., & Morel, F. M.: Revelle revisited: Buffer factors that quantify the response of ocean chemistry to changes in DIC and alkalinity. *Global Biogeochemical Cycles*, 24(1), 2010.
- 550 Feely, R. A., Alin, S. R., Carter, B., Bednaršek, N., Hales, B., Chan, F., ... & Juranek, L.: Chemical and biological impacts of ocean acidification along the west coast of North America. *Estuarine, Coastal and Shelf Science*, 183, 260-270, 2016.
- Fennel, K., Long, M. C., Algar, C., Carter, B., Keller, D., Laurent, A., Mattern, J. P., Musgrave, R., Oschlies, A., Ostiguy, J., Palter, J. B., and Whitt, D. B.: Modelling considerations for research on ocean alkalinity enhancement (OAE), in: *Guide to Best Practices in Ocean Alkalinity Enhancement Research*, edited by: Oschlies, A., Stevenson, A., Bach, L. T., Fennel, K., Rickaby, R. E. M., Satterfield, T., Webb, R., and Gattuso, J.-P., Copernicus Publications, State Planet, 2-oae2023, 9, <https://doi.org/10.5194/sp-2-oae2023-9-2023>, 2023.
- 555 Ferderer, A., Chase, Z., Kennedy, F., Schulz, K. G., & Bach, L. T.: Assessing the influence of ocean alkalinity enhancement on a coastal phytoplankton community. *Biogeosciences*, 19(23), 5375-5399, 2022.

- 560 Fuhr, M., Wallmann, K., Dale, A. W., Kalapurakkal, H. T., Schmidt, M., Sommer, S., ... & Geilert, S.: Alkaline mineral addition to anoxic to hypoxic Baltic Sea sediments as a potentially efficient CO₂-removal technique. *Frontiers in Climate*, 6, 1338556, 2024.
- Gadeken, K. J., Lockridge, G., & Dorgan, K. M.: An in situ benthic chamber system for improved temporal and spatial resolution measurement of sediment oxygen demand. *Limnology and Oceanography: Methods*, 21(11), 645-655, 2023.
- 565 Gimenez, I., Waldbusser, G. G., & Hales, B.: Ocean acidification stress index for shellfish (OASIS): Linking Pacific oyster larval survival and exposure to variable carbonate chemistry regimes. *Elem Sci Anth*, 6, 51, 2018.
- Goldenberg, S. U., Riebesell, U., Brüggemann, D., Börner, G., Sswat, M., Folkvord, A., Couret, M., Spjelkavik, S., Sánchez, N., Jaspers, C., and Moyano, M.: Viability of coastal fish larvae under ocean alkalinity enhancement: from organisms to communities, *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2024-286>, 2024.
- 570 Green, M. A., Waldbusser, G. G., Reilly, S. L., Emerson, K., & O'Donnell, S.: Death by dissolution: sediment saturation state as a mortality factor for juvenile bivalves. *Limnology and Oceanography*, 54(4), 1037-1047, 2009.
- Green, M. A., Waldbusser, G. G., Hubazc, L., Cathcart, E., & Hall, J.: Carbonate mineral saturation state as the recruitment cue for settling bivalves in marine muds. *Estuaries and Coasts*, 36, 18-27, 2013.
- Gustafsson, E., Hagens, M., Sun, X., Reed, D. C., Humborg, C., Slomp, C. P., & Gustafsson, B. G.: Sedimentary alkalinity generation and long-term alkalinity development in the Baltic Sea. *Biogeosciences*, 16(2), 437-456, 2019.
- 575 Hales, B., Takahashi, T., & Bandstra, L.: Atmospheric CO₂ uptake by a coastal upwelling system. *Global Biogeochemical Cycles*, 19(1), 2005.
- Hales, B., Suhrbier, A., Waldbusser, G. G., Feely, R. A., & Newton, J. A.: The carbonate chemistry of the “fattening line,” Willapa Bay, 2011–2014. *Estuaries and Coasts*, 40, 173-186, 2017.
- 580 Hangx, S. J., & Spiers, C. J.: Coastal spreading of olivine to control atmospheric CO₂ concentrations: A critical analysis of viability. *International Journal of Greenhouse Gas Control*, 3(6), 757-767, 2009.
- Hartmann, J., West, A. J., Renforth, P., Köhler, P., De La Rocha, C. L., Wolf-Gladrow, D. A., ... & Scheffran, J.: Enhanced chemical weathering as a geoengineering strategy to reduce atmospheric carbon dioxide, supply nutrients, and mitigate ocean acidification. *Reviews of Geophysics*, 51(2), 113-149, 2013.
- 585 Hofmann, A. F., Soetaert, K., Middelburg, J. J., & Meysman, F. J.: AquaEnv: An Aquatic Acid–Base Modelling Environment in R. *Aquatic Geochemistry*, 16, 507-546, 2010.
- Hofmann, G. E., Smith, J. E., Johnson, K. S., Send, U., Levin, L. A., Micheli, F., ... & Martz, T. R.: High-frequency dynamics of ocean pH: a multi-ecosystem comparison. *PloS One*, 6(12), e28983, 2011.
- Hu, X., & Cai, W. J.: An assessment of ocean margin anaerobic processes on oceanic alkalinity budget. *Global Biogeochemical Cycles*, 25(3), 2011a.
- 590 Hu, X., & Cai, W. J.: The impact of denitrification on the atmospheric CO₂ uptake potential of seawater. *Marine Chemistry*, 127(1-4), 192-198, 2011b.

- Huettel, M., Ziebis, W., Forster, S., & Luther Iii, G. W.: Advective transport affecting metal and nutrient distributions and interfacial fluxes in permeable sediments. *Geochimica et Cosmochimica Acta*, 62(4), 613-631, 1998.
- 595 Jahnke, R. A., Craven, D. B., & Gaillard, J. F.: The influence of organic matter diagenesis on CaCO₃ dissolution at the deep-sea floor. *Geochimica et Cosmochimica Acta*, 58(13), 2799-2809, 1994.
- Jourabchi, P., Van Cappellen, P., & Regnier, P.: Quantitative interpretation of pH distributions in aquatic sediments: A reaction-transport modeling approach. *American Journal of Science*, 305(9), 919-956, 2005.
- Jourabchi, P., Meile, C., Pasion, L. R., & Van Cappellen, P.: Quantitative interpretation of pore water O₂ and pH distributions
600 in deep-sea sediments. *Geochimica et Cosmochimica Acta*, 72(5), 1350-1364, 2008.
- Köhler, P., Abrams, J. F., Völker, C., Hauck, J., & Wolf-Gladrow, D. A.: Geoengineering impact of open ocean dissolution of olivine on atmospheric CO₂, surface ocean pH and marine biology. *Environmental Research Letters*, 8(1), 014009, 2013.
- Krumins, V., Gehlen, M., Arndt, S., Van Cappellen, P., & Regnier, P.: Dissolved inorganic carbon and alkalinity fluxes from coastal marine sediments: model estimates for different shelf environments and sensitivity to global change. *Biogeosciences*,
605 10(1), 371-398, 2013.
- Lueker, T. J., Dickson, A. G., & Keeling, C. D. (2000). Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for K₁ and K₂: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium. *Marine chemistry*, 70(1-3), 105-119.
- Luff, R., & Wallmann, K.: Fluid flow, methane fluxes, carbonate precipitation and biogeochemical turnover in gas hydrate-
610 bearing sediments at Hydrate Ridge, Cascadia Margin: numerical modeling and mass balances. *Geochimica et Cosmochimica Acta*, 67(18), 3403-3421, 2003.
- Meile, C., & Van Cappellen, P.: Global estimates of enhanced solute transport in marine sediments. *Limnology and Oceanography*, 48(2), 777-786, 2003.
- Meysman, F. J., & Montserrat, F.: Negative CO₂ emissions via enhanced silicate weathering in coastal environments. *Biology Letters*, 13(4), 20160905, 2017.
615
- Middelburg, J. J., Soetaert, K., Herman, P. M., & Heip, C. H.: Denitrification in marine sediments: A model study. *Global Biogeochemical Cycles*, 10(4), 661-673, 1996.
- Middelburg, J. J., Soetaert, K., & Herman, P. M.: Empirical relationships for use in global diagenetic models. *Deep Sea Research Part I: Oceanographic Research Papers*, 44(2), 327-344, 1997.
- 620 Millero, F. J.: Carbonate constants for estuarine waters. *Marine and Freshwater Research*, 61(2), 139-142, 2010.
- Montserrat, F., Renforth, P., Hartmann, J., Leermakers, M., Knops, P., & Meysman, F. J.: Olivine dissolution in seawater: implications for CO₂ sequestration through enhanced weathering in coastal environments. *Environmental Science & Technology*, 51(7), 3960-3972, 2017.
- Morris, A. W., & Riley, J. P.: The bromide/chlorinity and sulphate/chlorinity ratio in sea water. *Deep sea research and oceanographic Abstracts (Vol. 13, No. 4, pp. 699-705)*. Elsevier, 1966.
- 625 Morse, J. W., & Mackenzie, F. T.: *Geochemistry of sedimentary carbonates*. Elsevier, 1990.

- Morse, J. W., Arvidson, R. S., & Lüttge, A.: Calcium carbonate formation and dissolution. *Chemical Reviews*, 107(2), 342-381, 2007.
- Mucci, A.: The solubility of calcite and aragonite in seawater at various salinities, temperatures, and one atmosphere total
630 pressure. *American Journal of Science*, 283(7), 780-799, 1983.
- Mucci, A., Sundby, B., Gehlen, M., Arakaki, T., Zhong, S., & Silverberg, N.: The fate of carbon in continental shelf sediments of eastern Canada: a case study. *Deep Sea Research Part II: Topical Studies in Oceanography*, 47(3-4), 733-760, 2000.
- Myers, T. J.: Buffering Estuarine Sediments Against Acidification. Master's Thesis. 77p. Oregon State University. https://ir.library.oregonstate.edu/concern/graduate_thesis_or_dissertations/br86bb964, 2022.
- 635 Pacella, S. R., Brown, C. A., Waldbusser, G. G., Labiosa, R. G., & Hales, B.: Seagrass habitat metabolism increases short-term extremes and long-term offset of CO₂ under future ocean acidification. *Proceedings of the National Academy of Sciences*, 115(15), 3870-3875, 2018.
- Pacella, S. R., Brown, C. A., Labiosa, R. G., Hales, B., Mochon Collura, T. C., Evans, W., & Waldbusser, G. G.: Feedbacks between estuarine metabolism and anthropogenic CO₂ accelerate local rates of ocean acidification and hasten threshold
640 exceedances. *Journal of Geophysical Research: Oceans*, 129(3), e2023JC020313, 2024.
- Paul, A. J., Haunost, M., Goldenberg, S. U., Hartmann, J., Sánchez, N., Schneider, J., Suitner, N., and Riebesell, U.: Ocean alkalinity enhancement in an open ocean ecosystem: Biogeochemical responses and carbon storage durability, *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2024-417>, 2024.
- Rao, A. M., Malkin, S. Y., Montserrat, F., & Meysman, F. J.: Alkalinity production in intertidal sands intensified by lugworm
645 bioirrigation. *Estuarine, Coastal and Shelf Science*, 148, 36-47, 2014.
- Rassmann, J., Lansard, B., Pozzato, L., & Rabouille, C.: Carbonate chemistry in sediment porewaters of the Rhône River delta driven by early diagenesis (northwestern Mediterranean). *Biogeosciences*, 13(18), 5379-5394, 2016.
- Renforth, P., & Henderson, G.: Assessing ocean alkalinity for carbon sequestration. *Reviews of Geophysics*, 55(3), 636-674.
- Riley, J. P. (1965). The occurrence of anomalously high fluoride concentrations in the North Atlantic. *Deep Sea Research*
650 A, 12(2), 219-220, 2017.
- Rooze, J., Egger, M., Tsandev, I., & Slomp, C. P.: Iron-dependent anaerobic oxidation of methane in coastal surface sediments: Potential controls and impact. *Limnology and Oceanography*, 61(S1), S267-S282, 2016.
- Rooze, J., Peterson, L., Peterson, R. N., & Meile, C.: Porewater flow patterns in surficial cold seep sediments inferred from conservative tracer profiles and early diagenetic modeling. *Chemical Geology*, 536, 119468, 2020.
- 655 Santos, I. R., Eyre, B. D., & Huettel, M.: The driving forces of porewater and groundwater flow in permeable coastal sediments: A review. *Estuarine, Coastal and Shelf Science*, 98, 1-15, 2012.
- Soetaert, K., Herman, P. M., & Middelburg, J. J.: A model of early diagenetic processes from the shelf to abyssal depths. *Geochimica et Cosmochimica Acta*, 60(6), 1019-1040, 1996.
- Soetaert, K., Hofmann, A. F., Middelburg, J. J., Meysman, F. J., & Greenwood, J.: The effect of biogeochemical processes on
660 pH. *Marine Chemistry*, 106(1-2), 380-401, 2007.

- Soetaert, K., Petzoldt, T., & Meysman, F.: Marelac: Tools for aquatic sciences. <https://cran.r-project.org/package=marelac>, 2010.
- Soetaert, K., & Meysman, F.: Reactive transport in aquatic ecosystems: Rapid model prototyping in the open source software R. *Environmental Modelling & Software*, 32, 49-60, 2012.
- 665 Subhas, A. V., Rollins, N. E., Berelson, W. M., Erez, J., Ziveri, P., Langer, G., & Adkins, J. F.: The dissolution behavior of biogenic calcites in seawater and a possible role for magnesium and organic carbon. *Marine Chemistry*, 205, 100-112, 2018.
- Taylor, L. L., Quirk, J., Thorley, R. M., Kharecha, P. A., Hansen, J., Ridgwell, A., ... & Beerling, D. J.: Enhanced weathering strategies for stabilizing climate and averting ocean acidification. *Nature Climate Change*, 6(4), 402-406, 2016.
- Thamdrup, B., & Canfield, D. E.: Benthic respiration in aquatic sediments. In *Methods in Ecosystem Science* (pp. 86-103).
670 New York, NY: Springer New York, 2000.
- Thomas, H., Schiettecatte, L. S., Suykens, K. O. N. E., Koné, Y. J. M., Shadwick, E. H., Prowe, A. F., ... & Borges, A. V.: Enhanced ocean carbon storage from anaerobic alkalinity generation in coastal sediments. *Biogeosciences*, 6(2), 267-274, 2009.
- Thullner, M., Dale, A. W., & Regnier, P.: Global-scale quantification of mineralization pathways in marine sediments: A
675 reaction-transport modeling approach. *Geochemistry, geophysics, geosystems*, 10(10), 2009.
- Uppström, L. R.: The boron/chlorinity ratio of deep-sea water from the Pacific Ocean. *Deep Sea Research A*, 21(2), 161-162, 1974.
- Waldbusser, G. G., Steenson, R. A., & Green, M. A.: Oyster shell dissolution rates in estuarine waters: effects of pH and shell legacy. *Journal of Shellfish Research*, 30(3), 659-669, 2011.
- 680 Waldbusser, G. G., Powell, E. N., & Mann, R.: Ecosystem effects of shell aggregations and cycling in coastal waters: an example of Chesapeake Bay oyster reefs. *Ecology*, 94(4), 895-903, 2013.
- Waldbusser, G. G., & Salisbury, J. E.: Ocean acidification in the coastal zone from an organism's perspective: multiple system parameters, frequency domains, and habitats. *Annual Review of Marine Science*, 6, 221-247, 2014.
- Wang, Y., & Van Cappellen, P.: A multicomponent reactive transport model of early diagenesis: Application to redox cycling
685 in coastal marine sediments. *Geochimica et Cosmochimica Acta*, 60(16), 2993-3014, 1996.
- Wolf-Gladrow D.A., Zeebe, R.E., Klaas, C., Koertzing, A. and Dickson, A.G.: Total alkalinity: The explicit conservative expression and its application to biogeochemical processes. *Marine Chemistry*, 106, 287-300, 2007.
- Zeebe, R. E., & Wolf-Gladrow, D.: CO₂ in seawater: equilibrium, kinetics, isotopes (No. 65). Gulf Professional Publishing, 2001.