

GOSI9: UK Global Ocean and Sea Ice configurations

Catherine Guiavarc’h¹, David Storkey¹, Adam T Blaker², Ed Blockley¹, Alex Megann², Helene Hewitt¹, Michael J Bell¹, Daley Calvert¹, Dan Copsey¹, Bablu Sinha², Sophia Moreton¹, Pierre Mathiot^{3,1}, and Bo An⁴

¹Met Office, FitzRoy Road, Exeter EX1 3PB, UK

²Marine Systems Modelling, National Oceanography Centre, Southampton, SO14 3ZH, UK

³Univ. Grenoble Alpes/CNRS/IRD/G-INP, IGE, Grenoble, France

⁴State Key Laboratory of Numerical Modeling for Atmospheric Sciences and Geophysical Fluid Dynamics, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, 100029, China

Correspondence: Catherine Guiavarc’h (catherine.guiavarch@metoffice.gov.uk)

Abstract. The UK Global Ocean and Sea Ice configuration version 9 (GOSI9) is a new traceable hierarchy of three model configurations at 1° , $\frac{1}{4}^\circ$ and $\frac{1}{12}^\circ$ based on version 4.0.4 of the NEMO code. GOSI9 has been developed as part of the UK’s Joint Marine Modelling Programme (JMMP), a partnership between the Met Office, the National Oceanography Centre, the British Antarctic Survey, and the Centre for Polar Observation and Modelling. Following a seamless approach it will be used for a variety of applications across a wide range of spatial and temporal resolutions: short-range coupled NWP forecasts, ocean forecasts, seasonal and decadal forecasts, climate and earth system modelling. The GOSI9 configurations are described in detail with special focus on the updates since the previous version (GO6-GSI8). Results from 30-year ocean-ice integrations forced by CORE2 fluxes are presented for the three resolutions and the impacts of the updates are assessed using the $\frac{1}{4}^\circ$ integrations. The upgrade to NEMO 4.0.4 includes a new sea ice model SI³ (Sea Ice modelling Integrated Initiative) and faster integration achieved through the use of partially implicit schemes that allow a significant increase in the length of the time step. The quality of the simulations are generally improved compared to GO6-GSI8. The temperature and salinity drifts are largely reduced thanks to the upgrade to NEMO 4.0.4 and the adoption of 4th order horizontal and vertical advections helping to reduce the numerical mixing. To improve the representation of the Southern Ocean, a scale-aware form of the Gent-McWilliams parametrisation and the application of a partial slip lateral boundary condition on momentum in the Southern Ocean have been added, resulting in a stronger and more realistic Antarctic Circumpolar Current (ACC) transport and a reduction of the temperature and salinity biases along the shelf of Antarctica. In the Arctic, the representation of sea ice is improved, leading to a reduction in surface temperature and salinity biases. In particular, the excessive and unrealistic Arctic summer sea ice melt in GO6-GSI8 is significantly improved in GOSI9 and can be attributed to the change in the sea ice model and to the higher albedos which increased sea ice thickness.

1 Introduction

The Joint Marine Modelling Programme (JMMP), founded in 2018, is a partnership between the Met Office and UK research centres: the National Oceanography Centre, the British Antarctic Survey and the Centre for Polar Observation and Modelling. The JMMP's ambition is to provide national capability modelling infrastructure, configurations and model output to the UK community. Adopting the seamless forecasting approach (Brown et al., 2012), the JMMP global ocean and sea ice configurations are used for a variety of applications across a wide range of spatial and temporal resolutions: short-range coupled NWP forecasts, ocean forecasts, seasonal and decadal forecasts, climate and Earth system modelling.

This paper describes the latest Global Ocean and Sea Ice configuration, GOSI9, based on the NEMO ocean modelling framework (Madec and system team, 2019). As for the previous global ocean configuration, GO6 (Storkey et al., 2018), GOSI9 is a traceable hierarchy of three resolutions: 1° , $\frac{1}{4}^\circ$ and $\frac{1}{12}^\circ$. Previously, new configurations of the Global Ocean (GO, Storkey et al., 2018; Megann et al., 2014) and Global Sea Ice (GSI, Ridley et al., 2018) were released separately, in part due to the dependence of the two models on different source codes. From NEMO version 4 (Madec and system team, 2019), the new native sea ice model SI³ (Sea Ice modelling Integrated Initiative) is available as part of the NEMO ocean modelling framework, and from this release onwards will be used for JMMP configurations. In the early stages of this development cycle the configuration was referred to as GO8p[0-7], and some papers based on these pre-release versions may have used this naming convention. However, it was realised that the previous release of the sea ice component had already reached version 8, and a second release at version 8 could lead to confusion. To explicitly recognise the joint release of both ocean and sea ice components we have adopted a new nomenclature for joint releases, GOSI.

Development of new configurations is motivated primarily by the desire to improve the representation of the ocean and sea ice, and secondarily to deliver better (more efficient) computational performance. The latter often comes from adopting newer code, whereas the former more frequently requires a combination of expert assessment to identify the cause of biases and to identify or write improved numerical schemes and an element of tuning/calibration to optimise the solution. Warm SST biases in the Southern Ocean have been a long standing issue in the Met Office coupled systems. In the Met Office GC3 coupled configuration (HadGEM3) based on the GO6+GSI8 ocean and sea ice components, the Southern Ocean SST bias was reduced compared with GC2 thanks to improved representation of clouds Williams et al. (2018) and tuning of the isopycnal diffusion parameter Storkey et al. (2018). However a warm bias still remained (Roberts et al., 2019). Hewitt et al. (2016) and Roberts et al. (2019) looked at the representation of the Southern Ocean in the hierarchy of HadGEM3 coupled models. They highlight the large biases in the Antarctic Circumpolar Current (ACC) transport especially in $\frac{1}{4}^\circ$ configuration which has the lowest transport, reduced by 40% compared to observations.

GOSI9 is required to deliver improved performance in both forced (ocean and sea ice only) and coupled configurations. It will form the ocean and sea ice component of the forthcoming GC5 version of the Met Office Hadley centre coupled climate model (Xavier et al., 2023) and will be the physical basis of the next UKESM (Mulcahy et al., 2023). To mitigate the risk of having to introduce configuration changes late in the development cycle a new approach was adopted: GOSI9 $\frac{1}{4}^\circ$ was tested in a coupled configuration at an early stage during the development process. This revealed some key insights into potential mean

55 state biases that we were then able to address during the cycle. We describe this process later, but the main focus of this paper is to present the results and testing of GOSI9 forced by atmospheric surface forcing. Evaluation of the final coupled configuration is presented in (Xavier et al., 2023).

The paper is organised as follows. Section 2 provides a detailed description of the GOSI9 configurations and the developments introduced into the global ocean configurations since the previous release, GO6. Section 3 describes the model simulations presented in this study, including the forcing and initialisation used. In Section 4 we present the results from the global evaluation for the three resolutions comparing GOSI9 and GO6, and the impact of individual changes are assessed using the $\frac{1}{4}^\circ$ degree configuration. Section 5 provides a more detailed evaluation of three key regions of interest: North Atlantic, North West Pacific and Southern Ocean.

2 Model description

65 GOSI9 is based on NEMO v4.0.4 (Madec and system team, 2019) and the SI³ sea ice component (Blockley et al., 2024) and is built on the traceable hierarchy of three model configurations at 1° , $\frac{1}{4}^\circ$ and $\frac{1}{12}^\circ$ horizontal resolutions described in Storkey et al. (2018). For ease of reference, in this section we provide a complete description of GOSI9. Details of the configuration that remain common to GO6, namely the grids, bathymetries, free surface solution and advection, mixing and boundary conditions are described first. We then describe the developments that have been made since GO6. Together these define the GOSI9 configuration.

2.1 Model grids and bathymetries

The model grids and bathymetries are unchanged from GO6 (Storkey et al., 2018). For the three resolutions, the grids are based on the ORCA global grids available on the NEMO framework (Madec and system team, 2019). eORCA1, eORCA025 and eORCA12 have a nominal 1° , $\frac{1}{4}^\circ$ and $\frac{1}{12}^\circ$ resolution at the Equator and an isotropic Mercator grid (i.e. same zonal and meridional grid spacing). To avoid singularity point in the ocean, in the Northern Hemisphere, the grids are quasi-isotropic bipolar with two north mesh poles being introduced on lands in Siberia and Canada. To better represent the equatorial dynamics, from 20° N/S the eORCA1 model grid meridional resolution increases towards $\frac{1}{3}^\circ$ at the Equator. In the Southern Hemisphere, south of 67° S, the grid is extended to 85° S following the method described in Mathiot et al. (2017) to include ice shelf cavities.

80 The three models have the same vertical grid with 75 vertical levels. The level thickness is double tanh function of depth increasing from 1m thickness near the surface to 200m at 6000m depth. It provides a high resolution near the surface to resolve ocean responses to atmospheric forcing, including the diurnal cycle (Bernie et al., 2005) and a reasonable resolution at mid-depths for long term climate responses. Stewart et al. (2017) show that 75 levels is the minimum number capable of resolving the second baroclinic mode and the surface layer. Partial step topography is used, making the depth of the bottom cell variable and adjustable to the real depth of the ocean (Adcroft et al., 1997; Barnier et al., 2006).

Different data sets were used to produce the model bathymetries for each resolution. In this regard, the hierarchy of resolutions is not fully traceable. For the 1° resolution model, eORCA1, the bathymetry is derived from ETOPO2 data set (National Geophysical Data Center, 2006) with additional data from IBSCO (Arndt et al., 2013) on the Antarctic shelf. For the $\frac{1}{4}^\circ$ resolution model, eORCA025, the bathymetry is derived from the ETOPO1 data set (Amante and Eakins, 2009) with additional data from GEBCO (IOC et al., 2003) in coastal regions and from IBSCO (Arndt et al., 2013) on the Antarctic shelf. For the $\frac{1}{12}^\circ$ resolution model, eORCA12, the bathymetry is derived from GEBCO_2014 (Weatherall et al., 2015). The bathymetry for eORCA025 had grid-scale smoothing applied. This was not done for the eORCA1 or eORCA12 bathymetries. A traceable set of bathymetries derived from a common source is planned for the next development cycle.

2.2 Free surface solution and advection

As in GO6, in order to represent accurately the surface fresh water flux, the model uses a nonlinear free surface allowing cell thicknesses throughout the water column to vary with time (z^* coordinate as in Adcroft and Campin, 2004). A change from GO6 (Storkey et al., 2018), in which a time-filtering solution was used with the fastest waves being filtered (Roullet and Madec, 2000), is that the equation of the surface pressure gradient is solved using the split-explicit free surface formulation, also called time-splitting formulation, following Shchepetkin and McWilliams (2005). The time-splitting solution allows for an explicit representation of the fastest external gravity waves.

For momentum advection, GOSI9 uses the vector-invariant form in which the horizontal advection is separated into rotational and irrotational terms. The vorticity term (including the Coriolis term) is calculated using the energy and enstrophy conserving scheme of Arakawa and Lamb (1981). In NEMO, different options of this scheme are available differing in the way the topographic boundary condition is represented. We retain the option used in GO6, `nn_ee_n_e3f=0`, which tends to reinforce the topography of the flow (Madec and system team, 2019, Sect. 5.2.1). The irrotational part of the momentum advection is formulated according to Hollingsworth et al. (1983) in order to avoid near grid-scale horizontal numerical instabilities (Ducouso et al., 2017). Advection of tracers is performed using the total variance diminishing (TVD) scheme of Zalesak (1979) with 4^{th} order on horizontal and vertical directions. In GO6, the same scheme was used but with 2^{nd} order on horizontal and vertical directions.

2.3 Mixing and boundary conditions

Lateral diffusion of momentum is on geopotential surfaces and, as in GO6, uses a Laplacian viscosity in eORCA1 and a bi-Laplacian viscosity in eORCA025 and eORCA12. The coefficients are specified in Table 1. In the bi-Laplacian case, to prevent instabilities due to numerical diffusion, the viscosity coefficients reduce polewards with the cube of the grid length. For the eORCA1 model, the viscosity coefficients reduce linearly with the increased meridional grid spacing towards the Equator, but are constant poleward of 20° N/S. This was an error in the input file which remains unchanged from GO6, and was identified too late in the development cycle to redress. Grid space dependent values have been tested and found to give small improvements in model fidelity. A modification to the input file similar to that applied by Hutchinson et al. (2023) will be necessary for applications in which the Antarctic ice shelf cavities are opened.

Lateral diffusion of tracers is performed along isoneutral surfaces using Laplacian mixing with coefficients given in Table 1. A parameterisation of adiabatic eddy mixing (Gent and McWilliams, 1990) with a spatially varying coefficient (Held and Larichev, 1996; Tréguier et al., 1997) is used in eORCA1. A weak grid-scale aware Gent and McWilliams (1990) parameterisation is now applied at the higher resolutions (see section 2.5.6).

The mixing parameterisations have been mainly unchanged compared to GO6. The vertical mixing of tracers and momentum is parameterized using a modified version of the Gaspar et al. (1990) turbulent kinetic energy (TKE) scheme (Madec and system team, 2019). To represent unresolved mixing due to internal wave breaking, a background vertical eddy diffusivity and a background viscosity are both applied. The background vertical eddy diffusivity has a value of $1.2 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$, which decreases linearly from $\pm 15^\circ$ latitude to a value of $1.2 \times 10^{-6} \text{ m}^2 \text{ s}^{-1}$ at $\pm 5^\circ$ latitude (Gregg et al., 2003). The background viscosity is applied globally with a constant value of $1.2 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$. Other unresolved processes are represented using parameterisations available in NEMO. Surface wave breaking mixing is parametrised following Craig and Banner (1994), increasing the mixing at the surface. Axell (2002) is used to represent the Langmuir cell mixing. An ad-hoc parameterisation is used to represent the mixing due to near-inertial wave breaking (Rodgers et al., 2014) with a length scale which can be varied geographically. Extensive work was carried to tune this length scale in GO6 (Storkey et al., 2018). Additional tuning was carried out during the development of GOSI9 in order to reduce biases in the coupled configuration (see Section 2.5.8).

To parametrise the convection, an increased vertical diffusivity of $10 \text{ m}^2 \text{ s}^{-1}$ is applied where the water column becomes unstable. The double diffusive mixing is parametrised using Merryfield et al. (1999). A climatological geothermal heat flux (Stein and Stein, 1992) is added as a bottom boundary condition. A quadratic bottom friction is used globally with enhanced coefficient in the Indonesian Throughflow, Denmark Strait and Bab-el-Mandeb regions. Following Beckmann and Döscher (1997), an advective and diffusive bottom boundary layer scheme is used. The tidal mixing is parameterised following Simmons et al. (2004) with a special formulation for the Indonesian Throughflow as recommended by Koch-Larrouy et al. (2008).

2.4 Fresh water input from land

Fresh water flux from the river runoffs is applied to the ocean surface layer. The freshwater runoffs are considered as fresh and as having the same temperature as the local SST. In order to avoid instabilities caused by shallow fresh layers, the vertical mixing is increased to $2 \times 10^{-3} \text{ m}^2 \text{ s}^{-1}$ in the top 10 m of the water column at runoff locations. Along Antarctica, to represent the icesheet fresh water input, we apply a parameterisation of ice shelf basal melting. Following Mathiot et al. (2017), climatological fresh water input is prescribed at the edge of the ice shelves through depth, to mirror the effect of ice shelf basal melt on the circulation. Due to stability issues in GOSI9 we use iceberg melt climatology based on GO6 integrations instead of interactive icebergs (see section 2.5.7).

	eORCA1	eORCA025	eORCA12
Lateral diffusion of momentum	laplacian	bilaplacian	bilaplacian
Lateral viscosity	$20,000 \text{ m}^2 \text{ s}^{-1}$	$-1.5 \times 10^{11} \text{ m}^4 \text{ s}^{-1}$	$-1.25 \times 10^{11} \text{ m}^4 \text{ s}^{-1}$
Isopycnal tracer diffusion	$1000 \text{ m}^2 \text{ s}^{-1}$	$150 \text{ m}^2 \text{ s}^{-1}$	$125 \text{ m}^2 \text{ s}^{-1}$

Table 1. Parameter changes between eORCA1, eORCA025 and eORCA12 configurations. The coefficients are calculated at the Equator. In the bi-Laplacian case (eORCA025 and eORCA12), the viscosity coefficients reduce polewards with the cube of the grid length. For the eORCA1 model, the viscosity coefficients reduce linearly with the increased meridional grid spacing towards the Equator, but are constant poleward of 20° N/S (see Section 2.3). The coefficients are unchanged compared to GO6 (see Table 1 in Storkey et al. (2018)).

2.5 Development and changes since GO6

2.5.1 Upgrade to NEMO 4.0.4

150 For GOSI9, the version of the NEMO code has been upgraded to NEMO 4.0.4 (compared to NEMO 3.6 for GO6). NEMO 4.0 was released in 2019 (Madec and system team, 2019). GOSI9 uses a split-explicit free surface proposed by Shchepetkin and McWilliams (2005). With the time split solution, external gravity waves are explicitly represented. GO6 used the filtered free surface available at NEMO v3.6 (Roulet and Madec, 2000) where the fastest waves are filtered. With the time split solution the fast barotropic motions (such as tides) are also simulated with a better accuracy. In NEMO4.0, the lateral diffusion code
155 has been rewritten with a different formulation including scale aware setting of eddy viscosity and diffusivity.

The AeroBULK package (Brodeau et al., 2017) has been implemented in NEMO 4.0. With AeroBULK, four bulk formulations are available in NEMO. GOSI9 uses the NCAR formulation (formerly CORE) appropriate for forcing the model with the CORE2 dataset, the same formulation used in GO6 but AeroBULK provides some refinements with the computation of the air density and with the reduction of approximation in the estimation of surface specific humidity of saturation by adding a
160 dependence to the sea level pressure.

2.5.2 Sea Ice model SI³

The sea ice component of GOSI9 is based upon NEMO's new native sea ice model, SI³ (Sea Ice modelling Integrated Initiative). SI³ is a dynamic-thermodynamic continuum sea ice model that includes an ice thickness distribution (ITD), conservation of horizontal momentum, an elastic-viscous plastic (EVP) rheology, and energy-conserving halo-thermodynamics (Vancoppenolle et al., 2023). SI³ has been available in the NEMO code since version 4.0, having been created by merging functionality
165 from several different sea ice models used with NEMO (namely LIM, CICE, and GELATO), building upon the LIM3 model of Rousset et al. (2015). SI³ is fully embedded within NEMO and is invoked from within NEMO's Surface Boundary Code (SBC) module.

Aside from the change in model, the physics of the sea ice component of GOSI9 are largely similar as used in the previous
170 GO6+GSI8 configuration (Ridley et al., 2018; Storkey et al., 2018) based upon CICE5. GOSI9 uses 5 thickness categories to
model the sub-grid ITD, with an additional ice-free category for open water (Thorndike et al., 1975). Thermodynamic growth
and melt of the sea ice is modelled using multi-layer thermodynamics, with 4 layers of ice and one of snow based upon Bitz and
Lipscomb (1999). Sea ice dynamics are modelled using the elastic–viscous–plastic (EVP) rheology of Hunke and Dukowicz
(2002). The biggest difference compared with GO6+GSI8 (Storkey et al., 2018) is the use of the broadband albedo scheme in
175 SI3 instead of the dual-band scheme used in CICE. This has allowed us to tune the albedo independently of the coupled model
and we increased the values of the albedo relative to the SI3 defaults (ponded ice albedo is increased from 0.27 to 0.36, dry ice
albedo is increased from 0.60 to 0.70 and dry snow albedo is increased from 0.85 to 0.87). The sea ice component of GOSI9 is
run on the same model grid as the ocean and at every ocean time step. More details on the specifics of the sea ice configuration,
along with details of how SI³ was adapted to work in the HadGEM3 coupled model, can be found in Blockley et al. (2024).

180 **2.5.3 Equation of state TEOS-10**

In GOSI9, the equation of state was upgraded to the Thermodynamic Equation Of Seawater – 2010 (TEOS-10, Ioc et al.,
2010) instead of the previous standard, the 1980 equation of state (EOS-80). An important change is the use of absolute
salinity and conservative temperature instead of practical salinity and potential temperature for EOS-80. TEOS-10 provides
a complete thermodynamically consistent representation of all thermodynamic properties of seawater and allows for a more
185 accurate representation of the heat content.

However, the use of TEOS-10 has an impact for users. For the FOAM ocean forecasting system, the observations assimilated
in the system are EOS-80 variables meaning that the model TEOS-10 variables need to be converted to EOS-80 prior to being
passed to the observation operator. There is no practical impact on coupling with atmosphere and sea ice, as when using
NEMO with TEOS-10 equation of state, the sea surface conservative (TEOS-10) temperature is converted to potential (EOS-
190 80) temperature before coupling.

Note that for this paper, the conservative temperature and absolute salinity fields from GOSI9 have been converted to po-
tential temperature and practical salinity to facilitate the comparison with GO6. The temperature and salinity results presented
throughout this paper are the EOS-80 variables (potential temperature and practical salinity).

2.5.4 Time step and performance

195 A benefit from upgrading to NEMO 4 has been the implementation of the adaptive-implicit vertical advection (Shchepetkin,
2015). As for most ocean models, the time step in NEMO needs to satisfy multiple criteria to maintain numerical stability.
The vertical CFL criterion is commonly the most limiting and imposes time and space discretisation constraints. Treating
the vertical advection implicitly can reduce these restrictions but causes large dispersive errors. With adaptive-implicit vertical
advection, the implicit scheme is only used in targeted areas where potential breaches of vertical CFL condition occur. It allows
200 a much longer time step while retaining the accuracy of the explicit scheme.

Configuration	GO6			GOSI9		
Resolution	1	1/4	1/12	1	1/4	1/12
time step (s)	2700	1350	300	3600	1800	600
number of core	224	486	6237	156	344	6150
number of simulated year per day	18	2	-	24	2.8	2

Table 2. Summary of GO6 and GOSI9 performances for the 3 resolutions. The integrations used are the reference simulations described in this manuscript and in Storkey et al. (2018). The integrations were performed on a Cray XC40 supercomputer using Intel Broadwell processors. The number of cores are for NEMO-SI3 only, it does not include XIOS cores. Note that we were not able to perform an additional integration to test GO6 eORCA12 performance in parallel to GOSI9 eORCA12.

Adaptive-implicit vertical advection has been introduced in the GOSI9 configurations as well as an implicit sea ice drag (available from NEMO 4.0.4). These changes allow the use of a considerably longer time step (Table 2) without introducing any significant changes or biases in the ocean and sea ice (not shown). The performances from GO6 configurations, as described in Storkey et al. (2018), and from GOSI9 configurations, as described in this paper, are summarized in Table 2. All the integrations were performed on a Cray X40 supercomputer using Intel Broadwell processors. For the $\frac{1}{4}^\circ$ model eORCA025, the time step has been increased by 33% for the ocean-only configuration and by 50% for the coupled GC5 configuration. For the $\frac{1}{12}^\circ$ configuration eORCA12, the time step has been increased by 100%, allowing to produce 2 years of simulation per day on 6150 cores. For the 1° configuration eORCA1, the time step has been increased by 33%. Note that for GO6, the "land suppression" option available in NEMO (Madec and system team, 2019) was only used for the $\frac{1}{12}^\circ$ configuration while it was used for the three resolutions in GOSI9. The "land suppression" option excludes much of the global land area from the calculations allowing to reduce the number of cores required. This explains the significant reduction of the number of cores used to perform the GOSI9 integrations for the 1° and $\frac{1}{4}^\circ$ resolutions.

2.5.5 Numerical mixing

Numerical mixing, caused mainly by truncations in the tracer advection scheme, is a recognised problem in ocean models. In models such as NEMO that use quasi-Eulerian vertical coordinates it arises from errors in vertical advection associated with internal waves and tides, and it also results from horizontal tracer advection in regions with a high cell Reynolds number where the mesoscale is not well resolved, which are particularly extensive on a $\frac{1}{4}^\circ$ grid at mid and high latitudes. Megann (2018) estimated the numerical mixing in a $\frac{1}{4}^\circ$ GO5.0 NEMO configuration by evaluating an effective diffusivity from the density transformation rate, and demonstrated that the effective diffusivity was over five times as large as the explicit diffusivity calculated in the model mixing scheme. Megann and Storkey (2021) found that increasing the viscosity in a $\frac{1}{4}^\circ$ GO6, either by using larger values for the fixed biharmonic viscosity parameter or changing to the Smagorinsky viscosity formulation, led to a reduction by between 10 and 20% in the effective diffusivity over much of the ocean interior, along with comparable reductions in temperature and salinity biases. The sole disadvantage of increased viscosity is a reduction in stability: in GO6, tripling the fixed viscosity required the time step to be reduced by 50% from the default of 1,350 s, while GOSI9 requires a 50%

225 reduction in time step from the default of 1,800 s if either the fixed viscosity is doubled or the Smagorinsky scheme is selected.
This increase in running cost ruled out the use of viscosity as a tool to reduce numerical mixing in coupled applications of
this ocean configuration. The z time-filtered Arbitrary Lagrangian-Eulerian coordinate (Leclair and Madec, 2011) has been
shown to significantly reduce numerical mixing from internal waves and tides (Megann et al., 2022; Megann, 2024), and may
be considered for inclusion in future global Global Ocean and Sea Ice configurations, although this again requires a reduced
230 timestep at present for stability.

As an alternative approach to reducing the numerical mixing resulting from advection in poorly-resolved mesoscale flows,
the tracer advection in both horizontal and vertical directions was changed from second order (the default in GO6) to fourth
order. This incurred no significant penalties in run time, and did not require any reduction in time step.

2.5.6 Southern Ocean tuning

235 As highlighted in the Introduction, the HadGEM3 coupled models with eddy-permitting ($\frac{1}{4}^\circ$) or eddy-resolving ($\frac{1}{12}^\circ$) ocean
resolution have a history of large-scale biases in the Southern Ocean. The largest biases appear at eddy-permitting resolution,
which shows weak ACC transports (Hewitt et al., 2016; Roberts et al., 2019) as well as overly active subpolar gyres in the
Weddell and Ross Seas and biases in the temperatures and salinities on the Antarctic shelves. Experiments in the coupled
model showed that these biases are linked, and that a package of changes aimed at damping the overactive gyres also had the
240 effect of improving the ACC transport and the shelf temperature and salinity biases. The work in the coupled model is described
in more detail in Storkey et al. (2024); in this paper (Section 5.1) we show the impact in the forced GOSI9 configurations.

The so-called Southern Ocean package of changes to the eddying models consists of the introduction of a scale-aware form
of the Gent-McWilliams parametrisation and the application of a partial slip lateral boundary condition on momentum in the
Southern Ocean.

245 Hallberg (2013) discusses ocean models that resolve eddies in parts of the domain but not in others. This is the case for
the $\frac{1}{4}^\circ$ and $\frac{1}{12}^\circ$ models presented here, in which eddies are well resolved at low latitudes but not at high latitudes due to the
decrease of the Rossby radius of deformation with latitude (see Hallberg (2013) Figure 1). At GOSI9 we try to account for
the effect of the unresolved eddies at high latitudes by introducing the same space- and time-dependent version of the Gent-
McWilliams scheme (Tréguier et al., 1997) as used in the low resolution model but with a coefficient that is capped to be zero
250 at low latitudes, ramping up to a low value of $75 \text{ m}^2 \text{ s}^{-1}$ at high latitudes.

As with GO6, a free slip lateral boundary condition on momentum is applied at all resolutions. To further damp the Southern
Ocean gyres in the $\frac{1}{4}^\circ$ and $\frac{1}{12}^\circ$ configurations, we increase the topographic drag by introducing partial slip boundary condition
(see Madec and system team (2019) section 7.1) south of 50°S .

2.5.7 Iceberg climatology

255 GO6 (Storkey et al., 2018) uses a Lagrangian iceberg model (Bigg et al., 1997; Martin and Adcroft, 2010). The same Lagrangian
iceberg model was deployed in GOSI9 but interactive icebergs caused stability problems. An initial stability issue caused by
excess melting was resolved by changing the melting temperature of the iceberg to the freezing point.

Stability was especially an issue with the coupled configuration, GC5, where to conserve fresh water, excess precipitation over Antarctica is balanced through the iceberg calving. At times this resulted in double the number of icebergs compared with the forced GOSI9, and accumulation of icebergs along the Antarctic peninsula caused regular crashes. Work was carried out to improve realism and stability: implementation of calving distribution along the ice shelf rather than on a single point; and implementation of a speed limiter for icebergs to prevent icebergs from travelling more than half a grid cell into the processor halo region in one time step. These changes reduced the frequency of crashes but not to a satisfactory level. It is hoped that further development can be done to improve stability.

In the meantime a fresh water iceberg climatology is used instead of the Lagrangian iceberg model. Tests were carried out using the Antarctic icebergs melt climatology from Merino et al. (2016). However, Merino et al. (2016)'s climatology does not include melt contributions from the Northern Hemisphere. Therefore we built a monthly fresh water iceberg climatology using the iceberg fresh water outputs from 30 year integrations of GO6 (Storkey et al., 2018), which are initialised from EN4 climatology and start from a state of rest with no icebergs. It takes around 4 years for the number of icebergs and the iceberg melt to stabilise, so the first five years of the simulations were discarded to create the climatology. A fresh water iceberg melt climatology was created for each resolution: 1° , $\frac{1}{4}^\circ$ and $\frac{1}{12}^\circ$ from the corresponding GO6 runs. The heat content of the melt water into the ocean is calculated using the freezing point temperature and the latent heat of melting is extracted from the ocean.

Figure S1 shows the annual mean fresh water flux from icebergs calculated from GO6 integrations. The distribution of iceberg melt is similar between the different resolutions. As the distribution depends closely on the circulation, the distribution for the higher resolution configuration features more small scales with iceberg melt occurring closer to the coast. The eORCA1 iceberg melt climatology differs from the higher resolution in the Weddell Sea which could be a consequence of the difference in circulation in that region.

2.5.8 Coupled model GC5 tuning

During the early testing phase with the coupled model GC5, biases arose in the Indian Ocean, with too cold SST and too warm subsurface temperatures. To better understand and to try to reduce these biases several sensitivity experiments were carried out, and these resulted in two key changes: an increase to the globally uniform value of the chlorophyll concentration; and a reduction in the TKE mixing depth between 10° S and 40° S. Increasing the chlorophyll concentration acts to reduce the depth of the solar penetrating radiation, warming the surface layer and cooling the subsurface. The chlorophyll concentration was increased from 0.05 mg.m^{-3} at GO6 to 0.1 mg.m^{-3} , which better matches the observed climatological value in the tropics. As part of the GO6 development, extensive work was carried to tune the near surface mixing (Storkey et al., 2018). Tuning of the e-folding length scale nn_htau associated with the parameterisation of near-inertial wave breaking resulted in the choice of a larger length scale in the Southern Ocean (Storkey et al., 2018, Fig. 2) where summertime climatological (de Boyer Montégut et al., 2004) mixed layers are on average deeper than in the northern latitudes. Revisiting this work we found that reducing the mixing depth between 10° and 40° S improved the warm subsurface bias in the Indian Ocean. The changes were tested in the forced $\frac{1}{4}^\circ$ GOSI9 configuration before being implemented as standard in the three GOSI9 configurations. In future

configurations, we plan to use a seasonal climatology of the chlorophyll concentration which will better represent the large variations associated with spring blooms.

3 Integrations

295 For the experiments described in this paper, the model initial conditions for temperature and salinity are from monthly climatologies based on EN4 objective analysis (Good et al., 2013) for the years 1995-2014. For the 30 year integrations presented in Section 4, the sea ice is initialised with a restart produced after 1 year of integration. The 1 year integration is initialised with temperature and salinity from EN4 climatology and the sea ice initial state is calculated by SI³ from the initial temperature and salinity. The model is spun up from a state of rest.

300 GOSI9 integrations are forced over the period 1976-2005 by the CORE2 surface data set (Large and Yeager, 2009) using the NCAR bulk formulae (Large and Yeager, 2009). Relative wind stress between the wind and the ocean current is used. As temporal resolution of the CORE2 forcing is not sufficient to resolve the diurnal forcing (Bernie et al., 2007), an artificial diurnal cycle is imposed on the daily mean shortwave fluxes. A sea surface salinity (SSS) restoration towards monthly mean climatology is applied with a -33.333 mm/day/psu restoring coefficient. A monthly climatology is applied for the fresh water
305 flux from river runoff (Bourdallé-Badie and Treguier, 2006), icebergs (see Section 2.5.7) and ice shelves (Rignot et al., 2013).

Sensitivity experiments with the $\frac{1}{4}^{\circ}$ configuration have been carried out to test the impact of changes detailed in Section 2.5. The list of experiments is detailed in Table 3. For all experiments, initialisation and forcing follow the protocol used for GOSI9 and described above. Results from these experiments are assessed in section 4. Mixed layer depth climatology from the de Boyer Montégut et al. (2004) dataset, sea surface temperature climatology from ESA CCI (Merchant et al., 2014), temperature
310 and salinity climatologies from EN4 objective analyses (Good et al., 2013) are used for comparison. EN4 temperature and salinity are averaged over the period 1995-2014, covering a period where data from Argo floats are available.

4 Model evaluation and comparison with GO6

In this section, the results from the three GOSI9 integrations are evaluated and compared against the GO6 integrations and observation-based products.

315 4.1 Global drift and budget analysis

The first metrics used to evaluate the model performance are the drifts in the globally integrated temperature and salinity from the initial conditions (Fig. 1 and 2). Starting from an observed climatological state, the model should ideally exhibit no strong trend over time. Fig. 1 and 2 compare the drifts in GO6 (left) and GOSI9 (right) for 1° (a, b), $\frac{1}{4}^{\circ}$ (c, d), and $\frac{1}{12}^{\circ}$ (e, f).

GOSI9 represents a substantial improvement in both temperature and salinity biases over the top 1000 m compared with
320 GO6. The warm bias which established quickly in the upper 300 m in GO6 is reduced to around $\frac{1}{3}$ its size across the three resolutions, and the strong cooling that was manifest primarily in the $\frac{1}{4}^{\circ}$ and $\frac{1}{12}^{\circ}$ resolutions is far smaller. The improvements

Experiment	GOSI	NEMO version	Sea Ice model	SO pack.	Eq. of State	Adv. order	Chl. conc. (mg.m ⁻³)	TKE mixing depth	length
GO6	GO6	3.6	CICE	no	EOS80	2nd order	0.05	Storkey et al. (2018)	1976-2005
GO6 4.0	GO6	4.0	SI ³	no	EOS80	2nd order	0.05	Storkey et al. (2018)	1976-2005
GOSI9 2nd order	GOSI9	4.0.4	SI ³	yes	TEOS10	2nd order	0.1	sect. 2.5.8	1976-2005
GOSI9 nn_etau	GOSI9	4.0.4	SI ³	yes	TEOS10	4th order	0.1	Storkey et al. (2018)	1976-1984
GOSI9 chl	GOSI9	4.0.4	SI ³	yes	TEOS10	4th order	0.05	sect. 2.5.8	1976-1994
GOSI9	GOSI9	4.0.4	SI ³	yes	TEOS10	4th order	0.1	sect. 2.5.8	1976-2005

Table 3. List of experiments with eORCA025 configuration. “SO pack.” stands for Southern Ocean package including GM eddy parametrisation and increased topographic drag (see section 2.5.6). “Adv. order” stands for order of the advection used in the model. The same order is used for the vertical and the horizontal advection. “Chl. conc.” stands for chlorophyll concentration. For all integrations, a constant is used.

in the upper 300 m are largely attributable to the increase of chlorophyll concentration and the reduction of the mixing depth (see section 2.5.8), whilst reduction of the numerical mixing from using the 4th order advection scheme and the upgrade to NEMO 4.0.4 are the leading contributors to the reduction in cooling below 300 m. Figure 3 shows an evolution of the globally integrated temperature drift from GO6 (a) through to GOSI9 (f) in the $\frac{1}{4}^{\circ}$ configuration. Figures 3b-e show intermediate stages of the development process, enabling us to attribute relative contributions to the improvement to individual changes that have been made to the model configuration. The panels correspond to the list of experiments shown in table 3.

Similarly, the fresh bias which develops over the upper 500 m, centred around 200 m depth, in GO6 is reduced by 50% or more (Fig. 2). This is mainly due to the upgrade to NEMO 4.0.4 (Fig. 4b) and the more accurate bulk formulae, which produces a salinification in the Tropics. The other developments resulted in minimal impact on the globally integrated salinity trends (Fig. 4). While in GO6 the amplitude of the salinity and temperature drifts varies with the resolutions, all three GOSI9 models show consistent drift with little variation between resolutions.

This assessment of model drift in globally integrated quantities is very positive, suggesting significant improvements over the previous generation of the models. However, we now need to assess whether these improvements arise from a reduction in biases or whether they are the product of large competing regional biases.




Figure 1. Global mean potential temperature drift from initial conditions (K) for GO6 (left) and GOSI9 (right).

4.2 Large scale biases

4.2.1 Zonal mean biases

For the next assessment we compute zonal and time mean quantities of temperature and salinity using the period 1996-2005, the last decade of the forced simulations. Figure 5 shows anomalies of these metrics computed for the $\frac{1}{4}^\circ$ configuration from both GO6 and GOSI9 from a reference 20-year climatology of EN4 v1.1, as well as anomalies between GOSI9 and GO6.

Both models exhibit a tendency to develop cold biases in the tropics and the Arctic, and warm biases in the extra-tropics and Southern Ocean. The cold bias between 200 m and 700 m present in GO6 between 30° S and 40° N is still present in GOSI9 but is significantly reduced, especially in the $\frac{1}{4}^\circ$ resolution (Fig. 5). The near-surface warm bias in the tropics is also greatly reduced, and there is a modest reduction of the cold bias in the Arctic. There is a slight degradation in the Southern Hemisphere extra-tropics, with a cold bias developing near 30° S, 50-100 m depth, and a slight increase in the warm bias between 40° - 60° S




Figure 2. Global mean practical salinity drift from initial conditions for GO6 (left) and GOSI9 (right).

below 200 m. Salinity biases reduce at most latitudes and depths, though a modest positive salinity bias develops in the shallow subsurface tropics. Overall, GOSI9 shows reduced temperature and salinity biases at most latitudes and depths.

We can attribute these changes to each of the development steps by comparing the simulations for each of the intermediate steps (Table 3). Figure 6 shows the incremental temperature and salinity anomalies that arise from updating the NEMO code from v3.6 to v4.0 (a, b), increasing the global mean chlorophyll concentration value to $0.1 \text{ mg}\cdot\text{m}^{-3}$ (c, d), and adjusting the `nn_htau` parameter that controls TKE mixing penetration (e, f). Upgrading to NEMO 4.0 results in a warming between 100 and 700 m between 20° S and 40° N and in the Arctic (Fig. 6a), and is the change primarily responsible for reducing the cold bias observed in GO6 in these regions (Fig. 5a). It also acts to increase the salinity in the shallow tropical region (Fig. 6b), partially compensating the fresh biases in GO6 (Fig. 5b) and leading to the slight saline bias in GOSI9 (Fig. 5d). Increasing the chlorophyll concentration to $0.1 \text{ mg}\cdot\text{m}^{-3}$ has a cooling effect, which reduces subsurface warm biases in the tropics but slightly increases cold biases in the extra tropics (Fig. 6c). The increase in chlorophyll does not affect the salinity. Adjusting the `nn_htau` parameter cools the subsurface between 50 m and 300 m and reduces the warm bias present in these layers between 30° and




Figure 3. Global mean potential temperature drift from initial conditions (K) for the experiments with the $\frac{1}{4}^\circ$ resolution model. (a) GO6. (b) GO6 4.0: GO6 upgraded to NEMO 4.0.4, this configuration is the first experiment using the sea ice model SI³ and has large sea ice biases impacting the surface salinity. (c) GOSI9. (d) GOSI9 nn_etau: GOSI9 with GO6 TKE mixing depth. (e) GOSI9 chl: GOSI9 with GO6 chlorophyll concentration. (f) GOSI9 2nd order: GOSI9 with 2^{nd} order advection in horizontal and vertical. The experiments are detailed in Table 3.




Figure 4. Global mean practical salinity drift from initial conditions (K) for the experiments with the $\frac{1}{4}^\circ$ resolution model. (a) GO6. (b) GO6 4.0: GO6 upgraded to NEMO 4.0.4, this configuration is the first experiment using the sea ice model SI³ and has large sea ice biases impacting the surface salinity. (c) GOSI9. (d) GOSI9 nn_etau: GOSI9 with GO6 TKE mixing depth. (e) GOSI9 chl: GOSI9 with GO6 chlorophyll concentration. (f) GOSI9 2nd order: GOSI9 with 2^{nd} order advection in horizontal and vertical. The experiments are detailed in Table 3.

360 40° S but in the forced GOSI9 configuration it introduces a cold bias centered at 100 m. This change also has a negligible effect on salinity. Its isolated effect is presented in Figure 6e,f. Despite the mixed impact on the forced GOSI9 configuration, the nn_htau adjustment was implemented with the purpose of keeping the same ocean configuration for forced and coupled




Figure 5. Zonal-mean potential temperature (K) and salinity : anomalies against 20-year climatology of EN4 v1.1 (Good et al., 2013) for the $\frac{1}{4}^\circ$ configuration for GOSI9 and GO6 and differences between GOSI9 and GO6. Model fields are time-means over the third decade of integration.

applications. In forced integration, GOSI9, the impact on the surface is more limited than in the coupled model as the SST is constrained by the bulk formulae but the impact on the subsurface is significant.

The more accurate tracer advection was found to result in reductions in the effective diffusivity, as defined by Megann (2018), of around 10%, as well as robust improvements in model biases, which were mainly attributed to increasing the order of the horizontal advection. Figure 7 shows the impact of the 4th order on the temperature in the top 700 m in the $\frac{1}{4}^\circ$ configuration. Overall, using the 4th order tends to warm the ocean especially in the Atlantic basin. The impact is also seen on the western boundary currents (Gulf Stream and Kuroshio) where the steering is changed.

4.2.2 Spatial biases

Whilst the ability of a model to adequately represent globally or zonally integrated properties is important for applications such as climate projections and Earth system model studies, minimising regional biases is also critical, not only for those regions but the biases can also affect climate and weather downstream. In this section we systematically assess the geographical distribution of biases.




Figure 6. Zonal-mean potential temperature (K, left) and salinity (right) anomalies resulting from: updating the NEMO code from v3.6 to v4.0 (a, b); increasing the global mean chlorophyll concentration value to 0.1 mg.m^{-3} (c, d); adjustment of the nn_etau parameter that controls TKE mixing penetration (e, f). Model fields are time-means over the third decade of integration.

For both GO6 and GOSI9, the three resolutions show a similar distribution of large-scale temperature and salinity biases. In GO6, temperatures are too warm at the surface and subsurface (Fig. 8 and Fig. 9) in the Southern Ocean, in the Tropics
 375 and in the Arctic Ocean. Changes made in GOSI9 significantly reduce the biases in the Arctic especially for the 1° and $\frac{1}{4}^\circ$ resolutions. This improvement is linked to an improved representation of Arctic sea ice in GOSI9 especially in Summer where the melting has been reduced and is in much better agreement with the observations. The warm SST bias in the Tropics present in GO6 is reduced in GOSI9, however the warm biases in the coastal upwellings in the East Pacific and East Atlantic are still present. At 100 m depth, GOSI9 is significantly cooler than GO6 (Fig. 9 and Fig. 5) as a result of the increased chlorophyll
 380 concentration. This change is net positive globally, reducing the warm bias centred at 100 m and present at all resolutions in GO6 (Fig. 5). However it introduces regional cold biases, especially in the Indian Ocean. In the North Atlantic subpolar gyre, there is a dipole of cooling and warming, which is discussed in further details in section 5.2. Looking at the sea surface salinity, GO6 has large scale fresh biases between 40° S and 40° N (Fig. 10). Outside the Arctic, the largest biases occur in the locations where subtropical mode waters ventilate. These subtropical surface fresh biases extend to depths of 500 m in
 385 the $\frac{1}{4}^\circ$ and $\frac{1}{12}^\circ$ resolutions (Fig. 5). With the upgrade to NEMO4 and its more accurate bulk formulae, these fresh biases are significantly reduced in GOSI9 and do not extend as deep (Fig. 5). In Arctic, the fresh biases present in GO6 are reduced in




Figure 7. Potential temperature anomalies averaged over 0-700 m against EN4 v1.1 average 1995-2014 (Good et al., 2013) for GOSI9 and GOSI9 2nd order and difference between GOSI9 2nd order and GOSI9. Experiments GOSI9 and GOSI9 2nd order are detailed in Section 3 and Table 3. Model fields are time-means over the third decade of the integrations.

GOSI9. We note that comparison with EN4 in the Arctic is problematic due to the existence of artificial spatial patterns arising from the interpolation of sparse observational data. Figures 11 and 12 respectively show the annual minimum and maximum mixed layer depth biases compared to de Boyer Montégut et al. (2004). Summer minimum mixed layer depths are in good agreement with observations in all three models. There is an overall tendency towards a slight shallow bias of order 5-10 m, with deep biases at the Equator and in a few regions around the Southern Ocean. Storkey et al. (2018) noted the too deep winter mixed layer in the North Atlantic subpolar gyre and in the Greenland-Iceland-Norway seas and in the Southern Ocean west of Drake Passage. These biases are still present in GOSI9 but with reduced amplitude. For the 1° resolution, biases in the NE Atlantic change from positive to negative, whilst the strong bias in the Labrador Sea is almost eliminated. Apart from these




Figure 8. Sea surface temperature anomalies against 1996-2014 average ESA CCI (Merchant et al., 2014) for GOSI9 and differences between GOSI9 and GO6. Model fields are time-means over the third decade of the integrations.

395 regions, differences in MLD between GO6 and GOSI9 can be observed in a latitudinal band around 40° S. This corresponds to the region where the TKE mixing depth has been reduced in GOSI9 (Section 2.5.8). Shallowing of MLD between GOSI9 and GO6, at around 40° S and at around 60° N, are more clearly visible in Figure S2, showing the mixed layer depth zonal mean for both models.

4.2.3 Sea ice

400 The seasonal cycle of sea ice area and volume is much improved in GOSI9 in both the northern and southern hemispheres (Fig. 13). This represents a marked improvement over GO6, for which the Arctic summer minimum was significantly lower than observed, especially for the 1° and $\frac{1}{4}$ ° resolutions. Ice representation is similar across the resolutions in GOSI9, so for conciseness we show only the assessment of $\frac{1}{4}$ °. Given the change of ice model from CICE5 to SI³, it is difficult to attribute these improvements more precisely. However, one significant change is the increase in albedo (section 2.5.2). Tested separately,
 405 it resulted in increased sea ice thickness and reduced melt in Summer Arctic sea ice (not shown). This is consistent with work from Rae et al. (2014) that shows how sensitive the sea ice models are to parameter changes that affect the surface radiation and highlights the Arctic sea ice as the most sensitive to snow albedo. Figure 14a-c shows the September mean for Arctic sea




Figure 9. Potential temperature anomalies at 100 m against EN4 v1.1 average 1995-2014 (Good et al., 2013) for GOSI9 and differences between GOSI9 and GO6. Model fields are time-means over the third decade of the integrations.

ice concentration for the $\frac{1}{4}^\circ$ configuration at GOSI9, gridded observations from HadISST, and the $\frac{1}{4}^\circ$ configuration at GO6. The northern hemisphere distribution of sea ice in GOSI9 is in much better agreement with observations than GO6, though
 410 concentrations are still lower than observed over the central Arctic. Increasing the albedo in GOSI9 reduces the summer melt allowing a better representation of the minimum sea ice area. As a result of improving the sea ice area in Summer, the temperature and salinity biases are also reduced (Fig. 8 and Fig. 10). Sea ice distribution in the southern hemisphere is also improved (Fig. 14d-f), though concentrations close to Antarctica in the Weddell, Bellingshausen, Amundsen and Somov Seas are all lower than observed. In the model low concentrations of ice not present in the observations also extend east of the
 415 Weddell Sea into the Lazarev Sea.

Overall, GOSI9 exhibits a more consistent and realistic representation of sea ice across the different resolutions than GO6+GSI8, for which the $\frac{1}{12}^\circ$ configuration was more realistic than the other two resolutions.

5 Evaluation of model performance in specific regions of interest

Since GO6, concerns have been raised as to whether biases in specific regions of interest are limiting model performance,
 420 particularly in coupled forecasts and climate projections. In response JMMP have established process evaluation groups (PEGs)




Figure 10. Sea surface salinity anomalies against EN4 v1.1 average 1995-2014 (Good et al., 2013) for GOSI9 and differences between GOSI9 and GO6. Model fields are time-means over the third decade of the integrations.

whose remit is to identify the causes of biases and to advise or deliver improvements to the model. To facilitate validations against observations and model comparison, key metrics have been identified and developed in a set of validation tools. These validation tools, MARINE_VAL, are available at https://github.com/JMMP-Group/MARINE_VAL. In this section we evaluate the improvements in model performance in three key regions.

425 5.1 Southern Ocean and Antarctic Circumpolar Current

As described in Section 2.5.6, significant work has been done to try to reduce large scale biases in the Southern Ocean in the coupled models with eddying ocean model resolution. In this section we show the impact of these changes in the forced model. The coupled results will be described in detail in a separate publication but here we also show results from a recent version of the HadGEM3 coupled model for illustration and comparison with the forced model results.

430 The biases are characterised by scalar metrics, detailed in the caption to Figure 15, in order to capture their time evolution. The immediate goal of the Southern Ocean package of changes in GOSI9 is to damp the overactive subpolar gyres in the Southern Ocean. The gyres in the Weddell Sea and Ross Sea are both too active at GO6 eddying resolutions and reduced in strength at GOSI9 (Figure 15 (b) and (c)). This change is primarily due to the Southern Ocean package, although the change




Figure 11. Annual minimum mixed layer depth (metres): anomalies against de Boyer Montégut et al. (2004) climatology for GOSI9 and GO6 and differences between GOSI9 and GO6. Model fields are time-means over the third decade of integration.

to 4th order advection in the horizontal also has a beneficial impact (not shown). The results for the coupled model (eddy-
 435 permitting ocean resolution, present-day forcing) show that the biases are larger than for the forced models, due to coupled
 model feedbacks, and are significantly reduced by the Southern Ocean package of changes.

The metrics for the subpolar gyre strengths are based on the local maximum of the barotropic streamfunction. By construction, this is the sum of the transport in the circumpolar Antarctic Slope Current (ASC) and the recirculating transport in the gyre. A large part of the reduction in this transport metric is due to a reduction in the strength of the ASC. This has
 440 the additional effect of reducing an unrealistically large westward flow at the southern boundary of the Drake Passage and hence increasing the net eastward transport in the Drake Passage (Figure 15 (a)). The GO6 models at all resolutions have net Drake Passage transport which is too small compared to the estimate of Donohue et al. (2016), with the largest bias at eddy-permitting resolution. The biases are significantly reduced at GOSI9 for the eddying models. Again, the biases in the coupled eddy-permitting model are larger than in the equivalent forced model, and there is a similar reduction with the use of
 445 the Southern Ocean package.

The very strong ASC in the eddying models tends to act as a barrier to exchange of water and ice between the Antarctic shelf and the open ocean (as also noted by Beadling et al. (2022)). This tends to lead to biases in the water mass properties




Figure 12. Annual maximum mixed layer depth (metres): anomalies against de Boyer Montégut et al. (2004) climatology for GOSI9 and GO6 and differences between GOSI9 and GO6. Model fields are time-means over the third decade of integration.

on the shelves. Two examples are shown here: the deep shelf water in the western Weddell sea tends to be too fresh in the coupled model integrations. This is a region of deep water formation and the freshening indicates that the process of deep water formation is tending to shut down in the model. The Southern Ocean package reduces this bias to some extent. The forced models have a smaller fresh bias in this region, which is somewhat reduced at GOSI9 compared to GO6. The other example is in the Amundsen Sea where relatively warm and salty circumpolar deep water (CDW) impinges onto the shelf. In the coupled model at eddy permitting resolution, the strong ASC acts as a barrier to this and we see cold biases developing in this region, which are partially alleviated by the Southern Ocean package. In the forced models in this case we don't see the same biases: if anything the water in this region is slightly too warm, and there is little difference between GO6 and GOSI9.

5.2 North Atlantic

The North Atlantic and its subpolar gyre is a key region for European weather and climate, and is also where some of the largest biases on all three model resolutions occur. It is a dynamically active region where we see significant differences in ocean currents and temperature and salinity biases between the different resolutions. In the GO6 1° configuration a strong cold bias in excess of -3 K manifests to the north of the Gulf Stream and east of Newfoundland (Fig. 8). All three resolutions




Figure 13. Mean seasonal cycles for integrated sea ice area (a,c) and sea ice volume (b,d) for the Northern and Southern hemispheres for GOSI9 and GO6-GSI8.1 for eORCA025. The meaning period is 1995-2014. In the panels representing the sea ice area (a and c), HadISST analysis (Titchner and Rayner, 2014) is represented in grey. While in the Northern Hemisphere volume (b), PIOMAS reanalysis (Zhang and Rothrock, 2003) is represented in grey. Grey dotted lines show mean $\pm \sigma$ and grey dashed lines show the mean $\pm 20\%$.




Figure 14. September mean for Arctic sea ice concentration (top) and February mean for Southern Ocean (bottom) averaged for 1995-2014 in GOSI9 eORCA025 (left), GO6-GSI8.1 eORCA025 (right) and HadISST (middle) analysis (Titchner and Rayner, 2014).




Figure 15. Time series of Southern Ocean metrics for the 30 years of GOSI9 and GO6 integrations compared with 30 years of a present-day forcing integration of two prototypes of the GC5 coupled model (N216-eORCA025) with and without the Southern Ocean Package described in Section 2.5.6 . The plotted quantities are annual means. Observational estimates and uncertainties are plotted as the black dots and lines to the right of the time series plots. From top left: a) the net eastward transport in the Drake Passage compared to the estimate of Donohue et al. (2016); b) the transport of the Weddell gyre as indicated by the maximum streamfunction in the WG box, compared to the estimate of Klatt et al. (2005); c) the transport of the Ross gyre as indicated by the maximum streamfunction in the RG box, compare to the estimate of Dotto et al. (2018); d) the salinity below 400 m spatially averaged over the WWED box in the western Weddell Sea; e) the temperature below 400 m averaged over the AMU box in the Amundsen Sea. For d and e, observational estimates are calculated from EN4.2.2.g10 profile dataset (Good et al., 2013) using means and standard deviations for all the profiles with data below 400 m in the defined boxes.

exhibit a warm bias throughout the rest of the subpolar gyre, which is in excess of 3 K in the eastern lobe of the subpolar gyre. Changes introduced in GOSI9 have resulted in a substantial reduction of this warm bias in all three resolutions. The cold bias at 1° remains similar in extent and magnitude, and where the warm bias has been corrected a more muted version of this cold

bias is also now present at $\frac{1}{4}^\circ$. Strong SST gradients exist across the Gulf Stream separation and North Atlantic Current, so
465 biases of this magnitude can easily arise from biases in the position of the main current pathways in this region.

The Gulf Stream (GS) separation is known to be sensitive to resolution (Chassignet and Marshall, 2008; Marzocchi et al., 2015). Chassignet and Marshall (2008) states that a resolution on the order of at least $1/10^\circ$ is a necessary condition for a western boundary current to realistically separate from the coast. Figure 16f shows GS separation latitude for the three GOSI9 resolutions. The GS pathway is diagnosed using the 15°C isotherm at 200m following (Seidov et al., 2019), and the
470 latitude of the GS at 72°W is displayed. In the 1° configuration, the GS separates too far north at 38.5°N and fails to deflect northwards. The position of the North Atlantic Current (NAC) is calculated with a method similar to the one used to diagnose the GS position but using the 10°C isotherm at 50m depth. The latitude of the NAC at 41°W is shown in Fig. 16g. In the 1° configuration the NAC position is 7° further south than observations. The poor representation of the GS and NAC in the 1° model was already highlighted in GO6 (Storkey et al., 2018) and is a known issue in eddy-parameterising models (Zhang and Vallis, 2007). The lack of northward advection of warm and salty water from the GS results in fresh and cold biases off
475 Newfoundland (Fig. 8,9,10a and 16c). In the $\frac{1}{4}^\circ$, as in the 1° , the GS separates too far north but the position of the NAC is noticeably improved compared with the 1° (Fig. 16g). At $\frac{1}{12}^\circ$, the GS does not overshoot but separates further south than observations. The separation too far from the coast in the $\frac{1}{12}^\circ$ was already present in GO6 and in the coupled model based on GO6 (Grist et al., 2021). The pathway of the NAC in $\frac{1}{12}^\circ$ is better represented than in the lower resolution models and brings
480 warmer and saltier water into the North Atlantic subpolar gyre (Fig. 8 and 10). Compared with GO6, there is no significant change in the GS separation latitude. However, GOSI9 exhibits a southward shift of the NAC compared with GO6. In GO6, the position of the NAC in $\frac{1}{4}^\circ$ and $\frac{1}{12}^\circ$ varies between 52°N and 56°N (not shown) further north than observations. In GOSI9, the position of the NAC is in better agreement with observations, and is mainly due to the impact of the 4^{th} order advection on the steering around Grand Banks (not shown). The additional GM poleward of 50° and the upgrade to NEMO4 also contribute to
485 the southward shift but to a lesser extent (not shown). As a result of this southward shift of the NAC, the northward advection of heat and salt is reduced in GOSI9. It reduces the warm and salty bias present in the subpolar gyre in GO6 and highlighted in Treguier et al. (2005) and Marzocchi et al. (2015). Marzocchi et al. (2015) linked the warm and salty bias to the strength of the cyclonic subpolar gyre. However, the reduction of the temperature and salinity biases in GOSI9 is not associated with a change in the strength of the SPG (not shown).

490 The time series of the annual-mean AMOC at 26°S N are shown in Fig. 16b. Observations from the RAPID array (Moat et al., 2020) between 2004-2018 estimate the transport to be 17.1 Sv with an estimated annual-mean rms uncertainty of 1.5 Sv. Transport estimates prior to RAPID suggest that the AMOC was slightly stronger during the 1980s and 1990s, with net transport from a 1992 hydrographic section calculated as 19.4 Sv (Bryden et al., 2005). Annual mean AMOC transports in the $\frac{1}{4}^\circ$ and $\frac{1}{12}^\circ$ GO6 models were unrealistic (Storkey et al., 2018), peaking at 26 Sv and 29 Sv respectively in the mid 1990s. Values in
495 2005 remained outside of observations, at 21 Sv and 23 Sv. In contrast, 1° showed excellent agreement with observations, with a peak transport of around 20 Sv in the mid-late 1990s, reducing to 17 Sv in 2005. Improvements in GOSI9 have reduced the AMOC in $\frac{1}{4}^\circ$ and $\frac{1}{12}^\circ$ so that both now peak at 20-21 Sv in 1995 and reduce to values around 17 Sv in 2005. The reduction in AMOC is associated with a decrease of the deep bias in the subpolar gyre and GIN seas mixed layer depth (Fig. 12). However,




Figure 16. Time series of North Atlantic metrics for the 30 years of GOSI9 integrations. The plotted quantities are annual means. When available, observational estimates and uncertainties are plotted as the black dots and lines to the right of the time series plots. From top left: a) Subpolar gyre strength (Sv); b) AMOC at 26.5° N at maximum depth (Sv) compared to 2004-2008 RAPID array (Moat et al., 2020); c) Mean sea surface temperature over the box off Newfoundland ($^{\circ}$ C); d) Mean mixed layer depth in the Labrador Sea box (m) compared to de Boyer Montégut et al. (2004); e) Mean sea surface salinity in the Labrador Sea box ; f) Gulf Stream separation latitude at 72° W ($^{\circ}$ N) calculated using the 15° C isotherm at 200 m following Seidov et al. (2019). g) North Atlantic Current latitude at 41° W ($^{\circ}$ N) calculated using at 10° C isotherm at 50 m depth following similar method used for the Gulf Stream separation. For c,e,f and g, observational estimates are calculated from EN4.2.2.g10 profile dataset (Good et al., 2013) using means and standard deviation

the convective overturning in the Labrador Sea is still too high in both the $\frac{1}{4}^{\circ}$ and $\frac{1}{12}^{\circ}$ configurations (Fig 16d). This suggests
500 that the AMOC in $\frac{1}{4}^{\circ}$ and $\frac{1}{12}^{\circ}$ is driven too much by excessive deep mixing in the Labrador Sea as previously noted by Megann
et al. (2014). The excessive deep convection in the Labrador Sea (Fig 16d) is common in ocean numerical model (Courtois
et al., 2017). Chanut et al. (2008) highlights the crucial role played by eddies in the restratification process but even at $\frac{1}{12}^{\circ}$
these eddies are not fully resolved. Unfortunately, AMOC volume transport in 1° is now too weak, around 16 Sv in 1995 and
13-14 Sv in 2005. Sensitivity experiments were carried out with 1° to understand the changes in AMOC strength from GO6 to
505 GOSI9, and the changes in AMOC are mainly driven by the increased albedos in GOSI9 and the improved the sea ice cover
(not shown). We speculate that in GO6 1° may have been getting the right answer for the wrong reason, with the unrealistically

low sea ice coverage in late summer-early winter exposing the surface of the ocean and leading to strong heat loss, strong deep convection, and a more vigorous AMOC. There are less than two years of overlap between the RAPID observation-based estimates of the strength of the AMOC and the reference simulations. This will be addressed in the next development cycle, for which we will update the forcing product used. It is also worth noting that the strength of the AMOC in numerical simulations is known to be sensitive both to the choice of forcing product and to the strength of numerical mixing at tropical and subtropical latitudes (e.g. Megann and Storkey, 2021), so care should be taken not to try to overfit transport in forced ocean models to the observations.

5.3 North West Pacific

The Kuroshio is the Western boundary current in the North Pacific subtropical gyre and plays an important role in transporting heat poleward. Following methods described in Qiu et al. (2014) and Qiu and Chen (2005), the position and strength of the Kuroshio is calculated for the 30-y GOSI9 integrations at all resolution and compared with sea surface height observations from CMEMS for the period 1994-2006 (<https://doi.org/10.48670/moi-00149>). The position of the Kuroshio extension in the 1° configuration (38° N) and in the $\frac{1}{4}^\circ$ configuration (37° N) is further north than in the observations (35° N) while the $\frac{1}{12}^\circ$ configuration is in better agreement with the observations (Fig. 17a). An et al. (2023) shows that coupled models with eddy rich ocean models better depict the Kuroshio extension dynamic and thermal structure than models with lower resolution ocean. Guo et al. (2003) show that as the resolution increases, the path, intensity, and vertical structure of the Kuroshio improves. They observed a northwards overshoot with the lower resolution model. They highlight that with increased resolution, the better representation of topography results in better reproduction of the interaction between baroclinicity and bottom topography. In addition to a better representation of the position of the Kuroshio extension, in the $\frac{1}{12}^\circ$ configuration, the intensity of the current is in better agreement with the observations than in the 1° and $\frac{1}{4}^\circ$ configurations (Fig. 17b). It is noted that in the 1° , the intensity of the Kuroshio is very weak compared to the observations. The subarctic frontal zone formed by the convergence of cold water from the Oyashio and warm water from the Kuroshio is shifted northward in the two lower resolution models and this results in a warm anomaly (Fig. 9).

6 Summary and discussions

GOSI9 is the latest UK global ocean and sea ice configuration developed by the JMMP partnership (Met Office, National Oceanography Centre, British Antarctic Survey, and Centre for Polar Observation and Modelling), superseding GO6 (Storkey et al., 2018). GOSI9 is a traceable hierarchy of three horizontal resolutions 1° , $\frac{1}{4}^\circ$ and $\frac{1}{12}^\circ$, and is based on the NEMO version 4.0.4 code (Madec and system team, 2019). The upgrade to NEMO 4.0.4 includes a new sea ice model SI³ (Sea Ice modelling Integrated Initiative) and faster integration achieved through the use of partially implicit schemes that allow a significant increase in the length of the time step. Other developments include the upgrade to TEOS10 equation of state, reduction in numerical mixing, and improved representation of the Southern Ocean. The interactive icebergs used in GO6 (Bigg et al., 1997; Martin and Adcroft, 2010) are switched off in GOSI9 due to stability issues, especially while testing with the coupled




Figure 17. Time series of Kuroshio metrics for the 30 years of GOSI9 integrations and for 13 years of observations from Altimeter satellite gridded Sea Level Anomalies (E.U. Copernicus Marine Service Information (CMEMS). Marine Data Store (MDS). DOI: 10.48670/moi-00149). Annual (thick lines) and monthly (thin lines) are plotted. (a) Kuroshio extension zonal averaged latitude position in 140°-165°E calculated following Qiu et al. (2014); Qiu and Chen (2005). (b) Kuroshio extension strength calculated using the sea surface height difference across the Kuroshio extension as in Qiu et al. (2014).

model. It is replaced by an iceberg melt climatology. The impact of these changes, principally in the $\frac{1}{4}^\circ$ are presented. The
 540 upgrade to NEMO 4.0.4 has a large impact with a significant reduction of the temperature and salinity biases (Fig. 3b and Fig.
 4b). Adopting the 4th order horizontal and vertical advection reduces the numerical mixing, helping to minimise the cold bias
 developing below 200 m (Fig. 3c and f). While the changes introduced to improve the representation of the Southern Ocean in
 the $\frac{1}{4}^\circ$ and $\frac{1}{12}^\circ$ configurations have limited impact on the warm SST bias in the forced model, it results in a stronger and more
 realistic ACC transport and a reduction of the temperature and salinity biases along the shelf of Antarctica (Fig. 15).

545 Tests with the coupled model were carried out in the early phase of GOSI9 development. It allowed us to integrate further
 tuning required for the coupled model early during the development cycle. In particular, after testing with the coupled model,
 change in TKE mixing depth and increased chlorophyll concentration were introduced to reduce subsurface biases. These two
 changes have a positive impact on both forced and coupled models.

The results from the 30-year integrations forced by the CORE2 dataset are presented for the three resolutions and compared
 550 against the GO6 integrations. In GOSI9, significant reductions in temperature and salinity drifts from initial condition are
 realised. Below 200 m these are mainly due to the upgrade to NEMO 4.0.4 and 4th order advection and in the top 200 m

primarily as a result of the change in chlorophyll (Figure 3). The global warm SST bias is reduced in the tropics and Arctic whilst minimal changes are observed in the Southern Ocean. Large improvements in Arctic surface temperature and salinity are linked to the improved sea ice representation. In particular, the excessive and unrealistic Arctic summer sea ice melt in 555 GO6 is significantly improved in GOSI9 (Figure 14), and can be attributed to the change in the sea ice model to SI³ and to the higher albedos which increased sea ice thickness.

The next round of development will focus on improving model fidelity in the North Atlantic and will include work already carried out to improve the representation of the Nordic overflows. Bruciaferri et al. (2024) combined the idea of Colombo (2018) with the multi-envelope approach of Bruciaferri et al. (2018) to successfully implement localised terrain following 560 coordinates in the Nordic overflow region. Their generalised approach has been tested with the GOSI9 $\frac{1}{4}^\circ$ configuration and is shown to significantly improve the realism of the Nordic overflows in simulations, reducing spurious cross-isopycnal mixing in this region of strong gravity currents. However, large scale salinity biases along the bathymetry of the subpolar gyre impact the mass properties of the water cascading. Improving the salinity bias and using appropriate coordinates are both key to improving the representation of the Nordic overflows. Another key challenge is to realistically represent the separation latitude of the Gulf 565 Stream. In the GOSI9 configuration, increasing the resolution is shown to improve the realism of the Gulf Stream separation latitude and the North Atlantic Current path. However, even with the $\frac{1}{12}^\circ$ it remains challenging. Work looking at the sensitivity of the Gulf Stream to the vertical coordinate system is ongoing (Bruciaferri et al., 2022) with the same approach used for the Nordic overflows and in parallel we plan to develop a $\frac{1}{12}^\circ$ configuration with a two-way nested AGRIF zoom with $\frac{1}{36}^\circ$ in the North Atlantic.

In GOSI9, progress have been made in the Southern Ocean with increased ACC transport and reduced temperature and salinity biases on the Antarctic shelf but ice-shelf cavities have remained parameterised. For climate applications requiring open ice cavities, such as UK Earth System Model (UKESM) configurations (Smith et al., 2021), improving water mass properties on the shelf is essential. Using a 1° forced configuration, Hutchinson et al. (2023) show that explicitly simulating the circulation beneath the largest ice shelves improves the realism of the Antarctic continental shelf circulation. However, 575 they note that the impact on the representation of the Antarctic Bottom Water is limited by the absence of realistic overflows. Recent work under the NERC funded ORCHESTRA project showed the merits of using a vertical sigma-coordinate around the Antarctic shelf region to greatly reduce temperature and salinity biases on the shelf and to preserve the density of Antarctic bottom water (Meijers et al., 2023). For the next configuration, we plan to open the ice-shelf cavities, taking advantage of the developments available at NEMO 4.2, and adopt the strategy used in the Nordic overflows using local terrain following 580 coordinates to improve the Antarctic overflows.

In GOSI9, the bathymetries for each configuration originate from a different source (see 2.1). In this respect the hierarchy of resolutions is not fully traceable. For future GOSI configurations, work is ongoing to produce traceable bathymetry for each resolutions.

In GOSI9, the constant value used for the chlorophyll concentration was tuned to better match climatological values in the 585 Tropics. For future configurations, the constant value will be replaced by a monthly chlorophyll concentration climatology derived from ocean colour observations that will better account for the variation in the solar penetration due to the large

spatial and seasonal variability. For the coupled model, it will allow a consistent approach between the ocean models and the atmosphere models, where a varying chlorophyll concentration is already used to compute the albedo.

590 The vertical mixing closure schemes currently available in NEMO underestimate the impact of important sources of mixing such as Langmuir turbulence and maximum turbulence due to shear at the base of the mixed layer. These sources of mixing are important for the near surface mixing globally (Belcher et al., 2012). As part of the UK OSMOSIS project a more physically based mixing scheme has been developed and is being implemented in NEMO. We expect this will replace the existing TKE scheme in the next configuration.

Code and data availability. GOSI9 official release is available to download (Guiavarc’h and Storkey, 2024). For each configuration, it includes code, namelists, links to download the input files and scripts to run the configurations.

Each of the CORE2-forced 1976–2005 reference simulations have been archived to the CEDA archive (NERC’s Environmental Data Service) and are available to download: eORCA1 (Blaker et al., 2023), eORCA025 (Guiavarc’h et al., 2023a), eORCA12 (Guiavarc’h et al., 2023b).

600 This study has been conducted using E.U. Copernicus Marine Service Information: Global Ocean Gridded L 4 Sea Surface Heights And Derived Variables Nrt (<https://doi.org/10.48670/moi-00149>).

Author contributions. CG prepared the manuscript with contributions from co-authors. CG, DS, and AB performed and analysed the main assessment integrations. EB, AM, HH, MB, DCa, DCo, BS, PM, SM and BA were involved in the development of the GOSI9 configurations, performed sensitivity experiments and assisted with the evaluation of the main integrations.

Competing interests. At least one of the (co-)authors is a member of the editorial board of Geoscientific Model Development.

605 *Acknowledgements.* This work was developed as part of the Joint Marine Modelling Programme (JMMP), a partnership between the Met Office, National Oceanography Centre, British Antarctic Survey and Centre for Polar Observation and Modelling. The authors were supported by the Met Office Hadley Centre Climate Programme funded by DSIT. This work was funded by the Met Office Weather and Climate Science for Service Partnership (WCSSP) India project which is supported by the Department for Science, Innovation & Technology (DSIT) and by the Met Office Climate Science for Service Partnership (CSSP) China project under the International Science Partnerships Fund (ISPF). The contributions from the National Oceanography Centre were funded by the Natural Environment Research Council (NERC) under the Climate Linked Atlantic Sector Science (CLASS) marine research programme (NE/R015953/1), as part of the Joint Marine Modelling Programme (JMMP).

References

- Adcroft, A. and Campin, J. M.: Rescaled height coordinates for accurate representation of free-surface flows in ocean circulation models, *Ocean Modelling*, 7, <https://doi.org/10.1016/j.ocemod.2003.09.003>, 2004.
- Adcroft, A., Hill, C., and Marshall, J.: Representation of topography by shaved cells in a height coordinate ocean model, *Monthly Weather Review*, 125, [https://doi.org/10.1175/1520-0493\(1997\)125<2293:ROTBSC>2.0.CO;2](https://doi.org/10.1175/1520-0493(1997)125<2293:ROTBSC>2.0.CO;2), 1997.
- Amante, C. and Eakins, B. W.: ETOPO1 Global Relief Model converted to PanMap layer format, NOAA-National Geophysical Data Center, 2009.
- An, B., Yu, Y., Hewitt, H., Wu, P., Furtado, K., Liu, H., Lin, P., Luan, Y., and Chen, K.: The benefits of high-resolution models in simulating the Kuroshio Extension and its long-term changes. , *Clim Dyn*, <https://doi.org/https://doi.org/10.1007/s00382-023-06862-z>, 2023.
- Arakawa, A. and Lamb, V. R.: A potential enstrophy and energy conserving scheme for the shallow water equations., *Monthly Weather Review*, 109, [https://doi.org/10.1175/1520-0493\(1981\)109<0018:APEAEC>2.0.CO;2](https://doi.org/10.1175/1520-0493(1981)109<0018:APEAEC>2.0.CO;2), 1981.
- Arndt, J. E., Schenke, H. W., Jakobsson, M., Nitsche, F. O., Buys, G., Goleby, B., Rebesco, M., Bohoyo, F., Hong, J., Black, J., Greku, R., Udintsev, G., Barrios, F., Reynoso-Peralta, W., Taisei, M., and Wigley, R.: The international bathymetric chart of the Southern Ocean (IBCSO) version 1.0-A new bathymetric compilation covering circum-Antarctic waters, *Geophysical Research Letters*, 40, <https://doi.org/10.1002/grl.50413>, 2013.
- Axell, L. B.: Wind-driven internal waves and Langmuir circulations in a numerical ocean model of the southern Baltic Sea, *Journal of Geophysical Research: Oceans*, 107, <https://doi.org/10.1029/2001jc000922>, 2002.
- Barnier, B., Madec, G., Penduff, T., Molines, J. M., Treguier, A. M., Le Sommer, J., Beckmann, A., Biastoch, A., Böning, C., Dengg, J., Derval, C., Durand, E., Gulev, S., Remy, E., Talandier, C., Theetten, S., Maltrud, M., McClean, J., and De Cuevas, B.: Impact of partial steps and momentum advection schemes in a global ocean circulation model at eddy-permitting resolution, *Ocean Dynamics*, 56, 543–567, <https://doi.org/10.1007/s10236-006-0082-1>, 2006.
- Beadling, R. L., Krasting, J. P., Griffies, S. M., Hurlin, W. J., Bronselaer, B., Russell, J. L., MacGilchrist, G. A., Tesdal, J.-E., and Winton, M.: Importance of the Antarctic Slope Current in the Southern Ocean Response to Ice Sheet Melt and Wind Stress Change, *Journal of Geophysical Research: Oceans*, 127, e2021JC017608, <https://doi.org/https://doi.org/10.1029/2021JC017608>, e2021JC017608, 2022.
- Beckmann, A. and Döscher, R.: A method for improved representation of dense water spreading over topography in geopotential-coordinate models, *Journal of Physical Oceanography*, 27, [https://doi.org/10.1175/1520-0485\(1997\)027<0581:AMFIRO>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<0581:AMFIRO>2.0.CO;2), 1997.
- Belcher, S. E., Grant, A. L. M., Hanley, K. E., Fox-Kemper, B., Van Roekel, L., Sullivan, P. P., Large, W. G., Brown, A., Hines, A., Calvert, D., Rutgersson, A., Pettersson, H., Bidlot, J.-R., Janssen, P. A. E. M., and Polton, J. A.: A global perspective on Langmuir turbulence in the ocean surface boundary layer, *Geophysical Research Letters*, 39, <https://doi.org/https://doi.org/10.1029/2012GL052932>, 2012.
- Bernie, D. J., Woolnough, S. J., Slingo, J. M., and Guilyardi, E.: Modeling diurnal and intraseasonal variability of the ocean mixed layer, *Journal of Climate*, 18, <https://doi.org/10.1175/JCLI3319.1>, 2005.
- Bernie, D. J., Guilyardi, E., Madec, G., Slingo, J. M., and Woolnough, S. J.: Impact of resolving the diurnal cycle in an ocean-atmosphere GCM. Part 1: A diurnally forced OGCM, *Climate Dynamics*, 29, <https://doi.org/10.1007/s00382-007-0249-6>, 2007.
- Bigg, G. R., Wadley, M. R., Stevens, D. P., and Johnson, J. A.: Modelling the dynamics and thermodynamics of icebergs, *Cold Regions Science and Technology*, 26, [https://doi.org/10.1016/S0165-232X\(97\)00012-8](https://doi.org/10.1016/S0165-232X(97)00012-8), 1997.

- Bitz, C. M. and Lipscomb, W. H.: An energy-conserving thermodynamic model of sea ice, *Journal of Geophysical Research: Oceans*, 104, 650 <https://doi.org/10.1029/1999jc900100>, 1999.
- Blaker, A., Blockley, E., Copsey, D., Guiavarc'h, C., Hewitt, H., Megann, A., and Storkey, D.: Ocean and sea ice model output from GOSI9 eORCA1 (1976-2005), NERC EDS British Oceanographic Data Centre NOC, <https://doi.org/10.5285/053d6318-59b2-6aa8-e063-6c86abc06f23>, 2023.
- Blockley, E., Fiedler, E., Ridley, J., Roberts, L., West, A., Copsey, D., Feltham, D., Graham, T., Livings, D., Rousset, C., Schroeder, D., and 655 Vancoppenolle, M.: The sea ice component of GC5: coupling SI³ to HadGEM3 using conductive fluxes, *Geoscientific Model Development*, 17, 6799–6817, <https://doi.org/10.5194/gmd-17-6799-2024>, 2024.
- Bourdallé-Badie, R. and Treguier, A.: A climatology of runoff for the global ocean-ice model ORCA025, Mercator-Ocean report, MOO-RP-425-365-MER, 2006.
- Brodeau, L., Barnier, B., Gulev, S. K., and Woods, C.: Climatologically significant effects of some approximations in the bulk parameteriza- 660 tions of turbulent air-sea fluxes, *Journal of Physical Oceanography*, 47, <https://doi.org/10.1175/JPO-D-16-0169.1>, 2017.
- Brown, A., Milton, S., Cullen, M., Golding, B., Mitchell, J., and Shelly, A.: Unified modeling and prediction of weather and climate: A 25-year journey, *Bulletin of the American Meteorological Society*, 93, <https://doi.org/10.1175/BAMS-D-12-00018.1>, 2012.
- Bruciaferri, D., Shapiro, G. I., and Wobus, F.: A multi-envelope vertical coordinate system for numerical ocean modelling, *Ocean Dynamics*, 68, <https://doi.org/10.1007/s10236-018-1189-x>, 2018.
- 665 Bruciaferri, D., Hewitt, H. T., Bell, M. J., Guiavarc'h, C., Storkey, D., Roberts, M. J., and Jackson, L.: Sensitivity of the Western North Atlantic circulation to the vertical coordinate system in global ocean models, <https://usclivar.org/meetings/gulf-stream-workshop>, 2022.
- Bruciaferri, D., Guiavarc'h, C., Hewitt, H. T., Harle, J., Almansì, M., Mathiot, P., and Colombo, P.: Localized General Vertical Coordinates for Quasi-Eulerian Ocean Models: The Nordic Overflows Test-Case, *Journal of Advances in Modeling Earth Systems*, 16, e2023MS003 893, <https://doi.org/https://doi.org/10.1029/2023MS003893>, e2023MS003893 2023MS003893, 2024.
- 670 Bryden, H. L., Longworth, H. R., and Cunningham, S. A.: Slowing of the Atlantic meridional overturning circulation at 25N, *Nature*, 438, <https://doi.org/10.1038/nature04385>, 2005.
- Chanut, J., Barnier, B., Large, W., Debreu, L., Penduff, T., Molines, J. M., and Mathiot, P.: Mesoscale eddies in the Labrador Sea and their contribution to convection and restratification, *Journal of Physical Oceanography*, 38, <https://doi.org/10.1175/2008JPO3485.1>, 2008.
- Chassignet, E. P. and Marshall, D. P.: Gulf stream separation in numerical ocean models, in: *Geophysical Monograph Series*, vol. 177, 675 <https://doi.org/10.1029/177GM05>, 2008.
- Colombo, P.: Modelling dense water flows through sills in large scale realistic ocean models: demonstrating the potential of a hybrid geopotential/terrain-following vertical coordinate, Ph.D. thesis, Université Grenoble Alpes, <http://www.theses.fr/2018GREAU017>, 2018.
- Courtois, P., Hu, X., Pennelly, C., Spence, P., and Myers, P. G.: Mixed layer depth calculation in deep convection regions in ocean numerical models, *Ocean Modelling*, 120, <https://doi.org/10.1016/j.ocemod.2017.10.007>, 2017.
- 680 Craig, P. D. and Banner, M. L.: Modeling wave-enhanced turbulence in the ocean surface layer, *Journal of Physical Oceanography*, 24, [https://doi.org/10.1175/1520-0485\(1994\)024<2546:MWETIT>2.0.CO;2](https://doi.org/10.1175/1520-0485(1994)024<2546:MWETIT>2.0.CO;2), 1994.
- de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology, *Journal of Geophysical Research: Oceans*, 109, <https://doi.org/10.1029/2004JC002378>, 2004.
- 685 Donohue, K. A., Tracey, K. L., Watts, D. R., Chidichimo, M. P., and Chereskin, T. K.: Mean Antarctic Circumpolar Current transport measured in Drake Passage, *Geophysical Research Letters*, 43, <https://doi.org/10.1002/2016GL070319>, 2016.

- Dotto, T. S., Naveira Garabato, A., Bacon, S., Tsamados, M., Holland, P. R., Hooley, J., Frajka-Williams, E., Ridout, A., and Meredith, M. P.: Variability of the Ross Gyre, Southern Ocean: Drivers and Responses Revealed by Satellite Altimetry, *Geophysical Research Letters*, 45, 6195–6204, <https://doi.org/https://doi.org/10.1029/2018GL078607>, 2018.
- 690 Ducouso, N., Le Sommer, J., Molines, J.-M., and Bell, M.: Impact of the “Symmetric Instability of the Computational Kind” at mesoscale- and submesoscale-permitting resolutions, *Ocean Modelling*, 120, 18–26, <https://doi.org/https://doi.org/10.1016/j.ocemod.2017.10.006>, 2017.
- Gaspar, P., Grégoris, Y., and Lefevre, J.-M.: A simple eddy kinetic energy model for simulations of the oceanic vertical mixing: Tests at station Papa and long-term upper ocean study site, *Journal of Geophysical Research*, 95, <https://doi.org/10.1029/jc095ic09p16179>, 1990.
- 695 Gent, P. R. and McWilliams, J. C.: Isopycnal Mixing in Ocean Circulation Models, *Journal of Physical Oceanography*, 20, [https://doi.org/10.1175/1520-0485\(1990\)020<0150:imioem>2.0.co;2](https://doi.org/10.1175/1520-0485(1990)020<0150:imioem>2.0.co;2), 1990.
- Good, S. A., Martin, M. J., and Rayner, N. A.: EN4: Quality controlled ocean temperature and salinity profiles and monthly objective analyses with uncertainty estimates, *Journal of Geophysical Research: Oceans*, 118, <https://doi.org/10.1002/2013JC009067>, 2013.
- Gregg, M. C., Sanford, T. B., and Winkel, D. P.: Reduced mixing from the breaking of internal waves in equatorial waters, *Nature*, 422, 700 <https://doi.org/10.1038/nature01507>, 2003.
- Grist, J. P., Josey, S. A., Sinha, B., Catto, J. L., Roberts, M. J., and Coward, A. C.: Future Evolution of an Eddy Rich Ocean Associated with Enhanced East Atlantic Storminess in a Coupled Model Projection, *Geophysical Research Letters*, 48, <https://doi.org/10.1029/2021GL092719>, 2021.
- Guiavarc’h, C. and Storkey, D.: JMMP-Group/GO_RELEASES: GOSI9 release, <https://doi.org/10.5281/zenodo.13814369>, 2024.
- 705 Guiavarc’h, C., Blaker, A., Blockley, E., Copsey, D., Hewitt, H., Megann, A., and Storkey, D.: Ocean and sea ice model output from GOSI9 eORCA025 (1976-2005), NERC EDS British Oceanographic Data Centre NOC, <https://doi.org/10.5285/0b38ca70-9b8f-3660-e063-6c86abc0e6da>, 2023a.
- Guiavarc’h, C., Blaker, A., Blockley, E., Copsey, D., Hewitt, H., Megann, A., and Storkey, D.: Ocean and sea ice model output from GOSI9 eORCA12 (1976-2005), NERC EDS British Oceanographic Data Centre NOC, <https://doi.org/10.5285/0b38c7f9-439d-34fb-e063-6c86abc02eca>, 2023b.
- 710 Guo, X., Hukuda, H., Miyazawa, Y., and Yamagata, T.: A triply nested ocean model for simulating the Kuroshio - Roles of horizontal resolution on JEBAR, *Journal of Physical Oceanography*, 33, [https://doi.org/10.1175/1520-0485\(2003\)033<0146:ATNOMF>2.0.CO;2](https://doi.org/10.1175/1520-0485(2003)033<0146:ATNOMF>2.0.CO;2), 2003.
- Hallberg, R.: Using a resolution function to regulate parameterizations of oceanic mesoscale eddy effects, *Ocean Modelling*, 72, 715 <https://doi.org/10.1016/j.ocemod.2013.08.007>, 2013.
- Held, I. M. and Larichev, V. D.: A scaling theory for horizontally homogeneous, baroclinically unstable flow on a beta plane, *Journal of the Atmospheric Sciences*, 53, [https://doi.org/10.1175/1520-0469\(1996\)053<0946:astfhh>2.0.co;2](https://doi.org/10.1175/1520-0469(1996)053<0946:astfhh>2.0.co;2), 1996.
- Hewitt, H. T., Roberts, M. J., Hyder, P., Graham, T., Rae, J., Belcher, S. E., Bourdallé-Badie, R., Copsey, D., Coward, A., Guiavarch, C., Harris, C., Hill, R., Hirschi, J. J., Madec, G., Mizielinski, M. S., Neining, E., New, A. L., Rioual, J. C., Sinha, B., Storkey, D., Shelly, 720 A., Thorpe, L., and Wood, R. A.: The impact of resolving the Rossby radius at mid-latitudes in the ocean: Results from a high-resolution version of the Met Office GC2 coupled model, *Geoscientific Model Development*, 9, <https://doi.org/10.5194/gmd-9-3655-2016>, 2016.
- Hollingsworth, A., Kållberg, P., Renner, V., and Burridge, D. M.: An internal symmetric computational instability, *Quarterly Journal of the Royal Meteorological Society*, 109, <https://doi.org/10.1002/qj.49710946012>, 1983.

- Hunke, E. C. and Dukowicz, J. K.: The elastic-viscous-plastic sea ice dynamics model in general orthogonal curvilinear coordinates on a sphere-Incorporation of metric terms, *Monthly Weather Review*, 130, [https://doi.org/10.1175/1520-0493\(2002\)130<1848:TEVPSI>2.0.CO;2](https://doi.org/10.1175/1520-0493(2002)130<1848:TEVPSI>2.0.CO;2), 2002.
- Hutchinson, K., Deshayes, J., Éthé, C., Rousset, C., de Lavergne, C., Vancoppenolle, M., Jourdain, N. C., and Mathiot, P.: Improving Antarctic Bottom Water precursors in NEMO for climate applications, *EGUsphere*, 2023, 1–28, <https://doi.org/10.5194/egusphere-2023-99>, 2023.
- IOC, IHO, and BODC: Centenary Edition of the GEBCO Digital Atlas, published on CD-ROM on behalf of the Intergovernmental Oceanographic Commission and the International Hydrographic Organization as part of the General Bathymetric Chart of the Oceans, *British Oceanographic Data*, 2003.
- Ioc, Scor, and Iapso: The international thermodynamic equation of seawater – 2010: Calculation and use of thermodynamic properties, 2010. JMMP: The Joint Marine Modelling Programme, <https://metoffice.sharepoint.com/sites/JointMarineModellingProgrammeExt>.
- Klatt, O., Fahrbach, E., Hoppema, M., and Rohardt, G.: The transport of the Weddell Gyre across the Prime Meridian, *Deep Sea Research Part II: Topical Studies in Oceanography*, 52, 513–528, <https://doi.org/https://doi.org/10.1016/j.dsr2.2004.12.015>, direct observations of oceanic flow: A tribute to Walter Zenk, 2005.
- Koch-Larrouy, A., Madec, G., Blanke, B., and Molcard, R.: Water mass transformation along the Indonesian throughflow in an OGCM, *Ocean Dynamics*, 58, <https://doi.org/10.1007/s10236-008-0155-4>, 2008.
- Large, W. G. and Yeager, S. G.: The global climatology of an interannually varying air - Sea flux data set, *Climate Dynamics*, 33, <https://doi.org/10.1007/s00382-008-0441-3>, 2009.
- Leclair, M. and Madec, G.: Z^* -Coordinate, an Arbitrary Lagrangian-Eulerian coordinate separating high and low frequency motions, *Ocean Modelling*, 37, <https://doi.org/10.1016/j.ocemod.2011.02.001>, 2011.
- Madec, G. and system team, N.: Nemo Ocean Engine - version 4.0.1, Notes du Pôle de modélisation de l'Institut Pierre-Simon Laplace (IPSL): (27)., <https://doi.org/10.5281/zenodo.3878122>, 2019.
- Martin, T. and Adcroft, A.: Parameterizing the fresh-water flux from land ice to ocean with interactive icebergs in a coupled climate model, *Ocean Modelling*, 34, <https://doi.org/10.1016/j.ocemod.2010.05.001>, 2010.
- Marzocchi, A., Hirschi, J. J., Holliday, N. P., Cunningham, S. A., Blaker, A. T., and Coward, A. C.: The North Atlantic subpolar circulation in an eddy-resolving global ocean model, *Journal of Marine Systems*, 142, <https://doi.org/10.1016/j.jmarsys.2014.10.007>, 2015.
- Mathiot, P., Jenkins, A., Harris, C., and Madec, G.: Explicit representation and parametrised impacts of under ice shelf seas in the z_* -coordinate ocean model NEMO 3.6, *Geoscientific Model Development*, 10, <https://doi.org/10.5194/gmd-10-2849-2017>, 2017.
- Megann, A.: Estimating the numerical diapycnal mixing in an eddy-permitting ocean model, *Ocean Modelling*, 121, <https://doi.org/10.1016/j.ocemod.2017.11.001>, 2018.
- Megann, A.: Quantifying numerical mixing in a tidally forced global eddy-permitting ocean model, *Ocean Modelling*, 188, 102 329, <https://doi.org/https://doi.org/10.1016/j.ocemod.2024.102329>, 2024.
- Megann, A. and Storkey, D.: Exploring Viscosity Space in an Eddy-Permitting Global Ocean Model: Is Viscosity a Useful Control for Numerical Mixing?, *Journal of Advances in Modeling Earth Systems*, 13, <https://doi.org/10.1029/2020MS002263>, 2021.
- Megann, A., Storkey, D., Aksenov, Y., Alderson, S., Calvert, D., Graham, T., Hyder, P., Siddorn, J., and Sinha, B.: GO5.0: the joint NERC–Met Office NEMO global ocean model for use in coupled and forced applications, *Geoscientific Model Development*, 7, 1069–1092, <https://doi.org/10.5194/gmd-7-1069-2014>, 2014.
- Megann, A., Chanut, J., and Storkey, D.: Assessment of the z^* Time-Filtered Arbitrary Lagrangian-Eulerian Coordinate in a Global Eddy-Permitting Ocean Model, *Journal of Advances in Modeling Earth Systems*, 14, <https://doi.org/10.1029/2022MS003056>, 2022.

- Meijers, A. J. S., Meredith, M. P., Shuckburgh, E. F., Kent, E. C., Munday, D. R., Firing, Y. L., King, B., Smyth, T. J., Leng, M. J., George Nurser, A. J., Hewitt, H. T., Povl Abrahamsen, E., Weiss, A., Yang, M., Bell, T. G., Alexander Brearley, J., Boland, E. J. D., Jones, D. C., Josey, S. A., Owen, R. P., Grist, J. P., Blaker, A. T., Biri, S., Yelland, M. J., Pimm, C., Zhou, S., Harle, J., and Cornes, R. C.: Finale: impact of the ORCHESTRA/ENCORE programmes on Southern Ocean heat and carbon understanding, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 381, 20220070, <https://doi.org/10.1098/rsta.2022.0070>, 2023.
- 765 Merchant, C. J., Embury, O., Roberts-Jones, J., Fiedler, E., Bulgin, C. E., Corlett, G. K., Good, S., McLaren, A., Rayner, N., Morak-Bozzo, S., and Donlon, C.: Sea surface temperature datasets for climate applications from Phase 1 of the European Space Agency Climate Change Initiative (SST CCI), *Geoscience Data Journal*, 1, <https://doi.org/10.1002/gdj3.20>, 2014.
- 770 Merino, N., Sommer, J. L., Durand, G., Jourdain, N. C., Madec, G., Mathiot, P., and Tournadre, J.: Antarctic icebergs melt over the Southern Ocean : Climatology and impact on sea ice, <https://doi.org/10.1016/j.ocemod.2016.05.001>, 2016.
- Merryfield, W. J., Holloway, G., and Gargett, A. E.: A global ocean model with double-diffusive mixing, *Journal of Physical Oceanography*, 29, [https://doi.org/10.1175/1520-0485\(1999\)029<1124:AGOMWD>2.0.CO;2](https://doi.org/10.1175/1520-0485(1999)029<1124:AGOMWD>2.0.CO;2), 1999.
- 775 Moat, B., E., F.-W., D.A., S., D., R., W.E., J., M.O., B., Volkov, D., and Collins, J.: Atlantic meridional overturning circulation observed by the RAPID-MOCHA-WBTS (RAPID-Meridional Overturning Circulation and Heatflux Array-Western Boundary Time Series) array at 26N from 2004 to 2020 (v2020.2), <https://doi.org/10.5285/e91b10af-6f0a-7fa7-e053-6c86abc05a09>, 2020.
- Mulcahy, J. P., Jones, C. G., Rumbold, S. T., Kuhlbrodt, T., Dittus, A. J., Blockley, E. W., Yool, A., Walton, J., Hardacre, C., Andrews, T., Bodas-Salcedo, A., Stringer, M., de Mora, L., Harris, P., Hill, R., Kelley, D., Robertson, E., and Tang, Y.: UKESM1.1: development and evaluation of an updated configuration of the UK Earth System Model, *Geoscientific Model Development*, 16, 1569–1600, <https://doi.org/10.5194/gmd-16-1569-2023>, 2023.
- 780 National Geophysical Data Center, N.: National Geophysical Data Center, 2006. 2-minute Gridded Global Relief Data (ETOPO2) v2., 2006.
- Qiu, B. and Chen, S.: Eddy-induced heat transport in the subtropical North Pacific from Argo, TMI, and altimetry measurements, *Journal of Physical Oceanography*, 35, <https://doi.org/10.1175/JPO2696.1>, 2005.
- Qiu, B., Chen, S., Schneider, N., and Taguchi, B.: A coupled decadal prediction of the dynamic state of the kuroshio extension system, *Journal of Climate*, 27, <https://doi.org/10.1175/JCLI-D-13-00318.1>, 2014.
- 785 Rae, J. G., Hewitt, H. T., Keen, A. B., Ridley, J. K., Edwards, J. M., and Harris, C. M.: A sensitivity study of the sea ice simulation in the global coupled climate model, HadGEM3, *Ocean Modelling*, 74, <https://doi.org/10.1016/j.ocemod.2013.12.003>, 2014.
- Ridley, J. K., Blockley, E. W., Keen, A. B., Rae, J. G., West, A. E., and Schroeder, D.: The sea ice model component of HadGEM3-GC3.1, *Geoscientific Model Development*, 11, <https://doi.org/10.5194/gmd-11-713-2018>, 2018.
- 790 Rignot, E., Jacobs, S., Mouginot, J., and Scheuchl, B.: Ice-shelf melting around antarctica, *Science*, 341, <https://doi.org/10.1126/science.1235798>, 2013.
- Roberts, M. J., Baker, A., Blockley, E. W., Calvert, D., Coward, A., Hewitt, H. T., Jackson, L. C., Kuhlbrodt, T., Mathiot, P., Roberts, C. D., Schiemann, R., Seddon, J., Vanni ere, B., and Vidale, P. L.: Description of the resolution hierarchy of the global coupled HadGEM3-GC3.1 model as used in CMIP6 HighResMIP experiments, *Geoscientific Model Development*, 12, 4999–5028, <https://doi.org/10.5194/gmd-12-4999-2019>, 2019.
- 795 Rodgers, K. B., Aumont, O., Mikaloff Fletcher, S. E., Plancherel, Y., Bopp, L., De Boyer Mont egut, C., Iudicone, D., Keeling, R. F., Madec, G., and Wanninkhof, R.: Strong sensitivity of Southern Ocean carbon uptake and nutrient cycling to wind stirring, *Biogeosciences*, 11, <https://doi.org/10.5194/bg-11-4077-2014>, 2014.

- Roullet, G. and Madec, G.: Salt conservation, free surface, and varying levels: A new formulation for ocean general circulation models, *Journal of Geophysical Research: Oceans*, 105, <https://doi.org/10.1029/2000jc900089>, 2000.
- 800 Rousset, C., Vancoppenolle, M., Madec, G., Fichefet, T., Flavoni, S., Barthélemy, A., Benshila, R., Chanut, J., Levy, C., Masson, S., and Vivier, F.: The Louvain-La-Neuve sea ice model LIM3.6: Global and regional capabilities, *Geoscientific Model Development*, 8, <https://doi.org/10.5194/gmd-8-2991-2015>, 2015.
- Seidov, D., Mishonov, A., Reagan, J., and Parsons, R.: Resilience of the Gulf Stream path on decadal and longer timescales, *Scientific Reports*, 9, <https://doi.org/10.1038/s41598-019-48011-9>, 2019.
- 805 Shchepetkin, A. F.: An adaptive, Courant-number-dependent implicit scheme for vertical advection in oceanic modeling, *Ocean Modelling*, 91, 38–69, <https://doi.org/10.1016/j.ocemod.2015.03.006>, 2015.
- Shchepetkin, A. F. and McWilliams, J. C.: The regional oceanic modeling system (ROMS): A split-explicit, free-surface, topography-following-coordinate oceanic model, *Ocean Modelling*, 9, <https://doi.org/10.1016/j.ocemod.2004.08.002>, 2005.
- 810 Simmons, H. L., Jayne, S. R., St. Laurent, L. C., and Weaver, A. J.: Tidally driven mixing in a numerical model of the ocean general circulation, *Ocean Modelling*, 6, [https://doi.org/10.1016/S1463-5003\(03\)00011-8](https://doi.org/10.1016/S1463-5003(03)00011-8), 2004.
- Smith, R. S., Mathiot, P., Siahaan, A., Lee, V., Cornford, S. L., Gregory, J. M., Payne, A. J., Jenkins, A., Holland, P. R., Ridley, J. K., and Jones, C. G.: Coupling the U.K. Earth System Model to Dynamic Models of the Greenland and Antarctic Ice Sheets, *Journal of Advances in Modeling Earth Systems*, 13, e2021MS002520, <https://doi.org/10.1029/2021MS002520>, e2021MS002520
- 815 2021MS002520, 2021.
- Stein, C. A. and Stein, S.: A model for the global variation in oceanic depth and heat flow with lithospheric age, *Nature*, 359, <https://doi.org/10.1038/359123a0>, 1992.
- Stewart, K. D., Hogg, A. M. C., Griffies, S. M., Heerdegen, A. P., Ward, M. L., Spence, P., and England, M. H.: Vertical resolution of baroclinic modes in global ocean models, *Ocean Modelling*, 113, <https://doi.org/10.1016/j.ocemod.2017.03.012>, 2017.
- 820 Storkey, D., Blaker, A. T., Mathiot, P., Megann, A., Aksenov, Y., Blockley, E. W., Calvert, D., Graham, T., Hewitt, H. T., Hyder, P., Kuhlbrodt, T., Rae, J. G., and Sinha, B.: UK Global Ocean GO6 and GO7: A traceable hierarchy of model resolutions, *Geoscientific Model Development*, 11, <https://doi.org/10.5194/gmd-11-3187-2018>, 2018.
- Storkey, D., Mathiot, P., Bell, M. J., Copsey, D., Guiavarc’h, C., Hewitt, H. T., Ridley, J., and Roberts, M. J.: Resolution dependence of interlinked Southern Ocean biases in global coupled HadGEM3 models, *EGUsphere*, 2024, 1–29, [https://doi.org/10.5194/egusphere-](https://doi.org/10.5194/egusphere-2024-1414)
- 825 2024-1414, 2024.
- Thorndike, A. S., Rothrock, D. A., Maykut, G. A., and Colony, R.: The thickness distribution of sea ice, *Journal of Geophysical Research*, 80, <https://doi.org/10.1029/jc080i033p04501>, 1975.
- Titchner, H. A. and Rayner, N. A.: The met office hadley centre sea ice and sea surface temperature data set, version 2: 1. sea ice concentrations, *Journal of Geophysical Research*, 119, <https://doi.org/10.1002/2013JD020316>, 2014.
- 830 Tréguier, A. M., Held, I. M., and Larichev, V. D.: Parameterization of Quasigeostrophic Eddies in Primitive Equation Ocean Models, *Journal of Physical Oceanography*, 27, 567 – 580, [https://doi.org/10.1175/1520-0485\(1997\)027<0567:POQEIP>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<0567:POQEIP>2.0.CO;2), 1997.
- Treguier, A. M., Theetten, S., Chassignet, E. P., Penduff, T., Smith, R., Talley, L., Beismann, J. O., and Böning, C.: The North Atlantic subpolar gyre in four high-resolution models, *Journal of Physical Oceanography*, 35, <https://doi.org/10.1175/JPO2720.1>, 2005.
- Vancoppenolle, M., Rousset, C., Blockley, E., and Group, t. N. S. I. W.: SI3 – Sea Ice modelling Integrated Initiative – The NEMO Sea Ice
- 835 Engine, Tech. rep., <https://zenodo.org/record/7534900#.Y8GIF-xKg-Q>, 2023.

- Weatherall, P., Marks, K. M., Jakobsson, M., Schmitt, T., Tani, S., Arndt, J. E., Rovere, M., Chayes, D., Ferrini, V., and Wigley, R.: A new digital bathymetric model of the world's oceans, *Earth and Space Science*, 2, <https://doi.org/10.1002/2015EA000107>, 2015.
- Williams, K. D., Copsey, D., Blockley, E. W., Bodas-Salcedo, A., Calvert, D., Comer, R., Davis, P., Graham, T., Hewitt, H. T., Hill, R., Hyder, P., Ineson, S., Johns, T. C., Keen, A. B., Lee, R. W., Megann, A., Milton, S. F., Rae, J. G. L., Roberts, M. J., Scaife, A. A., Schiemann, R., Storkey, D., Thorpe, L., Watterson, I. G., Walters, D. N., West, A., Wood, R. A., Woollings, T., and Xavier, P. K.: The Met Office Global Coupled Model 3.0 and 3.1 (GC3.0 and GC3.1) Configurations, *Journal of Advances in Modeling Earth Systems*, 10, 357–380, <https://doi.org/https://doi.org/10.1002/2017MS001115>, 2018.
- Xavier, P., Willett, M., Graham, T. and Earnshaw, P., Copsey, D., Narayan, N., Marzin, C., Zhu, H., Sellar, A., Ackerley, D., Blockley, E., Bodas-Salcedo, A., Bushell, A., Choi, N., Chua, X. R., Guiavarc'h, C., Hassim, M., Heming, J., Hudson, D., Ineson, S., Jones, C., Keane, R. J., Kim, K., Kim, J., Kuhlbrodt, T., In Lee, M., Le, C., Martin, G., McCabe, A., Moise, A., Ridley, J., Robert, L., Sahany, S., Schiemann, R. K. H., Storkey, D., Tennant, W., Tomassini, L., Tsushima, Y., Weedon, G. P., West, A., Wheeler, M., Zhou, X., and Zhu, H.: Assessment of the Met Office Global Coupled model version 5 (GC5) configurations, Report. Met Office, Exeter, <https://centaur.reading.ac.uk/112173/>, 2023.
- Zalesak, S. T.: Fully multidimensional flux-corrected transport algorithms for fluids, *Journal of Computational Physics*, 31, [https://doi.org/10.1016/0021-9991\(79\)90051-2](https://doi.org/10.1016/0021-9991(79)90051-2), 1979.
- Zhang, J. and Rothrock, D. A.: Modeling global sea ice with a thickness and enthalpy distribution model in generalized curvilinear coordinates, *Monthly Weather Review*, 131, [https://doi.org/10.1175/1520-0493\(2003\)131<0845:MGSIIWA>2.0.CO;2](https://doi.org/10.1175/1520-0493(2003)131<0845:MGSIIWA>2.0.CO;2), 2003.
- Zhang, R. and Vallis, G. K.: The role of bottom vortex stretching on the path of the North Atlantic western boundary current and on the northern recirculation gyre, *Journal of Physical Oceanography*, 37, <https://doi.org/10.1175/JPO3102.1>, 2007.