



# An intercomparison of four gridded precipitation products over Europe using the three-cornered-hat method

Llorenç Lledó<sup>1</sup>, Thomas Haiden<sup>2</sup>, and Matthieu Chevallier<sup>2</sup>

<sup>1</sup>ECMWF, Bonn, Germany

<sup>2</sup>ECMWF, Reading, UK

**Correspondence:** Llorenç Lledó (llorenç.lledo@ecmwf.int)

**Abstract.** Precipitation is arguably one of the most relevant surface variables impacting human lives on the planet, but global-coverage, high-resolution and good-quality observations are not readily available. In particular, gridded observational datasets are much needed for model development and forecast quality assessment. Here we compare the quality of four types of gridded precipitation products over Europe, namely: a rain-gauge interpolation; a satellite-derived product; a radar composite; and a reanalysis. Each product has its own strengths and weaknesses, and since each precipitation estimate uses different measuring techniques, we can employ a triangulation method to estimate the error variance of each product with respect to the unknown true values. Results show that: a) the satellite product has limited quality over Europe and may be problematic to use in quantitative forecast evaluation and diagnostics; b) the radar composite has spurious features that need to be considered when used in verification; c) all products struggle in topographically complex areas; d) the rain-gauge interpolation is not free of errors, despite rain gauges being often treated as ground truth in the literature; and e) the reanalysis dataset produces in some cases the best available estimates, in particular over the European near-coastal waters.

## 1 Introduction

Surface precipitation affects many aspects of human life on Earth: rain, droughts and floods can threaten human lives and infrastructures, and impact agriculture and energy production, biodiversity, forest fires, air quality, availability of drinking water, health, businesses, and everyday lives. Unsurprisingly, precipitation forecasts are an essential part of weather prediction, and the importance of measuring precipitation globally has been widely recognized (Hou et al., 2014). In particular, gridded precipitation observations are needed to assess socio-economic impacts of precipitation hazards, better understand physical processes, calibrate numerical models, or verify weather forecasts.

While nowadays many observational precipitation products are available (e.g. see Beck et al. (2019) for a list of 26 alternatives), their spatial coverage, horizontal resolution, temporal sampling, and overall quality vary widely. Each product has its own strengths and weaknesses: for instance, some products have very good quality but are only available over land or for certain countries; other datasets have global coverage at the cost of lower resolution; some datasets include rain estimates over the oceans, where in-situ observations are not available; and some products are designed to be available close to real-time, while other products put more emphasis on quality control and homogeneity.



25 In view of this variety of products, it is crucial to assess both their uncertainty and the suitability for specific purposes. This study was guided by the need to verify kilometre-scale global forecasts produced under the framework of the Destination Earth programme (Hoffmann et al., 2023). In the case of forecast verification, the quality of the observational datasets affects the score results and can potentially alter the conclusions reached (Bowler, 2008; Candille and Talagrand, 2008; Duc and Saito, 2018; Bessac and Naveau, 2021; Ramon et al., 2023). In practical terms, high-resolution observations with a wide coverage over  
30 land and oceans should be preferred for the verification of global kilometre-scale forecasts. Precipitation is highly variable both in space and time, and its highly skewed distribution renders time or space aggregates non-representative of the most extreme conditions occurring within a grid box.

In this work we compare the quality of four independently derived precipitation products over Europe (see section 3.1). We employ a triangulation technique to estimate the error variance of each product (described in section 3.3). This method  
35 relies on the assumption that the precipitation estimates are made using independent observation and retrieval techniques. Here, we have extended the method to allow for a certain degree of inter-dependence in the datasets. Therefore, we review the existing measurement techniques in section 2 to understand their strengths and shortcomings. Results and conclusions follow in sections 4 and 5.

## 2 Four alternative ways of estimating precipitation

40 Several measurement techniques are available to estimate surface precipitation (Prigent, 2010; Sun et al., 2018). Rain gauges are the most direct way of observation, and if well maintained and installed they produce good-quality records. However, gauges have problems measuring when there are high winds, or when the precipitation is not in liquid phase, and they produce estimates that are only valid at the local scale. Moreover, their installation and maintenance are costly and therefore there is a limited number of such stations, mostly concentrated around populated areas. Gridded products can be produced from gauge  
45 networks, and their quality will be dictated by the number of stations within each grid cell.

Remote sensing provides an alternative to in-situ measurements of surface precipitation. Weather radars can measure precip-  
itation in the surroundings by sending horizontal beams of electromagnetic waves and measuring the energy reflected by water suspended in the air. Several types of radars with different frequency bands are operating in many countries. Radars produce high-resolution information down to kilometre-scale, but they have known issues: beam attenuation (quality degrades when  
50 moving away from the radar location); beam blocking by topography; the indirect nature of the measurement which introduces errors when transforming measured reflectivity to precipitation rates (Z-R relationship); spurious echoes and clutter due to wind turbines, bird migrations, planes, etcetera; or the inability to measure close to ground level, therefore confusing virga for precipitation or missing low-level precipitation. Weather radars are also a costly infrastructure, and they are only available for populated areas and in countries that can afford it. Gridded products can be made by combining information from multiple  
55 radars.

Satellite-mounted remote sensing devices allow measuring precipitation from space in regions where traditional observations or weather radars are not available, e.g. over the oceans or in remote areas. Many types of satellites and instruments can be



**Table 1.** Main characteristics of the observational datasets employed.

	E-OBS	IMERG	OPERA	ERA5
Product type	Rain gauges	Satellite	Radar	Reanalysis
Spatial resolution	0.1 degrees	0.1 degrees	~2 km	0.25 degrees
Coverage	Europe, only land	Global*	Europe, land and offshore	Global
Time sampling	Daily	30 min.	15 min.	hourly
Period	1950–present	2010–Sep 2021	March 2013–present	1940–present

\* Fewer instruments available above 60°N.

employed. Infrared imagers on geostationary satellites can estimate convective precipitation through brightness temperatures and cloud top height information. Low-Earth-orbiting satellites can also estimate precipitation by measuring passive microwave radiation or by employing active microwave instruments (e.g. on-board weather radars). The main sources of uncertainty for satellite-derived precipitation observations are: the indirect nature of the observations; the sparse or intermittent coverage of polar-orbiting satellites; the inhomogeneity of the techniques employed over land and sea, e.g. by passive microwave imagers; difficulties to detect warm-cloud (low-level) precipitation; and the algorithms needed to combine the available information from multiple instruments and satellites.

Finally, reanalyses combine numerical weather prediction models and data-assimilation techniques to estimate a gap-free and physically consistent state of the atmosphere. Current global reanalyses have coarser resolutions than other state-of-the-art precipitation observation systems, but they offer an additional perspective of surface precipitation. Precipitation estimates from reanalyses are not derived directly from precipitation measurements, but by observations of other atmospheric quantities such as temperature, humidity, or atmospheric circulation, which in turn shape precipitation in the model. Precipitation estimates from reanalyses inherit the weaknesses of the models employed (e.g. they rely on parametrizations), and have difficulties representing small-scale convective precipitation, but can do a good job reproducing larger-scale stratiform precipitation.

### 3 Data and methods

#### 3.1 Datasets

Four gridded datasets covering Europe have been employed. Table 1 summarises the main characteristics of each of them. While many other gridded precipitation datasets are available, for this study we have selected one product for each type of measurement technique to guarantee independence of products as much as possible.

##### 3.1.1 E-OBS

E-OBS (Comes et al., 2018) is a gridded product produced by KNMI and derived from in-situ rain-gauge measurements over Europe. As such, it only provides values over land. The station density within each grid cell is variable both in space and time,



80 impacting the quality of the precipitation estimates. We have employed version 26.0e of this dataset. As an in-situ product, it is sometimes used as a reference for calibrating or verifying other types of measurements, although that is only valid in areas of high station density. Moreover, this product is not free of errors. For instance, the daily accumulation periods of some gauges do not correspond to accumulations at midnight. All our analyses are agnostic and do not assume that one product is an error-free reference used to benchmark others.

### 85 **3.1.2 IMERG**


The Integrated Multi-satellitE Retrievals for Global Precipitation Mission (IMERG), (Huffman et al., 2023, 2020) is a satellite-derived product combining information from passive and active sensors aboard both geostationary and polar-orbiting satellites. Its main strength is that it provides global coverage (including oceans) with a 30-minute time sampling on a 0.1 degree grid. The IMERG algorithm is complex and merges infrared temperatures, passive microwave, and active radar soundings. We employ  
90 the Final Run IMERG version 07B product, which has been quality controlled and scaled so that monthly totals match those of rain gauges, where available.

### **3.1.3 OPERA**

The Operational Program on the Exchange of Weather Radar Information (OPERA), (Huuskonen et al., 2014; Saltikoff et al., 2019) produces a weather radar composite combining information from around 200 radars from 25 European countries. This  
95 radar network is very inhomogeneous in terms of radar technologies employed in different countries (bands, scanning strategy, etc. . . ), but the Opera data center (ODC, also known as Odyssey) produces the composites from individual information of each radar, which minimizes the differences due to post-processing. The hourly total precipitation composite is available on a grid of 2 km resolution every 15 minutes. As a very basic measure of quality control, we have set any values above 300 mm/h as invalid.

### 100 **3.1.4 ERA5**

ERA5 (Hersbach et al., 2023, 2020) is a state-of-the-art global reanalysis produced by ECMWF. By employing a frozen numerical weather prediction model and data assimilation scheme it recreates the state of the atmosphere during the past decades. A reanalysis constrains the solution to be physically consistent and close to existing observations available. Therefore, it produces accurate estimates of precipitation even in regions without direct observations. Indeed, ERA5 does not assimilate  
105 precipitation observations other than the NCEP stage IV precipitation over the US. This means that precipitation estimates from ERA5 in Europe derive mainly from other atmospheric quantities constrained by observations (e.g. humidity, geopotential height, winds. . . ) and their relationship to precipitation in the model. Total precipitation in ERA5 includes both parametrized and large-scale precipitation in forms of rain and snow, and takes into account some of the precipitation evaporation.



**Figure 1.** Data availability for each product, indicated as a percentage of the whole period.

### 3.2 Data collocation

110 In order to perform the comparative assessment, all the data has been collocated in a common space and time grid. First the data has been accumulated to daily total precipitation (00:00–24:00UTC). Then each dataset has been interpolated to a  $0.1^\circ \times 0.1^\circ$  latitude-longitude grid. In particular, the OPERA radar data has been interpolated from a finer grid using a conservative interpolation, and the ERA5 reanalysis has been interpolated from a coarser grid with a nearest-neighbour method. The E-OBS and IMERG products were already on a  $0.1^\circ$  grid.

115 The period employed covers October 2016 to September 2021, amounting to 5 years or 1826 days. The data has been cropped to a European domain covering  $20^\circ\text{W}$  to  $40^\circ\text{E}$  and  $30^\circ\text{N}$  to  $70^\circ\text{N}$ . However, some of the assessed products do not cover the whole domain, and a full comparison is only possible for those grid points where all four products are available. Figure 1 shows a map of data availability (in %) for each product. E-OBS is only available over land. The radar composite covers only a limited number of countries with an additional offshore buffer. ERA5 and IMERG cover all the region and for all times, although IMERG has a reduced satellite coverage above of  $60^\circ\text{N}$ .

The assessment is performed separately at each grid point. To ensure that the statistics (means, variances, covariances) computed for each product are comparable, only the days that are available for all four products are considered. Therefore, the exact periods at each location can slightly differ. ERA5, IMERG and E-OBS are available almost all the time, so the dates used at each grid point are dictated by the availability of OPERA data. Additionally, we have discarded any grid point with  
125 less than 75% of data availability, ensuring sample sizes of at least 1370 days. This is within the range of 500 to 5000 samples recommended by Sjoberg et al. (2021) for the 3CH method.



### 3.3 Three- and four-cornered hat error estimation

The three-cornered-hat (3CH) method was originally developed to estimate random errors of atomic clocks (see Sjoberg et al. (2021) for a comprehensive review of the method). The 3CH and the related triple collocation method have been used in the area of weather and climate to estimate observational errors of wave height (Janssen et al., 2007), sea surface temperature (O'Carroll et al., 2008), precipitation (Roebeling et al., 2012; Massari et al., 2017), surface wind (Abdalla and De Chiara, 2017), or radio occultation measurements (Anthes and Rieckh, 2018).

The method is based on a simple idea: the variance of the difference of two independent observation systems  $A$  and  $B$  is the sum of the error variance of each product:


$$\begin{aligned} \text{var}(A - B) &= \text{var}(A - t - (B - t)) = \text{var}(A - t) + \text{var}(B - t) - 2\text{cov}(A - t, B - t) \\ &\approx \text{var}(A - t) + \text{var}(B - t) \end{aligned} \quad (1)$$

where  $t$  represents the true unknown values. The approximation is only valid if the errors of product  $A$  and  $B$  are uncorrelated, i.e. the observations are derived independently. For example, the second panel in figure 2 shows the variance of differences between the gauge product and the radar composite. If we assume that radar and gauges measure the same unknown quantity  $t$  employing different techniques which should produce uncorrelated errors, we can interpret figure 2 as the sum of the individual error variances of both products. Areas with large values in figure 2 are caused by errors in either product  $A$  or  $B$  or both (notice that variances cannot be negative by definition, so there is no room for error compensation).

At this point we cannot say what proportion of the difference variance is attributable to each product error. To that end, if we have a third independent observation system  $C$ , we can repeat the above steps with differences between  $A$  and  $C$ , and  $B$  and  $C$  to obtain a system of three equations and three unknowns that can be solved to obtain a quantitative result:

$$\begin{cases} v_{A-B} = v_{A-t} + v_{B-t} - 2c_{AtBt} = v_{A-t} + v_{B-t} \\ v_{A-C} = v_{A-t} + v_{C-t} - 2c_{AtCt} = v_{A-t} + v_{C-t} \\ v_{B-C} = v_{B-t} + v_{C-t} - 2c_{BtCt} = v_{B-t} + v_{C-t} \end{cases} \quad (2)$$

In our case we have identified four observational systems, therefore we can constrain the computation a bit more. The four-cornered hat (4CH) method extends the system of equations with three more equations, (six in total, one for each pair of products), for only four unknowns, which results in an overdetermined system. This allows us to choose two covariance terms that we can compute explicitly instead of assuming they are zero.



**Figure 2.** Variance of the differences between pairs of products.

$$\left\{ \begin{array}{l}
 v_{A-B} = v_{A-t} + v_{B-t} - 2c_{AtBt} = v_{A-t} + v_{B-t} \\
 v_{A-C} = v_{A-t} + v_{C-t} - 2c_{AtCt} = v_{A-t} + v_{C-t} \\
 v_{A-D} = v_{A-t} + v_{D-t} - 2c_{AtDt} = v_{A-t} + v_{D-t} \\
 v_{B-C} = v_{B-t} + v_{C-t} - 2c_{BtCt} = v_{B-t} + v_{C-t} \\
 v_{B-D} = v_{B-t} + v_{D-t} - 2c_{BtDt} \\
 v_{C-D} = v_{C-t} + v_{D-t} - 2c_{CtDt}
 \end{array} \right. \quad (3)$$

Assessing which two covariance terms should be computed is a science-informed but subjective matter. The independence of two observational products reduces to identifying confounding variables that could have an impact on the measurements on both measuring techniques. Notice that independently derived products do not directly guarantee uncorrelated errors. For





example, if two measuring approaches are sensitive to rain drop size, or two products have difficulties to identify low-level  
155 precipitation, that will result in correlated errors in the data. Small sample sizes can also result in non-zero covariances due to  
sampling randomness. In our setting, the following assumptions have been made:


- Satellite and gauge errors are considered independent since measurement techniques are very different in nature. Al-  
though the final IMERG product is bias adjusted with monthly total precipitation of some gauge stations, the variability  
of daily total values is only weakly affected by that adjustment.
- 160 – Radar and gauge errors are considered independent since measurement techniques are very different in nature. OPERA  
does not use rain gauges to calibrate the composites.
- ERA5 and gauge errors are considered independent since ERA5 is not assimilating any rain gauge information beyond  
the US.
- ERA5 and radar errors are independent, since ERA5 is not assimilating any radar data over Europe.
- 165 – ERA5 assimilates radiances from many satellites, including microwave imagers, microwave sounders and infrared im-  
agers (see figure 5 in Hersbach et al. (2020)). Some of those instruments (e.g. the GPM microwave imagers) are also  
used by the IMERG algorithm (Huffman et al., 2020). Therefore, we have decided to explicitly compute this covariance  
term.
- Some GPM satellites used in IMERG incorporate a dual-band weather radar on-board. Although the overall characteris-  
170 tics of this on-board radars is very different from conventional ground-based radars, some aspects such as the equations  
employed to convert radar reflectivity to precipitation rates are common to both estimate techniques. Therefore, we have  
decided to explicitly compute this covariance term.

#### 4 Results

As a first step to intercompare the four datasets, the difference in the daily total mean precipitation is presented in figure 3  
175 for each pair of products. The largest differences can be seen over the ocean, with IMERG producing 1 to 2 mm/day more  
than ERA5, and ERA5 giving more precipitation than OPERA by 2 to 5 mm/day. Over land, ERA5 and IMERG have a better  
agreement, with important differences concentrated over high mountain ranges only. Both products are wetter than E-OBS and  
OPERA, although OPERA is the driest product in most locations. While these difference plots allow spotting spatial patterns  
in the data (e.g. beam attenuation in all OPERA panels, or systematic biases over Poland for EOBS) and getting an idea of the  
180 uncertainty range, the true values are unknown, and hence we cannot judge which of the products yields a closer estimate of  
the true mean precipitation.

Under the assumptions made in the previous section, the 4CH method allows estimating the error variances of each product  
and therefore ranking the products from worst to best. Figure 4 shows the 4CH error variances of each product (only available





**Figure 3.** Mean of the differences between pairs of products.

over the grid points covered by all four products). The IMERG product shows the largest errors overall, which tend to be worse in coastal areas of the Iberian Peninsula and the British Isles, and around the Alps. The OPERA composite also has very high error variances in some regions, clearly related to issues in particular radars, e.g. the radars in Madrid or Copenhagen, but in other areas such as Germany the quality is very good. Clutter can also be seen in some areas of eastern Europe. It is reassuring to see that the method can recreate the spatial error patterns produced by the radars without any knowledge of the radar locations. The E-OBS product emerges as a good product in almost all of the continent, although it has some quality issues in areas such as the Balkans. The ERA5 dataset, despite being a lower-resolution product, shows errors close to those of E-OBS, and even better in some regions such as Brittany or the British Isles (see figure A1 for a plot of the ranking of each product at each grid point). ERA5 has less quality in mountainous areas such as the Alps. Some localized spots in figure 4 have negative variance values. Those can be mainly seen for E-OBS over the Iberian Peninsula and in the north-east of Italy, with a mean value close to  $-1.6 \text{ mm}^2$  over those locations. Those values are unphysical, since error variances must be positive by definition, and result from the assumptions made in section 3 not being entirely correct in those grid points. Indeed, the error covariance between IMERG and ERA5 have unrealistic strong negative values in those regions (see figure A2). While some




authors have proposed ways forward to constrain the 3CH solutions to remain positive-definite (e.g. see Premoli and Tavella (1993)), we have not explored this avenue, because the issue is minor and restricted to few locations in our case.

200 Since convective and stratiform precipitation have very different characteristics in terms of spatial patterns, intensity, duration, or height in the atmosphere, it is reasonable to ask if the four products studied here represent well these two distinct types of precipitation. To answer that, we have stratified the results into two seasons, an extended winter (NDJFM, predominantly stratiform precipitation) and extended summer (MJJAS, mostly convective precipitation). The results in figure 5 show that E-OBS, OPERA and ERA5 have better accuracy in winter than in summer, whereas IMERG has better performance in summer than in winter. This confirms that IMERG is better at sensing convective precipitation events than stratiform precipitation, whereas the contrary can be said for the other products. Still, IMERG is not better than the other three products in  
205 itation, whereas the contrary can be said for the other products. Still, IMERG is not better than the other three products in summer. An interesting aspect to notice in figure 5 is that the quality issues we saw in figure 4 for OPERA almost disappeared in the winter/summer stratifications. A plausible explanation is that some outliers that are only present in some days of April or October (therefore excluded in both composites) had a strong influence on the overall result (see section 4 of Sjoberg et al. (2021)).

## 210 5 Conclusions


The choice of a verification dataset for global and high-resolution precipitation forecasts is challenging. From a technical perspective IMERG is a good choice because it covers the oceans, and it produces a relatively high resolution and mostly gap-free dataset. However, we have seen that its quality over Europe is not particularly good compared to existing alternatives. Moreover, its quality over the ocean is also not good (see figure A3). Both E-OBS and the OPERA radar composites provide  
215 a better alternative over European land. The OPERA data is a higher resolution product, and although it might be biased with respect to in-situ measurements, with a bit of additional quality control it might be the best way to go for verifying high-resolution simulations over Europe. Both products have the limitation of not providing information over the oceans, which might be a problem to obtain robust results (given that oceans cover two thirds of our planet) or for applying spatial verification techniques that rely on neighbourhood analyses. The only remaining alternative might then be using modelled products such as  
220 ERA5. Our results show that the quality of ERA5 is not far from that of E-OBS in Europe, especially in winter (in agreement with Lavers et al. (2022)). But since ERA5 is produced with a specific version of the IFS model, comparing IFS forecasts to ERA5 does not provide a very stringent evaluation, as forecast and analysis share very similar model limitations and systematic biases. Higher-resolution modelled datasets such as regional reanalyses could provide better results than ERA5, but we have not explored this option here. In short, the promise of a global-coverage, high-resolution and good-quality precipitation dataset  
225 measured from space is not yet fully realized, and therefore verifying global precipitation forecasts against reanalysis datasets is still a justified alternative.

Extending this study to other regions would be interesting. Specifically, the IMERG dataset is thought to be of better quality in the tropics due to the predominance of convective precipitation, and it has already been used there to assess the quality of convective-permitting simulations (Becker et al., 2021). Unfortunately, an analysis such as the one we presented is only possible



**Figure 4.** 4CH-derived error variances for the four products.

230 in areas with a good radar and gauge network. The US would be an ideal place to apply the 4CH method, but unfortunately there the ERA5 assimilates the stage IV (radar and gauges) product, making our assumptions invalid.




**Figure 5.** 4CH error variances in winter (top row) and summer (bottom row) for each observational product.

This study has only dealt with error variances, and therefore does not directly address quality aspects of the precipitation products that might be relevant for extreme events, such as performance under heavy rain (see e.g. Sun et al. (2018)). Additional analyses showing other quality aspects of the IMERG dataset in Europe can be found in Navarro et al. (2019) or O et al. (2017),  
 235 whereas the quality of IMERG over the US has also been studied in Beck et al. (2019). These studies take one reference dataset as the true values and evaluate other products. However, our agnostic analysis has confirmed that gauge-based products are not free of errors either.

*Code and data availability.* All the analyses were done with Python and Xarray (Hoyer and Hamman, 2017). A Jupyter Notebook containing all the analyses can be found at [https://github.com/lluritu/4CH\\_precip\\_comparison](https://github.com/lluritu/4CH_precip_comparison). E-OBS version 26.0e can be obtained from the Copernicus Climate Change Service at [https://surfobs.climate.copernicus.eu/dataaccess/access\\_eobs.php](https://surfobs.climate.copernicus.eu/dataaccess/access_eobs.php). The GPM IMERG v07B can be accessed from NASA GES DISC at <https://doi.org/10.5067/GPM/IMERG/3B-HH/07>. OPERA data is available from EUMETNET ([www.eumetnet.eu/opera](http://www.eumetnet.eu/opera)). ERA5 was downloaded from the Copernicus Climate Change Service (C3S) (2023) at <https://doi.org/10.24381/cds.adbb2d47>.  
 240







**Figure A1.** Ranking of the four products according to 4CH error variance.

## Appendix A

This appendix contains three additional figures. Figure A1 shows a ranking of the four products according to the 4CH analysis, from best to worst at each grid point, which helps realizing which datasets are better overall. Figure A2 displays the covariance terms that were computed as part of the 4CH analysis described in section 3.3. Finally figure A3 shows an additional 3CH analysis including only IMERG, OPERA and ERA5. In this case the errors are assumed uncorrelated. We can see that IMERG and OPERA datasets struggle to represent precipitation over the oceans, whereas ERA5 has negative values there. This indicates that IMERG and OPERA errors are indeed correlated over the ocean. The results over land are comparable to those in figure 4.



**Figure A2.** Error covariances of the 4CH method.



**Figure A3.** Error variance of the three products that cover the near-coastal European waters, under the assumption of uncorrelated errors.

250 *Author contributions.* Llorenç Lledó: conceptualization, methodology, data curation, software, visualization, writing - original draft preparation. Thomas Haiden: conceptualization, methodology, writing - review and editing. Matthieu Chevallier: writing - review and editing.

*Competing interests.* The authors declare that they have no conflict of interest.



*Acknowledgements.* The work presented in this paper has been produced in the context of the European Union's Destination Earth Initiative and relates to tasks entrusted by the European Union to the European Centre for Medium-Range Weather Forecasts implementing part of  
255 this Initiative with funding by the European Union. Views and opinions expressed are those of the author(s) only and do not necessarily reflect those of the European Union or the European Commission. Neither the European Union nor the European Commission can be held responsible for them. The authors want to thank Philippe Lopez and Tobias Becker for assistance with the pre-processing of OPERA and IMERG datasets respectively, and Estíbaliz Gascón for assessing the quality of IMERG for specific case studies. We acknowledge the E-OBS dataset from the EU-FP6 project UERRA (<https://www.uerra.eu>) and the Copernicus Climate Change Service, and the data providers in the  
260 ECA&D project (<https://www.ecad.eu>). The GPM IMERG data were provided by the NASA Goddard Space Flight Center's Precipitation Measurement Missions Science Team and Precipitation Processing System, which develop and compute the GPM IMERG as a contribution to GPM, including data archiving at the NASA GES DISC. ERA5 was downloaded from the Copernicus Climate Change Service (C3S) (2023). The results contain modified Copernicus Climate Change Service information. Neither the European Commission nor ECMWF is responsible for any use that may be made of the Copernicus information or data it contains.





## 265 References

- Abdalla, S. and De Chiara, G.: Estimating Random Errors of Scatterometer, Altimeter, and Model Wind Speed Data, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 10, 2406–2414, <https://doi.org/10.1109/JSTARS.2017.2659220>, 2017.
- Anthes, R. and Rieckh, T.: Estimating observation and model error variances using multiple data sets, *Atmospheric Measurement Techniques*, 11, 4239–4260, <https://doi.org/10.5194/amt-11-4239-2018>, 2018.
- 270 Beck, H. E., Pan, M., Roy, T., Weedon, G. P., Pappenberger, F., van Dijk, A. I. J. M., Huffman, G. J., Adler, R. F., and Wood, E. F.: Daily evaluation of 26 precipitation datasets using Stage-IV gauge-radar data for the CONUS, *Hydrology and Earth System Sciences*, 23, 207–224, <https://doi.org/10.5194/hess-23-207-2019>, 2019.
- Becker, T., Bechtold, P., and Sandu, I.: Characteristics of convective precipitation over tropical Africa in storm-resolving global simulations, *Quarterly Journal of the Royal Meteorological Society*, 147, 4388–4407, <https://doi.org/10.1002/qj.4185>, 2021.
- 275 Bessac, J. and Naveau, P.: Forecast score distributions with imperfect observations, *Advances in Statistical Climatology, Meteorology and Oceanography*, 7, 53–71, <https://doi.org/10.5194/ascmo-7-53-2021>, 2021.
- Bowler, N. E.: Accounting for the effect of observation errors on verification of MOGREPS, *Meteorological Applications*, 15, 199–205, <https://doi.org/10.1002/met.64>, 2008.
- Candille, G. and Talagrand, O.: Impact of observational error on the validation of ensemble prediction systems, *Quarterly Journal of the Royal Meteorological Society*, 134, 959–971, <https://doi.org/10.1002/qj.268>, 2008.
- 280 Cornes, R. C., van der Schrier, G., van den Besselaar, E. J. M., and Jones, P. D.: An Ensemble Version of the E-OBS Temperature and Precipitation Data Sets, *Journal of Geophysical Research: Atmospheres*, 123, 9391–9409, <https://doi.org/10.1029/2017JD028200>, 2018.
- Duc, L. and Saito, K.: Verification in the presence of observation errors: Bayesian point of view, *Quarterly Journal of the Royal Meteorological Society*, 144, 1063–1090, <https://doi.org/10.1002/qj.3275>, 2018.
- 285 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>,  
290 2020.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1940 to present, <https://doi.org/10.24381/cds.adbb2d47>, 2023.
- Hoffmann, J., Bauer, P., Sandu, I., Wedi, N., Geenen, T., and Thiemert, D.: Destination Earth – A digital twin in support of climate services, *Climate Services*, 30, 100 394, <https://doi.org/10.1016/J.CLISER.2023.100394>, 2023.
- 295 Hou, A. Y., Kakar, R. K., Neeck, S., Azarbarzin, A. A., Kummerow, C. D., Kojima, M., Oki, R., Nakamura, K., and Iguchi, T.: The Global Precipitation Measurement Mission, *Bulletin of the American Meteorological Society*, 95, 701–722, <https://doi.org/10.1175/BAMS-D-13-00164.1>, 2014.
- Hoyer, S. and Hamman, J.: xarray: N-D labeled arrays and datasets in Python, *Journal of Open Research Software*, 5,  
300 <https://doi.org/10.5334/jors.148>, 2017.



- Huffman, G., Stocker, E., Bolvin, D., Nelkin, E., and Tan, J.: GPM IMERG Final Precipitation L3 1 day 0.1 degree x 0.1 degree V07, <https://doi.org/10.5067/GPM/IMERGDF/DAY/07>, 2023.
- Huffman, G. J., Bolvin, D. T., Braithwaite, D., Hsu, K., Joyce, R., Kidd, C., Nelkin, E. J., Sorooshian, S., Tan, J., and Xie, P.: NASA Global Precipitation Measurement (GPM) Integrated Multi-satellite Retrievals for GPM (IMERG) Algorithm Theoretical Basis Document Version 06, Tech. rep., NASA, [https://gpm.nasa.gov/sites/default/files/2020-05/IMERG\\_ATBD\\_V06.3.pdf](https://gpm.nasa.gov/sites/default/files/2020-05/IMERG_ATBD_V06.3.pdf), 2020.
- 305 Huuskonen, A., Saltikoff, E., and Holleman, I.: The Operational Weather Radar Network in Europe, *Bulletin of the American Meteorological Society*, 95, 897–907, <https://doi.org/10.1175/BAMS-D-12-00216.1>, 2014.
- Janssen, P. A., Abdalla, S., Hersbach, H., and Bidlot, J. R.: Error Estimation of Buoy, Satellite, and Model Wave Height Data, *Journal of Atmospheric and Oceanic Technology*, 24, 1665–1677, <https://doi.org/10.1175/JTECH2069.1>, 2007.
- 310 Lavers, D. A., Simmons, A., Vamborg, F., and Rodwell, M. J.: An evaluation of ERA5 precipitation for climate monitoring, *Quarterly Journal of the Royal Meteorological Society*, 148, 3152–3165, <https://doi.org/10.1002/qj.4351>, 2022.
- Massari, C., Crow, W., and Brocca, L.: An assessment of the performance of global rainfall estimates without ground-based observations, *Hydrology and Earth System Sciences*, 21, 4347–4361, <https://doi.org/10.5194/hess-21-4347-2017>, 2017.
- Navarro, A., García-Ortega, E., Merino, A., Sánchez, J. L., Kummerow, C., and Tapiador, F. J.: Assessment of IMERG Precipitation Estimates over Europe, *Remote Sensing* 2019, Vol. 11, Page 2470, 11, 2470, <https://doi.org/10.3390/RS11212470>, 2019.
- 315 O, S., Foelsche, U., Kirchengast, G., Fuchsberger, J., Tan, J., and Petersen, W. A.: Evaluation of GPM IMERG Early, Late, and Final rainfall estimates using WegenerNet gauge data in southeastern Austria, *Hydrology and Earth System Sciences*, 21, 6559–6572, <https://doi.org/10.5194/hess-21-6559-2017>, 2017.
- O’Carroll, A. G., Eyre, J. R., and Saunders, R. W.: Three-Way Error Analysis between AATSR, AMSR-E, and In Situ Sea Surface Temperature Observations, *Journal of Atmospheric and Oceanic Technology*, 25, 1197–1207, <https://doi.org/10.1175/2007JTECHO542.1>, 2008.
- 320 Premoli, A. and Tavella, P.: A revisited three-cornered hat method for estimating frequency standard instability, *IEEE Transactions on Instrumentation and Measurement*, 42, 7–13, <https://doi.org/10.1109/19.206671>, 1993.
- Prigent, C.: Precipitation retrieval from space: An overview, *Comptes Rendus Geoscience*, 342, 380–389, <https://doi.org/10.1016/j.crte.2010.01.004>, 2010.
- 325 Ramon, J., Lledó, L., Ferro, C. A. T., and Doblas-Reyes, F. J.: Uncertainties in the observational reference: implications in skill assessment and model ranking of seasonal predictions, *Quarterly Journal of the Royal Meteorological Society*, <https://doi.org/10.1002/qj.4628>, 2023.
- Roebeling, R. A., Wolters, E. L. A., Meirink, J. F., and Leijnse, H.: Triple Collocation of Summer Precipitation Retrievals from SEVIRI over Europe with Gridded Rain Gauge and Weather Radar Data, *Journal of Hydrometeorology*, 13, 1552–1566, <https://doi.org/10.1175/JHM-D-11-089.1>, 2012.
- 330 Saltikoff, E., Haase, G., Delobbe, L., Gaussiat, N., Martet, M., Idziorek, D., Leijnse, H., Novák, P., Lukach, M., and Stephan, K.: OPERA the Radar Project, *Atmosphere*, 10, 320, <https://doi.org/10.3390/atmos10060320>, 2019.
- Sjoberg, J. P., Anthes, R. A., and Rieckh, T.: The Three-Cornered Hat Method for Estimating Error Variances of Three or More Atmospheric Datasets. Part I: Overview and Evaluation, *Journal of Atmospheric and Oceanic Technology*, 38, 555–572, <https://doi.org/10.1175/JTECH-D-19-0217.1>, 2021.
- 335 Sun, Q., Miao, C., Duan, Q., Ashouri, H., Sorooshian, S., and Hsu, K. L.: A Review of Global Precipitation Data Sets: Data Sources, Estimation, and Intercomparisons, *Reviews of Geophysics*, 56, 79–107, <https://doi.org/10.1002/2017RG000574>, 2018.