

# Influence of river runoffs and precipitation on the seasonal and interannual variability of Sea Surface Salinity in East Northern Tropical Atlantic

Clovis Thouvenin-Masson<sup>1</sup>, Jacqueline Boutin<sup>1</sup>, Vincent Échevin<sup>1</sup>, Alban Lazar<sup>1</sup>, Jean-Luc Vergely<sup>2</sup>

5 <sup>1</sup> Sorbonne Université, LOCEAN-IPSL, CNRS/IRD/MNHN, Paris, France

<sup>2</sup>ACRI-st, Guillaucourt, France

*Correspondence to:* Clovis Thouvenin-Masson (clovis.thouvenin-masson@locean.ipsl.fr) and Jacqueline Boutin (jb@locean.ipsl.fr)

## Abstract:

10 In tropical regions, the fresh water flux entering the ocean originates primarily from precipitation and, to a lesser extent when considering basin scale averages, from continental rivers. Nevertheless, at regional scale, river flows can have a significant impact on the surface ocean dynamics. Riverine fresh water modifies salinity, and therefore density, stratification and circulation. With its particular coastline, high cumulative river discharge, and the vicinity of Inter Tropical Convergence Zone (ITCZ), the eastern part of North Tropical Atlantic (e-NTA) region off Northwest Africa is a particularly interesting location  
15 to study the linkage between precipitation, river outflow and Sea Surface Salinity (SSS). Here we focus on the regional e-NTA SSS seasonal cycle and interannual variability. We quantify the impact of river runoff and precipitation on SSS by means of regional simulations forced by different interannual and climatological river runoffs and precipitation products. The simulated SSS are compared with the Climate Change Initiative (CCI) satellite SSS, *in situ* SSS from Argo, ships and a coastal mooring, and the GLORYS reanalysis SSS. The analysis of the salinity balance in the mixed layer is conducted to explore the dynamics  
20 influencing the SSS variability. Overall, the simulations reproduce well the seasonal cycle and interannual variability despite a positive mean model bias north of 15N. The seasonal cycle is impacted by the phasing of the different runoff products. The mixed layer SSS decrease during the rainy season is mainly driven by precipitation followed by runoff by means of horizontal advection and partly compensated by vertical mixing. In terms of interannual anomalies, river runoffs have a more direct impact on SSS than precipitation. This study highlights the importance of properly constraining river runoffs and precipitation  
25 to simulate realistic SSS, and the importance of observing SSS in coastal regions to validate such constraints.

## 1 Introduction

The upper layer of the ocean is where exchanges between the ocean and the atmosphere take place. Air sea forcing (e.g., wind, heat flux) generates turbulence in the surface layer, leading to the formation of a surface mixed layer from a few meters to hundreds of meters thick, with homogeneous characteristics (e.g., temperature and salinity), and whose bottom is characterized  
30 by a marked density gradient, the pycnocline. This layer receives various freshwater flows, such as precipitation, or river discharge. These low salinity water inputs make the density of the surface waters to decrease, which can lead to an increase of

the density gradient between the surface and subsurface waters. Freshwater inputs can also generate significant salinity gradients within the mixed layer (Mignot et al. 2007), leading to the formation of intermediate layers known as barrier layers. The latter isolates the surface layer from the deep ocean, inhibiting heat exchange between the ocean surface and subsurface (Vialard and Delecluse 1998). Such ocean-atmosphere interactions might impact the formation of water masses and their evolution, as well as air-sea exchanges of heat and gases. Since the 1980s, the quality and availability of *in situ* river discharge measurements have declined due to a lack of funding and to an unwillingness by states to share these data with the general public (Chandanpurkar et al. 2017, Durand et al. 2019). As a result, current Ocean General Circulation Model (OGCMs) such as used to generate the GLORYS reanalysis (Lellouche et al. 2021) typically utilize climatological river discharge products (Dai et al., 2009), which have not been updated for more than a decade. However, it has been shown that river discharges tend to vary strongly interannually. Gévaudan et al. (2022) found that the Amazon River runoff anomalies can reach values of the order of  $50\,000\text{ m}^3\text{s}^{-1}$  (25 % of the climatological value) and that these anomalies have a significant influence on the surface salinity of the tropical Atlantic Ocean. Chandanpurkar et al. (2022) studied the influence of river discharge interannual variability on salinity at the mouths of the world's major rivers. They found that river discharge interannual variability is responsible for a standard deviation of 1.3 to 3 pss of salinity, and that models that take the interannual variation of river discharge into account simulate SSS that are closer to satellite observations. At the scale of the global ocean, a recent study (Fournier et al. 2023) demonstrates that SSS variability, averaged in estuarine regions of major river plumes, is strongly correlated with the global water cycle variability, particularly in relation to the ENSO phenomenon.


In this paper, we focus on the eastern North Tropical Atlantic region (e-NTA,  $10^\circ\text{N}$ - $17^\circ\text{N}$ / $20^\circ\text{W}$ - $12^\circ\text{W}$ ; Figure 1), a region subject to high river discharge forcings, resulting from rainfall over the high mountain plateaus of Guinea. There is a geographical disparity in the river flows: to the north of Dakar ( $14.7^\circ\text{N}$ ), only the Senegal River has an average outflow of over  $500\text{ m}^3\text{s}^{-1}$  (Roudier et al. 2014), whereas to the south of Dakar, freshwater discharge takes place through numerous rivers along the coast, with the Gambia River being the most significant. River flow in e-NTA is highly seasonal with rivers that run almost dry during the boreal summer and peak in autumn after the rainy season. While their interannual variations are not well known due to a lack of data, they are expected to be strongly influenced by the West African monsoon with large interannual variations that are expected to grow of 10 – 28% with climate change (Akinsanola, 2020). Moreover, studies based on climate models (Ardoin-Bardin et al. 2009) predict a long-term decreasing trend for these river flows by up to 27 % for Senegal river and up to 37 % for Gambia river in 2080.

Cumulating all the river discharges of Senegal and Guinea [ $12^\circ\text{N}$ - $17^\circ\text{N}$ ] leads to an average monthly outflow of  $\sim 30\,000\text{ m}^3\text{s}^{-1}$  at its annual maximum in September. In comparison, the largest Amazon outflow in May is  $276\,000\text{ m}^3\text{s}^{-1}$ , and the largest outflow of the Congo River is  $56\,000\text{ m}^3\text{s}^{-1}$  in December (Wohl and Lininger 2022). However, e-NTA region is of particular interest because it is subject to both river discharge and intense precipitation linked to the meridional displacements of the InterTropical Convergence Zone (ITCZ). Furthermore, this region hosts the thriving Senegalese coastal upwelling, a region where human populations are highly dependent on small pelagic fisheries as a source of protein (Failler et al. 2014).

65 Concerning the impact of freshwater fluxes on salinity in e-NTA region, only the seasonal variation and their driving physical processes have been studied by Camara et al. (2015) using the Nucleus for European Modelling of the Ocean (NEMO) ocean model. They found that runoffs and precipitations were the main contributors of the freshening in the e-NTA, and that poleward advection of low salinity waters along the coasts was partly compensated by vertical diffusion of salinity.

The aforementioned studies demonstrate the usefulness of salinity as a tracer for variations in the water cycle, both from the perspective of the seasonal cycle and interannual variability. However, to our knowledge, no study has yet focused on interannual variability. This is the aim of this study, in which we aim to differentiate the effects of precipitation from those of river discharges on coastal salinity in the e-NTA region. To achieve this, the surface ocean dynamics is simulated by the Coastal and Regional Ocean Community (CROCO) model with various configurations of climatological or interannual forcings. The model results are compared with Mercator's GLORYS reanalysis, satellite and *in situ* SSS measurements (*e.g.*, merchant ships, Argo floats, buoys). We estimate the seasonal cycle and interannual variation in salinity for each configuration, intercompare these different configurations, and identify the physical drivers of SSS using a mixed-layer salinity balance, following a methodology similar to Camara et al. (2015).

Sect. 2 presents the data, and the methods used. Sect. 3 is dedicated to the results and includes a validation of the modeled SSS, an analysis of the observed and modeled SSS anomalies and a study of modeled SSS sensitivity to changes in freshwater flux forcings. Finally, a few points are raised for discussion and conclusions in sect. 4.



**Figure 1: CCI satellite SSS (color) averaged over October–November–December (over ocean); white contours indicate the averaged ERA5 E-P (evaporation minus precipitation) rate (in mm/d) over this period. Topography (color scale) is shown on land. Along the coast, colored squares indicate averaged ISBA river runoffs over September–October–November (color scale). The area delimited by the red line corresponds to the merged catchment areas of the rivers flowing into the area of study, extracted from the HydroSHEDS database (Hydrological data and maps based on Shuttle Elevation Derivatives). Black dotted boxes delimit the e-NTA and south of e-NTA regions. The yellow star represents the position of Melax buoy.**

## 2 Data & methods

85 The region we focus on is identical to the one studied in Camara et al. (2015), for the purpose of comparing the results obtained. We refer to it as the Eastern North Tropical Atlantic (e-NTA). This region is strongly impacted by river water outflows, as it includes major rivers (Senegal, Gambia, Casamance, big and Little Scarcies; see Figure 1). In the following, we study the salinity and fresh water forcings variables averaged over this region. We focus on the longest common period for which all salinity and forcing products are available, from January 1, 2010, before which satellite salinity products are not available, to

95 June 19, 2019, after which the ISBA-CTRIP product is no longer available.



## 2.1 Salinity data

### - Satellite maps

Three L-band radiometric satellite missions have measured SSS from space: SMOS (2010-present), SMAP (2015-present), and Aquarius (2012-2015). The version 3.2 of the SSS product generated as part of the Climate Change Initiative project (CCI) is used here, covering a period from 2010 to 2021 (Boutin et al. 2021a). These data are generated with a temporal optimal interpolation of the three satellites measurements, as described for version 2 in Boutin et al. (2021b). Developments between version 2 and version 3 are described in detail in Thouvenin-Masson et al. (2022). SSS fields are available on a 25 km Equal-Area Scalable Earth (EASE2) and they are used here at a weekly temporal resolution. Due to the spatial resolution of satellite SSS measurements, data taken at less than ~40 km from the coast are flagged as they must be considered with caution due to land contamination (e.g. (Zine et al. 2008)). This flag filtering is applied in the present study. In this satellite product, a correction is applied to remove the instantaneous effect of rain on the top surface satellite measurements to remain consistent with bulk salinity recorded by most *in situ* instruments (Supply et al. 2020).

### - GLORYS reanalysis

The GLORYS12V1 product is a Copernicus Marine Environment Monitoring Service (CMEMS) global ocean eddy-resolving reanalysis available at daily resolution from 1993 to 2023. These reanalyses are based on the NEMO ocean model forced by ERA5 data and by climatological river runoffs at the surface. Satellite sea level anomalies, Sea Surface Temperature (SST), sea ice concentration, *in situ* temperature, and salinity vertical profiles (but not satellite SSS) are assimilated using a reduced-order Kalman filter derived from a singular evolutive extended Kalman (SEEK) filter with a three-dimensional multivariate background error covariance matrix and a 7 day assimilation cycle (Lellouche et al. 2018, Lellouche et al. 2021). Model reanalysis output is available at daily temporal resolution on a regular 1/12° grid for 50 vertical levels. See Lellouche et al. (2021) for a complete description of the model.

### - *In situ* data

*In situ* data are used to evaluate the CROCO simulations. This involves measurements from thermosalinographs (TSG) installed on merchant and research vessels, from Argo floats near the surface, and from the Melax mooring.

*TSG*: The delayed mode data set from Laboratoire d'Etudes en Géophysique et Océanographie Spatiales (TSG-LEGOS-DM) is used. It is derived from voluntarily observing ships, collected, validated, archived and made freely available by the French Sea Surface Salinity Observation Service (Alory et al. 2015). Adjusted values, when available, and TSG data with quality flags

125 = 1 and 2 ('good' or 'probably good') only are selected. TSG data are available from 1993 to present, between 5 to 15 m depth, and we use the hourly resolution product.

*Argo profilers*: The Argo project is a set of about 4000 profilers moving in the global ocean. These instruments provide around 100 000 temperature and salinity measurements annually over the global ocean, and with an average spacing of 3 degrees  
130 between measurements (Argo 2023). These data are collected and made freely available by the international Argo project and the national programs that contribute to it. The Argo SSS data gathered in the Salinity-Pilot Exploitation Platform (Pi-Mep) database (Guimbard et al. 2021) is used. In this database Argo data from the Global Data Assembly Centre (GDAC) database (Argo 2023) with a quality index of 1 or 2 are selected. Argo measurements between 10 m and 0 m depth are considered as surface data (most of the Argo data resulting from this selection are taken at a depth of about 5 m).

135

*Melax*: The Melax mooring is equipped with oceanographic and atmospheric sensors. Moored at 36 m depth, it is located at 30 km from the coast. It measures the physical and biogeochemical parameters over the Senegalese shelf [ 14°20' N,17°14' W], south of the city of Dakar (Tall et al. 2021, Figure 1). The mooring captured surface salinity almost continuously from mid-February 2015 to August 2016. We use the Melax measurement averaged daily over this time period.

## 140 **2.2 Regional simulations: the CROCO model**

The ocean model CROCO (<https://www.CROCO-ocean.org/>;(Hilt et al. 2020) is used to simulate salinity variations in the e-NTA region. CROCO has vertical sigma coordinates, which are well suited for coastal studies. The slow mode and the fast barotropic mode are computed separately using a time-splitting algorithm (Shchepetkin and McWilliams 2009), improving the  
145 consistency, accuracy, and stability of the simulations. High-order numerical schemes enable the representation of small-scale structures such as mesoscale eddies and filaments. The AGRIF (adaptive refinement of the horizontal grid ; (Debreu et al. 2008) module is utilized, enabling the embedment of a sub-domain in which small-scales are more finely resolved. In the configuration used here, the parent grid covering [7°N-35°N; 30°W-10°W] has a resolution of 10 km, and the child grid used in the Senegal region [12°N-18°N; 20°W-15°W] has a resolution of 2 km. More details on the model configuration can be  
150 found in Ndoye et al. (2018). Daily outputs from the MERCATOR model output at 1/12° resolution (GLOBAL-ANALYSISFORECAST-PHY-001-024; downloaded from <http://marine.copernicus.eu/>) are used to force physical properties (temperature, salinity, velocity and sea level) at the open boundary conditions (OBC) of the parent grid.

Hourly atmospheric forcings (air temperature, relative humidity, 10 m wind, radiative fluxes) from the ERA5 reanalysis (see  
155 below) are used in all simulations. No surface salinity restoring to climatological observations (*e.g.*, Ndoye et al. (2018)) is used.

160 In order to estimate the relative importance of interannual variations in each of the freshwater fluxes forcings, five CROCO simulations are performed with different rain rate and river runoffs forcings. The other hourly forcing terms (air temperature, wind, radiative flux, etc.) are kept identical for all simulations. Three simulations are forced with synoptic freshwater flux forcings, including interannual variations. These simulations are called CROCOglofas, CROCOisba and CROCOimerg. Two simulations are forced with climatological rain rates or climatological river outflows respectively. They are named CROCOprclm and CROCOroclm. The forcings of each simulation are summarized in Table 1.

165

**Table 1: list of simulations and their freshwater flux forcings.**

Name	Precipitation	River discharge
<b>Interannual simulations</b>		
CROCOglofas	ERA5 hourly	GloFAS daily
CROCOisba	ERA5 hourly	ISBA daily
CROCOimerg	IMERG hourly	ISBA daily
<b>Climatological simulations</b>		
CROCOroclm	ERA5 hourly	GloFAS climatology
CROCOprclm	IMERG climatology	ISBA daily

170 **Precipitation and runoffs forcings**

*ERA-5* is a reanalysis produced by the European Centre for Medium-Range Weather Forecasts (ECMWF), which provides comprehensive modeling of atmospheric, continental surface and ocean wave variables (Hersbach et al. 2020). Based on the Cycle 41r2 Integrated Forecast System (IFS), ERA5 hourly fields are available over the period 1950-2023 at a horizontal resolution of 31 km.

175 The precipitation value used in the CROCO simulations is composed of the convective precipitation field (cp) produced by the IFS convection scheme, which represents precipitation at sub-grid scales, and the stratiform precipitation field (sp) produced by the IFS cloud model, which represents the formation and dissipation of clouds and large-scale precipitation due to changes in atmospheric variables such as pressure, temperature and humidity.

180 *Integrated Multi-satellitE Retrievals for Global Precipitation Measurement (IMERG)* is a rain rate product based on satellite precipitation measurements. It combines information from the Global Precipitation Measurement (GPM) satellite constellation with infrared (IR) satellite data taken by geostationary satellites to estimate precipitation over the majority of the Earth's surface

at a frequency of 30 minutes. The algorithm is based on the Climate Prediction Center Morphing (CMORPH; (Joyce et al. 2004) method, and takes advantage of the high repetition rate of IR satellites to track the movement of less frequent but more accurate microwave and radar-detected rainfall cells. IMERG data are available from 2000 to present, at a resolution of 0.1° every half-hour.

Over the ocean, these two precipitation products are consistent in terms of mean values and have similar climatologies and anomalies (Figure 5c, Figure 2b), after integration over the e-NTA. Nevertheless, IMERG rain rates are more variable locally and extend over a larger range of values than ERA5 rain rates (see Figures S.1 and S.3, blue and red curves).

190

*The Global Flood Awareness System* (GloFAS; <http://www.globalfloods.eu/>; (Harrigan et al. 2020) is one of the components of the Copernicus Emergency Management Service (CEMS). This system is designed to help prevent flooding on a global scale, notably by providing water level forecasts for river basins. It is based on satellite data, on soil temperature and humidity, on precipitation from ERA5, and on *in situ* data. These data are integrated into the Hydrology in the Tiled ECMWF Scheme for Surface Exchanges over Land (HTESSEL) continental surface model, which is part of the ECMWF's integrated forecasting system (IFS 41r2) via a terrestrial data assimilation system explained in de Rosnay et al. (2012). The resulting runoff is then integrated into the LISFLOOD runoff routing model. The GloFAS hydrological model simulations are available from 1979 to present at a daily and 0.1° resolution.

200

*ISBA-CTRIP* river discharge estimation is used in this study. ISBA-CTRIP combines two models: the Interaction Soil-Biosphere-Atmosphere (ISBA; <https://www.umr-cnrm.fr/isbadoc/model.html>) hydrological model developed by the Centre National de Recherches Météorologiques (CNRM) within the framework of the IPCC (see Decharme et al. (2019) for a full description of the model) and the CTRIP (CNRM version of Total Runoffs Integrating Pathway) model, which is an improved version of the TRIP model used to simulate river runoff to the ocean from the total runoff calculated by ISBA. In the configuration used here this model uses Tier-2 Water Resources Re-analysis precipitation at 0.25° resolution (WRR2) from the E2O project as forcing. The E2O dataset is directly based on the 3-hourly ERA-Interim reanalysis (<https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era-interim>) over the 1979-2014 period. Precipitations have been hybridized with observations using the Multi-Source Weighted-Ensemble Precipitation (MSWEP; <http://www.gloh2o.org>) data set (Beck et al. 2017). ISBA-CTRIP data is available daily from 1979 to June 19, 2019, at a 0.5° resolution.

210

GloFAS and ISBA runoffs, after summing the individual outflows for the region studied, have similar climatologies (maximum difference of  $1.10^8 \text{ m}^3/\text{d}$ , see Figure 2b). The simulated river runoffs exhibit strong interannual anomalies in this area (Figure 5c). These river runoff anomalies are highly correlated with monsoon anomalies (Figure 8) and can reach  $11\,000 \text{ m}^3\text{s}^{-1}$ , i.e., almost 30 % of the seasonal variation (Figure 2b). These interannual anomalies (see sect. 3.3 below) are sometimes of opposite

215

signs between the two products, with differences reaching  $12.10^8 \text{ m}^3/\text{d}$  (Figure 5c). These differences and their origins are discussed in section 4.

### Processes controlling the mixed-layer salinity budget

220

Diagnostics implemented in CROCO make it possible to isolate the various terms involved in the salinity balance of the mixed layer to identify the dynamical processes that modify salinity. In CROCO, the temporal variation of the salinity,  $\partial_t S$ , is expressed as follows, for each layer of the water column:

$$\partial_t S = -\partial_x(uS) - \partial_y(vS) - \partial_z(wS) + \partial_z(K_z \partial_z S) \quad (1)$$

225 With  $t$  the time dimension,  $x$ ,  $y$ ,  $z$  the zonal, meridional and vertical dimensions respectively,  $u$ ,  $v$ ,  $w$  the current on the  $x$ ,  $y$ , and  $z$  dimensions respectively,  $S$  the salinity,  $K_z$  the vertical diffusion coefficient.

The boundary conditions on salinity fluxes are:

- At surface ( $z = 0$ ):  $K_z \partial_z S = SSS(E - P)/\rho_0$

230 - At ocean bottom ( $z = -H$ ):  $K_z \partial_z S = 0$  (no exchanges through the bottom)

Over the mixed layer of height ( $h$ ), the salinity budget is computed as follows:

$$(1/h) \cdot \int_{-h}^0 (\partial_t S) \cdot dz = (1/h) \cdot \int_{-h}^0 \left( -\partial_x(uS) - \partial_y(vS) - \partial_z(wS) + \partial_z(K_z \partial_z S) \right) \cdot dz \quad (2)$$

Note  $S_m$  the depth-averaged salinity in the mixed layer:  $S_m = 1/h \cdot \int_{-h}^0 S dz$

235 The left-hand side can be expressed as the sum of the time variation of  $S_m$  and the entrainment term:

$$\begin{aligned} (1/h) \cdot \int_{-h}^0 (\partial_t S) \cdot dz &= \partial_t \left( (1/h) \cdot \int_{-h}^0 S \cdot dz \right) + \partial_t h/h \cdot \left( 1/h \int_{-h}^0 S \cdot dz - S(-h) \right) \\ \partial_t S_m + \partial_t h/h (S_m - S(-h))/h &= -(1/h) \cdot \int_{-h}^0 -\partial_x(uS) dz - (1/h) \int_{-h}^0 \partial_y(vS) dz \\ &\quad - (1/h) \int_{-h}^0 \partial_z(wS) dz + (1/h) \cdot \int_{-h}^0 \partial_z(K_z \partial_z S) \cdot dz \end{aligned} \quad (3)$$

240 The last term (related to vertical diffusion) is equal to:

$$(1/h) \int_{-h}^0 \partial_z(K_z \partial_z S) \cdot dz = (1/h) \cdot [K_z \partial_z S]_{-h}^0 = (1/h) \cdot SSS(E - P)/\rho_0 - (1/h)[K_z \partial_z S]_{-h} \quad (4)$$

CROCO computes online (for each time step)  $S_m$  and  $\partial_t S$ , so it can compute the entrainment term as a residual:

$$\partial_t h/h \cdot \left( 1/h \int_{-h}^0 S dz - S(-h) \right) = (1/h) \cdot \int_{-h}^0 (\partial_t S) \cdot dz - \partial_t \left( (1/h) \cdot \int_{-h}^0 S dz \right) \quad (5)$$

245 Thus, the final equation is:

$$\begin{aligned} \underbrace{\partial_t S_m}_{rate} = & - \underbrace{(1/h) \cdot \int_{-h}^0 -\partial_x(uS) dz}_{zonal\ advection} - \underbrace{(1/h) \cdot \int_{-h}^0 \partial_y(vS) dz}_{meridional\ advection} - \underbrace{(1/h) \cdot \int_{-h}^0 \partial_z(wS) dz}_{vertical\ advection} \\ & + \underbrace{(1/h) \cdot SSS(E - P)/\rho_0}_{forcing} - \underbrace{(1/h)[K_z \partial_z S]_{-h}}_{vertical\ mixing} - \underbrace{\partial_t h(S_m - S(-h))/h}_{entrainment} \end{aligned} \quad (6)$$

In this study, the mixing terms were found to be negligible compared to the other terms, and the horizontal and vertical advection terms were found to largely offset each other. Thus, in the following, the so-called advection term comprises the sum of the zonal, meridional and vertical advection terms. The runoff forcing is introduced via the zonal advection term  $-\partial_x(uS)$  as rivers are introduced as westward zonal flows at different locations of the west African coastline.

250

## 2.3 Comparison between SSS datasets and simulations

### 255 Analysis of cross-correlations

To study the relationship between the different variables involved in the salinity budget averaged over the e-NTA, we correlate forcing terms with salinities or other terms (such as Mixed Layer Depth (MLD)) using the determination coefficient ( $r^2$ ). To identify cause-and-effect relationships that can take weeks to establish, the correlation maximum by allowing a time delay (up to  $\pm 90$  days) between the variables is used.

260

### Climatological and interannual variability of the salinity budget

The different variables linked to the salinity budget are spatially averaged over the e-NTA, and the resulting time series are analyzed from January 2010 to July 2019, the time period over which ISBA runoff was available at the time of this study. The averaged seasonal and interannual signals are then extracted from the original signal, as follows:

265 *Seasonal signal:* a two-stage method is used to calculate a climatological seasonal variation. A daily climatology is first calculated by averaging data available on each day of the year between 2010 and 2019. To eliminate short-term fluctuations, the daily climatology is then smoothed using a 1-month moving-average filter.


*Interannual signal:* To remove the seasonal variations, the monthly climatology is subtracted from the original daily time series. A 3-month moving average is then applied to filter intraseasonal variability in order to focus on interannual variability at seasonal time scales.

### **Colocation methodologies when comparing in-situ and gridded data**

When comparing gridded SSS and Melax mooring or Argo floats SSS, a spatial bilinear interpolation of the gridded product and a selection of the nearest neighbor in time are used to collocate products represented on a grid with local data from Argo floats or Melax buoy. When comparing gridded SSS and TSG SSS, we first perform a smoothing of in-situ data with a gaussian window along the ship track to a resolution comparable to gridded datasets (e.g., model or satellite data type), given the high spatio-temporal sampling of TSG measurements. The standard deviation of this filter is set to one quarter of the spatial resolution of the model grid (95 % of the weight in a radius of half the spatial resolution). The resulting smoothed TSG data are compared to the nearest corresponding satellite or model pixel in time. Only the SSS in pixels that are common to modeled and satellite SSS are considered when comparing results obtained with both datasets.

## **3. Results**

### **3.1 Analysis and validation of the seasonal cycle of salinity**



285 **Figure 2: (a) Seasonal cycle of SSS in the e-NTA region: CCI (blue), GLORYS (cyan) and various CROCO simulations. The red area represents the range between the minimum and maximum value of the simulated SSS. (b) ERA5 and IMERG precipitation (orange, in  $\text{m}^3/\text{day}$ ), GloFAS and ISBA runoffs (black, in  $\text{m}^3/\text{day}$ ) climatologies. (c) Trends of the salinity balance equation (in  $\text{pss}/\text{day}$ ). The yellow line represents the effect of the atmospheric forcing, the blue line the lumped advection term (horizontal and vertical, including runoff forcing), the pink line the entrainment term at the base of the mixed layer. The red line is the SSS rate term, which corresponds approximately to the sum of the other three terms (vertical and horizontal diffusion are negligible). (d) Climatologies of the MLD (in m; blue) and salinity gradient at the base of the mixed layer (in  $\text{pss}/\text{m}$ ; red), for CROCOglofas simulation.**

290

300

A study of the salinity balance resulting from the different CROCO simulations was carried out to analyze the dynamical processes at the origin of the SSS seasonal cycle (Figure 2c). The SSS climatological variations are governed by the effect of precipitation during the summer rainy season. Rainfall accumulates on the continent during this period, generating intense river runoff 1-2 months later. The peak of river runoff is reached in September-October. This drives a continuous drop in salinity from the start of the rainy season onwards. These two effects also generate a strong vertical salinity gradient at the base of the mixed layer, and a decrease of the MLD (Figure 2d). The ocean transfers this freshwater input towards the ocean interior through vertical advection (not shown) and entrainment. During the dry season (January-May), the atmospheric forcing



is slightly positive (due to evaporation being larger than precipitation) and associated with a deeper mixed layer. There is also a strong vertical inflow of salt in the mixed layer by advection (not shown), which corresponds to the coastal upwelling particularly marked in March. Relatively salty upwelled waters are then redistributed by horizontal advection, notably by westward Ekman transport. This analysis is in line with Camara et al. (2015).


305

Although similar in shape and of the same order of magnitude, the seasonal cycles of the various SSS products present noticeable differences (Figure 2a): CROCO SSS have a seasonal cycle of smaller amplitude than that of CCI and GLORYS, and CCI SSS is in phase with CROCOglofas. On the other hand, CCI SSS lags CROCO SSS forced by ISBA runoffs (CROCOimerg and CROCOisba) and GLORYS SSS by ~2 weeks. Between the CROCO simulations, there is a difference of the order of 0.1 pss in amplitude, which may stem from a difference in the amplitude of the seasonal cycle of the precipitation products that were used, the amplitude of IMERG seasonal cycle being  $3 \cdot 10^8 \text{ m}^3/\text{day}$  larger than the one of ERA5 (Figure 2b).

310

### 3.2 Evaluation of the CROCO simulations using *in situ* measurements

#### 3.2.1 Coastal SSS from the Melax mooring off southern Senegal



315 **Figure 3: Surface salinity between March 2015 and August 2016 at the Melax mooring (black line), from CROCO simulations (red lines; red shading shows the range between the maximum and minimum simulated values), from satellite CCI data (blue), and from GLORYS data (cyan). Notice that the nearest satellite data pixel is further than 30 km apart from the mooring position and further than 50 km apart from the coast.**

Salinity measurements at the Melax mooring provide a useful time series for the evaluation of the simulation near the Senegalese coast. During its first two years of deployment, the mooring recorded an almost continuous time series, with a SSS oscillation of ~2-3 pss amplitude in November 2015 (Figure 3). CROCO SSS agree well with *in situ* SSS, during the dry seasons (March-September 2015 and March-July 2016) and at the time of the very strong oscillation in November. GLORYS also gives consistent results but underestimates the amplitude of the first observed oscillation of SSS (Mid-October – early November 2015). GLORYS SSS is also highly oscillatory and too low over the periods when *in situ* SSS is stable (before September 2015 and after May 2016). The CCI SSS is further from *in situ* SSS, which is expected as the pixel collocated with the mooring is 30 km offshore of the mooring (and 55 km from coast), due to the application of the coastal flag and the land contamination close to the coast. In addition, in such a coastal area with unresolved satellite SSS variability, an uncertainty arises from the sampling difference between a pointwise *in situ* measurements and a satellite measurement integrated over ~50km (Thouvenin-Masson, et al. 2022), which is greater than the GLORYS and CROCO horizontal resolutions.

The salinity balance is used to explain the origin of the strong oscillation detected in mid-2015 (see Figure S.5). Freshening is initiated in August 2015 by an event of intense precipitation and amplified by advection of freshwater from the coastal regions south of the mooring which collect a strong river runoff until November 2015. The observed oscillation in mid-November 2015 is caused by an oscillation of the zonal advection term, leading to an intensified westward (eastward) transport of relatively low (high) salinity waters (not shown).

335


### 3.2.2 Argo and TSG

Although most of the *in situ* measurements are taken at a depth between 5 and 10 meters, it has been chosen to compare them with the salinity in the top layer of the model, in order to be able to analyze these validations in the light of comparisons with satellite measurements taken at the first centimeter of the ocean (note also that there are no strong vertical salinity gradients in the top 5 m of the model water column).

340

Of the three types of gridded products, satellite observations are the closest to *in situ* data, with  $r^2$  values of 0.94 and 0.89 for comparisons with Argo and TSG data respectively (Table 2). Except for TSG measurements taken very close to the coast where the satellite data are slightly overestimated, the differences observed rarely exceed 0.2 pss in absolute value (Figure 4b,f).

345



**Figure 4:(top) Argo SSS (a) used as reference. Difference between Argo SSS and various SSS fields (CCI (b), GLORYS (c) and CROCOimerg (d)). (bottom) TSG SSS (e) used as reference. Difference between TSG SSS and various SSS fields (CCI (f), GLORYS (g) and CROCOimerg (h)). Comparisons were averaged over  $0.2^\circ$  boxes to ease visualization.**

350 Since the GLORYS reanalysis assimilates Argo data, the statistics of the comparisons with this dataset are very good, as expected. The  $r^2$  values are 0.91 when comparing to Argo, and 0.86 when comparing to TSG, only slightly lower than those obtained with CCI (Table 2). However, there is a negative bias of -0.8 to -1 psu with respect to Argo and TSG measurements taken on the continental shelf at the mouth of the Senegal River ( $16^\circ\text{N}$ ). These Argo data were taken at the end of 2012, which corresponds to a period when river outflows were particularly high (Figure 5c). It is therefore likely that these differences can  
 355 be explained by the use of climatological runoff in GLORYS (Figure 4c,g).

When CROCO SSS are compared with *in situ* SSS, a significant bias of up to +0.5 psu is observed, which is fairly systematic near the coast and north of  $14.7^\circ\text{N}$ . This positive bias is observed with respect to both Argo and TSG data and with all CROCO simulations, thus seems robust. South of  $14.7^\circ\text{N}$  and far from the coast ( $30^\circ\text{W}$ - $20^\circ\text{W}$ ), a negative bias of the order of -0.2 psu is observed, while the few TSG data available on the continental shelf show a positive bias (Figure 4d,h). There is a stronger  
 360 bias in CROCOprclm (Table 2), suggesting that the climatological precipitation field strongly reduces the effect of rainfall on SSS. Comparison statistics between the other simulations and *in situ* data are very close (with maximum absolute differences in terms of  $r^2$  of the order of 0.01), with slightly better results for CROCOimerg.


365 The seasonal variability of the CROCO SSS bias with respect to CCI shows similar patterns (positive bias near the north Senegalese – Mauritanian coast), regardless of the year (not shown). In the following, these simulations are therefore analyzed on a relative basis after removing the annual mean bias of SSS. The figures shown below are based on fields from which the mean SSS has been removed. The origin of the systematic SSS bias in CROCO is discussed in sect. 4.

370 The statistics calculated for the global and e-NTA regions are provided for reference in Table 2. There are significantly fewer co-located points in e-NTA, lower dynamics of SSS and a higher proportion of points close to the coast compared to the global region. Consequently, the statistics are consistently less favorable, with  $r^2$  values reaching 0.43 (0.67) compared to Argo floats (TSG).

375 **Table 2: Summary statistics of comparisons between various salinity products and in situ data over the global / e-NTA areas (see maps in Figure 4). “Std diff” stands for “standard difference”.**

	Argo (17378 / 902 points)			TSG (133139 / 8033 points)		
	$r^2$	Std diff	bias	$r^2$	Std diff	bias
CCI	0.94/0.70	0.16/0.30	-0.01/0.01	0.89/0.65	0.20/0.37	-0.01/-0.02
GLORYS	0.91/0.69	0.21/0.32	-0.02/0.01	0.86/0.64	0.23/0.38	-0.01/-0.02
CROCOglofas	0.81/0.43	0.30/0.45	0.07/0.19	0.77/0.65	0.30/0.36	0.14/0.14
CROCOroclm	0.80/0.25	0.31/0.62	0.06/0.14	0.77/0.66	0.30/0.35	0.14/0.13
CROCOisba	0.81/0.34	0.30/0.54	0.06/0.11	0.76/0.67	0.30/0.35	0.14/0.12
CROCOimerg	0.82/0.40	0.30/0.54	0.03/0.07	0.78/0.66	0.29/0.36	0.11/0.10
CROCOprclm	0.69/0.06	0.36/0.63	0.24/0.28	0.67/0.59	0.34/0.42	0.27/0.23

### 3.3 Origin of the main interannual variations




380 **Figure 5: Band-pass filtered SSS anomalies for a) CCI (blue), GLORYS (cyan) and the three interannual CROCO simulations (red;**  
**the red shading corresponds to the range between the minimum and maximum of the simulated SSS), b) CROCO simulation with**  
**interannual runoffs (purple solid line) and climatological runoffs (purple dotted line), CROCO simulation with interannual rain**  
**rate (magenta solid line) and climatological rain rate (magenta dotted line). Only pixels where CCI data are available have been**  
**considered in generating these curves. c) Anomalies of the different forcings, runoffs (brown, GLOFAS plain line, ISBA dashed line)**  
**and precipitation (orange, ERA5 plain line, IMERG dashed line). The y-axis has been reversed for easier comparison with SSS**  
385 **anomalies. Periods in grey shading correspond to large SSS anomalies for which a detailed analysis is given below.**

The interannual anomalies calculated for the different CROCO simulations are now compared to the interannual anomalies of the CCI and GLORYS products, used as references given their good agreement with *in situ* data (Table 2).

390 Interannual variations of SSS in the e-NTA are significant, oscillating between -0.4 pss and 0.4 pss and therefore of the same order of magnitude as the seasonal cycle (Figure 2). Overall, the different SSS estimates are in relatively good agreement. There is no long-term trend in the anomalies (Figure 5): for each year, anomalies are close to zero during the first half of the year (late winter-summer), and reach their extrema at the end of the year (fall-early winter), lagging by a few months the anomalies of rainfall and runoff (Figure 5c). The interannual variability of SSS derived from the CROCO simulations (Figure 5a) correctly represents the main variability compared to CCI and GLORYS. There are differences between CROCO 395 simulations, which can reach 0.2-0.3 pss during the rainy season at the end of the year (e.g., 2014, 2015). For the rest of the year, the differences remain negligible. The strongest CROCO SSS anomalies are generally produced by CROCOglofas. In 2011, 2015 and 2018 strong SSS anomalies were observed in the CCI and GLORYS products (see grey shading in Figure 5) and were well represented by the CROCO simulations (Figure 5a). The spatial distributions of SSS simulated by CROCOimerg and those of CCI SSS over the region for these three years are very similar during these time periods (Figure 400 6) and are now studied in more detail.

There is a significant disparity in the anomalies of the two river discharge forcing products, with anomalies sometimes having opposite signs (Figure 5c). This disparity is explored in the discussion (Section 4.2).





405 **Figure 6: SSS anomalies over the three analysed periods: late 2011 (a, b, c), late 2015 (d, e, f), and late 2018 (g, h, i). Left column (a,d,g): anomalies of the terms in the salinity balance equation (in pss/day) for CROCOimerg. The color code used is the same as that used in Figure 2a. Only pixels where CCI data are available have been considered in generating these curves. The black dotted line is the ISBA runoffs anomaly (the y-axis has been reversed). The grey shading indicates time periods of strong salinity variations.**

410 **Central column (b,e,h): simulated SSS maps (in pss) averaged over 3 months for CROCOimerg. Arrows show the surface currents anomalies. Right column (c, f, i): CCI SSS maps (in pss) averaged over 3 months. Grey contours depict IMERG precipitations (in mm/d, contour spacing is 1 mm/d; darker grey correspond to higher precipitation).**

### 3.3.1 Positive SSS anomaly in 2011

415 Around mid-2011, CROCO simulates a steep SSS increase and positive anomaly, which is in good agreement with the CCI anomaly (Figure 5a). The GLORYS anomaly displays the same variation albeit with a lower magnitude. Although all CROCO simulations reproduce the SSS increase, the CROCOisba anomaly is the closest to the CCI anomaly.

This anomaly is found in all CROCO simulations, and is therefore independent of the differences between precipitation and runoff forcing anomalies (Figure 5b,c). The fact that the SSS increase is also found in the simulation with climatological precipitations (CROCOprclm), and that the IMERG and ERA5 rainfall anomalies are of opposite sign (Figure 5c), suggests that this anomaly does not result from a rainfall anomaly. Moreover, the difference in runoff generates a difference in SSS anomalies that is only of second order. Consequently, the SSS increase must arise mainly from the ocean circulation. The salinity balance (Figure 6a) confirms that a positive anomaly of the advection term is the main cause of the SSS increase, while atmospheric forcing is compensated by entrainment. A closer analysis of the advection anomaly indicates that it is mainly due to the anomaly of currents ( $V'S$ , see Figure S.6, blue curve) related to an increase of the southward wind-driven coastal current (coastal jet) through the climatological poleward gradient of salinity (Figure 6b).

### 3.3.2 Negative SSS anomaly in 2015

430 Starting in mid-2015, most CROCO simulations show a significant freshening (from -0.4 pss for CROCOglofas to -0.2 pss for the other simulations) which is in good agreement with CCI and GLORYS (Figure 5a). In contrast, simulations forced by climatological precipitations (CROCOprclm) display no anomaly, while simulations forced by climatological (CROCOroclm) and ISBA (CROCOimerg) runoff display a freshening of 50% weaker than with GLOFAS runoff (CROCOglofas) (Figure 5b). This suggests that this freshening is initially due to the precipitation anomaly, followed by the subsequent runoff anomaly (Figure 5c). The analysis of the salinity balance equation confirms this hypothesis (Figure 6d,e,f): the freshening is due firstly to the rain intensification (October-December 2015). It is then slightly reinforced by the runoff in December-January and by an anomalous salty water outflow at the northern boundary of the e-NTA region (Figure S.6, blue curve). This anomalous advection of fresh water is amplified by the higher GLOFAS runoff in CROCOglofas. This effect is discussed in more details in sect. 4.



### 440 3.3.3 Positive SSS anomaly in 2018


In mid-2018, the SSS anomalies reach  $\sim 0.3$  pss (Figure 5a). As in 2011, all CROCO simulations, including those with climatological forcing, reproduce this positive anomaly, which cannot thus be attributed to one particular forcing anomaly. Analysis of the salinity balance (Figure 6g) reveals that the salty anomaly initially results from the strong negative precipitation anomaly (i.e., rain deficit) found in both ERA5 and IMERG products (Figure 5c; Figure 6i). The greater impact of the IMERG precipitation anomaly on CROCOimerg SSS (Figure 6i) could be due to a more localized and intense precipitation anomaly in IMERG than in ERA5 (see Figure S.1). This precipitation anomaly is accompanied by a large ISBA runoff negative anomaly (Figure 5c). This explains why CROCOprlm also simulates the positive SSS anomaly (albeit weaker than in CROCOimerg) without a precipitation anomaly. Because the GLOFAS runoff anomaly is positive (i.e. larger runoff), the simulations forced by ISBA runoff (CROCOimerg and CROCOisba) display a stronger anomaly than that forced by GLOFAS runoff (CROCOglofas).

In conclusion, the 2018 SSS anomaly is due to the combined effects of precipitation anomalies and river discharges. It is primarily caused by a strong precipitation negative anomaly (observed in both forcing datasets), which is not entirely compensated by entrainment (Figure 6c). This is then accompanied by a river discharge negative (positive) anomaly of ISBA (GloFAS) runoff, thereby accentuating (mitigating) the salinity anomaly through advection. This runoff anomaly explains the CROCOprlm SSS anomaly. The large GloFAS runoff is surprising as it is opposite to the rain deficit over the oceanic region during this period (Figure 5c).

### 3.4 SSS Sensitivity to the freshwater forcings

In this section, we investigate more thoroughly the sensitivity of the simulated SSS to a modulation of runoff forcing, all other model forcings being kept identical. We analyze the temporal variability of simulated salinity over the whole simulated period. Three test cases are set up using the simulations described in Table 1: the differences between simulations with a climatological runoff and a synoptic runoff (the GloFAS product) are first analyzed with regard to the interannual variability of forcing; then the difference in SSS induced by the use of the ISBA or GloFAS synoptic runoff products is analyzed. Last, the effects of SSS modulation by precipitation changes is presented.

The panels in Figure 7 depict these case studies. For each case, the difference in SSS between the studied simulations is shown in red, the differences in forcings in the e-NTA region in solid blue lines, and those occurring south of the e-NTA region ( $[5^{\circ}\text{N } 10^{\circ}\text{N} - 10^{\circ}\text{W } 20^{\circ}\text{W}]$ , Figure 1) in dashed lines. The region south of the e-NTA is indeed the site of strong freshwater influx, and the general oceanic circulation tends to advect these waters northward into the e-NTA region. Maximum of the cross-correlation function ( $r^2$ ) and corresponding temporal lag are determined over the e-NTA region and the region south of the e-NTA as explained in sect. 2.3. See Figure S.4 for more information about the cross-correlations and the lags between forcing differences and their effect on salinity.



475 **Figure 7:** a) Differences between GloFAS and ISBA runoffs anomalies (blue) and differences between SSS anomalies (red) (CROCOglofas-CROCOisba). b) GloFAS runoff anomalies (blue) and difference between SSS anomalies (red) (CROCOglofas-

CROCOroclm). Note that CROCOglofas, CROCOroclm CROCOisba are forced by the same ERA5 precipitation fields. In a) and b), blue lines show the sum of the runoff anomalies of rivers flowing directly into the e-NTA (local runoff, solid line) and south of the e-NTA (dashed line). c) Differences between IMERG and ERA5 precipitation anomalies (blue line) in e-NTA (solid line) and south of e-NTA (dashed line), and differences between SSS anomalies in e-NTA (red line) from simulations using IMERG (CROCOimerg) and ERA5 (CROCOisba) with the same ISBA runoff forcing. To ease the reading of the figure, the axis of precipitation and runoffs anomalies have been reversed. Maximum correlation ( $r^2$ ) and time lag between the time series of SSS anomaly and of the sum of the freshwater flux in and south of the e-NTA regions are indicated in the top left of the panels.

### 485 3.4.1 SSS sensitivity to interannual versus climatological runoff

The influence of GloFAS interannual runoff variability on salinity is investigated by comparing the CROCOroclm SSS to the CROCOglofas SSS. The other forcings (*i.e.*, ERA5) are kept identical between the two simulations (Table 1) so that the differences observed in SSS is mainly the consequence of the difference in river outflow forcing and its effect on nearshore ocean dynamics. Note that SSS differences can also arise from differences in mesoscale circulation due to dynamical (chaotic) nonlinearities or intrinsic variability unrelated to the forcings.

Interannual runoff variability has a significant effect on SSS (Figure 7a). The GloFAS runoff interannual variability is indeed correlated with the difference between CROCOglofas SSS and CROCOroclm SSS ( $r^2$  of 0.38 for a lag of 54 days; Figure 7a). Differences in SSS may be due to both a local and a remote runoff anomaly (*i.e.*, south of the e-NTA) (Figure 7a, dotted line). The salinity balance for CROCOglofas and CROCOroclm (not shown) indicates that, as for the climatological cycle (Figure 2c), a difference in runoff is partly compensated by a difference in entrainment of opposite sign. There is indeed a correlation of 0.98 between the runoff interannual anomaly and the entrainment terms difference, 75% of the runoff anomalies being compensated by entrainment on average. The lag time (54 days) between runoffs anomalies and SSS difference is likely due to a localized (nearshore) effect of the runoffs taking time to spread offshore and to modify the ocean surface layer over the entire e-NTA.

### 3.4.2 SSS sensitivity to a change in runoff interannual variability.

The differences between ISBA and GLOFAS runoffs anomalies (Figure 7b) are greater than the anomalies of either runoff taken independently, as the anomalies are frequently of opposite sign (Figure 5c). Consequently, the impact of the total runoff difference on salinity is larger than in the previous case study, with differences reaching 0.3 pss. A  $r^2$  of 0.56 is obtained between the difference in runoffs and the difference in SSS, with a time lag of 68 days; difference of runoffs is also offset by the entrainment at the bottom of the mixed layer: runoff difference and entrainment difference have indeed a correlation of 0.98 and entrainment compensates on average for 84 % of the runoff difference, with a time lag of 13 days.

In these two case studies, the effect of runoff is variable: small deviations from the climatology can generate significant differences in SSS as in 2015 (Figure 7a,b), while large runoff anomalies sometimes have a limited effect as in 2017 or 2018. These differences in behavior can be explained by surface current anomalies: in 2015, a northwesterly current transports the SSS anomaly linked to river flow, so that it has a greater impact on the mean SSS of the e-NTA region (Figure 6e). Thus, averaged SSS is particularly affected by a small change in runoff. Conversely, in 2018 (Figure 7b), the large difference ( $1.2 \cdot 10^9 \text{ m}^3 \text{ d}^{-1}$ ) between the two runoffs forcings with anomalies of opposite signs (Figure 5c), coincided with a northerly wind anomaly (not shown). The SSS anomaly produced by runoff in 2018 is therefore confined to the coasts south of the e-NTA, it does not spread northward, and it has a relatively little impact (0.22 pss) on the mean salinity of the area (Figure 6h).

### 3.4.3 SSS sensitivity to a change in rain rate interannual variability

The effect of a change in precipitation on the simulated salinity, more specifically on the difference induced by ERA5 (CROCOisba) and IMERG (CROCOimerg) synoptic precipitation products, is shown on Figure 7c.

The differences between the IMERG and ERA5 mean precipitation fields are small in comparison with the amplitude of their climatologies, (Figure 5c, Figure 2b), but the aggregated freshwater fluxes over the e-NTA are of the same order as the differences between runoffs forcings. The differences in SSS resulting from this difference in rain rate (Figure 7c) are weaker than those associated with runoff differences and not strongly driven by the direct effect of rain, in contrast to the effect of river discharge (sect. 3.4.2), as shown by the low correlation ( $r^2=0.19$ ). Note, however, that differences in SSS seem to be linked to differences in precipitation over certain periods (e.g., mid 2011, mid-2013, late 2014). As for runoff, the salinity balance of CROCOimerg shows that an anomaly in the rain rate forcing term is nearly totally compensated by an anomaly in the entrainment term (Figure 6a.d.g.). This adjustment is almost exactly correlated with the rainfall difference ( $r^2=0.99$ ), and the entrainment difference compensates in average 98 % of the forcing term difference. The weaker correlation with precipitation than with runoff anomalies could be explained by the fact that precipitation anomalies are weaker locally but spread over a larger region than runoffs. The influx of freshwater from rivers also occurs in coastal regions in the continental shelf region, where the mixed layer occupies the entire water column down to the bottom. This prevents the freshwater from being expelled by entrainment, forcing it to spread by advection, thus reducing the salinity of the mixing layer.

## 4 Conclusions and discussion

SSS are simulated over the period 2010-2019 using an ocean circulation regional model (CROCO) off the west African region forced by various precipitation and river runoff products. The simulated SSS are compared to various local and global datasets: the CCI satellite product, the GLORYS reanalysis, the ARGO floats database, a coastal mooring and TSG measurements. The

comparisons show that modelled SSS is systematically too high north of 15°N, but is quite consistent with observations in terms of anomalies. Moreover, comparison with a coastal mooring 30 km off the coasts of southern Senegal shows an excellent agreement. The simulation forced by IMERG precipitation and ISBA runoff is slightly closer to the observations than the other simulations, on average over the period of study.

The simulated SSS are analysed in terms of seasonal cycle and interannual anomalies averaged over the eastern North Tropical Atlantic (e-NTA). At first order, the amplitude and phase of the SSS seasonal cycle are only slightly modified by the different precipitation and runoff products used as model forcing. However, there is a time lag of about 2 weeks between the simulations, which corresponds to a shift in the seasonal cycles of the two runoff products. There is also a difference in amplitude of 0.1 pss, which may be due to a difference in precipitation forcing. The seasonal cycles of the CCI satellite data and the GLORYS reanalysis are also out of phase by about 2 weeks, which could originate from the climatological runoffs used in the GLORYS reanalysis. Analysis of the modelled mixed layer SSS budget indicates that the SSS decrease during the rainy season is driven initially by precipitation and a few weeks later by river runoff by means of horizontal advection of low salinity coastal waters. These negative trends are partly (nearly fully) compensated for runoffs (precipitation) by entrainment of relatively saline subsurface water into the mixed layer. This can be explained as follows: when precipitation occurs, surface waters become less saline, and a vertical gradient of salinity is formed in the surface layer. As the mixed layer depth deepens during night-time, saline subsurface water is incorporated (entrained) into the mixed layers, leading to an increased mixed layer salinity. This diurnal salinisation of the mixed layer occurs even when the mixed layer tends to decrease at a seasonal time scale (Figure 2d). Thus, the larger the precipitation, the larger the salinity vertical gradient, the larger the entrainment and compensation by salinisation of the mixed layer.

Despite the systematic model bias, modelled and observed SSS interannual variations are overall in good agreement. Large SSS anomalies are often correlated with large precipitation or runoffs anomalies within the e-NTA and from neighbouring regions whose surface waters are then advected by surface currents into the e-NTA. However, a propagation of the river plume is not systematic, and depends in particular on the wind-driven surface circulation patterns.

A study of the sensitivity of SSS to precipitation and runoff interannual variability shows a different response of the surface ocean to the two types of forcing. A difference in precipitation is almost totally compensated by entrainment, while a difference in runoff is compensated by between 75% and 84% on average. For a change in forcing of an equivalent order in terms of mean freshwater input, surface salinity is therefore more impacted by river runoffs than by precipitation.

565

#### 4.1 Uncertainties on SSS interannual variability

Over several time periods (*e.g.*, 2010, 2013, 2014 and 2016; Figure 5a), the CROCO SSS anomalies are markedly different from those of the reference products (GLORYS and CCI). In 2010, the CROCO anomalies appear to be underestimated compared with GLORYS and CCI. The latest shows a very strong freshening ( $\sim -0.4$  pss) during most of the year (Figure 5a) which is likely exaggerated. In 2010, the CCI dataset relies only on SMOS dataset which absolute calibration in 2010 is questionable (Boutin et al. 2021). The GLORYS product also shows a freshening ( $-0.15$  pss), but weaker than in CCI. The CROCOglofas simulation is the closest in 2010 to GLORYS. In 2013, there is a large disparity in SSS (up to 0.3 pss) between the various CROCO simulations. The CROCOglofas SSS displays a very weak anomaly, similar to CCI, while all the other CROCO simulations show a moderate freshening ( $-0.15$  pss to  $-0.28$  pss). These discrepancies could be explained by the relatively large difference in runoff during this period (Figure 5c). CROCOglofas is less biased than CROCOisba with respect to CCI and GLORYS, suggesting that the GloFAS 2013 anomalously low runoff is more realistic. In contrast, in 2014, CROCOglofas shows an unrealistic, high positive SSS anomaly, not found in the other simulations. This strong anomaly is associated with an anomalously low GloFAS runoff, in contrast to the ISBA anomalously high runoff driving the more realistic SSS anomaly in CROCOisba. In 2016, the CROCO simulations are in relatively good agreement with one another but display a much stronger freshening than found in both reference products (Figure 5a). Only CROCOprlm does simulate correctly the moderate freshening (Figure 5b, see magenta dotted line). Analysis of the salinity balance shows that advection drives the overly strong freshening (not shown), which may result from unrealistic surface currents in this period. Intrinsic variability may also play a role as all simulations including CROCOprlm are forced by the same ERA5 winds.


#### 4.2 Uncertainties on fresh water forcings

585 - Disparity between river discharge forcings interannual variability

This study highlights the disparity between two river discharge products available for West Africa. The GloFAS and the ISBA products are rather consistent in terms of seasonal cycle amplitude, with a maximum difference between seasonal cycles of  $3 \cdot 10^8$  m<sup>3</sup>/day (Figure 2b), but the phasing of the cycle is shifted by about 2 weeks and interannual anomalies are frequently of opposite sign (Figure 5c). This is in line with results from (Decharme et al. 2019) showing that runoffs simulated by different hydrological models driven by various precipitation products have a significant disparity in the intensities and phase of seasonal runoff cycles (*e.g.*, see Fig 14 of (Decharme et al. 2019)).

To understand the origin of these disparities, the runoff anomalies of each product are compared with the rainfall anomalies over the catchment areas of the rivers flowing through the study region (see Figure 1 for catchment delimitation). Runoff anomalies of each product are strongly correlated with the rainfall anomalies used in the hydrological models (Figure 8): GloFAS runoff anomalies are correlated with an  $r^2$  of 0.87 to the ERA5 rainfall anomalies over the catchment area, with a time lag of 22 days, and ISBA runoffs anomalies are correlated with the IMERG rainfall anomalies over the catchment area with

an  $r^2$  of 0.56, and with a time lag of 33 days. This suggests that the quality of runoffs estimation is highly dependent on the quality of the estimation of rainfall on land. Comparisons of modelled SSS with *in situ* SSS over the entire period (2010-2019) show slightly better results in simulations forced by ISBA but using the GloFAS (resp. ISBA) product leads to more accurate modelled SSS at the beginning (resp. end) of the studied period. In conclusion, the present study presents an indirect evaluation of runoff interannual variability by analyzing their impact on SSS.



605 **Figure 8: Runoff anomalies (brown) and watershed precipitation anomalies (orange), for GloFAS and ERA5 (solid lines) and ISBA and IMERG (dashed lines), over the catchment areas of the rivers flowing through the study region (see Figure 1 for catchment delimitation). Y-axis has been reversed.**

- Impact of a monthly climatological precipitation forcing

The CROCOprclm simulation was designed to suppress the effect of interannual precipitation variability on salinity. However, the calculation of a monthly climatological precipitation field by averaging monthly precipitation rates from various years drastically changes the distribution of precipitation (Figures S.2 and S.3) by smoothing and attenuating the highly localized precipitation phenomena. Climatological values do not exceed  $2 \cdot 10^{-2}$  m/day, which is too low for synoptic fields. So, rather than the effect of temporal precipitation anomalies, this simulation highlights the effect of a reduction of spatial rainfall variability. To reproduce a commonly observed forcing, it would be better to design a daily climatological field consisting of a succession of typical years with realistic precipitation distribution but scaled to a climatology in terms of total precipitation quantities. Such experiments are planned for future studies.

### 4.3 CROCO SSS biases

As seen in sect. 3.2.2, the CROCO SSS in all simulations were too high with respect to observations, mainly north of Cape Verde (15°N). This SSS bias is in fact associated with a positive temperature bias of ~0.5-1.5°C (Figure S.7 and text below). Such an SST bias is estimated to lead to an excess of evaporation that could explain about one third of the SSS bias (see histograms in Figure S.8). The remaining SSS biases could be due to a salinity bias in the subsurface waters transported to the surface layer by coastal upwelling and then offshore by Ekman currents.

Other processes neglected or misrepresented in the regional model may impact the SSS bias. First, the salinity of the river inflows (i.e., the runoff salinity), set to 15 pss in our study, may have an impact. This choice is debatable as it is expected that the salinity from rivers would be closer to 0 pss. For example, salinity gradually increases from ~0 pss at ~7 km from the coast to 10 pss at the estuary mouth of the Suwannee River in West Florida (Laurel-Castillo and Valle-Levinson (2023)).

Most West African river mouths have the particularity of being located near very flat coasts, which promotes the formation of large estuaries, despite their relatively low runoff (Descroix et al. 2020). These large estuaries allow the intrusion of seawater inland and facilitate water evaporation. Salinity in the Senegal River reach a minimum of 10 pss in October, at the peak of the flow, and 35 pss in winter (Mikhailov and Isupova (2008)). The Sine Saloum and Casamance rivers even have inverse estuaries, with estuarine salinity higher than that of the ocean (Pagès and Citeau 1990, Descroix et al. 2020).

The value of 15 pss was chosen considering mixing between river waters and seawater as well as evaporation inside the estuary. However, the effect of a change in CROCO runoff salinity can impact on SSS in the freshwater plume: a sensitivity study shows that setting the runoff salinity to 1 pss instead of 15 pss can lead to a decrease in SSS of about 1 pss in regions traversed by the plume (Figure S.9). Future studies should consider the seasonal variability of runoff salinity, when data is available. It should be noted that this effect on SSS from a reduction in runoff salinity has little impact offshore of the Senegal river (Figure S.9) and thus does not explain the positive SSS bias north of 15°N (Figure 4).

Last, tidal effects were not considered in our simulations, mainly to reduce computing time (a short model time step is required in the presence of strong tidal currents). However, it has been shown in the Amazon plume region that tides can impact plume propagation (Ruault et al. 2020) and conversely, that river flows can enhance tidal elevation (Durand et al. 2022). A sensitivity study of SSS to tides shows that tides can, in some cases, cause an increase in average SSS in the e-NTA region of about 1 pss due to increased vertical mixing (Figure S.10). Thus, including this effect would not reduce the positive bias north of 15°N and the impact on our results is likely to be weak. However, a more detailed evaluation of the tidal effect on river plume propagation in our region of interest would be needed to confirm these results.



Data availability: TSG data are available at <https://doi.org/10.6096/SSS-LEGOS>; Argo data have been downloaded from Pi-MEP database on <https://pimep.ifremer.fr/diffusion/data/cci-l4-esa-merged-oi-v3.2-7dr/argo/>. Melax data are available on <https://zenodo.org/records/4095436>. CCI data are available at The Centre for Environmental Data Analysis (CEDA) (<https://dx.doi.org/10.5285/5920a2c77e3c45339477acd31ce62c3c>). GLORYS reanalysis data are available on <https://doi.org/10.48670/moi-00021>; the ERA5 dataset provided by the ECMWF is available on <https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels?tab=overview>. CROCO model informations are available on <https://www.CROCO-ocean.org/> (Hilt et al. 2020). Given the large size of the modeling experiment outputs (~1.6 TB for each simulation), the dataset is not stored online and can be shared upon request to the corresponding authors. The ISBA-CTrip data are available from the authors (Decharme et al. 2019). IMERG data are available on [https://disc.gsfc.nasa.gov/datasets/GPM\\_3IMERGDF\\_07/summary?keywords=%22IMERG%20final%22](https://disc.gsfc.nasa.gov/datasets/GPM_3IMERGDF_07/summary?keywords=%22IMERG%20final%22); GloFAS data are available on <https://cds.climate.copernicus.eu/cdsapp#!/provider/provider-cems-without?tab=overview>.

Author contributions: CTM conducted the study and wrote the initial version of the paper. CTM, VE, JB designed the simulations, CTM prepared the forcing fields, VE performed the simulations and CTM analyzed them. CTM, JB, and JLV performed the satellite data processing and analysis. AL collected the Melax in-situ data. CTM, JB, VE, AL participated in the discussion of the results. All authors have read, improved, and agreed with the content of the paper.

Competing interests: The contact author has declared that none of the authors has any competing interests.

The Supplement related to this article is available online at doi:

Acknowledgements: This work is part of the PhD of Clovis Thouvenin-Masson funded by CNES and ACRI-st. We also received funding support from ESA CCI+SSS project (4000123663/18/I-NB) and CNES (the French National Center for Space Studies) through the TOSCA SMOS-Ocean. CROCO modelling experiments were conducted on the IDRIS HPC Jean-Zay under DARI projects A0110101140 and A0130101140. GLORYS global ocean model outputs were provided by the Copernicus Marine Environment Monitoring Service (CMEMS). We thank X. Perrot for discussions about rain rates products, M. Gevaudan, J. Jouanno and S. Biancamaria for providing precious information about ISBA-CTrip, and M. Marchesiello for insightful discussions about CROCO model.

- Akinsanola, A. A., W. Zhou, T. Zhou and N. Keenlyside (2020). "Amplification of synoptic to annual variability of West African summer monsoon rainfall under global warming." *npj Climate and Atmospheric Science* **3**(1): 21.
- Alory, G., T. Delcroix, P. Téchiné, D. Diverrès, D. Varillon, S. Cravatte, Y. Gouriou, J. Grelet, S. Jacquin, E. Kestenare, C. Maes, R. Morrow, J. Perrier, G. Reverdin and F. Roubaud (2015). "The French contribution to the voluntary observing ships network of sea surface salinity." *Deep Sea Research Part I: Oceanographic Research Papers* **105**: 1-18.
- 680 Ardoin-Bardin, S., A. Dezetter, E. Servat, J.-E. Paturel, G. Mahe, H. Niel and C. Dieulin (2009). "Using general circulation model outputs to assess impacts of climate change on runoff for large hydrological catchments in West Africa." *Hydrological Sciences Journal* **54**(1): 77-89.
- Beck, H. E., A. I. Van Dijk, V. Levizzani, J. Schellekens, D. G. Miralles, B. Martens and A. De Roo (2017). "MSWEP: 3-hourly 0.25 global gridded precipitation (1979–2015) by merging gauge, satellite, and reanalysis data." *Hydrology and Earth System Sciences* **21**(1): 589-615.
- 685 Boutin, J., N. Reul, J. Köhler, A. C. Martin, R. Catany, S. Guimbard, F. Rouffi, J.-L. Vergely, M. Arias, M. C. . and et al. (2021). "Satellite-based Sea Surface Salinity designed for Ocean and Climate Studies." *Earth and Space Science Open Archive*: 46.
- 690 Boutin, J. V., J.-L.; Reul, N.; Catany, R.; Koehler, J.; Martin, A.; Rouffi, F.; Arias, M.; Chakroun, M.; Corato, G.; Estella-Perez, V.; Guimbard, S.; Hasson, A.; Josey, S.; Khvorostyanov, D.; Kolodziejczyk, N.; Mignot, J.; Olivier, L.; Reverdin, G.; Stammer, D.; Supply, A.; Thouvenin-Masson, C.; Turiel, A.; Vialard, J.; Cipollini, P.; Donlon, C. (2021). "ESA Sea Surface Salinity Climate Change Initiative (Sea\_Surface\_Salinity\_cci): weekly and monthly sea surface salinity products, v03.21, for 2010 to 2020." *NERC EDS Centre for Environmental Data Analysis*, 08 October 2021.
- 695 Camara, I., N. Kolodziejczyk, J. Mignot, A. Lazar and A. T. Gaye (2015). "On the seasonal variations of salinity of the tropical Atlantic mixed layer." *Journal of Geophysical Research: Oceans* **120**(6): 4441-4462.
- Chandanpurkar, H. A., T. Lee, X. Wang, H. Zhang, S. Fournier, I. Fenty, I. Fukumori, D. Menemenlis, C. G. Piecuch, J. T. Reager, O. Wang and J. Worden (2022). "Influence of Nonseasonal River Discharge on Sea Surface Salinity and Height." *Journal of Advances in Modeling Earth Systems* **14**(2).
- 700 Chandanpurkar, H. A., J. T. Reager, J. S. Famiglietti and T. H. Syed (2017). "Satellite-and reanalysis-based mass balance estimates of global continental discharge (1993–2015)." *Journal of Climate* **30**(21): 8481-8495.
- Dai, A., T. Qian, K. E. Trenberth and J. D. Milliman (2009). "Changes in Continental Freshwater Discharge from 1948 to 2004." *Journal of Climate* **22**(10): 2773-2792.
- de Rosnay, P., G. Balsamo, C. Albergel, J. Muñoz-Sabater and L. Isaksen (2012). "Initialisation of Land Surface Variables for Numerical Weather Prediction." *Surveys in Geophysics* **35**(3): 607-621.
- 705 Debreu, L., C. Vouland and E. Blayo (2008). "AGRIF: Adaptive grid refinement in Fortran." *Computers & Geosciences* **34**(1): 8-13.
- Decharme, B., C. Delire, M. Minvielle, J. Colin, J. P. Vergnes, A. Alias, D. Saint-Martin, R. Sférian, S. Sénési and A. Voldoire (2019). "Recent Changes in the ISBA-CTrip Land Surface System for Use in the CNRM-CM6 Climate Model and in Global Off-Line Hydrological Applications." *Journal of Advances in Modeling Earth Systems* **11**(5): 1207-1252.
- 710 Descroix, L., Y. Sané, M. Thior, S.-P. Manga, B. D. Ba, J. Mingou, V. Mendy, S. Coly, A. Dièye, A. Badiane, M.-J. Senghor, A.-B. Diedhiou, D. Sow, Y. Bouaita, S. Soumaré, A. Diop, B. Faty, B. A. Sow, E. Machu, J.-P. Montoroi, J. Andrieu and J.-P. Vandervaere (2020). "Inverse Estuaries in West Africa: Evidence of the Rainfall Recovery?" *Water* **12**(3).
- Durand, F., C. G. Piecuch, M. Becker, F. Papa, S. V. Raju, J. U. Khan and R. M. Ponte (2019). "Impact of Continental Freshwater Runoff on Coastal Sea Level." *Surveys in Geophysics* **40**(6): 1437-1466.
- 715 Durand, F., L. Testut, J. Jouanno and A. C. Fassoni-Andrade (2022). "Role of the amazon outflow on the barotropic tide on the amazonian shelf." *Continental Shelf Research* **238**.
- Failler, P., H. El Ayoubi and A. Konan (2014). *Industrie des pêches et de l'aquaculture en Côte d'Ivoire*.
- Fournier, S., J. T. Reager, H. A. Chandanpurkar, M. Pascolini-Campbell and S. Jarugula (2023). "The Salinity of Coastal Waters as a Bellwether for Global Water Cycle Changes." *Geophysical Research Letters* **50**(24).
- 720 Gévaudan, M., F. Durand and J. Jouanno (2022). "Influence of the Amazon-Orinoco Discharge Interannual Variability on the Western Tropical Atlantic Salinity and Temperature." *Journal of Geophysical Research: Oceans* **127**(6).

- Guimbard, S., N. Reul, R. Sabia, S. Herlédan, Z. E. Khoury Hanna, J.-F. Piollé, F. Paul, T. Lee, J. J. Schanze, F. M. Bingham, D. Le Vine, N. Vinogradova-Shiffer, S. Mecklenburg, K. Scipal and H. Laur (2021). "The Salinity Pilot-Mission Exploitation Platform (Pi-MEP): A Hub for Validation and Exploitation of Satellite Sea Surface Salinity Data." Remote Sensing **13**(22): 4600.
- 725 Harrigan, S., E. Zsoter, L. Alfieri, C. Prudhomme, P. Salamon, F. Wetterhall, C. Barnard, H. Cloke and F. Pappenberger (2020). "GloFAS-ERA5 operational global river discharge reanalysis 1979–present." Earth System Science Data **12**(3): 2043-2060.
- 730 Hersbach, H., B. Bell, P. Berrisford, S. Hirahara, A. Horányi, J. Muñoz-Sabater, J. Nicolas, C. Peubey, R. Radu and D. Schepers (2020). "The ERA5 global reanalysis." Quarterly Journal of the Royal Meteorological Society **146**(730): 1999-2049.
- Hilt, M., F. Auclair, R. Benschila, L. Bordoio, X. Capet, L. Debreu, F. Dumas, S. Jullien, F. Lemarié, P. Marchesiello, C. Nguyen and L. Roblou (2020). "Numerical modelling of hydraulic control, solitary waves and primary instabilities in the Strait of Gibraltar." Ocean Modelling **151**.
- 735 Joyce, R. J., J. E. Janowiak, P. A. Arkin and P. Xie (2004). "CMORPH: A method that produces global precipitation estimates from passive microwave and infrared data at high spatial and temporal resolution." Journal of hydrometeorology **5**(3): 487-503.
- Laurel-Castillo, J. A. and A. Valle-Levinson (2023). "A Statistics-Based Analytical Framework for Intratidal and Tidally Averaged Salinity Distribution in Estuaries." Journal of Geophysical Research: Oceans **128**(12).
- 740 Lellouche, J.-M., E. Greiner, O. Le Galloudec, G. Garric, C. Regnier, M. Drevillon, M. Benkiran, C.-E. Testut, R. Bourdalle-Badie, F. Gasparin, O. Hernandez, B. Levier, Y. Drilllet, E. Remy and P.-Y. Le Traon (2018). "Recent updates to the Copernicus Marine Service global ocean monitoring and forecasting real-time 1/12° high-resolution system." Ocean Science **14**(5): 1093-1126.
- Lellouche, J.-M., E. Greiner, B.-B. Romain, G. Gilles, M. Angélique, D. Marie, B. Clément, H. Mathieu, L. G. Olivier, R. Charly, C. Tony, T. Charles-Emmanuel, G. Florent, R. Giovanni, B. Mounir, D. Yann and L. T. Pierre-Yves (2021). "The Copernicus Global 1/12° Oceanic and Sea Ice GLORYS12 Reanalysis." Frontiers in Earth Science **9**.
- 745 Mignot, J., C. de Boyer Montégut, A. Lazar and S. Cravatte (2007). "Control of salinity on the mixed layer depth in the world ocean: 2. Tropical areas." Journal of Geophysical Research **112**(C10).
- Mikhailov, V. N. and M. V. Isupova (2008). "Hypersalinization of river estuaries in West Africa." Water Resources **35**(4): 367-385.
- 750 Ndoye, S., X. Capet, P. Estrade, E. Machu, L. Kounta, B. Sow, M. Diakhaté and A. T. Gaye (2018). "A numerical modeling study of the Southern Senegal upwelling shelf: Circulation and upwelling source waters." African Journal of Environmental Science and Technology **12**(12): 487-500.
- Pagès, J. and J. Citeau (1990). "Rainfall and salinity of a Sahelian estuary between 1927 and 1987." Journal of Hydrology **113**(1-4): 325-341.
- 755 Roudier, P., A. Ducharne and L. Feyen (2014). "Climate change impacts on runoff in West Africa: a review." Hydrology and Earth System Sciences **18**(7): 2789-2801.
- Ruault, V., J. Jouanno, F. Durand, J. Chanut and R. Benschila (2020). "Role of the Tide on the Structure of the Amazon Plume: A Numerical Modeling Approach." Journal of Geophysical Research: Oceans **125**(2).
- 760 Shchepetkin, A. F. and J. C. McWilliams (2009). "Correction and commentary for "Ocean forecasting in terrain-following coordinates: Formulation and skill assessment of the regional ocean modeling system" by Haidvogel et al., J. Comp. Phys. **227**, pp. 3595–3624." Journal of Computational Physics **228**(24): 8985-9000.
- Supply, A., J. Boutin, G. Reverdin, J.-L. Vergely and H. Bellenger (2020). Variability of Satellite Sea Surface Salinity Under Rainfall. Satellite Precipitation Measurement: Volume 2. V. Levizzani, C. Kidd, D. B. Kirschbaum et al. Cham, Springer International Publishing: 1155-1176.
- 765 Tall, A. W., E. Machu, V. Echevin, X. Capet, A. Pietri, K. Corréa, S. M. Sall and A. Lazar (2021). "Variability of Dissolved Oxygen in the Bottom Layer of the Southern Senegalese Shelf." Journal of Geophysical Research: Oceans **126**(5).
- Thouvenin-Masson, C., J. Boutin, J.-L. Vergely, G. Reverdin, A. C. Martin, S. Guimbard, N. Reul, R. Sabia, R. Catany and O. Hembise Fanton-d'Andon (2022). "Satellite and in situ sampling mismatches: Consequences for the estimation of satellite sea surface salinity uncertainties." Remote Sensing **14**(8): 1878.
- 770 Wohl, E. and K. B. Lininger (2022). "Hydrology and discharge." Large Rivers: Geomorphology and Management, Second Edition: 42-75.

775 Zine, S., J. Boutin, J. Font, N. Reul, P. Waldteufel, C. Gabarro, J. Tenerelli, F. Petitcolin, J.-L. Vergely, M. Talone and S. Delwart (2008). "Overview of the SMOS Sea Surface Salinity Prototype Processor." IEEE Transactions on Geoscience and Remote Sensing **46**(3): 621-645.