

Millennial scale sea surface temperatures of the western Arabian Sea between 37 - 67 ka BP

Jennifer Scott^{1,2}, Douglas Coenen^{3,4}, Simon Jung²

¹ School of Engineering and Physical Science, Heriot Watt University, Edinburgh, EH14 4AS, United Kingdom

² School of Geosciences, University of Edinburgh, Edinburgh, EH9 3FE, United Kingdom

³ Institute of Geosciences, Goethe University Frankfurt, Frankfurt am Main, Germany,

⁴ Frankfurt Isotope and Element Research Center (FIERCE), Goethe University Frankfurt, Frankfurt am Main, Germany

Correspondence to: Jennifer Scott (js2092@hw.ac.uk) or Simon Jung (simon.jung@ed.ac.uk)

Abstract. The Asian monsoon system is a crucial part of the global climate system affecting a significant proportion of the world population. Understanding the controls for changes in the monsoon system is crucial for meaningful assessments of future climate change. The Arabian Sea is part of the wider Asian monsoon system and has been studied regarding controls of monsoon variability through time. In this study we present sea surface temperature data from 37 - 67 ka BP from sediment core NIOP 929 from the western Arabian Sea assessing the importance of northern/southern hemispheric climate change driving monsoon circulation in the Arabian Sea. Earlier work implies a straightforward link between monsoon variation in the Arabian Sea and northern hemisphere millennial scale climate change during glacial periods, as depicted in Greenland ice cores. We present a new millennial-scale Mg/Ca based sea surface temperature reconstruction based on the planktic foraminifera species *G. bulloides* and *G. ruber*. We use these data to calculate seasonal sea surface temperatures. The SST data are variable with a maximum short-term change of 8-9 °C. The variations in our SST records appear not related to change in either hemisphere in a straightforward fashion by not showing a phase-locked relation to millennial scale change in Greenland or Antarctic ice core records. We discuss these changes in the context of the Arabian Sea potentially being a “melting pot” with both the northern and the southern hemisphere exerting influence on a seasonal scale.

1. Introduction

The climate and surface ocean conditions of the Arabian Sea are controlled by the seasonally reversing Asian monsoon winds (Brock et al., 1992; Prell and Curry, 1981; Rixen et al., 1996; Shetye et al., 1994). The change from the summer southwest monsoon (SW monsoon) to the winter northeast monsoon (NE monsoon) entails seasonally

and spatially varying patterns in sea surface temperature/circulation, nutrient distribution and biological productivity (Anand et al., 2008; Conan and Brummer, 2000; Curry et al., 1992; Ivanochko et al., 2005; Jung et al., 2002; Peeters et al., 2002; Rostek et al., 1997; Saher et al., 2007b; Saher et al., 2007a; Saher et al., 2009; Schulz et al., 1998). The SW monsoon in particular is socio-economically significant because it drives rainfall on the Indian subcontinent, affecting the livelihood of billions of people through agriculture and water resources (Gadgil and Gadgil, 2006; Mishra et al., 2020; Webster et al., 1998).

The SW monsoon is caused by differential land-ocean heating and the resulting atmospheric pressure gradients between the Indian subcontinent and the Indian Ocean. During summer, development of a low-pressure cell over the Tibetan Plateau causes the intertropical convergence zone (ITCZ) to shift northward, inducing south-westerly winds in the western Arabian Sea (Schott and McCreary, 2001) (Fig. 1). These winds drive upwelling along the coast of Somalia and Oman through Ekman transport (Clemens et al., 1991; Wyrki, 1973) (Fig. 2a). Modern-day sea surface temperatures (SSTs) have been recorded as low as 16°C during the summer SW monsoon (Swallow and Bruce, 1966), reflecting the cold, nutrient rich waters surfacing from a depth of around 200m (Peeters et al., 2002) (Fig. 2a). During winter the pressure gradient reverses and the ITCZ shifts southward, inducing northeasterly monsoon winds in the Arabian Sea (Fig. 1). No upwelling occurs off Somalia during the winter and the inter-monsoon periods (Fig. 2b), entailing generally higher SSTs: typically, in the modern-day between 25–28°C but can reach over 30°C (Swallow and Bruce, 1966).

The seasonally changing wind interacting with surface waters results in coastal upwelling off Somalia and Oman during summer and no upwelling in these regions during winter (Clemens, 1991 #19762; Wyrki, 1973 #19763). During upwelling, modern sea surface temperatures (SSTs) are as low as 16°C during summer (Swallow, 1966 #19764), reflecting the cold, nutrient rich waters surfacing from a depth of around 200m (Peeters, 2002 #19756). During winter and the inter-monsoon periods, when upwelling ceases, SSTs are higher and can reach over 30°C (Swallow, 1966 #19764). The sedimentary record reflects the seasonally reversing upwelling signal and has been used to reconstruct monsoon variability on glacial-interglacial timescales, with a stronger SW monsoon being recorded as lower sea surface temperatures and increased productivity (Rostek et al., 1997). Productivity proxies used to investigate monsoon variability include abundance pattern in planktonic foraminifera (Anderson and Prell, 1993; Gupta et al., 2003; Kroon, 1991), $\delta^{18}\text{O}$ (Prell et al., 1992), total organic carbon content (Caley et al., 2013; Ivanochko et al., 2005; Schulz et al., 1998), $\delta^{15}\text{N}$ (Altabet et al., 1995; Ivanochko et al., 2005), Ba/Al ratios (Ivanochko et al., 2005), bromine records (Caley et al., 2013) and changes in coccolithophore assemblages (Beaufort et al., 1997). Several authors have studied the glacial-interglacial SST history of the western Arabian Sea, using alkenones (Rostek et al., 1997; Saher et al., 2009; Sonzogni et al., 1997) and Mg/Ca derived calcification

temperatures of planktonic foraminifera (Anand et al., 2008; Ganssen et al., 2011; Saher et al., 2007b; Saher et al., 2007a; Saher et al., 2009). These studies support the notion that summer monsoon induced upwelling may have been slightly weakened during the last glacial, but principally persisted throughout this period. Compared to glacial-interglacial SST reconstructions there is a lack of high-resolution, centennial to millennial scale records from the region, with the only available data sets covering the last deglaciation (Anand et al., 2008; Saher et al., 2007b; Saher et al., 2007a) and the previous interglacial (Saher et al., 2009). In this study we will address this shortfall by presenting a millennial scale Mg/Ca derived SST record for *Globigerinoides ruber* and *Globigerina bulloides* from sediment core NIOP 929 spanning 37 to 67 ka BP. A novel proxy is employed to reconstruct the winter and summer monsoon sea surface temperatures using these foraminiferal Mg/Ca based temperatures.

Formatted: Font: Italic

2. Methods

2.1. Core NIOP 929

Sediment core NIOP 929 was recovered during the 1992 - 1993 Netherlands Indian Ocean Programme (NIOP) (Van Hinte et al., 1995). The 16.15m long core was taken from the western Arabian Sea (Fig. 1) (13°42' 21" N, 53°14' 76" E) at 2490m water depth, with the sampling location being within the modern day summer monsoon induced upwelling area off the coast of Somalia (Rostek et al., 1997; Saher et al., 2007a).

2.2. $\delta^{18}\text{O}$ record

Sampling of core NIOP 929 involved continuous slicing of the top 210 cm of the sediment sequence in 0.5cm steps. Below this depth sampling continued at 1cm steps. The stable oxygen isotope data from the top 210 cm used in this study have been published previously (Saher et al., 2007b; Saher et al., 2007a). The remainder has been measured at the University of Edinburgh and are reported on here. Samples were freeze dried and wet-washed over a 63µm sieve before 25-30 specimens from the 250 -355 µm size fraction of both *G. ruber* and *G. bulloides* were picked for stable isotope analysis. In rare cases where samples did not contain sufficient specimen less than 25 specimen were analysed. Sample preparation for stable isotope analysis followed standard procedures. The stable isotope analyses for the top 210 cm of core NIOP 929 were carried out at the Vrije Universiteit Amsterdam using a Finnigan MAT252 equipped with a Kiel device. Deeper sections of the core used in this study were measured at the School of

Geosciences at the University of Edinburgh with a Thermo Electron Delta+ Advantage stable isotope mass spectrometer coupled to a Kiel Carbonate III preparation device. The reproducibility of the Amsterdam oxygen isotope data is 0.08‰ and that of the Edinburgh data is 0.1‰.

2.3. NIOP 929 age model

The age model for core NIOP 929 is based on a stepwise approach (compare Fig. 3). In step 1, the age model for the upper 210 cm of core NIOP 929 is established based on seven ^{14}C ages from (Saher et al., 2007b). The ^{14}C ages were recalibrated to calendar ages using the ^{14}C age calibration program Calib 8.2 (Stuiver and Reimer, 1993), using a ΔR value of 45 (± 67) (Southon et al., 2002) (Tab. 1, Fig. 3). In steps 2 and 3, the age model of samples deeper than 210 cm is based on oxygen isotope stratigraphy. The NIOP 929 *G. ruber* $\delta^{18}\text{O}$ record was initially tuned to the LR04 $\delta^{18}\text{O}$ global stack of (Lisiecki and Raymo, 2005) (step 2), as the population of *G. ruber* reflects annual average conditions of the study site (Conan and Brummer, 2000). The LR04 age control points are shown in Tab. 2. In the final (third) step, this initial age model was then refined by tuning to the Greenland ice core NGRIP $\delta^{18}\text{O}$ record (Andersen et al., 2006; Rasmussen et al., 2006; Svensson et al., 2006; Vinther et al., 2006; Wolff et al., 2010) (the combined set of tuning based age control points is shown in Tab. 3). This approach is supported by earlier findings implying that monsoonal change at the millennial scale in the Arabian Sea based on synchronous millennial-scale variations in the Arabian Sea and in phase with millennial scale change in the North Atlantic region (Altabet et al., 1995; Ivanochko et al., 2005; Jung et al., 2002; Schulz et al., 1998) (Fig. 23). Both the NGRIP $\delta^{18}\text{O}$ record and the initially age modelled NIOP 929 *G. ruber* $\delta^{18}\text{O}$ record were filtered to a 700-yr running mean before tuning to better constrain the expression of millennial scale climate variability in the records (Zeeden et al., 2020) (Fig. 32).

The uncertainties in our age model depend on the nature of the age control points. AMS ^{14}C based age estimates entail 1 sigma uncertainty ranges between ~200 and ~350 years (Tab. 1). The 2 sigma age envelopes range from ~410 to ~710 years (Tab. 1). The maximum layer counting error for the NGRIP record varies between 7% during the cold stadial events and 4% during the warm interstadial events (Andersen et al., 2006). The uncertainty in the LR04 record varies throughout the 5 Ma record. For the youngest section, i.e. 1-0 Ma it is around 4 ka (Lisiecki and Raymo, 2005). Whilst these uncertainties limit the accuracy of absolute dating of specific events beyond the capabilities of the individual dating method, the comparability of the sequence of events in our study is also depended on the quality of the visual alignment of events within the temporal framework determined by the reference record. It is difficult to quantify this uncertainty, but our best conservative guess that it is in the region of 2-3 ka. Whilst this

is potentially a reason for concern when comparing short-term events, in the absence of absolute dating methods covering the entirety of the record presented in this study, it is the best that can be done.

2.4. Mg/Ca paleothermometry

Sediment samples for Mg/Ca paleothermometry were obtained from core NIOP 929 from 387.5 - 643.5 cm depth (Van Hinte et al., 1995). Availability of samples did not allow for continuous sampling at 1cm spacing. No samples were available from 644.5 - 655 cm depth due to sample loss in core recovery (Van Hinte et al., 1995). Twenty-five tests of both *Globigerinoides ruber* (white) and *Globigerinoides bulloides* from the 250 - 355 µm size fraction were picked for Mg/Ca analysis. Samples were prepared using the method described by (Barker et al., 2003), with an additional MilliQ rinse and ultrasonification prior to clay removal instead of crushing samples due to high levels of sample loss. Samples were analysed at the School of GeoSciences at University of Edinburgh on a Varian ICP-OES Vista Pro (De Villiers et al., 2002). Instrumental precision was ± 1% based on replicate measurements of the ECRM 752-1 carbonate standard. The measured Mg/Ca concentrations were then converted to temperatures using Eq. (1) for *G. ruber* (Anand et al., 2003) and Eq. 2 for *G. bulloides* (Mashiotta et al., 1999):

$$Mg/Ca = 0.34 e^{0.827T}$$

(1)

$$Mg/Ca = 0.47 e^{0.107T}$$

(2)

where Mg/Ca is the measured Mg/Ca ratio of the sample and T the reconstructed temperature recorded in the foraminiferal calcite. The Mg/Ca temperature estimate of (Mashiotta et al., 1999) was used instead of (Vázquez Riveiros et al., 2016) as the temperature range of 10 - 25°C better suits the expected temperature range of the environment (Vázquez Riveiros et al., 2016).

2.5. Winter and summer temperature estimates

Previous work by (Saher et al., 2007a) used the δ¹⁸O values for both *G. ruber* and *G. bulloides* for core NIOP 929 in conjunction with Mg/Ca temperatures from *G. ruber* to reconstruct summer and winter monsoon temperatures for the past 20 ka BP. Part of their study involved analysing seasonal foraminiferal abundance data from nearby sediment traps (Conan and Brummer, 2000; Curry et al., 1992; Nair et al., 1989) in order to define the *G. ruber* and *G. bulloides*

flux throughout the year. A zero flux of both *G. ruber* and *G. bulloides* was assumed for the inter-monsoon periods,
 140 based on the sediment trap data. With *r* and *b* denoting the fractions of *G. ruber* and *G. bulloides* which calcify during
 the summer monsoon season, their analysis yielded results of 0.54 ± 0.08 for *r* and 0.91 ± 0.06 for *b* (Saher et al.,
 2007a).

As this study reconstructs both *G. ruber* and *G. bulloides* Mg/Ca temperatures, we can apply these seasonal
 abundance values to reconstruct winter and summer monsoon temperatures. As illustrated by (Saher et al., 2007a),
 145 a population of *G. ruber* will record a temperature which can be attributed as 0.5 summer temperature and 0.5 winter
 temperature (Fig. 2) (Conan and Brummer, 2000). *G. bulloides* is predominantly present during the summer monsoon
 period, however a small portion is present within the winter monsoon period meaning the temperature signal
 recorded in a population can be inferred as 0.9 summer temperature and 0.1 winter temperature (Fig. 2) (Conan and
 Brummer, 2000). Therefore, winter and summer monsoon temperatures can be calculated from *G. ruber* and *G.*
 150 *bulloides* Mg/Ca temperatures as shown in Figure 2.

~~As this study reconstructs both *G. ruber* and *G. bulloides* Mg/Ca temperatures, we can apply these seasonal
 abundance values to reconstruct winter and summer monsoon temperatures. As illustrated by (Saher, 2007 #19752),
 a population of *G. ruber* will record a temperature which can be attributed as 0.5 summer temperature and 0.5 winter
 temperature. *G. bulloides* is predominantly present during the summer monsoon period, however a small portion is
 155 present within the winter monsoon period meaning the temperature signal recorded in a population can be inferred
 as 0.9 summer temperature and 0.1 winter temperature. Therefore, winter and summer monsoon temperatures can
 be calculated as shown in Equation 3 and 4 respectively, with *R* = *G. ruber* Mg/Ca temperature, *B* = *G. bulloides*
 Mg/Ca temperature, *W* = winter monsoon temperature and *S* = summer monsoon temperature:~~

$$\frac{1.8R - B}{0.8} = W \quad (1.8R - B) / 0.8 = W$$

$$(B - 0.1W) / 0.9 = S$$

(4)

165 **8.3. Results**

8.1.3.1. $\delta^{18}\text{O}$ records

The $\delta^{18}\text{O}$ records of *G. ruber* and *G. bulloides* from core NIOP 929 cover the time interval from 1.3 ka BP to 126 ka BP on a centennial scale (Fig. 23). The $\delta^{18}\text{O}$ *G. ruber* record has a range of values from 0.4‰ at 21.4 ka BP to -2.3‰ at 1.8 ka BP. The $\delta^{18}\text{O}$ *G. bulloides* record has a range of values from 0.3‰ at 22.7 ka BP to -2.1‰ at 8.5 ka BP. Within the time window between 35 and 70 ka BP (Fig. 43) both stable oxygen isotope records are dominated by millennial to sub-centennial oscillations. In the stable isotope record of *G. ruber* this variability is superimposed on weakly defined long-term minima centered at ~60 ka BP, ~52-53 ka BP, ~45-46 ka BP, with maxima being recorded in between. Weakly defined long-term minima in the stable oxygen isotope record of *G. bulloides* are centered at ~57-58 ka BP, ~52 ka BP, ~44-46 ka BP and 38 ka BP, alternating with maxima occurring in between.

175 **8.2.3.2. Mg/Ca temperature reconstructions**

The Mg/Ca based temperature records of *G. ruber* and *G. bulloides* from core NIOP 929 cover the time interval from 37.4 ka BP to 67.7 ka BP on a sub millennial scale (Fig. 3-54-6). The *G. ruber* Mg/Ca temperature record has a range of 9.2°C from a low of 21.2°C at 49.0 ka BP, to a high of 30.4°C at 39.2 ka BP, with a mean temperature of 23.7°C (Fig. 4, 5, 56). The average temporal resolution for the *G. ruber* record over the reconstructed period is 203 yrs. The *G. bulloides* Mg/Ca temperature record has a similarly large range in temperatures of 9.9°C, with the lowest value of 16.0°C at 62.5 ka BP, the highest value of 26.8°C at 64.8 ka BP and a mean temperature of 21.2°C (Fig. 4, 5, 56). The average temporal resolution for the *G. bulloides* record is 216 yrs.

In Figure 34, we use 700 year box car filtered times series to focus on the most robust signal in our SST records. Both, the *G. ruber* and the *G. bulloides* based SST records display several maximum and minimum spikes at a millennial scale. Prominent maximum spikes in the *G. ruber* SST records are centred at ~61 ka BP, 57-54 ka BP (two sub peaks), ~51 ka BP, ~42 ka BP, ~40.5 ka BP and ~39.5 ka BP, with the highest temperatures of >27°C being linked with the youngest maxima. Pronounced minima in the *G. ruber* SST record are centred at ~63 ka BP, ~57-57.5 ka BP, ~52 ka BP, ~49 ka BP (broad maxima), ~46 ka BP, ~44 ka BP, ~41.5 ka BP, ~40 ka BP and ~37 ka BP. Some of the maximum spikes in the *G. ruber* record are roughly matched in the *G. bulloides* SST record (Fig. 43), but there are differences too. Distinct maxima in the *G. bulloides* SST record are centred at ~67 ka BP, ~65 ka BP,

Formatted: Font: Italic

Formatted: Font: Italic

~61 ka BP (broad maxima), ~52 ka BP, ~49-46 ka BP (two sub peaks), ~41.5 ka BP, ~40.5 ka BP and ~38.5 ka BP. Interestingly, contrasting the *G. ruber* SST record, the highest values the *G. bulloides* SST record appear near the beginning and the end of the record. Minima in the latter occur at ~66 ka BP, ~63 ka BP, ~57-57.5 ka BP, ~53.5 ka BP, ~49.5 ka BP, ~45 ka BP, ~42.5 ka BP and ~39.5 ka BP, with amplitudes diminishing in the in the upper part of the record.

In relation to orbitally driven climate change, whilst not being at the centre of this study, it is interesting to note that both records suggest some similarity with summer insolation at 30°N, with the *G. bulloides* SST record showing a slightly stronger affinity.

8.3.3.3. Winter and summer temperature reconstructions

The Mg/Ca based temperature records of *G. ruber* and *G. bulloides* were used to reconstruct winter and summer monsoon temperatures from 37.4 ka BP to 67.7 ka BP (Fig. 4.67). Winter temperature is lowest at 17.3°C at 64.8 ka BP and highest at 34.7°C at 37.5 ka BP, with an overall range of 17.4°C and mean temperature of 26.7°C (Fig. 7). Summer temperature is lowest at 15°C at 62.5 ka BP and highest at 27.7°C at 37.9 ka BP, with a range of 12.4°C and a mean temperature of 20.4°C (Fig. 7).

The general shapes of the 700 year box car filtered, season specific winter and summer temperature records are distinctly different (Fig. 43), by being dominated by (mostly) negative and positive excursions, respectively. For the majority of the winter SST record, maximum values are around 28 - 29°C in Fig. 43, with a maximum centred at 39.5 ka BP being a possible exception. Prominent minima in winter temperatures occur around 65 ka BP, ~52-52.5 ka BP, ~41.5 ka BP (single data point) and 38 ka BP. These minima are superimposed on a trend of generally increasing winter temperatures in the early part of the record. Whilst close to the detection limit of the record, the long-term component of the winter temperature time series resembles summer insolation at 65degN (Fig. 43).

For the majority of the summer temperature time series, minimum values are around 18 °C or slightly below. Prominent SST maxima occur at ~67 ka BP, ~65 ka BP, ~61-62 ka BP (broad), ~52 ka BP, ~49-46 ka BP (two sub peaks), ~42 ka BP, and ~38 ka BP. These short-term maxima are superimposed on a long-term oscillation with generally higher summer temperatures occurring near the beginning and the end of the sequence.

The longer-term variability in winter and summer temperatures can be subdivided into four periods (Fig. 43). In period I from 37 - 42 ka BP summer monsoon temperatures are on average ~3.7°C lower than the winter monsoon temperatures compared to a larger temperature difference in the preceding period. The reduced temperature difference in period I stems from two prominent excursions at ~41.5 ka BP and ~38 ka BP (Fig. 43) in both, summer

Formatted: Superscript

220 and winter temperatures, showing maximum and minimal values, respectively. During both events (one of which
only defined by one data point), the normal temperature gradient (sensu Swallow and Bruce, 1966) reversed with
summer temperatures exceeding winter values by 4-6 degree in Fig. 43.
Period II from 42 – ~51.7 ka BP is more consistent with the seasonal temperature gradient of modern monsoon
dynamics showing mean summer monsoon temperatures 7.4°C lower than winter values. During this period there
225 are no instances where summer temperatures are warmer-higher than winter temperatures, with winter temperatures
between 4.2 – 11.9°C higher than summer temperatures. During period III from 51.7 – 64.4 ka BP summer monsoon
temperatures were on average 7.5°C lower than winter temperatures, amid short term variability in particular in the
unsmoothed record (Fig. 67) with summer temperatures being up to 4.7°C higher than winter values. In the smoothed
record in Fig. 43, this is reflected by short periods with summer and winter temperature differences being close to
230 zero, most prominently depicted at ~52 ka BP. During Period IV from 64.4 – 67.7 ka BP mean summer temperatures
are ~0.3°C higher than winter values. There is, however, a large variability displayed during this period. Similar to
period I, there are two prominent temperature excursions in both, summer and winter temperatures, showing
maximum and minimal values, respectively. Both events, centred at ~65 ka BP and ~66.5 – 67 ka BP, entail a reversal
of the normal temperature gradient with summer temperatures exceeding winter values by ~4.5°C and ~1°C,
235 respectively.

9.4. Discussion

9.1.4.1. Robustness of Mg/Ca based SST estimates in the Arabian Sea

Here we present the most complete temperature history of core NIOP 929 to date (Fig. 43, 5). The rapid SST changes
reconstructed in this study are greater than those of any previous reconstruction for core NIOP 929 (Fig. 43, 65). The
240 9.2°C range of temperatures reported in the *G. ruber* record of this study is much greater than the 5.3°C range in
temperatures in *G. ruber* across the last deglaciation (Saher et al., 2007b; Saher et al., 2007a) and 4.2°C range across
the penultimate interglacial (Saher et al., 2009).

The high-resolution Mg/Ca temperature reconstructions of this study from 37.4 ka BP to 67.7 ka BP can be directly
compared to the alkenone based Uk'37 temperature record of (Rostek et al., 1997). Alkenone Uk'37 temperature
245 estimates have been shown to reflect annual average SSTs (Bijma et al., 2001; Budziak et al., 2000; Müller et al.,
1998; Sonzogni et al., 1997), and with roughly equal fluxes of *G. ruber* calcifying in the winter and summer in the

Arabian Sea (Conan and Brummer, 2000; Curry et al., 1992), the *G. ruber* Mg/Ca SST record can also be considered an annual average SST record. Overall, there is reasonably good agreement between the two proxies, with the alkenone Uk'37 record of (Rostek et al., 1997) tracking the mean value of the higher resolution sections of this study's *G. ruber* Mg/Ca record (Fig. 6S). For the lower resolution period 43 - 52 ka BP the records do not agree as well, with the alkenone Uk'37 temperature reports values around 2°C higher than those of the *G. ruber* Mg/Ca record (Fig. 6S).

Previous studies have noted the disparity in the values of alkenone Uk'37 and Mg/Ca based temperature records for core NIOP 929, with (Saher et al., 2007b) finding the alkenone Uk'37 based temperature estimates of (Rostek et al., 1997) around 2°C higher than their Mg/Ca *G. ruber* record for the last deglaciation, and (Saher et al., 2009) finding a near constant offset of 3.5°C between the alkenone Uk'37 record of (Sonzogni et al., 1997) and their *G. ruber* Mg/Ca record during MIS 5 – 6 (Fig. 6S). Similar offsets have been documented in the North Atlantic Ocean for the past 28 ka BP (Elderfield and Ganssen, 2000) and the Equatorial Atlantic Ocean for the last 270 ka BP (Nürnberg et al., 2000). The transfer function used to convert alkenone saturation to temperature and the equation to convert Mg/Ca ratio to temperatures can affect the offset between the two records (Saher et al., 2009). As the alkenone saturation values of (Rostek et al., 1997) are not available, changing the transfer function used to investigate the potential difference is not an option. Using the alternative Mg/Ca to temperature equation of (Elderfield and Ganssen, 2000) would only increase the temperature offset (Saher et al., 2009). An alternative explanation for the difference in the alkenone Uk'37 and Mg/Ca based temperature records is changes in carbonate chemistry through the carbonate ion effect (Elderfield et al., 2006; Lea et al., 1999), which may affect the recording of temperature in the different proxy carriers (Saher et al., 2009). This problem is only important for Mg/Ca ratios in foraminifera at temperatures below ~3°C (Elderfield et al., 2006) and at pH values lower than present (Russell et al., 2004). In core NIOP 929 however, planktic foraminifera calcified at SST's ~values are well above 3°C (Rostek et al., 1997) and pCO₂ values of the Vostok ice core indicate pH changes during the Quaternary having were likely in a range not significantly affecting Mg/Ca temperature estimates been above present day values (Russell et al., 2004) (Lea, 1999 #19797). It is therefore likely that the Mg/Ca records of core NIOP 929 have not been significantly affected by changes in carbonate chemistry through the carbonate ion effect (Saher et al., 2009). Another explanation is that as alkenone Uk'37 and Mg/Ca values are recorded by different organisms, differences in the seasonal or depth habitats between the two proxy carriers result in different SST estimates (Saher et al., 2009). The modern Arabian Sea alkenone producing algae reflect annual average conditions (Prah et al., 2000). Based on sediment trap data, the planktonic foraminifera *G. ruber* occurs in the modern Arabian Sea in roughly equal proportions during the winter and summer season (Conan and Brummer, 2000; Curry et al., 1992). This supports the notion that Mg/Ca- data based on sediments

(which contain a mix of *G. ruber* from both seasons) reflect annual average conditions too. Differences between Mg/Ca and Uk'37 -based SST estimates may entail shifts of the seasonal or depth preference of *G. ruber* and/or the alkenone producing species (Saher et al., 2009). An additional factor which may explain the higher alkenone Uk'37 temperature estimates in the 43 - 52 ka BP time window could be the preferential dissolution of high-Mg calcite in *G. ruber*, which would result in lower temperature estimates (Branson et al., 2013; Brown and Elderfield, 1996; Russell et al., 1994; Sadekov et al., 2005; Saher et al., 2009). Indicators of carbonate dissolution such as average shell weight, fragmentation, proportion of test loss during cleaning and CaCO₃ content in the bulk sediment (Tachikawa et al., 2008) are not available for the 37.4 ka BP to 67.7 ka BP section of core NIOP 929. ~~however, so the potential effects of carbonate dissolution cannot be quantified at this time. Planktic foraminifera in the region, however, are generally well preserved at the depth of core NIOP 929 (Conan et al., 2002) supporting the notion that carbonate dissolution has not significantly affected our Mg/Ca data. The difference in resolution between the alkenone Uk'37 and Mg/Ca temperature records in the 37.4 ka BP to 67.7 ka BP section could also potentially be masking the variability present in the alkenone Uk'37 record if it were reconstructed at a higher resolution (Fig. 5).~~ ~~Also,~~ despite the small areas of disagreement, the overall agreement between the two proxies across the studied period adds confidence to the *G. ruber* Mg/Ca SST record of this study.

The seasonal temperature reconstruction enables insights into the seasonal SST evolution in the western Arabian Sea. The outlined method does however assume a constant seasonal distribution of *G. ruber* and *G. bulloides* through time, which may not have been the case (Saher et al., 2007b). The seasonal abundance of *G. bulloides* reflects food availability by being present almost exclusively during the summer upwelling period (Conan and Brummer, 2000; Curry et al., 1992). In the past, however, during glacial periods with a weakened summer monsoon and strengthened winter northeast monsoon (Anderson and Prell, 1993; Prell and Campo, 1986; Saher et al., 2007b), highest food availability may have shifted to the winter monsoon season. In addition, colder winter monsoons may have induced vertical mixing bringing nutrients to the surface (Reichart et al., 1998). The latter model has been proposed for the eastern Arabian Sea, but the change involved in such a scenario for the western Arabian Sea is currently unclear. The seasonal difference between the winter and summer monsoon SSTs of this study (Fig. 43, 67) indicates strong summer monsoon winds generally persisting during the study period. This supports the notion that the glacial seasonal food distribution may have been broadly similar to the modern one, with the summer monsoon season driving the abundance of *G. bulloides*. Even if summer monsoon induced nutrient/food supply had dropped, during more extreme events, increased winter monsoon induced nutrient/food supply may not have occurred. This suggestion is based on results from the wider Arabian Sea showing that during the extreme Heinrich Events both, SW and NE monsoons were weakened (Singh et al., 2011). This supports the notion that NE monsoon winds could

not have driven nutrient/food supply in the western Arabian Sea even during extreme events such as Heinrich Events. This conclusion is in line with (Ganssen et al., 2011) in relation to HE's and on- in line with similar conclusions on glacial-interglacial time scales with (Jung et al., 2002). Hence, it is unlikely that, during periods with lower temperatures prevailing during summer, the proportion of *G. bulloides* present during the winter northeast monsoon was substantially different than today. This view is indeed in line with combined sediment core and model data from the northern Indian Ocean implying that the flux of *G. bulloides* is largely controlled by wind forcing (Bassinot et al., 2011), occurring during summer at the location of core NIOP 929.

Changes in the seasonal abundance of *G. ruber* are more difficult to assess. In the modern Arabian Sea close to site NIOP 929, *G. ruber* occurs during both seasons in roughly equal numbers (Conan and Brummer, 2000). This entails that *G. ruber* is rather insensitive to changes in both, temperature, and food/nutrient availability, implying that it is able to tolerate changes in both parameters without affecting its abundance. The data in Fig. 43 imply that even the lowest and highest SST estimates of 18 °C and 29-30 °C in our data broadly resemble the SST range in the modern Arabian Sea of 16°C (Swallow and Bruce, 1966) to 28.5 °C (Conan, 2006). These data are also in line with estimates of temperature ranges recorded over the last ~20 ka in core NIOP 905 (Ganssen et al., 2011), close to the location of core NIOP 929. These findings, in turn, suggest that *G. ruber* abundances may have been broadly stable, although additional research is required to assess the strength of this argument.

9.2.4.2. Long-term monsoon variability in the Arabian Sea in relation to periods I to IV

A large seasonal SST difference between summer and winter temperatures stems from low (upwelling driven) summer SSTs and high winter SSTs. In Figure 34, SSTs during winter are generally higher than during summer. The average difference in Fig. 43 for periods II and III, is around 8.5 °C. In periods I and IV the average seasonal SST differences are smaller. Broadly, this seasonal gradient is similar to the range in temperatures in the modern-day Arabian Sea (Swallow and Bruce, 1966). It is also comparable to the seasonal gradient of 5°C reported for the topmost section in core NIOP 929 (Saher et al., 2007b). This supports the notion of a generally strong SW monsoon and a relatively weak NE monsoon prevailing between 37 and 67 ka BP, similar to the modern setup.

The definition of periods I-IV rests on absences/occurrences of periods with a reversed seasonal temperature gradient, with these reversals being short-term in nature. We note that within the uncertainty of the age model, there is no obvious relationship to known short-term events such as Heinrich Events (HE's; positioning of HE's in reference to the NGRIP record follows (Bond et al., 1999)). The most pronounced temperature reversals at ~66.5 - 67 ka BP, ~65 ka BP, ~41.5 ka BP (with the caveat of this event being defined by one data point) and ~38 ka BP

Formatted: Font: Italic

occur before or after HE's. Also, there does not seem to be an obvious relationship with cold/warm oscillations in Greenland or Antarctic Ice cores (Fig. 43). This finding indicates that pronounced changes in seasonal monsoon strength occurred without an obvious phase-locked relationship with the millennial-scale change patterns prevailing elsewhere. We will explore this relationship further below.

The short maxima in summer SSTs and minima in winter SSTs in core NIOP 929 imply a seasonal change in monsoon strength. Enhanced summer SSTs point to a reduction in SW monsoon airflow. Reduced winter SSTs indicate a change during the NE monsoon season. It is difficult to pinpoint the specific cause of these cooling events. They could indicate cooler airmasses during the winter season affecting the Arabian Sea or alternatively, a stronger NE monsoon may have induced deeper mixing of the surface ocean allowing entrainment of cooler subsurface water into the mixed layer to be recorded in the planktic foraminifera record. A combination of both options is possible too (compare discussion in (Saher et al., 2007b)).

Seasonal variation in monsoon strength has been reported in previous studies (Anderson and Prell, 1993; Prell and Campo, 1986), albeit with a focus on orbitally driven monsoonal change. Supporting findings at the millennial scale have been described for the last glacial period at 13 - 20 ka BP in core NIOP 929 (Saher et al., 2007a), showing rapid changes in both summer and winter SSTs. Also, (Anand et al., 2008) reports *G. bulloides* and *G. ruber* based SSTs from western Arabian Sea core NIOP 905 covering the last 35ka. Interestingly, there is a difference between *G. bulloides* and *G. ruber* based SSTs between cores NIOP 905 (Anand et al., 2008) and NIOP 929 (this study). Core NIOP 905 does not seem to indicate a reversal of the temperature gradient between the species, but rather periods with a reduced gradient (Anand et al., 2008) in line with (Ganssen et al., 2011). This finding contrasts the data from core NIOP 929. Core NIOP 905 is located off the coast of Somalia, within the "Great Whirl", i.e. the central regional upwelling area. In the modern Arabian Sea core NIOP 929 is affected by upwelling too, but to a lesser extent. Summer upwelling indicative species *G. bulloides* occurred in core NIOP 905 over the last 35 ka BP even during Heinrich events (Jung et al., 2002), entailing coverage of full glacial conditions as well as superimposed short-term extremes, including a very strong Heinrich Event 1. The Heinrich Event-Dansgaard-Oeschger variability during the parts of MIS3 and MIS4 covered in this study probably does not comprise more extreme climate states than have occurred over the last 35 ka BP. This indicates that the SW monsoon winds varied in strength but persisted off the coast of Somalia throughout the study interval of this study. Accordingly, weakened, but persistent SW monsoon winds would have suppressed a larger change in the seasonal SST gradient at site NIOP 905. At the location of core NIOP 929, during periods with a temperature reversal, the suppression effect, i.e. maintaining some SW monsoon induced upwelling, would be smaller, possibly along with a reduction of the size of the Great Whirl. This in-turn supports

the notion of summer upwelling being more susceptible to disruptions, entailing larger SST changes at site NIOP 929, in line with the large swings in temperature in our data.

370 Similar to core NIOP 929, the time series for core SK17 from the eastern Arabian Sea (Anand et al., 2008) show short term reversals in the *G. bulloides* and *G. ruber* based SSTs over the last 35 ka BP. Off the west coast of India, periods of food/nutrient enhancement in surface waters, which *G. bulloides* responds to with high abundances (sensu, Curry et al., 1992), may occur during both winter and summer. A small amount of localised upwelling occurs during summer with similar conditions sporadically developing during winter (Cullen and Prell, 1984). In either
375 season, the upwelling is weak compared to the strong upwelling in the western Arabian Sea off Somalia. Without persistent strong upwelling it is probable that short-term disruptions of the monsoon circulation during MIS3/4, reflected in the temperature reversals in core NIOP 929, affected the eastern Arabian Sea in a similar way. Rather than involving a subtle change, monsoon circulation was more substantially affected, leading to reversed seasonal temperature gradients in the eastern Arabian Sea (sensu (Anand et al., 2008)). Principal support for the notion of
380 large changes in monsoonal circulation during MIS3 comes from the west Pacific warm pool (Stott et al., 2002), north China (Wang et al., 2021) and the Japan Sea (Sagawa et al., 2023), suggesting that the wider Asian monsoon circulation displayed significant variability during MIS 3/4.

9.3.4.3. Phasing of seasonal millennial scale SST variations in the Arabian Sea and beyond

One of the most striking findings of our study is the presence of short-term positive and negative temperature
385 excursions in core NIOP 929 (Fig. 43). Whilst other proxy records from the Arabian Sea (Altabet et al., 1995; Ivanochko et al., 2005; Schulz et al., 1998) revealed a systematic relationship with glacial northern hemisphere millennial climate change, our SST records do not display such a phase-locked persistent relationship. During HE6 for example in Fig. 43 the *G. bulloides* and *G. ruber* based SSTs as well the summer SST record indicate cooler conditions. During HE4 in contrast, the *G. ruber* based SST record indicates a warming event. Both, the *G. bulloides*
390 based SST and the summer temperature time series, however, show a short-term SST reduction embedded in an overall temperature rise during HE4. Comparable observations from the Arabian Sea are in (Anand et al., 2008) and to some degree in (Saher et al., 2007b) supporting the notion that temperature changes in the region are not phase locked and consistent with millennial scale climate oscillations in the northern hemisphere. Although there are a range of possible explanations involving local/regional changes in water properties (see discussions in (Anand et al.,
395 2008; Saher et al., 2007b)), the data from core NIOP 929 may offer an alternative view. This notion is based on the rough similarity between all summer monsoon affected SST records (*G. bulloides* and *G. ruber* based SSTs and the

derived summer SST's) with short-term changes in the Antarctic ice core $\delta^{18}\text{O}$ record in Fig. 43. It is not a perfect correlation, but within the uncertainty of the age models it does indicate that some of the temperature changes in core NIOB 929 do reflect a link with southern hemisphere climate change. Assessing the seasonal large-scale meteorological change driving monsoon circulation in the Arabian Sea, allows taking this notion further. It could be argued that the seasonal shift in the location of the Inter Tropical Convergence Zone (ITCZ) entails a switch in hemispheric "climate dominance" in the Arabian Sea. The ITCZ, the thermal equator, could be regarded as the dividing line/area, with northern hemisphere influence dominating north of it and southern hemispheric influence governing to the south. Within each meteorological hemisphere, polar climate changes could be transmitted relatively easily equatorward, with the ITCZ constraining further propagation. In view of the concept of the bipolar seesaw, i.e. phase-offset cold warm oscillations between the poles during MIS2-5 (Epica, 2006), this might shed light on the rather complex relationship of millennial scale temperature changes in the Arabian Sea with related change elsewhere. During northern hemisphere summer, the ITCZ reaches its northernmost position, i.e. being located over the Asian continent. During this season, southern hemisphere climate conditions would affect the Arabian Sea. Hence, millennial scale oscillations recorded over Antarctica (Epica, 2006) and references therein) could have affected monsoon circulation in the Arabian Sea chronicled as SSTs changes broadly following the southern hemisphere signal. During northern hemisphere winter, the ITCZ resides south of the equator, allowing northern hemisphere climate variations driving the monsoonal record in the Arabian Sea. This scenario entails that the Arabian Sea potentially records both northern and southern hemisphere millennial-scale change patterns. A non-straightforward and not phase-locked relation of the millennial scale SST records from the Arabian Sea with similar change elsewhere would result from this, as indicated in Fig. 43. Such a scenario would also explain why the winter SST record displays the least similarity with the southern hemisphere and its long-term change resembles insolation change at 65°N , suggesting a close relation with the north. This scenario is in line with a growing body of research challenging the paradigm of a solely northern hemispheric control on monsoon variability on millennial to orbital time scales (An et al., 2011; Caley et al., 2013; Rohling et al., 2009). In particular the comparison of the Hulu Cave data from continental China and ice core data from Antarctica (Rohling et al., 2009) and sediment data from the Arabian Sea (Caley et al., 2013) promote the idea of a stronger southern hemisphere monsoon control in relation to millennial scale change ~~to~~ during glacial periods. Our data add to this idea by supporting the notion that, periodically, the southern hemispheric influence is mainly exerted during northern hemispheric summer, similar to earlier suggestions (Wang et al., 2013). (~~Veres et al., 2013; Laskar et al., 2011~~)

10.5. Conclusions

The temperature reconstruction of core NIOP 929 using Mg/Ca paleothermometry of *G. ruber* and *G. bulloides* shows a range of 9°C in both records and rapid temperature changes from 37 - 67 ka BP. The novel approach of using the Mg/Ca SST records of *G. ruber* and *G. bulloides* to reconstruct winter and summer monsoon SSTs allows assessment of monsoonal controls at a seasonal scale. The results indicate that the summer southwest monsoon was strong for most of the period 37 - 67 ka BP but was significantly weakened multiple times, reversing the normal seasonal temperature gradient in the region. Also, the SST records do not show the expected phase-locked millennial-scale relationship with change in Greenland ice cores, nor do they show such a relationship with southern hemisphere millennial-scale climate change. The supports the notion of the Arabian Sea being a melting pot of climate controls, with both northern and southern hemisphere influence being recorded in sediments from the region. Further research is required, using a broader range of climate proxies such as productivity records, to verify the main implications of our study.

6. Competing Interests

S. Jung is co-editor of the special issue dedicated to Dick Kroon and will not be involved in the handling of this manuscript. The other authors declare that they have no conflict of interest.

11.7. Acknowledgements

We would like to thank Dr Laetitia Pichevin at the School of Geosciences of the University of Edinburgh for helping with the analytical side of the project. We are grateful for numerous final year students for helping to establish the stable isotope records for core NIOP 929. We are also grateful for support of this project by colleagues at Heriot Watt University - Dr Babette Hoogakker and Dr Katrina Nilsson-Kerr and to the members of the Heriot Watt Science Discussions group as well as to Dr Kate Darling for her insightful thoughts on foraminifera ecology.

Formatted: Heading 1

11.1.8. References

- 450 Altabet, M. A., Francois, R., Murray, D. W., and Prell, W. L.: Climate-related variations in denitrification in the Arabian Sea from sediment $15\text{N}/14\text{N}$ ratios, *Nature*, 373, 10.1038/373506a0, 1995.
- An, Z., Clemens, S. C., Shen, J., Qiang, X., Jin, Z., Sun, Y., Prell, W. L., Luo, J., Wang, S., Xu, H., Cai, Y., Zhou, W., Liu, X., Liu, W., Shi, Z., Yan, L., Xiao, X., Chang, H., Wu, F., Ai, L., and Lu, F.: Glacial-interglacial Indian summer monsoon dynamics, *Science*, 333, 10.1126/science.1203752, 2011.
- 455 Anand, P., Elderfield, H., and Conte, M. H.: Calibration of Mg/Ca thermometry in planktonic foraminifera from a sediment trap time series, *Paleoceanography*, 10.1029/2002PA000846, 2003.
- Anand, P., Kroon, D., Singh, A. D., Ganeshram, R. S., Ganssen, G., and Elderfield, H.: Coupled sea surface temperature-seawater $\delta^{18}\text{O}$ reconstructions in the Arabian Sea at the millennial scale for the last 35 ka, *Paleoceanography*, 10.1029/2007PA001564, 2008.
- 460 Andersen, K. K., Svensson, A., Johnsen, S. J., Rasmussen, S. O., Bigler, M., Röthlisberger, R., Ruth, U., Siggaard-Andersen, M. L., Peder Steffensen, J., Dahl-Jensen, D., Vinther, B. M., and Clausen, H. B.: The Greenland Ice Core Chronology 2005, 15-42 ka. Part 1: constructing the time scale, *Quaternary Science Reviews*, 25, 10.1016/j.quascirev.2006.08.002, 2006.
- Anderson, D. M. and Prell, W. L.: A 300 KYR Record of Upwelling Off Oman during the Late Quaternary: Evidence of the Asian Southwest Monsoon, *Paleoceanography*, 8, 10.1029/93PA00256, 1993.
- 465 Barker, S., Greaves, M., and Elderfield, H.: A study of cleaning procedures used for foraminiferal Mg/Ca paleothermometry, *Geochem. Geophys. Geosyst.*, 10.1029/2003GC000559, 2003.
- Bassinot, F. C., Marzin, C., Braconnot, P., Marti, O., Mathien-Blard, E., Lombard, F., and Bopp, L.: Holocene evolution of summer winds and marine productivity in the tropical Indian Ocean in response to insolation forcing: data-model comparison, *Climate of the past*, 7, 815-829, 10.5194/cp-7-815-2011, 2011.
- 470 Beaufort, L., Lancelot, Y., Camberlin, P., Cayre, O., Vincent, E., Bassinot, F., and Labeyrie, L.: Insolation cycles as a major control of equatorial Indian Ocean primary production, *Science*, 278, 10.1126/science.278.5342.1451, 1997.
- Bijma, J., Altabet, M., Conte, M., Kinkel, H., Versteegh, G. J. M., Volkman, J. K., Wakeham, S. G., and Weaver, P. P.: Primary signal: Ecological and environmental factors-Report from Working Group 2, *Geochem. Geophys. Geosyst.*, 2, 10.1029/2000gc000051, 2001.
- 475 Bond, G. C., Showers, W., Elliot, M., Evans, M., Lotti, R., Hajdas, I., Bonani, G., and Johnson, S.: The North Atlantic's 1-2 kyr climate rhythm; relation to Heinrich events, Dansgaard/Oeschger cycles and the Little Ice Age, in:

Formatted: Heading 1

- Mechanisms of global climate change at millennial time scales., edited by: Clark, P. U., Webb, R. S., and Keigwin, L. D., Geophysical Monograph, American Geophysical Union, Washington, DC, United States, 35-58, 1999.
- 480 Branson, O., Redfern, S. A. T., Tyliszczak, T., Sadekov, A., Langer, G., Kimoto, K., and Elderfield, H.: The coordination of Mg in foraminiferal calcite, *Earth and Planetary Science Letters*, 383, 10.1016/j.epsl.2013.09.037, 2013.
- Brock, J. C., McClain, C. R., Anderson, D. M., Prell, W. L., and Hay, W. W.: Southwest Monsoon Circulation and Environments of Recent Planktonic Foraminifera in the Northwestern Arabian Sea, *Paleoceanography*, 7, 485 10.1029/92PA01267, 1992.
- Brown, S. J. and Elderfield, H.: Variations in Mg/Ca and Sr/Ca ratios of planktonic foraminifera caused by postdepositional dissolution: Evidence of shallow Mg-dependent dissolution, *Paleoceanography*, 11, 10.1029/96PA01491, 1996.
- Budziak, D., Schneider, R. R., Rostek, F., Müller, P. J., Bard, E., and Wefer, G.: Late Quaternary insolation forcing 490 on total organic carbon and C37 alkenone variations in the Arabian Sea, *Paleoceanography*, 15, 10.1029/1999PA000433, 2000.
- Caley, T., Zaragosi, S., Bourget, J., Martinez, P., Malaizé, B., Eynaud, F., Rossignol, L., Garlan, T., and Ellouz-Zimmermann, N.: Southern Hemisphere imprint for Indo-Asian summer monsoons during the last glacial period as revealed by Arabian Sea productivity records, *Biogeosciences*, 10.5194/bg-10-7347-2013, 2013.
- 495 Clemens, S., Prell, W., Murray, D., Shimmield, G., and Weedon, G.: Forcing mechanisms of the Indian Ocean monsoon, *Nature*, 10.1038/353720a0, 1991.
- Conan, S. H. and Brummer, G. J. A.: Fluxes of planktic foraminifera in response to monsoonal upwelling on the Somalia Basin margin, *Deep-Sea Research Part II: Topical Studies in Oceanography*, 10.1016/S0967-0645(00)00022-9, 2000.
- 500 Conan, S. M. H.: Calibration of planktic foraminifera proxies for paleoproductivity and seasonal variability in the western Arabian Sea, PhD-Thesis - Research and graduation internal, 2006.
- Conan, S. M. H., Ivanova, E., and Brummer, G.-J. A.: Quantifying carbonate dissolution and calibration of foraminiferal dissolution indices in the Somali Basin, *Mar. Geol.*, 182, 325-349, 2002.
- Cullen, J. L. and Prell, W. L.: Planktonic foraminifera of the northern Indian Ocean; distribution and preservation in 505 surface sediments, *Mar. Micropaleontol.*, 9, 1-52, 1984.
- Curry, W. B., Ostermann, D. R., Gupta, M. V. S., and Ittekkot, V.: Foraminiferal production and monsoonal upwelling in the Arabian Sea: Evidence from sediment traps, *Geological Society Special Publication*, 64, 10.1144/GSL.SP.1992.064.01.06, 1992.

de Villiers, S., Greaves, M., and Elderfield, H.: An intensity ratio calibration method for the accurate determination
510 of Mg/Ca and Sr/Ca of marine carbonates by ICP-AES, *Geochem. Geophys. Geosyst.*, 10.1029/2001gc000169, 2002.

Elderfield, H. and Ganssen, G.: Past temperature and $\delta^{18}\text{O}$ of surface ocean waters inferred from foraminiferal Mg/Ca ratios, *Nature*, 405, 10.1038/35013033, 2000.

Elderfield, H., Yu, J., Anand, P., Kiefer, T., and Nyland, B.: Calibrations for benthic foraminiferal Mg/Ca
515 paleothermometry and the carbonate ion hypothesis, *Earth and Planetary Science Letters*, 250, 10.1016/j.epsl.2006.07.041, 2006.

EPICA: One-to-One coupling of glacial climate variability in Greenland and Antarctica, *Nature*, 444, 195-198, 2006.

Gadgil, S. and Gadgil, S.: The Indian monsoon, GDP and agriculture, *Economic & Political Weekly*, 2006.

Ganssen, G. M., Peeters, F. J. C., Metcalfe, B., Anand, P., Jung, S. J. A., Kroon, D., and Brummer, G. J. A.:
520 Quantifying sea surface temperature ranges of the Arabian Sea for the past 20 000 years, *Climate of the Past*, 10.5194/cp-7-1337-2011, 2011.

Gupta, A. K., Anderson, D. M., and Overpeck, J. T.: Abrupt changes in the Asian southwest monsoon during the holocene and their links to the North Atlantic Ocean, *Nature*, 421, 10.1038/nature01340, 2003.

Ivanochko, T. S., Ganeshram, R. S., Brummer, G. J. A., Ganssen, G., Jung, S. J. A., Moreton, S. G., and Kroon, D.:
525 Variations in tropical convection as an amplifier of global climate change at the millennial scale, *Earth and Planetary Science Letters*, 235, 10.1016/j.epsl.2005.04.002, 2005.

Jung, S. J. A., Ivanova, E., Reichert, G. J., Davies, G. R., Ganssen, G., Kroon, D., and van Hinte, J. E.: Centennial-millennial-scale monsoon variations off Somalia over the last 35 ka, *Geological Society Special Publication*, 10.1144/GSL.SP.2002.195.01.18, 2002.

Kroon, D.: Distribution of extant planktic foraminiferal assemblages in Red Sea and Northern Indian Ocean surface
530 waters, *Revista Espanola de Micropaleontologia*, 23, 37-74, 1991.

[Laskar, J., Fienga, A., Gastineau, M. & Manche, H. La2010: a new orbital solution for the long-term motion of the Earth. *Astronomy and astrophysics \(Berlin\)* 532. A89 \(2011\).](#)

Lea, D. W., Bijma, J., Spero, H. J., and Archer, D.: Implications of a Carbonate Ion Effect on Shell Carbon and
535 Oxygen Isotopes for Glacial Ocean Conditions, in: *Use of Proxies in Paleoclimatology*, 1999.

Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records, *Paleoclimatology*, 20, 10.1029/2004PA001071, 10.1029/2004PA001071, 2005.

Mashiota, T. A., Lea, D. W., and Spero, H. J.: Glacial-interglacial changes in Subantarctic sea surface temperature and $\delta^{18}\text{O}$ -water using foraminiferal Mg, *Earth and Planetary Science Letters*, 170, 10.1016/S0012-821X(99)00116-8, 1999.

Mishra, A. K., Dwivedi, S., and Das, S.: Role of Arabian Sea warming on the Indian summer monsoon rainfall in a regional climate model, *International Journal of Climatology*, 40, 10.1002/joc.6328, 2020.

Müller, P. J., Kirst, G., Ruhland, G., Von Storch, I., and Rosell-Melé, A.: Calibration of the alkenone paleotemperature index U37K based on core-tops from the eastern South Atlantic and the global ocean (60°N-60°S), *Geochim. Cosmochim. Acta*, 62, 1757-1772, 10.1016/S0016-7037(98)00097-0, 1998.

Nair, R. R., Ittekkot, V., Manganini, S. J., Ramaswamy, V., Haake, B., Degens, E. T., Desai, B. N., and Honjo, S.: Increased particle flux to the deep ocean related to monsoons, *Nature*, 338, 10.1038/338749a0, 1989.

Nürnberg, D., Müller, A., and Schneider, R. R.: Paleo-sea surface temperature calculations in the equatorial east Atlantic from Mg/Ca ratios in planktic foraminifera: A comparison to sea surface temperature estimates from U37K', oxygen isotopes, and foraminiferal transfer function, *Paleoceanography*, 15, 10.1029/1999PA000370, 2000.

Peeters, F. J. C., Brummer, G. J. A., and Ganssen, G.: The effect of upwelling on the distribution and stable isotope composition of *Globigerina bulloides* and *Globigerinoides ruber* (planktic foraminifera) in modern surface waters of the NW Arabian Sea, *Global Planet. Change*, 10.1016/S0921-8181(02)00120-0, 2002.

Prahl, F. G., Dymond, J., and Sparrow, M. A.: Annual biomarker record for export production in the central Arabian Sea, *Deep-Sea Research Part II: Topical Studies in Oceanography*, 47, 10.1016/S0967-0645(99)00155-1, 2000.

Prell, W. L. and Campo, E. V.: Coherent response of Arabian Sea upwelling and pollen transport to late Quaternary monsoonal winds, *Nature*, 323, 10.1038/323526a0, 1986.

Prell, W. L. and Curry, W. B.: Faunal and isotopic indices of monsoonal upwelling: Western Arabian Sea, *Oceanol. Acta*, 4, 1981.

Prell, W. L., Murray, D. W., Clemens, S. C., and Anderson, D. M.: Evolution and Variability of the Indian Ocean Summer Monsoon: Evidence from the Western Arabian Sea Drilling Program, in, 1992.

Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M., Clausen, H. B., Siggaard-Andersen, M. L., Johnsen, S. J., Larsen, L. B., Dahl-Jensen, D., Bigler, M., Röthlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M. E., and Ruth, U.: A new Greenland ice core chronology for the last glacial termination, *Journal of Geophysical Research Atmospheres*, 111, 10.1029/2005JD006079, 2006.

Reichart, G. J., Lourens, L. J., and Zachariasse, W. J.: Temporal variability in the northern Arabian Sea oxygen minimum zone (OMZ) during the last 225,000 years, *Paleoceanography*, 13, 10.1029/98PA02203, 1998.

Rixen, T., Haake, B., Ittekkot, V., Guptha, M. V. S., Nair, R. R., and Schlüssel, P.: Coupling between SW monsoon-related surface and deep ocean processes as discerned from continuous particle flux measurements and correlated satellite data, *Journal of Geophysical Research C: Oceans*, 101, 10.1029/96JC02420, 1996.

Rohling, E. J., Liu, Q. S., Roberts, A. P., Stanford, J. D., Rasmussen, S. O., Langen, P. L., and Siddall, M.: Controls on the East Asian monsoon during the last glacial cycle, based on comparison between Hulu Cave and polar ice-core records, *Quaternary Science Reviews*, 10.1016/j.quascirev.2009.09.007, 2009.

Rostek, F., Bard, E., Beaufort, L., Sonzogni, C., and Ganssen, G.: Sea surface temperature and productivity records for the last 240 kyr on the Arabian Sea, in: *Deep-Sea Research Part II: Topical Studies in Oceanography*, *Deep-Sea Research Part II: Topical Studies in Oceanography*, 1997, 10.1016/S0967-0645(97)00008-8, 1997.

Russell, A. D., Hönisch, B., Spero, H. J., and Lea, D. W.: Effects of seawater carbonate ion concentration and temperature on shell U, Mg, and Sr in cultured planktonic foraminifera, *Geochim. Cosmochim. Acta*, 68, 10.1016/j.gca.2004.03.013, 2004.

Russell, A. D., Emerson, S., Nelson, B. K., Erez, J., and Lea, D. W.: Uranium in foraminiferal calcite as a recorder of seawater uranium concentrations, *Geochim. Cosmochim. Acta*, 58, 10.1016/0016-7037(94)90497-9, 1994.

Sadekov, A. Y., Eggins, S. M., and De Deckker, P.: Characterization of Mg/Ca distributions in planktonic foraminifera species by electron microprobe mapping, *Geochem. Geophys. Geosyst.*, 6, 10.1029/2005GC000973, 2005.

Sagawa, T., Hasegawa, T., Narita, Y., Yokoyama, M., Kubota, Y., Okazaki, Y., Goto, A. S., Suzuki, Y., Ikehara, K., and Nakagawa, T.: Millennial-scale paleotemperature change in the Japan Sea during Marine Isotope Stage 3: Impact of meridional oscillation of the subpolar front, *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 626, 111713, 10.1016/j.palaeo.2023.111713, 2023.

Saher, M. H., Peeters, F. J. C., and Kroon, D.: Sea surface temperatures during the SW and NE monsoon seasons in the western Arabian Sea over the past 20,000 years, *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 10.1016/j.palaeo.2007.01.014, 2007a.

Saher, M. H., Jung, S. J. A., Elderfield, H., Greaves, M. J., and Kroon, D.: Sea surface temperatures of the western Arabian Sea during the last deglaciation, *Paleoceanography*, 10.1029/2006PA001292, 2007b.

Saher, M. H., Rostek, F., Jung, S. J. A., Bard, E., Schneider, R. R., Greaves, M., Ganssen, G. M., Elderfield, H., and Kroon, D.: Western Arabian Sea SST during the penultimate interglacial: A comparison of U δ^{37} and Mg/Ca paleothermometry, *Paleoceanography*, 10.1029/2007PA001557, 2009.

Schott, F. A. and McCreary, J. P.: The monsoon circulation of the Indian Ocean, *Prog. Oceanogr.*, 51, 1-123, 10.1016/s0079-6611(01)00083-0, 2001.

Schulz, H., Von Rad, U., and Erlenkeuser, H.: Correlation between Arabian Sea and Greenland climate oscillations of the past 110,000 years, *Nature*, 393, 10.1038/31750, 1998.

Shetye, S. R., Gouveia, A. D., and Shenoi, S. S. C.: Circulation and water masses of the Arabian Sea, *Proceedings of the Indian Academy of Sciences - Earth and Planetary Sciences*, 103, 10.1007/BF02839532, 1994.

Singh, A. D., Jung, S. J. A., Darling, K., Ganeshram, R., Ivanochko, T., and Kroon, D.: Productivity collapses in the Arabian Sea during glacial cold phases, *Paleoceanography*, 26, 10.1029/2009PA001923, 2011.

Sonzogni, C., Bard, E., Rostek, F., Lafont, R., Rosell-Mele, A., and Eglinton, G.: Core-top calibration of the alkenone index vs sea surface temperature in the Indian Ocean, in: *Deep-Sea Research Part II: Topical Studies in Oceanography*, *Deep-Sea Research Part II: Topical Studies in Oceanography*, 1997/01/01, 1445-1460, 10.1016/S0967-0645(97)00010-6, 1997.

Southon, J., Kashgarian, M., Fontugne, M., Metivier, B., and Yim, W. W. S.: Marine Reservoir Corrections for the Indian Ocean and Southeast Asia, *Radiocarbon*, 44, 167-180, 10.1017/S0033822200064778, 2002.

Stott, L., Poulsen, C. J., Lund, S., and Thunell, R.: Super ENSO and global climate oscillations at millennial time scales, *Science (American Association for the Advancement of Science)*, 297, 222-226, 10.1126/science.1071627, 2002.

Stuiver, M. and Reimer, P. J.: Extended 14C data base and revised CALIB 3.0 14C age calibration program, *Radiocarbon*, 35, 10.1017/S0033822200013904, 1993.

Svensson, A., Andersen, K. K., Bigler, M., Clausen, H. B., Dahl-Jensen, D., Davies, S. M., Johnsen, S. J., Muscheler, R., Rasmussen, S. O., Röthlisberger, R., Peder Steffensen, J., and Vinther, B. M.: The Greenland Ice Core Chronology 2005, 15-42 ka. Part 2: comparison to other records, *Quaternary Science Reviews*, 25, 10.1016/j.quascirev.2006.08.003, 2006.

Swallow, J. C. and Bruce, J. G.: Current measurements off the Somali coast during the southwest monsoon of 1964, *Deep-Sea Research and Oceanographic Abstracts*, 13, 10.1016/0011-7471(76)90908-6, 1966.

Tachikawa, K., Sépulcre, S., Toyofuku, T., and Bard, E.: Assessing influence of diagenetic carbonate dissolution on planktonic foraminiferal Mg/Ca in the southeastern Arabian Sea over the past 450 ka: Comparison between *Globigerinoides ruber* and *Globigerinoides sacculifer*, *Geochem. Geophys. Geosyst.*, 9, 10.1029/2007GC001904, 2008.

van Hinte, J. E., van Weering, T. C. E., and Troelstra, S. R.: Tracing a seasonal upwelling, report, 146, 1995.

Vázquez Riveiros, N., Govin, A., Waelbroeck, C., Mackensen, A., Michel, E., Moreira, S., Bouinot, T., Caillon, N., Orgun, A., and Brandon, M.: Mg/Ca thermometry in planktic foraminifera: Improving paleotemperature estimations

for *G. bulloides* and *N. pachyderma* left, *Geochem. Geophys. Geosyst.*, 17, 1249-1264, 10.1002/2015GC006234, 2016.

Veres, D. et al. The Antarctic ice core chronology (AICC2012): an optimized multi-parameter and multi-site dating approach for the last 120 thousand years. *Clim. Past* 9, 1733-1748 (2013).

Vinther, B. M., Clausen, H. B., Johnsen, S. J., Rasmussen, S. O., Andersen, K. K., Buchardt, S. L., Dahl-Jensen, D., Seierstad, I. K., Siggaard-Andersen, M. L., Steffensen, J. P., Svensson, A., Olsen, J., and Heinemeier, J.: A synchronized dating of three Greenland ice cores throughout the Holocene, *Journal of Geophysical Research Atmospheres*, 111, 10.1029/2005JD006921, 2006.

Wang, X., Sun, J., Longstaffe, F. J., Gu, X., Du, S., Cui, L., Huang, X., and Ding, Z.: Climatic quantification and seasonality of the late MIS 3 in north China; a perspective from carbon and oxygen isotopes of fossil mammal teeth, *Quaternary science reviews*, 272, 107222, 10.1016/j.quascirev.2021.107222, 2021.

Wang, Y. V., Leduc, G., Regenberg, M., Andersen, N., Larsen, T., Blanz, T., and Schneider, R. R.: Northern and southern hemisphere controls on seasonal sea surface temperatures in the Indian Ocean during the last deglaciation, *Paleoceanography*, 28, 619-632, <https://doi.org/10.1002/palo.20053>, 2013.

Webster, P. J., Magaña, V. O., Palmer, T. N., Shukla, J., Tomas, R. A., Yanai, M., and Yasunari, T.: Monsoons: Processes, predictability, and the prospects for prediction, *Journal of Geophysical Research: Oceans*, 10.1029/97JC02719, 1998.

Wolff, E. W., Chappellaz, J., Blunier, T., Rasmussen, S. O., and Svensson, A.: Millennial-scale variability during the last glacial: The ice core record, *Quaternary Science Reviews*, 29, 10.1016/j.quascirev.2009.10.013, 2010.

Wyrski, K.: Physical Oceanography of the Indian Ocean, in, 1973.

Zeeden, C., Obreht, I., Veres, D., Kaboth-Bahr, S., Hošek, J., Marković, S. B., Böskén, J., Lehmkuhl, F., Rolf, C., and Hambach, U.: Smoothed millennial-scale palaeoclimatic reference data as unconventional comparison targets: Application to European loess records, *Sci Rep*, 10, 5455, 10.1038/s41598-020-61528-8, 2020.

Altabet, M. A., Francois, R., Murray, D. W., and Prell, W. L.: Climate related variations in denitrification in the Arabian Sea from sediment $\delta^{15}\text{N}/\delta^{14}\text{N}$ ratios. *Nature*, 373, 10.1038/373506a0, 1995.

An, Z., Clemens, S. C., Shen, J., Qiang, X., Jin, Z., Sun, Y., Prell, W. L., Luo, J., Wang, S., Xu, H., Cai, Y., Zhou, W., Liu, X., Liu, W., Shi, Z., Yan, L., Xiao, X., Chang, H., Wu, F., Ai, L., and Lu, F.: Glacial-interglacial Indian summer monsoon dynamics. *Science*, 333, 10.1126/science.1203752, 2011.

Anand, P., Elderfield, H., and Conte, M. H.: Calibration of Mg/Ca thermometry in planktonic foraminifera from a sediment trap time series. *Paleoceanography*, 10.1029/2002PA000846, 2003.

- 660 Anand, P., Kroon, D., Singh, A. D., Ganeshram, R. S., Ganssen, G., and Elderfield, H.: Coupled sea surface temperature-seawater $\delta^{18}\text{O}$ reconstructions in the Arabian Sea at the millennial scale for the last 35 ka, *Paleoceanography*, 10.1029/2007PA001564, 2008.
- Andersen, K. K., Svensson, A., Johnsen, S. J., Rasmussen, S. O., Bigler, M., Röthlisberger, R., Ruth, U., Siggaard-Andersen, M. L., Peder Steffensen, J., Dahl-Jensen, D., Vinther, B. M., and Clausen, H. B.: The Greenland Ice Core Chronology 2005, 15-42 ka. Part 1: constructing the time scale, *Quaternary Science Reviews*, 25, 10.1016/j.quascirev.2006.08.002, 2006.
- 665 Anderson, D. M. and Prell, W. L.: A 300-KYR Record of Upwelling Off Oman during the Late Quaternary: Evidence of the Asian Southwest Monsoon, *Paleoceanography*, 8, 10.1029/93PA00256, 1993.
- Barker, S., Greaves, M., and Elderfield, H.: A study of cleaning procedures used for foraminiferal Mg/Ca paleothermometry, *Geochem. Geophys. Geosyst.*, 10.1029/2003GC000559, 2003.
- 670 Bassinot, F. C., Marzin, C., Braconnot, P., Marti, O., Mathien-Blard, E., Lombard, F., and Bopp, L.: Holocene evolution of summer winds and marine productivity in the tropical Indian Ocean in response to insolation forcing: data-model comparison, *Climate of the past*, 7, 815-829, 10.5194/cp-7-815-2011, 2011.
- Beaufort, L., Lancelot, Y., Camberlin, P., Cayre, O., Vincent, E., Bassinot, F., and Labeyrie, L.: Insolation cycles as a major control of equatorial Indian Ocean primary production, *Science*, 278, 10.1126/science.278.5342.1451, 1997.
- 675 Bijma, J., Altabet, M., Conte, M., Kinkel, H., Versteegh, G. J. M., Volkman, J. K., Wakeham, S. G., and Weaver, P. P.: Primary signal: Ecological and environmental factors-Report from Working Group 2, *Geochem. Geophys. Geosyst.*, 2, 10.1029/2000gc000051, 2001.
- Bond, G. C., Showers, W., Elliot, M., Evans, M., Lotti, R., Hajdas, I., Bonani, G., and Johnson, S.: The North Atlantic's 1-2 kyr climate rhythm: relation to Heinrich events, Dansgaard/Oeschger cycles and the Little Ice Age, in: *Mechanisms of global climate change at millennial time scales.*, edited by: Clark, P. U., Webb, R. S., and Keigwin, L. D., Geophysical Monograph, American Geophysical Union, Washington, DC, United States, 35-58, 1999.
- 680 Branson, O., Redfern, S. A. T., Tylliszczak, T., Sadekov, A., Langer, G., Kimoto, K., and Elderfield, H.: The co-ordination of Mg in foraminiferal calcite, *Earth and Planetary Science Letters*, 383, 10.1016/j.epsl.2013.09.037, 2013.
- 685 Brock, J. C., McClain, C. R., Anderson, D. M., Prell, W. L., and Hay, W. W.: Southwest Monsoon Circulation and Environments of Recent Planktonic Foraminifera in the Northwestern Arabian Sea, *Paleoceanography*, 7, 10.1029/92PA01267, 1992.

Brown, S. J. and Elderfield, H.: Variations in Mg/Ca and Sr/Ca ratios of planktonic foraminifera caused by
 postdepositional dissolution: Evidence of shallow Mg dependent dissolution, *Paleoceanography*, 11,
 10.1029/96PA01491, 1996.
 Budziak, D., Schneider, R. R., Rostek, F., Müller, P. J., Bard, E., and Wefer, G.: Late Quaternary insolation forcing
 on total organic carbon and C37 alkenone variations in the Arabian Sea, *Paleoceanography*, 15,
 10.1029/1999PA000433, 2000.
 Caley, T., Zaragosi, S., Bourget, J., Martinez, P., Malaizé, B., Eynaud, F., Rossignol, L., Garlan, T., and Ellouz-
 Zimmermann, N.: Southern Hemisphere imprint for Indo-Asian summer monsoons during the last glacial period as
 revealed by Arabian Sea productivity records, *Biogeosciences*, 10.5194/bg-10-7347-2013, 2013.
 Clemens, S., Prell, W., Murray, D., Shimmield, G., and Weedon, G.: Forcing mechanisms of the Indian Ocean
 monsoon, *Nature*, 10.1038/353720a0, 1991.
 Conan, S. H. and Brummer, G. J. A.: Fluxes of planktic foraminifera in response to monsoonal upwelling on the
 Somalia Basin margin, *Deep Sea Research Part II: Topical Studies in Oceanography*, 10.1016/S0967-
 0645(00)00022-9, 2000.
 Conan, S. M. H.: Calibration of planktic foraminifera proxies for paleoproductivity and seasonal variability in the
 western Arabian Sea, PhD-Thesis - Research and graduation internal, 2006.
 Conan, S. M. H., Ivanova, E., and Brummer, G. J. A.: Quantifying carbonate dissolution and calibration of
 foraminiferal dissolution indices in the Somali Basin, *Mar. Geol.*, 182, 325-349, 2002.
 Cullen, J. L. and Prell, W. L.: Planktonic foraminifera of the northern Indian Ocean: distribution and preservation in
 surface sediments, *Mar. Micropaleontol.*, 9, 1-52, 1984.
 Curry, W. B., Ostermann, D. R., Guptha, M. V. S., and Ittekkot, V.: Foraminiferal production and monsoonal
 upwelling in the Arabian Sea: Evidence from sediment traps, *Geological Society Special Publication*, 64,
 10.1144/GSL.SP.1992.064.01.06, 1992.
 de Villiers, S., Greaves, M., and Elderfield, H.: An intensity ratio calibration method for the accurate determination
 of Mg/Ca and Sr/Ca of marine carbonates by ICP-AES, *Geochem. Geophys. Geosyst.*, 10.1029/2001gc000169,
 2002.
 Elderfield, H. and Ganssen, G.: Past temperature and $\delta^{18}\text{O}$ of surface ocean waters inferred from foraminiferal
 Mg/Ca ratios, *Nature*, 405, 10.1038/35013033, 2000.
 Elderfield, H., Yu, J., Anand, P., Kiefer, T., and Nyland, B.: Calibrations for benthic foraminiferal Mg/Ca
 paleothermometry and the carbonate ion hypothesis, *Earth and Planetary Science Letters*, 250,
 10.1016/j.epsl.2006.07.041, 2006.

- EPICA: One-to-One coupling of glacial climate variability in Greenland and Antarctica, *Nature*, 444, 195–198, 2006.
- 720 Gadgil, S. and Gadgil, S.: The Indian monsoon, GDP and agriculture, *Economic & Political Weekly*, 2006.
- Ganssen, G. M., Peeters, F. J. C., Metcalfe, B., Anand, P., Jung, S. J. A., Kroon, D., and Brummer, G. J. A.: Quantifying sea surface temperature ranges of the Arabian Sea for the past 20 000 years, *Climate of the Past*, 10.5194/cp-7-1337-2011, 2011.
- Gupta, A. K., Anderson, D. M., and Overpeck, J. T.: Abrupt changes in the Asian southwest monsoon during the holocene and their links to the North Atlantic Ocean, *Nature*, 421, 10.1038/nature01340, 2003.
- 725 Ivanoenko, T. S., Ganeshram, R. S., Brummer, G. J. A., Ganssen, G., Jung, S. J. A., Moreton, S. G., and Kroon, D.: Variations in tropical convection as an amplifier of global climate change at the millennial scale, *Earth and Planetary Science Letters*, 235, 10.1016/j.epsl.2005.04.002, 2005.
- Jung, S. J. A., Ivanova, E., Reichert, G. J., Davies, G. R., Ganssen, G., Kroon, D., and van Hinte, J. E.: Centennial-millennial-scale monsoon variations off Somalia over the last 35 ka, *Geological Society Special Publication*, 10.1144/GSL.SP.2002.195.01.18, 2002.
- 730 Kroon, D.: Distribution of extant planktic foraminiferal assemblages in Red Sea and Northern Indian Ocean surface waters, *Revista Espanola de Micropaleontologia*, 23, 37–74, 1991.
- Lea, D. W., Bijma, J., Spero, H. J., and Archer, D.: Implications of a Carbonate Ion Effect on Shell Carbon and Oxygen Isotopes for Glacial Ocean Conditions, in: *Use of Proxies in Paleooceanography*, 1999.
- 735 Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}O$ records, *Paleoceanography*, 20, 10.1029/2004PA001071, 10.1029/2004PA001071, 2005.
- Mashietta, T. A., Lea, D. W., and Spero, H. J.: Glacial-interglacial changes in Subantarctic sea surface temperature and $\delta^{18}O$ -water using foraminiferal Mg, *Earth and Planetary Science Letters*, 170, 10.1016/S0012-821X(99)00116-8, 1999.
- 740 Mishra, A. K., Dwivedi, S., and Das, S.: Role of Arabian Sea warming on the Indian summer monsoon rainfall in a regional climate model, *International Journal of Climatology*, 40, 10.1002/joc.6328, 2020.
- Müller, P. J., Kirst, G., Ruhland, G., Von Storch, I., and Rosell-Melé, A.: Calibration of the alkenone paleotemperature index U_{37K} based on core tops from the eastern South Atlantic and the global ocean (60°N–60°S), *Geochim. Cosmochim. Acta*, 62, 1757–1772, 10.1016/S0016-7037(98)00097-0, 1998.
- 745 Nair, R. R., Ittekkot, V., Manganini, S. J., Ramaswamy, V., Haake, B., Degens, E. T., Desai, B. N., and Honje, S.: Increased particle flux to the deep ocean related to monsoons, *Nature*, 338, 10.1038/338749a0, 1989.

- Nürnberg, D., Müller, A., and Schneider, R. R.: Paleo-sea surface temperature calculations in the equatorial east Atlantic from Mg/Ca ratios in planktic foraminifera: A comparison to sea surface temperature estimates from U37K' oxygen isotopes, and foraminiferal transfer function, *Paleoceanography*, 15, 10.1029/1999PA000370, 2000.
- Peeters, F. J. C., Brummer, G. J. A., and Ganssen, G.: The effect of upwelling on the distribution and stable isotope composition of *Globigerina bulloides* and *Globigerinoides ruber* (planktic foraminifera) in modern surface waters of the NW Arabian Sea, *Global Planet. Change*, 10.1016/S0921-8181(02)00120-0, 2002.
- Prahl, F. G., Dymond, J., and Sparrow, M. A.: Annual biomarker record for export production in the central Arabian Sea, *Deep-Sea Research Part II: Topical Studies in Oceanography*, 47, 10.1016/S0967-0645(99)00155-1, 2000.
- Prell, W. L. and Campo, E. V.: Coherent response of Arabian Sea upwelling and pollen transport to late Quaternary monsoonal winds, *Nature*, 323, 10.1038/323526a0, 1986.
- Prell, W. L. and Curry, W. B.: Faunal and isotopic indices of monsoonal upwelling: Western Arabian Sea, *Oceanol. Acta*, 4, 1981.
- Prell, W. L., Murray, D. W., Clemens, S. C., and Anderson, D. M.: Evolution and Variability of the Indian Ocean Summer Monsoon: Evidence from the Western Arabian Sea Drilling Program, in: 1992.
- Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M., Clausen, H. B., Siggaard-Andersen, M. L., Johnsen, S. J., Larsen, L. B., Dahl-Jensen, D., Bigler, M., Röthlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M. E., and Ruth, U.: A new Greenland ice core chronology for the last glacial termination, *Journal of Geophysical Research Atmospheres*, 111, 10.1029/2005JD006079, 2006.
- Reichart, G. J., Lourens, L. J., and Zachariasse, W. J.: Temporal variability in the northern Arabian Sea oxygen minimum zone (OMZ) during the last 225,000 years, *Paleoceanography*, 13, 10.1029/98PA02203, 1998.
- Rixen, T., Haake, B., Ittekkot, V., Guptha, M. V. S., Nair, R. R., and Schlüssel, P.: Coupling between SW monsoon-related surface and deep ocean processes as discerned from continuous particle flux measurements and correlated satellite data, *Journal of Geophysical Research C: Oceans*, 101, 10.1029/96JC02420, 1996.
- Rohling, E. J., Liu, Q. S., Roberts, A. P., Stanford, J. D., Rasmussen, S. O., Langen, P. L., and Siddall, M.: Controls on the East Asian monsoon during the last glacial cycle, based on comparison between Hulu Cave and polar ice-core records, *Quaternary Science Reviews*, 10.1016/j.quascirev.2009.09.007, 2009.
- Rostek, F., Bard, E., Beaufort, L., Sonzogni, C., and Ganssen, G.: Sea surface temperature and productivity records for the last 240 kyr on the Arabian Sea, in: *Deep-Sea Research Part II: Topical Studies in Oceanography, Deep-Sea Research Part II: Topical Studies in Oceanography*, 1997, 10.1016/S0967-0645(97)00008-8, 1997.

- Russell, A. D., Hönisch, B., Spero, H. J., and Lea, D. W.: Effects of seawater carbonate ion concentration and temperature on shell U, Mg, and Sr in cultured planktonic foraminifera, *Geochim. Cosmochim. Acta*, 68, 10.1016/j.gca.2004.03.013, 2004.
- 780 Russell, A. D., Emerson, S., Nelson, B. K., Erez, J., and Lea, D. W.: Uranium in foraminiferal calcite as a recorder of seawater uranium concentrations, *Geochim. Cosmochim. Acta*, 58, 10.1016/0016-7037(94)90497-9, 1994.
- Sadekov, A. Y., Eggins, S. M., and De Deckker, P.: Characterization of Mg/Ca distributions in planktonic foraminifera species by electron microprobe mapping, *Geochem. Geophys. Geosyst.*, 6, 10.1029/2005GC000973, 2005.
- 785 Sagawa, T., Hasegawa, T., Narita, Y., Yokoyama, M., Kubota, Y., Okazaki, Y., Goto, A. S., Suzuki, Y., Ikehara, K., and Nakagawa, T.: Millennial-scale paleotemperature change in the Japan Sea during Marine Isotope Stage 3: Impact of meridional oscillation of the subpolar front, *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 626, 111713, 10.1016/j.palaeo.2023.111713, 2023.
- Saher, M. H., Peeters, F. J. C., and Kroon, D.: Sea surface temperatures during the SW and NE monsoon seasons in the western Arabian Sea over the past 20,000 years, *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 10.1016/j.palaeo.2007.01.014, 2007a.
- 790 Saher, M. H., Jung, S. J. A., Elderfield, H., Greaves, M. J., and Kroon, D.: Sea surface temperatures of the western Arabian Sea during the last deglaciation, *Paleoceanography*, 10.1029/2006PA001292, 2007b.
- Saher, M. H., Rostek, F., Jung, S. J. A., Bard, E., Schneider, R. R., Greaves, M., Ganssen, G. M., Elderfield, H., and Kroon, D.: Western Arabian Sea SST during the penultimate interglacial: A comparison of U K37 and Mg/Ca paleothermometry, *Paleoceanography*, 10.1029/2007PA001557, 2009.
- 795 Schott, F. A. and McCreary, J. P.: The monsoon circulation of the Indian Ocean, *Prog. Oceanogr.*, 51, 1-123, 10.1016/s0079-6611(01)00083-0, 2001.
- Schulz, H., Von Rad, U., and Erlenkeuser, H.: Correlation between Arabian Sea and Greenland climate oscillations of the past 110,000 years, *Nature*, 393, 10.1038/31750, 1998.
- 800 Shetye, S. R., Gouveia, A. D., and Shenoi, S. S. C.: Circulation and water masses of the Arabian Sea, *Proceedings of the Indian Academy of Sciences – Earth and Planetary Sciences*, 103, 10.1007/BF02839532, 1994.
- Singh, A. D., Jung, S. J. A., Darling, K., Ganeshram, R., Ivanochko, T., and Kroon, D.: Productivity collapses in the Arabian Sea during glacial cold phases, *Paleoceanography*, 26, 10.1029/2009PA001923, 2011.
- 805 Sonzogni, C., Bard, E., Rostek, F., Lafont, R., Rosell-Mele, A., and Eglinton, G.: Core-top calibration of the alkenone index vs sea surface temperature in the Indian Ocean, in: *Deep-Sea Research Part II: Topical Studies in*

- Oceanography, Deep-Sea Research Part II: Topical Studies in Oceanography, 1997/01/01, 1445-1460, 10.1016/S0967-0645(97)00010-6, 1997.
- 810 Southon, J., Kashgarian, M., Fontugne, M., Metivier, B., and Yim, W. W. S.: Marine Reservoir Corrections for the Indian Ocean and Southeast Asia, Radiocarbon, 44, 167-180, 10.1017/S0033822200064778, 2002.
- Stott, L., Poulsen, C. J., Lund, S., and Thunell, R.: Super ENSO and global climate oscillations at millennial time scales, Science (American Association for the Advancement of Science), 297, 222-226, 10.1126/science.1071627, 2002.
- 815 Stuiver, M. and Reimer, P. J.: Extended 14C data base and revised CALIB 3.0 14C age calibration program, Radiocarbon, 25, 10.1017/S0033822200013904, 1993.
- Svensson, A., Andersen, K. K., Bigler, M., Clausen, H. B., Dahl-Jensen, D., Davies, S. M., Johnsen, S. J., Muscheler, R., Rasmussen, S. O., Röthlisberger, R., Peder Steffensen, J., and Vinther, B. M.: The Greenland Ice Core Chronology 2005, 15-42 ka. Part 2: comparison to other records, Quaternary Science Reviews, 25, 10.1016/j.quascirev.2006.08.003, 2006.
- 820 Swallow, J. C. and Bruce, J. G.: Current measurements off the Somali coast during the southwest monsoon of 1964, Deep-Sea Research and Oceanographic Abstracts, 13, 10.1016/0011-7471(76)90908-6, 1966.
- Tachikawa, K., Sépulchre, S., Toyofuku, T., and Bard, E.: Assessing influence of diagenetic carbonate dissolution on planktonic foraminiferal Mg/Ca in the southeastern Arabian Sea over the past 450 ka: Comparison between Globigerinoides ruber and Globigerinoides sacculifer, Geochem. Geophys. Geosyst., 9, 10.1029/2007GC001904, 2008.
- 825 van Hinte, J. E., van Weering, T. C. E., and Troelstra, S. R.: Tracing a seasonal upwelling, report, 146, 1995.
- Vázquez Riveiros, N., Govin, A., Waelbroeck, C., Mackensen, A., Michel, E., Moreira, S., Bouinot, T., Caillon, N., Orgun, A., and Brandon, M.: Mg/Ca thermometry in planktic foraminifera: Improving paleotemperature estimations for G. bulloides and N. pachyderma left, Geochem. Geophys. Geosyst., 17, 1249-1264, 10.1002/2015GC006234, 2016.
- 830 Vinther, B. M., Clausen, H. B., Johnsen, S. J., Rasmussen, S. O., Andersen, K. K., Buchardt, S. L., Dahl-Jensen, D., Seierstad, I. K., Siggaard Andersen, M. L., Steffensen, J. P., Svensson, A., Olsen, J., and Heinemeier, J.: A synchronized dating of three Greenland ice cores throughout the Holocene, Journal of Geophysical Research Atmospheres, 111, 10.1029/2005JD006921, 2006.
- 835 Wang, X., Sun, J., Longstaffe, F. J., Gu, X., Du, S., Cui, L., Huang, X., and Ding, Z.: Climatic quantification and seasonality of the late MIS 3 in north China: a perspective from carbon and oxygen isotopes of fossil mammal teeth, Quaternary science reviews, 272, 107222, 10.1016/j.quascirev.2021.107222, 2021.

Wang, Y. V., Leduc, G., Regenberg, M., Andersen, N., Larsen, T., Blanz, T., and Schneider, R. R.: Northern and southern hemisphere controls on seasonal sea surface temperatures in the Indian Ocean during the last deglaciation, *Paleoceanography*, 28, 619–632, <https://doi.org/10.1002/palo.20053>, 2013.

Webster, P. J., Magaña, V. O., Palmer, T. N., Shukla, J., Tomas, R. A., Yanai, M., and Yasunari, T.: Monsoons: Processes, predictability, and the prospects for prediction, *Journal of Geophysical Research: Oceans*, 10.1029/97JC02719, 1998.

Wolff, E. W., Chappellaz, J., Blunier, T., Rasmussen, S. O., and Svensson, A.: Millennial scale variability during the last glacial: The ice core record, *Quaternary Science Reviews*, 29, 10.1016/j.quascirev.2009.10.013, 2010.

Wyrtki, K.: *Physical Oceanography of the Indian Ocean*, in: 1972.

Zeeden, C., Obrecht, I., Veres, D., Kaboth-Bahr, S., Hošek, J., Marković, S. B., Böskén, J., Lehmkuhl, F., Rolf, C., and Hambach, U.: Smoothed millennial scale palaeoclimatic reference data as unconventional comparison targets: Application to European loess records, *Sci Rep*, 10, 5455, 10.1038/s41598-020-61528-8, 2020.

Field Code Changed

12.9. Figure captions and table headings

Table 1: Original and updated ages from Saher et al., (2007) 14C ages for core NIOP 929, recalibrated to calendar ages using the 14C age calibration program Calib 8.2 (Stuiver and Reimer, 1993), using a ΔR value of 45 (± 67) (Southon et al., 2002).

Table 2: Age control points based on tuning to the LR04 $\delta^{18}O$ global stack of Lisiecki and Raymo (2005).

Tabel 3: Combined age control points based on tuning to the LR04 $\delta^{18}O$ global stack of Lisiecki and Raymo (2005) and the Greenland ice core NGRIP $\delta^{18}O$ record (Andersen et al., 2006; Rasmussen et al., 2006; Svensson et al., 2006; Vinther et al., 2006; Wolff et al., 2010).

Figure 1: Map of the Arabian Sea region. The locations of core NIOP 905, NIOP 929 and MD76-131 are indicated. Green arrow schematically represents summer monsoon winds. Grey arrow schematically represents winter monsoon winds.

Figure 2: Seasonal foraminiferal flux dynamics in the western Arabian Sea for *G. bulloides* and *G. ruber*. (a) Typical summer conditions and (b) typical winter conditions. (c) Winter and summer temperature estimates derived from Mg/Ca temperature estimates from *G. bulloides* and *G. ruber* and seasonal foraminiferal flux dynamics.

Figure 23: Oxygen isotopes for *G. ruber* (red) and *G. bulloides* (blue) compared to the NGRIP ice core data. Overview of the age model development of core NIOP 929. The records shown are (top down) the LR04 $\delta^{18}O$ global stack of Lisiecki and Raymo (2005), the Greenland ice core NGRIP $\delta^{18}O$ record (Andersen et al., 2006; Rasmussen et al.,

Formatted: Superscript

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

2006; Svensson et al., 2006; Vinther et al., 2006; Wolff et al., 2010), the $\delta^{18}\text{O}$ - record *G. ruber* of core NIOP 929 (700 year boxcar filtered record in bold, original record at reduced opacity) and the depth-age record for core NIOP 929. Pairs of triangles and vertical lines indicate origin of age control points; red = AMS¹⁴C, green = LR04 and blue = NGRIP. For uncertainties see text.

Formatted: Font: Italic

Formatted: Superscript

Figure 34: Summary figure for core NIOP 929 showing (top down): the Antarctic EPICA $\delta^{18}\text{O}$ record (Veres et al., 2013), summer temperature estimates (this work), winter temperature estimates (this work), difference between winter and summer SST estimates (winter minus summer; grey line depicts temperature equivalence, negative values indicate temperature reversals), SST estimates based on Mg/Ca data from *G. bulloides* (this work), SST estimates based on Mg/Ca data from *G. ruber* (this work), stable isotope values for *G. bulloides* and *G. ruber* and *G. bulloides* (this work), and the Greenland ice core NGRIP $\delta^{18}\text{O}$ record (Andersen et al., 2006; Rasmussen et al., 2006; Svensson et al., 2006; Vinther et al., 2006; Wolff et al., 2010), the respective Mg/Ca-based SSTs estimates, the derived winter and summer SST time series and the seasonal temperature difference (winter minus summer; grey line depicts temperature equivalence, negative values indicate temperature reversals). Grey bars indicate position of Heinrich Events (HE's). A 700 year box car filter has been applied to all SST timeseries. Periods I-IV, discussed in the text, are indicated. Positioning of HE's in reference to NGRIP follows Bond et al. (1999). Insolation data taken from Laskar et al. (2011).

Formatted: Font: Italic


Formatted: Font: Italic

Figure 45: Mg/Ca based temperature reconstructions for *G. ruber* (red) and *G. bulloides* (blue) from core NIOP 929. Dashed lines in the temperature records indicate the mean temperature values for each record across the reconstructed period.

Figure 56: Temperature history of core NIOP 929 to 126 ka BP including Mg/Ca temperatures, alkenone temperatures and stable oxygen isotope values. All records are placed on the age model created in this study. *G. ruber* Mg/Ca temperature records of this study from 37.4-67.7 ka BP, Saher et al., (2007) from 2.9-20 ka BP and Saher et al., (2009) from 95.5-124 ka BP in red. *G. bulloides* Mg/Ca temperature record of this study from 37.4-67.7 ka BP in blue. Alkenone temperature records of Rostek et al., (1997) in dark orange and Saher et al., (2009) in light orange. *G. ruber* $\delta^{18}\text{O}$ record of this study in red and *G. bulloides* $\delta^{18}\text{O}$ record of this study in blue. NGRIP $\delta^{18}\text{O}$ record in black.

Figure 67: Winter and summer temperature estimates from core NIOP 929. From top down, NGRIP $\delta^{18}\text{O}$ record (black), winter (green) and summer (yellow) temperature reconstructions from core NIOP 929 and the difference between winter and summer temperatures (black). Dashed lines in the temperature records indicate the mean temperature values for each record across the reconstructed period.


900




905




910



915



920



925

930 Altabet, M. A., Francois, R., Murray, D. W., and Prell, W. L.: Climate-related
 variations in denitrification in the Arabian Sea from sediment $^{15}\text{N}/^{14}\text{N}$ ratios, *Nature*,
~~273, 10.1028/273506a0, 1995.~~
 An, Z., Clemens, S. C., Shen, J., Qiang, X., Jin, Z., Sun, Y., Prell, W. L., Luo, J.,
 Wang, S., Xu, H., Cai, Y., Zhou, W., Liu, X., Liu, W., Shi, Z., Yan, L., Xiao, X.,
 935 Chang, H., Wu, F., Ai, L., and Lu, F.: Glacial-interglacial Indian summer monsoon
 dynamics, *Science*, 333, 10.1126/science.1203752, 2011.
 Anand, P., Elderfield, H., and Conte, M. H.: Calibration of Mg/Ca thermometry in
 planktonic foraminifera from a sediment trap time series, *Paleoceanography*,
~~10.1029/2002PA000846, 2003.~~
 940 Anand, P., Kroon, D., Singh, A. D., Ganeshram, R. S., Ganssen, G., and Elderfield,
 H.: Coupled sea surface temperature-seawater $\delta^{18}\text{O}$ reconstructions in the Arabian Sea
 at the millennial scale for the last 35 ka, *Paleoceanography*, 10.1029/2007PA001564,
~~2008.~~
 Andersen, K. K., Svensson, A., Johnsen, S. J., Rasmussen, S. O., Bigler, M.,
 Röthlisberger, R., Ruth, U., Siggaard-Andersen, M. L., Peder Steffensen, J., Dahl-
 945 Jensen, D., Vinther, B. M., and Clausen, H. B.: The Greenland Ice Core Chronology
 2005, 15–42 ka. Part 1: constructing the time scale, *Quaternary Science Reviews*, 25,
~~10.1016/j.quascirev.2006.08.002, 2006.~~
 Anderson, D. M. and Prell, W. L.: A 300 KYR Record of Upwelling Off Oman during the
 Late Quaternary: Evidence of the Asian Southwest Monsoon, *Paleoceanography*, 8,
 950 ~~10.1029/93PA00256, 1993.~~
 Barker, S., Greaves, M., and Elderfield, H.: A study of cleaning procedures used for
 foraminiferal Mg/Ca paleothermometry, *Geochem. Geophys. Geosyst.*,
~~10.1029/2003GC000559, 2003.~~
 Bassinot, F. C., Marzin, C., Braconnot, P., Marti, O., Mathien-Blard, E., Lombard,
 955 F., and Bopp, L.: Holocene evolution of summer winds and marine productivity in the
 tropical Indian Ocean in response to insolation forcing: data model comparison,
Climate of the past, 7, 815–829, 10.5194/cp-7-815-2011, 2011.
 Beaufort, L., Lancelot, Y., Camberlin, P., Cayre, O., Vincent, E., Bassinot, F., and
 Labeyrie, L.: Insolation cycles as a major control of equatorial Indian Ocean primary
 960 production, *Science*, 278, 10.1126/science.278.5342.1451, 1997.
 Bijma, J., Altabet, M., Conte, M., Kinkel, H., Versteegh, G. J. M., Volkman, J. K.,
 Wakeham, S. G., and Weaver, P. P.: Primary signal: Ecological and environmental
 factors Report from Working Group 2, *Geochem. Geophys. Geosyst.*, 2,
~~10.1029/2000gc000051, 2001.~~
 965 Bond, G. C., Showers, W., Elliot, M., Evans, M., Lotti, R., Hajdas, I., Bonani, G.,
 and Johnson, S.: The North Atlantic's 1–2 kyr climate rhythm: relation to Heinrich
 events, Dansgaard/Oeschger cycles and the Little Ice Age, in: *Mechanisms of global
 climate change at millennial time scales.*, edited by: Clark, P. U., Webb, R. S., and
 Keigwin, L. D., *Geophysical Monograph*, American Geophysical Union, Washington, DC,
 970 United States, 35–58, 1999.

Branson, O., Redfern, S. A. T., Tylicszczak, T., Sadekov, A., Langer, G., Kimoto, K.,
 and Elderfield, H.: The coordination of Mg in foraminiferal calcite, *Earth and
 Planetary Science Letters*, 282, 10.1016/j.epsl.2013.09.037, 2013.
 Brock, J. C., McClain, C. R., Anderson, D. M., Prell, W. L., and Hay, W. W.: Southwest
 975 Monsoon Circulation and Environments of Recent Planktonic Foraminifera in the
 Northwestern Arabian Sea, *Paleoceanography*, 7, 10.1029/92PA01267, 1992.
 Brown, S. J. and Elderfield, H.: Variations in Mg/Ca and Sr/Ca ratios of planktonic
 foraminifera caused by postdepositional dissolution: Evidence of shallow Mg-dependent
 dissolution, *Paleoceanography*, 11, 10.1029/96PA01491, 1996.
 980 Budziak, D., Schneider, R. R., Rostek, F., Müller, P. J., Bard, E., and Wefer, G.:
 Late Quaternary insolation forcing on total organic carbon and C37 alkenone variations
 in the Arabian Sea, *Paleoceanography*, 15, 10.1029/1999PA000433, 2000.
 Galey, T., Zaragosi, S., Bourget, J., Martinez, P., Malaizé, B., Eynaud, F.,
 Rossignol, L., Garlan, T., and Ellouz-Zimmermann, N.: Southern Hemisphere imprint for
 985 Indo-Asian summer monsoons during the last glacial period as revealed by Arabian Sea
 productivity records, *Biogeosciences*, 10.5194/bg-10-7347-2013, 2013.
 Clemens, S., Prell, W., Murray, D., Shimmield, G., and Weedon, G.: Forcing mechanisms
 of the Indian Ocean monsoon, *Nature*, 10.1038/353720a0, 1991.
 Conan, S. H. and Brummer, G. J. A.: Fluxes of planktic foraminifera in response to
 990 monsoonal upwelling on the Somalia Basin margin, Deep-Sea Research Part II: Topical
 Studies in Oceanography, 10.1016/S0967-0645(00)00022-9, 2000.
 Conan, S. M. H.: Calibration of planktic foraminifera proxies for paleoproductivity
 and seasonal variability in the western Arabian Sea, PhD Thesis — Research and
 graduation internal, 2006.
 995 Conan, S. M. H., Ivanova, E., and Brummer, G. J. A.: Quantifying carbonate dissolution
 and calibration of foraminiferal dissolution indices in the Somali Basin, *Mar. Geol.*,
 182, 325–349, 2002.
 Cullen, J. L. and Prell, W. L.: Planktonic foraminifera of the northern Indian Ocean,
 distribution and preservation in surface sediments, *Mar. Micropaleontol.*, 9, 1–52,
 1000 1984.
 Curry, W. B., Ostermann, D. R., Guptha, M. V. S., and Ittekkot, V.: Foraminiferal
 production and monsoonal upwelling in the Arabian Sea: Evidence from sediment traps,
Geological Society Special Publication, 64, 10.1144/GSL.SP.1992.064.01.06, 1992.
 1005 de Villiers, S., Greaves, M., and Elderfield, H.: An intensity ratio calibration
 method for the accurate determination of Mg/Ca and Sr/Ca of marine carbonates by ICP-
 AES, *Geochem. Geophys. Geosyst.*, 10.1029/2001gc000169, 2002.
 Elderfield, H. and Gansssen, G.: Past temperature and $\delta^{18}\text{O}$ of surface ocean waters
 inferred from foraminiferal Mg/Ca ratios, *Nature*, 405, 10.1038/35013033, 2000.
 1010 Elderfield, H., Yu, J., Anand, P., Kiefer, T., and Nyland, B.: Calibrations for
 benthic foraminiferal Mg/Ca paleothermometry and the carbonate ion hypothesis, *Earth
 and Planetary Science Letters*, 250, 10.1016/j.epsl.2006.07.041, 2006.
 EPICA: One to One coupling of glacial climate variability in Greenland and Antarctica,
Nature, 444, 195–198, 2006.
 1015 Gadgil, S. and Gadgil, S.: The Indian monsoon, GDP and agriculture, *Economic &
 Political Weekly*, 2006.

Ganssen, G. M., Peeters, F. J. C., Metcalfe, B., Anand, P., Jung, S. J. A., Kroon, D., and Brummer, G. J. A.: Quantifying sea surface temperature ranges of the Arabian Sea for the past 20 000 years, *Climate of the Past*, 10.5194/cp 7 1237 2011, 2011.

1020 Gupta, A. K., Anderson, D. M., and Overpeck, J. T.: Abrupt changes in the Asian southwest monsoon during the holocene and their links to the North Atlantic Ocean, *Nature*, 421, 10.1038/nature01340, 2003.

1025 Ivanochko, T. S., Ganeshram, R. S., Brummer, G. J. A., Ganssen, G., Jung, S. J. A., Moreton, S. G., and Kroon, D.: Variations in tropical convection as an amplifier of global climate change at the millennial scale, *Earth and Planetary Science Letters*, 235, 10.1016/j.epsl.2005.04.002, 2005.

1030 Jung, S. J. A., Ivanova, E., Reichert, G. J., Davies, G. R., Ganssen, G., Kroon, D., and van Hinte, J. E.: Centennial millennial scale monsoon variations off Somalia over the last 35 ka, *Geological Society Special Publication*, 10.1144/GSL.SP.2002.195.01.18, 2002.

1035 Kroon, D.: Distribution of extant planktic foraminiferal assemblages in Red Sea and Northern Indian Ocean surface waters, *Revista Espanola de Micropaleontologia*, 23, 37-74, 1991.

1040 Lea, D. W., Bijma, J., Spero, H. J., and Archer, D.: Implications of a Carbonate Ion Effect on Shell Carbon and Oxygen Isotopes for Glacial Ocean Conditions, in: *Use of Proxies in Paleoceanography*, 1999.

1045 Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}O$ records, *Paleoceanography*, 20, 10.1029/2004PA001071, 10.1029/2004PA001071, 2005.

1050 Mashiotta, T. A., Lea, D. W., and Spero, H. J.: Glacial interglacial changes in Subantarctic sea surface temperature and $\delta^{18}O$ water using foraminiferal Mg, *Earth and Planetary Science Letters*, 170, 10.1016/S0012-821X(99)00116-8, 1999.

1055 Mishra, A. K., Dwivedi, S., and Das, S.: Role of Arabian Sea warming on the Indian summer monsoon rainfall in a regional climate model, *International Journal of Climatology*, 40, 10.1002/joc.6328, 2020.

1060 Müller, P. J., Kirst, G., Ruhland, G., Von Storch, I., and Rosell-Melé, A.: Calibration of the alkenone paleotemperature index U_{37K} based on core tops from the eastern South Atlantic and the global ocean (60°N-60°S), *Geochim. Cosmochim. Acta*, 62, 1757-1772, 10.1016/S0016-7037(98)00097-0, 1998.

Nair, R. R., Ittekkot, V., Manganini, S. J., Ramaswamy, V., Haake, B., Degens, E. T., Desai, B. N., and Honjo, S.: Increased particle flux to the deep ocean related to monsoons, *Nature*, 338, 10.1038/338749a0, 1989.

Nürnberg, D., Müller, A., and Schneider, R. R.: Paleo sea surface temperature calculations in the equatorial east Atlantic from Mg/Ca ratios in planktic foraminifera: A comparison to sea surface temperature estimates from U_{37K}, oxygen isotopes, and foraminiferal transfer function, *Paleoceanography*, 15, 10.1029/1999PA000370, 2000.

Peeters, F. J. C., Brummer, G. J. A., and Ganssen, G.: The effect of upwelling on the distribution and stable isotope composition of *Globigerina bulloides* and *Globigerinoides ruber* (planktic foraminifera) in modern surface waters of the NW Arabian Sea, *Global Planet. Change*, 10.1016/S0921-8181(02)00120-0, 2002.

Prahl, F. G., Dymond, J., and Sparrow, M. A.: Annual biomarker record for export production in the central Arabian Sea, *Deep-Sea Research Part II: Topical Studies in Oceanography*, 47, 10.1016/S0967-0645(99)00155-1, 2000.

1065 Prell, W. L. and Campo, E. V.: Coherent response of Arabian Sea upwelling and pollen transport to late Quaternary monsoonal winds, *Nature*, 323, 10.1038/323526a0, 1986.

Prell, W. L. and Curry, W. B.: Faunal and isotopic indices of monsoonal upwelling: Western Arabian Sea, *Oceanol. Acta*, 4, 1981.

1070 Prell, W. L., Murrpay, D. W., Clemens, S. C., and Anderson, D. M.: Evolution and Variability of the Indian Ocean Summer Monsoon: Evidence from the Western Arabian Sea Drilling Program, in, 1992.

Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M., Clausen, H. B., Siggaard Andersen, M. L., Johnsen, S. J., Larsen, L. B., Dahl-Jensen, D., Bigler, M., Röthlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M. E., and Ruth, U.: A new Greenland ice core chronology for the last glacial termination, *Journal of Geophysical Research Atmospheres*, 111, 10.1029/2005JD006079, 2006.

1075 Reichant, G. J., Lourens, L. J., and Zachariasse, W. J.: Temporal variability in the northern Arabian Sea oxygen minimum zone (OMZ) during the last 225,000 years, *Paleoceanography*, 13, 10.1029/98PA02203, 1998.

1080 Rixen, T., Haake, B., Ittekkot, V., Gupta, M. V. S., Nair, R. R., and Schlüssel, P.: Coupling between SW monsoon-related surface and deep ocean processes as discerned from continuous particle flux measurements and correlated satellite data, *Journal of Geophysical Research C: Oceans*, 101, 10.1029/96JC02420, 1996.

Rohling, E. J., Liu, Q. S., Roberts, A. P., Stanford, J. D., Rasmussen, S. O., Langen, P. L., and Siddall, M.: Controls on the East Asian monsoon during the last glacial cycle, based on comparison between Hulu Cave and polar ice core records, *Quaternary Science Reviews*, 10.1016/j.quascirev.2009.09.007, 2009.

1085 Rostek, F., Bard, E., Beaufort, L., Sonzogni, C., and Ganssen, G.: Sea surface temperature and productivity records for the last 240 kyr on the Arabian Sea, in: *Deep-Sea Research Part II: Topical Studies in Oceanography, Deep-Sea Research Part II: Topical Studies in Oceanography*, 1997, 10.1016/S0967-0645(97)00008-8, 1997.

1090 Russell, A. D., Hönisch, B., Spero, H. J., and Lea, D. W.: Effects of seawater carbonate ion concentration and temperature on shell U, Mg, and Sr in cultured planktonic foraminifera, *Geochim. Cosmochim. Acta*, 68, 10.1016/j.gca.2004.03.013, 2004.

1095 Russell, A. D., Emerson, S., Nelson, B. K., Erez, J., and Lea, D. W.: Uranium in foraminiferal calcite as a recorder of seawater uranium concentrations, *Geochim. Cosmochim. Acta*, 58, 10.1016/0016-7037(94)90497-9, 1994.

Sadekov, A. Y., Eggins, S. M., and De Deckker, P.: Characterization of Mg/Ca distributions in planktonic foraminifera species by electron microprobe mapping, *Geochim. Geophys. Geosyst.*, 6, 10.1029/2005GC000973, 2005.

1100 Sagawa, T., Hasegawa, T., Narita, Y., Yokoyama, M., Kubota, Y., Okazaki, Y., Goto, A. S., Suzuki, Y., Ikehara, K., and Nakagawa, T.: Millennial scale paleotemperature change in the Japan Sea during Marine Isotope Stage 3: Impact of meridional oscillation of the subpolar front, *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 626, 111713, 10.1016/j.palaeo.2023.111713, 2023.

1105

Saher, M. H., Peeters, F. J. C., and Kroon, D.: Sea surface temperatures during the SW and NE monsoon seasons in the western Arabian Sea over the past 20,000 years, *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 10.1016/j.palaeo.2007.01.014, 2007a.

Saher, M. H., Jung, S. J. A., Elderfield, H., Greaves, M. J., and Kroon, D.: Sea surface temperatures of the western Arabian Sea during the last deglaciation, *Paleoceanography*, 10.1029/2006PA001292, 2007b.

Saher, M. H., Rostek, F., Jung, S. J. A., Bard, E., Schneider, R. R., Greaves, M., Ganssen, G. M., Elderfield, H., and Kroon, D.: Western Arabian Sea SST during the penultimate interglacial: A comparison of U K37 and Mg/Ca paleothermometry, *Paleoceanography*, 10.1029/2007PA001557, 2009.

Schott, F. A. and McCreary, J. P.: The monsoon circulation of the Indian Ocean, *Prog. Oceanogr.*, 51, 1-123, 10.1016/S0079-6611(01)00083-0, 2001.

Schulz, H., Von Rad, U., and Erlenkeuser, H.: Correlation between Arabian Sea and Greenland climate oscillations of the past 110,000 years, *Nature*, 393, 10.1038/31750, 1998.

Shetye, S. R., Gouveia, A. D., and Shenoi, S. S. C.: Circulation and water masses of the Arabian Sea, *Proceedings of the Indian Academy of Sciences - Earth and Planetary Sciences*, 103, 10.1007/BF02839532, 1994.

Singh, A. D., Jung, S. J. A., Darling, K., Ganeshram, R., Ivanochko, T., and Kroon, D.: Productivity collapses in the Arabian Sea during glacial cold phases, *Paleoceanography*, 26, 10.1029/2009PA001923, 2011.

Sonzogni, C., Bard, E., Rostek, F., Lafont, R., Rosell-Mele, A., and Eglinton, G.: Core top calibration of the alkenone index vs sea surface temperature in the Indian Ocean, in: *Deep Sea Research Part II: Topical Studies in Oceanography*, Deep Sea Research Part II: Topical Studies in Oceanography, 1997/01/01, 1445-1460, 10.1016/S0967-0645(97)00010-6, 1997.

Southon, J., Kashgarian, M., Fontugne, M., Metivier, B., and Yim, W. W. S.: Marine Reservoir Corrections for the Indian Ocean and Southeast Asia, *Radiocarbon*, 44, 167-180, 10.1017/S0033822200064778, 2002.

Stott, L., Poulsen, C. J., Lund, S., and Thunell, R.: Super ENSO and global climate oscillations at millennial time scales, *Science (American Association for the Advancement of Science)*, 297, 222-226, 10.1126/science.1071627, 2002.

Stuiver, M. and Reimer, P. J.: Extended 14C data base and revised CALIB 3.0 14C age calibration program, *Radiocarbon*, 35, 10.1017/S0033822200013904, 1993.

Svensson, A., Andersen, K. K., Bigler, M., Clausen, H. B., Dahl-Jensen, D., Davies, S. M., Johnsen, S. J., Muscheler, R., Rasmussen, S. O., Röthlisberger, R., Peder Steffensen, J., and Vinther, B. M.: The Greenland Ice Core Chronology 2005, 15-42 ka. Part 2: comparison to other records, *Quaternary Science Reviews*, 25, 10.1016/j.quascirev.2006.08.003, 2006.

Swallow, J. C. and Bruce, J. C.: Current measurements off the Somali coast during the southwest monsoon of 1964, *Deep-Sea Research and Oceanographic Abstracts*, 13, 10.1016/0011-7471(76)90008-6, 1966.

Tachikawa, K., Sépulchre, S., Toyofuku, T., and Bard, E.: Assessing influence of diagenetic carbonate dissolution on planktonic foraminiferal Mg/Ca in the southeastern Arabian Sea over the past 450 ka: Comparison between Globigerinoides ruber and

- ~~Globigerinoides sacculifer, *Geochem. Geophys. Geosyst.*, 9, 10.1029/2007GC001904, 2008.~~
- ~~van Hinte, J. E., van Weering, T. C. E., and Troelstra, S. R.: Tracing a seasonal upwelling, report, 146, 1995.~~
- 1155 ~~Vázquez Riveiros, N., Govin, A., Waelbroeck, C., Mackensen, A., Michel, E., Moreira, S., Bouinot, T., Caillon, N., Orgun, A., and Brandon, M.: Mg/Ca thermometry in planktic foraminifera: Improving paleotemperature estimations for *G. bulloides* and *N. pachyderma* left, *Geochem. Geophys. Geosyst.*, 17, 1249–1264, 10.1002/2015GC006234, 2016.~~
- 1160 ~~Vinther, B. M., Clausen, H. B., Johnsen, S. J., Rasmussen, S. O., Andersen, K. K., Buchardt, S. L., Dahl-Jensen, D., Seierstad, I. K., Siggaard-Andersen, M. L., Steffensen, J. P., Svensson, A., Olsen, J., and Heinemeier, J.: A synchronized dating of three Greenland ice cores throughout the Holocene, *Journal of Geophysical Research: Atmospheres*, 111, 10.1029/2005JD006921, 2006.~~
- 1165 ~~Wang, X., Sun, J., Longstaffe, F. J., Gu, X., Du, S., Cui, L., Huang, X., and Ding, Z.: Climatic quantification and seasonality of the late MIS 3 in north China: a perspective from carbon and oxygen isotopes of fossil mammal teeth, *Quaternary science reviews*, 272, 107222, 10.1016/j.quascirev.2021.107222, 2021.~~
- 1170 ~~Wang, Y. V., Leduc, G., Regenberg, M., Larsen, N., Blanz, T., and Schneider, R. R.: Northern and southern hemisphere controls on seasonal sea surface temperatures in the Indian Ocean during the last deglaciation, *Paleoceanography*, 28, 619–632, <https://doi.org/10.1002/palo.20053>, 2013.~~
- ~~Webster, P. J., Magaña, V. O., Palmer, T. N., Shukla, J., Tomas, R. A., Yanai, M., and Yasunari, T.: Monsoons: Processes, predictability, and the prospects for prediction, *Journal of Geophysical Research: Oceans*, 10.1029/97JC02719, 1998.~~
- 1175 ~~Wolff, E. W., Chappellaz, J., Blunier, T., Rasmussen, S. O., and Svensson, A.: Millennial-scale variability during the last glacial: The ice core record, *Quaternary Science Reviews*, 29, 10.1016/j.quascirev.2009.10.013, 2010.~~
- ~~Wyrtki, K.: Physical Oceanography of the Indian Ocean, in, 1973.~~
- 1180 ~~Zeebe, C., Obrecht, I., Veres, D., Kaboth Bahr, S., Hošek, J., Marković, S. B., Böken, J., Lehmkuhl, F., Rolf, C., and Hambach, U.: Smoothed millennial scale palaeoclimatic reference data as unconventional comparison targets: Application to European loess records, *Sci Rep*, 10, 5455, 10.1038/s41598-020-61528-8, 2020.~~
- 1185
- ~~Altabet, M. A., Francois, R., Murray, D. W., and Prell, W. L.: Climate related variations in denitrification in the Arabian Sea from sediment 15N/14N ratios, *Nature*, 373, 10.1038/373506a0, 1995.~~
- 1190 ~~An, Z., Clemens, S. C., Shen, J., Qiang, X., Jin, Z., Sun, Y., Prell, W. L., Luo, J., Wang, S., Xu, H., Cai, Y., Zhou, W., Liu, X., Liu, W., Shi, Z., Yan, L., Xiao, X., Chang, H., Wu, F., Ai, L., and Lu, F.: Glacial interglacial Indian summer monsoon dynamics, *Science*, 333, 10.1126/science.1203752, 2011.~~

- 1195 Anand, P., Elderfield, H., and Conte, M. H.: Calibration of Mg/Ca thermometry in planktonic foraminifera from a sediment trap time series, *Paleoceanography*, ~~10.1029/2002PA000846, 2002.~~
- Anand, P., Kroon, D., Singh, A. D., Ganeshram, R. S., Ganssen, G., and Elderfield, H.: Coupled sea surface temperature seawater $\delta^{18}\text{O}$ reconstructions in the Arabian Sea at the millennial scale for the last 35 ka, *Paleoceanography*, ~~10.1029/2007PA001564, 2008.~~
- 1200 Andersen, K. K., Svensson, A., Johnsen, S. J., Rasmussen, S. O., Bigler, M., Röthlisberger, R., Ruth, U., Siggaard-Andersen, M. L., Peder Steffensen, J., Dahl-Jensen, D., Vinther, B. M., and Clausen, H. B.: The Greenland Ice Core Chronology 2005, 15–42 ka, Part 1: constructing the time scale, *Quaternary Science Reviews*, ~~25, 10.1016/j.quascirev.2006.08.002, 2006.~~
- 1205 Anderson, D. M. and Prell, W. L.: A 300 Kyr Record of Upwelling Off Oman during the Late Quaternary: Evidence of the Asian Southwest Monsoon, *Paleoceanography*, ~~8, 10.1029/93PA00256, 1993.~~
- Barker, S., Greaves, M., and Elderfield, H.: A study of cleaning procedures used for foraminiferal Mg/Ca paleothermometry, *Geochem. Geophys. Geosyst.*, ~~10.1029/2003GC000559, 2003.~~
- 1210 Bassinot, F. C., Marzin, C., Braconnot, P., Marti, O., Mathien-Blard, E., Lombard, F., and Bopp, L.: Holocene evolution of summer winds and marine productivity in the tropical Indian Ocean in response to insolation forcing: data model comparison, *Climate of the past*, ~~7, 815–829, 10.5194/cp-7-815-2011, 2011.~~
- 1215 Beaufort, L., Lancelot, Y., Camberlin, P., Cayre, O., Vincent, E., Bassinot, F., and Labeyrie, L.: Insolation cycles as a major control of equatorial Indian Ocean primary production, *Science*, ~~278, 10.1126/science.278.5342.1451, 1997.~~
- Bijma, J., Altabet, M., Conte, M., Kinkel, H., Versteegh, G. J. M., Volkman, J. K., Wakeham, S. G., and Weaver, P. P.: Primary signal: Ecological and environmental factors Report from Working Group 2, *Geochem. Geophys. Geosyst.*, ~~2, 10.1029/2000gc000051, 2001.~~
- 1220 Bond, G. C., Showers, W., Elliot, M., Evans, M., Lotti, R., Hajdas, I., Bonani, G., and Johnson, S.: The North Atlantic's 1–2 kyr climate rhythm: relation to Heinrich events, Dansgaard/Oeschger cycles and the Little Ice Age, in: *Mechanisms of global climate change at millennial time scales*, edited by: Clark, P. U., Webb, R. S., and Keigwin, L. D., *Geophysical Monograph*, American Geophysical Union, Washington, DC, United States, ~~35–58, 1999.~~
- 1225 Branson, O., Redfern, S. A. T., Tylliszczak, T., Sadekov, A., Langer, G., Kimoto, K., and Elderfield, H.: The coordination of Mg in foraminiferal calcite, *Earth and Planetary Science Letters*, ~~282, 10.1016/j.epsl.2013.09.027, 2013.~~
- 1230 Brock, J. C., McClain, C. R., Anderson, D. M., Prell, W. L., and Hay, W. W.: Southwest Monsoon Circulation and Environments of Recent Planktonic Foraminifera in the Northwestern Arabian Sea, *Paleoceanography*, ~~7, 10.1029/92PA01267, 1992.~~
- 1235 Brown, S. J. and Elderfield, H.: Variations in Mg/Ca and Sr/Ca ratios of planktonic foraminifera caused by postdepositional dissolution: Evidence of shallow Mg-dependent dissolution, *Paleoceanography*, ~~11, 10.1029/96PA01491, 1996.~~

~~Budziak, D., Schneider, R. R., Rostek, F., Müller, P. J., Bard, E., and Wefer, G.: Late Quaternary insolation forcing on total organic carbon and C37 alkenone variations in the Arabian Sea, *Paleoceanography*, 15, 10.1029/1999PA000433, 2000.~~
 1240 ~~Caley, T., Zaragosi, S., Bourget, J., Martinez, P., Malaizé, B., Eynaud, F., Ressignol, L., Garlan, T., and Elouez Zimmermann, N.: Southern Hemisphere imprint for Indo-Asian summer monsoons during the last glacial period as revealed by Arabian Sea productivity records, *Biogeosciences*, 10.5194/bg-10-7247-2013, 2013.~~
 1245 ~~Clemens, S., Prell, W., Murray, D., Shimmield, G., and Weedon, G.: Forcing mechanisms of the Indian Ocean monsoon, *Nature*, 10.1038/353720a0, 1991.~~
~~Conan, S. H. and Brummer, G. J. A.: Fluxes of planktic foraminifera in response to monsoonal upwelling on the Somalia Basin margin, Deep-Sea Research Part II: Topical Studies in Oceanography, 10.1016/S0067-0645(00)00022-9, 2000.~~
 1250 ~~Conan, S. M. H.: Calibration of planktic foraminifera proxies for paleoproductivity and seasonal variability in the western Arabian Sea, PhD Thesis — Research and graduation internal, 2006.~~
~~Conan, S. M. H., Ivanova, E., and Brummer, G. J. A.: Quantifying carbonate dissolution and calibration of foraminiferal dissolution indices in the Somali Basin, *Mar. Geol.*, 182, 325–349, 2002.~~
 1255 ~~Cullen, J. L. and Prell, W. L.: Planktonic foraminifera of the northern Indian Ocean: distribution and preservation in surface sediments, *Mar. Micropaleontol.*, 9, 1–52, 1984.~~
~~Curry, W. B., Ostermann, D. R., Guptha, M. V. S., and Ittekkot, V.: Foraminiferal production and monsoonal upwelling in the Arabian Sea: Evidence from sediment traps, *Geological Society Special Publication*, 64, 10.1144/GSL.SP.1992.064.01.06, 1992.~~
 1260 ~~de Villiers, S., Greaves, M., and Elderfield, H.: An intensity ratio calibration method for the accurate determination of Mg/Ca and Sr/Ca of marine carbonates by ICP-AES, *Geochem. Geophys. Geosyst.*, 10.1029/2001gc000169, 2002.~~
 1265 ~~Elderfield, H. and Ganssen, G.: Past temperature and δ18O of surface ocean waters inferred from foraminiferal Mg/Ca ratios, *Nature*, 405, 10.1038/35013033, 2000.~~
~~Elderfield, H., Yu, J., Anand, P., Kiefer, T., and Nyland, B.: Calibrations for benthic foraminiferal Mg/Ca paleothermometry and the carbonate ion hypothesis, *Earth and Planetary Science Letters*, 250, 10.1016/j.epsl.2006.07.041, 2006.~~
 1270 ~~EPICA: One to One coupling of glacial climate variability in Greenland and Antarctica, *Nature*, 444, 195–198, 2006.~~
~~Gadgil, S. and Gadgil, S.: The Indian monsoon, GDP and agriculture, *Economic & Political Weekly*, 2006.~~
~~Ganssen, G. M., Peeters, F. J. C., Metcalfe, B., Anand, P., Jung, S. J. A., Kroon, D., and Brummer, G. J. A.: Quantifying sea surface temperature ranges of the Arabian Sea for the past 20 000 years, *Climate of the Past*, 10.5194/cp-7-1337-2011, 2011.~~
 1275 ~~Gupta, A. K., Anderson, D. M., and Overpeck, J. T.: Abrupt changes in the Asian southwest monsoon during the holocene and their links to the North Atlantic Ocean, *Nature*, 421, 10.1038/nature01340, 2002.~~
 1280 ~~Ivanochko, T. S., Ganeshram, R. S., Brummer, G. J. A., Ganssen, G., Jung, S. J. A., Moreton, S. G., and Kroon, D.: Variations in tropical convection as an amplifier of~~

global climate change at the millennial scale, *Earth and Planetary Science Letters*, 235, 10.1016/j.epsl.2005.04.002, 2005.

1285 Jung, S. J. A., Ivanova, E., Reichert, G. J., Davies, G. R., Ganssen, G., Kroon, D., and van Hinte, J. E.: Centennial-millennial scale monsoon variations off Somalia over the last 25 ka, *Geological Society Special Publication*, 10.1144/GSL.SP.2002.195.01.18, 2002.

Kroon, D.: Distribution of extant planktic foraminiferal assemblages in Red Sea and Northern Indian Ocean surface waters, *Revista Espanola de Micropaleontologia*, 23, 37-74, 1991.

1290 Laskar, J., Fienga, A., Gastineau, M., and Manche, H.: La2010: a new orbital solution for the long term motion of the Earth, *Astronomy and astrophysics (Berlin)*, 532, A89, 10.1051/0004-6261/201116836, 2011.

Lea, D. W., Bijma, J., Spero, H. J., and Archer, D.: Implications of a Carbonate Ion Effect on Shell Carbon and Oxygen Isotopes for Glacial Ocean Conditions, in: *Use of Proxies in Paleoceanography*, 1999.

1295 Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}O$ records, *Paleoceanography*, 20, 10.1029/2004PA001071, 10.1029/2004PA001071, 2005.

Mashiotto, T. A., Lea, D. W., and Spero, H. J.: Glacial interglacial changes in Subantarctic sea surface temperature and $\delta^{18}O$ water using foraminiferal Mg, *Earth and Planetary Science Letters*, 170, 10.1016/S0012-821X(99)00116-8, 1999.

1300 Mishra, A. K., Dwivedi, S., and Das, S.: Role of Arabian Sea warming on the Indian summer monsoon rainfall in a regional climate model, *International Journal of Climatology*, 40, 10.1002/joc.6328, 2020.

1305 Müller, P. J., Kirst, G., Ruhland, G., Von Storch, I., and Rosell-Melé, A.: Calibration of the alkenone paleotemperature index U_{37K} based on core tops from the eastern South Atlantic and the global ocean (60°N-60°S), *Geochim. Cosmochim. Acta*, 62, 1757-1772, 10.1016/S0016-7037(98)00097-0, 1998.

Nair, R. R., Ittekkot, V., Manganini, S. J., Ramaswamy, V., Haake, B., Degens, E. T., 1310 Desai, B. N., and Honjo, S.: Increased particle flux to the deep ocean related to monsoons, *Nature*, 338, 10.1038/338749a0, 1989.

Nürnberg, D., Müller, A., and Schneider, R. R.: Paleo sea surface temperature calculations in the equatorial east Atlantic from Mg/Ca ratios in planktic foraminifera: A comparison to sea surface temperature estimates from U_{37K}, oxygen isotopes, and foraminiferal transfer function, *Paleoceanography*, 15, 10.1029/1999PA000370, 2000.

1315 Peeters, F. J. C., Brummer, G. J. A., and Ganssen, G.: The effect of upwelling on the distribution and stable isotope composition of *Globigerina bulloides* and *Globigerinoides ruber* (planktic foraminifera) in modern surface waters of the NW Arabian Sea, *Global Planet. Change*, 10.1016/S0921-8181(02)00120-0, 2002.

1320 Prah, F. G., Dymond, J., and Sparrow, M. A.: Annual biomarker record for export production in the central Arabian Sea, *Deep Sea Research Part II: Topical Studies in Oceanography*, 47, 10.1016/S0967-0645(99)00155-1, 2000.

1325 Prell, W. L. and Campo, E. V.: Coherent response of Arabian Sea upwelling and pollen transport to late Quaternary monsoonal winds, *Nature*, 323, 10.1038/323526a0, 1986.

- Prell, W. L. and Curry, W. B.: Faunal and isotopic indices of monsoonal upwelling: Western Arabian Sea, *Oceanol. Acta*, 4, 1981.
- Prell, W. L., Murray, D. W., Clemens, S. C., and Anderson, D. M.: Evolution and Variability of the Indian Ocean Summer Monsoon: Evidence from the Western Arabian Sea Drilling Program, in, 1992.
- 1330 Rasmussen, S. O., Andersen, K. K., Svensson, A. M., Steffensen, J. P., Vinther, B. M., Clausen, H. B., Siggaard Andersen, M. L., Johnsen, S. J., Larsen, L. B., Dahl-Jensen, D., Bigler, M., Röthlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M. E., and Ruth, U.: A new Greenland ice core chronology for the last glacial termination, *Journal of Geophysical Research Atmospheres*, 111, 10.1029/2005JD006079, 2006.
- 1335 Reichert, G. J., Lourens, L. J., and Zachariasse, W. J.: Temporal variability in the northern Arabian Sea oxygen minimum zone (OMZ) during the last 225,000 years, *Paleoceanography*, 13, 10.1029/98PA02203, 1998.
- 1340 Rixen, T., Haake, B., Ittekkot, V., Gupta, M. V. S., Nair, R. R., and Schlüssel, P.: Coupling between SW monsoon-related surface and deep ocean processes as discerned from continuous particle flux measurements and correlated satellite data, *Journal of Geophysical Research C: Oceans*, 101, 10.1029/96JC02420, 1996.
- Rohling, E. J., Liu, Q. S., Roberts, A. P., Stanford, J. D., Rasmussen, S. O., Langen, P. L., and Siddall, M.: Controls on the East Asian monsoon during the last glacial cycle, based on comparison between Hulu Cave and polar ice-core records, *Quaternary Science Reviews*, 10.1016/j.quascirev.2009.09.007, 2009.
- 1345 Rostek, F., Bard, E., Beaufort, L., Sonzogni, C., and Ganssen, G.: Sea surface temperature and productivity records for the last 240 kyr on the Arabian Sea, in: Deep Sea Research Part II: Topical Studies in Oceanography, Deep Sea Research Part II: Topical Studies in Oceanography, 1997, 10.1016/S0967-0645(97)00008-8, 1997.
- 1350 Russell, A. D., Hönisch, B., Spero, H. J., and Lea, D. W.: Effects of seawater carbonate ion concentration and temperature on shell U, Mg, and Sr in cultured planktonic foraminifera, *Geochim. Cosmochim. Acta*, 68, 10.1016/j.gea.2004.02.013, 2004.
- 1355 Russell, A. D., Emerson, S., Nelson, B. K., Erez, J., and Lea, D. W.: Uranium in foraminiferal calcite as a recorder of seawater uranium concentrations, *Geochim. Cosmochim. Acta*, 58, 10.1016/0016-7037(94)90497-9, 1994.
- Sadekov, A. Y., Eggins, S. M., and De Deckker, P.: Characterization of Mg/Ca distributions in planktonic foraminifera species by electron microprobe mapping, *Geochim. Geophys. Geosyst.*, 6, 10.1029/2005GC000973, 2005.
- 1360 Sagawa, T., Hasegawa, T., Narita, Y., Yokoyama, M., Kubota, Y., Okazaki, Y., Goto, A. S., Suzuki, Y., Ikehara, K., and Nakagawa, T.: Millennial-scale paleotemperature change in the Japan Sea during Marine Isotope Stage 3: Impact of meridional oscillation of the subpolar front, *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 626, 111713, 10.1016/j.palaeo.2022.111713, 2022.
- 1365 Saher, M. H., Peeters, F. J. C., and Kroon, D.: Sea surface temperatures during the SW and NE monsoon seasons in the western Arabian Sea over the past 20,000 years, *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 10.1016/j.palaeo.2007.01.014, 2007a.

- 1370 Saher, M. H., Jung, S. J. A., Elderfield, H., Greaves, M. J., and Kroon, D.: Sea surface temperatures of the western Arabian Sea during the last deglaciation, *Paleoceanography*, 10.1029/2006PA001292, 2007b.
- Saher, M. H., Rostek, F., Jung, S. J. A., Bard, E., Schneider, R. R., Greaves, M., Ganssen, G. M., Elderfield, H., and Kroon, D.: Western Arabian Sea SST during the penultimate interglacial: A comparison of U- δ^{37} and Mg/Ca paleothermometry, *Paleoceanography*, 10.1029/2007PA001557, 2009.
- 1375 Schott, F. A. and McCreary, J. P.: The monsoon circulation of the Indian Ocean, *Prog. Oceanogr.*, 51, 1–123, 10.1016/S0079-6611(01)00083-0, 2001.
- Schulz, H., Von Rad, U., and Erlenkauer, H.: Correlation between Arabian Sea and Greenland climate oscillations of the past 110,000 years, *Nature*, 393, 10.1038/31750, 1998.
- 1380 Shetye, S. R., Gouveia, A. D., and Shenoi, S. S. C.: Circulation and water masses of the Arabian Sea, *Proceedings of the Indian Academy of Sciences Earth and Planetary Sciences*, 103, 10.1007/BF02839532, 1994.
- Singh, A. D., Jung, S. J. A., Darling, K., Ganeshram, R., Ivanochko, T., and Kroon, D.: Productivity collapses in the Arabian Sea during glacial cold phases, *Paleoceanography*, 26, 10.1029/2009PA001923, 2011.
- 1385 Sonzogni, C., Bard, E., Rostek, F., Lafont, R., Rosell-Mele, A., and Eglinton, G.: Core-top calibration of the alkenone index vs sea surface temperature in the Indian Ocean, in: *Deep Sea Research Part II: Topical Studies in Oceanography*, Deep Sea Research Part II: Topical Studies in Oceanography, 1997/01/01, 1445–1460, 10.1016/S0967-0645(97)00010-6, 1997.
- 1390 Southon, J., Kashgarian, M., Fontugne, M., Metivier, B., and Yim, W. W. S.: Marine Reservoir Corrections for the Indian Ocean and Southeast Asia, *Radiocarbon*, 44, 167–180, 10.1017/S0033822200064778, 2002.
- 1395 Stott, L., Poulsen, C. J., Lund, S., and Thunell, R.: Super ENSO and global climate oscillations at millennial time scales, *Science (American Association for the Advancement of Science)*, 297, 222–226, 10.1126/science.1071627, 2002.
- Stuiver, M. and Reimer, P. J.: Extended 14C data base and revised CALIB 3.0 14C age calibration program, *Radiocarbon*, 35, 10.1017/S0033822200013904, 1993.
- 1400 Svensson, A., Andersen, K. K., Bigler, M., Clausen, H. B., Dahl-Jensen, D., Davies, S. M., Johnsen, S. J., Muscheler, R., Rasmussen, S. O., Röthlisberger, R., Peder Steffensen, J., and Vinther, B. M.: The Greenland Ice Core Chronology 2005, 15–42 ka. Part 2: comparison to other records, *Quaternary Science Reviews*, 25, 10.1016/j.quascirev.2006.08.003, 2006.
- 1405 Swallow, J. C. and Bruce, J. G.: Current measurements off the Somali coast during the southwest monsoon of 1964, *Deep Sea Research and Oceanographic Abstracts*, 13, 10.1016/0011-7471(76)90908-6, 1966.
- Tachikawa, K., Sepulchre, S., Toyofuku, T., and Bard, E.: Assessing influence of diagenetic carbonate dissolution on planktonic foraminiferal Mg/Ca in the southeastern Arabian Sea over the past 450 ka: Comparison between *Globigerinoides ruber* and *Globigerinoides sacculifer*, *Geochem. Geophys. Geosyst.*, 9, 10.1029/2007GC001904, 2008.
- 1410

van Hinte, J. E., van Weering, T. C. E., and Troelstra, S. R.: Tracing a seasonal
 upwelling, *report*, 146, 1995.
 1415 Vázquez Riveiros, N., Govin, A., Waelbroeck, C., Mackensen, A., Michel, E., Moreira,
 S., Bouinot, T., Caillon, N., Orgun, A., and Brandon, M.: Mg/Ca thermometry in planktic
 foraminifera: Improving paleotemperature estimations for *G. bulloides* and *N.*
 pachyderma left, *Geochem. Geophys. Geosyst.*, 17, 1249–1264, 10.1002/2015GC006234,
 2016.
 1420 Veres, D., Bazin, L., Landais, A., Toyé Mahamadou Kele, H., Lemieux-Dudon, B.,
 Parrenin, F., Martinerie, P., Blayo, E., Blunier, T., Capron, E., Chappellaz, J.,
 Rasmussen, S. O., Severi, M., Svensson, A., Vinther, B., and Wolff, E. W.: The
 Antarctic ice core chronology (AICC2012): an optimized multi-parameter and multi-site
 dating approach for the last 120 thousand years, *Clim. Past*, 9, 1733–1748, 10.5194/cp-
 9-1733-2013, 2013.
 1425 Vinther, B. M., Clausen, H. B., Johnson, S. J., Rasmussen, S. O., Andersen, K. K.,
 Buchardt, S. L., Dahl-Jensen, D., Seierstad, I. K., Siggaard-Andersen, M. L.,
 Steffensen, J. P., Svensson, A., Olsen, J., and Heinemeier, J.: A synchronized dating
 of three Greenland ice cores throughout the Holocene, *Journal of Geophysical Research*
 1430 *Atmospheres*, 111, 10.1029/2005JD006921, 2006.
 Wang, X., Sun, J., Longstaffe, F. J., Gu, X., Du, S., Cui, L., Huang, X., and Ding,
 Z.: Climatic quantification and seasonality of the late MIS 3 in north China: a
 perspective from carbon and oxygen isotopes of fossil mammal teeth, *Quaternary science*
 reviews, 272, 107222, 10.1016/j.quascirev.2021.107222, 2021.
 1435 Wang, Y. V., Leduc, G., Regenber, M., Andersen, N., Larsen, T., Blanz, T., and
 Schneider, R. P.: Northern and southern hemisphere controls on seasonal sea surface
 temperatures in the Indian Ocean during the last deglaciation, *Paleoceanography*, 28,
 619–632, <https://doi.org/10.1002/pale.20053>, 2013.
 1440 Webster, P. J., Magaña, V. O., Palmer, T. N., Shukla, J., Tomas, R. A., Yanai, M.,
 and Yasunari, T.: Monsoons: Processes, predictability, and the prospects for
 prediction, *Journal of Geophysical Research: Oceans*, 10.1029/97JC02719, 1998.
 Wolff, E. W., Chappellaz, J., Blunier, T., Rasmussen, S. O., and Svensson, A.:
 Millennial scale variability during the last glacial: The ice core record, *Quaternary*
 Science Reviews, 29, 10.1016/j.quascirev.2009.10.013, 2010.
 1445 Wyrtki, K.: *Physical Oceanography of the Indian Ocean*, in, 1973.
 Zeeden, C., Obrecht, I., Veres, D., Kaboth-Bahr, S., Hošek, J., Marković, S. B.,
 Böskén, J., Lehmkuhl, F., Rolf, C., and Hambach, U.: Smoothed millennial scale
 palaeoclimatic reference data as unconventional comparison targets: Application to
 European loess records, *Sci Rep*, 10, 5455, 10.1038/s41598-020-61528-8, 2020.
 1450

Field Code Changed

Depth (cm)	Original age from Saher et al., (2007) (kaBP)	Recalibrated age - median (kaBP)	1 sigma age range (kaBP)	2 sigma age range (kaBP)	Recalibrated - original age (kaBP)
21.75	1.292	1.305	1.197 - 1.398	1.102 - 1.516	0.013
50.75	5.699	5.693	5.579 - 5.804	5.475 - 5.905	-0.006
76.75	8.929	8.928	8.780 - 9.069	8.632 - 9.209	-0.001
82.25	9.737	9.779	9.603 - 9.912	9.526 - 10.088	0.042
125.25	13.508	13.569	13.449 - 13.700	13.312 - 13.805	0.061
180.25	18.313	18.031	17.901 - 18.177	17.712 - 18.289	-0.282
210.25	19.92	20.035	19.865 - 20.212	19.663 - 20.375	0.115

Table 1

Depth (cm)	Age (kaBP)
317.5	31
496.5	45
669.5	62
832.5	87
958	107
1025.5	126

Table 2

depth (cm)	age (initial LR tuning; kaBP)	new age based on NGRIP tuning (kaBP)	note
251	24.13	23.14	NGRIP based age
317.5	31		LR04 based control point used
357.5	34.128	35.2	NGRIP based age
396.5	37.179	37.92	NGRIP based age
463.5	42.419	41.86	NGRIP based age
496.5	45		LR04 based control point used
571.5	52.37	53.88	NGRIP based age
669.5	62		LR04 based control point used
671.5	62.307	62.9	NGRIP based age
727.5	70.896	70.02	NGRIP based age
741.5	73.043	73.18	NGRIP based age
774.5	78.104	77.28	NGRIP based age
832.5	87		LR04 based control point used
958	107.28	109.52	LR04 control point with revised NGRIP based age
1025.5	126		LR04 based control point used

Table 3






Figure 1




(C)



G. bulloides

Summer = 0.1
Winter = 0.9

$B = 0.9S + 0.1W$



G. ruber

Summer = 0.5
Winter = 0.5

$R = 0.5S + 0.5W$

rearranged B equation to: $S = \frac{B}{0.9} - \frac{0.1W}{0.9}$

substitute this value of S into R equation and rearrange:

$$R = 0.5 * \left(\frac{B}{0.9} - \frac{0.1W}{0.9} \right) + 0.5W$$

$$2R = \frac{B}{0.9} - \frac{0.1W}{0.9} + W$$

$$2R - \frac{B}{0.9} = -\frac{0.1W}{0.9} + W$$

$$1.8R - B = -0.1W + 0.9W$$

$$1.8R - B = 0.8W$$

allowing for: $W = \frac{1.8R - B}{0.8}$

which can be used in: $S = \frac{B - 0.1W}{0.9}$

Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7