

Supporting Information for Occurrence, abundance, and formation of atmospheric tarballs from a wide range of wildfires in the western US

5 Kouji Adachi¹, Jack E. Dibb², Joseph M. Katich^{3,4#}, Joshua P. Schwarz³, Hongyu Guo^{4,5}, Pedro Campuzano-Jost^{4,5}, Jose L. Jimenez^{4,5}, Jeff Peischl³, Christopher D. Holmes⁶, James Crawford⁷

¹Department of Atmosphere, Ocean, and Earth System Modelling Research, Meteorological Research Institute, Tsukuba, Japan

²Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham, NH, USA

³Chemical Sciences Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO, USA

⁴Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, CO, USA

10 ⁵Department of Chemistry, University of Colorado Boulder, Boulder, CO, USA

⁶Earth, Ocean, and Atmospheric Science, Florida State University, Tallahassee, FL, USA


⁷NASA Langley Research Center, Hampton, VA, USA

[#]now at Ball Aerospace, Boulder, CO, USA

Correspondence to: Kouji Adachi (adachik@mri-jma.go.jp)

15

Supporting figures 1-3.



20

Figure S1. The flow chart for classifying individual particles into seven categories based on an image analysis technique (Deep learning) and STEM-EDS measurements.












Figure S2. Representative TEM images of samples containing TBs from each wildfire. Scale bars indicate 5 μ m. The header of each TEM image shows the sampling date and time (UTC) and the name of the wildfire.



30 **Figure S3. Fractions of samples having aggregated TBs for samples with various TB number fractions. The number of samples containing TBs was 179, while those containing aggregated TBs were 84. n = 54, 50, 36, 16, 10, and 13 for the number of samples with TB fractions of 0%–5%, 5%–10%, 10%–15%, 15%–20%, 20%–25%, and >25%, respectively.**