

1 **Concurrent** Bering Sea and Labrador Sea ice melt extremes in March
2 **2023: A confluence of meteorological events aligned with**
3 **stratosphere-troposphere interactions**

Deleted: Simultaneous

4 Thomas J. Ballinger¹, Kent Moore^{2,3}, Qinghua Ding⁴, Amy H. Butler⁵, James E. Overland⁶, Richard L.
5 Thoman¹, Ian Baxter⁴, Zhe Li⁴, and Edward Hanna⁷

6 ¹International Arctic Research Center, University of Alaska Fairbanks, Fairbanks, AK, USA

7 ²Department of Physics, University of Toronto, Toronto, Ontario, Canada

8 ³Department of Chemical and Physical Sciences, University of Toronto, Canada, Toronto, Ontario, Canada

9 ⁴Department of Geography, and Earth Research Institute, University of California, Santa Barbara, Santa Barbara, CA, USA

10 ⁵Chemical Sciences Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO, USA

11 ⁶Pacific Marine Environmental Laboratory, National Oceanic and Atmospheric Administration, Seattle, WA, USA

12 ⁷Department of Geography and Lincoln Climate Research Group, [University of Lincoln](#), Lincoln, UK

13

14 *Correspondence to:* Thomas J. Ballinger (tjballinger@alaska.edu)

15 **Abstract.** Today's Arctic is characterized by a lengthening of the sea ice melt season, but also by fast and at times unseasonal
16 melt events. Such anomalous melt cases have been identified in Pacific and Atlantic Arctic sector sea ice studies. Through
17 observational analyses, we document an unprecedented, ~~concurrent~~ marginal ice zone melt event in the Bering Sea and
18 Labrador Sea in March of 2023. Taken independently, variability in the cold season ice edge at synoptic time scales is common.
19 However, such anomalous, short-term ice loss over either region *during the climatological sea ice maxima* is uncommon, and
20 the tandem ice loss that occurred qualifies this as a rare event. The atmospheric setting that supported the unseasonal melt
21 events was preceded by a sudden stratospheric warming event that ~~led to positive tropospheric height anomalies across much~~
22 of the Arctic and the development of anomalous mid-troposphere ridges over the ice loss regions. These large-scale
23 anticyclonic centers funneled extremely warm and moist airstreams onto the ice causing melt. Further analysis identified the
24 presence of atmospheric rivers within these warm airstreams whose characteristics likely contributed to this bi-regional ice
25 melt event. Whether such a confluence of anomalous wintertime events associated with troposphere-stratosphere coupling may
26 occur more often in a warming Arctic remains a research area ripe for further exploration.

Deleted: simultaneous

Deleted: , along with ongoing La Niña teleconnections.

27 **1 Introduction**

28 Observational analyses of the Arctic atmosphere have noted warmer air temperatures and increased moisture content during
29 the last two decades relative to previous years (Ballinger et al., 2023; Boisvert et al., 2023). Periods of increased climate
30 variability (Hanna et al., 2015) can coincide with these atmospheric changes in the Arctic to produce extreme meteorological
31 phenomena, which may influence human and environmental systems both within and beyond the high northern latitudes.

35 Moreover, terrestrial Arctic snow and sea ice extent, area, and depth/thickness control heat exchange between the land, ocean,
36 and atmosphere (Serreze and Barry, 2011). With less snow and sea ice in a warming Arctic, instances of surface-to-atmosphere
37 heating perturbations can magnify impacts of synoptic circulation patterns on local and/or remote surface weather extremes
38 (Francis and Vavrus, 2015; Zhang et al., 2018; Tachibana et al., 2019; Bailey et al., 2021). Thus air-sea interactions resulting
39 in extreme events in today's Arctic are structurally complex (Walsh et al., 2020) and shaped by the surface condition/type and
40 prevailing weather pattern (Overland et al., 2021).

41
42 A key consideration of complex Arctic extreme events is their timing of occurrence within the annual cycle. As an example,
43 the Arctic Ocean's ice cover tends to thin and decline (thicken and increase) through the boreal summer (winter) months up to
44 the September minima (March maxima). However, analyses of satellite observations have shown a trend toward earlier melt
45 onset across most of the Arctic marginal seas (e.g., Stroeve and Notz, 2018) with unusually-timed and often isolated ice loss
46 events during winter or early spring interspersed on these trends. The North Atlantic Arctic region that includes marginal seas
47 around Greenland, Iceland, and northwest Europe has experienced several of these cases in recent times. During mid-April of
48 2013, a persistent anticyclone over Greenland coincided with record-early melt onset in the Baffin Bay, Davis Strait, and
49 Labrador Sea region that was ~8 weeks earlier than the 1981-2010 average (Ballinger et al., 2018). Above freezing air
50 temperatures at the North Pole during late December of 2015 led to a substantial loss of sea ice over the Arctic Ocean (Moore,
51 2016). In late February and early March of 2018, a polynya unexpectedly opened off the northern Greenland coast that was
52 driven by anomalously warm and strong southerly winds that were preceded by a sudden stratospheric warming (SSW) event
53 (Moore et al., 2018). In one of the most notable examples, an Arctic cyclone that registered record-low central pressure
54 traversed the Barents and Kara seas in late January of 2022 and caused record surface winds and attendant ice loss for the time
55 of year (Blanchard-Wrigglesworth et al., 2022). Unlike the previous cases, dynamical and ocean processes rather than
56 thermodynamics were attributed to this unseasonal ice loss event.

57
58 There is a large body of research into so-called compound extreme climate events such as the simultaneous occurrence in a
59 particular region of a drought and heat wave or a storm surge and fluvial flooding (e.g., Zscheischler et al., 2018; AghaKouchak
60 et al., 2020). Less well-studied are so-called concurrent climate extreme events where two or more spatially isolated regions
61 are subject to simultaneous or near-simultaneous extremes (Zhou et al., 2023). Compound events may be associated with a
62 single overarching phenomenon such as a hurricane, while concurrent events are typically associated with amplified Rossby
63 Waves (Kornhuber et al., 2020).

64
65 In this study, we have identified the first known observation of a concurrent climate extreme event in the Arctic as well as one
66 that is associated with a SSW. This concurrent event is marked by unusually-timed sea ice melt in the Bering Sea and Labrador
67 Sea during March of 2023. Our goals in this observationally-based case study are to describe the respective regional sea ice
68 conditions during March 2023, place them in historical spatial and temporal context, and evaluate the synoptic atmospheric

- Deleted: ¶
Here
- Deleted: w
- Deleted: e characterize a
- Deleted: simultaneous
- Deleted: and
- Deleted: cross-Arctic
- Deleted: event
- Deleted: that transpired in

78 mechanisms responsible for the ensuing melt extremes. As part of our analyses, we evaluate the probability of such sea ice
79 melt extremes amidst the period that encompassed the climatological Arctic sea ice maximum. We conclude with a discussion
80 of our findings that considers seasonal and synoptic meteorological anomalies that occurred during and around the time of
81 these melt events. Our conclusions also touch upon the implications of Arctic warming for analogous future melt events.

82 2. Data and Methods

83 2.1 Sea ice and atmospheric datasets

84 Daily sea ice concentration (SIC in %) is derived from the NOAA/NSIDC Climate Data Record (CDR) of passive microwave
85 SIC, version 4 (Meier et al., 2021, 2022). This dataset represents a blended product of the NASA Team algorithm (Cavalieri
86 et al., 1984) and NASA Bootstrap algorithm (Comiso, 1986), and is available daily on a 25 km² grid from 1979-onwards.

87
88 ECMWF fifth generation global atmospheric reanalysis (ERA5) data at their 31 km native resolution for 1979-2023 (Hersbach
89 et al., 2020) are used to evaluate atmospheric conditions across the Arctic region during and around the SSW event and ensuing
90 sea-ice melt extremes. ERA5 fields examined include 2-meter air temperature (T2m in °C), total column water vapor (in mm),
91 total precipitation, which is the sum of large-scale and convective precipitation including rain and snowfall, that reaches the
92 surface (in mm/day), [net and downward longwave radiation \(in W/m²\)](#), and geopotential heights (in m) over the atmospheric
93 column from 1000 hPa to 1 hPa. Unless otherwise stated, data are binned to daily means. Studies have shown ERA5 to be
94 effective at capturing Arctic weather and climate variability. As an example, during a research expedition in Fram Strait,
95 Graham et al. (2019) noted ERA5 air temperatures, humidity, and winds exhibited relatively strong correlations and low biases
96 in comparison with radiosonde observations and performed better overall than other modern atmospheric reanalyses in the
97 region. Numerous other studies have relied upon ERA5 data to understand the synoptic evolution and characteristics of
98 airstreams within the Arctic (e.g., Nygard et al., 2020; Papritz et al., 2022; Kirbus et al., 2023).

99
100 In addition to reanalysis fields, daily averaged T2m data from regional weather stations are evaluated (**Figure 1**). We
101 deliberately selected near-coastal weather stations based on several criteria, including multidecadal records that are relatively
102 complete (>95% of dates surveyed register a T2m value) for sites located north and south of both the early March long-term
103 mean and 2023 ice edge in the Bering Sea and Labrador Sea, respectively. Data from leap years are omitted as 2023 was not
104 one. For the Bering Sea region, we obtained T2m data from the National Centers for Environmental Information Applied
105 Climate Information System (NCEI ACIS) for Alaska terrestrial weather stations at St. Paul (57.16°N, 170.22°W) and
106 Kotzebue (66.89°N, 162.58°W). The St. Paul historical record is surveyed from 1916-2023, while the Kotzebue record is
107 assessed from 1923-2023. For the Labrador region two western Greenland weather station records, which are maintained by
108 the Danish Meteorological Institute (DMI), are obtained for Nuuk (64.17°N, 51.75°W) and Aasiaat (68.70°N, 52.75°W). Both
109 of these Greenland records span 1958 to 2023. We supplement NCEI ACIS and DMI observations with Programme for

110 Monitoring of the Greenland Ice Sheet (PROMICE) automatic, on-ice weather station temperatures, measured from a nominal
111 height of 2.7 m above the ice-sheet surface, for two sites: one is near Nuuk on a peripheral glacier (NUK_K; 64.16°N ,
112 51.36°W; 710 m asl) and the other is found within the lower ablation area of the Greenland Ice Sheet (GrIS) in the Qassimiut
113 region (QAS_L; 61.03°N, 46.85°W; 280 m asl) (Fausto et al., 2021). The PROMICE data records are relatively short, with
114 NUK_K established in 2015 and QAS_L in 2008, though both are 99% complete for the dates we surveyed and provide
115 valuable information on GrIS in situ air temperatures on the rather observationally sparse Greenland Ice Sheet.

116
117 Several atmospheric indices are analyzed and discussed in this work. The SSW compendium (Butler et al., 2017, updated), a
118 long-term archive of indicator climate indices associated with SSW events, confirmed the onset of the late-winter 2023 SSW
119 event (16 February). We examine one such metric of this archive that we term the Polar Vortex Index (PVI) that describes the
120 daily-mean, zonal-mean winds at 60°N and 10 hPa, where the timing of the shift from westerly to easterly stratospheric flow
121 between November and April signifies the SSW onset (Charlton and Polvani, 2007). The PVI is analyzed from 1979-2023.
122 SSWs are known to influence the mid-to-high latitude tropospheric circulation patterns and often precede a negative North
123 Atlantic Oscillation (NAO) regime and high-latitude anticyclonic blocking (Baldwin et al., 2020). Therefore, we elect to
124 analyze the daily NAO and region-specific Greenland Blocking Index (GBI) and Alaska Blocking Index (ABI). The NAO
125 used here extends from 1950 to 2023 and is defined as the leading, rotated principal component of standardized 500 hPa
126 geopotential height (z500) anomalies from 20-90°N (Barnston and Livezey, 1987). The GBI describes the mean z500 across
127 60-80°N, 20-80°W (Hanna et al., 2013), and the ABI depicts the averaged z500 from 55-75°N and 125-180°W (Ballinger et
128 al., 2022). These blocking indices are analyzed over the 1948 to 2023 period.

129

130 *2.2 Extreme event detection methods*

131 We examine moisture transport into the Arctic during our case study by employing an atmospheric river (AR) detection
132 algorithm developed by Guan and Waliser (2019). This algorithm is applied on 6-hourly ERA5 integrated water vapor transport
133 (IVT in kg/m/s) data, averaged from 1000hPa to 300 hPa on a 1.5° x 1.5° global grid. In this framework, ARs for each 6-hour
134 interval are defined when an IVT threshold exceeding the 85th percentile of climatological IVT is reached for a grid cell in the
135 domain of interest. Additionally, these ARs must meet specific criteria related to the orientation, length, and length-to-width
136 ratio of IVT, as outlined by Guan and Waliser (2019). Widely adopted in previous studies spanning the tropics to the high
137 latitudes including the Arctic and Antarctic, this algorithm serves as a reliable scheme for AR analysis (Collow et al., 2022).
138 We examine the duration of AR events passing through the Alaska and Greenland regional domains shown in **Figure 1** leading
139 up to, coinciding with, and following the Bering and Labrador melt events, respectively. AR duration is defined as the
140 percentage (%) of the day in which an AR resides within any portion of the respective domains. We also measure the intensity
141 of AR events, defined here as the mean IVT of all grid cells that cross into either domain associated with an AR.

142

143 Daily atmospheric indices and maps of the reanalysis data are presented, and values are identified that meet or exceed an
144 extreme value threshold (i.e., 95th or 99th percentile) relative to a specified number of days across the data records described in
145 Section 2.1. For example, a 99th percentile St. Paul, Alaska T2m value during the 90-day “winter” period from 1 January – 31
146 March 1916-1923 (where 9576 days registered a daily mean T2m reading) is 3.3°C. Use of the full historical period or select
147 portions of dataset’s records along with extended time windows (e.g., 1 January – 31 March) provided a larger sample size
148 from which to calculate extreme values relative to the period specified or season (e.g., 90 days) versus a singular date of
149 reference.

150 3. Results

151 3.1 Extreme and unusually-timed sea ice melt

152 The regional SIC means, variability, and anomalies around the peak of the melt events relative to 1-15 March 2000 to 2023
153 are shown in **Figure 1**. This subset of years is selected as winter months since 2000 have seen a large decline in sea ice
154 conditions (Stroeve and Notz, 2018). In the Labrador region, the 50% climatological ice edge tilted northeast to southwest
155 from Davis Strait into the Labrador Sea and transitioned in the marginal ice zone to nearly 100% SIC on the western flank of
156 this boundary (**Figure 1a**). In contrast, the Bering Sea ice edge exhibited a more zonal orientation and extended from ~61°N
157 in the western Bering Sea to ~59°N in the eastern Bering Sea (**Figure 1d**). From 2000 to 2023, interannual SIC variability for
158 the first half of March in these marginal ice zone areas was ~30% (**Figure 1b, e**), while early March 2023 saw SIC reductions
159 along the ice edge on the order of ~30% (**Figure 1c,f**).

160





Figure 1. Sea ice concentration (SIC in %) from the NOAA/NSIDC CDR dataset. Mean conditions for the period 1-15 March 2000-2023 for: (a) the Labrador Sea and d) the Bering Sea. The SIC standard deviation (%) for 1-15 March 2000-2023 is shown for: (b) the Labrador Sea and e) the Bering Sea. The sea ice concentration anomaly on 5 March 2023 relative to the 1-15 March 2000-2023 period is shown for (c) the Labrador Sea and (f) the Bering Sea. In (a) and (d) locations of the weather stations mentioned in the text are indicated with arrows.

161
 162 The SIC conditions in these areas of >30% variability are examined more closely with respect to the winter of 2023. Winter is
 163 loosely defined here as January through March. From mid-January through February, the daily Labrador SIC exceeded the
 164 2000-2023 mean, then abruptly plummeted to below-normal conditions in early March and remained below-average through
 165 the end of the month (**Figure 2a**). The Bering SIC showed more variability about the SIC day-of-year means with periods of
 166 slightly above and below-normal ice cover into early March and through the rest of the month (**Figure 2b**). While single day
 167 SIC departures through winter in both areas did not breach the 5th or 95th percentiles for the day of year, the largest 4-day
 168 changes (<20% SIC losses) occurred roughly at the same time and culminated on March 5th in the Labrador Sea and March 6th
 169 in the Bering Sea (see dashed red vertical lines in **Figure 2**). While day-to-day sea ice variability is not unusual throughout

170 winter, the day-of-year mean curves (thick black lines in **Figure 2**) suggest that ice growth tends to continue in both of these
 171 regions throughout most of March aligned with the typical pan-Arctic sea ice maximum (Meier et al., 2023).
 172





Figure 2. Time series (red curves) of the daily SIC averaged over the regions, a) Labrador Sea and b) Bering Sea, respectively in Figures 1b,e, where the standard deviation exceeds 30% for the period 1 January to 26 March 2023. The black line represents the daily mean value for the period 2000-2023 with the shading incorporating daily values between the 5th and 95th percentile values. The ending dates for the 4-day window with the largest change in ice concentration are shown with the dotted red lines.

173



174 Histograms provide additional probabilistic perspective on the likelihood of such 4-day ice loss events for the times of year
 175 they occurred in 2023 (**Figure 3**). Since 2000, both the Labrador Sea (red curve) and Bering Sea (blue curve) have shown
 176 quasi-normal SIC distributions over the 1-15 March period. The 2023 4-day changes in both areas, characterized by ~20% SIC
 177 reduction in the Bering Sea and ~27% SIC decline in the Labrador Sea, represent extreme outliers found in the far-left tails of
 178 their respective data distributions. The magnitude of these short-term SIC loss events is uncommon for the time of year, which
 179 prompts further investigation into the synoptic processes that drive, and potentially link, these rare, concurrent events.






Figure 3. Histograms of the change over a 4-day period during 1-15 March 2000-2023 for the Labrador Sea (red) and Bering Sea (blue) regions used in Figure 2. The shading represents the regions bounded by the 1st and 99th percentile values. The largest changes during March 2023 are indicated by the dashed lines.

180 *3.2 Synoptic mechanisms, part 1: The 2023 SSW event and its stratosphere-troposphere signatures*

181 On 16 February, a SSW occurred that appears to have largely initiated the synoptic environment conducive to the cross-Arctic
 182 melt events. **Figure 4** shows the winter-long evolution of the height anomalies with respect to the SSW event. In mid-January
 183 2023, positive tropospheric heights in the 1000-100 hPa layer preceded positive height anomalies aloft that developed toward
 184 late January and early February. The positive height anomalies indicate upward troposphere to stratosphere coupling that
 185 resulted in a minor stratospheric warming event at the end of January. Over the two weeks that followed, a second, stronger
 186 and positive (~2 sigma) geopotential height anomaly developed aloft within the upper stratosphere and peaked on 16 February

187 in conjunction with the day of the shift from westerly to easterly 10 hPa winds at 60°N found in the PVI (**Figure 5a**), which
 188 marked the date of SSW onset (Butler et al., 2017, updated). The PVI dipped to roughly the 1st percentile following SSW onset
 189 on 28 February and 1 March, characterizing this as an anomalously strong event for this time of year. The PVI reached a
 190 minimum wind speed of -18 m/s on 28 February, which places it as the 6th strongest reversal (out of 28 such events) of the
 191 polar vortex winds during a stratospheric warming from 1979-2023 (Lee and Butler 2019). As is the tendency with SSWs, the
 192 influence of the above-average, upper stratospheric air pressures and temperatures (latter not shown) descended during this
 193 time, yielding increased heights across the depth of the stratosphere through late February (**Figure 4**). By early March, the
 194 SSW warming signal propagated toward the surface and large positive height anomalies extended through the depth of the
 195 tropospheric column. The largest positive height anomalies within the lower troposphere and at the surface coincided with the
 196 Bering Sea and Labrador Sea melt events.

197




198

199

200 **Figure 4.** Polar cap (60-90°N) standardized geopotential height anomalies (unitless) from the surface to the upper stratosphere
 201 during winter 2023. The standardized anomalies are calculated at each pressure level by removing the daily climatology and
 202 dividing by the daily standard deviation. The standardized anomalies are shown relative to the day of year for the 1979-2023
 203 period of the ERA5 reanalysis.

204

205 In the two weeks that led up to this strong SSW event, the large-scale mid-tropospheric circulation was characterized by a
 206 positive NAO fluctuation between 0 and 1.5 sigma, indicative of stronger than normal westerly winds across the mid-to-high



Deleted:

208 latitudes (**Figure 5b**). Negative height anomalies (lower than normal pressure) across most of the polar cap troposphere
 209 between 1-15 February (**Figure 4**) support this assertion. After the SSW event on 16 February, the NAO slightly increased for
 210 two days then plummeted, reversed sign, and became strongly negative (~ -1 sigma) from 2-8 March around the melt events
 211 (**Figure 5b**). Zooming in on the study regions of interest, strong, lagged ridging responses are noted in the respective mid-
 212 tropospheric height fields. The z500 pattern atop the Labrador Sea area of ice loss described by the GBI is >100 m above-
 213 average from 1-12 March, including record-high day of year departures (since 1948) from 4-7 March when the GBI values
 214 exceeded the 99th percentile (**Figure 5c**). This period also corresponded with the strongest downward coupling of the SSW
 215 event to surface conditions (**Figure 4**). While comparatively not as extreme as those of the GBI, ABI values are also
 216 considerably higher-than-average during most of the same period (4-12 March), punctuated by >100 m anomalies from 5-11
 217 March (**Figure 5d**). As a precursor to the SSW event and subsequent pressure increase through much of the atmospheric
 218 column, initially there were ABI peak with values >150 m above-normal from 27-30 January capped by the 28 January ABI
 219 value (5496.83 m) falling just shy of the 99th percentile (5501.82 m). The Alaska ridge (not shown) associated with these
 220 elevated ABI values appears well-timed with upward coupling of the troposphere to the stratosphere (**Figure 4**). These late
 221 January and mid-March examples could suggest a slow forcing mechanism such as the background La Niña state rendering
 222 stronger blocking anticyclones over Alaska. We revisit this discussion in **Section 4**.

223






Figure 5. Daily atmospheric indices for 1 January – 31 March 2023 (purple lines) overlapping the multi-sectoral melt event for the a) Polar Vortex Index (m/s), b) North Atlantic Oscillation Index (standardized), c) Greenland Blocking Index (m), and d) Alaska Blocking Index (m). Considering all days from 1 January to 31 March for the respective indices, full periods of record (see Section 2.1), the mean of each variable (black line), 1st percentile (blue dashed line), and 99th percentile (red dashed line) are shown in each graphic. The sudden stratospheric warming event on 16 February 2023 is labeled with a green diamond, and to draw attention to the dates around the Labrador Sea and Bering Sea melt events, the period from 2-10 March 2023 is identified by yellow triangles.

Deleted: T2m


224
225 The evolution of the day-to-day z500 spatial pattern provides perspective to the values of the large-scale circulation and
226 regional blocking indices overlapping the melt events. The height pattern over Greenland, Baffin Bay and Labrador Sea is
227 above-normal and successively strengthens during 2-4 March (**Figure 6a-c**) before the peak in the short-term Labrador Sea
228 melt observed on 5 March when western Greenland and Baffin Bay is engulfed in >99th percentile height anomalies (**Figure**
229 **6d**). Meanwhile, below-normal mid-tropospheric pressure over Alaska and poleward of the central Bering Sea from 2-4 March
230 gave way to higher-than average pressure by 5 March and preceded the 6 March peak in the Bering Sea ice loss (**Figure 6e**).
231 A large-scale dipole structure is evident from 6-10 March, as the North American (Eurasian) high-latitudes spanning the
232 International Dateline (i.e., 180°W) to ~30°W (30°W-180°W) exhibited higher-than-normal (lower-than-normal) heights with
233 extreme departures around Greenland (**Figure 6e-i**) that are reflected in the magnitude of the daily GBI anomalies (**Figure**
234 **5c**). Midtropospheric ridging over high-latitude North America with larger anomalies over Greenland than Alaska represents
235 a common regional weather regime (Lee et al., 2023), however, the z500 anomalies observed during the latter portion of our
236 case study are relatively higher in magnitude. In terms of set-up, over the 9-day period, the blocking pattern developed initially
237 over the Iceland region before retrograding westwards over Greenland towards the Labrador Sea and Baffin Bay. Such
238 retrograde movements have been noted to occur in other cases of blocking development over the Greenland region (Hanna et
239 al., 2018). While the z500 pattern orientation and development are not uncommon, the strength of the anticyclonic anomalies
240 is notable in this case.
241



242

243

244 **Figure 6.** 500 hPa geopotential height (z500) anomaly (m) from the ERA5 at 0 GMT on: a) 2 March, b) 3 March, c) 4 March,
 245 d) 5 March, e) 6 March, f) 7 March, g) 8 March, h) 9 March, and i) 10 March 2023. The anomalies are presented with respect
 246 to the period 16 February – 15 March 1979-2023. Gridpoints where the anomalies are less than the 1st percentile (blue hues)
 247 or greater than the 99th percentile (red hues) based on the above period are indicated with the '+'.
 248



Deleted:

250 3.3 Synoptic mechanisms part 2: Thermodynamic effects

251 The synoptic set-up following the 2023 SSW event was characterized by mid-tropospheric height increases and development
252 of intense ridging patterns over the Labrador Sea and Bering Sea that spanned the respective regional ice loss events. Here, we
253 examine the thermodynamic environment overlapping the aforementioned atmospheric circulation anomalies. **Figure 7** shows
254 the daily pan-Arctic T2m anomaly field (shading) around the melt events; the 0°C isotherm (blue contour) is overlaid for
255 reference. During 2-4 March, air temperature anomalies over south central Greenland, Davis Strait, and northern Labrador
256 waters overlapping the ice edge were above-normal (**Figure 7a-c**). In particular, from the 2nd to the 3rd of March, the 0°C
257 isotherm abruptly migrated westward and encompassed much of the Labrador Sea including the ice edge (refer to **Figure 1a**).
258 During this time 99th percentile warm extremes were found across the northern Labrador Sea, the southern tip of Greenland,
259 and the southwestern Irminger Sea. Warm extremes persisted in the vicinity of the ice edge on 5 March (**Figure 7d**), then the
260 large temperature anomalies (~15-16°C) expanded to cover much of the area from the Labrador Sea through Baffin Bay on 6-
261 7 March (**Figure 7e,f**). While the warm air mass appeared to propagate westward into northeastern Canada in the days that
262 followed, T2m anomalies remained above-average in these areas until colder air moved into the region on 10 March (**Figure**
263 **7g-i**).





Figure 7. Two-meter air temperature anomaly ($^{\circ}\text{C}$) from the ERA5 at 0 GMT on: a) 2 March, b) 3 March, c) 4 March, d) 5 March, e) 6 March, f) 7 March, g) 8 March, h) 9 March, and i) 10 March 2023. The anomalies are shown with respect to the period 16 February – 15 March 1979–2023. Grid points where the anomalies are less than the 1st percentile or greater than the 99th percentile based on the above period are indicated with the '+'. The blue curves represent the 0°C isotherm.

265 A warm air incursion into the Bering Sea was also apparent during this same time. From the 3rd to the 4th of March, the 0°C
 266 isotherm migrated several degrees northward as anomalously warm air penetrated into the Bering region (Figure 7b,c). The
 267 general southwest to northeast trajectory of the mild airstream was apparent in the days that followed. The 0°C isotherm entered
 268 the northeastern Bering Sea and southwestern Alaska on 5 March as anomalous melt along the ice edge continued, while
 269 temperatures over the western Bering Sea and northeastern Siberia remained below normal (Figure 7d,e). Air temperatures
 270 remained above average to extreme in western and northern Alaska during the days that followed as the airmass propagated
 271 into the high Arctic over 6-10 March (Figure 7f-i).

272
 273 Despite the 31 km resolution of the ERA5 fields, the array of synoptic maps makes it challenging to ascertain the extent of the
 274 temperature extremes, especially along coastal areas and along the approximate ice edges. The daily T2m fields are therefore
 275 supplemented with weather station time series to provide additional perspective on the air temperatures. During the Labrador
 276 Sea ice loss event, above-average air temperatures at Nuuk, Greenland to the southeast of the ice edge were recorded with
 277 >0°C daily mean temperatures from 2-7 March with warm air temperature extremes observed on 3-4 March (Figure 8a).





Figure 8. Weather station two-meter air temperature (T2m in °C) 1 January – 31 March 2023 daily time series (gray lines) overlapping the multi-sectoral melt event for a) Nuuk, b) Aasiaat, c) St. Paul, and d) Kotzebue. Considering all days from 1 January to 31 March for the respective stations' full periods of record (see Section 2.1), the mean T2m (black line), 1st percentile

(blue dashed line), and 99th percentile (red dashed line) are shown in each graphic. The sudden stratospheric warming event on 16 February 2023 is labelled with a green diamond, and to draw attention to the dates around the Labrador Sea and Bering Sea melt events, the period from 2-10 March 2023 is identified by yellow triangles. For reference the weather stations are overlaid on **Figure 1**.

Likewise, above-freezing, extreme air temperatures were observed in the GrIS lower ablation zone in the Qassimiut region (QAS_L) and on a glacier tangential to the Nuuk DMI station (NUK_K) during this period (**Figure S1a,b**). Meanwhile, in Aasiaat, Greenland roughly ~500 km north of Nuuk, the air temperatures were above-normal during this time, but were not above-freezing or considered extreme by the criteria used here (**Figure 8b**). Over the Bering Sea, St. Paul Island observed a stint of above-freezing temperatures that ranked near the 99th percentile for 4-7 March (**Figure 8c**), while Kotzebue on Alaska's northwest coast saw near- to slightly-above normal air temperatures during the Bering ice loss period but the airstream neither exceeded 0°C nor the 99th percentile criteria (**Figure 8d**).


279 Further analysis into the thermodynamic environment revealed that the anomalously warm airstreams advected over both the
280 Labrador and Bering regions possessed extreme water vapor content around the time of their respective melt peaks shown in
281 time series in Figure 9a,b and Figure 10a,b and in maps presented in Figure S2. During these peaks, both seas experienced
282 anomalous net and downwelling radiation in excess of the 95th percentile (Figures 9c,d and Figures 10c,d) with that energy
283 likely driving ice loss through melt. To further investigate the hydrometeorological nature of these airstreams the Guan and
284 Waliser (2019) atmospheric river (AR) detection algorithm was run separately for the Labrador Sea and Bering Sea domains
285 shown in **Figure 1**. Warm, moist conditions that overlapped these melt events were associated with AR activity (**Figure 11**).
286 An AR resided over the Labrador Sea for >40% of the day on 3-4 March, and its residence time was extreme on 5 March
287 (~60% of the day; **Figure 11a**). Moisture within this AR (**Figure 11c**) and total precipitation from the AR (**Figure S4**) were
288 both above-average, but not extreme. Meanwhile, daily AR residence time within the Bering Sea exceeded 40% on 4-7 March,
289 with an AR duration extreme (>60% of the day) on 5 March preceding the short-term melt peak on 6 March when IVT was
290 also extreme (**Figure 11b,d**). Extreme ERA5 daily precipitation associated with the AR intrusion fell in the Bering region
291 from 5-8 March (**Figure S4**). These persistent and anomalously warm and wet airmasses contributed to these tandem melt
292 extremes. Further analysis is ongoing to examine the full surface energy budget, including the role of rainfall, toward shaping
293 the observed melt.
294

- Deleted:** high
- Deleted:** , especially at
- Deleted:** (5 March and 6 March, respectively)
- Formatted:** Font: Bold
- Formatted:** Font: Bold
- Deleted:**)
- Formatted:** Superscript
- Formatted:** Font: Bold
- Formatted:** Font: Bold
- Deleted:** 9
- Deleted:** 9
- Deleted:** 9
- Deleted:** 9
- Deleted:** A
- Deleted:** appeared to have
- Deleted:** exchange processes
- Deleted:** that may have induced the



307
308
309
310
311
312
313
314
315

Figure 9. Time series (red curves) of ERA5: a) two-meter air temperature ($^{\circ}\text{C}$), b) total column water vapor (mm), c) net longwave radiation (W/m^2), and d) downward longwave radiation (W/m^2) averaged over the Labrador Sea region, indicated in Figure 1b, for the period January 1 to March 26, 2023. The black line represents the climatological mean value for the period 2000-2023 with shading incorporating values between the 5th and 95th percentiles. The ending date for the 4-day window with the largest change in sea ice concentration is shown with the dotted red line.



316
317

318 **Figure 10.** Time series (red curves) of ERA5: a) two-meter air temperature ($^{\circ}\text{C}$), b) total column water vapor (mm), c) net
 319 longwave radiation (W/m^2), and d) downward longwave radiation (W/m^2) averaged over the Bering Sea region, indicated in
 320 Figure 1e, for the period January 1 to March 26, 2023. The black line represents the climatological mean value for the period
 321 2000-2023 with shading incorporating values between the 5th and 95th percentiles. The ending date for the 4-day window with
 322 the largest change in sea ice concentration is shown with the dotted red line.

Deleted: ¶




Figure 11. March atmospheric river (AR) duration (% of day AR in domain) and intensity (kg/m/s) for the Labrador region (a,c) and Bering region (b,d), respectively (teal lines). The AR data are calculated over the same domains as shown in Figure 1. The thick black line in each panel represents the 1979-2023 mean. Considering all days from 1 February to 31 March for the respective regions for the 1979 to 2023 period, the 99th percentile (red dashed lines) are shown in all panels while the 1st percentile represents AR non-occurrence, and therefore is not marked in these plots. The sudden stratospheric warming event on 16 February 2023 is labelled with a green diamond, and to draw attention to the dates around the Labrador Sea and Bering Sea melt events, the period from 2-10 March 2023 is identified by yellow triangles.


4. Discussion and conclusions

324 Tandem, unusually-timed sea ice melt extremes in the Bering Sea and Labrador Sea occurred in early March 2023. The retreat
 325 of the ice edge in both marginal seas was similarly driven by the confluence of anomalous meteorological phenomena
 326 following an SSW event that occurred in mid-February. Below we discuss the ice loss events and focus on the attendant
 327 atmospheric mechanisms that provided thermodynamic support for their occurrence.

328

329 4.1 Perspectives on ice losses during the maximum and supporting atmospheric processes

330 Amidst the decline of winter season ice coverage and thickness in the warming Arctic, the latitude of the ice edge can vary on
 331 daily timescales due to wind and melt-driven processes. However, the probability curves shown in Figure 3 suggest that such



Deleted:

Deleted: 9

Deleted: Greenland

Deleted: Alaska

332 short-term March 2023 sea ice losses in either the Bering or Labrador regions, taken independently, qualify as extreme events.
333 Both the magnitude of losses and the unusual timing of their anomalous occurrence aligned with the climatological Arctic sea
334 ice maximum may further qualify these melt extremes collectively as a rare synoptic ice loss event. We do not assess ice edge
335 changes in other marginal seas during the March historical record to establish whether other areas participated in this event.

336
337 The anatomy of the melt extremes can be described by a confluence of anomalous atmospheric phenomena that simultaneously
338 occurred over the Bering Sea and Labrador Sea. The melt period was preceded by an SSW event that led to a shift in the large-
339 scale mid-tropospheric circulation regime over the polar cap as evidenced by the rapid transition over two weeks from strong
340 positive to negative NAO conditions and lower to higher mid-tropospheric air pressure over the high Arctic. The noted shift
341 to negative NAO followed by the development of a Greenland block that supported southerly winds and warm advection across
342 the Labrador Sea following a SSW has been documented in previous studies (e.g., Charlton-Perez et al., 2018; Domeisen,
343 2019; Domeisen and Butler, 2020). While the set-up of the Greenland block is not unique to this event, its magnitude for the
344 time of year is remarkable as shown by the extremes highlighted in the GBI time series (**Figure 5c**) and z500 spatial plots
345 (**Figure 6**).

346
347 SSWs on average tend to elicit a weaker atmospheric dynamical response over the Bering region than the Labrador Sea. Smith
348 et al. (2018) analyzed data from the Whole Atmosphere Community Climate Model of NCAR's Community Earth System
349 Model and found that over the 40 days following SSW onset there were minimal sea-level pressure (SLP) changes over the
350 Bering Sea and greater Alaska, but there were large, positive SLP anomalies located northward and eastward of these areas
351 including around Greenland. Across SSW winters (JFM), the authors also found similar SLP signatures over Greenland, but
352 negative SLP anomalies and northerly winds over Alaska and the Bering Sea. The interpretation of the large-scale circulation
353 pattern is complicated by the La Niña phase that prevailed during winter 2023 (**Figure S3a**). La Niña favors North Pacific
354 ridging into the Gulf of Alaska with low pressure across the Bering Strait and Pacific Arctic (**Figure S3b**). However, compared
355 to the average conditions that occur after SSWs during La Niña winters, March 2023 was marked by a relatively stronger
356 North Pacific ridge that extended further north over Bering Sea and mainland Alaska. These anomalously high mid-
357 tropospheric heights, reflected by an increase in the ABI (**Figure 5d**) and shown in the positive z500 anomaly maps from 5-
358 10 March (**Figure 6d-i**), drove warm advection that caused Bering Sea ice to melt. In addition to the SSW event and La Niña
359 phase, factors such as internal variability of the climate system and air-sea interactions over the North Pacific Ocean may have
360 played a role in inducing the anomalously strong mid-tropospheric ridge extending from Greenland to Alaska.

361
362 In both the Labrador Sea and Bering Sea, anomalous atmospheric circulation characteristics, namely the stationary, extreme
363 blocking anticyclones, supported southerly advection of above-normal to extremely warm and moist air that led to these
364 thermodynamically-driven melt events (e.g., **Figures 7, 9-11, and Figure S2**). Additional investigation of the airstreams
365 revealed that anomalous ARs were present in both regions during this time and played a critical role in the simultaneous melt

Deleted: -to-extreme

Deleted: Figure

Formatted: Font: Not Bold

Deleted: ,

Deleted: and such events are known to slow sea ice growth in Arctic winter (Zhang et al., 2023). These ARs were unusual for the time of year,

Deleted: thus may have

Deleted: unexpected timing of

374 ~~extremes~~. The extreme duration of the AR over the peak Labrador Sea melt and extreme duration and intensity immediately
375 preceding the Bering Sea melt, both on 5 March, likely ~~enhanced~~ downwelling longwave energy transfer into the ice, causing
376 its short-term, yet remarkable, decline. ~~Past studies have likewise identified downward longwave radiative flux during AR~~
377 ~~passage as a key process that tends to decrease ice mass balance during summer (Mattingly et al., 2018; Wille et al., 2019;~~
378 ~~Francis et al., 2020) and slows wintertime ice growth (Hegyi and Taylor, 2018; Zhang et al., 2023). Here, we document the~~
379 ~~role of anomalous large-scale meteorological characteristics, including tandem AR events, that drove unprecedented and~~
380 ~~concurrent~~ sea ice melt at a time of year characterized by maximum ice extent.

381

382 4.2 Additional considerations emanating from this case study

383 This rare ice loss event ~~concurrently~~ encompassing the Bering Sea and Labrador Sea was shaped by a confluence of synoptic
384 extremes that aligned in time to induce thermodynamic melt of the sea ice edge. We look at this ice loss from a thermodynamic
385 perspective, though concede that in addition to supporting melt that southerly winds could have induced some sea ice
386 compaction in the marginal ice zones through convergence. If this event was examined through a sea ice budget lens, we
387 acknowledge that producing estimates of ice dynamical processes, such as wind-driven convergence and divergence, would
388 be important to gain a more complete understanding of the evolution of mechanisms responsible for these regional ice losses.
389 Follow-on work will take a broader view of thermodynamic processes, which may provide additional insight into ice loss
390 mechanisms elucidated in this case study. For example, resolution of the sea ice types and surface energy balance before,
391 during, and after the melt event may provide perspectives on ice-air interactions that shaped it.

392

393 Related to the surface energy balance processes, further analyses will delve deeper into the roles of latent heating and humidity
394 fluxes in shaping the ice melt event. Rainfall (<1 mm) was observed during 2-3 March in the rain gauges at the Nuuk and
395 Aasiaat DMI weather stations, and, if it were not for sporadic station outages from 2-10 March, rain on other days during this
396 period may have been documented (C. Drost Jensen 2024, personal communication). Nearby, separate near-coastal weather
397 stations maintained by Asiaq Greenland Survey also documented small amounts (<1 mm) of rainfall at Nuuk and Kobbefjord
398 (A. Ginnerup 2024, personal communication). Meanwhile, terrestrial weather stations at Kotzebue and Nome, Alaska, ~300
399 km to the southwest, saw >25 mm of cumulative rainfall during 4-6 March, which are 3-day total precipitation records for both
400 weather stations in March (R. Thoman 2023, personal communication). Spatial patterns of ERA5 total precipitation over this
401 period are consistent with these observations (Figure S4). In addition to rain measurements near the coast, rain on cold snow
402 was also detected in weather station observations found in the southwestern GrIS accumulation zone, which is rare for the time
403 of year (J. Box 2024, personal communication). Further diagnostic evaluation is needed to determine the extent, frequency,
404 amount, and impacts of rainfall on the cold snow cover on the GrIS and sea ice during this period. Thus, follow-on studies of
405 the surface energy exchange processes and precipitation characteristics may help to broaden our perspective of this complex
406 extreme event.

407

Deleted: contributed to

Deleted: Thus, the near-extreme to

Deleted: extreme

Deleted: regional circulation patterns (GBI and ABI) appear to have supported the advection of

Deleted: extremely warm and moist airstreams

Deleted: significant

Deleted: both

Deleted: simultaneously

417 It is clear from recent years that there are occurrences of a variety of extreme Arctic events that vary in location, season, and
418 type which meet or exceed previous records (Walsh et al. 2020). Philosophically, it is difficult to project let alone interpret the
419 future frequency of these events without detailed historical analogues. It has been proposed that the recent increase of Arctic
420 extremes is due to an overlap of steadily increasing Arctic warming that is constructively superimposed on the natural range
421 of atmospheric and oceanic dynamics, e.g., jet stream meanders, atmospheric blocking, storms, and upper-ocean heat content
422 (Overland 2022), which could themselves, at least in some cases, be influenced by anthropogenic global heating. This is
423 certainly the case with the concurrent examples from the Labrador Sea and Bering Sea in March 2023. Whether this extreme
424 event foreshadows a more frequent occurrence of similar events in the future is an open but intriguing question that merits
425 careful future investigation.

426
427 *Data availability.* Alaska weather station data are available from <https://xmacis.rcc-acis.org>. Greenland coastal weather station
428 records were obtained from Caroline Drost Jensen (DMI). PROMICE observations are from
429 <https://dataverse.geus.dk/dataset.xhtml?persistentId=doi:10.22008/FK2/IW73UU>. The NAO index was downloaded from
430 <https://www.cpc.ncep.noaa.gov/products/precip/CWlink/pna/nao.shtml>. ERA5 reanalysis fields are obtained from the
431 Copernicus Climate Data Store at [https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-](https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels?tab=overview)
432 [levels?tab=overview](https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels?tab=overview). Sea ice data are downloaded from NSIDC at <https://nsidc.org/data/g02202/versions/4>. The SSW
433 Compendium can be found at <https://csl.noaa.gov/groups/csl8/sswcompendium/majorevents.html>.

434
435 *Author contributions.* T.B. and G.W.K.M. conceived the study with input from Q.D., A.H.B., J.E.O., R.L.T., I.B., Z.L., and
436 E.H. as the study developed. R.L.T. provided assistance with data acquisition. All authors provided feedback on draft iterations
437 of the paper.

438
439 *Competing Interests.* None.

440
441 *Acknowledgements.* T.J.B. and Q.D. were funded by NSF Arctic System Science awards 2246600 and 2246601, respectively.
442 J.E.O. is supported by NOAA's GOMO Arctic Research Program. PMEL contribution #53XX. GWKM was funded by the
443 Natural Sciences and Engineering Research Council of Canada. E.H. was supported by NERC NE/W005875/1. The authors
444 wish to thank Caroline Drost Jensen and Anders Ginnerup for assistance obtaining and interpreting the Greenland weather
445 station records. We also thank numerous scientists for discussions on topics related to high-latitude precipitation and GrIS
446 meteorology, including Jason Box, Jakob Abermann, Matthew Sturm, and Melinda Webster.

447 **References**

- 448 [AghaKouchak, A., Chiang, F., Huning, L.S., Love, C.A., Mallakpour, I., Mazdiyasni, O., Moftakhari, H., Papalexiou, S.M.,](#)
449 [Ragno, E., and Sadegh, M.: Climate Extremes and Compound Hazards in a Warming World. *Ann. Rev. Earth Planet. Sci.*, **48**,](#)
450 <https://doi.org/10.1146/annurev-earth-071719-055228>, 2020.
- 451
- 452 Baldwin, M.P., Ayarzagüena, B. Birner, T., Butchart, N., Butler, A.H., Charlton-Perez, A.J., Domeisen, D.I.V., Garfinkel,
453 C.I., Garny, H., Gerber, E.P., Hegglin, M.I., Langematz, U., and Pedatella, N.M.: Sudden Stratospheric Warmings. *Rev.*
454 *Geophys.*, **59**, e2020RG000708, <https://doi.org/10.1029/2020RG000708>, 2020.
- 455
- 456 Bailey, H., Hubbard, A., Klein, E.S., Mustonen, K.-R., Akers, P.D., Marttila, H., and Welker, J.M.: Arctic sea-ice loss fuels
457 extreme European snowfall. *Nat. Geosci.*, **14**, 283-288, <https://doi.org/10.1038/s41561-021-00719-y>, 2021.
- 458
- 459 Ballinger, T.J., Hanna, E., Hall, R.J., Cropper, T.E., Miller, J., Ribergaard, M.H., Overland, J.E., and Høyer, J.L.: Anomalous
460 blocking over Greenland preceded the 2013 extreme early melt of local sea ice. *Ann. Glaciol.*, **59**, 181-190,
461 <https://doi:10.1017/aog.2017.30>, 2018.
- 462
- 463 Ballinger, T.J., Walsh, J.E., Alexeev, V.A., Bieniek, P.A., McLeod, J.T.: The Alaska Blocking Index, version 2: Analysis and
464 covariability with statewide and large-scale climate from 1948-2020. *Int. J. Climatol.*, **42**, 9767-9787,
465 <https://doi:10.1002/joc.7864>, 2022.
- 466
- 467 Ballinger, T.J., Overland, J.E., Wang, M., Bhatt, U.S., Brettschneider, B., Hanna, E., Hanssen-Bauer, I., Kim, S.-J., Thoman,
468 R.L., and Walsh, J.E.: Surface air temperature [in “State of the Climate in 2022”]. *Bull. Am. Meteor. Soc.*, **104**, S279-S281,
469 <https://doi:10.1175/10.1175/BAMS-D-23-0079.1>, 2023.
- 470
- 471 Barnston, A.G., and Livezey, R.E.: Classification, seasonality, and persistence of low-frequency atmospheric circulation
472 patterns. *Mon. Wea. Rev.*, **115**, 1083-1126, [https://doi.org/10.1175/1520-0493\(1987\)115<1083:CSAPOL>2.0.CO;2](https://doi.org/10.1175/1520-0493(1987)115<1083:CSAPOL>2.0.CO;2), 1987.
- 473
- 474 Blanchard-Wrigglesworth, E., Webster, M., Boisvert, L., Parker, C., and Horvat, C.: Record Arctic cyclone of January 2022:
475 Characteristics, impacts, and predictability. *J. Geophys. Res. – Atmos.*, **127**, e2022JD037161,
476 <https://doi.org/10.1029/2022JD037161>, 2022.
- 477
- 478 Boisvert, L., Parker, C., and Valkonen, E.: A warmer and wetter Arctic: Insights from a 20-years AIRS record. *J. Geophys.*
479 *Res. – Atmos.*, **128**, e2023JD038793, <https://doi.org/10.1029/2023JD038793>, 2023.

480
481 Butler, A.H., Sjoberg, J.P., Seidel, D.J., and Rosenlof, K.H.: A sudden stratospheric warming compendium. *Earth Syst. Sci.*
482 *Data*, 9, 63-76, <https://doi.org/10.5194/essd-9-63-2017>, 2017.
483
484 Cavalieri, D., Gloersen, P., and Campbell, W.J.: Determination of sea ice parameters with the NIMBUS-7 SMMR. *J. Geophys.*
485 *Res.*, 89 (D4), 5355-5369, <https://doi.org/10.1029/JD089iD04p05355>, 1984.
486
487 Charlton, A.J., and Polvani, L.M.: A new look at Stratospheric Sudden Warmings: Part I: Climatology and modeling
488 benchmarks. *J. Climate*, 20, 449-469, <https://doi.org/10.1175/JCLI3996.1>, 2007.
489
490 Charlton-Perez, A.J., Ferranti, L., and Lee, R.W.: The influence of the stratospheric state on North Atlantic weather regimes.
491 *Q.J.R. Meteorol. Soc.*, 144, 1140-1151, <https://doi.org/10.1002/qj.3280>, 2018.
492
493 Collow, A.B.M., Shields, C.A., Guan, B., Kim, S., Lora, J.M., McClenny, E.E., Nardi, K., Payne, A., Reid, K., Shearer, E.J.,
494 Tomé, R., Wille, J.D., Ramos, A.M., Gorodetskaya, I.V., Leung, L.R., O'Brien, T.A., Ralph, F.M., Rutz, J., Ullrich, P.A.,
495 Wehner, M.: An overview of ARTMIP's Tier 2 Reanalysis Intercomparison: Uncertainty in the detection of atmospheric rivers
496 and their associated precipitation. *J. Geophys. Res. – Atmos.*, 127, e2021JD036155, <https://doi.org/10.1029/2021JD036155>,
497 2022.
498
499 Comiso, J.C.: Characteristics of Arctic winter sea ice from satellite multispectral microwave observations. *J. Geophys. Res. –*
500 *Oceans*, 91 (C1), 975-994, <https://doi.org/10.1029/JC091iC01p00975>, 1986.
501
502 Domeisen, D.I.V.: Estimating the frequency of sudden stratospheric warming events from surface observations of the North
503 Atlantic Oscillation. *J. Geophys. Res. Atmos.*, 124, 3180-3194, <https://doi.org/10.1029/2018JD030077>, 2019.
504
505 Domeisen, D.I.V., and Butler, A.H.: Stratosphere drivers of extreme events at the Earth's surface. *Comm. Earth Environ.*, 1,
506 59, <https://doi.org/10.1038/s43247-020-00060-z>, 2020.
507
508 Fausto, R. S., van As, D., Mankoff, K. D., Vandecrux, B., Citterio, M., Ahlström, A. P., Andersen, S. B., Colgan, W., Karlsson,
509 N. B., Kjeldsen, K. K., Korsgaard, N. J., Larsen, S. H., Nielsen, S., Pedersen, A. Ø., Shields, C. L., Solgaard, A. M., and Box,
510 J. E.: Programme for Monitoring of the Greenland Ice Sheet (PROMICE) automatic weather station data, *Earth Syst. Sci. Data*,
511 13, 3819–3845, <https://doi.org/10.5194/essd-13-3819-2021>, 2021.
512

513 Francis, J.A., and Vavrus, S.J.: Evidence for a wavier jet stream in response to rapid Arctic warming. *Environ. Res. Lett.*, 10,
514 014005, <https://doi.org/10.1088/1748-9326/10/1/014005>, 2015.

515

516 [Francis, D., Mattingly, K.S., Temmi, M., Massom, R., and Heil, P.: On the crucial role of atmospheric rivers in the two major](#)
517 [Weddell Polynya events in 1973 and 2017 in Antarctica. *Sci. Adv.*, 6, <https://doi.org/10.1126/sciadv.abc2695>, 2020.](#)

518

519 Graham, R.M., Hudson, S.R., and Matarilli, M.: Improved performance of ERA5 in Arctic gateway relative to four global
520 atmospheric reanalyses. *Geophys. Res. Lett.*, 46, 6138-6147, <https://doi.org/10.1029/2019GL082781>, 2019.

521

522 Guan, B., and Waliser, D.E.: Tracking atmospheric rivers globally: spatial distributions and temporal evolution of life cycle
523 characteristics. *J. Geophys. Res.*, 124, 12523–12552, <https://doi.org/10.1029/2019JD031205>, 2019.

524

525 Hanna, E., Jones, J.M., Cappelen, J., Mernild, S.H., Wood, L., Steffen, K., and Huybrechts, P.: The influence of North Atlantic
526 atmospheric and oceanic forcing effects on 1900-2010 Greenland summer climate and ice melt/runoff. *Int. J. Climatol.*, 33,
527 862-880, <https://doi.org/10.1002/joc.3475>, 2013.

528

529 Hanna, E., Cropper, T.E., Jones, P.D., Scaife, A.A., and Allan, R.: Recent seasonal asymmetric changes in the NAO (a marked
530 summer decline and increased winter variability) and associated changes in the AO and Greenland Blocking Index. *Int. J.*
531 *Climatol.*, 35, 2540-2554, <https://doi.org/10.1002/joc.4157>, 2015.

532

533 Hanna, E., Hall, R.J., Cropper, T.E., Ballinger, T.J., Wake, L., Mote, T., and Cappelen, J.: Greenland blocking index daily
534 series 1851-2015: Analysis of changes in extremes and links with North Atlantic and UK climate variability and change. *Int.*
535 *J. Climatol.*, 38, 3546-3564, <https://doi.org/10.1002/joc.5516>, 2018.

536

537 [Hegyi, B.M., and Taylor, P.C.: The unprecedented 2016-2017 Arctic sea ice growth season: The critical role of atmospheric](#)
538 [rivers and longwave fluxes. *Geophys. Res. Lett.*, 45, 5204-5212, <https://doi.org/10.1029/2017GL076717>, 2018.](#)

539

540 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radum, R.,
541 Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M.,
542 De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A.,
543 Haimberger, L., Healy, S., Hogan, R.J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G.,
544 de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 Global Reanalysis. *Q.J. R. Meteorol.*
545 *Soc.*, 146, 1999-2049, <https://doi.org/10.1002/qj.3803>, 2020.

546

547 Kirbus, B., Tiedeck, S., Camplani, A., Chylik, J., Crewell, S., Dahlke, S., Ebell, K., Gorodetskaya, I., Griesche, H., Handorf,
548 D., Höschel, I., Lauer, M., Neggers, R., Rückert, J., Shupe, M.D., Spreen, G., Walbröl, A., Wendisch, M. and Rinke, A.:
549 Surface impacts and associated mechanisms of a moisture intrusion into the Arctic observed in mid-April 2020 during
550 MOSAiC. *Front. Earth Sci.*, 11, 1147848, <https://doi.org/10.3389/feart.2023.1147848>, 2023.

551
552 [Kornhuber, K., Coumou, D., Vogel, E., Lesk, C., Donges, J.F., Lehmann, J., and Horton, R.M.: Amplified Rossby waves](#)
553 [enhance risk of concurrent heatwaves in major breadbasket regions. *Nature Clim. Change*, 10, 48-53,](#)
554 <https://doi.org/10.1038/s41558-019-0637-z>, 2020.

555
556 Lee, S.H., and Butler, A.H.: The 2018-2019 Arctic stratospheric polar vortex. *Weather*, 75, 52-57,
557 <https://doi.org/10.1002/wea.3643>, 2019.

558
559 Lee, S.H., Tippett, M.K., and Polvani, L.M.: A new year-round weather regime classification for North America. *J. Clim.*, 36,
560 7091-7108, <https://doi.org/10.1175/JCLI-D-23-0214.1>, 2023.

561
562 [Mattingly, K.S., Mote, T.L., and Fettweis, X.: Atmospheric river impacts on the Greenland Ice Sheet surface mass balance. *J.*](#)
563 [Geophys. Res. – Atmos.](#), 123, 8538-8560, <https://doi.org/10.1029/2018JD028714>, 2018.

564
565 Meier, W.N., Fetterer, F., Windnagel, A.K., and Stewart, J.S.: NOAA/NSIDC Climate Data Record of Passive Microwave Sea
566 Ice Concentration, Version 4. NSIDC: National Snow and Ice Data Center. <https://doi.org/10.7265/efmz-2t65>, 2021.

567
568 Meier, W.N., Stewart, J. S., Windnagel, A., and Fetterer, F. M.: Comparison of hemispheric and regional sea ice extent and
569 area trends from NOAA and NASA passive microwave-derived climate records. *Remote Sens.*, 14, 619.
570 <https://doi.org/10.3390/rs14030619>, 2022.

571
572 Meier, W.N., Petty, A., Hendricks, S., Kaleschke, L., Divine, D., Farrell, S., Gerland, S. Perovich, D., Ricker, R., Tian-Kunze,
573 X., and Webster, M.: Sea Ice. NOAA Arctic Report Card 2023, R.L. Thoman, T.A. Moon, and M.L. Druckenmiller, Eds.,
574 <https://doi.org/10.25923/f5t4-b865>, 2023.

575
576 Moore, G.W.K.: The December 2015 North Pole warming event and the increasing occurrence of such events. *Sci. Rep.*, 6,
577 39084, <https://doi.org/10.1038/srep39084>, 2016.

578
579 Moore, G.W.K., Schweiger, A., Zhang, J., and Steele, M.: What caused the remarkable February 2018 north Greenland
580 polynya? *Geophys. Res. Lett.*, 45, 1342-13350, <https://doi.org/10.1029/2018GL080902>, 2018.

581
582 Nygård, T., Naakka, T., and Vihma, T.: Horizontal moisture transport dominates the regional moistening patterns in the Arctic.
583 *J. Clim.*, 33, 6793-6807, <https://doi.org/10.1175/JCLI-D-19-0891.1>, 2020.
584
585 Overland, J.E., Ballinger, T.J., Cohen, J., Francis, J.A., Hanna, E., Jaiser, R., Kim, B.-K., Kim, S.-J., Ukita, J., Vihma, T.,
586 Wang, M., and Zhang, X.: How do intermittency and simultaneous processes obfuscate the Arctic influence on midlatitude
587 winter extreme weather events? *Environ. Res. Lett.*, 16, 043002, <https://doi.org/10.1088/1748-9326/abdb5d>, 2021.
588
589 Overland, J.E.: Arctic climate extremes. *Atmos.*, 13, 1670, <https://doi.org/10.3390/atmos13101670>, 2022.
590
591 Papritz, L., Hauswirth, D., and Hartmuth, K.: Moisture origin, transport pathways, and driving processes of intense wintertime
592 moisture transport into the Arctic. *Weather Clim. Dyn.*, 3, 1-20, <https://doi.org/10.5194/wcd-3-1-2022>, 2022.
593
594 Serreze, M.C., and Barry, R.G. Processes and impacts of Arctic amplification: A research synthesis. *Glob. Planet. Change*, 77,
595 85-96, <https://doi.org/10.1016/j.gloplacha.2011.03.004>, 2011.
596
597 Smith, K.L., Polvani, L.M., and Tremblay, L.B.: The impact of stratospheric circulation extremes on minimum Arctic sea ice
598 extent. *J. Climate*, 31, 7169-7183, <https://doi.org/10.1175/JCLI-D-17-0495.1>, 2018.
599
600 Stroeve, J., and Notz, D.: Changing state of Arctic sea ice across all seasons. *Environ. Res. Lett.*, 13, 103001,
601 <https://doi.org/10.1088/1748-9326/aade56>, 2018.
602
603 Tachibana, Y., Komatsu, K.K., Alexeev, V.A., and Ando, Y.: Warm hole in Pacific Arctic sea ice cover forced mid-latitude
604 Northern Hemisphere cooling during winter 2017-18. *Sci. Rep.*, 9, 5567, <https://doi.org/10.1038/s41598-019-41682-4>, 2019.
605
606 Walsh, J.E., Ballinger, T.J., Euskirchen, E.S., Hanna, E., Mård, J., Overland, J.E., Tangen, H., and Vihma, T.: Extreme weather
607 and climate events in northern areas: A review. *Earth-Sci. Rev.*, 209, 103324, <https://doi.org/10.1016/j.earscirev.2020.103324>,
608 2020.
609
610 [Wille, J.D., Favier, V., Dufour, A., Gorodetskaya, I.V., Turner, J., Agosta, C., and Codron, F.: West Antarctic surface melt](#)
611 [triggered by atmospheric rivers. *Nature Geosci.*, 12, 911-916, <https://doi.org/10.1038/s41561-019-0460-1>, 2019.](#)
612
613 [Zhang, P., Wu, Y., Simpson, I.R., Smith, K.L., Zhang, X., De, B., and Callaghan, P.: A stratospheric pathway linking a colder](#)
614 [Siberia to Barents-Kara sea ice loss. *Sci. Adv.*, 4, eaat6025, <https://doi.org/10.1126/sciadv.aat6025>.](#)

615
616 Zhang, P., Chen, G., Ting, M., Leung, L.R., Guan, B., and Li, L.: More frequent atmospheric rivers slow the seasonal recovery
617 of Arctic sea ice. *Nature Clim. Change*, 13, 266-273, <https://doi.org/10.1038/s41558-023-01599-3>, 2023.
618
619 [Zhou, S., Yu, B., and Zhang, Y.: Global concurrent climate extremes exacerbated by anthropogenic climate change. *Sci. Adv.*,
620 9, eabo1638, <https://doi.org/10.1126/sciadv.abo1638>, 2023.](#)
621
622 [Zscheischler, J., Westra, S., van den Hurk, B.J.J.M., Seneviratne, S.I., Ward, P.J., Pitman, A., AghaKouchak, A., Bresch, D.N.,
623 Leonard, M., Wahl, T., and Zhang, X.: Future climate risk from compound events. *Nature Clim. Change*, 8, 469-477,
624 <https://doi.org/10.1038/s41558-018-0156-3>, 2018.](#)
625