

Position-specific kinetic isotope effects for nitrous oxide: A new expansion of the Rayleigh model

Elise D. Rivett^{1,2,*}, Wenjuan Ma³, Nathaniel E. Ostrom^{2,4,*}, Eric L. Hegg^{1,2,*}

¹ Department of Biochemistry & Molecular Biology, Michigan State University, East Lansing, Michigan 48824, United States

5 ²DOE Great Lakes Bioenergy Research Center, Michigan State University, East Lansing, MI, 48824, United States

³Center for Statistical Training and Consulting, Michigan State University, East Lansing, MI, 48824, United States

⁴Department of Integrative Biology, Michigan State University, East Lansing, Michigan 48824, United States

Correspondence to: Elise D. Rivett (rivettel@msu.edu), Nathaniel E. Ostrom (ostromn@msu.edu), Eric L. Hegg (erichegg@msu.edu)

10 **Abstract.** Nitrous oxide (N₂O) is a potent greenhouse gas and the most significant anthropogenic ozone-depleting substance currently being emitted. A major source of anthropogenic N₂O emissions is the microbial conversion of fixed nitrogen species from fertilizers in agricultural soils. Thus, understanding the enzymatic mechanisms by which microbes produce N₂O has environmental significance. Measurement of the ¹⁵N/¹⁴N isotope ratios of N₂O produced by purified enzymes or axenic microbial cultures is a promising technique for studying N₂O biosynthesis. Typically, N₂O-producing enzymes combine
15 nitrogen atoms from two identical substrate molecules (NO or NH₂OH). Position-specific isotope analysis of the central (N^α) and outer (N^β) nitrogen atoms in N₂O enables the determination of the individual kinetic isotope effects (KIEs) for N^α and N^β, providing mechanistic insight into the incorporation of each nitrogen atom. Previously, position-specific KIEs (and fractionation factors) were quantified using the Rayleigh distillation equation, *i.e.*, via linear regression of δ¹⁵N^α or δ¹⁵N^β against [-lnf/(1-f)], where f is the fraction of substrate remaining in a closed system. This approach, however, is inaccurate
20 for N^α and N^β because it does not account for fractionation at N^α affecting the isotopic composition of substrate available for incorporation into the β position (and vice versa). Therefore, we developed a new expansion of the Rayleigh model that includes specific terms for fractionation at the individual N₂O nitrogen atoms. By applying this Expanded Rayleigh model to a variety of simulated N₂O synthesis reactions with different combinations of normal/inverse/no KIEs at N^α and N^β, we demonstrate that our new model is both accurate and robust. We also applied this new model to two previously published
25 datasets describing N₂O production from NH₂OH oxidation in a methanotroph culture (*Methylosinus trichosporium*) and N₂O production from NO by a purified *Histoplasma capsulatum* (fungal) P450 NOR, demonstrating that the Expanded Rayleigh model is a useful tool in calculating position-specific fractionation for N₂O synthesis.

1 Introduction

Nitrous oxide (N₂O) is a greenhouse gas with a global warming potential approximately 300 times greater than that of CO₂
30 (100-year time horizon) Stocker (2013). In addition, N₂O is the primary source of stratospheric NO_x and therefore is a

significant ozone-depleting substance (Ravishankara et al., 2009). Levels of atmospheric N₂O are increasing by an average of ~0.29% annually (Lan et al., 2022), primarily due to rising anthropogenic N₂O emissions, which comprise approximately 40% of annual global N₂O emissions (Tian et al., 2020). In the US, at least 70% of annual anthropogenic N₂O emissions result from agricultural practices, such as the application of nitrogen-containing fertilizer (*e.g.*, ammonium and nitrate) to the field. Soil microbes can metabolize these nitrogen-containing compounds through processes such as nitrification and denitrification, producing N₂O either as a side-product (nitrification), intermediate (complete denitrification), or end-product (incomplete denitrification) (Kuypers et al., 2018). Understanding the underlying mechanisms of N₂O production for each metabolic pathway is critical for tracing and ultimately mitigating environmental N₂O emissions.

Many different types of enzymes catalyze N₂O formation in microbes. Examples of N₂O-producing enzyme families include cytochrome P460 (Caranto et al., 2016) and hydroxylamine oxidoreductase (Yamazaki et al., 2014) in nitrifiers, “bacterial” nitric oxide reductases (cNOR, qNOR, etc.) found in bacterial/archaeal denitrifiers, and fungal nitric oxide reductase (P450 NOR) from fungal denitrifiers (Yamazaki et al., 2014; Yang et al., 2014; Lehnert et al., 2018) as well as detoxifying enzymes such as flavodiiron nitric oxide reductases (Romão et al., 2016). Distinguishing among different sources of N₂O in the environment and identifying the relative contribution of each source is challenging and requires a combination of analytical methods. One useful technique for N₂O source apportionment is **stable isotope analysis**. This method takes advantage of the fact that many physical and chemical processes, such as enzyme-catalyzed reactions, generally have slightly different rate constants for different isotopes of the same element (*e.g.*, ¹⁴N and ¹⁵N). In an isotopically sensitive process, the isotope that reacts faster will be more prevalent (enriched) in the product relative to the substrate. Different processes generally have distinct patterns of isotopic enrichment and thus have unique isotopic signatures.

Stable isotope analysis is particularly useful for studying N₂O synthesis because the central (N^α) and outer (N^β) nitrogen atoms in N₂O are non-equivalent and non-exchangeable, and the isotope ratios of these two nitrogen atoms can be determined individually (Toyoda and Yoshida, 1999). These position-specific isotope ratios can aid in identifying which enzyme(s) produce N₂O in microbial cultures under varying growth conditions (Sutka et al., 2003, 2004; Sutka et al., 2006; Sutka et al., 2008; Toyoda et al., 2005; Frame and Casciotti, 2010; Yamazaki et al., 2014; Haslun et al., 2018). Additionally, because isotopic enrichment is typically dependent on the rate-limiting step(s) of the enzyme-catalyzed reaction, analyzing the isotopic signatures of individual enzymes can provide valuable insights into each enzyme’s catalytic mechanism.

Isotopic enrichment for a specific process can be quantified by determining a parameter known as the fractionation factor ($\alpha_{p/s}$). For reactions with first-order kinetics, $\alpha_{p/s}$ is equal to the rate constant of the heavy isotope (k_H) divided by the rate constant for the light isotope (k_L). Because $\alpha_{p/s}$ values are typically close to 1, $\alpha_{p/s}$ is often converted to its per mil equivalent, the enrichment factor ($\epsilon_{p/s}$) (Mariotti et al., 1981):

$$\epsilon_{p/s} = (\alpha_{p/s} - 1) * 1000 \quad (1)$$

Notably, $\alpha_{p/s}$ is also equal to the inverse of the kinetic isotope effect (KIE = k_L/k_H). The KIE and fractionation factor values for a specific reaction describe the isotopic preference of that reaction. Most commonly, the light isotope reacts faster than the

Commented [ER1]: Reviewer 1 comment (L46): Edited text as suggested

Deleted: compound-specific

Deleted: , or the measurement of isotope ratios in a particular chemical compound

Commented [ER2]: Community comment: Bigeleisen and Wolfsberg, 1958 ref. is replaced with the Mariotti et al., 1981 ref. Bigeleisen and Wolfsberg, 1958 is already cited in section 2.2.

Deleted: (Mariotti et al., 1981)

Commented [ER3]: It is also appropriate to cite Mariotti here.

heavy isotope (*i.e.*, $KIE > 1$), which is designated as a normal KIE. Conversely, when the heavy isotope reacts faster ($KIE < 1$), the reaction is said to have an inverse KIE. If the heavy and light isotopes react at the same rate ($KIE = 1$), the reaction has no kinetic isotope effect.

Mariotti and colleagues (1981) derived an approximation of the Rayleigh distillation equation (hereafter referred to as the (standard) Rayleigh equation) that is frequently used to determine enrichment factors in closed systems (Eq. (2)) (Mariotti et al., 1981).

$$\delta^{15}N^p = \delta^{15}N^{s0} - \epsilon_{p/s} * \frac{f \ln(f)}{1-f} \quad (2)$$

Here, $\delta^{15}N^p$ is the δ value of accumulated product (p) at time t , $\delta^{15}N^{s0}$ is the δ value of the substrate at $t = 0$, and f is the fraction of substrate remaining at time t ($f = N^s/N^{s0}$). (The δ values are directly related to isotope ratios; see “Isotope nomenclature” below.) Notably, the Rayleigh equation is based on the assumption that the element of interest is transferred from the substrate to a single position (or equivalent positions) in the product.

The Rayleigh equation (Eq. (2)) holds true for $\delta^{15}N^{bulk}$ (the average of $\delta^{15}N^{\alpha}$ and $\delta^{15}N^{\beta}$) because the overall reaction can be thought of as the unidirectional conversion of two identical substrate molecules (*e.g.*, two NO or two NH_2OH molecules) to one product molecule with two nitrogen atoms ($2S \rightarrow P_2$). In the absence of side reactions, this means that $\delta^{15}N^{bulk}$ can be calculated at any point during the reaction (*i.e.*, for any value of f) as long as the δ value of the initial substrate ($\delta^{15}N^{s0}$) and the enrichment factor (ϵ_{N-bulk}) are known. In practice, the ϵ_{N-bulk} value for a specific enzymatic reaction is found by plotting $\delta^{15}N^{bulk}$ versus $[-\ln f/(1-f)]$. Because it is assumed that changes in $\delta^{15}N^{bulk}$ are due solely to the bulk kinetic isotope effect of the reaction in question, the slope of this plot (determined by linear regression) is the bulk enrichment factor (ϵ_{N-bulk}), and the y-intercept corresponds to $\delta^{15}N^{s0}$.

Unfortunately, the relationship between $\delta^{15}N^{\alpha}$, $\delta^{15}N^{\beta}$, and their corresponding enrichment factors is more complex because the two N atoms are non-exchangeable and both steps that incorporate NO (or NH_2OH) are potentially isotopically sensitive. *In other words, fractionation at the α position alters the isotopic ratio of the remaining substrate ($\delta^{15}N^s$) that is available for incorporation into the β position, and vice versa.* Thus, simply determining the change in $\delta^{15}N^{\alpha}$ (or $\delta^{15}N^{\beta}$) as a function of $[-\ln f/(1-f)]$ over the course of the reaction (*i.e.*, finding the slope of a Rayleigh plot via linear regression of $\delta^{15}N^{\alpha}$ or $\delta^{15}N^{\beta}$ against $[-\ln f/(1-f)]$) is not an accurate method for calculating the enrichment factor for N^{α} (or N^{β}).

The complexity of calculating enrichment factors for a branched N_2O biosynthesis reaction can be demonstrated mathematically by substituting $\frac{1}{2}(\delta^{15}N^{\alpha} + \delta^{15}N^{\beta})$ for $\delta^{15}N^p$ and $\frac{1}{2}(\epsilon_{N-\alpha} + \epsilon_{N-\beta})$ for $\epsilon_{p/s}$ in the Rayleigh equation (Eq. (2)), yielding equation 3:

$$\frac{1}{2}(\delta^{15}N^{\alpha} + \delta^{15}N^{\beta}) = \delta^{15}N^{s0} - \frac{1}{2}(\epsilon_{N-\alpha} + \epsilon_{N-\beta}) \frac{f \ln(f)}{1-f} \quad (3)$$

Here, $\epsilon_{N-\alpha}$ and $\epsilon_{N-\beta}$ stand for the isotopic enrichment factors for the α and β nitrogen atoms, respectively. Equation 3 can then be rearranged to solve for $\delta^{15}N^{\alpha}$ (Eq. (4)):

$$\delta^{15}N^{\alpha} = 2\delta^{15}N^{s0} + \epsilon_{N-\alpha} \left(-\frac{f \ln(f)}{1-f} \right) + \epsilon_{N-\beta} \left(-\frac{f \ln(f)}{1-f} \right) - \delta^{15}N^{\beta} \quad (4)$$

Commented [ER4]: Fixed typo

Deleted: [[-lnf/(1-f)]]

The underlying difficulty in using Eq. (4) is that for a given value of f , $\delta^{15}\text{N}^\alpha$ depends not only on $\delta^{15}\text{N}^{\text{std}}$ and $\epsilon_{\text{N-}\alpha}$, but also on $\epsilon_{\text{N-}\beta}$ and the current $\delta^{15}\text{N}^\beta$ value. Put another way, we have one equation and two unknowns ($\epsilon_{\text{N-}\alpha}$ and $\epsilon_{\text{N-}\beta}$).

To address this issue, we developed an expanded version of the Rayleigh model that includes specific terms for the individual nitrogen atoms in N_2O . To test the robustness of this new Expanded Rayleigh model, we generated simulated isotopic data for each of five different potential catalytic mechanisms for N_2O formation. For each N_2O synthesis scenario, the size and skewness of the simulated measurement error was varied, and 1000 simulated datasets were generated for each combination of error level and skewness type. We then applied the standard Rayleigh model and Expanded Rayleigh model to each dataset to determine $\epsilon_{\text{N-bulk}}$, $\epsilon_{\text{N-}\alpha}$, and $\epsilon_{\text{N-}\beta}$ and compared the precision and accuracy of the two models. Finally, we compared the results of applying the standard Rayleigh model and Expanded Rayleigh model to previously published isotopic data.

2 Methods

2.1 Isotope nomenclature

The isotopic ratio, R , is defined as the abundance of the heavy isotope (^{15}N) divided by the abundance of the light isotope (^{14}N):

$$R = \frac{\text{Heavy Isotope}}{\text{Light Isotope}} = \left(\text{e.g. } \frac{^{15}\text{N}}{^{14}\text{N}} \right) \quad (5)$$

In the case of nitrogen, ^{14}N is naturally much more abundant than ^{15}N . (Atom percent of ^{15}N in atmospheric N_2 (air) is $0.3663 \pm 0.0004\%$, so $R = 0.0036765$) (Junk and Svec, 1958; Mariotti et al., 1981; Skrzypek and Dunn, 2020). The R value of a sample is usually measured relative to that of a standard ($R_{\text{sample}}/R_{\text{standard}}$). For nitrogen, the standard is atmospheric N_2 . Because there is naturally little variation in ^{15}N , $R_{\text{sample}}/R_{\text{standard}}$ values are typically close to 1, and isotopic ratios are usually converted to the δ notation, which is commonly expressed in the per mil form (‰) by multiplying the unitless form of δ by 1000:

$$\delta = \left(\frac{R_{\text{sample}}}{R_{\text{standard}}} - 1 \right) * 1000 \quad (6)$$

where R_{sample} is the isotopic ratio of the sample, and R_{standard} is the isotopic ratio of the analytical standard (Mariotti et al., 1981). The per mil form of δ is used throughout this manuscript.

N^{bulk} is defined as the total number of “equivalents” of nitrogen atoms in N_2O , or $2 * (\text{mol of } \text{N}_2\text{O})$. N^α and N^β are defined as the number of moles of N at the α (central) and β (outer) positions of N_2O , respectively.

$$0.5 * N^{\text{bulk}} = N^\alpha = N^\beta = \text{mol } \text{N}_2\text{O} \quad (7)$$

Thus, $^{15}\text{N}^{\text{bulk}}$ is defined as the total number of moles of ^{15}N at both the α and β positions:

$$^{15}\text{N}^{\text{bulk}} = ^{15}\text{N}^\alpha + ^{15}\text{N}^\beta \quad (8)$$

Similarly, $^{14}\text{N}^{\text{bulk}}$ is defined as the total number of moles of ^{14}N at the α and β positions of N_2O :

Commented [ER5]: Reviewer 1 comment (P4, L116): Updated Rstandard value

Deleted: ≈

Deleted: 0.003663

Commented [ER6]: Reviewer 1 comment (P4, L116): Changed text to clarify that the per mil δ notation is used throughout.

Deleted: the per mil equivalent,

Deleted: Throughout this manuscript,

Commented [ER7]: Reviewer 1 comment (P4, L121): Edited text to keep units consistent with Eq. (7).

Deleted: (or

Deleted:)

Deleted: atoms

$$^{14}N^{bulk} = ^{14}N^{\alpha} + ^{14}N^{\beta} \quad (9)$$

2.2 α , ϵ , and KIE

As discussed above, the isotopic fractionation factor ($\alpha_{p/s}$) is equal to the inverse of the kinetic isotope effect (KIE) for first-order reactions and for higher-order reactions when the rare (heavy) isotope is dilute (*i.e.*, ^{15}N or ^{13}C at natural abundance) (Eq. (10)) (Bigeleisen and Wolfsberg, 1958). KIE is defined as the rate constant of the light isotope, k_L , divided by the rate constant of the heavy isotope, k_H (Eq. (11)).

$$KIE = \frac{1}{\alpha_{p/s}} \quad (10)$$

$$KIE = \frac{k_L}{k_H} \quad (11)$$

Formally, $\alpha_{p/s}$ is defined as the isotopic ratio of instantaneously produced product (R_{pi}) divided by the isotopic ratio of residual substrate at the same instant (R_s) (Eq. (12)) (Mariotti et al., 1981).

$$\alpha_{p/s} = \frac{\frac{^{15}N^{pi}}{^{14}N^{pi}}}{\frac{^{15}N^s}{^{14}N^s}} = \frac{R_{pi}}{R_s} \quad (12)$$

As noted above, the fractionation factor can also be expressed in per mil (‰) terms by using Eq. (1) to convert α to the per mil enrichment factor, ϵ .

2.3 Assumptions

We used the Rayleigh equation (Eq. (2)) to determine α_{N-bulk} , the fractionation factor for N^{bulk} . The Rayleigh model depends on five key assumptions:

2.3.1 Assumption 1

The reaction is irreversible and occurs in a closed system in the absence of side-reactions (Mariotti et al., 1981).

This means that the final $\delta^{15}N^{bulk}$ value is equal to the initial $\delta^{15}N^s$ value ($\delta^{15}N^{s0}$).

$$\delta^{15}N^{bulk,final} = \delta^{15}N^{s0} \quad (13)$$

2.3.2 Assumption 2

The bulk fractionation factor ($\alpha_{p/s}$ or α_{N-bulk}) is constant.

This means that the isotopic ratio of bulk N produced in an infinitely short time period ($R_{bulk,i}$) divided by the isotopic ratio of the remaining substrate (R_s) is constant (Eq. (14)) (Mariotti et al., 1981).

$$\alpha_{N-bulk} = \frac{R_{bulk,i}}{R_s} \quad (14)$$

Commented [ER8]: Added Mariotti ref

This assumption should be valid as long as the rate-limiting step of the chemical transformation in question remains the same.

2.3.3 Assumption 3

The light isotope (^{14}N) is much more abundant than the heavy isotope (^{15}N), which is true for all experiments performed with ^{15}N at or near natural abundance.

This assumption was employed by Mariotti and colleagues to approximate total N abundance ($^{14}\text{N} + ^{15}\text{N}$) as ^{14}N in their definition of f (Mariotti et al., 1981). For N_2O isotopocules, the assumption that $^{14}\text{N} \gg ^{15}\text{N}$ also means that variations in apportionment of ^{15}N between the α and β positions will not significantly affect the abundance of ^{14}N at the α or β position. Thus, at or near natural abundance of ^{15}N , both $^{14}\text{N}^\alpha$ and $^{14}\text{N}^\beta$ may be approximated as $0.5 * ^{14}\text{N}^{\text{bulk}}$. This approximation underlies the assumption that $\delta^{15}\text{N}^{\text{bulk}}$ is approximately equal to the average of $\delta^{15}\text{N}^\alpha$ and $\delta^{15}\text{N}^\beta$ (Eq. (15)).

$$\delta^{15}\text{N}^{\text{bulk}} = \frac{1}{2} * (\delta^{15}\text{N}^\alpha + \delta^{15}\text{N}^\beta) \quad (15)$$

Equation 15 is routinely used in isotopic studies of N_2O to calculate $\delta^{15}\text{N}^\beta$ values using experimentally determined $\delta^{15}\text{N}^{\text{bulk}}$ and $\delta^{15}\text{N}^\alpha$ values (Toyoda and Yoshida, 1999).

2.3.4 Assumption 4

The values of δ_{s0} and δ_s are relatively close to 0‰.

Mariotti's approximation of the Rayleigh equation uses a simplification that is only valid if $\delta_s/1000$ and $\delta_{s0}/1000$ are small relative to 1 (Mariotti et al., 1981). In practice, for an N_2O synthesis reaction where $\epsilon_{\text{N-bulk}}$ is -20‰, a $\delta^{15}\text{N}^{s0}$ value of ± 100 ‰ will introduce an error of ~ 2 ‰ when $\epsilon_{\text{N-bulk}}$ is determined using this approximation (Table S1).

2.3.5 Assumption 5

The absolute value of ϵ (e.g., $\epsilon_{\text{N-bulk}}$) is relatively small.

The magnitude of $\delta^{15}\text{N}^s$ values depend on both $\delta^{15}\text{N}^{s0}$ and $\epsilon_{\text{N-bulk}}$. Thus, similar to the previous assumption, this assumption was introduced to facilitate the approximation that $\delta^{15}\text{N}^s/1000$ is small relative to 1 (Mariotti et al., 1981). As with assumption 4, the error introduced by this approximation increases as the absolute value of ϵ increases. For a reaction where $\delta^{15}\text{N}^{s0}$ is equal to 0‰ and $\epsilon_{\text{N-bulk}}$ is equal to ± 50 ‰, this approximation introduces an error of ~ 1 ‰ to ϵ values ($\epsilon_{\text{N-bulk}}$, $\epsilon_{\text{N-}\alpha}$, and $\epsilon_{\text{N-}\beta}$) while a reaction where $\epsilon_{\text{N-bulk}}$ is equal to ± 100 ‰ introduces an error of 5-6‰ (Table S2).

2.3.6 Assumption 6

For the purposes of modeling N_2O synthesis, we will use all of the well-established assumptions listed above. Additionally, to account for fractionation at N^α and N^β individually, we introduce one more assumption:

The individual fractionation factors for the α and β nitrogen atoms in N_2O ($\alpha_{\text{N-}\alpha}$ and $\alpha_{\text{N-}\beta}$) are essentially constant.

190 We define $\alpha_{N-\alpha}$ and $\alpha_{N-\beta}$ in a manner analogous to that of $\alpha_{N-\text{bulk}}$ (Eq. (14)). Thus, $\alpha_{N-\alpha}$ is defined as the isotopic ratio of N^α from instantaneously produced N_2O ($R_{\alpha,i}$) divided by R_s (Eq. (16)).

$$\alpha_{N-\alpha} = \frac{\frac{^{15}N^{\alpha,i}}{^{14}N^{\alpha,i}}}{\frac{^{15}N^s}{^{14}N^s}} = \frac{R_{\alpha,i}}{R_s} \quad (16)$$

Similarly, $\alpha_{N-\beta}$ is defined as the instantaneous isotopic ratio of N^β ($R_{\beta,i}$) divided by R_s (Eq. (17)).

$$\alpha_{N-\beta} = \frac{\frac{^{15}N^{\beta,i}}{^{14}N^{\beta,i}}}{\frac{^{15}N^s}{^{14}N^s}} = \frac{R_{\beta,i}}{R_s} \quad (17)$$

195 Put another way, we assume *a priori* that the processes that govern preferential incorporation of one isotope over another remain unchanged at both the α and β positions of N_2O during the course of the reaction. We expect this assumption to be valid throughout the reaction, as long as the rate-limiting steps of N_2O biosynthesis are unchanged.

2.4 The standard Rayleigh model and simulation of error-free $\delta^{15}N^{\text{bulk}}$ and $\delta^{15}N^s$ values

To compare the standard Rayleigh approach and our nonlinear expansion of the Rayleigh model, we simulated isotopic data for a number of different combinations of kinetic isotope effects for N^α and N^β (see Table 1 and Fig. 1). Each idealized, error-free dataset had three groups of equally spaced values of f (0.7-0.2), similar to a typical experiment with three replicates (Tables S4-S8). For each dataset, $\delta^{15}N^{s0}$ was set at 0‰, and $\epsilon_{N-\text{bulk}}$ was set at either -20‰ (normal KIE, KIE = 1.0204) or +20‰ (inverse KIE, KIE = 0.9804). Equation 2, Mariotti's approximation of the Rayleigh model for accumulated product (Mariotti et al., 1981), was used to calculate $\delta^{15}N^{\text{bulk}}$ values of N_2O for each value of f ($\delta^{15}N^{\text{bulk}} = \delta^{15}N^p$). Similarly, δ values for remaining substrate ($\delta^{15}N^s$) were calculated using Eq. (18), Mariotti's approximation of the Rayleigh model written in terms of substrate (Mariotti et al., 1981).

$$\delta^{15}N^s = \delta^{15}N^{s0} + \epsilon_{N-\text{bulk}} * \ln(f) \quad (18)$$

The $\delta^{15}N^{\text{bulk}}$ and $\delta^{15}N^s$ values simulated using the standard Rayleigh equation are shown in Fig. 1 in gray and orange, respectively.

2.5 Nonlinear expansion of the Rayleigh model

Simulated $\delta^{15}N^\alpha$ and $\delta^{15}N^\beta$ values were calculated based on the assumption that, like the fractionation factor for bulk N, the fractionation factors for the individual N^α and N^β atoms are constant (see Assumption 6 above). To relate $\alpha_{N-\alpha}$ and $\alpha_{N-\beta}$ to $\alpha_{N-\text{bulk}}$, we define the fraction of instantaneously converted $^{15}N^{\text{bulk}}$ ($^{15}N^{\text{bulk},i}$) that is incorporated into the α position as ρ (Eq. (19)).

$$\rho = \frac{^{15}N^{\alpha,i}}{^{15}N^{\text{bulk},i}} \quad (19)$$

215 Similarly, we define the fraction of instantaneously converted ^{14}N ($^{14}N^{\text{bulk},i}$) that goes to the α position as τ (Eq. (20)).

Commented [ER9]: Response to community comment: Added the KIE values that correspond to the ϵ values listed.

Deleted: Modeling was performed with R statistical software (R Core Team, 2022), and figures were produced with ggplot2 (Wickham, 2016).¶

$$\tau = \frac{{}^{14}\text{N}^{\alpha,i}}{{}^{14}\text{N}^{\text{bulk},i}} \quad (20)$$

220 In keeping with the premise that $\alpha_{\text{N-}\alpha}$ and $\alpha_{\text{N-}\beta}$ are constant, we assume *a priori* that ρ and τ are constant. Thus, the equation for $\alpha_{\text{N-}\alpha}$ may be written in terms of ρ , τ , and $\alpha_{\text{N-bulk}}$:

$$\alpha_{\text{N-}\alpha} = \frac{\frac{\rho \cdot {}^{15}\text{N}^{\text{bulk},i}}{\tau \cdot {}^{14}\text{N}^{\text{bulk},i}}}{\frac{{}^{15}\text{N}^{\alpha}}{{}^{14}\text{N}^{\alpha}}} = \frac{\rho}{\tau} * \alpha_{\text{N-bulk}} \quad (21)$$

In the absence of side-reactions, ${}^{15}\text{N}^{\text{bulk},i}$ must be equal to the sum of ${}^{15}\text{N}^{\alpha,i}$ and ${}^{15}\text{N}^{\beta,i}$. Similarly, ${}^{14}\text{N}^{\text{bulk},i}$ must be equal to the sum of ${}^{14}\text{N}^{\alpha,i}$ and ${}^{14}\text{N}^{\beta,i}$. Thus, the equation for $\alpha_{\text{N-}\beta}$ may also be written in terms of ρ , τ , and $\alpha_{\text{N-bulk}}$:

$$225 \alpha_{\text{N-}\beta} = \frac{\frac{(1-\rho) \cdot {}^{15}\text{N}^{\text{bulk},i}}{(1-\tau) \cdot {}^{14}\text{N}^{\text{bulk},i}}}{\frac{{}^{15}\text{N}^{\beta}}{{}^{14}\text{N}^{\beta}}} = \frac{1-\rho}{1-\tau} * \alpha_{\text{N-bulk}} \quad (22)$$

Equations 21 and 22 form the basis for the Expanded Rayleigh model.

To simulate $\delta^{15}\text{N}^{\alpha}$ and $\delta^{15}\text{N}^{\beta}$ values and determine the corresponding fractionation factors, ρ and τ must be related to accumulated isotope ratios, R_{α} and R_{β} . Although ρ is formally defined in Eq. (19) as the fraction of instantaneously converted ${}^{15}\text{N}$ apportioned to the α position, ρ can also be written in terms of *accumulated* ${}^{15}\text{N}$ values as long as ρ is constant:

$$230 \rho = \frac{{}^{15}\text{N}^{\alpha}}{{}^{15}\text{N}^{\text{bulk}}} \quad (23)$$

Similarly, as long as τ is constant, τ may be written in terms of accumulated ${}^{14}\text{N}$ values:

$$\tau = \frac{{}^{14}\text{N}^{\alpha}}{{}^{14}\text{N}^{\text{bulk}}} \quad (24)$$

Using the definition of ρ from Eq. (23), R_{α} can be written in terms of ρ by substituting ${}^{15}\text{N}^{\alpha}$ with $\rho \cdot {}^{15}\text{N}^{\text{bulk}}$ and replacing ${}^{14}\text{N}^{\alpha}$ with $0.5 \cdot ({}^{\text{N}^{\text{bulk}}}) - {}^{15}\text{N}^{\alpha}$ (Eq. (25)).

$$235 R_{\alpha} = \frac{{}^{15}\text{N}^{\alpha}}{{}^{14}\text{N}^{\alpha}} = \frac{\rho \cdot {}^{15}\text{N}^{\text{bulk}}}{0.5 \cdot \text{N}^{\text{bulk}} - \rho \cdot {}^{15}\text{N}^{\text{bulk}}} \quad (25)$$

The same approach can be used to define R_{β} in terms of ρ (Eq. (26)).

$$R_{\beta} = \frac{{}^{15}\text{N}^{\beta}}{{}^{14}\text{N}^{\beta}} = \frac{(1-\rho) \cdot {}^{15}\text{N}^{\text{bulk}}}{0.5 \cdot \text{N}^{\text{bulk}} - (1-\rho) \cdot {}^{15}\text{N}^{\text{bulk}}} \quad (26)$$

The term $0.5 \cdot \text{N}^{\text{bulk}}$ is equal to moles of N_2O (Eq. (7)), and ${}^{15}\text{N}^{\text{bulk}}$ can be calculated for any value of f using N^{bulk} and $\delta^{15}\text{N}^{\text{bulk}}$ (see SI). Equations 25 and 26 can be converted to δ notation as follows (see Eq. (6)):

$$240 \delta^{15}\text{N}^{\alpha} = \left(\frac{\rho \cdot {}^{15}\text{N}^{\text{bulk}}}{(0.5 \cdot \text{N}^{\text{bulk}} - \rho \cdot {}^{15}\text{N}^{\text{bulk}}) \cdot R_{\text{standard}}} - 1 \right) * 1000 \quad (27)$$

$$\delta^{15}\text{N}^{\beta} = \left(\frac{(1-\rho) \cdot {}^{15}\text{N}^{\text{bulk}}}{(0.5 \cdot \text{N}^{\text{bulk}} - (1-\rho) \cdot {}^{15}\text{N}^{\text{bulk}}) \cdot R_{\text{standard}}} - 1 \right) * 1000 \quad (28)$$

Deleted:]

Equations 27 and 28 can be used to determine $\delta^{15}\text{N}^\alpha$ and $\delta^{15}\text{N}^\beta$ based on $\delta^{15}\text{N}^{\text{bulk}}$ as long as ρ is constant and $\delta^{15}\text{N}^{\text{bulk}}$ can be approximated as the average of $\delta^{15}\text{N}^\alpha$ and $\delta^{15}\text{N}^\beta$ (see Eq. (15) and Table S3).

245 2.6 Simulation of error-free $\delta^{15}\text{N}^\alpha$ and $\delta^{15}\text{N}^\beta$ values

Idealized, error-free $\delta^{15}\text{N}^\alpha$ and $\delta^{15}\text{N}^\beta$ values (shown in Fig. 1) were simulated for the values of ρ listed in Table 1 by calculating $\delta^{15}\text{N}^\alpha$ and $\delta^{15}\text{N}^\beta$ using Eq. (27) and Eq. (28) (Tables S4-S8). As outlined in Table 1, five different scenarios with different combinations of isotope effects (none, normal, or inverse) for the α and β nitrogen atoms of N_2O were simulated. For Datasets 1 and 5, either $\epsilon^{15}\text{N}^\alpha$ or $\epsilon^{15}\text{N}^\beta$ was set to 0‰ (no isotope effect, $\text{KIE} = 1$), and the corresponding values of ρ and τ were back-calculated (see SI for details). The remaining values of ρ were chosen to produce datasets with the different combinations of normal and/or inverse isotope effects (Datasets 2-4). The ρ values tested (0.4898-0.5200) are all close enough to 0.5 such that the average of $\delta^{15}\text{N}^\alpha$ and $\delta^{15}\text{N}^\beta$ is approximately equal to $\delta^{15}\text{N}^{\text{bulk}}$ (Table S3). In all cases, discrepancies between $\delta^{15}\text{N}^{\text{bulk}}$ values calculated using Eq. (2) and $\delta^{15}\text{N}^{\text{bulk}}$ values calculated by averaging $\delta^{15}\text{N}^\alpha$ and $\delta^{15}\text{N}^\beta$ (Eq. (15)) are well below the typical analytical error of 0.5-0.7‰ (Yang et al., 2014).

255

Commented [ER12]: Response to community comment - Added reference to figure to show how calculations are connected to figures.

Deleted: Modeling was performed with R statistical software (R Core Team, 2022), and figures were produced with ggplot2 (Wickham, 2016).

260 **Table 1. Parameters for idealized, error-free datasets describing N₂O synthesis for five simulated scenarios (Datasets 1-5) with varying combinations of $\epsilon_{N\text{-bulk}}$ and ρ .**

| Simulated | | $\epsilon_{N\text{-bulk}}$ | δ_{50} | ρ ^b | τ ^c |
|-----------|-----------------------|----------------------------|---------------|---------------------|---------------------|
| dataset | Subset | (‰) ^a | (‰) | | |
| 1 | $0.3 \leq f \leq 0.7$ | -20 | 0 | 0.5102 | 0.49996 |
| 2 | $0.3 \leq f \leq 0.7$ | -20 | 0 | 0.5050 | 0.49998 |
| 3 | $0.3 \leq f \leq 0.7$ | -20 | 0 | 0.5200 | 0.49993 |
| 4 | $0.3 \leq f \leq 0.7$ | 20 | 0 | 0.5050 | 0.49998 |
| 5 | $0.3 \leq f \leq 0.7$ | -20 | 0 | 0.4898 | 0.50004 |

Deleted: ^a

Deleted: ^b

^a Note that an ϵ value of -20‰ corresponds to a KIE of 1.0204 and an ϵ value of +20‰ corresponds to a KIE of 0.9804.

Commented [ER14]: Response to community comment: Added footnote with the KIE values that correspond to ϵ values.

^b Parameter introduced in the Expanded Rayleigh model (Eq. (23)): $\rho = {}^{15}\text{N}^{\alpha/15}\text{N}^{\text{bulk}}$.

Deleted: ^a

^c Parameter introduced in the Expanded Rayleigh model (Eq. (24)): $\tau = {}^{14}\text{N}^{\alpha/14}\text{N}^{\text{bulk}}$. τ was calculated for each value of f ; the

Deleted: ^b

265 average value of τ is listed.





Figure 1. Simulated $\delta^{15}\text{N}$ values for N_2O synthesis in a closed system with different combinations of KIE $^{15}\text{N}^\alpha$ and KIE $^{15}\text{N}^\beta$.

$\delta^{15}\text{N}$ values of remaining substrate ($\delta^{15}\text{N}^\beta$) and accumulated product (N_2O) ($\delta^{15}\text{N}^{\text{bulk}}$, $\delta^{15}\text{N}^\alpha$, and $\delta^{15}\text{N}^\beta$) are plotted as a function of the fraction of substrate reduced (1-f) with reactions progressing from left to right. Values of $\delta^{15}\text{N}^{\text{bulk}}$ and $\delta^{15}\text{N}^\alpha$ were calculated using Eq (2) or Eq (18) (Mariotti's approximation of the Rayleigh equation (Mariotti et al., 1981)), where $\epsilon^{15}\text{N}^{\text{bulk}}$ is set to -20‰ (Datasets 1-3, 5) or +20‰ (Dataset 4) and the initial $\delta^{15}\text{N}^\alpha$ value ($\delta^{15}\text{N}^{\alpha 0}$) is set to 0‰. Values of $\delta^{15}\text{N}^\alpha$ and $\delta^{15}\text{N}^\beta$ were calculated using Eq. (27) and Eq. (28). For Datasets 1 and 5, either $\epsilon^{15}\text{N}^\alpha$ or $\epsilon^{15}\text{N}^\beta$ was set to 0‰ (no isotope effect, KIE = 1), and the corresponding value of ρ was back-calculated (see SI for details). For Datasets 2-4, values of $\delta^{15}\text{N}^\alpha$ and $\delta^{15}\text{N}^\beta$ were calculated using pre-determined values of ρ that were less than (Datasets 2 and 4) or greater than (Dataset 3) the value of ρ from Dataset 1. See Table 3 for the KIE values for each simulation.

2.7 Determination of ρ and τ for the Expanded Rayleigh model

To determine $\alpha_{\text{N-}\alpha}$ and $\alpha_{\text{N-}\beta}$ using Eq. (21) and Eq. (22), ρ and τ must be calculated using values that can be measured experimentally. Because the abundance of ^{14}N is so much greater than that of ^{15}N , variations in ^{15}N apportionment between N^α and N^β are negligible when determining the fraction of $^{14}\text{N}^{\text{bulk}}$ at the α position (*i.e.*, τ). For enzyme-catalyzed N_2O synthesis with substrate at or near natural abundance of ^{15}N , we have found that τ is always 0.5000 (see analysis of data from Sutka et al., 2006 and Yang et al., 2014 (below)). For the purposes of testing our new Expanded Rayleigh model, we calculated more



Deleted:

Commented [ER15]: Response to reviewer 1: Replaced Fig. 1 after recalculating simulated δ values using updated Rstandard value. (New calculations use Rstd = 0.0036765 instead of Rstd = 0.003663). Note that this change has essentially no effect on the calculated values.

Commented [ER16]: Corrected typo

Deleted: (below)

precise values of τ . To do this, τ was calculated for each value of f using Eq. (24), and the average τ value for each Dataset
 290 was used to calculate fractionation factors.

To determine ρ using values that can be measured experimentally, Eq. (15) is rewritten so that ρ is the only unknown parameter. This can be accomplished by replacing $\delta^{15}N^\alpha$ with the equivalent term from Eq. (27), producing Eq. (29), a nonlinear equation we have named nonlinear model 1.

$$\delta^{15}N^{bulk} = \frac{1}{2} * \left[\left(\frac{\rho *^{15}N^{bulk}}{(0.5 * N^{bulk} - \rho *^{15}N^{bulk}) * R_{standard}} - 1 \right) * 1000 + \delta^{15}N^\beta \right] \quad (29)$$

295 Alternatively, $\delta^{15}N^\beta$ could be replaced with a term that includes ρ (Eq. (28)), producing Eq. (30) (nonlinear model 2):

$$\delta^{15}N^{bulk} = \frac{1}{2} * \left[\delta^{15}N^\alpha + \left(\frac{(1-\rho) *^{15}N^{bulk}}{(0.5 * N^{bulk} - (1-\rho) *^{15}N^{bulk}) * R_{standard}} - 1 \right) * 1000 \right] \quad (30)$$

Both nonlinear models were used to find the optimal value of ρ via nonlinear least squares regression for each of our simulated datasets. ↓

Deleted: Nonlinear least squares regression was performed as previously described (Baty et al., 2015) using a starting ρ value of 0.5.





Figure 2. General workflow for using the Expanded Rayleigh model to determine fractionation factors (α) and kinetic isotope effects (KIEs) for the individual N atoms (N^{α} and N^{β}) in N_2O .

To determine the fractionation factors for the α and β atoms in N_2O ($\alpha_{N-\alpha}$ and $\alpha_{N-\beta}$), three separate values must be calculated. **1.** The fractionation factor for bulk N (α_{N-bulk}) is determined by finding the slope (ϵ_{N-bulk}) of a standard Rayleigh plot where $y = [-\ln f / (1-f)]$ and $x = \delta^{15}N^{bulk}$, and converting ϵ_{N-bulk} to α_{N-bulk} as has been previously described (Mariotti et al., 1981). **2.** Nonlinear modeling is used to determine the optimal value of ρ (the fraction of $^{15}N^{bulk}$ at the α position). For Expanded Rayleigh model 1 (shown here), $\delta^{15}N^{\alpha}$ is replaced in the top equation with an equivalent expression that includes ρ (bottom equation) to generate the model used for nonlinear least squares regression (Eq. (29)). For Expanded Rayleigh model 2, $\delta^{15}N^{\beta}$ is substituted, producing Eq. (30) (not shown). **3.** The value of τ (the fraction of $^{14}N^{bulk}$ at the α position) is determined by dividing $^{14}N^{\alpha}$ by $^{14}N^{bulk}$ for each step of the reaction and averaging the results. **4.** Finally, α_{N-bulk} , ρ , and τ are used to calculate $\alpha_{N-\alpha}$ and $\alpha_{N-\beta}$. **5-6.** These individual fractionation factors can be converted to the corresponding KIE values (Eq. (10)) or ϵ values (Eq (1)).

2.8 Generation of datasets with simulated error

To mimic experimental error, 1000 simulated datasets were generated from each error-free dataset by adding random error to N^s , $\delta^{15}N^{bulk}$, and $\delta^{15}N^{\alpha}$ using a modified version of the procedure outlined by Scott and colleagues (Scott et al., 2004). Error was propagated to f by recalculating f using the N^s values with error added ($f = N^s / N^{s0}$). Similarly, error was propagated to $\delta^{15}N^{\beta}$ by recalculating $\delta^{15}N^{\beta}$ using $\delta^{15}N^{bulk}$ and $\delta^{15}N^{\alpha}$ values with error added (Eq. (15)). For each simulation, fractionation factors and KIE values were calculated using the standard Rayleigh model and the Expanded Rayleigh model (versions 1 and



Deleted:

Commented [ER18]: Replaced Fig. 2 to correct a typo in panel 1 (“ $\alpha_{N-\alpha}$ ” was replaced with “ α_{N-bulk} ”).

2), and the KIE values from each set of 1000 simulations were averaged. Three “levels” of error (low, medium, and high) and three types of skewness (none, left-skewed, and right-skewed, see below) were used for a total of nine conditions (Table 2, Fig. 3, and Figs. S1-S4). For simulations with low error, the standard deviation of each randomly generated error term was set to the level of error expected in a typical experiment (Table 2). The standard deviation of N^s values was set to 1.5% of N^{s0} (the maximum value of N^s), which corresponds to the estimated level of error in gas volume measurements (see SI). The standard deviations of $\delta^{15}N^{bulk}$ and $\delta^{15}N^a$ were set to 0.5‰ and 0.7‰, respectively, which correspond to standard deviations typical for isotopic analysis of N_2O via IRMS (Yang et al., 2014). To further test the robustness of the linear and nonlinear models, medium- and high-level error terms were generated by multiplying the low-level error terms for N^s , $\delta^{15}N^{bulk}$, and $\delta^{15}N^a$ by the appropriate factor (Table 2). The multiplication factors for medium- and high-level error were chosen so that the average R^2 values produced by linear regression of $\delta^{15}N^{bulk}$ against $[-\ln f/(1-f)]$ were approximately 0.7 and 0.4, respectively. Similar R^2 values have been observed for standard Rayleigh linear regression plots for N_2O biosynthesis (e.g., (Yang et al., 2014), Figs. S2-S4).

335

Table 2. Standard deviation and skewness of random error added to simulated N^s , $\delta^{15}N^{bulk}$, and $\delta^{15}N^a$ values.

| Level of error ^a | N^s (nmol) | $\delta^{15}N^{bulk}$ (‰) | $\delta^{15}N^a$ (‰) | Skewness ^b | |
|-----------------------------|-------------------------------------|---------------------------|----------------------|-----------------------|------------------------------|
| Low | $0.015 * N^{s0}$ ^c | 0.5 | 0.7 | None | $ \text{Skewness} \leq 0.5$ |
| | | | | Left | Skewness < -0.5 |
| | | | | Right | Skewness > 0.5 |
| Medium | $2 * (0.015 * N^{s0})$ ^c | $2.5 * (0.5)$ | $2.5 * (0.7)$ | None | $ \text{Skewness} \leq 0.5$ |
| | | | | Left | Skewness < -0.5 |
| | | | | Right | Skewness > 0.5 |
| High | $3 * (0.015 * N^{s0})$ ^c | $5 * (0.5)$ | $5 * (0.7)$ | None | $ \text{Skewness} \leq 0.5$ |
| | | | | Left | Skewness < -0.5 |
| | | | | Right | Skewness > 0.5 |

^a “Level of error” refers to the standard deviations of N^s (nmol of remaining substrate), $\delta^{15}N^{bulk}$, and $\delta^{15}N^a$.

^b Skewness of the residuals for the standard Rayleigh model (linear regression of $\delta^{15}N^{bulk}$ against $[-\ln f/(1-f)]$).

340 ^c N^{s0} refers to nmol of substrate at time = 0 (i.e., the maximum amount of substrate). N^{s0} was set to 10,000 nmol in all simulations.

For each level of error, three sets of 1000 simulations with different types of skewness (no skew, left-skewed, or right-skewed) were performed to mimic types of skewness that might be observed in experimental data. For example, if N₂O leaks from a sample bottle prior to analysis, the quantity of N₂O produced will be underestimated, and the corresponding values of N^s and f will be overestimated. Additionally, since light isotopes have a higher rate of both diffusion and effusion, values of δ¹⁵N^{bulk}, δ¹⁵N^α, and δ¹⁵N^β will be overestimated. This scenario is represented by simulations where the random error values added to N^s, δ¹⁵N^{bulk}, and δ¹⁵N^α were derived from left-skewed distributions. Skewness classifications were then assigned based on the skewness of the residuals for the standard Rayleigh model where the dependent variable (y) is δ¹⁵N^{bulk} (Table 2). Simulations with residuals between -0.5 and 0.5 (inclusive) were designated “no skew,” meaning the distribution of these residuals did not differ significantly from the normal distribution. Simulations where standard Rayleigh residuals had a skewness value less than -0.5 were designated “left-skewed” (*i.e.*, left-tailed distribution), and simulations where skewness was greater than 0.5 were designated “right-skewed” (*i.e.*, right-tailed distribution).

To assess the accuracy of each model under different circumstances, average KIE values and standard deviations were calculated for each set of 1000 simulations with error. The absolute relative difference of each average KIE was calculated using Eq. (31),

$$\text{absolute relative difference} = \left| \frac{\text{estimate} - \text{actual}}{\text{actual}} \right| \quad (31)$$

where the estimate is the average KIE and actual values were calculated directly from the input values for simulations without error (see Table 3). Lower absolute relative difference values indicate higher accuracy. Goodness of fit was assessed for the (linear) standard Rayleigh model and nonlinear models 1 and 2 by calculating the average root mean square error (RMSE) for each set of 1000 simulations. For each simulation, p-values for ε_{N-bulk} and ρ (coefficients extracted from the (linear) standard Rayleigh model and nonlinear models 1 or 2, respectively) were determined using the one-sample *t*-test (two-sided) (Baty et al., 2015; Ritz and Streibig, 2008; Kalpić et al., 2011). The null hypothesis used to calculate these p-values is that there is no kinetic isotope effect. For the linear model, this means that the null hypothesis is ε_{N-bulk} = 0 (KIE ¹⁵N^{bulk} = 1). For the nonlinear models, the null hypothesis is that ρ = 0.5 (KIE ¹⁵N^α = 1 and KIE ¹⁵N^β = 1).

Deleted: The skewness function from the moments package in R was used to calculate skewness values (Komsta and Novomestky, 2022). In this package, skewness is defined as $(1/n) * \sum((x - \bar{x})/s)^3$, where *n* is the sample size, \bar{x} is the sample mean, and *s* is the sample standard deviation (Hippel, 2011).

Deleted: by finding the absolute value of the difference of the estimate (average KIE) and the actual value divided by the actual value

Deleted: (

Deleted:).

Commented [ER20]: Response to community comment: Reword sentence to streamline methods/eliminate redundancy.

Deleted: A




Figure 3. Example simulations derived from Dataset 1 with varying levels of error and types of skewness.

Each panel shows a single dataset (representative of 1000 simulated datasets) consisting of three replicates with five timepoints each. All simulations were derived from Dataset 1 ($\delta^{15}\text{N}^0 = 0\text{‰}$, $\epsilon_{\text{N-bulk}} = -20\text{‰}$, $(\text{KIE}^{15}\text{N}^{\text{bulk}} = 1.0204)$, $\text{KIE}^{15}\text{N}^{\alpha} = 1.0000$, and $\text{KIE}^{15}\text{N}^{\beta} = 1.0417$). Each level of error and type of skewness is described in Table 2. Example graphs derived from Datasets 2-4 are shown in Figs. S1-S4.


2.9 Analysis of previously published experimental data

The standard and Expanded Rayleigh models were applied to published data on N₂O produced from NH₂OH by *Methylophilus trichosporium* (ATCC 49243) (Sutka et al., 2006). We calculated values of f , $\delta^{15}\text{N}^{\alpha}$, and $\delta^{15}\text{N}^{\beta}$ using the N₂O concentrations, $\delta^{15}\text{N}^{\text{bulk}}$ values, and site preference (SP) values from replicate B measured by Sutka et al. (see Table S14 for original data and our calculated values, and Fig. 4 for a graphical representation of these data). SP is defined as the difference between $\delta^{15}\text{N}^{\alpha}$ and $\delta^{15}\text{N}^{\beta}$:

$$SP = \delta^{15}\text{N}^{\alpha} - \delta^{15}\text{N}^{\beta} \quad (32)$$

The data for replicate B was chosen because this data had a range of f values large enough for $\epsilon_{\text{N-bulk}}$ to be determined via the standard Rayleigh method (linear regression of $\delta^{15}\text{N}^{\text{bulk}}$ against $[-\ln f/(1-f)]$). See SI for details.

Deleted: Modeling and statistical analyses were performed with R statistical software (R Core Team, 2022), random numbers representing simulated error were generated using the rsn function from the skew-normal distribution package (Azzalini, 2023), skewness was calculated with the moments package (Komsta and Novomesky, 2022), and figures were produced with ggplot2 (Wickham, 2016).¶



Commented [ER22]: Response to reviewer 1: Replaced Fig. 3 after recalculating simulated δ values using updated Rstandard value. (New calculations use Rstd = 0.0036765 instead of Rstd = 0.003663). Note that this change has essentially no effect on the calculated values.

Commented [ER23]: Response to community comment: Added KIE value that corresponds to ϵ value.

Deleted: .


We also applied the standard and Expanded Rayleigh models to previously published isotopic data on N₂O production from purified *Histoplasma capsulatum* (fungal) P450 NOR (Fig. 5 and Table S16) (Yang et al., 2014). Because this dataset covered a wide range of *f* values (*f* = 0.42-0.87) and $\delta^{15}\text{N}^{\text{bulk}}$ varied linearly with $[-\ln f/(1-f)]$, we used the standard Rayleigh model (Eq. (2)) to determine $\epsilon_{\text{N-bulk}}$ via linear regression of $\delta^{15}\text{N}^{\text{bulk}}$ against $[-\ln f/(1-f)]$. $\epsilon_{\text{N-bulk}}$ was determined for each of three independent biological replicates (13 observations/replicate), and the results were averaged. Similarly, to determine standard Rayleigh values of $\epsilon_{\text{N-}\alpha}$ and $\epsilon_{\text{N-}\beta}$, we performed linear regression of $\delta^{15}\text{N}^{\alpha}$ or $\delta^{15}\text{N}^{\beta}$ against $[-\ln f/(1-f)]$ for each independent replicate and averaged the results.

To apply the Expanded Rayleigh model to the isotopic data on N₂O production by P450 NOR, we used a modified procedure because, in this case, ρ and τ vary linearly with $[-\ln f/(1-f)]$. **Using nonlinear regression to determine ρ in this case yields a value of ρ in between the more extreme values of ρ observed at the start and end of the reaction and does not represent the data from the overall reaction very well.** Therefore, instead of using nonlinear model 1 (Eq. (29) or model 2 (Eq. (30)) to determine ρ using data from the entire reaction, we calculated ρ for each individual timepoint (*i.e.*, each individual value of *f*) using Eq. (23). Similarly, τ was calculated for each individual observation using Eq. (24). These individual ρ and τ values were then used to calculate $\alpha_{\text{N-}\alpha}$ (Eq. (21)) and $\alpha_{\text{N-}\beta}$ (Eq. (22)) values for each value of *f*, which were then converted to KIE values using Eq. (10). This approach yielded KIE values that increased (KIE ¹⁵N^α) or decreased (KIE ¹⁵N^β) as the reaction progressed. To estimate the range of position-specific KIEs produced by the Expanded Rayleigh model, individual KIE ¹⁵N^α and KIE ¹⁵N^β values representing the early part of the reaction (*f* = 0.8 ± 0.03) or the latter part of the reaction (late reaction, *f* = 0.5 ± 0.03) were averaged (Table 6, Table S17).

Formatted: Justified, Line spacing: 1.5 lines

Commented [ER24]: Community comment 1: Added text to clarify why the Expanded Rayleigh model was applied to individual observations for the P450 NOR data.

Deleted: ¶
Modeling was performed with R statistical software (R Core Team, 2022), and $\delta^{15}\text{N}$ values were plotted with ggplot2 (Wickham, 2016). A linear model was used to determine if SP changed as a function of $[-\ln f/(1-f)]$. ¶



425 Figure 4. Experimentally measured $\delta^{15}\text{N}$ values for N_2O synthesis from NH_2OH in a pure culture of *Methylosinus trichosporium* (data from (Sutka et al., 2006)).

Data previously reported for *M. trichosporium* (*Methylocystis* sp.) replicate B (Sutka et al., 2006). $\delta^{15}\text{N}$ values of accumulated product (N_2O) ($\delta^{15}\text{N}^{\text{bulk}}$, $\delta^{15}\text{N}^{\alpha}$, and $\delta^{15}\text{N}^{\beta}$) are plotted as a function of the fraction of substrate reduced (1-f) with the reaction progressing from left to right. We back-calculated values of $\delta^{15}\text{N}^{\alpha}$, and $\delta^{15}\text{N}^{\beta}$ using the reported N_2O concentrations, $\delta^{15}\text{N}^{\text{bulk}}$ values, and SP values. The initial measured $\delta^{15}\text{N}$ value of NH_2OH ($\delta^{15}\text{N}^{\text{S}0}$), -2.3‰, is shown at $f = 1$.

430




Figure 5. Experimentally measured product $\delta^{15}\text{N}$ values and calculated $\delta^{15}\text{N}^{50}$ values for N_2O synthesis from NO by purified *H. capsulatum* P450 NOR (data from (Yang et al., 2014)).

435 Data previously reported for *H. capsulatum* (fungal) P450 NOR replicates A-C (Yang et al., 2014). $\delta^{15}\text{N}$ values of accumulated product (N_2O) ($\delta^{15}\text{N}^{\text{bulk}}$, $\delta^{15}\text{N}^{\alpha}$, and $\delta^{15}\text{N}^{\beta}$) are plotted as a function of the fraction of substrate reduced (1-f) with the reaction progressing from left to right. We back-calculated $\delta^{15}\text{N}^{50}$ for each replicate by finding the intercept of the corresponding standard Rayleigh plot (Eq. (2)) where $x = [-\ln f/(1-f)]$ and $y = \delta^{15}\text{N}^{\text{bulk}}$.

2.10 Modeling, statistical analysis, and figures

440 Modeling and statistical analyses of simulated and experimental δ values were performed with R statistical software (R Core Team, 2022), and figures were produced with ggplot2 (Wickham, 2016). To determine ρ for the Expanded Rayleigh model, nonlinear least squares regression was performed as previously described (Baty et al., 2015) using a starting ρ value of 0.5.

445 For datasets with simulated error, random numbers representing simulated error were generated using the rsn function from the skew-normal distribution package (Azzalini, 2023), and skewness was calculated with the moments package (Komsta and Novomestky, 2022). In the moments package, skewness is defined as $(1/n) * \sum((x - \bar{x})/s)^3$, where n is the sample size, \bar{x} is the sample mean, and s is the sample standard deviation (Hippel, 2011).

For previously published experimental data, a linear model was used to determine if SP, ρ , or τ varied as a function of $[-\ln f/(1-f)]$.

Commented [ER26]: Response to community comment: Consolidated information on plotting and calculations into one new section.

3 Results

450 3.1 Error-free simulations

To demonstrate that the standard Rayleigh model produces inaccurate results for the individual nitrogen atoms in N₂O, idealized, error-free datasets were simulated representing different scenarios with varying combinations of KIEs for N^α and N^β (Fig. 1, Table 1). Because these data were simulated assuming that the fractions of ¹⁵N and ¹⁴N apportioned to each position remain constant (i.e., constant ρ and τ), the distance between δ¹⁵N^α and δ¹⁵N^β over the course of each reaction is constant, and the rates of change of δ¹⁵N^α, δ¹⁵N^β, and δ¹⁵N^{bulk} with respect to reaction progress (1-f) are essentially equal (Fig. 1). Thus, when the standard Rayleigh model (Eq. 2) is applied to each dataset, the slopes (ε) of δ¹⁵N^α, δ¹⁵N^β, and δ¹⁵N^{bulk} against [-lnf/(1-f)] for each dataset are all approximately equal, as are the corresponding KIE ¹⁵N^{bulk}, KIE ¹⁵N^α, and KIE ¹⁵N^β values (Table 3). While the standard Rayleigh KIE ¹⁵N^{bulk} values match the actual KIE ¹⁵N^{bulk} values, the KIE ¹⁵N^α and KIE ¹⁵N^β values determined using the standard Rayleigh approach differ significantly from the actual KIEs calculated from simulation input values (Table 3). In each of the five simulated reactions (Datasets 1-5), the isotopic preference for N^α differs from that of N^β, which can be verified visually by noting that the δ¹⁵N^α values are significantly different than the δ¹⁵N^β values throughout each reaction (Fig. 1). However, in all five cases, the standard Rayleigh model produces KIE values for N^α and N^β that are approximately equal, highlighting the fact that the standard Rayleigh model inaccurately quantifies ¹⁵N apportionment between N^α and N^β. (If KIE ¹⁵N^α were equal to KIE ¹⁵N^β, the curves for δ¹⁵N^α, δ¹⁵N^β, and δ¹⁵N^{bulk} shown in Fig. 1 would all be on top of each other.)

The inadequacies of the standard Rayleigh model are most evident when N^α and N^β have opposing kinetic isotope effects, i.e., scenarios where there is an inverse isotope effect at one position and a normal isotope effect at the other position (Dataset 3), or when only one position has a non-zero fractionation factor (Datasets 1, 5). In these cases, the standard Rayleigh model fails to predict the correct type of isotope effect for one of the nitrogen atoms. For example, Dataset 1 represents a scenario where there is no isotopic preference at the α position, meaning instantaneous δ¹⁵N^α (δ¹⁵N^{αi}) is always equal to δ¹⁵N^ε, which can be verified visually by noting that accumulated δ¹⁵N^α ≈ δ¹⁵N^ε at the start of the reaction (Fig. 1). Under these circumstances, KIE ¹⁵N^α is equal to 1.0000. However, the standard Rayleigh model predicts a much higher KIE ¹⁵N^α value (1.0208), incorrectly indicating that there is a reasonably strong, normal KIE at the α position. The standard Rayleigh KIE ¹⁵N^β value calculated for Dataset 1 is also too low (1.0200 instead of 1.0417), underestimating the preference for ¹⁴N at the β position. Similarly misleading results are produced for Dataset 5, where there is no isotope effect for N^β, but the standard Rayleigh model predicts a normal isotope effect at this position. Along the same lines, the standard Rayleigh KIE values for Dataset 3 incorrectly indicate that both N^α and N^β have normal isotope effects when, in this case, N^α actually has an inverse isotope effect. Thus, even for a simulated dataset with no experimental error, the standard Rayleigh model produces results that are quantitatively—and, sometimes, qualitatively—incorrect.

480 **Table 3. KIE values for N^{bulk}, N^α, and N^β calculated directly from simulation input values (Actual) or from the standard Rayleigh model or Expanded Rayleigh model applied to error-free simulated datasets.**

Commented [ER27]: Response to community comment: Added text to highlight the connection between figures, calculations, and results.

Deleted: Application of the standard Rayleigh model to each dataset produces KIE ¹⁵N^{bulk} values that

Deleted: (calculated by converting the input ε_{N^{bulk}} value to a KIE value). However,

Deleted: .

Deleted: KIE ¹⁵N^α and KIE ¹⁵N^β.

Deleted: .

Commented [ER28]: Response to community comment: Added text to highlight the connection between calculations and results.

| Simulated dataset | | Model | | | |
|---|--|---------------------|--------------------------------|----------------------------------|----------------------------------|
| | | Actual ^a | Standard Rayleigh ^b | Expanded Rayleigh 1 ^c | Expanded Rayleigh 2 ^c |
| 1 No KIE $^{15}\text{N}^{\alpha}$ Normal KIE $^{15}\text{N}^{\beta}$ | KIE $^{15}\text{N}^{\text{bulk}}$ | 1.0204 | 1.0204 | 1.0204 | 1.0204 |
| | KIE $^{15}\text{N}^{\alpha}$ | 1.0000 | 1.0208 | 1.0000 | 1.0000 |
| | KIE $^{15}\text{N}^{\beta}$ | 1.0417 | 1.0200 | 1.0417 | 1.0417 |
| 2 Normal KIE $^{15}\text{N}^{\alpha}$ Normal KIE $^{15}\text{N}^{\beta}$ | KIE $^{15}\text{N}^{\text{bulk}}$ | 1.0204 | 1.0204 | 1.0204 | 1.0204 |
| | KIE $^{15}\text{N}^{\alpha}$ | 1.0103 | 1.0206 | 1.0103 | 1.0103 |
| | KIE $^{15}\text{N}^{\beta}$ | 1.0308 | 1.0202 | 1.0308 | 1.0308 |
| 3 Inverse KIE $^{15}\text{N}^{\alpha}$ Normal KIE $^{15}\text{N}^{\beta}$ | KIE $^{15}\text{N}^{\text{bulk}}$ | 1.0204 | 1.0204 | 1.0204 | 1.0204 |
| | KIE $^{15}\text{N}^{\alpha}$ | 0.9810 | 1.0212 | 0.9810 | 0.9810 |
| | KIE $^{15}\text{N}^{\beta}$ | 1.0631 | 1.0196 | 1.0631 | 1.0631 |
| 4 Inverse KIE $^{15}\text{N}^{\alpha}$ Inverse KIE $^{15}\text{N}^{\beta}$ | KIE $^{15}\text{N}^{\text{bulk}}$ | 0.9804 | 0.9804 | 0.9804 | 0.9804 |
| | KIE $^{15}\text{N}^{\alpha}$ | 0.9706 | 0.9802 | 0.9707 | 0.9706 |
| | KIE $^{15}\text{N}^{\beta}$ | 0.9903 | 0.9806 | 0.9903 | 0.9903 |
| 5 Normal KIE $^{15}\text{N}^{\alpha}$ No KIE $^{15}\text{N}^{\beta}$ | KIE $^{15}\text{N}^{\text{bulk}}$ | 1.0204 | 1.0204 | 1.0204 | 1.0204 |
| | KIE $^{15}\text{N}^{\alpha}$ | 1.0417 | 1.0200 | 1.0417 | 1.0417 |
| | KIE $^{15}\text{N}^{\beta}$ | 1.0000 | 1.0208 | 1.0000 | 1.0000 |

^a Actual values were calculated directly from the input values used to generate the dataset.

490 ^b KIE values were calculated from $\epsilon_{\text{N-bulk}}$, $\epsilon_{\text{N-}\alpha}$, or $\epsilon_{\text{N-}\beta}$ values obtained via linear regression of $\delta^{15}\text{N}^{\text{bulk}}$, $\delta^{15}\text{N}^{\alpha}$, or $\delta^{15}\text{N}^{\beta}$ against $[-\ln f/(1-f)]$ (see Eq. (2)).

^c For the Expanded Rayleigh model, $\alpha_{\text{N-bulk}}$ was determined with the standard Rayleigh approach, ρ was determined via nonlinear regression (nonlinear model 1 or 2, Eq. (29) or Eq. (30)), and τ was determined by averaging $^{14}\text{N}^{\alpha}/^{14}\text{N}^{\text{bulk}}$ for every step of the reaction.

495 The standard Rayleigh model produces inaccurate results for the individual N atoms in N_2O because the changes in $\delta^{15}\text{N}^{\alpha}$ depend on both KIE $^{15}\text{N}^{\alpha}$ and KIE $^{15}\text{N}^{\beta}$, as do changes in $\delta^{15}\text{N}^{\beta}$. Linear regression of standard Rayleigh plots for $\delta^{15}\text{N}^{\alpha}$ or $\delta^{15}\text{N}^{\beta}$ fails to capture this nuance. To address this issue, we developed the Expanded Rayleigh model that introduces two new parameters, ρ and τ , to define how $^{15}\text{N}^{\text{bulk}}$ and $^{14}\text{N}^{\text{bulk}}$, respectively, are apportioned between N^{α} and N^{β} . Fractionation factors for N^{α} and N^{β} are determined by combining $\alpha_{\text{N-bulk}}$, ρ , and τ (Eq. (21) and Eq. (22)). For experiments conducted at natural abundance of ^{15}N , the value of τ , which is equal to $^{14}\text{N}^{\alpha}/^{14}\text{N}^{\text{bulk}}$, should always be quite close to 0.5, as discussed above. Indeed, each simulated reaction step has a mean value of τ equal to 0.500 for Datasets 1-5 (Tables S9-S13), so the average value of $^{14}\text{N}^{\alpha}/^{14}\text{N}^{\text{bulk}}$ was used to calculate of $\alpha_{\text{N-}\alpha}$ and $\alpha_{\text{N-}\beta}$. To determine the value of ρ that fits the entire dataset as accurately as possible, we performed nonlinear least squares regression using the formula shown in Eq. (29) (nonlinear model 1) or Eq. (30) (nonlinear model 2). For error-free Datasets 1-5, nonlinear least-squares regression with either nonlinear model correctly predicted ρ . Combining these ρ and τ values with $\alpha_{\text{N-bulk}}$ values derived from the standard Rayleigh equation yields KIE $^{15}\text{N}^{\alpha}$

Deleted: have

Deleted: an expanded version of

Commented [ER29]: Reviewer 1 comment (P21, L432): Edited sentence to avoid redundancy.

and KIE $^{15}\text{N}^\beta$ values that are identical to the expected values out to three or four decimal places (Table 3). Thus, unlike the standard Rayleigh approach, the Expanded Rayleigh model successfully recapitulates the kinetic isotope effects for N^α and N^β in an error-free dataset.

3.2 Application of the Expanded Rayleigh model to simulations with error

To test the robustness of the Expanded Rayleigh model, we applied this model to simulated data with error at varying levels of size and skewness (Table 2, Fig. 3, and Figs. S1-S4) and averaged the results from 1000 simulations for each error category (Table 4, Tables S9-S13). These results indicate that at a low level of error (the level of error expected for a typical experiment), the KIE values produced by the Expanded Rayleigh model are quite accurate (absolute difference relative to true values is 0.001, or 0.1%, see Eq. (31)), regardless of the skewness of the standard Rayleigh model residuals (Fig. 6, Figs. S5-S8). The standard deviations for each average KIE $^{15}\text{N}^{\text{bulk}}$, KIE $^{15}\text{N}^\alpha$, or KIE $^{15}\text{N}^\beta$ value (0.0013-0.0014) are also similar to standard error values from typical experiments (~0.0004-0.0021) (Table 5, Table 6) (Sutka et al., 2006; Yang et al., 2014). As the absolute value of the simulated error increases, the average KIE values become less accurate, and the corresponding standard deviations increase. Nonetheless, even at the highest level of error, the maximum absolute relative difference of average KIE values is 0.005 (0.5% difference), and the range of values covered by average KIE \pm standard deviation includes the actual KIE value. Thus, the Expanded Rayleigh model provides a robust method for determining KIE values even when the datasets have a high level of error or skewness.

In contrast to the Expanded Rayleigh model, the KIE $^{15}\text{N}^\alpha$ and KIE $^{15}\text{N}^\beta$ values determined for simulations with error using the standard Rayleigh model alone are much less accurate (Fig. 6, Figs. S5-8). For all simulations derived from Datasets 1-5 at all levels of error and types of skewness, the absolute relative differences of average standard Rayleigh KIE $^{15}\text{N}^\alpha$ and KIE $^{15}\text{N}^\beta$ values ranged from 0.01-0.04 (1-4% difference). These absolute relative differences are 2-40 times higher than the corresponding differences of Expanded Rayleigh model estimates (Fig. 6, Figs. S5-8). Differences of this magnitude are large enough that the true KIE $^{15}\text{N}^\alpha$ and KIE $^{15}\text{N}^\beta$ values typically do not fall within the range covered by average KIE \pm standard deviation calculated using the standard Rayleigh model (data not shown). Thus, as discussed for the no-error simulations, standard Rayleigh KIE $^{15}\text{N}^\alpha$ and KIE $^{15}\text{N}^\beta$ values do not accurately quantify isotopic fractionation at the α and β positions of N_2O and can even lead to inaccurate designation of the type of KIE (*i.e.*, no KIE, normal KIE, or inverse KIE) at either N^α or N^β . In short, for all levels of error and types of skewness tested, the standard Rayleigh model is only accurate for bulk N. *To obtain KIE $^{15}\text{N}^\alpha$ and KIE $^{15}\text{N}^\beta$ values with the same level of accuracy as KIE $^{15}\text{N}^{\text{bulk}}$, the KIE $^{15}\text{N}^{\text{bulk}}$ value obtained via the standard Rayleigh model must be combined with ρ and τ , parameters from the Expanded Rayleigh model.*

The error for KIE $^{15}\text{N}^\alpha$ and KIE $^{15}\text{N}^\beta$ values is derived from three sources: error in $\alpha_{\text{N-bulk}}$, error in ρ , and error in τ . In all of our simulations with added error, the most significant source of error is the error associated with $\alpha_{\text{N-bulk}}$, which is derived from linear regression of $\delta^{15}\text{N}^{\text{bulk}}$ against $[-\ln f/(1-f)]$. As the level of error increases, the accuracy of $\epsilon_{\text{N-bulk}}$ (and thus the accuracy of $\alpha_{\text{N-bulk}}$ and KIE $^{15}\text{N}^{\text{bulk}}$) decreases more significantly than the accuracy of ρ and τ estimates (Tables S9-S13). The



Commented [ER30]: Response to community comment: Added reference to appropriate figures to highlight connection between figures and results.

same trend is apparent for other assessments of model fit. The RMSE values for the nonlinear regression portion of the Expanded Rayleigh model (1 or 2) are always slightly lower than the RMSE values for the standard Rayleigh linear model (Fig. 6, Figs. S5-S8), indicating that the linear and nonlinear models produce similar “goodness of fit” values. Similarly, the p-values for $\epsilon_{N\text{-bulk}}$, the slope of the linear regression model, are always greater than the p-values for ρ , the coefficient extracted from the nonlinear portion of the Expanded Rayleigh model. (In both cases, the null hypothesis is that $KIE = 1$; see Methods.)



545 Out of each set of 1000 simulations at high-level error, a small number of simulated datasets (14-19%) produce linear regression plots where confidence in the slope is low (*i.e.*, $\epsilon_{N\text{-bulk}}$ p-value > 0.05) (data not shown). In contrast, all of the simulations derived from all five datasets with varying types of KIEs produce ρ values that are likely to be significant (p-value < 0.001). Overall, application of the Expanded Rayleigh model to simulated datasets with varying types of error indicates that

550 this method of determining individual fractionation factors for N^{α} and N^{β} is robust even at high levels of error.

(a) Dataset 1: No KIE $^{15}\text{N}^\alpha$, Normal KIE $^{15}\text{N}^\beta$



(a) Dataset 1: No KIE $^{15}\text{N}^\alpha$, Normal KIE $^{15}\text{N}^\beta$



Deleted:

Commented [ER31]: Response to reviewer 1: Replaced Fig. 6 after recalculating simulated δ values using updated Rstandard value. (New calculations use Rstd = 0.0036765 instead of Rstd = 0.003663). Note that this change has essentially no effect on the calculated values.

Commented [ER32]: Response to community comment: Added KIE value.

Figure 6. Comparison of the accuracy and goodness of fit of the standard Rayleigh model and Expanded Rayleigh models 1 and 2.

A. Comparison of the accuracy of KIE $^{15}\text{N}^\alpha$ and KIE $^{15}\text{N}^\beta$ values. For each model, the absolute relative difference for KIE $^{15}\text{N}^\alpha$ and KIE $^{15}\text{N}^\beta$ values (average for 1000 simulated datasets derived from Dataset 1) are shown. Actual values for Dataset 1: $\delta^{15}\text{N}^{\alpha 0} = 0\%$, $\epsilon_{\text{N-bulk}} = -20\%$ (KIE $^{15}\text{N}^{\text{bulk}} = 1.0204$), KIE $^{15}\text{N}^\alpha = 1.0000$, and KIE $^{15}\text{N}^\beta = 1.0417$. Absolute relative difference (Eq. (31)) is the absolute value of the difference between the estimated value and actual value divided by the actual value ($(\text{estimate} - \text{actual})/\text{actual}$). B. Comparison of the average RMSE values for each set of 1000 simulated datasets derived from Dataset 1. Both the standard Rayleigh model and the Expanded Rayleigh model (1 and 2) use $\delta^{15}\text{N}^{\text{bulk}}$ as the dependent variable, so average RMSE values can be compared directly. Lower RMSE values indicate a better goodness of fit. At each error level (low, medium, and high), the absolute relative difference value (A) or average RMSE (B) is depicted with a symbol that represents skewness type as shown in the legend. Note that in most cases these symbols overlap.

565 **Table 4. Precision and accuracy of values calculated with Expanded Rayleigh model 1 using simulated datasets derived from Dataset 1 (no isotope effect for N^α, normal isotope effect for N^β).**

| Model | Level of error | Skewness | Standard Rayleigh R ² a, b | KIE ¹⁵ N ^{bulk} a | KIE ¹⁵ N ^α a | KIE ¹⁵ N ^β a |
|----------------------------------|----------------|----------|---------------------------------------|---------------------------------------|------------------------------------|------------------------------------|
| Actual ^c | None | No | NA | 1.0204 ± NA | 1 ± NA | 1.0417 ± NA |
| Expanded Rayleigh 1 ^d | Low | No | 0.95 ± 0.02 | 1.0202 ± 0.0013 | 0.9998 ± 0.0013 | 1.0414 ± 0.0014 |
| | | Left | 0.94 ± 0.02 | 1.0202 ± 0.0013 | 0.9998 ± 0.0013 | 1.0414 ± 0.0014 |
| | | Right | 0.94 ± 0.02 | 1.0202 ± 0.0013 | 0.9998 ± 0.0013 | 1.0415 ± 0.0014 |
| Expanded Rayleigh 1 ^d | Medium | No | 0.75 ± 0.09 | 1.0196 ± 0.0031 | 0.9992 ± 0.0030 | 1.0408 ± 0.0032 |
| | | Left | 0.74 ± 0.1 | 1.0197 ± 0.0032 | 0.9994 ± 0.0032 | 1.0409 ± 0.0033 |
| | | Right | 0.74 ± 0.1 | 1.0197 ± 0.0031 | 0.9993 ± 0.0031 | 1.0410 ± 0.0033 |
| Expanded Rayleigh 1 ^d | High | No | 0.44 ± 0.17 | 1.0186 ± 0.0060 | 0.9982 ± 0.0060 | 1.0398 ± 0.0061 |
| | | Left | 0.43 ± 0.17 | 1.0190 ± 0.0058 | 0.9987 ± 0.0058 | 1.0401 ± 0.0061 |
| | | Right | 0.43 ± 0.17 | 1.0187 ± 0.0059 | 0.9983 ± 0.0059 | 1.0400 ± 0.0062 |

^a Average value ± standard deviation calculated from 1000 values generated from 1000 simulated datasets.

^b Average R² value for linear regression of δ¹⁵N^{bulk} against [-flnf/(1-f)] (using Eq. (2), the standard Rayleigh equation).

^c Actual values were calculated directly from the input values used to generate the dataset.

570 ^d For Expanded Rayleigh model 1, α_{N-bulk} was determined with the standard Rayleigh approach, ρ was determined via nonlinear regression (nonlinear model 1, Eq. (29)), and τ was determined by averaging ¹⁴N^{α/14}N^{bulk} for every step of the reaction (Eq. (24)). The results for Expanded Rayleigh model 2 (Table S9) are very similar.

3.3 Application of the Expanded Rayleigh model to previously published isotopic data

575 As a proof-of-concept example, we applied the standard Rayleigh model and the Expanded Rayleigh model to previously published experimental data and compared the results. We chose data collected for N₂O biosynthesis by a culture of *M. trichosporium* (*Methylocystis* sp.) grown aerobically in the presence of NH₂OH (Sutka et al., 2006). *Methylocystis* species encode the gene for hydroxylamine oxidoreductase (HAO), an enzyme that not only converts NH₂OH to NO during nitrification, but also produces N₂O under certain conditions (Yamazaki et al., 2014). As *Methylocystis* species lack the gene for cytochrome P460 (which has also been shown to convert NH₂OH to N₂O) (Caranto et al., 2016), the isotopic fractionation observed in this experiment is assumed to be due to only one process: HAO-catalyzed conversion of NH₂OH to N₂O. Indeed, 580 the ε_{N-bulk} value for N₂O produced in this experiment (5.3‰ ± 0.4 for replicate B (KIE ¹⁵N^{bulk} = 0.9947 ± 4e-04, see Table S15) (Sutka et al., 2006)) is similar to the ε_{N-bulk} value of 2.0‰ (KIE ¹⁵N^{bulk} = 0.9980) reported for purified HAO from *Nitrosomonas europaea* (Yamazaki et al., 2014). The differences between δ¹⁵N^α and δ¹⁵N^β (SP) are also very similar for the two experiments, indicating that HAO was the primary source of N₂O production in the cultures of *M. trichosporium*, meaning 585 the Rayleigh model could be applied to this system.

Deleted: R²

Commented [ER33]: Community comment 1: Streamline calculation descriptions by removing redundant info.

Deleted: calculated from 1000 R² values generated from 1000 simulated datasets

Deleted: (

Commented [ER34]: Response to community comment: Added KIE values.

Deleted: very

Application of the standard Rayleigh model (Eq. (2)) to $\delta^{15}\text{N}^{\text{bulk}}$ via linear regression of $\delta^{15}\text{N}^{\text{bulk}}$ against $[-\ln f/(1-f)]$ indicated that $\epsilon_{\text{N-bulk}}$ was $5.3\% \pm 0.4$ and KIE $^{15}\text{N}^{\text{bulk}}$ was 0.9947 ± 0.0004 (inverse isotope effect) (Table 5) for N_2O production by *M. trichosporium*. Similarly, application of the standard Rayleigh model to $\delta^{15}\text{N}^{\alpha}$ and $\delta^{15}\text{N}^{\beta}$ produced a KIE $^{15}\text{N}^{\alpha}$ value of 0.9952 ± 0.0015 and a KIE $^{15}\text{N}^{\beta}$ value of 0.9942 ± 0.0021 . Thus, the standard Rayleigh model predicts an inverse isotope effect for both N^{α} and N^{β} , which theoretically means that both positions are enriched in ^{15}N relative to substrate. However, all of the $\delta^{15}\text{N}^{\beta}$ values (-11.7 to -13.3‰) are more negative than the measured $\delta^{15}\text{N}^{\text{S0}}$ value of -2.3‰ (Fig. 4 and Table S14), indicating that ^{15}N is *depleted* at the β position and that ^{14}N is preferred (normal isotope effect). The $\delta^{15}\text{N}^{\alpha}$ values (21.1 to 22.7‰), in contrast, are more positive than $\delta^{15}\text{N}^{\text{S0}}$, indicating that there is an inverse isotope effect at this position. The Expanded Rayleigh model produces position-specific KIEs that match this qualitative assessment: KIE $^{15}\text{N}^{\alpha} = 0.9779 \pm 0.0004$ (inverse), and KIE $^{15}\text{N}^{\beta} = 1.0121\text{-}1.0122 \pm 0.0004$ (normal) (Table 5). Additionally, the values of ρ and τ do not vary significantly with f (Table S15), indicating that fractionation at the α and β positions appears to be constant. Comparison of the RMSE values for the standard (linear) Rayleigh model and the nonlinear portion of the Expanded Rayleigh model also indicates that both models have a similar goodness of fit. Thus, the Expanded Rayleigh model can be applied to experimental data to accurately assess position-specific KIEs.

605

Table 5. Comparison of standard Rayleigh and Expanded Rayleigh KIE values \pm standard error for N_2O produced from NH_2OH by an axenic culture of *M. trichosporium* (*Methylocystis* sp.). Values were calculated using isotopic data previously published for *M. trichosporium* replicate B (Sutka et al., 2006).

| Model | Standard Rayleigh (linear) RMSE ^a | Nonlinear RMSE ^b | KIE $^{15}N^{bulk}$ ^c | KIE $^{15}N^{\alpha}$ ^c | KIE $^{15}N^{\beta}$ ^c |
|----------------------------------|--|-----------------------------|----------------------------------|------------------------------------|-----------------------------------|
| Standard Rayleigh ^d | 0.07 | NA | $0.9947 \pm 4e-04$ | 0.9952 ± 0.0015 | 0.9942 ± 0.0021 |
| Expanded Rayleigh 1 ^e | 0.07 | 0.16 | $0.9947 \pm 4e-04$ | $0.9779 \pm 4e-04$ | $1.0121 \pm 4e-04$ |
| Expanded Rayleigh 2 ^e | 0.07 | 0.16 | $0.9947 \pm 4e-04$ | $0.9779 \pm 4e-04$ | $1.0122 \pm 4e-04$ |

610 ^a RMSE (root mean square error) for standard Rayleigh model (Eq. (2), linear regression of $\delta^{15}N^{bulk}$ against $[-f]^{lnf}/(1-f)$)

^b RMSE for nonlinear model 1 or 2, Eq. (29) or Eq. (30)

^c Average value \pm standard error

^d KIE values were calculated from ϵ_{N-bulk} , $\epsilon_{N-\alpha}$, or $\epsilon_{N-\beta}$ values obtained via linear regression of $\delta^{15}N^{bulk}$, $\delta^{15}N^{\alpha}$, or $\delta^{15}N^{\beta}$ against $[-f]^{lnf}/(1-f)$.

615 ^e For the Expanded Rayleigh model, bulk values (α_{N-bulk} , ϵ_{N-bulk} , and KIE $^{15}N^{bulk}$) were determined with the standard Rayleigh approach, ρ was determined via nonlinear regression, and τ was determined by averaging $^{14}N^{\alpha}/^{14}N^{bulk}$ for every step of the reaction. Then $\alpha_{N-\alpha}$ and $\alpha_{N-\beta}$ were calculated with Eq. (21) or Eq. (22) and converted to KIE values using Eq. (10). The only difference between Expanded Rayleigh models 1 (Eq. (29)) and 2 (Eq. (30)) is which δ value ($\delta^{15}N^{\alpha}$ or $\delta^{15}N^{\beta}$) is substituted with a ρ -containing expression in the nonlinear model.

620 We also compared the results of applying the standard and Expanded Rayleigh models to previously published isotopic data on NO reduction to N_2O by purified *H. capsulatum* P450 NOR (Yang et al., 2014). In this experiment, $\delta^{15}N^{bulk}$ varied linearly with $[-f]^{lnf}/(1-f)$ ($R^2 = 0.68-0.94$), and averaging the slopes of the standard Rayleigh plots for each individual replicate indicated that ϵ_{N-bulk} was equal to $9.10\text{‰} \pm 1.42$ and KIE $^{15}N^{bulk}$ was equal to 0.9910 ± 0.0014 (Table 6 and Table S17). In other words, on average, N_2O became depleted in ^{15}N (*i.e.*, $\delta^{15}N^{bulk}$ decreased) as the reaction proceeded (Fig. 5 and Table S16). However, the N^{α} position became more enriched in ^{15}N as the reaction progressed (*i.e.*, $\delta^{15}N^{\alpha}$ increased as f decreased). Consequently, ρ increased (and τ decreased) over the course of the reaction (Table S17), and the observed values of $\alpha_{N-\alpha}$ and $\alpha_{N-\beta}$ also varied with f , in contrast to Assumption 6 (above). Thus, the Expanded Rayleigh model, which is based on the assumption that $\alpha_{N-\alpha}$ and $\alpha_{N-\beta}$ remain constant over the course of the reaction, does not fit this data optimally. Therefore, using nonlinear regression (Eq. 29 or Eq. 30) to determine ρ is not appropriate in this case. Using a modified version of the Expanded Rayleigh model, however, we calculated ρ for each individual timepoint and estimated apparent KIE $^{15}N^{\alpha}$ and KIE $^{15}N^{\beta}$ values for the beginning and end of the observed range of f values (see Methods section 2.9). Our estimated KIE $^{15}N^{\alpha}$ values ranged from 0.9823 ± 0.0016 (earlier in the reaction) to 0.9781 ± 0.0016 (later in the reaction), indicating that the N^{α} position is subject to an inverse isotope effect throughout the reaction. On the other hand, the KIE $^{15}N^{\beta}$ estimates ranged from 0.9998 ± 0.0015 to 1.0041 ± 0.0013 , suggesting that there is essentially no apparent isotope effect at the β position, especially

Commented [ER35]: Reviewer 1 comment (Table 5): Made model descriptions more specific to clarify that “Expanded” refers to the Expanded Rayleigh model.

Formatted Table

Deleted: (nonlinear model 1 or 2, Eq. (29) or Eq. (30))

Commented [ER36]: Reviewer 1 comment (Table 5): Added text to make the difference between Expanded Rayleigh models 1 and 2 clearer.

Deleted: ¶

Commented [ER37]: Response to community comment: Added text to clarify why the Expanded Rayleigh model was not applied to the whole dataset for the P450 NOR data.

early in the reaction. These results are consistent with a qualitative assessment of the δ values. For example, the $\delta^{15}\text{N}^\alpha$ values (-25.8 to -31.0‰) are more positive than the calculated $\delta^{15}\text{N}^{\text{S0}}$ values (*i.e.*, the intercept of the standard Rayleigh plots for $\delta^{15}\text{N}^{\text{bulk}}$), which range from -44.7‰ to -48.3‰. This indicates that the α position is enriched in ^{15}N relative to the substrate.

640 On the other hand, the $\delta^{15}\text{N}^\beta$ values near the start of the reaction (-45.2 to -48.3‰) are similar to the calculated $\delta^{15}\text{N}^{\text{S0}}$ values, suggesting that there is very little isotopic fractionation at the β position.

In contrast to the Expanded Rayleigh model estimates, the standard Rayleigh calculations yield a normal isotope effect for N^α ($\text{KIE } ^{15}\text{N}^\alpha = 1.0127 \pm 0.0030$) and an inverse isotope effect for N^β ($\text{KIE } ^{15}\text{N}^\beta = 0.9694 \pm 0.002$). (Note that

645 applying the standard Rayleigh model to individual observations and averaging KIEs for the early and later parts of the reaction yields similar results to applying the standard Rayleigh model to the entire reaction (Table 6, Table S17); see SI for details.)

While it is possible that N^β exhibits an apparent inverse isotope effect near the beginning of the reaction, the magnitude of this effect is likely much smaller (*i.e.*, early $\text{KIE } ^{15}\text{N}^\beta$ is much closer to 1) than the standard Rayleigh calculations suggest. More strikingly, the standard Rayleigh prediction that N^α is subject to a normal isotope effect (^{14}N preferred) is clearly incorrect, as the α position is enriched in ^{15}N relative to the initial substrate. Moreover, as ^{15}N is depleted from the substrate pool, the fraction of ^{15}N incorporated into the α position *increases*, indicating that the preference for ^{15}N at N^α increases (becomes more inverse) as the reaction proceeds. Overall, analysis of the P450 NOR data demonstrates that even for reactions where the apparent position-specific KIEs vary, the Expanded Rayleigh model produces considerably more accurate KIE estimates than the standard Rayleigh model.

655

Commented [ER38]: Response to community comment: Added results for applying the standard Rayleigh model to individual observations.

Table 6. Comparison of standard Rayleigh and Expanded Rayleigh KIE values \pm standard error for N₂O production from NO by purified *Histoplasma capsulatum* (fungal) P450 NOR [calculated using previously published isotopic data (Yang et al., 2014)].

| Model | Extent of reaction (range of f) | | KIE ¹⁵ N ^{bulk} a | KIE ¹⁵ N ^α a | KIE ¹⁵ N ^β a |
|--------------------------------|---------------------------------|-----------|---------------------------------------|------------------------------------|------------------------------------|
| Standard Rayleigh ^b | All | 0.42-0.87 | 0.9910 \pm 0.0014 | 1.0127 \pm 0.0030 | 0.9694 \pm 0.0022 |
| Standard Rayleigh ^c | Early | 0.77-0.81 | 0.9908 \pm 0.0013 | 1.0130 \pm 0.0031 | 0.9687 \pm 0.0030 |
| Standard Rayleigh ^c | Late | 0.47-0.52 | 0.9909 \pm 0.0015 | 1.0121 \pm 0.0029 | 0.9698 \pm 0.0018 |
| Expanded Rayleigh ^d | Early | 0.77-0.81 | 0.9910 \pm 0.0014 | 0.9823 \pm 0.0016 | 0.9998 \pm 0.0015 |
| Expanded Rayleigh ^d | Late | 0.47-0.52 | 0.9910 \pm 0.0014 | 0.9781 \pm 0.0016 | 1.0041 \pm 0.0013 |

Commented [ER39]: Response to community comment: Added values calculated by applying the standard Rayleigh model to individual observations and averaging the results from early or late observations.

Deleted: °

Commented [ER40]: Response to reviewer comment: Made model descriptions more specific to clarify that “Expanded” refers to the Expanded Rayleigh model.

Deleted: °

Deleted: calculated for values from three separate replicates (standard Rayleigh model KIEs and Expanded Rayleigh KIE ¹⁵N^{bulk}) or for six (early) or seven (late) individual observations (Expanded Rayleigh model KIE ¹⁵N^α and KIE ¹⁵N^β). Individual early-reaction or late-reaction KIEs from all three replicates were pooled and averaged.

660 ^a Average value \pm standard deviation.

^b KIE values were calculated from $\epsilon_{N\text{-bulk}}$, $\epsilon_{N\text{-}\alpha}$, or $\epsilon_{N\text{-}\beta}$ values obtained via linear regression of $\delta^{15}N^{\text{bulk}}$, $\delta^{15}N^{\alpha}$, or $\delta^{15}N^{\beta}$ against $[-\ln f/(1-f)]$. The standard Rayleigh model values presented here differ slightly from the previously published values (Yang et al., 2014) due to our exclusion of the earliest observation(s) from each replicate (*i.e.*, observations with the highest values of f were excluded).

665 ^c For the standard Rayleigh model applied to individual observations, $\epsilon_{N\text{-bulk}}$, $\epsilon_{N\text{-}\alpha}$, or $\epsilon_{N\text{-}\beta}$ values were determined using Eq. (S24); the y-intercept listed in that equation corresponds to the y-intercept of $\delta^{15}N^{\text{bulk}}$, $\delta^{15}N^{\alpha}$, or $\delta^{15}N^{\beta}$ against $[-\ln f/(1-f)]$ (determined by linear regression of the data from each replicate). KIE values for six (early) or seven (late) individual observations were pooled and averaged.

Deleted: °

670 ^d For the Expanded Rayleigh model applied to individual observations, bulk values ($\alpha_{N\text{-bulk}}$, $\epsilon_{N\text{-bulk}}$, and KIE ¹⁵N^{bulk}) were determined with the standard Rayleigh approach. ρ was calculated for each observation using Eq. (23) ($\rho = {}^{15}N^{\alpha}/{}^{15}N^{\text{bulk}}$), and τ was determined for every step of the reaction using Eq. (24) ($\tau = {}^{14}N^{\alpha}/{}^{14}N^{\text{bulk}}$). Then $\alpha_{N\text{-}\alpha}$ and $\alpha_{N\text{-}\beta}$ were calculated for each individual observation with Eq. (21) or Eq. (22) and converted to KIE values using Eq. (10). KIE ¹⁵N^α and KIE ¹⁵N^β values for six (early) or seven (late) individual observations were pooled and averaged.

4 Discussion

675 4.1 Proper application of the Expanded Rayleigh model

The Rayleigh model has been used to study kinetic isotope effects for decades and has been adapted many different times for use in specific scenarios. However, to our knowledge, this is the first examination of how the Rayleigh model applies to a process where two heavy atoms from the same substrate pool (*e.g.*, NO or NH₂OH) are incorporated into non-equivalent positions in the product (*e.g.*, N₂O). Our simulations of different N₂O-producing scenarios demonstrate that the original Rayleigh model should not be applied to the individual atoms in such cases as this produces inaccurate results. However, the new Expanded Rayleigh model presented here can be used to accurately determine individual kinetic isotope effects. This

expanded model allows for a more granular analysis than would be possible from determining only bulk kinetic isotope effects and SP.

In practice, when can the Expanded Rayleigh model be used? To develop the Expanded Rayleigh model, we employed the same assumptions that the original model is founded on. Therefore, any situations where the original Rayleigh model applies will also likely be suitable for the Expanded Rayleigh model. More specifically, the Expanded Rayleigh model will likely produce accurate results for any closed system where N₂O is produced from a substrate pool with ¹⁵N at or near natural abundance, *as long as the rate-determining step(s) for each atom remains constant*. Thus, the Expanded Rayleigh model is ideally suited for studies of isolated N₂O-synthesizing enzymes, as this experimental setup provides a closed system where N₂O is produced by a single catalytic process.

One method for confirming that the rate-determining steps of N₂O synthesis are constant is to determine if $\epsilon_{N\text{-bulk}}$ and SP are constant for a particular dataset. If enzymatic catalysis is rate-limiting throughout a time course experiment, $\epsilon_{N\text{-bulk}}$ will be constant, and the slope of a standard Rayleigh plot should be linear. In contrast, if substrate diffusion (or some other process) becomes rate-limiting, curvilinear behavior may be exhibited in a standard Rayleigh plot, as has been observed for multi-step N₂O production in microbial cultures (Sutka et al., 2008; Haslun et al., 2018). Of course, to use the Expanded Rayleigh model, $\epsilon_{N\text{-}\alpha}$ and $\epsilon_{N\text{-}\beta}$ must be constant as well. To verify that the individual fractionation factors are constant, one could plot $\delta^{15}N^{\alpha}$ and $\delta^{15}N^{\beta}$ against $[-\ln f/(1-f)]$ (*i.e.*, create standard Rayleigh plots for $\delta^{15}N^{\alpha}$ and $\delta^{15}N^{\beta}$) and confirm that these plots are linear and roughly parallel. Alternatively, SP (the difference of $\delta^{15}N^{\alpha}$ and $\delta^{15}N^{\beta}$, Eq. (32)) can be monitored over time. A lack of a significant trend in SP values is a reasonably good indication that $\epsilon_{N\text{-}\alpha}$ and $\epsilon_{N\text{-}\beta}$ are constant. In studies of microbial cultures where N₂O is predominantly produced by a single type of enzyme (*e.g.*, bacterial NOR, fungal NOR, or HAO), SP generally remains constant as substrate is consumed (Toyoda et al., 2005; Sutka et al., 2006; Sutka et al., 2008; Haslun et al., 2018). The fact that SP is constant for a variety of N₂O production pathways in a microbial cell suggests that enzyme-specific fractionation at N^α and N^β typically remains constant, meaning the Expanded Rayleigh model can likely be accurately applied to a variety of N₂O-producing enzymes.

At present, there is a scarcity of studies reporting N₂O fractionation by purified enzymes for multiple values of *f*, limiting our ability to test the Expanded Rayleigh model with published experimentally-derived datasets. We were, however, able to apply the Expanded Rayleigh model to previously published data on *M. trichosporium* cultures (Sutka et al., 2006), whose HAO-catalyzed oxidation of NH₂OH was the primary source of N₂O and displayed constant values of $\epsilon_{N\text{-bulk}}$ and SP. Thus, we were able to calculate observed values of KIE ¹⁵N^α and KIE ¹⁵N^β that serve as reasonable estimates of HAO-specific isotope effects (Table 5). Additionally, we have used the Expanded Rayleigh model to re-evaluate isotopic data our group has collected for purified fungal P450 NOR (Table 6) (Yang et al., 2014). The value of $\epsilon_{N\text{-bulk}}$ for P450 NOR appears to be constant, but SP increased by 14‰ as the fraction of NO reduced changed from 10% to 50% (Yang et al., 2014). Given the proposed P450 NOR reaction mechanism, the observed isotope effect for the first NO molecule is influenced both by an equilibrium isotope effect related to NO binding and by an intrinsic KIE associated with catalysis (Yang et al., 2014). Conversely, the rate-

determining step(s) for the second NO molecule are likely only associated with catalysis. Thus, the isotopic fractionation for the first nitrogen atom to be incorporated will be more sensitive to changes in NO concentration as the reaction progresses than the second N atom, leading to SP values that are not constant. Even in these circumstances, however, the Expanded Rayleigh model estimates for KIE $^{15}\text{N}^\alpha$ and KIE $^{15}\text{N}^\beta$ are a significant improvement on the previously published standard Rayleigh estimates, which incorrectly predict a normal isotope effect at the α position. Overall, the Expanded Rayleigh model fits the available data on isotopic fractionation during N_2O biosynthesis reasonably well, although we recommend collecting isotopic data across a range of f values to confirm that SP (and therefore KIE $^{15}\text{N}^\alpha$ and KIE $^{15}\text{N}^\beta$) remains constant when using this model.

As outlined above (“Assumptions”), the Expanded Rayleigh model is most accurate for reactions where the isotopic fractionation factors ($\epsilon_{\text{N-bulk}}$, $\epsilon_{\text{N-}\alpha}$, and $\epsilon_{\text{N-}\beta}$) and $\delta^{15}\text{N}^{\text{std}}$ values are not too extreme. This stipulation, which also applies to the standard Rayleigh equation, is necessary in part because an approximation introduced by Mariotti and colleagues (Mariotti et al., 1981) decreases in accuracy the farther $\delta^{15}\text{N}^{\text{std}}$ and $\delta^{15}\text{N}^{\text{s}}$ are from 0‰. Additionally, extremely high, positive values of $\delta^{15}\text{N}^{\text{std}}$ or $\epsilon_{\text{N-bulk}}$ would result in large, positive product δ values, corresponding to an increased abundance of ^{15}N in the product. As both the nonlinear Rayleigh model and the underlying calculation of $\delta^{15}\text{N}^\beta$ values depend on the assumption that the amount of ^{15}N in N_2O is close to natural abundance, the accuracy of the Expanded Rayleigh model would decrease in such circumstances. Thus, applying the Expanded Rayleigh model to experiments with spiked ^{15}N is not advised. Simulations of δ values calculated without using Mariotti’s approximation indicate that for absolute values of $\epsilon_{\text{N-bulk}}$ or $\delta^{15}\text{N}^{\text{std}}$ less than or equal to 50‰, the level of error introduced by Mariotti’s approximation is similar to the expected level of experimental error (*i.e.*, absolute relative difference in KIE values of ~ 0.001) (Tables S1-S2).

The magnitude of $\epsilon_{\text{N-}\alpha}$ and $\epsilon_{\text{N-}\beta}$ must also be considered when applying the Expanded Rayleigh model. In theory, if isotopic enrichment at N^α or N^β is very large, the assumption that essentially equal amounts of ^{14}N are apportioned to each position (*i.e.*, that $\tau \approx 0.5$) would be violated, and Eq. (15) would no longer hold. However, violating this assumption would require unusually large heavy-atom kinetic isotope effects or a spiked sample. As outlined in Table S3, for a simulated reaction where $\epsilon_{\text{N-bulk}}$ is set at -20‰ (KIE $^{15}\text{N}^{\text{bulk}} = 1.0204$) and $\delta^{15}\text{N}^{\text{std}}$ is 0‰, individual KIE values would have to be significantly lower than 0.96 or greater than 1.08 to introduce error greater than analytical error. These KIE $^{15}\text{N}^\alpha$ or KIE $^{15}\text{N}^\beta$ values correspond to p values less than 0.47 or greater than 0.53. In practice, our group has found that for two different types of N_2O -producing enzymes, p is nowhere near these extremes. For example, for purified P450 NOR (Yang et al., 2014) and for HAO-catalyzed NH_2OH oxidation in an axenic culture (Sutka et al., 2006), p does not exceed 0.509 (Table S15, Table S17). Overall, we anticipate that the Expanded Rayleigh model will be applicable to a wide variety of N_2O biosynthesis reactions.

Deleted: (KIE $^{15}\text{N}^{\text{bulk}} = 1.0204$)

4.2 Other practical considerations

The Expanded Rayleigh model presented here does not require measurement of $\delta^{15}\text{N}^{\text{std}}$. This is advantageous for N_2O synthesis reactions where NO is the substrate because directly measuring an accurate $\delta^{15}\text{N}$ value for NO is quite challenging

due to the reactivity of NO, the lack of isotopic standards for NO, and the fact that $^{14}\text{N}^{16}\text{O}$ has the same molecular weight as $^{15}\text{N}^{15}\text{N}$. However, $\delta^{15}\text{N}^{\text{S0}}$ can be directly measured when NH_2OH is used as the substrate for N_2O production. Determining $\delta^{15}\text{N}^{\text{S0}}$ experimentally may be useful in validating application of the standard Rayleigh model, as the intercept of the standard Rayleigh model is expected to be equal to $\delta^{15}\text{N}^{\text{S0}}$. For example, we determined that the intercept of the standard Rayleigh plot for N_2O produced by NH_2OH oxidation by *M. trichosporium* is $0.5 \pm 0.3\%$, which is similar to the measured $\delta^{15}\text{N}^{\text{S0}}$ value of -2.3% (Sutka et al., 2006), confirming that the standard Rayleigh model is applicable for this dataset. Thus, when feasible, measuring $\delta^{15}\text{N}^{\text{S0}}$ would complement the Expanded Rayleigh approach.

The nonlinear portion of the Expanded Rayleigh model (used to determine ρ) can be written in two forms, nonlinear model 1 (Eq. (29)) and nonlinear model 2 (Eq. (30)). Both models are based on the assumption that $\delta^{15}\text{N}^{\text{bulk}}$ is equal to the average of $\delta^{15}\text{N}^{\text{a}}$ and $\delta^{15}\text{N}^{\text{b}}$ (Eq. (15)). Therefore, both models are expected to produce very similar results as long as that assumption holds. Indeed, as we have demonstrated here, both models yield nearly identical values of ρ , KIE $^{15}\text{N}^{\text{a}}$, and KIE $^{15}\text{N}^{\text{b}}$ for simulated datasets (Fig. 6, Figs. S5-S8, Tables S9-S13), and we have obtained similar results for the available experimental data on N_2O production from NH_2OH (Table 5). Thus, either version of the Expanded Rayleigh model may be used. If there is any doubt about which version to use, we recommend selecting the model that has a lower RMSE value (and thus fits the data better).

5 Conclusions

In this manuscript, we presented the Expanded Rayleigh model, a novel adaptation of the Rayleigh distillation equation that describes the position-specific isotopic enrichment that occurs at N^{a} and N^{b} during N_2O synthesis. Using simulated datasets representing multiple different combinations of KIEs, we demonstrated that the Expanded Rayleigh model accurately recapitulates KIE $^{15}\text{N}^{\text{a}}$ and KIE $^{15}\text{N}^{\text{b}}$ for a wide variety of scenarios. Our simulations also demonstrate that this new model is robust even when applied to skewed data and/or data with a high level of error. Additionally, we have shown the Expanded Rayleigh model fits the experimentally-measured isotopic data for N_2O production from NH_2OH quite well (Sutka et al., 2006) and provides significantly improved estimates for the position-specific KIEs for P450 NOR-catalyzed N_2O synthesis. Thus, the Expanded Rayleigh model provides a reliable method for quantifying isotopic fractionation at N^{a} and N^{b} , and it promises to be a valuable tool for experimentally probing the catalytic mechanisms of N_2O -synthesizing enzymes. Finally, we note that the Expanded Rayleigh model will also likely be appropriate for the analysis of other reactions where two atoms from the same substrate pool are incorporated into distinct, non-exchangeable positions in the product.

Code and data availability

All code used in this manuscript is available at:
<https://github.com/GLBRC/position-specific-KIE>

The results for each of the 1000 simulations with error (derived from Datasets 1-5) can be found at:

<https://doi.org/10.5281/zenodo.13323399> |

Commented [ER41]: Updated link: New link points to simulated values calculated with updated Rstandard value.

790 **Author contribution**

EDR, NEO, and ELH conceived this study. EDR developed the model and analyzed the data with contributions from WM, NEO, and ELH. WM provided critical input for statistical analysis and coding. EDR prepared the original draft of the manuscript and figures, and all authors participated in editing and revision of the manuscript.

Deleted: <https://zenodo.org/records/10888931>

Competing interests

795 The authors declare that they have no conflict of interest.

Acknowledgements

We thank Dr. Julius Campeciño and Dr. Helen Kreuzer for a careful reading of a draft of this manuscript and for their insightful comments and questions. We also thank Dr. Julius Campeciño for helpful discussions during initial development of the Expanded Rayleigh model. We also acknowledge an anonymous reviewer whose comments first alerted us to the complications associated with position-specific application of the Rayleigh model.

This material is based upon work supported by the Great Lakes Bioenergy Research Center, U.S. Department of Energy, Office of Science, Biological and Environmental Research Program under Award Number DE-SC0018409.

References

- 805 Azzalini, A.: The R package 'sn': The Skew-Normal and Related Distributions such as the Skew-t and the SUN (version 2.1.1). URL <http://azzalini.stat.unipd.it/SN/>; <https://cran.r-project.org/package=sn> [code], 2023.
- Baty, F., Ritz, C., Charles, S., Brutsche, M., Flandrois, J.-P., and Delignette-Muller, M.-L.: A Toolbox for Nonlinear Regression in R: The Package nlstools, *J Stat Softw*, 66, 1 - 21, 10.18637/jss.v066.i05, 2015.
- 810 Bigeleisen, J. and Wolfsberg, M.: Theoretical and experimental aspects of isotope effects in chemical kinetics, *Adv Chem Phys*, 1, 15-76, 1958.

- Caranto, J. D., Vilbert, A. C., and Lancaster, K. M.: *Nitrosomonas europaea* cytochrome P460 is a direct link between nitrification and nitrous oxide emission, *PNAS*, 113, 14704-14709, 10.1073/pnas.1611051113, 2016.
- 815 Frame, C. H. and Casciotti, K. L.: Biogeochemical controls and isotopic signatures of nitrous oxide production by a marine ammonia-oxidizing bacterium, *Biogeosciences*, 7, 2695-2709, 10.5194/bg-7-2695-2010, 2010.
- Haslun, J. A., Ostrom, N. E., Hegg, E. L., and Ostrom, P. H.: Estimation of isotope variation of N₂O during denitrification by *Pseudomonas aureofaciens* and *Pseudomonas chlororaphis*: implications for N₂O source apportionment, *Biogeosciences*, 15, 3873-3882, 10.5194/bg-15-3873-2018, 2018.
- 820 Hippel, P. v.: Skewness, in: *International Encyclopedia of Statistical Science*, edited by: Lovric, M., Springer Berlin Heidelberg, Berlin, Heidelberg, 1340-1342, 10.1007/978-3-642-04898-2_525, 2011.
- Junk, G. A. and Svec, H. J.: Nitrogen isotope abundance measurements, US atomic energy commission. Office of technical information., ISC 1138, 1958.
- Kalpić, D., Hlupić, N., and Lovrić, M.: Student's t-Tests, in: *International Encyclopedia of Statistical Science*, edited by: Lovric, M., Springer Berlin Heidelberg, Berlin, Heidelberg, 1559-1563, 10.1007/978-3-642-04898-2_641, 2011.
- 825 Komsta, L. and Novomestky, F.: moments: Moments, Cumulants, Skewness, Kurtosis and Related Tests. R package version 0.14.1, <https://CRAN.R-project.org/package=moments>. [code], 2022.
- Kuypers, M. M. M., Marchant, H. K., and Kartal, B.: The microbial nitrogen-cycling network, *Nat Rev Microbiol*, 16, 263-276, 10.1038/nrmicro.2018.9, 2018.
- 830 Trends in globally-averaged CH₄, N₂O, and SF₆ determined from NOAA Global Monitoring Laboratory measurements. Version 2023-11, last access: 11/27/2023.
- Lehnert, N., Dong, H. T., Harland, J. B., Hunt, A. P., and White, C. J.: Reversing nitrogen fixation, *Nat Rev Chem*, 2, 278-289, 10.1038/s41570-018-0041-7, 2018.
- Mariotti, A., Germon, J. C., Hubert, P., Kaiser, P., Letolle, R., Tardieux, A., and Tardieux, P.: Experimental determination of nitrogen kinetic isotope fractionation: Some principles; illustration for the denitrification and nitrification processes, *Plant Soil*, 62, 413-430, 10.1007/BF02374138, 1981.
- 835 R Core Team: R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>. [code], 2022.
- Ravishankara, A. R., Daniel, J. S., and Portmann, R. W.: Nitrous oxide (N₂O): The dominant ozone-depleting substance emitted in the 21st century, *Science (New York, N.Y.)*, 326, 123-125, doi:10.1126/science.1176985, 2009.
- 840 Ritz, C. and Streibig, J. C.: *Nonlinear Regression with R*. [electronic resource], 1st ed. 2008., Use R!, Springer New York 2008.
- Romão, C. V., Vicente, J. B., Borges, P. T., Frazão, C., and Teixeira, M.: The dual function of flavodiiron proteins: oxygen and/or nitric oxide reductases, *J Biol Inorg Chem*, 21, 39-52, 10.1007/s00775-015-1329-4, 2016.

- 845 Scott, K. M., Lu, X., Cavanaugh, C. M., and Liu, J. S.: Optimal methods for estimating kinetic isotope effects from different forms of the Rayleigh distillation equation, *Geochim Cosmochim Acta*, 68, 433-442, 10.1016/s0016-7037(03)00459-9, 2004.
- Skrzypek, G. and Dunn, P. J. H.: Absolute isotope ratios defining isotope scales used in isotope ratio mass spectrometers and optical isotope instruments, *Rapid Communications in Mass Spectrometry*, 34, e8890, <https://doi.org/10.1002/rcm.8890>, 2020.
- 850 Stocker, T. F., Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley, Press, C. U. (Ed.): IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge, United Kingdom and New York, NY, USA2013.
- summary.lm: Summarizing Linear Model Fits. R package stats (version 3.6.2): <https://www.rdocumentation.org/packages/stats/versions/3.6.2/topics/summary.lm>, last access: 12/12/2023.
- 855 Sutka, R. L., Adams, G. C., Ostrom, N. E., and Ostrom, P. H.: Isotopologue fractionation during N₂O production by fungal denitrification, *Rapid Commun Mass Sp*, 22, 3989-3996, 10.1002/rcm.3820, 2008.
- Sutka, R. L., Ostrom, N. E., Ostrom, P. H., Gandhi, H., and Breznak, J. A.: Nitrogen isotopomer site preference of N₂O produced by *Nitrosomonas europaea* and *Methylococcus capsulatus* Bath, *Rapid Commun Mass Sp*, 17, 738-745, 860 10.1002/rcm.968, 2003.
- Sutka, R. L., Ostrom, N. E., Ostrom, P. H., Gandhi, H., and Breznak, J. A.: Nitrogen isotopomer site preference of N₂O produced by *Nitrosomonas europaea* and *Methylococcus capsulatus* Bath, *Rapid Commun Mass Sp*, 18, 1411-1412, <https://doi.org/10.1002/rcm.1482>, 2004.
- 865 Sutka, R. L., Ostrom, N. E., Ostrom, P. H., Breznak, J. A., Gandhi, H., Pitt, A. J., and Li, F.: Distinguishing nitrous oxide production from nitrification and denitrification on the basis of isotopomer abundances, *Appl Environ Microb*, 72, 638-644, 10.1128/aem.72.1.638-644.2006, 2006.
- Tian, H., Xu, R., Canadell, J. G., Thompson, R. L., Winiwarter, W., Suntharalingam, P., Davidson, E. A., Ciais, P., Jackson, R. B., Janssens-Maenhout, G., Prather, M. J., Regnier, P., Pan, N., Pan, S., Peters, G. P., Shi, H., Tubiello, F. N., Zaehle, S., Zhou, F., Arneeth, A., Battaglia, G., Berthet, S., Bopp, L., Bouwman, A. F., Buitenhuis, E. T., Chang, J., 870 Chipperfield, M. P., Dangal, S. R. S., Dlugokencky, E., Elkins, J. W., Eyre, B. D., Fu, B., Hall, B., Ito, A., Joos, F., Krummel, P. B., Landolfi, A., Laruelle, G. G., Lauerwald, R., Li, W., Lienert, S., Maavara, T., MacLeod, M., Millet, D. B., Olin, S., Patra, P. K., Prinn, R. G., Raymond, P. A., Ruiz, D. J., van der Werf, G. R., Vuichard, N., Wang, J., Weiss, R. F., Wells, K. C., Wilson, C., Yang, J., and Yao, Y.: A comprehensive quantification of global nitrous oxide sources and sinks, *Nature*, 586, 248-256, 10.1038/s41586-020-2780-0, 2020.
- 875 Toyoda, S. and Yoshida, N.: Determination of nitrogen isotopomers of nitrous oxide on a modified isotope ratio mass spectrometer, *Anal Chem*, 71, 4711-4718, 10.1021/ac9904563, 1999.

Toyoda, S., Mutobe, H., Yamagishi, H., Yoshida, N., and Tanji, Y.: Fractionation of N₂O isotopomers during production by denitrifier, *Soil Biol Biochem*, 37, 1535-1545, <https://doi.org/10.1016/j.soilbio.2005.01.009>, 2005.

Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag New York. [code], 2016.

880 Yamazaki, T., Hozuki, T., Arai, K., Toyoda, S., Koba, K., Fujiwara, T., and Yoshida, N.: Isotopomeric characterization of nitrous oxide produced by reaction of enzymes extracted from nitrifying and denitrifying bacteria, *Biogeosciences*, 11, 2679-2689, 10.5194/bg-11-2679-2014, 2014.

Yang, H., Gandhi, H., Ostrom, N. E., and Hegg, E. L.: Isotopic fractionation by a fungal P450 nitric oxide reductase during the production of N₂O, *Environ Sci Technol*, 48, 10707-10715, 10.1021/es501912d, 2014.

885