

# Spatial-temporal patterns of anthropogenic and biomass burning contributions on air pollution and mortality burden changes in India from 1995 to 2014

Bin Luo<sup>1</sup>, Yuqiang Zhang<sup>1</sup>, Tao Tang<sup>2,3</sup>, Hongliang Zhang<sup>4</sup>, Jianlin Hu<sup>5</sup>, Jiangshan Mu<sup>1</sup>, Wenxing Wang<sup>1</sup>, Likun Xue<sup>1</sup>

<sup>1</sup>Environment Research Institute, Shandong University, Qingdao 266237, China

<sup>2</sup>School of the Environment, Yale University, New Haven, CT 06511, USA

<sup>3</sup>Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

<sup>4</sup>Department of Environmental Science and Engineering, Fudan University, Shanghai 200438, China

<sup>5</sup>Jiangsu Key Laboratory of Atmospheric Environment Monitoring and Pollution Control, Collaborative Innovation Center of Atmospheric Environment and Equipment Technology, Nanjing University of Information Science & Technology, Nanjing 210044, China

*Correspondence to:* Yuqiang Zhang (yuqiang.zhang@sdu.edu.cn), Likun Xue (xuelikun@sdu.edu.cn)

**Abstract.** Anthropogenic (ANTHRO) and biomass burning (BB) emissions are the major sources of ambient air pollution. India has experienced a dramatic deterioration in air quality over the past few decades, but no systematic assessment has been made to investigate the individual contributions of ANTHRO and BB emissions changes over the long term in India, extension to non-urban areas. In this study, we conducted a comprehensive analysis of the long-term trends of particulate matter with aerodynamic diameters  $< 2.5 \mu\text{m}$  ( $\text{PM}_{2.5}$ ) and ozone ( $\text{O}_3$ ) in India and their mortality burden changes from 1995 to 2014, using a state-of-the-art high-resolution global chemical transport model (CAM-chem). Our simulations revealed a substantial nationwide increase in annual mean  $\text{PM}_{2.5}$  ( $6.71 \mu\text{g m}^{-3} \text{decade}^{-1}$ ) and  $\text{O}_3$  ( $7.08 \text{ ppbv decade}^{-1}$ ), with the Indo-Gangetic Plain (IGP) and eastern central India as hotspots for  $\text{PM}_{2.5}$  and  $\text{O}_3$  trend changes individually. Noteworthy substantial  $\text{O}_3$  decreases were observed in the northern IGP which were potentially linked to  $\text{NO}$  titration due to a surge in  $\text{NO}_x$  emissions. Sensitivity analyses highlighted ANTHRO emissions as primary contributors to rising  $\text{PM}_{2.5}$  and  $\text{O}_3$ , while BB played a prominent role in winter and spring. In years with high BB activity, the contributions from BB emissions on both  $\text{PM}_{2.5}$  and  $\text{O}_3$  changes were comparable with or even exceeding ANTHRO emissions in specific areas. We further estimated that the elevated air pollutants were associated with increased premature mortality attributable to  $\text{PM}_{2.5}$  and  $\text{O}_3$ , leading to 97.83 K and 73.91 K per decade. Despite a per capita decrease in the IGP region, the increased population offset its effectiveness.

## 1 Introduction

Air pollution is among the most detrimental environmental factors to human health. According to the World Health Organization (WHO) database, 99% of the global population lives in areas where air quality surpasses WHO guideline limits

(WHO: Air Pollution, World Health Organization, available at: [https://www.who.int/health-topics/air-pollution#tab=tab\\_1](https://www.who.int/health-topics/air-pollution#tab=tab_1), last access: 21 September 2024). The two most concerned pollutants, particulate matter with aerodynamic diameters  $< 2.5 \mu\text{m}$  ( $\text{PM}_{2.5}$ ) and ozone ( $\text{O}_3$ ), can cause significant damage to the human heart and lungs (Hoek et al., 2013; Hystad et al., 2013; Villeneuve et al., 2015), potentially leading to premature death when exposed over extended periods (Dedoussi et al., 2020; Fuller et al., 2022). The latest Global Burden Disease (GBD2019) study, a comprehensive research initiative that quantifies health loss due to diseases, injuries, and risk factors worldwide, estimated that exposure to air pollution, including both household and ambient pollution, led to 6.7 million premature deaths (95% confidence interval [CI], 5.9 to 7.5 million) worldwide in 2019 (GBD 2019 Risk Factors Collaborators., 2020). Thus, the urgency of dealing with air pollution has become one of the most pressing global challenges.

It is well-known that surface air pollution is usually unequally distributed in space, with higher levels in developing countries than in developed countries (GBD 2015 Risk Factors Collaborators., 2016). For example, India was ranked as the most polluted country in the world in 2021, with 63 of the world's 100 most polluted cities (IQAir: 2021 World Air Quality Report, available at: [https://lib.icimod.org/record/35767/files/HimalDoc2022\\_2021WorldAirQualityReport.pdf?type=primary](https://lib.icimod.org/record/35767/files/HimalDoc2022_2021WorldAirQualityReport.pdf?type=primary), last access: 21 September 2024). Previous modeling studies indicated that districts exceeding India's annual ambient standard of  $40 \mu\text{g m}^{-3}$  rose from 200 to 385 out of 640 from 1998 to 2020 (Guttikunda and Ka, 2022). The GBD2019 study estimated that premature deaths attributed to ambient  $\text{PM}_{2.5}$  and  $\text{O}_3$  pollution accounted for 10.4% (8.4-12.3) and 1.8% (0.9-2.7) of the total deaths in India in 2019, respectively, and the death rate per 100,000 people increased by 115.3% (28.3-344.4) and 139.2% (96.5-195.8) from 1990 to 2019, respectively (Pandey et al., 2021). However, the GBD2019 study did not separate the air quality changes due to various contribution factors, such as anthropogenic (ANTHRO) and biomass burning (BB). Meanwhile, the elevated chemical reaction rates in India, driven by intense sunlight and warm temperatures, create conditions conducive to ozone formation. Additionally, strong convection enhances the transport of ozone and its precursors, such as  $\text{NO}_y$ , to higher altitudes where the ozone lifetime is prolonged, facilitating accumulation. This phenomenon positions India as a hotspot for ozone pollution, significantly impacting air quality in downwind regions (Zhang et al., 2016, 2021a).

As seen from the Community Emissions Data System (CEDS) inventory (Hoesly et al., 2018), the increasing trends of ANTHRO emissions of major air pollutants, such as nitrogen oxides ( $\text{NO}_x$ ), carbon monoxide ( $\text{CO}$ ), and non-methane volatile organic compound (NMVOC), are significantly higher in India than those in other regions (Wang et al., 2022). Meanwhile, crop yields in India have significantly enhanced since the mid-1960s after the Green Revolution, contributing to increased BB emissions (Huang et al., 2022). The study showed that from 1950–51 to 2017–18, the crop residue burning in India increased from 18 million tonnes to 116 million tonnes in terms of total biomass burned (Venkatramanan et al., 2021). The frequency and intensity of forest fires in India have also increased in recent years due to persistent warmer temperatures and climate extremes (Vadrevu et al., 2019; Jain et al., 2021). These in turn could pose significant threats to ambient air

65 quality and human health because large amounts of compounds are emitted into the atmosphere, namely carbon dioxide  
(CO<sub>2</sub>), NO<sub>x</sub>, particulate matter (PM), and other chemical species (Crutzen and Andreae, 1990; Carvalho et al., 2011; Lan et  
al., 2022; Miranda et al., 2005). Previous studies have utilized observational and satellite data to assess the impacts of  
ANTHRO and BB sources on air quality trends in some Indian cities (Gurjar et al., 2016; Vohra et al., 2022). Additionally,  
70 model simulations have been employed to analyze source contributions to air pollution (Conibear et al., 2018a, b). However,  
there remains a lack of comprehensive assessments regarding the impacts of long-term ANTHRO and BB emissions changes  
on air quality, particularly in non-urban areas.

In this study, we aim to improve our understanding of the spatial-temporal distribution of major air pollutants, mainly  
surface PM<sub>2.5</sub> and O<sub>3</sub>, and the related mortality burden in India from 1995-2014 using a state-of-the-art global chemistry  
transport model. In addition, the individual contributions of changes in ANTHRO and BB emissions were further separated  
75 to better understand the causes of worsening air quality and escalating health risks in India. The selected period encompasses  
a dynamic phase of rapid changes in both ANTHRO and BB activities in India, thereby providing an ideal context for  
investigating their respective contributions to air pollution.

## 2 Methods

### 2.1 CAM-chem model configuration

80 We simulated surface PM<sub>2.5</sub> and O<sub>3</sub> concentrations over India between 1995 and 2014 using the global chemistry model  
CAM-chem, which is based on version 6 of the Community Atmosphere Model (CAM6), the atmospheric component of the  
Community Earth System Model (CESM2), as detailed by Danabasoglu et al. (2020) and Emmons et al. (2020). Following  
Emmons et al. (2020) the original model was run at 1.25° (longitude) × 0.9° (latitude) horizontal resolution with 32 vertical  
levels reaching ~45 km. We configured the Model of Ozone and Related Chemical Tracers Tropospheric and Stratospheric  
85 (MOZART-TS1) chemistry mechanism with various complexity choices for tropospheric and stratospheric chemistry  
(Emmons et al., 2020). The aerosol module adopted the four-mode version of the Modal Aerosol Model (MAM4), including  
sulfate, black carbon, primary organic matter, secondary organic aerosols, sea salt, and mineral dust. The first level of the  
model outputs was considered the surface level, and all the model outputs were then regridded to a finer resolution 0.5° ×  
0.5° to match the grid-cell population and baseline mortality rates datasets in performing the health impact assessment.

90 Global historical ANTHRO emissions were adopted from CEDS (version 2017-05-18), which provides monthly  
emissions of ANTHRO aerosol and precursor compounds at 0.5° × 0.5° from 1750 to 2014 and were used in the Coupled  
Model Intercomparison Project Phase 6 (CMIP6) experiments (Emmons et al., 2020; Hoesly et al., 2018). The ANTHRO  
emissions includes eight sectors: agriculture; energy; industrial; transportation; residential, commercial, other; solvents  
production and application; waste and international shipping (Hoesly et al., 2018). The air pollutants from the CEDS  
95 inventory, especially the NMVOC, were then re-specified to match the chemical species in the latest CESM2 model,

following the steps introduced by Emmons et al. (2020). Interpolation of the emission inventory from its original resolution ( $0.5^\circ \times 0.5^\circ$ ) to the target model resolution ( $0.9^\circ \times 1.25^\circ$ ) before being input into the model. Global historical BB emissions were sourced from van Marle et al. (2017) at monthly temporal resolution and  $0.5^\circ$  native resolution, with all emissions occurring at the surface. Additionally, the biogenic emissions were calculated using the Model of Emissions of Gases and Aerosols from Nature (MEGAN v2.1). More emissions used are described in Emmons et al., (2020).


## 2.2 Numerical experiments designs

The standard simulation (BASE) was driven by year-to-year variability ANTHRO and BB emissions from 1995 to 2014, as described above. To separate the contributions from these two emission sources, we then conducted two sensitivity simulations in which ANTHRO emissions (FixAN) and BB emissions (FixBB) were fixed at 1995 levels individually, while all other parameters were kept consistent with the BASE (Table 1). Subtracting the BASE from each sensitivity enables quantifying the influences of changes in ANTHRO and BB emissions on air quality and its associated health burden in India, respectively. We will discuss the air quality and mortality burden changes in six Indian regions based on meteorological conditions and aerosol variability (Fig. 1).

**Table 1. Model simulations performed in this study**

Simulation	Anthropogenic emissions	Biomass burning emissions
BASE	V	V
FixAN	1995	V
FixBB	V	1995

"V" indicates that particular input is subject to interannual variation in the simulation during the period 1995-2014, "FixAN" indicates that only global ANTHRO emissions were set to 1995 conditions in the simulation. "FixBB", indicates that only global BB emissions were set to 1995 level.



115 **Figure 1. A map of India marked into six regions based on meteorological conditions and aerosol variability (adapted from David et al. 2018).**

### 2.3 Trend estimation

In this study, we applied the Theil-Sen estimator (Theil, 1992; Sen, 1968) to calculate the magnitude of trends in surface  $PM_{2.5}$  and  $O_3$  concentrations and the attributed mortality burden spanning from 1995 to 2014. The Theil-Sen estimator is a robust non-parametric method for trend analysis based on the median slope, which is insensitive to outliers and highly competent in identifying the slope of non-normally distributed data, as described in eq 1. This method has been widely used to analyse temporal trends in air pollutants that are always non-normally distributed (e.g., Munir et al., 2013; Sarkar et al., 2019; Vanem and Walker, 2013; Wan et al., 2023).

120

$$Slope = Median \frac{(x_i - x_j)}{(t_i - t_j)} \quad (1)$$

Where  $x_i$  and  $x_j$  represent the concentrations of either PM<sub>2.5</sub>, O<sub>3</sub>, or attributed premature mortality at time  $t_i$  and  $t_j$  ( $i > j$ ), respectively, for the same parameter.  $Slope > 0$  indicates an increasing trend;  $Slope < 0$  indicates a decreasing trend.

125 In complement to the Theil-Sen estimator, we used the nonparametric Mann-Kendall test to assess the significance of temporal trends within the data series (Zhang et al., 2022a, b). According to previous studies, p-value less than 0.05 is most commonly treated as the absolute threshold of statistical significance (Christiansen et al., 2020; Wang et al., 2021; Zhou et al., 2017). The above methods were completed by implementing a Python program with the package “pymannkendall”, as detailed at <https://pypi.org/project/pymannkendall/>, last accessed on March 20, 2024.

#### 130 2.4 Mortality burdens of surface PM<sub>2.5</sub> and O<sub>3</sub> in India

Based on an integrated exposure-response function utilized in the most recent GBD studies, we estimated the mortality burden associated with long-term exposure to ambient annual PM<sub>2.5</sub> and 6-month running average of daily maximum 8-hr average (6mMDA8) O<sub>3</sub> in India spanning from 1995 to 2014, as described in eq 2.

$$\Delta Mort = y_0 \times AF \times pop = y_0 \times \left( \frac{RR - 1}{RR} \right) \times pop \quad (2)$$

135 Where  $\Delta Mort$  refers to the annual mortality burden attributed to long-term PM<sub>2.5</sub> or O<sub>3</sub> exposure, and  $y_0$  is the baseline mortality rate for a specific cause of disease.  $AF$  is the attributable fraction measuring the PM<sub>2.5</sub> or O<sub>3</sub> exposure attributable disease burden, which is represented by  $\frac{RR-1}{RR}$  ( $RR$  refers to relative risk).  $pop$  represents the exposed population above the age of 25 for each grid cell in the domain.

140 Following our previous work (Zhang et al., 2021b), we obtained the baseline mortality rate ( $y_0$ ) for each country and 5-year age group from 1995-2014 from the GBD2017 project (GBD 2017 Risk Factor Collaborators., 2018). The  $RR$  of long-term PM<sub>2.5</sub> exposure associated with mortality burden due to specific disease was estimated using an integrated exposure-response model (IER) constructed by Burnett et al. (2014) and updated in GBD2017. The  $RR$  for long-term O<sub>3</sub> exposure was obtained from Turner et al. (2016) which indicated an  $RR$  of 1.12 (95% confidence interval (CI): 1.08, 1.16) for respiratory disease. The recent GBD2019 reported a relatively lower  $RR$  for the chronic obstructive pulmonary disease (COPD), a subcategory of respiratory disease (1.06, with 95% CI: 1.03, 1.10). To be comparable with the GBD2019 results, we also  
145 estimated the O<sub>3</sub>-related mortality burden for the COPD in India during the same period. Population distribution with age stratification data ( $pop$ ) was retrieved from the GBD2017 with a horizontal resolution of 0.1°. The population-weighted (pop-weighted) average of specific air pollutants discussed in the results was calculated by weighting the population of all


grid cells inside each administrative region or country. Additionally, we calculated mortality rates per capita (avoid deaths per 100,000 people) in each administrative region to exclude the influence of varying populations.

## 150 **3 Results and discussion**

### **3.1 CAM-chem evaluation**

We performed a comprehensive model evaluation by comparing our simulated monthly concentrations from the BASE with multiple datasets, including ground-based observations in India, historical multi-model simulation from the CMIP6 project, and different versions of multi-year reanalysis data from the Atmospheric Composition Analysis Group (ACAG) at  
155 Washington University in St. Louis, hereinafter referred as 'Wustl Extracts' (van Donkelaar et al., 2021). We also compared our simulated PM<sub>2.5</sub> and O<sub>3</sub> with previously published studies in India using either global or regional chemical transport models (CTMs), as well as the concentration reported from the GBD2019. We selected available ground-level PM<sub>2.5</sub> observations over India from previous studies (Latha and Badarinath, 2005; Panwar et al., 2013; Reddy et al., 2012; Saradhi et al., 2008; Tiwari et al., 2009, 2013), which were also collected by the ACAG. The locations of these sites are listed in  
160 Table S1. Figure S1 indicated that the model exhibits good performance in capturing seasonal variations of surface PM<sub>2.5</sub> observations, especially during the peak months, with correlation coefficients (R) ranging from 0.59 to 0.91. Two exceptions are Mumbai (with R of -0.16), where the model showed a contrasting trend for the seasonal PM<sub>2.5</sub> characteristics (Fig. S1b), and Mukteshwar (with R of 0.45). One possible explanation is the potential underestimation of emission inventories, especially during early periods for developing regions, such as India (McDuffie et al., 2020; Wang et al., 2022; Agarwal et al., 2024). For O<sub>3</sub>, our model showed an even higher R when compared with the available surface observation sites in India  
165 from 1997 to 2011 (Fig. S2). Unlike the underestimations of surface PM<sub>2.5</sub> in India, the CAM-chem model tends to overestimate surface O<sub>3</sub>, which was not very uncommon for global CTM and also frequently discussed in previous studies (Hou et al., 2023; Tilmes et al., 2015; Young et al., 2018; Zhang et al., 2021b). The overestimation was partly caused by the coarse resolution, which leads to diluted emissions of O<sub>3</sub> precursors and then simulated high O<sub>3</sub> production. Figure 2  
170 compared our study with several previous studies and other publicly available PM<sub>2.5</sub> and O<sub>3</sub> datasets, as detailed in Tables S2 and S3. The comparisons indicated our simulated results using the CAM-chem agree very well with previous studies for both PM<sub>2.5</sub> and O<sub>3</sub>, based on either the various metrics, such as annual averages, pop-weighted averages or annual maximum of 6mMDA8 O<sub>3</sub>, consistent with the findings within the multiple CMIP6 models (Turnock et al., 2020). Figure S3 further compares the long-term trend of annual surface PM<sub>2.5</sub> concentrations from 1998 to 2014 in the BASE and Wustl Extracts  
175 dataset. A consistent increasing trend was found in both datasets, with temporal R of 0.86 and lower estimations in our model. The model performs better in eastern India than in western India, with R usually being larger than 0.9 and NMB lower than -25%. Similarly, compared to the simulated trend in our study with different versions of Wustl Extracts and the GBD2019, our simulated PM<sub>2.5</sub> concentration is lower, and the simulated O<sub>3</sub> is higher (Fig. S4). The underestimation of the

180 surface PM<sub>2.5</sub> was partly caused by the missing model representation of nitrate and ammonium (Ren et al., 2023) and the secondary organic aerosol (Liu et al., 2021).



**Figure 2. Comparison of annual PM<sub>2.5</sub> and O<sub>3</sub> concentrations in India with previous studies. Note that the metrics vary depending on the study.**

### 3.2 Spatial and temporal distribution of air pollution changes in India from 1995 to 2014

#### 185 3.2.1 Historical emissions in India from 1995 to 2014


We first assessed the interannual variation of ANTHRO and BB emissions of CO, NO<sub>x</sub>, NMVOC, sulfur dioxide (SO<sub>2</sub>), ammonia (NH<sub>3</sub>), black carbon (BC), and organic carbon (OC) in India between 1995 and 2014 from the CEDS. Figure S5 indicated an overall increase in ANTHRO emissions before slowly falling after 2011. Significant inter-annual variations for BB emissions, such as in 1999, 2006, and 2009, were mainly caused by climate change-induced hot and arid conditions (Sahu et al., 2015). Figure S6 showed that ANTHRO emissions occurred predominately in IGP and central India, significantly increasing across all regions. Unlike other administrative regions, northern and eastern India, such as Punjab and Manipur, features a higher ratio of BB emissions to ANTHRO emissions.

#### 3.2.2 The long-term trends of PM<sub>2.5</sub> and O<sub>3</sub> in India from 1995 to 2014

195 From the BASE simulation, we estimated that the annual mean pop-weighted PM<sub>2.5</sub> and O<sub>3</sub> for India in 1995 and 2014 were 29.88 µg m<sup>-3</sup> and 67.41 ppbv, respectively. Figure 3a, b showed that annual average PM<sub>2.5</sub> concentrations gradually rose from the south to the north, with high levels predominantly found in the IGP, mainly caused by high ANTHRO emissions (Fig. S6) and reduced ventilation due to obstruction by the Tibetan Plateau (Gao et al., 2018). Surface annual average O<sub>3</sub> concentrations gradually increase from west to east and south to north, with the highest levels concentrated in northern India and the eastern part of central India. The spatial patterns of the PM<sub>2.5</sub> and O<sub>3</sub> distribution in India were also



200 seen in several previous studies, though they only discussed one or several specific years (Jia et al., 2021; Pandey et al., 2021).



205 **Figure 3. Spatial distributions of PM<sub>2.5</sub> (top panel) and O<sub>3</sub> (bottom panel) for annual average in 1995 (a, f) and 2014 (b, g), with the trends from 1995 to 2004 (c, h), 2005 to 2014 (d, i), and 1995 to 2014 (e, j). The black dot denotes the areas where the trend is statistically significant ( $p < 0.05$ ). The units are  $\mu\text{g m}^{-3}$  for PM<sub>2.5</sub> (a,b) and ppbv for O<sub>3</sub> in (f, g), and  $\mu\text{g m}^{-3}$  per decade ( $\mu\text{g m}^{-3} \text{ decade}^{-1}$ ) for PM<sub>2.5</sub> trends (c,d,e), and ppbv per decade ( $\text{ppbv decade}^{-1}$ ) for O<sub>3</sub> trends (h,i,j).**


210 From Figure 3, we also found that both PM<sub>2.5</sub> and O<sub>3</sub> showed a statistically significant increasing trend all over the country from 1995 to 2014, with a nation-wide increasing rate of  $6.71 \mu\text{g m}^{-3} \text{ decade}^{-1}$  ( $p < 0.01$ ) for pop-weighted PM<sub>2.5</sub> and  $7.08 \text{ ppbv decade}^{-1}$  ( $p < 0.01$ ) for pop-weighted O<sub>3</sub>, respectively (Fig. S7), which was mainly driven by rapid industrialization and substantial economy development (Pandey et al., 2014; Sadavarte and Venkataraman, 2014). However, distinct spatial heterogeneity for the increasing trend was observed for the two air pollutants. PM<sub>2.5</sub> exhibited varying degrees of increase across India, with the most distinctive increase occurring in the IGP, where the maximum trend reached  $12.60 \mu\text{g m}^{-3} \text{ decade}^{-1}$ . This notable rise can be attributed to the increased regional ANTHRO emissions (Fig. S6). For O<sub>3</sub>, eastern central India experienced the highest O<sub>3</sub> increases, with an obvious increase in the eastern and the lowest increases in western India. One thing needs to be pointed out that in northern IGP, including New Delhi, significant O<sub>3</sub> decreases were also observed, which could be caused by the inhibited O<sub>3</sub> production due to NO titration as a result of dramatic increase in NO<sub>x</sub> emissions, as discussed in Karambelas et al. (2018). Splitting the trend into two periods (from 1995 to 2004 and from 2005 to 2014), we found a larger increasing trend in the latter period than that in the previous one for both PM<sub>2.5</sub> and O<sub>3</sub>,

220 which may be due to the rapid urbanization and growing transportation activities over populous regions (Fig. S8) in recent years in India (Gao et al., 2018).

### 3.3 Driving factor analysis for the air pollution changes in India

#### 3.3.1 Contributions to the annual and seasonal trends

Figure 4 showed the contributions of ANTHRO and BB emissions changes on area-weighted PM<sub>2.5</sub> and O<sub>3</sub> trends from 1995 to 2014. Not surprisingly that ANTHRO emissions changes dominated the PM<sub>2.5</sub> and O<sub>3</sub> deterioration in India. 225 Changes in ANTHRO emissions alone increased area-weighted PM<sub>2.5</sub> by 5.46 μg m<sup>-3</sup> decade<sup>-1</sup> (p < 0.01) and area-weighted O<sub>3</sub> by 2.71 ppbv decade<sup>-1</sup> (p < 0.01), accounting for 102.21% and 104.11% of the total changes, respectively. The contributions of changes in BB emissions were relatively minor, with distinct interannual variations and seasonal variations. Spatially, we found that both the long-term PM<sub>2.5</sub> and O<sub>3</sub> trends were mostly dominated by the ANTHRO emissions changes all over India (Fig. S9a, c). Changes in BB emissions led to a slight increasing trend of PM<sub>2.5</sub> in most of India and a 230 decreasing trend in eastern India, though neither of these trends was statistically significant. BB emissions seemed to increase O<sub>3</sub> in IGP and central India and decrease O<sub>3</sub> in western India, but the trends were insignificant either (Fig. S9b, d).





235 **Figure 4. Drivers for trends of area-weighted (a-c) PM<sub>2.5</sub> and (d-f) O<sub>3</sub> in India in 1995-2014. The yellow shadings in (a, d) showed the evolution of model-simulated PM<sub>2.5</sub> and O<sub>3</sub> concentrations in the FixAN simulation, with the red shadings illustrating the estimation of the PM<sub>2.5</sub> and O<sub>3</sub> concentrations resulting from changes in ANTHRO emissions compared to the 1995 level. (b, e) as**



for (a, d), but for impacts of changes in BB emissions. (c, f) denotes the estimated PM<sub>2.5</sub> and O<sub>3</sub> trends in India derived from the BASE simulation and impacts of ANTHRO and BB emissions, respectively.

It is well recognized that BB emissions usually featured a distinct seasonal trend, especially in India, where they were influenced by the monsoon. Hence, here we quantified the seasonal trend of PM<sub>2.5</sub> and O<sub>3</sub> from ANTHRO and BB emissions for DJF (December-January-February), MAM (March-April-May), JJA (June-July-August, monsoon season), and SON (September-October-November, post-monsoon season) from 1995 to 2014. From Fig. 5a-h, we found that the contributions of ANTHRO emissions had consistent spatial patterns for the seasonal PM<sub>2.5</sub> trend, with larger influences in the post-monsoon seasons (DJF and SON), which was estimated to be responsible for PM<sub>2.5</sub> enhancement by as high as 17.08  $\mu\text{g m}^{-3}$  decade<sup>-1</sup> because of decreased vertical dispersion and diffusion of aerosol caused by lower solar radiation during winter and surface wind speeds (Bran and Srivastava, 2017). The contributions of ANTHRO emissions during the MAM and JJA were modulated as a result of increased precipitation, strong air convergence, and uplift strong air convergence during the presence of the summer monsoon, which impeded the accumulation of PM<sub>2.5</sub> concentrations at ground level (Bran and Srivastava, 2017; Gao et al., 2020; Lu et al., 2018). Unlike PM<sub>2.5</sub>, the contributions of ANTHRO emissions changes on surface O<sub>3</sub> trend in India had a distinct spatial pattern across seasons (Fig. 5i-p). The ANTHRO emissions had a much stronger positive influence on the O<sub>3</sub> increases in northern, eastern central, and eastern India during JJA and SON, while it had the largest increases in southern India in the pre-monsoon season (MAM, Fig. 5j). It was reported that the stronger solar radiation and higher temperature in MAM were attributed to an increase in the photochemical efficiency of O<sub>3</sub> in the presence of NO<sub>x</sub> (Doherty et al., 2013; Jacob and Winner, 2009; Pusede et al., 2015). The decreased O<sub>3</sub> in the IGP was most pronounced in the DJF season (Fig. 5i), which was mainly attributed to lower solar radiation and titration of O<sub>3</sub> by higher NO<sub>x</sub> levels (Kumar et al., 2012). Additionally, the occurrence of winter monsoon led to extensive air subsidence in northern India, resulting in low net O<sub>3</sub> production and strong horizontal export, which ultimately leads to relatively low O<sub>3</sub> levels (Lu et al., 2018).

# PM<sub>2.5</sub>




# O<sub>3</sub>



260 **Figure 5. Seasonal patterns of (a-d) ANTHRO and (e-h) BB emissions contributions for the trends of PM<sub>2.5</sub> in India from 1995 to 2014 and (i-p) for O<sub>3</sub>. The units are  $\mu\text{g m}^{-3}$  per decade for PM<sub>2.5</sub> and ppbv per decade for O<sub>3</sub>. The dots in the plots indicate statistically significant trends, with p-values less than 0.05.**

### 3.3.2 Contributions to the seasonal air quality changes


Figure 6 showed the spatial distributions of BB contributions for seasonal PM<sub>2.5</sub> and O<sub>3</sub> changes between 1995 and 2014, respectively, as detailed in Table S4. The changes in BB emissions from 1995 to 2014 contributed significantly to the  
265 PM<sub>2.5</sub> increases in eastern India (over  $20 \mu\text{g m}^{-3}$ ) with a high incidence of forest fires (Jena et al., 2015). It also resulted in an increase of O<sub>3</sub> by more than 4 ppbv in eastern India in MAM. Contributions to seasonal PM<sub>2.5</sub> and O<sub>3</sub> changes from BB were comparable or even exceeding the those from ANTHRO in some regions, such as Manipur and Nagaland (Fig. S10). With a higher BB fraction in other years, such as 1999, these contributions could even be even higher, reaching up to  $46.03 \mu\text{g m}^{-3}$  and 6.46 ppbv for PM<sub>2.5</sub> and O<sub>3</sub>, respectively (Fig. S11). Therefore, we concluded that the BB emissions in India posed a  
270 great threat to the air quality and thus could not be overlooked.



**Figure 6. Spatial distributions of the BB contribution for seasonal (a-d) PM<sub>2.5</sub> and (e-h) O<sub>3</sub> changes from 1995 to 2014 for DJF, MAM, JJA, and SON. The contributions from BB were calculated as the differences between BASE and FixBB in 2014. The units are  $\mu\text{g m}^{-3}$  and ppbv.**

### 275 3.4 Long-term trends of premature mortality due to PM<sub>2.5</sub> and O<sub>3</sub> in India


We estimated that the national mortality burden attributable to ambient PM<sub>2.5</sub> exposure rose significantly from 698.29 thousand in 1995 to 893.33 thousand in 2014, at a rate of 97.83 thousand per decade ( $p < 0.01$ , Figure 7a). Similarly, the mortality burden attributable to O<sub>3</sub> exposure also notably rose from 414.50 thousand in 1995 to 580.03 thousand in 2014, being 73.91 thousand per decade ( $p < 0.01$ ). We observed that the hotspots of premature mortality attributable to PM<sub>2.5</sub> and O<sub>3</sub> exposure occurred in New Delhi and IGP regions in 1995 and 2014 (Fig. 7b-e), coincidentally with the dense population (Fig. S8). We found that Uttar Pradesh, Bihar, West Bengal, and Haryana, four states within the IGP region, accounted for 41.00% and 39.77% of the national premature mortality due to PM<sub>2.5</sub> and O<sub>3</sub> in 2014, respectively. Considering this heterogeneous spatial distribution, it is imperative for the IGP region to implement stronger air pollution control policies to safeguard human health, as discussed in Jia et al. (2021). Our estimations for the O<sub>3</sub>-related mortality burden were higher than those reported from the GBD2019 (Fig. S12) since we applied a higher RR and used larger baseline mortality rates (see Methods section 2.4). After recalculating the O<sub>3</sub>-related mortality burden using the GBD2019 metrics, we reported an increasing trend of 29.74 thousand deaths per decade<sup>-1</sup> for O<sub>3</sub>-related mortality, comparable to the GBD2019 estimation of 33.24 thousand deaths per decade<sup>-1</sup>. However, our estimated mortality burdens were still slightly higher than the GBD2019 due to the O<sub>3</sub> overestimation in our model (Fig. 2 and Fig. S4).



290

**Figure 7. Spatial-temporal change of mortality burden attributable to PM<sub>2.5</sub> and O<sub>3</sub>. (a) interannual variation from 1995 to 2014. The shaded area indicates the range of 95% confidence interval (gray indicates half of the range). (b-e) spatial distributions of the average annual premature mortality attributable to (b-c) PM<sub>2.5</sub> and (d-e) O<sub>3</sub> in 1995 and 2014.**

To isolate the effects of population heterogeneous among regions, we also quantified the mortality burden changes per capita (avoided deaths per 100,000 people) from 1995 to 2014 (Fig. 8). PM<sub>2.5</sub>-attributable premature mortality per capita was higher in the IGP and eastern India, with the highest in Chandigarh (427.2), followed by Sikkim (153.6), Meghalaya (140.3), and NCT of Delhi (126.1) in 1995 (Fig. S13). The spatial distribution of O<sub>3</sub>-attributable premature mortality per capita resembled that of PM<sub>2.5</sub>, but the values were relatively smaller, with the maximum value also appearing in Chandigarh (288.0), followed by Sikkim (120.2), Meghalaya (68.6), and NCT of Delhi (68.0) in 1995 (Fig. S13). Over the period from 1995 to 2014, PM<sub>2.5</sub>- and O<sub>3</sub>- attributable premature mortality per capita decreased in the north and increased in the south (Fig. 8), indicating that the increasing trend of premature mortality attributable to PM<sub>2.5</sub> and O<sub>3</sub> in the IGP region was mainly driven by the increased population (Fig. S8).




305 **Figure 8. Spatial distributions of premature mortality attributable to PM<sub>2.5</sub> or O<sub>3</sub> per capita (avoid deaths per 100,000 people) in (a, d) 1995, (b, e) 2014, and (c, f) changes from 1995 to 2014 in the state of India.**

Figure 9 showed that changes in ANTHRO emissions from 1995 to 2014 increased premature mortality per capita attributable to PM<sub>2.5</sub>, with the higher values located mainly in eastern IGP and central India. Changes in BB emissions increased premature mortality attributable to PM<sub>2.5</sub> per capita in eastern, western, and southern India and decreased in IGP and central India. The state with the largest increase was Manipur (2.55), followed by Nagaland (2.06), which was associated with the high incidence of wildfires in these regions. The state that experienced the largest decrease was Jharkhand (-1.71), with Bihar (-1.02) followed behind. To explore contribution changes from ANTHRO and BB emissions, we estimated the premature mortality attributable to PM<sub>2.5</sub> per capita in 2000, 2005, and 2010-2014 in Table S5, respectively, consistent with the demonstrations from GBD2017. There was a sharp rise in contributions to premature mortality attributable to PM<sub>2.5</sub> from changes in ANTHRO emissions from 1995 to 2014. Not surprisingly the premature mortality attributable to PM<sub>2.5</sub> from changes in BB emissions fluctuated greatly from 1995 to 2014. In 2000, a year with high BB emissions (Fig. S5), the contributions of changes in BB emissions to the premature mortality attributable to PM<sub>2.5</sub> in the states of Mizoram, Nagaland, Arunachal Pradesh, and Tripura reached 5.14, 4.90, 4.86, and 4.17, respectively, which exceeding the contributions of changes in ANTHRO emissions in that year (Table S5).

310

315





320 **Figure 9. Spatial distributions of contributions to premature mortality attributable to PM<sub>2.5</sub> per capita (avoided deaths per 100,000 people) from changes in (a) ANTHRO and (b) BB emissions from 1995 to 2014.**

#### 4 Conclusions

In this study, we applied a state-of-the-art global CTM (CAM-chem) to provide a detailed assessment of long-term trends of the ambient annual mean PM<sub>2.5</sub> and O<sub>3</sub> in India and their health burden from 1995 to 2014, as well as the driving factor from ANTHRO and BB emissions changes. The annual mean area-weighted PM<sub>2.5</sub> over India increased at 5.34 μg m<sup>-3</sup> decade<sup>-1</sup> (p < 0.01) from 1995 to 2014, dominated by the ANTHRO emissions (5.46 μg m<sup>-3</sup> decade<sup>-1</sup>, p < 0.01). The highest and fastest PM<sub>2.5</sub> growth was in the IGP regions due to the rapid industrialization, urbanization, and transportation growth. For annual mean area-weighted O<sub>3</sub>, the increase was 2.60 ppbv decade<sup>-1</sup> (p < 0.01), dominated by the ANTHRO emissions as well (2.71 ppbv decade<sup>-1</sup>, p < 0.01). We found that O<sub>3</sub> concentrations were highest in northern India, with the fastest growth occurring in northern, central, and eastern India. The contributions from BB emissions for the long-term trends were not significant for either PM<sub>2.5</sub> (0.09 μg m<sup>-3</sup> decade<sup>-1</sup>, p < 0.30) or O<sub>3</sub> (-0.01, p < 0.80), and showed significant seasonal variations due to large inter-annual variability features. However, when we examine the air quality changes in specific years, such as 1999 and 2014, when there were larger BB activities in India, we found that the contributions from BB could be comparable to or even exceed those from ANTHRO during DJF and MAM, reaching over 46.03 μg m<sup>-3</sup> and 6.46 ppbv for PM<sub>2.5</sub> and O<sub>3</sub>, respectively.

Further estimation of mortality burden showed a 27.93% (698.29 to 893.33 thousand) increase in premature mortality attributable to PM<sub>2.5</sub> between 1995 and 2014 (22.94% for 2005-2014), and a 39.93% (414.50 to 580.03 thousand) increase for O<sub>3</sub> (44.54% increasing during 2005-2014). Changes in ANTHRO and BB emissions were responsible for an enhancement of premature mortality attributable to PM<sub>2.5</sub> by 88.78% (97.83 thousand per decade, p < 0.01) and 0.02% (2.38 thousand per decade, p < 0.10). After removing the effect of population growth, our analysis revealed a notably higher

mortality burden per capita attributable to PM<sub>2.5</sub> in the IGP regions. However, it was noteworthy that the mortality burden per capita in these regions exhibited a significant decline over the period of 1995-2014, despite the increasing trend of premature mortality. This suggested that population growth is the primary factor driving the trend of premature mortality.

Our study was subject to several uncertainties and limitations. First of all, the coarser resolution in the global model (0.9° × 1.25°) was frequently found to be unable to realistically represent the complex physical and chemical processes of regional-scale air pollution, especially for O<sub>3</sub> (Yue et al., 2023). Moreover, missing chemical mechanisms in the model, such as the lack of representations of nitrate and ammonium (Ren et al., 2023) and the secondary organic aerosol (Liu et al., 2021), prevented the model from accurately simulating PM<sub>2.5</sub> concentration, especially during heavily polluted regions, such as China and India (Turnock et al., 2020). Another major uncertainty originated from the inaccurate emission inventory, especially for developing regions in early periods, as reported by the global datasets (Paulot et al., 2018; Wang et al., 2022). Zhang et al. (2021b) revealed that model performance with global CEDS inventory tends to predict lower bias for surface PM<sub>2.5</sub> and higher bias for surface O<sub>3</sub> compared with a regional emission inventory (MEIC) in China due to disparities in spatial allocation. Xie et al. (2024) also highlighted a significant underestimation of agricultural fires in the inventory. Moreover, the uncertainty from health functions ranging from the choice of the exposure-response functions (Ostro et al., 2018; Giani et al., 2020) and the uncertainties of the baseline mortality rates both had different impacts on human health (Lelieveld et al., 2015; Pozzer et al., 2023). Meanwhile, in estimating the mortality burden, we applied the RR derived from a global study instead of India which could potentially have a higher value. Thus, our estimations for the air pollution-related mortality burden could be conservative. More epidemiology studies should be carried out in India to retrieve their own RR. Finally, another limitation in our experimental design was that we set global fixed emissions for ANTHRO and BB instead of in India only, ignoring the impact of intercontinental transportation.

### **Code and data availability**

The CESM model code is available at [https://www.cesm.ucar.edu/models/cesm2/release\\_download.html](https://www.cesm.ucar.edu/models/cesm2/release_download.html) (last access: 12 March 2024). Observation data is available at <https://wustl.box.com/s/79pfex658crbq4dykxh51vvfdpksfhj5> (last access: 29 March 2024), which is collected by the Atmospheric Composition Analysis Group (ACAG) at Washington University in St. Louis.

### **Author contributing**

B.L. analyzed the simulation results and wrote the manuscript. Y.Z. and T.T. conceived the idea, designed and conducted the experiment. Y.Z., T.T., H.Z., J.H., J.M., W.W., and L.X. revised the original paper. All authors contributed to the manuscript.

## Competing interests

370 The authors declare that they have no conflict of interest.

## Acknowledgments

This work was supported by the National Natural Science Foundation of China (42375172). We would like to acknowledge the Atmospheric Composition Analysis Group, the Washington University in St. Louis (WUSTL), for providing open access to the satellite-derived PM<sub>2.5</sub> (<https://sites.wustl.edu/acag/datasets/surface-pm2-5/>, last accessed on March 18, 2024). We  
375 also acknowledge the High Performance Computer resources (2023-EL-PT-000184) from the National Key Scientific and Technological Infrastructure project “Earth System Numerical Simulation Facility” (EarthLab). This material is based upon work supported by the National Center for Atmospheric Research, which is a major facility sponsored by the NSF under cooperative agreement no. 1852977. We thank all the scientists, software engineers, and administrators who contributed to the development of CESM2.

## 380 References

- Agarwal, P., Stevenson, D. S., and Heal, M. R.: Evaluation of WRF-Chem-simulated meteorology and aerosols over northern India during the severe pollution episode of 2016, *Atmospheric Chemistry and Physics*, 24, 2239–2266, <https://doi.org/10.5194/acp-24-2239-2024>, 2024.
- Bran, S. H. and Srivastava, R.: Investigation of PM<sub>2.5</sub> mass concentration over India using a regional climate model,  
385 *Environmental Pollution*, 224, 484–493, <https://doi.org/10.1016/j.envpol.2017.02.030>, 2017.
- Burnett, R. T., Pope, C. A., Ezzati, M., Olives, C., Lim, S. S., Mehta, S., Shin, H. H., Singh, G., Hubbell, B., Brauer, M., Anderson, H. R., Smith, K. R., Balme, J. R., Bruce, N. G., Kan, H., Laden, F., Prüss-Ustün, A., Turner, M. C., Gapstur, S. M., Diver, W. R., and Cohen, A.: An Integrated Risk Function for Estimating the Global Burden of Disease Attributable to Ambient Fine Particulate Matter Exposure, *Environmental Health Perspectives*, 122, 397–403,  
390 <https://doi.org/10.1289/ehp.1307049>, 2014.
- Carvalho, A., Monteiro, A., Flannigan, M., Solman, S., Miranda, A. I., and Borrego, C.: Forest fires in a changing climate and their impacts on air quality, *Atmospheric Environment*, 45, 5545–5553, <https://doi.org/10.1016/j.atmosenv.2011.05.010>, 2011.
- Christiansen, A. E., Carlton, A. G., and Porter, W. C.: Changing Nature of Organic Carbon over the United States, *Environ. Sci. Technol.*, 54, 10524–10532, <https://doi.org/10.1021/acs.est.0c02225>, 2020.
- Conibear, L., Butt, E. W., Knote, C., Spracklen, D. V., and Arnold, S. R.: Current and Future Disease Burden From Ambient Ozone Exposure in India, *GeoHealth*, 2, 334–355, <https://doi.org/10.1029/2018GH000168>, 2018a.

- Conibear, L., Butt, E. W., Knote, C., Arnold, S. R., and Spracklen, D. V.: Residential energy use emissions dominate health impacts from exposure to ambient particulate matter in India, *Nat Commun*, 9, 617, <https://doi.org/10.1038/s41467-018-02986-7>, 2018b.
- 400
- Crutzen, P. J. and Andreae, M. O.: Biomass Burning in the Tropics: Impact on Atmospheric Chemistry and Biogeochemical Cycles, *Science*, 250, 1669–1678, <https://doi.org/10.1126/science.250.4988.1669>, 1990.
- Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliessner, B., Phillips, A. S., Sacks, W., 405 Tilmes, S., van Kampenhout, L., Vertenstein, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001916, <https://doi.org/10.1029/2019MS001916>, 2020.
- 410 David, L. M., Ravishankara, A. R., Kodros, J. K., Venkataraman, C., Sadavarte, P., Pierce, J. R., Chaliyakunnel, S., and Millet, D. B.: Aerosol Optical Depth Over India, *Journal of Geophysical Research: Atmospheres*, 123, 3688–3703, <https://doi.org/10.1002/2017JD027719>, 2018.
- Dedoussi, I. C., Eastham, S. D., Monier, E., and Barrett, S. R. H.: Premature mortality related to United States cross-state air pollution, *Nature*, 578, 261–265, <https://doi.org/10.1038/s41586-020-1983-8>, 2020.
- 415 Doherty, R. M., Wild, O., Shindell, D. T., Zeng, G., MacKenzie, I. A., Collins, W. J., Fiore, A. M., Stevenson, D. S., Dentener, F. J., Schultz, M. G., Hess, P., Derwent, R. G., and Keating, T. J.: Impacts of climate change on surface ozone and intercontinental ozone pollution: A multi-model study, *Journal of Geophysical Research: Atmospheres*, 118, 3744–3763, <https://doi.org/10.1002/jgrd.50266>, 2013.
- van Donkelaar, A., Hammer, M. S., Bindle, L., Brauer, M., Brook, J. R., Garay, M. J., Hsu, N. C., Kalashnikova, O. V., 420 Kahn, R. A., Lee, C., Levy, R. C., Lyapustin, A., Sayer, A. M., and Martin, R. V.: Monthly Global Estimates of Fine Particulate Matter and Their Uncertainty, *Environ. Sci. Technol.*, 55, 15287–15300, <https://doi.org/10.1021/acs.est.1c05309>, 2021.
- Emmons, L. K., Schwantes, R. H., Orlando, J. J., Tyndall, G., Kinnison, D., Lamarque, J.-F., Marsh, D., Mills, M. J., Tilmes, S., Bardeen, C., Buchholz, R. R., Conley, A., Gettelman, A., Garcia, R., Simpson, I., Blake, D. R., Meinardi, S., and 425 Pétron, G.: The Chemistry Mechanism in the Community Earth System Model Version 2 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001882, <https://doi.org/10.1029/2019MS001882>, 2020.
- Fuller, R., Landrigan, P. J., Balakrishnan, K., Bathan, G., Bose-O'Reilly, S., Brauer, M., Caravanos, J., Chiles, T., Cohen, A., Corra, L., Cropper, M., Ferraro, G., Hanna, J., Hanrahan, D., Hu, H., Hunter, D., Janata, G., Kupka, R., Lanphear, B., Lichtveld, M., Martin, K., Mustapha, A., Sanchez-Triana, E., Sandilya, K., Schaeffli, L., Shaw, J., Seddon, J., Suk, W., 430 Téllez-Rojo, M. M., and Yan, C.: Pollution and health: a progress update, *The Lancet Planetary Health*, 6, e535–e547, [https://doi.org/10.1016/S2542-5196\(22\)00090-0](https://doi.org/10.1016/S2542-5196(22)00090-0), 2022.

- Gao, M., Beig, G., Song, S., Zhang, H., Hu, J., Ying, Q., Liang, F., Liu, Y., Wang, H., Lu, X., Zhu, T., Carmichael, G. R., Nielsen, C. P., and McElroy, M. B.: The impact of power generation emissions on ambient PM<sub>2.5</sub> pollution and human health in China and India, *Environment International*, 121, 250–259, <https://doi.org/10.1016/j.envint.2018.09.015>, 2018.
- 435 Gao, M., Gao, J., Zhu, B., Kumar, R., Lu, X., Song, S., Zhang, Y., Jia, B., Wang, P., Beig, G., Hu, J., Ying, Q., Zhang, H., Sherman, P., and McElroy, M. B.: Ozone pollution over China and India: seasonality and sources, *Atmospheric Chemistry and Physics*, 20, 4399–4414, <https://doi.org/10.5194/acp-20-4399-2020>, 2020.
- GBD 2015 Risk Factors Collaborators: Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2015: a systematic analysis for the  
440 Global Burden of Disease Study 2015, *The Lancet*, 388, 1659–1724, [https://doi.org/10.1016/S0140-6736\(16\)31679-8](https://doi.org/10.1016/S0140-6736(16)31679-8), 2016.
- GBD 2017 Risk Factor Collaborators: Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017, *The Lancet*, 392, 1923–1994,  
445 [https://doi.org/10.1016/S0140-6736\(18\)32225-6](https://doi.org/10.1016/S0140-6736(18)32225-6), 2018.
- GBD 2019 Risk Factors Collaborators: Global burden of 87 risk factors in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019, *The Lancet*, 396, 1223–1249, [https://doi.org/10.1016/S0140-6736\(20\)30752-2](https://doi.org/10.1016/S0140-6736(20)30752-2), 2020.
- Giani, P., Castruccio, S., Anav, A., Howard, D., Hu, W., and Crippa, P.: Short-term and long-term health impacts of air  
450 pollution reductions from COVID-19 lockdowns in China and Europe: a modelling study, *The Lancet Planetary Health*, 4, e474–e482, [https://doi.org/10.1016/S2542-5196\(20\)30224-2](https://doi.org/10.1016/S2542-5196(20)30224-2), 2020.
- Gurjar, B. R., Ravindra, K., and Nagpure, A. S.: Air pollution trends over Indian megacities and their local-to-global implications, *Atmospheric Environment*, 142, 475–495, <https://doi.org/10.1016/j.atmosenv.2016.06.030>, 2016.
- Guttikunda, S. and Ka, N.: Evolution of India’s PM<sub>2.5</sub> pollution between 1998 and 2020 using global reanalysis fields  
455 coupled with satellite observations and fuel consumption patterns, *Environ. Sci.: Atmos.*, 2, 1502–1515, <https://doi.org/10.1039/D2EA00027J>, 2022.
- Hoek, G., Krishnan, R. M., Beelen, R., Peters, A., Ostro, B., Brunekreef, B., and Kaufman, J. D.: Long-term air pollution exposure and cardio- respiratory mortality: a review, *Environ Health*, 12, 43, <https://doi.org/10.1186/1476-069X-12-43>, 2013.
- 460 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O’Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geoscientific Model Development*, 11, 369–408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.

- 465 Hou, X., Wild, O., Zhu, B., and Lee, J.: Future tropospheric ozone budget and distribution over east Asia under a net-zero scenario, *Atmospheric Chemistry and Physics*, 23, 15395–15411, <https://doi.org/10.5194/acp-23-15395-2023>, 2023.
- Huang, T., Ma, J., Song, S., Ling, Z., Macdonald, R. W., Gao, H., Tao, S., Shen, H., Zhao, Y., Liu, X., Tian, C., Li, Y., Jia, H., Lian, L., and Mao, X.: Health and environmental consequences of crop residue burning correlated with increasing crop yields midst India’s Green Revolution, *npj Clim Atmos Sci*, 5, 1–9, <https://doi.org/10.1038/s41612-022-00306-x>,  
470 2022.
- Hystad, P., Demers, P. A., Johnson, K. C., Carpiano, R. M., and Brauer, M.: Long-term Residential Exposure to Air Pollution and Lung Cancer Risk, *Epidemiology*, 24, 762, <https://doi.org/10.1097/EDE.0b013e3182949ae7>, 2013.
- India State-Level Disease Burden Initiative Air Pollution Collaborators: Health and economic impact of air pollution in the states of India: the Global Burden of Disease Study 2019, *The Lancet Planetary Health*, 5, e25–e38, [https://doi.org/10.1016/S2542-5196\(20\)30298-9](https://doi.org/10.1016/S2542-5196(20)30298-9), 2021.  
475
- Jacob, D. J. and Winner, D. A.: Effect of climate change on air quality, *Atmospheric Environment*, 43, 51–63, <https://doi.org/10.1016/j.atmosenv.2008.09.051>, 2009.
- Jain, M., Saxena, P., Sharma, S., and Sonwani, S.: Investigation of Forest Fire Activity Changes Over the Central India Domain Using Satellite Observations During 2001–2020, *GeoHealth*, 5, e2021GH000528, <https://doi.org/10.1029/2021GH000528>, 2021.  
480
- Jena, C., Ghude, S. D., Pfister, G. G., Chate, D. M., Kumar, R., Beig, G., Surendran, D. E., Fadnavis, S., and Lal, D. M.: Influence of springtime biomass burning in South Asia on regional ozone (O<sub>3</sub>): A model based case study, *Atmospheric Environment*, 100, 37–47, <https://doi.org/10.1016/j.atmosenv.2014.10.027>, 2015.
- Jia, B., Gao, M., Zhang, X., Xiao, X., Zhang, S., and Lam Yung, K. K.: Rapid increase in mortality attributable to PM<sub>2.5</sub> exposure in India over 1998–2015, *Chemosphere*, 269, 128715, <https://doi.org/10.1016/j.chemosphere.2020.128715>,  
485 2021.
- Karambelas, A., Holloway, T., Kinney, P. L., Fiore, A. M., DeFries, R., Kiesewetter, G., and Heyes, C.: Urban versus rural health impacts attributable to PM<sub>2.5</sub> and O<sub>3</sub> in northern India, *Environ. Res. Lett.*, 13, 064010, <https://doi.org/10.1088/1748-9326/aac24d>, 2018.
- 490 Kumar, R., Naja, M., Pfister, G. G., Barth, M. C., Wiedinmyer, C., and Brasseur, G. P.: Simulations over South Asia using the Weather Research and Forecasting model with Chemistry (WRF-Chem): chemistry evaluation and initial results, *Geoscientific Model Development*, 5, 619–648, <https://doi.org/10.5194/gmd-5-619-2012>, 2012.
- Lan, R., Eastham, S. D., Liu, T., Norford, L. K., and Barrett, S. R. H.: Air quality impacts of crop residue burning in India and mitigation alternatives, *Nat Commun*, 13, 6537, <https://doi.org/10.1038/s41467-022-34093-z>, 2022.
- 495 Latha, K. M. and Badarinath, K.: Seasonal variations of PM<sub>10</sub> and PM<sub>2.5</sub> particles loading over tropical urban environment, *International Journal of Environmental Health Research*, 15, 63–68, <https://doi.org/10.1080/09603120400018964>, 2005.
- Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., and Pozzer, A.: The contribution of outdoor air pollution sources to premature mortality on a global scale, *Nature*, 525, 367–371, <https://doi.org/10.1038/nature15371>, 2015.

- Liu, Y., Dong, X., Wang, M., Emmons, L. K., Liu, Y., Liang, Y., Li, X., and Shrivastava, M.: Analysis of secondary organic aerosol simulation bias in the Community Earth System Model (CESM2.1), *Atmospheric Chemistry and Physics*, 21, 8003–8021, <https://doi.org/10.5194/acp-21-8003-2021>, 2021.
- Lu, X., Zhang, L., Liu, X., Gao, M., Zhao, Y., and Shao, J.: Lower tropospheric ozone over India and its linkage to the South Asian monsoon, *Atmospheric Chemistry and Physics*, 18, 3101–3118, <https://doi.org/10.5194/acp-18-3101-2018>, 2018.
- van Marle, M. J. E., Kloster, S., Magi, B. I., Marlon, J. R., Daniau, A.-L., Field, R. D., Arneth, A., Forrest, M., Hantson, S., Kehrwald, N. M., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Yue, C., Kaiser, J. W., and van der Werf, G. R.: Historic global biomass burning emissions for CMIP6 (BB4CMIP) based on merging satellite observations with proxies and fire models (1750–2015), *Geoscientific Model Development*, 10, 3329–3357, <https://doi.org/10.5194/gmd-10-3329-2017>, 2017.
- McDuffie, E. E., Smith, S. J., O'Rourke, P., Tibrewal, K., Venkataraman, C., Marais, E. A., Zheng, B., Crippa, M., Brauer, M., and Martin, R. V.: A global anthropogenic emission inventory of atmospheric pollutants from sector- and fuel-specific sources (1970–2017): an application of the Community Emissions Data System (CEDS), *Earth System Science Data*, 12, 3413–3442, <https://doi.org/10.5194/essd-12-3413-2020>, 2020.
- Miranda, A. I., Ferreira, J., Valente, J., Santos, P., Amorim, J. H., and Borrego, C.: Smoke measurements during Gestosa-2002 experimental field fires, *Int. J. Wildland Fire*, 14, 107, <https://doi.org/10.1071/WF04069>, 2005.
- Munir, S., Habeebullah, T. M., Seroji, A. R., Gabr, S. S., Mohammed, A. M. F., and Morsy, E. A.: Quantifying temporal trends of atmospheric pollutants in Makkah (1997–2012), *Atmospheric Environment*, 77, 647–655, <https://doi.org/10.1016/j.atmosenv.2013.05.075>, 2013.
- Ostro, B., Spadaro, J. V., Gumy, S., Mudu, P., Awe, Y., Forastiere, F., and Peters, A.: Assessing the recent estimates of the global burden of disease for ambient air pollution: Methodological changes and implications for low- and middle-income countries, *Environmental Research*, 166, 713–725, <https://doi.org/10.1016/j.envres.2018.03.001>, 2018.
- Pandey, A., Sadavarte, P., Rao, A. B., and Venkataraman, C.: Trends in multi-pollutant emissions from a technology-linked inventory for India: II. Residential, agricultural and informal industry sectors, *Atmospheric Environment*, 99, 341–352, <https://doi.org/10.1016/j.atmosenv.2014.09.080>, 2014.
- Panwar, T. S., Hooda, R. K., Lihavainen, H., Hyvarinen, A. P., Sharma, V. P., and Viisanen, Y.: Atmospheric aerosols at a regional background Himalayan site—Mukteshwar, India, *Environ Monit Assess*, 185, 4753–4764, <https://doi.org/10.1007/s10661-012-2902-8>, 2013.
- Paulot, F., Paynter, D., Ginoux, P., Naik, V., and Horowitz, L. W.: Changes in the aerosol direct radiative forcing from 2001 to 2015: observational constraints and regional mechanisms, *Atmospheric Chemistry and Physics*, 18, 13265–13281, <https://doi.org/10.5194/acp-18-13265-2018>, 2018.
- Pozzer, A., Anenberg, S. C., Dey, S., Haines, A., Lelieveld, J., and Chowdhury, S.: Mortality Attributable to Ambient Air Pollution: A Review of Global Estimates, *GeoHealth*, 7, e2022GH000711, <https://doi.org/10.1029/2022GH000711>, 2023.

- Pusede, S. E., Steiner, A. L., and Cohen, R. C.: Temperature and Recent Trends in the Chemistry of Continental Surface Ozone, *Chem. Rev.*, 115, 3898–3918, <https://doi.org/10.1021/cr5006815>, 2015.
- 535 Reddy, B. S. K., Kumar, K. R., Balakrishnaiah, G., Gopal, K. R., Reddy, R. R., Reddy, L. S. S., Ahammed, Y. N., Narasimhulu, K., Moorthy, K. K., and Babu, S. S.: Potential Source Regions Contributing to Seasonal Variations of Black Carbon Aerosols over Anantapur in Southeast India, *Aerosol Air Qual. Res.*, 12, 344–358, <https://doi.org/10.4209/aaqr.2011.10.0159>, 2012.
- Ren, F., Lin, J., Xu, C., Adeniran, J. A., Wang, J., Martin, R. V., Van Donkelaar, A., Hammer, M., Horowitz, L., Turnock, S. T., Oshima, N., Zhang, J., Bauer, S., Tsigaridis, K., Seland, Ø., Nabat, P., Neubauer, D., Strand, G., Van Noije, T., Le Sager, P., and Takemura, T.: Evaluation of CMIP6 model simulations of PM<sub>2.5</sub> and its components over China, <https://doi.org/10.5194/egusphere-2023-2370>, 2023.
- 540 Sadavarte, P. and Venkataraman, C.: Trends in multi-pollutant emissions from a technology-linked inventory for India: I. Industry and transport sectors, *Atmospheric Environment*, 99, 353–364, <https://doi.org/10.1016/j.atmosenv.2014.09.081>, 2014.
- 545 Sahu, L. K., Sheel, V., Pandey, K., Yadav, R., Saxena, P., and Gunthe, S.: Regional biomass burning trends in India: Analysis of satellite fire data, *J Earth Syst Sci*, 124, 1377–1387, <https://doi.org/10.1007/s12040-015-0616-3>, 2015.
- Saradhi, I. V., Prathibha, P., Hopke, P. K., Pandit, G. G., and Puranik, V. D.: Source Apportionment of Coarse and Fine Particulate Matter at Navi Mumbai, India, *Aerosol Air Qual. Res.*, 8, 423–436, <https://doi.org/10.4209/aaqr.2008.07.0027>, 2008.
- 550 Sarkar, C., Roy, A., Chatterjee, A., Ghosh, S. K., and Raha, S.: Factors controlling the long-term (2009–2015) trend of PM<sub>2.5</sub> and black carbon aerosols at eastern Himalaya, India, *Science of The Total Environment*, 656, 280–296, <https://doi.org/10.1016/j.scitotenv.2018.11.367>, 2019.
- Sen, P. K.: Estimates of the Regression Coefficient Based on Kendall’s Tau, *Journal of the American Statistical Association*, 63, 1379–1389, <https://doi.org/10.1080/01621459.1968.10480934>, 1968.
- 555 Theil, H.: A Rank-Invariant Method of Linear and Polynomial Regression Analysis, in: *Henri Theil’s Contributions to Economics and Econometrics*, vol. 23, edited by: Raj, B. and Koerts, J., Springer Netherlands, Dordrecht, 345–381, [https://doi.org/10.1007/978-94-011-2546-8\\_20](https://doi.org/10.1007/978-94-011-2546-8_20), 1992.
- Tilmes, S., Lamarque, J.-F., Emmons, L. K., Kinnison, D. E., Ma, P.-L., Liu, X., Ghan, S., Bardeen, C., Arnold, S., Deeter, M., Vitt, F., Ryerson, T., Elkins, J. W., Moore, F., Spackman, J. R., and Val Martin, M.: Description and evaluation of tropospheric chemistry and aerosols in the Community Earth System Model (CESM1.2), *Geoscientific Model Development*, 8, 1395–1426, <https://doi.org/10.5194/gmd-8-1395-2015>, 2015.
- 560 Tiwari, S., Srivastava, A. K., Bisht, D. S., Bano, T., Singh, S., Behura, S., Srivastava, M. K., Chate, D. M., and Padmanabhamurty, B.: Black carbon and chemical characteristics of PM<sub>10</sub> and PM<sub>2.5</sub> at an urban site of North India, *J Atmos Chem*, 62, 193–209, <https://doi.org/10.1007/s10874-010-9148-z>, 2009.
- 565



- Tiwari, S., Srivastava, A. K., Bisht, D. S., Parmita, P., Srivastava, M. K., and Attri, S. D.: Diurnal and seasonal variations of black carbon and PM<sub>2.5</sub> over New Delhi, India: Influence of meteorology, *Atmospheric Research*, 125–126, 50–62, <https://doi.org/10.1016/j.atmosres.2013.01.011>, 2013.
- 570 Turner, M. C., Jerrett, M., Pope, C. A., Krewski, D., Gapstur, S. M., Diver, W. R., Beckerman, B. S., Marshall, J. D., Su, J., Crouse, D. L., and Burnett, R. T.: Long-Term Ozone Exposure and Mortality in a Large Prospective Study, *Am J Respir Crit Care Med*, 193, 1134–1142, <https://doi.org/10.1164/rccm.201508-1633OC>, 2016.
- 575 Turnock, S. T., Allen, R. J., Andrews, M., Bauer, S. E., Deushi, M., Emmons, L., Good, P., Horowitz, L., John, J. G., Michou, M., Nabat, P., Naik, V., Neubauer, D., O'Connor, F. M., Olivie, D., Oshima, N., Schulz, M., Sellar, A., Shim, S., Takemura, T., Tilmes, S., Tsigaridis, K., Wu, T., and Zhang, J.: Historical and future changes in air pollutants from CMIP6 models, *Atmospheric Chemistry and Physics*, 20, 14547–14579, <https://doi.org/10.5194/acp-20-14547-2020>, 2020.
- Vadrevu, K. P., Lasko, K., Giglio, L., Schroeder, W., Biswas, S., and Justice, C.: Trends in Vegetation fires in South and Southeast Asian Countries, *Sci Rep*, 9, 7422, <https://doi.org/10.1038/s41598-019-43940-x>, 2019.
- 580 Vanem, E. and Walker, S.-E.: Identifying trends in the ocean wave climate by time series analyses of significant wave heightdata, *Ocean Engineering*, 61, 148–160, <https://doi.org/10.1016/j.oceaneng.2012.12.042>, 2013.
- Venkatramanan, V., Shah, S., Rai, A. K., and Prasad, R.: Nexus Between Crop Residue Burning, Bioeconomy and Sustainable Development Goals Over North-Western India, *Frontiers in Energy Research*, 8, 2021.
- 585 Villeneuve, P. J., Weichenthal, S. A., Crouse, D., Miller, A. B., To, T., Martin, R. V., van Donkelaar, A., Wall, C., and Burnett, R. T.: Long-term Exposure to Fine Particulate Matter Air Pollution and Mortality Among Canadian Women, *Epidemiology*, 26, 536, <https://doi.org/10.1097/EDE.0000000000000294>, 2015.
- Vohra, K., Marais, E. A., Bloss, W. J., Schwartz, J., Mickley, L. J., Van Damme, M., Clarisse, L., and Coheur, P.-F.: Rapid rise in premature mortality due to anthropogenic air pollution in fast-growing tropical cities from 2005 to 2018, *Science Advances*, 8, eabm4435, <https://doi.org/10.1126/sciadv.abm4435>, 2022.
- 590 Wan, L., Bento, V. A., Qu, Y., Qiu, J., Song, H., Zhang, R., Wu, X., Xu, F., Lu, J., and Wang, Q.: Drought characteristics and dominant factors across China: Insights from high-resolution daily SPEI dataset between 1979 and 2018, *Science of The Total Environment*, 901, 166362, <https://doi.org/10.1016/j.scitotenv.2023.166362>, 2023.
- 595 Wang, H., Lu, X., Jacob, D. J., Cooper, O. R., Chang, K.-L., Li, K., Gao, M., Liu, Y., Sheng, B., Wu, K., Wu, T., Zhang, J., Sauvage, B., Nédélec, P., Blot, R., and Fan, S.: Global tropospheric ozone trends, attributions, and radiative impacts in 1995–2017: an integrated analysis using aircraft (IAGOS) observations, ozonesonde, and multi-decadal chemical model simulations, *Atmospheric Chemistry and Physics*, 22, 13753–13782, <https://doi.org/10.5194/acp-22-13753-2022>, 2022.
- Wang, L., Zhang, X., and Ming, J.: Aerosol Optical Properties Measured Using a PAX in Central Asia from 2016 to 2019 and the Climatic and Environmental Outlooks, *ACS Earth Space Chem.*, 5, 95–105, <https://doi.org/10.1021/acsearthspacechem.0c00306>, 2021.

- Xie, X., Zhang, Y., Liang, R., Chen, W., Zhang, P., Wang, X., Zhou, Y., Cheng, Y., and Liu, J.: Wintertime Heavy Haze Episodes in Northeast China Driven by Agricultural Fire Emissions, *Environ. Sci. Technol. Lett.*, <https://doi.org/10.1021/acs.estlett.3c00940>, 2024.
- Young, P. J., Naik, V., Fiore, A. M., Gaudel, A., Guo, J., Lin, M. Y., Neu, J. L., Parrish, D. D., Rieder, H. E., Schnell, J. L., Tilmes, S., Wild, O., Zhang, L., Ziemke, J., Brandt, J., Delcloo, A., Doherty, R. M., Geels, C., Hegglin, M. I., Hu, L., Im, U., Kumar, R., Luhar, A., Murray, L., Plummer, D., Rodriguez, J., Saiz-Lopez, A., Schultz, M. G., Woodhouse, M. T., and Zeng, G.: Tropospheric Ozone Assessment Report: Assessment of global-scale model performance for global and regional ozone distributions, variability, and trends, *Elementa: Science of the Anthropocene*, 6, 10, <https://doi.org/10.1525/elementa.265>, 2018.
- Yue, M., Dong, X., Wang, M., Emmons, L. K., Liang, Y., Tong, D., Liu, Y., and Liu, Y.: Modeling the Air Pollution and Aerosol-PBL Interactions Over China Using a Variable-Resolution Global Model, *JGR Atmospheres*, 128, e2023JD039130, <https://doi.org/10.1029/2023JD039130>, 2023.
- Zhang, R., Wu, X., Zhou, X., Ren, B., Zeng, J., and Wang, Q.: Investigating the effect of improved drought events extraction method on spatiotemporal characteristics of drought, *Theor Appl Climatol*, 147, 395–408, <https://doi.org/10.1007/s00704-021-03838-z>, 2022a.
- Zhang, R., Qi, J., Leng, S., and Wang, Q.: Long-Term Vegetation Phenology Changes and Responses to Preseason Temperature and Precipitation in Northern China, *Remote Sensing*, 14, 1396, <https://doi.org/10.3390/rs14061396>, 2022b.
- Zhang, Y., Cooper, O. R., Gaudel, A., Thompson, A. M., Nédélec, P., Ogino, S.-Y., and West, J. J.: Tropospheric ozone change from 1980 to 2010 dominated by equatorward redistribution of emissions, *Nature Geosci*, 9, 875–879, <https://doi.org/10.1038/ngeo2827>, 2016.
- Zhang, Y., West, J. J., Emmons, L. K., Flemming, J., Jonson, J. E., Lund, M. T., Sekiya, T., Sudo, K., Gaudel, A., Chang, K.-L., Nédélec, P., and Thouret, V.: Contributions of World Regions to the Global Tropospheric Ozone Burden Change From 1980 to 2010, *Geophysical Research Letters*, 48, e2020GL089184, <https://doi.org/10.1029/2020GL089184>, 2021a.
- Zhang, Y., Shindell, D., Seltzer, K., Shen, L., Lamarque, J.-F., Zhang, Q., Zheng, B., Xing, J., Jiang, Z., and Zhang, L.: Impacts of emission changes in China from 2010 to 2017 on domestic and intercontinental air quality and health effect, *Atmos. Chem. Phys.*, 21, 16051–16065, <https://doi.org/10.5194/acp-21-16051-2021>, 2021b.
- Zhou, H., Zhou, C., Lynam, M. M., Dvonch, J. T., Barres, J. A., Hopke, P. K., Cohen, M., and Holsen, T. M.: Atmospheric Mercury Temporal Trends in the Northeastern United States from 1992 to 2014: Are Measured Concentrations Responding to Decreasing Regional Emissions?, *Environ. Sci. Technol. Lett.*, 4, 91–97, <https://doi.org/10.1021/acs.estlett.6b00452>, 2017.

630