

1 Reconstructed glacier area and volume changes in the European 2 Alps since the Little Ice Age

3 Johannes Reinthaler¹, Frank Paul¹

4 ¹Department of Geography, University of Zurich, Zurich, Switzerland

5 *Correspondence to:* Johannes Reinthaler (johannes.reinthaler@geo.uzh.ch)

6

7 **Abstract.** Glaciers in the European Alps have experienced drastic area and volume loss since the end of the Little
8 Ice Age (LIA) around the year 1850. How large these losses were is only poorly known as published estimates of
9 area loss are mostly based on simple up-scaling and alpine-wide reconstructions of LIA glacier surfaces are lacking.
10 For this study, we compiled all digitally available LIA glacier extents for the Alps and added missing outlines for
11 glaciers $>0.1 \text{ km}^2$ by manual ~~digitising~~digitizing. This was based on geomorphologic interpretation of moraines
12 and trimlines on very high-resolution images in combination with historic topographic maps and modern glacier
13 outlines. Glacier area changes are determined for all glaciers with LIA extents at a regional scale. Glacier surface
14 reconstruction with a Geographic Information System (GIS) was applied to calculate (a) glacier volume changes
15 for the entire region from the LIA until ~~today~~around 2015 and (b) total LIA glacier volume in combination with a
16 reconstructed glacier bed. The glacier area shrunk by ~~2405~~2438 km^2 (-57%) from ~~4211~~4244 km^2 at the LIA
17 maximum to 1806 km^2 in 2015 and volume was reduced from about ~~281~~280 km^3 around 1850 to 100 km^3 (-~~65~~64%)
18 in 2015, roughly in line with previous estimates. ~~In the mean~~On average, glacier surfaces lowered by -43.36 m until
19 2015 (-0.2526 m a^{-1}), which is three-times less than observed over the 2000 to 2015 period (-0.82 m a^{-1}). ~~The~~
20 ~~strongest volume losses occurred at around 1600 m and at least 1832 glaciers melted away completely.~~ Many
21 glaciers have now only remnants of their former coverage left ~~and at least 1938 glaciers melted away completely,~~
22 which led to deglaciation of entire catchments. The new datasets should support a wide range of studies related to
23 the ~~effects~~determination of climate change ~~impacts~~ in the Alps, ~~e.g. future glacier evolution, hydrology, land cover~~
24 ~~change, plant succession and emerging hazards.~~

25 1 Introduction

26 Glaciers in the European Alps are among the most intensely studied worldwide. During recent decades, increasing
27 temperatures caused accelerated glacier retreat and ~~downwasting~~down-wasting, impacting water supplies during

28 dry periods, glacier forefield ecosystems, slope stability and tourism (Brunner et al., 2019; Cannone et al., 2008;
29 Haeberli et al., 2007; Oppikofer et al., 2008). While it is crucial to determine the future evolution of glaciers and
30 its consequences, reconstructing past glacier extents and changes allows us to put possible future developments
31 into perspective. Direct observation of glacier extents (including pictorial evidence) and first measurements of front
32 variations in the Alps date back to pre-industrial times (e.g. Zumbühl and Nussbaumer ~~(, 2018)~~, whereas first
33 topographic maps with glacier extents were published in the 19th century for different Alpine regions (Table S1)
34 in the supplemental material). The large body of literature presenting outlines from historic glacier extents in the
35 Alps and elsewhere (e.g. Grove, 2001) in printed (analogue) form is a highly valuable source of information, but
36 hard to use in today's digital world and need thus to be digitized and geocoded first.

37 As an alternative, very high-resolution satellite or aerial images allow us to identify and delineate terminal and
38 lateral moraines or ~~trim lines~~trimlines (separating regions with a different density of vegetation cover) to
39 reconstruct Little Ice Age (LIA) glacier extents (e.g. Reinthaler and Paul, 2023; Lee et al., 2021). For the Alps,
40 numerous studies have already created LIA inventories for specific regions or countries (e.g. Colucci and Žebre,
41 2016; Fischer et al., 2015; ~~Garent and Deline, 2011; Gardent, 2014;~~ Knoll et al., 2009; Lucchesi et al., 2014; Maisch
42 et al., 2000; Nigrelli ~~et al.~~, 2015; Scotti and Brardinoni, 2018; Zanoner et al., 2017) and ~~the related LIA~~
43 from Switzerland and Austria are freely available from open repositories or the Global Land Ice Measurements
44 from Space (GLIMS) glacier database (Raup et al., 2007) (further details are listed in Table S2). However, for some
45 regions in the Alps (493% of all ~~glaciers~~glacier area according to the Randolph Glacier Inventory, v7.0; RGI
46 v7.0 Consortium, 2023) digitized LIA glacier extents were not available and have been newly digitised in this study
47 (Figure 1, Table S3). Whereas reconstructions of glacier extent and surfaces for the LIA maximum have been
48 compiled and published for many regions around the world, e.g. for Patagonia by Glasser et al. (2011), Greenland's
49 peripheral glaciers by Carrivick et al. (2023), the Himalaya by Lee et al. (2021) and for New Zealand by Carrivick
50 et al. (2020), this information was so far not available for the entire European Alps.

51 In the Alps, glaciers reached between 1250 and 1850/60 several ~~glaciers~~times rather similar maximum extents
52 ~~occurred between 1350 and 1850/60,~~ with the exact timing depending on the glacier (e.g. Zumbühl and Holzhauser,
53 1988; Nussbaumer et al., 2011). ~~From these, only, Nicolussi et al., 2022).~~ Especially for smaller glaciers, the
54 ~~moraines and trimlines from the last LIA maximum (around 1850) are sufficiently complete for digitizing. However,~~
55 ~~in contrast to other regions in the world, extent~~ could have been reached at any of the LIA advance periods (e.g.
56 1350, 1600, 1820, 1850). Extent differences (e.g. between 1850, 1820 or 1600) are the different maximum stages
57 were generally small in the Alps. For example, with older advances sometimes being slightly larger (Le Roy et al.,

58 2024. More specifically, most glaciers in the Italian and western Alps reached their last maximum extent around
59 1820, but re-advanced to almost the same position around 1850 (Solomina et al., 2015). In contrast, Austrian
60 glaciers reached their last maximum in the 1850s to 1860s (Ivy-Ochs et al., 2009). ~~Later~~However, only the moraines
61 and trimlines from the last maximum extent (around 1850) are sufficiently complete and have thus been used for
62 digitizing. In most regions of the Alps, later re-advances took place in the 1890s, 1920s and 1970s to ~~80s~~1980s.
63 Terminal and partly also lateral moraines from these re-advances can still be seen in several glacier ~~forefields~~.
64 forefields (e.g. Paul and Bolch, 2019). The study by Zemp et al. (2008) suggests a glacier area reduction of almost
65 50% between 1850 (4474 km²) and 2000 (2271.6 km²) ~~using~~based on a size-dependent extrapolation scheme to
66 obtain alpine-wide extents for 1850. The study by Hoelzle et al. (2003) used parameterisation schemes (e.g. to
67 derive mass balance from length changes) whereas Colucci and Žebre (2016) used volume-area scaling to derive
68 former glacier volume for the Julian Alps. ~~By~~However, only by reconstructing the former glacier surface directly,
69 distributed glacier thicknesses and elevation changes can be derived. ~~Whereas reconstructions of glacier extent and~~
70 ~~surfaces for the LIA maximum have been compiled and published for many regions around the world, for example~~
71 ~~Patagonia by Glasser et al. (2011), Greenlands peripheral glaciers by Carrivick et al. (2023), Himalaya by Lee et~~
72 ~~al. (2021) and New Zealand by Carrivick et al.~~
73 ~~(2020), this information was so far not available for the entire European Alps.~~
74 This study presents a first complete compilation of LIA maximum glacier extents from around 1850 along with a
75 reconstruction of their surfaces and calculation of their volumes for all glaciers in the Alps larger than 0.1 km².
76 Furthermore, we quantify changes in glacier area, volume and elevation between the LIA and around the year 2000
77 and analyse related spatial variations at the regional scale.

78


79 **2 Datasets and Methods**


80 **2.1 Regional subdivision**

81 **2.1 Study regions**

82 For the regional-scale calculations, we have adopted the ‘International Standardized Mountain Subdivision of the
83 AlpsAlps’ (Marazzi, 2004), which was previously also used by Sommer et al. (2020) to regionally aggregate recent
84 glacier mass changes. The dataset consists of two main subdivisions into the Eastern and Western Alps
85 and 14 subdivisions into smaller regions (see Figure 1). Regions with a very small glacier coverage (<5 km²) were
86 merged with neighbouring regions (Maritime with Cottian Alps; German Prealps with Austrian Prealps). Glacier
87 area and volume changes were also calculated per country and for five major river basins (Rhine, Rhone, Danube,
88 Po, SE Alps).

89






90
 91 **Figure 1: Study region of the European Alps.** In white are the 14 sub-regions, in yellow the existing LIA glacier outlines
 92 (3891 km²) from various sources (see Table S2), in blue the new LIA glacier outlines (329 km²) and in red the glaciers of the RGI
 93 v7.0 (<0.1 km²) without a LIA equivalent (6.8 km²). The orange squares denote the location of sub-regions shown in Figs. 2 and 5.
 94 The red dashed line marks the division between Eastern and Western Alps. Background image: ESRI, (2023b).

95 **2.2 Glacier outlines**


96 We have used glacier outlines representing maximum LIA extents from various sources (see Table S2), a second
 97 dataset from the year 2003 compiled by Paul et al. (2011) and available from RGI v7.0 and a third dataset from
 98 around 2015/16 described by Paul et al. (2020) and available from GLIMS. The 2003 inventory has been derived
 99 from Landsat 5 images and the 2015/16 inventory from Sentinel-2; both datasets were taken as they are and not
 100 modified. Due to differences in interpretation of glacier extents by different analysts for the two datasets, we will
 101 only present glacier changes at a regional scale rather than per glacier.

102 Missing LIA extents were digitized for important individual glaciers and glaciers larger than 0.1 km² in RGI v7.0
 103 based on the geomorphological interpretation of trimlines and frontal as well as lateral moraines as visible on very
 104 high-resolution (up to 0.5 m) images (Figure 2). These images were provided by web map services from ESRI

105 (world imagery, standard and clarity (ESRI, 2023b), Google (<https://earth.google.com/web/>) and Bing
106 (www.bing.com/maps/)) and used in combination with roughly geocoded historical maps (see Table S1 for details)
107 to aid in the interpretation. As a starting point for the LIA outline digitizing, we used reshaped outlines from
108 1967-1971 (for France according to Vivian, 1975) and the RGI v7.0 from 2003 for the other regions.
109 (RGI Consortium, 2023).



110



111
 112 **Figure 2:** For the example region of the **Dauphiné Ortler Alps** we show the new (**blue/white**) and existing (**yellow and black**) LIA
 113 outlines as well as glaciers smaller than 0.1 km² without LIA outlines (red). **The black box shows the location of the inset above it,**
 114 **illustrating multiple recession moraines and the resulting LIA outline.** Background image: Sentinel-2 true colour, acquired on
 115 1224.08.2022, source: Copernicus Sentinel data 2022.

116 The largest regions without published/available LIA outlines were the Italian region of Lombardy and parts of the
 117 Pennine Alps, Rhaetian Alps West and Rhaetian Alps South as well as the Dauphiné Alps (see Table S3 for a list
 118 of regions with previously missing LIA glacier extents). Some research was done in the Rhaetian Alps (e.g. Scotti
 119 et al., 2014; Hagg et al., 2017), but not all outlines were digitally available. For the glaciers in Germany, published
 120 maps (Hirtreiter, 1992) were combined with late 19th-century outlines (available from [https://www.bayerische-](https://www.bayerische-gletscher.de)
 121 [gletscher.de](https://www.bayerische-gletscher.de)) and extended to visible moraines. In total, around 767471 glaciers (in RGI v7.0) did not have a LIA
 122 equivalent of which 389218 now have one (216147 glaciers at LIA). The remaining 253 unconsidered glaciers are
 123 generally small (<0.1 km²) and are not expected to change the area and volume change calculation on a regional
 124 scale substantially (they have a total area of 12.17.7 km²) when neglecting them. The existing and new LIA datasets

125 combined cover 99.46% of the 2003 glacier area in RGI v7.0 (Figure 1). ~~A few~~The glaciers that melted away
126 before 2003 would lower this number by a few decimals.

127 2.3 GIS-based surface reconstruction

128 The reconstruction of glacier surfaces is based on elevation information along the LIA outlines and interpolation
129 of the area in between. The modern elevation was extracted from the 10 m resolution Copernicus Digital Elevation
130 Model (DEM) acquired by TanDEM-X between 2011 and 2015 (ESA, 2019) along points on the outlines with 100
131 m equidistance. The interpolation of the glacier surface is based on the up-scaling approach presented by Reinthaler
132 and Paul (~~in review.~~2024) that uses the pattern of recent elevation change rates for each glacier (Hugonnet et al.,
133 2021) to calculate glacier-specific elevation change gradients from bilinear interpolation through all points. The
134 method calculates a scaling factor by dividing the gradient by the LIA elevation change (from interpolating outline
135 points only). ~~using Natural Neighbor~~. The resulting scaling factor (median per region; Figure S1) is then applied
136 to the gradient to shift the modern DEM to the LIA elevation of the glacier. The surface for the area between the
137 modern and the LIA outline was interpolated using the Topo to Raster ~~(tool based on the ANUDEM method that~~
138 ~~has been~~ optimised for point input data ~~without enforcing drainage to derive hydrologically correct DEMs~~
139 ~~(Hutchinson, 1989)~~. For glaciers where no relationship between elevation change and elevation was found i.e. no
140 elevation change gradient, only the outline points were interpolated. The output result is a 30 m resolution DEM of
141 LIA glacier surfaces for nearly all glaciers in the Alps. From this DEM, topographic properties (e.g. median,
142 minimum elevation, slope) were extracted for each glacier.

143 2.4 Volume reconstruction and change assessment

144 In Table 1, the raster datasets used for the volume reconstruction and change assessment are listed. Calibrated
145 glacier bed datasets were used to calculate the contemporary total glacier volume for Switzerland
146 (Grab et al., 2021) and Austria (Helfricht et al., 2019). For the remaining regions, the glacier bed that was inverted
147 from modelled glacier thickness data by Millan et al. (2022) combined with the Copernicus DEM ~~was~~ used ~~to~~
148 ~~determine glacier volume~~. All glacier bed datasets were clipped to the 2015 glacier extent. Area, elevation and
149 volume changes were calculated since the LIA until around the year 2000 (DEM from 2000, outlines from 2003)
150 and around 2015 (change rates from Hugonnet et al. (2021) between 2000 and 2014, DEM and outlines from
151 2015/16). To simplify the presentation of changes, we refer to the time periods P1 (LIA-2000), P2 (2000-2015/16)
152 and P3 (LIA-2015/16) even though outlines, DEMs and change rates refer to slightly different years. Similarly, we
153 have used the year 1850 as the date of the end of the last LIA maximum ~~LIA~~ extent from which the change rates

154 ~~were calculated~~, even though individual glaciers ~~may have reached their maximum extent~~ started receding from this
 155 ~~position~~ at different times. ~~Glacier~~For glacier changes for time periods between the LIA and 2000 ~~were quantified~~
 156 ~~by Helfricht et al. (2019) for Austria and~~, results for Switzerland were compared to Mannerfelt et al. (2022) ~~for~~
 157 ~~Switzerland but so far not for the entire European Alps. As these~~(2022). Glacier change values from more local
 158 studies ~~used different~~ (e.g. Abermann et al. 2009) were not considered due to differences in the sample and input
 159 datasets ~~(outlines, DEMs) as a base and refer to different time periods, we do not compare our results with results~~
 160 ~~from these studies.~~

161
 162 The void-filled SRTM DEM (3-arc seconds) and the Copernicus DEM were used as the year 2000 and 2015 glacier
 163 surfaces to calculate volume and elevation changes for P1 and P3, respectively (ESA, 2019; NASA Shuttle Radar
 164 Topography Mission (SRTM), 2013). Elevation change rates for P2 were taken from Hugonnet et al. (2021) as it
 165 is widely used and because issues with radar penetration of the SRTM and Copernicus DEMs are much more
 166 prominent over the much shorter time period (e.g. Dehecq et al. 2016). This probably resulted in positive elevation
 167 changes in several accumulation areas (Figure S6).

168
 169 **Table 1: Raster datasets used for the glacier change assessment. P1 refers to the time period LIA to 2000, P2 is from around 2000 to**
 170 **around 2015 and P3 from the LIA to around 2015. The time periods for the glacier bed and thickness datasets refer to the time of**
 171 **DEM acquisition and velocity calculation, respectively.**

| Dataset | Reference | Region | Used for | Date | Time period |
|-----------------------|--|-------------------------------------|--------------------------------|------------|---------------------------|
| LIA surface DEM | This study | Alps | Volume/elevation change (rate) | LIA (1850) | P1 & P3 |
| Copernicus DEM | ESA, 2019 | Alps | Volume/elevation change (rate) | 2011-2015 | P3 |
| SRTM DEM | NASA Shuttle Radar Topography Mission (SRTM), 2013 | Alps | Volume/elevation change (rate) | 2000 | P1 |
| Elevation change rate | Hugonnet et al., 2021 | Alps | Volume/elevation change (rate) | 2000-2014 | P2 |
| Glacier bed | Grab et al., 2021 | Switzerland | Total glacier volume | N/A | 2015 |
| Glacier bed | Helfricht et al., 2019 | Austria | Total glacier volume | N/A | 2016 |
| Glacier thickness | Millan et al. 2022 | Alps except Austria and Switzerland | Total glacier volume | 2017-2018 | 2017-2018 |

172

173 2.5 Uncertainty assessment

174 We applied a simplified approach to quantify all relevant sources of uncertainty on the total LIA volume and
 175 volume changes rather than a glacier or cell-specific uncertainty assessment as used by Martín-Español et al.

176 (2016). The main reasons are our highly variable input datasets and methods as well as the focus on regional rather
177 than glacier-specific changes.

178 ~~The overall~~ Our LIA glacier volume calculation has three independent uncertainty components, glacier area
179 (outlines), surface reconstruction and bed topography/ice thickness. The area uncertainty of the digitized LIA
180 glacier outlines is overall about $\pm 5\%$ (Reinthal and Paul, 2023), but the relative area uncertainty is lower for
181 larger and higher for smaller glaciers (Paul et al., 2013). ~~Due~~ The surface reconstruction uncertainty is due to the
182 lack of reference data, ~~an uncertainty assessment of the reconstructed LIA surfaces is difficult to quantify~~. However,
183 for a case study in the Bernese Alps, the mean difference to a dataset derived by Paul (2010) from digitised historic
184 contour lines with 100 m equidistance could be obtained and the mean vertical error was quantified to 4.6 m
185 (Reinthal and Paul, in review.), ~~which gives an uncertainty of the 2024~~. Considering this would change the total
186 LIA volume ~~of by~~ 6.9%. Uncertainties of the bed topography ~~impact on the~~ directly relate to the uncertainty of the
187 calculated ice thickness and would change the total glacier volume ~~are in the range of 4.1 to by around~~ 5% for the
188 calibrated (Grab et al., 2021; Helfricht et al., 2019) and up to 30% for the un-calibrated datasets (Millan et al.,
189 2022). ~~Considering~~ When considering the proportions of the three datasets, the overall volume uncertainty
190 ~~regarding resulting from~~ the bed topography was quantified to 12.7% (see details in supplement). Combining the
191 ~~uncertainty three uncertainties~~ relating to glacier outlines area, surface reconstruction and bed topography, the total
192 random error of the glacier volume ~~derived here~~ is calculated as $\varepsilon = \sqrt{(5.05^2 + 6.9^2 + 12.7^2)}$ or 15.3%.

193 Excluding the glaciers without a reconstructed extent and the missing glaciers leads to a systematic underestimation
194 of the volume and volume change calculations, i.e. this introduces a bias. For the already existing LIA outline
195 datasets, almost all LIA glacier extents were digitised in the related studies (independent of their size), including
196 those that have since melted away. For the glaciers $>0.1 \text{ km}^2$ in RGI v7.0 that do not have LIA extents (total area
197 of ~~12.17.7~~ km^2), we have extrapolated their LIA area from the mean relative change of the size class smaller than
198 1 km^2 to ~~38.3724.9~~ km^2 with an estimated total volume of ~~0.2617~~ km^3 when using the parameterisation scheme by
199 Haeberli and Hoelzle (1995) and a constant mean ice thickness. For already disappeared glaciers that were not
200 mapped, the quantification of their area and volume is more challenging. According to
201 Parkes and Marzeion (2018), disappeared glaciers globally accounted for 4.4 mm (lower bound) of SLR sea level
202 rise compared to 89.1 mm for all glaciers in RGI v5.0 (4.9%). Using the lower bound, since many glaciers
203 disappeared were mapped in the Alps, this would lead to a total underestimation of the volume of around ~~14.1~~ km^3
204 ~~(5.0%)~~. For the volumes of 2003 and 2015, only the uncertainty regarding the bed topography is considered, but
205 ~~DEM accuracy and glacier mapping uncertainties add to the overall uncertainty~~. ~~13.3~~ km^3 (4.8%).

207 3.1 Glacier area changes


208 The total LIA glacier area of the Alps was ~~reconstructed to 4211±213~~estimated at 4244±214 km² of which 2119
 209 km² remained in 2003 (-50% or -0.~~3233~~3233% a⁻¹) and 1806 km² in 2015 (-57% or -0.35% a⁻¹). This is a loss of 313
 210 km² or 15% (-1.2% a⁻¹) for P2. In the eastern Alps (regions 9-14) the relative area loss for P3 is -64% compared to
 211 ~~-5253~~% for the western Alps. Highest area losses are found in the Cottian and Maritime Alps (Region 2) with -
 212 92.5%, Dolomites ~~and~~, Carnic ~~and Julian~~ Alps (Region 13) with -82% and the Lepontine Alps (Region 8) with -
 213 78%. The least affected regions are the Pennine Alps (Regions 5) ~~with -45%~~ and the Bernese Alps (Region 6) both
 214 with -44% (cf. Table 2 and Figure 3a, the changes per country are listed in Table S4). At least to some extent, the
 215 larger glaciers in Regions 5 and 6 caused the smaller relative area changes, but in absolute terms, they are higher (
 216 Figure 3a). The size dependency is also reflected by the glacier area changes per size class, where small glaciers
 217 have higher relative area losses than large glaciers (Figure S11). Glaciers smaller than 1 km² (in 1850) lost 74% of
 218 their area until 2015 whereas glaciers between 5 and 10 km² lost 46% and the two glaciers larger than 50 km² lost
 219 20% of their area.

220 For P2, the total glacier area ~~shrank~~shrank by 15% (-1.22% a⁻¹), but many of the mostly very small glaciers (287)
 221 had a larger area in 2015 than in 2003. This is caused by differences in interpretation from different analysts, sensor
 222 resolutions (Landsat vs. Sentinel-2) and mapping conditions (snow, clouds and shadow) rather than by growing
 223 glaciers: (cf. Paul et al. 2020). The given 2003 to 2015 area change rate should ~~thus~~ be considered as a lower bound,
 224 as correcting the 2015 outlines to the 2003 interpretation would have led to an even larger area loss.


225 3.2 Glacier elevation changes

226 Glaciers in the entire Alps experienced severe volume ~~loss~~losses since the LIA (Figure 3d). The mean elevation
 227 change for P3 over the entire Alps was -43.~~3~~3~~m~~7 m (regionally between -21.9 m and -51.0 m) without a significant
 228 difference between the eastern and the western Alps: ~~(-45.3 m vs. -42.6 m)~~. The highest ~~changes were~~thinning was
 229 observed in the Eastern (Region 10; ~~-50.2~~51.0 m) and southern Rhaetian Alps (Region 11; ~~-46.7~~47.2 m) and the
 230 Bernese Alps (Region 6; ~~-46~~47.4 m). Generally, elevation changes for P3 were largest at an elevation of around
 231 ~~1600~~1650 m ((-105 m)); dominated by Region 6 (western Alps) and decreasing towards higher elevations (Figure
 232 4). For P2, the maximum has shifted upward to 1750 m. The smaller elevation changes at ~~low~~the lowest elevations
 233 can be explained by the smaller ice thickness during the LIA. ~~The largest elevation changes (-105 m) were found~~
 234 ~~at 1650 m in the western (Figure 4a) and at 2250 m (-65 m) in and thus less ice available for melting. In~~ the eastern
 235 Alps, elevation changes for P3 were largest at 2250 m (-65 m) (Figure 4b), basically reflecting the larger) with a

236 shift down to 2050 m for P2. The east-west difference can be explained by glaciers in the eastern Alps not reaching
 237 furtheras far down as in the western Alps. InThe lowering of the point of highest elevation change for P2 in the
 238 eastern Alps could be related to artefacts, since very little glacier area is present at this elevation. At elevations
 239 between 2150 and 3950 m, elevation changes were very similar in the eastern and western Alps.



240



241


242 **Figure 3: Glacier change measures averaged per sub-region for a) to c) and as a raster product for d)-].** The panels show: a) Relative
 243 **area changes [%] in relation to total LIA area, d) elevation changes for P3, b) volume changes [%] in relation to total LIA volume**
 244 **for P3, c) acceleration of volume change rates for P1 compared to P2 (Hugonnet et al., 2021)-) and d) rasterized (4 km) elevation**
 245 **changes for P3.** All background images: ESRI, (2023a).

246 3.3 Glacier volume changes

247 The total glacier volume of the Alps at their LIA maximum extent is calculated as $281280 \pm 43 \text{ km}^3$ of which
 248 $99.6 \pm 12.6 \text{ km}^3$ remained in 2015 (-65.64%). Considering the uncertainty (15.3%) and a possible underestimation
 249 due to missing glaciers of 54.8%, the LIA volume could be as high as 338.4336 km^3 and as low as 238.237 km^3 .
 250 Thereby, the western Alps lost $105.87 \pm 9 \text{ km}^3$ (-58.5%), whereas the eastern Alps lost $75.61 \pm 6.4 \text{ km}^3$ (-75.30%).
 251 The total volume change was highest in regions 3, 5, and 6 (western Alps) as well as 10 and 12 (eastern Alps), i.e.
 252 the regions with the largest glaciers (Figure 3a). Relative volume change was **most dramatic/highest** in regions 1 (-
 253 $75.578.9\%$), 2 (-96.86%), 4 (-75.10%) and 8 (-81.94%) in the western Alps and regions 12 (-79.97%),

254 13 (-88.987.4%) and 14 (-78.21%) in the eastern Alps, i.e. apart from Region 12 those with the smallest glaciers
 255 (Figure 3b; values per country are listed in Table S4). Overall, volume change was highest in an altitude range
 256 between 2500 m and 3000 m (Figure S2), i.e. the elevation range with the largest area. This compensates for the
 257 lower mean elevation change at this altitude. 3D Oblique perspective views generated from a DEM and a hillshade
 258 visualizations of of it are visualized for the LIA and modern glacier surface can be seen in Figures S13-S16S15-S18.

259




260

261 **Figure 4: Elevation changes for P3 with elevation per sub-region for a) the western Alps (sub-regions 1-8) and b) the eastern Alps.**
 262 **(sub-regions 9-14). The regional means are shown in black and the mean of the entire Alps are in grey. The black dotted line**
 263 **indicates the LIA area (secondary x-axis) for the specific elevation band.**

264 3.4 Increase in glacier area, elevation and volume change rates

265 Area, elevation, and volume change rates were much higher in P2 compared to P1. The glacier area change rate
266 was nearly four times higher for P2 (-1.23% a⁻¹) compared to P1 (-0.~~3233~~% a⁻¹) (Table 2, Figure S3). Thereby, the
267 increase in the western Alps (4.~~9x8x~~) is ~~more than~~ two times larger compared to the eastern Alps (2.4x). In Region
268 12 (Tauern Alps West), the area change rates for P2 almost didn't change, beyond mapping uncertainties. In Region
269 4 (Savoy Prealps) fast melting glaciers led to the largest area change rate increase (~~11.9x12x~~), whereas Region 6
270 (Bernese Alps) experienced the lowest area change rate until 2000 (-0.22% a⁻¹) but is also showing a recent strong
271 increase (6.1x).

272 Overall, elevation change rates were 3.~~32~~ times higher for P2 as derived by Hugonnet et al. (2021) compared to
273 P1. Here, the increase was a bit larger in the western (3.~~5x4x~~) than in the eastern Alps (3.~~0x2.9x~~). Regionally, the
274 increase was largest in Regions 4 (5.6x), 7 (5.7x), ~~4 (5x0x)~~, 8 (4.~~5x3x~~) and 9 (4.~~2x1x~~) (Figure 3c). The change is
275 also dependent on the elevation with the elevation loss rate decreasing towards higher elevations (Figures S4 and
276 S5). Notable is the small increase in Region 13, which could be explained by the presence of mostly small glaciers
277 (partly only remnants left) with short response times that now experience only small changes. When calculating
278 the change rates for P2 with the data from Sommer et al. (2020) (-0.65 m a⁻¹) and the DEM difference between the
279 COP DEM and the SRTM DEM (-0.59 m a⁻¹) (Figures S6 and S7), the regional variability is similar, but the increase
280 in the elevation change rate is lower compared to the dataset from Hugonnet et al. (2021) (~~-0.82 m a⁻¹~~). Further
281 research is necessary to investigate what causes the differences among the available datasets. More detailed views
282 of elevation change patterns before and after the year 2000 are shown in Figures 5 and S10.



283
 284 Figure 5: Examples of elevation change rates between the LIA and 2000 (a and c) and 2000-2014 after Hugonnet et al. (2021) (b and
 285 d) for the Bernese Alps (a and b) and the Ötztal Alps (c and d) using the same colour legend for both periods. All background images:
 286 ESRI (2023a).

287 The absolute volume change rates increased by 3842% in P2 (from Hugonnet et al. 2021) compared to P1.
 288 Interestingly, whereas the western Alps experienced a strong increase in the volume change (4852%), the eastern
 289 Alps experienced only a slight increase (4418%). Nevertheless, some regions have shown a lower volume loss rate
 290 for P2 compared to P1 (Regions 2, 12, 13 and 14). The volume change rates for larger river basins increased by
 291 545%, for the RhôneRhine and 5554% for the RhineRhône. The other basins have about constant volume loss rates,
 292 even slightly decreasing after 2000 (-17%) in the south-eastern Alps (Adige, Piave, Brenta, Tagliamento and Soča).
 293 A table of country and basin-specific area and volume changes can be found in Tables S4 and S5.

294 3.5 Glaciers that melted away


295 Temperature increase has caused at least 1938 glaciers with a LIA area of 309 km² to melt away by 2015. This is
 296 a lower bound estimate because several glaciers that were not mapped in 2003 or 2015 were also not mapped with

297 their LIA extent. Most of the lost glaciers can be found in Regions 5 (Pennine Alps) and 6 (Bernese Alps) with 324
298 and 295 glaciers respectively. The largest area loss of glaciers that have completely melted away by 2015 was
299 found in Regions 3 (Graian Alps) and 9 (Rhaetian Alps West) with 44.06 km² and 54.48 km² respectively (see
300 Figure S14 for the glacier count and area for all regions). These regional differences have uncertainties because
301 different analysts have likely worked along a different rule set for the mapping LIA extents and might thus not have
302 included all disappeared glaciers (this also applies to the newly digitised LIA glaciers). Nevertheless, some
303 formerly glacierized catchments, such as large parts of the Engadin - Val Chamuera (Switzerland), Val Spöl
304 (Switzerland/Italy) and Samnaun Valley (Switzerland/Austria), parts of the Italian Dolomites (e.g. Val Gardena)
305 and the German Alps (e.g. Rein Valley) are now basically ice-free (Fig. S8).

306 **3.6 Change in topographic parameters**

307 The median glacier elevation, which can be used as a proxy for the balanced-budget ELA₀ (Braithwaite and Raper,
308 2009), increased from 2898 m during the LIA to 3040 m in 2015 (+142 m). The western Alps experienced a slightly
309 higher increase (146 m) than the eastern Alps (133 m). The change was largest in the Pennine Alps (Region 5; 158
310 m) and the Lepontine Alps (Region 8; 144 m). The smallest changes were observed in the Cottian and Maritime
311 Alps (Region 2; 23 m) and the North-eastern Alps (Region 14; 37 m).

312



313 -
 314 **Figure 5: Examples of elevation change rates between the LIA and 2000 (a and c) and 2000-2014 after Hugonnet et al. (2021) (b and**
 315 **d) for the Bernese Alps (a and b) and the Ötztal Alps (c and d) using the same colour legend for both periods. All background images:**
 316 **ESRI, (2023a).**

317
 318

319
320
321

Table 2: Glacier area, ~~and~~ volume and elevation changes for each region as well as total areas and volumes. Also listed are long term and recent change rates. P1 stands for the period ~~between~~-LIA andto around 2000, similarly P2 for 2000-~~too~~ 2015 and P3 for LIA-to 2015. Elevation change ~~raterates~~ for P2 are taken from Hugonnet et al. (2021).

| Region ID | Main division | Region name | Area | | Relative area change | | | Relative area change rate | | | Volume | | | Volume change | | | Mean elevation change | | | Elevation change rate | | | Change of median elevation (m) |
|-----------|---------------|----------------------------------|-------------------------------|-------------------------|-------------------------|---------------------------|--------------------------|---------------------------|-------------------------|-----------------------------|------------------------|-----------------------------|--------------------------------|-------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|--------|-----------------------|-------|-------|--------------------------------|
| | | | LIA (km ²) | 2003 (km ²) | 2015 (km ²) | P1 (%) | P3 (%) | P1 (% a ⁻¹) | P2 (% a ⁻¹) | P3 (% a ⁻¹) | LIA (km ³) | 2015 (km ³) | P3 (km ³) | P1 (m a ⁻¹) | P3 (m a ⁻¹) | P1 (m a ⁻¹) | P2 (m a ⁻¹) | P3 (m a ⁻¹) | P1 (m) | P3 (m) | P1/P2 | P2/P3 | |
| 1 | west | Dauphiné Alps | 34191.84 | 90.77 | 64.76 | 42.93 52.69 | 59.2866 74 | -0.2834 | -2.39 | 7.989 29 | 1.96 | 6.027 3 | 0.241 02 | -0.2426 | -0.2425 | -0.81 | 3.4210 | 425 127 | | | | | |
| 2 | west | Coitian & Maritime Alps | 20.6970 | 2.68 | 1.55 | -87.03 | -92.52 | -0.57 | -3.53 | 0.5855 | 0.02 | -0.5653 | 30.6329 57 | -0.4516 -0.4918 | -0.42 | 2.7358 | 23 | | | | | | |
| 3 | west | Graian Alps | 636.446 48.48 | 332.27 | 267.42 | 47.79 48.76 | 57.9858 76 | -0.3132 | -1.63 | 43.3542 78 | 15.76 | -27.602 | -0.25 | -0.1412 -0.1413 | -0.27 | 2.8996 | 105 | | | | | | |
| 4 | west | Savoy Prealps | 16.4541 | 8.25 | 4.4 | 49.84 | 73.2418 | -0.33 | -3.89 | 0.5149 | 0.13 | -0.3936 | 22.7421 92 | -0.1412 -0.1413 | -0.69 | 5.0158 | 76 75 | | | | | | |
| 5 | west | Pennine Alps | 703.036 90.48 | 431 | 387.67 | 38.69 37.58 | 4443.86 | -0.25 | -0.84 | 50.3149 72 | 21.51 | -28.821 | -41.771 | -0.24 | 3.02 | 158 | | | | | | | |
| 6 | west | Bernese Alps | 689.781 7 | 462.42 | 389.37 | 32.96 | 43.6550 | -0.22 | -1.32 | 64.4844 31.68 | 31.68 | -32.876 | -0.2628 -0.2829 | -1.03 | 3.8874 | 128 | | | | | | | |
| 7 | west | Glarus Alps | 107.524 | 53.23 | 41.49 | 50.54 | 61.4439 | -0.33 | -1.84 | 5.9197 | 2.46 | -3.4451 | 30.0732 17 | -0.4517 -0.4819 | -0.86 | 5.7104 | 98 97 | | | | | | |
| 8 | west | Leponthine Alps | 182.725 | 52.4 | 39.51 | 71.32 | 78.3736 | -0.47 | -2.05 | 7.5132 | 1.36 | -6.1559 | -33.8455 -0.4819 | -0.2420 | 4.6230 | 144 | | | | | | | |
| 9 | east | Rhaetian Alps West | 32354.49 | 147.96 | 118.38 | 58.04 | 66.4361 | -0.38 | -1.67 | 18.5309 | 5.52 | 13.011 2.57 | 37.4836 68 | -0.1920 -0.2322 | -0.82 | 4.4912 | 135 | | | | | | |
| 10 | east | Rhaetian Alps East | 470.856 | 207.99 | 185.97 | 55.83 | 60.549 | -0.36 | -0.88 | 31.4412 | 7.87 | 23.532 | 50.1851 03 | -0.2931 -0.331 | -0.82 | 2.7869 | 84 80 | | | | | | |
| 11 | east | Rhaetian Alps South | 284.481 | 122.53 | 100.59 | 56.93 | 64.6460 | -0.37 | -1.49 | 18.0617 86 | 4.9 | 13.161 2.96 | 46.6747 15 | -0.28 -0.2829 | -0.95 | 3.4334 | 44 108 | | | | | | |
| 12 | east | Tauern Alps West | 540.645 41.17 | 195.52 | 194.76 | 63.83 | 63.9864 | -0.42 | -0.03 | 30.7248 | 6.18 | 24.543 | 45.6146 21 | -0.2829 -0.28 | -0.61 | 2.2214 | 138 | | | | | | |
| 13 | east | Dolomites & Carnic & Julian Alps | 23.24 | 4.8 | 4.2 | 79.35 | -81.91 | -0.52 | -1.03 | 0.6154 | 0.07 | -0.5447 | 26.5223 7 | -0.15 -0.1614 | -0.24 | 1.5856 | 69 | | | | | | |
| 14 | east | Northeastern Alps | 23.6260 | 7.32 | 6.17 | 69.68 | 73.8987 | -0.45 | -1.32 | 1.00 | 0.22 | -0.78 | 32.4134 74 | -0.2123 -0.221 | -0.45 | 2.1419 | 37 | | | | | | |
| 15 | west | Western Alps | 22547.12 | 1433.62 | 1196.172 | 43.06 | 52.4653 | -0.2829 | -1.38 | 180.635 | 74.88 | 105.76 | 42.2660 -0.2425 | -0.26 | 3.5142 | 447 146 | | | | | | | |
| 16 | east | Eastern Alps | 1695.45 1697.33 | 686.12 | 610.07 | 59.53 | 64.0206 | -0.39 | -0.92 | 13199.09 25 | 24.76 | 75.567 4.33 | 44.8945 -0.2627 | -0.27 | 2.9688 | 134 133 | | | | | | | |
| 17 | All | Alps | 4244.12 4244.45 | 2119.141 | 1806.242 | 49.97 | 57.2744 | -0.3233 | -1.23 | 280.952 79.63 | 99.64 | 181.34 170.00 | -43.3266 -0.2526 | -0.26 | 3.320 | 143 142 | | | | | | | |

322 ~~3.51.1 Glaciers that melted away~~

323 ~~Temperature increase has caused at least 1832 glaciers with a LIA area of 292 km² to melt away. This is a lower~~
324 ~~bound estimate because several glaciers that were not mapped in 2003 or 2015 were also not mapped with their~~
325 ~~LIA extent. Most of these glaciers can be found in Regions 5 and 6 (Pennine and Bernese Alps) with the largest~~
326 ~~area loss in Regions 3 and 9 (Graian and Rhaetian Alps West). These regional differences have uncertainties~~
327 ~~because different analysts have likely worked along a different rule set for the mapping and might thus not have~~
328 ~~included all disappeared glaciers (this also applies to the newly digitised LIA glaciers). Nevertheless, some~~
329 ~~previously glacierized catchments like the Val Chamuera in the Engadin (Switzerland) are now basically ice free~~
330 ~~(Figure S8).~~

331 ~~3.61.1 Change in topographic parameters~~

332 ~~The median glacier elevation, which can be used as a proxy for the balanced-budget ELA₀ (Braithwaite and Raper,~~
333 ~~2009), increased from 2897 m during the LIA to 3040 m in 2015 (+143 m). The western Alps experienced a slightly~~
334 ~~higher increase (147 m) than the eastern Alps (134 m). The change was largest in the Pennine Alps (Region 5; 158~~
335 ~~m) and the Lepontine Alps (Region 8; 144 m). The smallest changes were observed in the Cottian and Maritime~~
336 ~~Alps (Region 2; 23 m) and the North-eastern Alps (Region 14; 37 m).~~

337 4 Discussion

338 4.1 Influence of methods on glacier volume change and comparison with other studies

339 Our estimate of the LIA glacier area is ~~263.2229.6~~ km² (~~65.1~~%) smaller than the value estimated by Zemp et al.
340 (2008) and thus outside our uncertainty range, even ~~if considering when including~~ already disappeared and not
341 digitised glaciers. It could thus be that the extrapolation method applied by Zemp et al. (2008) gives slightly too
342 large areas for the LIA. This is reasonable when considering that ~~the~~ area change rates ~~they used for extrapolation~~
343 have recently strongly increased. Applying them backwards would result in too large areas with this method.
344 Comparing the reconstructed volumes with the GIS-based method applied here with values calculated with the
345 parameterisation scheme by Haeberli and Hoelzle (1995), a large difference is visible. In total, the parameterisation
346 scheme results in a 25% lower total glacier volume for the LIA (224 km³ vs. ~~281280~~±43 km³ in our study). This is
347 also visible on a regional scale where the parameterisation scheme is lower in all but three regions (9, 13, and 14).
348 Especially Regions 3 and 6, where some of the largest glaciers in the Alps are located, had ~~4341~~% and ~~2526~~%
349 lower volumes with the parameterisations scheme. However, for 2015 the volume differences are only 1.2 km³ (or
350 1.2%) smaller with the parameterisation scheme (99.6±12.6 vs 98.4 km³). Although this ~~would could~~ lead to the

351 conclusion that the GIS-based surface reconstruction overestimates LIA glacier volumes, we speculate that the
352 approach by Haeberli and Hoelzle (1995) rather underestimates LIA volumes. For example, the mean slope of the
353 glaciers might have increased so that mean glacier thickness decreased. It also needs to be considered that the
354 parameterisation scheme has its limitations and works best if glacier extents are in balance with climatic conditions
355 (which is certainly not the case in 2015). When the GIS-based surface reconstruction overestimates glacier
356 volumes, this also applies to the calculated volume change rates and the recent acceleration of volume loss rates
357 found here would be even larger.


358 ~~Looking~~Recently published elevation and volume changes since 1931 by Mannerfelt et al. (2022) showed that in
359 regions 6 (Bernese Alps) and 7 (Glarus Alps) the mean elevation change (rate) was -49.2 m (-0.57 m a⁻¹) and -46.5
360 m (-0.54 m a⁻¹) respectively. In this study, we found a lower mean elevation change (rate) since the LIA with -47.4
361 m (-0.28 m a⁻¹) and -32.2 m (-0.17 m a⁻¹) for both regions respectively. Volume change values indicate that most
362 of the melt occurred after 1931, namely -29.4 km³ and -3.8 km³ (Mannerfelt et al. 2022) versus -32.8 km³ and -3.5
363 km³ (this study). Higher elevation change rate values were generally observed by Mannerfelt et al. (2022) at
364 ~~specific~~lower elevation, especially at some large glacier tongues (e.g. Great Aletsch Glacier, Unteraarglacier and
365 Rhone Glacier). At higher elevations, the estimate in this study gives slightly higher elevation change rate values,
366 which could mean that our reconstructed LIA surfaces still are too high in these regions (Figure S12).

367 When analysing glaciers with long observation periods, the volume changes published by GLAMOS (2022)
368 ~~published volume changes starting from~~for the period 1850-1900 (digitised from historic maps) ~~for~~are of some
369 Swiss glaciers ~~use~~. For most of ~~these~~them, volume changes are in good agreement with our ~~estimated~~estimate (e.g.
370 Great Aletsch Glacier: -6.8 km³ (1880-2017), vs. -6.6 km³ in P3). ~~Some~~However, also outliers exist, for example,
371 the Lower Grindelwald glacier. Here, GLAMOS (2022) estimated the volume change between 1861 and 2012 to
372 be -0.44 km³, whereas our calculations resulted in -1.2 km³ and the parameterisation scheme in -0.57 km³. The
373 Lower Grindelwald glacier is ~~one~~a glacier where the bi-linear elevation change gradient could not be calculated
374 due to the low correlation between elevation and elevation change rate, thus the surface was only reconstructed
375 using the outline points, leading to an overestimation of the LIA surface elevation, especially in the (comparably
376 large) accumulation area. However, as the differences could be positive or negative, we would argue that at the
377 granularity of the regional aggregation shown in Figure 1 and Table 2, the volume changes obtained here are likely
378 ~~very~~accurate (within 5% of the real value), but at the scale of individual glaciers deviations might reach 50% or
379 more, depending on the specific characteristics of a glacier- (see details in Reinthaler and Paul, 2024).


380 The difference in the LIA volumes between the parameterisation scheme and the GIS-based reconstruction
 381 increases with increasing glacier area and decreases with mean slope (Figure S9). Therefore, for large, flat glaciers
 382 like those found in Regions 3 and 6, the difference is greatest. The parameterisation scheme uses only mean slope
 383 (derived from glacier length and elevation range) to determine mean ice thickness and might thus underestimate
 384 volumes for large and bottom-heavy glaciers such as Aletsch, Unteraar or Gorner where a large part of the volume
 385 is stored in the lower, flat part. Also, Lüthi et al. (2010) found that the volume at the end of the LIA was larger
 386 relative to the length of the glaciers, confirming that the parameterisation scheme might underestimate glacier
 387 volume. The parameterisation scheme by Haeberli and Hoelzle (1995) might underestimate glacier volume and
 388 thus provide a minimum estimate of LIA glacier volumes.

389 On the other hand, the parameterisation scheme and GIS-based reconstruction gave very similar results for the
 390 thickness change rate. The mean elevation change rate for P3 using the GIS-based reconstruction is -0.26 m a^{-1}
 391 whereas it is -0.25 m a^{-1} with the parameterisation scheme. Regionally, the difference between the methods can be
 392 much larger, with the rate from the GIS-based method being 4544% higher in Region 14 (-0.3624 m a^{-1} vs -0.214
 393 m a^{-1}) and 2933% lower in Region 6 (-0.22 m a^{-1} vs. 0.2829 m a^{-1}) compared to the parameterisation scheme (Figure
 394 6). Results published by Hoelzle et al. (2003) also using the parameterisation scheme are in line with our results,
 395 giving $-0.11 \text{ m w.e. a}^{-1}$ for small glaciers and $-0.25 \text{ m w.e. a}^{-1}$ for large glaciers between 1850 and 1996 for the
 396 Swiss Alps.

397



398



399
 400 **Figure 6: a) Mean elevation change rate for each region (as calculated from the parameterisation (blue) and the GIS approach**
 401 **(yellow). Dashed lines indicate the Alpine-wide mean rate. b) Region ID and regional changelowering of the median glacier elevation**
 402 **between the LIA and 2015. Background image: ESRI, (2023a).**

403 **4.2 Influence of timing on glacier change rates**

404 The change rates since the LIA also depend on the date of the LIA maximum. Since this is a bit different for each
 405 glacier and only known for some of them, an approximate regional average of 1850 has been used. To assess the
 406 impact of the LIA maximum date on the calculated change rates, a 20-year upper and lower bound was applied.
 407 The area change rates would decrease from -0.35% a⁻¹ for 1850 to -0.31% a⁻¹ when using 1830 and increase
 408 to -0.3940% a⁻¹ when starting in 1870. Similarly, the elevation change rates would decrease from -0.26 m a⁻¹ to -
 409 0.2324 m a⁻¹ and -0.3 m a⁻¹, respectively. Thereby, the impact of the LIA starting date on elevation change rates is
 410 not linear but increases towards a smaller date range (Figure S12).

411 Figure S12). More details on the impact of the date on change rates can be found in Reinthaler and Paul (2023).

412 Finally, since P1 is much longer than P2, the rates have to be interpreted with caution. Between the LIA maximum
 413 and the year 2000 most glaciers in the Alps experienced at least two periods with glacier stagnation or even re-
 414 advances (1920s and 1980s), which results in a lower overall change rate compared to a period with a constant
 415 decrease, i.e. glaciers in the Alps were basically retreating and losing mass continuously since the year 2000.

416 **4.3 Climatic and hydrological implications**

417 The observed change in median elevation of 143142 m would translate to a temperature increase of 0.84 to 1.43
 418 °C, depending on the atmospheric lapse rate applied (Haeberli et al., 2019; Kuhn, 1989; Rolland, 2003; Zemp et
 419 al., 2007). This is lower than the 1.5° and 1.6° temperature increase determined by Begert and Frei (2018) and Auer
 420 et al. (2007) for Switzerland and the Alps, respectively. In the eastern Alps, the median elevation change (and thus

421 temperature increase) was slightly lower (133 m; 0.78-1.33° C) compared to the eastern Alps (147 m; 0.86-1.46°
422 C). Precipitation trends since the 19th century are inconclusive, but the Alpine region has become somewhat drier
423 and sunnier since the 1990s (Auer et al., 2007), both enhancing glacier melt. However, as glaciers are not in balance
424 with the current climate, their ablation regions will continue shrinking and thus shifting the median elevation further
425 up-wards. For the large glaciers with flat tongues, this effect is somewhat compensated by the ongoing surface
426 lowering.

427 The impact of long-term ice loss extends beyond the immediate glacierized landscape, affecting glacier runoff and
428 water availability. The excess melt (imbalance) of the glacier adds to the overall runoff with its usual seasonal
429 variations. Our calculations reveal that the absolute volume loss rate in the eastern Alps has only slightly increased
430 in P2 (+418%), indicating that the peak of the imbalance contribution is near. Indeed, some regions in the eastern
431 Alps (Regions 12-14) experienced a decreasing imbalance contribution, implying that peak water in those regions
432 might have occurred already. Moreover, the rivers in the south-eastern Alps flowing into the Adriatic Sea also
433 experienced a decreasing glacier imbalance contribution and the basins draining into the Po and Danube rivers
434 showed stagnating volume loss rates, indicating that peak water might be reached in the near future. Similarly,
435 Huss and Hock (2018) suggest that European basins may have already reached or be on the brink of reaching peak
436 water. On the other hand, the volume loss rates continued to increase dramatically until at least 2015 in the western
437 Alps (except in Region 2). Nevertheless, according to Huss et al. (2008) the peak run-off in highly glacierized
438 basins in the western Alps will be reached in the coming decades.

439 **5 Conclusion**

440 This study has calculated the massive glacier area and volume loss in the European Alps since the end of the Little
441 Ice Age. After the compilation of existing and manual digitising of missing LIA glacier outlines, we obtained a
442 99% areal coverage-by-area. For these all glaciers, the total area was 57% smaller in 2015 (1806 km²) compared to
443 the LIA maximum (4211±2134244±214 km²). The LIA glacier surface reconstruction with a GIS-based approach
444 resulted in an estimated volume loss of +81180±15.54 km³ or 6564% of the original 281280±43 km³. Despite the
445 strongly reduced glacier area by the year 2003, the post-2000 period (P2) witnessed about three times higher rates
446 of elevation loss than in the mean for the LIA to 2000 period (P1), indicating an increasing impact of climate
447 forcing. At the same time, the run-off contribution by glacier imbalance was decreasing after 2000 in some regions
448 of the eastern Alps, while still increasing in the western Alps.

449

450 Due to the temperature increase, at least ~~1832~~1938 glaciers melted away, with numerous others diminished to small
451 remnants of their previous extent. The median glacier elevation was ~~143~~142 m higher in 2015 than at the end of
452 the LIA and will further increase, as most glaciers have not yet adjusted their geometry to current climatic
453 conditions. The resulting deglaciation of entire mountain catchments with related effects on the Alpine landscape
454 will thus also continue. This has far-reaching implications for water resources, run-off, ecosystems, hydropower
455 production and tourism in the Alpine region and requires timely consideration. The here presented dataset will
456 certainly help in assessing the impacts of climate change on mountain landscapes in further detail.

457 **Competing interests**

458 The contact author has declared that none of the authors has any competing interests.

459 **Acknowledgement**

460 The work of J. R. is supported by PROTECT. This project has received funding from the European Union's Horizon
461 2020 research and innovation programme under grant agreement No 869304, PROTECT contribution number XX.

462 The work of F.P. has been performed in the framework of the ESA project Glaciers_cci+ (4000127593/19/I-NB).

463 We thank the editor and reviewers for their careful reading and constructive comments which helped to improving
464 the clarity of the paper. We also thank Melaine Le Roy, Riccardo Scotti and Renato R. Colucci, for pointing us to
465 unconsidered datasets which have been integrated.

466 **Authors contributions**

467 J. R. led the study and the writing of the paper and performed the glacier surface reconstruction as well as all data
468 analysis. F. P. provided ideas and comments and contributed to the writing of the paper ~~and~~as well as to the
469 digitising of outlines.

470

471 **Data availability statement**

472 LIA surface elevations and new outlines will be made available using an online repository (Zenodo). The new LIA
473 outlines will also be available from the GLIMS glacier database.

475 **References**

- 476 [Abermann, J., Lambrecht, A., Fischer, A., and Kuhn, M.: Quantifying changes and trends in glacier area and](#)
477 [volume in the Austrian Ötztal Alps \(1969-1997-2006\), *Cryosphere*, 3, 205–215, \[https://doi.org/10.5194/tc-3-\]\(https://doi.org/10.5194/tc-3-205-2009\)](#)
478 [205-2009, 2009.](#)
- 479 Auer, I., Böhm, R., Jurkovic, A., Lipa, W., Orlik, A., Potzmann, R., Schöner, W., Ungersböck, M., Matulla, C.,
480 Briffa, K., Jones, P., Efthymiadis, D., Brunetti, M., Nanni, T., Maugeri, M., Mercalli, L., Mestre, O., Moisselin,
481 J. M., Begert, M., Müller-Westermeier, G., Kveton, V., Bochnicek, O., Stastny, P., Lapin, M., Szalai, S.,
482 Szentimrey, T., Cegnar, T., Dolinar, M., Gajic-Capka, M., Zaninovic, K., Majstorovic, Z., and Nieplova, E.:
483 HISTALP - Historical instrumental climatological surface time series of the Greater Alpine Region,
484 *International Journal of Climatology*, 27, 17–46, <https://doi.org/10.1002/joc.1377>, 2007.
- 485 Begert, M. and Frei, C.: Long-term area-mean temperature series for Switzerland – Combining homogenized
486 station data and high resolution grid data, *International Journal of Climatology*, 38, 2792–2807,
487 <https://doi.org/https://doi.org/10.1002/joc.5460>, 2018.
- 488 Braithwaite, R. J. and Raper, S. C. B.: Estimating equilibrium-line altitude (ELA) from glacier inventory data, *Ann.*
489 *Glaciol*, 50, 127–132, <https://doi.org/10.3189/172756410790595930>, 2009.
- 490 Brunner, M. I., Farinotti, D., Zekollari, H., Huss, M., and Zappa, M.: Future shifts in extreme flow regimes in
491 Alpine regions, *Hydrol. Earth Syst. Sci*, 23, 4471–4489, <https://doi.org/10.5194/hess-23-4471-2019>, 2019.
- 492 Cannone, N., Diolaiuti, G., Guglielmin, M., and Smiraglia, C.: Accelerating climate change impacts on alpine
493 glacier forefield ecosystems in the European Alps, *Ecological Applications*, 18, 637–648,
494 <https://doi.org/10.1890/07-1188.1>, 2008.
- 495 Carrivick, J. L., James, W. H. M., Grimes, M., Sutherland, J. L., and Lorrey, A. M.: Ice thickness and volume
496 changes across the Southern Alps, New Zealand, from the Little Ice Age to present, *Sci. Rep.*, 10, 1–10,
497 <https://doi.org/10.1038/s41598-020-70276-8>, 2020.
- 498 Carrivick, J. L., Boston, C. M., Sutherland, J. L., Pearce, D., Armstrong, H., Bjørk, A., Kjeldsen, K. K., Abermann,
499 J., Oien, R. P., Grimes, M., James, W. H. M., and Smith, M. W.: Mass Loss of Glaciers and Ice Caps Across
500 Greenland Since the Little Ice Age, *Geophys. Res. Lett.*, 50, <https://doi.org/10.1029/2023GL103950>, 2023.
- 501 Colucci, R. R. and Žebre, M.: Late Holocene evolution of glaciers in the southeastern Alps, *J. Maps*, 12, 289–299,
502 <https://doi.org/10.1080/17445647.2016.1203216>, 2016.

503 Dehecq, A., Millan, R., Berthier, E., Gourmelen, N., Trouvé, E., and Vionnet, V.: Elevation Changes Inferred from
504 TanDEM-X Data over the Mont-Blanc Area: Impact of the X-Band Interferometric Bias, *IEEE J Sel Top Appl*
505 *Earth Obs Remote Sens*, 9, 3870–3882, <https://doi.org/10.1109/JSTARS.2016.2581482>, 2016.

506 ESA: Copernicus DEM EEA-10, <https://doi.org/10.5270/ESA-c5d3d65>, 2019.

507 ESRI: “Topographic” [basemap]. scale not given. “World Topographic Map”,
508 <https://www.arcgis.com/home/item.html?id=30e5fe3149c34df1ba922e6f5bbf808f>, 2023a.

509 ESRI: “World Imagery” [basemap]. scale not given. “world imagery vivid”.
510 <https://www.arcgis.com/home/item.html?id=10df2279f9684e4a9f6a7f08febac2a9>, 2023b.

511 Fischer, M., Huss, M., and Hoelzle, M.: Surface elevation and mass changes of all Swiss glaciers 1980-2010,
512 *Cryosphere*, 9, 525–540, <https://doi.org/10.5194/tc-9-525-2015>, 2015.

513 Gardent, M.: Inventaire et retrait des glaciers dans les Alpes françaises depuis la fin du Petit Age Glaciaire,
514 Université de Savoie, 444 pp., 2014.

515 Gardent, M. and Deline, P.: Stages of glacial retreat in the French Alps since the termination of the Little Ice Age,
516 8th EGU General Assembly, halsde-00879488, 2011.

517 GLAMOS: Swiss Glacier Volume Change, release 2022, <https://doi.org/10.18750/volumechange.2022.r2022>,
518 2022.

519 Glasser, N. F., Harrison, S., Jansson, K. N., Anderson, K., and Cowley, A.: Global sea-level contribution from the
520 Patagonian Icefields since the Little Ice Age maximum, *Nat. Geosci.*, 4, 303–307,
521 <https://doi.org/10.1038/ngeo1122>, 2011.

522 Grab, M., Mattea, E., Bauder, A., Huss, M., Rabenstein, L., Hodel, E., Linsbauer, A., Langhammer, L., Schmid,
523 L., Church, G., Hellmann, S., Deleze, K., Schaer, P., Lathion, P., Farinotti, D., and Maurer, H.: Ice thickness
524 distribution of all Swiss glaciers based on extended ground-penetrating radar data and glaciological modeling,
525 *Journal of Glaciology*, 67, 1074–1092, <https://doi.org/10.1017/jog.2021.55>, 2021.

526 Grove, A. T.: The “Little Ice Age” and its geomorphological consequences in Mediterranean Europe, *Climate*
527 *Change*, 48, 121–136, <https://doi.org/https://doi.org/10.1023/A:1005610804390>, 2001.

528 Haeberli, W. and Hoelzle, M.: Application of inventory data for estimating characteristics of and regional climate-
529 change effects on mountain glaciers: a pilot study with the European Alps, *Ann. Glaciol.*, 21, 206–212,
530 <https://doi.org/10.3189/s0260305500015834>, 1995.

- 531 Haeberli, W., Hoelzle, M., Paul, F., and Zemp, M.: Integrated monitoring of mountain glacier as key indicators of
532 global climate change: The European Alps, *Journal of Glaciology*, 46, 150–160,
533 <https://doi.org/10.3189/172756407782871512>, 2007.
- 534 Haeberli, W., Oerlemans, J., and Zemp, M.: The Future of Alpine Glaciers and Beyond, 1–38 pp.,
535 <https://doi.org/10.1093/acrefore/9780190228620.013.769>, 2019.
- 536 [Hagg, W., Scotti, R., Villa, F., Mayer, E., Heilig, A., Mayer, C., Tamm, W., and Hock, T.: Evolution of two cirque](#)
537 [glaciers in lombardy and their relation to climatic factors \(1962-2016\), *Geogr. Ann. Ser. A Phys. Geogr.*, 99,](#)
538 [371–386, <https://doi.org/10.1080/04353676.2017.1368834>, 2017.](#)
- 539 Helfricht, K., Huss, M., Fischer, A., and Otto, J. C.: Calibrated ice thickness estimate for all glaciers in Austria,
540 *Front Earth Sci (Lausanne)*, 7, 1–15, <https://doi.org/10.3389/feart.2019.00068>, 2019.
- 541 Hirtreiter, Gerhard.: Spät- und postglaziale Gletscherschwankungen im Wettersteingebirge und seiner Umgebung,
542 154, 1992.
- 543 Hoelzle, M., Haeberli, W., Dischl, M., and Peschke, W.: Secular glacier mass balances derived from cumulative
544 glacier length changes, *Glob. Planet. Change*, 36, 295–306, [https://doi.org/10.1016/S0921-8181\(02\)00223-0](https://doi.org/10.1016/S0921-8181(02)00223-0),
545 2003.
- 546 Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussailant,
547 I., Brun, F., and Käab, A.: Accelerated global glacier mass loss in the early twenty-first century, *Nature*, 592,
548 726–731, <https://doi.org/10.1038/s41586-021-03436-z>, 2021.
- 549 Huss, M. and Hock, R.: Global-scale hydrological response to future glacier mass loss, *Nat. Clim. Chang.*, 8, 135–
550 140, <https://doi.org/10.1038/s41558-017-0049-x>, 2018.
- 551 Huss, M., Farinotti, D., Bauder, A., and Funk, M.: Modelling runoff from highly glacierized alpine drainage basins
552 in a changing climate, *Hydrological Processes*, 22, 3888–3902, <https://doi.org/10.1002/hyp.7055> Modelling,
553 2008.
- 554 [Hutchinson, M. F.: A new procedure for gridding elevation and stream line data with automatic removal of spurious](#)
555 [pits, *J. Hydrol.*, 106, 211–232, \[https://doi.org/10.1016/0022-1694\\(89\\)90073-5\]\(https://doi.org/10.1016/0022-1694\(89\)90073-5\), 1989.](#)
- 556 Ivy-Ochs, S., Kerschner, H., Maisch, M., Christl, M., Kubik, P. W., and Schlüchter, C.: Latest Pleistocene and
557 Holocene glacier variations in the European Alps, *Quat. Sci. Rev.*, 28, 2137–2149,
558 <https://doi.org/10.1016/j.quascirev.2009.03.009>, 2009.
- 559 Kienholz, C., Rich, J. L., Arendt, A. A., and Hock, R.: A new method for deriving glacier centerlines applied to
560 glaciers in Alaska and northwest Canada, *Cryosphere*, 8, 503–519, <https://doi.org/10.5194/tc-8-503-2014>, 2014.

- 561 Knoll, C., Kerschner, H., Heller, A., and Rastner, P.: A GIS-based reconstruction of little ice age glacier maximum
562 extents for South Tyrol, Italy, *Transactions in GIS*, 13, 449–463, [https://doi.org/10.1111/j.1467-](https://doi.org/10.1111/j.1467-9671.2009.01173.x)
563 [9671.2009.01173.x](https://doi.org/10.1111/j.1467-9671.2009.01173.x), 2009.
- 564 Kuhn, M.: The Response of the Equilibrium Line Altitude to Climate Fluctuations: Theory and Observations, in:
565 *Glacier fluctuations and climatic change*, edited by: Oerlemans, J., Kluwer, Dodrecht, 407–417,
566 https://doi.org/10.1007/978-94-015-7823-3_26, 1989.
- 567 Lee, E., Carrivick, J. L., Quincey, D. J., Cook, S. J., James, W. H. M., and Brown, L. E.: Accelerated mass loss of
568 Himalayan glaciers since the Little Ice Age, *Sci. Rep.*, 11, 1–8, <https://doi.org/10.1038/s41598-021-03805-8>,
569 2021.
- 570 [Le Roy, M., Ivy-Ochs, S., Nicolussi, K., Monegato, G., Reitner, J. M., Colucci, R. R., Ribolini, A., Spagnolo, M.,
571 and Stoffel, M.: Holocene glacier variations in the Alps, 367–418 pp., \[https://doi.org/10.1016/b978-0-323-\]\(https://doi.org/10.1016/b978-0-323-99712-6.00018-0\)
572 \[99712-6.00018-0\]\(https://doi.org/10.1016/b978-0-323-99712-6.00018-0\), 2024.](https://doi.org/10.1016/b978-0-323-99712-6.00018-0)
- 573 Lucchesi, S., Fioraso, G., Bertotto, S., and Chiarle, M.: Little Ice Age and contemporary glacier extent in the
574 Western and South-Western Piedmont Alps (North-Western Italy), *J. Maps*, 10, 409–423,
575 <https://doi.org/10.1080/17445647.2014.880226>, 2014.
- 576 Lüthi, M. P., Bauder, A., and Funk, M.: Volume change reconstruction of Swiss glaciers from length change data,
577 *J. Geophys. Res.*, 115, 1–8, <https://doi.org/10.1029/2010JF001695>, 2010.
- 578 Maisch, M., Wipf, A., Denzler, B., Battaglia, J., and Benz, C.: Die Gletscher der Schweizer Alpen:
579 Gletscherhochstand 1850, aktuelle Vergletscherung, Gletscherschwundsenarien, in: *Schlussbericht NFP 31*,
580 second edition, Hochschulverlag ETH Zurich, Zurich, 373p, 2000.
- 581 Mannerfelt, E. S., Dehecq, A., Hugonnet, R., Hodel, E., Huss, M., Bauder, A., and Farinotti, D.: Halving of Swiss
582 glacier volume since 1931 observed from terrestrial image photogrammetry, *Cryosphere*, 16, 3249–3268,
583 <https://doi.org/10.5194/tc-16-3249-2022>, 2022.
- 584 Marazzi, S.: Die Orographischen Einteilungen der Alpen und die ‘IVOEA,’ in: *Die Gebirgsgruppen der Alpen.*
585 *Ansichten, Systematiken und Methoden zur Einteilung der Alpen*, Grimm, P, 69–96, 2004.
- 586 Martín-Español, A., Lapazaran, J. J., Otero, J., and Navarro, F. J.: On the errors involved in ice-thickness estimates
587 III: Error in volume, *Journal of Glaciology*, 62, 1030–1036, <https://doi.org/10.1017/jog.2016.95>, 2016.
- 588 Millan, R., Mouginot, J., Rabatel, A., and Morlighem, M.: Ice velocity and thickness of the world’s glaciers, *Nat.*
589 *Geosci.*, 15, 124–129, <https://doi.org/10.1038/s41561-021-00885-z>, 2022.

590 NASA Shuttle Radar Topography Mission (SRTM): Shuttle Radar Topography Mission (SRTM) Global, Open
591 Topography, <https://doi.org/https://doi.org/10.5069/G9445JDF>, 2013.

592 [Nicolussi, K., Roy, M. Le, Schlüchter, C., Stoffel, M., and Wacker, L.: The glacier advance at the onset of the Little
593 Ice Age in the Alps: New evidence from Mont Miné and Morteratsch glaciers, *Holocene*, 32, 624–638,
594 <https://doi.org/10.1177/09596836221088247>, 2022.](https://doi.org/10.1177/09596836221088247)

595 Nigrelli, G., Lucchesi, S., Bertotto, S., Fioraso, G., and Chiarle, M.: Climate variability and Alpine glaciers
596 evolution in Northwestern Italy from the Little Ice Age to the 2010s, *Theor. Appl. Climatol.*, 122, 595–608,
597 <https://doi.org/10.1007/s00704-014-1313-x>, 2015.

598 Nussbaumer, S. U., Steinhilber, F., Trachsel, M., Breitenmoser, P., Beer, J., Blass, A., Grosjean, M., Hafner, A.,
599 Holzhauser, H., Wanner, H., and Zumbühl, H. J.: Alpine climate during the Holocene: A comparison between
600 records of glaciers, lake sediments and solar activity, *J. Quat. Sci.*, 26, 703–713,
601 <https://doi.org/10.1002/jqs.1495>, 2011.

602 Oppikofer, T., Jaboyedoff, M., and Keusen, H. R.: Collapse at the eastern Eiger flank in the Swiss Alps, *Nat.
603 Geosci.*, 1, 531–535, <https://doi.org/10.1038/ngeo258>, 2008.

604 Parkes, D. and Marzeion, B.: Twentieth-century contribution to sea-level rise from uncharted glaciers, *Nature*, 563,
605 551–554, <https://doi.org/10.1038/s41586-018-0687-9>, 2018.

606 [Paul, F. and Bolch, T.: Glacier changes since the Little Ice Age. In: T. Heckmann and D. Morche \(eds.\),
607 *Geomorphology of Proglacial Systems, Geography of the Physical Environment, Springer Nature*, pp. 23-42:
608 \[https://doi.org/10.1007/978-3-319-94184-4_2\]\(https://doi.org/10.1007/978-3-319-94184-4_2\), 2019.](https://doi.org/10.1007/978-3-319-94184-4_2)

609 [Paul, F., Frey, H., and Bris, R. Le: A new glacier inventory for the European Alps from Landsat TM scenes of
610 2003: Challenges and results, *Ann. Glaciol.*, 52, 144–152, <https://doi.org/10.3189/172756411799096295>, 2011.](https://doi.org/10.3189/172756411799096295)

611 Paul, F., Barrand, N. E., Baumann, S., Berthier, E., Bolch, T., Casey, K., Frey, H., Joshi, S. P., Kononov, V., Le
612 Bris, R., Mölg, N., Nosenko, G., Nuth, C., Pope, A., Racoviteanu, A., Rastner, P., Raup, B., Scharer, K.,
613 Steffen, S., and Winsvold, S.: On the accuracy of glacier outlines derived from remote-sensing data, *Ann.
614 Glaciol.*, 54, 171–182, <https://doi.org/10.3189/2013AoG63A296>, 2013.

615 Paul, F., Rastner, P., Azzoni, R. S., Diolaiuti, G., Fugazza, D., Bris, R. Le, Nemec, J., Rabatel, A., Ramusovic, M.,
616 Schwaizer, G., and Smiraglia, C.: Glacier shrinkage in the Alps continues unabated as revealed by a new glacier
617 inventory from Sentinel-2, *Earth Syst. Sci. Data*, 12, 1805–1821, <https://doi.org/10.5194/essd-12-1805-2020>,
618 2020.

- 619 [Raup, B. H., Racoviteanu, A., Khalsa, S. J. S., Helm, C., Armstrong, R., and Arnaud, Y.: The GLIMS geospatial](#)
620 [glacier database: A new tool for studying glacier change, *Glob. Planet. Change*, 56, 101–110,](#)
621 <https://doi.org/10.1016/j.gloplacha.2006.07.018>, 2007.
- 622 Reinthaler, J. and Paul, F.: Using a Web Map Service to map Little Ice Age glacier extents at regional scales, *Ann.*
623 *Glaciol.*, 1–19, <https://doi.org/https://doi.org/10.1017/aog.2023.39>, 2023.
- 624 Reinthaler, J. and Paul, F.: [MethodsAssessment of methods](#) for reconstructing Little Ice Age glacier surfaces ~~in~~
625 [the examples of](#) Novaya Zemlya and the Swiss Alps: ~~Uncertainties and volume changes~~, *Geomorphology*, 461,
626 <https://doi.org/https://doi.org/10.2139/ssrn.4593636>, ~~in review~~1016/j.geomorph.2024.109321 Received, 2024.
- 627 [RGI Consortium: Randolph Glacier Inventory - A Dataset of Global Glacier Outlines, Version 7.0,](#)
628 <https://doi.org/10.5067/f6jmovy5navz>, 2023.
- 629 Rolland, C.: Spatial and seasonal variations of air temperature lapse rates in alpine regions, *J. Clim.*, 16, 1032–
630 1046, [https://doi.org/10.1175/1520-0442\(2003\)016<1032:SASVOA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<1032:SASVOA>2.0.CO;2), 2003.
- 631 [Scotti, R., Brardinoni, F., and Crosta, G. B.: Post-LIA glacier changes along a latitudinal transect in the Central](#)
632 [Italian Alps, *Cryosph.*, 8, 2235–2252, https://doi.org/10.5194/tc-8-2235-2014](#), 2014.
- 633 Solomina, O. N., Bradley, R. S., Hodgson, D. A., Ivy-Ochs, S., Jomelli, V., Mackintosh, A. N., Nesje, A., Owen,
634 L. A., Wanner, H., Wiles, G. C., and Young, N. E.: Holocene glacier fluctuations, *Quat. Sci. Rev.*, 111, 9–34,
635 <https://doi.org/10.1016/j.quascirev.2014.11.018>, 2015.
- 636 Sommer, C., Malz, P., Seehaus, T. C., Lippl, S., Zemp, M., and Braun, M. H.: Rapid glacier retreat and
637 downwasting throughout the European Alps in the early 21st century, *Nat. Commun.*, 11,
638 <https://doi.org/10.1038/s41467-020-16818-0>, 2020.
- 639 [Vivian, R.: *Les glacier des Alpes occidentales, Grenoble: Allier, 1975.*](#)
- 640 Zanoner, T., Carton, A., Seppi, R., Carturan, L., Baroni, C., Salvatore, M. C., and Zumiani, M.: Little Ice Age
641 mapping as a tool for identifying hazard in the paraglacial environment: The case study of Trentino (Eastern
642 Italian Alps), *Geomorphology*, 295, 551–562, <https://doi.org/10.1016/j.geomorph.2017.08.014>, 2017.
- 643 Zemp, M., Hoelzle, M., and Haeberli, W.: Distributed modelling of the regional climatic equilibrium line altitude
644 of glaciers in the European Alps, *Glob. Planet. Change*, 56, 83–100,
645 <https://doi.org/10.1016/j.gloplacha.2006.07.002>, 2007.
- 646 Zemp, M., Paul, F., Hoelzle, M., and Haeberli, W.: Glacier Fluctuations in the European Alps 1850-2000: an
647 overview and spatio-temporal analysis of available data, in: *Darkening Peaks: Glacier Retreat, Science, and*
648 *Society*, University of California Press, Berkeley, 152–167, 2008.

- 649 Zumbühl, H. J. and Holzhauser, H.: Alpengletscher in der Kleinen Eiszeit, Die Alpen, Sonderheft zum 125 jährigen
650 Jubiläum des SAC, 65, 129–322, 1988.
- 651 Zumbühl, H. J. and Nussbaumer, S. U.: Little ice age glacier history of the central and western Alps from pictorial
652 documents, Geographical Research Letters, 44, 115–136, <https://doi.org/10.18172/cig.3363>, 2018.
- 653