

Comparing an idealized deterministic-stochastic model (SUP model, version 1) of the tide-and-wind driven sea surface currents in the Gulf of Trieste to HF Radar observations

Sofia Flora^{1,2}, Laura Ursella², and Achim Wirth³

¹University of Trieste, Department of Mathematics, Informatics and Geosciences, Trieste, Italy

²National Institute of Oceanography and Applied Geophysics - OGS, Trieste, Italy

³Univ. Grenoble Alpes, CNRS, Grenoble INP, LEGI, 38000 Grenoble, France

Correspondence: Sofia Flora (sflora@ogs.it)

Abstract. In the Gulf of Trieste the sea surface currents are observed by High Frequency Radar for almost two years (2021-2022) at a temporal resolution of 30 min. We developed a hierarchy of idealized models to simulate the observed sea surface currents, combining a deterministic and a stochastic approach, in order to reproduce the externally forced motion and the internal variability, which is characterized by a fat-tailed statistics. The deterministic signal includes tidal and Ekman forcing and resolves the slowly varying part of the flow, while the stochastic signal represents the fast-varying small-scale dynamics, characterized by Gaussian or fat-tailed statistics, depending on the statistical used. This is done using Langevin equations and modified Langevin equations with a Gamma distributed variance parameter. The models were adapted to resolve the dynamics under nine tidal and wind Forcing Protocols in order to best fit the observed forced motion and internal variability Probability Density Function (PDF). The stochastic signal requires 2 stochastic degrees of freedom when the averaged tidal forcing is adopted, while it needs 1/2 stochastic degree of freedom when the complete tidal forcing is used. Despite its idealization, the deterministic-stochastic model with stochastic fat-tailed statistics captures the essential dynamics and permits to mimic the observed PDF. Moreover, a Fluctuation Response Relation is valid when the stochastic signal is perturbed, showing that the response to an external perturbation can be obtained by considering the fluctuations of the unperturbed system.

1 Introduction

The Gulf of Trieste is a shallow, semi-enclosed basin in the northern Adriatic Sea (Mediterranean Sea, Fig. 1). Its surface circulation is influenced by the broader Adriatic cyclonic (counter-clockwise) circulation and the basin's thermohaline stratification and circulation (Cosoli et al., 2012; Querin et al., 2021). Due to its shallowness and small scale (horizontal length scale of around 20 km and depth of around 25 m), its current dynamics highly depend on two external forcings: the wind forcing and the Isonzo/Soča river freshwater input (Querin et al., 2006; Cosoli et al., 2012, 2013; Querin et al., 2021).

The Bora and the Sirocco are the main strong wind patterns affecting the dynamics in the Gulf of Trieste. The Bora is an East-North-Easterly katabatic wind, bringing cold and dry air from the continent over the sea (Poulain and Raicich, 2001). Its strength is heavily influenced by local topography, making it particularly powerful over the Gulf of Trieste where it can reach

gusts up to 180 km/h. When it blows, surface water is pushed out of the Gulf towards the Adriatic basin, and a compensating bottom counter-current flows in, causing upwelling on the coastal side (Malačič et al., 2001; Querin et al., 2006; Reyes Suárez et al., 2022). The Sirocco is a moist, warm and relatively mild southerly wind, channeled by the Adriatic coastal mountains. It causes sea level rise in the northern Adriatic, leading to a strong southward return flow when the wind diminishes (Cosoli et al., 2012).

In recent years, the Gulf of Trieste has benefitted from an abundance of observational data, particularly High Frequency Radar (HFR) sea surface currents. The amount of data will strongly increase in the near future. This large amount of data allowed for a statistical approach to characterize the data (Flora et al., 2023). In particular, the authors found the analytical Probability Density Function (PDF) of the observed HFR sea surface velocity increments using a superstatistical (a superposition of statistics) analysis and the principle of maximum entropy. Superstatistical analysis (Beck et al., 2005) considers a system with a clear time scale separation: at fast time scale a Gaussian PDF is observed of which the variance evolves slowly, on the long time scale. In the Gulf of Trieste the fast time scale of around 2 h is consistent with the turbulence of the mixed layer dynamics. The slow time scale of almost 2 days is consistent with the variability of the synoptic wind forcing, and the variance is distributed according to a Gamma distribution with shape parameter equal to 2. The Gaussian is the minimum entropy PDF with a finite variance and the Gamma distribution is the minimum entropy PDF for a positive variable, as is the variance, with a specified mean. The total PDF of the sea surface velocity increments is a fat-tailed PDF, that means that extreme events occur more often than in a Gaussian statistics. The superstatistical analysis however is descriptive and has no predictive skills.

When modelling the turbulent ocean currents subject to a known wind forcing, there is part of the dynamics which can be described by a deterministic model and the unresolved turbulent processes that have to be described stochastically. This concept is reflected in Stochastic Differential Equations (SDEs) and the corresponding Fokker-Planck equations. The former consider the evolution of many possible realizations which allow to construct a time evolving PDF, while the latter describe the deterministic evolution of the PDF. The use of SDEs in air-sea interaction was pioneered by Hasselmann (1976), who initiated a strand of scientific research on stochastic climate models. Taking into account the coupled ocean-atmosphere-cryosphere-land system, he divided it into a fast varying "weather" system and a slowly responding "climate" system, clearly separated by a fast and a slow time scale. In his modelization the slow climate variables play the role of large particles interacting with an ensemble of smaller particles, the fast weather variables in the analogy, of the so-called Brownian motion problem. In this picture, the dispersion of the climate variables is inferred from the statistics of the weather variables with which they interact. Brownian motion was first analytically described by Einstein (1905) and Langevin et al. (1908) (see Einstein (1956) and Lemons and Gythiel (1997) respectively for an English translation). Since that time, the SDE of the motion of a Brownian particle is called the "Langevin equation". The SDE approach, adopted also in this paper, has been widely used in the past decades in a variety of scientific fields: statistical mechanics (Baldovin et al., 2018), condensed matter physics (Silveira and Aarão Reis, 2012), marine biology and oceanography (Brillinger and Stewart, 2010), space weather science (Alberti et al., 2018), biophysics (Ham et al., 2022) and finance (Wand et al., 2024) to name a few. Regarding climate science, the Langevin equation

has been widely used (Berglund and Gentz, 2002; Wirth, 2019; van den Berk et al., 2021) and great improvements have been achieved. Franzke et al. (2015) and Palmer (2019) show that including stochasticity into the parameterized representations of subgrid processes of physical climate systems has improved the skill of forecasts and reduced systematic model error. It is important to remark that the use of additive noise in linear SDE has limitations in terms of extreme events predictability, since it leads to a Gaussian statistics. Adopting multiplicative noise or nonlinear damping, taking into account non-linear interactions between resolved and unresolved modes of variability (Franzke et al., 2005; Wirth, 2018) leads to non-Gaussian statistics increasing the fatness of the PDF tails of the modelled variables (Sura, 2013).

Modern concepts of non-equilibrium statistical mechanics have been applied to environmental fluid dynamics and to components of the climate system. Often these considerations are limited to conceptual pertinence and tested on idealized models, such as the Lorenz models (Lorenz, 1996) and others, or on data from numerical models. In the present work we apply a hierarchy of idealized models for the sea surface current observations from the Gulf of Trieste, starting with a deterministic modelization and then adding increasingly complex stochasticity. For solving these models we rely on analytical calculations where possible and proceed with solutions of involved SDEs. This approach enables connecting observational data in a systematic way to the underlying principles of physics.

Our modelling approach is idealized and local: it considers one point in the space, evolving in time, with wind shear and tidal deterministic forcing and a stochastic signal that mimics all the smaller unresolved scale dynamics. The aim of the modelization is to simulate the evolution of the forced motion and of the observed analytical PDF. It is based on the slow evolution of a short time scale Gaussian and can be interpreted by a SDE. This enables us to understand which is the role of the stochasticity in the simulation of the Gulf of Trieste wind- and tide-driven circulation. The model is finally used to test the Fluctuation Response Relation (FRR), described by Lacorata and Vulpiani (2007), that relates the system's reaction to external perturbations to spontaneous fluctuations of the unperturbed system. An evaluation of the ocean currents forecasting based on the SDE predictability is also given.

In Sect. 2 the observed HFR sea surface data and the model forcing atmosphere data are presented. In Sect. 3 the modelization is explained with some FRR background and predictability evaluation methodology, while the computational results and their discussion are given in Sect. 4. The conclusions are reported in Sect. 5. In the Appendixes the details and methods of the analytical calculations are shown.

2 The data

Two classes of time series are used: the observed sea surface horizontal HFR current data and the forecasted wind data in the Gulf of Trieste (Northern Adriatic Sea, Fig. 1a). The data sets used are identical to those presented in Flora et al. (2023) and refer to a selected grid point in the Gulf of Trieste (point P and point WRF_P in Fig. 1b) and cover a time range of almost two years: from January 1, 2021 to October 18, 2022.




Figure 1. (a) Gulf of Trieste location (red rectangular) in the Adriatic Sea; (b) Gulf of Trieste zoom with percentage of available HFR data (in multiple colors) in the selected period. The HFR baseline is shown with the red line between Aurisina and Piran. The HFR "P" grid point is shown with the black asterisk and the closest WRF grid point is marked with the blue star and called "WRF_P" (Flora et al., 2023).

Sea surface currents are measured by two beamforming WELLEN RADAR stations operating in the Gulf of Trieste (Lorente et al., 2022) (Fig. 1b, red dots). The HFR system combines the radial sea surface currents of the two stations to obtain the longitudinal u and the latitudinal v components, with a space resolution of 1.5 km and a time resolution of 30 min. The data are open access, further details can be found in OGS et al. (2023). The quality control standards from the EU high-frequency node (Corgnati et al., 2018) were applied to the data set. In addition, any remaining spikes were removed. The measured u and v time series used in this article are from the P grid point (Fig. 1b).

The atmosphere data consist of the forecasted wind velocity field at 10 m above the surface in the Gulf of Trieste from the Weather Research and Forecasting (WRF) model (<https://www.mmm.ucar.edu/models/wrf>), version 4.2.1. The forecasting is performed daily by the Agenzia Regionale per la Protezione dell'Ambiente del Friuli Venezia Giulia (ARPA FVG) using initial and boundary conditions from the National Oceanic and Atmospheric Administration Global Forecasting System (<https://www.ncei.noaa.gov/products/weather-climate-models/global-forecast>) and provides the wind time series for the same day and the following one. The field has a spatial resolution of 2 km and a temporal resolution of 1 h. Additional technical details can be found in Goglio (2018). The wind components time series used in this article are from the WRF_P grid point (Fig. 1b) and account for the one-day forecasting.

This Section provides the model definition, the FRR background with details on how it is dealt within this study and some methodology to evaluate the predictability of the developed model.

3.1 The model hierarchy

The aim of the present work is to numerically simulate observed sea surface currents with a slow dynamics, that is governed by the slow components of the forcings and with fast variations that parameterise the remaining, unresolved, processes. This is achieved when the time averaged dynamics as well as the statistics of the time increments of the model agrees with the observation. In the following, the time series prefix " δ " gives the increment time-series, $\delta u(t) = u(t + \delta) - u(t)$, and we call the observed fat-tailed PDF of the sea surface current increments, from Flora et al. (2023), as the "Exp-Lin" PDF, since it is composed of an exponential and a linear term.

When modelling a natural process, as the currents, through a SDE, part of the dynamics is resolved and part of it is parameterized by noise. What is deterministic and what is noise depends on the degree of coarse-graining. In our previous work (Flora et al., 2023) all the dynamics was parameterized and the model has no predictability. The other extreme is a purely deterministic model that includes all the processes involved. Such model is unattainable. We implemented a hierarchy of three idealised local models (two velocity components at one point in space, evolving in time) for the sea surface current in the Gulf of Trieste. The first is the purely DETERministic (DET) model, taking into account the tidal signal and the wind-forced Ekman dynamics, while the unresolved processes are neglected. The second is the GAUSSian (GAU) model, that adds Gaussian additive stochastic noise for which the corresponding SDE can be solved analytically. The third is the SUPERstatistical (SUP) model that considers a deterministic signal with the most realistic stochastic noise. The stochastic part of this model is partially resolvable analytically. Furthermore, these models are forced by nine different Forcing Protocols (FPs) (see Sect. 4). The deterministic and stochastic models come in pairs, the more processes and time-scales are resolved deterministically, the less have to be parameterised stochastically.

More precisely, the total sea surface current vector $\mathbf{u}_o$ is given in terms of the tidal current $\mathbf{u}_M$, the wind-forced Ekman sea surface current $\mathbf{u}_E$ and the stochastic current $\mathbf{u}_S$, as detailed in Table 1 for the different models. The tidal signal is an analytic and linear combination of n tidal components $u_M(t) = \sum_{i=1}^n a_{u,i} \cos(\omega_i t - \phi_{u,i})$ (analogously for v_M), where a , ω and ϕ are the amplitude, the frequency and the phase of each tidal component, respectively.

The Ekman current, subject to the wind forcing, the Coriolis force and the friction with the underlying water-masses, is modelled by:

$$\frac{d}{dt} \begin{pmatrix} u_E \\ v_E \end{pmatrix} = \begin{pmatrix} -\frac{C_B}{h} \tilde{u} & f \\ -f & -\frac{C_B}{h} \tilde{u} \end{pmatrix} \begin{pmatrix} u_E \\ v_E \end{pmatrix} + \frac{1}{\tilde{h}} \begin{pmatrix} F_u \\ F_v \end{pmatrix}. \quad (1)$$

In Eq. (1), $C_B = \rho_o c_b$ where c_b is the underlying ocean layer drag coefficient, $\tilde{h} = \rho_o h$ where ρ_o is the ocean density and h is the considered ocean surface layer depth, $\tilde{u}$ is the total speed of the surface layer as defined in Table 1. The last does

not include the tidal current, since it is assumed that the tidal signal is constant along the water column and does not create shear, while the Ekman and stochastic signals are present in the surface layer only. Finally f is the Coriolis parameter and $\mathbf{F} = \rho_a c_a |\mathbf{u}_a - \mathbf{u}_o|(\mathbf{u}_a - \mathbf{u}_o)$ is the wind forcing (ρ_a is the atmosphere density, c_a the atmosphere drag coefficient and $\mathbf{u}_a$ the wind velocity at 10 m above the sea surface). Eq. (1) is an ordinary differential equation of the DET model, while it includes the stochastic quantity $\tilde{u}$ in the GAU and SUP models. If the wind forcing were to cease, the modelled mean flow would fall to the tidal signal.

The Ekman modelization in Eq. (1) is only valid on frequency scales comparable to the Coriolis parameter. The stochastic velocity, representing the fast unresolved turbulent dynamics and present in the GAU and SUP models, is defined by the following set of equations (for the purpose of the study, i.e. to reproduce the observed superstatistical statistics, it is sufficient to have uncorrelated stochastic velocity components):

$$d \begin{pmatrix} x \\ y \end{pmatrix} = -\gamma_x \begin{pmatrix} x \\ y \end{pmatrix} dt + \begin{pmatrix} \sqrt{Q_u} & 0 \\ 0 & \sqrt{Q_v} \end{pmatrix} \begin{pmatrix} dW_x \\ dW_y \end{pmatrix} \quad (2)$$

$$\frac{d}{dt} \begin{pmatrix} u_S \\ v_S \end{pmatrix} = -\gamma_u \begin{pmatrix} u_S \\ v_S \end{pmatrix} + \eta \begin{pmatrix} x \\ y \end{pmatrix} \quad (3)$$

In Eq. (2) and Eq. (3) the variables dW_x and dW_y are derived from independent Wiener stochastic processes. For every independent realization of the Wiener processes W_x and W_y , we obtain a solution of the dynamics, named a random walker. The collection of all the random walkers forms the stochastic ensemble and allows to construct a PDF. The coefficient $\gamma_x = 1/\tau$ is the inverse of the characteristic short time scale τ (short compared to the long time scale discussed below in connection with the SUP model) whose numerical value is determined from the observations (Flora et al., 2023), γ_u is a coefficient which is a posteriori adjusted to obtain the observed HFR initial decay of the autocorrelation function of the SUP model sea surface currents (Fig. 7) and η is a proper constant discussed below. The difference between the GAU and the SUP model lies in the terms Q . In the GAU model, that is a simplification of the SUP model, Q_u and Q_v are constants: they are the mean values of the SUP model Q variables as given in Table 1. The GAU model is a system of linear SDEs and all variables are Gaussian and are determined by their means and variances. All the dependencies for means and variances on the original parameters are given in Appendix A. Furthermore in Eq. (A32) the analytical condition on η for which the Gaussian variable δu_S has the same mean and variance as x (analogously δv_S with y), given a fixed γ_u , is shown. This also helps to adjust the parameters of the SUP model.

In the SUP model, the Q terms originate themselves from 2ν stochastic processes α_i : $Q = \sum_{i=0}^{2\nu} \alpha_i^2$. The variable ν gives the degrees of freedom (dof), according to the interpretation of Flora et al. (2023): the system has one dof if its positive variable characterizing the variability of the system maximizes the entropy, i.e. it is exponentially distributed. This corresponds to half

the value of Beck and Cohen (2003)'s interpretation. Each stochastic process α_i is governed by:

$$d\alpha_i = -\mu\alpha_i dt + \beta dW_i, \quad i = 1, \dots, 2\nu \quad \text{and} \quad \beta = \begin{cases} \beta_u & \text{for the } u \text{ component} \\ \beta_v & \text{for the } v \text{ component} \end{cases} \quad (4)$$

where the variables dW_i are derived from independent Wiener stochastic processes, β_u and β_v are constants whose values are determined empirically, $\mu = 1/T$ is the inverse of the characteristic long time scale T (i.e. $\tau \ll T$), corresponding to the variation time-scale of the variance in the superstatistical approach. Its numerical value is determined from the observations (Flora et al., 2023). The variables α_i are solutions of the Langevin equation: they are Ornstein-Uhlenbeck processes (we refer the reader to Kloeden and Platen (1999) for a pedagogical discussion of stochastic processes), characterized by a Gaussian statistics:

$$p_\alpha(\alpha_i) = \sqrt{\frac{\mu}{\pi\beta^2}} e^{-\mu\alpha_i^2/\beta^2}. \quad (5)$$

In the SUP model the variables Q depend on the variables ν and α_i , as shown in Table 1. The Q variables are distributed according to a Gamma distribution with shape parameter equal to ν , given in Eq. (B5) and Table B1. In Eq. (B6) we give the computation for its first moment $E[Q]$, used in the GAU model. In Appendix B the PDFs of the variable x (and y) are derived for $\nu = 2$ and $\nu = 1/2$. A decrease of the stochastic degrees of freedom ν leads to a reduction of the variability (non-maximal entropy) in the unresolved processes, represented by the stochastic variables. In the case of 2 dof, the variables x and y are Exp-Lin distributed with a PDF given by:

$$\begin{cases} p_{\nu=2}(x) = \frac{\sqrt{\mu\gamma_x}}{2\beta_u} e^{-2\sqrt{\mu\gamma_x}|x|/\beta_u} \left(\frac{2\sqrt{\mu\gamma_x}}{\beta_u}|x| + 1 \right) \\ p_{\nu=2}(y) = \frac{\sqrt{\mu\gamma_x}}{2\beta_v} e^{-2\sqrt{\mu\gamma_x}|y|/\beta_v} \left(\frac{2\sqrt{\mu\gamma_x}}{\beta_v}|y| + 1 \right). \end{cases} \quad (6)$$

In the case of 1/2 dof, the variables x and y are distributed according to a modified Bessel function of the second kind of zero order $K_0(|z|)$:

$$\begin{cases} p_{\nu=\frac{1}{2}}(x) = \frac{2\sqrt{\mu\gamma_x}}{\pi\beta_u} K_0\left(\frac{2\sqrt{\mu\gamma_x}}{\beta_u}|x|\right) \\ p_{\nu=\frac{1}{2}}(y) = \frac{2\sqrt{\mu\gamma_x}}{\pi\beta_v} K_0\left(\frac{2\sqrt{\mu\gamma_x}}{\beta_v}|y|\right). \end{cases} \quad (7)$$

The SDEs for the variables u_S and v_S in Eq. (3) contain a linear drag term and are forced by a coloured non-Gaussian noise. We do not know an analytical distribution of u_S and v_S . Nevertheless, thanks to the definition of the constant η in Eq. (A32) discussed in connection with the GAU model, the distributions of δu_S and δv_S are numerically similar to the PDFs of x and y (shown in Sect. 4.1).

As it can be seen in Eq. (6) and Eq. (7), the final stochastic PDFs depend on the β parameter, that is linked to the variable's second order moment, by $E[x^2] = \frac{\beta^2}{\mu\gamma_x}$ for $\nu = 2$. For this case, it will be shown in Sect. 4.1 that the PDF of the total velocity increment is given mainly by the stochastic velocity increment PDF, only. For this reason in the numerical simulations with 2 dof in the stochastic part of the model the value of β is therefore fixed by the observed variance s^2 of the HFR velocity

increment found in Flora et al. (2023): $\beta = s\sqrt{\mu\gamma/x}$. In the case of 1/2 dof in the stochastic signal, the tidal and Ekman signal
 195 contribute more to the variance of the total velocity increment, but their contribution is unknown analytically. For this reason
 in this case the β coefficient is increased empirically, in order to fit the observed variance in the total velocity increment PDF.
 In Table 1 the definition of the variables distinguishing the DET, GAU and SUP models is given with a summary of the main

	u_o	$\tilde{u}$	Q	x and y distribution	δu_S and δv_S distribution
DET	$u_E + u_M$	$ u_E $	no stochasticity	no stochasticity	no stochasticity
GAU	$u_E + u_M + u_S$	$ u_E + u_S $	$E[Q]_{SUP} = \nu\beta^2/\mu$	Gaussian, Eq. (A28)	Gaussian, Eq. (A30)
SUP	$u_E + u_M + u_S$	$ u_E + u_S $	$\sum_{i=1}^{2\nu} \alpha_i^2$	$\nu = 2$: Exp-Lin, Eq. (6) $\nu = 1/2$: Bessel, Eq. (7)	not analytical, but numerically similar to the x and y PDF

Table 1. Variables definition distinguishing the different models and x , y , δu_S and δv_S PDFs' characteristics. The variable ν indicates the degrees of freedom of the stochastic signal.

properties of the models' variables. For all the models we can consider, or not, the eddy depletion term in the wind stress term,
 i.e. the relative velocity of the wind with respect to the sea surface current, causing a reduction of kinetic energy injection into
 200 the ocean (Zhai et al., 2012), in contrast to the absolute wind velocity. We report the results just with the eddy depletion term,
 since it is the theoretically most correct formulation and it does not significantly burden the computational calculation.

In summary, we developed three types of models: the DET model is purely deterministic with tidal and Ekman currents, the
 GAU model adds a Gaussian stochastic signal, while the SUP model considers a stochastic sea surface current with fat-tailed
 increments instead of the Gaussian noise (Table 1).

205 3.2 The FRR background and methods

The FRR relates the response of a system perturbed by an external perturbation to internal fluctuations of the unperturbed
 system. Exploring the FRR is particularly challenging and interesting when considering natural systems where perturbation
 experiments cannot be performed and statistical ensembles are not available. Our SUP model is an example of a subcomponent
 of the climate system and it is instructive to explore the FRR for such model. We refer to Lacorata and Vulpiani (2007) for the
 210 theoretical background concerning FRRs. We show here some fundamentals of the theoretical concepts and the methodology
 we applied to the most involved of the SUP models (FP9, Table 2, Sect. 4).

In the present case, given the SUP model evolutionary system with the SUP variables $u_o = u_E + u_M + u_S$ and $v_o = v_E +$
 $v_M + v_S$, a small fixed perturbation Δu_S or either Δv_S is imposed at $t = t_0$ to each walker. Then the system is left free to
 215 evolve and, for each time, the separation $\Delta u_o(t|t_0) = u_{o,P}(t) - u_o(t)$ and $\Delta v_o(t|t_0) = v_{o,P}(t) - v_o(t)$ between the perturbed
 ($u_{o,P}$ and $v_{o,P}$) and the unperturbed (u_o and v_o) systems is computed. We emphasise that the methodology requires one ve-
 locity component to be perturbed and not both in the same perturbed simulation. It is possible to obtain the mean value of the

perturbed variables through a mean response function $R_{uu}(t)$ or $R_{vv}(t)$:

$$\begin{aligned}\langle \Delta u_o(t) \rangle &= R_{uu}(t) \Delta u_S && \text{when } u_S \text{ is perturbed} \\ \langle \Delta v_o(t) \rangle &= R_{vv}(t) \Delta v_S && \text{when } v_S \text{ is perturbed}\end{aligned}\tag{8}$$

220 where the angle brackets $\langle \dots \rangle$ indicate the ensemble mean. The FRR is concerned with expressing the mean response function in terms of correlation functions of the unperturbed system.

In practice, the mean response functions are computed as follows. After the first perturbation at t_0 , again at $t_k = t_0 + k\Delta t$ the selected variable, in the perturbed system, is perturbed with the same variation and the separation between the perturbed and the unperturbed systems is computed. The perturbation time is fixed to $\Delta t = 12$ h, allowing us to repeat the procedure

225 $M = 1311 \gg 1$ times, and $\Delta u_S = \Delta v_S = 8$ cm/s. The diagonal mean response functions are computed as follows:

$$\begin{aligned}R_{uu}(t) &= \frac{1}{M} \sum_{k=0}^{M-1} \frac{\langle \Delta u_o(t_k + t | t_k) \rangle}{\Delta u_S} && \text{when } u_S \text{ is perturbed} \\ R_{vv}(t) &= \frac{1}{M} \sum_{k=0}^{M-1} \frac{\langle \Delta v_o(t_k + t | t_k) \rangle}{\Delta v_S} && \text{when } v_S \text{ is perturbed.}\end{aligned}\tag{9}$$

According to Lacorata and Vulpiani (2007), the FRR holds if the mean response functions have a connection with some suitable correlation functions computed in the unperturbed system. In particular, in the case of multivariate Gaussian variables, the mean response functions are a linear combination of the variables correlations. In Sect. 4.2 the results with unperturbed
230 initial condition $\mathbf{u}_o(t=0) = 0$ and when the stochastic signal is initially perturbed, as described before, is shown. Some brief comments for the case when the Ekman system is initially perturbed are also provided.

3.3 Predictability evaluation methods

In order to test predictability capabilities of the SUP model, perturbation methods are adopted assuming that the HFR observations are not affected by observational uncertainty and represent the reality, i.e. modelled data are replaced by observations. In
235 the following this method is called "observation-based perturbation". In detail, every Δt time the perturbed system ($\mathbf{u}_{o,P}$) is updated to the observed HFR velocities (both the u_o and v_o components in the same simulation), in particular it is the stochastic signal to be perturbed. The method may appear equivalent to the FRR method, but the equivalence is not valid, since: (i) the initial perturbation is not a constant, but changes for each Δt time window and for each walker; (ii) we are perturbing both the u_o and v_o components (in the stochastic signal) in the same perturbed simulation. We define the following functions:

$$240 \quad \xi(t) = \sqrt{\frac{\frac{1}{M} \sum_{k=0}^{M-1} \langle (\mathbf{u}_{o,P}(t_k + t) - \mathbf{u}_o(t_k + t))^2 \rangle}{\frac{1}{M} \sum_{k=0}^{M-1} \langle (\mathbf{u}_{o,P}(t_{k-1} + \Delta t) - \mathbf{u}_{HFR}(t_k))^2 \rangle}}\tag{10}$$

$$\epsilon(t) = \sqrt{\frac{\frac{1}{M} \sum_{k=0}^{M-1} \langle (\mathbf{u}_{o,P}(t_k + t) - \mathbf{u}_{HFR}(t_k + t))^2 \rangle}{\frac{1}{M} \sum_{k=0}^{M-1} \langle (\mathbf{u}_{o,P}(t_{k-1} + \Delta t) - \mathbf{u}_{HFR}(t_k))^2 \rangle}}\tag{11}$$

where the time $t_{k-1} + \Delta t$ means t_k right before the perturbation. The function $\xi(t)$ quantifies the difference between the perturbed and the unperturbed system, while the function $\epsilon(t)$ computes the difference between the perturbed system and the observations. Both are normalized through the mean initial perturbation over all the time windows. In Sect. 4.3 the results are presented. Some brief comments for the case when the Ekman system is initially perturbed are also provided.

4 Results and discussion

We test the different models imposing different Forcing Protocols (FPs) in the tidal and wind time series (Table 2). In the most complete FP the complete tidal and wind forcing is included (FP9), in the other FPs the forcing signal is either averaged by a moving average over 12 h or completely set to zero.

Forcing Protocol		1	2	3 (n.s.)	4 (n.s.)	5	6 (n.s.)	7 (n.s.)	8	9
Forcings	tide	✗	✗	✗	$\overline{12h}$	$\overline{12h}$	$\overline{12h}$	✓	✓	✓
	wind	✗	$\overline{12h}$	✓	✗	$\overline{12h}$	✓	✗	$\overline{12h}$	✓
Results	dof ν	2	2	2	2	2	2	1/2	1/2	1/2
	$\frac{\beta}{s\sqrt{\mu\gamma_x}}$	1	1	1	1	1	1	1.5	1.5	1.5

Table 2. Settings and results of the experiments on the models: the ✗ symbol means that the forcing is set to 0, the $\overline{12h}$ symbol means that the forcing is filtered with a 12 h moving average, the ✓ symbol means that we use the original forcing. The variable s is the observed standard deviation of the HFR velocity increment found in Flora et al. (2023). The abbreviation n.s. stays for "not shown".

The numerical values of the coefficients common to all the models and to all the runs are described here. Tidal harmonic analysis calculations on the HFR sea surface currents were done using the MatLab programme `t_tide` (Pawlowicz et al., 2002). The resultant tidal ellipse parameters are shown in Table 3. The $S1$, $M2$ and $S2$ tidal components show a sufficiently high Signal-to-Noise Ratio (SNR) and are therefore included to be considered in our modelization. The conversion from the ellipse parameters to the sinusoidal tidal coefficients is applied according to Foreman (1978). The non-filtered tidal signal consists of the $S1$, $M2$ and $S2$ components, while the 12 h moving averaged signal is characterized by the diurnal oscillation only. The wind velocity u_a time series is linearly interpolated from its 1 h original resolution or from a 12 h moving average to the

tide	ω [h^{-1}]	SEMA [cm s^{-1}]	SEMI [cm s^{-1}]	INC [$^\circ$]	PHA [$^\circ$]	SNR
$S1$	0.0426667	6.732 ± 0.692	-2.522 ± 0.66	-26.46 ± 6.65	233.62 ± 7.64	95
$M2$	0.0805114	3.537 ± 0.407	0.296 ± 0.39	47.84 ± 6.05	151.27 ± 5.96	76
$S2$	0.0833333	3.389 ± 0.454	-0.167 ± 0.39	23.67 ± 7.22	130.69 ± 7.59	56

Table 3. Tidal ellipse parameters with 95 % confidence interval estimates, extracted from the HFR data.

integration time step of the numerical model (2.5 min). The determination of $\tilde{h}$ and C_B as best linear fit functions of the wind speed is given in Appendix C. The algorithm to determine the wind regimes Bora, Sirocco, Mistral and Low wind is explained

260 in Flora et al. (2023): if the daily wind speed is higher than the threshold value of 3 m/s, then it is defined as Bora, Sirocco or Mistral depending on its direction; if it is lower, it is classified as Low wind. Mistral occurs just for a few days in almost two years, it is therefore considered statistically not significant. In the numerical integration the parameters shown in Table 4 are used and a Runge-Kutta method of the second order is adopted for the Ekman system.

parameter	value
total integration time	656 days
Runge Kutta time step	3×10^2 s
δ	1.44×10^4 s
ρ_a	1.3 kg m^{-3}
c_a	10^{-3}
ensemble size	10^5
stochastic time step dt	1.5×10^2 s
T	1.746×10^5 s
τ	6.6×10^3 s
$\mu = 1/T$	$5.728 \times 10^{-6} \text{ s}^{-1}$
$\gamma_x = 1/\tau$	$1.515 \times 10^{-4} \text{ s}^{-1}$
s_u	$12.265 \times 10^{-2} \text{ m s}^{-1}$
s_v	$9.6696 \times 10^{-2} \text{ m s}^{-1}$
$s_u \sqrt{\mu \gamma_x}$	$3.612 \times 10^{-6} \text{ m s}^{-2}$
$s_v \sqrt{\mu \gamma_x}$	$2.848 \times 10^{-6} \text{ m s}^{-2}$
γ_u	$1.166 \times 10^{-4} \text{ s}^{-1}$
η	$1.660 \times 10^{-4} \text{ s}^{-1}$

Table 4. Involved parameters in the numerical modelization: total integration time, Runge Kutta time step, increment time δ (from the velocity increment definition), atmosphere density ρ_a , atmosphere drag coefficient c_a , ensemble size, stochastic time step dt , long time scale T , short time scale τ , α_i 's linear drag coefficient μ , x 's and y 's linear drag coefficient γ_x , observed HFR velocity increments standard deviations s_u and s_v , u_S 's and v_S 's linear drag coefficient γ_u and forcing coefficient η . The numerical values of δ , τ , T , s_u and s_v are taken from Flora et al. (2023).

4.1 Comparison of the DET, GAU and SUP models on different forcings time scales

265 All the models start from the initial condition $\mathbf{u}_o(t=0) = \mathbf{u}_{\text{HFR}}(t=0)$, where $\mathbf{u}_{\text{HFR}}$ is the observed HFR sea surface velocity, with $\mathbf{u}_S(t=0) = 0$. The obtained numerical sea surface currents are then compared to the HFR measurements. We start this section commenting FP1 which has no tides and no wind forcing (see Table 2) and compare the DET, GAU and SUP models in the FP5 and FP9 cases (see Table 2). We conclude this section with a discussion on the change of the stochastic dof when the tidal forcing is modified. We only show time series for the u -component of the sea surface current variables, results for the

270 v -component are similar.

The stochastic model has to be adapted to the FPs to best represent the dynamics unresolved by the deterministic part of the model. The FP1 does not take into account any physical forcings and the deterministic part vanishes. The DET model is therefore null, while the GAU and SUP models have a non-zero stochastic part. The SUP model best fits the observed data
275 when $\nu = 2$ dof are chosen (see Flora et al. (2023)), as we obtain a total velocity increment ($\delta u_o = \delta u_S$, because in FP1 $\delta u_M = \delta u_E = 0$) distributed according to the Exp-Lin PDF (as summarized in Table 1). The result is shown in Fig. 2 left column, in which SUP δu_o of FP1 coincides with SUP δu_S of FP5 (as shown in Table 2 and commented below, they both have $\nu = 2$ stochastic dof). The GAU model in FP1 (Eq. (A30) and Fig. 2 left column, in which GAU δu_o of FP1 coincides with GAU δu_S of FP5) fails in reproducing the fat-tailed Exp-Lin PDF.

280




Figure 2. PDFs of the stochastic variables of the GAU and SUP models for FP5 (left column) and FP9 (right column) in lin-lin plots (top row) and lin-log plots (bottom row). In particular, analytical solution of the GAU δu_S variable from Eq. (A30) (blue continuous line) and PDFs of the models variables: numerical GAU δu_S (blue dots), SUP x (yellow continuous line), SUP δu_S (red continuous line). PDFs of the analytical superstatistical Exp-Lin PDF from Eq. (6) (black dotted line with crosses) and of the analytical superstatistical Bessel function from Eq. (7) (black dotted line with circles) are also shown. Note that in FP5 (FP9) the SUP x , the SUP δu_S and the Exp-Lin (Bessel) lines almost superpose.

In the FP5 a 12 h moving-averaged tidal and wind forcing is applied to the models, while the FP9 includes the forcings without averaging (see Table 2). For the different types of forcings we choose the dof that best fitted the superstatistical Exp-Lin PDF from Flora et al. (2023). In particular, for FP5 $\nu = 2$, while for FP9 $\nu = 1/2$ give the best fits. In Fig. 2 the PDFs of the stochastic variables of the GAU and SUP models for FP5 and FP9 are compared to the analytical PDFs. The figures validate the choice of the parameters in our model hierarchy for the different forcings, as summarized in Table 1: (i) the GAU δu_S follows the analytical Gaussian from Eq. (A30); (ii) in FP5 the stochastic variable SUP x is Exp-Lin distributed while in FP9 it is Bessel distributed; (iii) the SUP δu_S are distributed as SUP x in both the simulations, showing the validity of the choice of η (fixed according to the analytical condition from the GAU model in Eq. (A32), its choice is discussed in Sect. 3.1 in

connection with the GAU and SUP models). This also validates the numerical model through the cases where the analytical
 290 solution is known.




Figure 3. Sea surface currents time series (u -component) during eleven days of April 2021 of the HFR observations with their measure accuracy (blue sticks with error bars) and from the models for FP5 (top) and for FP9 (bottom): DET model u_o with the Gaussian GAU model ensemble standard deviation of the stochastic velocity (yellow lines), GAU model ensemble mean of u_o with their ensemble standard deviation (blue lines), SUP model ensemble mean of $u_E + u_M$ with their ensemble standard deviation (light blue lines), SUP model ensemble mean of u_o with their ensemble standard deviation (red lines). The SUP u_o path of two random walkers is reported (light and dark green dotted lines). The symbol $\langle \dots \rangle$ in the legend stands for the ensemble mean. The blue and red shadings represent Bora and Sirocco wind regimes respectively, seen in Flora et al. (2023).




Figure 4. Normalised 2D histograms of the correlation between the HFR observations and the DET model u -component velocities from the simulated period (almost two years), from FP5 (top) and from FP9 (bottom). The r correlation coefficient is reported. The light blue line is the linear regression of the scattered data, while the red line is the bisector.

In Fig. 3 the time series of the observed and model velocities for some days of April 2021 are reported. The difference in the forcing time scale is clearly visible in the deterministic DET model velocities variability: smoother in FP5 than in FP9. They both follow the slow variability of the observed HFR currents. In fact, looking at Fig. 4, the linear regression of the scattered data has a lower slope with respect to the bisector, meaning that the DET model is not able to simulate the observed extremes.

295 Nevertheless, their 2D histograms have an elongated peak along the bisector and their correlation coefficient reaches 55 % in FP5 and 56 % in FP9 for the u -component (29 % and 35 % for the v -component, respectively). Moreover, it is possible to see that sometimes (e.g. 3rd of April 2021 from Fig. 3) the observed u_{HFR} sea surface current shows a negative peak at the beginning of Bora events. This behaviour is missing in the DET model u_o velocity. This fact can be due to the model instantaneous change of the values of $\tilde{h}$ and C_B with the wind speed, while the sea surface system in nature takes some time to adjust to the

300 wind forcing through vertical mixing (the thin surface layer suddenly accelerates, then thickens and the speed decreases).

Regarding the ensemble averaged sea surface currents of the GAU and SUP models, they follow closely the DET model time series. This shows that the stochastic models increase the variability to observed values, but have little influence on the averaged dynamics. We emphasise that if our model were perfect, the observations would be one of the walkers and therefore would at times be outside the SUP total velocity ensemble standard deviation σ_{u_o} (red) band, as is the case for the two random

305 walkers (green lines) reported in Fig. 3. The SUP ensemble standard deviation of the $\mathbf{u}_E + \mathbf{u}_M$ velocity (light blue band) is smaller than the SUP total velocity ensemble standard deviation σ_{u_o} (red band), as expected, since the tidal velocity is purely deterministic and in the Ekman velocity the stochasticity enters only through the drag. What is particularly interesting is that the GAU σ_{u_o} ensemble standard deviation (blue band) almost completely coincides with the SUP σ_{u_o} (red band) and not with

the GAU σ_{u_S} (yellow band) ensemble standard deviation. A clear example of that is visible on the 3rd, 4th, 6th and 13th of April 2021 in Fig. 3 for the u -component, where the wind forcing increases and the GAU and SUP σ_{u_o} standard deviations (blue and red bands respectively) reduce with respect to the GAU σ_{u_S} statistics (yellow band). In this case it originates from the stochasticity present in the Ekman drag term.




Figure 5. PDFs of the observed, analytical and numerical velocity-increments from FP5 (left column) and FP9 (right column) in lin-lin plots (top row) and lin-log plots (bottom row). In particular, we show the PDFs of the observed HFR velocity-increment (black crosses), of the analytical superstatistical PDF from Eq. (6) (black line), of the analytical Gaussian of the GAU δu_S variable from Eq. (A30) (blue dashed line) and of the models variables: DET δu_o (yellow line), GAU δu_o (blue line), SUP $\delta u_E + \delta u_M$ (dotted red line), SUP δu_o (full red line).

Looking at the PDFs of the velocity increments in Fig. 5, it can be seen that the DET model PDF in the FP5 is more peaked around zero, as compared to FP9 which is closer to the observations. This is because the latter model has a higher variability due to the inclusion of higher frequency forcing data. Despite the rather high correlation of the DET currents with the observations, the DET model does not explain most of the variability seen in the HFR data, especially the tails, the extreme events, are not well reproduced.

From Fig. 5 one can also observe that in FP5 the GAU δu_o PDF is similar to the analytical PDF $N(0, \sigma_{\delta u_S})$ from Eq. (A30), while in FP9 it deviates, showing a lower peak and fat tails. The reason of that is explained in more detail for the SUP model in the next paragraph, but in summary it is due to the capacity in the FP9 of the tidal and Ekman modelization to resolve large part of the variability. For both the FPs, although the first and second moments of the velocities are comparable to the SUP model (Fig. 3) and the shape of the PDF of the velocity increments has improved as compared to the results of the DET model (Fig. 5), the tails of the GAU velocity increment PDFs are (by construction) not fat enough to resemble the superstatistical Exp-Lin PDF and the occurrence of extreme events is strongly under estimated.

Regarding the SUP model, in FP5 the SUP $\delta u_E + \delta u_M$ PDF is very peaked, due to the 12 h forcing time scale that suppresses a large part of the variability. Hence, the SUP $\delta u_E + \delta u_M$ PDF contributes little to the SUP δu_o PDF, that is distributed similar to the SUP δu_S Exp-Lin PDF, having fat tails (compare Fig. 2 and Fig. 5). In FP9 the SUP $\delta u_E + \delta u_M$ PDF is distributed closely to the DET δu_o PDF (larger with respect to the FP5 case), meaning that a large part of the SUP δu_o variability is resolved by the deterministic model. The SUP δu_S PDF is given by the Bessel function of Eq. (7) and shown in Fig. 2. The SUP δu_o PDF, shown in Fig. 5, is a convolution of the SUP $\delta u_E + \delta u_M$ PDF (no analytical form available) and the Bessel function. This PDF shows a lower and shifted peak with respect to the observed and superstatistical Exp-Lin PDFs (probably due to the SUP $\delta u_E + \delta u_M$ PDF), but its tails slightly deviate from the observed and Exp-Lin ones.

The percentages of the observed HFR velocity increments inside the SUP $\sigma_{\delta u_o}$ (and $\sigma_{\delta v_o}$) ensemble standard deviation band are measured from the model results:

$$\text{FP5} \begin{cases} \Pr(\delta u_o - \sigma_{\delta u_o} \leq \delta u_{\text{HFR}} \leq \delta u_o + \sigma_{\delta u_o})_{\text{SUP}}^{\text{FP5}} \simeq 73\% \\ \Pr(\delta v_o - \sigma_{\delta v_o} \leq \delta v_{\text{HFR}} \leq \delta v_o + \sigma_{\delta v_o})_{\text{SUP}}^{\text{FP5}} \simeq 73\% \end{cases} \quad (12)$$

$$\text{FP9} \begin{cases} \Pr(\delta u_o - \sigma_{\delta u_o} \leq \delta u_{\text{HFR}} \leq \delta u_o + \sigma_{\delta u_o})_{\text{SUP}}^{\text{FP9}} \simeq 61\% \\ \Pr(\delta v_o - \sigma_{\delta v_o} \leq \delta v_{\text{HFR}} \leq \delta v_o + \sigma_{\delta v_o})_{\text{SUP}}^{\text{FP9}} \simeq 62\% \end{cases} \quad (13)$$

Repeating the calculation in Eq. (12) and Eq. (13) without the eddy depletion term, we obtain a difference of less than 1 % with respect to the shown case, in which the relative velocity between the sea and the atmosphere is considered. For FP5 the percentages compare well to the analytical superstatistical Exp-Lin values:

$$\Pr(-\sigma \leq x \leq \sigma) = \int_{-\sigma}^{+\sigma} \frac{e^{-2|x|/\sigma} \left(\frac{2|x|}{\sigma} + 1 \right)}{2\sigma} dx = \int_0^{+\sigma} \frac{e^{-2x/\sigma} \left(\frac{2x}{\sigma} + 1 \right)}{\sigma} dx = \frac{1}{2} \int_0^2 e^{-z} (z + 1) dz \simeq 73\% \quad (14)$$

This fact quantitatively confirms that the SUP model in FP5 needs $\nu = 2$ dof in the stochastic signal, producing a reliable PDF with respect to the observations. In this respect, the observations are therefore indistinguishable from a single walker of the SUP model ensemble. The percentages of the FP9 case, that has $\nu = 1/2$ dof in the stochastic signal, show small discrepancies with respect to the analytical ones. Despite the deviations from the analytical superstatistical Exp-Lin PDF, the SUP model in FP9 reproduces the PDF tails of the velocity increments with even greater fidelity to the observations than in the FP5 case (Fig. 5).




Figure 6. PDFs of the observed u_{HFR} (black crosses) and numerical models sea surface currents from FP5 (left column) and FP9 (right column) in lin-lin plots (top row) and lin-log plots (bottom row): DET u_o (yellow line), GAU u_o (blue line), SUP u_o (red line).

We move now our attention from the PDFs of the velocity increments to the PDFs of the velocities (Fig. 6). The DET model is not able to represent the observed PDFs, with a FP5 clearly missing the most of the variability and a FP9 (full forcing) improving the situation, but still not reproducing the fat tails. A stochastic part is needed to represent unresolved processes and to increase the variability. The GAU and SUP models can reproduce better the observations. The GAU model shows a good pattern around the peak, while it fails in the reproduction of the extreme events, with errors of around one and three orders of magnitude for the occurrence of extreme velocities of 0.5 m s^{-1} in the FP5 and FP9, respectively. The SUP model in FP5 confirms, also with this variable, to be able to catch the observed statistics. In FP9 the SUP model shows a higher peak, but demonstrates to reproduce some details that FP5 can not, as the PDF decay with two different slopes.




Figure 7. Temporal auto-correlation function of the u -component of the HFR sea surface velocities (black asterisks) and of the models variables from FP5 (left) and FP9 (right): auto-correlation of the DET u_o (continuous yellow line) and of the SUP u_o (continuous red line). The components of the DET u_o auto-correlation are reported: the auto-correlation of the DET u_M (dotted yellow line), of the DET u_E (dashed yellow line), and the cross-correlation of the DET u_M and u_E (yellow dotted line with dots), where the normalization is based on the DET u_o auto-correlation.

Regarding the deterministic DET model u_o temporal auto-correlation functions shown in Fig. 7, the periodicity of the peaks are consistent with what is observed from the HFR data, while the amplitude of the peaks is higher with respect to the observations. This fact is mainly due to the high auto-correlation of the tidal signal and the absence of stochastic variability.

360 The cross-correlation between the DET u_M and the DET u_E velocities shows a clear undamped periodicity of 1 day. This is due to the daily update of the forecast wind data set: it introduces a spurious artificial periodicity that perfectly correlates with the daily $S1$ -component tidal variability. The observed temporal decay (with correlation time scale of approximately 5 h) is well reproduced by the DET model u_o auto-correlation from FP9, while it is overestimated by FP5. This can be easily explained: FP5 has a 12 h forcing time scale that brings spurious memory to the system.

365 The time decay and the modulation of the SUP u_o and v_o temporal auto-correlation functions are well reproduced in both simulations. The fact that the SUP initial decay is comparable with the HFR pattern shows that the value of the γ_u coefficient shown in Table 4 is well chosen. The amplitudes in FP5 underestimate the observed ones, while in FP9 the daily peaks are present. In both the simulations the SUP auto-correlation is more damped in time with respect to the DET one, due to the introduction of the stochasticity, as expected. The long time variability of the correlation functions of the idealized models is

370 not expected to be similar to the observed ones, since the model space domain is zero-dimensional and the equations do not include the dynamics of eddies and other coherent two-dimensional structures.




Figure 8. PDFs of the HFR (black crosses) and SUP model total velocity increments δu (left column) and sea surface velocities u (right column) in lin-lin plots (top row) and log-lin plots (bottom row) from: FP2 (blue line), FP5 (orange line) and FP8 (yellow line). The analytical superstatistical PDF from Eq. (6) is also reported (black line).

We have seen that moving from longer forcing time averaging FP5 to no forcing averaging FP9, the stochastic model needs less dof. In particular, the number of stochastic dof must decrease when the unaveraged tidal components are considered, as reported in Table 2. This can be seen numerically in Fig. 8, where all the PDFs are able to fit appropriately the observations.

375 The rest of the FPs presented in Table 2 (FP3, FP4, FP6 and FP7) are not shown, because of the following reasons. FP4 and FP7 do not take into account any wind forcing, so they are not able to mimic the observed velocities. FP3 and FP6, looking at the PDFs of SUP δu_o and u_o , give very similar results to FP5, where the main differences are in the height of the PDFs peaks.




Figure 9. Ekman angle as a function of the daily WRF wind speed using the HFR (sticks with errorabars) and the models sea surface currents from FP5 (left) and FP9 (right): DET model Ekman angle (yellow line) and SUP model ensemble mean Ekman angle (black line with coloured dots, the colours indicate the present wind regime). The grey area indicates the SUP model ensemble standard deviation for the Ekman angle. The errors are calculated as shown in Appendix D. The horizontal black line is the observed HFR Ekman angle average.

In order to check if the models capture the observed veering angle, the angle between the daily averaged wind and the daily averaged selected sea surface current is computed (a positive Ekman angle means that the sea surface current is on the right with respect to the wind). As seen in Fig. 9, the observed HFR Ekman angle shows a large spread for low wind regimes, while it accumulates around the observed mean value with stronger wind speeds. In both simulations the Ekman angle obtained from the DET model (yellow line) collapses towards the same angle range for high wind speeds but with less variability with respect to the observations. The SUP mean Ekman angle is almost the same as for the DET model and converges towards the observed mean Ekman angle with increasing wind speed. The SUP modelled standard deviation (grey area) reflects the observed variability on the entire domain of the Ekman angle for low wind speeds and shows a decrease of variability for stronger wind speeds. The SUP modelled Ekman angle includes, most of the times, the observed Ekman angles inside the one standard deviation band. FP5 shows, both for the DET and SUP model, higher variability bands with respect to FP9 due to the higher number of stochastic dof.

4.2 The FRR and the SUP model

390 In this section the FRR is tested on the SUP model under the forcings of FP9. In Fig. 10 the perturbation methodology on the time series, described in Sect. 3.2, and the effects on the total sea surface velocity are visualized. In this figure, it is the u_S velocity component to be perturbed, while the independent v_S component remains unperturbed. The total SUP u_o velocity component is affected by a clearly visible perturbation that decays in time, while the total SUP v_o velocity component is slightly perturbed through the Ekman drag term.




Figure 10. SUP model FP9 stochastic velocities u_S (left); HFR (crosses) and SUP FP9 total sea surface velocities u_o (right): u -component in blue, v -component in red. The SUP model time series are from the first walker in the: unperturbed system (dashed lines) and perturbed system (continuous lines with circles).

395 In Fig. 11 the diagonal response functions $R_{uu}(t)$ and $R_{vv}(t)$ with the correlations of the observed, unperturbed Ekman, stochastic and total velocities and analytical functions discussed further are reported. The diagonal response functions show an exponential decay with time scale corresponding to the stochastic velocity drag coefficient γ_u (i.e. $1/\gamma_u \simeq 140$ min, see exponential fit in Fig. 11). When it is u_E or v_E that is perturbed, the diagonal response functions show an exponential decay with time scale corresponding to the mean Ekman drag term $\frac{C_B \bar{u}}{h}$ (i.e. $\frac{\bar{h}}{C_B \bar{u}} \simeq 65$ min, not shown). This reveals that any external
400 perturbation is decaying due to the corresponding model drag (γ_u if we perturb the stochastic velocity, $\frac{C_B \bar{u}}{h}$ if we perturb the Ekman velocity).




Figure 11. Diagonal response functions (thick dotted lines) and temporal autocorrelations in lin-lin plots (left) and lin-log plots (right) of HFR (asterisks) and unperturbed SUP model FP9 velocities: u_E (dashed and dotted line), u_S (thick dashed line), u_o (continuous line). The blue indicates the u -component, while the red the v -component. The analytical autocorrelation function of the stochastic velocity from Wirth (2019) is reported with the black continuous line. The exponential decay with the stochastic drag coefficient γ_u is shown with the black thin dotted line.

As expected, since u_o and v_o are not Gaussian variables and include terms that are not affected by the perturbation, as the tidal signal, the diagonal response functions' behaviour is different from the corresponding auto-correlations. It is then not possible to describe the response functions as linear combinations of the correlations in the unperturbed system.

405 What is particularly interesting to observe in Fig. 11 is that the SUP u_S and v_S auto-correlations coincide with the analytical auto-correlation function for a stochastic Gaussian variable, defined through a linear SDE with coloured Gaussian noise, found by Wirth (2019) in his Appendix C2:

$$C_{u_S}(t) = \frac{\gamma_u e^{-\gamma_x t} - \gamma_x e^{-\gamma_u t}}{\gamma_u - \gamma_x}, \quad (15)$$

showing a dependence on the drag coefficients only and not on the variance of the coloured Gaussian. This result seems
 410 surprising, since our stochastic velocities are obtained through a linear SDE with Bessel, and not Gaussian, coloured noise. Our interpretation of the fact is the following: the stochastic velocity drag time scale $1/\gamma_u \simeq 140$ min is one order of magnitude

smaller than the superstatistical gaussianity time scale $T \simeq 2$ days of the superstatistical analysis. This means that, under the correlation point of view, the stochastic velocity does not have the time to develop the non-gaussian characteristics, but operates at an almost constant variance. As a consequence, after the decay time of about 5 h, the SUP u_S and v_S auto-correlations
415 collapse exponentially with the same γ_u coefficient of the diagonal response functions. This can be checked also analytically, through Eq. (15) and knowing that $\gamma_u < \gamma_x$:

$$C_{u_S}(t) = \frac{\gamma_x}{\gamma_u - \gamma_x} \left(\frac{\gamma_u}{\gamma_x} e^{-\gamma_x t} - e^{-\gamma_u t} \right) \xrightarrow{t \rightarrow \infty} \frac{\gamma_x}{\gamma_x - \gamma_u} e^{-\gamma_u t} \propto R_{uu}(t) \quad (16)$$

So, knowing the auto-correlation function of the stochastic velocity, it is possible to obtain the response function of the total velocity. It is then possible to state that the FRR holds in the SUP model when the perturbation is applied to the stochastic
420 signal.

4.3 Predictability evaluation results

In Fig. 12 the observation-based perturbation methodology on the time series, described in Sect. 3.3, and the effects on the total sea surface velocities are visualized. In this figure, it is the u_S and v_S velocity components to be perturbed. Both the total SUP u_o and v_o show the observation-based perturbation subject to an evident decay in time.




Figure 12. SUP model FP9 stochastic velocities (left); HFR (crosses) and SUP FP9 total sea surface velocities (right): u -component in blue, v -component in red. The SUP model time series are from the first walker in the unperturbed system (dashed lines) and perturbed system using a data-assimilation method (continuous lines with circles).




Figure 13. Temporal autocorrelations in lin-lin (left) and lin-log plots (right) of the HFR (asterisks) and SUP model stochastic (thick dashed lines) and total (continuous lines) u (blue) and v (red) sea surface velocities with the ξ (purple thick continuous line) and ϵ (green thick continuous line) functions. The exponential fits of the ξ function are shown (purple thin continuous lines). The exponential decay with the stochastic drag coefficient γ_u is shown with the black thin dotted line.

425 The $\xi(t)$ and $\epsilon(t)$ functions are reported in Fig. 13 and represent the normalized distance between the perturbed-unperturbed and the perturbed-observed systems, respectively. The function $\xi(t)$ can be considered as a generalization of the response function in the FRR when both the components are perturbed of a quantity not known a priori. It is possible to see that the $\xi(t)$ function has an exponential decay with time scale $1/\gamma_u \simeq 140$ min. When it is the Ekman signal to be perturbed, we obtain two exponential decays: $\tau_1 \simeq 75$ min for $t < 2$ h and $\tau_2 \simeq 140$ min for $t > 4$ h (not shown). The external perturbation, due to
430 the observation-based methodology, decays with the stochastic drag coefficient when we perturb the stochastic velocity, while a clear correspondence with the drag coefficient is not found when the Ekman signal is perturbed. The $\epsilon(t)$ function increases its value in time from zero and, after about 5 hours, saturates to one, showing that after that time the perturbed system has completely lost the memory of the initialization to the observations. This time of 5 hours coincides with the auto-correlation time scale of the SUP stochastic and total velocities.

In the Gulf of Trieste the analytical superstatistical PDF of the observed HFR sea surface current increments is known (Flora et al., 2023) and is here called Exp-Lin PDF. The Exp-Lin PDF is non-Gaussian with fat-tails (the extreme events occur more often with respect to a Gaussian statistics), it provides all the statistical moments but does not give any predictive information. In this study we have developed a hierarchy of idealized models whose aim is to mimic the sea surface current time series, characterized by the observed statistics.

The hierarchy is organized as follows: the DET model is purely deterministic and includes tidal and Ekman signals, the GAU model adds a coloured-in-time Gaussian stochastic velocity, while the SUP model adds a stochastic sea surface velocity with superstatistical fat-tailed increments that simulate the unresolved dynamics. A variable is superstatistical if it is locally (in time) Gaussian with a variance evolving over a longer time scale. In our case the variance is obtained through the sum of 2ν squared (Gaussian) Ornstein-Uhlenbeck processes and results to be Gamma distributed. The parameter ν identifies the dof of the stochastic system. The models are then tested imposing nine FPs combining different types of tidal and wind forcings: complete, 12 h moving-averaged time series or omitted forcing (Table 2). The stochastic models are adjusted for each FP in order to best fit the observed PDF.

In the following we point out the general differences between the models' results. The DET model is able to simulate the slow variability of the observed HFR currents, reaching the 55-56% of correlation coefficient (Fig. 4). The GAU and SUP stochastic models increase the variability to observed values, but, since their ensemble means almost coincide with the DET model results, they show little influence on the slow dynamics (Fig. 3).

The main differences between the models are evident looking at the PDFs of their variables. Regarding the total velocity increment PDFs (Fig. 5), the DET model does not explain the variability of the observed superstatistical Exp-Lin PDF, which results in a stark underestimation of the variability. The GAU model improves the results, but without the observed fat tails, that is the occurrence of extreme events is underestimated. When we consider the SUP model, in order to fit the observed statistics, the stochastic modelization requires $\nu = 2$ stochastic dof when the tidal forcing is omitted or averaged, while it needs $\nu = 1/2$ stochastic dof when the complete tidal signal is considered (Table 2). This result affects the shape of the SUP stochastic velocity increment PDF (Fig. 2) that is Exp-Lin when $\nu = 2$ and Bessel when $\nu = 1/2$. The SUP model shows a total velocity increment PDF that is representative of observations (Fig. 5): it is almost Exp-Lin when the FP considers averaged forcings and $\nu = 2$ stochastic dof, while it has a shifted and slightly lower peak but tails that satisfactorily fit the observed data when the FP considers the complete forcings and $\nu = 1/2$ stochastic dof. The convergence to the PDF and, in particular, the enlargement of the PDF tails to the observed ones in the progression from the DET model to the GAU and then to the SUP model is also visible in the PDFs of the velocities (Fig. 6). In addition, we obtain that the FP with complete forcings (FP9) and, consequently, with $\nu = 1/2$ stochastic dof, enables the SUP model to catch the tails with a double slope in the velocity PDF, as seen from the observations.

The velocities temporal auto-correlation functions under the FP9 (Fig. 7) reveals that the SUP model, after a realistic time

decay of around 5 h (driven by the deterministic signal), allows for a better representation, with respect to the DET model,
470 of the temporal modulations. The Ekman dynamics analysis (Fig. 9) indicates that strong wind increases the likelihood of the
SUP model generating trajectories that match the observed average Ekman state. This pattern aligns with the HFR Ekman
angle data, where lower wind forcing results in more variability, while higher wind forcing brings values closer to the average.

The SUP model under the most complete forcings and, consequently, with $\nu = 1/2$ dof (FP9) is then explored applying exter-
475 nal perturbations to the stochastic signal. The FRR holds when the stochastic signal is perturbed, showing that the response
to an external perturbation can be obtained by considering the fluctuations of the unperturbed system. In particular, the SUP
model response function decays with the stochastic velocity drag time scale ($1/\gamma_u$, see Fig. 11 and Eq. (16)), indicating a clear
memory time scale for external stochastic perturbations.

The predictability evaluation through the observation-based perturbation shows that the SUP model, given any external correc-
480 tion to the HFR values, reaches an equilibrium distance from the observations, converges to the unperturbed realization, after
5 hours (Fig. 13). This time value is consistent with the auto-correlation decay of the observed and SUP stochastic and total
velocities.

From the predictive point of view, we should remark that our modelling is extremely idealised: it is local in space and does
485 not take into account other external forcings besides wind and tides, such as river discharge. The modelling can be further
developed including the Isonzo/Soča river influence, other air-sea interaction processes as wave generation, moving the space
domain from one point (0 dimension) to 2 dimensions including advection, eddy dynamics and also incorporating the broader
Adriatic circulation.

Despite its idealization, the SUP model presented in this study can reproduce part of the deterministic (tide-wind-forced) large
490 scale dynamics and the observed fat tails of the HFR velocity increments PDFs and is eventually able to simulate extreme
events. One of the most interesting results is that when we have a more detailed deterministic signal (given by the complete
forcings), a larger part of the dynamics is resolved and the stochastic part of the model must not only have less variability, it
must also decrease its dof. The SUP model can therefore be taken as an example for the modelling community to reflect on
how stochasticity can be used to reproduce extreme events and how its characteristics vary depending on the coarse graining
495 of the forcings. Various applications can be found to the SUP model, for example it is currently used to explore the role of
extreme events in the kinetic energy fluxes between the atmosphere and the sea. Additionally, evaluating the air-sea interaction
bulk formulas in CMEMS MFS model (or another full physics numerical model adopted for the Gulf of Trieste) in light of our
observations, and therefore of the SUP model findings, would be extremely interesting.

Code and data availability. The exact version of the model used to produce the results used in this paper is archived on Zenodo (Flora
500 et al., 2024), as are input data and scripts to run the model for all the simulations presented in this paper. In particular: `twd_model.f90`
is the code implementing the DET model, `twg_model.f90` is the code implementing the GAU model, `tws_model.f90` is the code

implementing the STO model with $\nu = 2$ and `twb_model.f90` is the code implementing the STO model with $\nu = 1/2$. Additionally, `twb_FRRD_model.f90`, `twb_FRRS_model.f90`, `twb_datassD_model.f90` and `twb_datassS_model.f90` are the codes implementing the STO model with $\nu = 1/2$ in the FRR and observation-based perturbation applications (applied to the deterministic or stochastic signals).

The HFR sea surface current data of the Gulf of Trieste are freely available from the European HFRadar node (OGS et al., 2023). The WRF forecasted wind field is obtainable upon request from ARPA FVG (<https://www.arpa.fvg.it>, last access: 29 October 2024, CRMA).

Appendix A: An analytical idealized case

This is the idealized case of the stochastic system described in Sect. 3, Eq. (2) and Eq. (3) in which:

$$\sqrt{\Sigma_{i=1}^{2\nu} \alpha_i^2} = \sqrt{Q} \simeq \text{constant} \quad (A1)$$

This approximation and the following analysis permits to find the analytical distribution of the variables x , u_S and δu_S . Considering the u -components, independent from the v -components, Eq. (2) and Eq. (3) can be written as follows:

$$\begin{cases} dx = -\gamma_x x dt + \sqrt{Q} dW_x \\ du_S = -\gamma_u u_S dt + \eta x dt \end{cases}$$

$$d \begin{pmatrix} x \\ u_S \end{pmatrix} = \begin{pmatrix} -\gamma_x & 0 \\ \eta & -\gamma_u \end{pmatrix} \begin{pmatrix} x \\ u_S \end{pmatrix} dt + \begin{pmatrix} \sqrt{Q} \\ 0 \end{pmatrix} dW_x \quad (A2)$$

$$= M \begin{pmatrix} x \\ u_S \end{pmatrix} dt + \begin{pmatrix} \sqrt{Q} \\ 0 \end{pmatrix} dW_x$$

where $M = \begin{pmatrix} -\gamma_x & 0 \\ \eta & -\gamma_u \end{pmatrix}$ can be diagonalized:

$$\det(M - mI) = \det \begin{pmatrix} -\gamma_x - m & 0 \\ \eta & -\gamma_u - m \end{pmatrix} = (\gamma_x + m)(\gamma_u + m) = 0 \quad (A3)$$

obtaining two eigenvalues $m_1 = -\gamma_x$ and $m_2 = -\gamma_u$ and the following eigenvectors (with $\gamma_x \neq \gamma_u$):

$$M\mathbf{e}_1 = m_1\mathbf{e}_1$$

$$\begin{pmatrix} -\gamma_x & 0 \\ \eta & -\gamma_u \end{pmatrix} \begin{pmatrix} e_{1x} \\ e_{1y} \end{pmatrix} = -\gamma_x \begin{pmatrix} e_{1x} \\ e_{1y} \end{pmatrix}$$

$$\begin{cases} -\gamma_x e_{1x} = -\gamma_x e_{1x} \\ \eta e_{1x} - \gamma_u e_{1y} = -\gamma_x e_{1y} \end{cases}$$

$$\begin{cases} e_{1y} = -\frac{\eta}{\gamma_x - \gamma_u} e_{1x} \\ e_{1x}^2 + e_{1y}^2 = 1 \end{cases}$$

$$e_{1x}^2 \left(1 + \frac{\eta^2}{(\gamma_x - \gamma_u)^2} \right) = 1$$

$$\begin{cases} e_{1x} = -\frac{\gamma_x - \gamma_u}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} \\ e_{1y} = \frac{\eta}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} \end{cases}$$

$$\mathbf{e}_1 = \begin{pmatrix} -\frac{\gamma_x - \gamma_u}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} \\ \frac{\eta}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} \end{pmatrix}$$

$$M\mathbf{e}_2 = m_2\mathbf{e}_2$$

$$\begin{pmatrix} -\gamma_x & 0 \\ \eta & -\gamma_u \end{pmatrix} \begin{pmatrix} e_{2x} \\ e_{2y} \end{pmatrix} = -\gamma_u \begin{pmatrix} e_{2x} \\ e_{2y} \end{pmatrix}$$

$$\begin{cases} -\gamma_x e_{2x} = -\gamma_u e_{2x} \\ \eta e_{2x} - \gamma_u e_{2y} = -\gamma_u e_{2y} \end{cases}$$

$$\begin{cases} e_{2x} = 0 \\ e_{2y} = 1 \end{cases}$$

$$\mathbf{e}_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

(A4)

For $\gamma_x \neq \gamma_u$, the system in Eq. (A2) is diagonalizable and it is possible to write $M = T \cdot D \cdot T^{-1}$ where:

$$T = \begin{pmatrix} -\frac{\gamma_x - \gamma_u}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} & 0 \\ \frac{\eta}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} & 1 \end{pmatrix} \quad D = \begin{pmatrix} -\gamma_x & 0 \\ 0 & -\gamma_u \end{pmatrix} \quad T^{-1} = \begin{pmatrix} -\frac{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{\gamma_x - \gamma_u} & 0 \\ \frac{\eta}{\gamma_x - \gamma_u} & 1 \end{pmatrix} \quad (A5)$$

520 and so, hypothesizing $\gamma_u < \gamma_x$ (the final distributions do not change with $\gamma_x < \gamma_u$, what is changing is just the sign of the Wiener processes in the diagonalised system):

$$d \begin{pmatrix} x \\ u_S \end{pmatrix} = T \cdot D \cdot T^{-1} \begin{pmatrix} x \\ u_S \end{pmatrix} dt + \begin{pmatrix} \sqrt{Q} \\ 0 \end{pmatrix} dW_x \quad (A6)$$

$$d \left[T^{-1} \begin{pmatrix} x \\ u_S \end{pmatrix} \right] = D \left[T^{-1} \begin{pmatrix} x \\ u_S \end{pmatrix} \right] dt + T^{-1} \begin{pmatrix} \sqrt{Q} \\ 0 \end{pmatrix} dW_x \quad (A7)$$

525

$$\begin{pmatrix} \tilde{x} \\ \tilde{u}_S \end{pmatrix} = T^{-1} \begin{pmatrix} x \\ u_S \end{pmatrix} = \begin{pmatrix} -\frac{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{\gamma_x - \gamma_u} x \\ u_S + \frac{\eta}{\gamma_x - \gamma_u} x \end{pmatrix}; \quad \begin{pmatrix} -\sqrt{Q_x} \\ \sqrt{Q_u} \end{pmatrix} = T^{-1} \begin{pmatrix} \sqrt{Q} \\ 0 \end{pmatrix} = \begin{pmatrix} -\frac{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{\gamma_x - \gamma_u} \sqrt{Q} \\ \frac{\eta}{\gamma_x - \gamma_u} \sqrt{Q} \end{pmatrix} \quad (A8)$$

obtaining:

$$d \begin{pmatrix} \tilde{x} \\ \tilde{u}_S \end{pmatrix} = D \begin{pmatrix} \tilde{x} \\ \tilde{u}_S \end{pmatrix} dt + \begin{pmatrix} -\sqrt{Q_x} \\ \sqrt{Q_u} \end{pmatrix} dW_x$$

$$\begin{cases} d\tilde{x} = -\gamma_x \tilde{x} dt - \sqrt{Q_x} dW_x \\ d\tilde{u}_S = -\gamma_u \tilde{u}_S dt + \sqrt{Q_u} dW_x \end{cases} \quad (\text{A9})$$

The approximation for which $\sqrt{Q} = \text{constant}$ leads to $\sqrt{Q_x}$ and $\sqrt{Q_u}$ also constants. This fact leads to the Gaussianity of the variables $\tilde{x}$ and $\tilde{u}_S$ (they are Ornstein-Uhlenbeck processes):

$$\begin{cases} \tilde{x} \sim N(0, \sigma_{\tilde{x}}^2) = N\left(0, \frac{Q_x}{2\gamma_x}\right) = N\left(0, \frac{\eta^2 + (\gamma_x - \gamma_u)^2}{(\gamma_x - \gamma_u)^2} \frac{Q}{2\gamma_x}\right) \\ \tilde{u}_S \sim N(0, \sigma_{\tilde{u}_S}^2) = N\left(0, \frac{Q_u}{2\gamma_u}\right) = N\left(0, \frac{\eta^2}{(\gamma_x - \gamma_u)^2} \frac{Q}{2\gamma_u}\right) \end{cases} \quad (\text{A10})$$

From Eq. (A8) it is possible to write x and u_S in terms of $\tilde{x}$ and $\tilde{u}_S$, resulting as linear combinations of Gaussian variables:

$$\begin{cases} x = -\frac{\gamma_x - \gamma_u}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} \tilde{x} \\ u_S = \tilde{u}_S + \tilde{r} \\ \tilde{r} = -\frac{\eta}{\gamma_x - \gamma_u} x = \frac{\eta}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} \tilde{x} \end{cases} \quad (\text{A11})$$

It leads to the Gaussianity of the variables x , $\tilde{r}$ and u_S with the following variances:

$$\begin{cases} x \sim N(0, \sigma_x^2) \\ \tilde{r} \sim N(0, \sigma_{\tilde{r}}^2) \\ u_S \sim N(0, \sigma_{u_S}^2) \end{cases} \quad (\text{A12})$$

$$\begin{cases} \sigma_x^2 = \frac{(\gamma_x - \gamma_u)^2}{\eta^2 + (\gamma_x - \gamma_u)^2} \sigma_{\tilde{x}}^2 = \frac{Q}{2\gamma_x} \\ \sigma_{\tilde{r}}^2 = \frac{\eta^2}{\eta^2 + (\gamma_x - \gamma_u)^2} \sigma_{\tilde{x}}^2 = \frac{\eta^2}{(\gamma_x - \gamma_u)^2} \frac{Q}{2\gamma_x} \\ \sigma_{u_S}^2 = \sigma_{\tilde{u}_S}^2 + \sigma_{\tilde{r}}^2 + 2 \lim_{t \rightarrow \infty} \text{corr}(\tilde{u}_S(t), \tilde{r}(t)) \\ = \frac{\eta^2}{(\gamma_x - \gamma_u)^2} \frac{Q}{2\gamma_u} + \frac{\eta^2}{(\gamma_x - \gamma_u)^2} \frac{Q}{2\gamma_x} + 2 \frac{\eta}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} \lim_{t \rightarrow \infty} \text{corr}(\tilde{u}_S(t), \tilde{x}(t)) \\ = \frac{\eta^2 (\gamma_x + \gamma_u)}{(\gamma_x - \gamma_u)^2} \frac{Q}{2\gamma_x \gamma_u} - \frac{2\eta^2}{(\gamma_x - \gamma_u)^2 (\gamma_x + \gamma_u)} Q \\ = \frac{\eta^2}{(\gamma_x - \gamma_u)^2} \frac{(\gamma_x + \gamma_u)^2 - 4\gamma_x \gamma_u}{2\gamma_x \gamma_u (\gamma_x + \gamma_u)} Q \\ = \frac{\eta^2}{2\gamma_x \gamma_u (\gamma_x + \gamma_u)} Q \end{cases} \quad (\text{A13})$$

where, since $\tilde{x}$ and $\tilde{u}_S$ are Ornstein-Uhlenbeck processes:

$$\begin{cases} \tilde{x}(t) = -e^{-\gamma_x t} \int_0^t e^{\gamma_x t'} \sqrt{Q_x} F(t') dt' \\ \tilde{u}_S(t) = e^{-\gamma_u t} \int_0^t e^{\gamma_u t'} \sqrt{Q_u} F(t') dt' \\ E[F(t') F(t'')] = \delta(t' - t'') \end{cases} \quad (\text{A14})$$

540 we have that:

$$\begin{aligned}
\text{corr}(\tilde{u}_S(t), \tilde{x}(t)) &= -e^{-(\gamma_x + \gamma_u)t} \sqrt{Q_x Q_u} \int_0^t \int_0^t e^{\gamma_x t'} e^{\gamma_u t''} E[F(t') F(t'')] dt' dt'' \\
&= -\frac{\eta \sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{(\gamma_x - \gamma_u)^2} Q e^{-(\gamma_x + \gamma_u)t} \frac{e^{(\gamma_x + \gamma_u)t} - 1}{\gamma_x + \gamma_u} \\
&= -\frac{\eta \sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{(\gamma_x - \gamma_u)^2} \frac{Q}{\gamma_x + \gamma_u} \left(1 - e^{-(\gamma_x + \gamma_u)t}\right) \\
&\xrightarrow{t \rightarrow \infty} -\frac{\eta \sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{(\gamma_x - \gamma_u)^2} \frac{Q}{\gamma_x + \gamma_u}
\end{aligned} \tag{A15}$$

Now, it is possible to express the δu_S variable, starting from its definition and using Eq. (A13), in terms of the diagonalized variables $\tilde{x}$ and $\tilde{u}_S$:

$$\begin{aligned}
\delta u_S(t) &= u_S(t + \delta) - u_S(t) \\
&= \tilde{u}_S(t + \delta) + \frac{\eta}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} \tilde{x}(t + \delta) - \tilde{u}_S(t) - \frac{\eta}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} \tilde{x}(t) \\
&= [\tilde{u}_S(t + \delta) - \tilde{u}_S(t)] + \frac{\eta}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} [\tilde{x}(t + \delta) - \tilde{x}(t)] \\
&= \delta \tilde{u}_S(t) + \frac{\eta}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} \delta \tilde{x}(t)
\end{aligned} \tag{A16}$$

545 where:

$$\begin{cases} \delta \tilde{u}_S(t) = \tilde{u}_S(t + \delta) - \tilde{u}_S(t) \\ \delta \tilde{x}(t) = \tilde{x}(t + \delta) - \tilde{x}(t) \end{cases} \tag{A17}$$

Since $\delta \tilde{u}_S$ and $\delta \tilde{x}$ are linear combinations of Gaussian variables, they are Gaussian variables themselves:

$$\begin{cases} \delta \tilde{u}_S \sim N(0, \sigma_{\delta \tilde{u}_S}^2) \\ \sigma_{\delta \tilde{u}_S}^2 = 2\sigma_{\tilde{u}_S}^2 + 2 \lim_{t \rightarrow \infty} \text{corr}(\tilde{u}_S(t + \delta), \tilde{u}_S(t)) \\ \quad = 2 \frac{\eta^2}{(\gamma_x - \gamma_u)^2} \frac{Q}{2\gamma_u} (1 - e^{-\gamma_u \delta}) \\ \delta \tilde{x} \sim N(0, \sigma_{\delta \tilde{x}}^2) \\ \sigma_{\delta \tilde{x}}^2 = 2\sigma_{\tilde{x}}^2 + 2 \lim_{t \rightarrow \infty} \text{corr}(\tilde{x}(t + \delta), \tilde{x}(t)) \\ \quad = 2 \frac{\eta^2 + (\gamma_x - \gamma_u)^2}{(\gamma_x - \gamma_u)^2} \frac{Q}{2\gamma_x} (1 - e^{-\gamma_x \delta}) \end{cases} \tag{A18}$$

where:

$$\begin{aligned}
\text{corr}(\tilde{u}_S(t+\delta), \tilde{u}_S(t)) &= \frac{\eta^2 Q}{(\gamma_x - \gamma_u)^2} e^{-2\gamma_u t} e^{-\gamma_u \delta} \int_0^{t+\delta} \int_0^t e^{\gamma_u t'} e^{\gamma_u t''} \delta(t' - t'') dt' dt'' \\
&= \frac{\eta^2 Q}{(\gamma_x - \gamma_u)^2} e^{-2\gamma_u t} e^{-\gamma_u \delta} \int_0^t e^{2\gamma_u t'} dt' \\
&= \frac{\eta^2 Q}{(\gamma_x - \gamma_u)^2} e^{-\gamma_u \delta} e^{-2\gamma_u t} \frac{e^{2\gamma_u t} - 1}{2\gamma_u} \\
&= \frac{\eta^2 Q}{(\gamma_x - \gamma_u)^2} \frac{e^{-\gamma_u \delta}}{2\gamma_u} (1 - e^{-2\gamma_u t}) \\
&\xrightarrow{t \rightarrow \infty} \frac{\eta^2}{(\gamma_x - \gamma_u)^2} \frac{Q}{2\gamma_u} e^{-\gamma_u \delta}
\end{aligned} \tag{A19}$$

$$\begin{aligned}
\text{corr}(\tilde{x}(t+\delta), \tilde{x}(t)) &= \frac{\eta^2 + (\gamma_x - \gamma_u)^2}{(\gamma_x - \gamma_u)^2} Q e^{-2\gamma_x t} e^{-\gamma_x \delta} \int_0^{t+\delta} \int_0^t e^{\gamma_x t'} e^{\gamma_x t''} \delta(t' - t'') dt' dt'' \\
&= \frac{\eta^2 + (\gamma_x - \gamma_u)^2}{(\gamma_x - \gamma_u)^2} Q e^{-2\gamma_x t} e^{-\gamma_x \delta} \int_0^t e^{2\gamma_x t'} dt' \\
&= \frac{\eta^2 + (\gamma_x - \gamma_u)^2}{(\gamma_x - \gamma_u)^2} Q e^{-\gamma_x \delta} e^{-2\gamma_x t} \frac{e^{2\gamma_x t} - 1}{2\gamma_x} \\
&= \frac{\eta^2 + (\gamma_x - \gamma_u)^2}{(\gamma_x - \gamma_u)^2} \frac{Q}{2\gamma_x} e^{-\gamma_x \delta} (1 - e^{-2\gamma_x t}) \\
&\xrightarrow{t \rightarrow \infty} \frac{\eta^2 + (\gamma_x - \gamma_u)^2}{(\gamma_x - \gamma_u)^2} \frac{Q}{2\gamma_x} e^{-\gamma_x \delta}
\end{aligned} \tag{A20}$$

Now, according to Eq. (A16) and Eq. (A18), since δu_S is a linear combination of Gaussian variables, it is Gaussianly distributed itself:

$$\begin{cases} \delta u_S \sim N(0, \sigma_{\delta u}^2) \\ \sigma_{\delta u}^2 = \sigma_{\delta \tilde{u}}^2 + \frac{\eta^2}{\eta^2 + (\gamma_x - \gamma_u)^2} \sigma_{\delta \tilde{x}}^2 + \frac{2\eta}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} \lim_{t \rightarrow \infty} \text{corr}(\delta \tilde{u}_S(t), \delta \tilde{x}(t)) \end{cases} \tag{A21}$$

where:

$$\begin{aligned}
\lim_{t \rightarrow \infty} \text{corr}(\delta \tilde{u}_S(t), \delta \tilde{x}(t)) &= \lim_{t \rightarrow \infty} \text{corr}(\tilde{u}_S(t+\delta) - \tilde{u}_S(t), \tilde{x}(t+\delta) - \tilde{x}(t)) \\
&= \lim_{t \rightarrow \infty} [\text{corr}(\tilde{u}_S(t+\delta), \tilde{x}(t+\delta)) - \text{corr}(\tilde{u}_S(t+\delta), \tilde{x}(t)) - \text{corr}(\tilde{u}_S(t), \tilde{x}(t+\delta)) + \text{corr}(\tilde{u}_S(t), \tilde{x}(t))]
\end{aligned} \tag{A22}$$

$$\begin{aligned}
\lim_{t \rightarrow \infty} \text{corr}(\tilde{u}_S(t + \delta), \tilde{x}(t + \delta)) &= \lim_{t \rightarrow \infty} \text{corr}(\tilde{u}_S(t), \tilde{x}(t)) \\
&= -\frac{\eta \sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{(\gamma_x - \gamma_u)^2} \frac{Q}{\gamma_x + \gamma_u}
\end{aligned} \tag{A23}$$

560

$$\begin{aligned}
\text{corr}(\tilde{u}_S(t + \delta), \tilde{x}(t)) &= -\frac{\eta \sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{(\gamma_x - \gamma_u)^2} Q e^{-\gamma_u(t + \delta)} e^{-\gamma_x t} \int_0^{t + \delta} \int_0^t e^{\gamma_u t'} e^{\gamma_x t''} \delta(t' - t'') dt' dt'' \\
&= -\frac{\eta \sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{(\gamma_x - \gamma_u)^2} Q e^{-\gamma_u \delta} e^{-(\gamma_x + \gamma_u)t} \int_0^t e^{(\gamma_x + \gamma_u)t'} dt' \\
&= -\frac{\eta \sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{(\gamma_x - \gamma_u)^2} \frac{Q}{\gamma_x + \gamma_u} e^{-\gamma_u \delta} (1 - e^{-(\gamma_x + \gamma_u)t}) \\
&\xrightarrow{t \rightarrow \infty} -\frac{\eta \sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{(\gamma_x - \gamma_u)^2} \frac{Q}{\gamma_x + \gamma_u} e^{-\gamma_u \delta}
\end{aligned} \tag{A24}$$

$$\begin{aligned}
\text{corr}(\tilde{u}_S(t), \tilde{x}(t + \delta)) &= -\frac{\eta \sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{(\gamma_x - \gamma_u)^2} Q e^{-\gamma_u t} e^{-\gamma_x(t + \delta)} \int_0^{t + \delta} \int_0^t e^{\gamma_u t'} e^{\gamma_x t''} \delta(t' - t'') dt' dt'' \\
&\xrightarrow{t \rightarrow \infty} -\frac{\eta \sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{(\gamma_x - \gamma_u)^2} \frac{Q}{\gamma_x + \gamma_u} e^{-\gamma_x \delta}
\end{aligned} \tag{A25}$$

And so, continuing Eq. (A22):

$$\lim_{t \rightarrow \infty} \text{corr}(\tilde{u}_S(t + \delta), \tilde{x}(t + \delta)) = -\frac{\eta \sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{(\gamma_x - \gamma_u)^2} \frac{Q}{\gamma_x + \gamma_u} [2 - (e^{-\gamma_x \delta} + e^{-\gamma_u \delta})] \tag{A26}$$

565

It is now possible the calculation started in Eq. (A21):

$$\begin{aligned}
\sigma_{\delta u}^2 &= \sigma_{\delta u}^2 + \frac{\eta^2}{\eta^2 + (\gamma_x - \gamma_u)^2} \sigma_{\delta x}^2 + \frac{2\eta}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} \lim_{t \rightarrow \infty} \text{corr}(\delta \tilde{u}_S(t), \delta \tilde{x}(t)) \\
&= \frac{2\eta^2}{(\gamma_x - \gamma_u)^2} \frac{Q}{2\gamma_u} (1 - e^{-\gamma_u \delta}) + \frac{\eta^2}{\eta^2 + (\gamma_x - \gamma_u)^2} \frac{2 \left[\eta^2 + (\gamma_x - \gamma_u)^2 \right]}{(\gamma_x - \gamma_u)^2} \frac{Q}{2\gamma_x} (1 - e^{-\gamma_x \delta}) + \\
&\quad + \frac{2\eta}{\sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}} (-) \frac{\eta \sqrt{\eta^2 + (\gamma_x - \gamma_u)^2}}{(\gamma_x - \gamma_u)^2} \frac{Q}{\gamma_x + \gamma_u} [2 - (e^{-\gamma_x \delta} + e^{-\gamma_u \delta})] \\
&= \frac{2\eta^2 Q}{(\gamma_x - \gamma_u)^2} \left[\frac{1}{2\gamma_u} (1 - e^{-\gamma_u \delta}) + \frac{1}{2\gamma_x} (1 - e^{-\gamma_x \delta}) - \frac{1}{\gamma_x + \gamma_u} (1 - e^{-\gamma_x \delta}) - \frac{1}{\gamma_x + \gamma_u} (1 - e^{-\gamma_u \delta}) \right] \\
&= \frac{2\eta^2 Q}{(\gamma_x - \gamma_u)^2} \left[\frac{\gamma_x - \gamma_u}{2\gamma_u (\gamma_x + \gamma_u)} (1 - e^{-\gamma_u \delta}) - \frac{\gamma_x - \gamma_u}{2\gamma_x (\gamma_x + \gamma_u)} (1 - e^{-\gamma_x \delta}) \right] \\
&= \frac{\eta^2 Q}{\gamma_x^2 - \gamma_u^2} \left(\frac{1 - e^{-\gamma_u \delta}}{\gamma_u} - \frac{1 - e^{-\gamma_x \delta}}{\gamma_x} \right)
\end{aligned} \tag{A27}$$




Figure A1. Gaussian PDFs of the stochastic variables x (in blue), u_S (in red) and δu_S (in green): the continuous lines are the analytical Gaussian PDFs in Eq. (A28), Eq. (A29) and Eq. (A30), while the points are the histograms of the simulated variables described in Eq. (A2) and Eq. (A16) in the case $Q = \text{constant}$. The used parameters are shown in Table A1, the simulation with $\eta = \eta_1$ results are shown in (a) and (b), while in (c) and (d) the results with $\eta = \eta_2$ are shown. In (a) and (c) the plots are lin-lin, while in (b) and (d) the plots are lin-log.

parameter	value
ensemble size	10^5
total integration time	~ 656 days
time step dt	1.5×10^2 s
τ	6.6×10^3 s
$\gamma_x = 1/\tau$	$1.515 \times 10^{-4} \text{ s}^{-1}$
Q	$2.6 \times 10^{-6} \text{ m}^2 \text{ s}^{-3}$
γ_u	$5.152 \times 10^{-5} \text{ s}^{-1}$
η_1	$1.333 \times 10^{-4} \text{ s}^{-1}$
$\eta_2 = \sqrt{\frac{\gamma_x^2 - \gamma_u^2}{2 \left[\frac{\gamma_x}{\gamma_u} (1 - e^{-\gamma_u \delta}) - (1 - e^{-\gamma_x \delta}) \right]}}$	$1.247 \times 10^{-4} \text{ s}^{-1}$
δ	1.44×10^4 s

Table A1. Involved parameters in the idealized analytical stochastic modelization.

In a summary, in the idealized case of the stochastic system, in which Q is constant, the variables x , u_S and δu_S are Gaussianly distributed:

$$570 \quad x \sim N(0, \sigma_x^2) = N\left(0, \frac{Q}{2\gamma_x}\right) \quad (\text{A28})$$

$$u_S \sim N(0, \sigma_{u_S}^2) = N\left(0, \frac{\eta^2 Q}{2\gamma_x \gamma_u (\gamma_x + \gamma_u)}\right) \quad (\text{A29})$$

$$\delta u_S \sim N(0, \sigma_{\delta u}^2) = N\left(0, \frac{\eta^2 Q}{\gamma_x^2 - \gamma_u^2} \left(\frac{1 - e^{-\gamma_u \delta}}{\gamma_u} - \frac{1 - e^{-\gamma_x \delta}}{\gamma_x} \right)\right) \quad (\text{A30})$$

575 It is possible to find a condition for which δu_S is distributed as x :

$$\begin{aligned} \sigma_{\delta u}^2 &= \frac{\eta^2 Q}{\gamma_x^2 - \gamma_u^2} \left(\frac{1 - e^{-\gamma_u \delta}}{\gamma_u} - \frac{1 - e^{-\gamma_x \delta}}{\gamma_x} \right) \\ &= \frac{Q}{2\gamma_x} \frac{2\eta^2}{\gamma_x^2 - \gamma_u^2} \left[\frac{\gamma_x}{\gamma_u} (1 - e^{-\gamma_u \delta}) - (1 - e^{-\gamma_x \delta}) \right] \\ &= \sigma_x^2 \frac{2\eta^2}{\gamma_x^2 - \gamma_u^2} \left[\frac{\gamma_x}{\gamma_u} (1 - e^{-\gamma_u \delta}) - (1 - e^{-\gamma_x \delta}) \right] \end{aligned} \quad (\text{A31})$$

$$\begin{aligned} \sigma_{\delta u}^2 = \sigma_x^2 &\iff \frac{2\eta^2}{\gamma_x^2 - \gamma_u^2} \left[\frac{\gamma_x}{\gamma_u} (1 - e^{-\gamma_u \delta}) - (1 - e^{-\gamma_x \delta}) \right] = 1 \\ &\iff \eta = \sqrt{\frac{\gamma_x^2 - \gamma_u^2}{2 \left[\frac{\gamma_x}{\gamma_u} (1 - e^{-\gamma_u \delta}) - (1 - e^{-\gamma_x \delta}) \right]}} \end{aligned} \quad (\text{A32})$$

All these facts are confirmed by the numerical simulations results, shown in Fig. A1 using the numerical parameters in Table A1.

Appendix B: Ornstein–Uhlenbeck processes and colored noise

Let's consider the stochastic part for the u -component (analogously for the v -component) of the SUP model:

$$d\alpha_i = -\mu\alpha_i dt + \beta_u dW_i \quad (\text{B1})$$

$$dx = -\gamma_x x dt + \sqrt{\sum_{i=1}^{2\nu} \alpha_i^2} dW_x \quad (\text{B2})$$

$$du_S = -\gamma_u u_S dt + \eta x dt \quad (\text{B3})$$

Since β and μ are constants, Eq. (B1) is a Langevin equation and the variables α_i are Ornstein–Uhlenbeck processes, so their distribution in the stationary state is Gaussian:

$$p_\alpha(\alpha_i) = \sqrt{\frac{\mu}{\pi\beta_u^2}} e^{-\mu\alpha_i^2/\beta_u^2} \quad (\text{B4})$$

Defining the new variable Q as the sum of 2ν squared identically distributed Gaussian variables $Q = \sum_{i=1}^{2\nu} \alpha_i^2$, it results as a variable distributed with a Gamma distribution with shape parameter equal to ν in the stationary state:

$$p_Q(Q) = \Gamma_{\nu, \tilde{\lambda}}(Q) = \frac{\tilde{\lambda}^\nu Q^{\nu-1} e^{-\tilde{\lambda}Q}}{\Gamma(\nu)} = \left(\frac{\mu}{\beta_u^2}\right)^\nu \frac{Q^{\nu-1}}{\Gamma(\nu)} e^{-\mu Q/\beta_u^2} \quad (\text{B5})$$

where $\tilde{\lambda} = \frac{\mu}{\beta_u^2}$ and with first moment:

$$E[Q] = \int_0^\infty \left(\frac{\mu}{\beta_u^2}\right)^\nu \frac{Q^\nu}{\Gamma(\nu)} e^{-\mu Q/\beta_u^2} dQ = \frac{\nu\beta_u^2}{\mu}. \quad (\text{B6})$$

It is possible to write Eq. (B2) in the following way:

$$dx = -\gamma_x x dt + \sqrt{Q} dW_x \quad (\text{B7})$$

If Q were a constant, then the equation above would be a Langevin equation and x would represent an Ornstein-Uhlenbeck process. But for construction Q is slowly varying respect x (because $\tau = \frac{1}{\gamma_x} \ll T = \frac{1}{\mu}$). So it is possible to assume that locally in time Q is constant and so locally x has a Gaussian distribution, given a certain variance $\sigma_x^2 = \frac{Q}{2\gamma_x}$ determined by the value of Q :

$$p(x|\sigma_x^2) = N(0, \sigma_x^2) = \frac{1}{\sqrt{2\pi\sigma_x^2}} e^{-x^2/2\sigma_x^2} = \sqrt{\frac{\gamma_x}{\pi Q}} e^{-\gamma_x x^2/Q} \quad (\text{B8})$$

In order to obtain the total x PDF, the σ_x^2 PDF $f(\sigma_x^2)$ is needed. Thanks to its dependence with Q , it is possible to obtain it:

$$f(\sigma_x^2) = \Gamma_{\nu, \lambda}(\sigma_x^2) = \frac{\lambda^\nu \sigma_x^{2(\nu-1)} e^{-\lambda \sigma_x^2}}{\Gamma(\nu)} = \left(\frac{2\mu\gamma_x}{\beta_u^2} \right)^\nu \frac{\sigma_x^{2(\nu-1)}}{\Gamma(\nu)} e^{-\frac{2\mu\gamma_x}{\beta_u^2} \sigma_x^2} \quad (\text{B9})$$

605 where $\lambda = \frac{2\mu\gamma_x}{\beta_u^2}$. It follows that:

$$\begin{aligned} p_\nu(x) &= \int_0^\infty f(\sigma_x^2) p(x|\sigma_x^2) d\sigma_x^2 \\ &= \frac{\lambda^\nu}{\Gamma(\nu)\sqrt{2\pi}} \int_0^\infty \sigma_x^{2(\nu-1)} e^{-\lambda \sigma_x^2} \frac{1}{\sqrt{\sigma_x^2}} e^{-x^2/2\sigma_x^2} d(\sigma_x^2) \\ &= \frac{\lambda^\nu}{\Gamma(\nu)\sqrt{2\pi}} \int_0^\infty z^{\nu-3/2} e^{-\lambda z - x^2/2z} dz \end{aligned} \quad (\text{B10})$$

that, for $\nu = 2$, it gives the following result (Gradshteyn and Ryzhik (2014) page 369, Eq. (3.471.16), called, in this article, Exp-Lin PDF:

$$\begin{aligned} p_{\nu=2}(x) &= -\frac{\lambda^2}{\sqrt{2}} \frac{\partial}{\partial \lambda} \left(\lambda^{-1/2} e^{-\sqrt{2\lambda}|x|} \right) \\ &= \frac{\sqrt{2\lambda} e^{-\sqrt{2\lambda}|x|} (\sqrt{2\lambda}|x| + 1)}{4} \\ &= \frac{\sqrt{\mu\gamma_x}}{2\beta_u} e^{-2\sqrt{\mu\gamma_x}|x|/\beta_u} \left(\frac{2\sqrt{\mu\gamma_x}}{\beta_u} |x| + 1 \right) \end{aligned} \quad (\text{B11})$$

610 while, for $\nu = 1/2$ (Gradshteyn and Ryzhik (2014) page 370, Eq. 3.478.4):

$$\begin{aligned} p_{\nu=1/2}(x) &= \frac{\sqrt{2\lambda}}{\pi} K_0 \left(\sqrt{2\lambda}|x| \right) \\ &= \frac{2\sqrt{\mu\gamma_x}}{\pi\beta_u} K_0 \left(\frac{2\sqrt{\mu\gamma_x}}{\beta_u} |x| \right) \end{aligned} \quad (\text{B12})$$

where $K_0(z)$ is the modified Bessel function of the second kind of 0 order. A summary of the results of this Appendix for the cases ν , 2 and 1/2 degrees of freedom (dof) is given in Table B1. Finally, Eq. (B3) does not present any analytic solution.

Appendix C: The Ekman system of the deterministic model - the coefficients linear relation with the wind speed

615 Defining the energy $E = \tilde{h} \frac{|\mathbf{u}_o|^2}{2}$ for the DET model in Eq. (1) and supposing, as a first approximation, the tidal signal negligible $\mathbf{u}_o \simeq \mathbf{u}_E$ (daily time scale), a slowlyly varying sea surface layer depth $\partial_t \tilde{h} = 0$ and neglecting the eddy depletion term $\mathbf{F} = \rho_a c_a |\mathbf{u}_a| \mathbf{u}_a$, so the energy variation in time can be expressed as follows:

$$\partial_t E = \tilde{h} \partial_t \frac{|\mathbf{u}_E|^2}{2} = \tilde{h} (u_E \partial_t u_E + v_E \partial_t v_E) = -C_B |\mathbf{u}_E|^3 + u_E F_u + v_E F_v \quad (\text{C1})$$

dof	$p_Q(Q)$	$E[Q]$	$p(x)$
ν	$\left(\frac{\mu}{\beta_u^2}\right)^\nu \frac{Q^{\nu-1}}{\Gamma(\nu)} e^{-\mu Q/\beta_u^2}$	$\nu\beta^2/\mu$	$\frac{(2\mu\gamma_x/\beta^2)^\nu}{\Gamma(\nu)\sqrt{2\pi}} \int_0^\infty z^{\nu-3/2} e^{-\frac{2\mu\gamma_x}{\beta^2}z - \frac{x^2}{2z}} dz$
2	$\left(\frac{\mu}{\beta_u^2}\right)^2 Q e^{-\mu Q/\beta_u^2}$	$2\beta^2/\mu$	$\frac{\sqrt{\mu\gamma_x}}{2\beta_u} e^{-2\sqrt{\mu\gamma_x} x /\beta_u} \left(\frac{2\sqrt{\mu\gamma_x}}{\beta_u} x + 1\right)$
$\frac{1}{2}$	$\sqrt{\frac{\mu}{\pi\beta_u^2}} \frac{e^{-\mu Q/\beta_u^2}}{\sqrt{Q}}$	$\beta^2/2\mu$	$\frac{2\sqrt{\mu\gamma_x}}{\pi\beta_u} K_0\left(\frac{2\sqrt{\mu\gamma_x}}{\beta_u} x \right)$

Table B1. Analytical PDFs and means of the variable Q and analytical PDFs of the variable x for ν , 2 and 1/2 degrees of freedom.

from which:

$$620 \quad \frac{C_B}{\tilde{h}} = \frac{u_E F_u + v_E F_v}{\tilde{h}|\mathbf{u}_E|^3} - \frac{\partial_t |\mathbf{u}_E|^2}{2|\mathbf{u}_E|^3} \quad (\text{C2})$$

Discretizing Eq. (C1) and Eq. (C2) around the time step t_i (where $\Delta t = t_{i+1} - t_i$) and using the centered scheme for the temporal derivative $\partial_t |\mathbf{u}_E|^2 = \frac{|\mathbf{u}_E(t_{i+1})|^2 - |\mathbf{u}_E(t_{i-1})|^2}{2\Delta t}$, it results:

$$\tilde{h} \frac{|\mathbf{u}_E(t_{i+1})|^2 - |\mathbf{u}_E(t_{i-1})|^2}{4} = -C_B(t_i) |\mathbf{u}_E(t_i)|^3 \Delta t + [u_E(t_i) F_u(t_i) + v_E(t_i) F_v(t_i)] \Delta t \quad (\text{C3})$$

and:

$$625 \quad \begin{aligned} \frac{C_B(t_i)}{\tilde{h}} &= \frac{u_E(t_i) F_u(t_i) + v_E(t_i) F_v(t_i)}{\tilde{h}|\mathbf{u}_E(t_i)|^3} - \frac{|\mathbf{u}_E(t_{i+1})|^2 - |\mathbf{u}_E(t_{i-1})|^2}{4|\mathbf{u}_E(t_i)|^3 \Delta t} \\ &= \frac{\alpha(t_i)}{\tilde{h}} - \zeta(t_i) \end{aligned} \quad (\text{C4})$$

where $\alpha(t_i) = \frac{u_E(t_i) F_u(t_i) + v_E(t_i) F_v(t_i)}{|\mathbf{u}_E(t_i)|^3}$, $\zeta(t_i) = \frac{|\mathbf{u}_E(t_i + \Delta t)|^2 - |\mathbf{u}_E(t_i - \Delta t)|^2}{4|\mathbf{u}_E(t_i)|^3 \Delta t}$ and $\tilde{h}$ has a low dependence on the time. Approximating the daily-averaged observed sea surface current as the daily deterministic Ekman sea surface current, it is possible to calculate the coefficients $\alpha(t_i)$ and $\zeta(t_i)$ directly from the observations and the daily-averaged wind forcing.

Eq. (1) results in:

$$630 \quad \begin{pmatrix} \partial_t u_E \\ \partial_t v_E \end{pmatrix} = \begin{pmatrix} -\left(\frac{\alpha}{h} - \zeta\right) |\mathbf{u}_E| & f \\ -f & -\left(\frac{\alpha}{h} - \zeta\right) |\mathbf{u}_E| \end{pmatrix} \begin{pmatrix} u_E \\ v_E \end{pmatrix} + \begin{pmatrix} \frac{F_u}{h} \\ \frac{F_v}{h} \end{pmatrix} \quad (\text{C5})$$

$$\begin{cases} \partial_t u_E - \zeta |\mathbf{u}_E| u_E - f v_E = \frac{F_u - \alpha |\mathbf{u}_E| u_E}{h} \\ \partial_t v_E - \zeta |\mathbf{u}_E| v_E + f u_E = \frac{F_v - \alpha |\mathbf{u}_E| v_E}{h} \end{cases} \quad (\text{C6})$$

From which it is possible to obtain the slowly varying $\tilde{h}$:

$$\begin{cases} \tilde{h} = \frac{F_u - \alpha |\mathbf{u}_E| u_E}{\partial_t u_E - \zeta |\mathbf{u}_E| u_E - f v_E} \\ \tilde{h} = \frac{F_v - \alpha |\mathbf{u}_E| v_E}{\partial_t v_E - \zeta |\mathbf{u}_E| v_E + f u_E} \end{cases} \quad (C7)$$

635 The discretization gives:

$$\begin{cases} \tilde{h}(t_i) = \frac{F_u(t_i) - \alpha(t_i) |\mathbf{u}_E(t_i)| u_E(t_i)}{[u_E(t_i + \Delta t) - u_E(t_i - \Delta t)] / 2\Delta t - \zeta(t_i) |\mathbf{u}_E(t_i)| u_E(t_i) - f v_D(t_i)} \\ \tilde{h}(t_i) = \frac{F_v(t_i) - \alpha(t_i) |\mathbf{u}_E(t_i)| v_E(t_i)}{[v_E(t_i + \Delta t) - v_E(t_i - \Delta t)] / 2\Delta t - \zeta(t_i) |\mathbf{u}_E(t_i)| v_E(t_i) + f u_E(t_i)} \end{cases} \quad (C8)$$

The two expressions in Eq. (C8) should give a unique value for $\tilde{h}$ in function of the time directly from the daily-averaged observed currents and the daily-averaged wind forcing. In practice, they almost agree and we calculated $\tilde{h}(t_i)$ as the average of these two expressions. From $\tilde{h}(t_i)$ it is possible to obtain also $C_B(t_i)$:

$$640 \quad C_B(t_i) = \alpha(t_i) - \zeta(t_i) \cdot \tilde{h}(t_i) \quad (C9)$$

Having the $\tilde{h}$, the C_B and the wind speed time series, it is possible to calculate the mean values of $\tilde{h}$ and C_B in function of the wind speed, obtaining a linear fit after deleting the values exceeding one standard deviation (Fig. C1).




Figure C1. Dependence on the wind speed of the model coefficients C_B and $\tilde{h}$, calculated accordingly to Eq. (C9) and to the mean of Eq.s (C8) respectively, and their linear regressions.

Appendix D: On the Ekman angle standard deviation

In this study the Ekman angle is defined as the angle between the daily wind and the daily sea surface current, it is positive if the sea surface current is on the right with respect to the wind direction. For simplicity it is here assumed that the wind does

645

not contribute with any error and is perfectly known. It follows that the standard deviation of the Ekman angle coincides with the standard deviation of the daily sea surface current angle. Under this assumption, the given Ekman angle standard deviation is an underestimation of its real error. So, in order to obtain this estimation of the Ekman angle standard deviation, starting from the modelled (HFR measured) sea surface current components with 5 min (30 min) time resolution and their stochastic variability (measure accuracy), two points must be followed: (i) the daily average transformation and (ii) the polar coordinates transformation.

Let's take into consideration the sea surface currents time series of a single day $x = \{u_1, \dots, u_n, v_1, \dots, v_n\}$, where $n = 288$ for the 5 min time resolution modelled sea surface currents and $n = 48$ for the 30 min HFR observed sea surface currents, with their $2n \times 2n$ symmetric covariance matrix:

$$\Sigma^x = \begin{pmatrix} \sigma_{u_1}^2 & \cdots & \sigma_{u_1 u_n} & \sigma_{u_1 v_1} & \cdots & \sigma_{u_1 v_n} \\ \vdots & \ddots & \vdots & \vdots & & \vdots \\ \sigma_{u_n u_1} & \cdots & \sigma_{u_n}^2 & \sigma_{u_n v_1} & \cdots & \sigma_{u_n v_n} \\ \sigma_{v_1 u_1} & \cdots & \sigma_{v_1 u_n} & \sigma_{v_1}^2 & \cdots & \sigma_{v_1 v_n} \\ \vdots & & \vdots & \vdots & \ddots & \vdots \\ \sigma_{v_n u_1} & \cdots & \sigma_{v_n u_n} & \sigma_{v_n v_1} & \cdots & \sigma_{v_n}^2 \end{pmatrix} = \begin{pmatrix} \Sigma_{uu}^x & \Sigma_{uv}^x \\ \Sigma_{uv}^x & \Sigma_{vv}^x \end{pmatrix} \quad (D1)$$

Here the sea surface currents are the ensemble averaged values for each time and the covariance matrix values are the relative ensemble variances and covariances, while for the observations the sea surface currents are simply the independent HFR measured sea surface currents (each measurement is independent from the others) with their HFR accuracy and 0 covariances (so Σ^x reduces to a diagonal matrix).

Calling $f = \begin{pmatrix} \bar{u} \\ \bar{v} \end{pmatrix}$ the daily average sea surface currents for a single day, its linear transformation (here the transformation matrix is called A) can be written as:

$$f = A \cdot x \quad (D2)$$

$$\begin{pmatrix} \bar{u} \\ \bar{v} \end{pmatrix} = \frac{1}{n} \begin{pmatrix} \underbrace{1 \cdots 1}_n & \underbrace{0 \cdots 0}_n \\ \underbrace{0 \cdots 0}_n & \underbrace{1 \cdots 1}_n \end{pmatrix} \begin{pmatrix} u_1 \\ \vdots \\ u_n \\ v_1 \\ \vdots \\ v_n \end{pmatrix} \quad (D3)$$

So, the covariance matrix of the transformed variable $\Sigma^f = \begin{pmatrix} \sigma_{\bar{u}}^2 & \sigma_{\bar{u}\bar{v}} \\ \sigma_{\bar{u}\bar{v}} & \sigma_{\bar{v}}^2 \end{pmatrix}$ can be obtained in the following way:

$$\begin{aligned}
\Sigma^f &= A \cdot \Sigma^x \cdot A^T \\
&= \frac{1}{n^2} \begin{pmatrix} 1 & \cdots & 1 & 0 & \cdots & 0 \\ 0 & \cdots & 0 & 1 & \cdots & 1 \end{pmatrix} \begin{pmatrix} \sigma_{u_1}^2 & \cdots & \sigma_{u_1 u_n} & \sigma_{u_1 v_1} & \cdots & \sigma_{u_1 v_n} \\ \vdots & \ddots & \vdots & \vdots & & \vdots \\ \sigma_{u_n u_1} & \cdots & \sigma_{u_n}^2 & \sigma_{u_n v_1} & \cdots & \sigma_{u_n v_n} \\ \sigma_{v_1 u_1} & \cdots & \sigma_{v_1 u_n} & \sigma_{v_1}^2 & \cdots & \sigma_{v_1 v_n} \\ \vdots & & \vdots & \vdots & \ddots & \vdots \\ \sigma_{v_n u_1} & \cdots & \sigma_{v_n u_n} & \sigma_{v_n v_1} & \cdots & \sigma_{v_n}^2 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \vdots & \vdots \\ 1 & 0 \\ 0 & 1 \\ \vdots & \vdots \\ 0 & 1 \end{pmatrix} \\
&= \frac{1}{n^2} \begin{pmatrix} \sigma_{u_1}^2 + \cdots + \sigma_{u_1 u_n} + \cdots + \sigma_{u_n u_1} + \cdots + \sigma_n^2 & \sigma_{u_1 v_1} + \cdots + \sigma_{u_1 v_n} + \cdots + \sigma_{u_n v_1} + \cdots + \sigma_{u_n v_n} \\ \sigma_{v_1 u_1} + \cdots + \sigma_{v_1 u_n} + \cdots + \sigma_{v_n u_1} + \cdots + \sigma_{v_n u_n} & \sigma_{v_1}^2 + \cdots + \sigma_{v_1 v_n} + \cdots + \sigma_{v_n v_1} + \cdots + \sigma_{v_n}^2 \end{pmatrix} \\
&= \frac{1}{n^2} \begin{pmatrix} \text{sum}(\Sigma_{uu}^x) & \text{sum}(\Sigma_{uv}^x) \\ \text{sum}(\Sigma_{uv}^x) & \text{sum}(\Sigma_{vv}^x) \end{pmatrix} \tag{D4}
\end{aligned}$$

670 Once obtained the daily sea surface current components $\begin{pmatrix} \bar{u} \\ \bar{v} \end{pmatrix}$ and their covariance matrix $\begin{pmatrix} \sigma_{\bar{u}}^2 & \sigma_{\bar{u}\bar{v}} \\ \sigma_{\bar{u}\bar{v}} & \sigma_{\bar{v}}^2 \end{pmatrix}$, it is possible to change from cartesian to polar coordinates:

$$\begin{cases} r = \sqrt{\bar{u}^2 + \bar{v}^2} \\ \theta = \arctan\left(\frac{\bar{v}}{\bar{u}}\right) \end{cases} \tag{D5}$$

The transformation now is not linear and one must resort to the jacobian matrix:

$$J = \begin{pmatrix} \frac{\partial r}{\partial \bar{u}} & \frac{\partial r}{\partial \bar{v}} \\ \frac{\partial \theta}{\partial \bar{u}} & \frac{\partial \theta}{\partial \bar{v}} \end{pmatrix} = \begin{pmatrix} \frac{\bar{u}}{\sqrt{\bar{u}^2 + \bar{v}^2}} & \frac{\bar{v}}{\sqrt{\bar{u}^2 + \bar{v}^2}} \\ -\frac{\bar{v}}{\bar{u}^2 + \bar{v}^2} & \frac{\bar{u}}{\bar{u}^2 + \bar{v}^2} \end{pmatrix} \tag{D6}$$

675 to obtain the new covariance matrix (θ and σ_θ are in radians):

$$\begin{aligned}
\begin{pmatrix} \sigma_r^2 & \sigma_{r\theta} \\ \sigma_{r\theta} & \sigma_\theta^2 \end{pmatrix} &= J \begin{pmatrix} \sigma_{\bar{u}}^2 & \sigma_{\bar{u}\bar{v}} \\ \sigma_{\bar{u}\bar{v}} & \sigma_{\bar{v}}^2 \end{pmatrix} J^T \\
&= \begin{pmatrix} \frac{\bar{u}}{\sqrt{\bar{u}^2 + \bar{v}^2}} & \frac{\bar{v}}{\sqrt{\bar{u}^2 + \bar{v}^2}} \\ -\frac{\bar{v}}{\bar{u}^2 + \bar{v}^2} & \frac{\bar{u}}{\bar{u}^2 + \bar{v}^2} \end{pmatrix} \begin{pmatrix} \sigma_{\bar{u}}^2 & \sigma_{\bar{u}\bar{v}} \\ \sigma_{\bar{u}\bar{v}} & \sigma_{\bar{v}}^2 \end{pmatrix} \begin{pmatrix} \frac{\bar{u}}{\sqrt{\bar{u}^2 + \bar{v}^2}} & -\frac{\bar{v}}{\bar{u}^2 + \bar{v}^2} \\ \frac{\bar{v}}{\sqrt{\bar{u}^2 + \bar{v}^2}} & \frac{\bar{u}}{\bar{u}^2 + \bar{v}^2} \end{pmatrix} \\
&= \begin{pmatrix} \frac{\bar{u}^2 \sigma_{\bar{u}}^2 + 2\bar{u}\bar{v}\sigma_{\bar{u}\bar{v}} + \bar{v}^2 \sigma_{\bar{v}}^2}{\bar{u}^2 + \bar{v}^2} & \frac{-\bar{u}\bar{v}\sigma_{\bar{u}}^2 + (\bar{u}^2 - \bar{v}^2)^2 \sigma_{\bar{u}\bar{v}} + \bar{u}\bar{v}\sigma_{\bar{v}}^2}{(\bar{u}^2 + \bar{v}^2)^{3/2}} \\ \frac{-\bar{u}\bar{v}\sigma_{\bar{u}}^2 + (\bar{u}^2 - \bar{v}^2)^2 \sigma_{\bar{u}\bar{v}} + \bar{u}\bar{v}\sigma_{\bar{v}}^2}{(\bar{u}^2 + \bar{v}^2)^{3/2}} & \frac{\bar{v}^2 \sigma_{\bar{u}}^2 - 2\bar{u}\bar{v}\sigma_{\bar{u}\bar{v}} + \bar{u}^2 \sigma_{\bar{v}}^2}{(\bar{u}^2 + \bar{v}^2)^2} \end{pmatrix}. \tag{D7}
\end{aligned}$$

The Ekman angle standard deviation is then taken as follows:

$$\sigma_{\theta_E} = \min(\sigma_{\theta}, \pi). \quad (\text{D8})$$

680 *Author contributions.* SF performed the major part of the research and the writing of the manuscript. LU and AW contributed to the research and the writing.

Competing interests. The contact author has declared that none of the authors has any competing interests.

685 *Acknowledgements.* The forecasts, analyses, and related services are based on data and products of the Regional Center for Environmental Modelling (CRMA), which is a sector of the Friuli Venezia Giulia Environmental Agency (ARPA FVG) ITALY (<https://www.arpa.fvg.it>, last access: 29 October 2024). The Piran HFR station has been operated by the Slovenian Environment Agency since July 2022. Between 2015 and July 2022 it was operated by the National Institute of Biology. ChatGPT was used for text refinement of Sect. 5.

References

- Alberti, T., Consolini, G., De Michelis, P., Laurenza, M., and Marcucci, M. F.: On fast and slow Earth's magnetospheric dynamics during geomagnetic storms: a stochastic Langevin approach, *Journal of Space Weather and Space Climate*, 8, A56, 2018.
- Baldovin, M., Puglisi, A., and Vulpiani, A.: Langevin equation in systems with also negative temperatures, *Journal of Statistical Mechanics: Theory and Experiment*, 2018, 043 207, 2018.
- Beck, C. and Cohen, E. G.: Superstatistics, *Physica A: Statistical mechanics and its applications*, 322, 267–275, 2003.
- Beck, C., Cohen, E. G. D., and Swinney, H. L.: From time series to superstatistics, *Phys. Rev. E*, 72, 056 133, <https://doi.org/10.1103/PhysRevE.72.056133>, 2005.
- Berglund, N. and Gentz, B.: Metastability in simple climate models: pathwise analysis of slowly driven Langevin equations, *Stochastics and Dynamics*, 2, 327–356, 2002.
- Brillinger, D. R. and Stewart, B. S.: Stochastic modeling of particle movement with application to marine biology and oceanography, *Journal of Statistical Planning and Inference*, 140, 3597–3607, <https://doi.org/https://doi.org/10.1016/j.jspi.2010.04.026>, 2010.
- Corgnati, L., Mantovani, C., Novellino, A., Rubio, A., and Mader, J.: Recommendation Report 2 on improved common procedures for HFR QC analysis. JERICO-NEXT WP5-Data Management, Deliverable 5.14, Version 1.0., <https://doi.org/http://dx.doi.org/10.25607/OBP-944>, 2018.
- Cosoli, S., Gačić, M., and Mazzoldi, A.: Surface current variability and wind influence in the northeastern Adriatic Sea as observed from high-frequency (HF) radar measurements, *Continental Shelf Research*, 33, 1–13, 2012.
- Cosoli, S., Ličer, M., Vodopivec, M., and Malačič, V.: Surface circulation in the Gulf of Trieste (northern Adriatic Sea) from radar, model, and ADCP comparisons, *Journal of Geophysical Research: Oceans*, 118, 6183–6200, 2013.
- Einstein, A.: Über die von der molekularkinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssigkeiten suspendierten Teilchen, *Annalen der physik*, 4, 1905.
- Einstein, A.: Investigations on the Theory of the Brownian Movement, Courier Corporation, 1956.
- Flora, S., Ursella, L., and Wirth, A.: Superstatistical analysis of sea surface currents in the Gulf of Trieste, measured by high-frequency radar, and its relation to wind regimes using the maximum-entropy principle, *Nonlinear Processes in Geophysics*, 30, 515–525, 2023.
- Flora, S., Wirth, A., and Ursella, L.: Codes: Comparing an idealized deterministic-stochastic model (SUP model, version 1) of the tide-and-wind driven sea surface currents in the Gulf of Trieste to HF Radar observations, <https://doi.org/10.5281/zenodo.14562025>, 2024.
- Foreman, M.: Manual for Tidal Currents Analysis and Prediction, Pacific Marine Science Report 78-6, Institute of Ocean Sciences, Patricia Bay, Sidney, British Columbia, 70, 1978.
- Franzke, C., Majda, A. J., and Vanden-Eijnden, E.: Low-order stochastic mode reduction for a realistic barotropic model climate, *Journal of the atmospheric sciences*, 62, 1722–1745, 2005.
- Franzke, C. L., O’Kane, T. J., Berner, J., Williams, P. D., and Lucarini, V.: Stochastic climate theory and modeling, *Wiley Interdisciplinary Reviews: Climate Change*, 6, 63–78, 2015.
- Goglio, A. C.: Progetto NAUSICA, https://www.arpa.fvg.it/export/sites/default/tema/crma/pubblicazioni/docs_pubblicazioni/2018gen01_arpa_fvg_crma_nausicarap2018_001.pdf, 2018.
- Gradshteyn, I. S. and Ryzhik, I. M.: Table of integrals, series, and products, Academic press, 2014.
- Ham, L., Coomer, M. A., and Stumpf, M. P.: The chemical Langevin equation for biochemical systems in dynamic environments, *The Journal of Chemical Physics*, 157, 2022.

- Hasselmann, K.: Stochastic climate models part I. Theory, *tellus*, 28, 473–485, 1976.
- Kloeden, P. E. and Platen, E.: Numerical Solution of Stochastic Differential Equations, Springer, 1999.
- 725 Lacorata, G. and Vulpiani, A.: Fluctuation-Response Relation and modeling in systems with fast and slow dynamics, *Nonlinear Processes in Geophysics*, 14, 681–694, 2007.
- Langevin, P. et al.: Sur la théorie du mouvement brownien, *CR Acad. Sci. Paris*, 146, 530, 1908.
- Lemons, D. S. and Gythiel, A.: Paul Langevin's 1908 paper "On the Theory of Brownian Motion" [" Sur la thiorie du mouvement brownien," *CR Acad. Sci.(Paris)* 146, 530-533 (1908)], *American Journal of Physics*, 65, 1079–1081, 1997.
- 730 Lorente, P., Aguiar, E., Bondoni, M., Berta, M., Brandini, C., Cáceres-Euse, A., Capodici, F., Cianelli, D., Ciruolo, G., Corgnati, L., et al.: Coastal high-frequency radars in the Mediterranean–Part 1: Status of operations and a framework for future development, *Ocean Science*, 18, 761–795, 2022.
- Lorenz, E. N.: Predictability: A problem partly solved, in: *Proc. Seminar on predictability*, vol. 1, Reading, 1996.
- Malačič, V., Petelin, B., Gačić, M., Artegiani, A., and Orlić, M.: *Regional Studies*, pp. 167–216, Springer Netherlands, Dordrecht, ISBN 978-94-015-9819-4, https://doi.org/10.1007/978-94-015-9819-4_6, 2001.
- 735 OGS, NIB, ARSO, and ARPAFVG: HFR-NAdr (High Frequency Radar NAdr network), <https://doi.org/https://doi.org/10.57762/8RRE-0Z07>, 2023.
- Palmer, T.: Stochastic weather and climate models, *Nature Reviews Physics*, 1, 463–471, 2019.
- Pawlowicz, R., Beardsley, B., and Lentz, S.: Classical tidal harmonic analysis including error estimates in MATLAB using T_TIDE, *Computers & geosciences*, 28, 929–937, 2002.
- 740 Poulain, P.-M. and Raicich, F.: *Forcings*, pp. 45–65, Springer Netherlands, Dordrecht, ISBN 978-94-015-9819-4, https://doi.org/10.1007/978-94-015-9819-4_2, 2001.
- Querín, S., Crise, A., Deponte, D., and Solidoro, C.: Numerical study of the role of wind forcing and freshwater buoyancy input on the circulation in a shallow embayment (Gulf of Trieste, Northern Adriatic Sea), *Journal of Geophysical Research: Oceans*, 111, 2006.
- 745 Querín, S., Cosoli, S., Gerin, R., Laurent, C., Malačič, V., Pristov, N., and Poulain, P.-M.: Multi-platform, high-resolution study of a complex coastal system: The TOSCA experiment in the Gulf of Trieste, *Journal of Marine Science and Engineering*, 9, 469, 2021.
- Reyes Suárez, N. C., Tirelli, V., Ursella, L., Ličer, M., Celio, M., and Cardin, V.: Multi-platform study of the extreme bloom of the barrel jellyfish *Rhizostoma pulmo* (Cnidaria: Scyphozoa) in the northernmost gulf of the Mediterranean Sea (Gulf of Trieste) in April 2021, *Ocean Science*, 18, 1321–1337, 2022.
- 750 Silveira, F. A. and Aarão Reis, F. D. A.: Langevin equations for competitive growth models, *Phys. Rev. E*, 85, 011601, <https://doi.org/10.1103/PhysRevE.85.011601>, 2012.
- Sura, P.: *Stochastic Models of Climate Extremes: Theory and Observations*, pp. 181–222, Springer Netherlands, Dordrecht, ISBN 978-94-007-4479-0, https://doi.org/10.1007/978-94-007-4479-0_7, 2013.
- van den Berk, J., Drijfhout, S., and Hazeleger, W.: Characterisation of Atlantic meridional overturning hysteresis using Langevin dynamics, *Earth System Dynamics*, 12, 69–81, 2021.
- 755 Wand, T., Wiedemann, T., Harren, J., and Kamps, O.: Estimating stable fixed points and Langevin potentials for financial dynamics, *Physical Review E*, 109, 024226, 2024.
- Wirth, A.: A fluctuation-dissipation relation for the ocean subject to turbulent atmospheric forcing, *Journal of Physical Oceanography*, 48, 831–843, 2018.

- 760 Wirth, A.: On fluctuating momentum exchange in idealised models of air–sea interaction, *Nonlinear Processes in Geophysics*, 26, 457–477, <https://doi.org/10.5194/npg-26-457-2019>, 2019.
- Zhai, X., Johnson, H. L., Marshall, D. P., and Wunsch, C.: On the Wind Power Input to the Ocean General Circulation, *Journal of Physical Oceanography*, 42, 1357 – 1365, <https://doi.org/10.1175/JPO-D-12-09.1>, 2012.